

U. S. Bureau of Commercial Fisheries

UNITED STATES DEPARTMENT OF COMMERCE

R. P. LAMONT, Secretary

BUREAU OF FISHERIES
HENRY O'MALLEY, Commissioner

Rare Bk

S4

11

A15

1928

pt. 2

REPORT

OF THE

**UNITED STATES
COMMISSIONER OF FISHERIES**

FOR THE FISCAL YEAR 1928

WITH

APPENDIXES

LIBRARY

Part II

N.O.A.A.
U.S. Dept of Commerce



**MARINE AND EARTH
SCIENCES LIBRARY**

OCT 16 1972

N.O.A.A.
U. S. Dept. of Commerce

UNITED STATES
GOVERNMENT PRINTING OFFICE
WASHINGTON : 1930

For sale by the Superintendent of Documents, Washington, D. C. Price \$1.65

1 '72 5515

National Oceanic and Atmospheric Administration

Report of the United States Commissioner of Fisheries

ERRATA NOTICE

One or more conditions of the original document may affect the quality of the image, such as:

Discolored pages
Faded or light ink
Binding intrudes into the text

This has been a co-operative project between the NOAA Central Library and the National Marine Fisheries Service (NOAA Fisheries). To view the original document, please contact the NOAA Central Library in Silver Spring, MD at (301) 713-2607 x124 or www.reference@nodc.noaa.gov.

LASON
Imaging Contractor
12200 Kiln Court
Beltsville, MD 20704-1387
November 19, 2004

C O N T E N T S

	Page
CHECK LIST OF THE FISHES AND FISHLIKE VERTEBRATES OF NORTH AND MIDDLE AMERICA NORTH OF THE NORTHERN BOUNDARY OF VENEZUELA AND COLOMBIA. By David Starr Jordan, Barton Warren Evermann, and Howard Walton Clark. (Document No. 1055. Issued February, 1930)-----	1-670 III

DEPARTMENT OF COMMERCE
BUREAU OF FISHERIES

HEADQUARTERS STAFF, 1927-28

Commissioner

HENRY O'MALLEY

Deputy Commissioner.—LEWIS RADCLIFFE.

Assistants in Charge of Divisions:

Office.—IRVING H. DUNLAF.

Fish Culture.—GLEN C. LEACH.

Inquiry Respecting Food Fishes.—ELMER HIGGINS.

Fishery Industries.—OSCAR E. SETTE.

Alaska Service.—WARD T. BOWER.

Architect and Engineer.—GEORGE A. SCHNEIDER.

Accountant.—CHARLES W. SCUDDER.

Superintendent Central Station and Aquaria.—L. G. HARRON.

Editor.—ANNA B. BROWN.

CHECK LIST OF THE FISHES AND FISHLIKE VERTEBRATES OF NORTH AND MIDDLE AMERICA NORTH OF THE NORTHERN BOUNDARY OF VENEZUELA AND COLOMBIA¹

By

DAVID STARR JORDAN, BARTON WARREN EVERMANN, and
HOWARD WALTON CLARK

INTRODUCTION

The present memoir comprises a list of all the fishes and fishlike vertebrates known to occur in the waters of north and middle America; more specifically all of continental America north of the Isthmus of Panama, and the outlying islands including the West Indies, the Greater and the Lesser Antilles, Greenland, Iceland, and the islands off the Pacific coast of Central America, Mexico, the United States, and Alaska. The salt-water species on the northern coasts of Colombia and Venezuela have been included. Those of the Galapagos, the Sea of Okhotsk, and the west coast of Kamchatka are not included. Oceanic species in general within the 1,000-fathom curve have been admitted, although it is not always easy to draw the line.

The majority of the species included will be found described in Fishes of North and Middle America, a 4-volume work by Jordan and Evermann published in 1896 to 1900 as Bulletin 47 of the United States National Museum. No other general work restricted to this continent and containing descriptions of all known species has been written. On that work was based the Check List of 1896 by Jordan and Evermann, of which the present Check List may be regarded as a revision.

In the present list are given, in as natural sequence as our present knowledge permits, (1) the names of all species and subspecies that the authors admit as valid, (2) as many of the vernacular or common names for each species as have been readily obtainable, (3) the known geographic distribution of each species, (4) the reference to the original description of the species, and (5) the reference to every real synonym. If a species has been described as new but once, there will, of course, be but one reference (as, for example, *Photocorynus spiniceps* on p. 509); if it has been described as new six times there will be six references (as, for example, *Eupomotis gibbosus* on p. 302). Thus, this work is really a check list of all the scientific names that have ever been applied to any American fish.

As 4,139 species and subspecies are admitted as valid, and as there are more than 3,000 real synonyms, it appears that each American fish has been described as new about two times on an average.

The first attempt at a Synopsis of the Fishes of North America was published by David Humphreys Storer in the Memoirs of the Ameri-

¹ Appendix X to the Report of the U. S. Commissioner of Fisheries for 1928. B. F. Doc. 1055.

can Academy of Arts and Sciences of Boston in 1846. That work enumerates 739 species distributed in 221 genera. As the fishes of the Rocky Mountain region, Mexico, the Pacific coast, and the West Indies were then scarcely known, this list was very incomplete and the nomenclature naturally was imperfect.

After an elaborate survey of the Pacific coast fish fauna in 1880, David Starr Jordan and the late Charles Henry Gilbert published a Synopsis of the Fishes of North America (1882), including all species known north of the Mexican boundary. The 1,340 species names in that synopsis were arranged in 487 genera; a few other species appeared in the appendix.

Jordan's Catalogue of the Fishes Known to Inhabit the Waters of North America North of the Tropic of Cancer (1887) was based on this synopsis and includes a number of additional species.

In the present memoir we have followed implicitly the rules of nomenclature laid down by the International Commission for Zoological Nomenclature, of which the senior author has been a member since 1908. One single case now under discussion (*Carcharias*) has not been disposed of fully as yet, and the view recommended by Jordan is adopted here tentatively. An important rule set forth by Jordan and Evermann in 1895 has not been acted upon. This rule reads: "We regard all generic names as different unless actually spelled alike." In accordance with decisions of the commission, another rule, which we have followed, is this: "No change is to be made in current nomenclature unless the case is fairly clear; as, for example, a matter of priority." In general we have followed current practice rather than accept questionable changes. One doubtful opinion should not be exchanged for another. The original orthography of all names (misprints aside) has been retained. Some supposed misprints (*Cypsilurus*, *Pachynathus*, etc.) have been referred to the International Commission, whose verdicts have been followed.

In accordance with the decision of the International Commission, we have disregarded (unless revived by later authors) the polynomial generic names of Gronow, Klein, Browne, and other able early writers, who failed to adopt the Linnæan code simply because they had not heard of it. This decision rests not on consideration of justice but upon that of permanence.

The most serious problem in taxonomy is that of the assignment of generic types to the genera of authors who had no conception of types. The "rule of the first reviser" was adopted by the commission at Boston in 1908, and the present writers have endeavored to follow it.

By far the most complete work on North American fishes is that by Jordan and Evermann, mentioned above—the Fishes of North and Middle America. In that work the authors recognized 1,113 genera, 3,263 species, and 133 subspecies—a total of 3,396 species and subspecies. The check list of Fishes and Fishlike Vertebrates of North and Middle America, by Jordan and Evermann, based upon the above and published in 1896, enumerates 3,100 species arranged in 1,053 genera, these being about double those in the Jordan catalogue of 1887, the additions being mainly from Mexico, the West Indies, and the deep seas.

The late modern period in systematic ichthyology is from 1880 to the present time and is especially signalized by the explorations of

the deep sea, of most of the continents, and of fossil beds. The unexampled explorations of Eigenmann in South America belong to this period, as also the records of Boulenger of the fishes of South Africa and the varied researches of Regan in the rich material of the British Museum. In this period the faunas of Australia, Japan, and Siberia have been studied critically and successfully. Our knowledge of fossil fishes is assuming definite form through the work of Woodward, Zittel, Hay, Errol Ivor White, and others who have given vertebrate paleontology, as related to ichthyology, a constructive turn.

We may again emphasize the distinction between the ancient and modern periods in animal taxonomy. Beginning with 1758, the date of Linnæus's *Systema Naturæ*, for 60 years a genus was usually regarded merely as a convenient pigeonhole into which species were thrown. With Cuvier's *Règne Animal* (1817) came the conception of arrangement "according to organization," or comparative anatomy. With Darwin's *Origin of Species*, 40 years later (1859), zoologists began, though slowly at first, to realize that a classification must be more than an inventory; that its basis must be genetic, and that the problems involved in natural grouping are vastly more complex than even great morphologists like Cuvier and Agassiz had realized. To paraphrase a sentence of Elliott Coues, "genera and species are but larger and smaller twigs of a tree, which we try to arrange as nearly as possible in accordance with nature's ramifications." This view, applied to taxonomy, involves the extensive subdivision of accepted genera, many of which were in the highest degree unnatural. Subdivision as practiced in the present paper may go too far in some cases, but our material must be analyzed first before we can group its parts in natural synthesis. No system of naming can progress beyond the knowledge on which it rests. We may emphasize here a statement attributed to Aristotle that "one should never expect a degree of accuracy which the subject matter does not permit."

Each tendency in evolution, large or small, diverging from the main current, should properly stand as a separate genus.

The problem of subspecies among fishes is especially incomplete and confused. Most of the marine species are clearly definable and not often subject to intergradations. Among the fresh-water forms, all sorts of modifications occur. A species, in general, is a definable group of related individuals that have run the gauntlet of life and have endured. Among our river fishes we find species of different types. There are numerous cases of species of compact though often wide distribution, in which all individuals are, to all appearances, similar. Examples of this type are the common perch (*Perca fluviatilis*), the common sucker (*Catostomus commersonii*), and the minnows (*Ericymba buccata* and *Exoglossum maxillingua*). Other forms, which we have elsewhere called geminate or twin species, may be distinct enough, having a few tangible characters of difference in spite of general resemblance. These, for the most part, are set apart from each other by barriers, their distinctness proportional to the efficiency of the barrier. Their origin may be due to the crossing of these barriers by individuals from the mass of their kind, thus acquiring a different environment with a different type of selection. Illustrations of geminate species set off by isolation may be found in abundance, as among the trout of western America, and other species,

especially in the Panama region, where many geminate species are found on each side of the Isthmus, which has split its species from their mates since Miocene times. The Miocene fossil fishes of California are largely geminate in origin, twins or parents of species now appearing on the coast. In most cases such forms are clearly distinct but to all appearance ancestral, the existing distinctions apparently due to varying incidents of selection in varying environment. Geminate species rarely inhabit the same region, though exceptions are known, attributable to reinvasion.

In many cases the distinctions between geminate forms are not complete, intergradations occurring along the line of the barriers or at different sections in climatic range. Types thus connected are known as subspecies. Subspecies are often nascent species; but in most cases a large amount of material is necessary before subspecies can be separated from geminate species. Among fresh-water fishes, each river basin shows some features of distinction, large or small, as does each oceanic island with regard to marine fishes. In most families of American river fishes studies of species have not been extensive enough to establish facts of intergradation. While in birds, reptiles, and mammals the subspecies are usually well defined and named, among the fishes this is still largely a matter of guess-work. Many authors base the recognition of subspecies not on actual intergradation but on guesses based on degrees of probability, a practice we have not followed. It has seemed to us better to regard related forms as distinct species until intergradation has been shown.

We find also among river fishes, and sometimes among marine forms (as in *Lutianus*), occasional individuals that to all appearance are hybrids. These are marked by a distinct blending of specific traits. They are extremely rare, however, often known only from a single example. A few hybrids are known or suspected among certain minnows, sunfishes, and darters. The best-known hybrids thus far recorded in America occur among the Centrarchidæ, or sunfishes, in which group they have been studied critically by Mr. Hubbs. Two common species (*Apomotis cyanellus* and *Eupomotis gibbosus*) figure particularly as probable parents. Hybrids may give rise to distinct species, but no example of this is known among our fishes.

Among fishes there are also occasional monstrosities, freaks, or "mutations," in which certain traits may be absent or exaggerated. Whether such qualities may be inherited is not known. If so, it is likely that they would soon be swamped by breeding with the mass unless especially protected by segregation. It does not appear that any species of fishes have arisen by mutation or from hybridization, though the latter possibility is greater than the former.

We may observe that compilations such as the present paper should constitute an exercise in modesty. Such work can never be made absolutely complete nor free from errors. It deals with the observations and thoughts of great minds, as also with a degree of ignorance, carelessness, and perversity, by which names have been multiplied without corresponding extension of ideas. Synonymy (the use of different names for the same group) in general represents the lack of exact knowledge or else a failure (often unavoidable) to deal justly with the work of others. It is, therefore, as has been said by Doctor Coues, often "a burden and disgrace to science."

Only those who have had experience in cataloguing, in a similar way, the species in some large group of animals or plants can appreciate the difficulties involved. In the first place, in order to make the list complete by the inclusion of all valid names and all real synonyms it was necessary to make a careful search of all ichthyological literature. It is too much to expect that we have been entirely successful in this search; some names probably have been overlooked. The verification of the thousands of references as to the publication, volume, page, date, spelling, and type locality, was even more exacting and arduous; but most difficult of all were the multitude of taxonomic problems involved. In many cases existing knowledge is not adequate for final settlement. It is not improbable that some of our decisions as to the taxonomic relationships of certain species and the validity of others may not meet the approval of all of our coworkers, but that is inevitable where knowledge is incomplete.

The present senior author, as a lifelong teacher of biology, may be pardoned a personal reference. Himself a student of Agassiz and Gill of the earlier period, he has had the honor of being also a friend, to some extent a protégé, of Baird, Günther, Vaillant, Lütken, Hilgendorf, and Cope. In addition, he has numbered among his own students many of those active in systematic ichthyology during the late modern period. Among these are James Francis Abbott, Willis Stanley Blatchley, Charles Harvey Bollman, Alembert Winthrop Brayton, Charles Victor Burke, Howard Walton Clark*, Herbert Edson Copeland, Frank Cramer, George Bliss Culver, Bradley Moore Davis, Mary Cynthia Dickerson, Charles Lincoln Edwards, Carl H. Eigenmann, Rosa Smith Eigenmann, Barton Warren Evermann, Bert Fesler, Morton William Fordice, Henry Weed Fowler, Charles Henry Gilbert, James Zaccheus Gilbert, David Kop Goss, Ulysses Simpson Grant IV, Arthur White Greeley, Joseph Grinnell, Harold Hannibal, Edmund Heller, Albert William Christian Theodore Herre, Jennie Horning, Carl Leavitt Hubbs, Elizabeth G. Hughes, Oliver Peebles Jenkins, Eric Knight Jordan, Philip Henry Kirsch, Richard Crittenden McGregor, Ernest Alexander McGregor, Charles Leslie McKay, Seth Eugene Meek, Charles William Metz, Heraclio R. Montalban, George Sprague Myers, Edward William Nelson, Robert Newland, Masamitsu Oshima, Keinosuke Otaki, Charles J. Pierson, Cora D. Reeves, Robert Earl Richardson, Cloudsley Rutter*, Norman Bishop Scofield, Alvin Seale, Michitaro Sindo, Tage Skogsberg, Robert Evans Snodgrass, John Otterbein Snyder, Edwin Chapin Starks, Joseph Swain, Shigeho Tanaka, Lionel William Wiedey, Wilbur Wilson Thoburn, Joseph C. Thompson, William Francis Thompson, Deogracias Villamin Villadolid, Yojiro Wakiya, Frank Walter Weymouth, Thomas Marion Williams, and Albert Jefferson Woolman.*

Those marked with an asterisk have also been students of Doctor Evermann, as have also the following: Frederick Morton Chamberlain, Ulysses Orange Cox, Samuel Frederick Hildebrand, Chancey Juday, William Converse Kendall, Clarence Hamilton Kennedy, William J. Moenkhaus, John Treadwell Nichols, Lewis Radcliffe, Josiah Thomas Scovell, and Alfred Cleveland Weed.

Many of these have made important contributions to our knowledge of fishes.

The preparation of this Check List has extended over a period of more than 20 years. It was begun in 1906 by the senior authors,

who worked upon it from time to time, as their other duties permitted, until 1925, when they were joined by the junior author, who thereafter gave most of his time to the work until its completion. The senior authors also have devoted much of their time to it since 1925.

During the progress of the work the authors have endeavored to consult and utilize every record bearing on the fish fauna of America. In addition to this they have received invaluable assistance and suggestions from virtually every present-day student of American fishes, among whom special mention should be made of the late Dr. Charles Henry Gilbert and Prof. John Otterbein Snyder of Stanford University, Dr. William Converse Kendall of the United States Bureau of Fisheries, Dr. Carl Leavitt Hubbs of the University of Michigan, George Sprague Myers of Stanford University, Barton Appler Bean of the United States National Museum, William Francis Thompson of the International Fisheries Commission, Dr. Arthur W. Henn of the Carnegie Museum, John Treadwell Nichols and Dr. E. W. Gudger of the American Museum of Natural History, and many others. To all these we express our indebtedness and grateful appreciation. For clerical assistance of one kind or another, particularly in looking up and verifying references, we are greatly indebted to Miss Jessie E. Drayton, Harry J. Christoffers, and Henry D. Aller, of the United States Bureau of Fisheries. Without the assistance so generously given by these and others, this memoir could hardly have been completed.

DAVID STARR JORDAN.
BARTON WARREN EVERMANN.
HOWARD WALTON CLARK.

STANFORD UNIVERSITY,
CALIFORNIA ACADEMY OF SCIENCES,
January 4, 1928.

Class I. LEPTOCARDII. Lancelets

Order A. CIRROSTOMI. Cirrostomes

Family 1. BRANCHIOSTOMIDÆ. Lancelets

Genus 1. BRANCHIOSTOMA Costa.

Branchiostoma Costa, Cenni Zoologici Napol., 49, 1834 (*B. lubricum* Costa = *lanceolatus* Pallas).
Amphioxus Yarrell, Hist. Brit. Fishes, 1836, 468 (*Limax lanceolatus* Pallas).
Amphioxides ² Gill, Amer. Nat., XXIX, 1895, 458 (*Branchiostoma pelagicum* Günther).

1. *Branchiostoma virginiae* ³ Hubbs. *Virginia lancelet*.
Shallow waters, burying itself in the sand, from Chesapeake Bay to Florida; Beaufort, Cape Lookout, etc.
Branchiostoma virginiae Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 105, 8, July 6, 1922, Sewall's Point, Va.
2. *Branchiostoma floridæ* Hubbs. *Florida lancelet*.
Coasts of Florida; Tampa, Pensacola, Cape Romans, St. Martin's Reef, Snapper Banks, etc.
Branchiostoma floridæ Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 105, 7, July 6, 1922, Tampa Bay, Fla.
3. *Branchiostoma bermudæ* Hubbs. *Bermuda lancelet*.
Bermuda Islands.
Branchiostoma bermudæ Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 105, 9, July 6, 1922, Bermuda.
4. *Branchiostoma caribæum* ⁴ Sundevall. *West Indian lancelet*.
West Indies.
Branchiostoma caribæum Sundevall, Vet. Akad. Förhandl., XI, 1853, 12, St. Thomas.
5. *Branchiostoma californiense* Gill. *California lancelet*; *California amphioxus*.
Pacific coast of America from San Diego Bay southward to San Luis Gonzales Bay, Gulf of California; reported also from Panama. Abundant in shallow water, buried in the sand.
Branchiostoma californiense Gill, in Andrews, Studies Biol. Lab. Johns Hopkins Univ., V, 1893, 235, San Diego.

Family 2. EPIGONICHTHYIDÆ

Genus 2. ASYMMETRON Andrews.

Asymmetron Andrews, Studies Biol. Lab. Johns Hopkins Univ., V, 1893, 237 (*A. lucayanum* Andrews).

6. *Asymmetron lucayanum* Andrews. *Bahama lancelet*; *Bahama amphioxus*.
West Indies; Bemini and Nassau, Bahamas; Culebra Island; Humacao, Porto Rico; south to Brazil. Common.
Asymmetron lucayanum Andrews, Studies Biol. Lab. Johns Hopkins Univ., V, 1893, 237, Bemini, Bahamas.

¹ Based on larvae showing no buccal cirri, found in the open sea near Hawaii.

² Until the recent studies of Hubbs, this species was wrongly identified with the European form (*B. lanceolatum*), which, however, does not occur in America.

³ Represented on the coast of Brazil (Rio Janeiro, etc.) by *B. platei* Hubbs.

Class II. CYCLOSTOMI. Lampreys

Order B. HYPEROTRETA. Hagfishes

Family 3. MYXINIDÆ. Hagfishes

Genus 3. MYXINE Linnaeus.

Myxine Linnaeus, Syst. Nat., ed. X, 1758, 650 (*M. glutinosa* L.).*Gastrobranchus* Bloch, Ichth., XII, 1797, 51, tab. 413 (*G. cæcus* Bloch).*Murænoblenna* Lacépède, Hist. Nat. Poiss., V, 1803, 647 (*M. olivacea* Lacépède).*Anopsis Rafinesque*, Anal. Nat., 1815, 93 (*M. olivacea* Lacépède).

- 7.
- Myxine limosa*
- Girard.
- Hagfish; Borer; Sucker; Slimefish.*

Newfoundland, south to Cape Cod; Grand Manan Island, New Brunswick; replacing in the western Atlantic the European *M. glutinosa* L.; not abundant.*Myxine limosa* Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 224, Grand Manan.

Family 4. EPTATRETIDÆ. Borers

Genus 4. POLISTOTREMA Gill.

Polistotrema Gill, Proc. U. S. Nat. Mus., IV, 1881, 30 (*Gastrobranchus dombeyi* Cuvier).

- 8.
- Polistotrema stoutii*
- (Lockington).
- California hagfish; Lamperina.*

Coast of California; common in Monterey Bay; Eel River, Humboldt Co., Calif., and other streams of northern California; San Francisco Bay.

Bdellostoma stoutii Lockington, Amer. Nat., XII, December, 1878, 793, Eel River, Calif.

- 9.
- Polistotrema deani*
- Evermann and Goldsborough.

Alaska to Santa Barbara, Calif., in deep water.

Polistotrema deani Evermann and Goldsborough, Bull. U. S. Bur. Fish., XXVI, 1906, 225, fig. 1, Albatross station 4238 in Behm Canal off Nose Point in 229 to 231 fathoms.*Polistotrema curtiss-jamesii* Townsend and Nichols, Bull. Amer. Mus. Nat. Hist., LII, 1925, 4, off California, lat. 34° to 36° in 440 to 585 fathoms.

Order C. HYPEROARTIA. Lampreys

Family 5. PETROMYZONIDÆ. Lampreys

Genus 5. PETROMYZON Linnaeus.

Petromyzon (Artedi) Linnaeus, Syst. Nat., ed. X, 1758, 230 (*P. marinus* L.).*Ammocætes* Duméril in Cuvier, Régne Anim., ed. I, 1817, 119, variant spelling of *Ammocetus*, a name given to a larval form (*branchialis*).*Bathymyzon* Gill, Proc. U. S. Nat. Mus., VI, 1883, 25 (*Bathymyzon bairdii* Gill); probably an old example that has lost teeth.*Oceanomyzon* Fowler, Proc. Acad. Nat. Sci. Phila., LIX, 1907, 461 (*O. wilsoni* Fowler); like the preceding, supposed by Creaser and Hubbs to be an old individual from the Gulf Stream.

- 10.
- Petromyzon marinus*
- Linnaeus.
- Great sea lamprey; Lamprey eel; Sucker; Lamprey; River lamprey; Green lamprey; Shad lamprey; Nine-eyes.*

Atlantic coasts of Europe and America, southward in America to Chesapeake Bay; abundant northward; ascending streams to spawn.

Petromyzon marinus Linnaeus, Syst. Nat., ed. X, 1758, 230, European seas.*Petromyzon americanus* LeSueur, Trans. Amer. Phil. Soc. Phila., I, 1818, 383, coast of Massachusetts.*Petromyzon nigricans* LeSueur, loc. cit., 385, Massachusetts; small example from fresh water.*Ammocetes bicolor* LeSueur, loc. cit., 386, Northampton, Mass. (larva).? *Petromyzon* (*Bathymyzon*) *bairdii* Gill, Proc. U. S. Nat. Mus., VI, 1883, 254, Gulf Stream, north lat. 40° 02', west long. 68° 50' 30", in 547 fathoms.? *Oceanomyzon wilsoni* Fowler, Proc. Acad. Nat. Sci. Phila., LIX, 1907, 461; open sea.

- 11. *Petromyzon marinus dorsatus* Wilder.** *Lake lamprey; Mud lamprey.*
Landlocked in Cayuga Lake and other lakes of western and central New York.
Petromyzon marinus dorsatus Wilder, in Jordan and Gilbert, Synop. Fishes North Amer., 1883, 869, Cayuga Lake, N. Y.
- Genus 6. ICHTHYOMYZON Girard.** *River lampreys.*
Ichthyomyzon Girard, Pac. R. R. Surv., X, 1858, 381 (*Petromyzon argenteus* Kirtland).
Scolecosoma Girard, loc. cit., 385 (*Ammocetes concolor* Kirtland); larva of *argenteus*.
- 12. *Ichthyomyzon concolor* (Kirtland).** *Silver lamprey.*
Great Lakes; upper Mississippi Valley; tributaries of Hudson Bay; south to Arkansas and Louisiana.
Petromyzon argenteus Kirtland, Zool. Ohio, 170, 1838, 197, Big Miami River, Ohio; name preoccupied.
Ammocetes concolor Kirtland, Jour. Bost. Soc. Nat. Hist., III, 1840, 473, plate, Mahoning River; Scioto River; larva.
Petromyzon bdellium Jordan, Cat. Fishes North Amer., 1885, 4; after Kirtland.
Ichthyomyzon castaneus Girard, Pac. R. R. Surv., X, 1858, 381, Galena, Minn.
Ichthyomyzon hirudo Girard, loc. cit., 382, Fort Smith, Ark.
- Genus 7. REIGHARDINA Creaser and Hubbs.**
Reighardina Creaser and Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 120, 1922, 4 (*Ichthyomyzon fossor* Reighard and Cummins = *Ammocetes unicolor* DeKay).
- 13. *Reighardina unicolor* (DeKay).**
Ammocetes unicolor DeKay, N. Y. Fauna: Fishes, 1842, 383, pl. 79, fig. 250; Lake Champlain.
Ammocetes borealis Agassiz, Lake Superior, 1850, 252, Michigan River, Mich.
Ichthyomyzon fossor Reighard and Cummins, Occ. Papers Mus. Zool. Univ. Mich., 31, 1916, 1 to 12, pl. 1, 2, Michigan.
- Genus 8. LETHENTERON Creaser and Hubbs.**
Lethenteron Creaser and Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 120, 1922, 6 (*Lampetra wilderi* Gage = *Petromyzon appendix* DeKay).
- 14. *Lethenteron appendix* (DeKay).** *Small black lamprey; Brook lamprey.*
New York to Iowa, in tributaries of the Great Lakes and of the Ohio and Mississippi, south to Virginia.
Petromyzon nigrum Rafinesque, Ichth. Ohiensis, 1820, 84, falls of Ohio River; name preoccupied.
Petromyzon appendix DeKay, N. Y. Fauna: Fishes, 1842, 381, pl. 64, fig. 211, Providence; Hudson Bay.
Lampetra wilderi Gage, in Jordan and Evermann, Fishes North and Mid. Amer., 1896, 13, Cayuga Lake, N. Y.
- Genus 9. TETRAPLEURODON Creaser and Hubbs.**
Tetrapleurodon Creaser and Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 120, 1922, 9 (*Lampetra spadicea* Bean).
- 15. *Tetrapleurodon spadiceus* (Bean).** *Mexican lamprey.*
Known only from the Rio Lerma, near Guanajuato, and from Lago de Chapala, Mexico.
Lampetra spadicea Bean, Proc. U. S. Nat. Mus., X, 1887, 374, Pl. XX, fig. 6, Rio Lerma, Guanajuato, Mexico.
- Genus 10. ENTOSPHENUS Gill.**
Entosphenus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 331 (*Petromyzon tridentatus* Gairdner).

16. *Entosphenus tridentatus* (Gardner). *Three-toothed lamprey; Western lamprey.*
 Pacific coast of America, from southern California to Unalaska; ascending streams even to the headwaters of the Columbia for spawning purposes.
Petromyzon tridentatus Gardner, in Richardson, Fauna Bor.-Amer., 1836, 293, falls of the Willamette River, Oreg.
Petromyzon lividus Girard, Pac. R. R. Surv., X, 1858, Pt. 4, Fishes, 379, Willamette River, Oreg.
Petromyzon astori Girard, loc. cit., 380, Astoria, Oreg.
Ammocetes cibarius Girard, loc. cit., 383 (larva).
17. *Entosphenus ciliatus* (Ayres).
 Streams of northern California.
Petromyzon ciliatus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 43, San Francisco.
Petromyzon epihexodon Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 331, Fort Reading, Calif.
- Genus 11. **LAMPETRA** (Bonnaterre) Gray. *Brook lampreys.*
Lampetra Bonnaterre, Enc. Meth., 1788; no paging; no type, used for all.
Lampreda Rafinesque, Anal. Nat., 1815, 94 (no species named; no description).
Pricus Rafinesque, loc. cit. (no species named); no description.
Lampetra Gray, Proc. Zool. Soc. Lond., 1851, 236 (*Petromyzon fluvialis* L.).
18. *Lampetra ayresii* (Günther). *Western lamprey.*
 Pacific coast of North America from the Fraser River southward to the Sacramento; very close to *L. fluviatilis* of Europe.
Petromyzon plumbeus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 27, San Francisco; name preoccupied.
Petromyzon ayresii Günther, Cat., VIII, 1870, 505, San Francisco; substitute for *plumbeus* Ayres, preoccupied.
19. *Lampetra borealis* (Girard). *Arctic lamprey.*
 Streams of northern Alaska and Kamchatka.
Petromyzon borealis Girard, Pac. R. R. Surv., X, 1858, Pt. 4, Fishes, 377, Great Slave Lake; not *Ammocetes borealis* Agassiz.
Ammocetes aureus Bean, Proc. U. S. Nat. Mus., IV, 1881, 159, Yukon River.
- Genus 12. **OKKELBERGIA** Creaser and Hubbs.
Okkelbergia Creaser and Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 120, 1922, 8 (*Petromyzon lamotteri* LeSueur).
20. *Okkelbergia lamotteri* (LeSueur).
Petromyzon lamotteri LeSueur, New York Fauna: Fishes, 1842, 382; pl. 79, fig. 249; no locality.
Ammocetes epyptera C. C. Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 327 (type redescribed by Fowler, *ibid.*, 1901, 328; 1907, 466).

Class III. ELASMOBRANCHII. (Selachii, Plagiostomata, Chondropterygii, Antacca). Sharks; Skates and rays; Chimeras

Subclass SELACHII. Sharks

Order D. CESTRACIONTES

Suborder PROARTHRI

Family 6. HETERODONTIDÆ. Bullhead sharks

Genus 13. GYROPLEURODUS Gill.

Gyropleurodus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 331 and 489 (*Cestracion francisci* Girard).

Tropidodus Gill, loc. cit., 489 (*Cestracion pantherinus* Valenciennes = *Cestracion quoyi* Fréminville).

21. *Gyropleurodus francisci* (Girard). *Bullhead shark.*

Coast of California; abundant south of Point Conception.

Cestracion francisci Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 196, Monterey Bay, Calif.

Order E. NOTIDANI. (Diplospondyli)

Suborder OPISTHARTHRI

Family 7. HEXANCHIDÆ. Cow sharks

Genus 14. HEXANCHUS Rafinesque.

Hexanchus Rafinesque, Caratteri, etc., 1810, 14 (*Squalus griseus* Bonnaterre).

Notidanus Cuvier, Règne Anim., ed. 1, 1817, 128 (*Squalus griseus* L.).

Monopterhinus Blainville, Faune Française, 1828, 77 (*Squalus griseus* Gmelin).

22. *Hexanchus griseus* (Bonnaterre). *Cow shark; Caffa bota.*

Mediterranean to Scotland; Cuba.

Squalus griseus Bonnaterre, Ichth. R. 1788, 9; Mediterranean; after *le griset* Bonnaterre.

Squalus griseus Gmelin, Syst. Nat., I, 1795, 1789; after Broussonet.

23. *Hexanchus corinus* Jordan and Gilbert. *Shovel-nosed shark.*

Pacific coast of America; Monterey Bay to Puget Sound.

Hexanchus corinus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880 (December 21), 352, Neah Bay, Wash., and Soquel, Calif.

Genus 15. NOTORYNCHUS Ayres.

Notorynchus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 73 (*Notorynchus maculatus* Ayres).

24. *Notorynchus maculatus*⁵ Ayres.

Pacific coast of the United States, from Monterey Bay northward to Washington; common northward, especially in Humboldt and San Francisco Bays.

Notorynchus maculatus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 72, San Francisco Bay.

Notorhynchus borealis Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 150, Nisqually, Wash.

Order F. EUSELACHII

Suborder GALEI. True sharks

Family 8. SCYLLIORHINIDÆ. Cat sharks

Genus 16. SCYLLIORHINUS Blainville. *Rousselles.*

Catulus Valmont, Dict. Raisonné, IV, 1768, 451 (*C. saxatilis* Valmont = *Squalus stellaris* L.); non binomial.

Catulus Smith, Proc. Zool. Soc. Lond., 1837, 85 (*Scyllium capense* Smith); name preoccupied by *Catulus* Kniphof, 1759, a genus of insects.

Scylliorhinus Blainville, Jour. Phys., 1816, 263 (*Squalus canicula* L., as restricted); spelled *Scylloirhinus* by Garman.

Scylium Cuvier, Règne Anim., ed. I, 1817, 124 (*Squalus canicula* L., as restricted).

25. *Scylliorhinus retifer* (Garman).

Gulf Stream in deep water; occasionally off South Atlantic coast.

Scylium retiferum Garman, Bull. Mus. Comp. Zool., VIII, 1881, 233, off coast of Virginia in deep water.

26. *Scylliorhinus cephalus* (Gilbert).

Gulf of California and southward in deep water.

Catulus cephalus Gilbert, Proc. U. S. Nat. Mus., XV, 1891, 541, deep water near the Revillagigedo Islands and in the Gulf of California.

Genus 17. CEPHALOSCYLIUM Gill.

Cephaloscylium Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 407, 412 (*Scylium laticeps* A. Duméril).

⁵ Regarded by Garman as identical with *N. platycephalus* (Tenore) of the Mediterranean.

27. *Cephaloscyllium uter* (Jordan and Gilbert). *Swell shark.*
Monterey to San Diego; very abundant in Santa Barbara channel.
Catulus uter Jordan and Gilbert, in Jordan and Evermann, Fishes North
and Mid. Amer., 1896, 25; Monterey, San Diego, and Santa Barbara
Channel.
- Genus 18. *APRISTURUS* Garman.
Apristurus Garman, Plagiostomia, Mem. Mus. Comp. Zool., XXXVI,
1913, 96 (*Scylliorhinus indicus* Brauer).
28. *Apristurus brunneus* (Gilbert).
*Gulf of California and northward in deep water; Monterey Bay; Puget
Sound; Hood Canal.*
Catulus brunneus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 542, Gulf of
California.
29. *Apristurus profundorum* (Goode and Bean).
North Atlantic, in deep water.
Scylliorhinus profundorum Goode and Bean, Oceanic Ichth., 1896, 17, Gulf
Stream.
- Genus 19. *PRISTIURUS* (Bonaparte) Müller and Henle.
Pristiurus Bonaparte, Prospetto del Sistema d'Ittiologia Generale, 1831,
121 (*Galeus melanostomus* Rafinesque); name only; defined in Icono-
graphia della Fauna Italica, 1841 (no paging).
Pristiurus Müller and Henle, Mag. Nat. Hist., III, Jan., 1838, 341 (*Galeus
melanostomus* Rafinesque).
Pristidurus Bonaparte, Mem. Soc. Sci. Neufchâtel, II, 1839, 11, variant
spelling.
30. *Pristiurus xaniurus* (Gilbert).
Pacific coast of southern and Lower California; abundant in deep water.
Catulus xaniurus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 540, off
southern and Lower California in 184 to 684 fathoms.
31. *Pristiurus areæ* Nichols.
Off the coast of Florida.
Pristiurus areæ Nichols, Amer. Mus. Novit., No. 256, March 12, 1927, 1,
off the coast of Florida.
- Family 9. GINGLYMOSOMIDÆ. Nurse sharks
- Genus 20. *GINGLYMOSOMA* Müller and Henle.
Ginglymostoma Müller and Henle, Wiegmann's Archiv, I, 1837, 396
(*Squalus cirratus* Gmelin).
32. *Ginglymostoma cirratum* (Gmelin). *Nurse shark; Gata.*
*West Indies and the west coast of Mexico; Cape Verde Islands; occasional
on our South Atlantic coast.*
Squalus cirratus Gmelin, Syst. Nat., I, 1492, 1788, American seas; after
Broussonet.
Squalus punctatus Bloch and Schneider, Syst. Ichth., 1801, 134, Cuba;
after *Gata Hispanus* of Parra.
Squalus punctulatus Bloch and Schneider, loc. cit., 549, Cayenne.
Squalus argus Bancroft, Zool. Jour., V, 1832, 82, West Indies.
Ginglymostoma fulvum Poey, Memorias, II, 1861, 342, Habana; young.
Ginglymostoma caboverdianus Capello, Jour. Sci. Math. Phys. Nat. Lisbon,
1867, 167, Cape Verde Islands.
- Family 10. PSEUDOTRIAKIDÆ
- Genus 21. *PSEUDOTRIAKIS* Capello.
Pseudotriakis Capello, Jour. Sci. Math. Phys. Nat. Lisbon, IV, 1867, 321
(*Pseudotriakis microdon* Capello).
33. *Pseudotriakis microdon* Capello. *Peixe carago.*
Portugal; Amagansett, Long Island.
Pseudotriakis microdon Capello, Jour. Sci. Math. Phys. Nat. Lisbon, 1868,
321, pl. 5, fig. 1, Portugal.

Family 11. CARCHARIIDÆ. Gray sharks

Genus 22. CYNIAS Gill.

Cynias Gill, Proc. U. S. Nat. Mus., XXVI, 1903, 960 (*Squalus canis* Mitchell); embryos without placental attachment.

Mustelus Valmont de Bomare, Dict. Raisonnée, ed. II, 1768, 786 (*Galeus asterias* Valmont = *Squalus canis* Mitchell; not given as a scientific name but as Latin vernacular).

34. *Cynias canis*^a (Mitchill). *Dog shark; Dogfish; Smooth dogfish; Blue dog.* Cape Cod to Cuba; abundant on the Atlantic coast of both continents. *Squalus canis* Mitchell, Trans. Lit. Phil. Soc. N. Y., 1815, 486, New York. *Mustelus asterias* Cloquet, Dict. Raisonnée, 1820, 407, Europe. *Mustellus stellatus* Risso, Europ. Mérid., III, 1826, 1126, Nice. *Mustelus plebejus* Bonaparte, Fauna Italica, Pesci, Fascl., Tome III, 1834, 43, pl. 132, Italy.

35. *Cynias lunulatus* (Jordan and Gilbert). *Gato.*

West coast of Mexico; common in Gulf of California.

Mustelus lunulatus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 108, Mazatlan, Mexico.

Genus 23. MUSTELUS Linck.

Mustelus Linck, Magazin für Neueste aus Physik und Naturgeschichte, 1790, 31 (*Squalus mustelus* Linnaeus); not of Valmont de Bomare.

Galeus^b Rafinesque, Caratteri, 1801, 13 (*Squalus mustelus* L., as restricted by Jordan and Gilbert).

Mustelus Leach, Observ. Squalus, 1813, 62 (*Squalus mustelus* L.).

Mustelus Cuvier, Règne Anim., ed. I, 1817, 128 (*Squalus mustelus* Cuv.).

Pleuracromylon Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 148, (*Mustelus lœvis* Risso); embryo attached by a placenta.

36. *Mustelus mustelus* (Linnaeus). *Smooth hound; Boce dulce.*

Southern Europe; occasional in America, lately taken on coast of New Jersey.

Squalus mustelus Linnaeus, Syst. Nat., ed. X, 1758, 235, Europe.

Mustelus lœvis Risso, Europ. Mérid., 1826, 327, Nice; and of European writers generally.

Galeus mustelus Rafinesque, Caratteri, 1910.

37. *Mustelus californicus* Gill.

California, north to Cape Mendocino.

Mustelus californicus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 148, San Francisco.

38. *Mustelus dorsalis* Gill.

Panama and neighboring waters, north to Gulf of California.

Mustelus dorsalis Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 149, Panama.

Mustelus mento Cope, Proc. Amer. Phil. Soc., XVII, 1877, 47, Pacific Ocean at Pacasmayo, Peru.

Genus 24. RHINOTRIACIS Gill.

Rhinotriacus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 486 (*Rhinotriacus henlei* Gill).

^a The European synonymy of *Mustelus mustelus* and *Cynias canis* is in a state of great confusion, as few of the earlier writers recognized any difference between the two species externally very much alike.

^b The genus *Galeus* of Rafinesque was based on species with spiracles, differing from *Squalus* by the presence of an anal fin. One new species is described, *Galeus melanostomus*, later type of Bonaparte's genus *Pristurus*; but on the same page and the next page Rafinesque speaks of other species—*Galeus vulpecula*, *G. catulus*, and *G. mustelus*. It is plain that *Squalus galeus* L. was his intended type, after the general custom of writers who elevated species to the rank of genus; but he does not mention that species in the paper nor in his Indice, and as *Squalus galeus* was therefore not available as type, Jordan and Gilbert selected *Squalus mustelus* for that purpose, thus making *Galeus* Rafinesque a synonym of *Mustelus* Leach, rather than to displace *Pristurus*. Apparently this procedure was allowable, unless some earlier author had chosen either *Pristurus melanostomus* or *Scylliorhinus catulus* (*canicula*) for that purpose.

39. Rhinotriacis henlei Gill.

Coast of California from Humboldt Bay to Monterey; rather rare.

Rhinotriacis henlei Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 486,
San Francisco.

Genus 25. TRIAKIS Müller and Henle.

Triakis Müller and Henle, Mag. Nat. Hist., II, 1838, 36 (*Triakis scyllium*);
corrected to *Triakis* by authors.

40. Triakis semifasciatum Girard. *Cat shark; Leopard shark.*

Cape Mendocino to San Diego; common.

Triakis semifasciatum Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854,
196, Presidio de San Francisco.

Mustelus felis Ayres, Proc. Calif. Acad. Nat. Sci., 1854, 17, San Francisco.

Genus 26. GALEORHINUS Blainville.

Galeorhinus Blainville, Bull. Soc. Philom., 1816, 121 (*Squalus galeus* L.;
restricted by Gill).

Eugaleus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 148 (*Squalus
galeus* L.); the first species named by Blainville being *Squalus mustelus*,
for which reason Garman uses *Galeorhinus* as the name of the genus
here called *Mustelus*.

41. Galeorhinus zyopterus Jordan and Gilbert. *Oil shark; Soup-fin shark.*

Coast of southern and Lower California, from San Francisco to Cedros
Island; very abundant.

Galeorhinus zyopterus Jordan and Gilbert, Synop. Fishes North Amer.,
1883, 871, San Pedro, Calif.

Genus 27. TRIENODON Müller and Henle.

Trienodon Müller and Henle, Plagiostomen, 1841, 55 (*Carcharias obesus*
Rüppell).

42. Trienodon obesus (Rüppell).

Tropical Pacific; reported from Cocos Island.

Carcharias obesus Rüppell, Neue Wirbel., Fisch., 1835, 64, pl. 18, fig. 2,
Djidda, Arabian Coast of Red Sea.

Genus 28. GALEOCERDO Müller and Henle.

Galeocerdo Müller and Henle, Plagiostomen, 1838, 59 (*Galeocerdo tigrinus*
Müller and Henle).

Boreogaleus Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 411 (*Squalus
arcticus* Faber).

43. Galeocerdo arcticus (Faber). *Spotted shark; Leopard shark; Tiger shark;
Tigrene.*

Tropical seas; occasionally to Iceland, Cape Cod, and San Diego.
Squalus arcticus Faber, Fische Islands, 1829, 17, Iceland and neighboring
seas.

Galeocerdo tigrinus Müller and Henle, Plagiostomen, 1838, 59, Pondicherry.
Galeus maculatus Ranzani, De Novis Speciebus Piscium, Dissert. Prima,
in Novi Com. Acad. Boron., IV, 1840, 68, Brazil.

Genus 29. HEMIGALEUS Bleeker.

Hemigaleus Bleeker, Verh. Bot. Gen., XIV, 1852, 45 (*Hemigaleus mi-
crostoma* Bleeker); not *Hemigalea*, an earlier genus of mammals.

44. Hemigaleus pectoralis Garman.

Coast of New England, probably in the Gulf Stream.

Hemigaleus pectoralis Garman, Bull. Mus. Comp. Zool., XLIV, 1906, 203,
"Aquarial Gardens," the fishes in which were taken off Massachusetts
and Rhode Island.

Genus 30. SCOLIODON Müller and Henle.

Scoliodon Müller and Henle, Wiegmann's Archiv, I, 1837, 398 (*Scoliodon
laticaudus* Müller and Henle).

45. Scoliodon terra-novæ (Richardson). *Sharp-nosed shark; Newfoundland
shark.*

Cape Cod to Florida; very common southward along the Atlantic coast.
Squalus (Carcharias) terra-novæ Richardson, Fauna Bor.-Amer., III, 1836,
289, Newfoundland; an error, probably for Florida.

Squalus punctatus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 483,
New York; not of Bloch and Schneider.

- 46. Scoliodon lalandi** (Müller and Henle).
West Indies to Brazil.
Carcharias lalandi Müller and Henle, *Plagiostomen*, 1838, 30, Rio Janeiro; Martinique; Guadeloupe.
- 47. Scoliodon porosus** (Poey).
Cuba.
Squalus porosus Poey, *Memorias*, II, 1861, 339, tab. 19, L. 11, 12, Habana.
- 48. Scoliodon longurio** (Jordan and Gilbert).
Pacific coast of Mexico.
Carcharias longurio Jordan and Gilbert, *Proc. U. S. Nat. Mus.*, V, 1882, 106, Mazatlan, Mexico.
- Genus 31. APRIONODON** Gill.
Aprion Müller and Henle, *Plagiostomen*, 1838-1841, 31 (*Carcharias isodon* Müller and Henle); preoccupied.
Aprionodon Gill, *Ann. Lyc. Nat. Hist. N. Y.*, VII, 1862, 411 (*Aprionodon punctatus* Gill; not *Squalus punctatus* Mitchell= *Carcharias isodon* Müller and Henle).
- 49. Aprionodon isodon** (Müller and Henle).
Atlantic Ocean; recorded from New York, Virginia, and Cuba; probably rare.
Carcharius isodon Müller and Henle, *Plagiostomen*, 1838, 32, New York.
Carcharius punctatus Günther, *Cat.*, VIII, 1870, New York; not *Squalus punctatus* Mitchell, which is *Scoliodon terre-novæ*.
- Genus 32. CARCHARIAS** Rafinesque. *Gray shark; Requin.*
Carcharias Rafinesque, *Caratteri*, 1810, 10 (*Squalus carcharias* L. intended, but *Carcharias taurus* the only species mentioned).
Carcharhinus Blainville, *Bull. Soc. Philom.*, 1816, 264 (*Squalus commersonii* Blainville, apparently the same as *Carcharias lamia* Rafinesque).
Carcharias Cuvier, *Règne Anim.*, ed. I, 1817, 125 ("*Squalus carcharias*," *Carcharias lamia* intended); not of Rafinesque.
Eulamia Gill, *Ann. Lyc. Nat. Hist. N. Y.*, VII, 1861, 401 (*Carcharias lamia* Rafinesque; *Squalus commersonii* Blainville being intended= *Carcharias lamia* Rafinesque was *Carcharodon carcharias* Rafinesque, the specific name being a substitute for *Squalus carcharias* L.).
Platypodon Gill, loc. cit. (*Carcharias menisorrah* Müller and Henle).
Isogomphodon Gill, loc. cit. (*Carcharias oxyrhynchus* Müller and Henle).
Lamioips Gill, loc. cit. (*Carcharias temmincki* Müller and Henle).
Isoplaidodon Gill, loc. cit. (*Carcharias sorrah* Müller and Henle).
- 50. Carcharias commersonii** (Blainville). *Cub shark; Requin; Requiem; Lamia; Common gray shark.*
Tropical parts of the Atlantic; common northward to the Florida Keys and in the Mediterranean; rarely straying northward to New York in the Gulf Stream.
Squalus commersonii Blainville, *Bull. Soc. Philom.*, 1816, 121, Mediterranean.
Carcharias lamia Risso, *Europ. Mérid.*, III, 1826, 19, Nice; not of Rafinesque.
Squalus longimanus Poey, *Memorias*, II, 1861, 338, Cuba.
- 51. Carcharias lamiella** Jordan and Gilbert. *Bay shark.*
San Diego Bay and southward along the Mexican coast.
Carcharias lamiella Jordan and Gilbert, *Proc. U. S. Nat. Mus.*, V, 1882, 110, San Diego; regarded by Garman as identical with *C. commersonii* of the Atlantic.
- 52. Carcharias platyrhynchos** (Gilbert).
Magdalena Bay, Lower California, to Galapagos Islands and the Revillagigedo Islands.
Eulamia platyrhynchus Gilbert, *Proc. U. S. Nat. Mus.*, XIV, 1891, 543, Clarion Island; Magdalena Bay; regarded by Garman as identical with *C. commersonii*.
- 53. Carcharias platyodon** (Poey). *Cub shark.*
Cuba to Texas; abundant in the Gulf of Mexico.
Squalus platyodon Poey, *Memorias*, II, 1861, 336, Habana.
Squalus obtusus Poey, loc. cit., 337, Habana.

54. *Carcharias milberti* Valenciennes. *Small blue shark; White shark.*
Cape Cod to Florida; not rare; recorded from the Mediterranean.
Carcharias (Prionodon) milberti Valenciennes, in Müller and Henle,
Plagiostomen, 1838, 38, New York.
Carcharias caruleus DeKay, N. Y. Fauna: Fishes, 1842, 49, New York.
Lamna caudata DeKay, loc. cit., 354, Brenton Reef, R. I.
55. *Carcharias obscurus* (LeSueur). *Dusky shark.*
North and Middle Atlantic; frequently on our coast.
Squalus obscurus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 223, New
York.
Carcharias falcipinnis Lowe, Proc. Zool. Soc. Lond., 1839, 90, Madeira.
Prionodon obvelatus Valenciennes, Poiss. îles Canaries, 1839, 103, Canaries.
56. *Carcharias azureus* Gilbert and Starks.
Carcharias azureus Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904,
11, Pl. II, fig. 5, Panama.
57. *Carcharias nicaraguensis* (Gill and Bransford). *Nigrone; Fresh-water
shark.*
Lake Nicaragua (landlocked); also in the Bay of Panama.
Eulamia nicaraguensis Gill and Bransford, Proc. Acad. Nat. Sci. Phila.,
XXIX, 1877, 190, Lake Nicaragua.
58. *Carcharias henlei* Valenciennes.
Coast of Brazil, northward to Guiana.
Carcharias henlei Valenciennes, in Müller and Henle, Plagiostomen,
1838, 46, Cayenne.
Carcharias porosus Ranzani, De Novis Speciebus Piscium, Dissert Prima.;
in Novi Com. Acad. Bonon., IV, 1840, 70, Brazil.
59. *Carcharias cerdale* (Gilbert).
Abundant at Panama.
Carcharhinus cerdale Gilbert, in Jordan and Evermann, Fishes North
and Mid. Amer., III, 1898, 2746, Panama; united by Garman with
C. menisorrah from the East Indies.
60. *Carcharias falciformis* Bibron. *Cazon de playa (wharf) shark.*
Cuba and neighboring waters.
Carcharias falciformis Bibron, in Müller and Henle, Plagiostomen, 1838,
47, Cuba.
Squalus tiburo Poey, Memorias, II, 1861, 331, Habana.
61. *Carcharias remotus* Valenciennes.
West Indies and Brazil.
Carcharias remotus Valenciennes, in Duméril, Hist. Nat. Poiss., I, 1865,
374.
Platypodon perezi Poey, Anal. Soc. Española Hist. Nat., V, 194, pl. 9,
figs. 2, 3, Cuba.
62. *Carcharias acronotus* (Poey).
Cuba.
Squalus acronotus Poey, Memorias, II, 1861, 335, Cuba.
63. *Carcharias velox* (Gilbert).
Panama.
Carcharhinus velox Gilbert, in Jordan and Evermann, Fishes North and
Mid. Amer., 1898, 2747, Panama.
64. *Carcharias oxyrhynchus* Müller and Henle.
Surinam.
Carcharias oxyrhynchus Müller and Henle, Plagiostomen, 1838, 41,
Surinam.
65. *Carcharias limbatus* Müller and Henle. *Carconetta.*
Tropical seas north to Florida; a stray example taken at Woods Hole,
Mass.
Carcharias (Prionodon) limbatus Müller and Henle, Plagiostomen, 1838,
49, Martinique.
Isogomphodon maculipinnis Poey, Repertorio, I, 1866, 191, Cuba.
Carcharias (Prionodon) müllerii Steindachner, Sitzungs. Kais. Akad.
Wiss. Wien, LVI, 1867, 356, West Indies.
Carcharias microps Lowe, Proc. Zool. Soc. Lond., 1840, 38, Madeira.
Prionodon cucuri Castelnau, Anim. Amer. Sud. Poiss., 1855, 99, Bahia.

- 66. *Carcharias æthalorus* Jordan and Gilbert.**
Mazatlan to Panama.
Carcharias æthalorus Jordan and Gilbert, Proc., U. S. Nat. Mus., V, 1882, 104, Mazatlan; regarded by Garman as inseparable from *C. limbatus*.
- 67. *Carcharias natator* (Meek and Hildebrand).**
Gulf of Panama.
Carcharhinus natator Meek and Hildebrand, Marine Fishes of Panama, Field Mus. Nat. Hist., Zool., XV, Pt. I, Dec. 20, 1923, 40, pl. 1, fig. 1, Panama City fish market.
- 68. *Carcharias fronto* Jordan and Gilbert. *Tiburon.***
Pacific coast of Mexico.
Carcharias fronto Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 102, Mazatlan. Regarded by Garman as inseparable from *C. milberti*.
- Genus 33. HYPOPRION Müller and Henle.⁸**
Hypopprion Müller and Henle, Plagiostomen, 1838, 34 (*Carcharias macloti* Müller and Henle).
Hypopprionodon Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 401 (*Carcharias hemiodon* Valenciennes).
- 69. *Hypopprion brevirostris* Poey.**
West Indies, north to Charleston and Pensacola.
Hypopprion brevirostris Poey, Repertorio, II, 1868, 451, Tab. 4, Cuba.
- 70. *Hypopprion signatus* Poey.**
Cuba.
Hypopprion signatus Poey, Synop., 1868, 452, Cuba.
Hypopprion longirostris Poey, Enumeratio, 1876, 199, Cuba.
- Genus 34. PRIONACE⁹ Cantor.**
Galeus Valmont de Bomare, Dict. Raisonnée, I, 1768, 371, "Cagnot bleu, *Galeus glaucus* Valmont"; Latin vernacular.
Prionodon Müller and Henle, Plagiostomen, 1838, 36 (*Squalus glaucus* L.); name preoccupied.
Prionace Cantor, Malayan Fishes, 1850, 399, substitute for *Prionodon* (*Squalus glaucus* Linnaeus).
Cynocephalus Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 401 (*Squalus glaucus* L.); after Klein, whose names are not binomial.
- 71. *Prionace glauca* (Linnaeus). *Great blue shark.***
A large shark of the warm seas, occasionally taken on our coasts, both Atlantic and Pacific.
Squalus glaucus Linnaeus, Syst. Nat., ed. X, 1758, 235, Seas of Europe.
Squalus cæruleus Blainville, Faune Française, 1828, 91, Mediterranean.
Squalus hirundinaceus Valenciennes, in Müller and Henle, Plagiostomen, 1838, 37, Brazil.

Family 12. SPHYRNIDÆ. Hammer-headed sharks

- Genus 35. RENICEPS Gill.**
Reniceps Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1862, 403 (*Squalus tiburo* Linnaeus).
- 72. *Reniceps tiburo* (Linnaeus). *Shovelhead shark; Bonnethead shark.***
Atlantic Ocean, abundant from Long Island southward; Pacific Ocean ranging from Mazatlan and Bay of Panama to China.
Squalus tiburo Linnaeus, Syst. Nat., ed. X, 1758, 234, America.
- Genus 36. PLATYSQUALUS Swainson.**
Platysqualus Swainson, Class. Anim., II, 1839, 318 (*Zygæna tudes* Cuvier, in Valenciennes); misnamed *Zygæna tiburo*.

⁸ Müller and Henle's Systematische Beschreibung der Plagiostomen was published from 1838 to 1841. Garman, probably correctly, gives this and other genera of sharks the date of 1841. Some of them appeared earlier in periodicals.

⁹ The reasons for not accepting the names of Valmont de Bomare in scientific nomenclature are given by Jordan and Evermann, Genera of Fishes, I, p. 27. These names (*Galeus*, *Vulpes*, *Catulus*, and *Mustelus*) are binomial only by accident, and not accepted by the author himself as generic in his list of genera (1791). These are, however, accepted by Garman (Plagiostomia, 1913).

73. *Platysqualus tudes* (Cuvier).

Gulf of California to Bay of Panama; West Indies; Mediterranean Sea and Indian Ocean.

Zygæna tudes Cuvier, in Valenciennes, Mém. Mus. Hist. Nat., IX, 1822, 225, Nice; after *Pantouflier* of Risso.

Genus 37. *SPHYRNA* Rafinesque.

Sphyraña Rafinesque, Indice, 1810, 60 (*Squalus zygæna* Linnæus).

Cestrorhinus Blainville, Jour. Phys., 1816, 264 (*Squalus zygæna* Linnæus).

Zygæna Cuvier, Règne Anim., ed. I, 1817, 127 (*Squalus zygæna* Linnæus); name preoccupied.

Sphyrichthys Thienemann, Lehrbuch der Zoologie, III, 1828, 408 (*Squalus zygæna* Linnæus).

Eusphyra Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 412 (*Squalus zygæna* Linnæus).

Cestracion Garman, Plagiostomen, 1813, 133 (*Squalus zygæna* L.); after Klein; not binomial; not of Cuvier.

74. *Sphyraña zygæna* (Linnæus). Hammerhead shark; Hammerhead; Shovelnose shark.

Cape Cod and Point Conception southward; Bay of Panama and Mazatlan in Pacific; and across to Hawaii and Japan; abundant.

Squalus zygæna Linnæus, Syst. Nat., ed. X, 1758, 234, Europe; America.

Zygæna malleus Valenciennes, Mém. Mus. Hist. Nat., IX, 1822, 223, pl. 11, fig. 1.

Zygæna lewini Lord, in Griffith, Animal Kingdom, X, 1834, 640, pl. 50, New Holland.

Zygæna subarcuata Storer, Proc. Bost. Soc. Nat. Hist., III, 1848, 71, Cape Cod.

Family 13. RHINEODONTIDÆ. Whale sharks

Genus 38. *RHINEODON* Smith.

Rhincodon Smith, Zool. Jour. Lond., No. XVI, 1829, 443 (*R. typicus* Smith); misprint for *Rhineodon*.

Micristodus Gill, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 177 (*M. punctatus* Gill).

Rhinodon and *Rhincodon* of authors.

75. *Rhineodon typicus* Smith. Whale shark; Mhor; Chagrin; Tiburon ballenas; Tint reva; East Indian basking shark.

Coast of Florida; Gulf of California; Cape of Good Hope; Japan to China; India; rare but widely diffused.

Rhincodon typicus Smith, Zool. Jour. Lond., No. XVI, 1829, 443, 444, Table Bay, Cape of Good Hope.

Micristodus punctatus Gill, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 177, Gulf of California.

Family 14. ALOPHIDÆ. Thresher sharks

Genus 39. *ALOPIAS* Rafinesque.

Vulpecula Valmont de Bomare, Dict. Raisonnée, III, 1768, 740 (*V. marina* Valmont); Latin vernacular not intended as a generic name.

Alopias Rafinesque, Caratteri, 1810, 12 (*Alopias macrourus* Rafinesque = *Alopecias vulpes* Günther).

Alopecias Müller and Henle, Plagiostomen, 1838, 74, amended orthography.

76. *Alopias vulpinus* (Bonnaterre). Thresher; Fox shark; Swingletail; Longtail shark; Fish shark; Sea fox; Sea ape; Pez zorro; Pesca pavone.

Mediterranean, Atlantic, and Pacific.

Vulpecula marina Valmont, Dict. Raisonnée, III, 740; not eligible.

Squalus vulpinus Bonnaterre, Tableau Encycl. Ichth., 1788, 9, Mediterranean; after Pennant.

Squalus vulpes Gmelin, Syst. Nat., I, 1789, 1496, Mediterranean; after Pennant.

Alopias macrourus Rafinesque, Caratteri, 1810, 12, Sicily.

Galeus vulpecula Rafinesque, Caratteri, 1810, 15; passing notice.

Family 15. ODONTASPIDÆ. Sand sharks

Genus 40. ODONTASPIS Agassiz.

Odontaspis Agassiz, Poiss. Fossiles, III, 1836, 87 (*Squalus ferox* Risso).

77. **Odontaspis littoralis** (Mitchill). *Sand shark; Shovelnose shark; Horned dogfish; Picked dogfish; Dogfish; Bone-dog; Skittle-dog; Hoe.*
Atlantic coast between Cape Cod and Cape Hatteras; regarded by Garman as not separable from *Carcharias taurus* Rafinesque of the Mediterranean.
Squalus americanus Mitchell, Trans. Lit. and Phil. Soc. N. Y., I, 1815, 483, New York; not of Shaw.
Squalus littoralis and *Squalus macroodus* Mitchell, Amer. Month. Mag., II, 1818, 328, New York.
Carcharias griseus Ayres, Bost. Jour. Nat. Hist., IV, 1843, 288, Long Island.

Family 16. LAMNIDÆ. Mackerel sharks; Man-eaters

Genus 41. LAMNA Cuvier. *Porbeagles.*

Lamna Cuvier, Règne Anim., ed. I, 1817, 126 (*Squalus cornubicus* Gmelin).
Lamia Risso, Europ. Mérid., III, 1826, 123 (*Squalus cornubicus* Gmelin); name preoccupied.

Selanonius Fleming, Brit. Anim., 1828, 169 (*Selanonius walkeri* Fleming = *Squalus cornubicus* Gmelin).

78. **Lamna nasus** (Bonnaterre). *Porbeagle; Mackerel shark; Salmon shark.*
North Atlantic and North Pacific; abundant in central and southeast Alaska.
Squalus nasus Bonnaterre, Tableau Encycl., Ichth., 1788, 10, shores of Cornwall.
Squalus cornubicus Gmelin, Syst. Nat., I, 1789, 1497, shores of Cornwall.
Squalus pennanti Walbaum, Artedi Pisc., 1792, 517, Cornwall; after Pennant.
Squalus monensis Shaw, Gen. Zool., V, 1804, 350, Anglesea.
Selanonius walkeri Fleming, Brit. Anim., 1828, 169, Lochfyne, Argyleshire.

Genus 42. ISURUS Rafinesque.

Isurus Rafinesque, Caratteri, 1810, 11 (*Isurus oxyrinchus* Rafinesque).
Oxyrhina Agassiz, Poiss. Fossiles, III, 1836, 276 (*Isurus spallanzani* Rafinesque = *Isurus oxyrinchus* Rafinesque).

Isuropsis Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1862, 398, 409 (*I. dekayi* Gill).

79. **Isurus punctatus** (Storer). *Mackerel shark.*

Atlantic coast of United States; common.

Lamna punctata Storer, Rept. Fishes Mass., 1839, 185, pl. 3, fig. 2, Massachusetts.

Oxyrhina dekayi Gill, Cat. Fish. E. Coast N. Amer., Suppl. Proc. Acad. Nat. Sci. Phila., XIII, 1861, 60, after Storer; name misprint for *dekeyi*.

80. **Isurus tigris** (Atwood). *Mackerel shark; Mackerel porbeagle; Dentuda.*

Cape Cod to West Indies; not generally common except in the Gulf of Mexico.

Lamna punctata DeKay, N. Y. Fauna: Fishes, 1842, 352, New York; not *Squalus punctatus* Mitchell.

Isuropsis dekayi Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1862, 409; after DeKey; not *Oxyrhina dekayi* Gill, 1861.

Carcharias tigris Atwood, Proc. Bost. Soc. Nat. Hist., XII, 1869, 268, Provincetown, Mass.

81. **Isurus oxyrinchus** Rafinesque. *Mackerel shark; Pesce tondo; Cane de mare.*

Mediterranean Sea and neighboring parts of the Atlantic; occasional on the Atlantic coast of America.

Isurus oxyrinchus Rafinesque, Caratteri, 1810, 12, Palermo.

Isurus spallanzani Rafinesque, Indice, 1810, 60, Sicily; after Spallanzani.

Oxyrhina gomphodon Müller and Henle, Plagiostomen, 1838, 68, Atlantic Ocean.

Squalus rostratus Macrì, Mem. Ac. Sci. Napoli, I, 1819, 55, Tab. 1, fig. 2, Naples.

Genus 43. CARCHARODON¹⁰ Smith. *Man-eater sharks; Great white sharks.*
Carcharodon Andrew Smith, in Müller and Henle, Mag. Nat. Hist., II,
 second article, January, 1838, 37 (*Carcharodon capensis* Smith =
Squalus carcharias Linnaeus).

- 82. *Carcharodon carcharias* (Linnaeus).** *Man-eater shark; Great white shark.*
 Temperate and tropical parts of Atlantic and Pacific; occasionally on our
 coasts, north to New York and Monterey Bay. A dangerous shark.
Squalus carcharias Linnaeus, Syst. Nat., ed. X, 1758, 235, Europe; after
 Aristedi.
Carcharias lamia Rafinesque, Indice, 1810, 44, after Linnaeus; not of
 Risso and later authors.
Carcharodon verus Agassiz, Poiss. Fossiles, III, 1838, 91.
Carcharodon rondeleti Müller and Henle, Plagiostomen, 1838, 70, Medi-
 teranean Sea and Atlantic Ocean.
Carcharodon Smith, in Bonaparte, Mem. Soc. Sci. Neufchâtel, II, 1839, 9
 (errata), New York and Monterey Bay.
Carcharias atwoodi Storer, Proc. Bost. Soc. Nat. Hist., III, 1848, 72,
 Provincetown.
Carcharodon capensis Smith, Ills. Zool. S. Africa, Pisces, 1849, pl. 4,
 Cape of Good Hope.

Family 17. CETORHINIDÆ. Basking sharks

Genus 44. CETORHINUS Blainville.

Cetorhinus Blainville, Bull. Soc. Philom., 1810, 169 (*Squalus gunneri*
 Blainville = *Squalus maximus* Gunner).
Selache Cuvier, Règne Anim., ed. I, 1817, 129 (*Squalus maximus* Gunner).
Polyprosopus Couch, Hist. Brit. Fishes, I, 1862, 67 (*Squalus rashleighanus*
 Couch = *Squalus maximus* Gunner).

- 83. *Cetorhinus maximus* (Gunner).** *Basking shark; Pélérin; Elephant shark;*
Bone shark; Sailfish; Sunfish; Hoe-mother; Capidoli or oilfish.
 Arctic seas, straying southward to Portugal, Virginia, and California.
Squalus maximus Gunner, Trondhjem Skrift., III, 1765, 33, coast of
 Norway.
Squalus gunneri Blainville, Bull. Sci. Philom., 1810, 256; after Gunner.
Squalus pelegrinus Blainville, loc. cit., 257, Europe.
Squalus homianus Blainville, loc. cit., 257, Europe.
Cetorhinus shavianus Blainville, loc. cit., 264, Europe.
Squalus isodus Saverino Macri, Mem. della R. Ac. Sci. Napoli, I, 1819, 55,
 pls. 1 and 2, Naples.
Squalus elephas LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 343, New
 Jersey.
Squalus rashleighanus Couch, Trans. Linn. Soc., XIV, 1825, 91, Cornwall.
Squalus cetaceus Gronow, Cat. Fishes, 1854, 6, Norway.
Polyprosopus macer Couch, Hist. Brit. Fishes, I, 1877, 68, England.

¹⁰ As to the authorship of the name *Carcharodon*, we have this information from the late Allan R. McCulloch, of the Australian Museum:
 "The name *Carcharodon* is not used by Smith. The earliest reference to it is that of Müller and Henle, as quoted above. These authors saw Smith's specimens (p. 33) and perhaps heard his paper read at the Zoological Society in September, 1837. They, therefore, quoted the name as of Smith. But Smith's paper was not published until February 13, 1838, he having read Müller and Henle's paper about a month previously, in which *Carcharodon* appeared, and he evidently withdrew it with names he refers to in his paper." (In lit. June 28, 1920.)

Order G. TECTOSPONDYLI.¹¹ Tectospondyloous sharks

Suborder SQUALOIDEI. (Sharks without anal fin)

Family 18. SQUALIDÆ. Dogfishes

Genus 45. SQUALUS Linnaeus.

Squalus (Artedi) Linnaeus, Syst. Nat., ed. X, 1758, 233 (includes all sharks; *Squalus acanthias* Linnaeus made type by subsequent writers, following Rafinesque, 1810).
Acanthorhinus Blainville, Jour. Phys., 1816, 263 (*Squalus acanthias* Linnaeus).
Acanthias Risso, Europ. Mérid., III, 1826, 131 (*Squalus acanthias* Linnaeus).

84. *Squalus acanthias* Linnaeus. Spiny dogfish; Cod shark; Codfish shark; Thornback shark; Picked dogfish; Bone-dog; Dogfish; Skittle-dog.
 North Atlantic, on both coasts and about Cuba; abundant along shores of Northern and Middle States.
Squalus acanthias Linnaeus, Syst. Nat., ed. X, 1758, 233, coast of Europe.
Acanthias vulgaris Risso, Europ. Mérid., III, 1826, 131, Nice.
Acanthias americanus Storer, Synop. N. Amer. Fishes, in Mem. Am. Acad., II, 1846, 506, Massachusetts.

85. *Squalus suckleyi* (Girard). California dogfish; Dogfish.
 Pacific coast of America, from Santa Barbara to the Aleutian Islands; abundant in southeast and middle Alaska, through to Kamchatka.
Spinax (Acanthias) suckleyi Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 196, Fort Steilacoom, Wash.

Genus 46. CENTROSCYMNUS Bocage and Capello.

Centroscymnus Bocage and Capello, Proc. Zool. Soc. Lond., 1864, 263 (*C. cælolepis* Bocage and Capello).

86. *Centroscymnus cælolepis* Bocage and Capello.
 Coast of Portugal and neighboring parts of the Atlantic; Gloucester, Mass., and Nova Scotia Banks.
Centroscymnus cælolepis Bocage and Capello, Proc. Zool. Soc. Lond., 1864, 263, Portugal.

Genus 47. CENTROSCYLLIUM Müller and Henle.

Centroscyllium Müller and Henle, Plagiostomen, 1838, 191 (*Spinax fabricii* Reinhardt).

87. *Centroscyllium fabricii* (Reinhardt).
 Greenland seas; Gloucester and the Nova Scotia Banks.
Spinax fabricii Reinhardt, Dansk. Vidensk. Selsk. Forhandl., III, 1828, XVI, Greenland.

Genus 48. ETMOPTERUS Rafinesque.

Etmopterus Rafinesque, Caratteri, 1810, 14 (*Etmopterus aculeatus* Rafinesque = *Squalus spinax* L.).
Acanthidium Lowe, Proc. Zool. Soc. Lond., 1839, 91 (*Acanthidium pusillum* Lowe, as restricted by Jordan and Evermann, 1896; restricted by Garman, 1912, to *A. calcea*, a species of *Deania* Jordan and Snyder).

88. *Etmopterus hillianus* (Poey).
 West Indies; Cuba; St. Kitts.
Spinax hillianus Poey, Memorias, II, 1861, 340, Cuba.

Family 19. SOMNIOSIDÆ. Sleeper sharks

Genus 49. SOMNIOSUS LeSueur.

Somnirosus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 222 (*Somnirosus brevipinna* LeSueur = *Squalus microcephalus* Bloch and Schneider).
Læmargus Müller and Henle, Plagiostomen, 1838, 93 (*Squalus borealis* Scoresby = *Squalus microcephalus* Bloch and Schneider).
Leiodon Wood, Proc. Bost. Soc. Nat. Hist., II, 1847, 174 (*Leiodon echinatum* Wood = *Squalus microcephalus* Bloch and Schneider).
Rhinocynmus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 264 (*Scymnus rostratus* Risso).

¹¹ We group together under this name those sharks that, like the rays, lack the anal fin and that have also simplified or degenerate vertebrae, in which the hard material is arranged in rings without radiation.

89. *Somniosus microcephalus* (Bloch and Schneider). *Sleeper shark; Nurse shark.*

Arctic seas, south to Cape Cod, Oregon, San Francisco Bay, and France.
Squalus microcephalus Bloch and Schneider, Syst. Ichth., 1801, 135,
 northern seas.

Somniosus brevipinna LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 222,
 Massachusetts.

Squalus borealis Scoresby, Arct. Reg., I, 1820, 538, pl. 15, Arctic.

Squalus norvegianus Blainville, Faune Française, 1828, 61, Norway.

Squalus glacialis Faber, Natur der Fische Islands, 1829, 23, Iceland.

Family 20. ECHINORHINIDÆ. Bramble sharks

Genus 50. ECHINORHINUS Blainville.

Echinorhinus Blainville, Bull. Soc. Philom., 1816, 121 (*Squalus spinosus* Gmelin).

Goniodus Agassiz, Poiss. Fossiles, III, 1836, 183 (*Squalus spinosus* Gmelin).

90. *Echinorhinus brucus* (Bonnaterre). *Bramble shark.*

Atlantic coasts of Europe, America, and Africa; scarce in the western Atlantic.

Squalus brucus Bonnaterre, Tableau Encycl., Ichth., II, 1788, 1500,
 "the ocean."

Squalus spinosus Gmelin, Syst. Nat., 1789, 1500, "the ocean."

Echinorhinus obesus Smith, Ills. Zool. So. Africa, Fishes, 1845, Pl. I, Cape
 of Good Hope.

Suborder SQUATINOIDEI

Family 21. SQUATINIDÆ. Angel sharks

Genus 51. SQUATINA Duméril.

Squatina Valmont de Bommare, Dict. Raisonnée, I, 1768, 117; not eligible.

Rhina Klein, Neuer Schauplatz, II, 1776, 587; non-binomial.

Squatina Duméril, Zool. Analyt., 1806, 102 (*Squatina angelus* Duméril =
Squalus squatina Linnæus).

Rhina Rafinesque, Caratteri, 1810, 14 (*Squalus squatina* L.).

91. *Squatina duméril*¹² LeSueur. *Angelfish; Monkfish.*

Atlantic coast of United States; Newport, R. I.

Squatina duméril LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 225, pl.
 10, probably Florida.

**92. *Squatina squatina* (Linnæus). *Monkfish; Angelfish; Squato; Shark ray;*
*Monkeyfish; Kingston; Fiddlefish; Little bullhead shark.***

Atlantic on both coasts, north to southern England and Cape Cod.

Squalus squatina Linnæus, Syst. Nat., ed. X, 1758, 233, coasts of Europe.

Squatina lavis Cuvier, Règne Anim., ed. I, 1817, 131; after Linnæus.

Squatina vulgaris Risso, Ichth. Nice, 1810, 45, Nice.

Squatina fimbriata Müller and Henle, Plagiostomen, 1838, 101, Mediterranean Sea.

Squatina oculata Bonaparte, Faun. Ital., Pesc., XXVIII, 1840, Fasci. 147,
 pl. 143, Italy.

Squatina lewisi Couch, Trans. Linn. Soc. Lond., XIV., 1825, 90, Cornwall.

93. *Squatina californica* Ayres. *California angel shark.*

Pacific coast of America, from southeastern Alaska southward; not rare
 on coast of California.

Squatina californica Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 29, San
 Francisco.

¹² As this species has not been distinguished generally from its near relative, *Squatina squatina* L., it is possible that all western Atlantic records belong to it.

Order H. BATOIDÆ. Rays

Suborder SARCURA. Thick-tailed rays

Family 22. PRISTIDÆ. Sawfishes

Genus 52. PRISTIS Linck.

Pristis Klein, Neuer Schauplatz, VII, 1779, 403; non-binomial.*Pristis* Linck, Mag. Neueste Physik und Naturgeschichte, 1790, 31
(*Squalus pristis* L.).*Pristis* Latham, Trans. Linn. Soc. Lond., II, 1794, 276 (*Squalus pristis* L.).

- 94.
- Pristis pectinatus*
- Latham:
- Common sawfish; Pez sierra.*

Tropical Atlantic, north to West Indies and Florida; occasionally at Cape May and Grassy Sound.

Pristis pectinatus Latham, Trans. Linn. Soc. Lond., II, 1794, 278, "in the ocean."*Pristis granulosa* Bloch and Schneider, Syst. Ichth., 1801, 352, Habana; after Parra.*Pristis mississippiensis* Rafinesque, Ichth. Ohiensis, 1820, 86, lower Mississippi.*Pristis megalodon* Duméril, Elasmobranchs, 1870, 476, pl. 9, Cayenne.*Pristis acutirostris* Duméril, loc. cit., 479, Martinique.*Pristis occa* Duméril, loc. cit., 479, locality unknown.

- 95.
- Pristis zephyreus*
- ¹²
- Jordan and Starks.
- Pez de espada.*

Tropical seas entering rivers, north to Mazatlan on the Pacific coast.

Pristis zephyreus Jordan and Starks, Fishes of Sinaloa, in Proc. Calif. Acad. Sci., V, 1895, 383, Mazatlan, Mexico; regarded by Garman as inseparable from *Pristis microdon* Latham.

Family 23. RHINOBATIDÆ. Guitarfishes

Genus 53. RHINOBATUS Linck. *Guitarfishes.**Rhinobatos* Klein, Neuer Schauplatz, II, 1776, 593 (*Rhinobatos seu= Squatina raja*; non-binomial).*Rhinobatos* Linck, Magazin Neuestes, 1790, 32; type not specified.*Rhinobatos* Bloch and Schneider, Syst. Ichth., 1801, 353 (*Rhinobatus rhinobatus*).*Syrrhina* Müller and Henle, Plagiostomen, 1838, 113 (*Rhinobatus columnae= Rhinobatos rhinobatus*).*Glaucostegus* Bonaparte, Catalogo Metodico, 1846, 14 (*Rhinobatus rhinobatus*).

- 96.
- Rhinobatos horkeli*
- (Müller and Henle).

Brazil north to Windward Islands; St. Eustatius.

Rhinobatos horkeli Müller and Henle, Plagiostomen, 1838, 122, pl. 41.

- 97.
- Rhinobatos lentiginosus*
- (Garman).
- Guitarro.*

Atlantic from Charleston, S. C., to Yucatan.

Rhinobatos lentiginosus Garman, Bull. Mus. Comp. Zool., VI, 1880, 168, coast of Florida.

- 98.
- Rhinobatos percellens*
- (Walbaum).
- Fiddlerfish; Puraque.*

West Indies to Jamaica and southern Brazil.

Raja percellens Walbaum, Artedi Pisc., 1792, 525, Brazil; after *Puraque* Marcgrave.*Rhinobatos electricus* Bloch and Schneider, Syst. Ichth., 1801, 356, Brazil.*Rhinobatos undulatus* Ölfers, Torpedo, 1831, 22, Brazil.*Rhinobatos glaucoinctus* Ölfers, loc. cit., 1831, 22, Brazil.*Rhinobatos marcgravii* Henle, Ueber Narcine, 1834, 34, Brazil.

- 99.
- Rhinobatos stellio*
- (Jordan and Rutter).

West Indies.

Rhinobatos stellio Jordan and Rutter, Proc., Acad. Nat. Sci. Phila., XLIX, 1897, 91, Kingston, Jamaica.¹² The African species, *P. perrotteti* Valenciennes, probably does not occur in America.

100. *Rhinobatos spinosus* (Günther).

Mexico.

Rhinobatos spinosus Günther, Cat., VIII, 1870, 518, Mexico.101. *Rhinobatos productus* (Ayres). *Guitarfish*.

San Francisco to San Diego.

Rhinobatos productus Ayres, in Girard, Proc., Acad. Nat. Sci. Phila., VII, 1854, 196, Presidio of Monterey, Calif.102. *Rhinobatos glaucostigma* (Jordan and Gilbert). *Guitarro*.

Gulf of California and southward.

Rhinobatos glaucostigma Jordan and Gilbert, Proc. U. S. Nat. Mus., VI, 1883, 210, Mazatlan.103. *Rhinobatos leucorhynchus* (Günther).

Panama and vicinity.

Rhinobatos leucorhynchus Günther, Proc. Zool. Soc. Lond., 1866, 604, Panama.Genus 54. *ZAPTERYX* Jordan and Gilbert.*Zapteryx* Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 53 (*Platyrrhina exasperata* Jordan and Gilbert).104. *Zapteryx exasperatus* (Jordan and Gilbert).

San Diego Bay; abundant.

Platyrrhina exasperata Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 32, San Diego.*Trigonorrhina alveata* Garman, Bull. Mus. Comp. Zool., VI, 1880, 169, San Diego.105. *Zapteryx xyster* Jordan and Evermann.

Panama.

Zapteryx xyster Jordan and Evermann, Fishes North and Mid. Amer., 1896, 65, Panama.

Family 24. PLATYRHINIDÆ

Genus 55. PLATYRHINOIDIS Garman.

Platyrrhinoidis Garman, Proc. U. S. Nat. Mus., III, 1880, 522 (*Platyrrhina triseriata* Jordan and Gilbert).106. *Platyrrhinoidis triseriatus* (Jordan and Gilbert).

Coast of California, from Point Conception southward.

Platyrrhina triseriata Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 36, Santa Barbara, Calif.

Family 25. RAJIDÆ. Skates and rays

Genus 56. RAJA Linnaeus.

Raja (Artedi) Linnaeus, Syst. Nat., ed. X, 1758, 231 (*Raja batis* Linnaeus).*Dipturus* Rafinesque, Caratteri, 1810, 16 (*Raja batis* L.).*Platopterus* Rafinesque, Anal. Nat. 1815, 93 (*Raja batis* L.).*Dasybatus* Blainville, Jour. Phys. 1816, 260 (*communis*).*Propterygia* Otto, Nova Acta Acad. Cæs. Leop. Carol. Nat. Curios., 1824, 777 (*hyposticta*; monstrous example with fins not adnate to head).*Læviraja* Bonaparte, Faun. Ital., XXV, 1839, 130 (*oxyrhynchus*).*Hieroptera* Fleming, Edinburgh New Philos. Jour., XXVI, 1841, 236 (*H. abredonensis* Flem.=*R. clavata* L.).*Batis* Bonaparte, Cat. Metod., 1846, 12 (*radula*; no description).*Raia* of various authors; merely change of spelling.*Amblyraja* Malm, Goteborg og Bohusian's Fauna, 1877, 607 (*Raia circularis* Malm=*Raja radiata* Donovan).*Lencoraja* Malm, loc. cit., 609 (*Raja fullonica* L.).*Alpharaia* Leigh-Sharpe,¹⁴ Jour. Morph., XXXIX, 1924, 568 (*Raja circularis* Couch).

¹⁴ Mr. Leigh-Sharpe (Jour. Morphology, XXXIX, December, 1924) divides the genus *Raja* into nine "pseudogenera" on the morphology of the genital organs. To settle the question of the standing of pseudogenera, Jordan (in Copeia, 1925) accepts all as subgenera. Most of those not named above (*Epsilonoraja*, *Zdaria*, *Elaria*, *Thetaria*, and *Iotaria*), outside of our limits, may prove valid genera.

- Betaria* Leigh-Sharpe, loc. cit., 568 (*Raja clavata* Linnaeus, a synonym of *Heiroptera* Fleming).
- Gammaria* Leigh-Sharpe, loc. cit., 568 (*Raja batis* Linnaeus, a synonym of *Raja* Linnaeus).
- Deltaria* Leigh-Sharpe, loc. cit., 568 (*Raja radiata* Linnaeus, a synonym of *Amblyraja* Malm).
- 107. *Raja lævis* Mitchell.** *Barndoor skate; Winter skate; Peck-nosed skate.*
Massachusetts to Florida, common northward.
Raja lævis Mitchell, Amer. Month. Mag., II, 1817, 327, New York; name perhaps preoccupied by eligible authors (Gronow, 1763; Valmont, 1765; Duhamel, 1782).
Raja stabuliforis Garman, Plagiostomia, Mem. Mus. Comp. Zool., XXXVI, 1913, 341, Pl. XXII, fig. 2, Pl. XLIV, figs. 4-6, New England coast (substitute for *lævis*, regarded as preoccupied, a matter to be questioned).
- 108. *Raja radiata* Donovan.**
Northern Iceland, southward on both coasts.
Raia radiata Donovan, Hist. Brit. Fish., V, 1820, pl. 114, Great Britain.
- 109. *Raja scabrata* Garman.**
Raia scabrata Garman, Plagiostomia, Mem. Mus. Comp. Zool., XXXVI, 1913, 340, pl. 21, pl. 44, figs. 1-3, Massachusetts Bay. Long confused with *Raja radiata*.
- 110. *Raja granulata* Gill.**
Banks of Newfoundland.
Raia granulata Gill, in Goode and Bean, Bull. Essex Inst., XI, 1879, 28, La Have Bank.
- 111. *Raja fyllæ* Lütken.**
Davis Straits, Greenland.
Raja fyllæ Lütken, Vid. Medd. Naturh. Foren. Kjøbenhavn, 1887, 1, pl. 1, Davis Straits.
- 112. *Raja eglanteria* Bosc.** *Skate; Brier ray; Possum ray; Sea possum; Bob-tailed skate; Clear-nosed skate; Summer skate.*
Cape Cod, southward to Florida; rather common.
Raia eglanteria Bosc, in Lacépède, Hist. Nat. Poiss., II, 1800, 104, Charleston, S. C.
Raia desmarestia LeSueur, Jour. Acad. Nat. Sci. Phila., IV, 1824, 100, Florida.
Raia chantenay LeSueur, loc. cit., 106, America.
Raia americana DeKay, N. Y. Fauna: Fishes, 1842, 368, pl. 66, fig. 215, Staten Island.
- 113. *Raja erinacea* Mitchell.** *Hedgehog ray; Common skate; Little skate; Tobaccoz; Prickly skate; Summer skate.*
Nova Scotia to the Carolinas; very abundant from Virginia to Maine.
Raia erinacea Mitchell, Amer. Jour. Sci. Arts, IX, 1825, 290, pl. 6, New York.
- 114. *Raja diaphanes* Mitchell.** *Clear ray.*
East Coast of United States.
Raia ocellata Mitchell, Trans. Lit. Phil. Soc. N. Y., 1, 1803, 877, New York; not of Russell.
Raia diaphanes Mitchell, loc. cit., 478, New York.
- 115. *Raja senta* Garman.**
Banks of Newfoundland to Cape Cod; in deep water.
Raja senta Garman, Proc. U. S. Nat. Mus., VIII, 1885, 43, Cape Cod Bay; La Have Bank.
- 116. *Raja microtrachys* Osburn and Nichols.**
Off west coast of Lower California.
Raja microtrachys Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1917, 142, fig. 1, off Guadalupe Island in deep water.
- 117. *Raja ackleyi* Garman.**
Yucatan Banks.
Raja ackleyi Garman, Bull. Mus. Comp. Zool., VIII, 1881, 234, Yucatan Banks.

- 118. *Raja ornata* Garman.**
Coast of South Carolina and Florida.
Raja ornata Garman, Bull. Mus. Comp. Zool., VIII, 1881, 235, off
 South Carolina.
- 119. *Raja plutonia* Garman.**
Deep water off South Carolina.
Raja plutonia Garman, Bull. Mus. Comp. Zool., VIII, 1881, 236, off
 South Carolina.
- 120. *Raja rhina* Jordan and Gilbert.**
Monterey to Alaska; not rare.
Raja rhina Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 251,
 Monterey; San Francisco.
- 121. *Raja parmifera* Bean.**
Common on coast of Alaska; Bristol Bay.
Raja parmifera Bean, Proc. U. S. Nat. Mus., IV, 1881, 157, Unalaska.
- 122. *Raja aleutica* Gilbert.**
Unalaska, Aleutian Islands.
Raja aleutica Gilbert, Rept. U. S. Fish Com., XIX, 1893 (1895), 397,
 pl. 21, Sannak Pass, Aleutian Islands, in 81 fathoms.
- 123. *Raja rosispinis* Gill and Townsend.**
Bering Sea.
Raja rosispinis Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897,
 231, Bering Sea.
Raja obtusa Gill and Townsend, loc. cit., 231, Bering Sea.
- 124. *Raja interrupta* Gill and Townsend.**
Raja interrupta Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897,
 232, Bering Sea.
- 125. *Raja binoculata* Girard. *Big skate of California.***
Pacific coast; abundant from Monterey to Sitka.
Raja binoculata Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 196,
 San Francisco.
Raja cooperi Girard, Pac. R. R. Surv., X, Pt. 4, Fishes, 1858, 372,
 Shoalwater Bay, Wash.
- 126. *Raja stellulata* Jordan and Gilbert.**
Coast of California and northward to Unalaska.
Raja stellulata Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880,
 133, Monterey Bay.
- 127. *Raja inornata* Jordan and Gilbert. *Common skate (California.)***
Coast of California; very abundant from San Diego to Cape Mendocino.
Raja inornata Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880,
 457, San Francisco.
Raja jordani Garman, loc. cit., VIII, 1885, 43, San Francisco.
Raja inornata inermis Jordan and Gilbert, loc. cit., IV, 1881, 74, Santa
 Barbara.
- 128. *Raja montereyensis* Gilbert.**
Monterey Bay.
Raja montereyensis Gilbert, Proc., U. S. Nat. Mus., XLVIII, 1915, 307,
 Monterey Bay off Santa Cruz, Calif., in 26 to 28 fathoms.
- 129. *Raja equatorialis* Jordan and Bollman.**
West coast of Colombia, between Panama and Galapagos Islands.
Raja equatorialis Jordan and Bollman, Proc. U. S. Nat. Mus., XII,
 1889, 150, off Colombia.
- 130. *Raja trachura* Gilbert.**
Santa Barbara Islands; south of Shumagin Islands, Central Alaska.
Raja trachura Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 539, off Santa
 Barbara, at *Albatross* station 2923 in 822 fathoms.
- 131. *Raja abyssicola* Gilbert.**
Queen Charlotte Island, British Columbia, in deep water.
Raja abyssicola Gilbert, Rept. U. S. Fish Com., XIX, 1893 (1895), 396,
 pl. 20, off Queen Charlotte Island, at *Albatross* station 3342 in 1,588
 fathoms.

Genus 57. DACTYLOBATUS Bean and Weed.

Dactylobatus Bean and Weed, Proc. U. S. Nat. Mus., XXVI, 1909, 459
(*D. armatus* Bean and Weed).

132. *Dactylobatus armatus* Bean and Weed.

Deep water off South Carolina.

Dactylobatus armatus Bean and Weed, Proc. U. S. Nat. Mus., XXXVI, 1911, 459, off Charleston in 270 fathoms.

Suborder NARCACIONTES

Family 26. TORPEDINIDÆ. Electric rays

Genus 58. TETRANARCE¹⁵ Gill. Great torpedo.

Tetranarce Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1862, 387 (*Torpedo occidentalis* Storer); misprinted *Tetronarce*.

Gymnotorpedo Fritsch, Arch. Anal. Phys., 1886, 365 (*Torpedo occidentalis* Storer).

133. *Tetranarce occidentalis* (Storer). Electricfish; Crampfish; Torpedo; Numbfish.

Atlantic coast of United States, Cape Cod to Cuba; not very common. *Torpedo occidentalis* Storer, Amer. Jour. Sci. Arts, XLV, 1843, 165, pl. 3, Massachusetts. Garman regards this species as inseparable from *Torpedo nobiliana* Bonaparte of the Mediterranean.

134. *Tetranarce californica* (Ayres). California torpedo; Crampfish.

Coast of California; not recorded from south of Monterey; recently reported from Cape Flattery, Wash.

Torpedo californica Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 74, San Francisco.

Genus 59. NARCINE Henle.

Narcine Henle, Ueber Narcine, 1834, 31 (*Torpedo brasiliensis* von Ölfers).

Syraxis Jourdain, in Bonaparte, Fauna Italica, fasc. 160, 1841 (*Narcine indica* Henle).

Cyclonarce Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 387 (*Raja timlei* Schneider).

Gonianarce Gill, loc. cit., 387 (*Narcine indica* Henle).

135. *Narcine brasiliensis* (Ölfers). Torpedofish; Small electric ray; Trembler. West Indies and Brazil, occasionally northward to Key West and the coast of North Carolina.

Torpedo brasiliensis von Ölfers, Torpedo, 1831, 19, tab. 2, fig. 4, Brazil.

Torpedo bancrofti Griffith, Anim. King., X, 1834, 649, Pl. XXXIV, Jamaica.

Narcine nigra Duméril, Revue Zool., 272, 1852, Brazil.

Torpedo pictus Gronow, Cat. Fishes, 13, 1854, Antilles.

Narcine brasiliensis corallina Garman, Bull. Mus. Comp. Zool., VIII, 1881, 234, Florida.

Narcine umbrosa Jordan, Proc. U. S. Nat. Mus., VII, 1884, 105, Key West.

136. *Narcine entemedor* Jordan and Starks. Entemedor.

Pacific coast of tropical America; Mazatlan, La Paz, Panama.

Narcine entemedor Jordan and Starks, Fishes of Sinaloa, Proc. Calif. Acad. Sci., V, 1895, 387, Mazatlan, Mexico.

Genus 60. DISCOPYGE Heckel.

Discopyge Heckel, in Tschudi, Fauna Peruana, 1846, 32 (*D. tschudii* Tschudi).

137. *Discopyge ommata* Jordan and Gilbert.

Panama; rare.

Discopyge ommata Jordan and Gilbert, Proc. U. S. Nat. Mus., XII, 1889, 151, Panama.

¹⁵ In the Old World genus *Torpedo* Forskål, as then understood, the spiracles are provided with a frieze of filaments, at least in joints. In *Tetranarce* these are absent. The following is the synonymy of *Torpedo*: *Torpedo* Forskål, Fauna Arabica, 1776, 16, in part (*Raja torpedo* L., confused with *Malapterurus electricus*, the electric catfish of the Nile).

Narcacion Klein, Neuer Schauplatz, IV, 1777, 726, nonbinomial (*Raja torpedo* L.).

Fimbriotorpedo Fritsch, loc. cit., 365 (*Torpedo marmorata* Risso = *Raja torpedo* L.).

Suborder MASTICURA. Whip-tailed rays

Family 27. DASYATIDÆ. Stingrays

Genus 61. PASTINACHUS Rüppell.

Pastinachus Rüppell, Atlas, Reise Nord Africa, 1828, 82 (*Raia sephen* Forskål); tail rounded above, with a keel below.

Hypolophus Müller and Henle, Plagiostomen, 1838, 170 (*Raia sephen* Forskål).

138. *Pastinachus centrourus* (Mitchill). *Stingaree; Stingray; Devilfish; Pigeon stingray; Clam-cracker.*
Coast of Maine to Cape Hatteras, abundant; the same or a related species in the Mediterranean.
Raja centroura Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 479, New York.

139. *Pastinachus schmardæ* (Werner).

West Indies.

Dasyatis schmardæ Werner, Zool. Jahrb., XXI, 1904, 298, Jamaica.

140. *Pastinachus torrei* (Garman). *Sabina; Lebisa.*

West Indies.

Dasyatis torrei Garman, Plagiostomia, Mem. Mus. Comp. Zool., XXXVI, 1913, 386, Tunas de Zaza, Cuba.

Genus 62. DASYATIS Rafinesque. *Stingrays*

Dasybatus Klein, Neuer Schauplatz, I, 1775, 191; not binomial.

Dasyatis Rafinesque, Caratteri, 1810, 16 (*Dasyatis ujo* Rafinesque = *Raja pastinaca* L.).

Uroxis Rafinesque, Indice, 1810, 61 (*U. ujo* Rafinesque).

Trigonobatus Blainville, Jour. Phys., 1816, 261 (*T. vulgaris* Blainville = *Pastinaca* L.).

Trygon Adanson, in Cuvier, Règne Anim., ed. I, 1817, 135 (*Raja pastinaca* L.).

Pastinaca Swainson, Class. Anim., 1839, 319 (*Pastinaca olivacea* Swainson).

Pastinaca DeKay, N. Y. Fauna: Fishes, 1842, 372 (*Raja pastinaca* L.).
Dasibatis Garman, corrected spelling.

141. *Dasyatis longus* (Garman).

Gulf of California to Panama: Galapagos.

Trygon longa Garman, Bull. Mus. Comp. Zool., VI, 1880, 170, Acapulco, Mexico.

142. *Dasyatis hastatus* (DeKay). *Whip stingray; Kit; Stingaree.*

West Indies to Brazil, north to Florida, North Carolina and Rhode Island.

Pastinaca hastata DeKay, N. Y. Fauna: Fishes, 1842, 373, pl. 65, fig. 214, Rhode Island.

143. *Dasyatis guttatus* (Schneider).

Surinam to Brazil; recorded from Granada.

Raia guttata Schneider, Syst. Ichth., 1801, 361, no locality given.

Trygon gymnura J. Müller, in Erdmann's Reise um die Erde, 1830, 25, taf. 13, Brazil.

Raja tuberculata Lacépède, Hist. Nat. Poiss., II, 1800, 106, pl. 4, fig. 1, Cayenne; not of Bonnaterre.

Genus 63. AMPHOTISTIUS Garman.

Amphotistius Garman, Plagiostomia, Mem. Mus. Comp. Zool., 1913, 375 (*Trygon sinensis* Steindachner); tail with a distinct fold above and below.

144. *Amphotistius say* (LeSueur). *Southern stingray.*

New Jersey to Brazil, common in Florida; occasionally northward to New York.

Raja say LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 42, New Jersey.

145. *Amphotistius sabinus* (LeSueur).
 Streams and estuaries of Florida; abundant on both coasts.
Trygon sabina LeSueur, Jour. Acad. Nat. Sci. Phila., IV, 1824, 109,
 Florida.
Trygon osteosticta Müller, in Erdmann's Reise um die Erde, 1830, 25,
 pl. 14, Brazil.
146. *Amphotistius dipterurus* (Jordan and Gilbert).
 Bay of San Diego and southward; locally abundant.
Dasybatis dipterurus Jordan and Gilbert, Proc. U. S. Nat. Mus., III,
 1880, 31, San Diego; identified by Garman with *Dasyatis brevis*
 Garman, from Peru.
- Genus 64. **PTEROPLATEA** Müller and Henle.
Pteroplatea Müller and Henle, Plagiostomen, 1838, 168 (*Raja altavela* L.)
147. *Pteroplatea altavela* (Linnaeus).
 Tropical Atlantic, occasionally northward to Mediterranean and in
 Gulf Stream; Beaufort, N. C.; Woods Hole, Mass.; taken at Penzance
 Point in Buzzards Bay (Nichols).
Raja altavela Linnaeus, Syst. Nat., ed. X, 1758, 232, no locality.
Dasyatis attavilla Rafinesque, Indice, 1810, 49, Sicily.
Pteroplatea canariensis Valenciennes, Ichth. Canaries, 1843, 100, pl. 23,
 fig. 9, Canary Islands.
Pteroplatea valenciennei A. Duméril, Elasmobranches, 1865, 612, no
 locality.
Pteroplatea vaillanti Rochebrune, Bull. Sci. Philom., IV, 1880, 159, no
 locality.
148. *Pteroplatea micrura* (Schneider). *Butterfly ray.*
 Rhode Island to Brazil.
Raia micrura Schneider, Syst. Ichth., 1801, 360, Surinam.
Raia macrura LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 41, Rhode
 Island.
149. *Pteroplatea marmorata* Cooper.
 Coast of California, from Point Conception southward to Cedros Island;
 common.
Pteroplatea marmorata Cooper, Proc. Calif. Acad. Nat. Sci., III, 1863,
 112, fig. 25, San Diego.
150. *Pteroplatea crebripunctata* Peters. *Manta raia colorado; Manta raia.*
 Gulf of California, southward along the west coast of Mexico; common.
Pteroplatea crebripunctata Peters, Monatsber. Akad. Wiss. Berlin, 1869,
 703, Mazatlan.
Pteroplatea rava Jordan and Starks, in Jordan, Fishes of Sinaloa, Proc.
 Calif. Acad. Sci., Ser. 2, V, 1895, 390, Mazatlan, Mexico.
- Genus 65. **UROBATIS** Garman.
Urobatis Garman, Plagiostomia, Mem. Mus. Comp. Zool., XXXVI,
 1913, 401 (*Leiobatus sloani* Blainville).
151. *Urobatis sloani* (Blainville). *Maid.*
 West Indies, generally common; occasionally northward; once (perhaps
 doubtfully) recorded from New Jersey.
Leiobatus sloani Blainville, Bull. Sci. Philom., 1816, 121, Jamaica; after
Pastinaca marina Sloan.
Raja jamaicensis Cuvier, Règne Anim., ed. I, 1817, 137, Jamaica.
Trygonobatus torpedinus Desmarest, Prem. Dec. Ichth., I, 1823, 6,
 Cuba.
152. *Urobatis vermiculatus* (Garman).
 West Indies.
Urobatis sloani vermiculatus Garman, Plagiostomia, Mem. Mus. Comp.
 Zool., XXXVI, 1913, 802, Progreso, Yucatan.
153. *Urobatis halleri* (Cooper). *Round stingray.*
 Coast of California, from Point Conception southward.
Urolophus halleri Cooper, Proc. Calif. Acad. Nat. Sci., III, 1863, 95,
 San Diego.

154. *Urobatis umbrifer* (Jordan and Starks).

West coast of Mexico.

Urolophus umbrifer Jordan and Starks, in Jordan, Fishes Sinaloa, Proc. Calif. Acad. Sci., Ser. 2, V, 1895, 389, Mazatlan, Mexico; regarded by Garman as inseparable from *Urobatis halleri*.

155. *Urobatis nebulosus* (Garman).

Gulf of California to Panama.

Urolophus nebulosus Garman, Proc. U. S. Nat. Mus., VIII, 1885, 41, Colima, Mexico.

156. *Urobatis concentricus* Osburn and Nichols.

Gulf of California.

Urobatis concentricus Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 144, fig. 2, Esteban Island, Gulf of California.

Genus 66. *UROTRYGON* Gill.

Urotrygon Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 173 (*Urotrygon mundus* Gill).

157. *Urotrygon mundus* Gill.

West coast of Mexico.

Urotrygon mundus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 173, Panama.

Urolophus asterias Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 579, Mazatlan; Panama.

Urolophus rogersi Jordan and Starks, in Jordan, Fishes of Sinaloa. Proc. Calif. Acad. Sci., Ser. 2, V, 1895, 388, Mazatlan.

158. *Urotrygon goodei* (Jordan and Bollman).

Panama and vicinity.

Urolophus goodei Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 151, near Panama.

159. *Urotrygon aspidurus* (Jordan and Gilbert).

Panama.

Urolophus aspidurus Jordan and Gilbert, Bull. U. S. Fish Com., IV, 1881, 307, Panama.

Family 28. AËTOBATIDÆ. Eagle rays

Genus 67. *STOASODON* Cantor.

Aëtobatis Müller and Henle, Plagiostomen, 1841, 180 (*Raia narinari* Euphrasen); not *Aëtobatus* Blainville.

Stoasodon Cantor, Cat. Malayan Fishes, 1850, 434 (*Raia narinari* Euphrasen).

Goniobatis Agassiz, Proc. Bost. Soc. Nat. Hist., VI, 1859, 385 (*flagellum*).

160. *Stoasodon narinari* (Euphrasen). Spotted stingray; Bishop ray; Obispo; Lady ray.

Tropical seas; north on the Atlantic coast to Virginia; Pacific coast, Gulf of California to Panama.

Raia narinari Euphrasen, Vedensk. Akad. Nya Handl., XI, 1790, 217, Brazil.

Raia flagellum Bloch and Schneider, Syst. Ichth., 1801, 361, Coromandel.

Myliobatis eeltenkee Rüppell, Neu. Wirb., 1835, 70, Red Sea.

Goniobatis macroptera McClelland, Calcutta Jour. Nat. Hist., I, 1841, 60, Bengal.

Aëtobatus laticeps Gill, Ann. Lyc. Nat. Hist. N. Y., VIII, 1865, 137, California.

Genus 68. *PTEROMYLÆUS* Garman.

Pteromylæus Garman, Plagiostomia, Mem. Mus. Comp. Zool., XXXVI, 1913, 437 (*Myliobatis asperrimus* Gilbert).

161. *Pteromylæus asperrimus* (Gilbert).

Panama.

Myliobatis asperrimus Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2754, Panama.

- Genus 69. *AëTOBATUS*¹⁰ Blainville. *Eagle rays.*
Aëtobatus Blainville, Jour. Phys., LXXXIII, 1816, 261 (*Raia aquila* as
Aëtobatus vulgaris Blainville).
Myliobatis Duméril in Cuvier, Règne Anim., ed. I, 1817, 137 (*Raia aquila*).
Holorhinus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 331 (*Rhinoptera vespertilio* Girard = *Myliobatus californicus* Gill).
162. *Aëtobatus fréminvillii* (LeSueur). *Stingaree; Sharp-nosed ray; Long-tailed stingray.*
Cape Cod to Brazil.
Myliobatis freminvillii LeSueur, Jour. Acad. Nat. Sci. Phila., IV, 1824, 111, Rhode Island.
Myliobatis bispinosus Storer, Proc. Bost. Soc. Nat. Hist., I, 1841, 53, Massachusetts.
Myliobatis acuta Ayres, loc. cit., I, 1842, 65, Brookhaven, Long Island, N. Y.
163. *Aëtobatus goodei* (Garman).
Central America to Brazil; Rio Grande do Sul.
Myliobatis goodei Garman, Proc. U. S. Nat. Mus., VIII, 1885, 39, Central America.
164. *Aëtobatus californicus* (Gill). *California stingray; Batfish.*
California from Cape Medocino southward.
Rhinoptera vespertilio Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 137, Tomales Bay; not *Myliobatis vespertilio* Bleeker.
Myliobatis californicus Gill, Ann. Lyc. Nat. Hist. N. Y., VIII, 1865, 137; after Girard.

Family 29. RHINOPTERIDÆ

- Genus 70. *RHINOPTERA* Kuhl.
Rhinoptera Kuhl in Cuvier, Règne Anim. ed. II, II, 1829, 401 (*Myliobatis marginatus* Goeffroy).
Zygotaxis Agassiz, Poiss. Fossiles, III, 1836, 79 (*Z. jussieui* Agassiz).
Mylorhina Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1865, 136 (*Rhinoptera lalandi* Müller and Henle).
Micromesus Gill, loc. cit., 136 (*Rhinoptera adspersa* Müller and Henle).
165. *Rhinoptera bonasus* Mitchell. *Cowfish; Cow-nosed ray.*
Cape Cod to Florida.
Rhinoptera bonasus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 479, New York; description inadequate.
Raia quadriloba LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 44, New Jersey.
166. *Rhinoptera steindachneri* Evermann and Jenkins. *Gabilan.*
Gulf of California.
Rhinoptera steindachneri Evermann and Jenkins, Proc. U. S. Nat. Mus., XIV, 1891, 130, pl. 1, fig. 1, Guaymas, Sonora, Mexico.
167. *Rhinoptera encenadæ* Rosa Smith.
Todos Santos Bay, Lower California.
Rhinoptera encenadæ Rosa Smith, Proc. U. S. Nat. Mus., IX, 1886, 220, Ensenada de Todos Santos, Lower California. Jaw only.

Family 30. MOBULIDÆ. Sea devils

- Genus 71. *MOBULA* Rafinesque.
Mobula Rafinesque, Indice, 1810, 61 (*M. auriculata* = *Squalus edentulus* Brunnich).
Apterurus Rafinesque, Indice, 1810, 62 (*A. fabroni* Rafinesque).
Cephalopterus Riso, Ichth. Nice, 1810, 14 (*Raja giorna* Lacépède); not of Goeffroy St. Hilaire, 1809, a genus of birds.
Dicerobatus Blainville, Bull. Soc. Philom., 1816, 116 (*Raja mobular* Lacépède).
Pterocephala Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 321, (*Raja giorna* Lacépède).

¹⁰ Müller and Henle correctly referred *Aëtobatus* Blainville to the type of *Raia aquila*, spelling the name, however, as *Altobatis*. They then created a new genus, *Aëtobatis* for *Raia narinari*, an arrangement without warrant.

168. *Mobula hypostoma* (Bancroft).
Jamaica.
Cephalopterus hypostomus Bancroft, Proc. Comm. Zool. Soc., 1830, 134,
Jamaica.
169. *Mobula olfersi* (Müller and Henle). *Small devilfish.*
Cape Lookout, N. C., to Brazil.
Cephalopterus olfersi Müller and Henle, Abhandl. Akad. Wiss. Berlin,
1834, 311, Brazil; regarded by Garman as identical with *M. thy-*
postoma.
- Genus 72. **CERATOBATIS** Boulenger.
Ceratobatis Boulenger, Ann. Mag. Nat. Hist., XX, 1897, 227 (*C. robertsi*
Boulenger).
170. *Ceratobatis robertsi* Boulenger.
Jamaica.
Ceratobatis robertsi Boulenger, Ann. Mag. Nat. Hist., XX, 1897, 227,
Jamaica.
- Genus 73. **MANTA** Bancroft.
Manta Bancroft, Zool. Jour., IV, 1828-29, 444 (*Cephalopterus manta*
Bancroft = *Raia birostris* Walbaum, 1792).
Ceratoptera Müller and Henle, Syst. Besch., Plagiostmen, 1838, 186
(*Cephaloptera vampyrus* Mitchell).
Brachioptilon Newman, Zoologist, 1849, 74 (*B. hamiltoni*).
Diabolichthys Holmes, Proc. Elliott Soc. Nat. Hist., I, 1856, 39 (*D. ellioti* Holmes).
171. *Manta birostris* (Walbaum). *Sea devil; Devilfish; Manta; Blanketfish.*
Tropical waters of America; north to New Jersey and San Diego; not
rare on the Florida coast.
Raia birostris Walbaum, Artedi Pisc., 1792, 535, after *Diabolus marinus*
Willughby, etc.
Raia manata Schneider, Syst. Ichth., 1801, 364, tropical America.
Raia imbricata Lacépède, Hist. Nat. Poiss., IV, 1803, 677, Gulf Stream.
Cephalopterus vampyrus Mitchell, Ann. Lyc. Nat. Hist. N. Y., I, 1823,
23, Delaware Bay.
Cephalopterus giorna LeSueur, Jour. Acad. Nat. Sci. Phila., IV, 1824,
115, Georgia.
Cephaloptera manta or *Manta americana* Bancroft, Zool. Jour., IV,
1828, 444, Jamaica.
Cephaloptera johnii Müller and Henle, Syst. Besch. Plagiostomen, 1838,
186, West Indies.
Diabolichthys ellioti Holmes, Proc. Elliott Soc. Nat. Hist., I, 1856, 39,
Charleston, S. C.
172. *Manta hamiltoni* (Newman).
San Diego; Gulf of California to Panama, not rare; not known to be
distinct from *Manta birostris*.
Brachioptilon hamiltoni Newman, Zoologist, 1849, 74, Gulf of California.
- Subclass HOLOCEPHALI. Chimæras**
- Order I. CHIMÆROIDEI. Chimæroids**
- Family 31. CHIMÆRIDÆ. Chimæras**
- Genus 74. **CHIMÆRA** Linnæus. *Elephantfishes.*
Chimæra Linnæus, Syst. Nat., ed. X, 1758, 236 (*C. monstrosa* Linnæus).
173. *Chimæra monstrosa* Linnæus.
Deep waters of the Atlantic; off Portugal and off the American coast
from Cape Cod northward.
Chimæra monstrosa L., Syst. Nat., ed. X, 1758, 236, Atlantic.
Chimæra mediterranea Risso, Europ. Mérid., III, 1826, 168, Nice.
Chimæra cristata Faber, Natur. der Fische Islands, 1829, 45, Iceland.
Callorhynchus centrina and *Callorhynchus atlanticus* Gronow, Cat.
Fishes, 15, 16, 1854.

Genus 75. BATHYALOPEX Collett.

Bathyalopex Collett, Arch. Math. Natur., XXXIII, 1901, 125 (*Chimæra mirabilis* Collett).

174. Bathyalopex plumbeus (Gill).

Gulf Stream in deep water.

Chimæra plumbea Gill, Bull. Phil. Soc. Wash., Dec., 1877, 182, La Have Bank, lat. $42^{\circ} 40' N.$, long. $63^{\circ} 23' W.$, in 350 fathoms.

175. Bathyalopex abbreviatus (Gill).

Chimæra abbreviata Gill, Proc. U. S. Nat. Mus., VI, 1883, 254, at $40^{\circ} N.$, long. $67^{\circ} W.$, in 1,290 fathoms.

Genus 76. PSYCHICHTHYS Fowler.

Psychichthys Fowler, Proc. Acad. Nat. Sci. Phila., LIX, 1907, 419 (*Hydrolagus waitei* Fowler).

176. Psychichthys affinis (Capello).

Open sea, from Cape Cod to Portugal.

Chimæra affinis Capello, Jour. Soc. Math., Lisbon, IV, 1868, 314, Portugal in deep water.

Genus 77. HYDROLAGUS Gill.

Hydrolagus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 331 (*Chimæra colliei* Lay and Bennett).

177. Hydrolagus colliei (Lay and Bennett). Ratfish; Elephantfish.

Pacific coast, Monterey to Alaska; abundant in cold water at no great depth.

Chimæra colliei Lay and Bennett, Beechey's Voy., Zool., 1839, 71, North Pacific.

Family 32. RHINOCHIMÆRIDÆ**Genus 78. HARRIOTTA Goode and Bean.**

Harriotta Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1895, 471 (*H. raleighana* Goode and Bean).

Anteliochimæra Tanaka, Jour. Coll. Sci., Imp. Univ. Tokyo, XXVIII, art. 8, 1909, Japan (*A. chætirhamphus* Tanaka).

178. Harriotta raleighana Goode and Bean.

Deep water off the north Atlantic coast of the United States.

Harriotta raleighana Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894 (Jan. 26, 1895), 472, pl. 19, Gulf Stream at $39^{\circ} N.$, $70^{\circ} W.$.

179. Harriotta curtiss-jamesi Townsend and Nichols.

Deep water off Lower California.

Harriotta curtiss-jamesi Townsend and Nichols, Bull. Amer. Mus. Nat. Hist., LII, 1925, 6, fig. 2, off Lower California, lat. 26° , in 645 fathoms.

Class IV. PISCES. True fishes**Subclass ACTINOPTERI****Order J. GLANIOSTOMI****Family 33. ACIPENSERIDÆ. Sturgeons****Genus 79. ACIPENSER¹⁷ Linnaeus. Sturgeons.**

Acipenser Linnaeus, Syst. Nat., ed. X, 1758, 237 (*A. sturio* L.).

Ichthocolla Geoffroy, Descr. de 719 Plantes, etc., 1767, 399 (presumably *A. huso* L.).

Sterleta Güldenstädt, Nov. Com. Sci. Petropol., XVI, 1772, 533 (*A. ruthenus* L.).

Sturio Rafinesque, Indice, 1810, 41 (*Sturio vulgaris* Rafinesque = *A. sturio* L.).

Sterletus Rafinesque, Ichth. Ohiensis, 1820, 80 (*A. serotinus* Rafinesque = *A. fulvescens* Rafinesque).

¹⁷ This group ultimately may need subdivision into distinct genera, but none of the many proposed is adequately defined.

- Helops* Brandt and Ratzeburg, Medizinische Zoologie, II, 1833, 3, 349
 (*A. stellatus* Pallas = *A. halops* Pallas).
Sterletus Brandt and Ratzeburg, loc. cit., 349 (*A. ruthenus* L.).
Sturio Müller, Anab. Myxinoïden, 1835, 77 (*A. sturio* L.).
Antaceus Heckel and Fitzinger, Ann. Wiener Mus., I, 1836, 293 (*A. schypa* Eichwald).
Lioniscus Heckel and Fitzinger, loc. cit., 370 (*A. glabra* Fitzinger).
Sterletus Bonaparte, Cat. Metod., 1845, 21 (*A. vultharius* L.).
Ellops Gistel, Naturg. Thierreichs, IX, 1848 (*A. helops* Pallas).
Schipa Brandt, Bull. Acad. Sci. Petersb., VII, 1850, 113 (*A. schypa* Eichwald).

- 180. *Acipenser transmontanus* Richardson.** *White sturgeon; Oregon sturgeon; Sacramento sturgeon.*
 Pacific coast of North America, from Monterey to Alaska; once abundant, now nearly extinct; largest of the sturgeons, reaching a weight of 1,000 pounds. A record sturgeon, *Acipenser transmontanus*, taken at Astoria, weighed 1,900 pounds.
Acipenser transmontanus Richardson, Fauna Bor.-Amer., III, 1836, 278, Fort Vancouver.
Acipenser brachyrhynchus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 15, San Francisco.
Acipenser aleutensis Fitzinger and Heckel, Ann. Wien. Mus. Naturg., 1836, Aleutian Islands, after Pallas.
Acipenser caryi Duméril, Nouv. Arch. Mus., III, 1867, 169, San Francisco.
Acipenser ayresi Duméril, loc. cit., 171, San Francisco.
Acipenser putnami Duméril, loc. cit., 178, San Francisco.

- 181. *Acipenser oxyrhynchus* Mitchell.** *Common sturgeon.*
 New York to Carolina, once abundant along the shores; very close to the common sea sturgeon of Europe (*Acipenser sturio* L.), but with fewer lateral plates and rougher ossification; the two perhaps identical.
Acipenser oxyrhynchus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1814, 462, New York.
Acipenser mitchilli Duméril, Nouv. Arch. Mus., III, 1867, New York.
Acipenser kennicotti Duméril, loc. cit., James River.
Acipenser girardi Duméril, loc. cit., Maryland.
Acipenser macrorhinus Duméril, loc. cit., New York.
Acipenser milberti Duméril, loc. cit., New York.
Acipenser bairdi Duméril, loc. cit., Maryland.
Acipenser storeri Duméril, loc. cit., Boston.
Acipenser holbrookii Duméril, loc. cit., Charleston.
Acipenser lecontei Duméril, loc. cit., New York.

- 182. *Acipenser brevirostris* LeSueur.** *Short-nosed sturgeon.*
 Cape Cod to Florida.
Acipenser brevirostrum LeSueur, Trans. Amer. Phil. Soc., I, 1818, 290, Atlantic coast.
Acipenser microrhynchus Duméril, Hist. Nat. Poiss., II, 1870, 105 and 164, New York.
Acipenser lesueuri Valenciennes, in Duméril, loc. cit., 105 and 166, Atlas, pl. 16, figs. 1 and 1a, New York.
Acipenser dekayi Duméril, loc. cit., 105 and 168, New York.

- 183. *Acipenser acutirostris* Ayres.** *Green sturgeon.*
 Pacific coast of America, ascending rivers from Monterey northward; not rare.
Acipenser acutirostris Ayres, Proc., Calif. Acad. Nat. Sci., I, 1854, 14, San Francisco.
Acipenser mediostriatus Ayres, loc. cit., 15, San Francisco.
Acipenser (Antaceus) agassizii Duméril, Nouv. Arch. Mus. Paris, III, 1867, 181, Pl. XI, figs. 2, 2a, 2b, San Francisco.
Acipenser (Antaceus) alexandri Duméril, loc. cit., 183, Pl. XV, figs. 1, 1a, 1b, San Francisco.
Acipenser (Antaceus) oligopeltis Duméril, loc. cit., 184, Pl. XV, figs. 2, 2a, 2b, San Francisco.

184. *Acipenser fulvescens* Rafinesque. *Lake sturgeon; Ohio sturgeon; Red sturgeon.*

Mississippi Valley, Great Lakes and northward, ascending rivers, but not entering the sea; varying much with age.

Acipenser fulvescens Rafinesque, Amer. Month. Mag., I, August, 1817, 288, Great Lakes.

Acipenser rubicundus LeSueur, Trans. Amer. Phil. Soc., I, 1818, 388, Lakes Ontario, Erie, and all the upper lakes.

Acipenser maculosus LeSueur, loc. cit., 393, Ohio River; young.

Acipenser serotinus Rafinesque, Ichth. Ohiensis, 1820, 80, Ohio River.

Acipenser ohiensis Rafinesque, loc. cit., 81, Ohio River.

Acipenser macrostomus Rafinesque, loc. cit., 81, Ohio River.

Dinectus truncatus Rafinesque, loc. cit., 82; erroneous, based on a drawing by Audubon.

Acipenser rupertianus Richardson, Fauna Bor.-Amer., III, 1836, 311, Albany River, Rupert Land.

Acipenser lærvis Agassiz, Lake Superior, 1850, 267; Lake Superior.

Acipenser carbonarius Agassiz, loc. cit., 271, Lake Superior.

Acipenser rhynchæus Agassiz, loc. cit., 276, Lake Superior.

Acipenser liopeltis Günther, Cat. Fishes, VIII, 1870, 341, Mississippi.

Acipenser (Huso) copei Duméril, Hist. Nat. Poiss., 1870, 105 and 108, supposedly upper Missouri.

Acipenser (Huso) rauchi Duméril, loc. cit., 105 and 118, Osage River, Mo.

Acipenser (Huso) richardsoni Duméril, loc. cit., 105 and 120, upper Mississippi.

Acipenser (Huso) anasimos Duméril, loc. cit., 105 and 122, Missouri River at St. Louis.

Acipenser (Huso) paranasimos Duméril, loc. cit., 105 and 124, Huntsville, Ala.

Acipenser (Huso) anthracinus Duméril, loc. cit., 105 and 126, Lake Erie.

Acipenser (Huso) megalaspis Duméril, loc. cit., 105 and 133, Lake Champlain.

Acipenser (Huso) lamarii Duméril, loc. cit., 105 and 139, Mississippi River or Lake Erie.

Acipenser (Huso) atelaspis Duméril, loc. cit., 105 and 141, Saskatchewan River.

Acipenser (Huso) rafinesquii Duméril, loc. cit., 105 and 143, Ohio River.

Acipenser (Huso) rosarium Duméril, loc. cit., 105 and 152, Atlas, pl. 17, figs. 2 and 2a, upper Mississippi or Lake Erie.

Acipenser (Huso) platyrhinus Duméril, loc. cit., 105 and 154, upper Mississippi or Lake Erie.

Acipenser (Huso) kirtlandi Duméril, loc. cit., 105 and 161, Lake Erie.

Acipenser (Huso) nertiniianus Duméril, loc. cit., 105 and 162, Michigan.

Acipenser (Antaceus) cincinnatus Duméril, loc. cit., 220 and 228, Ohio River.

Acipenser (Antaceus) buffalo Duméril, loc. cit., 220 and 231, Lake Erie. Buffalo.

Genus 80. SCAPHIRHYNCHUS Heckel. *Shovelnose sturgeons.*

Scaphirhynchus Heckel, Ann. Wien. Mus. Naturg., I, 1835, 71 (*S. rafinesquei* Heckel = *Acipenser platyrhynchus* Rafinesque); not *Scaphorhynchus* Maximilian, a genus of birds.

Scaphirhynchops Gill, Trans. Amer. Phil. Soc., V, 1863, 12, 178 (*Acipenser platyrhynchus* Rafinesque).

185. *Scaphirhynchus platyrhynchus* (Rafinesque). *Shovelnose sturgeon; Switchtail; Hackleback; Sand sturgeon.*

Mississippi Valley; common in the larger streams.

Acipenser platyrhynchus Rafinesque, Ichth. Ohiensis, 1820, 80, Ohio River.

Genus 81. PARASCAPHIRHYNCHUS Forbes and Richardson.

Parascaphirhynchus Forbes and Richardson, Bull. Ill. Lab. Nat. Hist., VII, 1905, 38 (*P. albus* Forbes and Richardson).

186. *Parascaphirhynchus albus* Forbes and Richardson. *White shovelnose sturgeon; White sturgeon.*

Mississippi River; rare.

Parascaphirhynchus albus Forbes and Richardson, Bull. Ill. Lab. Nat. Hist., VII, 1905, 38, Mississippi River at Grafton, Ill.

Order K. SELACHOSTOMI

Family 34. POLYODONTIDÆ. Paddlefishes

Genus 82. POLYODON Lacépède.

Polyodon Lacépède, Bull., Soc. Philom., 1797, 49 (*P. folium* Lacépède).
Spathularia Shaw, Gen. Zool., V, 1804, 362 (*S. reticulata*=*Squalus spathula* Walbaum).

Platirostra LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817 (1818), 228 (*P. edentula*).

Planirostra Rafinesque, Ichth. Ohiensis, 1820, 83 (*P. edentula*).

Proceros Rafinesque, loc. cit., 87 (*P. maculatus*).

187. *Polyodon spathula* (Walbaum). *Paddlefish; Spoonbill cat; Duckbill cat; Spadefish.*

Mississippi Valley and south in the larger streams, reaching a weight of 173 pounds; rare in Lake Erie.

Squalus spathula Walbaum, Artedi Pisc., 1792, 522; after *Spathula* Rosier, Jour. Phys., 1774, 384, pl. 2; no locality given.

Polyodon feuille Lacépède, Hist. Nat. Poiss., I, 1798, 402, locality unknown.

Polyodon folium Bloch and Schneider, Syst. Ichth., 1801, 457; after Lacépède.

Spatularia reticulata Shaw, Gen. Zool., V, 1804, 362, after Lacépède.

Platirostra edentula LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 227, Ohio River.

Acipenser lagenarius Rafinesque, Ichth. Ohiensis, 1820, 86, Ohio River.

Proceros maculatus Rafinesque, loc. cit., 87, Mississippi River at St. Genevieve, Mo.

Proceros vittatus Rafinesque, loc. cit., 1820, 86, 87, Lake Ontario.

Order L. HOLOSTEI

Suborder GINGLYMODI

Family 35. LEPISOSTEIDÆ. Gar pikes

Genus 83. LEPISOSTEUS Lacépède. *Gar pikes.*

Lepisosteus Lacépède, Hist. Nat. Poiss., V, 1803, 331 (*L. gavialis* Lacépède=*Esox osseus* Linnaeus).

Sarchirus Rafinesque, Jour. Acad. Nat. Sci. Phila., I, 1818, 419 (*S. vittatus* Rafinesque).

Lepidosteus Agassiz; corrected spelling.

188. *Lepisosteus osseus* (Linnaeus). *Long-nosed gar; Billfish; Common gar pike; Ohio gar; Ohio garfish; Billy gar.*

Maryland to Florida and Louisiana, in coastwise streams; also (*huronensis*) Great Lakes and the Mississippi Valley; the most abundant of thegars.

Esox osseus Linnaeus, Syst. Nat., ed. X, 1758, 313, after Artedi; based on *Acus maxima squamosa viridis*, the green gar fish of Catesby, 1738, pl. 30, Virginia.

Lepisosteus gavialis Lacépède, Hist. Nat. Poiss., V, 1803, 333, "lakes and rivers of both Indies."

Lepisosteus longirostris Rafinesque, Ichth. Ohiensis, 1820, 74, Muskingum River, Ohio; head.

Macrognathus loricatus Gronow, Cat. Fishes, 1854, 148, after Linnaeus.

Lepidosteus crassus Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, Bombay Hook, Delaware River.

Lepidosteus louisianensis Duméril, Hist. Nat. Poiss., 1870, 343, New Orleans.

Lepisosteus huronensis Richardson, Fauna Bor.-Amer., III, 1836, 237, Penitanguishene, Lake Huron.

?*Sarchirus vittatus* Rafinesque, loc. cit., 69, Ohio River; young; not identifiable.

?*Sarchirus argenteus* Rafinesque, loc. cit., 86, Ohio River; young; not identifiable.

Lepidosteus semiradiatus Agassiz, Poiss. Fossiles, II, 1836, 2, pl. 2.

- Lepisosteus bison* DeKay, New York Fauna: Fishes, 1842, 271, Buffalo, N. Y.
Lepisosteus lineatus Thompson, Hist. Vermont, 1842, 145, Lake Champlain.
Lepidosteus leptorhynchus Girard, Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 351, Devil River, Tex.
Lepidosteus olarius Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 86 "Platte River near Fort Riley"; but Fort Riley was on the Kansas River.
Lepidosteus milberti Duméril, Hist. Nat. Poiss., 1870, 323 and 328, New York.
Lepidosteus harlani Duméril, loc. cit., 323 and 329, Wabash River.
Lepidosteus smithi Duméril, loc. cit., 323 and 330, upper Mississippi.
Lepidosteus ayresi Duméril, loc. cit., 323, 331, Wabash River.
Lepidosteus copei Duméril, loc. cit., 323, 332, northern North America.
Lepidosteus lesueuri Duméril, loc. cit., 323, 335, Wabash River.
Lepidosteus elizabeth Duméril, loc. cit., 323, 336, no locality.
Lepidosteus troostii Duméril, loc. cit., 323, 339, United States.
Lepidosteus piquitanus Duméril, loc. cit., 323, 340, Lake Erie.
Lepidosteus thompsoni Duméril, loc. cit., 323, 342, Upper Mississippi.
Lepidosteus lamarii Duméril, loc. cit., 323, 337, northern North America.
?Lepidosteus clintonii Duméril, loc. cit., 323, 338, no locality.
Lepidosteus horatii Duméril, loc. cit., 323, 341, northern North America.

Genus 84. CYLINDROSTEUS Rafinesque. *Short-nosed gars.*

Cylindrosteus Rafinesque, Ichth. Ohiensis, 1820, 72 (*Lepisosteus platostomus* Rafinesque).

189. *Cylindrosteus platostomus* (Rafinesque).

Mississippi Valley and Great Lakes; common.

Lepisosteus platostomus Rafinesque, Ichth. Ohiensis, 1820, 72, Ohio River.¹⁸

Lepisosteus oxyurus Rafinesque, loc. cit., 73, Ohio River.

Lepisosteus albus Rafinesque, loc. cit., 73, Ohio River.

Lepidosteus grayi, Agassiz, Poiss. Fossiles, II, 1836, 2, Louisiana.

Lepidosteus (*Cylindrosteus*) *oculatus* Winchell, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 183, Huron River, Mich.

Cylindrosteus zedockti Duméril, Hist. Nat. Poiss., II, 1870, 348, 353, Upper Mississippi.

Cylindrosteus rafinesquii Duméril, loc. cit., 348, 352, United States.

190. *Cylindrosteus scabriceps* Fowler.

Lower Missouri River.

Cylindrosteus scabriceps Fowler, Proc. Acad. Nat. Sci. Phila., LXII, 1910, 607, Pl. XXXVIII, figs. 8, 9, Leavenworth, Kans.; a doubtful species.

191. *Cylindrosteus agassizii* Duméril.

Missouri Basin.

Cylindrosteus agassizii Duméril, Hist. Nat. Poiss., II, 1870, 351, St. Louis.

Cylindrosteus productus Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 86, San Antonio, Tex.

192. *Cylindrosteus castelnauddii* Duméril.

Florida and Louisiana.

Cylindrosteus castelnauddii Duméril, Hist. Nat. Poiss., II, 1870, 355, Lake Lafayette, Florida.

Cylindrosteus bartoni Duméril, loc. cit., 356, New Orleans.

Lepidosteus latirostris Girard, Pac. R. R. Surv., X, pl. 4, Fishes, 1858, 352, Rio Pecos, Tex.

193. *Cylindrosteus megalops* Fowler.

Florida.

Cylindrosteus megalops Fowler, Proc. Acad. Nat. Sci. Phila., LXII, 1910, 609, Pl. XXXVIII, figs. 15, 16, Bayport, Fla.; a doubtful species.

¹⁸ Many other tributary streams mentioned.

Genus 85. ATRACTOSTEUS Rafinesque. *Alligator gars.*

Litholepis Rafinesque, Amer. Month. Mag., III, 1818, 447 (*L. adamantis* Rafinesque), based on a fancy painting by Audubon of an imaginary fish.

Atractosteus Rafinesque, Ichth. Ohiensis, 1820, 72 (*Lepisosteus ferox* Rafinesque).

194. *Atractosteus spatula* (Lacépède). *Mississippi alligator gar.*

Lower Mississippi Valley in the larger streams, reaching a length of 5 feet or more.

Lepisosteus spatula Lacépède, Hist. Nat. Poiss., V, 1803, 333, no locality. *Lepisosteus (Atractosteus) ferox* Rafinesque, Ichth. Ohiensis, 1820, 73. Ohio River.

Lepidosteus (Atractosteus) berlandieri Girard, Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 353, Tamaulipas.

Atractosteus lucius Duméril, Hist. Nat. Poiss., 1870, 364, Tampico, Mexico.

195. *Atractosteus tristechus* (Bloch and Schneider). *Cuban alligator gar; Manjuari.*

Rivers of Cuba.

Esox tristechus Bloch and Schneider, Syst. Ichth., 1801, 395, Cuba, after Manjuari of Para.

Lepidosteus manjuari Poey, Memorias, I, 1860, 273, Cuba.

196. *Atractosteus tropicus* Gill.

Central America; Nicaragua.

Atractosteus tropicus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 172, streams near Panama.

Order M. HALECOMORPHI**Family 36. AMIIDÆ. Bowfins****Genus 86. AMIA Linnaeus.**

Amia Linnaeus, Syst. Nat., ed. XII, 1766, 500 (*A. calva*); not of Gronow, 1763, which is a nonbinomial name for *Apogon*.

Amiatus Rafinesque, Anal. Nat., 1815, 88 (*A. calva*); substitute for *Amia* regarded as too short for symmetry.

197. *Amia calva* Linnaeus. *Bowfin; John A. Grindle; Mudfish; Lawyer; Poisson de marais; Choupique (Louisiana).*

Great Lakes, Mississippi Valley to Virginia, Florida, and Texas, in lakes and sluggish streams.

Amia calva Linnaeus, Syst. Nat., ed. XII, 1766, 500, Charleston, S. C. *Amia ocellicauda* Richardson, Fauna Bor.-Amer., III, 1836, 236, Lake Huron, male.

Amia occidentalis DeKay, New York Fauna: Fishes, 1842, 269, St. Marys River, Mich.

Amia marmorata Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 412, New Orleans.

Amia ornata LeSueur, in Cuvier and Valenciennes, loc. cit., 420, Chibault, Mississippi River.

Amia viridis LeSueur, loc. cit., 421, New Orleans.

Amia canina Cuvier and Valenciennes, loc. cit., 424, Lake Erie; after Kirtland.

Amia lentiginosa Cuvier and Valenciennes, loc. cit., 426; after Bonnaterre; no locality.

Amia subcærulea Cuvier and Valenciennes, loc. cit., 427, New Orleans.

Amia cinerea Cuvier and Valenciennes, loc. cit., 430, Charleston, S. C.

Amia reticulata LeSueur, loc. cit., 431, Wabash River.

Amia thompsoni Duméril, loc. cit., II, 1870, 419, Lake Champlain.

Amia piquotii Duméril, loc. cit., 1870, 423, upper Mississippi.

Superorder TELEOSTEI. Bony fishes

Order N. ISOSPONDYLI (Physostomi; Malacoptyerigii). Soft-rayed fishes

Suborder ELOPOIDEA

Family 37. ELOPIDÆ. Tenpounders

Genus 87. ELOPS Linnaeus.

Elops Linnaeus, Syst. Nat., ed. XII, 1766, 518 (*E. saurus*).

Mugilomorus Lacépède, Hist. Nat. Poiss., V, 1803, 398 (*M. anna-carolina*).

Trichonotus Rafinesque, Anal. Nat., 1815, 88 (*Mugilomorus anna-carolina*); substitute for *Mugilomorus*, considered objectionable.

198. *Elops saurus* Linnaeus. Tenpounder; John Mariggle; Bonyfish; Bonefish; Big-eyed herring; Matajuelo real; Lisa francesa.

Tropical Atlantic; West Indies, north to Long Island; common.

Elops saurus Linnaeus, Syst. Nat., ed. XII, 1766, 518, Carolinas.

Argentina carolina Linnaeus, loc. cit., 519, Carolinas.

Mugilomorus anna-carolina Lacépède, Hist. Nat. Poiss., V, 1803, 398, South Carolina.

Elops inermis Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 445, New York.

199. *Elops hawaiiensis* Regan. Awa.

Hawaiian and Pacific coast of Mexico north to Gulf of California; Hawaii.

Elops hawaiiensis Regan, Ann. Mag. Nat. Hist., 9th ser., III, 1909, 39, Hawaii.

Family 38. MEGALOPIDÆ. Tarpons

Genus 88. TARPON Jordan and Evermann. Grande écaille.

Tarpon Jordan and Evermann, Fishes North and Mid. Amer., 1896, 409 (*Megalops atlanticus* Cuvier and Valenciennes).

200. *Tarpon atlanticus* (Cuvier and Valenciennes). Tarpon; Tarpum; Grande écaille; Silverfish; Sabalo; Savanilla.

Long Island to Brazil; abundant off Florida, breeding in the West Indies.

Megalops atlanticus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 398, Guadeloupe; San Domingo; Martinique; Porto Rico.

Megalops elongatus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 224, Long Island.

Suborder ALBULOIDEI

Family 39. ALBULIDÆ. Ladyfishes

Genus 89. ALBULA (Gronow) Bloch and Schneider. Ladyfishes.

Conorhynchus Nozeman, Act. Select., III, 1757, 382; non-binomial.

Albula Gronow, Zoophyl., 1763, 102; non-binomial.

Albula Bloch and Schneider, Syst. Ichth., 1801, 432 (*A. conorrhynchus* = *Esox vulpes* L.).

Butyrinus Lacépède, Hist. Nat. Poiss., V, 1803, 45 (*B. banana* = *Esox vulpes* L.).

Glossodus Cuvier, in Agassiz, Spix, Pisc. Brasil., 1829, 48 (*A. forskali* = *Esox vulpes* L.).

201. *Albula vulpes* (Linnaeus). Ladyfish; Bonefish; Macabi; Sanducha; Bananafish.

Tropical seas, ranging northward to San Diego; Long Island and southern Japan; perhaps more than one species included.

Unbarana Margrave, Hist. Brazil, 1648, Brazil.

Vulpes bahamensis Catesby, Nat. Hist. Carolinas, etc., 1737, Pl. II, fig. 1, Bahamas; non-binomial.

Esox vulpes Linnaeus, Syst. Nat., ed. X, 1758, 313, Bahamas; based on the bonefish, *Vulpes bahamensis* of Catesby.

- Argentina glossodonta* Forskål, Descr. Anim., 1775, 68, Djidda, Arabia.
Macabi Parra, Diff. Piezas Cuba, 1787, 88, pl. 35, fig. 1, Cuba; based on *Unbarana* of Marcgrave.
Synodus argenteus Bloch and Schneider, Syst. Ichth., 1801, 398, Asia.
Clupea brasiliensis Bloch and Schneider, loc. cit., 427, Brazil.
Albula conorhynchus Bloch and Schneider, loc. cit., 432, Antilles; after Gronow and Plumier; called *Albula plumieri* on plate 86.
Amia immaculata Bloch and Schneider, loc. cit., Central America; after *Macabi* of Parra.
Butyrinus banana Lacépède, Hist. Nat. Poiss., V, 1803, 46, Ile de France.
Clupea macrocephala Lacépède, loc. cit., 426, Martinique; on a drawing by Plumier.
Glossodus forskldli Agassiz, in Spix, Pisc. Brasil., 1829, 49, Bahia; called *Engraulis sexicus* and *Engraulis bahiensis* on plates 22 and 24.
Albula parræ Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 339, Martinique; Bahia; Rio de Janeiro.
Albula goreensis Cuvier and Valenciennes, loc. cit., 342, Gorea.
Albula neoguinaica Cuvier and Valenciennes, loc. cit., 350, New Guinea.
Albula seminuda Cuvier and Valenciennes, loc. cit., 351, New Guinea.
Albula erythrocheilus Cuvier and Valenciennes, loc. cit., 352, Friendly Islands.
Albula forsteri Cuvier and Valenciennes, loc. cit., 354, Tahiti.
Albula rostrata Gronow, Cat. Fishes, 1854, 189, American Ocean.

Genus 90. DIXONINA Fowler.

Dixonina Fowler, Proc. Acad. Nat. Sci. Phila., LXII, 1910, 651 (*D. nemoptera* Fowler).

202. *Dixonina nemoptera* Fowler.

San Domingo, West Indies.

Dixonina nemoptera Fowler, Proc. Acad. Nat. Sci. Phila., LXII, 1910, 652, San Domingo.

Suborder CLUPEOIDEI

Family 40. CHANIDÆ. Milkfishes

Genus 91. CHANOS Lacépède.

Chanos Lacépède, Hist. Nat. Poiss., V, 1803, 395 (*C. arabicus*).
Lutodeira Van Hasselt, in Rüppell, Atlas Reise Nord Africa, 1828, 17 (*L. indica* Rüppell = *Mugil chanos* L.)
Ptycholepis Gray, in Dieffenbach's Travels in New Zealand, II, about 1842, 218 (*Mugil salmoneus* Forster).

203. *Chanos cyprinella* Cuvier and Valenciennes. Milkfish; *Sabalo*; *Awa*; *Chani*; *Anged*.

Gulf of California to Panama and Hawaii; replaced in the western Pacific by *Chanos chanos* Forskål.

Chanos cyprinella Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 194, 198, Hawaiian Islands.

Family 41. HIODONTIDÆ. Mooneyes

Genus 92. HIODON LeSueur. Mooneyes.

Hiodon LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 334 (*H. tergisus* LeSueur).

Glossodon Rafinesque, Amer. Month. Mag., II, September, 1818, 354 (*G. harenoides*).

Hydon, corrected orthography.

204. *Hiodon tergisus* LeSueur. Mooneye; Toothed herring.

Great Lakes and Mississippi Valley, north to Assiniboine River.

Hiodon tergisus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 364, Ohio River.

Glossodon harenoides Rafinesque, Amer. Month. Mag., II, September, 1818, 354, Ohio River.

Glossodon heterurus Rafinesque, loc. cit., 354, Falls of Ohio River.

Hiodon clodalis LeSueur, loc. cit., Ohio River at Pittsburgh.

Hydon vernalis Rafinesque, Ichth. Ohiensis, 1820, 42, Ohio River.

- 205. *Hiodon selenops* Jordan and Bean.** *Southern mooneye; Southern toothed herring.*

Tennessee, Cumberland, and Alabama Rivers.

Hyodon selenops Jordan and Bean, Bull. U. S. Nat. Mus., X, 1877, 67, Chattanooga, Tenn.

Genus 93. AMPHIODON Rafinesque. *Goldeyes.*

Amphiodon Rafinesque, Jour. Phys., Paris, 1819, 421, Ohio River (*A. alosoides*; misprinted *alveoides*).

Elattonistius Gill and Jordan, Bull. U. S. Nat. Mus., X, 1877, 68 (*Hyodon chrysopsis* Richardson = *Amphiodon alosoides* Rafinesque).

- 206. *Amphiodon alosoides* Rafinesque.** *Goldeye; La Quesche; Naccaysh.*

Ohio River and north to Saskatchewan; abundant and much valued in Manitoba.

Amphiodon alosoides (misprinted *alveoides*) Rafinesque, Jour. Phys., Paris, 1819, 421, Ohio River.

Hyodon amphiodon Rafinesque, Ichth. Ohiensis, 1820, 42, Ohio River near the falls.

Hyodon chrysopsis Richardson, Fauna Bor.-Amer., III, 1836, 232, Cumberland House, Saskatchewan River.

Cyprinus (Abramis) smithii Richardson, loc. cit., 110, Richelieu River.

Family 42. CLUPEIDÆ. HERRINGS

Genus 94. CLUPEA Linnaeus. *Herrings.*

Clupea Linnaeus, Syst. Nat., ed. X, 1758, 317 (*C. harengus* L.).

Rogenia Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 340 (*R. alba*, "the whitebait," the young of *harengus*).

- 207. *Clupea harengus* Linnaeus.** *Common herring.*

North Atlantic Ocean, south on our coast to Sandy Hook; Massachusetts; New Hampshire.

Clupea harengus Linnaeus, Syst. Nat., ed. X, 1758, 317, seas of Europe.

Clupea membras Pallas, Zoogr. Rosso.-Asiat., III, 1811, 211, Baltic Sea.

Clupea halec Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 451, New York.

Clupea vittata Mitchell, loc. cit., 456, New York.

Clupea cerulea Mitchell, loc. cit., 457, New York.

Clupea elongata LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 234, Massachusetts.

Clupea leachi Yarrell, Zool. Jour., V, 1829, 277, pl. 12, England.

Clupea minima Peck, in Storer, Rept. Fish. Mass., 1839, 113, New Hampshire.

Clupea lineolata Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 256, locality unknown.

Rogenia alba Cuvier and Valenciennes, loc. cit., 340, England; the young form known as whitebait.

- 208. *Clupea pallasii* Cuvier and Valenciennes.** *California herring.*

Pacific coast, from Kamchatka to San Diego.

Clupea harengus var. *Pallas*, Zoogr. Rosso.-Asiat., III, 1811, 209, Kamchatka.

Clupea pallasii Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 253, Kamchatka; based on Pallas's specimens.

Clupea mirabilis Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 138, San Francisco.

Spratelloides bryoporus Cope, Proc. Amer. Phil. Soc., XIII, 1873, 25, Sitka; young.

Genus 95. POMOLOBUS Rafinesque. *Alewives.*

Pomolobus Rafinesque, Ichth. Ohiensis, 38, 1820 (*P. chryschloris* Rafinesque).

Alausella Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 36 (*Clupea parvula* Mitchell = *Clupea mediocris* Mitchell).

Subgenus ALAUSELLA Gill.

- 209. *Pomolobus pseudoharengus* (Wilson).** *Alewife; Branch herring; Gas-pereau; Wall-eyed herring; Big-eyed herring; Sawbelly; Ellwife; Bang.* Atlantic coast of the United States; Lake Ontario and landlocked in the "Finger Lakes" of New York.
Clupea pseudoharengus Wilson, in Rees's Encycl., IX, about 1811, probably Philadelphia.
Clupea vernalis Mitchell, Rept. Fish. N. Y., 1814, 22, and Trans., Lit. Phil. Soc. N. Y., I, 1815, 454, New York.
Clupea megalops Rafinesque, Amer. Month. Mag., 1818, 206, Delaware River.
Meletta venosa Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 374, New York.
Pomolobus pseudoharengus lacustris Jordan, Man. Vert., ed. I, 1876, 265, Cayuga Lake, New York; specimens landlocked.
- 210. *Pomolobus mediocris* (Mitchill).** *Hickory shad; Hickory jack; Fall herring; Tailor herring; Mattowacca.* Atlantic coast of United States from Maine to Florida.
Clupea mediocris Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 450, New York.
Clupea mattowacca Mitchell, loc. cit., 451, pl. 5, fig. 8, New York.
? *Clupea parvula* Mitchell, loc. cit., 452, New York.
? *Clupea pusilla* Mitchell, loc. cit., 452, New York.
Clupea fasciata LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 233, Massachusetts.
Clupea virescens DeKay, New York Fauna: Fishes, 1842, 242, New York.
Alosa lineata Storer, Proc., Bost. Soc. Nat. Hist., II, 1847, 242, Massachusetts.
- 211. *Pomolobus aestivalis* (Mitchill).** *Glut herring; Blueback; Black-belly; Summer herring; Kyack; Sawbelly.* Atlantic coast.
Clupea aestivalis Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 456, New York.
Alosa cyanonotum Storer, Hist. Fish. Mass., 1867, 161, Provincetown, Mass.
- Subgenus POMOLOBUS Rafinesque.**
- 212. *Pomolobus chrysocloris* Rafinesque.** *Skipjack; Blue herring; Nailrod.* Gulf of Mexico and Mississippi Valley; ascending streams, and introduced through the canals into Lake Erie and Lake Michigan.
Pomolobus chrysocloris Rafinesque, Ichth. Ohiensis, 1820, 38, Ohio River.
Meletta sueurii Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 375, Wabash River.
- Genus 96. ALOSA Linck.** *Shad.*
Alosa Linck, Mag. Neuestes Phys. Naturgesch., 1790, 35 (scarcely defined; type not specified; *Clupea alosa* L. = *Alosa* Cuvier).
Alosa Cuvier, Régne Anim., ed. II, 1829, 319 (*Clupea alosa* L.).
Alausa Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 389 (*Clupea alosa* L.).
- 213. *Alosa sapidissima* (Wilson).** *Common shad; American shad; North River shad; Potomac shad; Connecticut River shad.* Atlantic coast of United States, from Miramichi River to Florida; from Monterey northward on the Pacific coast, as an introduced species.
Clupea sapidissima Wilson, in Rees's New Cyclopaedia, IXG., neither pagination nor date, but prior to 1812; no locality, but probably Philadelphia.
Clupea indigena Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 454, New York.
Alosa praestabilis DeKay, New York Fauna: Fishes, 1842, 255, pl. 15, fig. 4, New York.
- 214. *Alosa ohioensis* Evermann.** *Ohio shad.* Mississippi River basin; Ohio River.
Alosa ohioensis Evermann, Rept. U. S. Fish Com., XXVII, 1901 (1902), 277, fig. 2, Ohio River at the falls.

215. *Alosa alabamae* Jordan and Evermann. *Alabama shad.*
 Gulf coast of United States; ascending streams.
Alosa alabamae Jordan and Evermann, Rept. U. S. Fish Com., XXI,
 1895 (1896), 203, Black Warrior River, Tuscaloosa, Ala.
- Genus 97. **SARDINIA** Poey. *True sardines.*
Sardinia Poey, Memorias, II, 1861, 311 (*S. pseudohispanica* Poey; not
 of Jordan and Evermann, which is *Sardinella anchovia* Cuvier and
 Valenciennes).
Sardina Antipa, Denkschr. Akad. Wiss. Wien, LXVIII, 1905 (*Clupea*
pilchardus L.).
216. *Sardinia cœrulea* (Girard). *California sardine.*
 Pacific coast, from Puget Sound to Magdalena Bay; very abundant.
Meletta cœrulea Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 138,
 San Francisco.
Alausa californica Gill, loc. cit., XIV, 1862, 281, California.
217. *Sardinia anchovia* (Cuvier and Valenciennes). *Sardinia de España.*
 West Indies, north to Cuba.
Sardinella anchovia Cuvier and Valenciennes, Hist. Nat. Poiss., XX,
 1847, 269, Rio Janeiro; Martinique.
*Sardina*¹⁹ *pseudohispanica* Poey, Memorias, II, 1861, 311, Cuba.
- Genus 98. **SARDINELLA** Cuvier and Valenciennes. *Smooth-headed sar-*
dines.
Sardinella Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 261,
 (*S. aurita* Cuvier and Valenciennes).
218. *Sardinella aurita* Cuvier and Valenciennes. *Smooth sardine;* *Sardina*
de España.
 Woods Hole to Florida; recorded from France; usually in rather deep
 water.
Sardinella aurita Cuvier and Valenciennes, Hist. Nat. Poiss., XX,
 1847, 261, coast of France.
- Genus 99. **HARENGLA** Cuvier and Valenciennes. *Scaled sardines.*
Harengula Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 280
 (*H. latula* Cuvier and Valenciennes).
219. *Harengula sardina* Poey. *Sardina de Ley.*
 West Indian fauna, north to Key West.
Harengula sardina Poey, Memorias, II, 1860, 310, Cuba.
Harengula callolepis Goode, Proc. U. S. Nat. Mus., II, 1879, 152,
 Bermuda.
220. *Harengula macrophthalmus* (Ranzani). *Sardina Escamuda;* *White-bill;*
Pincho.
 West Indies, Florida to Brazil.
Clupea macrophalma Ranzani, Nov. Com. Acad. Sci. Bonon., V, 1842,
 320, Brazil.
Harengula humeralis Cuvier and Valenciennes, Hist. Nat. Poiss., XX,
 1847, 293, Rio Janeiro; Bahia; Guadeloupe; San Domingo.
Alausa striata Cuvier and Valenciennes, loc. cit.; XX, 1847, 429,
 Guadeloupe; Bahia.
Alosa bishopi Müller and Troschel, in Schomburgk, Hist. Barbados,
 1848, 675, Barbados.
221. *Harengula clupeola* Cuvier and Valenciennes. *Cailleu.*
 West Indies.
Harengula clupeola Cuvier and Valenciennes, Hist. Nat. Poiss., XX,
 1847, 289, Guadeloupe Island, West Indies.
222. *Harengula maculosa* Cuvier and Valenciennes.
 West Indies.
Harengula maculosa Cuvier and Valenciennes, Hist. Nat. Poiss., XX,
 1847, 292, Martinique.
Alosa apicalis Müller and Troschel, in Schomburgk, Hist. Barbados,
 1848, 675, Barbados.

¹⁹ Recent examination of Poey's type by Alvin Seale shows that this fish is a *Sardinia* apparently identical with *Clupea anchoria* Cuvier and Valenciennes, not the *Sardinella pseudohispanica* of Jordan and Evermann, which apparently is the same as *Sardinella aurita* Cuvier and Valenciennes.

223. *Harengula pensacolæ* Goode and Bean.
Florida.
Harengula pensacolæ Goode and Bean, Proc. U. S. Nat. Mus., II, 1879,
152, Pensacola, Fla.
224. *Harengula thrissina* (Jordan and Gilbert).
Gulf of California.
Clupea thrissina Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882,
353, Cape San Lucas.
- Genus 100. *LILE* Jordan and Evermann.
Lile Jordan and Evermann, Fishes North and Mid. Amer., 1896, 431
(*Clupea stolifera* Jordan and Gilbert).
225. *Lile stolifera* (Jordan and Gilbert).
Gulf of California to Panama; Mazatlan, Mexico.
Clupea stolifera Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881,
339, Mazatlan.
- Genus 101. *OPISTHONEMA* Gill. *Thread herring*.
Opisthonema Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 37 (*Clupea*
thrissa Broussonet = *Megalops oglina* LeSueur); not of Osbeck.
226. *Opisthonema oglinum* (LeSueur). *Thread herring; Machuelo; Cailleu*
tassart; Sprat.
West Indies, ranging regularly north to Florida and Carolina; Longport,
N. J.; Newport, R. I.; Fortress Monroe.
Clupea thrissa Broussonet, Ichthyologie, fasc. I, 1782, Carolina; Jamaica;
not of Osbeck, 1757, which is a Chinese species of *Clupanodon*.
Megalops oglina LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 359,
Newport, R. I.
Megalops notata LeSueur, loc. cit., 359, Guadeloupe.
Chatoëssus signifer DeKay, N. Y. Fauna: Fishes, 1842, 264, New York.
Chatoëssus eumorpha Gosse, Naturalist's Sojourn in Jamaica, 1851,
290, Jamaica.
227. *Opisthonema libertate* (Günther). *Sardina machete*.
Pacific coast of Mexico and Central America.
Meletta libertate Günther, Proc. Zool. Soc. Lond., 1866, 603, Libertad,
Central America.
- Genus 102. *BREVOORTIA* Gill. *Menhadens*.
Brevoortia Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 37 (*Alosa*
menhaden Storer = *Clupea tyrannus* Latrobe).
228. *Brevoortia tyrannus* (Latrobe). *Menhaden; Mossbunker; Bonyfish;*
Whitefish; Bugfish; Fatback; Pogy.
Nova Scotia to West Indies; abundant.
Clupea tyrannus Latrobe, Trans. Amer. Phil. Soc. Phila., V, 1802, 77,
pl. 1, Chesapeake Bay.
Clupea menhaden Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 453,
New York.
Clupea neglecta Rafinesque, Amer. Month. Mag., 1818, 208, Long Island.
Clupea carolinensis Gronow, Cat. Fishes, 1854, 140, South Carolina.
229. *Brevoortia tyrannus brevicaudata* Goode. *Short-tailed menhaden*.
South Atlantic north to Connecticut.
Brevoortia tyrannus brevicaudata Goode, Proc. U. S. Nat. Mus., I, 1878,
34, Noank, Conn.
230. *Brevoortia patronus* Goode. *Gulf menhaden*.
Gulf of Mexico; coast of Texas.
Brevoortia patronus Goode, Proc. U. S. Nat. Mus., I, 1878, 39, Brazos
Santiago, Tex.
- Genus 103. *ILISHA* Gray.
Platygaster Swainson, Class. Anim., II, 1839, 294 (*Clupea africana*
Bloch); name preoccupied.
Ilisha Gray, in Richardson, Ichthy. China, in Proc. Brit. Assoc., 1845
(1846), 306 (*I. abnormis*); no description.
Pellona Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 300
(*P. orbignyanæ* = *Pristigaster flavipinnis* Valenciennes).

- 281. *Ilisha caribbaea* Meek and Hildebrand.**
Caribbean Sea.
Ilisha caribbaea Meek and Hildebrand, Marine Fishes of Panama, Pt. 1, Field Mus. Nat. Hist., Zool., XV, Dec. 20, 1923, 191, Pl. X, fig. 1, Porto Bello, Panama.
- Genus 104. CHIROCENTRODON Günther.**
Chirocentrodon Günther, Cat., VII, 1868, 463 (*C. tenuatus* Günther).
Medipellona Jordan and Seale, Mem. Mus. Comp. Zool., LXVII, No. 11, May, 1926, 417 (*Pellona bleekeriana* Poey).
- 282. Chirocentrodon tenuatus Günther.**
Jamaica.
Chirocentrodon tenuatus Günther, Cat., VII, 1868, 463, Jamaica.
- 283. Chirocentrodon bleekeriana (Poey).** *Anchoa pelada.*
Matanzas, Cuba.
Pellona bleekeriana Poey, Repertorio, II, 1867, 242, Matanzas, Cuba.
- Genus 105. OPISTHOPTERUS Gill.**
Opisthopterus Gill, Proc. Acad. Nat. Sci. Phila., XII, 1861, 38 (*Pristigaster tartoon* Cuvier and Valenciennes).
- 284. Opisthopterus dovii (Günther).**
Panama.
Pristigaster dovii Günther, Cat., VII, 1861, 461, Panama.
- 285. Opisthopterus macrops (Günther).**
Panama.
Pristigaster macrops Günther, Proc. Zool. Soc. Lond., 1866, 603, Panama.
- Genus 106. PLIOSTEOSTOMA Norman.**
Pliosteostoma Norman, Ann. Mag. Nat. Hist., XI, 1923, 21 (*Pristigaster lutipinnis* Jordan and Gilbert).
- 286. Pliosteostoma lutipinnis (Jordan and Gilbert).**
Pacific coast of Mexico.
Pristigaster lutipinnis Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 340, Mazatlan, Mexico.
- Genus 107. ODONTOGNATHUS Lacépède.**
Odontognathus Lacépède, Hist. Nat. Poiss., II, 1799, 221 (*O. mucronatus*).
Gnathobolus Bloch and Schneider, Syst. Ichth., 1801, 556 (*Odontognathus mucronatus* Lacépède.).
- 287. Odontognathus mucronatus Lacépède.**
Coast of Guiana.
Odontognathus mucronatus Lacépède, Hist. Nat. Poiss., 1799, 221, pl. 7, fig. 2, Cayenne.
- 288. Odontognathus panamensis (Steindachner).**
Panama.
Pristigaster (Odontognathus) panamensis Steindachner, Ichth. Beitr., V, 1876, 24, Panama.
- Genus 108. PRISTIGASTER Cuvier.**
Pristigaster Cuvier, Règne Anim., ed. I, 1817, 176 (type not named; later fixed as *P. cayanus* Cuvier).
- 289. Pristigaster cayanus Cuvier.**
Coast of Guiana and northern Brazil.
Pristigaster cayanus Cuvier, Règne Anim., ed. I, 1817, pl. 10, fig. 3, Cayenne.
Pristigaster martinii Agassiz, in Spix, Pisc. Brasil., 1829, 55, Brazil.
Pristigaster phaeon Cuvier and Valenciennes, Hist. Nat. Poiss., XX, 1847, 338, mouth of Amazon.

Family 43. DUSSUMIERIIDÆ. Round herrings

- Genus 109. JENKINSIA Jordan and Evermann.**
Jenkinsia Jordan and Evermann, Fishes North and Mid. Amer., 1896, 418 (*Dussumieriella stolifera* Jordan and Gilbert).

- 240. Jenkinsia stolifera** (Jordan and Gilbert).
 Gulf of Mexico from Florida to Yucatan.
Dussumieria stolifera Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 25, Key West, Fla.
- 241. Jenkinsia lamprotænia** (Gosse).
 Jamaica.
Clupea lamprotænia Gosse, Naturalist's Sojourn in Jamaica, 1851, 291, pl. 1, fig. 2, Jamaica.
- Genus 110. HALECULA** Jordan.
Halecula Jordan, Fossil Fishes of the Calif. Miocene, VIII, 1925, (*Etrumeus acuminatus* Gilbert).
- 242. Halecula acuminata** (Gilbert).
 Gulf of California.
Etrumeus acuminatus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 56, Gulf of California at Albatross station 3012 in 22 fathoms.
- Genus 111. ETRUMEUS** Bleeker.
Etrumeus Bleeker, Verh. Bat. Gen. Japan, XXV, 1853, 58, (*Clupea micropus* Temminck, and Schlegel, a Japanese species).
Perkinsia Rosa Smith Eigenmann, Amer. Nat., XXV, 1891, 153 (*P. othonops* R. S. Eigenmann).
- 243. Etrumeus sadina** (Mitchill). *Round herring.*
 Cape Cod to Gulf of Mexico.
Clupea sadina Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 457, New York.
Alosa teres DeKay, New York Fauna: Fishes, 1842, 262, New York.
- 244. Etrumeus micropus** Temminck and Schlegel.
 Coast of Japan; Hawaii; found once off Point Loma in California.
Etrumeus micropus Temminck and Schlegel, Fauna Japonica, Poiss., 1850, 236, pl. 107, fig. 2, Japan.
Perkinsia othonops Rosa Smith Eigenmann, Amer. Nat., XXV, 1891, 153, Point Loma, San Diego, Calif.
- Family 44. DOROSOMIDÆ. Gizzard shad**
- Genus 112. DOROSOMA** Rafinesque. *Gizzard shads.*
Dorosoma Rafinesque, Ichth. Ohiensis, 1820, 39 (*D. notata* Rafinesque).
Chaloëssus Cuvier, Règne Anim., ed. 2, II, 1829, 320; in part (includes *Opisthonema*, etc.; restricted by Cuvier and Valenciennes to *Megalops cepedianus* LeSueur).
- 245. Dorosoma cepedianum** (LeSueur). *Gizzard shad; Hickory shad.*
 Cape Cod to Texas, entering all streams and variously landlocked.
Megalops cepedianus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, Delaware and Chesapeake Bays.
Clupea heterura Rafinesque, Amer. Month. Mag., 1818, 354, Ohio River.
Dorosoma notata Rafinesque, Ichth. Ohiensis, 1820, 39, Ohio River.
Chaloëssus ellipticus Kirtland, Rep. Zool. Ohio, 1839, 169, Ohio River.
Dorosoma insociabilis Abbot, Proc. Acad. Nat. Sci. Phila., XII, 1860, 365, Sturgeon Pond, Trenton, N. J.
- 246. Dorosoma cepedianum exile** Jordan and Gilbert.
 Streams of Texas.
Dorosoma cepedianum exile Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 585, Galveston, Tex.
- Genus 113. SIGNALOSA** Evermann and Kendall.
Signalosa Evermann and Kendall, Bull. U. S. Fish Com., XVII, 1897 (Feb. 9, 1898), 127 (*S. atchafalayæ* Evermann and Kendall).
- 247. Signalosa atchafalayæ** Evermann and Kendall.
 Coast of Louisiana and southward.
? *Chaloëssus mexicanus* Günther, Cat., VII, 1868, 409, Mexico.
Signalosa atchafalayæ Evermann and Kendall, Bull. U. S. Fish Com., XVII, 1897 (Feb. 9, 1898), 127, pl. 7, fig. 4, Atchafalaya River, Melville, La.

248. *Signalosa atchafalayæ vanhyningi* Weed.

Near Gainesville, Fla.

Signalosa atchafalayæ vanhyningi Weed, Field Mus. Nat. Hist., Zool., XII, October, 1925, 145, Prairie Creek 6 miles southeast of Gainesville, Fla.**249. *Signalosa mexicana* (Günther).**

New Orleans to Central America.

Chatoëssus mexicanus Günther, Cat., VII, 1868, 409, Mexico.**250. *Signalosa mexicana campi* Weed.**

Streams about Brownsville, Tex.

Signalosa mexicana campi Weed, Field Mus. Nat. Hist., Zool., XII, October, 1925, 143, Resaca de la Guerra, near Brownsville, Tex.**251. *Signalosa petenensis* (Günther).**

Lake Peten.

Meletta petenensis Günther, Proc. Zool. Soc. Lond., 1866, 603, Lake Peten, Yucatan.*European anchovy* (?) *Engraulis encrasicholus*
Family 45. ENGRAULIDÆ. Anchovies

Genus 114. ENGRAULIS Cuvier. Anchovies.

Engraulis Cuvier, Règne Anim., ed. I, 1817, 174 (*Clupea encrasicholus* L.).*Engrasicholus* Fleming, Brit. Anim., 1828, 183 (*Clupea encrasicholus* L.).**252. *Engraulis mordax* Girard. California anchovy.**

Pacific coast of America, from Vancouver Island to Lower California.

Engraulis mordax Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 138, and in Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 334, Shoalwater Bay, Wash.**253. *Engraulis nanus* Girard. Bay anchovy.**

San Francisco Bay; not known elsewhere.

Engraulis nanus Girard, Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 335, San Francisco.Genus 115. ANCHOVIELLA²⁰ Fowler. American anchovies.*Anchoviella* Fowler, Proc. Acad. Nat. Sci. Phila., LXIII, 1911, 211 (*Engraulis perasciatus* Poey).*Anchoa* Jordan and Evermann, Proc. Calif. Acad. Sci., 4th Ser., XV, April 27, 1927, 502 (*Engraulis compressus* Girard).

Subgenus ANCHOVIELLA Fowler.

254. *Anchoviella perasciata* (Poey). Grubber broadhead.

Florida Keys to Jamaica.

Engraulis perasciatus Poey, Memorias, II, 1861, 313, Cuba.**255. *Anchoviella miarchus* (Jordan and Gilbert).**

Pacific coast of America, from Mazatlan to Panama.

Stolephorus miarchus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 344, Mazatlan; perhaps a larval form.**256. *Anchoviella exigua* (Jordan and Gilbert).**

Pacific coast of Mexico at Mazatlan.

Stolephorus exigua Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 343, Mazatlan.**257. *Anchoviella perthecata* (Goode and Bean).**

Pensacola, Fla.

Stolephorus perthecatus Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 434, Pensacola, Fla.

²⁰ *Anchoviella* differs from *Stolephorus* Lacépède and related Old World genera in the entire absence of ventral scales. The many species cover a variety of forms; rostrals and spinifer with its relatives being perhaps generically distinct from *perasciatus*, *epsetus*, and related forms, having many more gillrakers; these latter species constitute the subgenus *Anchova*, the others the subgenus *Anchoviella* Fowler.

- 258. Anchoviella epsetus** (Bonnaterre). *Striped anchovy; Manjua; Anchovy fry.*
 Cape Cod to Brazil.
Piquitinga Marcgrave, Hist. Brasil., 1648, 159, Brazil.
Menidia Browne, Hist. Jamaica, 1756, 441, Jamaica; nonbinomial.
Esox epsetus Bonnaterre, Tabl. Ichth., 1788, 175, Jamaica; after Browne.
Atherina brownii Gmelin, Syst. Nat., 1789, 1397, Jamaica; after Browne.
Engraulis tricolor Agassiz, in Spix, Pisc. Brasil., 1829, 51, Bahia, Pará.
Engraulis lemniscatus Cuvier, Règne Anim., ed, 2, II, 1829, 323, Brazil;
 after *Piquitinga* of Marcgrave.
Stolephorus hiulcus Goode and Bean, Proc. U. S. Nat. Mus., II, 1879,
 343, Clearwater Harbor, Fla.; depth 5½; A, 22.
- 259. Anchoviella cubana** (Poey). *Bocon.*
 Cuba and Porto Rico.
Engraulis cubanus Poey, Synopsis, 1868, 420, Cuba.
- 260. Anchoviella ischana** (Jordan and Gilbert).
 Pacific coast of America, from Mazatlan to Panama.
Stolephorus ischanus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV,
 1881, 340, Mazatlan.
Stolephorus starksi Gilbert and Pierson, Fishes of Panama, in Jordan
 and Evermann, Fishes North and Mid. Amer., 1898, 2813, Panama.
- 261. Anchoviella cultrata** (Gilbert).
 Santa Margarita Island, Lower California.
Stolephorus cultratus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 544,
 Santa Margarita Island, Lower California.
- 262. Anchoviella delicatissima** (Girard).
 San Diego Bay and southward on coast of Lower California.
Engraulis delicatissimus Girard, Proc. Acad. Nat. Sci. Phila., VII,
 1854, 154, and Pac. R. R. Surv., X, Pt. 4, Fishes, 1858, 335, San
 Diego.
- 263. Anchoviella helleri** Hubbs.
 Gulf of California.
Anchoviella helleri Hubbs, Proc. Biol. Soc., Wash., XXXIV, Mar. 31,
 1921, 47, San Felipe Bay, gulf coast of Lower California.
- 264. Anchoviella choerostoma** (Goode). *Hogmouth fry.*
 Bermuda Islands; common in Hamilton Harbor.
Engraulis charostomus Goode, Amer. Jour. Sci. Arts, August, 1874,
 125, Bermuda.
- 265. Anchoviella argyrophana** (Cuvier and Valenciennes).
 Gulf Stream; Woods Hole, Mass.
Engraulis argyrophanus Cuvier and Valenciennes, Hist. Nat. Poiss.,
 XXI, 1848, 49, equatorial Atlantic.
Stolephorus eurystole Swain and Meek, Proc. Acad. Nat. Sci. Phila.,
 XXXVI, 1884, 34. Woods Hole, Mass.
- 266. Anchoviella curta** (Jordan and Gilbert).
 Mazatlan, Mexico.
Stolephorus curtus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881,
 343, Mazatlan, Mexico.
- 267. Anchoviella mitchilli** (Cuvier and Valenciennes).
 Cape Cod to Texas on sandy shores, entering rivers; common.
Engraulis mitchilli Cuvier and Valenciennes, Hist. Nat. Poiss., XXI,
 1848, 50, New York; Carolina; Lake Ponchartrain.
Engraulis louisiana LeSueur, in Cuvier and Valenciennes, Hist. Nat.
 Poiss., XXI, 1848, 51, Lake Ponchartrain, La.
Engraulis duodecim Cope, Trans. Amer. Phil. Soc., 1866, 405, Beasley's
 Point, N. J. (dorsal fin said to be entirely anterior to the long and
 deeply concave anal, which is not true in *S. mitchilli*; the description
 otherwise agrees).
- 268. Anchoviella lucida** (Jordan and Gilbert). *Sardina.*
 Mazatlan, Mexico.
Stolephorus lucidus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV,
 1881, 341, Mazatlan.

- 269. Anchoviella naso** (Gilbert and Pierson).
Panama; common.
Stolephorus naso Gilbert and Pierson, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2813, Panama.
- 270. Anchoviella robertsi** (Jordan and Rutter).
West Indies.
Stolephorus robertsi Jordan and Rutter, Proc. Acad. Nat. Sci. Phila., XLIX, 1897, 95, Jamaica.
- 271. Anchoviella astilbe** (Jordan and Rutter).
West Indies.
Stolephorus astilbe Jordan and Rutter, Proc. Acad. Nat. Sci. Phila., XLIX, 1897, 95, Jamaica.
- 272. Anchoviella scofieldi** (Jordan and Culver).
Astillero at Mazatlan; not very abundant.
Stolephorus scofieldi Jordan and Culver, in Jordan, Fishes of Sinaloa, Proc. Calif. Acad. Sci., 3d Ser., V, 1895, 410, Mazatlan, Mexico.
- 273. Anchoviella lyolepis** (Evermann and Marsh).
Porto Rico.
Stolephorus lyolepis Evermann and Marsh, Bull. U. S. Fish Com., XX, 1900 (Dec. 29), 89, fig. 13, Culebra Island, Porto Rico.
- 274. Anchoviella branchiomelas** (Eigenmann).
North coast of Colombia.
Stolephorus branchiomelas Eigenmann, Proc. Am. Phil. Soc., LVI, 1917, 682, mouth of Rio Dagua, Colombia.
- 275. Anchoviella arenicola** (Meek and Hildebrand).
Panama.
Anchovia arenicola Meek and Hildebrand, Marine Fishes of Panama, in Field Mus. Nat. Hist., Zool., XV, 1923, 201, Pl. XIII, fig. 1, Taboga Island, Panama.
- 276. Anchoviella scitula** (Fowler).
Coast of California; San Diego.
Anchovia scitula Fowler, Proc. Acad. Nat. Sci. Phila., LXIII, 1911, 211, fig. 2, San Diego, Calif.
- 277. Anchoviella parva** (Meek and Hildebrand).
Colon.
Anchovia parva Meek and Hildebrand, Marine Fishes of Panama, in Field Mus. Nat. Hist., Zool., XV, 1923, 202, Pl. X, fig. 2, Porto Bello, Panama.
- 278. Anchoviella elongata** (Meek and Hildebrand).
Canal Zone.
Anchovia elongata Meek and Hildebrand, Marine Fishes of Panama, in Field Mus. Nat. Hist., Zool., XV, 1923, 198, Pl. XII, fig. 2, Míndi Cut, Canal Zone.
- 279. Anchoviella cayorum** (Fowler).
Florida Keys.
Anchovia charostoma cayorum Fowler, Proc. Acad. Nat. Sci. Phila., LVII, 1906, 85, fig. 4, Hailer's Rock, Florida Keys.
- 280. Anchoviella platyargyrea** (Fowler).
West Indies.
Anchovia platyargyrea Fowler, Proc. Acad. Nat. Sci. Phila., LXIII, 1911, 216, fig. 4, St. Martin's, West Indies.
- 281. Anchoviella lepidentostole** (Fowler).
Coast of Surinam.
Anchovia lepidentostole Fowler, Proc. Acad. Nat. Sci. Phila., LXIII, 1911, 214, fig. 3, Surinam.
- 282. Anchoviella guianensis** (Eigenmann).
Coast of Guiana.
Stolephorus guianensis Eigenmann, Mem. Carnegie Mus., V, 1912, 448, Pl. LXII, fig. 5, Bartica Rocks, British Guiana; misspelled *Stoleferus* in plate.

- 283. Anchoviella trinitatis** (Fowler).
 British West Indies.
Anchovia trinitatis Fowler, Proc. Acad. Nat. Sci. Phila., LXVII, 1915,
 527, fig. 3, Port of Spain, Trinidad, British West Indies.
- 284. Anchoviella filifera** (Fowler).
 British West Indies.
Anchovia filifera Fowler, Proc. Acad. Nat. Sci. Phila., LXVII, 1915,
 534, fig. 2, Port of Spain, Trinidad, British West Indies.
- 285. Anchoviella abbotti** (Fowler).
 British West Indies.
Anchovia abbotti Fowler, Proc. Acad. Nat. Sci. Phila., LXVII, 1915,
 522, fig. 1, Port of Spain, Trinidad, British West Indies.
- Subgenus ANCHOA Jordan and Evermann.
- 286. Anchoviella compressa** (Girard).
 Point Conception to Lower California.
Engraulis compressus Girard, Pac. R. R. Surv., X, 1858, 336, San Diego.
- 287. Anchoviella spinifera** (Cuvier and Valenciennes).
 Coast of Guiana; recorded from Panama by Steindachner, probably an
 error.
Engraulis spinifer Cuvier and Valenciennes, Hist. Nat. Poiss., XXI,
 1848, 39, Cayenne.
- 288. Anchoviella rastralis** (Gilbert and Pierson).
 Panama; abundant.
Stolephorus rastralis Gilbert and Pierson, in Jordan and Evermann,
 Fishes North and Mid. Amer., 1898, 2811, Panama.
- 289. Anchoviella mundeola** (Gilbert and Pierson).
 Panama; abundant.
Stolephorus mundeolus Gilbert and Pierson, in Jordan and Evermann,
 Fishes North and Mid. Amer., 1898, 2812, Panama.
- 290. Anchoviella panamensis** (Steindachner).
 Panama.
Engraulis panamensis Steindachner, Ichth. Beitr., IV, 1875, 39, Panama.
- 291. Anchoviella eigenmannia** (Meek and Hildebrand).
 Panama.
Anchovia eigenmannia Meek and Hildebrand, Marine Fishes of Panama,
 in Field Mus. Nat. Hist., Zool., XV, 1923, 20, Taboga Island, Canal
 Zone.
- Genus 116. ANCHOVIA** Jordan and Evermann.
Anchovia Jordan and Evermann, Fishes North and Mid. Amer., 1896,
 449, (*Engraulis macrolepidotus* Kner and Steindachner).
- 292. Anchovia macrolepidota** (Kner and Steindachner). *Sardina bocona*.
 Gulf of Mexico to Panama.
Engraulis macrolepidotus Kner and Steindachner, Abhandl. Bayer.
 Akad. Wiss., X, 1864, 21, Pl. III, fig. 2, Rio Bayano, Panama..
- 293. Anchovia clupeoides** (Swainson). *Hechudo; Grubber broadhead*.
 Coast of Cuba and southward; common, ascending streams.
Engraulis clupeoides Swainson, Nat. Hist. Fish., II, 1839, 388, Pernam-
 buco.
Stolephorus surinamensis Bleeker, Nederl. Tijdschr. Dierk., 1866, 178,
 Surinam.
Engraulis productus Poey, Repertorio, 1866, 380, Cuba.
- Genus 117. AMPLOVA** Jordan and Seale.
Amplova Jordan and Seale, Copeia, No. 141, 1925, 31 (*Stolephorus*
brevirostrum Meek and Hildebrand; not *Engraulis brevirostris*,
 Günther, also a species of *Amplova*).
- 294. Amplova balboæ** Jordan and Seale.
 Panama.
Amplova balboæ Jordan and Seale, Copeia, No. 141, Apr. 30, 1925,
 31, substitute for *Stolephorus brevirostrum* Meek and Hildebrand.

Anchovia brevirostrum Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, Pt. I, 198, Pl. XV, fig. 1; not *Engraulis brevirostris* Günther from Brazil, also a species of *Amplova*.

Genus 118. **CETENGRAULIS** Günther.

Cetengraulis Günther, Cat., VII, 1868, 383 (*Engraulis edentulus* Cuvier).

295. **Cetengraulis edentulus** (Cuvier). *Bocon; Tarpon fry.*
West Indies to Brazil; common in Cuba.
Sprat Sloane, Hist. Jamaica, II, 1727, 282, Jamaica; after Sloane.
Engraulis edentulus Cuvier, Règne Anim., ed. 2, II, 1829, 323, Jamaica.
Engraulis brevis Poey, Repertorio, I, 1866, 379, Cuba.

296. **Cetengraulis giberti** (Evermann and Marsh).
Coast of Porto Rico.
Stolephorus giberti Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 352, Palo Seco near San Juan, Porto Rico.

297. **Cetengraulis garmani** (Evermann and Marsh). *Porto Rican anchovy.*
Porto Rico.
Stolephorus garmani Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899, (Dec. 19), 352, Puerto Real, Porto Rico.

298. **Cetengraulis mysticetus** (Günther).
Pacific coast of tropical America.
Engraulis mysticetus Günther, Proc. Zool. Soc. Lond., 1866, 604, Panama.
Stolephorus opercularis Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 275, Punta San Felipe, Gulf of California.

299. **Cetengraulis engymen** Gilbert and Pierson.
Panama Bay.
Cetenengraulis engymen Gilbert and Pierson, in Jordan and Evermann, Fishes North and Mid. Amer. 1898, 2815, Panama Bay.

Genus 119. **PTERENGRAULIS** Günther.

Pterengraulis Günther, Cat., VII, 1868, 398, (*Clupea atherinoides* L.).

300. **Pterengraulis atherinoides** (Linnaeus).
Coast of Guiana and Brazil; ascending rivers.
Clupea atherinoides Linnaeus, Syst. Nat., ed. XII, 1766, 523, Surinam.

Genus 120. **LYCENGRAULIS** Günther.

Lycengraulis Günther, Cat., VII, 1868, 399 (*Engraulis grossidens* Cuvier).

301. **Lycengraulis grossidens** (Cuvier).
Coast of Guiana and Brazil.
Engraulis grossidens Cuvier, in Agassiz, Spix, Pisc. Brasil., 1828, 50, Rio Janeiro.
Engraulis janeiro Agassiz, loc. cit., pl. 24, fig. 1, Rio Janeiro.
Engraulis dentex Cuvier and Valenciennes, Hist. Nat. Poiss., XXI, 1848, 28, Rio Janeiro.

302. **Lycengraulis poeyi** (Kner and Steindachner).
Rio Bayano, Panama.
Engraulis poeyi Kner and Steindachner, Abhandl. Bayer. Akad. Wiss., X, 1864, 23, with plate; Rio Bayano.

Family 46. ALEPOCEPHALIDÆ

Genus 121. **ALEPOCEPHALUS** Risso.

Alepocephalus Risso, Mém. Acad. Nat. Sci. Turin, XXV, 1820, 270 (*A. rostratus* Risso).

303. **Alepocephalus agassizii** Goode and Bean.
Gulf Stream.
Alepocephalus agassizii Goode and Bean, Bull. Mus. Comp. Zool., X, 1883, 318, Gulf Stream.

304. **Alepocephalus productus** Gill.
Gulf Stream.
Alepocephalus productus Gill, Proc. U. S. Nat. Mus., VI, 1883, 256, Gulf Stream at Albatross station 2035 in 1,362 fathoms.

- 305. *Alepocephalus tenebrosus* Gilbert.**
Santa Barbara Channel, California.
Alepocephalus tenebrosus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891,
 545, Santa Barbara Channel.
- 306. *Alepocephalus asperifrons* Garman.**
Off Panama in deep water.
Alepocephalus asperifrons Garman, Mem. Mus. Comp. Zool., XXIV,
 1899, 291, Pl. LIX, fig. 1, Panama.
- 307. *Alepocephalus converifrons* Garman**
Off Pacific coast of Mexico, in deep water.
Alepocephalus converifrons Garman, Mem. Mus. Comp. Zool., XXIV,
 1899, 292, Pl. LIX, fig. 2; off Pacific coast of Mexico.
- 308. *Alepocephalus fundulus* Garman.**
Pacific coast of Panama, in deep water.
Alepocephalus fundulus Garman, Mem. Mus. Comp. Zool., XXIV,
 1899, 293, Pl. LVII, fig. 2; off Panama in deep water.
- Genus 122. *MITCHILLINA* Jordan and Evermann.**
Mitchillina Jordan and Evermann, Fishes North and Mid. Amer.,
 1896, 453, (*Alepocephalus bairdii* Goode and Bean).
- 309. *Mitchillina bairdii* (Goode and Bean).**
Grand Banks of Newfoundland.
Alepocephalus bairdii Goode and Bean, Proc. U. S. Nat. Mus., II, 1879,
 55, Grand Banks of Newfoundland.
- Genus 123. *BATHYTROCTES* Günther.**
Bathytroctes Günther, Ann. Mag. Nat. Hist., II, 5th ser., 1878, 249
 (*B. microlepis* Günther).
- 310. *Bathytroctes stomias* Gilbert.**
Coast of Oregon.
Bathytroctes stomias Gilbert, Proc. U. S. Nat. Mus., XII, 1890, 53,
 coast of Oregon in 877 fathoms, *Albatross* station 3074.
- 311. *Bathytroctes inspector* Garman.**
Pacific coast of Panama, in deep water.
Bathytroctes inspector Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
 288, Pl. M., fig. 1; Panama.
- 312. *Bathytroctes alveatus* Garman.**
Pacific coast of Panama in deep water.
Bathytroctes alveatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
 Pl. LVIII, fig. 1, 287; Panama.
- 313. *Bathytroctes alvifrons* Garman.**
Pacific coast of Panama in deep water.
Bathytroctes alvifrons Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
 286, Pl. LVIII, figs. 2, 2a; Panama.
- Genus 124. *XENOGNATHUS* Gilbert.**
Xenognathus Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 311, (*X. profundorum* Gilbert).
- 314. *Xenognathus profundorum* Gilbert.**
Coast of California, in deep water, 1,350 to 2,182 fathoms.
Xenognathus profundorum Gilbert, Proc. U. S. Nat. Mus., XLVIII,
 1915, 311, off Santa Catalina Island, California.
- Genus 125. *NARCETES* Alcock.**
Narcetes Alcock, Ann. Mag. Nat. Hist., VI, 1890, 305 (*N. erimelas*
 Alcock).
- 315. *Narcetes stomias* (Gilbert).**
*Off Pacific coast of America, from Oregon to southern California in
 deep water.*
Bathytroctes stomias Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 53,
 coast of Oregon in 877 fathoms, *Albatross* station 3074.
- 316. *Narcetes pluriserialis* Garman.**
Pacific coast of Panama in deep water.
Narcetes pluriserialis Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
 289, Pl. LVII, fig. 3; Panama.

Genus 126. LEPTOCHILICHTHYS Garman.

Leptochilichthys Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 284
(*L. agassizii* Garman).

317. *Leptochilichthys agassizii* Garman.

Pacific coast of Panama, in deep water.

Leptochilichthys agassizii Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 286, Pl. LVIII, fig. 3; off Panama.

Genus 127. PLATYTROCTES Günther.

Platytroctes Günther, Ann. Mag. Nat. Hist., II, 1878, 249 (*P. apus* Günther).

318. *Platytroctes apus* Günther.

Mid-Atlantic and the Arabian Sea.

Platytroctes apus Günther, Ann. Mag. Nat. Hist., II, 1878, 249, Mid-Atlantic.

Genus 128. TALISMANIA Goode and Bean.

Talismania Goode and Bean, Oceanic Ichth., 1895, 41 (*Bathytroctes homopterus* Vaillant).

319. *Talismania antillarum* Goode and Bean.

Caribbean Sea.

Talismania antillarum Goode and Bean, Oceanic Ichth., 1895, 44, fig. 49, Albatross station 2394, in Caribbean Sea, lat. 28° 38' 30" N., long. 87° 2' W., 420 fathoms.

320. *Talismania aequatoris* Goode and Bean.

Coast of Ecuador.

Talismania aequatoris Goode and Bean, Oceanic Ichth., 1895, 44, fig. 50, off coast of Ecuador.

Genus 129. CONOCARA Goode and Bean.

Conocara Goode and Bean, Oceanic Ichth., 1895, 39 (*C. macdonaldi* Goode and Bean).

321. *Conocara macdonaldi* Goode and Bean.

Gulf of Mexico.

Conocara macdonaldi Goode and Bean, Oceanic Ichth., 1895, 39, fig. 48, Gulf of Mexico.

322. *Conocara macroptera* (Vaillant).

Deep seas, coast of Morocco to Soudan, banc d'Arguin, and from the Canaries. A specimen 8 inches in length was obtained by the Albatross at station 2751, lat. 16° 57' N., long. 63° 12' W., in 68 fathoms.

Alepocephalus macropterus Vaillant, Expéd. Sci. Travailleur et Talisman, Poissons, 1888, 150, Pl. XI, figs. 2a, 2b, 2c, coast of Morocco, Soudan, and Canaries.

Genus 130. BAJACALIFORNIA Townsend and Nichols.

Bajacalifornia Townsend and Nichols, Bull. Amer. Mus. Nat. Hist., LII, 1925, 8 (*B. burragei* Townsend and Nichols).

323. *Bajacalifornia burragei* Townsend and Nichols.

West coast of Lower California in deep water.

Bajacalifornia burragei Townsend and Nichols, Bull. Am. Mus. Nat. Hist., LII, 1925, 8, fig. 3, off Todos Santos Bay, Lower California, in 590 fathoms.

Genus 131. XENODERMICHTHYS Günther.

Xenodermichthys Günther, Ann. Mag. Nat. Hist., II, ser. 5, 1878, 250 (*X. nodulosus* Günther).

Aleposomus Gill, Amer. Nat., XVIII, 1884, 433 (*A. copei* Gill).

324. *Xenodermichthys copei* (Gill).

Gulf Stream.

Aleposomus copei Gill, Amer. Nat., XVIII, 1884, 433, Gulf Stream, lat. 37° 12' 20" N., long. 69° 39' W.

Genus 132. ERICARA Gill and Townsend.

Ericara Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897, 232 (*E. salmonaea* Gill and Townsend).

325. *Ericara salmonaea* Gill and Townsend.

Bering Sea, southwest of Pribilof Islands.

Ericara salmonaea Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897, 232, Bering Sea.

Genus 133. *ATOPICHTHYS* Garman.

*Atopichthys*²¹ Garman, Mem., Mus. Comp. Zool., XXIV, 1899, 325 (*A. esunculus* Garman).

326. *Atopichthys esunculus* Garman.

Pacific coast of Mexico.

Atopichthys esunculus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 327, Pl. LXV, figs. 2, 2a; off Acapulco, Mexico.

Suborder SALMONOIDEI

Family 47. SALMONIDÆ. Salmon; Trout

Genus 134. *ONCORHYNCHUS* Suckley. *Pacific salmon.*

Oncorhynchus Suckley, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 313 (*Salmo scouleri* Richardson = *Salmo gorbuscha* Walbaum).

Hypsafario Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 330 (*Salmo kennnerlyi* Gill).

Subgenus. *ONCORHYNCHUS* Suckley.327. *Oncorhynchus gorbuscha* (Walbaum). *Humpback salmon; Pink salmon; Haddo; Holia; Gorbuscha.*

Pacific coast and rivers of North America and Asia, from California northward.

Salmo gorbuscha Walbaum, Artedi Pisc. 1792, 69, Kamchatka; after Gorbuscha of Pennant and Krasheninnikof.

Salmo gibber Bloch and Schneider, Syst. Ichth., 1801, 409, Kamchatka; after Krasheninnikof.

Salmo proteus Pallas, Zoogr. Rosso-Asiat., III, 1811, 376, Bering Sea.

Salmo scouleri Richardson, Fauna Bor.-Amer., III, 1836, 158; Observatory Inlet.

Salmo tschawytschiformis Smitt, I Riksmusecum Befintliga Salmonider, 1886, 161; Port Clarence.

328. *Oncorhynchus keta* (Walbaum). *Dog salmon; Hayko; Le kai salmon; Chum; Calico salmon.*

Pacific coast of America, from San Francisco to Kamchatka.

Salmo keta vel kayko Walbaum, Artedi Pisc., 1792, 72, rivers of Kamchatka; after *keta* or *kayko* of Pennant and Krasheninnikof.

Salmo lagocephalus Pallas, Zoogr. Rosso-Asiat., III, 1811, 372; Bering Sea.

?*Salmo japonicus* Pallas, loc. cit., 382, Curile Islands; Amur River.

Salmo dermatus Richardson, Voy. Herald, Zool., 1854, 167; Yukon River.

Salmo consuetus Richardson, loc. cit., 168; Yukon River.

Salmo canis Suckley, Ann. Lyc. Nat. Hist. N. Y., VII, 1858, 9, and Monogr. Salmo, 101, 1861 (1874); Puget Sound.

329. *Oncorhynchus kisutch* (Walbaum). *Silver salmon; Kisutch; Medium red salmon; Skowitz; Hoopid salmon; Coho salmon; Bielaya ryba; Quisutsch; Tschaviche.*

From San Francisco northward, especially in Puget Sound and the Alaskan fjords; south on the Asiatic coast to Japan.

?*Salmo milktschitsch* Walbaum, Artedi Pisc., 1792, 70, Bering Sea; after *milktschutsch* or *miltschitsch* of Pennant and Krasheninnikof; probably the young of *kisutch*.

Salmo kisutch Walbaum, Artedi Pisc., 1792, 70, rivers and lakes of Kamchatka; after the *kisutch* of Pennant.

?*Salmo striatus* Bloch and Schneider, Syst. Ichth., 1801, 407, Kamchatka; after *milktschitsch* of Krasheninnikof.

²¹ This name is given not as a genus, but rather as a group name of larval fishes related to the oels or to the Alepocephalidae, commonly known as *Leptocephalus*; but *Leptocephalus morrisi* (type of *Leptocephalus*) is now known to be a young conger. A number of these larvae (*A. sicarius*, *cinctus*, *dentatus*, *falcidens*, *acutus*, *opichthys*, *cingulus*, *lynchus*, *obtusus*, and *longidens*) are described by Garman (Deep Sea Fishes), but they need not be specifically recognized here. The species called *Atopichthys ensunculus* Garman, perhaps belonging to the Alepocephalidae, has been designated as the type of this group.

Salmo kisutch Bloch and Schneider, loc. cit., 407, Kamchatka; after Pennant.

Salmo sanguinolentus Pallas, Zoogr. Rosso-Asiat., III, 1811, 379, Bering Sea.

Salmo tsuppitch Richardson, Fauna Bor.-Amer., III, 1836, 224, Columbia River.

Subgenus HYPASFARIO Gill.

330. *Oncorhynchus nerka* (Walbaum). *Blueback salmon; Redfish; Fraser River salmon; Nerka; Sau-qui salmon; Sockeye or Sukkegh salmon; Krashnaya ryba.*

Klamath River and Rogue River to northern Alaska, Kamchatka, and Japan.

Salmo nerka Walbaum, Artedi Pisc., 1792, 71; after the *nerka* of Pennant, *nerka* of Krasheninnikof; rivers and seas of Kamchatka.

Salmo lycodon Pallas, Zoogr. Rosso-Asiat., III, 1811, 370, Okhotsk Sea, Kamchatka.

Salmo paucidens Richardson, Fauna Bor.-Amer., III, 1836, 222, Columbia River.

Salmo tapdisma Cuvier and Valenciennes, Hist. Nat. Poiss., XXI, 1848, 365, Kamchatka; on a drawing.

Salmo arabatch Cuvier and Valenciennes, loc. cit., Kamchatka; on a drawing.

Salmo melanopterus Cuvier and Valenciennes, loc. cit., Kamchatka; on a drawing.

Fario aurora Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 218; Girard, Pac. R. R. Surv., X, 1858, 308, pl. 68, Astoria, Oreg.

331. *Oncorhynchus tshawytscha* (Walbaum). *Chinook salmon; Tchaviche; King salmon; Columbia River salmon; Spring salmon; Sacramento River salmon; Tyee salmon; Tschawytscha; Quinnat salmon.*

Alaska, Oregon, and California; southward to Monterey, and to northern China.

Salmo tshawytscha Walbaum, Artedi Pisc., 1792, 71, rivers of Kamchatka; after the *tshawytscha* of Krasheninnikof, Descr. Kamchatka, 1768, 178, and the *tshawytscha* of Pennant, 1792.

Salmo orientalis Pallas, Zoogr. Rosso-Asiat., III, 1811, 367, Kamchatka.

Salmo quinnat Richardson, Fauna Bor.-Amer., III, 1836, 219, Columbia River; and of many writers.

Fario argyreus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 218, Cape Flattery; Fort Steilacoom.

Salmo confluentus Suckley, Ann. Lyc. Nat. Hist. N. Y., December, 1858, and Pacific R. R. Surv., XII, pt. 2, 1860, 334, Puyallup River near Fort Steilacoom.

Salmo cooperi Suckley, Notices New Spec. N. Amer. Salmon, New York, June, 1861, Okanogan River.

Salmo warreni Suckley, loc. cit., June, 1861, and loc. cit., 1861, 117 (1874) Fraser River, British Columbia.

Salmo richardi Suckley, loc. cit., June, 1861, and loc. cit., 1861, 117, Fraser and Skagit Rivers.

332. *Oncorhynchus kenneryi* (Suckley).

Lakes of Oregon, Washington, Idaho, British Columbia, Alaska, and Hokkaido. A dwarf landlocked form. Probably several forms should be recognized.

Salmo kenneryi Suckley, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 307, Lake Chiloweyuck, Washington.

Genus 135. SALMO Linnaeus. *Salmon and trout.*

Salmo Linnaeus, Syst. Nat., ed. X, 1758, 308 (*S. salar* Linnaeus); after Artedi.

Trutta Linnaeus, loc. cit., 308 (*trutta*, etc.: "Truttæ corpore variegato"). *Trutta* Geoffroy, Descr. de 719 Plantes, etc., 1767, 719 (*Salmo trutta* Linnaeus).

Fario Cuvier and Valenciennes, Hist. Nat. Poiss., XXI, 1848, 277 (*Fario argenteus*=*Salmo fario* L.).

Salar Cuvier and Valenciennes, loc. cit., 314 (*Salmo ansonii*=*Salmo fario*). *Trutta* Siebold, Süswasserfische Mitteleuropas, 1863, 280 (*Salmo trutta* L.).

Subgenus SALMO Linnaeus.

- 333. *Salmo salar* Linnæus.** *Common Atlantic salmon.*
 North Atlantic; ascending all suitable rivers in northern Europe and region north of Cape Cod; formerly abundant in the Hudson and occasional in the Delaware, its northern limit being about lat. 50° N., and southwest Greenland.
Salmo salar Linnæus, Syst. Nat., ed. X, 1758, 308, seas of Europe.
Salmo mas Walbaum, Artedi Pisc., 1792, 58, Germany; based on "Der Hakelachs" of Bloch; probably not intended as a binomial name.
Salmo gloveri Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 86, Union River, Me.
- 334. *Salmo sebago* Girard.** *Landlocked salmon; Sebago salmon.*
 Lakes of Maine, New Hampshire, New Brunswick, and Nova Scotia.
Salmo sebago Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 380, Sebago Lake, Me.
- 335. *Salmo ouananiche* McCarthy.** *Ouananiche; Winninish.*
 Lake St. John, Saguenay River, and northward.
Salmo salar ouananiche Eugene McCarthy, Forest and Stream, March 10, 1894, 206, and in Jordan and Evermann, Fishes North and Mid. Amer., 1896, 487, Saguenay River, Canada.
- 336. *Salmo mykiss* Walbaum.** *Mykiss; Somka; Kamchatka salmon trout; Muikichi.*
 Kamchatka; probably not occurring in America.
Mykiss Pennant, Arctic Zool., 1792, 126, Kamchatka; after Krasheninnikof, etc.
Salmo mykiss Walbaum, Artedi Pisc., 1792, 59, Kamchatka; based on *mykiss* of Pennant.
Salmo pensinensis Pallas, Zoogr. Rosso-Asiat., III, 1811, Gulf of Penshin.
Salmo purpuratus Pallas, loc. cit., 374, Bering Sea.
- 337. *Salmo regalis* Snyder.** *Royal silver trout.*
 Lake Tahoe.
Salmo regalis Snyder, Bull. U. S. Bur. Fish., XXXII, 1912 (Dec. 31), 28, Lake Tahoe near Brockway, Nev.
- Subgenus TRUTTA Linnæus.**
- 338. *Salmo clarkii* Richardson.** *Columbia River trout; Cut-throat trout.*
 Coastwise streams, from Puget Sound south to Elk River, Humboldt County, California.
Salmo clarkii Richardson, Fauna Bor.-Amer., III, 1836, 225, Cathlapooitl River.
Fario stellatus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 219, Fort Steilacoom, Shoalwater Bay.
Salmo brevicauda Suckley, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 308, Puget Sound.
- 339. *Salmo lewisi* (Girard).** *Yellowstone trout; Cut-throat trout.*
 Snake River basin above the Shoshone Falls, through Two-Ocean Pass to the headwaters of the Yellowstone and other affluents of the upper Missouri; Yellowstone Lake.
Salar lewisi Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 219, Falls of Missouri River.
Salmo carinatus Cope, Hayden's Geol. Surv. Mont., 1871 (1872); locality unknown, probably Yellowstone River.
- 340. *Salmo bouvieri* (Bendire).** *Waha Lake trout.*
 Waha Lake, Washington.
Salmo purpuratus bouvieri Bendire, in Jordan and Gilbert, Synopsis, 1883, 315, Waha Lake, Wash.
- 341. *Salmo utah* Suckley.** *Cut-throat trout of Utah Lake.*
 Lakes and streams west of the Wasatch range, especially in Bear, Provo, Jordan, and Sevier Rivers and in Utah Lake.
Salmo utah Suckley, Monogr. Salmo, 1861 (1874), 136, Utah Lake; pale specimens from the lake.

- 342. *Salmo gibbsii* Suckley.** Middle Columbia River basin.
Fario tsuppitch Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 218,
 Fort Dalles, Oreg.; not of Richardson.
Salmo gibbsii Suckley, Ann. Lyc. Nat. Hist. N. Y., VII, 1858, 1, Fort
 Dalles; also in the Yakima, John Day, and Boise Rivers; also in the
 Columbia at The Dalles, Oreg.
- 343. *Salmo henshawi* Gill and Jordan.** *Lake Tahoe trout; Silver trout; Truckee
 trout; Pyramid Lake trout.* Basin of the post-tertiary Lake Lahontan; Lake Tahoe, Pyramid Lake,
 WENNEMUCA Lake, Webber Lake, Donner Lake, Independence Lake,
 Truckee River, Humboldt River, Carson River, and most streams of
 the eastern slope of the Sierra Nevada; also the headwaters of
 Feather River, west of the Sierra Nevada; probably introduced from
 Nevada.
Salmo henshawi Gill and Jordan, in Jordan, Man. Vert., ed. 2, 1878,
 258, Lake Tahoe.
- 344. *Salmo tahoensis* (Jordan and Evermann).** *Great trout of Lake Tahoe.*
 Deep waters of Lake Tahoe.
Salmo clarkii tahoensis Jordan and Evermann, Fishes North and Mid.
 Amer., 1898, 2870, Lake Tahoe.
- 345. *Salmo jordani* (Meek).** *Jordan's trout; Spotted trout of Lake Southerland.*
 Lake Southerland, west of Puget Sound.
Salmo clarkii jordani Meek, Publ. Field Col. Mus., Zool., I, 1897
 (1899), 229, Lake Southerland, Wash.
- 346. *Salmo declivifrons* (Meek).** *Salmon trout of Lake Southerland.*
 Lake Southerland, west of Puget Sound.
Salmo clarkii declivifrons Meek, Field Col. Mus., Zool., I, 1899, 230,
 Lake Southerland, Wash.
- 347. *Salmo virginalis* (Girard).** *Rio Grande trout.*
 Upper Rio Grande and southward into the mountains of Chihuahua.
Salar virginalis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 220,
 Utah Creek and at Sangre de Cristo Pass, Colo.
Salmo spilurus Cope, Hayden's Geol. Surv. Mont., 1871 (1872), 470;
 Sangre de Cristo Pass, Colo.
- 348. *Salmo pleuriticus* Cope.** *Colorado River trout.*
 Colorado River basin.
Salmo pleuriticus Cope, Hayden's Geol. Surv. Mont., 1871 (1872),
 471; headwaters of Green River, Wyo.
- 349. *Salmo stomias* Cope.** *Greenback trout.*
 Headwaters of Arkansas and Platte Rivers.
Salmo stomias Cope, Hayden's Geol. Surv. Wyo., 1870 (1871), 433;
 South Platte River; locality unknown.
- 350. *Salmo macdonaldi* (Jordan and Evermann).** *Yellowfin trout.*
 Headwaters of the Arkansas River.
Salmo mykiss macdonaldi Jordan and Evermann, Proc. U. S. Nat.
 Mus., XII, 1889 (January 20, 1890), 453, Twin Lakes, Colo.
- 351. *Salmo crescentis* (Jordan and Beardslee).** *Blueback trout of Lake Crescent.*
 Crescent Lake, Clallam County, Washington.
Salmo gairdneri crescentis Jordan and Beardslee, Proc. Calif. Acad.
 Sci., 2d ser., VI, 1896, 207, pl. 22, Crescent Lake, Clallam County,
 Wash.
- 352. *Salmo bathæcetor* Meek.** *Long-headed trout of Crescent Lake.*
 Deep water of Crescent Lake, Washington.
Salmo bathæcetor Meek, Field Col. Mus., Zool., I, 1899, 227, Crescent
 Lake, Wash.
- 353. *Salmo beardslei* (Jordan and Seale).** *Speckled trout of Lake Crescent.*
 Crescent Lake, Clallam County, Wash.
Salmo gairdneri beardslei Jordan and Seale, Proc. Calif. Acad. Sci.,
 2d Ser., VI, 1896, 209, pl. 23, Crescent Lake, Clallam County, Wash.

354. *Salmo eremogenes* Evermann and Nichols.

Streams of central Wash.

Salmo eremogenes Evermann and Nichols, Proc. Biol. Soc. Wash., XXII, June 25, 1909, 93, pl. 2, Crab Creek at Rocky Ford near Ritzville, Wash.

355. *Salmo smaragdus* Snyder. *Emerald trout.*

Truckee Basin, Nev.

Salmo smaragdus Snyder, Bull. U. S. Bur. Fish., XXXV, 1915-16 (Sept. 28, 1917), 80, fig. 9, Pyramid Lake, Nev.

356. *Salmo aquilarum* Snyder. *Eagle Lake trout.*

Lakes of northeastern California.

Salmo aquilarum Snyder, Bull. U. S. Bur. Fish., XXXV, 1915-16 (Sept. 28, 1917), 77, fig. 7, Eagle Lake near mouth of Pine Creek, Modoc Co., Calif.

357. *Salmo irideus* Gibbons. *Rainbow trout; Coast range trout.*

Coastwise mountain streams of the Coast Ranges from the northern coast of California southward to San Diego County, Calif.; entering the sea where it grows to 6 or 10 pounds, returning as a steelhead.

Salmo irideus Gibbons, Proc. Calif. Acad. Nat. Sci., I, 1855, 35, San Leandro Creek, Alameda County, Calif.

Salmo rivularis Ayres, loc. cit., 42, Martinez, Calif.; steelhead form.

358. *Salmo gairdnerii* Richardson. *Brook trout of western Oregon to southern Alaska; Steelhead; Salmon trout; Kamloops trout; Stil-tse.*

Southern Alaska and Puget Sound to southern Oregon; in streams of the Coast Ranges and the Cascades, descending to the sea as a "steelhead"; perhaps not separable from *Salmo irideus*. Kamloops Lake, Okanagan Lake, and other lakes tributary to Frazer River or to the upper Columbia.

Salmo gairdnerii Richardson, Fauna Bor.-Amer., III, 1836, 221, Columbia River at Fort Vancouver.

Salmo masoni Suckley, Pac. R. R. Surv., XII, part 2, 1860, 345, Cathlapoochee River.

Salmo truncatus Suckley, Ann. Lyc. Nat. Hist. N. Y., VII, 1862, 3, New Dungeness, straits of Fuca.

Oncorhynchus kamloops Jordan, Forest and Stream, XXXIX, No. 19, November 10, 1892, 405, Kamloops Lake, British Columbia.

359. *Salmo shasta* (Jordan). *McCloud River rainbow trout; Rainbow trout of fish culturists.*

Streams of the Sierra Nevadas from Mount Shasta southward; known from McCloud River; the limits of its range not well known. Introduced into the Truckee River and many eastern streams, as well as in Europe and New Zealand.

Salmo irideus or rainbow trout of most writers and especially of fish-culturists; not *Salmo irideus* Gibbons, which is based on coastwise specimens.

Salmo gairdneri shasta Jordan, Thirteenth Biennial Rept. Calif. Fish Com., 1894, 142, with plate; McCloud River at Baird, Shasta County, Calif.

360. *Salmo stonei* (Jordan). *Nissuee trout; No-shee trout.*

McCloud River and upper Sacramento River.

Salmo gairdneri stonei Jordan, Thirteenth Biennial Rept. Calif. Fish Com., 1894, 142, with plate; McCloud River at Baird, Calif.; perhaps a variant of *Salmo shasta*.

361. *Salmo newberrii* Girard.

Klamath River.

Fario or *Salmo newberrii* Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 225; new name for fish misidentified and described as "*Salmo gairdneri* Richardson" in Pacific R. R. Survey, X, 1859, 313, Pl. LXXI, figs. 1-4, Klamath River, Oreg.

362. *Salmo gilberti* (Jordan). *Kern River trout.*

Kern River, Calif.

Salmo gairdneri gilberti Jordan, Thirteenth Biennial Rept. Calif. Fish Com., 1894, 143, with plate; Kern River at Soda Springs, Calif.

363. *Salmo rosei* Jordan and McGregor. *Rose trout.*
 Certain headwaters of Kern River, Calif.
Salmo rosei Jordan, Proc. Acad. Nat. Sci. Phila., LXXVI., 1924, 19,
 Pl. II, Lake Culver, Calif., and neighboring lakes.
364. *Salmo agua-bonita* (Jordan). *Golden trout of south fork of Kern River.*
 South fork of Kern River, Calif.; introduced into Cottonwood Lakes,
 east of Mount Whitney.
Salmo mykiss agua-bonita Jordan, Proc. U. S. Nat. Mus., XV, 1892,
 481, "Volcano or Whitney Creek, California," by error; the original
 type came from the Cottonwood Lakes on the east side of Mount
 Whitney, into which the species had been introduced from the south
 fork of Kern River. This species does not occur in Volcano Creek
 nor in Whitney Creek.
365. *Salmo whitei* Evermann. *Golden trout of western tributaries of Kern
 River; Stewart White trout.*
 Western tributaries of Kern River, Calif.
Salmo whitei Evermann, Bull. U. S. Bur. Fish., XXV, 1905 (May 19,
 1906), 20, Pl. XVI; south fork of Kaweah River, Calif., into which
 the species had been transplanted from Coyote Creek, a western
 tributary of Kern River.
366. *Salmo roosevelti* Evermann. *Golden trout of Volcano Creek; Roosevelt
 trout; Volcano Creek trout.*
 Native only to Volcano Creek, an eastern tributary of Kern River,
 Calif.
Salmo roosevelti Evermann, Bull. U. S. Bur. Fish., XXV, 1905 (May 19,
 1906), 26, Pl. I, Volcano Creek, Kern River Basin, Calif.
367. *Salmo evermanni* Jordan and Grinnell. *San Bernardino trout; Evermann's
 trout.*
 Headwaters of Santa Ana River, Calif., above the falls.
Salmo evermanni Jordan and Grinnell, Proc. Biol. Soc. Wash., XXI,
 1908, 31, Pl. II, south fork of the Santa Ana River, Calif., at 8,200
 feet altitude.
368. *Salmo nelsoni* Evermann. *Lower California trout; Nelson's trout.*
 Mountain streams of northern Lower California.
Salmo nelsoni Evermann, Proc. Biol. Soc. Wash., XXI, 1908, 26, Pl. I,
 San Ramon River, San Pedro Martir Mountains, Lower California.
- Genus 138. CRISTIVOMER Gill and Jordan. *Great Lakes trout.*
 Cristivomer Gill and Jordan, in Jordan, Man. Vert., ed. 2, 1878, 358.
 (*Salmo namaycush* Walbaum).
369. *Cristivomer namaycush* (Walbaum). *Great Lakes trout; Mackinaw
 trout; Longue (Vermont); Togue (Maine); Namaycush; Masamacush.*
 Great Lakes region and lakes of northern New York, New Hampshire,
 and Maine; headwaters of Columbia and Fraser Rivers; streams of
 Vancouver Island, and north to the Arctic Circle.
Namaycush salmon Pennant, Arctic Zool., 1792, 191, Hudson Bay.
Salmo namaycush Walbaum, Artedi Pisc., 1792, 68, Hudson Bay; based
 on the *Namaycush salmon* of Pennant.
Salmo pallidus Rafinesque, Amer. Month. Mag., December, 1817,
 120, Lake Champlain.
Salmo amethystinus Mitchell, Jour. Acad. Nat. Sci. Phila., I, 1818, 410,
 Sault Ste. Marie.
Salmo confinis DeKay, N. Y. Fauna: Fishes, 1842, 238, Louis Lake,
 Hamilton County, N. Y.; Silver Lake, Pa.
Salmo symmetricus Prescott, Am. Jour. Sci. Arts, XI, 1851, 340, Lake
 Winnipiseogee, N. H.
Salmo toma Hamlin, The Togue, in Holmes, 2d Ann. Rept., Maine
 Fish Com., 1862, 109; lakes of Maine.
Salmo adirondacus Norris, Amer. Angler's Book, 1865, 255; Adirondack
 lakes.

370. *Cristivomer siscowet* (Agassiz). *Siscowet.*

Lake Superior, in deep water, where it feeds on the fat ciscoes; a deep-water variant of *C. namaycush*.

Salmo siscowet Agassiz, Lake Superior, 1850, 333, Lake Superior.

Salmo siskawitz Agassiz, in Herbert Frank Forester's Fish and Fishing, 1850, 112, with plate; Lake Superior.

Salmo ursinus Barnston, Rept. Fish. Canada, Lake Superior; reference uncertain.

Genus 137. *SVELVINUS* Richardson. *Charrs.*

Salvelinus Nilsson, Prodr. Icth. Scand., 1832, 7 (*alpinus*); group name.

Salvelinus Richardson, Fauna Bor.-Amer., III, 1836, 169 (*Salmo salvelinus* L.); after Nilsson.

Baione DeKay, N. Y. Fauna: Fishes, 1842, 244 (*B. fontinalis* DeKay).

Umbla Rapp, Fische Bodensees, 1854, 32 (*Salmo umbla*=*alpinus* L.).

371. *Salvelinus fontinalis* (Mitchill). *Brook trout; Speckled trout; Eastern brook trout.*

Maine to the Saskatchewan and northward to Labrador, southward in the Alleghenies to headwaters of the Savannah, Chattahoochee, Catawba, and French Broad Rivers; largely introduced into western streams but not native west of the Mississippi except in Minnesota and Iowa. Some unnamed variants included.

Salmo fontinalis Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 435, near New York City.

Salmo alleghanensis Rafinesque, Ichth. Ohiensis, 1820, 44, brooks falling into the Allegheny and Monongahela Rivers.

Salmo nigrescens Rafinesque, Ichth. Ohiensis, 1820, 45, near the Laurel Hills, Pa.

Salmo canadensis Hamilton Smith, in Griffith's Cuvier, X, 1834, 474, Canada; dots blood red, each "in white circular spot."

Salmo hudsonicus Suckley, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 310, Hudson Bay and tributaries; Labrador; Newfoundland.

372. *Salvelinus agassizii* Garman. *Dublin Pond trout.*

Dublin Pond, N. H.

Salmo agassizii Garman, Nineteenth Rept., Mass. Fish Com., 1885, 20, Dublin Pond (Lake Monadnock), N. H.; Center Pond, N. H.

373. *Salvelinus timagamensis* Henn and Rinkenbach. *Aurora trout.*

White Pine Lake, Clearwater Lake, Timigami region, Ontario, of the St. Lawrence Basin.

Salvelinus timagamensis Henn and Rinkenbach, Ann. Carnegie Mus., XVI, 1925, 131, figs. 1, 2, and 3; White Pine Lake.

374. *Salvelinus aureolus* Bean. *Sunapee trout; American saibling; Golden trout of Sunapee Lake.*

Sunapee Lake, N. H.; Dan Hole Pond, Carroll County, N. H., tributary to Saco River; and Floods Pond, Otis, Me., tributary to Union River; also in Averill Pond, Vt.

Salvelinus aureolus Bean, Proc. U. S. Nat. Mus., X, 1887, 628, Sunapee Lake, N. H.

375. *Salvelinus rossii* (Richardson). *Ross's arctic salmon.*

Lakes of Greenland and Boothia Felix.

?*Salmo hearnei* Richardson, Franklin's First Voy. 1823, 706, and in Fauna Bor.-Amer., III, 1836, 167, Bloody Fall, Coppermine River, Lat. 67°; not certainly recognizable, description imperfect.

Salmo rossii Richardson, App., Ross's Voy., LVI, 1835; and in Fauna Bor.-Amer., III, 1836, 163, Regent's Inlet, Boothia Felix.

Salmo alipes Richardson, loc. cit., LVII, 1835, and in Fauna Bor.-Amer., III, 1836, 169, Lakes in Regent's Inlet, Boothia Felix.

Salmo nitidus Richardson, loc. cit., LVII, 1835, and Fauna Bor.-Amer., 1836, 171; Lake of Boothia Felix.

376. *Salvelinus stagnalis* (Fabricius). *Greenland charr.*

Waters of Greenland, Boothia, and neighboring regions.

Salmo stagnalis Fabricius, Fauna Grænlandica, 1780, 175; Alpine ponds of Greenland; not migratory.

Salmo rivalis Fabricius, loc. cit., 176, Alpine brooks of Greenland; not entering the sea.

Salmo immaculatus H. R. Storer, Jour. Bost. Soc. Nat. Hist., VI, 1850, 364, lower St. Lawrence; Canadian "salmon trout." Name preoccupied. Sea-run forms.

377. *Salvelinus naresi* (Günther).
 Lakes of Arctic America, Discovery Bay, and Cumberland Gulf.
Salmo naresi Günther, Proc. Zool. Soc. Lond., 1877, 476, plate, freshwater lakes near Discovery Bay.
378. *Salvelinus arcturus* (Günther). *Arctic charr*.
 Victoria Lake and Flœberg Beaches of Arctic America, latitude 82° 34'.
Salmo arcturus Günther, Proc. Zool. Soc. Lond., 1877, 294, Pl. XXXII, Victoria Lake, Flœberg Beach.
379. *Salvelinus malma* (Walbaum). *Malma trout*.
 Waters of Kamchatka and Alaska, everywhere abundant; descending to the sea; apparently intergrading southward with *Salvelinus spectabilis*.
Salmo malma Walbaum, Artedi Pisc., 1792, 66, Kamchatka; based on *Malma* of Pennant.
Salmo curilis Pallas, Zoogr. Rosso-Asiat., III, 1811, 251, Curile Islands.
Salmo callaris Pallas, loc. cit., 353, Bering Sea.
Salmo levigatus Pallas, loc. cit., 385, Curile Islands.
Salmo nummifer Cuvier and Valenciennes, Hist. Nat. Poiss., XXI, 1848, 365, Kamchatka; on a drawing by Mertens.
Salmo erythrourhchos Cuvier and Valenciennes, loc. cit., 367, Kamchatka.
Salmo tudes Cope, Proc. Amer. Phil. Soc. Phila., XIII, 1873, 24, Captains Harbor, Unalaska.
Salmo pluvius Hilgendorf, Mitth., Deutsch. Ges. Ostasiens, XI, 1876, 25; clear streams throughout Japan.
380. *Salvelinus malma spectabilis* (Girard). *Dolly Varden trout*; *Oregon charr*; *Bull trout*; *Red-spotted trout*; *Golet*.
 Streams west of the Cascade Range, from the upper Sacramento to Montana and Alaska; in the Columbia basin as far up as Montana and Idaho.
Salmo spectabilis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 218; Fort Dalles, Oreg.
Salmo bairdii Suckley, loc. cit., 309, tributary of Flathead River, Mont.
Salmo parkei Suckley, Ann. Lyc. Nat. Hist. N. Y., VII, 1861, 309, Kootenay River.
Salmo campbelli Suckley, loc. cit., 313, Fort Dalles, Oreg.; substitute for *spectabilis*, preoccupied in *Fario*, not in *Salmo* or *Salvelinus*.
Salmo lordii Günther, Cat., VI, 1866, 148, Skagit River; dwarf specimens.
381. *Salvelinus oxyessa* (Girard). *Oxyessa trout*; *Blueback trout*; *Quasky*.
 Rangeley Lakes, western Maine.
Salmo oxyessa Girard, Proc. Bost. Soc. Nat. Hist., IV, 1854, 262, Oquassa or Oquosoc Lake, Me.
382. *Salvelinus martoni* (Garman). *Lac de Marbre trout*; *Marston trout*.
Salmo martoni Garman, Science, July 14, 1893, Lac de Marbre, Ottawa County, Quebec.

Family 48. COREGONIDÆ. Whitefishes

- Genus 138. *STENODUS* Richardson. *Inconnu*.
Stenodus Richardson, in Back's Narrative Arctic Land Exped., back append., 1836, 521 (*Salmo mackenzii* Richardson).
383. *Stenodus mackenzii* (Richardson). *Inconnu*.
 Delta of MacKenzie River.
Salmo mackenzii Richardson, Franklin's Jour., 1823, 707, MacKenzie River.
- Genus 139. *LEUCICHTHYS* Dybowski. *Lake herrings and ciscoes*.
Argyrosomus Agassiz, Lake Superior, 1850, 330 ("clupeiformis" of DeKay, not of Mitchell = *Salmo harengus* Rich); not of M. de la Pylaie, which is *Sciæna aquila* Cuvier.
Leucichthys Dybowski, Fische Baikal-Wassersysteme, Verk. KK. Zool.-Bot. Gesell. Wien, Bd. XXIV, 1874, 300 (*Salmo omul* Pallas).
Allosomus Jordan, Man. Vert., ed. 2, 1878, 361 (*Coregonus tullibee* Richardson).

Thrissomimus Gill, in Jordan and Evermann, Bull. U. S. Bur. Fish., XXIX, 1909 (1911), 3 (*Coregonus artedi* LeSueur).
Cisco Jordan and Evermann, Bull. U. S. Bur. Fish., XXIX, 1909 (1911), 3 (*Argyrosomus nigripinnis* Gill).

Subgenus **THRISSOMIMUS** Gill.

384. **Leucichthys harengus** (Richardson). *Saginaw Bay herring; Georgian Bay herring.*
 Shallow bays of Lakes Huron and Michigan; Georgian Bay, Saginaw Bay, Green Bay, etc.
Salmo (Coregonus) harengus Richardson, Fauna Bor.-Amer., III, 1836, 210, Pl. XC, fig. 2; Georgian Bay at Penetanguishene, Ontario.
Coregonus clupeiformis Agassiz, Lake Superior, 1850, 339, the Pic;
 Shallow Bays, Michipicoten Island; not *clupeiformis* of Mitchell.
Coregonus albus Agassiz; loc. cit., p. 342, the Pic; not of LeSueur.
385. **Leucichthys osmeriformis** (Smith). *Seneca Lake herring; Seneca Lake smelt.*
 Lakes of central New York, tributary to Lake Ontario.
Coregonus hoyi Bean, Proc. U. S. Nat. Mus., V, 1882, 658; not of Gill.
Coregonus osmeriformis Smith, Bull. U. S. Fish. Com., XIV, 1894, Pl. I, II, Seneca Lake, Skaneateles Lake, N. Y.
386. **Leucichthys sisco** (Jordan). *Cisco of Lake Tippecanoe.*
 Small lakes of Wisconsin and northern Indiana.
Argyrosomus sisco Jordan, Amer. Nat., 1875, 135, Lake Tippecanoe near Warsaw, Ind.
387. **Leucichthys sisco huronius** (Jordan and Evermann). *Lake Huron herring; Blueback or Michigan herring.*
 Lakes Huron and Michigan. The common form from which *L. sisco* is derived.
Argyrosomus huronius Jordan and Evermann, Proc. U. S. Nat. Mus., XXXVI, March 3, 1909, 167, fig. 2, Port Stanley, Ontario.
388. **Leucichthys ontariensis** Jordan and Evermann.
 Lake Ontario and Cayuga Lake, N. Y.
Coregonus clupeiformis DeKay, New York Fauna: Fishes, 1842, 248, pl. 60, fig. 198, Lake Ontario; not *clupeiformis* of Mitchell.
Leucichthys ontariensis Jordan and Evermann, Bull. U. S. Bur. Fish., XXIX, 1909 (1911), 13, Lake Ontario and Cayuga Lake, N. Y.
389. **Leucichthys lucidus** (Richardson). *Great Bear Lake herring.*
 Mackenzie River and tributaries; Great Bear Lake and River.
Salmo (Coregonus) lucidus Richardson, Fauna Bor.-Amer., III, 1836, 207, Pl. XC, fig. 1, with figure, Great Bear Lake.
390. **Leucichthys laurettae** (Bean). *Lauretta whitefish.*
 Lakes and streams of northern and western Alaska.
Coregonus laurettae Bean, Proc. U. S. Nat. Mus., IV, 1881, 156, Point Barrow, Alaska.
391. **Leucichthys alascanus** (Scofield). *Arctic lake herring.*
Argyrosomus alascanus Scofield, in Jordan and Evermann, Fish. North and Mid. Amer., 1898, 2817, northern Alaska near Bering Straits.
392. **Leucichthys pusillus** (Bean). *Least lake herring.*
 Lakes of Yukon basin and Alaska generally.
Coregonus pusillus Bean, Proc. U. S. Nat. Mus., XI, 1888, 526, Kobuk River, Alaska.
393. **Leucichthys athabascæ** Harper and Nichols. *Athabasca herring.*
 Saskatchewan River.
Leucichthys athabascæ Harper and Nichols, Bull. Amer. Mus. Nat. Hist., XLI, 1919, 269, Saskatchewan River.
394. **Leucichthys macrognathus** Harper and Nichols. *Mackenzie River herring.*
 Mackenzie River.
Leucichthys macrognathus Harper and Nichols, Bull. Amer. Mus. Nat. Hist., XLI, 1919, 269, Mackenzie River.

- 395. *Leucichthys entomophagus* Harper and Nichols.**
Mackenzie River.
Leucichthys entomophagus Harper and Nichols, Bull. Amer. Mus. Nat. Hist., XLI, 1919, 267, Mackenzie River.
- 396. *Leucichthys gemmifer* Snyder.** *Cisco of Lake Bonneville.*
Bear River Basin.
Leucichthys gemmifer Snyder, Bull. U. S. Bur. Fish., XXXVI, 1817-18 (May 7, 1919), 4, Bear Lake, near Fish Haven, Idaho.
- 397. *Leucichthys artedi* (LeSueur).** *Lake herring; Erie herring; Common lake herring; Grayback.*
Lake Erie and Lake St. Clair; Lake Huron at Port Huron.
Coregonus artedi LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 231, Lake Erie (at Buffalo) and Niagara River (Lewiston).
- 398. *Leucichthys artedi bisselli* (Bollman).** *Rawson Lake herring; Bissell's herring.*
Glacial lakes of southern Michigan once tributary to Lake Erie.
Coregonus tullibee bisselli Bollman, Bull. U. S. Fish Com., VIII, 1888, 223, Rawson Lake and Howard Lake, Mich.
- 399. *Leucichthys eriensis* (Jordan and Evermann).** *Jumbo herring; Erie great herring.*
Lake Erie and Lake Huron.
Argyrosomus eriensis Jordan and Evermann, Proc. U. S. Nat. Mus., XXXVI, March 3, 1909, 165, fig. 1, Lake Erie at Port Stanley.
- 400. *Leucichthys reighardi* Koelz.**
Lake Michigan.
Leucichthys reighardi Koelz, Occ. Papers, Mus. Zool. Univ. Mich., No. 146, March 22, 1924, 5; Lake Michigan 18 miles off Michigan City, Ind.
- 401. *Leucichthys alpenæ* Koelz.**
Lake Michigan and Lake Huron, including Georgian Bay.
Leucichthys alpenæ Koelz, Occ. Papers, Mus. Zool. Univ. Mich., No. 146, I, March 22, 1924; Lake Michigan off Isle aux Galets.
- Subgenus CISCO Jordan and Evermann.**
- 402. *Leucichthys supernas* Jordan and Evermann.** *Cisco of Lake Superior.*
Deep waters of Lake Superior.
Leucichthys supernas Jordan and Evermann, Bull. U. S. Bur. Fish., XXIX, 1909 (1911), 22, Lake Superior.
- 403. *Leucichthys prognathus* (H. M. Smith).** *Cisco of Lake Ontario; Ontario longjaw; Bloater; Chub.*
Deep waters of Lake Ontario, in depths of 60 fathoms and more.
Coregonus prognathus Smith, Bull. U. S. Fish Com., XIV, 1894, 4, pl. 1, fig. 3, Lake Ontario at Wilson, N. Y.
- 404. *Leucichthys johannæ* (Wagner).** *Lake Michigan cisco; Bloater of Lake Michigan.*
Northwestern part of Lake Huron in deep water, and also for the whole length of Lake Michigan.
Argyrosomus johannæ Wagner, Science, N. S., XXXI, No. 807, June 17, 1910, 957-958; Lake Michigan in about 25 fathoms, some 18 miles off Racine, Wis.
- 405. *Leucichthys nigripinnis* (Gill).** *Blackfin of Lake Michigan.*
Deep waters of Lake Michigan and certain small lakes in Wisconsin.
Argyrosomus nigripinnis Gill, in Hoy, Trans. Wis. Acad. Sci., I, 1872, 100, Lake Michigan off Racine; name only.
Coregonus nigripinnis Jordan, Man. Vert., ed. 2, 1878, 275; description.
- 406. *Leucichthys cyanopterus* Jordan and Evermann.**
Deep waters of Lake Superior.
Leucichthys cyanopterus Jordan and Evermann, Bull. U. S. Bur. Fish., XXIX, 1909 (1911), 27, fig. 13, Lake Superior, off Marquette, Mich.

- 407. *Leucichthys hoyi* (Gill). *Cisco of Lake Michigan; Kiyi; Chub; Mooneye cisco.***
 Lake Michigan in deep water.
Argyrosomus hoyi Gill, in Hoy, Trans. Wis. Acad. Sci., I, 1872, 100,
 Lake Michigan off Racine; no description.
Argyrosomus hoyi Gill, in Jordan, Amer. Nat., March, 1875, 135, Lake Michigan off Racine, Wis.; description same specimens.
- 408. *Leucichthys kiyi* Koelz. *Kiyi; Mooneye; Waterbelly.***
 Lake Michigan.
Leucichthys kiyi Koelz, Occ. Papers, Mus. Zool. Univ. Mich., No. 104,
 Dec. 31, 1921; Lake Michigan, 12 miles E. by S. of the mouth of Sturgeon Bay ship channel in about 60 fathoms.
- 409. *Leucichthys birgei* Wagner. *Cisco of Green Lake.***
 Green Lake, Wisconsin.
Leucichthys birgei Wagner, Bull. Wis. Nat. Hist. Soc., IX, 1911, 73,
 Green Lake, Wis., 28 meters below the surface.
- 410. *Leucichthys zenithicus* (Jordan and Evermann). *Longjaw of Lake Superior.***
 Lake Superior, in deep water; possibly in other lakes.
Argyrosomus hoyi Milner, Rept. U. S. Fish Com., II, 1872-73 (1874),
 86, Lake Superior at Outer Island, Wis.; not of Gill, Hoy, or Jordan,
 and not original type.
Argyrosomus zenithicus Jordan and Evermann, Proc. U. S. Nat. Mus., XXXVI, March 3, 1909, 169, fig. 3, Lake Superior between Duluth and Isle Royale.
- Subgenus ALLOSOMUS Jordan.**
- 411. *Leucichthys manitoulinus* Jordan and Evermann. *Manitoulin tullibee.***
 North channel of Lake Huron, and probably lakes of Minnesota.
Argyrosomus tullibee, Evermann ar i Smith, Rept. U. S. Fish Com., XX,
 1894, 320, pl. 28; in part, the plate being tullibee.
Leucichthys manitoulinus Jordan and Evermann, Bull. U. S. Bur. Fish.,
 XXIX, 1909, 31, fig. 16, Blind River, North channel, Lake Huron.
- 412. *Leucichthys tullibee* (Richardson). *Tullibee; Tulipi.*
 Lake Winnipeg Basin, perhaps entering Lake Superior Basin.
Salmo (Coregonus) tullibee Richardson, Fauna Bor.-Amer., III, 1836,
 201, Cumberland House, Pine Island Lake, near Lake Winnipeg.
Leucichthys nippigon Koelz, Occ. Papers Mus. Zool. Univ. Mich., No.
 158, April 6, 1925; Lake Nipigon, Lake Winnipeg.**
- Genus 140. COREGONUS Linnaeus. Whitefishes; Lavarets.**
Coregonus Artedi, Genera Pisc., 1738, 9.
Coregonus Linnaeus, Syst. Nat., Ed. X, 1758, 310 (*Salmo lavaretus* L.).
Tripteronotus Lacépède, Hist. Nat. Poiss., V, 1803, 48 (*T. hauin*
Lacépède=*Salmo lavaretus* L.).
Coregonus Lacépède, loc. cit., X, 263 (*Salmo lavaretus* L.).
Coregonus Cuvier, Régne Anim. Ed. I, 1817, 162 (*thymallus*, *lavaretus*,
etc.).
- 413. *Coregonus clupeaformis* (Mitchill). *Whitefish; Labrador whitefish;*
Lake Superior whitefish; Manitoba whitefish; Musquaw River white-
fish; Whiting of Lake Winnepeaukee; Shad of Lake Champlain.
 Great Lakes, excepting Lake Erie; and neighboring waters.
Salmo clupeaformis Mitchell, Amer. Month. Mag., II, 1818, 321, Falls
 of St. Mary, northern extremity of Lake Huron.
Salmo otsego Clinton, account of the *Salmo otsego* or the Otsego basse,
 1822, 1, with plate, Otsego Lake.
Coregonus labradoricus Richardson, Fauna Bor.-Amer., III, 1836, 206,
 Musquaw River, Labrador.
Salmo (Coregonus) sapidissimus Agassiz, Lake Superior, 1850, 344,
 Lake Superior.
Salmo latior Agassiz, loc. cit., 348, Lake Superior.
Coregonus neohantoni Prescott, Amer. Jour. Sci. Arts, XI, 1851, 343,
 Lake Winnepeaukee, N. H.
?Coregonus richardsonii Günther, Cat., VI, 1866, 185; Arctic America.**

414. *Coregonus clupeaformis albus* (LeSueur). *Lake Erie whitefish; Common whitefish.*
 Lake Erie and Lake St. Clair; Lake Ontario and various lakes to western New York and New Hampshire. New York; probably not separable from *Coregonus clupeaformis*, although local differences appear.
Coregonus albus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 232, Lake Erie.
415. *Coregonus nelsonii* Bean. *Humpback whitefish; Nelson's whitefish.*
 Alaska, from Bristol Bay northward.
Coregonus nelsonii Bean, Proc. U. S. Nat. Mus., VII, 1884, 48, Nulato, Alaska.
- Genus 141. **PROSOPIUM** Milner.
Prosopium Milner, in Jordan, Man. Vert., ed. 2, 1878, 361 (*Coregonus quadrilateralis* Richardson).
416. *Prosopium quadrilaterale* (Richardson). *Menominee whitefish; Pilotfish; Round whitefish; Shadwaite.*
 Alaska and upper Great Lakes to New England, in lakes.
Coregonus quadrilateralis Richardson, Franklin's Narrative, 1823, 714, Pl. XXV, fig. 2, Fort Enterprise, British America.
Coregonus novae-angliae Prescott, Amer. Jour. Sci. Arts, XI, 1851, 342, Lake Winnipesaukee, N. H.
417. *Prosopium kennicotti* (Milner). *Kennicott's whitefish.*
 Mackenzie River, Canada; Yukon River and other streams of the Alaska region; Lake Bennett, Alaska.
Coregonus kennicotti Milner, in Jordan and Gilbert, Synopsis, 1883, 298, Fort Good Hope, British America.
418. *Prosopium preblei* (Harper and Nichols).
 Mackenzie River, Canada.
Coregonus preblei Harper and Nichols, Bull. Amer. Mus. Nat. Hist., XLI, 1919, 266, Pl. XV, fig. 2, Mackenzie River.
419. *Prosopium stanleyi* (Kendall). *Stanley's whitefish.*
 Lakes of northern Maine.
Coregonus stanleyi Kendall, Bull. U. S. Fish Com., XXII, 1902 (1904), 366, with figure, thoroughfare between Mud and Cross Lakes, Aroostook County, Me.
420. *Prosopium williamsoni* (Girard). *Rocky Mountain whitefish.*
 Rivers of Sierra Nevada and west slope of the Rocky Mountains, from the Fraser and the Columbia to the Truckee and other streams of the Lahontan Basin of Nevada; abundant, especially in lakes of northern Idaho, western Montana, Washington, and Utah.
Coregonus williamsoni Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 136, Des Chutes River, Oreg.
421. *Prosopium williamsoni cismontanus* (Jordan). *Yellowstone whitefish.*
 Streams of the Rocky Mountain region tributary to the upper Missouri.
Coregonus williamsoni cismontanus Jordan, Bull. U. S. Fish Com., IX, 1889, 49, pl. 9, figs. 8, 9, Horsethief Creek, Madison River, Mont.
 Hardly distinct from *P. williamsoni*.
422. *Prosopium coulteri* (Eigenmann and Eigenmann). *Coulter's whitefish.*
 Headwaters of the Columbia.
Coregonus coulteri Eigenmann and Eigenmann, Amer. Nat., Nov., 1892, 961, Kicking Horse River at Field, British Columbia.
423. *Prosopium couesi* (Milner).
 Headwaters of Saskatchewan River.
Coregonus couesi Milner, Rept. U. S. Fish Com., II, 1872-73 (1874), 88, Chief Mountain Lake, Mont.
424. *Prosopium spilonotus* (Snyder). *Bonneville whitefish.*
 Bear River Basin.
Coregonus spilonotus Snyder, Bull. U. S. Bur. Fish., XXXVI, 1917-18 (May 7, 1919), 6, Bear Lake, near Fish Haven, Idaho.

- 425. *Prosopium abyssicola* (Snyder).** *Bear Lake whitefish.*
Bear Lake, Idaho.
Coregonus abyssicola Snyder, Bull. U. S. Bur. Fish., XXXVI, 1917-18
(May 7, 1919), 8, Bear Lake, near Fish Haven, Idaho.
- Genus 142. *IRILLION* Jordan.**
Irillion Jordan, Proc. Acad. Nat. Sci. Phila., LXX, 1918, 342 (*Coregonus oregonius* Jordan and Snyder).
- 426. *Irillion oregonius* (Jordan and Snyder).** *Chiselmouth jack.*
Mountain streams of Oregon tributary to the Columbia.
Coregonus oregonius Jordan and Snyder, Proc. U. S. Nat. Mus., XXXVI, 1911, 425, with fig., McKenzie River, Oreg.

Family 49. THYMALLIDÆ. Graylings

- Genus 143. *THYMALLUS* Cuvier.** *Graylings.*
Thymallus Cuvier, Règne Anim., ed. 2, II, 1829, 306 (*Salmo thymallus* Linnaeus); not *Thymalus* Latreille, 1802, a genus of Coleoptera.
Choregonus Minding, Lehrbuch Naturgesch. Fische, 1832, 119 (*Salmo thymallus* L.) substitute for *Thymallus*, regarded as preoccupied.
Orthocolus Gistel, Thiereich, 1848, XI, 1838 (*Salmo thymallus* L.); substitute for *Thymallus*, regarded as preoccupied.
- 427. *Thymallus signifer* (Richardson).** *Arctic grayling; Poisson bleu.*
Mackenzie River to Alaska and the Arctic Ocean.
Coregonus signifer Richardson, Franklin's Jour., 1823, 711; Winter Lake near Fort Enterprise.
Coregonus thymalloides Richardson, Fauna Bor.-Amer., III, 1836, 714, Winter River.
- 428. *Thymallus tricolor* Cope.** *Michigan grayling.*
Streams of northern Michigan; formerly abundant in Au Sable River, Jordan River, and other streams in the southern peninsula, and Otter Creek, near Keweenaw, in the northern peninsula.
? *Thymallus ontariensis* Cuvier and Valenciennes, Hist. Nat. Poiss., XXI, 1848, 452; said to have been brought by Milbert from Lake Ontario; most likely from Lake Geneva, Switzerland.
Thymallus tricolor Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 80, Au Sable River, Mich.
- 429. *Thymallus montanus* Milner.** *Montana grayling.* A hardy and attractive aquarium fish.
Madison and Gallatin Rivers, Montana.
Thymallus montanus Milner, Rept. U. S. Fish Com., II, 1872-73 (1874), 741, tributary of Missouri River at Camp Baker.

Family 50. OSMERIDÆ. Smelts

- Genus 144. *MALLOTUS* Cuvier.** *Capelins.*
Mallotus Cuvier, Règne Anim., ed. 2, II, 1829, 305 (*Clupea villosa* Müller).
- 430. *Mallotus villosus* (Müller).** *Capelin; Lodde.*
Arctic America, south to Cape Cod and southern Alaska; Kamchatka.
Clupea villosa Müller, Prodr. Zool. Dan., 1777, 245, Greenland.
Salmo arcticus Fabricius, Fauna Greenlandica, 1780, 177, Greenland.
Salmo grönlandicus Bloch, Ichthyologie, VIII, 1794, pl. 99, Greenland.
Salmo socialis Pallas, Zoogr. Ross.-Asiat., III, 389, islands between Asia and America.
Osmerus microdon Cuvier and Valenciennes, Hist. Nat. Poiss., XXI, 1848, 385, no locality.
- Genus 145. *THALEICHTHYS* Girard.** *Eulachon.*
Thaleichthys Girard, Pac. R. R. Surv., X, 1858, Fishes, 325 (*T. stevensi* Girard = *Salmo* (*Mallotus*) *pacificus* Richardson).
- 431. *Thaleichthys pacificus* (Richardson).** *Eulachon; Candlefish; Oolachan.*
Oregon to Alaska.
Salmo (*Mallotus*) *pacificus* Richardson, Fauna Bor.-Amer., III, 1836, 226, Columbia River.
Thaleichthys stevensi Girard, Pac. R. R. Surv., X, 1858, 325, pl. 75, figs. 1-4, Puget Sound.

432. *Thaleichthys albatrossis* (Jordan and Gilbert). *Kodiak smelt.*
 Off Kodiak Island, Alaska.
Osmerus albatrossis Jordan and Gilbert, in Jordan and Evermann,
 Fishes North and Mid. Amer., 1898, 2823, Kodiak Island; probably
 not distinct from *Thaleichthys pacificus*.
- Genus 146. **SPIRINCHUS** Jordan and Evermann.
Spirinchus Jordan and Evermann, Fishes North and Mid. Amer., I,
 1896, 522 (*Osmerus thaleichthys* Ayres).
433. *Spirinchus thaleichthys* (Ayres).
 Pacific coast of America from San Francisco northward to Bristol Bay,
 Alaska.
Osmerus thaleichthys Ayres, Proc. Calif. Acad. Nat. Sci., II, 1860, 62,
 San Francisco.
- Genus 147. **ALLOSMERUS** Hubbs.
Allosmerus Hubbs, Proc. Biol. Soc. Wash., XXXVIII, 1925, 53 (*Osmerus attenuatus* Lockington).
434. *Allosmerus attenuatus* (Lockington).
 Coast of California, from San Francisco northward.
Osmerus attenuatus Lockington, Proc. U. S. Nat. Mus., III, 1880, 66,
 San Francisco.
- Genus 148. **OSMERUS Lacépède.** *Smelts.*
Osmerus Lacépède, Hist. Nat. Poiss., V, 1803, 229 (*Salmo eperlanus* L.).
Eperlanus Gaimard, Voy. Island and Grönland, 1851 (*Salmo eperlanus* L.).
435. *Osmerus dentex* Steindachner. *Rainbow herring.*
 Coast of Alaska and south on the Asiatic coast to northern China.
Osmerus dentex Steindachner, Sitzungsber. Kais. Akad. Wiss. Wien, LXI,
 1870, 429, northern China.
436. *Osmerus mordax* (Mitchill). *American smelt; Icefish.*
 Atlantic coast of United States, from Virginia northward to Gulf of
 St. Lawrence; Lakes Champlain and Memphremagog.
Atherina mordax Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 446,
 New York.
Osmerus viridescens Le Sueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 230,
 coast of Maine.
437. *Osmerus mordax spectrum* (Cope). *Wilton smelt.*
 Wilton Pond, Kennebec County, Me.
Osmerus spectrum Cope, Proc. Amer. Phil. Soc., XI, 1870, 490, Wilton
 Pond, Kennebec County, Me.
438. *Osmerus mordax abbotti* (Cope). *Cobessicontic smelt.*
 Landlocked in Cobessicontic Lake, Kennebec County, Me.
Osmerus abbotti Cope, Proc. Amer. Phil. Soc., XI, 1870, 490, Cobessi-
 contic Lake, Me.
439. *Osmerus sergeanti* ²² Norris.
 New York Bay and southward.
Osmerus sergeanti Norris, Proc. Acad. Nat. Sci. Phila., XX, 1868, 93,
 Schuylkill River, N. J.
- Genus 149. **HYPOMESUS** Gill. *Surf smelts.*
Mesopus ²³ Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 14 (*Argentina pretiosa* Girard).
Hypomesus Gill, loc. cit., 15 (*Argentina pretiosa* Girard).
440. *Hypomesus pretiosus* (Girard). *Surf smelt.*
 Coast of California and Oregon, from Monterey northward.
Argentina pretiosa Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 155,
 San Francisco.
Osmerus elongatus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 17, San
 Francisco.

²² The fish is described, but not named, in Proc. Acad. Nat. Sci. Phila., XIII, 1861, 58. It is named in Proc. Acad. Nat. Sci. Phila., XX, 1868, 93.

²³ *Mesopus*, a name left in type through failure to correct proof sheets. *Hypomesus* of same date, a page lower, chosen by author as name.

441. *Hypomesus olidus* (Pallas). *Pond smelt.**Alaska and Kamchatka.**Salmo (Osmerus) olidus* Pallas, Zoogr. Rosso.-Asiat., III, 1811, 391,
lakes and rivers of Kamchatka.*?Osmerus oligodon* Kner, Denkschr. Akad. Wiss. Wien, 1865, 185, tab.
4, fig. 1.Genus 150. *THEROBROMUS* Lucas. *Ghost smelt.**Therobromus* Lucas, in Jordan and Gilbert, Fishes of Bering Sea, 1898,
440 (*T. callorhini* Lucas).442. *Therobromus callorhini* Lucas. *Ghost smelt.**Bering Sea.**Therobromus callorhini* Lucas, in Jordan and Gilbert, Fishes of Bering
Sea, in Jordan, Fur Seals and Fur-Seal Islands, Pt. III, 1898, 440,
Bering Sea, in southern portion; known only from skeletons and
bones found in the stomachs of fur seals.

Family 51. ARGENTINIDÆ. Argentines

Genus 151. *ARGENTINA* Linnaeus. *Argentines.**Argentina* (Artedi) Linnaeus, Syst. Nat., Ed. X, 1758, 315 (*A. sphyræna*
L.).*Silus Reinhardt*, Bemærk., Skandinav. Ichthy., 1833, 11 (*Argentina silus*
Ascanius).*Acantholepis Kröyer*, Danmarks Fiske, III, 1846-49, 98 (*Argentina silus*
Ascanius).443. *Argentina silus* Ascanius.Northern Europe; occasionally taken on the Grand Banks and off the
coast of Maine.*Argentina silus Ascanius*, Icon. Rev. Nat., 1763, Pl. III, 3, 24. Norway.*Silus ascanii* Reinhardt, Bemærk., Skandinav. Ichth., 1833, 11, Norway.*Argentina syrtensium* Goode and Bean, Proc. U. S. Nat. Mus., I, 1878,
261, off Sable Island in 200 fathoms; from stomach of *Phycis tenuis*.444. *Argentina striata* Goode and Bean.

Gulf of Mexico.

Argentina striata Goode and Bean, Oceanic Ichth., 1895, 52, fig. 62,
Gulf Stream.445. *Argentina sialis* Gilbert.

Coast of California.

Argentina sialis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 56, Albatross
station 3017 in 58 fathoms.

Genus 152. LEUROGLOSSUS Gilbert.

Leuroglossus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 57 (*L. stilbius*
Gilbert).446. *Leuroglossus stilbius* Gilbert.

Coast of California, in deep water.

Leuroglossus stilbius Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 57,
from Albatross stations 2997 and 2998 in 221 and 40 fathoms.

Family 52. MICROSTOMIDÆ

Genus 153. NANSENIA Jordan and Evermann.

Nansenia Jordan and Evermann, Fishes North and Mid. Amer., 1896,
528 (*Microstomus grænlandica* Reinhardt).447. *Nansenia grænlandica* (Reinhardt).

Greenland.

Microstomus grænlandica Reinhardt, Vidensk. Selsk.. Naturv. Math.
Afhandl., VIII, 1841, LXXIV, Greenland.

Family 53. BATHYLAGIDÆ

Genus 154. BATHYLAGUS Günther.

Bathylags Günther, Ann. Mag. Nat. Hist., II, 1878, 248 (*B. antarcticus*
Günther).

448. *Bathylagus benedicti* Goode and Bean.
Gulf Stream.
Bathylagus benedicti Goode and Bean, Oceanic Ichth., 1895, 55, fig. 64,
Gulf Stream.
449. *Bathylagus euryops* Goode and Bean.
Gulf Stream.
Bathylagus euryops Goode and Bean, Oceanic Ichth., 1895, 55, fig. 63,
Gulf Stream, in about lat. 40° N. and long. 70° W.
450. *Bathylagus pacificus* Gilbert.
Coast of Washington, in deep water.
Bathylagus pacificus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 55,
Albatross stations 3071 and 3074, off coast of Washington.
451. *Bathylagus borealis* Gilbert.
Coast of Alaska, in deep water.
Bathylagus borealis Gilbert, Rept. U. S. Fish Com., XIX, 1893 (1896),
402, north of Unalaska Island at *Albatross* station 3327 in 322
fathoms.
452. *Bathylagus milleri* Jordan and Gilbert.
Off coast of southern California in deep water.
Bathylagus milleri Jordan and Gilbert, in Jordan and Evermann, Fishes
North and Mid. Amer., 1898, 2825, Cortez Banks, off San Diego, Calif.
- Genus 155. *XENOPHTHALMICHTHYS* Regan.
Xenophthalmichthys Regan, Ann. Mag. Nat. Hist., ser. 9, XV, 1925, 59
(*X. danæ* Regan).
453. *Xenophthalmichthys danæ* Regan.
Depths of the Caribbean Sea.
Xenophthalmichthys danæ Regan, Ann. Mag. Nat. Hist., ser. 9, XV,
1925, 59, off Santa Lucia in deep water.

Suborder STOMIATIOIDEI

Family 54. STOMIATIDÆ

- Genus 156. *STOMIAS* Cuvier.
Stomias Cuvier, Règne Anim., Ed. I, 1817, 184 (*Esox boa* Risso).
454. *Stomias ferox* Reinhardt.
Greenland and southward; Gulf Stream, from Bahama Channel to the
Grand Banks.
Stomias ferox Reinhardt, Vidensk. Selsk. Naturv. Math. Afhandl., X,
1842, 77, Greenland.
455. *Stomias affinis* Günther.
Sombrero Island, West Indies.
Stomias affinis Günther, Challenger Rept., XXII, 1887, 205, Pl. LIV,
Fig. A; Sombrero Island, West Indies.
456. *Stomias colubrinus* Garman.
Off Panama in deep water.
Stomias colubrinus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
275, Pl. LVII, fig. 7, station 3380, lat. 6° 17' N., long. 82° 5' W. in
1,872 fathoms.
457. *Stomias hexagonatus* Garman.
Eastern tropical Pacific in deep water.
Stomias hexagonatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
Pl. LVI, fig. 5; various stations in the Pacific.
458. *Stomias atriventer* Garman.
Eastern Pacific in deep water.
Stomias atriventer Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 277,
Pl. LVI, fig. 4; *Albatross* station 3433, lat. 25° 76' 15" N., long. 109°
48' W., in 1,218 fathoms.
- Genus 157. *DACTYLOSTOMIAS* Garman.
Dactylostomias Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 279
(*D. filifer* Garman).

459. Dactylostomias filifer Garman.

Off Panama in deep water.

Dactylostomias filifer Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 279, Pl. LVI, fig. 6, *Albatross* stations 3413, 3414, and 3418.

Genus 158. ZASTOMIAS Gilbert.

Zastomias Gilbert, Proc., U. S. Nat. Mus., XLVIII, 1915, 322 (*Z. scintillans* Gilbert).

460. Zastomias scintillans Gilbert.

Coast of California; Monterey Bay in deep water.

Zastomias scintillans Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 322, pl. 15, off Monterey Bay, Calif.

Genus 159. ECHIOSTOMA Lowe.

Echiostoma Lowe, Proc. Zool. Soc. Lond., 1843, 87 (*E. barbatum* Lowe).

Hyperchoristus Gill, Proc. U. S. Nat. Mus., VI, 1883, 256 (*H. tanneri* = *Echiostoma barbatum*).

461. Echiostoma barbatum Lowe.

Madeira; Gloucester, Mass.; and southward in the Gulf Stream to the old Bahama Channel.

Echiostoma barbatum Lowe, Proc. Zool. Soc. Lond., 1843, 88, Madeira.

Hyperchoristus tanneri Gill, Proc. U. S. Nat. Mus., VI, 1883, 256, Gulf Stream at *Albatross* station 2083, at 40° 26' 40" N., 60° 58' W., in 956 fathoms.

462. Echiostoma margaritum Goode and Bean.

Middle of the Gulf of Mexico.

Echiostoma margarita Goode and Bean, Oceanic Ichth., 1895, 109, fig. 131, Gulf of Mexico.

Genus 160. GRAMMATOSTOMIAS Goode and Bean.

Grammatostomias Goode and Bean, Oceanic Ichth., 1895, 110 (*G. dentatus* Goode and Bean).

463. Grammatostomias dentatus Goode and Bean.

Gulf Stream.

Grammatostomias dentatus Goode and Bean, Oceanic Ichth., 1895, 110, fig. 133, Gulf Stream.

Genus 161. PHOTONECTES Günther.

Lucifer Döderlein, Weigmann's Archiv, 1882, 26 (*L. albipinnis* Döderlein); name preoccupied in Crustacea.

Photonectes Günther, Challenger Rept., XXII, 1887, 212 (*L. albipinnis* Döderlein).

464. Photonectes gracilis Goode and Bean.

Martinique.

Photonectes gracilis Goode and Bean, Oceanic Ichth., 1895, 112, fig. 137, off Martinique at *Blake* station XL in 472 fathoms.

Family 55. IDIACANTHIDÆ.**Genus 162. IDIACANTHUS Peters.**

Idiacanthus Peters, Monatsber. Akad. Wiss. Berlin, 1876, 846 (*I. fasciola* Peters).

Bathyophis Günther, Ann. Mag. Nat. Hist., ser. 5, II, 1878, 181 (*B. ferox* Günther).

465. Idiacanthus ferox (Günther).

North Atlantic, nearly midway, in 2,750 fathoms.

Bathyophis ferox Günther, Ann. Mag. Nat. Hist., ser. 5, II, 1878, 181, middle of North Atlantic in 2,750 fathoms.

466. Idiacanthus antrostomus Gilbert.

Coast of southern California in deep water.

Idiacanthus antrostomus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 54, off southern California, at *Albatross* station 2980 in 603 fathoms.

Family 56. MALACOSTEIDÆ

Genus 163. MALACOSTEUS Ayres.

Malacosteus Ayres, Jour. Bost. Soc. Nat. Hist., 1849, 53 (*M. niger* Ayres).

467. *Malacosteus niger* Ayres.

Open sea; Gulf Stream and southward to Barbados.

Malacosteus niger Ayres, Jour. Bost. Soc. Nat. Hist., 1849, 53, Gulf Stream, 42° N., 60° W.

Family 57. ASTRONESTHIDÆ

Genus 164. ASTRONESTHES Richardson.

Astronesthes Richardson, Ichth. Voy. *Sulphur*, 1845, 97 (*A. nigra*, Richardson).

Phænodon Lowe, Proc. Zool. Soc. Lond., XVIII, 1850, 250 (*P. ringens* Lowe).

468. *Astronesthes richardsoni* (Poey).

Deep waters of West Indies; Indian and Pacific Oceans.

Chauliodus richardsoni Poey, Memorias, I, 1852, 176, Cuba.

469. *Astronesthes gemmifer* Goode and Bean.

Gulf Stream.

Astronesthes gemmifer Goode and Bean, Oceanic Ichth., 1895, 105, fig. 124, Grand Banks, lat. 44° 25' N., long 53° 12' W., in 300 fathoms, in halibut stomach.

470. *Astronesthes niger* Richardson.

Deep waters of all seas.

Astronesthes niger Richardson, Ichth. Voy. *Sulphur*, 1845, 97, Atlantic.

Family 58. STYLOPTHALMIDÆ

Genus 165. STYLOPTHALMUS Brauer.

Stylophthalmus Brauer, Zool. Anzeiger, XXV, 1902, 298 (*S. paradoxus* Brauer).

471. *Stylophthalmus paradoxus* Brauer.

Open sea, north to southern California (San Pedro, W. F. Thompson).

Stylophthalmus paradoxus Brauer, Tiefseefische Valdivia Exped. (Zool. Anzeiger, XXV), 1902, 298; deep sea off the Cape of Good Hope and the Indian Ocean.

Family 59. CHAULIODONTIDÆ

Genus 166. CHAULIODUS Bloch and Schneider.

Chauliodus Bloch and Schneider, Syst. Ichth., 1801, 430 (*C. sloani* Bloch and Schneider).

Leptodes Swainson, Nat. Hist. Anim., II, 298 (*Chauliodes sloani* Bloch and Schneider).

472. *Chauliodus sloani* Bloch and Schneider.

Mediterranean and deep waters of the Atlantic; Georges Banks.

Chauliodus sloani Bloch and Schneider, Syst. Ichth., 1801, 430, Atlantic; after *Vipera marina* of Catesby.

473. *Chauliodus macouni* Bean.

Coast of California to British Columbia and Queen Charlotte Islands.

Chauliodus macouni Bean, Proc. U. S. Nat. Mus., XIII, 1890, 44, off Cape St. James, Queen Charlotte Islands.

474. *Chauliodus barbatus* Garman.

Off Pacific coast of Panama.

Chauliodus barbatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 271, Pl. K, figs. 2, 2a; off Panama.

Family 60. GONOSTOMIDÆ

- Genus 167. **GONOSTOMA** Rafinesque.
Gonostoma Rafinesque, Indice, 1810, 64 (*G. denudatum* Rafinesque).
475. **Gonostoma denudatum** Rafinesque.
 Gulf of Mexico at *Albatross* station 2665; off Morocco and Cape Verdea; Mediterranean Sea.
Gonostoma denudata Rafinesque, Indice, 1810, 65, Palermo.
- Genus 168. **ZAPHOTIAS** Goode and Bean.
Bonapartia Goode and Bean, Oceanic Ichth., 1895, 102 (*B. pedaliota* Goode and Bean); name preoccupied.
Zaphotias Goode and Bean, in Jordan and Evermann, Fish. North and Mid. Amer., 1898, 2826 (*Bonapartia pedaliota* Goode and Bean).
476. **Zaphotias pedaliotus** (Goode and Bean).
 Gulf Stream.
Bonapartia pedaliota Goode and Bean, Oceanic Ichth., 1895, 102, fig. 120, *Albatross* station 2642, in $25^{\circ} 20' 30''$ N. Lat., $79^{\circ} 58'$ W. Long., depth 217 fathoms.
- Genus 169. **CYCLOTHONE** Goode and Bean.
Cyclothona Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1883, 221 (*C. lusca* Goode and Bean).
Sigmops Gill, Proc. U. S. Nat. Mus., VI, 1883, 256 (*S. stigmaticus* Gill).
Neostoma Filhol, La Nature, 1884, 184 (*N. bathyphilum* Filhol).
- Subgenus **CYCLOTHONE** Goode and Bean.
477. **Cyclothona microdon** (Günther).
 Bermuda; Atlantic, Pacific, and Antarctic Oceans; Pacific coast from Oregon to Panama.
Gonostoma microdon Günther, Ann. Mag. Nat. Hist., II, 1878, 188, near Bermuda.
Cyclothona lusca Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1883, 221, Gulf Stream.
478. **Cyclothona bathyphila** Vaillant.
 Gulf of Gascony and off the Azores; western Atlantic.
Neostoma bathyphilum Filhol, La Nature, 1884, 184; name and rough figure only.
Cyclothona bathyphila Vaillant, Expéd. Sci. Travailleur et Talisman, Poissons, 1888, 96, pl. 8, figs. 1, 1a; Gulf of Gascony, Azores.
479. **Cyclothona acclinidens** Garman.
 Off California in deep water.
Cyclothona acclinidens Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 247, Pl. T, fig. 4; numerous stations of *Albatross* off California coast.
480. **Cyclothona signata** Garman.
 Off Panama in deep water.
Cyclothona signata Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 246, Pl. T, fig. 3; *Albatross* station 3382, off Panama.
- Subgenus **SIGMOPS** Gill.
481. **Cyclothona elongata** (Günther).
 New Guinea; Banda; American coast; Gulf Stream; Indian Ocean.
Gonostoma elongatum Günther, Ann. Mag. Nat. Hist., II, 1878, 187, New Guinea; Banda.
Sigmops stigmaticus Gill, Proc. U. S. Nat. Mus., VI, 1883, 256, Gulf Stream, in $38^{\circ} 19' 26''$ N., $68^{\circ} 20' 20''$ W.
- Genus 170. **YARRELLA** Goode and Bean.
Yarrella Goode and Bean, Oceanic Ichth., 1895, 103 (*Y. blackfordi* Goode and Bean).
482. **Yarrella blackfordi** Goode and Bean.
 Gulf Stream.
Yarrella blackfordi Goode and Bean, Oceanic Ichth., 1895, 103, fig. 121, Gulf Stream at *Albatross* station 2376, in 324 fathoms.
- Genus 171. **LYCHNOPOLES** Garman.
Lychnópoles Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 244 (*L. argenteolus* Garman).

483. *Lychnopoles argenteolus* Garman.

Off Panama in deep water.

Lychnopoles argenteolus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 244, Pl. LIII, figs. 4, 4a; off Panama.

Family 61. MAUROLICIDÆ

Genus 172. MAUROLICUS Cocco.

Maurolicus Cocco, Nuovi Ann. Sci. Nat., II, 1838, 32 (*M. amethystina punctatus* Cocco).484. *Maurolicus pennanti* (Walbaum). *Pearlsides*.

Open seas, widely distributed; New England coast at Nahant, Province-town, Woods Hole, etc.

Argentina pennanti Walbaum, Artedi Pisc., 1792, 47, England; after "sheppy Argentine" of Pennant.485. *Maurolicus oculatus* Garman.

Deep water off Pacific coast of Mexico.

Maurolicus oculatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 241, Pl. LIII, fig. 3; off Pacific coast of Mexico.

Genus 173. VINCIGUERRIA Jordan and Evermann.

Vinciguerria Jordan and Evermann, in Goode and Bean, Oceanic Ichth., 1895, 513 (*V. attenuata* Cocco).*Zalarges* Jordan and Williams, in Jordan and Starks, Proc. Calif. Acad. Sci., ser. 2, V, 1895, 793 (*Z. nimbarius* Jordan and Williams).486. *Vinciguerria attenuata* (Cocco).

Open Atlantic, west to the Bahama regions.

Maurolicus attenuatus Cocco, Nuovi Ann. Sci. Nat., II, 1838, 33, coast of Italy.487. *Vinciguerria lucetia* (Garman).

Eastern Pacific, in deep water.

Maurolicus lucetius Garman, Deep Sea Fishes, in Mem. Mus. Comp. Zool., XXIV, 1899, 242, Pl. T, fig. 2; various Albatross stations, eastern Pacific.488. *Vinciguerria nimbaria* (Jordan and Williams).

Open sea between Oregon and Hawaii.

Zalarges nimbarius Jordan and Williams, Proc. Calif. Acad. Sci., ser. 2, V, 1895 (1896), 793, open Pacific.

Genus 174. VALENCIENELLUS Jordan and Evermann.

Valenciennellus Jordan and Evermann, Fishes North and Mid. Amer., 1896, 577 (*Maurolicus tripunctulatus* Esmark).489. *Valenciennelus tripunctulatus* (Lütken).

Madagascar; Denmark Straits between Greenland and Iceland.

Maurolicus tripunctulatus Esmark, Christiania Vid. Selsk. Forh., 1870, 488, Madagascar.490. *Valenciennellus stellatus* Garman.

Deep water off California.

Valenciennellus stellatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 239, Pl. LIII, fig. 2; off California coast, lat. $30^{\circ} 31' 35''$ N., long. $140^{\circ} 5' 30''$ W. in 300 fathoms.

Family 62. STERNOPTICHIDÆ

Genus 175. STERNOPTIX Herrmann.

Sternoptix Herrmann, Naturforscher, 2 Stück, XVI, 1781, 8 (*S. diaphana* Herrmann).*Sternoptyx*, authors' amended spelling.491. *Sternoptix obscura* (Garman).

Eastern Pacific in deep water.

Sternoptix obscura Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 232, Pl. LIII, fig. 1; numerous stations in eastern Pacific in depths ranging from 421 to 1,823 fathoms.

- 492. *Sternoptix diaphana* Herrmann.**
 Atlantic; Gulf Stream, from Santa Cruz Island to the Grand Banks.
Sternoptix diaphana Herrmann, Naturforscher, XVI, 1781, 8, Jamaica.
- Genus 178. ARGYROPELECUS Cocco.**
Argyropelecus Cocco, Arch. del R. Acad. Peloritano, 1829, 146 (*A. hemigymnus* Cocco).
Pleurothyris Lowe, Hist. Fish. Madeira, 1843, 64 (*Sternoptyx olfersii* Cuvier).
- 493. *Argyropelecus hemigymnus* Cocco.**
 Atlantic and Mediterranean; Gulf Stream off Southern New England.
Argyropelecus hemigymnus Cocco, Arch. del R. Acad. Peloritano, 1829, 146, coast of Italy.
Sternoptyx mediterraneus Cocco, Giorn. Faro de Messina, IV, 1838, 7, coast of Italy.
- 494. *Argyropelecus olfersii* (Cuvier).**
 Open Atlantic; coast of Norway to Brazil and Cape of Good Hope;
 Grand Banks southward in the Gulf Stream.
Sternoptyx olfersii Cuvier, Règne Anim., ed. 2, II, 1829, 316, near Cape of Good Hope.
Argyropelecus durvillii Cuvier and Valenciennes, Hist. Nat. Poiss., XXII, 1849, 405, open Atlantic.
- 495. *Argyropelecus lychnus* Garman.**
 Western Pacific in deep water.
Argyropelecus lychnus Garman, Deep Sea Fishes, in Mem. Mus. Comp. Zool., XXIV, 1899, 234, Pl. T, fig. 1, 16; many stations in eastern Pacific.
- 496. *Argyropelecus affinis* Garman.**
 Tropical Atlantic in deep water.
Argyropelecus affinis Garman, Deep Sea Fishes, in Mem. Mus. Comp. Zool., XXIV, 1899, 237; sta. 2117, N. lat. $15^{\circ} 24' 40''$, W. long. $63^{\circ} 31' 30''$, in 683 fathoms.
- Genus 177. POLYIPNUS Günther.**
Polyipnus Günther, Challenger Rept., 1887, 170 (*P. spinosus* Günther).
- 497. *Polyipnus laternatus* Garman.**
 Off Barbados.
Polyipnus laternatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 238; Blake station 280, off Barbados in 221 fathoms.
- 498. *Polyipnus spinosus* Günther.**
 Gulf of Mexico; between the Philippines and Borneo.
Polyipnus spinosus Günther, Challenger Rept., XXII, 1887, 170, from the surface near the Philippine Islands.
- Order O. LYOPOMI**
- Family 63. HALOSAURIDÆ**
- Genus 178. *HALOSAURUS* Johnson.**
Halosaurus Johnson, Ann. Mag. Nat. Hist., XIV, 1863, 74 (*H. oweni* (*owenii*) Johnson).
Halosaurichthys Alcock, loc. cit., 1889, 454, (*H. carinicauda* Alcock).
- 499. *Halosaurus güntheri* Goode and Bean.**
 Gulf Stream.
Halosaurus güntheri Goode and Bean, Oceanic Ichth., 1895, 131, Gulf Stream, in lat. $39^{\circ} 13' N.$, long. $72^{\circ} 1' W.$, Albatross station 2722.
- 500. *Halosaurus radiatus* Garman.**
 Off Panama in deep water.
Halosaurus radiatus Garman, Deep Sea Fishes, in Mem. Mus. Comp. Zool., XXIV, 1899, 298, Pl. LX, figs. 2, 2a, Pl. LXXXIV, figs. 3-6; several stations off Panama in depths from 259 to 511 fathoms.

501. *Halosaurus attenuatus* Garman.

Off Panama in deep water.

Halosaurus attenuatus Garman, Deep Sea Fishes, in Mem. Mus. Comp. Zool., XXIV, 1899, 296, Pl. LX, figs. 1, 1a, *Albatross* station 3413, lat. 2° 34' N., long. 92° 06' W., in 1,360 fathoms.

Genus 179. *ALDROVANDIA* Goode and Bean.

Aldrovandia Goode and Bean, Oceanic Ichth., 1895, 132 (*Halosaurus rostrata* Günther).

Halosauropsis Collett, Poiss. Prov. Comp. l'Hirondelle, Fasc. *X., Monaco, 1896, 143 (*Halosaurus macrochir*).

502. *Aldrovandia rostrata* (Günther).

Mid-Atlantic.

Halosaurus rostratus Günther, Ann. Mag. Nat. Hist., II, 1878, 252, Mid-Atlantic.

503. *Aldrovandia macrochir* (Günther).

Marion Island; *Blake* stations 308 and 325.

Halosaurus macrochir Günther, Ann. Mag. Nat. Hist., II, 1878, 251, between Cape of Good Hope and Kerguelen Island.

504. *Aldrovandia goodei* (Gill).

Gulf Stream, at *Albatross* stations 2037 in 1,731 fathoms; 2051 in 1,106 fathoms; 2035 in 1,362 fathoms, and 2052 in 1,098 fathoms.

Halosaurus goodei Gill, Proc. U. S. Nat. Mus., VI, 1883, 257, Gulf Stream off South Carolina.

505. *Aldrovandia gracilis* Goode and Bean.

Gulf Stream and Gulf of Mexico.

Aldrovandia gracilis Goode and Bean, Oceanic Ichth., 1895, 134, fig. 157, off Guadeloupe and in Gulf Stream at *Albatross* stations 2380 and 2381.

506. *Aldrovandia pallida* Goode and Bean.

Gulf of Mexico, in deep water.

Aldrovandia pallida Goode and Bean, Oceanic Ichth., 1895, 135, fig. 158, Gulf of Mexico, at *Albatross* station 173.

Order P. HETEROMI. Spiny eels

Family 64. LIPOGENYIDÆ

Genus 180. *LIPOGENYS* Goode and Bean.

Lipogenys Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894 (1895), 469 (*L. gillii* Goode and Bean).

507. *Lipogenys gillii* Goode and Bean.

Gulf Stream.

Lipogenys gillii Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894 (1895), 469, pl. 18, fig. 3, and in Oceanic Ichth., 173, 1895, station 2742, Gulf Stream.

Family 65. NOTACANTHIDÆ

Genus 181. *NOTACANTHUS* Bloch.

Notacanthus Bloch, Abhandl., Böhmischen Gesellsch. Wiss., I, 1787, 278 (*N. chemnitzi* Bloch).

Acanthonotus Bloch, Ichthyologia, XII, 1797, 113 (*A. nasus* Bloch = *N. chemnitzi* Bloch).

Campylacion (Fabricius, 1798) Reinhardt, Vidensk. Selsk. Naturv. Math. Afhandl., 1838, 120 (*C. fabricii* Reinhardt).

508. *Notacanthus chemnitzi* Bloch.

Greenland; West Indies; Iceland; Scandinavia; South Greenland.

Notacanthus chemnitzi Bloch, Abhandl., Böhmischen Gesellsch. Wiss., I, 1787, 278, northern seas.

Notacanthus nasus Bloch, Ichthyologia, XII, 1795, 113, pl. 431, "India."

Campylodon fabricii Reinhardt, Vidensk. Selsk. Naturv. Math. Afhandl.,

1838, 120, Greenland.

509. *Notacanthus analis* Gill.

Gulf Stream.

Notacanthus analis Gill, Proc. U. S. Nat. Mus., VI, 1883, 255, lat. 40° N., long. 69° W., Albatross stations 2677 in 478 fathoms and 2676 in 407 fathoms.510. *Notacanthus phasganorus* Goode.

Grand Banks, Newfoundland.

Notacanthus phasganorus Goode, Proc. U. S. Nat. Mus., III, 1880, 535, Grand Banks of Newfoundland.511. *Notacanthus spinosus* Garman.

Off Panama.

Notacanthus spinosus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 301, Pl. L, figs. 4, 4a, 4b; Albatross stations 3384 and 3354, 322 and 458 fathoms.Genus 182. **POLYACANTHONOTUS** Bleeker.*Polyacanthonotus* Bleeker, Versl. Akad. Amsterdam (2), VIII, 368 (*Notacanthus rissoanus* Filippi and Verany).*Zanotacanthus* Gill, Johnson's Cyclopædia, III, 1876, 883 (*Notacanthus rissoanus* Filippi and Verany).*Macdonaldia* Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894 (1895), 467 (*Notacanthus rostratus* Collett).512. *Polyacanthonotus challengerii* (Vaillant).

North Pacific and Bering Sea.

Notacanthus rissoanus Günther, Fishes, Challenger, XXII, Part LVII, Pl. LXI, fig. 13 (not of Filippi and Verany).*Notacanthus challengerii* Vaillant, Expéd. Sci. Travailleur et Talisman, 1888, south of Yeddo; based on Günther's description.513. *Polyacanthonotus rostratus* (Collett).

Gulf Stream.

Notacanthus rostratus Collett, Bull. Soc. Zool. France, 1889, 307, off Newfoundland, at Albatross station 2216 in 963 fathoms, and station 2553 in 551 fathoms.

Order Q. SYMBRANCHIA. Symbranchoid eels

Suborder HOLOSTOMI

Family 66. SYNBRANCHIDÆ

Genus 183. **SYNBRANCHUS** Bloch.*Synbranchus* Bloch, Ichthyologia, IX, 1795, 87 (*Synbranchus marmoratus* Bloch).*Unibranchiatura* Lacépède, Hist. Nat. Poiss., V, 1803, 658 (*Synbranchus marmoratus*).*Ophisternon* McClelland, Calcutta Jour. Nat. Hist., V, 1845, 197 (*O. bengalensis*).*Tetrabranchus* Bleeker, Nat. Tijds. Ned. Ind., II, 1862, 69 (*T. micropthalmus* McClelland).*Symbranchus* of authors generally; corrected spelling.514. *Synbranchus marmoratus* Bloch.

Tropical America, in streams from the Amazon northward to southern Mexico and St. Lucia; generally abundant; recorded northward from Vera Cruz, Trinidad, St. Lucia, Lake Peton, Huamuchal, Rio Chisoy, etc.

Synbranchus marmoratus Bloch, Ichthyologia, IX, 1795, 87, pl. 418, tropical America.*Synbranchus immaculatus* Bloch, loc. cit., pl. 419, fig. 1.*Synbranchus transversalis* Bloch and Schneider, loc. cit., 1801, 524, Guinea; after Gronow.*Unibranchiatura grisea*, *marmorata*, *immaculata*, and *lineata* Lacépède, Hist. Nat. Poiss., V, 1803, 658, Surinam.*Synbranchus fuliginosus* Ranzani, Nov. Com. Acad. Sci. Bonon, IV, 1840, 75, pl. 11, fig. 1, Brazil.*Muraena lumbriacus* Gronow, Cat. Fish., 1854, 18, Sea of Guinea.*Syphobranchus vittatus* Castelnau, Anim. Amer. Sud, Poiss., 84, 1855, pl. 44, fig. 3, Rio de Janeiro.

Order R. CARENCHELI. Long-necked eels

Family 67. DERICHTHYIDÆ

Genus 184. DERICHTHYS Gill.

Derichthys Gill, Amer. Nat., XVIII, 1884, 433 (*D. serpentinus* Gill).

515. *Derichthys serpentinus* Gill.

Gulf Stream.

Derichthys serpentinus Gill, Am. Nat., XVIII, 1884, 433, Gulf Stream at Albatross station 2094 in 1,022 fathoms.

Genus 185. GORGASIA Meek and Hildebrand.

Gorgasia Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, 133 (*G. punctata* Meek and Hildebrand).

516. *Gorgasia punctata* Meek and Hildebrand.

Panama.

Gorgasia punctata Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, 133, Chame Point, Panama.

Order S. APODES

Suborder ENCHELYCEPHALI

Series ANGUILLOIDEI

Family 68. ANGUILLIDÆ. True eels

Genus 186. ANGUILLA Shaw. Eels.

Anquilla Shaw, Gen. Zool., IV, 1803, 15 (*A. anguilla*).

Muræna Bleeker, Poey, etc. (taking as type *Muræna anguilla*, the first species mentioned by Artedi under *Muræna*).

Tribanchus (Peters) Müller, Abhandl., Akad. Wiss. Berlin, 1844 (*T. anguillaris* Peters = *Anguilla australis* Rich.).

517. *Anguilla rostrata* (LeSueur). American eel; Fresh-water eel.

Atlantic coast of the United States; very abundant from Maine to Mexico, ascending all rivers south of Canada and east of the Rocky Mountains, individuals sometimes resident throughout the Mississippi Valley; common in the West Indies.

Muræna rostrata LeSueur, Jour. Acad. Nat. Sci. Phila., 1817, 81, Cayuga Lake.

Muræna bostonensis LeSueur, loc. cit., 81, Boston.

Muræna serpentina LeSueur, loc. cit., 82, Newport, R. I.

Muræna macrocephala LeSueur, loc. cit., 82, Saratoga, N. Y.

Muræna argentea LeSueur, loc. cit., 82, Boston Bay.

Anguilla chrysopa Rafinesque, Amer. Month. Mag. and Crit. Rev., 1817, 120, Lake George; Hudson River; Lake Champlain.

Anguilla blephura Rafinesque, loc. cit., 120, Long Island.

Anguilla laticeps Rafinesque, loc. cit., 445, Ohio River.

Anguilla aterrima Rafinesque, Ichth. Ohiensis, 1820, 78, Tennessee and Cumberland Rivers.

Anguilla xanthomelas Rafinesque, loc. cit., 78, Ohio River.

Anguilla lutea Rafinesque, loc. cit., 78, Ohio River.

Anguilla tenuirostris DeKay, N. Y. Fauna: Fishes, 1842, 310, New York.

Anguilla novæorleansis Kaup, Apodal Fishes, 1856, 43, fig. 33, New Orleans, La.

Anguilla punctatissima Kaup, loc. cit., 44, Niagara River.

Anguilla cubana Kaup, loc. cit., 44, Cuba.

Anguilla novæterra Kaup, loc. cit., 45, fig. 35, Newfoundland.

Anguilla texana Kaup, loc. cit., 45, fig. 36, Texas.

Anguilla wabashensis Kaup, loc. cit., 46, Wabash River.

Anguilla tyrannus Girard, U. S. Mex. Bound. Surv., Ichth., 1859, 75, Rio Grande.

Family 69. SIMENCHELYIDÆ. Snub-nosed eels

Genus 187. SIMENCHELYS Gill.

Simenchelys Gill, in Goode and Bean, Bull. Essex Inst., 1879, 27 (*S. parasiticus* Gill).

Conchognathus Collett, Bull. Soc. Zool. France, 1889, 123 (*C. grimaldii* Collett).

518. *Simenchelys parasiticus* Gill.

Offshore banks, in deep water, south of Newfoundland; also recorded from the Azores.

Simenchelys parasiticus Gill, in Goode and Bean, Bull. Essex Inst., 1879, 27, Newfoundland banks.

Conchognathus grimaldii Collett, Bull. Soc. Zool. France, 1889, 122, Azores.

Family 70. SYNAPHOBRANCHIDÆ

Genus 188. SYNAPHOBRANCHUS Johnson.

Synaphobranchus Johnson, Proc. Zool. Soc. Lond., 1862, 169 (*S. kaupii* Johnson = *Muræna pinnata* Gronow).

519. *Synaphobranchus pinnatus* (Gronow).

North Atlantic and western Pacific; common about the Madeiras, Canaries, etc., and also about the banks of Newfoundland.

Muræna pinnata Gronow, Cat. Fish., 1854, 19, locality unknown.

Synaphobranchus kaupii Johnson, Proc. Zool. Soc. Lond., 1862, 169, Madeira.

Genus 189. HISTIOBRANCHUS Gill.

Histiobranchus Gill, Proc. U. S. Nat. Mus., VI, 1883, 255 (*H. infernalis* Gill).

520. *Histiobranchus bathybius* (Günther).

Northern and western Pacific in deep water; Bering Strait; off Japan and off Cape of Good Hope; one specimen obtained by Doctor Gilbert in Bering Sea in 1890.

Synaphobranchus bathybius Günther, Ann. Mag. Nat. Hist., XX, 1877, 445, and in Voy. Challenger, 254, Pl. LXII, fig. 6, 1887, off Yeddo; North Pacific; Kerguelen Island.

521. *Histiobranchus infernalis* Gill.

Gulf Stream.

Histiobranchus infernalis Gill, Proc. U. S. Nat. Mus., VI, 1883, 255, Gulf Stream, at Albatross station 2037, 38° 30' N., 69° W., in 1,731 fathoms.

Family 71. ILYOPHIDÆ. Ooze eels

Genus 190. ILYOPHIS Gilbert.

Ilyophis Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 351 (*I. brunneus* Gilbert).

522. *Ilyophis brunneus* Gilbert.

One specimen from Chatham Island, Galapagos Archipelago, in 634 fathoms.

Ilyophis brunneus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 352, Chatham Island, Galapagos Archipelago, at Albatross station 2808 in 654 fathoms.

Family 72. MURÆNESOCIDÆ

Genus 191. MURÆNESOX McClelland.

Murænesox McClelland, Calcutta Jour. Nat. Hist., V, 1844, 408 (*M. tricuspidata* McClelland).

Branchyconger Bleeker, Nederl. Tijdschr. Dierk., II, 1865, 236 (*Muræna savanna* Cuvier).

523. *Murænesox savanna* (Cuvier).

Cuba to Rio Janeiro, not common; occasional in the Mediterranean Sea.

Muræna savanna Cuvier, Règne Anim., ed. 2, II, 1829, 350, Martinique.

Conger brasiliensis Ranzani, De Novis Spec. Pisc. Diss. Prima., IV, 1838, 17, pl. 13, fig. 1, Brazil.

Congrus curvidens Richardson, Voy. Erebus and Terror, 1844, 111, no locality.

Conger limbatus Castelnau, Anim. Amér. Sud, 1855, 83, pl. 43, fig. 3, Rio Janeiro.

524. *Muraenesox coniceps* Jordan and Gilbert.

Cape San Lucas to the coast of Colombia; generally common.

Muraenesox coniceps Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 348, Mazatlan, Mexico.

Family 73. CONGRIDÆ (LEPTOCEPHALIDÆ). Conger eels

Genus 192. *Conger* Cuvier. Conger eels.a. Adult forms (*Conger*):

Conger Cuvier, Oken, in Règne Anim., Ed. I, 1817, 231 ("Les Congres" Cuvier, Oken, 1882) (*Muraena conger* Linnæus).

Congrus Richardson, Voy. Erebus and Terror, 1844, 107 (*Muraena conger* Linnæus).

b. Larval forms (*Leptocephalus*):

Leptocephalus Scopoli, Int. Hist. Nat., 1777, 453 (*L. morrissi* Gmelin).

Oxyurus Rafinesque, Caratteri, 1810, 19 (*O. vermiciformis* Lacépède).

Helmictis Rafinesque, Indice, 1810, 62.

Helminchthys Costa, Fauna Napoli, Pesci, tavoli 31, 1854 (*H. diaphanus* Costa).

? *Leptocephalichthys* Bleeker, Act. Soc. Sci. Indo-Neder., I, 69 (*L. hypselosoma* Bleeker, a larval form allied to *Conger*).

? *Diaphanichthys* Peters, Sitzungsber., Akad. Wiss. Berlin, 1864, 399 (*D. brevicaudus* Peters); larva allied to *Conger*.

525. *Conger conger* (Linnæus). Conger eel.

Atlantic Ocean; generally common on both coasts, from Cape Cod to Brazil, also on coasts of Asia and Africa.

a. Adult forms:

Muraena supremo marginæ pinnæ dorsalis nigro Artedi, Synon., 40, 1738, 2, Mediterranean.

Muraena conger Linnæus, Syst. Nat., ed. X, 1758, 245; based on Artedi.

Muraena nigra Risso, Ichth. Nice, 1810, 93 (black variety), Nice.

Anquilla oceanica Mitchell, Jour. Acad. Nat. Sci. Phila., 1818, 407, off New York.

Conger verus Risso, Europ. Mérid., III, 1826, 201, Nice.

Conger vulgaris Cuvier, Règne Anim., ed. 2, II, 1829, 350, France.

Conger rubescens Ranzani, De Novis Spec. Pisc. Diss. Prima, 1838, 19, Pl. V, fig. 5, Mediterranean.

Ophisoma obtusa Swainson, Nat. Hist. Fish., Reptiles, and Amphib., II, 1839, 395, Sicily.

Conger orbignyanus Valenciennes, D'Orbigny, Voy. Am. Merid., Poiss., 1839, pl. 12, 1, South America.

Conger verreauxi Kaup, Apodal Fishes, 1856, 115, no locality.

Conger esculentus Poey, Memorias, II, 1861, 346, Cuba.

b. Larval forms:

Leptocephalus morrisi Gmelin, Syst. Nat., 1150, 1877, Holyhead, England.

Ophidium pellucidum Couch, Lond. Mag. Nat. Hist., V, 1832, 3113, 742, England.

Leptocephalus gracilis Storer, Mem. Amer. Acad., II, 1839, 524, Massachusetts.

Leptocephalus spallanzani, *candidissimus*, etc., of European writers.

526. *Conger caudilimbatus* (Poey).

Tropical parts of Atlantic; Pensacola to Cuba and Madeira.

Echelus caudilimbatus Poey, Repertorio, II, 1867, 249, Cuba.

Conger macrops Günther, Cat., VIII, 1870, 40, Madeira, Bahamas.

Conger caudicula Bean, Proc. U. S. Nat. Mus., V, 1882, 436, Pensacola.

Genus 193. *ARIOSOMA* Swainson.

Ariosoma Swainson, Nat. Hist. Fish., I, 1838, 220; II, 1839, 195 (no type named, logotype as restricted by Bleeker and by Swainson *Ophisoma acuta* Swainson = *Muraena balearica* De la Roche).

Ophisoma Swainson, loc. cit., 1839; *Ophisoma acuta* Swainson; substitute for *Ariosoma*; the name preoccupied.

Congermuræna Kaup, Apodal Fishes, 1856, 71 (108) (*Muraena balearica* De la Roche, as restricted by Bleeker).

Congrellus Ogilby, Proc. Linn. Soc. New South Wales, XXIII, 1898, 288 (*Muraena balearica* De la Roche).

527. Ariosoma opisthoptalma (Ranzani).

West Indies to Brazil.

Conger opisthoptalmus Ranzani, De Novis Spec. Pisc. Diss. Prima, 1838, 16, Pl. V, fig. 1, Bahia.*Conger microstomus* Castelnau, Anim. Amér. Sud, 1855, 83, pl. 42, fig. 4, Rio Janeiro.*Conger analis* Poey, Memorias, II, 1861, 318, Habana.*Conger impressus* Poey, loc. cit., 318, Cuba.**528. Ariosoma giberti (Ogilby).**

Panama; probably also at the Galapagos and Cape San Lucas.

Ophisoma balearicum Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 349,

Albatross station 2797, Panama Bay, 33 fathoms; not of De la Roche.

Congrellus giberti Ogilby, Proc. Linn. Soc. New South Wales, 1898, 288, after Gilbert.**529. Ariosoma caudale (Garman).**

Off Panama in deep water.

Congermuræna caudalis Garman, Mus. Comp. Zool., XXIV, 1899, 305; Albatross station 3354, lat. $7^{\circ} 9' 45''$ N., long. $80^{\circ} 50'$ W., in 322 fathoms.Genus 194. ANAGO²⁴ Jordan and Hubbs.*Anago* Jordan and Hubbs, Mem. Carnegie Mus., X, 1925, 191 (*Conger anago* Schlegel).**530. Anago nitens (Jordan and Bollman).**One specimen dredged at Albatross station 2801, off Panama, $8^{\circ} 47'$ N., $79^{\circ} 29' 30''$ W., in 14 fathoms.*Ophisoma nitens* Jordan and Boiluan, Proc. U. S. Nat. Mus., XII, 1889, 153, Panama.**531. Anago macrurus (Gilbert).**

Gulf of California.

Ophisoma macrurum Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 351, Gulf of California, at Albatross station 3105.

Genus 195. HILDEBRANDIA Jordan and Evermann.

Hildebrandia Jordan and Evermann, Proc. Calif. Acad. Sci., 4th Ser., XVI, Apr. 27, 1927, 502 (*Congermuræna flava* Goode and Bean).**532. Hildebrandia flava (Goode and Bean).**

Albatross stations 2121 and 2122 in 31 and 34 fathoms; station 2402 in 111 fathoms; also Blake station 264 in 84 fathoms; Gulf Stream.

Congermuræna flava Goode and Bean, Oceanic Ichth., 1895, 138, fig. 159, Gulf Stream.

Genus 196. GNATHOPHIS Kaup.

Gnathophis Kaup, Abhandl., Natur. Verein Hamburg, 1859 (1860), IV, Abth. 2, 7 (*G. heterognathus* Kaup); teeth blunt.**533. Gnathophis guppai (Norman).**

West Indies.

Congromuræna guppai Norman, Ann. Mag. Nat. Hist., ser. 9, XV, 1925, 313, Tobago.**534. Gnathophis proriger (Gilbert).**

Panama to Ecuador.

Ophisoma prorigerum Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 350, coast of Ecuador, at Albatross station 2792 in 401 fathoms; also at station 2799, near Panama.

Genus 197. CONGROSOMA Garman.

Congrosoma Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 308 (*Congrosoma evermanni* Garman).**535. Congrosoma evermanni Garman.**

Off Panama.

Congrosoma evermanni Garman, Mem. Mus. Comp. Zool., XXIV, 1899, Pl. LXII, fig. 1, Albatross station 3355, lat. $7^{\circ} 12' 20''$ N., long. $80^{\circ} 55'$ W. in 182 fathoms.

²⁴ The shore species named below may not, any of them, belong to *Anago* or, in fact, to the same genus. They seem nearer the Japanese form *Anago* than to *Ariosoma*.

Genus 198. UROCONGER Kaup.

Uroconger Kaup, Apodal Fishes, 1856, 110 (*U. lepturus* Kaup).536. *Uroconger vicinus* Vaillant.Deep waters off the coast of northern Africa; a young individual taken by the *Albatross* at station 2161 in 146 fathoms.*Uroconger vicinus* Vaillant, Expéd. Sci. Travailleur et Talisman, 1883, 86, Pl. VI, fig. 1, Band d'Arguin, off Soudan, off Cape Verde Islands.537. *Uroconger varidens* Garman.

Off Panama in deep water.

Uroconger varidens Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 304, Pl. LXI, fig. 1, *Albatross* stations 3396 and 3357 in 259 and 782 fathoms.

Genus 199. MICROCONGER Fowler.

Microconger Fowler, Proc. Acad. Nat. Sci. Phila., LXIV, 1912, 9 (*Leptocephalus caudalis* Fowler).538. *Microconger caudalis* (Fowler).

Coast of Lower California.

Leptocephalus (*Microconger*) *caudalis* Fowler, Proc. Acad. Nat. Sci. Phila., LXIV, 1912, 9; off Lower California.

Genus 200. MAYERINA Silvester.

Mayerina Silvester, Carnegie Inst. Year Book, XIV, 1915, 214 (*Mayerina mayeri* Silvester).539. *Mayerina mayeri* Silvester.

Porto Rico.

Mayerina mayeri Silvester, Carnegie Inst. Year Book, XIV, 1915, 214, Porto Rico.

Genus 201. XENOMYSTAX Gilbert.

Xenomystax Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 348 (*X. atrarius* Gilbert).540. *Xenomystax atrarius* Gilbert.

Coast of Ecuador.

Xenomystax atrarius Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 348, 1° S., 81° W., at *Albatross* station 2792 in 401 fathoms.541. *Xenomystax rictus* Garman.

Off Panama.

Xenomystax rictus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 315, Pl. N, several *Albatross* stations (3384, 3394, 3354, 3404, 3417) in from 322 to 511 fathoms.

Genus 202. HOPLUNNIS Kaup.

Hoplunnis Kaup, Aalen. Fische Hamburg Mus., 1860, 19 (*H. schmidii* Kaup).542. *Hoplunnis schmidti* Kaup.

Caribbean Sea at Puerto Cabello, near Aspinwall; Colon.

Hoplunnis schmidti Kaup, Aalen. Fische Hamburg Mus., 1859, 19, pl. 2, fig. 4, Puerto Cabello, near Aspinwall, Colon.543. *Hoplunnis diomedianus* Goode and Bean.A single individual obtained by the *Albatross* in the Gulf of Mexico.
Hoplunnis diomedianus Goode and Bean, Oceanic Ichth., 146, 1895, fig. 163, Gulf of Mexico, *Albatross* station 2402, 28° 36' N., 86° 50' W., 111 fathoms.

Genus 203. NEOCONGER Girard.

Neoconger Girard, Proc. Acad. Nat. Sci. Phila., X, 1858 (1859), 171 (*N. mucronatus* Girard).544. *Neoconger mucronatus* Girard.

Coast of Texas.

Neoconger mucronatus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858 (1859), 171, St. Joseph Island, Texas.545. *Neoconger vermicularis* Gilbert.

Lower California to Panama.

Neoconger vermicularis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 57, off Lower California, at *Albatross* station 3035 in about 30 fathoms.

Genus 204. LEPTOCONGER Poey.

Leptoconger Poey, Anal. Soc. Espaniola Hist. Nat., Madrid, 1880, 250
(*L. perlongus* Poey).

546. *Leptoconger perlongus* (Poey).

Matanzas, Cuba.

Neoconger perlongus Poey, Ann. Lyc. Nat. Hist. N. Y., XI, 1874, 67,
pl. 9, figs. 3 and 4, Matanzas, Cuba.

Family 74. NETTASTOMIDÆ. Duckbill eels; Sorcerers**Genus 205. CHLOPSIS Rafinesque.**

Chlopsis Rafinesque, Indice, 1810, 58 (*C. bicolor* Rafinesque).

Sauvanchelys Peters, Sitzungsber. Akad. Wiss. Berlin, 1864, 397 (*S. cancrivora* Peters).

547. *Chlopsis equatorialis* Gilbert.

Coast of Ecuador.

Chlopsis equatorialis Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 347,
off coast of Ecuador, 1° S., 81° W., at *Albatross* station 2792 in 401
fathoms.

548. *Chlopsis giberti* Garman.

Off Panama and San Diego.

Chlopsis giberti Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
Pl. LXII, fig. 2, various stations of the *Albatross*, from 322 to 511
fathoms.

Genus 206. VENEFICA Jordan and Davis.

Venefica Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 651 (*Nettastoma procerum* Goode and Bean).

549. *Venefica procera* (Goode and Bean).

Gulf Stream; also taken off San Pedro, Calif.

Nettastoma procerum Goode and Bean, Bull. Mus. Comp. Zool., X,
1883, 224, Gulf Stream, at *Albatross* station 325.

550. *Venefica ocella* Garman.

Off Panama in deep water.

Venefica ocella Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 318,
Pl. LXI, fig. 2, *Albatross* station 3366, lat. 5° 30' N., long. 86° 45' W.,
in 1,067 fathoms.

551. *Venefica tentaculata* Garman.

Off Panama in deep water; San Diego.

Venefica tentaculata Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
319, Pl. M, fig. 2, various *Albatross* stations at depths ranging from
770 to 978 fathoms.

Genus 207. TÆNIOCONGER Herre.

Tænioconger Herre, Philippine Jour. Sci., XXIII, No. 2, August, 1923,
152 (*T. chapmani* Herre).

552. *Tænioconger digueti* Pellegrin.

Gulf of California.

Tænioconger digueti Pellegrin, Compt. Rend. Acad. Sci. Paris, CLXXVII,
1923; 789, near Island of Espiritu Santo, Gulf of California.

Family 75. ECHELIDÆ. Worm eels**Genus 208. AHLIA Jordan and Davis.**

Ahlia Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 639 (*Myrophis egmontis* Jordan).

553. *Ahlia egmontis* (Jordan).

Florida Keys.

Myrophis egmontis Jordan, Proc. Acad. Nat. Sci. Phila., XXXVI, 1884,
44, Egmont Key, Fla.

Genus 209. MYROPHIS Lütken.

Myrophis Lütken, Vid. Med. Naturh. Foren. Kjøbenhavn, 1, 1851
(*M. punctatus* Lütken).

554. *Myrophis punctatus* Lütken.

West Indies; coast of Texas to Surinam; common along our Gulf coast.
Myrophis punctatus Lütken, Vid. Med. Naturh. Foren. Kjøbenhavn, 1, 1851, 9, West Indies.

Myrophis microstigmus Poey, Repertorio, II, 1867, 250, Cuba.

Myrophis lumbucus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 261, Galveston, Tex.

555. *Myrophis vafer* Jordan and Gilbert.

Pacific coast of tropical America, from Guaymas to Panama.

Myrophis vafer Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 645, Panama.

Genus 210. CHILORHINUS Lütken.

Chilorhinus Lütken, Vid. Med. Naturh. Foren. Kjøbenhavn, 1, 1851 (*C. suensonii* Lütken).

556. *Chilorhinus suensonii* Lütken.

St. Croix, West Indies.

Chilorhinus suensonii Lütken, Vid. Med. Naturh. Foren. Kjøbenhavn, 1, 1851, St. Croix.

Family 76. NEMICHTHYIDÆ. Thread eels; Snipe eels**Genus 211. NEMICHTHYS Richardson.**

Nemichthys Richardson, Voy. Samarang, 1848, 25 (*N. scolopaceus* Richardson).

Leptorhynchus Lowe, Ann. Mag. Nat. Hist., X, 1852, 54 (*L. leuchtenbergi* Lowe); preoccupied.

Belonopsis Brandt, Mem. Acad. Sci. St. Petersb., Savans Etrangères, 1854, 174 (*Leptorhynchus leuchtenbergi* Lowe).

557. *Nemichthys scolopaceus* Richardson. Snipe eel.

Atlantic Ocean, in deep water; many specimens taken off the New England coast and off the Grand Banks; common about Madeira.

Nemichthys scolopacea Richardson, Voy. Samarang, 1848, 25, South Atlantic.

Leptorhynchus leuchtenbergii Lowe, Mem. Acad. Sci. St. Petersb., Savans Etrangères 1854, 171, Madeira.

Belonopsis leuchtenbergii Brandt, loc. cit., 1854, 174 (with good plate).

558. *Nemichthys avocetta* Jordan and Gilbert.

Puget Sound near Seattle; only the type known.

Nemichthys avocetta Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 409, harbor of Port Gamble, near Seattle.

559. *Nemichthys fronto* Garman

Off Pacific coast of Mexico, in deep water.

Nemichthys fronto Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 324, Pl. LXV, fig. 1, Albatross stations 3384 and 3434, in 458 and 1,588 fathoms.

Genus 212. AVOCETTINA Jordan and Davis.

Avocettina Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 655 (*Nemichthys infans* Günther).

560. *Avocettina infans* (Günther).

Known from West Indies, and middle Atlantic off Pernambuco.

Nemichthys infans Günther, Ann. Mag. Nat. Hist., 5th Ser., II, 1878, 251, and in Challenger Rept., XXII, 1887, 264, mid. Atlantic in 2,500 fathoms.

561. *Avocettina gilli* (Bean).

Coast of southeastern Alaska in deep water.

Labichthys gilli Bean, Proc. U. S. Nat. Mus., XIII, 1890, 45, east of Prince of Wales Island, Alaska.

562. *Avocettina elongata* (Gill and Ryder).

Gulf Stream.

Labichthys elongatus Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883, 262, Gulf Stream at Albatross station 2100, 39° 22' N., 68° 34' W.

Genus 213. **SERRIVOMER** Gill and Ryder.

Serrivomer Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883, 260 (*S. beanii* Gill and Ryder).

563. **Serrivomer beanii** Gill and Ryder.

Gulf Stream; also taken in the Gulf of California.

Serrivomer beanii Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883, 261, Gulf Stream, lat. $41^{\circ} 40' 30''$, long. $65^{\circ} 28' 30''$ in 855 fathoms.

564. **Serrivomer sector** Garman.

Gulf of California to Panama in deep water.

Serrivomer sector Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 320, Pl. LXIII, numerous stations of the Albatross and off Guaymas, 50 miles south, surface to 700 fathoms.

Genus 214. **SPINIVOMER** Gill and Ryder.

Spinivomer Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883, 261 (*S. goodei* Gill and Ryder).

565. **Spinivomer goodei** Gill and Ryder.

Gulf Stream.

Spinivomer goodei Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883, 261, Gulf Stream, at lat. $38^{\circ} 19' 26''$, long. $68^{\circ} 20' 20''$ in 2,361 fathoms.

Genus 215. **LABICHTHYS** Gill and Ryder.

Labichthys Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883, 261 (*L. carinatus* Gill and Ryder).

566. **Labichthys carinatus** Gill and Ryder.

Gulf Stream.

Labichthys carinatus Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883, 253, 255, 261, Gulf Stream, $41^{\circ} 13' N.$, $65^{\circ} 33' W.$

567. **Labichthys bowersi** Garman.

Off Panama in deep water.

Labichthys bowersi Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 823, Pl. XIV, fig. 1, Albatross stations 3414, 3388, and 3361, in 1,168 to 2,232 fathoms.

Family 77. **OPHICHTHYIDÆ. Snake eels**Genus 216. **SPHAGEBRANCHUS** Bloch.

Sphagebranchus Bloch, Ichthyologia, IX, 1795, 88, pl. 419 (*S. rostratus* Bloch).

Cæcilia Lacépède, Hist. Nat. Poiss., II, 1800, 135 (*C. brandieriana* Lacépède = *Muraena cæca* Linnæus); not *Cæcilia* L. a genus of *Batrachia*.

Apterichthys Duméril, Zool. Anal., 112, 1806, 331 (*Muraena cæca* Linnæus).

Branderius Rafinesque, Anal. Nat., 1815, 93; substitute for *Cæcilia* Lacépède.

Ichthyapus Brisout de Barneville, Revue Zool., 1847, 219 (*I. acutirostris*).

Ophisuraphis Kaup, Apodal Fishes, 1856, 29 (*O. gracilis* Kaup).

568. **Sphagebranchus anguiformis** (Peters).

Open Atlantic, near the West Indies.

Ophichthys (*Sphagebranchus*) *anguiformis* Peters, Monatsber. Akad. Wiss. Berlin, 1876, 849, Atlantic Ocean, $15^{\circ} 40' N.$, $23^{\circ} 5' W.$

569. **Sphagebranchus selachops** (Jordan and Gilbert).

Rocks about Cape San Lucas.

Apterichthys selachops Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 356, Cape San Lucas.

570. **Sphagebranchus ophioneus** Evermann and Marsh.

Porto Rico.

Sphagebranchus ophioneus Evermann and Marsh, Fish. Porto Rico, Bull. U. S. Fish Com., XX, 1900, 73, fig. 7, off Mayaguez, Porto Rico.

Genus 217. **VERMA** Jordan and Evermann.

Verma Jordan and Evermann, Fishes North and Mid. Amer., 1896, 374 (*Sphagebranchus kendalli* Gilbert).

571. *Verma kendalli* (Gilbert).
 Coast of Florida, in rather deep water.
Sphagebranchus kendalli Gilbert, Bull. U. S. Fish Com., IX, 1889 (1891), 310, off west coast of Florida, 25° 34' N., 82° 50' W., in 25 fathoms.
- Genus 218. CRYPTOPTERENCHELYS Fowler.
Cryptopterus Kaup, Aalen. Fische Hamburg Mus., II, 1860 (*C. puncticeps* Kaup); name preoccupied.
Cryptopterenchelys Fowler, Amer. Mus. Novitates, CLXII, March 31, 1925 (*Cryptopterus puncticeps* Kaup); substitute for *Cryptopterus* Kaup, preoccupied.
572. *Cryptopterenchelys puncticeps* (Kaup).
 Caribbean Sea.
Cryptopterus puncticeps Kaup, Aalen. Fische Hamburg Mus., II, 1860, pl. 1, fig. 2, Puerto Cabello.
573. *Cryptopterenchelys frontalis* (Garman).
 Off Panama.
Ophichthys (Cryptopterus) frontalis Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 309, Albatross stations 3386, 3389, 3391.
- Genus 219. MYRICHTHYS Girard.
Pisodonophis Kaup, Apodal Fishes, 1856, 15 (in part; not type, as restricted by Bleeker, which is *P. cancrivorus*).
Myrichthys Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 58 (*M. tigrinus* Girard).
Ophisurus Swainson, Bleeker, etc. (not of Lacépède, nor of Risso, nor Kaup, who restrict the name to *O. serpens*).
574. *Myrichthys tigrinus* Girard.
 Pacific coast of Mexico; rather common about Mazatlan, and occasionally ranging northward.
Myrichthys tigrinus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 58, Adair Bay, Oregon; doubtless an error.
Ophisurus xysturus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 346, Mazatlan.
575. *Myrichthys xysturus* (Jordan and Gilbert).
 Pacific coast of Mexico; common among the rocks about Mazatlan.
Ophichthys xysturus Jordan and Gilbert, Proc. U. S. Nat. Mus., IX, 1881, 346, Mazatlan, Mexico.
576. *Myrichthys oculatus* (Kaup).
 Tropical Atlantic; Cuba to Surinam and Cape Verde Islands.
Pisodonophis oculatus Kaup, Apodal Fishes, 1856, 22, Curaçao.
Ophisurus latimaculatus Poey, Repertorio, II, 1867, 252, pl. 3, fig. 1, Cuba.
577. *Myrichthys acuminatus* (Gronow).
 West Indies, occasionally northward to Florida Keys.
Muraena acuminata Gronow, Fish. Brit. Mus., 1854, 21, Insula Div., Eustachii.
Pisodonophis guttulatus Kaup, Apodal Fishes, 1856, 21, fig. 10, Martinique.
Ophisurus longus Poey, Repertorio, II, 1867, 254, Cuba.
Ophichthys pisavarius Poey, Anal. Soc. Española Hist. Nat., 1875, 196, Cuba.
578. *Myrichthys pantostigmus* Jordan and McGregor.
 Revillagigedo Islands.
Myrichthys pantostigmus Jordan and McGregor, Rept. U. S. Fish Com., XXIV, 1898 (1899), 274, pl. 4, Clarion Island.
- Genus 220. LETHARCHUS Goode and Bean.
Letharchus Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 436 (*L. velifer* Goode and Bean).
579. *Letharchus pacificus* Osburn and Nichols.
 Gulf of California.
Letharchus pacificus Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 146, fig. 3, Cape San Lucas.

580. *Letharchus velifer* Goode and Bean.

Coast of Florida, in rather deep water; known only from the Snapper Banks off Pensacola and Tampa.

Letharchus velifer Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 437, West Florida.

Genus 221. *PISODONOPHIS* Kaup.

Pisodonophis Kaup, Apodal Fishes, 1856, 17 (*P. cancrivorus* Kaup as restricted by Bleeker; later written *Pisoodonophis* by Kaup and since spelled *Pisodontophis* by purists).

581. *Pisodonophis daspilotus* Gilbert.

Panama.

Pisodonophis daspilotus Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2803, Panama.

Genus 222. *OMOCHELYS* Fowler.

Omochelys Fowler, Proc. Acad. Nat. Sci. Phila., LXX, 1918, 3 (*Pisodonophis cruentifer* Goode and Bean).

Styletor Jordan, loc. cit., LXX, 1918 (1919), 343 (*Pisoodonophis cruentifer* Goode and Bean); name preoccupied.

582. *Omochelys cruentifer* (Goode and Bean).

Two specimens at station 1035 of U. S. Fish Commission steamer *Fish Hawk*, 39° 57' N., 69° 28' W., in 120 fathoms; four others at nearly same region in 245 fathoms.

Pisodonophis cruentifer Goode and Bean, Oceanic Ichth., 1895, 147, fig. 166, Gulf Stream.

Genus 223. *CALLECHELYS* Kaup.

Callechelys Kaup, Apodal Fishes, 1856, 28 (*C. guichenoti* Kaup).

583. *Callechelys muræna* Jordan and Evermann.

Snapper Banks off Pensacola.

Callechelys muræna Jordan and Evermann, Proc. U. S. Nat. Mus., IX, 1886, 466, Snapper Banks.

Genus 224. *BASCANICHTHYS* Jordan and Davis.

Bascanichthys Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 621 (*Cæcula bascanium* Jordan).

584. *Bascanichthys bascanium* (Jordan).

Egmont Key, Florida.

Cæcula bascanium Jordan, Proc. Acad. Nat. Sci. Phila., XXXVI, 1884, 48, Egmont Key, Fla.

585. *Bascanichthys scuticaris* (Goode and Bean).

West coast of Florida.

Sphagebranchus scuticaris Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 343, Cedar Key.

Sphagebranchus teres Goode and Bean, loc. cit., V, 1882, 436, west Florida.

586. *Bascanichthys cylindricus* Meek and Hildebrand.

Panama.

Bascanichthys cylindricus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, 152, Pl. VIII, Chame Point, Panama.

587. *Bascanichthys panamensis* Meek and Hildebrand.

Panama.

Bascanichthys panamensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, Pl. VII, 151, Chame Point, Panama.

588. *Bascanichthys bascanoides* Osburn and Nichols.

Coast of Lower California.

Bascanichthys bascanoides Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 147, fig. 4, San Cristobal, from the stomach of *Myceteropera venadorum*.

589. *Bascanichthys peninsulæ* (Gilbert).

La Paz Bay, Gulf of California.

Callechelys peninsulæ Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 548, La Paz Bay, Lower California.

Genus 225. QUASSIREMUS Jordan and Davis.

Quassiremus Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 622
(*Ophichthus evionthas* Jordan and Bollman).

590. *Quassiremus nothochir* (Gilbert).

San Josef Island, Gulf of California.

Ophichthys nothochir Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 58,
San Josef Island, Gulf of California.

591. *Quassiremus evionthas* (Jordan and Bollman).

Hood Island, Galapagos Archipelago.

Ophichthus evionthas Jordan and Bollman, Proc. U. S. Nat. Mus., XII,
1889, 154, Hood Island, Galapagos Archipelago.

Genus 226. OPHICHTHUS Thunberg and Ahl.

Ophichthus Ahl, De Muræna et Ophichtho, 1789 (*Muræna ophis* L.).

Cogrus Rafinesque, Caratteri, 1810, 62 (*C. maculatus* Rafinesque).

Centrurophis Kaup, Apodal Fishes, 1856, 2 (*C. spadiceus* Kaup).

Pæciolcephalus Kaup, loc. cit., 5 (*P. bonapartei* Kaup).

Microdontophis Kaup, loc. cit., 6 (*M. alipinnis* Kaup).

Cæciliophis Kaup, loc. cit., 6 (*C. compar* Kaup).

Herpetoichthys Kaup, loc. cit., 7 (*H. ornatissimus* Kaup).

Elapsopspis Kaup, loc. cit., 9 (*E. versicolor* Kaup).

Murænopsis Kaup, loc. cit., 11 (*Murænophis ocellatus* LeSueur); the name wrongly credited to LeSueur.

Scyhalophis Kaup, loc. cit., 13 (*S. magnioculis* Kaup).

Leptorhinophis Kaup, loc. cit., 14 (*Ophisurus gomesi* Castelnau).

Uranichthys Poey, Repertorio, II, 1867, 256 (*Muræna havannensis* Bloch and Schneider).

Oxyodontichthys Poey, Anales Soc. Espanola Hist. Nat., 1880, 254 (*macrurus*).

Ophichthys Bleeker, Günther, and of recent authors generally.

592. *Ophichthus ophis* (Linnaeus).

West Indies; apparently not common.

? *Serpens marinus maculosus* Willoughby, Hist. Pisc., 1686, Tab. G., 9, no locality.

? *Muræna ophis* Linnaeus, Syst. Nat., ed. X, 1758, 244; after Willoughby; may be identical with *Ophichthius regius*, a St. Helena species likewise spotted with black.

Innominado Parra, Dif. Piezas Hist. Nat., 1787, pl. 37, fig. 2, Habana.
Muræna havannensis Bloch and Schneider, Syst. Ichth., 1801, 491; after Parra.

? *Ophisurus guttatus* Cuvier, Règne Anim., ed. 1, 1817, 232, Surinam; after Bloch, pl. 154.

? *Muræna maculosa* Cuvier, loc. cit., Surinam; after *Ophisurus ophis* Lacépède, which is based on Bloch, pl. 154.

Herpetoichthys sulcatus Kaup, Apodal Fishes, 1856, 8, fig. 5, locality unknown.

Uranichthys brachycephalus Poey, Repertorio, II, 1867, 257, Cuba.

593. *Ophichthus retropinnis* Eigenmann.

Snapper Banks off Pensacola, Fla.

Ophichthys retropinnis Eigenmann, Proc. U. S. Nat. Mus., X, 1887, 116, Snapper Banks off Pensacola.

594. *Ophichthus guttifer* (Bean and Dresel).

Snapper Banks off Pensacola, Fla.

Ophichthys guttifer Bean and Dresel, Proc. Biol. Soc. Wash., II, 1884, 100, Snapper Banks.

595. *Ophichthus ocellatus* (LeSueur).

West Indies, south to Brazil and north to Pensacola.

Murænophis ocellatus LeSueur, Jour. Acad. Nat. Sci. Phila., V, 1825, 108, pl. 4, fig. 3, South America.

- 596. *Ophichthys triserialis* (Kaup).**
 Pacific coast of tropical America, rather common from Lower California to the Galapagos.
Murænopsis triserialis Kaup, Apodal Fishes, 1856, 12, Pacific.
Herpetoichthys callisoma Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 475, locality unknown.
Ophisurus californiensis Garrett, Proc. Calif. Acad. Nat. Sci., III, 1863, 66, Margarita Bay, Lower California.
Ophichthus rugifer Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 155, Charles Island, Galapagos; young with longer pectorals.
- 597. *Ophichthus biserialis* (Garman).**
 Galapagos Islands.
Ophichthys biserialis Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 311, Chatham Island.
- 598. *Ophichthus chamensis* Meek and Hildebrand.**
 Panama.
Ophichthus chamensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, 155, Pl. VIII, fig. 2, Chame Point, Panama.
- Subgenus SCYTALOPHIS Kaup.**
- 599. *Ophichthus gomesii* (Castelnau).** *Sea serpent; Sea eel; Whipsnake eel.*
 South Carolina to Rio Janeiro; generally common, especially about the Florida Keys and Cuba.
Ophisurus gomesii Castelnau, Anim. Amér. Sud, 1855, 84, pl. 44, fig. 2; Rio Janeiro.
Ophisurus chrysops Poey, Memorias, II, 1861, 321, Habana.
Oxydorichthys brachyurus Poey, Synopsis, 1868, 426; and Poey, Enumeratio, II, 1875, 155, Habana.
Oxydorichthys macrurus Poey, Anal. Soc. Espaniola Hist. Nat., 1880, 254, Habana.
Oxydorichthys limbatus Poey, loc. cit., 254, Habana; name a substitute for *brachyurus*.
- 600. *Ophichthus zophochir* (Jordan and Gilbert).**
 Pacific Coast of Mexico, Guaymas to Acapulco.
Ophichthys zophochir Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 347.
- 601. *Ophichthus magnioculus* (Kaup).**
 West Indies to Brazil; Aspinwall.
Scytalophis magnioculus Kaup, Apodal Fishes, 1856, 13, fig. 7, St. Croix; Brazil.
- 602. *Ophichthus parilis* (Richardson).**
 Cuba to Brazil.
Ophisurus parilis Richardson, Voy. *Erebus* and *Terror*, 1844, 105, West Indies.
Ophichthys pauciporus Poey, Repertorio, II, 1868, 255, pl. 3, fig. 5.
- Genus 227. MYSTRIOPHIS Kaup.**
Mystriophis Kaup, Apodal Fishes, 1856, 10 (*Ophisurus rostellatus* Richardson).
Crotolopsis Kaup, Abhandl. Natur. Verein Hamburg, IV, 1860, 12 (*C. quinquefasciata* Kaup).
Echiopsis Kaup, loc. cit., 13, (*Ophisurus intertinctus* Richardson).
Macrodonophis Poey, Repertorio, II, 1867, 251 (*M. mordax ophiurus* Poey).
- Subgenus ECHIOPSIS Kaup**
- 603. *Mystriophis intertinctus* (Richardson).**
 West Indies, north to Pensacola, Fla.
Ophisurus intertinctus Richardson, Voy. *Erebus* and *Terror*, Fishes, 1844, 102, West Indies.
Ophisurus sugillatus Richardson, loc. cit., 103, probably West Indies.
Crotalopsis punctifer Kaup, Abhandl. Natur. Verein Hamburg, IV, 1860, 2, 12, pl. 1, fig. 3, Puerto Cabello.
Conger mordax Poey, Memorias, II, 1861, 319, Cuba.
Ophichthys schneideri Steindachner, Ich. Beitr., VIII, 1879, 66, Brazil.

Genus 228. SCYTALICHTHYS Jordan and Davis.

Scytalichthys Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 635 (*Ophichthys miurus* Jordan and Gilbert).

604. *Scytalichthys miurus* (Jordan and Gilbert).

Cape San Lucas, Lower California.

Ophichthys miurus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 357, Cape San Lucas.

Genus 229. BRACHYSOMOPHIS Kaup.

Brachysomophis Kaup, Apodal Fishes, 1856, 9 (*B. horridus* Kaup).

?*Achrophichthys* Bleeker, Poisson Ined. Murènes, Ned. Tijdschr. Dierk., II, 242, 1865, (*A. typus*), young.

605. *Brachysomophis crocodilinus* (Bennett).

East Indies; a specimen recorded by Günther from the Galapagos.

Ophisurus crocodilinus Bennett, Proc. Zool. Soc. Lond., 1833, 32, Mauritius.

Brachysomophis horridus Kaup, Apodal Fishes, 1856, 9, fig. 6, Otaheite [Tahiti].

?*Achiropichthys typus* Bleeker, Nederl. Tijdschr. Dierk., II, 1865, 242, Celebes.

Family 78. ACANTHENCHELYIDÆ

Genus 230. ACANTHENCHELYS Regan.

Acanthenchelys Régan, Ann. Mag. Nat. Hist., X, 1922, 296 (*A. spinicauda* Régan).

606. *Acanthenchelys spinicauda* Régan.

West Indies.

Acanthenchelys spinicauda Régan, Ann. Mag. Nat. Hist., X, 1922, 296, Island of Tobago.

Family 79. MORINGUIDÆ

Genus 231. MORINGUA Gray.

Moringua Gray, Zool. Misc., 1831, 9 (*M. lateralis* Gray).

Rataboura Gray, Illust. Ind. Zool., 1831, 95 (*Muræna ratibouri* Hamilton = *Ratabouri hardwickei* Gray).

Pachyurus Swainson, Nat. Hist. Fish, Reptiles and Amphib., II, 1839, 335 (*Moringua linearis* Gray; not of Agassiz, 1829).

Pseudomoringua Bleeker, Nederl. Tijdschr. Dierk., II, 1865, 114 (*Moringua lumbricoidea* Richardson).

607. *Moringua boekei* Metzelaar.

Caribbean Sea.

Moringua boekei Metzelaar, Trop. Atl. Visschen, 1919, 17, Curaçao.

Genus 232. APHTHALMICHTHYS Kaup.

Aphthalmichthys Kaup, Apodal Fishes, 1856, 68 (*A. javanicus* Kaup).

608. *Aphthalmichthys caribbeus* Gill and Smith.

Porto Rico.

Aphthalmichthys caribbeus Gill and Smith, Science, XI, No. 286, 1900, 974, San Geronimo, near San Juan, Porto Rico.

Genus 233. STILBISCUS Jordan and Bollman.

Stilbiscus Jordan and Bollman, Proc. U. S. Nat. Mus., XI, 1888, 549 (*S. edwardsi* Jordan and Bollman).

609. *Stilbiscus edwardsi* Jordan and Bollman.

Green Turtle Cay, one of the Bahamas.

Stilbiscus edwardsi Jordan and Bollman, Proc. U. S. Nat. Mus., XI, 1888, 549, Green Turtle Cay, Bahamas.

Genus 234. GORDIICHTHYS Jordan and Davis.

Gordiichthys Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 644 (*G. irretitus* Jordan and Davis).

610. *Gordiichthys irretitus* Jordan and Davis.

Snapper Banks off west Florida in rather deep water.

Gordiichthys irretitus Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 644, off Pensacola, Fla.

Order T. COLOCEPHALI

Family 80. MURÆNIDÆ. Morays

Genus 235. ENCHELYCORE Kaup.

Enchelycore Kaup, Apodal Fishes, 1856, 72 (*Muræna nigricans* Bonnaterre).

611. *Enchelycore nigricans* (Bonnaterre).

West Indies; rather common.

Muræna unicolor maxillis elongatus teretiusculus, inferiore longiore, etc., Gronow, Zoophy., 1, 1763, 163, South America.

Muræna nigricans Bonnaterre, Encycl. Meth. Ichth., 1788, 34; after Gronow.

Muræna anguina Gronow, Cat. Fish., 1854, 18, South America.

Enchelycore euryrhina Kaup, Apodal Fishes, 1856, 73, no locality.

612. *Enchelycore nigrocastaneus* (Cope).

West Indies.

Gymnothorax nigrocastaneus Cope, Trans. Amer. Phil. Soc., XIV, 1871, 483, St. Martin.

?*Gymnothorax umbrosus* Poey, Ann. Lyc. Nat. Hist. N. Y., XI, 1874, 67, Habana, Cuba.

Genus 236. MURÆNA Linnæus. Morays.

Muræna Araldi, Gen. Pisc., 1738, 23, in part; includes all eels.

Muræna Klein, Hist. Pisc., 1742, 28, in part; includes all eels without pectoral fins.

Muræna Linnæus, Syst. Nat., ed. X, 1758, 244 (*helena*, etc.); includes all eels.

Muræna Thunberg and Ahl, De Muræna et Ophichthus, 1789, 6 (restricted to *helena*, etc.), includes species without pectoral fins.

Muræna Günther and of authors generally; not of Bleeker.

Murænophis Lacépède, Hist. Nat. Poiss., V, 1803, 630 (*helena*, etc.).

Limamuræna Kaup, Apodal Fishes, 1856, 95 (*L. guttata* Kaup).

613. *Muræna melanotis* (Kaup).

Tropical Atlanta, from Africa to South America.

Limamuræna melanotis Kaup, Aalen. Fische Hamburg Mus., 1859, 27, pl. 4, fig. 3.

614. *Muræna lentiginosa* Jenyns. *Morena pinta*.

Pacific coast of America, from Gulf of California to Galapagos.

Muræna lentiginosa Jenyns, Voy. Beagle, Zool., 1842, 143, Galapagos Islands.

Muræna pinta Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 345, Mazatlan.

615. *Muræna retifera* Goode and Bean.

Coast of South Carolina, in rather deep water; one taken at Tuckernuck Island, Mass.

Muræna retifera Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 435, off Charleston.

616. *Muræna argus* (Steindachner).

Gulf of California.

Gymnothorax (*Limamuræna*) *argus* Steindachner, Ichth. Notizen, X, 1870, 17, Pl. IV, Altata, Gulf of California.

617. *Muræna insularum* Jordan and Davis.

Galapagos Islands.

Muræna insularum Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 609, Chatham Island.

618. *Muræna myrialeucosticta* Fowler.

West Indies.

Muræna myrialeucosticta Fowler, Proc. Acad. Nat. Sci. Phila., LXIV, 1912, 18, St. Martin.

619. *Muræna clepsydra* Gilbert.

Panama.

Muræna clepsydra Gilbert, in Jordan and Evermann, Fish. North and Mid. Amer., 1898, 2805, Panama.

Genus 237. PYTHONICHTHYS Poey.

Pythonichthys Poey, Repertorio, II, 1867, 265 (*P. sanguineus* Poey).

620. *Pythonichthys sanguineus* Poey.

Coast of Cuba, in rather deep water.

Pythonichthys sanguineus Poey, Repertorio, II, 1867, 265, pl. 2, fig. 7, Cuba.

Genus 238. *RABULA* Jordan and Davis.

Rabula Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 589
(*R. aquæ-dulcis* Jordan and Davis = *Rabula davisi* Fowler, not
Muræna aquæ-dulcis Cope).

621. *Rabula davisi* Fowler.

Said to be from San Diego, which is doubtful.

Rabula davisi Fowler, Proc. Acad. Nat. Sci. Phila., LXIV, 1912, 21;
after *Rabula aquæ-dulcis* Jordan and Davis; not of Cope.

622. *Rabula marmorata* (Valenciennes).

Galapagos Islands.

Murænophis marmoratus Valenciennes, Voy. Vénus, Zool., 1855, 347, pl.
10, fig. 1, Galapagos.

623. *Rabula panamensis* (Steindachner).

Pacific coast of Central America.

Muræna panamensis Steindachner, Ichth. Beitr., V, 1876, 19, Panama.

624. *Rabula longicauda* (Peters).

Tropical Atlantic, off the West Indies.

Muræna longicauda Peters, Monatsberichte, 1876, 850, open Atlantic,
with *Sphagebranchus anguiformis*.

Genus 239. *GYMNOTHORAX* Bloch. *Morays.*

Gymnothorax Bloch, Naturgesch. Ausl. Fische, IX, 1795, 83, restricted
to allies of *M. afra* by Günther (*G. muræna* Bloch).

Lycodontis McClelland, Calcutta Jour. Nat. Hist., V, No. XVIII,
1844, 173 (*L. literata* = tile).

Throndontis McClelland, loc. cit., V, No. XVIII, 1844, 174 (*T. reticulata* = *tessellata*).

Eurymyctera Kaup, Apodal Fishes, 1856, 72, (*E. crudelis* Kaup).

Polyuranodon Kaup, loc. cit., 96 (*Muræna polyuranodon* Bleeker).

Tæniophis Kaup, Aalen. Fische Hamburg Mus., Nachträge, 1859, 10
(*T. westphali* Kaup).

Priodontophis Kaup, loc. cit., 1859, 22 (*Gymnothorax ocellatus* Agassiz).

Neomuræna Girard, U. S. Mex. Bound. Surv., Fishes, 1859, 76 (*N.
nigromarginata* Girard = *ocellatus*).

Pseudomuræna Johnson, Proc. Zool. Soc. Lond., 1860, 167 (*P. madeirensis* Johnson).

625. *Gymnothorax mordax* (Ayres). *Conger eel of California.*

Point Conception to Cedros Island; abundant about Santa Barbara
Islands.

Muræna mordax Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 30,
Cerros [Cedros] Island.

626. *Gymnothorax moringa* (Cuvier). *Common spotted moray; Hamlet;
Bel.*

Gulf of Mexico, Pensacola, and West Indies to Rio Janeiro and St.
Helena.

Muræna maculata nigra; the black moray, Catesby, Nat. Hist. Caro-
linas, etc., 1738, pl. 21, Bahamas.

?*Gymnothorax afer* Bloch, Syst. Ichth., 1795, pl. 417, Africa.

Muræna moringa Cuvier, Règne Anim., ed. 2, II, 1829, 352; after
Catesby.

Gymnothorax rostratus Agassiz, Spix, Pisc. Brasil., 1830, 91, pl. 50a,
Brazil.

Muræna moringua Richardson, Voy. Erebus and Terror, Fishes, 1844, 89,
Jamaica.

Muræna punctata Gronow, Cat. Fish., 1854, 18, North America.

Murænophis curvilineata Castelnau, Anim. Amér. Sud, Poiss., 1855, 81, pl.
42, fig. 2, Rio Janeiro.

Murænophis caramuru Castelnau, loc. cit., 85, pl. 43, Bahia.

Gymnothorax flavoscriptus Poey, Enumeratio, 1875, 158, Cuba.

Gymnothorax picturatus Poey, Anal. Soc. Espaniola Hist. Nat., 1880,
257, Cuba.

627. *Gymnothorax funebris* Ranzani. *Black moray; Morena verde.*
 Tropical America; Rio Janeiro; West Indies.
Muræna maculata nigra et viridis (the moray) Catesby, Nat. Hist. Carolinas, etc., 1738, pl. 20, Bahamas.
Gymnothorax funebris Ranzani, Nov. Com. Acad. Sci. Inst. Bonon., IV, 1840, 76, Brazil.
Muræna lineopinnis Richardson, Voy. *Erebus* and *Terror*, Fishes, 1844, 89, Puerto Cabello.
Muræna infernalis Poey, Memorias, II, 1861, 347, 354, Cuba.
Thyrsodea concolor Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 479, Vera Cruz.
Muræna eebus Poey, Memorias, II, 1861, 426, Cuba.
628. *Gymnothorax castaneus* (Jordan and Gilbert).
 Pacific coast of Mexico; common about Mazatlan; representing *G. funebris* of the West Indies.
Sidera castanea Jordan and Gilbert, Proc. U. S. Nat. Mus., V. 1882, 647, Mazatlan.
629. *Gymnothorax sanctæ-helenæ* (Günther).
 Tropical Atlantic; recorded from St. Helena and the Bermudas.
Muræna sanctæ-helenæ Günther, Cat., VIII, 1870, 115, St. Helena.
630. *Gymnothorax vicinus* (Castelnau).
 Tropical Atlantic, Cuba to Africa and Brazil.
Murænopsis vicina Castelnau, Anim. Amér. Sud, Poiss., 1855, 81, pl. 42, fig. 4, Bahia.
Gymnothorax versipunctatus Poey, Enumeratio, 1875, 156, Cuba.
Thyrsodea maculipinnis Kaup, Apodal Fishes, 1856, 86, Gold Coast.
631. *Gymnothorax punctarenæ* Fowler.
 Puntarenas, Costa Rica, one specimen known.
Gymnothorax punctarenæ Fowler, Copeia, No. 39, Jan. 24, 1917, 3, Puntarenas, Costa Rica.
632. *Gymnothorax brunneus* Nichols.
 Bermudas.
Gymnothorax brunneus Nichols, Proc. Biol. Soc. Wash., XXXIII, 1920, 59, Bermuda.
633. *Gymnothorax virescens* Poey.
 Cuba.
Gymnothorax virescens Poey, Enumeratio, 1875, 156, Cuba.
634. *Gymnothorax polygonius* Poey.
 Cuba.
Gymnothorax polygonius Poey, Ann. Lyc. Nat. Hist. N. Y., XI, 1870, 68, Habana.
635. *Gymnothorax dovii* (Günther). *Morena pintita.*
 Gulf of California to the Galapagos Archipelago.
Muræna dovii Günther, Cat., VIII, 1870, 103, Panama.
Muræna pintita Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 346, Mazatlan.
636. *Gymnothorax conspersus* Poey.
 Cuba to Rio Janeiro.
Murænopsis punctata Castelnau, Anim. Amér. Sud, 1855, 82, Rio Janeiro; not *Gymnothorax punctatus* Bloch.
Gymnothorax conspersus Poey, Repertorio, II, 1868, 259, Cuba.
637. *Gymnothorax jordani* (Evermann and Marsh).
 Porto Rico.
Lycondontis jordani Evermann and Marsh, Fish. Porto Rico, Bull. U. S. Fish Com., XX, 1900 (1902), pt. 1, 78, pl. 2; off Mayaguez, Porto Rico.
638. *Gymnothorax miliaris* (Kaup).
 West Indies.
Thyrsodea miliaris Kaup, Apodal Fishes, 1856, 90, Martinique.
Muræna multocellata Poey, Memorias, II, 1861, 324, Cuba; based on specimens with smaller spots than usual.
Gymnothorax scriptus Poey, Repertorio, II, 1868, 261, Cuba.
Muræna myrialeucostictus Fowler, Proc. Acad. Nat. Sci. Phila., LXIV, 1912, 18, St. Martins, West Indies.

639. *Gymnothorax elaboratus* (Poey).
Cuba.
Muræna elaborata Poey, Memorias, II, 1861, 323, Cuba.
640. *Gymnothorax obscuratus* Poey.
Cuba.
Gymnothorax obscuratus Poey, Ann. Lyc. Nat. Hist. N. Y., IX, 1870, 320, Cuba.
641. *Gymnothorax chlevastes* (Jordan and Gilbert).
Galapagos Islands.
Sidera chlevastes Jordan and Gilbert, Proc. U. S. Nat. Mus., VI, 1883, 208, Galapagos.
642. *Gymnothorax albimentis* (Evermann and Marsh).
Porto Rico.
Lycodontis albimentis Evermann and Marsh, Fish. Porto Rico, Bull. U. S. Fish Com., XX, 1900 (1902), pt. 1, 78, fig. 9; off Culebra Island.
643. *Gymnothorax concolor* (Abbott).
West Indies.
Thyrsoidea concolor Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 479, Vera Cruz.
? *Muræna erebus* Poey, Memorias, II, 1861, 426, Cuba.
644. *Gymnothorax aquædulcis* (Cope).
Rio Grande, Costa Rica.
Muræna aquæ-dulcis Cope, Rept., U. S. Geol. Surv., 1871 (1872), 474; said to be from Rio Grande, Costa Rica. It is evidently not a freshwater fish. According to Fowler, it is not a *Rabula*, the dorsal beginning well before the gill opening.
645. *Gymnothorax carcinognathus* Fowler.
West Indies.
Gymnothorax carcinognathus Fowler, Proc. Acad. Nat. Sci. Phila., LXIV, 1912, 22, St. Martins, West Indies.
646. *Gymnothorax verilli* (Jordan and Gilbert).
Panama; one specimen known.
Sidera verrilli Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 623, Panama.
647. *Gymnothorax ocellatus* Agassiz. *Spotted moray; Moray eel.*
Pensacola to Rio Janeiro.
Gymnothorax ocellatus Agassiz, Spix, Pisc. Brasil., 1828, 91, pl. 50b, Brazil.
648. *Gymnothorax ocellatus saxicola* Jordan and Davis.
Snapper Banks off Pensacola and deep water off Cuba.
Muræna meleagris Quoy and Gaimard, Voy. Freycinet, Zool., 1824, 245, pl. 52, fig. 2; not of Shaw.
Gymnothorax ocellatus saxicola Jordan and Davis, Rept. U. S. Fish Com., XVI, 1888, 606, Snapper Banks off Pensacola.
649. *Gymnothorax ocellatus nigromarginatus* (Girard).
Coast of west Florida and Texas.
Neomuræna nigromarginata Girard, U. S. Mex. Bound. Surv., 1859, 76, pl. 41, St. Joseph Island, Tex.
- Genus 240. *SIDERA* Kaup.
Siderea Kaup, Uebersicht der Aale, 1856, 59 (*Muræna pfeifferi* Bleeker = *Muræna picta* Ahl); name later written *Sidera*.
650. *Siderea picta* (Ahl).
South Seas; East Indies; Hawaii; Revillagigedos; taken at Clarion Island.
Muræna picta Ahl, De Muræna et Ophichthœ, VI, 1787, 6, tab. 2.
Murænophis pantherina Lacépède, Hist. Nat. Poiss., V, 1803, 628.
Muræna variegata Quoy and Gaimard, Voy. Uranie, 1824, 246, pl. 52.
Muræna lita Richardson, Voy. Erebus and Terror, 1844, 84, Moluccas.
Muræna siderea Richardson, loc. cit. 85, pl. 48, Australia.
Muræna pfeifferi Bleeker, Nat. Tijds. Ned. Ind., V, 173, Celebes.

Genus 241. AHYNNODONTOPHIS Fowler.

Ahynnodontophis Fowler, Proc. Acad. Nat. Sci. Phila., LXIV, 1912, 25 (*A. stigmanotus* Fowler).

651. *Ahynnodontophis stigmanotus* Fowler.

West Indies (?)

Ahynnodontophis stigmanotus Fowler, Proc. Acad. Nat. Sci. Phila., LXIV, 1912, 25; supposed to be from the West Indies.

Genus 242. ECHIDNA Forster.

Echidna Forster, Bibliotheca Banksiae, 1777, 181 (*E. variegata* Forster).
Gymnomuræna Lacépède, Hist. Nat. Poiss., V, 1803, 648 (*G. doliatæ* Lacépède).

Gymnopsis Rafinesque, Anal. Nat., 1815, 93 (substitute for *Gymnomuræna*).

Megaderia Rafinesque, loc. cit., 93 (substitute for *Echidna* Forster).
Molarii Richardson, Voy. *Erebus* and *Terror*, 1844, 79 (*Muræna ophis* = *Echidna nebulosa*).

Pæciolophis Kaup, Apodal Fishes, 1856, 98 (*Gymnothorax catenatus* Bloch).

652. *Echidna catenata* (Bloch).

West Indies, from Bermuda to Surinam; our specimens from Santa Lucia.

Muræna seu conger brasiliensis Seba, Thesaurus, II, 1738, 72, pl. 69, figs. 4, 5, Brazil.

Gymnothorax catenatus Bloch, Naturgesch. Ausl. Fische, XII, 1795, 74, pl. 415, fig. 1, Coromandel; an error.

Muræna sordida Cuvier, Règne Anim., ed. 1, 1817, 233, after Seba.

Muraenophis catenula Lacépède, Hist. Nat. Poiss., V, 1803, 628 and 641, Palmerston Island; after Bloch.

Echidna fuscomaculata Poey, Repertorio, II, 1868, 263, Cuba.

Echidna flavoscripta Poey, loc. cit., 264, Cuba.

653. *Echidna nocturna* (Cope).

Pacific coast of Mexico, the two known specimens from the Rio Grande, in Costa Rica and from Cape San Lucas.

Pæciolophis nocturnus Cope, Rept. U. S. Geol. Surv. Mont., 1871 (1872), 474, Rio Grande at San José, Costa Rica.

654. *Echidna scabra* Garman

Cocos Island.

Echidna scabra Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 314, Cocos Island.

655. *Echidna cocosa* Garman.

Cocos Island.

Echidna cocosa Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 313, Cocos Island.

656. *Echidna chionostigma* Fowler.

Gulf of California (?).

Echidna chionostigma Fowler, Proc. Acad. Nat. Sci. Phila., LXIV, 1912, 28; supposed to be from the Gulf of California.

Genus 243. UROPTERYGIUS Rüppell.

Gymnomuræna Lacépède, Hist. Nat. Poiss., V, 1803, 648 (*G. doliatæ* Lacépède); restricted first by Kaup, in 1856, to *doliata*, which is a species of *Echidna*.

Ichthyophis Lesson, Voy. Coquille, II, 1830, 120 (*I. pantherinus* Lesson); not of Fitzinger, 1829, a genus of reptiles.

Uropterygius Rüppell, Neue Wirbelthiere, Fische, 1835, 83 (*Nebrius concolor* Rüppell).

Muraenoblenna Kaup, Apodal Fishes, 1856, 67, (*Ichthyophis tigrina* Lesson); not of Lacépède, 1803, which is *Myxine*.

Scuticaria Jordan and Evermann, Fish. North and Mid. Amer., 1896, 403, (*Uropterygius necturus* Jordan and Gilbert).

Subgenus SCUTICA Jordan and Evermann.

657. *Uropterygius necturus* (Jordan and Gilbert).

Gulf of California.

Gymnomuræna nectura Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 356, Cape San Lucas.

Genus 244. CHANNOMURÆNA Richardson.

Channo-Muræna Richardson, Voy. Erebus and Terror, 1844, 96 (*Ichthyo-*
phis vittata Richardson).

658. *Channomuræna vittata* (Richardson).

Coast of Cuba.

Raro Parra, Dif. Piezas Hist. Nat., 1780, 66, pl. 30, fig. 3, Habana.

Ichthyothis vittatus Richardson, Voy. Sulphur, Fishes, 1844, 114, pl. 53,
figs. 7-9; locality uncertain.

Channomuræna cubensis Poey, Repertorio, II, 1867, 266, pl. 3, fig. 6,
Cuba.

Order U. LYOMERI

Family 81. SACCOPHARYNGIDÆ. Gulpers

Genus 245. SACCOPHARYNX Mitchell.

Saccopharynx Mitchell, Ann. Lyc. Nat. Hist. N. Y., 1824, 86 (type afterwards called *S. flagellum* Mitchell).

Ophiognathus Harwood, Philos. Trans., 1827, 277 (*O. ampullaceus* Harwood).

659. *Saccopharynx ampullaceus* (Harwood).

Atlantic.

Ophiognathus ampullaceus Harwood, Philos. Trans., 1827, 52, Atlantic.

Saccopharynx flagellum Cuvier, Règne Anim., ed. 2, 1829, 355, Atlantic;
after Mitchell.

Saccopharynx chordatus Storer, Synopsis, 1846, 237, open Atlantic; after
Mitchill.

Family 82. EURYPHARYNGIDÆ. Swallowers

Genus 246. GASTROSTOMUS Gill and Ryder.

Gastrostomus Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883, 271
(*G. bairdii* Gill and Ryder).

660. *Gastrostomus bairdii* Gill and Ryder.

Deep water off Newfoundland Banks and Davis Strait.

Gastrostomus bairdii Gill and Ryder, Proc. U. S. Nat. Mus., VI, 1883,
271, off the Grand Banks of Newfoundland.

661. *Gastrostomus pacificus* Bean.

Pacific Ocean in deep water.

Gastrostomus pacificus Bean, Smiths. Misc. Coll., XLV, 1904, 254,
fig. 31; halfway between Midway Island and Guam.

Series OSTARIOPHYSI

Order V. HETEROGNATHI

Family 83. CHARACINIDÆ. Characins

Genus 247. PHENAGONIATES Eigenmann and Wilson.

Phenagoniates Eigenmann and Wilson, Indiana Univ. Studies, XIX,
1914, 2 (*P. wilsoni* Eigenmann = *Ræboides macrolepis* Meek and
Hildebrand).

662. *Phenagoniates macrolepis* (Meek and Hildebrand).

Atlantic slope of Panama.

Ræboides macrolepis Meek and Hildebrand, Field Mus. Nat. Hist.,
Zool., X, 1913, '84, Rio Cupe, Boca de Cupe, Tuyra River Basin.

Genus 248. COMPSURA Eigenmann.

Compsura Eigenmann, Mem. Carnegie Mus., VII, 1915, 60 (*Compsura*
heterura Eigenmann).

663. *Compsura gorgonæ* (Evermann and Goldsborough).

Both slopes of Panama.

Cheirodon gorgonæ Evermann and Goldsborough, Proc. Biol. Soc. Wash., XXII, 1909, 99, figs. 1, 3, Gorgona, Atlantic slope, Panama Canal Zone.

Genus 249. *PSEUDOCHIRODON* Meek and Hildebrand.

Pseudocheirodon Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 275 (*P. affinis* Meek and Hildebrand).

664. *Pseudocheirodon affinis* Meek and Hildebrand.

Both slopes of Panama.

Cheirodon insignis Evermann and Goldsborough, Proc. Biol. Soc. Wash., XXII, 1909, 98, Tabernilla, Atlantic slope; not of Steindachner.

Pseudocheirodon affinis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 275, Pl. XVIII, Rio Gatun, Monte Liria, Canal Zone, Panama.

Genus 250. *GEPHYROCHARAX* Eigenmann.

Gephyrocharax Eigenmann, Indiana Univ. Studies, No. 16, 1912, 23 (*G. chocöensis* Eigenmann).

665. *Gephyrocharax atricaudatus* (Meek and Hildebrand).

Both slopes, Canal Zone, Panama.

Deuterodon atricaudata Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1912, 68, Rio Frijoles, Canal Zone.

Gephyrocharax atricaudatus Eigenmann, Indiana Univ. Studies, No. 20, 1914, 41, Panama Canal Zone.

666. *Gephyrocharax intermedius* Meek and Hildebrand.

Rio Chama, Panama.

Gephyrocharax intermedius Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 278, Rio Chama.

Genus 251. *ASTYANAX* Baird and Girard.

Astyianax Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 26 (*A. argentatus* Baird and Girard = *Tetragonopterus mexicanus* Filippi).

Pacilurichthys Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 417 (*P. brevoortii* Gill).

All the species mentioned below belong to the subgenus or genus *Astyianax*, distinguished from *Tetragonopteroides* Cuvier by the absence of teeth on the maxillary and by the shorter anal of less than 40 rays.

667. *Astyianax mexicanus* (Filippi).

Rio Balass and Rio Tehuacan to tributaries of the Rio Grande in Texas, the northernmost species of the multitude of Characins.

Tetragonopterus mexicanus Filippi, Guerin's Rev. Mag. Zool., 1853, 166, Mexico.

Astyianax argentatus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 27, Comanche Spring and Brownsville, Tex.

Tetragonopterus fulgens Bocourt, Ann. Sci. Nat. Zool., IX, 62, 1868.

Tetragonopterus nitidus Vaillant and Pellegrin, Bull. Mus. Hist. Nat. Paris, 1904, 324.

Tetragonopterus brevimanus Günther, Cat., V, 1864, 325, Yzabel; Rio San Gerónimo, Guatemala.

668. *Astyianax aeneus* (Günther).

Oaxaca, Mexico, and Rio Chagres, and other streams about Panama, south to Brazil.

Tetragonopterus aeneus Günther, Proc. Zool. Soc. Lond., 1860, 319, Oaxaca, Mexico,

Tetragonopterus microphthalmus Günther, Cat., V, 1864, 324, Lake Amatitlan, Guatemala; Rio Rimac.

Tetragonopterus panamensis Günther, loc. cit., 324, Pacific coast of Panama; Yzabel.

Tetragonopterus humilis Günther, loc. cit., 327, Lake Amatitlan, Guatemala.

Tetragonopterus aerstedi Krøyer, in Lütkin, Ict. Bidrag., III, 1874, 229, Rio San Juan, Nicaragua.

- 669. *Astyanax petenensis* (Günther).**
 Nicaragua to western Ecuador; southeast to Argentine Republic.
Tetragonopterus petenensis Günther, Cat., V, 1864, 326, Lake Peten,
 Nicaragua.
- 670. *Astyanax scabripinnis* (Jenyns).**
 Jamapa, Mexico, to Rio Janeiro.
Tetragonopterus scabripinnis Jenyns, Voy. *Beagle*, Zool., Fishes, 1842,
 125, Rio Janeiro, Brazil.
- 671. *Astyanax macrophthalmus* (Regan).**
 Streams of eastern Mexico.
Tetragonopterus macrophthalmus Regan, Biol. Centrali-Amer., 1915,
 171, eastern Mexico; Monzorangi, Vera Cruz.
- 672. *Astyanax nasutus* Meek.**
Astyanax nasutus Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 108,
 Lake Managua, Nicaragua.
- 673. *Astyanax fasciatus* (Cuvier).**
 Coasts of Brazil, from Para to Napo and on the Pacific side from
 Panama to Ecuador.
Tetragonopterus Fischeri Steindachner, Flussfische Südamerikas, I, 1879,
 166, Mamo River at Chepo, Colombia.
- 674. *Astyanax ruberrimus* Eigenmann.**
 Panama, both slopes.
Astyanax ruberrimus Eigenmann, Indiana Univ. Studies, XVIII, 1913,
 25, Isthmeria, Atlantic slope.
- 675. *Astyanax brevoortii* (Gill).**
 Island of Trinidad.
Pacilurichthys brevoortii Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 417.
 Trinidad.
Pacilurichthys tenuirus Gill, loc. cit., 418, Trinidad.
Pacilurichthys pulcher Gill, loc. cit., 419, Trinidad.
- Genus 252. *BRYCONAMERICUS* Eigenmann.**
Bryconamericus Eigenmann, Ann. Carnegie Mus., IV, 1907, 139, (*B. exodon* Eigenmann).
- 676. *Bryconamericus emperador* (Eigenmann and Ogle).**
 Streams of both slopes, Panama.
Astyanax emperador Eigenmann and Ogle, Proc. U. S. Nat. Mus.,
 XXXIII, 1907, 26, Empire Station, Panama.
- 677. *Bryconamericus cascajalensis* Meek and Hildebrand.**
 Panama.
Bryconamericus cascajalensis Meek and Hildebrand, Field Mus. Nat.
 Hist., Zool., X, 1916, 284, Rio Cascajal, Puerto Bello, Panama.
- Genus 253. *HEMIBRYCON* Günther.**
Hemibrycon Günther, Cat., V, 1864, 318 (*Tetragonopterus (Hemibrycon) polyodon* Günther).
- 678. *Hemibrycon dariensis* Meek and Hildebrand.**
 Basin of Rio Tuyra, Panama.
Hemibrycon dariensis Meek and Hildebrand, Field Mus. Nat. Hist.,
 Zool., X, 1916, 285, Pl. XX, mouth of Rio Yape, Panama.
- Genus 254. *HYPHESSOBRYCON* Durbin.**
Hypessobrycon Durbin, in Eigenmann, Bull. Mus. Comp. Zool., LII,
 1908, 100, (*Hemigrammus compressus* Meek).
Dermatocheir Durbin, Ann. Carnegie Mus., 1909, 55 (*D. calablepta* Durbin).
- 679. *Hypessobrycon panamensis* Durbin.**
 Rio Chagres Basin, Panama.
Hypessobrycon panamensis Durbin, in Eigenmann, Bull. Mus. Comp.
 Zool., LII, 1908, 101, Rio Boqueron, Atlantic slope, Panama.
Hemigrammus minutus Meek and Hildebrand, Field Mus. Nat. Hist.,
 Zool., X, 1912, 67, Agua Clara, Panama, Atlantic side of Isthmus of
 Panama.

Genus 255. HEMIGRAMMUS Gill.

Hemigrammus Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 420 (*H. unilineatus* Gill).

680. *Hemigrammus unilineatus* (Gill).

Streams of Trinidad.

Pacilurichthys unilineatus Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1854, 420, Trinidad.

Genus 256. CREAGRUTUS Günther.

Creagrutus Günther, Cat., V, 1864, 339 (*Leporinus mulleri* Günther).

681. *Creagrutus notropoides* Meek and Hildebrand.

Rio Chagres, Panama.

Creagrutus notropoides Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1912, 68, Rio Indio, Chagres basin.

682. *Creagrutus simus* Meek and Hildebrand.

Rio Tuyra Basin, Panama.

Creagrutus simus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1912, 85, Rio Cupe, Cituro, Panama.

Genus 257. RÖBOIDES Günther.

Raboides Günther, Cat., V, 1864, 347 (*Epicyrtus microlepis* Reinhardt).

683. *Raboides guatemalensis* (Günther).

Rio Chagres and streams of Guatemala.

Anacyrtus guatemalensis Günther, Cat., V, 1864, 347, Rio Chagres; Huamuchal.

684. *Raboides salvadoris* Hildebrand. *Plateado*; *Sardina*; *Alma seca*; *Ulmina*.

Lakes Guija and Metapan and Rio Sucio and Lempa, El Salvador.

Raboides salvadoris Hildebrand, Bull. U. S. Bur. Fish., XLI, 1925, 246, fig. 9, Rio Sucio, Sitio del Niño.

685. *Raboides occidentalis* Meek and Hildebrand.

Pacific slope of Panama and southward.

Raboides occidentalis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 293, Pl. XXIII, Rio Cardenas, Corozal, Panama.

Genus 258. BRYCON Müller and Troschel.

Brycon Müller and Troschel, Horae Ichth., I, 1845, 15 (*B. falciatus* Müller and Troschel).

Chalcinopsis Kner, Sitzungsb. Akad. Wiss. München, 1863, 226 (*C. striatulus* Kner).

Megalobrycon Günther, Proc. Zool. Soc. Lond., 1869, 423 (*Brycon cephalus* Günther).

686. *Brycon striatulus* (Kner).

Rio Chagres and streams about Panama.

Chalcinopsis striatulus Kner, Sitzungsb. Akad. Wiss. München, 1863, 226, Panama.

Chalcinopsis chagresensis Kner, loc. cit., 1863, 226, Rio Chagres.

687. *Brycon guatemalensis* Regan.

Brycon guatemalensis Regan, Biol. Centrali-Amer., 1908, 168, lakes and streams of Guatemala and Nicaragua.

688. *Brycon petrosus* Meek and Hildebrand. *Pipon*.

Chagres River basin, Atlantic slope of Panama.

Brycon petrosus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 84, Pl. XXIV, 1916; upper Chagres, Panama.

689. *Brycon dentex* Günther.

Yucatan to Ecuador.

Brycon dentex Günther, Proc. Zool. Soc. Lond., 1860, 240, Esmeraldas, Ecuador.

690. *Brycon argenteus* Meek and Hildebrand.

Rio Charrrera, and in Rio Bayano and Rio Tuyra basins, Pacific slope of Panama.

Brycon argenteus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 84, Pl. XXV, 298, 1916; Rio Aruza, Aruza, Panama.

691. *Brycon oligolepis* Regan.

Streams of Panama.

Brycon oligolepis Regan, Ann. Mag. Nat. Hist., IX, 1913, 462, Rio Condoto, Panama.

Genus 259. *BRAMOCHARAX* Gill.

Bramocharax Gill, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 189 (*B. transfordi* Gill).

692. *Bramocharax transfordi* Gill.

Lake Nicaragua.

Bramocharax transfordi Gill, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 190, Lake Nicaragua.

693. *Bramocharax elongatus* Meek. *Sabalito*.

Bramocharax elongatus Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 110, Lake Managua, Nicaragua.

Genus 260. *PIABUCINA* Cuvier and Valenciennes.

Piabucina Cuvier and Valenciennes, Hist. Nat. Poiss., XXII, 1849, 122 (161) (*P. erythrinoides* Cuvier and Valenciennes).

694. *Piabucina panamensis* Gill.

Tripoli, Atlantic slope of the Isthmus of Panama.

Piabucina panamensis Gill, Proc. Acad. Nat. Sci. Phila., XXVIII, 1876, 336, Rio Tripoli, Isthmus of Panama.

695. *Piabucina festae* Boulenger.

Panama.

Piabucina festae Boulenger, Bull. Mus. Torino, XIV, 1899, 346, I, Rio Sabina, Panama.

Genus 261. *HOPLIAS* Gill.

Macrodon Müller, in Wiegmann's Archiv, 1843, 163, (*Macrodon trahira* =

Synodus malabaricus Schneider); name preoccupied in *Sciaenidae*.

Hoplias Gill, Proc. U. S. Nat. Mus., XXVI, 1903, 1016 (*Synodus malabaricus* Schneider).

696. *Hoplias malabaricus* (Bloch).

Rio Tuya Basin; both slopes of Colombia; Ecuador, and south to Brazil.

Esox malabaricus Bloch, Naturgesch. Ausl. Fische, pt. 8, 1794, 149, pl. 392, Tranquebar; locality wrongly given.

Macrodon tereira Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 508, Bahia; San Francisco; Amazon; Maracaibo.

Erythrinus trahira Agassiz, in Spix, Pisc. Brasil., 1829, pl. 18, Bahia; San Francisco; Amazon; Maracaibo.

Erythrinus macrodon Agassiz, loc. cit., 43, pl. 18, Alma, Prov. Bahia, San Francisco.

Erythrinus microcephalus Agassiz, loc. cit., 44, San Francisco.

Erythrinus brasiliensis Agassiz, loc. cit., 45, pl. 20, Peruaguacu.

Macrodon auritus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 519, Montevideo.

Macrodon teres Cuvier and Valenciennes, loc. cit., 521, Lake Maracaibo.

Macrodon palana Cuvier and Valenciennes, loc. cit., 522, Cayenne.

Macrodon cimara Cuvier and Valenciennes, loc. cit., 586, Cayenne.

Macrodon ferox Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 51, Trinidad.

697. *Hoplias microlepis* (Günther).

Panama to Ecuador.

Macrodon microlepis Günther, Cat., V, 1864, 282, western Ecuador, Guatemala.

Family 84. GASTEROPELICIDÆ.

Genus 262. *GASTEROPELECUS* Gronow.

Gasteropeleucus Gronow, Mus. Ichth., II, 1763, 7 (type given in Gen. Fish. as *Clupea sternica* L.).

698. *Gasteropeleucus maculatus* Steindachner. *Peché-peche*.

Rio Mamoni, near Panama.

Gasteropeleucus maculatus Steindachner, Flussfische Südamerikas, 1879, 168, Rio Mamoni, at Chepo, Panama.

699. *Gasteropelecus sternicla* Pallas.

Surinam; Brazil; Essequibo.

Gasteropelecus sternicla Pallas, Spic. Zool., 1780, fasc. 8, p. 50, tab. 3,
fig. 4.

Family 85. XIPHOSTOMIDÆ

Genus 263. *CTENOLUCIUS* Gill.*Ctenolucius* Gill, Suppl., Proc. Acad. Nat. Sci. Phila., XIII, 1861, 8
(*Luciocharax insculptus* Steindachner).*Luciocharax* Steindachner, Denkschr. Akad. Wiss. Wien, XXXIX,
1878 (1879), 67 (*L. insculptus* Steindachner).700. *Ctenolucius insculptus* (Steindachner).

Rio Magdalena and Rio Momoni near Panama.

Luciocharax insculptus Steindachner, Denkschr. Akad. Wiss. Wien,
XXXIV, 1878 (1879), 67 Rio Magdalena, and Rio Mamoni near
Panama.701. *Ctenolucius beani* (Fowler).

Pacific coast of Panama; both slopes of Colombia.

Belonocharax beani Fowler, Proc. Acad. Nat. Sci. Phila., LVIII, 1906,
464, Rio Truando, a tributary of Rio Atrato, Panama.*Luciocharax striatus* Boulenger, Ann. Mag. Nat. Hist., ser. 8, VII,
1911, 312, Boca de Calmia, Choco.

Family 86. ANOSTOMIDÆ. (Curimatidae)

Genus 264-265. *CURIMATA* Walbaum.*Curimata* Walbaum, Artedi Pisc., 1792, 80 (*Salmo marcgravii* Walbaum,
based on *Charax maxilla superiore longiore* Gronow, which is *Salmo cyprinoides* Linnaeus).*Les Curimates* Cuvier, Règne Anim., ed. 1, II, 1817, 165 (*Salmo edentulus*
Bloch).*Curimatus* of authors generally; altered spelling.*Curimatella* Eigenmann and Eigenmann, Ann. N. Y. Acad. Sci., IV,
1889, 415 (*C. lepidurus* Eigenmann and Eigenmann).*Semitapicus* Eigenmann and Eigenmann, loc. cit., 415 (*Curimatus spilurus* Günther).701a. *Curimata magdalense* Steindachner. *Sardina blanca*.

Rio Magdalena and Rio Momoni, Colombia.

Curimatus magdalense Steindachner, Denkschr. Akad. Wiss. Wien,
XXXIX, 1878 (1879), 34, Rio Magdalena, Colombia.

Family 87. HEMIODONTIDÆ

Genus 266. *CHARACIDIUM* Reinhardt.*Characidium* Reinhardt, Overs. Dansk. Forh. Kjøbenhavn, 1866, 56,
(*C. fasciatum* Reinhardt).702. *Characidium marshi* Breder.

Tuyra Basin, Pacific slope of Panama.

Characidium marshi Breder (C. M.), Amer. Mus. Novitates, CLXXX,
July 5, 1925, 5, fig. 5, Rio Sucubti, Darien, Panama.Genus 267. *APAREIODON* Eigenmann.*Apareiodon* Eigenmann, Ann. Carnegie Mus., X, 1916, 71 (*Parodon piracicabae* Eigenmann).703. *Apareiodon compressus* Breder.

Tuyra Basin, Pacific slope of Panama.

Apareiodon compressus C. M. Breder, Jr., Amer. Mus. Novitates,
CLXXX, July 5, 1925, 4, fig. 3, 4, Rio Chucunaque, Panama.704. *Apareiodon duriensis* (Meek and Hildebrand).

Rio Cupe, Panama.

Parodon duriensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool.,
X, 1913, 83, Pl. XVII, 1916, Rio Cupe, Pacific slope of Panama.

Order W. GYMNONOTI

Family 88. GYMNOTIDÆ

Genus 268. GYMNOTUS Linnaeus.

Gymnotus Linnaeus, Syst. Nat., Ed. X, 1758, 246 (*G. carapo* L.).
Carapus Rafinesque, Indice, 37, 1810, 57 (*Gymnotus acus* Linnaeus, prior to *Fierasfer* Cuvier, not *Carapus* Cuvier = *Giton* Kaup).
Carapus Müller and Troschel, Horæ Zool., III, 1849, 13 (*Gymnotus fasciatus* Pallas = *Gymnotus carapo* L.).

705. *Gymnotus carapo* Linnaeus.

Carapo Marcgrave, Hist. Pisc. Brasil., 1648, 170, Brazil.
Gymnotus carapo Linnaeus, Syst. Nat., Ed. X, 1758, 246.
Gymnotus fasciatus Pallas, Spic. Zool., VII, 1769, 35, fresh waters of America; anal rays about 180.
Gymnotus albus Pallas, loc. cit., 36, Surinam.
Gymnotus putaol Lacépède, Hist. Nat. Poiss., II, 1801, 176, Brasil.

Genus 269. APTERONOTUS Lacépède.

Apteronotus Lacépède, Hist. Nat. Poiss., II, 1800, 208 (*A. passan* Lacépède = *Gymnotus albifrons* L.).
Sternarchus Schneider, Syst. Ichth., Bloch, 1801, 497 (*Gymnotus albifrons* L.).
Memarchus Kaup in Duméril, Ichth. Anal., 1856, 201 (*Gymnotus albifrons* L.).

706. *Apteronotus rostratus* (Meek and Hildebrand).

Streams of Panama, in Atlantic drainage, and of Colombia (Rio Cauca).
Stenarchus rostratus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 85, Rio Grande near Cana, Panama.

Genus 270. STERNOPYGUS Müller and Troschel.

Sternopygus Müller and Troschel, Horæ Zool., III, 1849, 13 (*Gymnotus macrurus* Bloch and Schneider).
Carapus Cuvier, Règne Anim., Ed. I, 1817, 237 (*Gymnotus macrourus* Bloch and Schneider); not *Carapus* Rafinesque, 1810.
Carapo Oken, Isis, 1817, 1183; after Cuvier.
Giton Kaup, in Duméril, Ichth. Anal., 1856, 201 (*Gymnotus fasciatus* Pallas).

707. *Sternopygus dariensis* Meek and Hildebrand.

Streams of Panama, Atlantic slope.
Sternopygus dariensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 309, Rio Tuyra at Marrigante.

Genus 271. HYPOPOMUS Gill.

Hypopomus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 152 (*Rhamphichthys mulleri* Kaup).
Brachyrhamphichthys Günther, Cat., VIII, 1890, 6 (*Rhamphichthys artedi* Kaup).

708. *Hypopomus brevirostris* (Steindachner).

Panama, on both slopes to Brazil and Rio de la Plata.
Rhamphichthys brevirostris Steindachner, Sitzungsber. Akad. Wiss. Wien, LVIII, 1868, 254, fig. 2, Garapore, Amazon, at Santarem.
Hypopomus occidentalis Regan, Ann. Mag. Nat. Hist., XIV, 1914, (8), 32, Condoto River, Colombia.

Genus 272. EIGENMANNIA Jordan and Evermann.

Cryptops Eigenmann, Ann. N. Y. Acad. Sci., VII, 1894, 626 (*Sternopygus humboldti* Steindachner); name preoccupied.
Eigenmannia Jordan and Evermann, Fish. North and Mid. Amer., 1896, 341 (*Sternopygus humboldti* Steindachner = *Sternarchus virescens* Valenciennes).

709. *Eigenmannia virescens* (Valenciennes).

Brazil, north in Atlantic drainage to Panama.
Sternarchus virescens Valenciennes, D'Orbigny Voyage, V, 1847, 11.
Sternopygus lineatus Müller and Troschel, Schomburgk's Reise, III, 1848, 640, British Guiana.
Sternopygus tumifrons Müller and Troschel, Horæ Zool., III, 1849, 14.
Sternopygus humboldti Steindachner, Denkschr. Akad. Wiss. Wien, 1878, 55, Rio Magdalena.

Order X. EVENTOGNATHI. Carps

Family 89. CATOSTOMIDÆ. Suckers and buffalofishes

Genus 273. CYCLEPTUS Rafinesque. *Blackhorse.*

Cycleptus Rafinesque, Jour. Phys., LXXXVIII, 1819, 421 (*C. nigrescens* Rafinesque = *Catostomus elongatus* LeSueur).

Rhytidostomus Heckel, Fisch. Syriens, in Russegger's Reisen, 1842, 1023 (*Catostomus elongatus* LeSueur).

710. *Cycleptus elongatus* (LeSueur). *Blackhorse; Gourdseed sucker; Missouri sucker; Suckerel.*

Mississippi valley; rather common in large streams.

Catostomus elongatus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 103, Ohio River.

Cycleptus nigrescens Rafinesque, Jour. Phys., LXXXVIII, 1819, 421, Ohio River.

Genus 274. MEGASTOMATOBUS Fowler. *Big-mouthed buffalofish.*

Megastomatobus Fowler, Proc., Acad. Nat. Sci. Phila., LXV, 1913, 45 (*Sclerognathus cyprinella* Cuvier and Valenciennes, the chef de filo of these authors for *Sclerognathus* being *Sclerognathus cyprinus*).

Ichthyobus Agassiz, Amer. Jour. Sci. Arts, 1855, 192 (*I. bubalus* Agassiz; not *Catostomus bubalus* Rafinesque; revised spelling of *Ictiobus*).

711. *Megastomatobus cyprinella* (Cuvier and Valenciennes). *Redmouth buffalofish; Common buffalofish; Bigmouth buffalo; Gourdseed buffalo.* Mississippi Valley to Red River of the North; generally abundant in larger streams.

Sclerognathus cyprinella Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 355 (477), Lake Pontchartrain, La.

Genus 275. ICTIOBUS Rafinesque. *Buffalofishes.*

Ictiobus Rafinesque, Ichth. Ohiensis, 1820, 55 (*Amblodon bubalus* Rafinesque).

Bubalichthys Agassiz, Amer. Jour. Sci. Arts, XVIII, 1855, 192 (*Carpiodes urus* Agassiz).

Subgenus BUBALICHTHYS Agassiz.

712. *Ictiobus urus* (Agassiz). *Mongrel buffalo; Round buffalo; Black buffalo, Chopper.*

Mississippi Valley; less common than the others in the large streams; the largest and least common of the buffalo fish.

?*Carpiodes niger* Rafinesque, Ichth. Ohiensis, 1820, 56, Ohio River; may be *M. cyprinella*.

Carpiodes urus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 355, Tennessee River, Ala.

Bubalichthys urus Agassiz, loc. cit., Ohio River.

?*Bubalichthys niger* Agassiz, loc. cit., 193, Ohio River.

?*Bubalichthys bonusus* Agassiz, loc. cit., 195, Osage River, Mo.

Subgenus ICTIOBUS Rafinesque.

713. *Ictiobus bubalus* (Rafinesque). *Smallmouth buffalo; Razor-backed buffalo; Suckermouth buffalo; Quillback buffalo; Brown buffalo; Rooter.* Mississippi Valley and southward; generally abundant.

?*Amblodon bubalus* Rafinesque, Jour. Phys., LXXXVIII, 1819, 421, Ohio River.

?*Carpiodes taurus* Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 355, Mobile River.

?*Carpiodes vitulus* Agassiz, loc. cit., 356, Wabash River at New Harmony.

Ichthyobus rauchi Agassiz, loc. cit., XIX, 1855, 192.

Ichthyobus cyanellus Nelson, Bull. Ill. Lab. Nat. Hist., I, 1876, 49, Illinois River.

Bubalichthys altus Nelson, in Jordan, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 73, Cairo, Ill.

Bubalichthys bubalinus Jordan, Bull. U. S. Nat. Mus., IX, 1877, 50, Ohio River.

- Genus 276. CARPIOIDES Rafinesque.** *Carp suckers.*
- Carpoides* Rafinesque, Ichth. Ohiensis, 1820, 56 (*Catostomus cyprinus* LeSueur).
- Sclerognathus* Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 472 (*Catostomus cyprinus* and *Sclerognathus cyprinella* Cuvier and Valenciennes; restricted to the former by Agassiz, 1854, to the latter by Günther, 1868).
- 714. *Carpoides cyprinus* (LeSueur).** *Carp sucker; American carp; Carp; Quillback.*
Streams about Chesapeake Bay; common in the Chesapeake and Potomac, rare in the Delaware.
Catostomus cyprinus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 91, Elk River and other tributaries of Chesapeake Bay.
Carpoides vacca Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 356, Susquehanna River.
- 715. *Carpoides carpio* (Rafinesque).** *Carp sucker; Common river carp; Sucker.*
Ohio Valley and southwest to central Texas; generally abundant and variable.
Catostomus carpio Rafinesque, Ichth. Ohiensis, 1820, 56, falls of the Ohio River.
Carpoides nummifer Cope, Proc. Amer. Phil. Soc., XI, 1870, 484, Wabash River.
- 716. *Carpoides cutisanserinus* Cope.**
Allegheny Basin.
Carpoides cutisanserinus Cope, Proc., Amer. Phil. Soc. Phila., XI, 1870, 481, Kiskiminitas River, Pennsylvania (Allegheny Basin).
Carpoides selene Cope, loc. cit., 1870, 41; supposed to be from Root River, Mich.
- 717. *Carpoides difformis* Cope.** *Blunt-nosed river carp; Sucker.*
Ohio valley and westward; generally common.
Carpoides difformis Cope, Proc., Amer. Phil. Soc., XI, 1870, 480, Wabash River.
- 718. *Carpoides bison* Agassiz.**
Osage River.
Carpoides bison Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 356, Osage River, Mo.
- 719. *Carpoides thompsoni* Agassiz.** *Lake carp; Drum; Shad (Sandusky, Ohio).*
Great Lakes region from Lake Champlain to Lake of the Woods, chiefly northward.
Carpoides thompsoni Agassiz, Amer. Jour. Sci. Arts, XVIII, 1855, 191, Lake Champlain.
- 720. *Carpoides velifer* (Rafinesque).** *Silver carp; Quillback; Spearfish; Sailfish; Skimback; Sailing sucker; Sailorfish.*
Mississippi Valley and westward to the upper Missouri; more common northward.
Catostomus velifer Rafinesque, Ichth. Ohiensis, 1820, 56, Ohio.
Carpoides damalis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 170, Milk River, Mont.
Carpoides grayi Cope, Proc., Amer. Phil. Soc., XI, 1870, 482, Western States.
- 721. *Carpoides tumidus* Baird and Girard.** *Metalote; Buffalo.*
Lowland streams and lagoons, from the Rio Grande to the Rio Panuco and Tampico.
Carpoides tumidus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 28, Rio Grande at Fort Brown, Tex.; perhaps not separable from *C. velifer*.
- 722. *Carpoides labiosus* Meek.**
Streams of eastern Mexico.
Carpoides labiosus Meek, Field Col. Mus., Zool., V, 1904, 29, fig. 6, Valles, San Luis Potosi, Mexico.

- 723. *Carpiodes elongatus* Meek.**
 Streams of the east coast of Mexico.
Carpiodes elongatus Meek, Field Col. Mus., Zool., V, 1904, 28, fig. 5,
 Rio San Juan, Rio Conchos, and Rio Sarto Maruta in Tamaulipas,
 Linares, Nueva Leon.
- 724. *Carpiodes microstomus* Meek.**
 Tributaries of the Rio Grande, on the Mexican plateau.
Carpiodes microstomus Meek, Field Col. Mus., Zool., V, 1904, 27, fig.
 4, Rio Conchos in Chihuahua.
- 725. *Carpiodes meridionalis* (Günther).**
 Southern Mexico and Guatemala.
Sclerognathus meridionalis Günther, Cat., VII, 1868, 23, Rio Usumacinta, Guatemala.
- Genus 277. NOTOLEPIDOMYZON Fowler.**
Notolepidomyzon Fowler, Proc. Acad. Nat. Sci. Phila., LXV, 1913, 47
 (*Pantosteus arizoneæ* Gilbert = *Catostomus clarki* Baird and Girard).
- 726. *Notolepidomyzon plebeius* (Baird and Girard).**
 Basin of the Rio Grande, Colorado to Chihuahua; Rio Piedras, Rio Casas Grandes; very common.
Catostomus plebeius Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 28, Rio Mimbres, a tributary of Lake Guzman, Chihuahua, Mexico.
Catostomus (Acomus) guzmaniensis Girard, loc. cit., VIII, 1856, 173, Rio Janos, tributary to Lago de Guzman, Chihuahua, Mexico.
Catostomus nebuliferus Garman, Bull. Mus. Comp. Zool., VIII, 1881, 89, Rio Nazas, Coahuila, Mexico.
- 727. *Notolepidomyzon clarkii* (Baird and Girard).**
 Rio Gila, Salt River, and their tributaries in Arizona.
Catostomus clarkii Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 27, Rio Santa Cruz, Gila Basin, Ariz.
Pantosteus arizoneæ Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1896, 170, Salt River, Tempe, Ariz.
- 728. *Notolepidomyzon sanctæ-anæ* (Snyder).**
 Streams of southern California.
Pantosteus sanctæ-anæ Snyder, Proc. U. S. Nat. Mus., XXXIV, 1908, 33, Santa Ana River, at Riverside, Calif.
- 729. *Notolepidomyzon generosus* (Girard). Mountain sucker.**
 Colorado Basin.
Catostomus (Acomus) generosus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 174, Cottonwood Creek; not in Utah, "no doubt from Colorado" (Snyder).
- Genus 278. PANTOSTEUS Cope.** Mountain suckers.
Pantosteus Cope, Wheeler's Surv., V, 1875, 673 (*Minomus platyrhynchus* Cope).
- 730. *Pantosteus delphinus* (Cope).** Bluehead sucker.
 Mountain streams tributary to the Colorado River in western Colorado and Wyoming; very abundant.
Minomus delphinus Cope, Hayden's Geol. Surv. Wyo., 1870, 435; probably Henry Fork of Green River, Wyo.
?Minomus bardus Cope, loc. cit., 436; probably Green River, Wyo.
- 731. *Pantosteus platyrhynchus* (Cope).**
 Streams of Utah Basin (Provo and Sevier Rivers).
Minomus platyrhynchus Cope, Proc. Am. Phil. Soc., XIV, 1874, 134, Utah Lake at Provo.
?Minomus jarrovii Cope, loc. cit., 135; "near Provo"; the type lost.
- 732. *Pantosteus jordani* Evermann.**
 Upper Missouri and Columbia River Basins.
Pantosteus jordani Evermann, Bull. U. S. Fish Com., XII, 1892 (Jan. 27, 1893), 51, fig. on p. 53; Whitewood Creek and other streams of the Black Hills, S. Dak.

733. *Pantosteus columbianus* Eigenmann.
Columbia Basin.

Pantosteus columbianus Eigenmann, Amer. Nat., Feb. 4, 1893, 151,
Columbia River; perhaps identical with *Pantosteus jordani*.

734. *Pantosteus virescens* Cope.

Tributaries of Lake Bonneville basin of Utah; abundant in Weber and
Bear Rivers; largest of the hard-mouthed or mountain suckers.

Pantosteus virescens Cope, Wheeler's Surv., Zool., 1875, 675; by error
ascribed to Arkansas River at Pueblo; must have come from the
basin of Utah.

735. *Pantosteus lahontan* Rutter.

Streams of western Nevada; Lahontan Basin.

Pantosteus lahontan Rutter, Bull. U. S. Fish Com., XXII, 1902, 146,
with figure; Susan River, a tributary of Honey Lake, Little Truckee
River, Prosser Creek; misprint for *lahontan*.

Genus 279. *CATOSTOMUS* LeSueur. Fine-scaled sucker.

Catostomus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 89 (*Cyprinus*
catostomus Forster).

Decactylus Rafinesque, Ichth. Ohiensis, 1820, 60 (*Catostomus bostoniensis*
Rafinesque = *Cyprinus commersoni* Lacépède).

Stomocatus Bonaparte, Fauna Italica, 1840, Fasc. 129 (*Cyprinus catostomus* Forster).

Minomus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 173 (*Catostomus insignis* Baird and Girard).

Acomus Girard, loc. cit., 173 (*Catostomus forsterianus* Richardson =
Cyprinus catostomus Forster).

Subgenus DECACTYLUS Rafinesque.

**736. *Catostomus commersonii* (Lacépède). Common sucker; White sucker;
Brook sucker; Fine-scaled sucker; Black mullet (Sandusky, Ohio); Fine-scaled mullet; Bay mare.**

Quebec and the Great Lakes to Montana, Colorado, southward to Missouri and Georgia; the commonest of the suckers, excessively abundant from Massachusetts west to Kansas.

Cyprinus commersonii Lacépède, Hist. Nat. Poiss., V, 1803, 503, locality unknown; wrongly accredited to Commerson, the description and figure drawn from an old example long in alcohol.

Cyprinus teres Mitchell, Trans. Lit. Phil. Soc. N. Y., 1815, 458, New York.

Catostomus communis LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 95, 106, Delaware River.

Catostomus bostoniensis LeSueur, loc. cit., 106, Boston.

Catostomus flexuosus Rafinesque, Ichth. Ohiensis, 1820, 59, Ohio River.

Catostomus reticulatus Richardson, Fauna Bor.-Amer., Fishes, 1836, 303, Albany River.

Catostomus gracilis Kirtland, Rept. Zool. Ohio, 1838, 169, Cleveland, Ohio.

Catostomus pallidus DeKay, New York Fauna: Fishes, 1842, 200, New York.

Catostomus sucklili Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 175, Milk River, Mont.

Catostomus chloropteron Abbott, loc. cit., XII, 1860, 473, Kansas.

?*Catostomus alticolus* Cope, Proc. Amer. Phil. Soc., XIV, 1874, 138, Twin Lake, Colo.

Moxostoma trisignum Cope, Wheeler's Surv., V, Zool., 1875, 679, Arkansas River.

?*Catostomus utawana* Mather, Twelfth Rept., N. Y. Fish Com., Surv. of Adirondacks, 1884, 36, Blue Mountain Lakes, N. Y.; dwarf form.

737. *Catostomus ardens* Jordan and Gilbert. Mullet of Utah Lake.

Great basin of Utah and upper Snake River, Idaho; locally abundant.

Catostomus ardens Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 464, Utah Lake, Provo, Utah.

738. *Catostomus occidentalis* Ayres. *Sacramento sucker.*
 Streams of northern and central California and southern Oregon.
Catostomus occidentalis Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 18,
 San Francisco.
Catostomus labiatus Ayres, loc. cit., 1855, 32, San Joaquin River, Stock-
 ton, Calif.
Catostomus arzeopus Jordan, Bull. U. S. Nat. Mus., XII, 1878, 173,
 South Fork of Kern River, Calif.
739. *Catostomus tsiltcoosensis* Evermann and Meek.
 Coastal streams of Oregon.
Catostomus tsiltcoosensis Evermann and Meek, Bull. U. S. Fish Com.,
 XVII, 1897, 66, fig. 1, Tsilcoos Lake, Lane Co., Oreg.
740. *Catostomus snyderi* Gilbert.
 Klamath Basin, California and Oregon.
Catostomus snyderi Gilbert, Bull. U. S. Fish Com., XVII, 1897 (Jan. 6,
 1898), 3, Upper Klamath Lake, Oreg.
741. *Catostomus microps* Rutter.
 Streams of the lava beds of California.
Catostomus microps Rutter, Bull. U. S. Bur. Fish., XXVII, 1907 (1908),
 120, fig. 1; Rush Creek, a tributary of Ash Creek, Aden, Modoc Co.,
 Calif.
742. *Catostomus macrocheilus* Girard. *Columbia River sucker; Biglip sucker.*
 Columbia River Basin, as far up as Flathead Lake, Montana; Upper
 Salmon Falls and the headwaters of Salmon River, Idaho.
Catostomus macrocheilus Girard, Proc. Acad. Nat. Sci. Phila., VIII,
 1856, 175, Astoria, Oreg.
743. *Catostomus insignis* Baird and Girard.
 Gila Basin, Ariz.
Catostomus insignis Baird and Girard, Proc. Acad. Nat. Sci. Phila.,
 VII, 1854, 28, Rio San Pedro, Ariz.
Catostomus gila Kirsch, Proc. U. S. Nat. Mus., XI, 1888, 555, Rio Gila,
 Fort Thomas, Ariz.
744. *Catostomus bernardini* Girard.
 San Bernardino Creek, a tributary of Rio Yaqui, on the boundary of
 Arizona and Sonora.
Catostomus bernardini Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856,
 175, San Bernardino Creek, Ariz.
745. *Catostomus sonoriensis* Meek.
 Rivers of Sonora on the Pacific slope.
Catostomus sonoriensis Meek, Field Col. Mus., Zool., Ser. III, No. 6,
 May, 1902, 75, Pl. XVI, Rio Paphigochic, Miñaca, Chihuahua,
 Mexico.
- Subgenus **CATOSTOMUS** LeSueur.
746. *Catostomus catostomus* (Forster). *Long-nosed sucker; Northern sucker;*
Red sucker; Black sucker (Sandusky, Ohio); *Sturgeon sucker.*
 Great Lakes; upper Missouri River; Ohio Basin; West Virginia;
 upper Columbia; eastward to Maine and northwestward to Alaska;
 very abundant northward.
Cyprinus catostomus Forster, Philos. Trans., LXIII, 1773, 155, tab. 6,
 streams about Hudson Bay.
Catostomus longirostrum and *hudsonius* LeSueur, Jour. Acad. Nat. Sci.
 Phila., I, 1817, 102, 107, Vermont; after Forster.
Catostomus forsterianus Richardson, Franklin's Jour., 1823, 720, Lake
 Huron; Great Slave Lake.
Catostomus aurora Agassiz, Lake Superior, 1850, 360, Lake Superior.
Catostomus nanomyzon Mather, Twelfth Rep., N. Y. Fish Com., Surv.
 of Adirondacks, 1884, 36, Big Moose Lake, northern New York;
 dwarf form.
747. *Catostomus richardsoni* Harper and Nichols. *Gray sucker.*
 Streams of northwestern Canada.
Catostomus richardsoni Harper and Nichols, Bull., Amer. Mus. Nat.
 Hist., XLI, 1919, 263, Pl. XV, fig. 1; type locality not designated.

748. *Catostomus griseus* Girard.

Platte River and upper Missouri region.

Catostomus (Acomus) griseus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 174, Sweetwater Fork of Platte River.

?*Catostomus (Acomus) lactarius* Girard, loc. cit., 174, Milk River, Mont.

749. *Catostomus retropinnis* Jordan.

Northern Montana.

Catostomus retropinnis Jordan, Bull. U. S. Nat. Mus., XII, 1878, 178, Milk River, Mont.

750. *Catostomus pocatello* Gilbert and Evermann. *Moo-gad-ee of the Fort Hall Indians.*

Upper Snake River.

Catostomus pocatello Gilbert and Evermann, Bull., U. S. Fish Com., XIV, 1894, 189, pl. 21, fig. 1, Ross fork of Snake River, near Pocatello, Idaho.

751. *Catostomus tahoensis* Gill and Jordan. *Tahoe sucker.*

Lake Tahoe and Humboldt River, Nev.

Catostomus tahoensis Gill and Jordan, in Jordan, Bull., U. S. Nat. Mus., XII, 1878, 173, Lake Tahoe, Calif.

752. *Catostomus discobolus* Cope.

Upper Colorado Basin.

Catostomus discobolus Cope, Hayden's Geol. Surv. Wyo., 1870 (1872), 435, Green River, Wyo.

753. *Catostomus chamberlaini* (Rutter).

Chasmistes chamberlaini Rutter, Bull. U. S. Fish Com., XXII, 1902 (1904), 147, Eagle Lake (a lake without outlet in the Lahontan Basin); apparently a variant of *C. tahoensis*.

754. *Catostomus latipinnis* Baird and Girard. *Flannel-mouthed sucker.*

Rio Colorado and Rio Gila and their tributaries.

Catostomus latipinnis Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 388, Rio San Pedro, Gila Basin, Ariz.

755. *Catostomus conchos* Meek.

Catostomus conchos Meek, Field Col. Mus., Zool., III, 1902, 75, Rio Conchos, Jimenez, Chihuahua, Mexico.

Genus 280. *HYPENTELIUM* Rafinesque.

Hypentelium Rafinesque, Jour. Acad. Nat. Sci. Phila., II, 1818, 421 (*H. macroplectum* Rafinesque = *Catostomus nigricans* LeSueur).

Eurystomus Rafinesque, Ichth. Ohiensis, 1820, 59 (*Catostomus megastomus* Rafinesque), based on an incorrect drawing of *Hypentelium nigricans*. *Hylomyzon* Agassiz, Amer. Jour. Sci. Arts, 1855, 205 (*Catostomus nigricans* LeSueur).

756. *Hypentelium nigricans* (LeSueur). *Black sucker; Mud sucker; Hog sucker; Stoneroller; Totter; Crawl-a-bottom; Hammerhead; Stonelugger; Hogmolly; Bootleg sucker; Spotted sucker; Niggerhead.*

New York to Minnesota, Kansas, Arkansas, and the Carolinas; abundant in swift or rocky streams.

Catostomus nigricans LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 102, Lake Erie.

Catostomus maculosus LeSueur, loc. cit., 103, Pipe Creek, Md.

Hypentelium macropterum Rafinesque, loc. cit., I, 1817, 420, Ohio River.

Catostomus xanthopus Rafinesque, Ichth. Ohiensis, 1820, 57, Ohio River.

Catostomus nigricans etowanus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876 (1877), 345, Etowah River, Rome, Ga.

Catostomus megastomus Rafinesque, Ichth. Ohiensis, 1820, 59; based on incorrect drawing.

Catostomus planiceps Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 334 (450), 1844, Wabash River.

Genus 281. *THOBURNIA* Jordan and Snyder.

Thoburnia Jordan and Snyder, Copeia, October 4, 1917, 88 (*Catostomus rhothaeus* Thoburn).

757. *Thoburnia rhothœca* (Thoburn).

Headwaters of Tennessee River.

Catostomus rhothœcus Thoburn, in Jordan and Evermann, Fishes North and Mid. Amer., 1896, 181, upper tributaries of Tennessee River; supposed to be from the French Broad River at Wolf Creek, Tenn.

Genus 282. *CHASMISTES* Jordan.

Chasmistes Jordan, Bull., Hayden's Geol. Surv. Terr., IV, No. 2, 1878, 417 (*C. secundus* Jordan; not *Catostomus secundus* Cope and Yarrow = *Chasmistes litorus* Jordan).

Lipomyzon Cope, Amer. Nat., XV, 1881, 59 (*Chasmistes brevirostris* Cope).

Pithecomyzon Fowler, Proc. Acad. Nat. Sci. Phila., LXV, 1913, 54 (*Chasmistes cujus* Cope).

Subgenus LIPOMYZON Cope.

758. *Chasmistes fecundus* (Cope and Yarrow). *Webug sucker.*

Utah Lake.

Catostomus fecundus Cope and Yarrow, Wheeler's Surv., Zool., V, 1875, 678, Utah Lake.

759. *Chasmistes brevirostris* Cope.

Klamath Lake, Oreg.

Chasmistes brevirostris Cope, Amer. Nat., XIII, 1879, 785, Klamath Lake, Oreg.

760. *Chasmistes stomias* Gilbert.

Upper Klamath Lake, Oreg.

Chasmistes stomias Gilbert, Bull., U. S. Fish Com., XVII, 1897 (1898), 5, text figure, Upper Klamath Lake, Oreg.

761. *Chasmistes litorus* Jordan. *June sucker.*

Utah Lake.

Chasmistes litorus Jordan, Bull., U. S. Nat. Mus., XII, 1878, 249, Utah Lake at Provo, Utah.

762. *Chasmistes cujus* Cope. *Couia; Kweewee; Quiwee.*

Pyramid Lake, Nevada; locally abundant.

Chasmistes cujus Cope, Proc. Acad. Nat. Sci. Phila., XXXV, 1883, 149, Pyramid Lake, Nev.

763. *Chasmistes copei* Evermann and Meek. *Tswam.*

Klamath Lakes, Oreg.

Chasmistes copei Evermann and Meek, Bull., U. S. Fish Com., XVII, 1894 (1898), 70, fig. 3; Pelican Bay, Upper Klamath Lake, Oreg.

Genus 283. *DELTISTES* Seale.

Deltistes Seale, Proc. Calif. Acad. Sci., ser. 2, VI, 1896, 269 (*Chasmistes luxatus* Cope).

764. *Deltistes luxatus* (Cope). *Tswam; Lost River sucker.*

Klamath Lake and River, Oreg; abundant.

Chasmistes luxatus Cope, Amer. Nat., XIII, 1879, 784, Klamath Lake, Oreg.

Catostomus rex Eigenmann, loc. cit., XXV, July, 1891, 667, Lost River, Oreg.

Genus 284. *XYRAUCHEN* Eigenmann and Kirsch. *Razorback suckers.*

Kyrauchen Eigenmann and Kirsch, Proc. U. S. Nat. Mus., XI, 1888, 556 (*Catostomus cypho* Lockington = *Catostomus texanus* Abbott).

765. *Kyrauchen texanus* (Abbott). *Razorback sucker; Humpback sucker.*

Basin of the Colorado and Gila Rivers (not in Texas).

Catostomus texanus Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 473, Colorado and New Rivers, Ariz.

Catostomus cypho Lockington, loc. cit., 1880, 237, Colorado River at mouth of the Gila, Ariz.

Kyrauchen unciniphaga Jordan and Evermann, Bull. U. S. Fish Com., 1889, 26, Uncompahgre River delta, Colo.; young example, probably the same.

Genus 285. *ERIMYZON* Jordan. *Chub suckers.*

Erimyzon Jordan, Bull. Buffalo Soc. Nat. Hist., III, 1876, 95 (*Cyprinus oblongus* Mitchell).

- 766. *Erimyzon suetta* (Lacépède).** *Chub; Chub sucker; Creekfish; Creek sucker.*
 Coastwise streams, Pennsylvania and south; very abundant southward.
Cyprinus suetta Lacépède, Hist. Nat. Poiss., V, 1803, 606, South Carolina.
Moxostoma kenneryi Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 171, Dry Creek, Victoria, Tex.
Moxostoma campbelli Girard, loc. cit., 172, Live Oak Creek and Devil River, Tex.
Moxostoma tenue Agassiz, Amer. Jour. Sci. Arts, 1855, 203, Mobile, Ala.
Erimyzon goodei Jordan, Bull. U. S. Nat. Mus., XII, 1878, 148, St. Johns River, Fla.
- 767. *Erimyzon oblongus* (Mitchill).** *Chub sucker; Sweet sucker; Mullet; Creek fish; Pin minnow.*
 Great Lakes region to Maine and the Dakotas, south to Virginia and Oklahoma; very abundant; replacing the deeper-bodied *sucetta* northward; the two probably intergrading.
Cyprinus oblongus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 459, New York.
Catostomus gibbosus LeSueur, Connecticut River, Northampton; *tuberculatus*, Germantown, Pa.; and *vittatus*, Wissahickon River, Pa.; Jour. Acad. Nat. Sci. Phila., I, 1817, 92, 93, 104.
Catostomus fasciolaris Rafinesque, Ichth. Ohiensis, 1820, 58, Ohio River.
Labeo elegans, New York; *esopus*, New York; and *elongatus*, Mohawk River; DeKay, N. Y. Fauna: Fishes, 1842, 192, 195, 394.
Moxostoma clavigiformis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 171, Coal Creek, Canadian River, Indian Territory [Okla.]
- Genus 286. *MINYTREMA* Jordan.** *Spotted suckers.*
Minytrema Jordan, Man. Vert., ed. 2, 1878, 318 (*Catostomus melanops* Rafinesque).
- 768. *Minytrema melanops* (Rafinesque).** *Spotted sucker; Striped sucker; Winter sucker; Sand sucker; Black winter sucker.*
 Great Lakes region to North Carolina and west to Texas and Montana; rather common westward.
Catostomus melanops Rafinesque, Ichth. Ohiensis, 1820, 57, Ohio River.
Catostomus fasciatus LeSueur, in Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 333 (449), Mississippi River.
Moxostoma victorinum Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 171, Dry Creek, Victoria, Tex.
Ptychostomus hajdeni Girard, loc. cit., 172, Yellowstone River; Missouri River at Fort Pierre, Neb.
- Genus 287. *MOXOSTOMA* Rafinesque.** *Redhorse.*
Moxostoma Rafinesque, Ichth. Ohiensis, 1820, 54 (*Catostomus anisurus* Rafinesque).
Terelulus Rafinesque, loc. cit., 57 (*Catostomus aureolus* LeSueur).
Ptychostomus Agassiz, Amer. Jour. Sci. Arts., XIX, 1855, 203 (*Catostomus aureolus* LeSueur).
Myxostoma Jordan; corrected orthography.
- 769. *Moxostoma papillosum* (Cope).** *White mullet.*
 Lowland streams, from the Dismal Swamp in Virginia to the Ocmulgee River, Ga.
Ptychostomus papillosum Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 470, Catawba and Yadkin Rivers, N. C.
- 770. *Moxostoma anisurum* (Rafinesque).** *White-nosed sucker.*
 Ohio River, Great Lakes region and Lake of the Woods; not very common, but widely distributed.
Catostomus anisurus Rafinesque, Ichth. Ohiensis, 1820, 54, Ohio River.
Catostomus carpio Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 457, Lake Ontario, not of Rafinesque.
Ptychostomus velatus Cope, Proc. Amer. Phil. Soc., XI, 1870, 471, Youghiogheny River, Pa.
Morostoma valenciennei Jordan, Proc. U. S. Nat. Mus., VIII, 1885, 73; substitute for *Catostomus carpio* Cuvier and Valenciennes; preoccupied.

771. *Moxostoma buccó* (Cope).
 Missouri River at St. Joseph.
Ptychostomus buccó Cope, Hayden's Geol. Surv. Wyo., 1870, 437, St. Joseph, Mo.; a doubtful species allied to *M. anisurum*.
772. *Moxostoma pidiense* (Cope).
 Great Pedee Basin, the Carolinas.
Ptychostomus pidiensis Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 471, Yadkin River, N. C.
773. *Moxostoma collapsum* (Cope).
 Lowland streams of North Carolina.
Ptychostomus collapsus Cope, Proc. Amer. Phil. Soc., XI, 1870, 471, Neuse, Yadkin, and Catawba Rivers, N. C.
774. *Moxostoma coregonus* (Cope). *Blue mullet*.
 Catawba and Yadkin Rivers, N. C.
Ptychostomus coregonus Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 472, Catawba and Yadkin Rivers.
775. *Moxostoma album* (Cope).
 Rivers of North Carolina.
Ptychostomus albus Cope, Proc., Amer. Phil. Soc. Phila., XI, 1870, 472, Catawba River, N. C.
776. *Moxostoma thalassinum* (Cope).
 Yadkin River, N. C.
Ptychostomus thalassinus Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 472, Yadkin River, N. C.
777. *Moxostoma congestum* (Baird and Girard). *Texas redhorse*.
 Streams of Texas.
Catostomus congestus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 27, Rio Salado, Tex.
Ptychostomus albodus Girard, loc. cit., VIII, 1856, 172, Rio San Juan, Monterey, Nuevo Leon, Mexico.
778. *Moxostoma austrinum* (Bean).
 Rio de Santiago, Mexico; known from Michoacan, Mexico (west of Sierra Madre); Rio Lerna at Salamanca.
Moxostoma austrina Bean, Proc. U. S. Nat. Mus., II, 1879, 302, Piedad in Morelia, Michoacan, Mexico.
779. *Moxostoma mascotæ* Regan.
 Streams of Jalisco (Pacific drainage), Mexico.
Moxostoma mascotæ Regan, Biol. Centrali-Amer., 1907, 147, Tab. XXIV, fig. 2, Rio Mascota, Mexico.
780. *Moxostoma parvidens* Regan.
 Rio Grande Basin.
Moxostoma parvidens Regan, Biol. Centrali-Amer., 1907, 147, Tab. XXIV, fig. 3, Rio Grande at Juarez, Mexico.
781. *Moxostoma aureolum* (LeSueur). *Common redhorse*; *Mullet*; *White sucker*; *Large-scaled sucker*; *Sucker*.
 Lake Ontario and Lake Michigan to the Missouri River and Lake of the Woods, south to Arkansas and Georgia; very abundant west of Alleghenies.
Catostomus aureolus LeSueur, Jour., Acad. Nat. Sci. Phila., I, 1817, 95, Lake Erie near Buffalo.
Catostomus duquesnii LeSueur, loc. cit., I, 105, 1817, Ohio River at Pittsburgh, Pa.
Catostomus erythrus Rafinesque, Amer. Month. Mag., III, 1818, 354, 355, Ohio River.
Catostomus oneida DeKay, N. Y. Fauna: Fishes, 1842, 198, Oneida Lake.
Rutilus melanurus Rafinesque, Ichth. Ohiensis, 1820, 51, Ohio River.
Myxostoma euryops Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 348, Oostanaula River, Rome, Ga.; probably a monstrosity of this species.

782. *Moxostoma lesueuri* (Richardson). *Piconou.*
 Tributaries of Hudson Bay and westward.
Catostomus lesueuri Richardson, Franklin's Jour., 1823, 1772, Pine
 Island Lake, British America.
783. *Moxostoma macrolepidotum* (LeSueur). *Mullet; Red mullet; Eastern
 redhorse; River sucker.*
 Streams about Chesapeake and Delaware Bays and southward to
 North Carolina; east of the Alleghenies.
Catostomus macrolepidotus LeSueur, Jour. Acad. Nat. Sci. Phila., I,
 1817, 94, Delaware River.
Ptychostomus lachrymatus Cope, Proc., Amer. Phil. Soc., XI, 1870, 474,
 Neuse River at Newbern, N. C.
784. *Moxostoma robustum* (Cope).
 Yadkin River, N. C.
Ptychostomus robustus Cope, Proc. Amer. Phil. Soc., XI, 1870, 473,
 Yadkin River, N. C.
785. *Moxostoma crassilabre* (Cope).
 Streams of eastern North Carolina.
Ptychostomus crassilabris Cope, Proc. Amer. Phil. Soc., XI, 1870, 477,
 Neuse River near Raleigh, N. C.
786. *Moxostoma conus* (Cope).
 Neuse and Yadkin Rivers, N. C.
Ptychostomus conus Cope, Proc. Amer. Phil. Soc., XI, 1870, 478, Yadkin
 River, N. C.
787. *Moxostoma breviceps* (Cope). *Short-headed redhorse.*
 Ohio Valley and Great Lakes region; abundant in Lake Erie.
Ptychostomus breviceps Cope, Proc. Amer. Phil. Soc., XI, 1870, 478,
 Youghiogheny River, Pa.
788. *Moxostoma pectilurum* (Jordan).
 Southern Mississippi to eastern Texas; Pearl, Tangipahoa, and Sabine
 Rivers.
Myxostoma pectilura Jordan, Bull. U. S. Nat. Mus., X, 1877, 66, Tangi-
 pahoa River, La.
789. *Moxostoma alleghaniensis* Nichols.
 Mountain streams of North Carolina.
Moxostoma alleghaniensis Nichols, Bull. Amer. Mus. Nat. Hist.,
 XXX, 1911, 275, Pl. XI, fig. 1, Marshall, N. C.
- Genus 288. **SCARTOMYZON** Fowler.
Scartomyzon Fowler, Proc. Acad. Nat. Sci. Phila., LXV, 1913, 59
 (*Ptychostomus cervinus* Cope).
790. *Scartomyzon rupiscartes* (Jordan and Jenkins). *Jumprocks.*
 Rivers of the Carolinas and Georgia, from the Catawba to the Chatta-
 hoochee; locally abundant.
Moxostoma rupiscartes Jordan and Jenkins, Proc. U. S. Nat. Mus.,
 XI, 1888, 353, Catawba River at Morgantown, and various other
 streams, representing *Scartomyzon cervinus* southward.
791. *Scartomyzon cervinus* (Cope). *Jumping mullet; Crawl-a-bottom.*
 Rivers of the South Atlantic States, from the James to the Neuse.
Teretulus cervinus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 236,
 headwaters of the Roanoke and James Rivers, Va.
- Genus 289. **PLACOPHARYNX** Cope.
Placopharynx Cope, Proc. Amer. Phil. Soc., XI, 1870, 467 (*P. carinatus*
 Cope).
792. *Placopharynx carinatus* Cope. *White sucker; Big-toothed redhorse.*
 Michigan to Tennessee, Georgia, and Arkansas; abundant in large
 streams, especially in the Ozarks.
Placopharynx carinatus Cope, Proc. Amer. Phil. Soc., XI, 1870, 467,
 Wabash River.

Genus 290. LAGOCHILA Jordan and Brayton. *Rabbitmouth suckers.*
Lagochila Jordan and Brayton, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 280, (*L. lacera* Jordan and Brayton); not *Lagocheilus*, an earlier genus of mollusks.
Quassialabia Jordan and Brayton, Man. Vert., Ed. II, 1878, 406 (*Lagochila lacera* Jordan and Brayton); substitute for *Lagochila*, regarded as preoccupied.

793. Lagochila lacera Jordan and Brayton. *Harelip sucker; Cut-lips; Split-mouth sucker; May sucker; Rabbitmouth sucker; Pealip sucker.*
 Mississippi Valley; Ozark Mountains; Scioto, Tippecanoe, Clinch, Cumberland, and Chickamauga Rivers, and the White River of Arkansas; nowhere very common.
Lagochila lacera Jordan and Brayton, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 280, Chickamauga River, Ringgold, Ga.

Family 90. CYPRINIDÆ. Daces and minnows

In arranging the genera of this by far the largest natural family of fishes, we have begun the series with the forms having closest affinity with Asiatic and then with European types—these having largely five teeth in the main row on the pharyngeals. Later follow the characteristically American forms, more or less degenerate as compared with the possible European ancestors, the series closing with aberrant forms from the same stock.

Genus 291. ORTHODON Girard.
Orthodon Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 182 (*Gila microlepidota* Ayres).

794. Orthodon microlepidotus (Ayres). *Blackfish.*
 Larger rivers and lakes of central and northern California; generally abundant.
Gila microlepidota Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 21, Sacramento and San Joaquin Rivers, Calif.

Genus 292. ACROCHEILUS Agassiz. *Chiselmouths.*
Acrocheilus Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 96 (*A. alutaceus* Agassiz and Pickering).

795. Acrocheilus alutaceus Agassiz and Pickering. *Chiselmouth; Square-mouth; Hardmouth.*
 Lakes and rivers of southeastern Oregon; lower Columbia and tributaries as far up as Spokane and Shoshone Falls.
Acrocheilus alutaceus Agassiz and Pickering, Amer. Jour. Sci. Arts, XIX, 1855, 99, Willamette Falls and Walla Walla River, Oreg.

Genus 293. MYLOCHEILUS Agassiz.
Mylocheilus Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 229 (*M. lateralis* Agassiz and Pickering).

796. Mylocheilus lateralis Agassiz and Pickering. *Columbia River chub.*
 Columbia and Frazer River Basins; ascending to Flathead Lake and Shoshone Falls.
Mylocheilus lateralis Agassiz and Pickering, Amer. Jour. Sci. Arts, 1855, 231, Columbia River at Fort Vancouver; confused by recent writers with *Leuciscus cairinus* Richardson, which is a near relative of *Richardsonius balteatus*.

797. Mylocheilus fraterculus Girard.
Mylocheilus fraterculus Girard, Proc., Acad. Nat. Sci. Phila., VIII, 1856, 169, "Monterey, Calif"; a doubtful record, as no *Mylocheilus* has been yet found in California; scales about 85; 77 in *Mylocheilus lateralis*.

Genus 294. MYLOPHARODON Ayres.
Mylopharodon Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 33 (*M. robustus* Ayres = *Mylopharodon conocephalus* Girard).

798. *Mylopharodon conocephalus* (Baird and Girard). *Kaweah chub.*
 Sacramento and San Joaquin Rivers, Calif.; southeastern Oregon, Pit River and Big Valley.
Gila conocephala Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 135, San Joaquin River, Calif.
Mylopharodon robustus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 33, San Joaquin River, Calif.
- Genus 295. **STYPODON** Garman.
Styptodon Garman, Bull. Mus. Comp. Zool., VIII, 1881, 90 (*S. signifer* Garman).
799. **Styptodon signifer** Garman.
 Lago de Parras, an isolated lagoon in Coahuila, Mexico.
Styptodon signifer Garman, Bull. Mus. Comp. Zool., VIII, 1881, 90, Lago de Parras, Coahuila, Mexico.
- Genus 296. **CHROSOMUS** Rafinesque.
Chrosomus Rafinesque, Ichth. Ohiensis, 1820, 47 (*Luxilus erythrogaster* Rafinesque).
800. **Chrosomus erythrogaster** Rafinesque. *Red-bellied dace.*
 New Brunswick to Ohio, Wisconsin, Colorado, and northern Alabama; St. Lawrence River; chiefly or entirely west of the Alleghenies; very abundant northward.
Luxilus or *Chrosomus erythrogaster* Rafinesque, Ichth. Ohiensis, 1820, 47, Ohio River.
Chrosomus pyrrhogaster Jordan, Man. Vert., Ed. I, 1876, 284, White River, Ind.
801. **Chrosomus eos** Cope.
 Streams east of the Alleghenies, north to the Susquehanna; scarce.
Chrosomus eos Cope, Proc., Acad. Nat. Sci. Phila., XIII, 1861, 523, Meshoppen Creek, Susquehanna River, Pa.; perhaps a variation of *Chrosomus erythrogaster*.
802. **Chrosomus oreas** Cope.
 Headwaters of Roanoke River, and clear tributaries of the Tennessee River.
Chrosomus oreas Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 233, head of Roanoke River, Montgomery County, Va.
803. **Chrosomus dakotensis** Evermann and Cox.
Chrosomus dakotensis Evermann and Cox, Rept. U. S. Fish Com., XIX, 1895, Crow Creek, Chamberlain, S. Dak.
- Genus 297. **LAVINIA** Girard.
Lavinia Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 137 (*L. exilicauda* Baird and Girard).
Lucilinus Jordan, Cat. Fishes N. Amer., 1885, 33 (*Laucosomus occidentalis* Baird and Girard); young.
804. **Lavinia exilicauda** Baird and Girard. *Hitch; Chigh.*
 Streams of the coast ranges about San Francisco and Monterey; locally common as far north as Clear Lake, Calif.
Lavinia exilicauda Baird and Girard, Proc., Acad. Nat. Sci. Phila., VII, 1854, 137, Sacramento River, Calif.
Leucosomus occidentalis Baird and Girard, loc. cit., VII, 1854, 137, Poso Creek; Four Creeks, Calif.; young of *L. exilicauda*.
Lavinia harengus Girard, loc. cit., VIII, 1856, 184, Monterey, Calif.
- Genus 298. **OXYGENEUM** Forbes.
Oxygeneum Forbes, Bull. Ill. Lab. Nat. Hist., II, 1885, 136 (*O. pulverulentum* Forbes).
805. **Oxygeneum pulverulentum** Forbes.
 Illinois River, Ill.
Oxygeneum pulverulentum Forbes, Bull. Ill. Lab. Nat. Hist., II, 1885, 136, Illinois River, Ill.
- Genus 299. **PTYCHOCEILUS** Agassiz. *Squawfish.*
Ptychocheilus Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 227 (*P. gracilis* Agassiz = *Cyprinus* (*Leuctrus*) *oregonensis* Richardson).

- 806. *Ptychocheilus oregonensis* (Richardson).** *Squawfish; Chappau; White-fish.*
 Rivers of Oregon and Washington, in the Columbia Basin to Montana and Idaho, and northward to British Columbia.
Cyprinus (Leuciscus) oregonensis Richardson, Fauna Bor.-Amer., III, 1836, 305, Columbia River.
Ptychocheilus gracilis Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 229, Willamette River.
- 807. *Ptychocheilus grandis* (Ayres).** *Sacramento pike; Sacramento squawfish.*
 Rivers of California and southeastern Oregon; Pit River.
Gila grandis Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 18, San Francisco.
Ptychocheilus major Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 229, San Francisco.
Ptychocheilus harfordi Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 72, Sacramento River; scales 90 to 95, a variant with smaller scales.
- 808. *Ptychocheilus lucius* Girard.** *White salmon of the Colorado; Colorado River squawfish.*
 Colorado River Basin, in all larger streams from Rio Uncompahgre to the Gulf of California; largest of American *Cyprinidae*, reaching 80 pounds.
Ptychocheilus lucius Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 209, Rio Colorado.
- 809. *Ptychocheilus umpqua* Snyder.**
 Coastal streams of Oregon.
Ptychocheilus umpqua Snyder, Bull. U. S. Bur. Fish., XXVII, 1907 (1908), 170, fig. 2, in part, Callopoia Creek, Oakland, Oreg.
- Genus 300. *GILA* Baird and Girard.**
Gila Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 368 (*G. robusta* Baird and Girard)..
- 810. *Gila elegans* Baird and Girard.** *Bony-tail; Gila trout; Verde trout.*
 Colorado and Gila Rivers; abundant in river channels.
Gila elegans Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 369, Zufi, Colorado, and Gila Rivers.
Gila emoryi Baird and Girard, loc. cit., 388, Gila River.
- 811. *Gila robusta* Baird and Girard.** *Roundtail.*
 Tributaries of the Rio Colorado and Rio Gila; very common up to the foot of the mountains in Colorado.
Gila robusta Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 368, Zufi River.
Gila gracilis Baird and Girard, loc. cit., 369, Zufi River.
Gila grahami Baird and Girard, loc. cit., 389, Rio San Pedro, tributary to Rio Gila.
Ptychocheilus vorax Girard, loc. cit., 1856, 209, locality unknown.
Leuciscus zunnensis Günther, Cat., VII, 1868, 241, Zuñi River; substitute for *gracilis*, preoccupied in *Leuciscus*.
Gila affinis Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 474, probably erroneously ascribed to Kansas River.
Gila naevia Cope, Hayden's Geol. Surv. Wyo., 1870 (1871), 441, tributary of Green River, Wyo.; young, perhaps of *G. elegans*.
- 812. *Gila seminuda* Cope and Yarrow.**
 Rio Virgen, in Utah and Nevada.
Gila seminuda Cope and Yarrow, Wheeler's Surv., Zool., V, 1875, 666, Rio Virgen, Nev.
- 813. *Gila minacea* Meek.**
 Streams of Chihuahua, Mexico.
Gila minacea Meek, Field Col. Mus., Zool., III, May, 1902, 80, Rio Paphigochic, Mifiaca, Chihuahua, Mex.
- Genus 301. *NOTEMIGONUS* Rafinesque.**
Notemigonous Rafinesque, Jour. Phys., LXXXVIII, 1819, 420, (*N. auratus* Rafinesque = *Cyprinus crysoleucus* Mitchell).
Hemiplus Rafinesque, Anal. Nat., I, 1820, 6 (*H. lacustris* Rafinesque).

- 814. *Notemigonus crysoleucas* (Mitchill).** *Roach; Sunfish; Dace; Bitter-head; Shiner; Chub; Bream; Gudgeon; Young shad; Windfish.*
 Nova Scotia and Maryland to the Dakotas.
Cyprinus crysoleucas Mitchill, Rept., Fish. N. Y., 1814, 23, New York.
Cyprinus hemiplus Rafinesque, Amer. Month. Mag., II, 1817, 121,
 Lake George, Saratoga, N. Y.
Notemigonus auratus Rafinesque, Jour. Phys., LXXXVIII, 1819, 421,
 Ohio River.
Hemiplus lacustris Rafinesque, Anal. Nat., I, 1820, 6, Lake George.
Abramis versicolor DeKay, N. Y. Fauna: Fishes, 1842, 191, Connecticut
 and Hudson Rivers; Pecskill.
- 815. *Notemigonus boscii* (Cuvier and Valenciennes).**
 Rivers of the South Atlantic States; abundant from the James to the
 Altamaha.
Leuciscus boscii Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844,
 232 (313), Carolina, Pennsylvania, and New York.
Cyprinus americanus Linneaus, Syst. Nat., ed. XII, 1766, 530, Charles-
 town; name preoccupied.
Notemigonus ischanus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876,
 364, South Fork Ocmulgee River, Flat Shoals, Ga.
Luxilus leptosomus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 203,
 Dry Creek, Victoria, Tex.
Luxilus seco Girard, loc. cit., 203, Rio Seco, tributary of Rio Nueces, Tex.
- 816. *Notemigonus gardoneus* (Cuvier and Valenciennes).**
 South Carolina.
Leuciscus gardoneus Cuvier and Valenciennes, Hist. Nat. Poiss., XVII,
 1844, 235 (316), Charleston, S. C.; perhaps a hybrid.
- Genus 302. OPSOPŒODUS Hay.**
Opsopœodus Hay, Proc. U. S. Nat. Mus., III, 1880, 507 (*O. emiliae* Hay).
Trychærodon Forbes, in Jordan and Gilbert, Bull. U. S. Nat. Mus., XVI,
 1882, 247 (*T. megalops* Forbes).
- 817. *Opsopœodus emiliae* Hay.**
 Lake Erie and southern Indiana and Illinois to Georgia and Mississippi.
Opsopœodus emiliae Hay, Proc. U. S. Nat. Mus., III, 1880, 507, Artesia,
 Macon, and Enterprise, Miss.
- 818. *Opsopœodus osculus* Evermann.**
 Rio Neches and other streams about Palestine and Houston, Tex.
Opsopœodus osculus Evermann, Bull. U. S. Fish Com., XI, 1891 (1892),
 82, Neches River near Palestine, Tex.
- 819. *Opsopœodus borealis* Harper and Nichols.** *Athabaska minnow.*
 Streams of Alberta.
Opsopœodus borealis Harper and Nichols, Bull. Amer. Mus. Nat. Hist.,
 XLI, 266, Lake Athabaska at Fort Chipewyan, Alberta, Canada.
- 820. *Opsopœodus megalops* (Forbes).**
 Western Ohio to Illinois.
Trychærodon megalops Forbes, in Jordan and Gilbert, Synopsis, 1883,
 247, Illinois River at Pekin and Peoria, Ill.
- Genus 303. OPSOPŒA Jordan and Evermann.**
Opsopœa Jordan and Evermann, Fishes North and Mid. Amer., 1896,
 247 (*Opsopœodus bollmani* Gilbert).
- 821. *Opsopœa bollmani* (Gilbert).**
 Lowland swamps of the Southern States; known from Satilla River,
 Waycross, Ga., and Obion River, Cypress, Tenn.
Opsopœodus bollmani Gilbert, Bull. U. S. Fish Com., VIII, 1888 (1890),
 226, Buckland Creek, Millen, Ga.
- Genus 304. ALGANSEA Girard.**
Algansea Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 182 (*Leuciscus*
tincella Cuvier and Valenciennes).
- 822. *Algansea tincella* (Cuvier and Valenciennes).**
 Lakes about the City of Mexico and in streams of central Mexico.
Leuciscus tincella Cuvier and Valenciennes, Hist. Nat. Poiss., XVII,
 1844, 240 (323), City of Mexico.

823. *Algansea dugesi* Bean.

Central Mexico, in streams tributary to Rio de Lerma, Lake Yuriria in Guanajuato.

Algansea dugesi Bean, Proc. U. S. Nat. Mus., XV, 1892, 283, Lake Yuriria, Guanajuato, Mex.

?*Algansea tarascorum* Steindachner, Einige Fischarten Mexicos, 1895, 11, pl. 3, figs. 2-2c, Lake Patzcuaro, Mex.

824. *Algansea lacustris* Steindachner.

Lake Patzcuaro, Jalisco, Mexico.

Algansea lacustris Steindachner, Einige Fischarten Mexicos, 1895, 10, pl. 3, figs. 1 and 1b, Lake Patzcuaro.

825. *Algansea sallaei* (Günther).

Streams of central Mexico.

Ceratichthys sallaei Günther, Cat., VII, 1868, 484, Cuernavaca, Mex.

Zophendum australe Jordan, Proc. U. S. Nat. Mus., II, 1879, 300, Lake Tupataro, Mex., Rio Lerma.

826. *Algansea rubescens* Meek.

Lake Patzcuaro, Michoacan.

Algansea rubescens Meek, Field Col. Mus., Zool., V, 1905, 46, fig. 9, Lake Patzcuaro.

827. *Algansea stigmatura* Regan.

Northern Mexico.

Algansea stigmatura Regan, Biol. Centrali-Amer., 1907, 152, Tab. XXV, fig. 2, Rio Grande de Santiago, Jalisco.

828. *Algansea affinis* Regan

Upper tributaries of Rio Lerma and Rio Panuco, Mexico.

Algansea affinis Regan, Biol. Centrali-Amer., 1907, 151, Tab. XXV, fig. 1, Aguas Calientes, Salamanca, and San Juan del Rio, Mexico.

Genus 305. *POGONICHTHYS* Girard.

Pogonichthys Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 136 (*P. inaequilobus* Baird and Girard = *Leuciscus macrolepidotus* Ayres).

Symmetrurus Jordan, Hayden's Geol. Surv. Terr., IV, 1878, 788, (*Pogonichthys argyreiosus* Girard); young.

829. *Pogonichthys macrolepidotus* (Ayres). *Splittail*.

Streams of central California; very common in the Sacramento and San Joaquin Rivers.

Leuciscus macrolepidotus Ayres, Proc. Calif. Acad. Nat. Sci., I, in Placer Times and Transcript, May 30, 1854, San Francisco, Calif.

Pogonichthys inaequilobus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 136, San Joaquin Valley, Calif.

Pogonichthys argyreiosus Girard, loc. cit., 153, Presidio, Calif.

Genus 306. *MARGARISCUS* Cockerell.

Margariscus Cockerell, Proc. Biol. Soc. Wash., XXII, 1909, 217 (*Clinostomus margarita* Cope).

As a small barbel is normally present in this genus and placed above the tip of the maxillary, as in *Semotilus*, the genus must be placed near *Leucosomus*.

830. *Margariscus margarita* (Cope). *Minnow*; *Goldthread shiner*; *Pearl minnow*.

Lake Erie, south to James River and headwaters of the Kanawha.

Clinostomus margarita Cope, Trans. Amer. Phil. Soc., XIII, 1866, 377, Conestoga River, Lancaster, Pa.

831. *Margariscus nachtriebi*²⁵ (Cox).

North Dakota to New Brunswick.

Leuciscus nachtriebi U. O. Cox, Rept. U. S. Fish Com., XX, 1894 (1896), 815, Minnesota, Mille Lacs Lake.

Couesius plumbeus rubrilateralis Philip Cox, Bull. Nat. Hist., 2, 1901, 42, Miramichi, Canada.

Leuciscus carletoni Kendall, Bull. U. S. Fish Com., XXII, 1902, 357, fig. 1, Penobscot River, Me.

²⁵"With much doubt we leave this species (*M. nachtriebi*) with *Margariscus*. The actual relation of this and other related forms is yet to be determined." (Jordan.)

Genus 307. LEUCOSOMUS Heckel. *Fallfishes.*

Leucosomus Heckel, Russegger's Reisen, 1, 1843, 1042 (*L. chryssoleucus* Heckel = *Cyprinus corporalis* Mitchell); not *Leucosoma* Gray, a genus of Salangidae.

Cheilonemus Baird, in Storer, Hist. Fish. Mass., 1855, 285 (*Leuciscus pulchellus* Storer = *Cyprinus corporalis* Mitchell).

832. *Leucosomus corporalis* (Mitchill). *Chub; Dace; Fallfish; Silver chub; Windfish; Corporal; Chivin.*

Eastern United States, from the St. Lawrence to James River, Virginia; largest of our eastern Cyprinidae; common northward, east of the Alleghenies.

Cyprinus corporalis Mitchell, Amer. Month. Mag., I, 1817, 289, Wallkill River, N. Y.

Cyprinus bullaris Rafinesque, loc. cit., II, Dec., 1817, 120, Hudson River.

Leuciscus argenteus Storer, Hist. Fish. Mass., 1839, 90, Worcester Co., Mass.

Leuciscus pulchellus Storer, loc. cit., 90, Walpole, Mass.

Leuciscus nitidus DeKay, N. Y. Fauna: Fishes, 1842, 209, Lake Champlain.

Leuciscus chrysopterus DeKay, loc. cit., 211, N. Y. Harbor.

Leuciscus rotengulus Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 236 (318), Carolinas.

Leucosomus cataractus Baird, Iconog. Encyl., II, 216, and in Cope, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 523, Susquehanna River.

Leucosomus rhotheus Cope, loc. cit., XIII, 1861, 523, Brandywine Creek and Rancocas Creek, Pa.

Squalius hyalope Cope, loc. cit., XVI, 1864, 280, Conestoga River, Pa.

Genus 308. SEMOTILUS Rafinesque. *Horned dace.*

Semotilus Rafinesque, Ichth. Ohiensis, 1820, 49 (*S. dorsalis* Rafinesque = *Cyprinus atromaculatus* Mitchell).

833. *Semotilus atromaculatus* (Mitchill). *Horned dace; Creek chub; Chub; Silver chub; Mud chub.*

Maine to New Jersey; westward to Kansas and Wyoming.

Cyprinus atromaculatus Mitchell, Amer. Month. Mag., II, 1818, 324, Wallkill River, N. Y.

Semotilus dorsalis Rafinesque, Ichth. Ohiensis, 1820, 49, Kentucky River.

Semotilus cephalus Rafinesque, loc. cit., 49, Ohio River.

Semotilus diploemia Rafinesque, loc. cit., 50, Ohio River.

Leuciscus iris Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 190, (255), New York and Carolina.

Leuciscus storeri Cuvier and Valenciennes, loc. cit., 237 (255), New York.

Leuciscus pulchelloides Ayres, Proc. Bost. Soc. Nat. Hist., III, 1849, 157, Connecticut.

Leucosomus pallidus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 190, Antelope Creek, Ark.

Leucosomus incrassatus Girard, loc. cit., 1856, 190, Choctaw Agency.

Semotilus macrocephalus Girard, loc. cit., 1856, 204, Fort Pierre, Nebr.

Semotilus speciosus Girard, loc. cit., 1856, 204, Sweetwater River, Nebr.

Semotilus hammondii Abbott, loc. cit., XII, 1860, 474, Kansas.

834. *Semotilus atromaculatus thoreauanus* (Jordan).

Flint River, Ga.

Semotilus thoreauanus Jordan, Bull., U. S. Nat. Mus., X, 1877, 63, Flint River, Ga.

Genus 309. RICHARDSONIUS Girard.

Richardsonius Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 201 (*Cyprinus (Abramis) baleatus* Richardson).

* The various American allies of *Leuciscus* Cuvier (type *Cyprinus leuciscus* L. of Europe, *Squalius*), we now divide among several genera, these corresponding to the subgenera in the previous check list. No species belonging to *Leuciscus*, as strictly defined, occurs in America or Asia. Many of the species, both in the groups allied to *Leuciscus* and *Rutilus* as well as in those related to *Hybopsis* and *Notropis*, need comparison and verification.

Richardsonius is here restricted to the single species, *baleatus*, distinguished, with its variant, by the very long anal fin. The genus *Cheonda* is provisionally recognized for species more or less similar to *Richardsonius* but with the anal fin shorter. This group seems to merge into *Tymora*, species with the anal fin short and the scales rather loosely imbricated. One species, *cavifrons*, forms another intermediate type with short anal and very small scales. Any arrangement of these fishes must be still wholly provisional.

- 835. *Richardsonius siuslawi* (Evermann and Meek).**
 Streams of western Oregon.
Leuciscus siuslawi Evermann and Meek, Bull. U. S. Fish Com., XVII, 1897 (1898), 72, Siuslaw River, Mapleton, Oreg.
- 836. *Richardsonius lateralis* Girard.**
 Streams of Washington and Montana, a doubtful species, nominally distinguished by the smaller number of anal rays.
Richardsonius lateralis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 202, Fort Steilacoom, Wash.
Leuciscus gilli Evermann, Bull., U. S. Fish Com., XI, 1891, 44, Browns Gulch Creek, Silver Bow, Mont.
- 837. *Richardsonius balteatus* (Richardson).** *Red-sided bream.*
 Columbia and Fraser Rivers, and streams about Puget Sound.
Cyprinus (Abramis) balteatus Richardson, Fauna Bor.-Amer., III, 1836, 301, Columbia River.
- 838. *Richardsonius thermophilus* Evermann and Cockerell.**
 Warm Springs, Silver Creek, and Silvies River, Oreg.
Richardsonius thermophilus Evermann and Cockerell, Proc. Biol. Soc. Wash., XXII, December 8, 1909, 185, Warm Springs, Harney Co., Oreg.
- Genus 310. CLARKINA Jordan and Evermann.**
Cyprinus (Leuciscus) caurinus Richardson is an ally of the genus *Richardsonius*, differing in the short anal fin of 9 rays only. The body is elongate and the scales (86) very small, well imbricated, differing from the type of *Tigoma*.
Clarkina Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, 1927 (April 27), 502 (*Cyprinus caurinus* Richardson).
- 839. *Clarkina caurina* (Richardson).**
Cyprinus (Leuciscus) caurinus Richardson, Fauna Bor.-Amer., III, 1836, 304, Columbia River at Fort Vancouver.
Leuciscus caurinus Snyder, Bull. U. S. Fish Com., XXVII, 1907, 173; not *Mylocheilus caurinus* Agassiz; Willamette River near Corvallis, Oreg.
- Genus 311. CHEONDA Girard.**
Cheonda Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 207 (*C. cooperi* Girard); scales moderate; anal rays 8 to 11, usually 9.
- 840. *Cheonda cooperi* Girard.**
 Lower Columbia River Basin.
Cheonda cooperi Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 207, Columbia River, Fort Vancouver, Wash.
- 841. *Cheonda microdon* (Snyder).**
Richardsonius microdon Snyder, Bull. U. S. Bur. Fish., XXXV, 1915-16 (1918), 58; scales 55; anal rays 8 or 9.
- 842. *Cheonda egregia* (Girard).**
 Nevada to northern California; known from the Humboldt, Truckee, and Sacramento-San Joaquin Basins and Napa Valley, Calif.
Tigoma egregia Girard, Pac. R. R. Surv., X, 1858, 291, locality Humboldt River at Imlay, Nev.
Gila ardesiaca Cope, Wheeler Surv., Zool., V, 1875, 660, Pl. XXX, figs. 1, 1a, locality unknown, probably Nevada.
Squalius galtae Cope, Proc. Acad. Nat. Sci. Phila., XXXV, 1883, 148, Pyramid Lake, Nev.
Phoxinus clevelandi C. H. and R. S. Eigenmann, West Amer. Scientist, 1889, 149, Aetna Springs, Napa County, Calif. (locality probably erroneous).
- 843. *Cheonda evermanni* (Judson).**
 South Platte basin.
Leuciscus evermanni Juday, Bull. U. S. Bur. Fish., XXIV, 1904 (1905), 226, with plate, Boulder Creek, Boulder, Colo.; A. 8; scales 47; teeth 2, 5-5, 1.

- 844. *Cheonda hydrophlox* (Cope).** *Silverside minnow; Po-he-wa.*
 Salt Lake basin and upper Snake River.
Clinostomus hydrophlox Cope, Hayden's Geol. Surv. Mont., 1871, 475,
 Blackfoot Creek, Idaho.
?Clinostomus tænia Cope, Proc., Amer. Phil. Soc., XIV, 1874, 133,
 Utah Lake, Provo, Utah.
- 845. *Cheonda humboldti* (Girard).**
 Locality doubtful.
Tigoma humboldti Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 206,
 Humboldt River, Nev.; may be identical with *C. hydrophlox*, but this
 seems unlikely.
- 846. *Cheonda montana* (Cope).**
Clinostomus montanus Cope, Hayden's Geol. Surv. Mont., 1871, 476,
 Grass Creek, Idaho.
?Clinostomus tænia Cope, Proc., Amer. Phil. Soc., XIV, 1874, 133,
 Utah Lake, Provo, Utah.
 May prove to be the same as *C. hydrophlox*.
- 847. *Cheonda copei* (Jordan and Gilbert).** *Leather-sided minnow.*
 Great Basin of Utah and Wood River, Idaho.
?Tigoma gracilis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 206,
 locality unknown; species unidentifiable, name preoccupied in
Leuciscus.
Squalius copei Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 461,
 Bear River, Evanston; not *Leuciscus copei* Günther, a *Notropis*.
Squalius aliciae Jouy, loc. cit., IV, 1881, 19, Provo River near Utah
 Lake, Utah.
- Genus 312. TIGOMA Girard.**
Tigoma Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 205 (*Gila*
puschella Baird and Girard).
Protoporus Cope, Hayden's Geol. Surv. Mont., 1871 (1872), 473 (*P.*
dominicus Cope, young of *Tigoma atraria* Girard).
Myloleucus Cope, loc. cit., 475 (*M. pulverulentulus* Cope); the teeth of
Siphateles accidentally placed in the pharynx of a *Tigoma*.
Temeculina Cockerell, Proc. Biol. Soc. Wash., XXII, 1909, 216 (*Rich-*
ardsonius orcutti Eigenmann and Eigenmann).
- 848. *Tigoma atraria* (Girard).**
 Great Basin of Utah, and Snake River Basin above Shoshone Falls.
?Tigoma lineata Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 206,
 locality unknown; not identifiable.
Siboma atraria Girard, loc. cit., 208, Fish Springs, Toole Co., Utah,
 near Desert.
Tigoma squamata Gill, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 42,
 Great Salt Lake Basin.
Protoporus dominicus Cope, Hayden's Geol. Surv. Mont., 1871, 473,
 Snake River, Fort Hall, Idaho.
Hybopsis bivittatus Cope, loc. cit., 474, Warm Springs, Utah.
Hybopsis timpanogenensis Cope, Proc. Amer. Phil. Soc., XIV, 1874, 134,
 Timpanogos, Utah; young.
Squalius cruentatus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880,
 460, Provo River, Utah.
Squalius rhomaleus Jordan and Gilbert, loc. cit., 461, Utah Lake, Utah.
- 849. *Tigoma gibbosa* (Baird and Girard).**
 Rio Gila, Ariz.
Gila gibbosa Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854,
 28, Rio Santa Cruz, Ariz.
Tigoma intermedia Girard, loc. cit., VIII, 1856, 206, Rio San Pedro of
 the Gila, Ariz.
Gila nigra Cope, Wheeler's Surv., Zool., V, 1875, 663, Ash Creek and
 San Carlos, Ariz.
Squalius lemmoni Rosa Smith, Bull. Calif. Acad. Sci., I, 1884, 3, Rillito
 Creek, near Tucson, Ariz.
- 850. *Tigoma purpurea* Girard.**
 San Bernardino Creek in southern Arizona, a tributary of Rio Yaqui.
Tigoma purpurea Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 206,
 San Bernardino Creek, Ariz.

851. *Tigoma pulchella* (Girard). *Pescadito; Chub of the Rio Grande.*

Rio Grande and other Mexican rivers.

Gila pulchella Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 29, Rio Mimbres, Chihuahua; not *Leuciscus pulchella* Storer, a *Cheilonomus*.

Tigoma nigrescens Girard, loc. cit., VIII, 1856, 207, Boca Grande and Rio Janos, Chihuahua, Mexico.

Tigoma pulchra Girard, loc. cit., 207, Boca Grande and Rio Janos; Chihuahua River, Mexico.

Clinostomus pandora Cope, Hayden's Geol. Surv. Mont., 1871, 475, tributaries Rio Grande, Sangre de Cristo Pass.

Gila gila Cope, Wheeler's Surv., Zool., V, 1875, 661, Rio de Acama and near Fort Wingate, N. Mex.

Cheonoda modesta Garman, Bull. Mus. Comp. Zool., VIII, 1881, 92, Rio Salinas, Saltillo, Coahuila.

Gila conspersa Garman, loc. cit., VIII, 1881, 91, Rio Nazas, Coahuila, Mexico.

852. *Tigoma bicolor* Girard.

Klamath Lake, Oreg.

Tigoma bicolor Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 206, Klamath Lake, Oreg.

Cheonoda cœrulea Girard, loc. cit., 207, Lost River, Oreg.

853. *Tigoma pulverulenta* (Cope).

Myloleucus pulverulentus Cope, Hayden's Geol. Surv. Mont., 1871, 475, Warm Springs, Utah.

Myloleucus is a synonym of *Tigoma*. In his study of these species, according to Fowler, Professor Cope by accident replaced the single-rowed teeth of a *Siphateles* in his type example of *Myloleucus pulverulentus*, a species having teeth in two rows.

Subgenus TEMECULINA Cockerell.

854. *Tigoma orcutti* (Eigenmann and Eigenmann).

Temecula, San Luis Rey, San Jacinto, and Santa Ana Rivers, California.
Phoxinus orcutti Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2nd ser., III, 1890, 2, Temecula River, Riverside County, Calif.

Genus 313. *SIBOMA* Girard.

Siboma Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 209 (*S. crassicauda* Girard).

855. *Siboma crassicauda* (Baird and Girard). *Sacramento chub.*

Rivers of California; generally abundant in the Sacramento and San Joaquin Rivers.

Lavinia crassicauda Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 137, San Joaquin River.

Tigoma crassa Girard, loc. cit., VIII, 1856, 207, Sacramento River.

856. *Siboma conformis* (Baird and Girard).

Tulare Valley and Sacramento-San Joaquin Basin, Calif.

Lavinia conformis Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 137; Poso Creek, Tulare County, Calif.

Genus 314. *CLINOSTOMUS* Girard.

Clinostomus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 211 (*Luxilus elongatus* Kirtland).

857. *Clinostomus elongatus* (Kirtland). *Red-sided shiner.*

Great Lakes, upper Mississippi Valley, and New York.

Luxilus elongatus Kirtland, Rept. Zool. Ohio, 1836, 169, 193, Mahoning River, Ohio, and Lake Erie near Cleveland.

Leuciscus productus Storer, Synopsis, 1846, 416, Wabash River.

?*Leuciscus longirostris* Kirtland, The Family Visitor, Mar. 14, 1850, Cleveland, Ohio, and Hudson River.

Squalius proriger Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 280, Michigan.

858. *Clinostomus vandoisulus* (Cuvier and Valenciennes). *Minnow; Long-mouthed minnow; Rosy dace; Pike shiner.*
Leuciscus vandoisulus Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 236 (317), South Carolina.
Clinostomus affinis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 212, James River, Va.
Clinostomus funduloides (Potomac River at Washington) and *carolinus* (Salem, N. C.) Girard, loc. cit., 212.
Gila estor Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 66, Elk River, Estill Springs, Tenn.
- Genus 315. **HEMITREMIA** Cope.
Hemitremia Cope, Proc. Amer. Phil. Soc., XI, 1870, 462 (*H. vittata* Cope = *Phoxinus flammatus* Jordan and Gilbert).
859. **Hemitremia vittata** Cope.
Tributaries of the Tennessee River; common in clear streams in northern Alabama.
Hemitremia vittata Cope, Proc. Amer. Phil. Soc., XI, 1870, 462, Holston River, Knoxville, Tenn.; *vittatus* is preoccupied in *Leuciscus*.
Phoxinus flammatus Jordan and Gilbert, in Jordan, Man. Vert., ed. 2, 1878, 303, Elk River, Tenn.
860. **Hemitremia milneriana** (Cope).
Upper Missouri River Basin.
Phoxinus milnerianus Cope, Amer. Nat., July, 1879, 440, upper Missouri River Basin; probably Battle Creek, Mont.
- Genus 316. **PFRILLE** Jordan.
Pfrille Jordan, Copeia, No. 32, 1924, 71 (*Phoxinus neogaeus* Cope).
861. **Pfrille neogaea** (Cope). *Bronze minnow.*
Upper Mississippi Valley and neighboring waters; also Maine and Connecticut lakes.
Phoxinus neogaeus Cope, Trans. Amer. Phil. Soc., XIII, 1866, 375, New Hudson, Mich.
?Leuciscus burtonianus LeSueur, in Vaillant, Bull. Soc. Philom., VIII, 1896, 28, with plate, Burton Mine, Mo.; perhaps a distinct species.
- Genus 317. **IOTICHTHYS** Jordan and Evermann.
Iotichthys Jordan and Evermann, Fishes North and Mid. Amer., 1896, 228 (*Clinostomus phlegethontis* Cope).
862. **Iotichthys phlegethontis** (Cope).
Tributaries of Great Salt Lake and Sevier Lake; basin of Lake Bonneville.
Clinostomus phlegethontis Cope, Proc. Amer. Phil. Soc., XIV, 1874, 137, Beaver River, Utah.
- Genus 318. **HESPEROLEUCUS** Snyder.
Hesperoleucus Snyder, Bull. U. S. Bur. Fish., XXXII, 1912, 63 (*Pogonichthys symmetricus* Girard).
Myloleucus Cope, Hayden's Geol. Surv. Mont., 1871, 475 (*M. pulverulentus* Cope = *Pogonichthys symmetricus* Baird and Girard, in part, not type, the teeth having been misplaced).
This genus and others related, having the teeth 4-5 or 5-5, were placed in the former check list and the Fishes of North and Middle America, in the European genus *Rutilus* Rafinesque; but no American or Asiatic dace is congeneric with *Rutilus rutilus* L.
863. **Hesperoleucus symmetricus** (Baird and Girard).
Rivers of California and Nevada and waters of Lake County, Oreg.
Pogonichthys symmetricus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 136, Fort Miller, San Joaquin Valley, Calif.
Myloleucus parovanus Cope, Proc. Amer. Phil. Soc., XIV, 1874, 131, Beaver River, Utah.
864. **Hesperoleucus navarroensis** Snyder. *Navarro roach.*
Navarro Basin on the western coast of northern California.
Hesperoleucus navarroensis Snyder, Bull. U. S. Bur. Fish., XXXII, 1912 (1914), 65, Navarro River near Philo, Calif.

- 865. *Hesperoleucus parvipinnis* Snyder.** *Short-finned roach.*
Gualala River Basin, Calif.
**Hesperoleucus parvipinnis* Snyder, Bull. U. S. Bur. Fish., XXXII, 1912*
(1914), 66, Gualala River, Sonoma County, Calif.
- 866. *Hesperoleucus venustus* Snyder.** *Venus roach.*
Russian River and the streams entering San Pablo, Suisun, and San
Francisco Bays.
**Hesperoleucus venustus* Snyder, Bull. U. S. Bur. Fish., XXXII, 1912*
(1914), 64, fig. 2, Coyote Creek near Gilroy Hot Springs, Calif.
- 867. *Hesperoleucus mitrulus* Snyder.** *Northern roach.*
Streams tributary to the north end of Goose Lake, Oreg.
**Hesperoleucus mitrulus* Snyder, Bull. U. S. Bur. Fish., XXXII, 1912,*
(1914), 67, Drew Creek, Lake County, Oreg.
- 868. *Hesperoleucus subditus* Snyder.** *Monterey roach.*
Pajaro River system in the region of foothills and lower mountain
ranges of coast ranges.
**Hesperoleucus subditus* Snyder, Bull. U. S. Bur. Fish., XXXII, 1902*
(1904), 67, fig. 3, Uvas Creek, Pajaro River Basin, Santa Clara
County, Calif.
- 869. *Hesperoleucus mearnsi* (Snyder).**
Rio Yaqui.
**Notropis mearnsi* Snyder, Proc. U. S. Nat. Mus., XLIX, 1916 (Dec.*
23, 1915), 582, fig., Rio San Bernardino, Yaqui Basin, Sonora,
Mexico.
- 870. *Hesperoleucus anticus* (Cope).**
Texas.
**Algansea antica* Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 282,*
Texas.
- Genus 319. *SIPHATELES* Cope.**
**Siphateles* Cope, Proc. Acad. Nat. Sci. Phila., XXXV, 1883, 146 (S.*
vittatus Cope); young example.
- 871. *Siphateles olivaceus* (Cope).**
Lake Tahoe; Pyramid Lake; Eagle Lake and Willow Creek, Calif.
**Leucus olivaceus* Cope, Proc. Acad. Nat. Sci. Phila., XXXV, 1883, 145,*
Pyramid Lake, Nev.
- 872. *Siphateles obesus* (Girard).**
Lakes and streams of Lahontan Basin.
**Algansea obesa* Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 183,*
Humboldt River, Nev.
**Leucus dimidiatus* Cope, loc. cit., XXXV, 1883, 145, Pyramid Lake,*
Nev.
**Siphateles vittatus* Cope, loc. cit., 146, Pyramid Lake, Nev.*
- 873. *Siphateles bicolor* (Girard).**
Lakes of southeastern Oregon (Klamath, Warner, Goose, etc.); Sacra-
mento-San Joaquin Basin; Pit River.
**Algansea bicolor* Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 183,*
Klamath Lake.
- 874. *Siphateles formosus* (Girard).**
**Algansea formosa* Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 183,*
Merced and Mohave Rivers, Calif.
**Myloleucus thalassinus* Cope, loc. cit., XXXV, 1883, 144, Goose Lake,*
Oreg.
- 875. *Siphateles columbianus* (Snyder).**
Lakes of southeastern Oregon.
**Rutilus columbianus* Snyder, Bull. U. S. Bur. Fish., XXVII, 1907, 92,*
fig. 4, Warm Springs near Harney Lake, Harney County, Oreg.
- 876. *Siphateles oregonensis* (Snyder).**
Lakes of southeastern Oregon.
**Rutilus oregonensis* Snyder, Bull. U. S. Bur. Fish., XXVII, 1907, 87,*
fig. 3, XL, Spring, Abert Lake, Oreg.

877. Siphateles mohavensis Snyder.

Mohave River, Calif.

Siphateles mohavensis Snyder, Proc. U. S. Nat. Mus., LIV, 1919 (Mar. 15, 1918), 298, figure, Mohave River near Victor, Calif.**Genus 320. LEUCIDIUS Snyder.***Leucidius* Snyder, Bull. U. S. Bur. Fish., XXXV, 1917, 64 (*L. pectinifer* Snyder).**878. Leucidius pectinifer Snyder. Lake minnow.**

Lahontan system, Nevada and northeastern California.

Leucidius pectinifer Snyder, Bull. U. S. Bur. Fish., XXXV, 1916-17 (Sept. 28, 1917), 64, figs. 5 and 6, Pyramid Lake, Nev.**Genus 321. NOTROPIS Rafinesque.***Notropis* ²⁷ Rafinesque, Amer. Month. Mag., II, 1818, 204 (*Notropis atherinoides* Rafinesque = *Alburnus rubellus* Agassiz).*Minnilus* Rafinesque, Ichth. Ohiensis, 1820, 45 (*M. dinemus* Rafinesque).*Alburnellus* Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856 (*A. dilectus* Girard).*Photogenis* Cope, Trans. Amer. Phil. Soc., XIII, 1867, 378 (*Squalius photogenis* Cope = *Photogenis leucops* Cope).**879. Notropis atherinoides Rafinesque. Shiner; Emerald minnow; Rosy minnow.**

Great Lakes region and Ohio and Mississippi Valleys, and north to Winnipeg.

Notropis atherinoides Rafinesque, Amer. Month. Mag., II, 1818, 204, Lake Erie.*Minnilus dinemus* Rafinesque, Ichth. Ohiensis, 1820, 46, Ohio River.*Alburnus rubellus* Agassiz, Lake Superior, 1850, 364, pl. 3, figs. 1-3, Sault Ste. Marie and the Pic, Lake Superior.*Alburnus nitidus* Kirtland, Cleveland Ann. Sci., 1854, 44, tributaries of Lake Erie.*Alburnellus jaculus* Cope, Trans., Amer. Phil. Soc., XIII, 1866 (1869). 387, St. Joseph River and Dowagiac River, Mich.*Leuciscus copii* Günther, Cat., VII, 1868, 255, Virginia and Michigan; ^{as b} titillifer *jaculus*, preoccupied.**880. Notropis photogenis (Cope). Silvery minnows.**

Allegheny region.

Rutilus compressus Rafinesque, Ichth. Ohiensis, 1820, 51, Monongahela River; not certainly identifiable; may be *Luxilis cornutus*.*Squalius photogenis* Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 280, Youghiogheny River, Pa.*Photogenis leucops* Cope, loc. cit., XIX, 1867, 164, Sinking Creek, and near Austinville, Va.*Photogenis leucops engraulinus* Cope, loc. cit., 164, tributaries of Kanawha River, Austinville, Va.**881. Notropis amoenus (Abbott).**

Clear streams of the Alleghanies from the Raritan to the Neuse.

Alburnellus amoenus Abbott, Amer. Nat., VIII, 1874, 334, Raritan River, N. J.**882. Notropis scepticus (Jordan and Gilbert).**

Rivers of the Carolinas from Cape Fear to the Santee.

Notropis photogenis ("pale variety"), Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 23.*Minnilus scepticus* Jordan and Gilbert, Synopsis, 1883, 200, Saluda River, S. C..

²⁷ The name *Notropis atherinoides* was given by Rafinesque to a small and very slender minnow from Lake Erie. It was apparently a semi-dried example having therefore a ridge on the back as implied in the name *Notropis*. It had a long anal fin of 11 rays, and the dorsal inserted behind the ventrals. Among the small minnows of the Great Lakes it could hardly be anything else than *rubellus*; yet the original description is very incomplete and perhaps not identifiable. As this is the oldest name for any of the small American minnows with four teeth in the main row of the pharyngeals, and as distinctions of genera among these is especially difficult, the writers in 1898 extended the name to include them all. But this assemblage is hardly a natural one, as the species diverge in various directions. Hence, in this catalogue the former subgenera are given generic rank.

883. *Notropis telescopus* (Cope).

Tennessee River Basin; very abundant in cold mountain streams.
Photogenes telescopus Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867
 165, Holston River, Va.

884. *Notropis arcansanus* (Meek).

Ozark region.

Notropis telecopus arcansanus Meek, Bull. U. S. Fish Com., IX, 1889
 (1891), 133, White River, Eureka Springs, Ark.

885. *Notropis universitatis* Evermann and Cockerell.

Boulder Creek, Boulder, Colo.

Notropis notatus var., Univ. of Colo. Stud., Apr., 1908, 170.

Notropis universitatis Evermann and Cockerell, Proc. Biol. Soc. Wash.,
 XXII, Dec., 1909, 187, Boulder Creek, Boulder, Colo.

886. *Notropis welaka* Evermann and Kendall.

St. Johns River, Fla.

Notropis welaka Evermann and Kendall, Bull. U. S. Fish Com., XVII,
 1897 (Feb. 9, 1898), 128, pl. 6, fig. 2, St. Johns River near Welaka,
 Fla.

887. *Notropis leuciodus* (Cope).

Tennessee River Basin, in mountain streams.

Photogenes leuciodus Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867,
 165, Holston River, Va.

888. *Notropis micropteryx* (Cope).

Headwaters of Cumberland and Tennessee Rivers; also in Ozark region.
Alburnellus micropteryx Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868,
 233, Holston River, Va.

889. *Notropis metallicus* Jordan and Meek.

Swamp streams in Georgia and Florida, from the Suwanee Basin to
 the Escambia.

Notropis metallicus Jordan and Meek, Proc. U. S. Nat. Mus., VII,
 1884, 475, Allapaha River, Nashville, Ga.

890. *Notropis louisianæ* Evermann.

Atchafalaya River, La.

Notropis louisianæ Evermann, in Jordan and Evermann, Fishes North
 and Mid. Amer., 1898, 2801, Atchafalaya River, Melville, La.

891. *Notropis socius* (Girard).

Live Oak Creek, southwestern Texas.

Alburnus socius Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 193,
 Live Oak Creek, Tex.

892. *Notropis amabilis* (Girard).

Rio Leona, a tributary of Rio Nueces, Tex.

Alburnus amabilis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856,
 193, Rio Leona, Tex.

893. *Notropis argenteus* (Cope).

Upper Mississippi Valley.

Alburnellus argenteus Cope, Trans. Amer. Phil. Soc., XIII, 1866, 388, Detroit
 River or St. Joseph River, Mich.

894. *Notropis pilosus* Fowler.

Notropis pilosus Fowler, Proc. Acad. Nat. Sci. Phila., LVI, 1904, 245,
 figure, Rogers, White River Basin, Ark.

895. *Notropis rubrifrons* (Cope). *Rosy-faced minnow.*

New York and western Pennsylvania to southern Michigan, Kansas,
 and Kentucky.

Alburnus rubrifrons Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 85,
 Kiskiminitas River, a tributary of Allegheny River.

Alburnellus percobromus Cope, Hayden's Geol. Surv. Wyo., 1870, 440,
 Missouri River at St. Joseph, Mo.

896. *Notropis dilectus* (Girard).

Lower Ohio to the Rio Grande; abundant in Arkansas and eastern Texas in sandy streams.

Alburnus dilectus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 193, Arkansas River, Fort Smith, Ark.

Alburnus lepidulus Girard, loc. cit., 212, Black Warrior River, Ala.

Alburnus oligaspis Cope, loc. cit., XVI, 1864, 282, Kansas.

Alburnellus jemezanus Cope, Wheeler Surv., Zool., V, 1875, 650, Rio Grande, San Ildefonso, Mexico.

897. *Notropis stilbius* (Jordan).

Alabama River and tributaries.

Nototropis stilbius Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 343, Etowah and Oostanaula Rivers, Rome, Ga.

898. *Notropis notemigonoides* Evermann.

Rio Neches near Palestine, Tex., and streams about Houston, Tex.

Notropis notemigonoides Evermann, Bull. U. S. Fish Com., XI, 1891 (1892), 81, Neches River, Palestine, Tex., and Sims Bayou, Houston, Tex.

899. *Notropis fumeus* Evermann.

Hunter Creek, Houston, Tex.

Notropis fumeus Evermann, Bull. U. S. Fish Com., XI, 1891 (1892), 81, Hunter Creek, Houston, Tex.

Genus 322. *PARANOTROPIS* Fowler.

Episema Cope and Jordan, Bull. U. S. Nat. Mus., X, 1877, 64 (*Photogenis scabriceps* Cope, name preoccupied).

Paranotropis Fowler, Proc. Acad. Nat. Sci. Phila., LVI, 1904, 245 (*Photogenis scabriceps* Cope).

900. *Paranotropis scabriceps* (Cope).

Kanawha River, W. Va.

Photogenis scabriceps Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867, 166, Sinking and Walker Creeks, tributaries of Kanawha River.

901. *Paranotropis ariommus* (Cope).

Ohio and Tennessee Valleys.

Photogenus ariommus Cope, Trans. Amer. Phil. Soc., XIII, 1866, 378, no locality given.

902. *Paranotropis jejunus* (Forbes).

Western Pennsylvania to Kansas and north to Winnipeg.

Episema jejuna Forbes, Bull. Ill. Lab. Nat. Hist., I, 1878, 60, Illinois River.

903. *Paranotropis megalops* (Girard).

Rivers of Texas from the Colorado westward.

Alburnus megalops Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 193, San Felipe Creek, Tex.; the name preoccupied in *Notropis (Luxilus)* but not in *Paranotropis*.

Genus 323. *LYTHRURUS* Jordan.

Lythrurus Jordan, Man. Vert., ed. 1, 1876, 272 (*Notropis lythrurus* Jordan = *Semotilus diplæmius* of authors, not of Rafinesque).

904. *Lythrurus umbratilis* (Girard).

Arkansas River and streams of Kansas and southwestern Iowa.

Alburnus umbratilis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 193, Sugar Loaf Creek, tributary of Poteau River, Ark.

Luxilus lucidus Girard, loc. cit., 203, Coal Creek, tributary of South Fork of Canadian River.

Minnius nigripinnis Gilbert, Bull., Washburn Lab. Nat. Hist., I, 1884, 14, Shunganunga Creek near Topeka, Kans.

?*Notropis macrolepidotus* Forbes, Bull. Ill. Lab. Nat. Hist., II, 1885, 138, Illinois.

- 905. *Lythrurus cyanocephalus* Copeland.**
 Rivers of southern Wisconsin to Ohio and Kentucky and Indiana.
Lythrurus cyanocephalus Copeland, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 70, Racine River, Wis.
Rutilus ruber Rafinesque, Ichth. Ohiensis 1820, 52, Elkhorn and Kentucky Rivers; named but not described.
Notropis lythrurus Jordan, Proc. U. S. Nat. Mus., VII, 1884, 476, White River, Indianapolis; a rather more slender southern form.
- 906. *Lythrurus atripes* Jordan.** *Blackfin; Compressed redfin.*
 Southern Illinois and Iowa.
Lythrurus atripes Jordan, Bull. Ill. Lab. Nat. Hist., No. 2, 1878, 59, streams of Union and Johnson Counties, Ill.
- 907. *Lythrurus ardens* (Cope).**
 Roanoke River, Va.
Hypsilepis ardens Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867, 163, headwaters of Roanoke River, Montgomery County, Va.
- 908. *Lythrurus fasciolaris* (Gilbert).**
 Southern bend of Tennessee River, in limestone streams.
Notropis umbratilis fasciolaris Gilbert, Bull. U. S. Fish Com., IX, 1889 (1891), 148, streams about Florence, Ala.
- 909. *Lythrurus matutinus* (Cope).**
 Neuse and Pamlico Rivers; common in sandy brooks.
Alburnellus matutinus Cope, Proc. Amer. Phil. Soc., XI, 1870, 465, Neuse River, Wake County, N. C.
- 910. *Lythrurus punctulatus* (Hay).**
 Tributaries of Big Hatchee River, northern Mississippi.
Minnius punctulatus Hay, Proc. U. S. Nat. Mus., III, 1880, 508, Tuscmibia River, a tributary of the Big Hatchee, near Corinth, Miss.
- 911. *Lythrurus bellus* (Hay).**
 Tombigbee River, Miss.
Minnius bellus Hay, Proc. U. S. Nat. Mus., III, 1880, 510, Tombigbee River, Artesia and Macon, Miss.
- 912. *Lythrurus roseipinnis* (Hay).**
 Sandy streams of Gulf States, from Escambia River to the Chickasawha.
Notropis roseipinnis Hay, in Jordan, Cat. Fishes N. Amer., 1885, 27, Chickasawha River, Enterprise, Miss.; substitute for *rubripinnis*, preoccupied.
- 913. *Lythrurus lirus* (Jordan).**
 Alabama River Basin; abundant in sandy streams.
Notropis lirus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 342, Etowah River, Rome, Ga.
Notropis alabamæ Jordan and Meek, Proc. U. S. Nat. Mus., VII, 1884, 476, Alabama River, Montgomery, Ala.
- Genus 324. *ORCELLA* Jordan and Evermann**
Orcella Jordan and Evermann, Fish. North and Mid. Amer., 1896, 254 (*N. orca* Woolman); not "*Orcælla*," a genus of Cetacea.
Orcula Jordan and Evermann, loc. cit., 1900, 3140, substitute for *Orcella*, regarded as preoccupied by *Orcælla*.
- 914. *Orcella nazas* (Meek).**
 Headwaters of the Rio Nazas in Durango, Mexico.
Notropis nazas Meek, Field Col. Mus., Zool., V, 1904, 70, fig. 19, Santiago Papasquiaro, Durango, Mexico.
- 915. *Orcella orca* (Woolman).**
 Rio Grande at El Paso, Tex.
Notropis orca Woolman, Bull. U. S. Fish Com., XIV, 1894, 56, Rio Grande, El Paso, Tex.
- Genus 325. *HYDROPHLOX* Jordan and Brayton**
Hydrophlox Jordan and Brayton, Bull. U. S. Nat. Mus., XIII, 1878, 32 (*Hybopsis rubricroceus* Cope).

- 916. *Hydrophlox rubricroceus* (Cope).** *Red fallfish.*
 Headwaters of the Tennessee and Savannah Rivers.
Hybopsis rubricroceus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 231,
 Tumbling Creek, tributary of the North Fork of Holston River, Va.
- 917. *Hydrophlox chlorocephalus* (Cope).**
 Santee Basin, in North and South Carolina.
Hybopsis chlorocephalus Cope, Proc. Amer. Phil. Soc., XI, 1870, 461,
 Catawba River, N. C.
- 918. *Hydrophlox lutipinnis* Jordan and Brayton.**
 Santee and Oconee Rivers, Ga.
Hydrophlox lutipinnis Jordan and Brayton, Bull. U. S. Nat. Mus.,
 XII, 1878, 36, Oconee River, Halls Springs, Ga.
- 919. *Hydrophlox chiliticus* (Cope).**
 Basin of Great Pee Dee River, N. C.; common in upland streams.
Hybopsis chiliticus Cope, Proc. Amer. Phil. Soc., XI, 1870, 462, Yadkin
 River, Rowan County, N. C.
- 920. *Hydrophlox altipinnis* (Cope).**
 Yadkin, Great Pee Dee, and Cape Fear Rivers, N. C.
Alburnellus altipinnis Cope, Proc. Amer. Phil. Soc., XI, 1870, 464,
 Yadkin River, Rowan County, N. C.
- 921. *Hydrophlox roseus* (Jordan).**
 Lowland streams of Gulf States from Ogeechee River to the Mississippi.
Luzilus roseus Jordan, Bull. U. S. Nat. Mus., X, 1877, 61, Notalbany
 River near Tikfaw, La.
- 922. *Hydrophlox stonei* (Fowler).**
 Streams of South Carolina.
Notropis stonei Fowler, Proc. Acad. Nat. Sci. Phila., LXXII, 1920, 391,
 figure, Pocataligo River, Manning, S. C.
- 923. *Hydrophlox chalybeus* (Cope).**
 Delaware River to Ogeechee River.
Hybopsis chalybeus Cope, Trans., Amer. Phil. Soc., XIII, 1866, 383,
 Schuylkill River, Pa.
- 924. *Hydrophlox abbotti* (Fowler).**
Notropis chalybeus abbotti Fowler, Proc. Acad. Nat. Sci. Phila., LVI,
 1904, 239, Pl. XVII, upper figure, Batsto River, N. J.
- 925. *Hydrophlox chrosomus* (Jordan).**
 Alabama Basin; very abundant in clear streams and outlets of springs.
Hybopsis chrosomus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 333,
 Etowah River, etc., Rome, Ga.
- 926. *Hydrophlox xenocephalus* (Jordan).**
 Georgia to Mississippi; common in streams of the pine woods, descending
 to brackish water; abundant in Perdido Bay, Pensacola, Fla.
Hybopsis xenocephalus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876,
 334, Etowah River, etc., Rome, Ga.
- Genus 326. COCCOTIS Jordan**
Coccotis Jordan, Rept. Zool. Ohio, 1882, 852 (*Hypsilepis coccogenis*
 Cope).
Coccogenia Cockerell, Proc. Biol. Soc. Wash., XXII, 1909, 190 (*Hypsi-*
lepis coccogenis Cope).
- 927. *Coccotis coccogenis* (Cope).**
 Cumberland, Tennessee, and Savannah Rivers.
Hypsilepis coccogenis Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867,
 160, Holston River, Va.
- 928. *Coccotis brimleyi* (B. A. Bean).**
Notropis brimleyi B. A. Bean, Proc. U. S. Nat. Mus., XXVI, 1903, 913,
 Cane River, N. C.
- 929. *Coccotis macdonaldi* (Jordan and Jenkins).**
 Mountain streams of Virginia tributary to the James and Shenandoah.
Notropis macdonaldi Jordan and Jenkins, Proc. U. S. Nat. Mus., XI,
 1888, 354, plate 44, fig. 4, Shenandoah River, Luray, Va.

930. *Coccotis zonatus* (Agassiz).

Mountain streams in the Ozark region of Missouri and Arkansas.
Alburnus zonatus Agassiz, Bull. Mus. Comp. Zool., I, 1863, 9, Osage River, Mo.

931. *Coccotis zonistius* (Jordan).

Chattahoochee River.

Luxilus zonistius Jordan, Proc. U. S. Nat. Mus., II, 1879, 239, Suwannee Creek, tributary of Chattahoochee River in northern Georgia.

Genus 327. *LUXILUS* Rafinesque.

Luxilus Rafinesque, Ichth. Ohiensis, 1820, 47 (*L. chryscephalus* Rafinesque = *Cyprinus cornutus* Mitchell).

Hypselepis Baird, Amer. Jour. Sci. Arts, XVII, 1854, 359 (*Cyprinus cornutus* Mitchell); later written *Hypselepis*.

932. *Luxilus cornutus* (Mitchill). *Shiner; Dace; Redfin.*

Entire region east of the Rocky Mountains excepting the south Atlantic States and Texas; very abundant; also Potomac River and tributaries; not found south of the James.

Cyprinus cornutus Mitchell, Amer. Month. Mag., I, July, 1817, 289, Wallkill River, N. Y.; preliminary notice, and II, 1818, 324, Westchester County, N. Y.

Cyprinus megalops Rafinesque, loc. cit., II, December, 1817, 121, Hudson River, above the falls.

Cyprinus melanurus Rafinesque, loc. cit., 121, Hudson River, above the falls.

?*Luxilus interruptus* Rafinesque, loc. cit., 49, Ohio River; may be *Notropis rubrifrons*.

Rutilus plargyrus Rafinesque, Ichth. Ohiensis, 1820, 50, Ohio River in Kentucky.

Rutilus compressus Rafinesque, loc. cit., 51, Monongahela River, etc., not certainly identifiable; may be *Notropis photogenes*.

Leuciscus vittatus DeKay, N. Y. Fauna: Fishes, 1842, 212, Chittenonga Creek, tributary of Mohawk River.

Argyreas rubripinnis Heckel, Russegger's Reisen, 1843, 1040.

Leuciscus spiralingulus Cuvier and Valenciennes, Hist. Nat. Poiss., XVII, 1844, 239 (321), New Jersey; New Harmony, Ind.

Leuciscus obesus Storer, Proc. Bost. Soc. Nat. Hist., II, 1845, 48, Florence, Ala.

Plargyrus typicus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 196, after Rafinesque.

Plargyrus bowmani Girard, loc. cit., 196, Sweetwater River, Nebr.

Plargyrus argentatus Girard, loc. cit., 212, James River, Va.

Alburnops plumbeolus Cope, loc. cit., XVI, 1864, 282, Saginaw River, Flint, Mich.

Hypselepis cornutus cyaneus Cope, loc. cit., XIX, 1867, 160, Montreal River, Keweenaw, Mich.

Leuciscus frontalis Agassiz, Lake Superior, 1850, 368, Montreal River, eastern shore of Lake Superior.

Leuciscus gracilis Agassiz, loc. cit., 370, Lake Huron.

Hypselepis cornutus gibbus Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867, 158, Monroe County, Mich.

933. *Luxilus cornutus isolepis* (Hubbs and Brown).

Oklahoma and Arkansas.

Notropis cornutus isolepis Hubbs and Brown, Proc. Okla. Acad. Sci., Pt. I, VI, Jan. 1, 1927, 129, Mountain Fork River, McCurtain County Okla.

934. *Luxilus cornutus chryscephalus* (Rafinesque).

Ohio River and southward; the southern form of *cornutus*.

Luxilus chryscephalus Rafinesque, Amer. Month. Mag., 1820, 47, Ohio River.

935. *Luxilus cerasinus* (Cope).

Roanoke River Basin, in mountain brooks.

Hypselepis cornutus cerasinus Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867, 159, headwaters of Roanoke River, Va.

936. *Luxilus albeolus* (Jordan).

Roanoke, Tar, and Neuse Rivers.

Notropis megalops albeolus Jordan, Bull. U. S. Fish Com., VIII, 1888 (1891), 123, Roanoke River near Roanoke, Va.

937. *Luxilus lacertosus* (Cope).

Headwaters of Tennessee River.

Hybopsis lacertosus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 232, Bear Creek, tributary of the Middle Fork of Holston River.

Genus 328. *EROGALA* Jordan and Brayton

Erogala Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1898, 20 (*Photogenis stigmaturus* Jordan).

938. *Erogala pyrrhomelas* (Cope).

North and South Carolina in the Santee Basin.

Photogenis pyrrhomelas Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 463, Catawba River, N. C.

939. *Erogala formosa* (Putnam).

Southern Georgia and Alabama.

Alburnus formosus Putnam, Bull. Mus. Comp. Zool., I, 1863, 9, Mobile; not *Moniana formosa* Girard.

Leuciscus hypselopterus Günther, Cat., VII, 1868, 255, Mobile; substitute for *formosus*, preoccupied in *Leuciscus* but not in *Erogala*.

Photogenis grandipinnis Jordan, Bull. U. S. Nat. Mus., X, 1877, 62, Flint River, Ga.

940. *Erogala whippili* (Girard). *Silverfin; Lemonfin; Satinfin; Steel-colored minnow.*

Central and northern New York to Minnesota, northern Alabama, and Arkansas, in clear streams.

Cyprinella whippili Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 198, Sugarloaf Creek, tributary Poteau River, Ark.

Photogenis spilopterus Cope, Trans. Amer. Phil. Soc., XIII, 1866, 378, St. Joseph River, southwestern Michigan.

Hybopsis fretensis Cope, loc. cit., 382, Detroit, Mich.; young of *C. whippili*, sive Hubbs.

941. *Erogala analostana* (Girard). *Silverfin; Satinfin; Lacefin; Shiner.*

Streams about Delaware and Chesapeake Bays; representing *Erogala whippili* east of the Alleghenies.

Cyprinella analostana Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 59, at Analostan Island, Washington, D. C.

942. *Erogala galactura* (Cope).

Ozark region to Cumberland, Tennessee, and Savannah Rivers; abundant in mountain streams.

Hypsilepis galacturus Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867, 160, Holston River, Va.

943. *Erogala stigmatura* (Jordan). *Spotted-tail minnow.*

Alabama Basin; very abundant.

Photogenis stigmaturus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 337, Etowah River, Rome, Ga.

Cyprinella calliura Jordan, Bull. U. S. Nat. Mus., X, 1877, 61, Black Warrior River, Ala.

944. *Erogala camura* (Jordan and Meek).

Arkansas River and tributaries.

Cliola camura Jordan and Meek, Proc. U. S. Nat. Mus., VII, 1884, 474, Arkansas River, Fort Lyon, Colo.

945. *Erogala nivea* (Cope).

Southern Virginia to South Carolina.

Hybopsis niveus Cope, Proc. Amer. Phil. Soc., XI, 1870, 460, Catawba River, N. C.

946. *Erogala chloristia* (Jordan and Brayton).

Santee River and tributaries.

Codoma chloristia Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 21, tributary of Saluda River, Greenville, S. C.

- 947. *Erogala trichroistia* (Jordan and Gilbert).**
 Alabama River Basin.
Codoma trichroistia Jordan and Gilbert, Bull. U. S. Nat. Mus., XII, 1878, 50, Etowah River, etc., Rome, Ga.
- 948. *Erogala callistia* (Jordan).**
 Alabama River Basin.
Photogenis callistius Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 337, Etowah River, etc., Rome, Ga.
- 949. *Erogala cærulea* (Jordan).**
 Alabama River Basin.
Photogenis cæruleus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 338, Etowah River, Rome, Ga.
- 950. *Erogala eurystoma* (Jordan).**
 Chattahoochee River.
Photogenis eurystomus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 356, Nancys Creek, tributary to Chattahoochee River, near Atlanta, exclusive of the smaller specimens mentioned, which are *Notropis zonistius*; not *Codoma eurystoma* Jordan.
Photogenis leucopus Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 41, Chattahoochee River, Gainesville, Ga.
- 951. *Erogala cercostigma* (Cope).**
 Sandy streams tributary to the Gulf of Mexico, from Pearl River and Red River to Rio Nueces; abundant.
Cyprinella cercostigma Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867, 157, Pearl River, Monticello, Miss.
Luxilus chickasawensis Hay, Proc. U. S. Nat. Mus., III, 1880, 506, Chickasawha River, Enterprise, Miss.
- 952. *Erogala xænura* (Jordan).**
 Altamaha River Basin, Ga.
Minnilus xænurus Jordan, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 79, Ocmulgee River, Flat Shoals, Ga.
- 953. *Erogala notata* (Girard).**
 Streams of Texas from Austin westward.
Cyprinella notata Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 198, Rio Seco, Tex.
- 954. *Erogala venusta* (Girard).**
 Streams of Texas.
Cyprinella venusta Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 198, Rio Sabinal, Tex.
- 955. *Erogala texana* (Girard).**
 Rivers of Texas from Trinity River to Rio Salado.
Cyprinella texana Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 198, Rio Salado and Turkey Creek, Tex.
- Genus 329. CYPRINELLA Girard.**
Cyprinella Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 196 (*Leuciscus bubalinus* Baird and Girard).
- 956. *Cyprinella bubalina* (Baird and Girard).**
 Streams of Arkansas and westward; not common.
Leuciscus bubalinus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 391, Otter Creek, Ark.
Cyprinella umbrosa Girard, loc. cit., VIII, 1856, 197, Coal Creek, tributary of South Fork Canadian River.
Cyprinella gunnisonii Girard, loc. cit., 197, Cottonwood Creek of the Rio Grande, Colo.
Cyprinella beckwithi Girard, loc. cit., 197, Arkansas River, near Fort Makee.
Cyprinella lepida Girard, loc. cit., 197, Rio Frio, Tex.
- 957. *Cyprinella ludibunda* Girard.**
 Rio Grande Basin.
Cyprinella ludibunda Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 199, locality now known to be Cottonwood Creek, Colo.

958. *Cyprinella macrostoma* Girard.

Rio Grande region, northward perhaps to Kansas.

Cyprinella macrostoma Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 198, Devils River, Tex.

?*Cyprinella luxiloides* Girard, loc. cit., 198, San Pedro Creek, Tex.

?*Cyprinella lugubris* Girard, loc. cit., 199, Cottonwood Creek, Colo.

?*Notropis umbrifer* Hay, Proc. U. S. Nat. Mus., X, 1887, 245, Solomon River at Beloit and Saline River near Wakeeney, Kans.

959. *Cyprinella forlonensis* (Meek).

Streams of Tamaulipas, Mexico.

Nototropis forlonensis Meek, Field Col. Mus., Zool., V, 1904, 70, fig. 20, Rio Forlon, Tamaulipas, Mexico.

960. *Cyprinella rubripinna* Garman.

Tributaries of Lago del Muerto, Coahuila, Mexico.

Cyprinella rubripinna Garman, Bull. Mus. Comp. Zool., VIII, 1881, 91, Lago del Muerto; preoccupied.

Notropis garmani Jordan, Cat. Fish. N. Amer., 1885, 813; substitute for *rubripinna*; preoccupied in *Luxilus* but not in *Cyprinella*.

Genus 330. *MONIANA* Girard.

Moniana Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 199 (*Leuciscus lutrensis* Baird and Girard); probably inseparable from *Cyprinella*.

961. *Moniana lutrensis* (Baird and Girard). *Redfin*.

Southern Illinois to South Dakota, Kansas, and the Rio Grande; very abundant in clear brooks southward.

Leuciscus lutrensis Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 391, Otter Creek, tributary of North Fork of Red River, Ark.

Cyprinella suavis Girard, loc. cit., VIII, 1856, 197, San Antonio, Tex.

Moniana leonina Girard, loc. cit., 199, Leon River, tributary to Rio San Antonio, Tex.

Moniana complanata Girard, loc. cit., 200, Brownsville, Tex.

Moniana luteabilis Girard, loc. cit., 200, Hurah Creek, tributary of Rio Pecos.

Moniana pulchella Girard, loc. cit., 200, Sugar Loaf Creek, tributary to the Poteau River.

Moniana couchi Girard, loc. cit., 201, China, Nuevo Leon, Mexico.

Moniana rutila Girard, loc. cit., 201, Cadereita, Nuevo Leon, Mexico.

Moniana gracilis Girard, loc. cit., 201, Monterrey, Nuevo Leon, Mexico.

Moniana gibbosa Girard, loc. cit., 201, Brownsville, Tex.

Cyprinella billingsiana Cope, Hayden's Geol. Surv. Wyo., 1870, 439, St. Joseph, Mo.

Moniana jugalis Cope, loc. cit., 439, St. Joseph, Mo.

Hypsilepis iris Cope, Wheeler Surv., Zool., V, 1875, 653, Rio Grande, San Ildefonso, N. Mex.

Cyprinella forbesi Jordan, Bull. Ill. State Lab. Nat. Hist., I, 1878, 57, clear streams of southern Illinois.

Chiola montiregis Cope, Proc. Amer. Phil. Soc., XII, 1885, 168 (read Dec. 7, 1884), Monterey, Nuevo Leon, Mexico.

962. *Moniana formosa* Girard.

Rio Mimbres, Chihuahua, Mexico.

Moniana formosa Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 201, Rio Mimbres, Chihuahua, Mexico.

963. *Moniana frigida* Girard.

Rio Frio and tributaries of Rio San Antonio, Tex., and Lake Santa Maria, Chihuahua, Mexico.

Moniana frigida Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 200, Rio Frio, Tex.

964. *Moniana santamariae* (Evermann and Goldsborough).

Rio Grande Basin in Mexico.

?*Notropis santamariae* Evermann and Goldsborough, Bull., U. S. Fish Com., XXI, 1901 (1902), 147, fig. 2, Lake Santa Maria in Chihuahua, Mexico.

965. *Moniana proserpina* Girard.

Rio Grande region.

Moniana proserpina Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 200, Devils River, Tex.

Moniana aurata Girard, loc. cit., 200, Piedra Painte, N. Mex.
? *Moniana tristis* Girard, loc. cit., 1856, 201, locality unknown.

966. *Moniana boucardi* (Günther). *Salmichi*.

Central Mexico.

Leuciscus boucardi Günther, Cat., VII, 1868, 485, Cuernavaca, Mexico.
Graodus nigrotæniatus Günther, loc. cit., 485, Atlisco, Mexico.

967. *Moniana robusta* (Meek).

Santa Rosalia and Jimenez, Mexico.

Notropis robustus Meek, Field Col. Mus., Zool., III, 1902, 82, Rio Conchos, Jimenez, Chihuahua, Mexico, figured as *Nototropis robustus*, in Field Col. Mus., Zool., V, 1904, 66.

968. *Moniana chihuahua* (Woolman).

Streams of Chihuahua, Mexico.

Notropis chihuahua Woolman, Amer. Nat., March, 1892, 260, Rio de los Conchos, Chihuahua, Mexico.

969. *Moniana callisema* (Jordan).

Ocmulgee River, Ga.

Episema callisema Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 363, South Fork of Ocmulgee River, Flat Shoals, Ga.

Genus 331. *HUDSONIUS* Girard.

Hudsonius Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 210 (*Clupea hudsonia* Clinton)

970. *Hudsonius chamberlaini* (Evermann).

Atchafalaya River, La.

Notropis chamberlaini Evermann, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2800, Atchafalaya River, Melville, La.

971. *Hudsonius hudsonius* (De Witt Clinton). *Spawneater; Spot-tailed minnow; Shiner; Smelt*.

The Dakotas and Lake Superior to New York, and southward; abundant in the Great Lakes.

Clupea hudsonia De Witt Clinton, Ann. Lyc. Nat. Hist. N. Y., I, 1824, 49, Hudson River.

Hudsonius fluviatilis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 210, Hudson River.

Luxilus selene Jordan, Bull. U. S. Nat. Mus., X, 1877, 60, Lake Superior, Bayfield, Wis.

Notropis scopiferus Eigenmann and Eigenmann, Amer. Nat., 1893, 153. Winnipeg, Brandon, Fort Qu'Appelle and Medicine Hat, Assiniboina.

972. *Hudsonius amarus* Girard. *Gudgeon; Spawneater; Silverfin; Shiner; Smelt shiner; Silvery minnow; Blue-lined minnow*.

Delaware and Potomac Rivers.

Hudsonius amarus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 210, Potomac River at Washington.

Hybopsis phaenæna Cope, loc. cit., 1864, 279, Delaware River, Trenton, N. J.

973. *Hudsonius saludanus* (Jordan and Brayton).

Coastwise streams from the James to the Ocmulgee.

Alburnops saludanus Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 16, a tributary of Saluda River, Greenville, S. C.

Hudsonius euryopa Bean, Proc. U. S. Nat. Mus., II, 1879, 285, McBean Creek, tributary of Savannah River, Ga.

974. *Hudsonius aletes* Jordan and Evermann.

Southern Indiana.

Hudsonius aletes Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, 1927 (Apr. 27), 502, Switz City Swamp, Green County, Ind.

Genus 332. HYBOPSIS Agassiz.

- Hybopsis* Agassiz, Amer. Jour. Sci. Arts, XVII, 1854 (*H. gracilis* Agassiz = *Rutilus amblops* Rafinesque?).
Alburnops Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 194 (*Hybopsis gracilis* Agassiz = *A. blennius* Girard).
Graodus Günther, Cat., VII, 1868, 485 (*Graodus nigrotæniatus* Girard, specimen with pharyngeal teeth lost = *Alburnops blennius* Girard).
Chriope Jordan, Hayden's Geol. Surv. Terr., IV, 1878 (*Hybopsis bifrenatus* Cope).
Minilius Jordan, Man. Vert., ed. 5, 1888, 56 (*Hybopsis procne* Cope).

975. *Hybopsis gracilis* Agassiz.

Tennessee River.

Hybopsis gracilis Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 358, Huntsville, Ala.

976. *Hybopsis blennius* (Girard). *Straw-colored minnow.*

Ohio, Michigan, and New York to Tennessee, Dakota, and Kansas, thence southwestward to Texas.

?*Minnilus microstomus* Rafinesque, Ichth. Ohiensis, 1820, 47, Kentucky River; not certainly identifiable.

?*Rutilus amblops* Rafinesque, loc. cit., 51; not certainly identifiable.

Alburnops blennius Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 194, Arkansas River, near Fort Smith, Ark.

Hybognathus stramineus Cope, loc. cit., XVI, 1864, 283, Detroit River, Grosse Isle.

Hybopsis missouriensis Cope, Hayden's Geol. Surv. Wyo., 1870, 437, Missouri River, St. Joseph, Mo.

Notropis reticulatus Eigenmann and Eigenmann, Amer. Nat., 1893, 152, Fort Qu'Appelle, Assiniboia.

977. *Hybopsis winchelli* Girard.

Black Warrior River, Ala.

Hybopsis winchelli Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 211, Black Warrior River, Ala.

This species and the preceding may prove to be identical with *Hybopsis gracilis*. The synonymy of these little fishes is still unsettled.

978. *Hybopsis procne* (Cope). *Minnow; Minnie; Swallowminnow shiner.*

Delaware River and Cayuga Lake Basin southward in coastwise streams as far as the Neuse River, N. C.

Hybognathus procne Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 283, Conestoga River, a tributary of the Susquehanna.

Hybopsis longiceps Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 231, headwaters of James and Roanoke Rivers, Va.

979. *Hybopsis cummingsi* (Myers).

Streams near Wilmington, N. C.

Notropis cummingsi Myers, Amer. Mus. Novitates, No. 168, April 23, 1925, 1, fig. 1, upper Burnt Mill Creek, Wilmington, N. C.

980. *Hybopsis cayuga* (Meek).

Connecticut River, northern New York, and upper Mississippi Valley.

Notropis cayuga Meek, Ann. Acad. Sci. N. Y., IV, 1888, 305, Cayuga Lake, N. Y.

981. *Hybopsis atrocaudalis* (Evermann).

Rio Neches, Palestine, Tex., also in Rio Comal at New Braunfels, Tex.

Notropis cayuga atrocaudalis Evermann, Bull. U. S. Fish Com., XI, 1891 (1892), 76, Neches River, Palestine, Tex.

982. *Hybopsis keimi* (Fowler).

Allegheny River, western Pennsylvania.

Notropis keimi Fowler, Proc. Acad. Nat. Sci. Phila., LX, 1908, 533, Plate XXVII, tributary of Allegheny River, Port Allegheny, Pa.

983. *Hybopsis volucella* (Cope).

Michigan, Wisconsin, and northern Indiana.

Hybognathus volucellus Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 283, Grosse Isle, Detroit River, Mich.

- 984. *Hybopsis phenacobius* (Forbes).**
 Streams of central Illinois.
Notropis phenacobius Forbes, Bull. Ill. Lab. Nat. Hist., II, 1885, 2, 137,
 Peoria Ill.
- 985. *Hybopsis spectrunculus* Cope.**
 Headwaters of Tennessee River in Tennessee, Virginia, and North Carolina.
Hybopsis spectrunculus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868,
 231, tributary of Middle Fork of Holston River, Va.
- 986. *Hybopsis shumardi* (Girard).**
 Missouri region and southwestward.
Alburnops shumardi Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856,
 194, Arkansas River, Fort Smith, Ark.
Alburnops illecebrosus Girard, loc. cit., 194, Arkansas River, Fort Smith, Ark.
- 987. *Hybopsis boops* (Gilbert). *Bigeye minnow.***
 Maumee River, west to Iowa and Missouri.
Notropis boops Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 201, Salt Creek, Brown County, Ind., and Flat Rock Creek, Decatur County, Ind.
- 988. *Hybopsis kanawha* (Jordan and Jenkins).**
 Tributaries of Kanawha River, southwestern Virginia.
Notropis kanawha Jordan and Jenkins, Proc. U. S. Nat. Mus., XI, 1888, 354, Plate 44, fig. 5, Reed Creek, Wytheville, Va.
- 989. *Hybopsis piptolepis* (Cope).**
 North Platte River.
Photogenis piptolepis Cope, Hayden's Geol. Surv. Wyo., 1870, 438, Red Cloud Creek, a tributary of North Platte River.
- 990. *Hybopsis gibberti* (Jordan and Meek).**
 Illinois to eastern Colorado.
Notropis gibberti Jordan and Meek, Proc. U. S. Nat. Mus., VIII, 1885, 4, Des Moines River and Village Creek, Ottumwa, Iowa; perhaps not distinct from *Hybopsis piptolepis*.
- 991. *Hybopsis horatii* (Cockerell).**
 Platte Basin.
Notropis horatii Cockerell, Science, XXXIV, No. 879, Nov. 3, 1911, 614, Julesburg, Colo.
- 992. *Hybopsis deliciosa* (Girard). *Straw-colored minnow.***
 Michigan, and southwest to Arkansas.
? *Minnilus microstomus* Rafinesque, Ichth. Ohiensis, 1820, 47, Ohio River; not certainly identifiable.
? *Hybopsis dorsalis* Agassiz, Amer. Jour. Sci. Arts., XVII, 1854, 358, Burlington, Iowa; unidentifiable.
Moniana deliciosa Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 199, Leon River, a tributary of the San Antonio River, Tex.
- 993. *Hybopsis deliciosa straminea* Cope. *Northern straw-colored minnow.***
Great Lakes region, mostly in small brooks, the northern form of *H. deliciosa*.
Hybopsis stramineus Cope, Trans. Amer. Phil. Soc., XIII, 1886, 381, Grosse Isle, Detroit River.
- 994. *Hybopsis calientis* (Jordan and Snyder).**
Streams of central Mexico.
Notropis calientis Jordan and Snyder, Bull. U. S. Fish. Com., XIX, 1899 (1900), 122, fig. 4, Rio Verde at Aguas Calientes, Mexico.
- 995. *Hybopsis buchanani* (Meek).**
Poteau River, Okla.
Notropis buchanani Meek, Bull. U. S. Fish Com., XV, 1895 (1896), 342, Poteau River near Poteau, Ind. Ter.
- 996. *Hybopsis heterolepis* (Eigenmann and Eigenmann).**
Notropis heterolepis Eigenmann and Eigenmann, Amer. Nat., 1893, 152, Cayuga Lake.
Notropis muskoka Meek, Field Col. Mus., Zool., I, 1898, 308, Muskoka Lake near Georgian Bay, Ontario.

- 997. *Hybopsis kendalli* (Evermann and Cockerell).**
 Cross Lake Thoroughfare, Me.
Notropis kendalli Evermann and Cockerell, Proc. Biol. Soc. Wash., XXII, Dec. 8, 1909, Cross Lake Thoroughfare, Me.
- 998. *Hybopsis ozarcanus* (Meek).**
 White River, Ark.
Notropis ozarcanus Meek, Bull., U. S. Fish Com., IX, 1889 (1891), 129, North Fork of White River, Ark.
- 999. *Hybopsis scylla* Cope.**
 Illinois River to western Kansas and Montana; abundant in the shallow sandy streams of the plains.
?Alburnus lineolatus Agassiz, in Putnam, Bull., Mus. Comp. Zool., I, 1863, 9, Osage River, Mo.; unidentifiable.
Hybopsis scylla Cope, Hayden's Geol. Surv. Wyo., 1870, 438, Red Cloud Creek, tributary of Platte River.
Cliola chloris Jordan, Bull. Hayden's Geol. Surv. Terr., IV, 1878, 791, upper Missouri River.
- 1000. *Hybopsis longirostris* (Hay).**
 Chickasawha River, Miss., to Escambia River, Fla.
Alburnops longirostris Hay, Proc., U. S. Nat. Mus., III, 1880, 504, Chickasawha River at Enterprise, Miss.
- 1001. *Hybopsis nocomis* (Evermann).**
 Trinity, San Marcos, and Comal Rivers, Tex.
Notropis nocomis Evermann, Bull. U. S. Fish Com., XI, 1891 (1892), 78, Trinity River, Palestine, Tex., and San Marcos River, San Marcos, Tex.
- 1002. *Hybopsis ertenburgeri* (Hubbs.)**
 Mountain Fork River, Okla.
Notropis ertenburgeri Hubbs, Proc. Okla. Acad. Sci., pt. 1, VI, Jan. 1, 1927, 127, Mountain Fork River, McCurtain County, Okla.
- 1003. *Hybopsis nux* (Evermann).**
 Neches and Trinity Rivers near Palestine, Tex.
Notropis nux Evermann, Bull. U. S. Fish Com., XI, 1891 (1892), 77, Neches River, Palestine, Tex.
- 1004. *Hybopsis simus* (Cope).**
 Rio Grande at San Ildefonso, N. Mex.
Alburnellus simus Cope, Wheeler Surv., V, Zool., 1875, 649, San Ildefonso, N. Mex.
- 1005. *Hybopsis sabine* (Jordan and Gilbert).**
 Sabine River, Tex.
Notropis sabine Jordan and Gilbert, Proc. U. S. Nat. Mus., IX, 1886, 15, Sabine River at Longview, Tex.
- 1006. *Hybopsis nitida* (Girard).**
 Tributaries of Rio San Juan at Cadereita, Nuevo Leon, Mexico.
Moniana nitida Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 201, Cadereita, Mexico; not *Alburnus nitidus* Kirtland, also a *Notropis* but not a *Hybopsis*.
Notropis braytoni Jordan and Evermann, Fishes North and Mid. Amer., 1896, 264, Cadereita, Mexico; substitute for *M. nitida* Girard, preoccupied.
- Subgenus CHIROPE Jordan.**
- 1007. *Hybopsis bifrenata* Cope. Bridled minnow; Minnie.**
 Maine to Maryland.
Hybopsis bifrenatus Cope, Trans., Amer. Phil. Soc., XIII, 1867, Schuylkill River, Conshohocken, Pa.
- 1008. *Hybopsis maculata* (Hay).**
 Chickasawha River, Miss.
Hemitremia maculata Hay, Proc. U. S. Nat. Mus., III, 1880, 1505, Chickasawha River, Enterprise, Miss.

- 1009. *Hybopsis anogenus* (Forbes).**
Western New York (Cayuga Lake) to northern Illinois.
Notropis anogenus Forbes, Bull. Ill. Lab. Nat. Hist., II, 1885, Art. 2,
138, Fox River, McHenry, Ill.
- 1010. *Hybopsis heterodon* (Cope).**
New York to Michigan and Kansas; common.
Ablurnops heterodon Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864,
281, Lansing; Grosse Isle, Mich.
Notropis germanus Hay, Proc. U. S. Nat. Mus., X, 1887, 252, Smoky
Hill River, Wallace, Kans.
- 1011. *Hybopsis heterodon richardsoni* (Hubbs and Greene).**
Fox River System, tributary to Green Bay, Wis.
Notropis heterodon richardsoni Hubbs and Greene, Univ. Mich. Mus.
Zool., Misc. Pub. 15, 1926, 39, Pl. III, fig. 1, Fox River, opposite
Lock 25, Columbia County, Wis.
- 1012. *Hybopsis jordani* (Eigenmann and Eigenmann).**
South Saskatchewan River, Medicine Hat, Assiniboia.
Notropis albeolus Eigenmann and Eigenmann, Amer. Nat., 1893, 152,
Medicine Hat, Assiniboia; name preoccupied.
Notropis jordani Eigenmann and Eigenmann, loc. cit., 592, substitute
for *Chriope albeolus*, preoccupied in *Notropis* but not in *Chriope*.
- Genus 333. PLATYGOBIO Gill.**
Platygobio Gill, Trans. Amer. Phil. Soc., V, 1863, 178 (*Pogonichthys*
communis Girard = *Cyprinus* (*Leuciscus*) *gracilis* Richardson).
- 1013. *Platygobio gracilis* (Richardson). Flat-headed chub.**
East slope of the Rocky Mountains, from the Missouri and Yellowstone
Rivers to the Saskatchewan; abundant in river channels as far south
as Kansas City.
Cyprinus (*Leuciscus*) *gracilis* Richardson, Fauna Bor.-Amer., Fishes,
1836, 120, Saskatchewan River at Carleton House.
Coregonus angusticeps Cuvier and Valenciennes, Hist. Nat. Poiss.,
XXI, 1848, 380-390 (534), Saskatchewan River.
Pogonichthys communis Girard, Proc. Acad. Nat. Sci. Phila., VIII,
1856, 188, Yellowstone River, Fort Pierre, Nebr., etc.
Pogonichthys (*Platygobio*) *gulonellus* Cope, loc. cit., XVI, 1864, 277,
near Bridgers Pass, Wyo.
- 1014. *Platygobio physignathus* (Cope).**
Upper waters of Arkansas River; very abundant in the channels; the
most common fish at Pueblo, Colo.
Ceratichthys physignathus Cope, Wheeler Surv., V, Zool., 1875 (1876),
651, Arkansas River, Pueblo, Colo.
- 1015. *Platygobio pallidus* Forbes.**
One specimen from Ohio River at Cairo, Ill.
Platygobio pallidus Forbes, in Jordan and Gilbert, Synopsis, 1883, 220,
Ohio River at Cairo, Ill.
- Genus 334. COUESIUS Jordan.**
Couesius Jordan, Hayden's Geol. Surv. Terr., IV, 1878, 785 (*Nocomis*
milneri Jordan = *Gobio plumbeus* Agassiz).
- 1016. *Couesius squamilentus* (Cope).**
Henry Fork of Green River in southwestern Wyoming.
Ceratichthys squamilentus Cope, Hayden's Geol. Surv. Wyo., 1870, 442,
Henry Fork of Green River, Wyo., Colorado Basin.
- 1017. *Couesius plumbeus* (Agassiz). Chub minnow; Lake chub; Plumbeus
minnow; Moose Lake minnow.**
Streams and lakes from Lake Superior east to the Adirondack region
and New Brunswick; not very common except northward.
Gobio plumbeus Agassiz, Lake Superior, 1850, 366, Lake Superior.
Ceratichthys prosthemius Cope, Trans. Amer. Phil. Soc., XIII, 1867,
Montreal River, Keweenaw Point, Lake Superior.
Nocomis milneri Jordan, Bull. U. S. Nat. Mus., X, 1877, 64, Lake
Superior.

- 1018. *Couesius greeni* Jordan.**
 Stuart Lake, headwaters of Fraser River, British Columbia, and Lake Pend d'Oreille, Idaho.
Couesius greeni Jordan, Proc. U. S. Nat. Mus., XVI, 1893, 313, Stuarts Lake, British Columbia.
- 1019. *Couesius dissimilis* (Girard).**
 Upper Missouri and Black Hills region.
Leucosomus dissimilis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 189, Milk River and Little Muddy River, Mont.
- 1020. *Couesius adustus* Woolman.**
 Rio de los Conchos, Chihuahua, Mexico.
Couesius adustus Woolman, Bull. U. S. Fish Com., XIV, 1894, 57, Rio de los Conchos, Chihuahua, Mexico.
- Genus 335. *NOCOMIS* Girard.**
Nocomis Girard, Proc. Acad. Nat. Sci. Phila., 1856, 190 (*N. nebrascensis* Girard = *Luxilus kentuckiensis* Rafinesque).
Ceratichthys Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 391 (*C. vigilax* Baird and Girard; not of Baird and Girard, 1856); no description.
- 1021. *Nocomis kentuckiensis* (Rafinesque).** *Chub; Hornyhead; River chub; Jerker; Indian chub.*
 Pennsylvania to Wyoming and Alabama, on both sides of the Alleghenies; abundant in the larger streams.
Luxilus kentuckiensis Rafinesque, Ichth. Ohiensis, 1820, 48, Ohio River.
?Catostomus melanotus Rafinesque, loc. cit., 58, Ohio River at the falls.
Semotilus biguttatus Kirtland, Bost. Jour. Nat. Hist., III, 1840, 344, Pl. V, Yellow Creek, Ohio.
Nocomis nebrascensis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 190, Sweetwater River, Nebr.
Ceratichthys leptcephalus Girard, loc. cit., 213, Salem, N. C.
Nocomis bellicus Girard, loc. cit., 213, Black Warrior River.
Ceratichthys cyclotis Cope, loc. cit., XVI, 1864, 277, Grosse Isle, Waterford; Clinton River and Bruce, Mich.
Ceratichthys stigmatus Cope, loc. cit., 1864, 278, Michigan.
- 1022. *Nocomis micropogon* (Cope).**
 Streams of Pennsylvania.
Ceratichthys micropogon Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 277, Conestoga River near Lancaster, Pa.
- Genus 336. *YURIRIA* Jordan and Evermann**
Yuriria Jordan and Evermann, Fishes North and Mid. Amer., 1896, 321 (*Hudsonius altus* Jordan).
- 1023. *Yuriria alta* (Jordan).** *Pesca blanca.*
 Lakes and streams of Guanajuato, tributary to Rio Lerma, Mexico, in Pacific drainage.
Hudsonius altus Jordan, Proc. U. S. Nat. Mus., II, 1879, 301, Lake Tupataro, Guanajuato, Mexico.
- Genus 337. *ERINEMUS* Jordan**
Erinemas Jordan, Man. Vert., 1876, 279 (*Ceratichthys hyalinus* Cope).
- 1024. *Erinemas storarianus* (Kirtland).** *Storer's chub; Lake minnow.*
 Great Lakes and westward to Wyoming, Tennessee, and Arkansas; abundant in the larger streams and lakes, especially in Iowa.
Rutilus storarianus Kirtland, Proc. Bost. Soc. Nat. Hist., I, 1842, 71, Lake Erie (name only); description given as *Leuciscus storarianus* in Bost. Jour. Nat. Hist., V, 1845, 30, Pl. I, fig. 2.
Gobio vernalis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 189, Arkansas River, Fort Smith, Ark.
Ceratichthys lucens Jordan, Proc. U. S. Nat. Mus., II, 1879, 238, Falls of Ohio River near New Albany, Ind.
- 1025. *Erinemas hyalinus* (Cope).** *Silver chub.*
Ceratichthys hyalinus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 226, tributary of Holston River.

- 1026. *Erinemus labrosus* (Cope).**
 Basin of the Santee in North and South Carolina; common.
Ceratichthys labrosus Cope, Proc. Amer. Phil. Soc., XI, 1870, 458,
 tributaries of Catawba River, McDowell and Burke Counties, N. C.
Ceratichthys zanemus Jordan and Brayton, Bull. U. S. Nat. Mus., XII,
 1878, 24, Saluda River, Greenville, N. C.
- 1027. *Erinemus hypsinotus* (Cope).**
 Santee Basin, in North and South Carolina.
Ceratichthys hypsinotus Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870,
 458, tributaries of Catawba River, McDowell County, N. C.
- 1028. *Erinemus rubrifrons* (Jordan).**
 Basin of the Altamaha, Ga.
Nocomis rubrifrons Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876,
 330, Ocmulgee River, Flat Rock, Ga.
- Genus 338. ERIMYSTAX Jordan.**
Erimystax Jordan, Rept. Geol. Surv. Ohio, IV, 1882, 858 (*Luxilus dissimilis* Kirtland).
- 1029. *Erimystax dissimilis* (Kirtland). *Spotted shiner.***
 Lake Erie to the headwaters of the Tennessee and west to Arkansas
 and Iowa.
Luxilus dissimilis Kirtland, Bost. Jour. Nat. Hist., III, 1840, 341, pl.
 4, fig. 2, Mahoning River, Ohio, and Lake Erie near Cleveland,
 Ohio.
- 1030. *Erimystax watauga* (Jordan and Evermann).**
 Holston River, Va.; Watauga River, Tenn.; White River, Ark.;
 Big Barren River, Ky.; and Tippecanoe River, Ind.
Hybopsis watauga Jordan and Evermann, Proc. U. S. Nat. Mus., XI,
 1888, 355, Watauga River, Elizabethtown, Tenn.; North Fork of
 Holston River, Saltville, Va.
- Genus 339. ERIMONAX Jordan.**
Erimonax Jordan, Copeia, 1924, 52 (*Ceratichthys monacus* Cope).
- 1031. *Erimonax monacus* (Cope).**
 Tennessee Basin in the river channels.
Ceratichthys monacus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 227,
 Holston River, Washington County, Va.
- Genus 340. OREGONICHTHYS Hubbs.**
Oregonichthys Hubbs, Occ. Papers, Mus. Zool. Univ. Mich., 1927
 (*Hybopsis crameri* Snyder).
- 1032. *Oregonichthys crameri* (Snyder).**
Hybopsis crameri Snyder, Bull. U. S. Bur. Fish., XXVII, 1907, 181,
 fig. 5, Willamette River at Oregon City, Oreg.
- 1033. *Oregonichthys cumingii* (Günther).**
 California; only the type known.
Ceratichthys cumingii Günther, Cat., VII, 1868, 177, California; type
 locality uncertain.
- Genus 341. MACRHYBOPSIS Cockerell and Alliston.**
Macrhybopsis Cockerell and Alliston, Proc. Biol. Soc. Wash., XXII,
 1909, 162 (*Gobio gelidus* Girard).
- 1034. *Macrhybopsis aestivalis* (Girard).**
 Arkansas River to the Rio Grande; abundant in sandy river channels,
 not in small brooks.
Gobio aestivalis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 189,
 Rio San Juan, near Cadereita, Nuevo Leon.
- 1035. *Macrhybopsis marconis* (Jordan and Gilbert).**
 Abundant in Rio San Marcos, San Marcos, Tex.; also known from the
 Guadalupe River near San Marcos and the Rio Comal at New
 Braunfels, Tex.
Hybopsis aestivalis marconis Jordan and Gilbert, Proc. U. S. Nat. Mus.,
 IX, 1886, 22, Rio San Marcos, San Marcos, Tex.

1036. *Macrhybopsis sterletus* (Cope).

Rio Grande.

Ceratichthys sterletus Cope, Wheeler Surv., V, Zool., 1875 (1876), 652,
Rio Grande, at San Ildefonso, New Mexico.

1037. *Macrhybopsis gelida* (Girard).

Middle Missouri River Basin, from Wyoming to eastern Nebraska.

Gobio gelidus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 188,
Milk River, Mont.

1038. *Macrhybopsis montana* (Meek).

Type supposed to be from the upper Missouri.

Hybopsis montanus Meek, Proc. U. S. Nat. Mus., VII, 1884, 526,
locality unknown but collected by Dr. F. V. Hayden, probably in
the upper Missouri Basin.

1039. *Macrhybopsis meeki* (Jordan and Evermann).

Missouri River at St. Joseph and elsewhere in the river channel.

Hybopsis meeki Jordan and Evermann, Fish. North and Mid. Amer.,
1896, 317, Missouri River at St. Joseph, Mo.

1040. *Macrhybopsis hyostoma* (Gilbert).

Indiana to Iowa, and south to the Alabama River.

Nocomis hyostomus Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 203,
White River, Bedford, Ind.

Genus 342. EXTRARIUS Jordan.

Extrarius Jordan, Proc. Acad. Nat. Sci. Phila., LXX, 1918, 342 (*Hy-
bopsis tetranemus* Gilbert).

1041. *Extrarius tetranemus* (Gilbert).

Tributaries of Arkansas River in Kansas and Arkansas.

Hybopsis tetranemus Gilbert, Bull. Washburn Lab. Nat. Hist., 1886,
208, Elm and Spring Creeks, Medicine Lodge, Kans.

Genus 343. RHINICHTHYS Agassiz.

Argyreus Heckel, Russiger's Reisen, 1842 (*Cyprinus atronasus* Mitchell;
name preoccupied).

Rhinichthys Agassiz, Lake Superior, 1850, 353 (*Cyprinus atronasus*
Mitchell).

**1042. *Rhinichthys cataractæ* (Cuvier and Valenciennes). *Long-nosed dace;*
*Black-nosed dace; Black minnow; Niagara gudgeon.***

New England to Virginia and Wisconsin.

Gobio cataractæ Cuvier and Valenciennes, Hist. Nat. Poiss., XVI, 1842,
240 (315), pl. 483, Niagara Falls.

Leuciscus nasutus Ayres, Jour. Bost. Soc. Nat. Hist., IV, 1843, 299,
West Hartford, Conn.

Rhinichthys marmoratus Agassiz, Lake Superior, 1850, 354, Pl. II, figs.
1 and 2, Sault Ste. Marie, Lake Superior.

1043. *Rhinichthys bowersi* Goldsborough and Clark.

Rhinichthys bowersi Goldsborough and Clark, Bull. U. S. Bur. Fish.,
XXVII, 1907, 36, Dry Fork, Herman, W. Va.

1044. *Rhinichthys dulcis* (Girard).

Upper Missouri, Platte, Arkansas, Rio Grande, and Colorado Rivers;
Columbia River; Utah Basin; and coastwise streams of Washington
and Oregon.

Argyreus dulcis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 185,
Sweetwater River, Nebr.

Rhinichthys maxillosus Cope, loc. cit., XVI, 1864, 278, Kansas.

Rhinichthys transmontanus Cope, Amer. Nat., 1879, 441, tributaries of
Rio Grande in Colorado and New Mexico.

Rhinichthys ocella Garman, Science Observer, III, 1881, 59, northeastern
Wyoming and Montana.

Rhinichthys luteus Garman, Bull. Mus. Comp. Zool., VIII, 1881, 87,
Bear River, Ogden, Utah.

- 1045. *Rhinichthys atronasus* (Mitchill).** *Dace; Black-nosed dace; Brook minnow; Striped dace.*
 New England to Minnesota, northern Alabama, and Virginia; very abundant in clear brooks and mountain streams.
Cyprinus atronasus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 460, Wallkill River; brooks of New York.
Cyprinus vittatus Rafinesque, Amer. Month. Mag., II, 1817, 121, upper Hudson River.
- 1046. *Rhinichthys croceus* (Storer).**
 Tennessee Basin; abundant in clear brooks.
Leuciscus croceus Storer, Proc. Bost. Soc. Nat. Hist., 1845, 48, Florence, Ala.
Rhinichthys obtusus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 357, Huntsville, Ala.
Rhinichthys badius Garman, Science Observer, III, 1881, 60, Clinch River, Tenn.
- 1047. *Rhinichthys lunatus* Cope.**
 Lakes and brooks of Michigan, Indiana, Wisconsin, and Minnesota, and in Rainy River.
Rhinichthys lunatus Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 278, Grosse Isle, Mich.
Rhinichthys arenatus Garman, Science Observer, 1881, 62, Sand Hill River, Minn.
- 1048. *Rhinichthys meleagris* Agassiz.**
 Illinois and Iowa.
Rhinichthys meleagris Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 357, Burlington, Iowa.
- 1049. *Rhinichthys simus* Garman.**
 Coahuila, Mexico.
Rhinichthys simus Garman, Science Observer, III, 1881, 61, Coahuila, Mexico.
- Genus 344. AGOSIA Girard.**
Agosia Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 186 (*Agosia chrysogaster* Girard).
Zophendum Jordan and Gilbert, Synopsis, 1883, 155 (*Hybognathus siderius* Cope = *Agosia chrysogaster* Girard).
- 1050. *Agosia chrysogaster* Girard.**
 Tributaries of the Gila River; common in the mountain brooks.
Agosia chrysogaster Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 187, Rio Santa Cruz, Sonora.
Agosia metallica Girard, loc. cit., 1856, 187, Rio San Pedro, Ariz.
Hyborhynchus (*Hybognathus*) *siderius* Cope, Wheeler Surv., V, Zool., 1875, 670, Pl. XXXI, figs. 6 and 6a, Camp Lowell, Ariz.
- Genus 345. APOCOPE Cope.**
Apocope Cope, Hayden's Geol. Surv. Mont., etc., 1871 (1872), 472 (*A. carringtoni* Cope).
Eritrema Cope, Wheeler's Surv., V, Zool., 1876, 645 (*Apocope henshavii* Cope).
- 1051. *Apocope nubila* (Girard).**
 Basin of the Columbia River from western Idaho, below Shoshone Falls of Snake River, to the coast, and in coastwise streams from Washington southward into Oregon.
Argyreus nubilus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 186, Fort Steilacoom, Puget Sound.
- 1052. *Apocope robusta* (Rutter).**
 Upper Sacramento Basin.
Agosia robusta Rutter, Bull. U. S. Fish Com., XXII, 1902, 148, Prosser Creek, Calif.

1053. *Apocope carringtonii* Cope.

Upper Snake River Basin to Heart Lake in Yellowstone Park, thence extending southward in the Great Basin and the Coast Ranges; southeastern Oregon.

Apocope carringtonii Cope, Hayden's Geol. Surv. Mont., 1871, 472, Warm Springs, Utah.

Apocope vulnerata Cope, loc. cit., 473, Logan, Utah.

Tigoma rhinichthyooides Cope, loc. cit., 473, Logan, Utah.

Rhinichthys henshawi Cope, Proc. Amer. Phil. Soc., XIV, 1874, 133, Provo, Utah.

Agosia novemradiata Cope, Proc. Acad. Nat. Sci. Phila., XXV, 1883, 141, Weber River at Echo, Utah.

1054. *Apocope oscula* (Girard).

Lower Colorado and Gila Rivers; not rare.

Argyreus osculus Girard, Proc. Acad. Nat. Sci. Phila., 1856, 186, Babocomori Creek, a tributary of the Rio San Pedro, Ariz.

Argyreus notabilis Girard, loc. cit., 1856, 186, Rio Santa Cruz, Sonora, a tributary of the Rio San Pedro, Ariz.

Ceratichthys ventricosus Cope, Proc. Amer. Phil. Soc., XXV, 1874, 136, Ariz.

1055. *Apocope couesii* Yarrow.

Colorado River Basin.

Apocope couesii Yarrow, Field and Forest, 1876, and Wheeler Surv., V, Zool., 1875, 648, Camp Apache, Ariz.

1056. *Apocope yarrowi* (Jordan and Evermann).

Streams of Colorado in the Colorado River Basin.

Agosia yarrowi Jordan and Evermann, Bull. U. S. Fish Com., IX, 1889 (1891), 28, Tomichi Creek and Gunnison River, Gunnison, Colo.

1057. *Apocope adobe* (Jordan and Evermann).

Sevier River, Utah; locally abundant.

Agosia adobe Jordan and Evermann, Bull. U. S. Fish Com., IX, 1889 (1891), Sevier River, Juab, Utah.

1058. *Apocope klamathensis* (Evermann and Meek).

Agosia klamathensis Evermann and Meek, Bull. U. S. Fish Com., XVII, 1897 (1898), 75, 9, fig. 5, Pelican Bay, Upper Klamath Lake, Oreg.

1059. *Apocope nevadensis* (Gilbert).

Warm springs in the deserts of southwestern Nevada; Ash Meadows, Indian Creek; and Vegas Creek, Nev.

Rhinichthys (Apocope) nevadensis Gilbert, Death Valley Exped., 1893, 230, pl. 6, fig. 1, Ash Meadows, Amargosa Desert, Nev.

1060. *Apocope velifera* (Gilbert).

Pahranagat Valley, southwestern Nevada.

Rhinichthys (Apocope) velifer Gilbert, Death Valley Exped., 1893, 229, pl. 6, fig. 2, Pahranagat Valley, Nev.

1061. *Apocope falcata* (Eigenmann and Eigenmann).

Columbia River Basin; locally abundant; Boise River at Caldwell, Idaho; Payette River at Payette, Idaho; Columbia River at Pasco and Umatilla; and Mill Creek at Walla Walla.

Agosia falcata Eigenmann and Eigenmann, Amer. Nat., XXVII, Feb. 4, 1893, 153, Boise River, Caldwell, Idaho.

1062. *Apocope umatilla* (Gilbert and Evermann).

Columbia River at Umatilla, Oreg., and Payette and Salmon Rivers, Idaho.

Agosia umatilla Gilbert and Evermann, Bull. U. S. Fish Com., XIV, 1894, 192, pl. 9, fig. 2, Columbia River, Umatilla, Oreg.

Genus 346. ERICYMBIA Cope.

Ericympba Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 88 (*E. buccata* Cope).

1063. *Ericymba buccata* Cope. *Silver-mouthed minnow.*
 Michigan and western Pennsylvania to Kansas and southward to western Florida.
Ericymba buccata Cope, Proc., Acad. Nat. Sci. Phila., XVII, 1865, 88, Kiskiminitas River, a tributary of the Monongahela, western Pennsylvania.
- Genus 347. **PHENACOBIUS** Cope.
Phenacobius Cope, Proc., Acad. Nat. Sci. Phila., XIX, 1867, 96 (*P. teretulus* Cope).
Sarcidium Cope, Hayden's Geol. Surv. Wyo., 1870, 363 (*S. scopiferum* Cope).
1064. *Phenacobius teretulus* Cope.
 Kanawha River, W. Va.
Phenacobius teretulus Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867, 96, Kanawha River, Eggleston Springs, W. Va.
1065. *Phenacobius mirabilis* (Girard). *Sucker-mouthed minnow.*
 Illinois to Arkansas.
Exoglossum mirabile Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 191, Arkansas River, Fort Smith, Ark.
1066. *Phenacobius scopifer* (Cope).
 Illinois to the Rio Grande.
Sarcidium scopiferum Cope, Hayden's Geol. Surv. Wyo., 1870 (1871), 440, Missouri River near St. Joseph, Mo.
Phenacobius teretulus iosternus Nelson, Bull. Ill. Lab. Nat. Hist., I, 1876, 46, brooks of McLean County, Ill.
1067. *Phenacobius uranops* Cope.
 Upper Tennessee Basin.
Phenacobius uranops Cope, Proc. Acad. Nat. Sci. Phila., XIX, 1867, 96, Holston River, Saltville, Va.
1068. *Phenacobius catostomus* Jordan.
 Alabama River Basin.
Phenacobius catostomus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 332, Silver Creek, Etowah River, Rome, Ga.
- Genus 348. **FALCULARIUS** Jordan and Snyder.
Falcula Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 124 (*Falcula chapalæ* Jordan and Snyder); name preoccupied.
Falcularius Jordan, Amer. Nat., XXXVII, 1903, 360 (*Falcula chapalæ* Jordan and Snyder).
1069. *Falcularius chapalæ* (Jordan and Snyder).
 Lake Chapala, Mexico.
Falcula chapalæ Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1901), 124, Lake Chapala at Ocotlan, Mexico.
- Genus 349. **XYSTROSUS** Jordan and Snyder.
Xystrosus Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1901), 123 (*X. popoche* Jordan and Snyder).
1070. *Xystrosus popoche* Jordan and Snyder.
 Basin of the Rio Lerma, Mexico.
Xystroesus popoche Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1901), 123, Lake Chapala at Ocotlan, Jalisco, Mexico.
- Genus 350. **EVARRA** Woolman.
Evarra Woolman, Bull. U. S. Fish Com., XIV, 1894, 64 (*E. eigenmanni* Woolman).
1071. *Evarra eigenmanni* Woolman.
 Central Mexico.
Evarra eigenmanni Woolman, Bull. U. S. Fish Com., XIV, 1894, 64, City of Mexico.
1072. *Evarra tlahuacensis* Meek.
 Chalco Lake, Tlahuac, Mexico.
Evarra tlahuacensis Meek, Field Col. Mus., Zool., III, 1902, 85, Pl. XVIII, lower figure, Chalco Lake, Tlahuac, Mexico.

Genus 351. TIAROGA Girard.

Tiaroga Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 204 (*T. cobitis* Girard).

1073. **Tiaroga cobitis** Girard.

Gila Basin, Ariz.

Tiaroga cobitis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 204, Rio San Pedro, Ariz.

Genus 352. AZTECULA Jordan and Evermann.

Azteca Jordan and Evermann, Fishes North and Mid. Amer., 1896, 254 (*Notropis aztecus* Woolman); name preoccupied.

Aztecula Jordan and Evermann, loc. cit., 1898, 2799, (*Notropis aztecus* Woolman = *Codoma vittata* Girard).

1074. **Aztecula vittata** (Girard).

Valley of Mexico and headwaters of the Rio Balsas at Pueblo, Mexico.

Codoma vittata Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 195, Valley of Mexico.

Notropis aztecus Woolman, Bull. U. S. Fish Com., XIV, 1894, 63, pl. 2, canals about City of Mexico.

1075. **Aztecula mexicana** Meek.

Basin of Rio San Juan on the Mexican plateau.

Aztecula mexicana Meek, Field Col. Mus., Zool., III, 1902, 81, figured; loc. cit., V, 1904, 62, fig. 15; San Juan del Rio, Mexico.

1076. **Aztecula lermæ** (Evermann and Goldsborough).

Basin of Rio Lerma.

• *Notropis lermæ* Evermann and Goldsborough, Bull. U. S. Fish. Com., XXI, 1901 (1902), 147, fig. 3, Lake Lerma, at Lerma, Mexico.

Genus 353. CODOMA Girard.

Codoma Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 194 (*C. ornata* Girard).

1077. **Codoma ornata** Girard.

Headwaters of the Rio Nezquital and Rio Nazas in Durango to upper tributaries of the Rio Yaqui and Rio Conchos in Chihuahua, Mexico.

Codoma ornata Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 195, Chihuahua River, Mexico.

1078. **Codoma topeka** (Gilbert).

Western Iowa and eastern South Dakota to Kansas; locally abundant in the prairie streams.

Notropis topeka Gilbert, Bull. Washburn Lab. Nat. Hist., I, 1884, 13, tributary of Kansas River, Topeka, Kans.

Notropis æneolus Hay, Proc. U. S. Nat. Mus., X, 1887, 245, Solomon River, Wakeeney, Kans.

Genus 354. HYBOGNATHUS Agassiz.

Hybognathus Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 223 (*H. nuchalis* Agassiz).

1079. **Hybognathus nuchalis** Agassiz. *Gudgeon; Smelt; Silvery minnow; Smelt minnow.*

Clear streams from the upper Missouri southward to Georgia and Texas, west of the Alleghenies.

Hybognathus nuchalis Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 224, Quincy, Ill.

Hybognathus evansi Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 182, Fort Pierre, Nebr.

Hybognathus placitus Girard, loc. cit., 1856, 182, sluices of the Arkansas River.

1080. **Hybognathus regius** Girard. *Gudgeon; Silvery minnow.*

Streams New Jersey to Virginia, east of the Alleghenies.

Hybognathus regius Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 209, Potomac River.

Hybognathus osmerinus Cope, Proc. Amer. Phil. Soc., XI, 1870, 466, Raritan River, N. J.

1081. *Hybognathus argyritis* Girard.

Upper Missouri region and Red River of the North.

Hybognathus argyritis Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 182, Milk River, Mont.

1082. *Hybognathus hayi* Jordan.

Lower Mississippi Valley and neighboring rivers in Mississippi.

Hybognathus hayi Jordan, Proc. U. S. Nat. Mus., VII, 1884, 548, Pearl River, Jackson, Miss.; this and three preceding are all very close to *Hybognathus nuchalis*.

Genus 355. *DIONDA* Girard.

Dionda Girard, Proc. Acad. Nat. Sci. Phila., VIII, 176 (*D. episcopa* Girard).

Algoma Girard, loc. cit., 180 (*A. amara* Girard).

1083. *Dionda episcopa* Girard.

Southern Texas and northern Mexico; south to the Rio Mezquital in Durango and the Rio San Juan in Nuevo Leon.

Dionda episcopa Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 177, headwaters of Pecos River and Comanche Spring, tributary to the Rio Grande, Tex.

Dionda terrensis Girard, loc. cit., 177, Nueces River, Tex.

Dionda papalis Girard, loc. cit., 178, Delaware Creek, a tributary of Pecos River, Tex.

Dionda argentosa Girard, loc. cit., 178, San Felipe Creek and Devils River, tributaries of Rio Grande.

Dionda chrysitis Girard, loc. cit., 178, Live Oak Creek, a tributary of the Pecos.

Dionda couchi Girard, loc. cit., 178, Guajuco, Monterey, and Cadereita, Nuevo Leon.

Hybognathus flavipinnis Cope, Bull. U. S. Nat. Mus., XVII, 1880, 36, Johnsons Fork of Llano River, Tex.

Hybognathus nigroteniata Cope, loc. cit., 37, Wallace Creek, a tributary of Medina River, Tex.

Hybognathus (Dionda) punctifer Garman, Bull. Mus. Comp. Zool., VIII, 1881, 89, Parras and spring near Saltillo, Mexico.

Hybognathus civilis Cope, Trans. Amer. Phil. Soc., 1884, 167, Monterey, Nuevo Leon.

1084. *Dionda serena* Girard.

Sabinal River, Tex.

Dionda serena Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 177, Sabinal River, Tex.

1085. *Dionda amara* (Girard)

Near Fort Brown, Tex.

Algoma amara Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 180, Lagoon near Fort Brown, Tex.

1086. *Dionda melanops* Girard.

Coahuila, Mexico.

Dionda melanops Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 178, Buena Vista, Coahuila.

1087. *Dionda nubila* (Forbes).

Western Illinois, west to Wyoming and southwest to the Ozark region. *Alburnops nubilus* Forbes, Bull. Ill. Lab. Nat. Hist., I, 1878, 56, Rock River, Ogle County, Ill.

1088. *Dionda plumbea* Girard.

Tributaries of Arkansas River, in Oklahoma.

Dionda plumbea Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 178, Canadian River, Okla.

Dionda spadicea Girard, loc. cit., 178, Fort Smith, Ark.

Dionda grisea Girard, Pac. R. R. Surv., X, 1858, 230, Pl. LII, figs. 6-10, near Choctaw Agency.

Genus 356. *CERATICHTHYS* Baird and Girard.

Ceraticthys Baird and Girard, Proc., Acad. Nat. Sci. Phila., VI, 1853, 391 (*C. vigilax*; misprinted *Ceraticthys*, not of Girard, 1856).

Cliola Girard, loc. cit., VIII, 1856, 192 (*Ceraticthys vigilax* Baird and Girard).

Hypargyrus Forbes, in Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 200 (*Hybopsis tuditanus* Cope).

- 1089. *Ceratichthys vigilax* Baird and Girard.** *Bullhead minnow; Fathead minnow.*
 Ohio to Georgia, Iowa, and Texas.
Ceratichthys vigilax Baird and Girard, Proc., Acad. Nat. Sci. Phila., VIII, 1853, 391, Otter Creek, North fork of Red River, Ark.
Cliola velox Girard, loc. cit., VIII, 1856, 192, San Pedro Creek and
Cliola vivax, Rio Leon, tributaries of Rio San Antonio, Tex.
Hybopsis tuditanus Cope, Trans., Amer. Phil. Soc., XIII, 1866, 381, Detroit, Mich.
Alburnops taurcephalus Hay, Proc. U. S. Nat. Mus., 1880, 503, Chic-kasawha River, Miss.
- 1090. *Ceratichthys smithi* (Evermann and Cox).**
 Southern South Dakota; known only from Prairie Creek near Scotland.
Cliola smithii Evermann and Cox, Rept. U. S. Fish Com., XIX, 1894 (1896), 400, Prairie Creek, Scotland, S. Dak.
- Genus 357. *HYBORHYNCHUS* Agassiz.**
Hyborhynchus Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 223 (*Minnilus notatus* Rafinesque).
- 1091. *Hyborhynchus notatus* (Rafinesque).** *Blunt-nosed minnow; Spotted minnow.*
 Quebec to Delaware; Kentucky, Alabama, Arkansas, and northward to the Dakotas; generally very abundant in small streams west of the Alleghenies.
Minnilus notatus Rafinesque, Ichth. Ohiensis, 1820, 47, Ohio River.
Hyborhynchus perspicuus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 179, Arkansas River, Fort Smith.
? *Hyborhynchus tenellus* Girard, loc. cit., 179, Choctaw Agency.
Hyborhynchus puniceus Girard, loc. cit., 179, Antelope Creek, Canadian River.
Hybopsis haematurus Cope, Trans. Amer. Phil. Soc., XIII, 1866, 382, St. Joseph River, Michigan City, Ind.
? *Hyborhynchus superciliosus* Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 234, Kanawha River.
- Genus 358. *PIMEPHALES* Rafinesque.** *Fatheads.*
Pimephales Rafinesque, Ichth. Ohiensis, 1820, 52 (*P. promelas* Rafinesque).
Coliscus Cope, Hayden's Geol. Surv. Wyo., 1872, 437 (*Coliscus parietalis* Cope; young of *Pimephales promelas*).
Spinicephalus LeSueur, Bull. Sci. Philom., VIII, 1896, 14, pl. 26 (*S. fibulatus* LeSueur = *Pimephales promelas* Rafinesque).
- 1092. *Pimephales promelas* Rafinesque.** *Fathead minnow; Blackhead minnow.*
 Lake Champlain to Kentucky, Dakota, and the Rio Grande; abundant in sluggish brooks, especially in the Missouri Basin.
Pimephales promelas Rafinesque, Ichth. Ohiensis, 1820, 53, pond near Lexington, Ky.
Pimephales maculosus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 180, sluices of the Arkansas River, Fort Mabee, Ark.
Pimephales fasciatus Girard, loc. cit., 180, Yellowstone River.
Plargyrus melanocephalus Abbott, loc. cit., XII, 1860, 325, Lake Whittlesey, Minn.
Pimephales milesi Cope, loc. cit., XVI, 1864, 282, Grosse Isle, Detroit River, Mich.
Pimephales agassizii Cope, Trans. Amer. Phil. Soc., XIII, 1866, 394, tributary of Ohio River, near Richmond, Ind.
Coliscus parietalis Cope, Hayden's Geol. Surv. Wyo., 1871, 437, St. Joseph, Mo.; young.
Hyborhynchus nigrellus Cope, Wheeler's Surv., V, Zool., 1875, 671, Arkansas River.
Spinicephalus fibulatus LeSueur, Bull. Soc. Philom., VIII, 1896, pl. 26; no locality.
- 1093. *Pimephales confertus* (Girard).**
 Tributaries of the Rio Grande.
Hyborhynchus confertus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 179, Hurrah Creek, a tributary of the Pecos River, Tex.

1094. **Pimephales anuli** Kendall.

Thoroughfare between Mud and Grass Lakes, Me.

Pimephales anuli Kendall, Bull. U. S. Fish Com., XXII, 1902, 257,
between Mud Lake and Grass Lake, Me.

Genus 359. **COCHLOGNATHUS** Baird and Girard.

Cochlognathus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII,
1854, 158 (*C. ornatus* Baird and Girard).

1095. **Cochlognathus ornatus** Baird and Girard.

Rio Grande Basin.

Cochlognathus ornatus Baird and Girard, Proc. Acad. Nat. Sci. Phila.,
VII, 1854, 158, Brownsville, Tex.

1096. **Cochlognathus biguttatus** Cope.

Trinity River, Tex.

Cochlognathus biguttatus Cope, Bull. U. S. Nat. Mus., XVII, 1880, 37,
Trinity River, Fort Worth, Tex.

Genus 360. **EXOGLOSSUM** Rafinesque.

Exoglossum Rafinesque, Jour. Acad. Nat. Sci. Phila., I, 1818, 419
(*E. lesueurianum* Rafinesque = *Cyprinus maxillingua* LeSueur).

Maxillingua Rafinesque, loc. cit., 421 (*Cyprinus maxillingua* LeSueur).

1097. **Exoglossum maxillingua** (LeSueur). *Cut-lips; Nigger chub; Nigger
dick; Butter chub; Niggerfish.*

Lake Ontario, St. Lawrence River, Lake Champlain, Hudson River,
and Cayuga Lake, and southward to Virginia; abundant in the
basins of the Susquehanna, Hudson, Potomac, James, Roanoke, and
Kanawha, but not widely distributed.

Cyprinus maxillingua LeSueur. Jour. Acad. Nat. Sci. Phila., I, 1817, 85,
Pipe Creek, Md.

Exoglossum annulatum Rafinesque, loc. cit., 1818, 421, Hudson River.

Exoglossum nigrescens Rafinesque, loc. cit., 422, Hudson River.

Exoglossum lesurianum Rafinesque, loc. cit., 420; after LeSueur.

Genus 361. **CAMPOSTOMA** Agassiz. *Stonerollers.*

Campostoma Agassiz, Amer. Jour. Sci. Arts, XIX, 1855, 218 (*Rutilus
anomalus* Rafinesque).

1098. **Campostoma anomalum** (Rafinesque). *Stoneroller; Greased chub;
Blue sucker; Rotgut minnow; Mammy; Doughbelly; Steel-backed chub.*

Central New York to Tennessee, Wyoming, and Texas.

Rutilus anomalus Rafinesque, Ichth. Ohiensis, 1820, 52, Licking River,
Ky.

Exoglossum spiniceps Cuvier and Valenciennes, Hist. Nat. Poiss.,
XVII, 1844, 363 (489), Wabash River.

Leuciscus prolrixus Storer, Proc. Bost. Soc. Nat. Hist., II, 1845, 48,
Alabama.

Exoglossum dubium Kirtland, Bost. Jour. Nat. Hist., V, 1845, 272,
Yellow Creek, Ohio.

Chondrostoma pullum Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 357,
Burlington, Iowa.

Campostoma nasutum Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856,
176, Cadereita, and near Monterey, Nuevo Leon, Mexico.

Campostoma callipteryx Cope, loc. cit., XVI, 1864, 284, Flint River,
Mich.

Campostoma mormyrus Cope, loc. cit., 284, Bruce, Macomb County, Mich.
Campostoma gobionium Cope, loc. cit., 284, Bruce, Macomb County, and

Grosse Isle, Mich.

Campostoma hippops Cope, loc. cit., 284, Platte River, Nebr.

Campostoma brevis Haseman, Proc. Ind. Acad. Sci., 1905 (1906), 161,
Wabash River near Wabash, Ind.

1099. **Campostoma ornatum** Girard.

Chihuahua River, and tributaries of the Rio Grande in Mexico.

Campostoma ornatum Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856,
176, Chihuahua, Mexico.

1100. *Campostoma pricei* Jordan and Thoburn.
 Rio Yaqui, Sonora and neighboring streams, tributary to the Gulf of California.
Campostoma pricei Jordan and Thoburn, in Jordan and Evermann, Fishes North and Mid. Amer., 1896, 203; Rucker Canyon, Chiricahua Mountains, Ariz., very near *C. ornatum*.

1101. *Campostoma formosulum* Girard.
 Texas; San Antonio River to the Rio Grande.
Campostoma formosulum Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 176, Rio Sabinal, near San Antonio, Tex.

Family 91. MEDIDÆ

- Genus 362. *LEPIDOMEDA* Cope.
Lepidomeda Cope, Proc. Amer. Phil. Soc. Phila., XIV, 1874, 131 (*L. vittata* Cope).
1102. *Lepidomeda vittata* Cope.
 Known only from the Colorado Chiquito River, Ariz., and from Pahrang-agat Valley, Nev.
Lepidomeda vittata Cope, Proc. Amer. Phil. Soc. Phila., XIV, 1874, 131, Rio Colorado Chiquito, Ariz.
1103. *Lepidomeda jarrovii* Cope.
 Colorado Chiquito River, Ariz.; also in southern Nevada, in springs in the desert.
Lepidomeda jarrovii Cope, Proc. Amer. Phil. Soc. Phila., XIV, 1874, 132, Rio Colorado Chiquito, Ariz.
- Genus 363. *MEDA* Girard.
Meda Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 191 (*M. fulgida* Girard).
1104. *Meda fulgida* Girard.
 Rio Gila Basin, Ariz.
Meda fulgida Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 192, Rio San Pedro, Ariz.
- Genus 364. *PLAGOPTERUS* Cope.
Plagopterus Cope, Proc. Amer. Phil. Soc., XIV, 1874, 130 (*P. argentissimus* Cope).
1105. *Plagopterus argentissimus* Cope.
 Colorado Basin in western Colorado; Fort Yuma.
Plagopterus argentissimus Cope, Proc. Amer. Phil. Soc., XIV, 1874, 130, San Luis Valley, Colo.

Order Y. NEMATOGNATHI.²⁸

Family 92. ARIIDÆ. Sea catfishes

- Genus 365. *BAGRE* Oken. *Gafftopsail catfishes*.
Les Bagres Cuvier, Règne Anim., II, 1817 (*Silurus bagre* L.).
Bagre Oken, Isis, 1817, 1183 (after les Bagres of Cuvier; type by tautonomy *Silurus bagre* Linnaeus).
Breviceps Swainson, Nat. Hist. Anim., I, 1838, 328 (*Breviceps filamentosus* Swainson); name preoccupied.
Felichthys Swainson, loc. cit., II, 1839, 305 (*Breviceps filamentosus* Swainson); substitute for *Breviceps*.
Ailurichthys Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 26 (*Silurus marinus* Mitchell).
Ælurichthys, corrected spelling.

²⁸ Until recently all members of this order have been usually included in the single family *Siluridæ*. As properly restricted the *Siluridæ* is a small group of large river fishes of Europe and Asia, resembling our larger catfishes, but without adipose fin, the anal joined to the caudal.

1106. *Bagre marina* (Mitchill). *Sea catfish; Gafflopsail.*

Cape Cod to Texas.

Silurus marinus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 433, New York.

In the investigation of Linnaean types at Upsala Dr. Einar Lönnberg has found an example of this species labeled *Silurus felis*. On the strength of this discovery this species has been called *Ælurichthys felis*. But Linnaeus's own description shows clearly that this was not his type, which must have been our common sea cat here called *Galeichthys felis*.

1107. *Bagre isthmensis* (Regan).

Atlantic coast of Panama; Colon.

Ælurichthys isthmensis Regan, Biol. Centrali-Amer., 1907, 117, Tab. XV, fig. 1, and Tab. XIX, fig. 1, Atlantic coast of Panama, Colon.

1108. *Bagre panamensis* (Gill).

Mazatlan to Panama.

Ælurichthys panamensis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 172, Panama.

1109. *Bagre pinnimaculatus* (Steindachner).

Mazatlan to Panama.

Ailurichthys pinnimaculatus Steindachner, Ichth. Beitr., IV, 1875, 15, Panama, Altata, Costa Rica.

1110. *Bagre eydouxii* (Cuvier and Valenciennes).

Guayaquil.

Galeichthys eydouxii Cuvier and Valenciennes, Hist. Nat. Poiss., XV, 1840, 43, Guayaquil.

1111. *Bagre scutatus* (Regan).

Coast of Panama and Ecuador.

Ælurichthys scutatus Regan, Biol. Centrali-Amer., 1907, 116, Tab. XV, fig. 2, and Tab. XIX, fig. 2, Pacific coast of Panama.

1112. *Bagre bahiensis* (Castelnau).

Mexico to Bahia.

Galeichthys bahiensis Castelnau, Anim. Amér. Sud, 1855, 37, Bahia.

Ælurichthys longispinis Günther, Cat., V, 1864, 178; Mexico; South America.

Genus 366. SCIADEICHTHYS Bleeker.

Sciadeichthys Bleeker, Act. Soc. Sci. Indo-Neerl., V, 1858, 99 (*Sciades pictus* Müller and Troschel); not *Sciadeichthys* Bleeker, 1863.

1113. *Sciadeichthys troscheli* (Gill).

Gulf of California to Panama.

Sciades troscheli Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 171, Panama.

Arius brandti Steindachner, Ichth. Beitr., IV, 1876, 21, Altata, Panama.

1114. *Sciadeichthys emphysetus* (Müller and Troschel).

Surinam.

Bagrus (*Sciades*) *emphysetus* Müller and Troschel, Horæ Ichth., III, 1849, 8, Surinam.

1115. *Sciadeichthys proops* (Cuvier and Valenciennes).

West Indies from Porto Rico to Surinam and south to Pernambuco.

Bagrus proops Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 457, Guiana; Porto Rico.

Genus 367. SELENASPIS Bleeker.

Selenaspis Bleeker, Vissch. Ind. Archipel., I, 1858, 62 (*Silurus hertzbergi* Bloch).

Leptarius Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 170 (*L. dowii* Gill).

1116. *Selenaspis dowii* (Gill).

Panama to Guayaquil.

Leptarius dowii Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 170, Panama.

Arius alatus Steindachner, Ichth. Beitr., IV, 1876, 19, Panama.

Genus 368. NETUMA Bleeker.

Netuma Bleeker, Vissch. Ind. Archipel., I, 1858, 62 (*N. thalassina*, Bleeker = *N. nasuta* Bleeker).
Guirilinga Bleeker, loc. cit., 62 (*Arius commersonii* Bleeker).
Notarius Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 171 (*Arius grandicassis* Cuvier and Valenciennes).

1117. Netuma planiceps (Steindachner).

Gulf of California to Panama; not rare.
Arius planiceps Steindachner, Ichth. Beitr., IV, 1876, 26, Altata, Panama.

1118. Netuma kessleri (Steindachner).

Gulf of California to Panama, not rare.
Arius kessleri Steindachner, Ichth. Beitr., IV, 1876, 24, Altata, Panama.

1119. Netuma insculpta (Jordan and Gilbert).

Panama.

Arius insculturatus Jordan and Gilbert, Bull. U. S. Fish Com., II, 1882 (1883), 41, Panama.

1120. Netuma platypogon (Günther).

Gulf of California to Peru; generally common.
Arius platypogon Günther, Cat., V, 1864, 147, San Jose de Guatemala.

1121. Netuma oscula (Jordan and Gilbert).

Panama; a single example known.

Arius osculus Jordan and Gilbert, Bull. U. S. Fish Com., II, 1882 (1883), 46, Panama.

1122. Netuma elattura (Jordan and Gilbert).

Panama; scarce.

Arius elaturus Jordan and Gilbert, Bull. U. S. Fish Com., II, 1882 (1883), 45, Panama.

Genus 369. GALEICHTHYS Cuvier and Valenciennes. Sea catfish.

Galeichthys Cuvier and Valenciennes, Hist. Nat. Poiss., XV, 1840, 28 (*G. feliceps* Cuvier and Valenciennes).

Hexanemeticthys Bleeker, Act. Soc. Sci. Indo-Neerl., V, 1858, 2 (*Bagrus sunducus* Bleeker).

Cephalocassis Bleeker, Vissch. Ind. Archipel., I, 1858, 62 (*Arius truncatus* Cuvier and Valenciennes).

Ariopsis Gill, Suppl. Proc. Acad. Nat. Sci. Phila., XIII, 1861, 56 (*Arius milberti* = *Silurus belis* L.).

Hemiarius Bleeker, Nederl. Tijdschr. Dierk., I, 1863, 7 (*Cephalocassis stormi* Bleeker).

Subgenus GALEICHTHYS ²⁹ Cuvier and Valenciennes.**1123. Galeichthys lentiginosus (Eigenmann and Eigenmann).**

Panama.

Tachisurus lentiginosus Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888, 139, Panama.

1124. Galeichthys peruvianus Lütken.

Pacific coast of Mexico to Peru.

Galeichthys peruvianus Lütken, Vid. Med. Naturh. Foren. Kjøbenhavn, 1874, 204, Callao, Peru.

Subgenus HEXANEMATICHTHYS ²⁰ Bleeker.**1125. Galeichthys felis (Linnæus). Sea cat.**

Cape Cod to Texas.

Silurus felis Linnæus, Syst. Nat., ed. XII, 1766, 503, Charleston, S. C.

Arius milberti Cuvier and Valenciennes, Hist. Nat. Poiss., XV, 1840, 74, New York; Charleston, S. C.

?*Arius equestris* Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1855, 26, Indianola, Tex.

1126. Galeichthys guentheri Regan.

Gulf of Mexico.

Galichthys guentheri Regan, Biol. Centrali-Amer., 1907, 124, Tab. XVI, fig. 1, and Tab. XIX, fig. 3, Gulf of Mexico.

* Top of head covered with soft skin, hiding the rugose plates.

** Top of head with thin skin which does not hide the rugose plates.

- 1127.** *Galeichthys surinamensis* (Bleeker).
Surinam.
Hexanematicthys surinamensis Bleeker, Versl. Akad. Amsterdam, 1862, 380, Surinam.
- 1128.** *Galeichthys seemani* (Günther).
Panama.
Arius seemani Günther, Cat., V, 1864, 147, "Central America."
- 1129.** *Galeichthys gilberti* Jordan and Williams.
Coast of Sinaloa.
Galeichthys gilberti Jordan and Williams, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 395, pl. 26, Mazatlan, Mexico.
- 1130.** *Galeichthys jordani* (Eigenmann and Eigenmann).
Panama.
Tachisurus jordani Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888, 142, Panama.
- 1131.** *Galeichthys cærulescens* (Günther).
West coast of Guatemala.
Arius cærulescens Günther, Cat., V, 1864, 149, Rio Huamuchal, Guatemala.
- 1132.** *Galeichthys azureus* Jordan and Williams.
Mazatlan, Mexico.
Galeichthys azureus Jordan and Williams, Proc. Calif. Acad. Sci., 2d ser., IV, 1895, 398, pl. 27, Mazatlan, Mexico.
- 1133.** *Galeichthys xenauchen* Gilbert.
Panama.
Galeichthys xenauchen Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2777, Panama.
- 1134.** *Galeichthys guatemalensis* (Günther).
Mazatlan to Central America.
Arius guatemalensis Günther, Cat., V, 1864, 145, Guatemala; Chiapas.
- 1135.** *Galeichthys assimilis* (Günther).
Atlantic coast of Central America, ascending streams.
Arius assimilis Günther, Cat., V, 1864, 146, Lake Yzabel, Guatemala.
- 1136.** *Galeichthys dasycephalus* (Günther).
Panama (wrongly ascribed to the Hawaiian Islands).
Arius dasycephalus Günther, Cat., V, 1864, 157, Oahu; apparently an error.
- 1137.** *Galeichthys longicephalus* (Eigenmann and Eigenmann).
Panama.
Tachisurus longicephalus Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888, 143, Panama.
- Genus 370. ARIUS** Cuvier and Valenciennes.
? *Tachysurus* Lacépède, Hist. Nat. Poiss., V, 1803, 151 (*T. chinensis* Lacépède; from a Chinese drawing, not identifiable as to species; may be a fresh-water fish or a *Netuma*).
Arius Cuvier and Valenciennes, Hist. Nat. Poiss., XV, 1840, 52 (*A. arius* Cuvier and Valenciennes, type by autonomy).
Ariodes Müller and Troschel, Horæ Ichth., III, 1849, 9 (*Arius arenarius* Müller and Troschel).
Pseudarius Bleeker, Nederl. Tijdschr. Dierk., I, 1862, 91 (*Arius arius* Cuvier and Valenciennes).
- 1138.** *Arius spixii* (Agassiz).
Trinidad to Brazil.
Pimelodus albifrons Spix, in Agassiz, Gen. Spec. Pisc. Brasil., 1829, 19, equatorial Brazil; name preoccupied.
Pimelodus spixii Agassiz, loc. cit., 19; after Spix.
Arius arenatus Cuvier and Valenciennes, Hist. Nat. Poiss., XV, 1840, 106, Cayenne.
Arius laticeps Günther, Cat., V, 1864, 171, British Guiana, Trinidad.
- 1139.** *Arius melanopus* Günther.
Pacific coast of Central America.
Arius melanopus Günther, Cat., V, 1864, 172, Rio Motagua.

1140. *Arius furthi* Steindachner.
 Panama.
Arius furthii Steindachner, Ichth. Beitr., IV, 1876, 29, Panama.
1141. *Arius steindachneri* (Gilbert and Starks).
 Pacific coast of Panama.
Tachysurus steindachneri Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 29, Pl. V, fig. 9, Panama Bay.
1142. *Arius evermanni* (Gilbert and Starks).
 Panama.
Tachysurus evermanni Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 32, Pl. V, fig. 10, Panama.
1143. *Arius liropus* (Bristol).
 Coast of Sonora, Mexico.
Tachysurus liropus Susan Brown Bristol, in Gilbert, Proc. U. S. Nat. Mus., XIX, 1896 (Feb. 5, 1897), 438, San Juan Lagoon, near mouth of Rio Ahome, Sonora, Mexico.
1144. *Arius emmelane* (Gilbert).
 Panama.
 • *Tachysurus emmelane* Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2785, Panama.
1145. *Arius taylori* Hildebrand. *Bagre*.
 Coast of El Salvador.
Arius taylori Hildebrand, Bull. U. S. Bur. Fish., XLI, 1925, 250, fig. 10, Rio Lempa, San Marcos.
1146. *Arius multiradiatus* Günther.
 Streams of Isthmus of Panama, Rio Bayano, Panama.
Arius multiradiatus Günther, Cat., V, 1864, 173, Rio Bayano, Panama.
- Genus 371. **CATHOROPS** Jordan and Gilbert.
Cathorops Jordan and Gilbert, Bull. U. S. Fish Com., II, 1882, 54 (*Arius hypophthalmus* Steindachner).
1147. *Cathorops hypophthalma* (Steindachner).
 Panama; rather scarce.
Arius hypophthalmus Steindachner, Ichth. Beitr., IV, 1875, 31, Pl. X, Panama.
1148. *Cathorops gulosa* (Eigenmann and Eigenmann).
 Panama.
Tachisurus gulosus Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d. ser., I, 1888, 146, Panama.
- Family 93. AMEIURIDÆ. Horned pouts; Channel cats
- Genus 372. **ISTLARIUS** Jordan and Snyder.
Istlaricus Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 118 (*I. balsanus* Jordan and Snyder).
1149. *Istlaricus balsanus* Jordan and Snyder.
 Balsas River Basin, Pacific slope of Mexico.
Istlaricus balsanus Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 118; tributary of Rio de las Balsas at Puente de Ixtla, Guerrero, Mexico (Pacific drainage).
- Genus 373. **ICTALURUS** Rafinesque. *Channel cats*.
Ictalurus Rafinesque, Ichth. Ohiensis, 1820, 61 (*Silurus punctatus* Rafinesque).
Elliops Rafinesque, loc. cit., 62 (*Pimelodus maculatus* Rafinesque).
Synechoglanis Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 39 (*S. beadlei* Gill = *Ictalurus furcatus* Rafinesque).
Ichthyælurus, corrected spelling.

- 1150. *Ictalurus furcatus*** (Cuvier and Valenciennes). *Blue cat; Great fork-tailed cat; Mississippi cat; Poisson bleu.*
Pimelodus furcatus LeSueur, in Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1840, 136, New Orleans.
Pimelodus affinis Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 26, Rio Grande.
Ameiurus ponderosus Bean, Proc. U. S. Nat. Mus., II, 1879, 286, St. Louis.
Ictalurus robustus Jordan, Bull. U. S. Nat. Mus., X, 1877, 76, Cairo, Ill.
- 1151. *Ictalurus anguilla*** Evermann and Kendall. *Eel cat; Willow cat; Fulton cat.*
 Mississippi River, occasionally as far north as Louisville, Ky., Muscatine and Fairport, Iowa, where it is known as the "Fulton cat."
Ictalurus anguilla Evermann and Kendall, Bull. U. S. Fish Com., XVII, 1897 (Feb. 9, 1898), 125, pl. 6, fig. 1, Atchafalaya River, La.
- 1152. *Ictalurus punctatus*** (Rafinesque). *Channel cat; White cat; Fiddler.*
 Rivers of the Great Lakes region, the Mississippi Valley, and streams tributary to the Gulf of Mexico.
Silurus punctatus Rafinesque, Amer. Month. Mag., 1818, 359, Ohio River.
Pimelodus caudafurcatus LeSueur, Mem. Mus., V, 1819, 152, Wabash River; Mississippi River.
Pimelodus maculatus, pallidus, argentinus, argyrus, with varieties *erythroptera, marginatus, lateralis, leucoplera* Rafinesque, Quarterly Jour. Sci., London, 1820, and Ichth. Ohiensis, 1820, 62, etc., Ohio River.
Pimelodus furcifer Cuvier and Valenciennes, Hist. Nat. Poiss., XV, 1840, 139, Surinam, by error.
Pimelodus gracilis Hough, Ann. Rept. Regent's, Albany, 1852, 26, Somerville, N. Y.
Pimelodus vulpes Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 170, rivers of Texas.
Synechoglanis beadlei Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1859, 40, St. Catherines, Ont.
Pimelodus houghi Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 159, Somerville; Ogdensburg; Foxburg, Pa.
Pimelodus megalops Girard, loc. cit., 161, Black Warrior River, Bigsby River, Ala.
Pimelodus graciosus Girard, loc. cit., 161, Prairie Mer Rouge, La.
Pimelodus hammondi and *notatus* Abbott, loc. cit., XII, 1860, 568, 569, Fort Riley, Kans.
Ictalurus simpsoni Gill, Proc. Bost. Soc. Nat. Hist., VIII, 1862, 43, Platte River.
- 1153. *Ictalurus meridionalis*** (Günther).
 Rio Usumacinta, Guatemala.
Amiurus meridionalis Günther, Cat., V, 1864, 102, and in Trans. Zool. Soc. Lond., 1868, 473, Rio Usumacinta, Guatemala.
- Genus 374. HAUSTOR** Jordan and Evermann.
Haustor Jordan and Evermann, Fishes North and Mid. Amer., 1896, 137 (*Gadus lacustris* Walbaum = *Pimelodus nigricans* LeSueur).
Villarius Rutter, Proc. Calif. Acad. Sci., VI, Ser. 2, June 23, 1896, 258 (*V. pricei* Rutter).
- 1154. *Haustor lacustris*** (Walbaum).
 Saskatchewan River and the Great Lakes.
Gadus lacustris Walbaum, Artedi Pisc., 1792, 144, Arctic America; based on the mathemeg or land cod of Pennant.
Pimelodus nigricans LeSueur, Mem. Mus., V, 1819, 153, Lake Erie, Lake Ontario.
Pimelodus borealis Richardson, Fauna Bor.-Amer., III, 1836, 135, Pine Island Lake.
- 1155. *Haustor okeechobensis*** (Heilprin).
 Florida.
Ictalurus okeechobensis Heilprin, Trans. Wagner Inst. Sci. Phila., I, 1887, pl. 18, Kissimee River, Fla.
Ameiurus lacustris okeechobensis Fowler, Proc. Acad. Nat. Sci. Phila., LI, 1899, 480; same type.

1156. *Haustor catus* (Linnaeus). *White cat; Channel cat of the Potomac.*

Silurus catus Linnaeus, Syst. Nat., ed. X, 1758, 305; based on the catfish of Catesby (*Bagre secundæ speciei marcgravei affinis*), which is a rude figure of the adult or "lophius" form of this species; northern part of America.

Pimelodus albifrons LeSueur, Mem. Mus., V, 1819, 148, Delaware River.
Pimelodus lynx Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 160, Potomac River.

Amiurus niveiventris Cope, Proc. Amer. Phil. Soc., XI, 1870, 486, Neuse River.

Ictalurus macaskeyi and *I. kevinskii* Stauffer, Hist. Lancaster County, Pa., 1869, 578, Lancaster County, Pa.

1157. *Haustor lophius* (Cope). *Big-mouthed cat.*

Potomac River.

Amiurus lophius Cope, Proc. Amer. Phil. Soc., XI, 1870, 486, Potomac River; not observed elsewhere. Known by its very wide flat head; apparently not the adult of *Ameiurus catus*, as has been supposed.

1158. *Haustor lupus* (Girard).

Rio Neuces and Rio Pecos, Tex.

Pimelodus lupus Girard, Pac. R. R. Surv., X, 1858, Pt. 4, Fishes, 211, Rio Pecos, Tex.

1159. *Haustor pricei* (Rutter).

Southern Arizona.

Villarius pricei Rutter, Proc., Calif. Acad. Sci., 2d ser., VI, 1896, 257, San Bernardino Creek, a tributary of the Yaqui River, southern Arizona.

1160. *Haustor dugesi* (Bean).

Rio Turbio, Guanajuato, Mexico, west of the Sierra Madre.

Ameiurus dugesi Bean, Proc. U. S. Nat. Mus., II, 1879, 304, Rio Turbio, Guanajuato, Mexico.

Genus 375. AMEIURUS Rafinesque. Horned pout.

Ameiurus Rafinesque, Ichth. Ohiensis, 1820, 65 (*Silurus cupreus* Rafinesque = *Pimelodus natalis* LeSueur).

Amiurus, corrected spelling.

1161. *Ameiurus ephippiatus* Jordan.

Coast swamps and black water streams from New Jersey to Florida.

Ameiurus ephippiatus Jordan, Bull. U. S. Nat. Mus., X, 1877, 85, St. John's River, Fla.

1162. *Ameiurus natalis* (LeSueur). Yellow cat.

Great Lakes region to Virginia and southward.

Pimelodus natalis LeSueur, Mem. Mus., 1819, 154, North America.

Silurus lividus and var. *fucus* Rafinesque, Quarterly Jour. Sci. Lond., 1820, 48, Ohio River.

Silurus cupreus Rafinesque, loc. cit., 1820, 51, Ohio River.

Pimelodus cupreus Rafinesque, Ichth. Ohiensis, 1820, 65, Ohio River.

Pimelodus caninus Richardson, Fauna Bor.-Amer., III, 1836, 132, Lake Huron.

Pimelodus felinus et *antonii* Girard, Pac. R. R. Surv., X, 1858, 209, Canadian and San Antonio Rivers, Tex.

Pimelodus cupresoides Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 159, Aux Plaines, Ill.

Ameiurus natalis analis Jordan, Bull. U. S. Nat. Mus., X, 1877, 81, Arkansas River.

Ameiurus belli Cope, Bull. U. S. Nat. Mus., XX, 1880, 35, Little Wichita River, Tex.

1163. *Ameiurus vulgaris* (Thompson).

Vermont to Minnesota and Illinois, chiefly northward. A doubtful species.

Pimelodus vulgaris Thompson, Hist. Vermont, 1842, 138, Lake Champlain.

Pimelodus dekayi Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 160, Oswego, N. Y.

Pimelodus ailiurus Girard, Pac. R. R. Surv., X, 1858, 210, Lake Amelia, Minn.

- 1164. *Ameiurus nebulosus* (LeSueur).** *Horned pout; Common bullhead; Small catfish; Schuylkill cat; Sacramento cat.*
 Great Lakes, Ohio Valley, eastward to Maine, southwestward to Texas and southwestward to Florida; introduced about 1877 into the Humboldt, Sacramento, San Joaquin, and Gila Rivers, where it is now excessively abundant.
Pimelodus nebulosus LeSueur, Mem. Mus., V, 1819, 149, Lake Ontario.
Pimelodus atrarius DeKay, N. Y. Fauna: Fishes, 1842, 185, Wappingers Creek, Dutchess County, N. Y.
Pimelodus felis Agassiz, Lake Superior, 1850, 281, Lake Superior.
Pimelodus vulpeculus Girard, Proc., Acad. Nat. Sci. Phila., XI, 1859, 160, Charleston, S. C.
Pimelodus hoyi Girard, loc. cit., 159, Racine, Wis.
Ameiurus mississipiensis Cope, Proc., Amer. Phil. Soc., 1870, 486, Mississippian Creek, Del.
- 1165. *Ameiurus nebulosus catulus* (Girard).**
 Streams of Texas; abundant.
Pimelodus catulus Girard, Pac. R. R. Surv., X, 1858, Pt. 4, Fishes, 208, Fort Smith, Ark.
- 1166. *Ameiurus prosthistius* Cope.**
 New Jersey.
Amiurus prosthistius Cope, Proc. Acad. Nat. Sci. Phila., XXXV, 1883, 133, Batsto River, N. J.
- 1167. *Ameiurus platycephalus* (Girard).** *Mud cat; Brown cat.*
 Streams of the Carolinas and eastern Georgia, Cape Fear to the Chattahoochee.
Pimelodus platycephalus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 161, Anderson, S. C.
Amiurus brunneus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1877, 366, Ocmulgee River, Flat Shoals, Ga.; the young of *platycephalus*.
- 1168. *Ameiurus brachyacanthus* Cope.**
 Streams of Texas.
Amiurus brachyacanthus Cope, Bull. U. S. Nat. Mus., XVII, 1880, 35, Wallace Creek, Bandera County, Tex.
- 1169. *Ameiurus melas* (Rafinesque).** *Black bullhead; Small catfish.*
 Northern New York (Genesee River) to Kansas and Texas; common especially west of the Mississippi.
Silurus melas Rafinesque, Quarterly Jour. Sci. Lond., 1820, 51, Ohio River.
Silurus xanthocephalus Rafinesque, loc. cit., 1820, 51, Ohio River.
Pimelodus pullus DeKay, N. Y. Fauna: Fishes, 1842, 184, Lake Pleasant, Janet, N. Y.
Pimelodus confinis Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 159, Racine, Wis.
Amiurus obesus Gill, Proc. Bost. Soc. Nat. Hist., 1862, 45, Nebraska.
Amiurus cragini Gilbert, Bull., Washburn Lab. Nat. Hist., I, 1884, 10, Garden City, Kans.
- 1170. *Ameiurus marmoratus* (Holbrook).**
 Lowland streams and swamps from southwestern Indiana (Switz City) southward and eastward to Florida.
Pimelodus marmoratus Holbrook, Jour. Acad. Nat. Sci. Phila., III, 1855, 54, South Carolina.
- Genus 376. *GRONIAS* Cope.** *Blind catfishes.*
Gronias Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 231 (*G. nigri-labris* Cope).
- 1171. *Gronias nigrilabris* Cope.**
 Cave streams tributary to Conestoga River in eastern Pennsylvania.
Gronias nigrilabris Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 231, Conestoga River, Pa.

Genus 377. OPLADELUS Rafinesque.

?*Pilodictis* Rafinesque, Jour. Phys., 1819, 422 (*P. limosa* Rafinesque, after Audubon; drawing of an imaginary catfish).

Leptops Rafinesque, Ichth. Ohiensis, 1820, 64 (*Pimelodus viscosus* Rafinesque = *Silurus olivaris* Rafinesque).

Opladelus Rafinesque, loc. cit., 64 (*Pimelodus nebulosus* Rafinesque = *Silurus olivaris* Rafinesque); name chosen by first reviser.

Ilicis Rafinesque, loc. cit., 66 (*Pimelodus nebulosus* Rafinesque = *Silurus olivaris* Rafinesque).

Hopladelus and *Pelodichthys*, corrected spelling.

- 1172. Opladelus olivaris** (Rafinesque). *Mud cat; Yellow cat; Bashaw; Russian cat; "Hoosier" at New Boston, Ill.; Goujon.*

Silurus olivaris Rafinesque, Amer. Month. Mag., 1818, 355, Ohio River.

Silurus nebulosus, viscosus and *limosus* Rafinesque, Quarterly Jour. Sci. Lond., 1820, 50, Ohio River.

Pylodictis limosus Rafinesque, Ichth. Ohiensis, 1820, 67, Ohio and Mississippi Rivers; based on a drawing by Audubon.

Pimelodus punctulatus Cuvier and Valenciennes, Hist. Nat. Poiss., XV, 1840, 134, New Harmony, Ind.

Pimelodus æneus Cuvier and Valenciennes, loc. cit., 135, New Orleans.

Genus 378. NOTURUS Rafinesque. Stone cats.

Noturus Rafinesque, Amer. Month. Mag., November, 1818, 41 (*N. flavus* Rafinesque).

- 1173. Noturus flavus** Rafinesque. *Stone cat.*

Great Lakes region, westward to Montana and Wyoming and south to Texas.

Noturus flavus Rafinesque, Amer. Month. Mag., 1818, 41, falls of Ohio River.

Noturus luteus Rafinesque, Jour. Phys., 1819, 421, Ohio River.

Noturus occidentalis Gill, Proc. Bost. Soc. Nat. Hist., VIII, 1862, 45, Platte River.

Noturus platycephalus Günther, Cat., V, 1864, 104, North America.

Genus 379. SCHILBEODES Bleeker. Tadpole cats.

Schilbeodes Bleeker, Ichth. Archip. Indici. Prodr. Siluri, 1858, 258 (*Silurus gyrinus* Mitchell).

- 1174. Schilbeodes gyrinus** (Mitchell).

Hudson River and westward through almost the entire Mississippi Valley, and Great Lakes region.

Silurus gyrinus Mitchell, Amer. Month. Mag., 1818, 322, Walkill River, N. Y.

Noturus sialis Jordan, Bull. U. S. Nat. Mus., X, 1877, 102, White River, Ind.

Genus 380. TROGLOGLANIS Eigenmann.

Trogloglanis Eigenmann, Proc. Amer. Phil. Soc., LVIII, 1919, 397 (*T. pattersoni* Eigenmann).

- 1175. Trogloglanis pattersoni** Eigenmann.

Artesian well in San Antonio, Tex.

Trogloglanis pattersoni Eigenmann, Proc. Amer. Phil. Soc., LVIII, 1919, 397, figs. 1 and 2.

Genus 381. RABIDA Jordan and Evermann. Mad toms.

Rabida Jordan and Evermann, Fishes North and Mid. Amer., 1896, 146 (*Noturus furoiosus* Jordan and Meek).

Pimelodon (LeSueur) Vaillant, Bull. Soc. Philom., 1896, 28 (*Pimelodon insignarius* LeSueur, unpublished plate of 1819; probably *Pimelodus insigne* Richardson, the *lemniscatus* = *Pimelode insigne* of LeSueur).

Subgenus PIMELODON.

- 1176. Rabida leptacantha** (Jordan).

Gulf States, in sandy streams; basins of Ogeechee, Alabama, and Pearl Rivers.

Noturus leptacanthus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 852, Silver Creek, branch of Etowah River, Rome, Ga.

- 1177. *Rabida nocturna* (Jordan and Gilbert).**
 Sandy streams, among weeds, in lower Mississippi Valley; known from the lower Wabash in Indiana and from the Poteau, Washita, and Saline Rivers, Ark.
Noturus nocturnus Jordan and Gilbert, Proc. U. S. Nat. Mus., IX, 1886, 6, Saline River at Benton, Ark.
- 1178. *Rabida funebris* (Jordan and Swain).**
 Tributary of Black Warrior River, Tuscaloosa, Ala.
Noturus funebris Jordan and Swain, Bull. U. S. Fish Com., IX, 1889, 153, North River, Tuscaloosa, Ala.
- 1179. *Rabida exilis* (Nelson).**
 Wisconsin to Missouri and Kansas; common in prairie streams.
Noturus exilis Nelson, Bull. Ill. Lab. Nat. Hist., 1876, 51, Illinois River.
Noturus classochir Swain and Kalb, Proc. U. S. Nat. Mus., V, 1882, 639, Naperville, Ill.
- 1180. *Rabida insignis* (Richardson). *Mad tom.***
 Pennsylvania to South Carolina; very common in streams east of the Alleghenies.
Pimelodus insignis Richardson, Fauna Bor.-Amer., III, 1836, 32, based on *Pimelode livree* LeSueur, Mem. Mus., V, 1819, 155, locality not known.
Noturus marginatus Baird, in Cope, Jour. Acad. Nat. Sci. Phila., VI, 1869, 237, Pa.
Pimelodon insignarius LeSueur, in Vaillant, Bull. Soc. Philom., 1896, 28.
- 1181. *Rabida giberti* (Jordan and Evermann).**
 Roanoke River in southern Virginia, locally abundant.
Noturus giberti Jordan and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 352, Roanoke River, Roanoke, Va.
- 1182. *Rabida miura* (Jordan).**
 Mississippi Valley and tributaries of Lake Michigan, south to Louisiana; common in the Ohio Valley.
Noturus miurus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1877, 371, White River, Ind.
- 1183. *Rabida eleuthera* (Jordan).**
 Mississippi Valley, rare, but widely distributed; tributaries of French Broad River (Big Pigeon River, Clifton, Tenn., French Broad at Hot Springs, N. C.); White River, Gosport, Ind.; Green River, Greensburg, Ky.; Schriner Lake, Columbia City, Ind.; White River, Fayetteville, Ark.; Sallisaw River, Indian Territory [Okla.]. Only a few specimens known.
Noturus eleutherus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1877, 371, Big Pigeon River, Clifton, Tenn.
- 1184. *Rabida furiosa* (Jordan and Meek).**
 Eastern North Carolina; known from the Neuse, Tar, and Little Rivers.
Noturus furiosus Jordan and Meek, Proc. U. S. Nat. Mus., XI, 1888, 351, Neuse River, N. C.
- Family 94. PIMELODIDÆ. South American catfishes; Bagres de Rio**
- Genus 382. *CONORHYNCHOS* Bleeker.**
Conorhynchos Bleeker, Visch. Ind. Archipel. Siluri, 1858, 191 (*Pimelodus conirostris* Cuvier and Valenciennes); not *Conorhynchus* Nozeman, a pre-Linnæan name (1757) revived by Gill (1861).
Conorhynchichthys Regan, *Conorhynchos* being regarded as preoccupied.
- 1185. *Conorhynchos nelsoni* Evermann and Goldsborough.**
 Rio Usumacinta, Yucatan.
Conorhynchus nelsoni Evermann and Goldsborough, Bull. U. S. Fish Com., XXI, 1901 (1902), 140, fig. 1, Rio Usumacinta at Monte Cristo, Yucatan.
- Genus 383. *RHAMDIA* Bleeker. *Bagres de Rio.***
Pteronotus Swainson, Nat. Hist. Fishes, Amphib., etc., II, 1839, 309 (*Heterobranchus sextentaculatus* Agassiz, preoccupied in mollusks).
Rhamdia Bleeker, Visch. Ind. Archipel.: Siluri, 1858, 197 (*Pimelodus quelon* Quoy and Gaimard).
Rhamdella Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888, 172 (*R. eriarcha* Eigenmann and Eigenmann).

- 1186. *Rhamdia wagneri* (Günther). *Barbudo.***
 Rivers of Central America, on both sides of the mountains; very abundant.
Pimelodus wagneri Günther, Fish. Cent. Amer., 1869, 474, Atlantic and Pacific rivers of Panama.
Rhamdia bransfordi Gill, Proc. Acad. Nat. Sci. Phila., XXVIII, 1876, 337, Panama.
- 1187. *Rhamdia oaxacæ* Meek. *Bagre.***
 Streams of Mexico on the Atlantic side south of Vera Cruz.
Rhamdia oaxacæ Meek, Field Col. Mus., Zool., III, 74 (figured in plate V, loc. cit., V, 1905) Rio Quiotepec, Cuicatlan, Mexico.
- 1188. *Rhamdia baronis-mülleri* (Troschel).**
 "Pacific Ocean" in Mexico.
Pimelodus baronis-mülleri Troschel, in Müller's Beitr. Gesch. Statis. und Zool. Mexico, III, 1865, 636, no definite locality.
- 1189. *Rhamdia motaguensis* (Günther).**
 Rio Motagua, Guatemala.
Pimelodus motaguensis Günther, Cat., V, 1864, 127, Rio Motagua, Guatemala.
- 1190. *Rhamdia guatemalensis* (Günther).**
 Huamuchal, Guatemala.
Pimelodus guatemalensis Günther, Cat., V, 1864, 122, Rio Huamuchal, Guatemala.
Pimelodus brachypterus Cope, Trans. Amer. Phil. Soc., XIII, 1866, 404, Orizaba, Mexico.
- 1191. *Rhamdia salvini* (Günther).**
 Rio San Gerónimo, Guatemala.
Pimelodus salvini Günther, Cat., V, 1864, 130, Rio San Gerónimo, Guatemala.
- 1192. *Rhamdia hypselura* (Günther).**
 Mexico.
Pimelodus hypselurus Günther, Cat., V, 1864, 126, Mexico.
- 1193. *Rhamdia brachycephala* (Regan).**
 Mexico.
Pimelodus brachycephalus Regan, Ann. Mag. Nat. Hist., XIX, 1907, 258, Guatemala, Rio Nacasil.
- 1194. *Rhamdia laticauda* (Heckel).**
 Mexico.
Pimelodus laticaudus Heckel, in Kner, Sitzungsb Kais. Akad. Wiss. Wien, XXVI, 1857, 420, Mexico.
- 1195. *Rhamdia godmani* (Günther).**
 Valley of Rio Usumacinta, Guatemala.
Pimelodus godmani Günther, Cat., V, 1864, 124, Rio Usumacinta, Guatemala.
- 1196. *Rhamdia nicaraguensis* (Günther).**
 Lake Nicaragua.
Pimelodus nicaraguensis Günther, Cat., V, 1864, 125, Lake Nicaragua, Nicaragua.
- 1197. *Rhamdia microptera* (Günther).**
 Rio San Gerónimo, Guatemala.
Pimelodus micropterus Günther, Cat., V, 1864, 124, Rio San Gerónimo, Guatemala.
- 1198. *Rhamdia managuensis* (Günther).**
 Lake Managua, Nicaragua.
Pimelodus managuensis Günther, Fish. Cent. Amer., 1869, 474, Lake Managua, Nicaragua.
- 1199. *Rhamdia cabreræ* Meek.**
 Lake Amatitlan, Guatemala.
Rhamdia cabreræ Meek, Field Mus. Nat. Hist., Zool., VII, 1906, 93, Lake Amatitlan, Nicaragua.

1200. *Rhamdia barbata* Meek. *Chuchin.*
Lakes of Nicaragua.
Rhamdia barbata Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 106,
 San Francisco, Nicaragua.
1201. *Rhamdia sacrificii* Barbour and Cole.
Chichen-Itza, Yucatan.
Rhamdia sacrificii Barbour and Cole, Bull. Mus. Comp. Zool., L, 1906,
 No. 5, 156, Sacrificial Cenote, near Chichen-Itza, Yucatan.
1202. *Rhamdia depressa* Barbour and Cole.
Near Chichen-Itza, Yucatan.
Rhamdia depressa Barbour and Cole, Bull. Mus. Comp. Zool., L, 1906,
 No. 5, 155, pl. 1, Ikil Cenote, near Chichen-Itza, Yucatan.
1203. *Rhamdia heteracantha* Regan.
Costa Rica.
Rhamdia heteracantha Regan, Biol. Centrali-Amer., Pisces, 1907, 134,
 Juan Viñas, Costa Rica.
1204. *Rhamdia polycaula* (Günther).
Rio San Geronimo.
Pimelodus polycaulus Günther, Cat., V, 1864, 131, Rio San Geronimo.
1205. *Rhamdia petenensis* (Günther).
Lake Peten, Yucatan.
Pimelodus petenensis Günther, Cat., V, 1864, 126, Lake Peten, Yucatan.
1206. *Rhamdia parryi* Eigenmann.
Rio Zanalenco, Chiapas.
Rhamdia parryi Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888,
 130, Rio Zanalenco, Chiapas.
1207. *Rhamdia underwoodi* Regan.
Costa Rica.
Rhamdia underwoodi Regan, Biol. Centrali-Amer., Pisces, 1907, 135,
 pl. 23, fig. 4, Juan Viñas, Costa Rica.
1208. *Rhamdia rogersi* (Regan).
Pimelodus rogersi Regan, Ann. Mag. Nat. Hist., XIX, 1907, 259,
 Irazu, Costa Rica.
Rhamdia regani Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 144,
 Rio Tiribi, San José.
- Genus 384. **CÆCORHAMDIA** Norman.
Cæcorhamdia J. R. Norman, Ann. Mag. Nat. Hist., ser. 9, XVIII, 1926,
 324 (*C. urichi* Norman).
1209. *Cæcorhamdia urichi* Norman.
Guacharo Cave, Trinidad.
Cæcorhamdia urichi J. R. Norman, Ann. Mag. Nat. Hist., ser. 9,
 XVIII, 1926, 324, Guacharo Cave, Trinidad.
- Genus 385. **PIMELODELLA** Eigenmann and Eigenmann.
Pimelodella Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d
 ser., I, 1888, 131 (*Pimelodus cristatus* Müller and Troschel).
1210. *Pimelodella chagresi* (Steindachner).
Rio Chagres.
Pimelodus chagresi Steindachner, Ichth. Beitr., IV, 1876, 34, Rio
 Chagres at Obispo, Canal Zone.
- Genus 386. **PIMELODUS** Lacépède
Pimelodus Lacépède, Hist. Nat. Poiss., V, 1803, 94 (*P. maculatus*
 Lacépède, as restricted by authors = *Silurus clarias* Bloch).
Pseudariodes Bleeker, Nederl. Tijdschr. Dierk., 1863, 99 (*Silurus clarias*
 Bloch).
Pseudorhamdia Bleeker, Versl. Akad. Amsterdam, XIV, 1862, 384
 (*Pimelodus ascita* Bleeker).

1211. *Pimelodus clarias* (Bloch).

Rio Mamoni (Panama) and southeastward to Rio de la Plata.

Silurus clarias Linnaeus, Syst. Nat., ed. X, 1758, 306, in part, not the type.

Pimelodus maculatus Lacépède, Hist. Nat. Poiss., V, 1803, 94, 107, Rio de la Plata.

Piramutana blockii Günther, Cat., V, 1864, III, Cayenne; Surinam.

Pimelodus rigidus Spix, in Agassiz, Pisc. Brasil., 1829, 19, Brazil.

Pimelodus macronema Bleeker, Silures de Suriname, 1864, 79, pl. 14, Surinam.

Pseudaroides pantherinus Lütken, Meddel. Nat. Hist. Forem., 1874, Nos. 1, 2, p. 194, Venezuela.

Pseudorhamdia piscatrix Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 569, Pebas.

Piramutana macrospila Günther, Ann. and Mag. Nat. Hist., 5th ser., VI, July, 1880, 10, Pl. II, Pirana, Rio de la Plata.

Megalonema punctatum Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 77, Rio Tuyra, Marrigante, Panama.

Megalonema robustum Meek and Hildebrand, loc. cit., 78, Rio Tuyra, Marrigante, Panama.

Genus 387. **TRACHCORYSTES** Bleeker.

Trachcorystes Bleeker, Vissch. Ind. Archipel. Siluri, 1858, 20 (*T. typus* Bleeker).

Parauchenipterus Bleeker, Nederl. Tijdschr. Dierk., I, 1863, 7 (*Silurus galeatus* L.).

1212. *Trachcorystes amblops* (Meek and Hildebrand).

Rio Tuyra Basin, Panama.

Felichthys amblops Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 77, Rio Tuyra, Marrigante, Panama.

Genus 388. **AGENEIOSUS** Lacépède.

Ageneiosus Lacépède, Hist. Nat. Poiss., V, 1803, 132 (*A. armatus* Lacépède = *Silurus militaris* Linnaeus).

Pseudageneiosus Bleeker, Nederl. Tijdschr. Dierk., I, 1863, 14 (*Ageneiosus brevifilis* Cuvier and Valenciennes).

1213. *Ageneiosus caucanus* Steindachner.

Lower Rio Tuyra, Marrigante, Panama.

Ageneiosus caucanus Steindachner, Denkschr. Kais Akad. Wiss. Wien, XLI, 1880, 61, Pl. VI, figs. 1, 2, Rio Cauca, Panama.

Family 95. PYGIDIIDÆ

Genus 389. **PYGIDIUM** Meyen.

Pygidium Meyen, Reise in Peru, I, 1835, 475 (*P. fuscum* Meyen).

1214. *Pygidium striatum* Meek and Hildebrand.

Upper Tuyra Basin, Panama.

Pygidium striatum Meek and Hildebrand, Field. Mus. Nat. Hist., Zool., X, 1913, 78, Rio Cana, Cana, Panama.

Family 96. ASTROBLEPIDÆ

Genus 390. **ASTROBLEPUS**³¹ Humboldt. *Cave catfishes.*

Astroblepus Humboldt, Obs. Zool. Phil. Mag., XXIV, 1806, 19, 331 (*A. grizalvii* Humboldt).

Cyclopium Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 305 (*Pimelodus cyclopium* Humboldt = *Cyclopium humboldti* Swainson).

Arges Cuvier and Valenciennes, Hist. Nat. Poiss., XV, 1840, 333 (*A. sabalo* Cuvier and Valenciennes).

Brontes Cuvier and Valenciennes, loc. cit., 341 (*B. prenadiilla* Cuvier and Valenciennes, preoccupied).

Strephon Gistel, Thierreich., VIII, 1848 (*Brontes prenadiilla* Cuvier and Valenciennes, preoccupied).

Stygogenes Günther, Cat., V, 1864, 223 (*S. humboldti* Günther).

³¹ According to Eigenmann, all these names belong to the same genus.

1215. *Astroblepus pirrensis* (Meek and Hildebrand).

Tuyra River Basin.

Cyclopium pirrense Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 83, Rio Cana, Cana, Panama (Pl. XVI, Field Mus. Nat. Hist., Zool., X).

Family 97. CALLICHTHYIDÆ

Genus 391. *HOPLOSTERNUM* Gill.

Hoplosternum Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 395 (*Callichthys lavigatus* Cuvier and Valenciennes = *Callichthys littoratus* Hancock).

1216. *Hoplosternum punctatum* Meek and Hildebrand.

Rio Marte Arnade, Panama.

?*Callichthys (Hoplosternum) thoracatus* (not Cuvier and Valenciennes), Steindachner, Sitzungsbs., Kais. Akad. Wiss. Wien, XLI, 1880, 66, Cauca, near Caceres.

Hoplosternum punctatum Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 264, Rio Marte Arnade, 6 miles east of Panama City.

Family 98. LORICARIIDÆ. Mailed catfishes

Genus 392. *HYPOSTOMUS* (Lacépède).

Plecostomus Gronow, Zoophyl., 1763, 127 (*Plecostomus dorso dipterygio* Gronow); not binomial.

Hypostomus Lacépède, Hist. Nat. Poiss., V, 1803, 144 (*Hypostomus guacari* Lacépède).

1217. *Hypostomus plecostomus* (Linnaeus).

Both slopes of Panama, south to Uruguay.

Loricaria plecostomus Linnaeus, Syst. Nat., ed. XII, I, 1766, 508, America.

Hypostomus guacari Lacépède, Hist. Nat. Poiss., V, 1803, 145, America.

Loricaria flava Shaw, Gen. Zool., V, 1805, 38, pl. 101.

Hypostomus robinii Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 46, Trinidad.

Hypostomus brasiliensis Bleeker, Silures de Suriname, 1864, 7, Surinam.

Plecostomus seminudus Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2nd ser., I, 1888, 169, Brazil.

Plecostomus boulengeri Eigenmann and Kennedy, Proc. Acad. Nat. Sci. Phila., LV, 1903, 502, Paraguay.

Genus 393. *THYSANOCARA* Regan.

Thysanocara Regan, Ann. Mag. Nat. Hist., ser. 7, XVII, 1906, 95 (*Hypostomus cirrhosus* Cuvier and Valenciennes).

1218. *Thysanocara spinosa* (Meek and Hildebrand).

Rare, male taken at mouth of Rio Yape, the female (type) in Rio Calobre, Panama.

Ancistrus spinosus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 252, Rio Calobre, tributary of the Rio Bayano, Panama.

Genus 394. *XENOCARA* Regan.

Xenocara Regan, Trans. Zool. Soc. Lond., XVII, 1903, 251 (*Chæstostomus latifrons* Günther).

1219. *Xenocara chagresi* (Eigenmann and Eigenmann).

Rio Chagres, Panama.

Ancistrus chagresi Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2nd ser., II, 1889, 47, Rio Chagres, Panama.

Genus 395. *CHÆSTOSTOMA* Heckel.

Chæstoma Heckel, in Tschudi, Fauna Peruana, 1846, 25 (*C. loborhynchus* Heckel).

Chæstostomus Kner and authors generally; needlessly altered spelling.

- 1220. *Chætostoma fischeri* (Steindachner). *Couchu.***
Rio Mamoni, near Panama.
Chætostomus fischeri Steindachner, Flussfische Südamerikas, I, 1879, 14,
 Rio Mamoni, Panama.
- Genus 396. *LASIANCISTRUS* Regan.**
Lasiancistrus Regan, Trans. Zool. Soc. Lond., XVII, Pt. III, 1904, 224
 and 237 (*Chætostomus heteracanthus* Günther).
- 1221. *Lasiancistrus planiceps* (Meek and Hildebrand).**
Rio Tuyra Basin, Panama.
Ancistrus planiceps Meek and Hildebrand, Field Mus. Nat. Hist., Zool.,
 X, 1913, 79, Pl. X; loc. cit., 1916, Rio Tuyra, Boca de Cupe, Panama.
- Genus 397. *LEPTOANCISTRUS* Meek and Hildebrand.**
Leptoancistrus Meek and Hildebrand, Field. Mus. Nat. Hist., Zool.,
 X, 1916, 254 (*Acanthias canescens* Meek and Hildebrand).
- 1222. *Leptoancistrus canensis* (Meek and Hildebrand).**
Mountain streams near Cana, Panama.
Acanthicus canensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool.,
 X, 1913, 80, Rio Cana, Cana, Panama.
- Genus 398. *STURISOMA* Swainson.**
Sturisoma Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 304
 (*Loricaria rostrata* Spix).
Hemiodon Kner, Panzerwelse, Denkschr., Kais. Akad. Wiss. Wien, 1854,
 65 (*H. depressus* Kner).
- 1223. *Sturisoma rostrata* (Spix).**
Rivers of eastern South America, from Panama to the Paraguay;
widely distributed and abundant.
Loricaria rostrata Spix, Pisc. Brasil., 1829, 5, Pl. III, Rio Blanco, Brazil.
Loricaria acuta Cuvier and Valenciennes, Hist. Nat. Poiss., XV, 1840,
 349 (472), pl. 452, Barra di Rio Negro.
?Loricaria barbata Kner, Panzerwelse, Denkschr. Kais. Akad. Wiss.
 Wien, 1853, 87, Rio Cujuba.
- Genus 399. *ANCISTRUS* Bleeker.**
Ancistrus Kner, Denkschr., Kais. Akad. Wiss. Wien, VII, 1854, 272
 (abstract issued in 1853; *Ancistrus medians* Kner, as first restricted
 by Gill).
Hemiancistrus Bleeker, Nederl. Tijdschr. Dierk., I, 1862 (1863), 78
 (*Ancistrus mediana* Kner).
- 1224. *Ancistrus guacharote* (Cuvier and Valenciennes).**
Porto Rico.
Hypostomus guacharote Cuvier and Valenciennes, Hist. Nat. Poiss., XV,
 1840, 375 (508), Porto Rico.
- 1225. *Ancistrus aspidolepis* (Günther).**
Veraqua, Central America.
Chætostomus aspidolepis Günther, Trans. Zool. Soc. Lond., VI, 1868
 (Nov. 27), 477, Veraqua.
- Genus 400. *LORICARIA* Linnæus.**
Loricaria Linnæus, Syst. Nat., ed. X, 1758, 307 (*L. cataphracta* Linnæus).
Hemiodon Kner, Denkschr. Akad. Wiss. Wien, 1854, about 65 (*H. de-
 pressus* Kner).
Loricarichthys Bleeker, Nederl. Tijdschr. Dierk., I, 1863, 3 (*Loricaria
 maculata* Bloch)
Pseudoloricaria Bleeker, loc. cit., 3 (*Loricaria lœviuscula* Cuvier and
 Valenciennes).
Parahemiodon Bleeker, loc. cit., 3 (*P. typus* Bleeker).
Hemiloricaria Bleeker, loc. cit., 3 (*H. caracasenses* Bleeker).
Pseudohemiodon Bleeker, loc. cit., 3 (*Hemiodon platycephalus* Kner).
- 1226. *Loricaria variegata* Steindachner.**
Rio Mamoni, Panama.
Loricaria variegata Steindachner, Flussfische Südamerikas, I, 1879, 15,
 Rio Mamoni, Panama.

- 1227. *Loricaria uracantha* Kner and Steindachner.**
 Rio Chagres, Panama.
Loricaria uracantha Kner and Steindachner, Abhandl., Bayer. Acad. Wiss., 1865, 56, Rio Chagres, Panama.
Loricaria brasfordi Gill, Proc. Acad. Nat. Sci. Phila., XXVIII, 1876, 338, Empire Station, Isthmus Railway, Panama.
- 1228. *Loricaria latiura* Eigenmann and Vance.**
 Rio Tuyra and Rio Atrato Basins, Panama; probably not in Pacific streams.
Loricaria filamentosa latiura Eigenmann and Vance, Ind. Univ. Stud., No. 16, 1912, 13, Boca de Certegai, Panama.
Loricaria tuyrensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 81, Rio Capeti, Panama.
- 1229. *Loricaria capetensis* Meek and Hildebrand.**
 Rio Tuyra Basin, Panama.
Loricaria capetensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 80, Pl. XII; loc. cit., 1916; Rio Capeti, tributary Rio Tuyra, Panama.
- 1230. *Loricaria fimbriata* Eigenmann and Vance.**
 Rio Tuyra Basin, Panama.
Locaria fimbriata Eigenmann and Vance, Ind. Univ. Stud., No. 16, 1912, 12; Boca de Certegai and Bernal Creek, Colombia.
- 1231. *Loricaria altipinnis* Breder.**
 Tuyra Basin, Pacific slope of Panama.
Loricaria altipinnis C. M. Breder, Jr., Amer. Mus. Novitates, CLXXX, 1925, 1, fig. 1, 2, Rio Chucunaque, Panama.
- Genus 401. RINELORICARIA Bleeker.**
Rineloricaria Bleeker, Nederl. Tijdschr. Dierk., I, 1862 (1863), 80 (*Loricaria lima* Kner).
- 1232. *Rineloricaria lima* (Kner).**
 Coastwise streams of Brazil, from the Parahyba northward to Panama; widely distributed and abundant in rocky brooks.
Loricaria lima Kner, Panzerwelse, Denkschr. Kais. Akad. Wiss. Wien, 1853, 89, Brazil.
?Loricaria strigilata Hensel, Wiegmann's Archiv, 1868, 368, Santa Cruz.
- Genus 402. OXYLORICARIA Bleeker.**
Oxyloricaria Bleeker, Nederl. Tijdschr. Dierk., I, 1863, 5 (*Loricaria barbata* Kner).
- 1233. *Oxyloricaria panamensis* (Eigenmann and Eigenmann).**
 Pacific slope of Panama and Colombia, south to Ecuador.
Loricaria rostrata (non Spix) Steindachner, Denkschr. Kais. Akad. Wiss. Wien, XLI, 1879, 165, Rio Mamoni, Panama.
Loricaria panamensis Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., ser. 2, II, 1889, 34, Panama.
Loricaria aurea Steindachner, Anz. Kais. Akad. Wiss. Wien, 1900, 206, Rio Magdalena.
Loricaria frenata Boulenger, Ann. Mag. Nat. Hist., ser. 7, IX, 1902, 69, northwestern Ecuador. *L. aurea* and *L. frenata* are extralimital and may not belong here.
Oxyloricaria dariensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 81, Rio Tuyra and Rio Bayano Basins, Panama.
- 1234. *Oxyloricaria citurensis* Meek and Hildebrand.**
 Rio Bayano and Rio Tuyra Basins, Panama.
Oxyloricaria citurensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 82, Pl. XII, loc. cit., 1916, Rio Cituro, Panama.

Order Z. INIOMI

Family 99. SYNOdontidæ. Lizardfishes

Genus 403. TRACHINOCEPHALUS Gill.

Trachinocephalus Gill, Cat. Fish. East Coast N. Amer., Suppl., Proc. Acad. Nat. Sci. Phila., XIII, 1861, 53 (*Saurus myops* Cuvier and Valenciennes).

1235. *Trachinocephalus myops* (Forster). *Ground spearing; Lagarto; Snake-fish.*

Tropical parts of the western Atlantic; West Indies and Brazil to coast of South Carolina; occasionally northward to Woods Hole.
Salmo myops Forster, in Bloch and Schneider, Syst. Ichth., 1801, 421, St. Helena.

Osmerus lemniscatus Lacépède, Hist. Nat. Poiss., V, 1803, 236, Martinique; after Plumier.

Saurus truncatus Agassiz, Pisc. Brasil., 1828, 82, Brazil.

Saurus brevirostris Poey, Memorias, II, 1861, 305, Cuba; erroneously stated to have 10 rays.

Genus 404. SYNODUS Gronow. Lizardfishes.

Synodus Gronow, in Scopoli, Int. Hist. Nat., 1777, 449 (*S. synodus* L.).
Tirus Rafinesque, Caratteri, 1810, 56 (*T. marmoratus* Rafinesque = *Esox synodus* Linnaeus).

Saurus Cuvier, Règne Anim., ed. 1, 1817, 169 (*Salmo saurus* Linnaeus).
Alpismaris Risso, Europ. Mérid., III, 1826, 458 (*A. risso* Risso, very young).

1236. *Synodus synodus* (Linnaeus).

'Coast of Brazil and Lesser Antilles.

?*Esox synodus* Linnaeus, Syst. Nat., ed. XII, 1766, 516, America; based on *Synodus* of Gronow.

1237. *Synodus saurus* (Linnaeus). *Tarentola; Lacerto; Tiru.*

Coasts of southern Europe and neighboring islands, recorded by Doctor Goode from the Bermudas.

Salmo saurus Linnaeus, Syst. Nat., Ed. XII, 1766, 511, Mediterranean; after Artedi.

Saurus griseus Lowe, Trans., Zool. Soc. Lond., II, 1841, 188, Madeira.
Saurida mediterranea Swainson, Nat. Hist. Anim., II, 1839, 287, Mediterranean; name only.

Tirus marmoratus Rafinesque, Caratteri, 1810, 56, Palermo.

Alpismaris risso Risso, Europ. Mérid., III, 1826, 458; very young.

Synodus lacerta Goode, Bull. U. S. Nat. Mus., V, 1876, 68, Bermuda specimen; not of Risso.

1238. *Synodus foetens* (Linnaeus). *Lizardfish; Gallwasps; Lagarto; Soapfish.*

Cape Cod to Brazil, very common from South Carolina southward on sandy coasts.

Salmo foetens Linnaeus, Syst. Nat., ed. XII, 1766, 513, South Carolina.

Osmerus alboides Lacépède, Hist. Nat. Poiss., V, 1803, 229, Carolinas; after Linnaeus.

Coregonus ruber Lacépède, loc. cit., 263, Martinique; after Plumier.

Esox salmoninus Mitchell, Trans., Lit. Phil. Soc. N. Y., I, 1815, 442, New York.

Saurus longirostris Agassiz, Pisc. Brasil., 1828, pl. 43, Brazil.

Saurus mexicanus Cuvier, Règne Anim., ed. 2, 1829, 314, Mexico.

Saurus spixianus Poey, Memorias, II, 1861, 304, Cuba.

1239. *Synodus lucioceps* (Ayres).

San Francisco to Santa Barbara, rather common in summer.

Saurus lucioceps Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 69, San Francisco.

1240. *Synodus jenkinsi* Jordan and Bollman.

Gulf of California to Galapagos Islands in 16 to 33 fathoms; common.

Synodus jenkinsi Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 153, off coast of Colombia, Galapagos Islands.

1241. **Synodus scituliceps** Jordan and Gilbert.
 Cape San Lucas to Panama.
Synodus scituliceps Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881,
 344, Mazatlan, Mexico.
1242. **Synodus lacertinus** Gilbert.
 Acapulco, Mexico; one specimen known.
Synodus lacertinus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 55,
 Acapulco, Mexico.
1243. **Synodus poeyi** Jordan.
 Cuba; abundant.
Synodus poeyi Jordan, Proc. U. S. Nat. Mus., IX, 1886, 526, Habana.
1244. **Synodus intermedius** (Agassiz). *Sand-diver.*
 Coast of southern Florida to Brazil; not rare.
Saurus intermedius Agassiz, Spix, Pisc. Brasil., 1828, 81, Brazil.
Saurus anolis Cuvier and Valenciennes, Hist. Nat. Poiss., XXII, 1849,
 483, Bahia, Martinique.
1245. **Synodus evermanni** Jordan and Bollman.
 Mazatlan to coast of Colombia; abundant.
Synodus evermanni Jordan and Bollman, Proc. U. S. Nat. Mus., XII,
 1889, 152, Pacific Ocean off coast of Colombia in 33 fathoms.
1246. **Synodus simulans** Garman.
 Off Cocos Island.
Synodus simulans Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 251,
 Pl. L, fig. 3; towings at surface off Cocos Island.
1247. **Synodus acutus** Garman.
 Off Panama.
Synodus acutus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 252,
 off coast of Panama.
- Genus 405. BATHYSAURUS** Günther.
Bathysaurus Günther, Ann. Mag. Nat. Hist., ser. 5, II, 1878, 181 (*B. ferox* Günther).
1248. **Bathysaurus agassizii** Goode and Bean.
 Gulf Stream, in deep water.
Bathysaurus agassizii Goode and Bean, Bull. Mus. Comp. Zool., X,
 1882, No. 5, 215, Gulf Stream, 33° N., in 647 fathoms.
1249. **Bathysaurus mollis** Günther.
 Southern Pacific to coast of Lower California.
 Found off Lower California, 23° N., 1,760 fathoms (only N. Amer.
 record; Townsend and Nichols, Bull. Amer. Mus. Nat. Hist., LII,
 1925, 10).
Bathysaurus mollis Günther, Ann. Mag. Nat. Hist., 5th ser., II, 1878,
 182, middle of South Pacific off Yeddo, in 1,875 and 2,385 fathoms.
- Genus 406. BATHYLACO** Goode and Bean.
Bathylaco Goode and Bean, Oceanic Ichth., 1895, 57 (*B. nigricans* Goode
 and Bean).
1250. **Bathylaco nigricans** Goode and Bean.
 Gulf of Mexico; off Santa Cruz, at a depth of 2,393 fathoms; a single
 specimen known.
Bathylaco nigricans Goode and Bean, Oceanic Ichth., 1895, 57, fig. 69,
 off Santa Cruz, Mexico.
- Genus 407. SAURIDA** Cuvier and Valenciennes.
Saurida Cuvier and Valenciennes, Hist. Nat. Poiss., XXII, 1849, 499
 (*Salmo tumbil* Bloch).
1251. **Saurida nebulosa** Cuvier and Valenciennes.
 West Indies north to St. Eustatius.
Saurida nebulosa Cuvier and Valenciennes, Hist. Nat. Poiss., XXII,
 1849, 504, pl. 649.

Family 100. CHLOROPHTHALMIDÆ

Genus 408. CHLOROPHTHALMUS Bonaparte.

Chlorophthalmus Bonaparte, Icon. Fauna Italica, III, 1834, 144 (*C. agassizi* Bonaparte).

Hyphalonedrus Goode, Proc. U. S. Nat. Mus., III, 1880, 483 (*H. chalybeius* Goode).

1252. *Chlorophthalmus agassizii* Bonaparte.

Atlantic and Mediterranean; western Atlantic.

Chlorophthalmus agassizii Bonaparte, Fauna Italica, 1840, pl. 121, Italy.

1253. *Chlorophthalmus chalybeius* (Goode).

Gulf Stream, in from 85 to 167 fathoms.

Hyphalonedrus chalybeius Goode, Proc. U. S. Nat. Mus., III, 1880 (Feb. 16, 1881), 484, Gulf Stream off Rhode Island.

1254. *Chlorophthalmus truculentus* Goode and Bean.

One specimen, *Blake* station LII, off Barbados.

Chlorophthalmus truculentus Goode and Bean, Oceanic Ichth., 1895, 61, fig. 72, off Barbados in 158 fathoms.

1255. *Chlorophthalmus mento* Garman.

Off Panama.

Chlorophthalmus mento Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 254, 3, Pl. LIV, figs. 1-1c, *Albatross* stations 3385, 3386, and 3389 in 210 to 286 fathoms.

Family 101. SUDIDÆ

Genus 409. SUDIS Rafinesque.

Sudis Rafinesque, Caratteri, 1810, 60 (*S. hyalina* Rafinesque).

1256. *Sudis ringens* Jordan and Gilbert.

Santa Barbara Channel, Calif., the only specimen known taken from the stomach of a hake, itself found in the stomach of a long-finned albicore (*Germeo*).

Sudis ringens Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 273, Santa Barbara Channel.

1257. *Sudis intermedium* (Poey).

Matanzas, Cuba; one specimen known.

Paralepis intermedium Poey, Repertorio, II, 1867, 416, Mantanzas, Cuba.

Genus 410. ARCTOZENUS Gill.

Arctozenus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 188 (*Paralepis borealis* Reinhardt).

1258. *Arctozenus borealis* (Reinhardt).

Deep waters of Arctic America.

Paralepis borealis Reinhardt, Vidensk. Selsk. Naturv. Math. Afhandl., VII, 1832, 115, Greenland.

1259. *Arctozenus coruscans* (Jordan and Gilbert).

Puget Sound; one specimen known.

Paralepis coruscans Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 411, Port Townsend, Wash.

Genus 411. PARALEPIS Risso.

Paralepis Risso, in Cuvier, Règne Anim., ed. 1, 1817 (*Coregonus paralepis* Risso = *Paralepis coregonoides* Risso).

1260. *Paralepis coregonoides* Risso.

Mediterranean, rare; Gulf stream at *Albatross* station 2393.

Paralepis coregonoides Risso, Europ. Mérid., III, 1826, 472, Pl. VII, fig. 15, Nice.

1261. *Paralepis barracudina* Fowler and Phillips.

Coast of New Jersey.

Paralepis barracudina Fowler and Phillips, Proc. Acad. Nat. Sci. Phila., LXII, 1910, 403, with figure, Corsons Inlet, Cape May County, N. J.

Genus 412. LESTIDIOPS Hubbs.

Lestidiops Hubbs, Univ. Calif. Publ. Zool., XVI, 1916, 154 (*L. sphyraenops* Hubbs).

1262. *Lestidiops sphyraenops* Hubbs.

Lestidiops sphyraenops Hubbs, Univ. Calif. Publ. Zool., XVI, 1916, 155, pl. 18, Avalon Bay, Santa Catalina Island.

Family 102. BATHYPTEROIDÆ**Genus 413. BATHYPTEROIS Günther.**

Bathypterois Günther, Ann. Mag. Nat. Hist., ser. 5, II, 1878, 83 (*B. longifilis* Günther).

Synapteretmus Goode and Bean, Oceanic Ichth., 1895, 64 (*Bathypterois quadrifilis* Günther).

Subgenus SYNAPTERETMUS Goode and Bean.**1263. *Bathypterois quadrifilis* Günther.**

Brazil; St. Vincent.

Bathypterois quadrifilis Günther, Ann. Mag. Nat. Hist., II, 1878, 184, off coast of Brazil in 500 to 770 fathoms.

1264. *Bathypterois pectoralis* Garman.

Off Central America in deep water.

Bathypterois pectoralis Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 257, Pl. K, fig. 1, Albatross stations 3376, 3393, 3407, and 3431 in 885 to 1,132 fathoms.

1265. *Bathypterois ventralis* Garman.

Off west coast of Mexico.

Bathypterois ventralis Garman, Mem., Mus. Comp. Zool., XXIV, 1899, Pl. LV, fig. 1, 256, Albatross stations 3418 and 3425 in 660 to 680 fathoms.

Family 103. BENTHOSAURIDÆ**Genus 414. BENTHOSAURUS Goode and Bean.**

Benthosaurus Goode and Bean, Bull. Mus. Comp. Zool., XII, 1886, No. 5, 168 (*B. grallator* Goode and Bean).

1266. *Benthosaurus grallator* Goode and Bean.

Gulf of Mexico and Gulf Stream, in lat. 24° 33' N., long. 84° 23' W., and lat. 39° 3' 15" N.; long 70° 50' 45" W.

Benthosaurus grallator Goode and Bean, Bull. Mus. Comp. Zool., XII, 1886, No. 5, 168, Gulf Stream.

Family 104. MYCTOPHIDÆ³² Lanternfishes**Genus 415. MACROSTOMA Risso.**

Macrostoma Risso, Europ. Mérid., III, 1826, 447 (*Macrostoma angustidens* Risso); not *Macrostomus* Wied., a genus of insects.

Notoscopelus Günther, Cat., V, 1864, 405 (*Lampanyctus resplendens* Richardson), to replace *Macrostoma*, regarded as preoccupied.

Catalemlle Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., III, 1890, 3, 24, (*C. brachycheir* Eigenmann and Eigenmann).

1267. *Macrostoma angustidens* Risso. Prickfish; *Maire d'Amplora*.

Atlantic and Indian Oceans, Norway and Greenland.

Macrostoma angustidens Risso, Europ. Mérid., III, 1826, 448, off Nice.

1268. *Macrostoma quercinum* (Goode and Bean).

Western Atlantic; Mediterranean.

Notoscopelus quercinus Goode and Bean, Oceanic Ichth., 1895, 83, fig. 97, Grand Banks, Newfoundland.

1269. *Macrostoma margariferum* (Goode and Bean).

Grand Banks of Newfoundland.

Notoscopelus margaritiferus Goode and Bean, Oceanic Ichth., 1895, 84, fig. 98, off Banquereau.

* Several species recently described as *Myctophum* may prove to belong to other genera.

- 1270. *Macrostoma castaneum* (Goode and Bean).**
 Grand Banks.
Notoscopelus castaneus Goode and Bean, Oceanic Ichth., 1895, 84,
 fig. 95, Grand Banks, Newfoundland.
- Genus 416. CERATOSCOPELUS Günther.**
Ceratoscopelus Günther, Cat., V, 1864, 405 and 412 (*Scopelus madeirensis*
 Lowe).
- 1271. *Ceratoscopelus madeirensis* (Lowe).**
 Atlantic and Mediterranean; west to the Grand Banks.
Scopelus madeirensis Lowe, Proc. Zool. Soc. Lond., 1839, 87, Madeira.
- Genus 417. LAMPANYCTUS Bonaparte.**
Lampanyctus Bonaparte, Fauna Italica, Fasc. XXVII, 1840 (*L. bona-*
partei Bonaparte).
- 1272. *Lampanyctus guntheri* Goode and Bean.**
 Grand Banks.
Lampanyctus guntheri Goode and Bean, Oceanic Ichth., 1895, 79, fig. 90,
 Grand Banks.
- 1273. *Lampanyctus gemmifer* Goode and Bean.**
 Gulf Stream.
Lampanyctus gemmifer Goode and Bean, Oceanic Ichth., 1895, 80,
 fig. 88, Grand Banks, in lat. $39^{\circ} 40'$ N., long. $71^{\circ} 35'$ W., in 538
 fathoms.
- 1274. *Lampanyctus lacerta* Goode and Bean.**
 Gulf of Mexico.
Lampanyctus lacerta Goode and Bean, Oceanic Ichth., 1895, 81, fig. 89,
 Gulf of Mexico, in lat. $28^{\circ} 38' 30''$ N., long. $85^{\circ} 52' 30''$ W.
- 1275. *Lampanyctus ritteri* Gilbert.**
 Coast of California; Monterey Bay; San Clemente.
Lampanyctus ritteri Gilbert, Proc. U. S. Nat. Mus., XLVII, 1915, 318,
 pl. 15, fig. 3, Monterey Bay; San Clemente Island, Calif.
- Genus 418. NYCTIMASTER Jordan.**
Nyctimaster Jordan, Proc. U. S. Nat. Mus., LIX, 1921, 645 (*Lampanyc-*
tus jordani Gilbert).
 Most of the species hitherto referred to *Lampanyctus*, having the scales
 of the lateral line not larger than the others, should be referred to
Nyctimaster.
- 1276. *Nyctimaster townsendi* (Eigenmann and Eigenmann).**
 Cortez Banks near San Diego, in 45 fathoms.
Myclophum townsendi Eigenmann and Eigenmann, West Amer. Sci-
 entist, 1889, 125, Cortez Banks, off San Diego, Calif.
- 1277. *Nyctimaster alatus* (Goode and Bean).**
 Gulf of Mexico.
Lampanyctus alatus Goode and Bean, Oceanic Ichth., 1895, 79, Gulf
 Stream.
- Genus 419. LAMPADENA Goode and Bean.**
Lampadena Goode and Bean, Oceanic Ichth., 1895, 85 (*L. speculigera*
 Goode and Bean).
- 1278. *Lampadena speculigera* Goode and Bean.**
 Gulf Stream.
Lampadena speculigera Goode and Bean, Oceanic Ichth., 1895, 85,
 fig. 99, Gulf Stream, lat. $39^{\circ} 48'$ N., long. $70^{\circ} 36'$ W., in 551 fathoms.
- Genus 420. NANNOBRACHIUM Günther.**
Nannobrachium Günther, Deep Sea Fishes, Challenger, 1887, 199, (*N.*
nigrum Günther).
- 1279. *Nannobrachium nanochir* (Gilbert).**
 Alaska to Santa Barbara Islands, in deep water.
Myclophum nanochir Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 51,
 off Washington, Albatross stations 2925, 3072, and others, in 266 to
 685 fathoms.

- 1280.** *Nannobrachium leucopsarum* (Eigenmann and Eigenmann).
Pacific coast of America, from Alaska to San Diego.
Myctophum (Stenobrachius) leucopsarum Eigenmann and Eigenmann,
Proc. Calif. Acad. Sci., 2d ser., III, 1890, 5, off Point Loma, Calif.,
in stomachs of *Sebastodes*.
- 1281.** *Nannobrachium macdonaldi* Goode and Bean.
Gulf Stream.
Nannobrachium macdonaldi Goode and Bean, Oceanic Ichth., 1895, 94,
fig. 110, Gulf Stream.
- 1282.** *Nannobrachium regale* (Gilbert).
Santa Barbara Channel, in deep water.
Myctophum regale Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 544,
Santa Barbara Channel, Albatross station 2923, in 822 fathoms.
- 1283.** *Nannobrachium mexicanum* (Gilbert).
Coast of Lower California, in deep water.
Myctophum mexicanum Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 51,
off Lower California, at Albatross stations 3008 and 3009 in 306
and 857 fathoms.
- Genus 421. *AETHOPRORA* Goode and Bean.
Aethoprrora Goode and Bean, Oceanic Ichth., 1895, 86 (*Scopelus meto-*
poclampus Cocco).
- 1284.** *Aethoprrora effulgens* Goode and Bean.
Western Atlantic.
Aethoprrora effulgens Goode and Bean, Oceanic Ichth., 1895, 87, fig.
103, Browns Bank and Albatross station 2127, 19° 45' N., 75° W.,
in 689 fathoms.
- 1285.** *Aethoprrora lucida* Goode and Bean.
Gulf Stream.
Aethoprrora lucida Goode and Bean, Oceanic Ichth., 1895, 87, fig. 102,
Gulf Stream, lat. 19° 45' N., long. 75° 4' W.
- Genus 422. *COLLETTIA* Goode and Bean.
Collettia Goode and Bean, Oceanic Ichth., 1895, 88 (*Myctophum rafinesquei*
Cocco).
- 1286.** *Collettia rafinesquei* (Cocco).
Mediterranean and Atlantic; Gulf Stream off the New England coast.
Nyctophus rafinesquei Cocco, Aleuni Salmon, etc., 1820, 20, Messina.
- 1287.** *Collettia nocturna* (Poey).
Coast of Cuba.
Myctophum nocturnum Poey, Memorias, II, 1861, 426, Cuba.
- Genus 423. *DIAPHUS* Eigenmann and Eigenmann.
Diaphus Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser.,
III, 1890, 3 (*Scopelus engraulis* Günther).
- 1288.** *Diaphus theta* Eigenmann and Eigenmann.
Point Loma, near San Diego, Calif., to coast of Oregon.
Diaphus theta Eigenmann and Eigenmann, Proc. Calif. Acad. Sci.,
2d. ser., III, March, 1890, 4, Point Loma, near San Diego.
Myctophum protocolus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 52,
July 2, coast of Washington, in 584 fathoms, Albatross station 3072,
in 584 fathoms.
- Genus 424. *RHINOSCOPELUS* Lütken.
Rhinoscopelus Lütken, Vidensk. Selsk. Naturv. Copenhagen, VII,
1892, 237 (*Scopelus coccoi* Cocco).
Alysia Lowe, Proc. Zool. Soc. Lond., 1839, 87 (*A. loricata* Lowe =
Scopelus coccoi Cocco; name preoccupied).
- 1289.** *Rhinoscopelus andreae* (Lütken).
Open Atlantic and Indian Ocean; Gulf Stream.
Scopelus andreae Lütken, Spolia Atlantica, 1892, 25, North Atlantic.
- 1290.** *Rhinoscopelus rarus* (Lütken).
Open Atlantic, west to 50° W., 33° N.
Scopelus rarus Lütken, Spolia Atlantica, II, 1892, 26, North Atlantic.

- 1291. *Rhinoscopelus coccoi* (Cocco).**
West Atlantic, Gulf Stream, Mediterranean, ranging from Newfoundland to Africa.
Scopelus coccoi Cocco, Giorn. Sci. Litt. Art. Sicilia (No. 77), Palermo, 1829, 143 ("Scopelo de Cocco"), Palermo.
Alysia loricata Lowe, Proc. Zool. Soc. Lond., 1839, 87, Madeira.
- Genus 425. *MYCTOPHUM* Rafinesque. *Lanternfishes.***
Myctophum Rafinesque, Indice, 1810, 56 (*M. punctatum* Rafinesque).
Scopelus Cuvier, Règne Anim., ed. 1, 1817, 56 (*Gasteropelecus humboldtii* Risso).
Nyctophus Cocco, Giorn. Sci. Litt. Art. Sicil., 1829, 44 (*N. rafinesquei*, substitute for *Myctophum*).
- 1292. *Myctophum punctatum* Rafinesque.**
Warmer parts of the Atlantic, Grand Banks, and from the Gulf Stream to the Mediterranean.
Myctophum punctatum Rafinesque, Indice, 1810, 56, Pl. II, fig. 5, Palermo.
Scopelinus caninianus Cuvier and Valenciennes, Hist. Nat. Poiss., XXII, 1849, 445, coast of Italy.
- 1293. *Myctophum affine* (Lütken).**
Open Atlantic west to 63° W., 38° N., and in the open Pacific. Perhaps more than one species included.
Scopelus affine Lütken, Spolia Atlantica, II, 1892, 32, open Atlantic.
Myctophum nitidulum Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 266, Pl. LVI, fig. 3, lat. 27° 50' N., long. 145° 45' 30" W.
Rhinocephalus oceanicus Jordan and Evermann, Bull. U. S. Fish Com., XX, 1902, 168, Hawaiian Islands.
Myctophum margaritatum Gilbert, loc. cit., XXIII, pt. 2, 1905, p. 596, pl. 68, fig. 2, off south coast of Molokai, Hawaiian Islands.
- 1294. *Myctophum humboldti* (Risso).**
Open seas, Mediterranean, and Atlantic on both shores.
Gasteropelecus humboldtii Risso, Ichth. Nice, 1810, 358, Nice.
- 1295. *Myctophum californiense* Eigenmann and Eigenmann.**
Cortez Banks, San Diego, Calif.
? *Myctophum boops* Richardson, Voy. Erebus and Terror, 1845, 39, open sea between Australia and New Zealand.
Myctophum californiense Eigenmann and Eigenmann, West Amer. Scientist, Nov. 9, 1889, 124, San Diego, Calif.
- 1296. *Myctophum gracile* (Lütken).**
Open Atlantic, west to 48° W., 23° N.
Scopelus gracilis Lütken, Spolia Atlantica, II, 1892, 35, open Atlantic.
? *Myctophum hians* Richardson, Voy. Erebus and Terror, 1845, 41, pl. 27, open sea, locality unknown.
- 1297. *Myctophum benoiti* (Cocco).**
Mediterranean to Norway and Greenland.
Scopelus benoiti Cocco, Lett. su Salmon., 1838, 12, pl. 2, fig. 4, Messina.
- 1298. *Myctophum hygomi* (Lütken).**
North Atlantic, occasional in our waters.
Scopelus hygomi Lütken, Spolia Atlantica, 1892, 257, North Atlantic.
Myctophum remiger Goode and Bean, Oceanic Ichth., 1895, 75, western Atlantic.
- 1299. *Myctophum oculatum* Garman.**
Off Panama in deep water.
Myctophum oculatum Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 260, Pl. LVI, fig. 2, numerous stations of the Albatross.
- 1300. *Myctophum atratum* Garman.**
Off western Mexico in the open sea.
Myctophum atratum Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 268, Albatross station 3433, lat. 25° 26' 15" N., long. 109° 48' W., in 1,218 fathoms.

1301. *Myctophum laternatum* Garman.

Off Pacific coast of Mexico in open sea.

Myctophum laternatum Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 267, Pl. LVI, fig. 7, several stations of the *Albatross*.

1302. *Myctophum aurolaternatum* Garman.

Off Panama.

Myctophum aurolaternatum Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 264, Pl. LV, fig. 3, *Albatross* station 3382, lat. $6^{\circ} 21' N.$, long. $80^{\circ} 41' W.$ at surface.

1303. *Myctophum luminosum* Garman.

Off Panama.

Myctophum luminosum Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 263, Pl. LV, fig. 2, *Albatross* station 3042, lat. $0^{\circ} 57' 30'' S.$, long. $89^{\circ} 3' 30'' W.$, in 421 fathoms.

1304. *Myctophum tenuiculum* Garman.

Off Panama in deep water.

Myctophum tenuiculum Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 262, pl. 5, fig. 5, *Albatross* station 3382, lat. $6^{\circ} 21' N.$, $80^{\circ} 41' W.$, at surface.

Genus 426. *BENTHOSEMA* Goode and Bean.

Benthosema Goode and Bean, Oceanic Ichth., 1895, 76 (*Scopelus muelleri* Gmelin).

1305. *Benthosema arcticum* (Lütken).

Davis Straits, Greenland.

Scopelus arcticus Lütken, Spolia Atlantica, 1892, 29, Davis Straits.

1306. *Benthosema muelleri* (Gmelin).

North Atlantic, occasional from Norway to Greenland, more abundant southward to South Carolina; probably always in very deep water.

Salmo müllerii Gmelin, Syst. Nat., 1788, 1378, Norway; after Strom.

Scopelus glacialis Reinhardt, Overs., 1835, 36, Greenland.

Genus 427. *SCOPELENGYS* Alcock.

Scopeleengys Alcock, Ann. Mag. Nat. Hist., ser. 6, VI, 1890, 302 (*S. tristis* Alcock).

1307. *Scopeleengys dispar* Garman.

Off Panama in deep water.

Scopeleengys dispar Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 254, Pl. LIV, figs. 2-2d, *Albatross* stations 3353, 3360, 3382, and 3383, from 695 to 1,832 fathoms.

Genus 428. *TARLETONBEANIA* Eigenmann and Eigenmann.

Tarletonbeania Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., IV, 1891, 6 (*T. tenua* Eigenmann and Eigenmann).

1308. *Tarletonbeania crenularis* (Jordan and Gilbert).

San Diego and vicinity.

Myctophum crenulare Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 274, Santa Barbara, Calif.

Myctophum procellarium Bean, in Jordan and Gilbert, loc. cit., 467, off Straits of Fuca.

Tarletonbeania tenua Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., IV, 1890, 7, Coronado Islands near San Diego, Calif.

Family 105. IPNOPIDÆ

Genus 429. *IPNOPS* Günther.

Iplops Günther, Ann. Mag. Nat. Hist., II, 1878, 187 (*I. murrayi* Günther).

1309. *Iplops murrayi* Günther.

Bottom fish, at about 2,000 fathoms; coast of Brazil, Tristan da Cunha, Celebes, in tropical America ($24^{\circ} 36' N.$, $84^{\circ} W.$) and off Bequia, Windward Islands.

Iplops murrayi Günther, Ann. Mag. Nat. Hist., II, 1878, 187, coast of Brazil, etc.

1310. *Ipnops agassizi* Garman.

Pacific coast of Central America in deep water.

Ipnops agassizii Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 259,
Pl. H, figs. 2, 2a, *Albatross* station 3413, lat. 2° 34' N., long. 92° 6'
W., in 1,360 fathoms.

Family 106. OMOSUDIDÆ

Genus 430. OMOSUDIS Günther.

Omosudis Günther, Challenger Rept., XXII, 1887, 201 (*O. lowii* Günther).

1311. *Omosudis lowii* Günther.

Open sea; Gulf Stream at *Albatross* station 2392.

Omosudis lowii Günther, Challenger Rept., XXII, 1887, 201, pl. 52, figs.
C, C', Philippine Islands, Magdalena.

Family 107. ALEPISAURIDÆ. Handsawfishes

Genus 431. ALEPISAURUS Lowe.

Plagyodus Steller, in Pallas, Zoogr. Rosso-Asiat., III, 1811, 383 (no type named, doubtless *P. aesculapius* Bean); given in oblique form only, as *Plagyodontem*.

Alepisaurus Lowe, Proc. Zool. Soc. Lond., I, 1833 (*A. ferox* Lowe).

1312. *Alepisaurus ferox* Lowe. *Lancetfish*.

Deep waters of the Atlantic; occasional off the coasts of Nova Scotia, Massachusetts, and on the Grand Banks.

Alepisaurus ferox Lowe, Trans. Zool. Soc. Lond., I, 1833, 395, Madeira.

Alepisaurus azureus Cuvier and Valenciennes, Hist. Nat. Poiss., XXII, 1849, 530, Canaries.

1313. *Alepisaurus aesculapius* (Bean). *Wolfish; Subotka*.

Coast of Alaska southward to California.

Plagyodus Steller in Pallas, Zoogr. Rosso.-Asiat., III, 1811, 383, Kurile Islands.

Alepidosaurus aesculapius Bean, Proc. U. S. Nat. Mus., V, 1882, 661, Iluliuk, Unalaska.

Genus 432. CAULOPUS Gill. *Handsawfishes*.

Caulopus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 128 (*Alepidosaurus borealis* Gill).

1314. *Caulopus altivelis* (Poey).

Cuba, in deep water.

Alepisaurus altivelis Poey, Memorias, II, 1861, 302, Cuba.

Alepidosaurus (*Caulopus*) *poeyi* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 181, Cuba.

1315. *Caulopus borealis* (Gill). *Handsawfish; Serra*.

Pacific coast of America from Alaska to Puget Sound.

Alepidosaurus borealis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 128, Puget Sound.

Alepidosaurus serra Gill, loc. cit., 129, "Monterey, Lower California."

Order AA. CETUNCULI

Family 108. RONDELETIIDÆ

Genus 433. RONDELETIA Goode and Bean.

Rondeletia Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894, 454
(*R. bicolor* Goode and Bean).

1316. *Rondeletia bicolor* Goode and Bean.

Gulf Stream.

Rondeletia bicolor Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894
(1895), 454, pl. 17, fig. 7, and Oceanic Ichth., 1895, 68, fig. 77, at
Albatross station 2724 in 1,641 fathoms.

Family 109. CETOMIMIDÆ

Genus 434. CETOMIMUS Goode and Bean.

Cetomimus Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894, 453
(*C. gillii* Goode and Bean).

1317. *Cetomimus gillii* Goode and Bean.

Gulf Stream.

Cetomimus gillii Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894
(1895), 452, pl. 17, fig. 2, and in Oceanic Ichth., 1895, 69, fig. 78,
Gulf Stream, in lat. $39^{\circ} 35'$ N., long. $71^{\circ} 24' 30''$ W.

1318. *Cetomimus storeri* Goode and Bean.

Gulf Stream.

Cetomimus storeri Goode and Bean, Proc. U. S. Nat. Mus., XVII, 1894
(1895), 453, pl. 17, fig. 3, and in Oceanic Ichth., 1895, 69, fig. 79,
Gulf Stream at Albatross station 2222 in 1,535 fathoms.

Order BB. XENOMI

Family 110. DALLIIDÆ. Alaska blackfish

Genus 435. DALLIA Bean.

Dallia Bean, Proc. U. S. Nat. Mus., II, 1879, 358 (*D. pectoralis* Bean).

1319. *Dallia pectoralis* Bean. *Alaska blackfish; Charnia ryba.*

Streams and ponds of northern Alaska and Siberia.

Dallia pectoralis Bean, Proc. U. S. Nat. Mus., II, 1879, 358, St.
Michaels, Alaska.

Umbrä delicatissima Smitt, Offer. Kongl. Vetensk. Akad. Forh. Arg.,
Stockholm, 1881, 38, pl. 5, fig. 1, Pittlekag, Siberia.

Order CC. HAPLOMI

Family 111. ESOCIDÆ. Pikes

Genus 436. ESOX Linnæus.

Esox Linnæus, Syst. Nat., ed. X, 1758, 313 (*E. lucius* Linnæus).

Lucius Geoffroy, Descr. de 719 Plantes, etc., 1767, 407 (*Esox lucius*
Linnæus).

Lucius Rafinesque, Caratteri, 1810, 59 (*Esox lucius* Linnæus = *L. vorax*
Rafinesque).

Picorellus Rafinesque, Ichth. Ohiensis, 1820, 70 (*Esox vittatus* Rafinesque,
a mythical species).

Masalongus Jordan, Klippart's rept., Ohio Fish Com., 1878, 92 (*Esox*
masquinongy Mitchell = *Esox nobilior* Thompson).

Kenoza Jordan and Evermann, Fishes North and Mid. Amer., 1896,
626 (*Esox americanus* Gmelin).

Subgenus MASCALONGUS Jordan.

1320. *Esox masquinongy* Mitchell. *Muskallunge;*¹³ *Muskellunge; Maskinongy;*
Muscalonge; Great pike.

Great Lakes region; Upper Mississippi Valley and northward.

Esox masquinongy Mitchell, "Mirror," 1824, 297, Lake Erie.

Esox masquinongy Mitchell, in Kirtland, Fishes of Ohio, 1838, 194,
Lake Erie.

Esox nobilior Thompson, Proc. Bost. Soc. Nat. Hist., III, 1850, 163,
Lake Champlain.

Esox nobilis Kirtland, Proc. Cleveland Acad. Nat. Sci., 1854, 84, Lake
Erie; name a slip of the pen for *Esox nobilior*.

1321. *Esox masquinongy ohiensis* Kirtland.

Ohio River and its tributaries; Chautauqua Lake, N. Y.

Esox salmonœus Rafinesque, Amer. Month. Mag., III, 1818, 447, Ohio
River; not *Esox salmonœus* Mitchell 1815 = *Synodus fœtens*.

Esox ohiensis Kirtland, Proc. Cleveland Acad. Nat. Sci., Feb. 7, 1854,
Mahoning River, Ohio.

¹³ The common names of the fishes of the pike family, and of the pike perches, with which they are confused, are too numerous to be given here. As Weed, who gives many of them (Field Mus. Nat. Hist., Zool. Leaflet No. 9, 1927) remarks, "There are as many as 80 names for 1 fish and 7 fish for one name."

1322. ***Esox masquinongy*** *immaculatus* Garrard. *Great northern pike.*
 Lakes of Wisconsin and Minnesota.
Esox immaculatus Garrard, MS.; noticed in various fishing journals;
 Eagle Lake, northern Wisconsin.
- Subgenus **ESOX** Linnæus.
1323. ***Esox estor**** LeSueur. *American pike.*
 Northern North America; eastern United States south to New York
 and the Ohio River.
Esox estor LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 413, Lake
 Erie at Buffalo.
??*Esox villatus* Rafinesque, Amer. Month. Mag., III, 1818, 447, Ohio
 River; mythical; described from hearsay.
??*Esox australis* Cuvier and Valenciennes, Hist. Nat. Poiss., XVII,
 1846, 323, locality unknown; thought to be Van Diemen's Land
 (Tasmania).
Esox depraudus LeSueur in Cuvier and Valenciennes, Hist. Nat. Poiss.,
 XVIII, 1846, 336, Wabash River, New Harmony, Ind.
Esox lugubris LeSueur, loc. cit., 338, Crab Orchard, Ky.
Esox boreus Agassiz, Lake Superior, 1850, 317, Lake Superior.
Esox lucoides Agassiz and Girard, in Herbert Forrester's Fish and
 Fishing, 1850, 154, Lake Superior.
- Subgenus **KENOZA** Jordan and Evermann.
1324. ***Esox niger*** LeSueur. *Eastern pickerel; Green pike; Jack.*
 Massachusetts to Florida, in lowland streams and swamps.
Esox reticulatus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1818, 414,
 Connecticut River, Adams, Mass.; Philadelphia; name preoccupied
 in Europe.
? *Esox phaleratus* Say in LeSueur, loc. cit., 416, St. Augustine, Fla.;
 unidentifiable.
Esox niger LeSueur, loc. cit., 415, South Carolina.
Esox tridecemlineatus Mitchell, "Mirror," 1825, 361, Oneida Lake,
 N. Y.
1325. ***Esox vermiculatus*** LeSueur. *Little pickerel; Grass pike.*
 Mississippi Valley and tributaries of Lake Erie and Lake Michigan,
 south to Mississippi and Arkansas.
Esox vermiculatus LeSueur in Cuvier and Valenciennes, Hist. Nat.
 Poiss., XVIII, 1846, 333, Wabash River, New Harmony, Ind.
Esox lineatus LeSueur, loc. cit., 335, Wabash River, New Harmony, Ind.
Esox crassus Agassiz, Amer. Jour. Sci. Arts, 1854, 308, Tennessee
 River, Huntsville, Ala.
Esox umbrosus Kirtland, Proc. Cleveland Acad. Nat. Sci., 1854, 79,
 small Bay of Rocky River, Rockport, near Cleveland, Ohio.
Esox cypho Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 78, Water-
 ford, Oakland County, Mich.
Esox porosus Cope, Trans. Amer. Phil. Soc. Phila., XIII, 1866, 408,
 Waterford, Oakland County, Mich.; substitute for *cypho* regarded
 as an inept name.
1326. ***Esox americanus*** Gmelin. *Barred pickerel.*
Esox lucius americanus Gmelin, Syst. Nat., 1788, 1390, Long Island,
 N. Y.; after Schöpf.
Esox scomberius Mitchell, Amer. Month. Mag., 1818, 322, Murderers
 Creek, N. Y.
Esox fasciatus DeKay, New York Fauna: Fishes, 1842, 224, Murderers
 Creek and other streams near New York.
Esox ornatus Girard, "Proc. Bost. Soc. Nat. Hist., V, 1854, 41," Proc.
 Acad. Nat. Sci. Phila., XXXVII, 1885, 369, 41; Charles River,
 Mass.
Esox raveneli Holbrook, Ichth. South Carolina, 1860, 201, Charleston,
 S. C.

* Not clearly separated from *Esox lucius* L. of Europe.

Family 112. UMBRIDÆ. Mud minnows

Genus 437. UMBRA Müller.

Umbræ Krämer, MS., in Müller, Anim. Austr. Infer., 1756; Müller, Abhandl. Akad. Wiss. Wien, 1842, 188 (*crameri*).

Melanura Agassiz, Amer. Jour. Sci. Arts, XVI, 1853, 135 (*M. annulata* Agassiz, not *Exoglossum annulatum* Rafinesque, etc.=*Leuciscus pygmæus* DeKay).

1327. *Umbra limi* (Kirtland). *Mud minnow; Dogfish.*

Great Lakes region and upper Mississippi Valley in weedy streams and ponds; Lake Champlain to Minnesota. A hardy and attractive aquarium fish.

Hydrargira lima Kirtland, Bost. Jour. Nat. Hist., III, 1840, 277, streams in northern Ohio.

Hydrargira fusca Thompson, Hist. Vermont, 1842, 137, Lake Champlain.

Hydrargira atricauda DeKay, N. Y. Fauna: Fishes, 1842, 220, Lake Champlain.

1328. *Umbra pygmæa* (DeKay). *Striped mud minnow; Eastern mud minnow.* Lowland streams and swamps coastwise, from Long Island to the Neuse River, N. C.; locally abundant.

Leuciscus pygmæus DeKay, New York Fauna: Fishes, 1842, 214, Tappan, Rockland County, N. Y.

Fundulus fuscus Ayres, Bost. Jour. Nat. Hist., IV, 1843, 296, Brookhaven, Long Island.

Umbra pygmæa bilineata Shufeldt, Aquatic Life, July, 1917, tributaries of Chesapeake Bay; slight color variation.

Order DD. CYPRINODONTES. (Microcyprini)

Family 113. CYPRINODONTIDÆ. Killifishes

Genus 438. PROFUNDULUS Hubbs.

Profundulus Hubbs, Misc. Pub. 13, Mus. Zool. Univ. Mich., Jan. 18, 1924, 12 (*Fundulus punctatus* Günther).

1329. *Profundulus punctatus* (Günther).

Southern Mexico and Central America in streams and lakes, south to Ecuador.

Fundulus punctatus Günther, Cat., VI, 1866, 320, Chiapas.

Fundulus guatemalensis Günther, loc. cit., 321, Lake Duenas.

Fundulus pachycephalus Günther, loc. cit., 321, Lake Atitlan, Guatemala.

1330. *Profundulus labialis* (Günther).

Rivers of Guatemala in Atlantic drainage, Lake Yzabal, Rio Chisoy and Rio San Gerónimo.

Fundulus labialis Günther, Cat., VI, 1866, 319, Rio San Gerónimo, Guatemala, Yzabal.

1331. *Profundulus oaxacæ* (Meek).

Streams of Oaxaca, Mexico.

Fundulus oaxacæ Meek, Pub., Field Col. Mus., Zool., III, 1902, 90, Oaxaca, Mexico.

1332. *Profundulus candalaricus* Hubbs.

Profundulus candalaricus Hubbs, Misc. Pub. 13, Mus. Zool., Univ. Mich., Jan. 18, 1924, 15, Candalaria, Guatemala, Rio Chiapas Basin, Atlantic drainage.

Genus 439. CUBANICHTHYS Hubbs.

Cubanicthys Hubbs, Misc. Pub. Univ. Mich. Mus. Zool., No. 16, July 9, 1926, 4 (*Fundulus cubensis* Eigenmann).

1333. *Cubanicthys cubensis* (Eigenmann).

Streams of Pinar del Rio, Cuba.

Fundulus cubensis Eigenmann, Bull. U. S. Fish Com., XXII, 1902 (1903), 222, Pinar del Rio, Cuba.

Genus 440. LEPTOLUCANIA Myers.

Leptolucania Myers, Amer. Mus. Novitates, June 6, 1924, 8 (*Heterandria ommata* Jordan= *Zygonectes manni* Hay).

1334. *Leptolucania ommata* (Jordan).

Florida; Indian River and Santa Fe River, and their tributaries.
Heterandria ommata Jordan, Proc. U. S. Nat. Mus., VII, 1884, 323,
 Indian River, Fla.
Zygonectes manni Hay, loc. cit., VIII, 1885, 555, Yellow Water River,
 Fla.

Genus 441. *CHRIOPEOPS* Fowler.

Chriopeops Fowler, Proc. Acad. Nat. Sci. Phila., LXVIII, 1916, 425
 (*Lucania goodei* Jordan).

1335. *Chriopeops goodei* (Jordan).

Waters of the Florida Everglades; locally abundant.
Lucania goodei Jordan, Proc. U. S. Nat. Mus., II, 1879, 240, Arlington
 River, Fla., a tributary of the St. Johns.

Genus 442. *LUCANIA* Girard.

Lucania Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 118 (*Limia*
venusta Girard); not *Lucanus*, a genus of beetles.

1336. *Lucania parva* (Baird and Girard). *Rainwaterfish; Little killifish.*

Atlantic coast from Connecticut to Key West, Texas, and Mexico.
Cyprinodon parvus Baird and Girard, Ninth Smithson. Rept., 1854
 (1855), 345, Long Island.
Limia venusta Girard, U. S. Mex. Bound. Surv., Ichth., 1859, 71, pl.
 39, figs. 20-23, Indianola, Tex.
Lucania affinis Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 118,
 Matamoras, Mexico.

Genus 443. *FUNDULUS* Lacépède.

Fundulus Lacépède, Hist. Nat. Poiss., V, 1803, 37 (*Fundulus mudfish*
 Lacépède = *Cobitis heteroclitus* Linnæus).
Hydrargyra Lacépède, loc. cit., 378 (*H. swampina* Lacépède = *Fundulus*
majalis Walbaum).
Borboryx (Broussonet) Goode and Bean, Proc. U. S. Nat. Mus., VIII,
 1885, 205 (*Cobitis heteroclitus* Linnæus).
Gambusinus Jordan and Evermann, Fish. North and Mid. Amer.,
 1896, 633 (*Fundulus rathbuni* Jordan and Meek).
Galasaccus Fowler, Proc. Acad. Nat. Sci. Phila., LXVIII, 1916, 417
 (*Hydrargyra similis* Baird and Girard).

1337. *Fundulus heteroclitus* (Linnæus). *Common killifish; Minnow; Pike*
minnow; Mummichog; Mud-minnow; Muddabbler; Cobbler.
 Coast of Maine to the Rio Grande; Gulf coast; common in brackish
 waters.

Cobitis heteroclitus Linnæus, Syst. Nat., ed. XII, 1766, 500, Charleston,
 S. C.
Pæcilia cœnicola Bloch and Schneider, Syst. Ichth., 1801, 452, Carolinas.
Fundulus mudfish Lacépède, Hist. Nat. Poiss., V, 1803, 37, Carolinas.
Hydrargyra swampina Lacépède, loc. cit., 378, South Carolina.
Fundulus nigerius Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 456,
 reported from Gaboon, West Africa, appears to be *heteroclitus*, wrong
 locality.

1338. *Fundulus heteroclitus fonticola* Cuvier and Valenciennes.

West Indies.

Fundulus fonticola Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII,
 1846, 148 (198), mountain springs of Porto Rico.
Fundulus antillarum Fowler, Proc. Acad. Nat. Sci. Phila., LXVIII,
 1916, 418, fig. 1, St. Martins, West Indies.

1339. *Fundulus heteroclitus bermudæ* Günther. *Mangrove minnow.*

Bermudas.

Fundulus bermudæ Günther, Ann. Mag. Nat. Hist., ser. 4, XIV, 1874,
 370, Bermudas.
Fundulus rhizophoræ Goode, Amer. Jour. Sci. Arts, XIII, 1877, 298,
 Basden Pond, Bermudas.

- 1340. *Fundulus heteroclitus macrolepidotus* (Walbaum).** *Common cobbler; Killifish; Catfish; Dogfish; Killy; Killie; River minnow; Mummichog; Blunthead.*
 Maine to Virginia; common in brackish waters. Hardly distinct from *F. heteroclitus*, of which it is the northern representative.
Cobitis macrolepidota Walbaum, Artedi Pisc., III, 1792, 11, Long Island; after Schöpf.
Cobitis killifish Walbaum, loc. cit., 12, Long Island; after Schöpf.
Esox pisciculus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 440, New York.
Esox pisculentus Mitchell, loc. cit., 441, New York.
Hydrargyra nigrofasciata LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 133, Newport, R. I.
Fundulus viridescens DeKay, New York Fauna: Fishes, 1842, 217, New York.
Fundulus zebra DeKay, loc. cit., 218, New York.
Hydrargyra ornata LeSueur, loc. cit., 131, Delaware River, near Philadelphia.
- 1341. *Fundulus heteroclitus grandis* Baird and Girard.**
 Gulf coast of the United States.
Fundulus grandis Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 389, Indianola, Tex.
Fundulus floridensis Girard, loc. cit., XI, 1859, 157, Charlotte Bay, Fla.
Zygoneutes funduloides Evermann, Bull. U. S. Fish Com., XI, 1891, 85, pl. 35, fig. 3, Dickinson Bayou, Dickinson, Tex.
- 1342. *Fundulus heteroclitus badius* Garman.**
 Gulf of St. Lawrence.
Fundulus heteroclitus badius Garman, Mem. Mus. Comp. Zool., XIX, 1895, Grand Manan.
- 1343. *Fundulus majalis* (Walbaum).** *Killifish; Mayfish; Rockfish; Catfish; Bass killy.*
 Cape Cod to Florida, chiefly marine in shallow bays.
Cobitis majalis Walbaum, Artedi Pisc., II, 1792, 12, Long Island; after *Mayfish* Schöpf.
Esox flavulus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 439, New York.
Esox zonatus Mitchell, loc. cit., 443, New York.
Hydrargyra trifasciata Storer, Jour. Bost. Soc. Nat. Hist., I, 1837, 417, Massachusetts.
Hydrargyra vernalis Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 154 (206), pl. 531; probably South Carolina.
- 1344. *Fundulus ocellaris* Jordan and Gilbert.**
 Pensacola to Fort Myers, Fla.
Fundulus ocellaris Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 254, Pensacola, Fla.
- 1345. *Fundulus parvipinnis* Girard.**
 Coast of California from Point Conception southward to Lower California.
Fundulus parvipinnis Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 154, San Diego, Calif.
- 1346. *Fundulus parvipinnis brevis* Osburn and Nichols.**
 Magdalena Bay, Lower California.
Fundulus parvipinnis brevis Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXV, 1916, 150, fig. 5, Magdalena Bay, Lower California.
- 1347. *Fundulus pallidus* Evermann.**
 Galveston Bay near Swan Lake, Tex.
Fundulus pallidus Evermann, Bull. U. S. Fish Com., XI, 1891 (1892), 84, pl. 35, Galveston Bay near Swan Lake, Tex.
- 1348. *Fundulus vinculus* Jordan and Gilbert.**
 Coast of Lower California.
Fundulus vinculus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 355, Cape San Lucas, Lower California.

- 1349. *Fundulus confluentus* Goode and Bean.**
 Streams of Florida.
Fundulus confluentus Goode and Bean, Proc. U. S. Nat. Mus., II, 1879,
 118, Lake Monroe, Fla.
- 1350. *Fundulus albolineatus* Gilbert.**
 Tennessee Basin in Alabama.
Fundulus albolineatus Gilbert, Bull. U. S. Fish Com., IX, 1889 (1891),
 149, Spring Creek, Huntsville, Ala.
- Subgenus GALASACEUS Fowler.**
- 1351. *Fundulus similis* (Baird and Girard). *Sac-à-lait.***
 Coast of the Gulf States.
Hydrargyre similis Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI,
 1853, 389, Indianola, Tex.
- Genus 444. ZYGONECTES Agassiz.**
Zygonectes Agassiz, Amer. Jour. Sci. Arts, 1854, 135 (*Semotilus olivaceus*
 Raf.=*Semotilus notatus* Raf.).
- 1352. *Zygonectes notatus* (Rafinesque). *Top minnow.***
 Michigan to Alabama, Mississippi, and Texas; generally abundant in
 ponds and canals.
Semotilus notatus Rafinesque, Ichth. Ohiensis, 1826, 86, tributaries of
 Ohio River in Kentucky; vaguely described but recognizable.
Pæcilia olivacea Storer, Proc. Bost. Soc. Nat. Hist., II, 1845 (1848),
 51, Florence, Ala.
Fundulus tenellus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI,
 1853, 389, Prairie Mer Rouge, La., and Russellville, Ky.
Zygonectes lateralis Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 353,
 Mobile, Ala.
Zygonectes zonatus Agassiz, loc. cit., 353, St. Louis, Mo.
Zygonectes pulchellus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859,
 113, Sugar Loaf Creek, Ark.
Fundulus aureus Cope, loc. cit., XVII, 1865, 78, Sugar Loaf Creek, Ark.
- 1353. *Zygonectes sciadicus* (Cope).**
 Middle Missouri River Basin; abundant in ponds and sluggish creeks
 in eastern Nebraska and southeastern South Dakota.
Fundulus sciadicus Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 78,
 Platte River, Nebr.
Haplochilus floripinnis Cope, Proc. Amer. Phil. Soc., XIV, 1874, 138,
 Platte River, at Denver, Colo.
Zygonectes lineatus Garman, Bull. Mus. Comp. Zool., VIII, 1881, 88,
 northeastern Wyoming.
Zygonectes macdonaldi Meek, Bull. U. S. Fish Com., IX, 1889, 122, pl.
 42, fig. 1, Jones Creek, Dixon, Mo., Osage fork of Gasconade River
 at Mansfield, Mo.
Fundulus scartes Meek, loc. cit., XV, 1895, 347, St. Francis River, Big
 Bay, Ark.
- 1354. *Zygonectes chrysotus* (Holbrook).**
 Coastwise swamps, South Carolina to Florida; rather common.
Fundulus chrysotus Holbrook, in Günther, Cat., VI, 1866, 317, Char-
 leston, S. C.
Zygonectes henshalli Jordan, Proc. U. S. Nat. Mus., II, 1879, 237, San
 Sebastian River, Fla.
Gambusia arlingtonia Goode and Bean, loc. cit., 118, Arlington River,
 Fla.
- 1355. *Zygonectes cingulatus* (Cuvier and Valenciennes).**
 New Jersey to Florida, in coastwise swamps, not rare; Escambia River
 at Flomaton, Ala.
Fundulus cingulatus Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII,
 1846, 147 (197), United States.
Zygonectes rubrifrons Jordan, Proc. U. S. Nat. Mus., II, 1879, 237, San
 Sebastian River, Fla.
Zygonectes auroguttatus Hay, loc. cit., VIII, 1885, 556, Westville, Fla.

1356. *Zygonectes dispar* Agassiz. *Top minnow.*
 Northern Ohio to Illinois, south to Mississippi; Maumee, Wabash, Big Black and Pearl Rivers.
Zygonectes dispar Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 353, creeks opposite St. Louis; Beardstown, Ill.
1357. *Zygonectes nottii* Agassiz. *Star-headed minnow; Top minnow.*
 Swamps and streams of Florida and neighboring States; not rare.
Zygonectes nottii Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 353, Mobile, Ala.
Zygonectes lineolatus Agassiz, loc. cit., 353, Augusta, Ga.
Zygonectes hieroglyphicus Agassiz, loc. cit., 353, Mobile, Ala.
Fundulus zonatus Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 146 (196), interior of South Carolina; not *Esox zonatus* Mitchell.
Zygonectes craticula Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 433, Elbow Creek, a tributary of Indian River, eastern Florida.
Zygonectes zonifer Jordan and Meek, loc. cit., VII, 1884, 482, Allapaha River, Nashville, Ga.
1358. *Zygonectes guttatus* Agassiz.
 Alabama.
Zygonectes guttatus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 353, Mobile, Ala.
Zygonectes escambiae Bollman, Proc. U. S. Nat. Mus., IX, 1886, 463, Escambia River, Flomaton, Ala.
1359. *Zygonectes luciae* (Baird).
 Atlantic coast from Long Island to Virginia; rare.
Hydrargyra luciae Baird, Ninth Smithson. Rept., 1854, 344, Beesleys Point, N. J.
1360. *Zygonectes pulvereus* Evermann.
 Coast of Texas.
Zygonectes pulvereus Evermann, Bull. U. S. Fish Com., XI, 1891 (1892), 85, Dickinson Bayou; Buffalo Bayou at Houston; Oso Creek at Corpus Christi, Tex.
1361. *Zygonectes jenkinsi* Evermann.
 Coast of Texas.
Zygonectes jenkinsi Evermann, Bull. U. S. Fish Com., XI, 1891 (1892), 86, Dickinson Bayou, arm of Galveston Bay, Tex.
1362. *Zygonectes limi* (Vaillant).
Fundulus limi Vaillant, Bull. Soc. Philom. (8), VI, 1894, 71, Lower California.
Fundulus meeki Evermann, Proc. Biol. Soc. Wash., XXI, 1908 (Jan. 23), 25, fig. 1, San Ignacio, central Lower California.
1363. *Zygonectes melapleurus* (Gosse). *Ticky-ticky.*
 Streams of Jamaica.
Pæclia melapleura Gosse, Naturalists' Sojourn in Jamaica, 1851, 84, Jamaica.
- Subgenus FONTICOLA Jordan and Evermannn.
1364. *Zygonectes seminolis* (Girard).
 Florida.
Fundulus seminolis Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 60, Palatka, eastern Florida.
- Subgenus MENONA Jordan and Evermann.
1365. *Zygonectes diaphanus* (LeSueur).
 Maine to North Carolina in river mouths; New York to northern Illinois in lakes.
Hydrargyra diaphana LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 130, Saratoga Lake, N. Y.
Hydrargyra multifaciata LeSueur, loc. cit., 131, Saratoga Lake, N. Y.
1366. *Zygonectes diaphanus menona* (Jordan and Copeland).
 Streams and lakes of Ohio to Iowa, Minnesota, and South Dakota.
Fundulus menona Jordan and Copeland, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 68, Catfish River, Dane County, Wis., outlet of Lake Menona, Madison, Wis.

Subgenus GAMBUSINUS Jordan and Evermann.

1367. *Zygonectes rathbuni* (Jordan and Meek).
Eastern North Carolina, abundant in spring brooks, not descending to salt water.
Fundulus rathbuni Jordan and Meek, Proc. U. S. Nat. Mus., XI, 1888, 356, Reedy Fork, Allemane Creek, Buffalo Creek, and other tributaries of the Cape Fear River, about Greensboro, N. C.
1368. *Zygonectes extensus* (Jordan and Gilbert).
Coast of Lower California.
Fundulus extensus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 355, Cape San Lucas, Lower California.
- Genus 445. *XENISMA* Jordan.
Xenisma Jordan, Bull. Buffalo Soc. Nat. Hist., 1876, 142 (*X. stellifera* Jordan).
1369. *Xenisma catenatum* (Storer). *Studfish*.
Tennessee and Cumberland Rivers and clear streams of the Ozark Mountains.
Pacilia catenata Storer, Synopsis, 1846, 430, Tennessee River, Florence, Ala.
1370. *Xenisma stellifer* Jordan. *Studfish*.
Alabama River and tributaries.
Xenisma stellifera Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 322, Etowah and Oostanaula Rivers, Rome, Ga.
- Genus 446. *PLANCTERUS* Garman.
Plancterus Garman, Mem. Mus. Comp. Zool., XIX, 1895, 96 (*Fundulus kansæ* Garman).
1371. *Plancterus zebra* (Girard).
Kansas, western Iowa and South Dakota to Texas and New Mexico, in upper tributaries of Kansas and Red Rivers.
Hydrargyre zebra Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 61, tributaries of Rio Grande "between Fort Defiance and Fort Union, N. Mex." name preoccupied in *Fundulus*.
Fundulus adinæ Jordan and Gilbert, Synopsis, 1883, 335, Rio Grande at Brownsville, Tex.
1372. *Plancterus kansæ* (Garman).
Kansas.
Fundulus kansæ Garman, Mem. Mus. Comp. Zool., XIX, 1895, 103, pl. 2, fig. 10, Kansas.
Fundulus zebrinus Jordan and Gilbert, Synopsis, 1882, 891; not of Girard.
- Genus 447. *OXYZYGONECTES* Fowler.
Oxyzygonectes Fowler, Proc. Acad. Nat. Sci. Phila., LXVIII, 1916, 425 (*Haplochilus dovii* Günther).
1373. *Oxyzygonectes balboæ* (Fowler).
Panama; only one example known.
Fundulus balboæ Fowler, Proc. Acad. Nat. Sci. Phila., LXVIII, 1916, 423, fig. 2, Panama.
1374. *Oxyzygonectes dovii* (Günther).
Coast of Costa Rica.
Haplochilus dovii Günther, Cat., VI, 1866, 316, Punta Arenas, Costa Rica.
- Genus 448. *RIVULUS* Poey.
Rivulus Poey, Memorias, II, 1861, 307, (*R. cylindraceus* Poey).
Cynodonichthys Meek, Field Col. Mus. Zool., V, 1904, 101 (*C. tenuis* Meek).
1375. *Rivulus cylindraceus* Poey.
Cuba.
Rivulus cylindraceus Poey, Memorias, II, 1861, 308, stream at Mardazo, near Habana, Cuba.
Rivulus marmoratus Poey, Anál. Soc. Espaniola Hist. Nat., IX, 1880, 248, Cuba.

1376. *Rivulus heyiei* Nichols.
Haiti.
Rivulus heyiei Nichols, Bull. Amer. Mus. Nat. Hist., XXXII, 1914, 143, Haiti.
1377. *Rivulus isthmensis* Garman.
Streams of Panama and Costa Rica.
Rivulus isthmensis Garman, Mem. Mus. Comp. Zool., XIX, 1895, 140, Rio San Jose, Costa Rica.
1378. *Rivulus brunneus* Meek and Hildebrand.
Panama.
Rivulus brunneus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 86, Toro Point, Canal Zone, Panama.
1379. *Rivulus flabellicauda* Regan.
Streams of Costa Rica.
Rivulus flabellicauda Regan, Ann. Mag. Nat. Hist., XIX, 1907, 64, Juan Vinas, Costa Rica.
1380. *Rivulus godmanni* Regan.
Rivulus godmanni Regan, Ann. Mag. Nat. Hist., 7th ser., XIX, 1907, 65, Guatemala.
1381. *Rivulus tenuis* (Meek).
Cynodontichthys tenuis Meek, Field Col. Mus., Zool., V, 1904, 101, El Hule, Oaxaca, Mexico.
1382. *Rivulus hartii* (Boulenger). *Rainfish; Small guabin.*
Mountain streams of Trinidad and Granada in rain-water pools.
Haplochilus hartii Boulenger, Proc. Zool. Soc. Lond., I, 1906, 389, Pl. XXI, fig. 2, Trinidad.
1383. *Rivulus chucunaque* Breder.
Tuya Basin, Pacific slope of Panama.
Rivulus chucunaque Breder (C. M.), Amer. Mus. Novitates, No. CLXXX, July 6, 1925, fig. 6, Rio Chucunaque, near Yavisa, Darien, Panama.
Rivulus chucunaque succubti Breder, loc. cit., 8, fig. 7, Rio Sucubti, Darien, Panama.
- Genus 449. *ADINIA* Girard.
Adinia Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 117 (*A. multifasciata* Girard).
1384. *Adinia multifasciata* Girard.
Gulf Coast, west Florida to Texas; locally abundant in shallow lagoons.
Adinia multifasciata Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 118, Galveston; St. Joseph Island, Indianola, Tex.
Fundulus xenicus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 255, Laguna Grande, Pensacola, Fla.; substitute for *multifasciata*, preoccupied in *Fundulus* but not in *Adinia*.
- Genus 450. *FLORIDICHTHYS* Hubbs.
Floridichthys Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 16, July 9, 1926, 16 (*Cyprinodon carpio* Günther).
1385. *Floridichthys carpio* (Günther).
Florida to Texas.
Cyprinodon carpio Günther, Cat., VI, 1866, 306, America.
Cyprinodon myodus Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 433, Pensacola, Fla.
- Genus 451. *JORDANELLA* Goode and Bean.
Jordanella Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 117 (*Jordanella floridæ* Goode and Bean).
1386. *Jordanella floridæ* Goode and Bean.
Streams from Florida to Yucatan.
Jordanella floridæ Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 117, Lake Monroe, Fla.
- Genus 452. *CYPRINODON* Lacépède.
Cyprinodon Lacépède, Hist. Nat. Poiss., V, 1803, 486 (*C. variegatus* Lacépède).
Prinodon Rafinesque, Anal. Nat., 1815, 88 (*Cyprinodon variegatus* Lacépède).
Encrotes Gistel, Thierreich, IX, 1848 (*Cyprinodon variegatus* Lacépède).
Trifarcus Poey, Memorias, II, 1861, 306 (*T. riverendi* Poey).

- 1387.** *Cyprinodon variegatus* Lacépède. *Killifish; Pursy minnow; Sheep-head minnow; Variegated minnow; Mummichog.*
 Cape Cod to Rio Grande, in bays, rarely entering streams.
Cyprinodon variegatus Lacépède, Hist. Nat. Poiss., V, 1803, 486, South Carolina.
Esox ovinus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 441, New York.
Lebias ellipsoidea LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 6, Florida.
Lebias rhomboidalis Valenciennes, in Humboldt, Obs. Zool. Phil. Mag., II, 1810 (1811), 100, Lake Pontchartrain, La.
Cyprinodon gibbosus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 390, Indianola, Tex.
- 1388.** *Cyprinodon riverendi* (Poey).
 Cuba and Florida Keys; common at Key West in shallow shore waters.
Trifarcus riverendi Poey, Memorias, II, 1861, 306, Habana; scarcely distinct from *C. variegatus*.
- 1389.** *Cyprinodon dearborni* Meek.
 Curaçao, Dutch West Indies.
Cyprinodon dearborni Meek, Field Mus. Nat. Hist., Zool., VII, 1909, 208, Willemstad, Curaçao, Dutch West Indies.
- 1390.** *Cyprinodon felicianus* (Poey).
 Cuba.
Trifarcus felicianus Poey, Synopsis, 1867, 412, Habana, Cuba.
- 1391.** *Cyprinodon latifasciatus* Garman.
 Parras, Coahuila, Mexico.
Cyprinodon latifasciatus Garman, Bull. Mus. Comp. Zool., VIII, 1881, No. 3, 92, Parras, Coahuila, Mexico.
- 1392.** *Cyprinodon bovinus* Baird and Girard.
 Leon Spring in southwestern Texas.
Cyprinodon bovinus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 389, Leon Spring, Tex.
- 1393.** *Cyprinodon martae* Steindachner.
 Santa Marta, coast of Colombia, South America.
Cyprinodon martae Steindachner, Ichth. Beitr., IV, 1875, 61, Santa Marta, Rio Cauca.
- 1394.** *Cyprinodon rubrofluviatilis* Fowler.
 Streams of Texas.
Cyprinodon bovinus rubrofluviatilis Fowler, Proc. Acad. Nat. Sci. Phila., LXVIII, 1916, 430, fig. 4, Rio Brazos, Seymour, Tex., and Red River, Tulip and Clarenden, Tex.
- 1395.** *Cyprinodon elegans* Baird and Girard.
 Tributaries of the Rio Grande.
Cyprinodon elegans Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 389, Comanche Spring, Tex.
Cyprinodon eximius Girard, loc. cit., XI, 1859, 158, Chihuahua River, Mexico.
- 1396.** *Cyprinodon macularius* Baird and Girard.
 Springs and streams of the desert, from southern Nevada to Sonora, locally abundant, often found in Artesian water and desert water holes in southwestern California.
Cyprinodon macularius Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 389, Rio San Pedro, Ariz.
Cyprinodon californiensis Girard, loc. cit., XI, 1859, 158, San Diego County, Calif., probably from Salt Springs in the desert (now Imperial County).
Lucania browni Jordan and Richardson, Proc. U. S. Nat. Mus., XXXIII, 1908, 319, hot spring in southeastern California.
- 1397.** *Cyprinodon nevadensis* Eigenmann.
 Death Valley.
Cyprinodon nevadensis Eigenmann, Proc. Calif. Acad. Sci., I, 1889, 270, Saratoga Springs, Death Valley, Inyo County, Calif.

1398. *Cyprinodon baileyi* Gilbert.

Pahranagat Valley, Nev.

Cyprinodon macularius baileyi Gilbert, Death Valley Exped., 1893, 233,
Pahranagat Valley, Nev.

Family 114. EMPETRICHTHYIDÆ

Genus 453. EMPETRICHTHYS³⁴ Gilbert.

Empetrichthys Gilbert, Death Valley Exped., Fishes, 1893, 233 (*E. merriami* Gilbert).

1399. *Empetrichthys merriami* Gilbert.

Death Valley in southeastern California.

Empetrichthys merriami Gilbert, Death Valley Exped., in North American Fauna, No. 7, May 31, 1893, 234, Ash Meadows, Amargosa Desert, on boundary between California and Nevada.

Family 115. GOODEIDÆ

Genus 454. ZOOGONETICUS Meek.

Zoogeneticus Meek, Field Col. Mus., Zool., III, 1902, 91 (*Z. diazi* Meek).

1400. *Zoogeneticus dugesii* (Bean).

Guanajuato, Mexico.

Fundulus dugesii Bean, Proc. U. S. Nat. Mus., X, 1887, 373, pl. 20, Guanajuato, Mexico.

1401. *Zoogeneticus diazi* Meek.

Lakes of Michoacan, Mexico.

Zoogeneticus diazi Meek, Field Col. Mus., Zool., III, 1902, 93, figured in Field Col. Mus., Zool., V, 1904, 114, fig. 32, Lake Patzcuaro, Mexico.

Zoogeneticus miniatius Meek, Field Mus. Nat. Hist., Zool., V, 1914, 115, fig. 33, Lake Zirahuén, Mexico.

1402. *Zoogeneticus robustus* (Bean).

Rio Lerma, Guanajuato.

Fundulus robustus Bean, Proc. U. S. Nat. Mus., XV, 1892, 285, pl. 44, fig. 2, Guanajuato, Mexico.

Zoogeneticus maculatus Regan, Ann. Mag. Nat. Hist. (7), XIII, 1904, 256, Rio Santiago (Lerma), Mexico.

1403. *Zoogeneticus quitzeoensis* (Bean).

Rio Lerma, Mexico.

Platypacilus quitzeoensis Bean, Proc. U. S. Nat. Mus., XXI, 1899 (Nov. 21, 1898), 540, fig. 1, Lake Quitzco, Lerma Basin.

Genus 455. GIRARDINICHTHYS Bleeker.

Girardinichthys Bleeker, Act Soc. Sci. Indo-Nederl., VII, 1860, 481 (*G. innominatus* Bleeker, after a "Lucania sp." of Girard).

Limnurgus Günther, Cat., VI, 1866, 309 (*L. variegatus* Günther = *Girardinichthys* Bleeker).

1404. *Girardinichthys innominatus* Bleeker.

Valley of Mexico.

Lucania sp., Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 118, City of Mexico.

Girardinichthys innominatus Bleeker, Act. Soc. Sci. Indo-Nederl., VII, 1860, 481, City of Mexico.

Limnurgus variegatus Günther, Cat., VI, 1866, 309, City of Mexico.

Characodon geddesi Regan, Ann. Mag. Nat. Hist., (7), XIII, 1904, 257, Xochomilco, Morelos, Mexico.

Genus 456. CHARACODON Günther.

Characodon Günther, Cat., VI, 1866, 308 (*C. lateralis* Günther).

³⁴This genus apparently a desert derivative of the extinct genus, *Parafundulus*, has probably no affinity with *Orestias* of Peru.

- 1405. *Characodon variatus* Bean.**
 Tributaries of Rio Lerma about Guanajuato, Mexico; locally common.
Characodon variatus Bean, Proc. U. S. Nat. Mus., X, 1887, 370, pl. 20,
 fig. 1, Guanajuato, Mexico.
Characodon ferrugineus Bean, loc. cit., 372, pl. 20, figs. 3 and 4, Guanajuato, Mexico.
Characodon eiseni Rutter, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 266,
 branch of the Rio Grande de Santiago at Tepic, Mexico.
- 1406. *Characodon lateralis* Günther.**
 Streams of Central America.
Characodon lateralis Günther, Cat., VI, 1866, 308, Central America.
- 1407. *Characodon garmani* Jordan and Evermann.**
 Tributaries of Rio Grande in Coahuila, Mexico.
Characodon garmani Jordan and Evermann, Fish. North and Mid.
 Amer., III, 1898, 2831, Parras, Coahuila, Mexico.
- 1408. *Characodon furcidens* Jordan and Gilbert.**
 Cape San Lucas southward to Colima, Mexico.
Characodon furcidens Jordan and Gilbert, Proc. U. S. Nat. Mus., V,
 1882, 354, Cape San Lucas, Lower California.
- Genus 457. *LERMICHTHYS* Hubbs.**
Lermichthys Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 16, July 9,
 1926, 18 (*Characodon multiradiatus* Meek).
- 1409. *Lermichthys multiradiatus* (Meek).**
 Rio Lerma Basin, Mexico.
Characodon multiradiatus Meek, Field Col. Mus., Zool., V, 1904, 119,
 Rio Lerma, Mexico.
Girardinichthys limnurgus Jordan and Evermann, Proc. Calif. Acad.
 Sci., 4th ser., XVI, Apr. 27, 1927, 502, Lake Lerma, Mexico.
- Genus 458. *BALSADICHTHYS* Hubbs.**
Balsadichthys Hubbs, Univ. Mich. Mus. Zool., Misc. Pub. No. 16, July
 9, 1926, 19 (*Goodea whitei* Meek).
- 1410. *Balsadichthys whitei* (Meek).**
 Rio de las Balsas Basin, Mexico.
Goodea whitei Meek, Field Col. Mus., Zool., V, 1904, 137, fig. 40, Rio
 Balsas, Yantepet, Cuautla, Mexico.
- Genus 459. *SKIFFIA* Meek.**
Skiffia Meek, Field Col. Mus., Zool., III, 1902, 102 (*S. lermæ* Meek).
- 1411. *Skiffia variegata* Meek.**
 Lake Zirahuen, Mexico.
Skiffia variegata Meek, Field Col. Mus., Zool., V, 1904, 143, Lake
 Zirahuen, Mexico.
- 1412. *Skiffia lermæ* Meek.**
Skiffia lermæ Meek, Field Col. Mus., Zool., III, 1902, 102, Lake Patzcuaro (Rio Lerma), Mexico.
- 1413. *Skiffia bilineata* (Bean).**
 Rio Lerma, Guanajuato, Mexico.
Characodon bilineatus Bean, Proc. U. S. Nat. Mus., X, 1887, 371,
 Guanajuato, Rio Lerma, Mexico.
- 1414. *Skiffia multipunctata* (Pellegrin).**
 Rio Lerma system; Lake Chapala, Mexico.
Xenendum multipunctatum Pellegrin, Bull. Mus. Hist. Nat., Paris, 1901,
 205, Jalisco, Mexico.
- Genus 460. *GOODEA* Jordan.**
Goodea Jordan, Proc. U. S. Nat. Mus., II, 1879, 299 (*G. atripinnis*
 Jordan).
Xenendum Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 128,
 fig. 9 (1906) (*X. xaleicone*=*Characodon luitpoldi* Steindachner).
- 1415. *Goodea atripinnis* Jordan.**
 Tributaries of Rio Lerma, Pacific slope, Mexico.
Goodea atripinnis Jordan, Proc. U. S. Nat. Mus., II, 1879, 299, Leon,
 Guanajuato, Mexico.

1416. **Goodea calientis** (Jordan and Snyder).

Streams of central Mexico.

Xenendum caliente Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 127, fig. 8, Aguas Calientes, Mexico.

1417. **Goodea luitpoldi** (Steindachner).

Basin of Rio Lerma, Mexico.

Characodon luitpoldi Steindachner, Einige Fischarten Mexicos, 1895, 12,

pl. 3, figs. 3-36, Lago de Patzcuaro, Michoacan, Mexico.

Xenendum xaliscone Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1901), 128, fig. 9, Laguna de Chapala, near Ocotlan, Jalisco, Mexico.

1418. **Goodea toweri** Meek.

Rio Panuco, San Luis Potosi, Mexico.

Goodea toweri Meek, Field Col. Mus., Zool., V, 1904, 138, fig. 41, Rio Verde, San Luis Potosi, Mexico.

1419. **Goodea captiva** Hubbs.

Panuco Basin, Mexico.

Goodea captiva Hubbs, Occ. Papers, Mus. Zool., Univ. Mich., No. 148, 1924, 4; upper tributary of the Rio Panuco at Jesus Maria, Mexico.

Family 116. POECILIIDÆ. Top minnows

Genus 461. **BELONESOX** Kner.

Belonesox Kner, Sitzungsber., Kais. Akad. Wiss. Wien, 1860, XL, 419 (*B. belizanus* Kner).

1420. **Belonesox belizanus** Kner.

Southern Mexico, Honduras, and Guatemala.

Belonesox belizanus Kner, Sitzungsber., Kais. Akad. Wiss. Wien, XL, 1860, 419, with figure, Belize; largest of the Poeciliidæ.

Genus 462. **CHAPALICHTHYS** Meek.

Chapalichthys Meek, Field Col. Mus., Zool., III, 1902, 97 (*Characodon encaustus* Jordan and Snyder).

1421. **Chapalichthys encaustus** (Jordan and Snyder).

Lake Chapala, Mexico.

Characodon encaustus Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1901), 126, Laguna de Chapala, near Ocotlan, Jalisco, Mexico.

Genus 463. **HETEROPHALLUS** Regan.

Heterophallus Regan, Ann. Mag. Nat. Hist., (8), XV, 1914, 66 (*H. rachovii* Regan).

1422. **Heterophallus rachovii** Regan.

Streams of eastern Mexico.

Heterophallus rachovii Regan, Ann. Mag. Nat. Hist., (8), XV, 1914, 66, Fig. A, Vera Cruz, Mexico.

Genus 464. **GAMBUSIA** Poey. Guajacones.

Gambusia Poey, Memorias, I, 1854, 382 (*G. punctata* Poey).

Hemixiphophorus Bleeker, Nat. Tijdschr. Neder.-Indie, XX, 1854, 440; type not named.

Paragambusia Meek, Field Col. Mus., Zool., V, 1904, 133 (*Gambusia nicaraguensis* Günther).

Heterophallina Hubbs (subgenus), Univ. Mich. Mus. Zool. Misc. Pub. No. 16, July 9, 1926 (*Gambusia regani* Hubbs).

Arthrophallus Hubbs (subgenus), loc. cit., 38 (*Heterandria patruleis* Baird and Girard).

Schizophallus Hubbs (subgenus), loc. cit., 40 (*Gambusia holbrookii* Girard).

- 1423. *Gambusia patruelis* (Baird and Girard).** *Top minnow; Mosquitofish.*
 Ditches, marshes, and lagoons, South Atlantic and Gulf States, lowland streams of the southwest; lower Mississippi from St. Louis to Louisiana and the Rio Grande. Introduced from Galveston into Hawaii, California, Formosa, Luzon, etc., a very effective mosquito killer.
Heterandria patruelis Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 390, Rio Sabinal, Rio Nueces, and Elm Creek, Tex.
Heterandria affinis Baird and Girard, loc. cit., 390, Rio Medina and Rio Salado, Tex.
?Gambusia speciosa Girard, loc. cit., 1859, 121, Rio San Diego, Nuevo Leon, Mexico.
?Gambusia gracilis Girard, loc. cit., 1859, 121, Matamoros, Mexico; name preoccupied.
Gambusia humilis Günther, Cat., VI, 1866, 334; substitute for *gracilis*.
Haplochilus melanops Cope, Proc. Amer. Phil. Soc., 1870, 457, Neuse River, N. C.
Zygonectes (Micristius) brachypterus Cope, Bull., U. S. Nat. Mus., XX, 1880, 34, Trinity River, Fort Worth, Tex.
Zygonectes inurus Jordan, Proc., U. S. Nat. Mus., 1882, 143, Cache Creek, southern Illinois.
Gambusia modesta Ahl, Blätter für Aquarium- und Terrariakunde, XXXVI, 1923, 220,
Gambusia myersi Ahl, loc. cit., 1925, 36, with figure (substitute for *modesta* preoccupied).
- 1424. *Gambusia holbrooki* (Girard).** *Holbrook's top minnow.*
 New Jersey to Florida and Alabama.
Heterandria holbrooki Agassiz, in Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 61, Charleston, S. C.
Haplochilus melanops Cope, Proc. Amer. Phil. Soc., XI, 1870, 457, Neuse River, N. C.
Zygonectes atrilatus Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 84, Neuse River, Goldsboro, N. C.
- 1425. *Gambusia nobilis* (Baird and Girard).**
Heterandria nobilis Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 390, Leona and Comanche Spring, Rio Grande, Tex.
- 1426. *Gambusia senilis* Girard.**
 Rio Grande Basin.
Gambusia senilis Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 122, Chihuahua, Mexico.
- Subgenus HEMIXIPHOPHORUS Bleeker.**
- 1427. *Gambusia gracilis* (Heckel).**
 Lowland streams of eastern Mexico.
Xiphophorus gracilis Heckel, Sitzungsber. Kais. Akad. Wiss. Wien, I, 1848, 300, taf. 9, figs. 3 and 4, Orizaba, Mexico.
Gambusia heckeli Bleeker, Fam. Cypr., 1860, 485.
- Subgenus GAMBUSIA Poey.**
- 1428. *Gambusia punctata* Poey.** *Guajacon.*
 Cuba.
Gambusia punctata Poey, Memorias, I, 1854, 384, Rio Almendares, Cuba.
Gambusia picturata Poey, Repertorio, 1868, 410, San Diego de los Baños, Cuba.
Gambusia punctata punctulata Garman, Mem. Mus. Comp. Zool., XIX, 1895, 87, Remedios, Cuba.
- 1429. *Gambusia puncticulata* Poey.**
 Streams of Cuba.
Gambusia puncticulata Poey, Memorias, I, 1854, 386, Cuba.
- 1430. *Gambusia melapleura* (Gosse).** *Ticky-ticky.*
 Streams of Jamaica; locally common.
Pæcilia melapleura Gosse, Naturalist's Sojourn in Jamaica, 1851, 84, Jamaica.

1431. *Gambusia nicaraguensis* Günther.
 Southern Mexico and Central America.
Gambusia nicaraguensis Günther, Cat., VI, 1866, 366, Lake of Nicaragua.
Gambusia dovi Regan, Proc. Zool. Soc. Lond., 1913, 986, Lake Nicaragua.
Gambusia yucatana Regan, Ann. Mag. Nat. Hist., 8th ser., XIV, 1914, 67, Progreso, Yucatan.
1432. *Gambusia mcnieri* Fowler.
 Panama.
Gambusia mcnieri Fowler, Proc. Acad. Nat. Sci. Phila., LXVIII, 1916, 433, Panama.
1433. *Gambusia wrayi* Regan.
 Jamaica.
Gambusia wrayi Regan, Proc. Zool. Soc. Lond., 1913, 988, Pl. XCIX, figs. 3, 4 and text fig. 168B, Jamaica.
1434. *Gambusia caymanensis* Regan.
 West Indies.
Gambusia caymanensis Regan, Proc. Zool. Soc. Lond., 1913, 990, Grand Cayman, West Indies.
1435. *Gambusia gracilior* Regan.
 Jamaica.
Gambusia gracilior Regan, Proc. Zool. Soc. Lond., 1913, 989, Pl. XCIX, figs. 5, 6, and text fig. 168C, Jamaica.
1436. *Gambusia dominicensis* Regan.
 Haiti.
Gambusia dominicensis Regan, Proc. Zool. Soc. Lond., 1913, 989, Pl. XCIX, fig. 7 and text fig. 169C, San Domingo.
1437. *Gambusia oligosticta* Regan.
 Jamaica.
Gambusia oligosticta Regan, Proc. Zool. Soc. Lond., 1913, 988, Pl. XCIX, figs. 1, 2, and text fig. 169B, Jamaica.
1438. *Gambusia manni* Hubbs.
 Bahamas.
Gambusia manni Hubbs, Copeia, 1927, 62, New Providence, one of the Bahamas.
 Subgenus HETEROPHALLINA Hubbs.
1439. *Gambusia regani* Hubbs.
 Rio Panuco, northwestern Mexico.
Gambusia regani Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 16, July 9, 1926, 28, Forlon, Tamaulipas, Mexico, in the River Forlon.
1440. *Gambusia vittata* Hubbs.
 Santa Engracia, Tamaulipas, Mexico.
Gambusia vittata Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 16, July 9, 1926, 26, Rio Forlon, at Forlon, Tamaulipas, Mexico.
1441. *Gambusia panuco* Hubbs.
 Rio Panuco, Mexico.
Gambusia panuco Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 16, July 9, 1926, 30, Rio Valles, at Valles, San Luis Potosi, Mexico.
- Genus 465. **BRACHYRHAPHIS** Regan.
Brachyrhaphis Regan, Proc. Zool. Soc. Lond., 1913, 997 (*Gambusia rhabdophora* Regan).
1442. *Brachyrhaphis terrabensis* (Regan).
 Pacific streams of Costa Rica.
Gambusia terrabensis Regan, Ann. Mag. Nat. Hist., (7), XIX, 1907, 260, Rio Grande de Terraba, Costa Rica.
1443. *Brachyrhaphis rhabdophora* (Regan).
 Costa Rica.
Gambusia rhabdophora Regan, Ann. Mag. Nat. Hist., (8), II, 1908, 457
 Volcano of Tenorio and Rio Grande de Terraba, Costa Rica.

- 1444. *Brachyrhaphis olomina* (Meek).**
Valley of the Rio Grande de Tarcoles, Costa Rica.
Priapichthys olomina Meek, Field Mus. Nat. Hist., Zool., X, 1912, 114,
Las Cañas, Alajuela, Costa Rica.
- 1445. *Brachyrhaphis episcopi* (Steindachner).**
Isthmus of Panama at Obispo station, and streams of the Pacific slope
of Panama.
Gambusia episcopi Steindachner, Ichth. Beitr., VI, 1878, 9, Obispo near
Panama.
Gambusia latipunctata Meek and Hildebrand, Field Mus. Nat. Hist.,
Zool., X, 1913, 87, Arrijan, Panama.
- 1446. *Brachyrhaphis cascajalensis* (Meek and Hildebrand).**
Streams of Atlantic slope of Panama.
Gambusia cascajalensis Meek and Hildebrand, Field Mus. Nat. Hist.,
Zool., X, 1913, 86, Rio Cascajal, Panama.
- 1447. *Brachyrhaphis parismina* (Meek).**
Costa Rica.
Gambusia parismina Meek, Field Mus. Nat. Hist., Zool., X, 1912, 71,
Costa Rica.
- Genus 466. TRIGONOPHALLUS Hubbs.**
Trigonophallus Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 16,
July 9, 1926, 48 (*T. punctifer* Hubbs).
- 1448. *Trigonophallus punctifer* Hubbs.**
Streams of Pacific coast of Panama.
Trigonophallus punctifer Hubbs, Univ. Mich. Mus. Zool. Misc. Pub.
No. 16, July 9, 1926, 49, Quibari Creek, Rio Secamola at Kakintin.
- Genus 467. PRIAPICHTHYS Regan.**
Priapichthys Regan, Proc. Zool. Soc. Lond., 1913, 991 (*Gambusia*
annectens Regan).
- 1449. *Priapichthys annectens* (Regan).**
Streams of Atlantic slope of Costa Rica.
Gambusia annectens Regan, Ann. Mag. Nat. Hist., (7), XIX, 1907,
250, Costa Rica.
- 1450. *Priapichthys annectens hesperis* Hubbs.**
Streams of the Pacific slope of Costa Rica.
Priapichthys annectens hesperis Hubbs, Univ. Mich. Mus. Zool. Misc.
Pub. No. 13, Jan. 18, 1924, 22, Rio Maria Aquilar, Rio Grande de
Tancos, and their mountain tributaries.
- 1451. *Priapichthys huberi* Fowler.**
Nicaragua.
Priapichthys huberi Fowler, Proc. Acad. Nat. Sci. Phila., LXXV, 1923,
27, Marceligo Creek, tributary of the Tunku River, at Miranda,
Nicaragua.
- Genus 468. PANAMICHTHYS Hubbs.**
Panamicthys Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 13,
Jan. 18, 1924, 8 (*Priapichthys panamensis* Meek and Hildebrand).
- 1452. *Panamicthys panamensis* (Meek and Hildebrand).**
Priapichthys panamensis Meek and Hildebrand, Field Mus. Nat. Hist.,
Zool., X, 1916, 322, Panama.
- Genus 469. PSEUDOXIPHOPHORUS Bleeker.**
Pseudoxiphophorus Bleeker, Ichth. Indice Prodr. Cypr., in Acata Societ.
Scient. Indo-Neer., VII, 1859, 483 (*Xiphophorus bimaculatus* Heckel).
Pacilodes Steindachner, Sitzungsber. Kais. Akad. Wiss. Wien, XLVIII,
1863, 176 (*P. bimaculatus* Steindachner).
- 1453. *Pseudoxiphophorus bimaculatus* (Heckel).**
Central Mexico; Rio Blanco, Orizaba.
Xiphophorus bimaculatus Heckel, Sitzungsber. Kais. Akad. Wiss. Wien,
1, 1848, pt. 3, 169, taf. 6, figs. 1 and 2, Mexico.
Pseudoxiphophorus reticulatus Troschel, in Miller's Reisen Vereinigten
Staaten, Canada, Mexico, III, 1865, 638, no locality.

- 1454. *Pseudoxiphophorus bimaculatus tæniatus* Regan.**
 Streams of the Atlantic slope of southeastern Mexico.
Pseudoxiphophorus bimaculatus tæniatus Regan, Ann. Mag. Nat. Hist., (7), XVI, 1905, 303, Rio Touto and San Domingo de Guzman.
- 1455. *Pseudoxiphophorus bimaculatus jonesi* (Günther).**
 Streams of southeastern Mexico.
Mollisiania jonesi Günther, Ann. Mag. Nat. Hist., (4), XIV, 1874, 371, Lake Alcuhuaca, Huamanita, Mexico.
Pseudoxiphophorus pauciradiatus Regan, Ann. Mag. Nat. Hist., (7), X, 1904, 14, 296, Orizaba, Mexico.
- Genus 470. *HETERANDRIA* Agassiz.**
Heterandria Agassiz, Amer. Jour. Sci. Arts, XVI, 1853, 135 (*H. formosus* Agassiz MS.; *H. holbrooki* Agassiz MS.).
- 1456. *Heterandria formosa* Agassiz.**
 Streams of South Carolina to Florida; one of the smallest of fishes.
Heterandria formosa Agassiz, in Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 62, Charleston, S. C.; Palatka, Fla.
- Genus 471. *PRIAPELLA* Regan.**
Priapella Regan, Proc. Zool. Soc. Lond., 1913, 992 (*Gambusia bonita* Meek).
- 1457. *Priapella bonita* (Meek).**
 Streams of western Mexico; Rio Papaloapam, Motzorongo.
Gambusia bonita Meek, Field Col. Mus., Zool., V, 1904, 132, fig. 39, 104, Refugio, Vera Cruz, Mexico.
- Genus 472. *ALLOGAMBUSIA* Hubbs.**
Allogambusia Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 13, Jan. 18, 1924, 8 (*Gambusia tridentiger* Garman).
- 1458. *Allogambusia cana* (Meek and Hildebrand).**
 Streams of the Pacific slope of Panama.
Gambusia cana Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 87, Rio Satiganti, Cana, Panama.
- 1459. *Allogambusia tridentiger* (Garman).**
 Rivers of Panama.
Gambusia tridentiger Garman, Mem. Mus. Comp. Zool., XIX, 1895, 89 (Pl. IV, fig. 10, teeth), rivers of Panama.
- Genus 473. *TOXUS* Eigenmann.**
Toxus Eigenmann, Bull. U. S. Fish Com., XXII, 1902 (1903), 226 (*T. riddlei* Eigenmann).
- 1460. *Toxus riddlei* Eigenmann.**
 Streams of western Cuba.
Toxus riddlei Eigenmann, Bull. U. S. Fish Com., XXII, 1902 (1903), 226, San Cristobal, Cuba.
- 1461. *Toxus cubensis* (Eigenmann).**
 Streams of western Cuba.
Heterandria cubensis Eigenmann, Bull. U. S. Fish Com., XXII, 1902 (1903), 227, Los Palacios, Cuba.
- Genus 474. *GLARIDICHTHYS* Garman.**
Glaridodon Garman, Mem. Mus. Comp. Zool., XIX, 1895, 40 (*latidens*); name preoccupied.
Glaridichthys Garman, Amer. Nat., XXX, 1896, 232 (same type).
- 1462. *Glaridichthys uninotatus* (Poey).**
 Cuba.
Girardinus uninotatus Poey, Memorias, II, 1861, 309, Cuba.
- 1463. *Glaridichthys torralbasi* Eigenmann.**
 Streams of Pinar del Rio, Cuba.
Glaridichthys torralbasi Eigenmann, Bull. U. S. Fish Com., XXII, 1902 (1903), 225, fig. 4, Pinar del Rio, Cuba.
- 1464. *Glaridichthys falcatus* Eigenmann.**
 Streams of western Cuba.
Glaridichthys falcatus Eigenmann, Bull. U. S. Fish Com., XXII, 1902 (1903), 224, San Cristobal, Palacios, Pinar del Rio, Cuba.

Genus 475. GIRARDINUS Poey.*Girardinus* Poey, Memorias, I, 1854, 383 (*G. metallicus* Poey).

- 1465. *Girardinus metallicus* Poey.**

Streams of Cuba.

Girardinus metallicus Poey, Memorias, I, 1854, 387, Cuba.*Girardinus denticulatus* Garman, Mem. Mus. Comp. Zool., XIX, 1895, 47, Remedios, Cuba.*Girardinus creolus* Garman, loc. cit., 47, Cuba.*Girardinus garmani* Eigenmann, Bull. U. S. Fish Com., XXII, 1902 (1903), 296, Pinar del Rio, Palacios, Cuba.**Genus 476. DARIENICHTHYS Hubbs.***Darienichthys* Hubbs, Univ. Mich. Mus. Zool., Misc. Pub. No. 13, Jan. 18, 1924, 8 (*Gambusia darienensis* Meek and Hildebrand).

- 1466. *Darienichthys darienensis* (Meek and Hildebrand).**

Streams of Pacific slope, Panama.

Gambusia darienensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 116, 88, Rio Capeti (Tuyra River Basin), Panama.**Genus 477. PÆCILIOPSIS Regan.***Pæciliopsis* Regan, Proc. Zool. Soc. Lond., 1913, 996 (*Pæcilia presidionis* Jordan and Culver).*Leptorhaphis* Regan, loc. cit., 980, 998 (*Gambusia infans* Woolman).

- 1467. *Pæciliopsis occidentalis* (Baird and Girard).**

Gila Basin, Ariz., and basins of Yaqui and Sonora Rivers, Sonora.

Heterandria occidentalis Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 390, Rio Santa Cruz, near Tucson, Ariz.*Girardinus sonoriensis* Girard, loc. cit., XI, 1859, 120, San Bernardino Creek, a tributary of the Rio Yaqui, Sonora.

- 1468. *Pæciliopsis infans* (Woolman).**

Rio Lerma, Salamanca, Mexico.

Gambusia infans Woolman, Bull. U. S. Fish Com., XIV, 1894, 62, Rio Lerma, Salamanca, Mexico.

- 1469. *Pæciliopsis balsas* Hubbs.**

Balsas Basin, Mexico.

Pæciliopsis balsas Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 16, July 9, 1926, 66, Rio Balsas, Guerrero, Mexico.

- 1470. *Pæciliopsis latidens* (Garman).**

Chihuahua.

Glariodon latidens Garman, Mem. Mus. Comp. Zool., XIX, 1895, 40, Chihuahua, Mexico.

- 1471. *Pæciliopsis fasciata* (Meek).**

Atlantic streams of southern Mexico.

Gambusia fasciata Meek, Field Col. Mus., Zool., V, 1904, 129, fig. 37, San Geronimo, Oaxaca, Mexico.

- 1472. *Pæciliopsis turrubarensis* (Meek).**

Costa Rica.

Gambusia turrubarensis Meek, Field Mus. Nat. Hist., Zool., X, 1912, 71, Rio Turrubara, Costa Rica.*Heterandria colombiana* Eigenmann and Henn, Indiana Univ. Stud., No. 8, 1912, 27, brackish water, mouth of Rio Dagua, Colombia.*Priapichthys fosteri* Hildebrand, Bull. U. S. Bur. Fish., XLI, 1925, 260, figs. 14 and 15, Rio Lempa, San Marcos, El Salvador, Central America.

- 1473. *Pæciliopsis presidionis* (Jordan and Culver).**

Rio Presidio, Sinaloa, Mexico.

Pæcilia presidionis Jordan and Culver, in Jordan, Fishes of Sinaloa, Proc., Calif. Acad. Sci., V, 1895, 413, pl. 29, Rio Presidio, Sinaloa, Mexico.**Genus 478. PÆCILISTES Hubbs.***Pæciliistes* Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 16, July 9, 1926, 68 (*Heterandria lutzi* Meek).

1474. *Pœciliistes pleurospilus* (Günther).
 Streams of Guatemala.
Girardinus pleurospilus Günther, Cat., VI, 1866, 353, Lake Dueñas,
 Guatemala.
Heterandria lutzi Meek, Field Col. Mus., Zool., V, 1904, 148, fig. 47,
 Orizaba, Mexico.
Priapichthys letonai Hildebrand, Bull. U. S. Bur. Fish., XLI, 1925,
 258, Rio San Miguel, San Salvador.
- Genus 479. *AULOPHALLUS* Hubbs.
Aulophallus Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. No. 16, July
 9, 1926, 69 (*Pæcilia elongata* Günther).
1475. *Aulophallus elongatus* (Günther).
 Panama, a large marine species quite unlike *Pæcilia* in habits.
Pæcilia elongata Günther, Cat., VI, 1866, 342, Panama.
- Genus 480. *PHALLICHTHYS* Hubbs.
Phallichthys Hubbs, Univ. Mich. Mus. Zool. Misc. Pub. 13, Jan. 18,
 1924, 10, (*Pæcilopsis isthmensis* Regan).
1476. *Phallichthys amates* (Miller).
 Motagua Basin, Guatemala.
Pæcilia amates Miller, Bull. Amer. Mus. Nat. Hist., XXIII, 1907, 108,
 fig. 1, pond and its outlet, Los Amates, an affluent of the Motagua
 River, Guatemala.
1477. *Phallichthys pittieri* (Meek).
 Costa Rica.
Pæcilia pittieri Meek, Field Mus. Nat. Hist., Zool., X, 1912, 71, La
 Junta, Costa Rica.
Pæcilopsis isthmensis Regan, Proc., Zool. Soc. Lond., 1913, 997, Pl.
 C, figs. 3, 4, and text fig. 171B, Colon, Panama.
- Genus 481. *XENOPHALLUS* Hubbs.
Xenophallus Hubbs, Mus. Zool. Univ. Mich. Misc. Pub. 13, Jan. 18,
 1924, 10 (*Gambusia umbratilis* Meek.)
1478. *Xenophallus umbratilis* (Meek).
 Streams of Costa Rica.
Gambusia umbratilis Meek, Field Mus. Nat. Hist., Zool., X, 1912, 70,
 Costa Rica.
- Genus 482. *PŒCILIA* Schneider.
Pæcilia Schneider, ed. Bloch, Syst. Ichth. 1801, 452 (*P. vivipara* Bloch
 and Schneider); not *Pæcilius* of genus of beetles.
Alazon Gistel, Thierreich, X, 1848 (*Pæcilia vivipara* Schneider); sub-
 stitute for *Pæcilia*, regarded as preoccupied.
Acropæcilia Hilgendorf, Sitzungsb. Naturf. Freunde, 1889, 52 (*Pæcilia*
tridens Hilgendorf).
Neopæcilia Hubbs, Univ. Mich. Mus. Zool., Misc. Pub. 13, 1924 (*N.*
holacanthus Hubbs).
1479. *Pæcilia vivipara* Bloch and Schneider.
 Rivers of Brazil; Guiana; Martinique; Porto Rico.
Pæcilia vivipara Bloch and Schneider, Syst. Ichth., 1801, 452, pl. 86,
 fig. 2, Surinam.
Pæcilia surinamensis Valenciennes, Humboldt, Observ. Zool., II, 1817,
 158, pl. 51, fig. 1, Surinam.
Pæcilia schneideri Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII,
 1846, 102 (135), Surinam.
Neopæcilia holacanthus Hubbs, Univ. Mich. Mus. Zool., Misc. Pub.
 No. 13, Jan. 18, 1924, 11, based on material recorded as *Pæcilia vivi-
 para* Evermann and Marsh, Bull., U. S. Fish Com., XX, 1900, first
 part (1902), 97, from Porto Rico (Ponce, Fijaro, Arroyo, and Hu-
 cares).
1480. *Pæcilia tridens* Hilgendorf.
 Streams of Haiti.
Pæcilia (*Acropæcilia*) *tridens* Hilgendorf, Sitzungsb. Naturf. Freunde,
 1889, 52, Port au Prince.

Genus 483. LIMIA Poey.

Limia Poey, Memorias, I, 1854, 388 (*L. cubensis* Poey = *Pæcilia vittata* Guichenot).

1481. *Limia heterandria* Regan.

Santo Domingo.

Limia heterandria Regan, Proc. Zool. Soc. Lond., 1913, 1017, Pl. CI, "Venezuela"; locality probably wrong.

Heterandria zonata Nichols, Bull. Amer. Mus. Nat. Hist., XXXIV, 1915, 603, fig. 3, Santo Domingo.

1482. *Limia vittata* (Guichenot). *Faguito; Guajica.*

Streams of Cuba.

Pæcilia vittata Guichenot, in Ramon de la Sagra, Hist. Nat. Cuba, Poiss., 1853, 146, pl. 5, fig. 1, Cuba.

Limia cubensis Poey, Memorias, I, 1854, 388, Habana.

Limia pavonina Poey, Enumeratio, 1876, 142, Habana.

1483. *Limia perugiae* (Evermann and Clark).

Santo Domingo.

Platypæcius perugiae Evermann and Clark, Proc. U. S. Nat. Mus., XXX, 1906, 851, Santo Domingo.

1484. *Limia caudofasciata* Regan.

Jamaica.

Limia caudofasciata Regan, Proc., Zool. Soc. Lond., 1913, 1017, Jamaica.

1485. *Limia nigrofasciata* Regan.

Limia nigrofasciata Regan, Proc., Zool. Soc. Lond., 1913, 1015, Pl. CI, figs. 1, 2, 1913, Haiti.

Limia arnoldi Regan, loc. cit., 1016, Miragoâne, Haiti; young of *nigrofasciata*.

1486. *Limia ornata* Regan.

Limia ornata Regan, Proc. Zool. Soc. Lond., 1913, 1016, Haiti.

1487. *Limia dominicensis* (Cuvier and Valenciennes).

Santo Domingo and Barbados.

Pæcilia dominicensis Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 99 (131), pl. 525, Santo Domingo.

Pæcilia melanogaster Günther, Cat., VI, 1866, 345, locality unknown.

Platypæcius dominicensis Evermann and Clark, Proc. U. S. Nat. Mus., XXX, 1906, 852, fig. 2, streams in San Francisco Mountains, Santo Domingo.

1488. *Limia melanonotata* Nichols and Myers.

Santo Domingo.

Limia melanonotata Nichols and Myers, Amer. Mus. Novitates, No. 79, 1923, 1, Hispaniola.

1489. *Limia versicolor* (Günther).

Rivers of Haiti.

Girardinus versicolor Günther, Cat., VI, 1866, 352, Santo Domingo.

Genus 484. LEBISTES De Filippi

Lebistes de Filippi, Arch. Zool. Anat. Fisiol., I, 1862, 69 (*L. paciloides* de Filippi).

Acanthophacelus Eigenmann, Proc. U. S. Nat. Mus., XXXII, 1907, 426 (*Pæcilia reticulatus* Peters).

1490. *Lebistes reticulatus* (Peters). *Millions; Bellyfish; Guppy and rainbow-fish of aquarists.*

Pools and streams of Barbados, Trinidad, and Venezuela.

Pæcilia reticulata Peters, Monatsber. Akad. Wiss. Berlin, 1859, 412, Caracus, Venezuela.

Lebistes paciloides de Filippi, Arch. Zool. Anat. Fisiol., I, 1861, 69, Pl. IV, fig. 6, Barbados.

Girardinus guppri Günther, Cat., VI, 1866, 353, Trinidad; Venezuela.

Acanthophacelus melanzonus Eigenmann, Ann. Carnegie Mus., VI, 1909, 51, Georgetown Trenches, British Guiana.

Genus 485. MOLLIENESIA LeSueur.

Mollienesia LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 2 (*Mollinesia latipinna* LeSueur).

- 1491. *Mollienesia latipinna* LeSueur.** *Mudfish.*
 South Carolina to northern Mexico, in lowland swamps and streams.
Mollinesia latipinna LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 3,
 New Orleans.
Pæcilia multilineata LeSueur, loc. cit., 4, New Orleans.
Pæcilia lineolata Girard, U. S. Mex. Bound. Surv., Ichth., 1859, 70,
 pl. 35, figs. 9 to 11, Rio Grande near Brownsville, Tex.
Limia pæciloidæ Girard, loc. cit., 70, pl. 38, figs. 8 to 14, Indianola, Tex.
Limia matamorensis Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859,
 117, Matamoras.
- 1492. *Mollienesia gracilis* Regan.**
 Lake Peten, Guatemala.
Mollinesia petenensis Günther, Cat., VI, 1866, 348, Lake Peten; not
Pæcilia petenensis Günther, also a *Mollienesia*.
Mollinesia gracilis Regan, Proc. Zool. Soc. Lond., 1913, 1012; substitute
 name.
- 1493. *Mollienesia petenensis* (Günther).**
 Lake Peten, Guatemala.
Pæcilia petenensis Günther, Cat., VI, 1866, 342, Lake Peten, Guatemala;
 not *Mollinesia petenensis* Günther, loc. cit., 348.
- 1494. *Mollienesia formosa* (Girard).**
 Mexico; Lake Alcohuaca, Huamantla, Mexico; Palo Alto.
Limia formosa Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 115, Palo
 Alto, Mexico.
- 1495. *Mollienesia spilura* (Günther).**
 Streams of Central America.
Pæcilia spilurus Günther, Cat., VI, 1866, 345, Central America.
- 1496. *Mollienesia velifera* Regan.**
 Yucatan.
Mollinesia velifera Regan, Ann. Mag. Nat. Hist., 8th ser., XIII, 1914,
 338, Progreso, Yucatan.
- 1497. *Mollienesia sphenops* (Cuvier and Valenciennes).**
 Lowland streams of Mexico and Central America.
Pæcilia sphenops Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII,
 1836, 98 (130), 525, Vera Cruz.
Molinessa fasciata Müller and Troschel, Monatsber. Akad. Wiss. Berlin,
 1844, 36, Mexico.
Pæcilia mexicana Steindachner, Sitzungsbs., Kais. Akad. Wiss. Wien,
 XLVIII, 1863, 178, taf. 4, fig. 2, southern Mexico.
Pæcilia thermalis Steindachner, loc. cit., 181, warm springs in Central
 America.
Xiphophorus gilli Kner and Steindachner, Abhandl. Bayer. Akad.
 Wiss., X, 1864, 28, taf. 4, fig. 1 (1866), Rio Chagres, Panama.
Gambusia modesta Troschel, in Müller's Reisen Vereinigten Staaten,
 Canada and Mexico, III, 1865, 639, Mexico.
Pæcilia chisoyensis Günther, Cat., VI, 1866, 342, Rio Chisoy, Vera
 Paz.
Gambusia plumbea Troschel, loc. cit., III, 1865, 640, Mexico.
Pæcilia dovii Günther, loc. cit., VI, 1866, 344, Lake Nicaragua; Lake
 Amatitlan.
Platypoecilus mentalis Gill, Proc. Acad. Nat. Sci. Phila., XXVIII, 1876,
 335, Isthmus of Panama.
Pæcilia boucardi Steindachner, Ichth. Beitr., VI, 1878, 8, Colon (or
 Aspinwall).
Pæcilia limantouri Jordan and Snyder, Bull. U. S. Fish Com., XIX,
 1899, 129, fig. 10, Rio Tamesin, Tampico, Mexico.
Pæcilia butleri Jordan, Proc. U. S. Nat. Mus., XI, 1888, 330, Rio
 Presidio, Mazatlan.
Platypoecilus tropicus Meek, Field Col. Mus., Zool., VII, 1907, 146, Tur-
 rialba, Costa Rica.
Pæcilia tenuis Meek, loc. cit., 147, San Jose, Costa Rica.
Pæcilia salvatoris Regan, Ann. Mag. Nat. Hist., (7), XIX, 1907, 65,
 Salvador.
Pæcilia spilonota Regan, loc. cit., 8th ser., II, 1908, 460, San Jose.

1498. **Mollienesia sphenops cuneata** (Garman).
Turbo, Gulf of Darien.
Pacilia cuneata Garman, Mem., Mus. Comp. Zool., XIX, 1895, 62, Pl. V, fig. 3, teeth, *Turbo*, Gulf of Darien.
1499. **Mollienesia sphenops vandepolli** (Van Lindth de Jeude).
 Tropical America, abundant in fresh-water pools.
Pacilia vandepolli Van Lindth de Jeude, Notes, Leyden Mus., IX, 1887, 137, Freshwater, Curaçao.
Pacilia vandepolli arubensis Van Lindth de Jeude, loc. cit., 138, Aruba, one of the Leeward Islands.
- Genus 486. ALFARO Meek.**
Petalosoma Regan, Ann. Mag. Nat. Hist., (8), II, 1908, 462 (*P. cultratum* Regan, name preoccupied).
Alfaro Meek, Field Mus. Nat. Hist., Zool., X, September, 1912, 72 (*A. acutiventris* Meek).
Petalurichthys Regan, Ann. Mag. Nat. Hist., X, November 1912, 494 (*Petalosoma cultratum* Regan; substitute for *Petalosoma* preoccupied).
1500. **Alfaro cultratus** (Regan).
 Streams of Costa Rica.
Petalosoma cultratum Regan, Ann. Mag. Nat. Hist., XI, 1908, 458, Rio Iroquois, Costa Rica.
Alfaro acutiventralis Meek, Field Mus. Nat. Hist., Zool., X, 1912, 72, Costa Rica.
- Genus 487. PLATYPŒCILUS Günther.**
Platypæcilius Günther, Cat., VI, 1866, 350 (*P. maculatus* Günther).
1501. **Platypœcilius maculatus** Günther. *Moonfish of the aquarists.*
 Mexico.
Platypæcilius maculatus Günther, Cat., VI, 1866, 350, Mexico.
Platypæcilius variatus Meek, Field Col. Mus., Zool., V, 1904, 145, 146, San Luis Potosi, Mexico.
Platypæcilius maculatus immaculatus Myers, Copeia, No. 113, Dec 20, 1922, 89, a form developed in an aquarium, therefore without standing in scientific nomenclature.
1502. **Platypœcilius couchianus** (Girard).
 Rio Grande Basin.
Limia couchiana Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 116, Rio San Juan, at Cadereita and Monterey, in Nuevo Leon, Mexico.
- Genus 488. XIPHOPHORUS Heckel.**
Xiphophorus Bleeker, Sitzungsbs. Kais. Akad. Wiss. Wien, I, 1848, pt. 3, 163 (*X. helleri* Heckel).
1503. **Xiphophorus helleri** Heckel. *Swordtail.*
 Lowland streams of southern Mexico and Guatemala, of Atlantic drainage.
Xiphophorus helleri Heckel, Sitzungsbs. Kais. Akad. Wiss. Wien, I, 1848, pt. 3, 163, pl. 5, Rio Chisoy, Guatemala.
Xiphophorus jalapae Meek, Field Col. Mus., Zool., III, 1902, 107, Jalapa.
1504. **Xiphophorus helleri guntheri** (Jordan and Evermann).
 Rivers of eastern slope of Guatemala.
Xiphophorus guntheri Jordan and Evermann, Fish. North and Mid. Amer., 1896, 702, Rio Chisoy (Usumacinta), Guatemala.
1505. **Xiphophorus helleri strigatus** (Regan).
 Atlantic slope of southern Mexico.
Xiphophorus strigatus Regan, Ann. Mag. Nat. Hist., (7), XIX, 1907, 65; streams of Atlantic slope of southern Mexico; Cordoba, Oaxaca, etc.
1506. **Xiphophorus helleri rachovii** (Regan).
 Atlantic slope of Guatemala.
Xiphophorus rachovii Regan, Ann. Mag. Nat. Hist., (8), VIII, 1911, 374, Puerto Barrios between Lake Yzabal and Rio Motagua.

1507. **Xiphophorus helleri brevis** (Regan).
British Honduras.
Xiphophorus brevis Regan, Ann. Mag. Nat. Hist., (7), XIX, 1907, 65,
British Honduras.
1508. **Xiphophorus montezumæ** Jordan and Snyder.
Rio Panuco, eastern Mexico.
Xiphophorus montezumæ Jordan and Snyder, Bull. U. S. Fish Com.,
XIX, 1899 (1900), 131, fig. 11, Rio Panuco, Mexico.

Family 117. ANABLEPIDÆ. Four-eyed fishes

- Genus 489. ANABLEPS** (Gronow) Scopeli.
Anableps Gronow, in Scopeli, Introd. Hist. Nat., 1777, 450 (no type
mentioned; *A. anableps* L. understood).
1509. **Anableps anableps** Linnæus. *Cuatro ojos; Four-eyed fish.*
Trinidad to Brazil; in streams.
Anableps anableps Linnæus, Syst. Nat., ed. 9, 1756, 55.
Anableps tetraphthalmus Bloch, Naturgesch. Ausl. Fische, VIII, 1794,
7, pl. 361.
1510. **Anableps dowei** Gill. *Cuatro ojos.*
Streams of Central America, locally common (Salvador-Chiapas,
Guatemala).
Anableps dowei Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 4, La
Union, Salvador.

Family 118. AMBLYOPSIDÆ. Cave blindfishes

- Genus 490. CHOLOGASTER** Agassiz.
Chologaster Agassiz, Amer. Jour. Sci. Arts, XVI, 1853, 135 (*C. cornutus*
Agassiz).
1511. **Chologaster cornutus** Agassiz. *Fish of the rice ditches of South Atlantic
States.*
Lowland swamps of the Southern States, from the Dismal Swamp to
the Okefinokee Swamp; locally very abundant.
Chologaster cornutus Agassiz, Amer. Jour. Sci. Arts, XVI, 1853, 135,
ditch in a rice field in South Carolina.
1512. **Chologaster avitus** Jordan and Jenkins. *Fish of Dismal Swamp.*
Streams and swamps from southeast Virginia southward, probably
varying into *C. cornutus*.
Chologaster avitus Jordan and Jenkins, Proc. U. S. Nat. Mus., XI, 1888,
356, pl. 44, fig. 8, outlet of Lake Drummond, Dismal Swamp, near
Suffolk, Va.

- Genus 491. FORBESELLA** Jordan and Evermann.
Forbesella Jordan and Evermann, Proc., Calif. Acad. Sci., 4th ser., XVI,
1927 (Apr. 27), 503 (*Chologaster papilliferus* Forbes).
This genus stands between the ancient type of *Chologaster* well colored
and with developed eyes, and the colorless blind fishes of the caves
of the middle west. From *Chologaster*, *Forbesella* differs in the pres-
ence of tactile papillæ as in *Amblyopsis*.

1513. **Forbesella papillifera** (Forbes). *Cavefish.*
Cave springs, Union and Pope Counties, Ill.
Chologaster papilliferus Forbes, Amer. Nat., XVI, January, 1882, 1,
cave spring in Union County, southern Illinois.
1514. **Forbesella agassizii** (Putnam).
Underground streams of Tennessee and Kentucky.
Chologaster agassizii Putnam, Amer. Nat., VI, 1872, 30, Pl. 2, figs.
4, 4a, 4b, from a well at Lebanon, Tenn.

- Genus 492. TYPHLOCHTHYS** Girard.
Typhlichthys Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 63 (*T.
subterraneus* Girard).

1515. *Typhlichthys subterraneus* Girard. *Small blindfish.*
Subterranean streams in limestone regions of Indiana, Kentucky,
Tennessee, Missouri, and Alabama.
Typhlichthys subterraneus Girard, Proc. Acad. Nat. Sci. Phila., XI,
1859, 63, well at Bowling Green, Ky.
1516. *Typhlichthys osborni* Eigenmann.
Horse Cave, near Mammoth Cave, Ky.
Typhlichthys osborni Eigenmann, Biol. Bull., VIII, 1905, 65, figs. 3 and
4, Horse Cave, Ky.
1517. *Typhlichthys wyandotte* Eigenmann.
Well near Corydon, Ind.
Typhlichthys wyandotte Eigenmann, Biol. Bull., VIII, 1905, 63, Corydon,
Ind.
- Genus 493. **TROGLICHTHYS** Eigenmann.
Troglichthys Eigenmann, A Case of Convergence, Science, new series,
IX, 1899, 282 (*Typhlichthys rosæ* Eigenmann).
1518. *Troglichthys rosæ* (Eigenmann). *Missouri cavefish.*
Typhlichthys rosæ Eigenmann, Science, new series, IX, 1899, No. 217,
282, caves in Missouri.
- Genus 494. **AMBLYOPSIS** DeKay.
Amblyopsis DeKay, New York Fauna: Fishes, 1842, 187 (*A. spelæus*
DeKay).
1519. *Amblyopsis spelæus* DeKay. *Mammoth Cave blindfish.*
Subterranean streams of the limestone regions of Kentucky and Indiana;
River Styx, Mammoth Cave; the largest species of the group.
Amblyopsis spelæus DeKay, New York Fauna: Fishes, 1842, 187,
Mammoth Cave, Ky.

Order EE. SYNENTOGNATHI

Family 119. BELONIDÆ. Houndfishes; Needlefishes

- Genus 495. **STRONGYLURA** Van Hasselt.
Strongylura Van Hasselt, Bull. Sci. Nat. Ferussac., II, 1824, 374 (*S. caudimaculata* Van Hasselt = *Belone strongylura* Van Hasselt, 1823).
1520. *Strongylura notata* (Poey). *Needlefish; Longjaws; Gar.*
West Indies, north to Pensacola; common about Key West.
Belone notata Poey, Memorias, II, 1861, 293, Cuba.
1521. *Strongylura scapularis* (Jordan and Gilbert).
Panama.
Tylosurus scapularis Jordan and Gilbert, Bull. U. S. Fish Com., I,
1881, 307, Panama.
1522. *Strongylura timucu* (Walbaum). *Timucu; Peixe agulha.*
Florida Keys to Brazil; not rare.
Timucu or *Peixe agulha* Marcgrave, Hist. Pisc. Brasil., 1648, 168,
Brazil.
Esox timucu Walbaum, Artedi Pisc., III, 1792, 88, Brazil; after *Timucu*
of Marcgrave.
Belone subtruncata Poey, Memorias, II, 1860, 295, Habana.
Belone depressa Poey, loc. cit., 296, Cuba.
Tylosurus sagitta Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884,
25, Key West.
Belone ardeola Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII,
1846, 315 (425), Porto Rico.
Belone cigonella Cuvier and Valenciennes, loc. cit., 323 (436), Porto
Rico.
Belone almeida Quoy and Gaimard, Voy. Uranie, Zool., 1824, 226,
Bay of Rio Janeiro.
Belone timucu Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII,
1846, 316 (426), Rio de Janeiro.

- 1523. *Strongylura argalus* (LeSueur).**
West Indies.
Belone argalus LeSueur, in Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 326 (439), Guadeloupe, West Indies.
- 1524. *Strongylura euryops* (Bean and Dresel). *Longjaw*.**
Cuba and Jamaica.
Tylosurus euryops Bean and Dresel, Proc. U. S. Nat. Mus., VII, 1884, 168, Jamaica.
- 1525. *Strongylura diplotænia* (Cope).**
St. Martins Island, West Indies.
Belone diplotænia Cope, Trans. Amer. Phil. Soc., XIV, 1871, 481, St. Martins.
- 1526. *Strongylura microps* (Günther).**
Coast of Guiana.
Belone microps Günther, Cat., VI, 1866, 237, Surinam, British Guiana.
- 1527. *Strongylura angusticeps* (Günther).**
Coast of Ecuador.
Belone angusticeps Günther, Cat., VI, 1866, 238, Ecuador.
- 1528. *Strongylura stolzmanni* (Steindachner). *Sierrita*.**
Pacific coast of America from Guaymas and Mazatlan to Peru.
Belone stolzmanni Steindachner, Ichth. Beitr., VII, 1878, 21, Tumbez, Peru.
Tylosurus sierrita Jordan and Gilbert, Proc. U. S. Nat. Mus.; IV, 1881, 458, Mazatlan.
- 1529. *Strongylura exilis* (Girard). *Needlefish; California garfish*.**
Coast of southern California, from Point Conception southward to Cedros Island, Mexico.
Belone exilis Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 149, San Diego.
- 1530. *Strongylura pterura* (Osburn and Nichols).**
Gulf of California.
Tylosurus pterurus Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 152, fig. 6, Carmen Island, Gulf of California.
- 1531. *Strongylura marina* (Walbaum).** *Garfish; Billfish; Needlefish; Agujón; Salt-water gar; Silver gar; Swordfish; Green gar; Skippick; Sea pike; Harvest pike; Silvery gar.*
Coast of Maine to Texas; Atlantic and Gulf coasts, entering rivers.
Esox marinus Walbaum, Arredi Pisc., III, 1792, 88; after Schöpf.
Esox longirostris Mitchell, Amer. Month. Mag., II, 1818, 322, Hudson River.
Belone truncata LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 126, New York Bay.
Belone scutator Girard, U. S. Mex. Bound. Surv., Ichth., 1859, 30, pl. 13, Brazos Santiago and St. Joseph Island, Tex.
- 1532. *Strongylura guianensis* (Müller and Troschel).**
Tylosurus guianensis Müller and Troschel in Schomburgk's Reise, III, 1848, 626, coast of Guiana.
Genus 496. **TYLOSURUS Cocco.**
Tylosurus Cocco, Giorn. Sci. Litt. Art. Sicilia, XVII, 1829, 18 (*T. cantrainei* Cocco = *Sphyraena acus* Lacépède).
- 1533. *Tylosurus fodiator* Jordan and Gilbert. *Agujón*.**
Pacific coast of Mexico.
Tylosurus fodiator Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 459, Mazatlan.
- 1534. *Tylosurus raphidoma* (Ranzani).** *Houndfish; Aguja de Casta; Guardfish.*
West Indies, Florida Keys to Brazil; north to Ocean City, N. J., and North Carolina.
Belone raphidoma Ranzani, Nov. Com. Acad. Sci. Bonon., V, 1842, 359, pl. 37, fig. 1, Brazil.
Belone gerania Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 325 (437), Martinique.
Belone crassa Poey, Memorias, II, 1861, 291, Cuba.
Belone melanochira Poey, loc. cit., 294, Cuba.
Tylosurus gladius Bean, Proc. U. S. Nat. Mus., V, 1882, 239, 430, Pensacola, Fla.

1535. *Tylosurus galeatus* (Cuvier and Valenciennes).
 Cayenne, French Guiana.
Belone galeata Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 319 (429), Cayenne, French Guiana.
1536. *Tylosurus pacificus* (Steindachner).
 Acapulco to Panama.
Belone pacifica Steindachner, Ichth. Beitr., III, 1875, 65, Panama.
1537. *Tylosurus acus* (Lacépède). *Houndfish; Agujón.*
 West Indies and occasionally northward.
Sphyraena acus Lacépède, Hist. Nat. Poiss., V, 1803, 6, pl., 1 fig. 3, Martinique.
?Esox imperialis Rafinesque, Caratteri, 1810, 59, Palermo.
?Tylosurus cantrinii Cocco, Giorn. Sci. Litt. Art. Sicilia, XVII, 1829, 18, pl. 1, fig. 4, Messina.
Belone latimanus Poey, Memorias, II, 1861, 290, Habana.
Belone jonesi Goode, Amer. Jour. Sci. Arts., XIII, 1877, 295, Bermuda.
1538. *Tylosurus caribbaeus* (LeSueur).
 West Indies.
Belone caribbea LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 127, Caribbean Sea.
Belone altipinnis Poey, Memorias, II, 1861, 293, Cuba.
- Genus 497. ABLENNES Jordan and Fordice.
Athlenes Jordan and Fordice, Proc. U. S. Nat. Mus., IX, 1886, 342
 (*Belone hians* Cuvier and Valenciennes); error of transcription for Ablennies.
1539. Ablennies hians (Cuvier and Valenciennes).
 West Indies, ranging from Florida to Brazil; generally common; also recorded from Acapulco and Hawaii.
Belone hians Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 321 (432), pl. 548, Bahia; common at Habana.
Belone maculata Poey, Memorias, II, 1861, 290, Cuba.

Family 120. SCOMBERESOCIDÆ

- Genus 498. SCOMBERESOX Lacépède. *Sauries.*
Scomberesox Lacépède, Hist. Nat. Poiss., V, 1803, 344 (*S. camperii* Lacépède).
Sayris Rafinesque, Caratteri, 1810, 60 (*S. recurvirostra* Rafinesque = *Esox saurus* Walbaum).
Grammiconus Costa, Ann. Mus. Zool. Univ. Napoli, I, 1862, 55 (*G. bicolor* Costa; said to lack the air bladder).
1540. *Scomberesox saurus* (Walbaum). *Saurie; Skipper; Billfish; Skipjack.*
 Temperate parts of Atlantic Ocean; common in schools especially north of Cape Cod and France.
Esox saurus Walbaum, Artedi Pisc., III, 1792, 93, Cornwall; after the saury pike of Pennant.
Scomberesox camperii Lacépède, Hist. Nat. Poiss., V, 1803, 345, locality unknown.
Sayris recurvirostra Rafinesque, Caratteri, 1810, 61; substitute for *camperi*.
Sayris hians Rafinesque, loc. cit., 61, Palermo.
Sayris bimaculatus Rafinesque, loc. cit., 62, Palermo.
Scomberesox scutellatum LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821 (1823), 132, Newfoundland.
Scomberesox storeri DeKay, N. Y. Fauna: Fishes, 1842, 229, pl. 35, fig. 3, Banks of Newfoundland; coast of Massachusetts; New York.
Scomberesox equirostrum LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821 (1823), 132, locality unknown.
Scomberesox rondeletii Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 350 (472), Mediterranean.
Grammiconus bicolor Costa, Ann. Mus. Zool. Univ. Napoli, I, 1862, 55, Naples.
?Scomberesox forsteri Cuvier and Valenciennes, Hist. Nat. Poiss., XVIII, 1846, 357 (481), New Zealand.
Sayris serrata Rafinesque, Caratteri, 1810, 61, after Rondelet.

Genus 499. *COLOLABIS* Gill.

Cololabis Gill, Proc. U. S. Nat. Mus., XVIII, 1895, 176 (*Scomberesox brevirostris* Peters).

1541. *Cololabis brevirostris* (Peters).

Coast of California, south to San Diego in the open sea; often in schools.
Scomberesox brevirostris Peters, Monatsberichte Akad. Wiss. Berlin, July, 1866, 521, Tomales Bay, Calif.; perhaps not separable from the Japanese *Cololabis saira* Brevoort.

Family 121. **HEMIRAMPHIDÆ. Halfbeaks**Genus 500. *EULEPTORHAMPHUS* Gill.

Euleptorhamphus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 156 (*E. brevoortii* Gill = probably *E. velox* Poey).

1542. *Euleptorhamphus longirostris* (Cuvier).

East Indies; Galapagos Islands.

Hemiramphus longirostris Cuvier, Règne Anim., ed. 2, II, 1829, 286, warm seas of both hemispheres.

1543. *Euleptorhamphus velox* Poey. *Slender halfbeak.*

West Indies, occasionally northward in the Gulf Stream to Massachusetts.

? *Euleptorhamphus brevoortii* Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 156, no locality.

Euleptorhamphus velox Poey, Synopsis, 1867, 383, Cuba; perhaps identical with the preceding.

Genus 501. *HYPORHAMPHUS* Gill. *Halfbeaks.*

Hyporhamphus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 131 (*H. tricuspidatus* Gill = *Hemiramphus unifasciatus* Ranzani).

1544. *Hyporhamphus unifasciatus* (Ranzani). *Common halfbeak; Escribano; Pajarito.*

Tropical seas on both coasts of America from Brazil north to Rhode Island, New Jersey, Guaymas, and Mazatlan; generally common from Key West to Rio de Janeiro; also from Guaymas to the Galapagos; variable; but the distinction into two or more species not trustworthy.

Hemiramphus unifasciatus Ranzani, Nov. Com. Acad. Bonon., V, 1842, 326, Brazil.

? *Hemiramphus picarti* Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 18 (25), Bono, Algiers.

Hemiramphus richardi Cuvier and Valenciennes, loc. cit., 19 (26), St. Croix, Antilles.

Hyporhamphus tricuspidatus Gill. Proc. Acad. Nat. Sci. Phila., XI, 1859, 131, Barbados.

Hemiramphus fasciatus Poey, Memorias, II, 1861, 299, Cuba; not of Bleeker.

1545. *Hyporhamphus roberti* ^{as} (Cuvier and Valenciennes).

Both coasts of tropical America; West Indies and Brazil north to Rhode Island, and on the Pacific coast from Gulf of California to the Galapagos.

Hemiramphus roberti Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 18 (24), Cayenne.

1546. *Hyporhamphus gilli* Meek and Hildebrand.

Acapulco to the Galapagos.

Hyporhamphus gilli Meek and Hildebrand, Field Mus. Nat. Hist. Pub. 215, Zool. Ser., XV, Dec. 20, 1923, 240, Pl. XVII, fig. 1, Panama Bay, Balboa.

1547. *Hyporhamphus roseæ* (Jordan and Gilbert).

Coast of southern California and probably southward.

Hemiramphus roseæ Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 335, San Diego, Calif.

* This species is slender than *H. unifasciatus*, with longer beak and more numerous gill rakers, 28 to 31 on lower limb (21 in *H. unifasciatus*).

1548. Hyporhamphus snyderi Meek and Hildebrand.

Lower California to Panama.

Hyporhamphus snyderi Meek and Hildebrand, Field Mus. Nat. Hist. Pub. 215, Zool., Ser. XV, Dec. 20, 1923, 240, Pl. XVII, fig. 2, Balboa, Panama Bay.

Genus 502. HEMIRAMPHUS Cuvier.

Hemiramphus Cuvier, Règne Anim., ed. 1, II, 1817, 186 (*Esox brasiliensis* Linnæus).

1549. Hemiramphus brasiliensis (Linnæus). *Balao; Escribano; Red-tailed balao.*

West Indies, generally abundant from Key West southward to Bahia.

Esox brasiliensis Linnæus, Syst. Nat., ed. X, 1758, 314, Jamaica; after Browne; the *Timucu* of Macgrave wrongly included in the synonymy; Bloch, Ichth., 1801, 391, corrected synonymy and description.

Hemiramphus brownii Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 10 (13), Guadeloupe, Martinique.

Hemiramphus pleii Cuvier and Valenciennes, loc. cit., 15 (19), Martinique, San Domingo.

Macronathus brevirostris Gronow, Cat. Fish., 1854, 148, America; after Browne.

Hemiramphus filamentosus Poey, Memorias, II, 1861, 297, Cuba.

1550. Hemiramphus balao LeSueur. *Balao; Piper; Blue-tailed balao.*

Cuba; also known from Panama; Guadeloupe; Martinique.

Hemiramphus balao LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1823, 136, Lesser Antilles, Guadeloupe, Martinique.

Hemiramphus macrochirius Poey, Memorias, II, 1861, 299, Cuba.

1551. Hemiramphus saltator Gilbert and Starks.

Panama; Galapagos Islands.

Hemiramphus saltator Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 53, Pl. IX, fig. 16, Panama.

Genus 503. CHRIODORUS Goode and Bean. *Hardheads.*

Chriodus Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 431 (*C. atherinoides* Goode and Bean).

1552. Chriodus atherinoides Goode and Bean. *Hardhead.*

Florida Keys and vicinity.

Chriodus atherinoides Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 432, Key West, Fla.

Genus 504. HEMIEXOCETUS Fowler.

Hemiexocetus Fowler, Proc. Acad. Nat. Sci. Phila., LIII, 1901, 293 (*H. caudimaculatus* Fowler).

1553. Hemiexocetus caudimaculatus Fowler.

Mazatlan, Mexico.

Hemiexocetus caudimaculatus Fowler, Proc. Acad. Nat. Sci. Phila., LIII, 1901, 294, Mazatlan, Mexico.

Family 122. EXOCETIDÆ. Flyingfishes**Genus 505. FODIATOR Jordan and Meek.**

Fodiator Jordan and Meek, Proc. U. S. Nat. Mus., VIII, 1885, 45 (*Exocetus acutus* Cuvier and Valenciennes).

1554. Fodiator acutus (Cuvier and Valenciennes). *Sharp-nosed flyingfish.*

Both shores of tropical America; Gulf of California; San Luis Gonzales; Panama; Nice.

Exocetus acutus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 91 (125), Surinam; Nice.

Genus 506. EVOLANTIA Heller and Snodgrass.

Evolantia Heller and Snodgrass, Proc. Wash. Acad. Sci., V, 1903, 189 (*Exocetus micropterus* Cuvier and Valenciennes).

1555. Evolantia microptera (Cuvier and Valenciennes). *Malolo.*

Galapagos Islands.

Exocetus micropterus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 92 (127), Port King George, New Holland, etc.

Genus 507. PAREXOCETUS Bleeker.

Parexocetus Bleeker, Nederl. Tijdschr. Dierk., III, 1866, 126 (*Exocetus mento* Cuvier and Valenciennes).

1556. Parexocetus mesogaster (Bloch).

Tropical seas; East Indies; West Indies; Hawaiian Islands; Newport, Rhode Island; Carolina coast.

Exocetus mesogaster Bloch. Naturgesch. Ausl. Fisch., IX, 1795, 17, Martinique; on a drawing by Plumier, in which the pectorals and ventrals are much too long.

Exocetus orbignianus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 95 (131), Montevideo.

Exocetus hillianus Gosse, Naturalist's Sojourn in Jamaica, II, 1851, Pl. I, fig. 1, Jamaica.

Exocetus gryllus Klunzinger, Verhandl. Zool. Botan. Gesell. Wien, 1871, 586, Red Sea.

Genus 508. EXOCETUS Linnaeus.

Exocetus Linnaeus, Syst. Nat., ed. X, 1758, 316 (*E. volitans*).

1557. Exocetus volitans Linnaeus.

Warm seas; Atlantic coast north to Newfoundland; England; Hawaiian Islands and Japan, largely pelagic.

Exocetus volitans Linnaeus, Syst. Nat., ed. X, 1758, 316, Atlantic Ocean; not of Cuvier and Valenciennes.

Exocetus evolans Linnaeus, loc. cit., ed. XIII, 1766, 521; after Gronow.

Exocetus splendens Abel, Narr. Voy. China, 1818, 4, China.

Exocetus georgianus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 139, equatorial America.

Exocetus chilensis Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 472, Chile.

1558. Exocetus obtusirostris Günther.

Tropical Atlantic; recorded from Cape Verde Islands, New Orleans, and Miami, Fla.

Exocetus obtusirostris Günther, Cat., VI, 1866, 283, Cape Verde Islands, etc.; according to Nichols (Copeia, No. 108, 1922, 49), this is different from *E. volitans*.

Genus 509. EXONAUTES Jordan and Evermann.

Exonautes Jordan and Evermann, Rept. U. S. Fish Com., 1895, 322 (*Exocetus exsiliens* P. L. S. Müller).

1559. Exonautes exsiliens (P. L. S. Müller).

Open seas; rare on our Atlantic coast.

Exocetus exsiliens Philip Ludwig Stafius Müller, Nürnberg, ed. Linnaeus, Syst. Nat., 1776, 209, Carolinas.

Exocetus fasciatus LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 8, pl. 4, fig. 2, Atlantic.

Exocetus lamellifer Kner and Steindachner, Neue Fische Godeffroy Mus., 1866, 29, Atlantic 12° S., 39° W.

1560. Exonautes rondeletii (Cuvier and Valenciennes).

Tropical seas; Florida; France; Acapulco; West Indies; southern Europe; perhaps the adult of *E. exsiliens*.

Exocetus rondeletii Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 84 (115), pl. 562, Naples; Sicily; Canaries.

Exocetus brachycephalus Günther, Cat., VI, 1866, 297, China.

Exocetus volador Jordan, Proc. U. S. Nat. Mus., VII, 1884, 34 Pensacola, Fla.

1561. Exonautes vinciguerræ (Jordan and Meek).

Open Atlantic; southern Europe; Newfoundland; St. Martins; Gulf of Mexico.

Exocetus vinciguerræ Jordan and Meek, Proc. U. S. Nat. Mus., VIII, 1885, 56, open sea off Newfoundland, 46° N., 61° W.

- 1562. *Exonautes rubescens* (Rafinesque).**
 Atlantic Ocean, north to Newfoundland; also in the Pacific if *E. speculiger* is the same.
Exocetus rubescens Rafinesque, Amer. Month. Mag., 1818, 205, Banks of Newfoundland.
? *Exocetus speculiger* Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 69 (63), Friendly Islands, Pacific coast of South America, etc.
Exocetus roberti Müller and Troschel, Schomburgk's Hist. Barbados, 1848, 675, Barbados.
? *Exocetus quadriremus* Gronow, Cat. Fish., 1854, 145, Spain and India.
Exocetus volitans of various authors; not of Linnaeus.
- 1563. *Exonautes affinis* (Günther).**
 West Indies to Africa; locally common.
Exocetus affinis Günther, Cat., VI, 1866, 288, West Africa; Cuba.
- 1564. *Exonautes rufipinnis* (Cuvier and Valenciennes).**
 Tropical America; Payta; Panama; Tabasco; Barbados.
Exocetus rufipinnis Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 72 (99), Payta, Peru.
Exocetus dowii Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 167, near Panama.
Exocetus scylla Cope, Trans. Amer. Phil. Soc., XIV, 1871, 481, Tabasco, Mexico.
- Genus 510. *CYPSELURUS* Swainson.**
Cypsilurus Swainson, Class. Fishes, etc., II, 1839, 296 (*Exocetus nuttalli* LeSueur); misprint for *Cypselurus*, according to decision Int. Com. Zool. Nom.
- 1565. *Cypselurus heterurus* (Rafinesque).** *Flyingfish; Single-bearded flying-fish.*
 Atlantic Ocean, generally common southward to Newfoundland and to England.
Exocetus heterurus Rafinesque, Caratteri, 1810, 58, Palermo.
Exocetus comatus Mitchell, Trans. Lit. Phil. Soc. N. Y., 1815, 448, pl. 5, fig. 1, New York.
Exocetus noveboracensis Mitchell, Amer. Month. Mag., II, 1818, 333, New York.
Exocetus appendiculatus Wood, Jour. Acad. Nat. Sci. Phila., IV, 1824, 283, pl. 17, fig. 2, locality unknown.
Exocetus melanurus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 74 (101), New York.
- 1566. *Cypselurus lutkeni* (Jordan and Evermann).**
 Coast of Cuba.
Exocetus lutkeni Jordan and Evermann, Fishes North and Mid. Amer., 1896, 736, Cape San Antonio, Cuba.
- 1567. *Cypselurus furcatus* (Mitchill).** *Double-bearded flyingfish.*
 Warm seas, north to Cape Cod and the Mediterranean, generally common.
Exocetus furcatus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 449, New York.
Exocetus nuttalli LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 10, Pl. IV, fig. 1, Gulf of Mexico.
Exocetus (Cypsilurus) procne De Filippie Verany, Mem. Acad. Sci. Torino, ser. 2, XVIII, 1857, 10, fig. 5, Nice.
Exocetus maculipinnis Vinciguerra, Ann. Mus. Genov., XVIII, 1883, pl. 1, fig. 6, 577, Tunis.
- 1568. *Cypselurus nigricans* (Bennett).**
 Tropical seas, north to Cuba, Central America, and France.
Exocetus nigricans Bennett, Whaling Voyage, II, 1840, 287, taken in both the Atlantic and Pacific Oceans, in lat. 5° N.
Exocetus bicolor Cuvier and Valenciennes, Hist. Nat. Poiss., XIX, 1846, 81 (111), Atlantic.
Exocetus spilopus Cuvier and Valenciennes, loc. cit., 86 (118), La Rochelle; St. Helena; West Indies; India; Arabia; De Witt Land.

1569. *Cypselurus xenopterus* (Gilbert).
 Clarion Island, one of the Revillagigedo Islands.
Exocetus xenopterus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 58,
 Clarion Island.
1570. *Cypselurus lineatus* (Cuvier and Valenciennes).
 Tropical Atlantic; Gorea; the Canaries; Madeira and Bermuda.
Exocetus lineatus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX,
 1846, 67 (92), Gorea; Canaries.
1571. *Cypselurus cyanopterus* (Cuvier and Valenciennes).
 Coasts of Brazil and Caribbean Sea; not common.
Exocetus cyanopterus Cuvier and Valenciennes, Hist. Nat. Poiss., XIX,
 1846, 71 (98), American seas.
Exocetus albidaetus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 167,
 Caribbean Sea.
1572. *Cypselurus bahiensis* (Ranzani).
 Tropical Atlantic; north to Cuba.
Exocetus bahiensis Ranzani, Nov. Com. Acad. Sci. Bonon., V, 1842,
 362, pl. 38, Bahia.
Exocetus vermiculatus Poey, Memorias, II, 1861, 300, Cuba.
? *Exocetus parrae* Poey, Synopsis, 1868, 385, Cuba.
1573. *Cypselurus spilonotopterus* (Bleeker).
 Tropical Pacific, north to Hawaii and Galapagos Islands.
Exocetus spilonotopterus Bleeker, Nederl. Tijdschr. Dierk., III, 1863,
 113, Sumatra.
1574. *Cypselurus californicus* (Cooper). *Great flyingfish; Volador.*
 Southern California; Point Conception to Cape San Lucas; Santa
 Barbara Islands; locally very abundant.
Exocetus californicus Cooper, Proc. Calif. Acad. Nat. Sci., III, 1863, 93,
 fig. 20, Santa Catalina Island.
1575. *Cypselurus callopterus* (Günther).
 Panama.
Exocetus callopterus Günther, Cat., VI, 1866, 292, Panama.
1576. *Cypselurus fucorum* (Zugmayer).
 Mid-Atlantic.
Exocetus fucorum Zugmayer, Bull. 193, Inst. Oceanogr., Jan. 20, 1911,
 in 31° 38' N., 42° 38' W., at surface.
1577. *Cypselurus holubi* (Steindachner).
Exocetus holubi Steindachner, in Collett, Voy. Hirondelle, 1896, 123,
 Pl. IV, fig. 18, Newfoundland.
1578. *Cypselurus gibbifrons* (Cuvier and Valenciennes).
 Atlantic Ocean; Newport, R. I.
Exocetus gibbifrons Cuvier and Valenciennes, Hist. Nat. Poiss., XIX,
 1846, 86 (118), Atlantic.

Order FF. ANACANTHINI

Family 123. MACROURIDÆ (Coryphaenoididæ). Grenadiers; Rat-tails

Genus 511. GADOMUS Regan.

Gadomus Regan, Ann. Mag. Nat. Hist., XII, 1903, 459, 466 (*Bathygadus longifilis* Goode and Bean).

1579. *Gadomus arcuatus* (Goode and Bean).

Gulf of Mexico, in deep water.

Bathygadus arcuatus Goode and Bean, Bull. Mus. Comp. Zool., XII,
 1886, No. 5, 158, off Martinique in 334 fathoms.

1580. *Gadomus longifilis* (Goode and Bean).

Deep waters of the Gulf of Mexico.

Bathygadus longifilis Goode and Bean, Proc. U. S. Nat. Mus., VIII,
 1885, 599, Gulf of Mexico at Albatross station 2392, lat. 28° 47' 30'
 N., long. 87° 27' W., in 724 fathoms.

Genus 512. BATHYGADUS Günther.

Bathygadus Günther, Ann. Mag. Nat. Hist., II, 1878, 23 (*B. cottooides* Günther).

Melanobranchus Regan, loc. cit., (7), XI, 1903, 459 (*Bathygadus melanobranchus* Vaillant).

1581. *Bathygadus favosus* Goode and Bean.

Off Martinique, in deep water.

Bathygadus favosus Goode and Bean, Bull. Mus. Comp. Zool., XII, 1886, No. 5, 160, off Martinique at *Blake* station 80 in 472 fathoms.

1582. *Bathygadus macrops* Goode and Bean.

Deep waters of the Gulf of Mexico.

Bathygadus macrops Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 598, at *Albatross* station 2396 in lat. $28^{\circ} 34' N.$, long. $86^{\circ} 48' W.$, in 335 fathoms.

Genus 513. STEINDACHNERIA Goode and Bean.

Steindachneria Goode and Bean, in Agassiz, Three Cruises of the *Blake*, II, 1888, 26 (diagnosis but no type named); not *Steindachneria* Eigenmann 1888 = *Steindachneridion* Eigenmann and Eigenmann, 1919.

Steindachnerella Eigenmann and Eigenmann, Amer. Nat., XXXI, 1897, 159 (*Steindachneria argentea* Goode and Bean).

1583. *Steindachneria argentea* Goode and Bean.

Gulf of Mexico.

Steindachneria argentea Goode and Bean, Oceanic Ichth., 1896, 419, fig. 351, off delta of Mississippi River in 68 fathoms.

Genus 514. TRACHYRINCUS Giorna.

Trachyrincus Giorna, Mem. Acad. Imp. Turin, XVI, 1805, 4, 179, (*Lepidolepsus trachyrincus* Risso, not mentioned).

1584. *Trachyrincus helolepis* (Gilbert).

Off the Pacific coast of Central America.

Trachyrhynchus helolepis Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 562, Pacific coast of Central America, at *Albatross* station 2818, in deep water.

Genus 515. DOLLOA Jordan.

Moseleya Goode and Bean, Oceanic Ichth., 1895, 417 (*Macrurus longifilis* Günther); name preoccupied.

Dolloa Jordan, Amer. Nat., XXXIV, 1900, 877 (*Macrurus longifilis* Günther).

1585. *Dolloa cyclolepis* (Gilbert).

Coast of British Columbia.

Nematonurus cyclolepis Gilbert, Rept. U. S. Fish Com., XIX, 1893 458, off Queen Charlotte Island in 1,588 fathoms.

Genus 516. NEMATONURUS Günther.

Nematonurus Günther, Challenger Rept., Deep Sea Fishes, XXII, 1887, 124, 150 (*N. armatus* Hector).

1586. *Nematonurus goodii* (Günther).

Gulf Stream from Cape Cod to Habana; generally abundant.

Macrurus asper Goode and Bean, Bull. Mus. Comp. Zool., X, 1883, No. 5, 196, Gulf Stream, south of New England in 1,242 fathoms.

Macrurus goodii Günther, Challenger Report, XXII, 1887, 36; substitute for *Macrurus asper*, preoccupied.

1587. *Nematonurus suborbitalis* (Gill and Townsend).

Bering Sea.

Macrurus suborbitalis Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897, 234, Bering Sea, southwest of Pribilof Islands, in 1,771 fathoms.

1588. *Nematonurus acrolepis* (Bean).

Straits of Juan de Fuca; off Bogoslof Island, north of Aleutian Islands; once taken at Monterey (W. F. Thompson).

Macrurus acrolepis Bean, Proc. U. S. Nat. Mus., VI, 1883, 362, Port Townsend, Wash.

1589. *Nematonurus abyssorum* Gilbert.

Off southern California in deep water.

Nematonurus abyssorum Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 374, pl. 21, off Santa Catalina in 1,350 to 2,182 fathoms.

Genus 517. **ALBATROSSIA** Jordan and Evermann.

Albatrossia Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2573 (*Macrourus pectoralis* Gilbert).

1590. *Albatrossia pectoralis* (Gilbert).

Off coast of Oregon.

Macrourus (Malacocephalus) pectoralis Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 563, off coast of Oregon, at *Albatross* stations 3071, 3074, and 3075, in 685 to 877 fathoms.

Macrurus (Nematonurus) magnus Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897, 234, Bering Sea, southwest of Pribilof Islands.

Genus 518. **BOGOSLOVIUS** Jordan and Evermann.

Bogoslovius Jordan and Evermann, Fish. North and Mid. Amer., 1898, 2574 (*B. clarki* Jordan and Gilbert).

1591. *Bogoslovius clarki* Jordan and Gilbert.

Bering Sea.

Bogoslovius clarki Jordan and Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2575, Bering Sea off Bogoslof Island in 664 fathoms.

1592. *Bogoslovius firmisquamis* (Gill and Townsend).

Bering Sea.

Macrurus firmisquamis Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897, 234, Bering Sea southwest of Pribilof Islands.

Genus 519. **CHALINURA** Goode and Bean.

Chalinura Goode and Bean, Bull. Mus. Comp. Zool., X, 1882, No. 5, 198 (*C. simula* Goode and Bean).

Chalinurus Günther, altered spelling.

1593. *Chalinura occidentalis* (Goode and Bean).

Atlantic Ocean off Cape Hatteras and Caribbean Sea.

Malacocephalus occidentalis Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 597, off Cape Hatteras in 132 fathoms.

1594. *Chalinura serrula* Bean.

East of Prince of Wales Island in deep water.

Chalinura serrula Bean, Proc. U. S. Nat. Mus., XIII, 1890, 37, east of Prince of Wales Island, at *Albatross* station 2859 in 1,569 fathoms.

1595. *Chalinura filifera* Gilbert.

Off Queen Charlotte Island, in deep water.

Chalinura filifera Gilbert, Rept. U. S. Fish Com., XIX, 1893 (1896), 458, off Queen Charlotte Island, at *Albatross* station 3342.

1596. *Chalinura simula* Goode and Bean.

Gulf Stream.

Chalinura simula Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1883, 199, Gulf Stream, about latitude 40°.

1597. *Chalinura brevibarbis* Goode and Bean.

Atlantic Basin.

Chalinura brevibarbis Goode and Bean, Oceanic Ichth., 1895, 413, Atlantic Ocean in 1,255 fathoms.

1598. *Chalinura spinulosa* Gilbert and Burke.

East coast of Kamchatka.

Chalinura spinulosa Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 92, fig. 36, Avatcha Bay.

Genus 520. **CORYPHÆNOIDES** ²⁶ Gunner.

Coryphænoides Gunner, Trondhj. Selsk. Skrift., III, 1765, 50 (*C. rupestris* Gunner).

²⁶ The name *Coryphænoides* is older than *Macrourus*, the term usually applied to the species of this genus. The distinction between *Coryphænoides* and *Macrourus* is rather slight, the mouth in *Macrourus* being smaller and more inferior. Gilbert and Hubbs unite the two groups as *Coryphænoides*. Many species have been described from the deep waters of America. These we leave in *Coryphænoides* or in *Macrourus* as their describers have placed them. Some of the genera accepted here are of questionable value.

1599. **Coryphænoides rupestris** Gunner.
Banks of Newfoundland to Norway.
Coryphænoides rupestris Gunner, Trondhj. Selsk. Skrift., III, 1765, 50.
Lepidolepus norvegicus Nilsson, Prodr. Ichth. Scand., 1832, 51, Norway.
1600. **Coryphænoides carapinus** Goode and Bean.
Gulf Stream.
Coryphænoides carapinus Goode and Bean, Bull. Mus. Comp. Zool., X, 1883, No. 5, 197, Gulf Stream, about lat. 40° , in deep water.
- Genus 521. MACROURUS**³⁷ Bloch.
Macrourus Bloch, Naturgesch. Ausl. Fische, II, 1786, 150 (*Coryphæna rupestris* Fabricius = *Macrourus berglax* Lacépède); not *Coryphænoides rupestris* Gunner.
Macrurus Günther, and of most authors; corrected spelling.
1601. **Macrourus berglax** Lacépède.
Massachusetts to Greenland and Norway.
Corypha rupestris Fabricius, Fauna Grœnlandica, 1780, 154, Greenland.
Macrourus berglax Lacépède, Hist. Nat. Poiss., III, 1802, 170, Greenland.
1602. **Macrourus bairdi** Goode and Bean. *Common rat-tail.*
West Indies to Massachusetts Bay; excessively common north in deep water.
Macrourus bairdi Goode and Bean, Amer. Jour. Sci. Arts., XIII, 1877, 471, Massachusetts Bay.
1603. **Macrourus stelgidolepis** Gilbert.
Pacific Ocean off Point Conception, Calif.
Macrurus (Macrourus) stelgidolepis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 116, Albatross station 2960, lat. $34^{\circ} 10' 45''$ N., long. $120^{\circ} 16' 45''$ W., in 267 fathoms.
1604. **Macrourus cinereus** Gilbert. *Popeye.*
North Pacific and Bering Sea; very abundant.
Macrurus cinereus Gilbert, Rept. U. S. Fish Com., XIX, 1893 (1896), 457, at Albatross stations 3307, 3329, and 3340, lat. 54° to 55° N. and long. 155° to 171° W., in 399 to 1,033 fathoms.
1605. **Macrourus bulbiceps** Garman.
Off Cocos Islands in deep water.
Macrurus bulbiceps Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 194, Pl. XLIV, off Cocos Islands.
1606. **Macrourus bucephalus** Garman.
Off Galapagos Islands to Gulf of California, in deep water.
Macrurus bucephalus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 195, Pl. XLIV, Gulf of California.
1607. **Macrourus barbiger** Garman.
Gulf of California in deep water.
Macrurus barbiger Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 197, Pl. XLV, Gulf of California.
1608. **Macrourus capito** Garman.
Off Panama in deep water.
Macrurus capito Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 200, off Bay of Panama.
1609. **Macrourus leucophæus** Garman.
Off Panama in deep water.
Macrurus leucophæus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 201, off Bay of Panama.
1610. **Macrourus boöps** Garman.
Off Panama in deep water.
Macrurus boöps Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 202, off Bay of Panama.

³⁷ The many deep-sea forms described by Garman are nearly all referred to *Macrourus* without reference to the closely related minor genera proposed for forms allied to *Macrourus* and *Coryphænoides*, the two principal genera of the family, and themselves barely separable.

1611. *Macrourus fragilis* Garman.
 Off Panama in deep water.
Macrourus fragilis Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 203,
 Pl. XLV, off coast of Panama.
1612. *Macrourus carminifer* Garman.
 Off Panama in deep water.
Macrourus carminifer Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
 204, Pl. XLVI, off Bay of Panama.
1613. *Macrourus gracilicauda* Garman.
 Off Panama in deep water.
Macrourus gracilicauda Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
 206, Pl. H, off Bay of Panama.
1614. *Macrourus orbitalis* Garman.
 Off Panama in deep water.
Macrourus orbitalis Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 207,
 Pl. XLVII, off Bay of Panama.
1615. *Macrourus cuspidatus* Garman.
 Gulf of California in deep water.
Macrourus cuspidatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
 209, Gulf of California.
1616. *Macrourus convergens* Garman.
 Off Panama in deep water.
Macrourus convergens Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
 210, Pl. XLVIII, off Bay of Panama.
1617. *Macrourus latirostratus* Garman.
 Off Panama in deep water.
Macrourus latirostratus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
 211, Pl. XLVIII, off Bay of Panama.
1618. *Macrourus latinusutus* Garman.
 Gulf of California in deep water.
Macrourus latinusutus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
 214, Gulf of California.
1619. *Macrourus anguliceps* Garman.
 From Galapagos Islands to Gulf of California in deep water.
Macrourus anguliceps Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
 212, Pl. G; off Bay of Panama.
1620. *Macrourus trichurus* Garman.
 Off Panama in deep water.
Macrourus trichurus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
 215, off Bay of Panama.
1621. *Macrourus tenuicauda* Garman.
 Off Panama in deep water.
Macrourus tenuicauda Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
 216, Pl. XLIX, off Bay of Panama.
1622. *Macrourus canus* Garman.
 Off Panama in deep water.
Macrourus canus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 217,
 Pl. XLIX, off Bay of Panama.
1623. *Macrourus albatrossus* Townsend and Nichols.
 Off Lower California in deep water.
Macrourus albatrossus Townsend and Nichols, Bull. Amer. Mus. Nat.
 Hist., LII, 1925, 17, lat. 31° W., southwest of San Diego in 1,076
 fathoms.
1624. *Macrourus ingolfi* Lütken.
 Iceland and Greenland in deep water.
Macrourus ingolfi Lütken, Ichth. Ingolf Exped., 1898, 27, south of Iceland.

1625. *Macrourus lepturus* Gill and Townsend.
 Bering Sea, St. Paul Island, Yunaska Island.
Macrourus lepturus Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897, 233, Bering Sea, southwest of the Pribilof Islands, in 1,401 fathoms.
Macrurus dorsalis Gill and Townsend, loc. cit., 233, Bering Sea, southwest of Pribilof Islands, in 1,401 fathoms.
- Genus 522. *CÆLORHYNCHUS* Giorna.
Cælorhynchus Giorna, Mem. Acad. Nat. Sci. Turin, XVI, 1805, 179, pl. 1, figs. 3 and 4 (*C. laville* Giorna).
1626. *Cælorhynchus occa* (Goode and Bean).
 Gulf of Mexico in deep water.
Macrurus occa Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 595, Gulf of Mexico in 335 fathoms.
1627. *Cælorhynchus carminatus* (Goode). *Grenadier*.
 Caribbean Sea, north in the Gulf Stream to Long Island.
Macrurus carminatus Goode, Proc. U. S. Nat. Mus., III, 1880, 346, Gulf Stream, 40° 02' 54" N., 70° 23' 40" W., in 115 fathoms.
1628. *Cælorhynchus caribbaeus* (Goode and Bean).
 West Indies to northern part of Gulf of Mexico.
Macrurus caribbaeus Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 594, northern part of Gulf of Mexico in 210 fathoms.
1629. *Cælorhynchus scaphopsis* (Gilbert).
 Gulf of California in deep water.
Macrurus (Cælorhynchus) scaphopsis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 115, Gulf of California at *Albatross* station 3015, lat. 29° 19' N., long. 112° 50' W., in 145 fathoms.
- Genus 523. *TRACHONURUS* Günther.
Trachonurus Günther, Challenger Rept., Deep-sea Fishes, XXII, 1887, 124 (*Macrurus villosus* Günther).
1630. *Trachonurus sulcatus* (Goode and Bean).
 West Indies and Gulf of Mexico in deep water.
Coryphaenoides sulcatus Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 596, Gulf of Mexico in 420 fathoms.
- Genus 524. *LIONURUS* Günther.
Lionurus Günther, Challenger Rept., Deep-sea Fishes, XXII, 1887, 124 (*Coryphaenoides filicauda* Günther).
1631. *Lionurus liolepis* (Gilbert).
 Off southern California in deep water.
Macrurus (Lionurus) liolepis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 117, coast of California at *Albatross* station 2980, lat. 33° 49' 45" N., long. 119° 24' 30" W. in 600 fathoms.
- Genus 525. *HYMENOCEPHALUS* Giglioli and Issel.
Hymenocephalus Giglioli and Issel, Pelagos, Genoa, 1884, 228 (*H. italicus* Giglioli and Issel).
1632. *Hymenocephalus cavernosus* (Goode and Bean).
 Gulf of Mexico in deep water.
Bathygadus cavernosus Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 598, *Albatross* station 2398, lat. 28° 45' N., long. 86° 26' W. in 227 fathoms.
- Genus 526. *ATELEOBRACHIUM* Gilbert and Burke.
Ateleobrachium Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 94 (*A. pterotum* Gilbert and Burke).
1633. *Ateleobrachium pterotum* Gilbert and Burke.
 East coast of Kamchatka.
Ateleobrachium pterotum Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 94, Avatcha Bay.

Family 124. BREGMACEROTIDÆ

Genus 527. BREGMACEROS (Cantor) Thompson.

Bregmaceros Thompson, in Charlesworth's Mag. Nat. Hist., IV, 1840, I, 184 (B. macclllandii Thompson).

Calloptilum Richardson, Voy. Sulphur, 94, 1843, 16 (C. mirum Richardson).

Asthenurus Tickell, Jour. Asiat. Soc. Bengal, 1865, 32 (A. atripinnis Tickell).

1634. *Bregmaceros bathymaster* Jordan and Bollman.

South Pacific and Indian Oceans; also off coast of Panama.

?*Bregmaceros macclllandii* Cantor, in Charlesworth's Mag. Nat. Hist., IV, 1840, 184, fig. 185, India.

?*Asthenurus atripinnis* Tickell, Jour., Asiat. Soc. Bengal, 1865, 32, with plate, Bay of Bengal.

Bregmaceros bathymaster Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 173, southwest of Panama.

1635. *Bregmaceros atlanticus* Goode and Bean.

Blake stations 99, off Granada, 113, off Nevis; and 185.

Bregmaceros atlanticus Goode and Bean, Bull., Mus. Comp. Zool., XII, 1886, No. 5, 165, off Nevis and Granada, West Indies.

1636. *Bregmaceros longipes* Garman.

Off coast of Acapulco, Mexico, in shallow water.

Bregmaceros longipes Garman, Mem., Mus. Comp. Zool., XXIV, 1899, 191, Pl. XLIII, off Pacific coast of Mexico.

Family 125. GAIDROPSARIDÆ. Rocklings

Genus 528. GAIDROPSARUS Rafinesque. Three-bearded rocklings.

Gaidropsarus Rafinesque, Indice, II, 1810, 51 (G. mustellurus = *Enchelyopus mediterraneus* Schneider).

Dropsarus Rafinesque, Anal. Nat., 1815, 88, shortened form; same type.

Les Mustèles Cuvier, Règne Anim., ed. 1, II, 1817, 215 (*tricirrhatus* = *mediterraneus*).

Mustela Oken, Isis, 1817 (for *les Mustèles*, not *Mustela* L., a genus of weasels).

Onos Risso, Hist. Europ. Méríd., III, 1826, 214 (*Gadus mustela* Bloch).

Mustela Stark, Elem. Nat. Hist., I, 1828, 425; after *les Mustèles*.

Motella Cuvier, Règne Anim., ed. 2, II, 1829, 334 (*Gadus mustela* L.); substitute for *Mustela* Cuvier.

Onus Günther, corrected spelling; not of Rafinesque.

1637. *Gaidropsarus ensis* (Reinhardt). Rockling; Three-bearded rockling.

New York to Greenland.

Motella ensis Reinhardt, Dansk. Vidensk. Selsk. Afhandl., VII, 1838, 116, 128, Greenland.

Onos rufus Gill, Proc. U. S. Nat. Mus., VI, 1883, 259, Gulf Stream.

1638. *Gaidropsarus septentrionalis* (Collett).

Coasts of Norway and Greenland.

Motella septentrionalis Collett, Ann. Mag. Nat. Hist., 4th ser., XV, 1875, 82, Lofoten, Floro, Bergen, coast of Norway and Bodö, lat. 67° 15' N., in 30 fathoms.

Genus 529. CILIATA Couch. Five-bearded rocklings.

Ciliata Couch, Mag. Nat. Hist., V, 1822, 15 (*C. glauca* Couch, the "mackerel midge," larva of *Gadus mustela* L.).

Couchia Thompson, Nat. Hist. Ireland, IV, 1856, 288 (*C. minor* Thompson = larva of *Gadus mustela* L.).

Molvella Kaup, Archiv Natur., XXIV, 1858, I, 90 (*Molvella borealis* Kaup = *Gadus mustela* L.).

1639. *Ciliata argentata* (Reinhardt). Silvery rockling; Mackerel midge; Five-bearded rockling.

Coasts of Greenland, south to Faroe and Bear Islands.

Motella argentata Reinhardt, Dansk. Vidensk. Selsk. Afhandl., VII, 1838, 116, 128, Greenland.

Motella reinhardti Kröyer, in Collett, Forhandl. Vidensk. Selsk. Chr., No. 14, 1878, 83, Greenland.

- Genus 530. ENCHELYOPUS**²⁸ Bloch and Schneider. *Four-bearded rocklings.*
Enchelyopus Bloch and Schneider, Syst. Ichth., 1801, 50 (*Gadus cimbrius* L.); not of Gronow, which is nonbinomial.
Rhinonemus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 241 (*Gadus cimbrius* L.).
- 1640. Enchelyopus cimbrius** (Linnaeus). *Four-bearded rockling.*
 North Atlantic on both coasts, south in deep water to the Gulf Stream; common in Massachusetts Bay.
Gadus cimbrius Linnaeus, Syst. Nat., ed. 12, I, 1766, 440, Atlantic Ocean; Scania.
Motella caudacuta Storer, Proc. Bost. Soc. Nat. Hist., III, 1848, 5, Long Point, Provincetown, Mass.
- Family 126. GADIDÆ. Codfishes**
- Genus 531. BOREOGADUS** Günther.
Boreogadus Günther, Cat., IV, 1862, 336 (*Gadus fabricii* Richardson = *Gadus saida* Lepechin).
- 1641. Boreogadus saida** (Lepechin).
 Arctic seas, Greenland to Alaska and northern Russia; common in the far north.
Gadus saida Lepechin, Nov. Comm. Acad. Sci. Petropol., XVIII, 1774, 512, Russia.
Merlangus polaris Sabine, Suppl. Parry's Voy., CCXI, 1824, Baffin's Bay.
Gadus fabricii Richardson, Fauna Bor.-Amer., III, 1836, 245, northern bays of Greenland.
Gadus agilis Reinhardt, Danske Vidensk. Selsk. Afhandl., VII, 1838, 126, Greenland.
- Genus 532. POLLACHIUS** Nilsson. *Pollacks.*
Pollachius Nilsson, in Bonaparte, Cat. Metod. Pesci. Europ., 1846, 45 (*Gadus pollachius* Linnaeus).
- 1642. Pollachius virens** (Linnaeus). *Pollack; Coalfish; Green cod.*
 North Atlantic; common northward on both coasts, south to Cape Cod.
Gadus virens Linnaeus, Syst. Nat., ed. X, 1758, 253, oceans of Europe.
Gadus carbonarius Linnaeus, loc. cit., 254, seas of Europe.
Gadus colinus Lacépède, Hist. Nat. Poiss., II, 1800, 416, England, etc.
Gadus purpureus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 370, New York.
Merlangus leptoccephalus DeKay, New York Fauna: Fishes, 1842, 288, pl. 45, fig. 146, Long Island.
- Genus 533. THERAGRA** Lucas.
Theragra Lucas, in Jordan and Gilbert, Rept. Fur-seal Invest., III, 1896, 486 (*Gadus chalcogrammus* Pallas).
- 1643. Theragra chalcogramma** (Pallas). *Alaska pollack.*
 North Pacific, south to Sitka.
Gadus chalcogrammus Pallas, Zoogr. Rosso.-Asiat., III, 1811, 198, Kamchatka.
Gadus periscopius Cope, Proc. Amer. Phil. Soc., XIII, 1873, 30, Unalaska.
- 1644. Theragra fucensis** (Jordan and Gilbert). *Wall-eyed pollack; Puget Sound pollack.*
 Puget Sound.
Pollachius chalcogrammus fucensis Jordan and Gilbert, Proc. U. S. Nat. Mus., XVI, 1893, 315, Puget Sound at Tacoma, Wash.
- Genus 534. ELEGINUS** Fischer.
Eleginus Fischer, Mem. Soc. Nat. Moscow, 2d ed., V, 1813, 4, 252-257 (*Gadus nauaga* Kölreuter).

²⁸ The name *Enchelyopus* ("eels without ventral fins") has been used variously since first proposed by Gronow; Gronow applied the name to *Zoarces*, Klein to *Trichinurus*; as both these writers were nonbinomial we recognize its first binomial use, that of Schneider, for *Gadus cimbrius* L.

- 1645. *Eleginus nawaga* (Kölreuter). *Wachna cod.***
 Puget Sound to Bering Sea.
Gadus nawaga Kölreuter, Nov. Com. Acad. Sci. Petropol., XIV, 1770, 484, pl. 12, coast of northern Russia.
Gadus gracilis Tilesius, Mém. Acad. Sci. Imp. St. Petersb., II, 1810, 354, Kamchatka.
Gadus wachna Pallas, Zoogr. Ross.-Asiat., III, 1811, 182, Kamchatka.
- Genus 535. *MICROGADUS* Gill. *Tomcods.***
Microgadus Gill, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 69 (*Gadus proximus* Girard).
- 1646. *Microgadus tomcod* (Walbaum). *Tomcod; Frostfish.***
 East coast of North America, from Virginia to Labrador.
Gadus tomcod Walbaum, Artedi Pisc., III, 1792, 133, Long Island; after Schöpf.
Gadus frost Walbaum, loc. cit., 134, North America.
Gadus tomcodus pruinosus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 269, New York.
Gadus tomcodus fuscus Mitchell, loc. cit., 369, New York.
Gadus tomcodus luteus Mitchell, loc. cit., 369, New York.
Gadus tomcodus mixtus Mitchell, loc. cit., 369, New York.
Gadus polymorphus Mitchell, loc. cit., 369, New York.
Morrhua americana Storer, Rept. Fish. Mass., 1839, 120, coast of Massachusetts.
- 1647. *Microgadus proximus* (Girard). *California tomcod.***
 West coast of North America, from Monterey to Alaska.
Gadus proximus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 141, San Francisco.
Morrhua californica Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 8, San Francisco.
- Genus 536. *GADUS* Linnaeus. *Codfishes.***
Gadus (Artedi) Linnaeus, Syst. Nat., ed. X, 1758, 251 (*Gadus morrhua* L.).
Morrhua Oken, Isis, 1817, 1182, after "Les Morrhues" Cuvier (*Gadus morrhua* L.).
Cephus Swainson, Nat. Hist. Class. Fish., II, 1839, 300 (*Gadus macrocephalus* Tilesius).
Cerdo Gistel, Thierreich, X, 1848 (*Gadus morrhua* L., by implication).
- 1648. *Gadus morrhua* Linnaeus. *Common codfish.***
 North Atlantic, south to Virginia.
Gadus callarias Linnaeus, Syst. Nat., ed. X, 1758, 252, Baltic Sea and ocean off Europe; young.
Gadus morrhua Linnaeus, loc. cit., seas of Europe; name preferred by first reviser.
Gadus vertagus Walbaum, Artedi Pisc., III, 1792, 143; after Jägershen.
?Gadus heteroglossus Walbaum, loc. cit., 144; after Hornbogen of Klein.
Gadus arenosus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 368, New York.
Gadus rupestris Mitchell, loc. cit., 368, New York.
?Gadus nanus Faber, Natur. Fische Islands, 1829, 112, 182, Iceland.
- 1649. *Gadus macrocephalus* Tilesius. *Alaska codfish; Pacific cod.***
 Bering Sea, south to coast of Oregon.
Gadus macrocephalus Tilesius, Mem. Acad. Sci. St. Petersb., II, 1810, 360, Bering Sea.
Gadus pygmaeus Pallas, Zoogr. Ross.-Asiat., III, 1811, 199, Kamchatka.
Gadus auratus Cope, Proc. Amer. Phil. Soc., XIII, 1873, 30, Unalaska.
- 1650. *Gadus ogac* Richardson. *Greenland codfish; Ogac.***
 Greenland; Godhaven.
Gadus ogac Richardson, Fauna Bor.-Amer., III, 1836, 246, Greenland.
- Genus 537. *MELANOGRAMMUS* Gill.**
Melanogrammus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 280 (*Gadus æglefinus* Linnaeus).
Æglefinus Malm, Göteborg och Bohusläns Fauna, 1877, 481 (*Gadus æglefinus* L.=*Æglefinus linnæi* Malm).

1651. **Melanogrammus aeglefinus** (Linnæus). *Haddock.*

North Atlantic, on both coasts.

Gadus aeglefinus Linnæus, Syst. Nat., ed. X, 1758, 251, ocean off Europe.

Morrhua puctatus Fleming, Brit. Anim., 1828, 191.

Aeglefinus linnæi Malm, Göteborg Bohusläns Fauna, 1877, 481.

Genus 538. **LEPIDION** Swainson.

Lepidion Swainson, Nat. Hist. Class'n Anim., I, 1838, 318, and II, 1839, 300 (*L. rubescens* Swainson = *Gadus lepidion* Risso); not

Lepidion Savigny, 1817,

Haloporphyrus Günther, Cat., IV, 1862, 358 (*Gadus lepidion* Risso).

1652. **Lepidion verecundum** Jordan and Cramer.

Off Pacific coast of Mexico.

Lepidion verecundum Jordan and Cramer, Proc. U. S. Nat. Mus., XIX, 1896, 456, Clarion Island, Mexico.

Genus 539. **MICROLEPIDIUM** Garman.

Microlepidium Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 180, (*M. grandiceps* Garman).

1653. **Microlepidium grandiceps** Garman.

Gulf of California in deep water.

Microlepidium grandiceps Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 181, Pl. XLIII, Gulf of California.

Genus 540. **ANTIMORA** Günther.

Antimora Günther, Ann. Mag. Nat. Hist., 5th ser., II, 1878, 18 (*A. rostrata* Günther).

1654. **Antimora viola** (Goode and Bean).

Banks of Newfoundland and southward.

Haloporphyrus viola Goode and Bean, Proc. U. S. Nat. Mus., I, 1878, 257, LaHave Bank in 400 fathoms.

1655. **Antimora microlepis** Bean.

Pacific Ocean, off Queen Charlotte Islands.

Antimora microlepis Bean, Proc. U. S. Nat. Mus., XIII, 1890, 38, lat. 51° 00' 23" N., long. 130° 34' W., off Cape St. James, Queen Charlotte Islands, in 875 fathoms.

1656. **Antimora rhina** Garman.

Off Panama in deep water.

Antimora rhina Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 185, off Bay of Panama.

Genus 541. **GADELLA** Lowe.

Gadella Lowe, Proc. Zool. Soc. Lond., 1843, 91 (*G. gracilis* Lowe).

Uraleptus Costa, Wiegmann's Archiv Naturgesch., 1858, 87 (*Gadus maraldi* Risso).

1657. **Gadella maraldi** (Risso).

Nice; Madeira; Naples and Catania; also off island of Nevis in the West Indies.

Gadus maraldi Risso, Ichth. Nice, 1810, 123, pl. 6, fig. 13, Nice.

Gadella gracilis Lowe, Proc. Zool. Soc. Lond., 1843, 91, Madeira.

Genus 542. **LOTELLA** Kaup.

Lotella Kaup, Wiegmann's Archiv, 1858, 88 (*Lota phycis* Temminck and Schlegel).

1658. **Lotella maxillaris** Bean.

Gulf Stream.

Lotella maxillaris Bean, Proc. U. S. Nat. Mus., VII, 1884, 241, lat. 39° 55' N., long. 70° 28' W., in 396 fathoms.

Genus 543. **PHYSICULUS** Kaup.

Physiculus Kaup, Wiegmann's Archiv, 1858, 88 (*P. dalwigkii* Kaup).

1659. **Physiculus kaupi** Poey.

Cuba.

Physiculus kaupi Poey, Repertorio, I, 1865, 186, Matanzas, Cuba.

1660. *Physiculus fulvus* Bean.

Caribbean Sea and north in the Gulf Stream to latitude 35°.

Physiculus fulvus Bean, Proc. U. S. Nat. Mus., VII, 1884, 240, lat. 40° 1' N., long. 69° 56' W., 79 fathoms.

1661. *Physiculus nematopus* Gilbert.

On both coasts of Lower California, in deep water.

Physiculus nematopus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 114, on both coasts of Lower California, at *Albatross* stations 2997, 3011, 3015, and 3016, in 71 to 221 fathoms.

1662. *Physiculus rastrelliger* Gilbert.

Off Panama to southern California in deep water.

Physiculus rastrelliger Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 113, coast of southern California.

1663. *Physiculus longipes* Garman.

Off Panama in deep water.

Physiculus longipes Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 188, Pl. XLII, off Bay of Panama.

Genus 544. *LOTA* Cuvier. *Burbots*.

Les Lottes Cuvier, Règne Anim., ed. 1, II, 1817, 215 (*Gadus lota* L.).

Lota Oken, Isis, 1817, 1182 (*Gadus lota* L.).

1664. *Lota maculosa* (LeSueur). *Burbot; Lake lawyer; Ling; Alekey trout; Eel pout and Long-tailed cat* (trade names at Sandusky, Ohio); *Mud blower*.

Rivers and lakes of New England and the Great Lakes region, north to Arctic seas, and west to the upper Missouri and Columbia River Basins.

Gadus maculosus LeSueur, Jour. Acad. Nat. Sci. Phila., I, 1817, 83, Lake Erie.

Gadus compressus LeSueur, loc. cit., 84, Connecticut River.

Gadus lacustris Mitchell, Amer. Month. Mag., II, 1818, 244, Sebago Lake, Maine; not of Walbaum.

Molva huntia LeSueur, Mém. Mus. Hist. Nat. Paris, V, 1819, 161, Connecticut River.

Lota inornata DeKay, New York Fauna: Fishes, 1842, 283, pl. 45, fig. 145, Hudson River, Lansingburgh, N. Y.

Lota brosmiana Storer, Bost. Jour. Nat. Hist., IV, 1841, 58, pl. 5, fig. 1, Lake Winnipissiogee, N. H.

Genus 545. *MOLVA* Fleming. *Lings*.

Molva Fleming, Hist. Brit. Anim., 1828, 192 (*Gadus molva* L. = *Molva vulgaris* Fleming).

Molva Nilsson, Scandinavian Fauna, IV, 1832, 573 (*Gadus molva* L.).

1665. *Molva molva* (Linnaeus). *Ling; Iverksaak*.

Spitzbergen to Gulf of Gascony, Arcachon; San Juan de Luz; Iceland; Greenland; Faroe Islands.

Gadus molva Linnaeus, Syst. Nat., ed. X, 1758, 254, seas of Europe.

Genus 546. *PHYCIS* Röse. *Codlings*.

Phycis Röse, Synonomia, etc., ed. II, 1793 (*Phycis* Aristotle = *Phycis tinca* Schroeder); not *Phycis* Fabricius, 1798, a genus of insectes.

Phycis Rafinesque, Amer. Month. Mag., 1818, 242 (*Phycis marginatus* Rafinesque = *Blennius chisso* Walbaum).

1666. *Phycis earlli* Bean.

Charleston, S. C.

Phycis earlli Bean, Proc. U. S. Nat. Mus., III, 1880, 69, Charleston, S. C.

1667. *Phycis cirratus* Goode and Bean.

Deep waters of the Gulf of Mexico.

Phycis cirratus Goode and Bean, Oceanic Ichth., 1896, 358, Gulf of Mexico, at lat. 29° 03' 15'' N., long. 88° 16' W.

Genus 547. UROPHYCIS Gill.

Phycis Bloch and Schneider, Syst. Ichth., 1801, 56 (*tinca* = *blennioides*), not *Phycis* Fabricius, 1798, a genus of Lepidoptera.

Urophycis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 240 (*Blennius regius* Walbaum).

Emphyodus Jordan and Evermann, Fish. North and Mid. Amer., 1898, 2552 (*Gadus tenuis* Mitchell).

1668. *Urophycis floridanus* (Bean and Dresel).

Pensacola, Fla.

Phycis floridanus Bean and Dresel, Proc. Biol. Soc. Wash., II, 1884, 100, Pensacola, Fla.

1669. *Urophycis regius* (Walbaum). *Spotted codling; Spotted hake.*

North Atlantic, south to Cape Fear.

Blennius regius Walbaum, Artedi Pisc., III, 1792, 186; after Schöpf.

Enchelyopus regalis Bloch and Schneider, Syst. Ichth., 1801, 53; after Schöpf.

Gadus punctatus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 372, New York.

1670. *Urophycis tenuis* (Mitchill). *Codling; Squirrel hake; Hake; Ling, Thimble-eyed ling; White hake; Old English hake.*

North Atlantic, south to Virginia; abundant northward.

Gadus tenuis Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 371, New York.

Phycis dekayi Kaup, Wiegmann's Archiv, 1858, Pt. I, 89, North America.

?*Phycis rostratus* Günther, Cat., IV, 1862, 353, no locality.

1671. *Urophycis chuss* (Walbaum). *Codling; Squirrel hake; Thimble-eyed ling; Old English hake; Chuss; Fork-beard; Red hake.*

Atlantic coast, chiefly northward.

Blennius chuss Walbaum, Artedi Pisc., III, 1792, 186, Long Island; after Schöpf.

Enchelyopus americanus Bloch and Schneider, Syst. Ichth., 1801, 53; after Schöpf.

Gadus longipes Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 372, pl. 1, fig. 4, New York.

Phycis marginatus Rafinesque, Amer. Month. Mag., II, 1818, 205, Point Judith, R. I.

1672. *Urophycis chesteri* (Goode and Bean).

Off Massachusetts, in deep water; abundant.

Phycis chesteri Goode and Bean, Proc. U. S. Nat. Mus., I, 1878, 256, off Cape Ann, in 140 fathoms.

Genus 548. LÆMONEMA Günther.

Læmonema Günther, Cat., IV, 1862, 356 (*Phycis yarrellii* Lowe).

1673. *Læmonema barbatulum* Goode and Bean.

Gulf Stream.

Læmonema barbatula Goode and Bean, Bull. Mus. Comp. Zool., X, 1883, No. 5, 204, Gulf Stream, in lat. $32^{\circ} 43' 25''$ N., long. $77^{\circ} 20' 30''$ W., in 230 fathoms, and $38^{\circ} 35'$ N., $73^{\circ} 13'$ W.

1674. *Læmonema melanurum* Goode and Bean.

Caribbean Sea, north to New York.

Læmonema melanurum Goode and Bean, Oceanic Ichth., 1895, 363, lat. $30^{\circ} 44'$ N., long. $79^{\circ} 26'$ W., in 440 fathoms.

1675. *Læmonema gracillipes* Garman.

Galapagos Islands to Bay of Panama in deep water.

Læmonema gracillipes Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 187, Pl. XLII; off Galapagos Islands.

Genus 549. BROSME Oken. *Cusk.*

Brosme Oken, Isis, 1817, 1182 (*Gadus brosme* Müller); after Les Brosmes Cuvier.

Brosmerus LeSueur, Mém. Mus. Hist. Nat. Paris, 1819, 158, Pl. XVI (*Brosmerus flavescens* LeSueur = *Gadus brosme* Müller).

Brosmius Cuvier, Régne Anim., ed. 2, II, 1829, 354 (*Gadus brosme* Müller).

1676. *Brosme brosme* (Müller). *Cusk; European cusk; Nejorpallujak; Torsk; Tusk.*

Polar regions, south to Cape Cod.

Gadus brosme Müller, Prodr. Zool. Dan., 1776, 41, Denmark.

Gadus torsk Bonnaterre, Encycl. Meth. Ichth., 1788, 51, Sundmer, Norway; after Ström.

Brosmerus flavesny LeSueur, Mém., Mus. Hist. Nat. Paris, V, 1819, 158, Banks of Newfoundland; Marblehead, Mass.

Family 127. MERLUCHIDÆ. Hakes

Genus 550. MERLUCCIUS Rafinesque. *Hakes.*

Merluccius Rafinesque, Caratteri, 1810, 25 (*M. smiridus* Rafinesque = *Gadus merluccius* L.).

Onus Rafinesque, Indice, 1810, 12; substitute for *Merluccius*.

Merlangus Rafinesque, loc. cit., 167, another substitute for *Merluccius*.

Merlus Gay, in Guichenot, Pesces de Chile, II, 1848, 238 (*Merlus gayi* Guichenot).

Epicopus Günther, Cat., II, 1860, 248 (*Merlus gayi* Guichenot).

1677. *Merluccius merluccius* (Linnæus). *European hake.*

Coasts of Europe, straying to Greenland.

Gadus merluccius Linnæus, Syst. Nat., ed. X, 1758, 254, no locality; Europe.

Gadus ruber Lacépède, Hist. Nat. Poiss., V, 1803, 673, Scotland; Dieppe.

Gadus merlus Risso, Ichth. Nice, 1810, 122, Nice; after Linnæus.

Onus riali Rafinesque, Indice, 1810, 12, 67; substitute for *Gadus merluccius*.

Merlucius esculentus Risso, Europ. Mérid., III, 1826, 220, Nice.

Merluccius ambiguus Lowe, Proc. Zool. Soc. Lond., 1840, 37, Madeira.

Merlucius lanatus Gronow, Cat. Fishes, ed. Gray, 1854, 130, Mediterranean.

1678. *Merluccius bilinearis* (Mitchill). *Silver hake; New England hake; Whiting; Old England hake; Silverfish; American hake; Stockfish.*

Coast of New York and northward.

Stomodon bilinearis Mitchill, Rept. Fish. N. Y., 1814, 7, New York.

Gadus albifrons Mitchill, Jour. Acad. Nat. Sci. Phila., I, 1817, 409, New York.

1679. *Merluccius productus* (Ayres).

Pacific coast, from San Diego northward to Washington.

Merlangus productus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 64 (65), no locality given; types from San Francisco market, probably from San Francisco Bay.

Homalopomus troubridgii Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 132, Astoria, Oreg.

1680. *Merluccius angustimanus* Garman.

Off Panama, in deep water.

Merluccius angustimanus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 183, Pl. XLI, fig. 1, off Bay of Panama.

Order GG. SALMOPERCÆ

Family 128. PERCOPSIDÆ. Sand rollers; Trout perch

Genus 551. PERCOPSIS Agassiz.

Percopsis Agassiz, Lake Superior, 1850, 284 (*P. guttatus* Agassiz = *Salmo omiscomaycus* Walbaum).

Salmoperca Thompson, App. Hist. Vermont, 1853, 33 (*S. pellucida* Thompson = *Salmo omiscomaycus* Walbaum).

- 1681.** *Percopsis omiscomaycus* (Walbaum). *Sand roller; Trout perch.*
Lake Champlain and Delaware River to Ohio River, Kansas, and northward; Great Lakes; Hudson Bay; Red River of the North; Saskatchewan River.
Salmo omiscomaycus Walbaum, Artedi Pisc., 1792, 65, Hudson Bay.
Percopsis guttatus Agassiz, Lake Superior, 1850, 286, Lake Superior.
Salmoperca pellucida Thompson, App. Hist. Vermont, 1853, 33, Lake Champlain.
Percopsis hammondii Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 151, Kansas.
- Genus 552. COLUMBIA** Eigenmann and Eigenmann.
Columbia Eigenmann and Eigenmann, Science, XX, Oct. 21, 1892, 233 (*C. transmontana* Eigenmann and Eigenmann).
- 1682.** *Columbia transmontana* Eigenmann and Eigenmann.
Columbia River Basin; Umatilla and Wallula Rivers.
Columbia transmontana Eigenmann and Eigenmann, Science, XX, 1892, 233, mouth of Umatilla River, Oreg.

Order HH. XENARCHI

Family 129. APHREDODERIDÆ. Pirate perches

- Genus 553. APHREDODERUS** LeSueur. *Pirate perches.*
Aphredoderus LeSueur, in Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 329 (445), (*A. gibbosus* LeSueur = *Scolopsis sayanus* Gilliams).
Sternotremia Nelson, Bull. Ill. Lab. Nat. Hist., I, 1876, 39 (*S. isolepis* Nelson); young with the vent posterior.
Asternotremia Nelson, in Jordan, Bull. U. S. Nat. Mus., X, 1877, 51. *Aphododerus*, attempted correction.
- 1683.** *Aphredoderus sayanus* (Gilliams). *Pirate perch; Pirate; Mud perch.*
From New York in coastwise streams to Texas; also in the Mississippi Basin, north to Michigan; abundant in sluggish waters.
Scolopsis sayanus Gilliams, Jour. Acad. Nat. Sci. Phila., IV, 1824, 81, near Philadelphia.
Aphredoderus gibbosus LeSueur, in Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 331 (448), Lake Pontchartrain, La.
Sternotremia isolepis Nelson, Bull. Ill. Lab. Nat. Hist., I, 1876, 39. Calumet River, Ill.; young with vent posterior.
Aphododerus cookianus Jordan, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 60, Sawyers Creek, Kendallville, Ind.
Asternotremia mesotrema Jordan, Bull. U. S. Nat. Mus., X, 1877, 52, Flint River, Ga.

Order II. ALLOTRIOGNATHI

Suborder HISTICHTHYES

Family 130. LOPHOTIDÆ

- Genus 554. LOPHOTES** Giorna.
Lophotes Giorna, Mém. Acad. Sci. Torino, XVI, 1805, 19 (*L. cepedianus* Giorna).
Leptopus Rafinesque, Specchio Scienze Palermo, I, 1814, 16 (*L. peregrinus* Rafinesque = *Lophotes cepedianus* Giorna).
Podolepites Rafinesque, Anal. Nat., 1815, 93; a substitute for *Leptopus*; same type.
- 1684.** *Lophotes cepedianus* Giorna.
Open seas north to Long Beach, Calif. (W. F. Thompson), and to Japan, if *Lophotes capellei* is the same.
Lophotes cepedianus Giorna, Mém. Acad. Sci. Torino, XVI, 1803, 19, Mediterranean.
Leptopus peregrinus Rafinesque, Specchio Scienze Palermo, I, 1814, 16, Palermo.
?Lophotes capellei Schlegel, Fauna Japonica, 1845, 132, pl. 71, Japan.

Suborder TÆNIOSOMI

Family 131. TRACHIPTERIDÆ. King of the herrings

Genus 555. TRACHIPTERUS Gouan.

Trachipterus Gouan, Hist. Nat. Poiss., 1770, 103 (*T. gouani* Gouan = *Cephalo trachyptera* Gmelin).

Trachipterus of authors; corrected spelling.

Bogmarus Schneider, Syst. Ichth., 1801, 518 (*B. islandicus* Schneider = *Trachipterus arcticus*).

Vogmarus Reid, Ann. Mag. Nat. Hist., III, 1849, 456 (*B. islandicus* Schneider); substitute name as the Danish name is Vaagnær.

1685. *Trachipterus arcticus* (Brünnich). *Dealfish*; *Vaagmær*.

Arctic Sea, Iceland to Norway.

Gymnogaster arcticus Brünnich, Nya Handl., Dansk. Vidensk. Selsk. Skrifter, III, 1788, 408, Tab. B, figs. 1-3.

Bogmarus islandicus Schneider, Syst. Ichth., 1801, 518, Tab. 101.

Trachipterus bogmarus Cuvier and Valenciennes, Hist. Nat. Poiss., X, 1827?, 346.

Trachipterus vogmarus Reinhardt, Dansk. Vidensk. Selsk. Skrifter, VII, 1838, 71, with fig.

1686. *Trachipterus rex-salmonorum* Jordan and Gilbert. *King of the salmon*.

Off coast of California.

Trachipterus rex-salmonorum Jordan and Gilbert, Proc. Calif. Acad. Sci., 2d ser., IV, 1894, 144, pl. 9, open sea, outside Bay of San Francisco.

Trachipterus seleniris Snyder, Proc. Acad. Nat. Sci. Phila., LX, 1908, 319, Monterey; adult form.

1687. *Trachipterus trachyurus* Poey.

Cuba.

Trachipterus trachyurus Poey, Memorias, II, 1861, 420, Cuba.

Suborder ATELAXIA

Family 132. STYLEPHORIDÆ

Genus 556. STYLEPHORUS Shaw.

Stylephorus Shaw, Trans. Linn. Soc. Lond., I, 1791, 90 (*S. chordatus* Shaw).

1688. *Stylephorus chordatus* Shaw.

Open sea, Atlantic; very rare.

Stylephorus chordatus Shaw, Trans., Linn. Soc. Lond., I, 1791, 90, pl. 6, Atlantic, between Cuba and Martinique.

Order JJ. SELENICHTHYES

Family 133. LAMPRIDÆ. Opah; Moonfish

Genus 557. LAMPRIS Retzius.

Lampris Retzius, Vetensk. Akad. Nya Handl., XX, 1799, 91 (*Zeus guttatus* Brünnich = *Zeus regius* Bonnaterre).

Chrysotus Lacépède, Hist. Nat. Poiss., III, 1803 (*Zeus luna* Gmelin). *Echimythes* Gistel, Thierreich, X, 1848 (*Zeus luna* Gmelin).

1689. *Lampris regius* (Bonnaterre). *Mariposa*; *Opah*; *Moonfish*; *San Pedro fish*; *Cravo*; *Jerusalem haddock*; *Glancefish*; *Gudlax*; *Poisson lune*.

Open seas, Atlantic and Pacific; Madeira; Newfoundland; Maine; Cuba; Monterey; Hawaii; Japan; recently taken off San Pedro.

*Zeus regius** Bonnaterre, Encycl. Meth. Ichth., 1788, 72, pl. 39, Torbay, England.

Zeus guttatus Brünnich, Danske Skrifter, III, 1788, 403, Elsinore, Denmark.

Zeus luna Gmelin, Syst. Nat., ed. XIII, 1789, 1225, Normandy.

* The name *regius* certainly has priority over *luna* and probably over *guttatus* also.

- Zeus stroemii* Walbaum, Artedi Pisc., ed. 2, 1792, 398; Norway; after Ström.
Scomber gunneri Bloch and Schneider, Syst. Ichth., 1801, 38, Throndhjem.
Zeus imperialis Shaw, Nat. Misc., IV, 1803, pl. 140, Torbay; after Pennant.
Zeus opah Shaw, Gen. Zool., IV, 1803, 287, Torbay; after Pennant.
Lampris lauta Lowe, Fishes Madeira, 1843, 27, Madcira.

Order KK. HETEROSOMATA. Flounders; Flatfishes

Family 184. BOTHIDÆ

Genus 558. LOPHOPSETTA Gill. Windowpanes.

Lophopsetta Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 220 (*Pleuronectes maculatus* Mitchell).

1690. *Lophopsetta maculata* (Mitchill). *Windowpane*.
 Atlantic coast of United States, from Casco Bay to South Carolina; common.
Pleuronectes maculatus Mitchell, Rept. in part, Fish. N. Y., 1814, 9, New York.
Pleuronectes aquosus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 389, pl. 2, fig. 3, New York.

Genus 559. PLATOPHRYS Swainson.

Platophrys Swainson, Nat. Hist. Fish., Reptiles and Amphib., II, 1839, 302 (*Pleuronectes ocellatus* Spix and Agassiz).

Peloria Cocco, Intorn. Aleuni Pesci, etc., 1844, 21 (*P. heckeli* Cocco).
Coccolus Bonaparte loc. cit., 21 (*C. annectens* Cocco, a larva of *Platophrys*).

Bothus Bonaparte, Cat. Metod., 1846, 49 (*podas*); not of Rafinesque.
Rhomboichthys Bleeker, Act. Soc. Sci. Indo-Neder., I, 1856, 67 (*R. myriaster* Bleeker).

1691. *Platophrys spinosus* (Poey).

West Indies.

Rhomboichthys spinosus Poey, Synopsis, 1868, 409, Cuba.

1692. *Platophrys constellatus* Jordan.

Galapagos Islands.

Platophrys constellatus Jordan, in Jordan and Goss, Rept. U. S. Fish Com., XIV, 1886 (1889), 266, James Island, Galapagos Islands.

1693. *Platophrys ocellatus* (Agassiz).

Tropical America; sandy shores from Long Island to Rio Janeiro.

Rhombus ocellatus Agassiz, in Spix, Pisc. Brasil., 1829, 85, pl. 46, Brazil.

Rhombus bahianus Castelnau, Anim. Amer. Sud, 1855, Bahia.

Platophrys nebularis Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 31, Key West, Fla.

1694. *Platophrys maculifer* (Poey).

West Indies.

?*Pleuronectes maculiferus* Poey, Memorias, II, 1861, 316, Cienfuegos, Cuba.

1695. *Platophrys ellipticus* (Poey).

West Indies.

Pleuronectes ellipticus Poey, Memorias, II, 1861, 315, Cuba.

1696. *Platophrys lunatus* (Linnæus). *Peacock flounder*.

West Indies.

Solea lunata et punctata (the sole) Catesby, Nat. Hist. Carolinas, 1725, Tab. 27, Bahamas.

Pleuronectes lunatus Linnaeus, Syst. Nat., ed. X, 1758, 269, Bahamas; based on Catesby and the various copyists.

Pleuronectes argus Bloch, Ichthyologia, 1783, Tab. 48, Martinique; after Plumier.

?*Pleuronectes surinamensis* Bloch and Schneider, Syst. Ichth., 1801, 156, Surinam; *satis parva et glabra*.

1697. **Platophrys leopardinus** (Günther).

Gulf of California.

Rhomboideichthys leopardinus Günther, Cat., IV, 1862, 434, locality unknown.

Platophrys leopardinus Jordan, Proc. U. S. Nat. Mus., VII, 1884, 260, specimen from Guaymas, Sonora, Mexico.

Genus 560. **SYACIUM** Ranzani.

Syacium Ranzani, Nov. Spec. Pisc., 1840, 20 (*S. micrurum* Ranzani).

Hemirhombus Bleeker, Versl. Akad. Amsterdam, XIII, 1862, 425 (*H. guineensis* Bleeker).

Aramaca Jordan and Goss, in Jordan, Cat. Fish. N. Amer., 1885, 133 (*Citharichthys pætulus* Jordan and Gilbert = *Pleuronectes aramaca* Dönnendorf = *Pleuronectes papillosum* Linnæus).

1698. **Syacium papillosum** (Linnæus).

West Indies.

Aramaca Marcegrave, Hist. Pisc. Brasil., 1648, 181, Brazil.

Pleuronectes papillosum Linnæus, Syst. Nat., ed. X, 1758, 271, Brazil; based on Marcegrave.

Rhombus soleæformis Agassiz, in Spix, Pisc. Brasil., 1829, 86, Tab. 47, Atlantic Ocean.

1699. **Syacium micrurum** Ranzani.

West Indies; Key West to Rio Janeiro.

Syacium micrurum Ranzani, Nov. Spec. Pisc. Diss. Sec., 1840, 20, pl. 5, Brazil.

Hippoglossus ocellatus Poey, Memorias, II, 1861, 314, Cuba.

Hemirhombus aramaca Günther, Cat., IV, 1862, 42, Cuba; Jamaica; not *Rhombus aramaca* Cuvier.

Citharichthys ethalion Jordan, Proc. U. S. Nat. Mus., IX, 1886, 52, Habana.

1700. **Syacium latifrons** (Jordan and Gilbert).

Pacific coast of tropical America; Panama.

Citharichthys latifrons Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 334, Panama.

1701. **Syacium ovale** (Günther).

Pacific coast of tropical America; Mazatlan to Panama.

Hemirhombus ovalis Günther, Proc. Zool. Soc. Lond., 1864, 154, Panama.

Genus 561. **CITHARICHTHYS** Bleeker. Whiffs.

Citharichthys Bleeker, Versl. Akad. Amsterd., XIII, 1862, 427 (*C. cayennensis* Bleeker = *C. spilopterus* Günther).

Orthopsetta Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 330 (*Psettichthys sordidus* Girard).

Metoponops Gill, loc. cit., XVI, 1864, 198 (*M. cooperi* Gill = *Psettichthys sordidus* Girard).

Subgenus **ORTHOPSETTA** Gill.1702. **Citharichthys sordidus** (Girard).

Pacific coast of North America, in water of moderate depth from British Columbia to Lower California.

Psettichthys sordidus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 142, San Francisco; Tomales Bay.

Metaponops cooperi Gill, loc. cit., XVI, 1864, 198, Santa Barbara; shriveled specimen.

1703. **Citharichthys fragilis** Gilbert.

Gulf of California.

Citharichthys fragilis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 120, Gulf of California, east coast of Lower California.

1704. **Citharichthys xanthostigma** Gilbert.

Magdalena Bay, west coast of Lower California to Hermosa Beach, southern California, and in Gulf of California.

Citharichthys xanthostigma Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 120, Gulf of California, west coast of Lower California, and Magdalena Bay.

1705. *Citharichthys stigmatus* Jordan and Gilbert.
 Coast of southern California.
Citharichthys stigmatus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 410, 411, Santa Barbara, Calif.
- Subgenus CITHARICHTHYS.
1706. *Citharichthys dinoceros* Goode and Bean.
 Deep waters of Gulf of Mexico.
Citharichthys dinoceros Goode and Bean, Bull. Mus. Comp. Zool., XII, No. 5, 157, off Martinique, St. Lucia, and Barbados.
1707. *Citharichthys platophrys* Gilbert.
 Southwest of Panama; one specimen known.
Citharichthys platophrys Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 454, Albatross station 2799, southwest of Panama.
1708. *Citharichthys arctifrons* Goode.
 Deep waters of Gulf Stream.
Citharichthys arctifrons Goode, Proc. U. S. Nat. Mus., III, 1880, 341, 472, Gulf Stream off southern coast of New England.
1709. *Citharichthys unicornis* Goode.
 Deep waters of Gulf Stream.
Citharichthys unicornis Goode, Proc. U. S. Nat. Mus., III, 1880, 342, Gulf Stream off southeast of New England.
1710. *Citharichthys uhleri* Jordan.
 West Indies.
Citharichthys uhleri Jordan, Rept. U. S. Fish Com., XIV, 1886, 273, Haiti.
1711. *Citharichthys macrops* Dresel.
 South Atlantic and Gulf coasts of the United States.
Citharichthys macrops Dresel, Proc. U. S. Nat. Mus., VII, 1884, 539, Pensacola, Fla.
1712. *Citharichthys spilopterus* Günther.
 Both coasts of tropical America, north to New Jersey and Mazatlan.
Citharichthys spilopterus Günther, Cat., IV, 1862, 421, New Orleans, San Domingo, Jamaica.
Citharichthys cayennensis Bleeker, Compt. Rend. Acad. Sci. Amsterdam, XIII, 1862, 6, Cayenne, name only.
Citharichthys guatemalensis Bleeker, Nederl. Tijdschr. Dierk., 1864, 73, Guatemala.
Hemirhombus fuscus Poey, Synopsis, 1868, 406, Habana.
1713. *Citharichthys arenaceus* Evermann and Marsh.
 Porto Rico and Santo Domingo.
Citharichthys arenaceus Evermann and Marsh, Bull. U. S. Fish Com., XX, first part, 1900 (1902), 326, fig. 106, Mayaguez, Porto Rico.
1714. *Citharichthys gibberti* Jenkins and Evermann.
 Pacific coast of tropical America, from Gulf of California to Panama.
Citharichthys gibberti Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 157, Guaymas, Mexico.
Citharichthys sumichrasti Jordan and Goss, Rept. U. S. Fish Com., XIV, 1886 (1889) 276, Rio Zanatenco, Chiapas, Mexico.
- Genus 562. *ETROPOUS* Jordan and Gilbert.
Etropus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 364 (*E. crossotus* Jordan and Gilbert).
1715. *Etropus microstomus* (Gill).
 Coast of New Jersey.
Citharichthys microstomus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 223, Beesleys Point, N. J.
1716. *Etropus micros* (Fowler).
Citharichthys micros Fowler, Proc. Acad. Nat. Sci. Phila., LXIII, 1911, 200, with figure, Corsons Inlet, Cape May, N. J.

1717. *Etropus rimosus* Goode and Bean.

Gulf of Mexico.

Etropus rimosus Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 593, coast of Florida between Pensacola and Cedar Keys, in 21 fathoms.

1718. *Etropus crossotus* Jordan and Gilbert.

Tropical America on both coasts, north to Guaymas, Mexico, and Cape Hatteras, N. C.

Etropus crossotus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 364, Mazatlan, Mexico.

Genus 563. *PERISSIAS* Jordan and Evermann.

Perissias Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2667 (*Platophrys tæniopterus* Gilbert).

1719. *Perissias tæniopterus* (Gilbert).

Gulf of California and western coast of Lower California.

Platophrys tæniopterus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 118, Gulf of California north of La Paz, at *Albatross* station 2298, lat. 24° 51' N., long. 110° 39' W., in 40 fathoms.

Genus 564. *ENGYOPHRYS* Jordan and Bollman.

Engyophrys Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889 (1890), 176 (*E. sancti-laurentii* Jordan and Bollman).

1720. *Engyophrys sancti-laurentii* Jordan and Bollman.

Pacific Ocean off coast of Colombia.

Engyophrys sancti-laurentii Jordan and Bollman, Proc., U. S. Nat. Mus., XII, 1889, 176, Pacific Ocean off coast of Colombia, at *Albatross* station 2805, 7° 56' N., 79° 41' 30" W., and station 2795, lat. 7° 57' N., long. 78° 55' W., in 33 and 51 fathoms.

Genus 565. *TRICHOPSETTA* Gill.

Trichopsetta Gill, Proc. U. S. Nat. Mus., XI, 1888, 601 (*Arnoglossus ventralis* Goode and Bean).

1721. *Trichopsetta ventralis* (Goode and Bean).

Gulf of Mexico, in deep water.

Citharichthys ventralis Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 592, Gulf of Mexico, *Albatross* station 2386, lat. 29° 15' N., long. 88° 6' W., in 60 fathoms.

Genus 566. *MONOLENE* Goode.

Monolene Goode, Proc., U. S. Nat. Mus., III, 1880, 338 (*M. sessilicauda* Goode).

Thyris Goode, loc. cit., 344 (*T. pellucidus* Goode); larva of *Monolene*. *Delothyrus* Goode, loc. cit., VI, 1883, 110; substitute for *Thyris*, pre-occupied.

1722. *Monolene sessilicauda* Goode.

Deep waters of the Gulf Stream.

Monolene sessilicauda Goode, Proc. U. S. Nat. Mus., III, 1880, 337, 338, deep sea south of New England.

Thyris pellucidus Goode, loc. cit., 344, Gulf Stream off the coast of Rhode Island.

1723. *Monolene atrimana* Goode and Bean.

Deep waters of the Caribbean Sea.

Monolene atrimana Goode and Bean, Bull. Mus. Comp. Zool., XII, 1886, 155, deep waters off Barbados.

Genus 567. *CYCLOPSETTA* Gill.

Cyclopsetta Gill, Proc. U. S. Nat. Mus., XI, 1888, 601 (*Hemirhombus fimbriatus* Goode and Bean).

1724. *Cyclopsetta fimbriata* (Goode and Bean).

Deep waters of the Gulf of Mexico.

Hemirhombus fimbriatus Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 591, deep waters of the Gulf of Mexico, between the Mississippi Delta and Cedar Keys, Fla.

1725. *Cyclopsetta quernea* (Jordan and Bollman).
 Pacific coast of America, off coast of Colombia.
Azevia quernea Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889,
 174, Pacific Ocean off coast of Colombia, at *Albatross* station 2802,
 lat. $8^{\circ} 38' N.$, long. $79^{\circ} 31' 30'' W.$
1726. *Cyclopsetta chittendeni* B. A. Bean.
 Trinidad Island.
Cyclopsetta chittendeni B. A. Bean, Proc. U. S. Nat. Mus., XVII, 1894,
 635, Trinidad Island.
- Genus 568. *AZEVIA* Jordan and Goss.
Azevia Jordan and Goss, Rept. U. S. Fish Com., XIV, 1886 (1888),
 271 (*Citharichthys panamensis* Steindachner).
1727. *Azevia panamensis* (Steindachner).
 Pacific coast of Central America.
Citharichthys panamensis Steindachner, Ichth. Beitr., III, 1875, 62,
 Panama.
- Genus 569. *PLATOTICHTHYS* Nichols.
Platotichthys Nichols, Bull. Amer. Mus. Nat. Hist., XLIV, 1921, 21
 (*P. chartes* Nichols).
1728. *Platotichthys chartes* Nichols.
 Bahamas.
Platotichthys chartes Nichols, Bull. Amer. Mus. Nat. Hist., XLIV,
 1921, 21, pl. III, Turk Island, Bahamas.
- Family 135. HIPPOGLOSSIDÆ. Halibuts
- Genus 570. *ATHERESTHES* Jordan and Gilbert.
Atheresthes Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 51
 (*Platysomatischthys stomias* Jordan and Gilbert).
1729. *Atheresthes stomias* (Jordan and Gilbert). *Arrow-toothed halibut.*
 San Francisco to Alaska.
Platysomatischthys stomias Jordan and Gilbert, Proc. U. S. Nat. Mus.,
 III, 1880, 301, off San Francisco, Calif.
- Genus 571. *REINHARDTIUS* Gill.
Reinhardtius Gill, Cat. Fishes East Coast N. Amer., Suppl. Proc. Acad.
 Nat. Sci. Phila., XIII, 1861, 50 (*Pleuronectes cynoglossus* Fabricius; not
 of Linnaeus = *Pleuronectes hippoglossoides* Walbaum).
Platysomatischthys Bleeker, Versl. Akad. Amsterdam, XIII, 1862, 426
 (*Pleuronectes pinguis* Fabricius).
1730. *Reinhardtius hippoglossoides* (Walbaum). *Greenland halibut.*
 Arctic parts of the Atlantic, south to Finland and Grand Banks.
Pleuronectes cynoglossus Fabricius, Fauna Greenlandica, 1780, 163,
 Greenland; not of Linnaeus.
Pleuronectes hippoglossoides Walbaum, Artedi. Pisc., 1792, 115; based
 on Fabricius.
Pleuronectes pinguis Fabricius, Zoologiske Bidrag., 1824, 43, Greenland.
Hippoglossus grænlandicus Günther, Cat., IV, 1862, 404, Greenland.
- Genus 572. *HIPPOGLOSSUS* Cuvier. *Halibut.*
Hippoglossus Cuvier, Règne Anim., ed. 1, II, 1817, 221 (*Pleuronectes*
hippoglossus Linnaeus).
1731. *Hippoglossus hippoglossus* (Linnaeus). *Common halibut.*
 All northern seas, southward in deep waters to France, Sandy Hook,
 and northern California.
Pleuronectes hippoglossus Linnaeus, Syst. Nat., ed. X, 1758, 269,
 European oceans.
Hippoglossus maximus Gottsche, Archiv für Naturgesch., 1835, 164,
 no locality.
Hippoglossus gigas Swainson, Nat. Hist. Class'n Anim., II, 1839, 302,
 no locality.
Hippoglossus ponticus Bonaparte, Cat. Mctod., 1846, 47, Black Sea;
 after Pallas.

Genus 573. LYOPSETTA Jordan and Goss.

Lyopsetta Jordan and Goss, in Jordan, Cat. Fish. N. Amer., 1885, 135
 (*Hippoglossoides exilis* Jordan and Gilbert).

1732. **Lyopsetta exilis** (Jordan and Gilbert).

North Pacific, in rather deep water; San Francisco to Alaska.
Hippoglossoides exilis Jordan and Gilbert, Proc. U. S. Nat. Mus., III,
 1880, off San Francisco, between the Golden Gate and Point Reyes,
 Calif.

Genus 574. HIPPOGLOSSOIDES Gottsche.

Hippoglossoides Gottsche, Archiv für Naturgesch., 1835, 164 (*Pleuronectes limanda* Linnæus).

Citharus Reinhardt, König. Dansk. Vid Selsk., 1838, 116 (*Pleuronectes platessoides* Fabricius; not *Citharus* Bleeker, 1862).

Drepanopsetta Gill, Cat. Fish. East Coast N. Amer., Suppl. Proc. Acad. Nat. Sci. Phila., XIII, 1861, 50 (*Pleuronectes platessoides* Fabricius).

Pomatopsetta Gill, Proc., Acad. Nat. Sci. Phila., XVI, 1864, 217 (*Platessa dentata* = *Pleuronectes platessoides* Fabricius).

1733. **Hippoglossoides platessoides** (Fabricius). *Sand dab; Dab.*

North Atlantic, south to Cape Cod, and coasts of England and Scandinavia.

Pleuronectes linguatula Müller, Prodr. Zool. Dan., 1776, 45; not of Linnæus.

Pleuronectes platessoides Fabricius, Fauna Grænlandica, 1780, 164, Greenland.

Pleuronectes limandooides Bloch, Naturgesch. Ausl. Fische, III, 1787, 24, Tab. 186, Europe.

1734. **Hippoglossoides classodon** Jordan and Gilbert.

Puget Sound and northward.

Hippoglossoides classodon Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 278, Puget Sound at Seattle and Tacoma, Wash.

1735. **Hippoglossoides robustus** Gill and Townsend.

Bering Sea.

Hippoglossoides robustus Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17), Bering Sea, lat. 56° 14' N., long. 164° 08' W., Albatross station 3541 in 49 fathoms.

1736. **Hippoglossoides hamiltoni** Jordan and Gilbert.

Off Dalnoi Point, Kamchatka.

Hippoglossoides hamiltoni Jordan and Gilbert, Rept. Fur-seal Invest., 1898, 489, Dalnoi Point, Avatcha Bay, Kamchatka, in 16 fathoms.

Genus 575. EOPSETTA Jordan and Goss.

Eopsetta Jordan and Goss, in Jordan, Cat. Fish. N. Amer., 1885, 135
 (*Hippoglossoides jordani* Lockington).

1737. **Eopsetta jordani** (Lockington). *California sole.*

West coast of United States, from Puget Sound to Monterey.

Hippoglossoides jordani Lockington, Proc. U. S. Nat. Mus., II, 1879, 73, San Francisco.

Genus 576. PSETTICHTHYS Girard.

Psettichthys Girard, Proc. Acad. Nat. Sci. Phila., VI, 1854, 140 (*P. melanostictus* Girard).

1738. **Psettichthys melanostictus** Girard.

Pacific coast of North America from Alaska south to Monterey.

Psettichthys melanostictus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 140, San Francisco; Astoria, Oreg.

Family 136. PARALICHTHYIDÆ. Bastard halibuts**Genus 577. HIPPOGLOSSINA Steindachner.**

Hippoglossina Steindachner, Ichth. Beitr., V, 1876, 13 (*H. macrops* Steindachner).

1739. **Hippoglossina atomata** Eigenmann and Eigenmann.
Off San Diego, Calif.
Hippoglossina atomata Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., III, 1890, 22, San Diego, Calif.
1740. **Hippoglossina macrops** Steindachner.
Pacific coast of Mexico at Mazatlan.
Hippoglossina macrops Steindachner, Ichth. Beitr., V, 1876, 13, pl. 3, Mazatlan, Mexico.
1741. **Hippoglossina bollmani** Gilbert.
Pacific coast off Colombia.
Hippoglossina macrops Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 175; not of Steindachner.
Hippoglossina bollmani Gilbert, loc. cit., XIII, 1890, 122, off coast of Colombia at Albatross station 2805, lat. $7^{\circ} 56' N.$, long. $79^{\circ} 41' 30'' W.$ in $51\frac{1}{2}$ fathoms.
1742. **Hippoglossina sabanensis** Boulenger.
Isthmus of Panama.
Hippoglossina sabanensis Boulenger, Bolletino Mus. Zool. Univ. Torino, XIV, 1899, 4, Rio Sabana, Darien.
- Genus 578. LIOGLOSSINA** Gilbert.
Lioglossina Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 122 (*L. tetraphthalma* Gilbert).
1743. **Lioglossina tetraphthalma** Gilbert.
Gulf of California.
Lioglossina tetraphthalmus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 122, Gulf of California, Albatross stations 3014 and 3016, in 29 and 76 fathoms.
- Genus 579. VERASPER** Jordan and Gilbert.
Verasper Jordan and Gilbert, Rept. Fur-seal Invest., 1898, 490 (*V. moseri* Jordan and Gilbert).
1744. **Verasper moseri** Jordan and Gilbert.
Kuril Islands.
Verasper moseri Jordan and Gilbert, Rept. Fur-seal Invest., 1898, 490, Pl. LXXXV, Shana Bay, Iturup Island, Kuril Group.
- Genus 580. XYSTREURYS** Jordan and Gilbert.
Xystreurus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 34 (*X. liolepis* Jordan and Gilbert).
1745. **Xystreurus liolepis** Jordan and Gilbert.
Coast of California, from Point Conception southward.
Xystreurus liolepis Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 34, Santa Barbara, Calif.
- Genus 581. PARALICHTHYS** Girard. *Bastard halibut*.
Paralichthys Girard, Pac. R. R. Surv., X, 1859, 146 (*P. maculosus* Girard).
Chænopsetta Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 218 (*Platesa oblonga* Storer).
Pseudorhombus Bleeker, Compt. Rend. Acad. Sci. Amsterdam, XIII, 1862, 5 (*Rhombus polytypus* Bleeker).
Uropsetta Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 330 (*Hippoglossus californicus* Ayres).
1746. **Paralichthys californicus** (Ayres). *Bastard halibut; Monterey halibut*.
Coast of California, Tomales Bay to San Diego.
Hippoglossus californicus Ayres, Proc. Calif. Acad. Nat. Sci., ser. 1, II, 1859, 29, and 1860, fig. 10, San Francisco.
1747. **Paralichthys magdalenæ** Abbott.
Coast of Lower California.
Paralichthys magdalenæ Abbott (James Francis), in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2871, Magdalena Bay, Lower California.

1748. **Paralichthys astuarius** Gilbert and Scofield.
 Mouth of Colorado River, Gulf of California.
Paralichthys astuarius Gilbert and Scofield, Proc. U. S. Nat. Mus., XX, 1898, 499, Pl. XXIX, Shoal Point, mouth of the Colorado River.
1749. **Paralichthys brasiliensis** (Ranzani).
 Atlantic coast of South America; said to range northward to Guatemala.
Hippoglossus brasiliensis Ranzani, Nov. Spec. Pisc., 1840, 10, Tab. 3, Brazil.
Pseudorhombus vorax Günther, Cat., IV, 1862, 429, South America.
1750. **Paralichthys woolmani** Jordan and Williams.
 Guaymas to Panama.
Paralichthys woolmani Jordan and Williams, in Gilbert, Proc. U. S. Nat. Mus., XIX, 1896, 457, Galapagos Islands.
1751. **Paralichthys sinaloæ** Jordan and Abbott.
 Pacific coast of Mexico.
Paralichthys sinaloæ Jordan and Abbott (J. F.), in Jordan and Evermann, Fish. North and Mid. Amer., 1898, 2872, Mazatlan and La Paz.
1752. **Paralichthys dentatus** (Linnæus). *Summer flounder; Fluke.*
 Atlantic coast of United States from Cape Cod to Florida.
Pleuronectes dentatus Linnæus, Syst. Nat., ed. XII, 1766, and of numerous copyists.
Pleuronectes melanogaster Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 390, New York; doubled example.
Platessa ocellaris DeKay, N. Y. Fauna: Fishes, 1842, 300, p. 47, fig. 152, New York.
Paralichthys ophryas Jordan and Gilbert, Synopsis, 1883, 822, Charleston, S. C.
1753. **Paralichthys lethostigmus** Jordan and Gilbert. *Southern flounder.*
 South Atlantic and Gulf coasts of the United States, north to New York.
Platessa oblonga DeKay, N. Y. Fauna: Fishes, 1842, 299, pl. 48, fig. 156, New York; not *Pleuronectes oblongus* Mitchell.
Paralichthys lethostigma Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 237, Jacksonville, Fla.
1754. **Paralichthys oblongus** (Mitchill). *Four-spotted flounder.*
 Coasts of New England and New York.
Pleuronectes oblongus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 391, New York.
1755. **Paralichthys squamilentus** Jordan and Gilbert.
 South Atlantic and Gulf coasts of United States.
Paralichthys squamilentus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 303, Pensacola, Fla.
1756. **Paralichthys alboguttus** Jordan and Gilbert. *Gulf flounder.*
 South Atlantic and Gulf coasts of the United States.
Paralichthys albogutta Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 302, Pensacola, Fla., and Beaufort, N. C.
- Genus 582. ANCYLOPSETTA** Gill.
Ancylopsetta Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 224 (*A. quadrocellatus* Gill).
1757. **Ancylopsetta quadrocellatus** Gill.
 South Atlantic and Gulf coasts of the United States.
Ancylopsetta quadrocellatus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 224, Pensacola; not *Platessa quadrocellata* Storer.
Paralichthys ommatus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 616, Charleston, S. C.
- Genus 583. RAMULARIA** Jordan and Evermann.
Ramularia Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2633 (*Ancylopsetta dendritica* Gilbert).
1758. **Ramularia dendritica** (Gilbert).
 Gulf of California.
Ancylopsetta dendritica Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 121, Gulf of California, at Albatross station 3022, lat. $30^{\circ} 58' 30''$ N., long. $113^{\circ} 17' 15''$ W., in 11 fathoms.

Genus 584. NOTOSEMA Goode and Bean.

Notosema Goode and Bean, Bull. Mus. Comp. Zool., X, 1882, 193 (*N. dilecta* Goode and Bean).

1759. *Notosema dilectum* Goode and Bean.

Gulf Stream.

Notosema dilecta Goode and Bean, Bull. Mus. Comp. Zool., X, 1883, 193, Gulf Stream off the coast of South Carolina.

Genus 585. GASTROPSETTA B. A. Bean.

Gastropsetta Bean, Proc. U. S. Nat. Mus., XVII, 1894, 633 (*G. frontalis* Bean).

1760. *Gastropsetta frontalis* B. A. Bean.

Key West and Apalachicola, Fla.

Gastropsetta frontalis B. A. Bean, Proc. U. S. Nat. Mus., XVII, 1894, 633, Key West and Apalachicola, at Albatross stations 2317 and 2783, in 45 fathoms.

Family 137. PLEURONECTIDÆ

Genus 586. PLEURONICHTHYS Girard.

Pleuronichthys Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1854, 139 (*P. cænosus* Girard).

Heteroprosopon Bleeker, Versl. Akad. Amsterdam, XIII, 1862 (*Pleuronectes cornutus* Temminck and Schlegel).

Parophrys Günther, Cat., IV, 1862, 454; not of Girard.

1761. *Pleuronichthys decurrens* Jordan and Gilbert.

Pacific coast of United States, south to Monterey.

Pleuronichthys decurrens Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 453, San Francisco; Monterey Bay.

1762. *Pleuronichthys verticalis* Jordan and Gilbert.

Coast of California, in deep water.

Pleuronichthys verticalis Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 49, San Francisco.

1763. *Pleuronichthys cænosus* Girard.

Alaska, southward to San Diego; most common about Puget Sound.

Pleuronichthys cænosus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 139, San Francisco.

1764. *Pleuronichthys nephelus* Starks and Thompson.

Puget Sound.

Pleuronichthys nephelus Starks and Thompson, Proc. U. S. Nat. Mus., XXXVIII, 1911, 282, fig. 1, San Juan Islands, Wash.

1765. *Pleuronichthys ritteri* Starks and Morris.

San Diego Bay.

Pleuronichthys ritteri Starks and Morris in Starks & Thompson, Proc. U. S. Nat. Mus., XXXVIII, 1911, 287, San Diego, Calif.

1766. *Pleuronichthys ocellatus* Starks and Thompson.

Gulf of California.

Pleuronichthys ocellatus Starks and Thompson, Proc. U. S. Nat. Mus., XXXVIII, 1911, 285, fig. 2, Gulf of California.

Genus 587. HYPSSOPSETTA Gill.

Hypsopsetta Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 330 (*Pleuronichthys guttulatus* Girard).

1767. *Hypsopsetta guttulata* (Girard). *Diamond flounder*.

Coast of California, Cape Mendocino to Magdalena Bay.

Pleuronichthys guttulatus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 137, Tomales Bay, Calif.

Parophrys ayresi Günther, Cat., IV, 1862, 457, San Francisco.

Genus 588. PAROPHYSYNS Girard.

Parophrys Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 139 (*P. vetulus* Girard).

1768. *Parophrys vetulus* Girard.

Pacific coast of North America, Alaska to Santa Barbara.

Parophrys vetulus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 140, California.

Pleuronectes diagrammus Günther, Cat., IV, 1862, 445, Victoria, British Columbia.

Parophrys hubbardii Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 281, San Francisco.

Genus 589. *PLEURONECTES* Linnaeus. *Plaice.*

Pleuronectes Artedi, Genera Pisc., in part, 1738, 16.

Pleuronectes Linnaeus, Syst. Nat., ed. X, 1758, 268; after Artedi (*P. platessa* Linnaeus).

Platea Cuvier, Règne Anim., ed. I, II, 1817, 220 (*Pleuronectes platessa* L.).

1769. *Pleuronectes quadrituberculatus* Pallas.

Coasts of Alaska and Kamchatka.

Pleuronectes quadrituberculatus Pallas, Zoogr. Rosso.-Asiat., III, 1811, 423, sea between Kamchatka and Alaska.

Pleuronectes pallasi Steindachner, Ichth. Beitr., VIII, 1879, Kamchatka.

Genus 590. *PLATICHTHYS* Girard. *Starry flounders.*

Platichthys Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 139 (*P. rugosus* Girard = *Pleuronectes stellatus* Pallas).

1770. *Platichthys stellatus* (Pallas). *Great flounder; California flounder.*

Pacific coast of America, from Point Conception to the Arctic Ocean, and south to Sakhalin Island.

Pleuronectes stellatus Pallas, Zoogr. Rosso.-Asiat., III, 1811, 416, Kamchatka, Aleutian and Kuril Islands.

Platichthys rugosus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 139, 155, San Francisco, Presidio; Petaluma, Calif.

Genus 591. *INOPSETTA* Jordan and Goss.

Inopsetta Jordan and Goss, Rept. U. S. Fish Com., 1885 (1887), 924 (136 of inside paging) (*Parophrys ischyurus* Jordan and Gilbert).

1771. *Inopsetta ischyra* (Jordan and Gilbert).

Puget Sound, probably northward to Alaska.

Parophrys ischyurus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 276, 453, Puget Sound.

Genus 592. *LEPIDOPSETTA* Gill.

Lepidopsetta Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 330 (*Psetichthys umbrosus* Girard).

1772. *Lepidopsetta bilineata* (Ayres).

Pacific coast of North America, Alaska to Monterey and northern Japan.

Platea bilineata Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 40 (39 of reprint), San Francisco.

Platichthys umbrosus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 136, Puget Sound.

Pleuronectes perarcuatus Cope, Proc. Amer. Phil. Soc. Phila., XIII, 1873, 32, Unalaska.

Genus 593. *ISOPSETTA* Lockington.

Isopetta Lockington, in Jordan and Gilbert, Bull. U. S. Nat. Mus., XVI, 1882, 832 (*Lepidopsetta isolepis* Lockington).

1773. *Isopsetta isolepis* (Lockington).

Puget Sound to Point Conception, in rather deep water.

Lepidopsetta isolepis Lockington, Proc. U. S. Nat. Mus., III, 1880, 325, San Francisco.

Genus 594. *LIMANDA* Gottsche. *Mud dabs.*

Limanda Gottsche, Archiv für Naturgesch., 1835, 100 (*Pleuronectes limanda* Linnaeus).

Myzopsetta Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 217 (*Platea ferruginea* Storer).

1774. **Limanda ferruginea** (Storer).
 Atlantic coast of North America, Labrador to New York.
Platessa ferruginea D. H. Storer, Rept. Fish. Mass., 1839, 141, pl. 2,
 Cape Ann.
Platessa rostrata H. R. Storer, Bost. Jour. Nat. Hist., VI, 1850, 268,
 Labrador.
1775. **Limanda aspera** (Pallas).
 Coasts of Alaska and Kamchatka.
Pleuronectes asper Pallas, Zoogr. Rosso.-Asiat., III, 1811, 425, east coast
 of Siberia.
1776. **Limanda proboscidea** Gilbert.
 Bristol and Herendeen Bays, Alaska.
Limanda proboscidea Gilbert, Rept. U. S. Fish Com., XIX, 1893 (1896),
 460, pl. 33, Bristol and Herendeen Bays, Alaska.
1777. **Limanda beanii** Goode.
 Deep water, off coast of New England.
Limanda beanii Goode, Proc. U. S. Nat. Mus., III, 1880 (Feb. 16, 1881),
 473 southern coast of New England, *Fish Hawk* stations 875, 876.
- Genus 595-596. **PSEUDOPLEURONECTES** Bleeker. *Winter flounders.*
Pseudopleuronectes Bleeker, Versl. Akad. Amsterdam, XIII, 1862, 428
 (*Platessa plana* Mitchell = *Pleuronectes americanus* Walbaum).
1778. **Pseudopleuronectes americanus** (Walbaum). *Common flatfish; Winter
 flounder; Roughback.*
 Atlantic coast of North America, from Labrador to Cape Lookout.
Flounder Schöpf, Schrift. Gesellsch. Naturf. Freunde, VIII, 1788, 148,
 New York.
Pleuronectes americanus Walbaum, Artedi Pisc., 1792, 113; based on
 the *Flounder* of Schöpf.
Pleuronectes planus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 387,
 New York.
Platessa pusilla DeKay, New York Fauna: Fishes, 1842, 296, pl. 47,
 fig. 153, New York.
1779. **Pseudopleuronectes dignabilis** Kendall. *Lemon-yellow flounder.*
 North Atlantic coast, Newfoundland to Cape Cod.
Pseudopleuronectes dignabilis Kendall, Bull. U. S. Fish Com., XXX,
 1910 (1912), 392, Georges Bank.
- Genus 597. **LIOPSETTA** Gill.
Liopsetta Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 216 (*Platessa
 glabra* Storer).
Euchalarodus Gill, loc. cit., 221 (*E. putnami* Gill, the male of *Liopsetta
 glabra* Storer).
1780. **Liopsetta glacialis** (Pallas). *Arctic flounder.*
 Arctic Ocean and Bering Sea south to St. Michaels, Alaska.
Pleuronectes glacialis Pallas, Itin., III, App., 1776, 706; mouth of the
 Obi River.
Pleuronectes cicatricosus Pallas, Zoogr. Rosso.-Asiat., III, 1811, 424,
 sea between Kamchatka and Alaska; male.
? *Platessa dvinensis* Lilljeborg, Veb. Ab. Handl., 1850, 360, Tab. 20,
 mouth of River Dwina.
Pleuronectes franklini Günther, Cat., IV, 1862, 442, Arctic seas of
 America; female.
1781. **Liopsetta putnami** (Gill). *Eel-back flounder.*
 Atlantic coast of North America, from Cape Cod northward to Labrador
 and beyond.
Platessa glabra Storer, Proc. Bost. Soc. Nat. Hist., 1843, 130, Massa-
 chusetts; female; not *Platessa glabra* of Rathke, 1837, a species of
Flesus.
Euchalarodus putnami Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864,
 216, 221, Salem, Mass.

1782. *Liopsetta obscura* (Herzenstein).

Sea of Okhotsk.

Pleuronectes obscurus Herzenstein, Mélanges Biologiques, 1890, 127, Manchuria.

Genus 598. **MICROSTOMUS** Gottsche. *Smear dabs.*

Microstomus Gottsche, Archiv für Naturgesch., 1835, 150 (*M. latidens* Gottsche); not *Microstoma* Risso, 1826.

Cynicoglossus Bonaparte, Fauna Italica, 1837, fasc. XIX (*Pleuronectes cynoglossus* Nilsson); not of Linnæus = *Pleuronectes kitt* Walbaum.

Cynoglossa Bonaparte, Cat. Metod., 1845, 48 (*Pleuronectes cynoglossus* Linnaeus).

Brachyprosopon Bleeker, Versl. Akad. Amsterdam, XIII, 1862, 428 (*Pleuronectes microcephalus* Donovan = *P. kitt* Walbaum).

Cynicoglossus Jordan and Gilbert, Synopsis, 1883, 460 (*Pleuronectes microcephalus* Donovan), *microstomus* being regarded as preoccupied.

1783. *Microstomus kitt* (Walbaum). *Smear dab.*

Arctic Europe, west to the sea between Greenland and Iceland.

Rhombus levius cornubiensis Jago in Ray, "Syn. Pisc.", 1713, 162, Tab. 1, fig. 1."

Smear dab Pennant, British Zoology, III, 1776, 230, pl. 41.

Pleuronectes kitt Walbaum, Arcted Pisc., III, 1792, 120; after Ray.

Pleuronectes quenseli Hölböll, Bohusläns, Sweden.

Pleuronectes quadridens Fabricius, Afhandl., Kongl. Dansk. Vid. Selsk., I, 1824, 39, Iceland.

Microstomus latidens Gottsche, Archiv für Naturgesch., 1835, 150, Zealand.

Pleuronectes gilli Steindachner, Ichth. Notizen, VII, 1868, polar sea, north of Iceland.

1784. *Microstomus pacificus* (Lockington).

Pacific coast of North America, Monterey to Vancouver Island and probably northward.

Glyptocephalus pacificus Lockington, Rept. Calif. Com. Fish., 1878-79, 43, off Point Reyes, Calif.

Genus 599. **EMBASSICHTHYS** Jordan and Evermann.

Embassichthys Jordan and Evermann, Check List, 1895 (1896), 506 (*Cynicoglossus bathybius* Gilbert).

1785. *Embassichthys bathybius* (Gilbert).

Coast of California, south of Point Conception.

Cynicoglossus bathybius Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 123, Santa Barbara Channel at Albatross station 2980, lat. 33° 49' 45" N., long. 119° 24' 30" W., in 603 fathoms.

Genus 600. **GLYPTOCEPHALUS** Gottsche.

Glyptocephalus Gottsche, Archiv für Naturgesch., 1835, 156 (*Pleuronectes saxicola* Fabricius = *P. cynoglossus* Linnaeus).

1786. *Glyptocephalus cynoglossus* (Linnaeus). *Craig fluke; Pole flounder; Gray sole; Lemon sole.*

North Atlantic, chiefly in deep water, south to Cape Cod and France.

Pleuronectes cynoglossus Linnaeus, Syst. Nat., ed. X, 1758, 269; after Gronow.

Pleuronectes saxicola Faber, Tidsskr. für Naturv., 5 B., 1828, 244, Denmark.

Glyptocephalus acadianus Gill, Proc. Acad. Nat. Sci. Phila., XXV, 1873, 360, Nova Scotia.

Genus 601. **ERREX** Jordan.

Errex Jordan, Proc. Acad. Nat. Sci. Phila., LXXI, 1919, 343 (*Glyptocephalus zachirus* Lockington).

1787. *Errex zachirus* (Lockington). *Rex sole; Long-finned sole.*

Waters of moderate depth of the northern Pacific; from San Francisco northward.

Glyptocephalus zachirus Lockington, Proc. U. S. Nat. Mus., II, 1879, 88, San Francisco.

Family 138. ACHIRIDÆ. Broad soles

Subfamily GYMNACHIRINÆ

Genus 602. GYMNACHIRUS Kaup.

Gymnachirus Kaup, Wiegmann's Archiv, I, 1858, 101 (*Gymnachirus nudus* Kaup).

1788. *Gymnachirus nudus* Kaup.

Coast of Brazil; once taken at Tisbury Great Pond, Martha's Vineyard Island, Mass. (Vinal N. Edwards).

Gymnachirus nudus Kaup, Wiegmann's Archiv, I, 1858, 101, Brazil

1789. *Gymnachirus fasciatus* Günther.

West Indies; very rare, north to Long Key, Fla. (W. C. Kendall).

Gymnachirus fasciatus Günther, Cat., IV, 1862, 488, locality unknown.

Subfamily ACHARINÆ

Genus 603. ACHIRUS Lacépède.

Achirus Lacépède, Hist. Nat. Poiss., IV, 1803, 658 (*Pleuronectes achirus* Linnæus).

Trinectes Rafinesque, Atlantic Jour. and Friend of Knowledge, I, 1832 (*T. scabra* Rafinesque = *Pleuronectes mollis* Mitchell).

Grammichthys Kaup, Arch. Naturgesch., 1858, 94 (*Achirus lineatus* Linnæus).

Baiostoma Bean, Proc. U. S. Nat. Mus., V, 1882, 413 (*B. brachialis* Bean).

Subgenus ACHIRUS Lacépède.

1790. *Achirus achirus* (Linnæus).

Atlantic coasts of tropical America.

Pleuronectes oculis dextris corpore glabro, pinnis pectoralibus nullus Gronow, Museum, I, No. 42, Surinam.

Pleuronectes achirus Linnæus, Syst. Nat., ed. X, 1758, 268, Surinam; based on Gronow.

Solea gronovii Günther, Cat., IV, 1862, 472, Surinam.

1791. *Achirus inscriptus* Gosse.

West Indies, north to Key West.

Achirus inscriptus Gosse, Nat. Sojourn Jamaica, 1851, 52, pl. 1, fig. 4, Jamaica.

Monochir reticulatus Poey, Memorias, II, 1861, 317, Cuba.

Subgenus GRAMMICHTHYS Kaup.

1792. *Achirus klunzingeri* (Steindachner).

Pacific coast of tropical America, Panama to Guayaquil.

Solea klunzingeri Steindachner, Zur Fische des Cauca und der Flüsse bei Guayaquil, 1879, 44, Guayaquil.

Subgenus TRINECTES Rafinesque.

1793. *Achirus fasciatus* Lacépède. *American sole; Hogchoker.*

Atlantic coast of United States, from Cape Cod to Texas; often ascending streams.

Achirus fasciatus Lacépède, Hist. Nat. Poiss., IV, 1803, 659, Charleston, S. C.

Pleuronectes mollis Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 388, pl. 2, fig. 4, New York.

Pleuronectes apoda Mitchell, Amer. Month. Mag. and Crit. Rev., Feb., 1818, 244, Straits of Bahama.

Trinectes scabra Rafinesque, Atlantic Jour. and Friend of Knowledge, I, 1832, Pennsylvania, in fresh water.

Solea browni Günther, Cat., IV, 1862, 477, New Orleans; Texas.

1794. *Achirus scutum* (Günther).

Pacific coast of Central America.

Solea scutum Günther, Cat., IV, 1862, 474, Gulf of Fonseca, Panama.

- 1795. *Achirus fonsecensis* (Günther).**
 Pacific coast of tropical America, Mazatlan to Fonseca.
Solea fonsecensis Günther, Cat., IV, 1862, 475, Gulf of Fonseca.
Solea panamensis Steindachner, Ichth. Beiträge, V, 1876, 10, Taf. II,
 Panama.
Solea fischeri Steindachner, Beiträge Kenntniss Fluss-Fische Sudamer.,
 I, 1879, 13, Rio Mamone near Panama.
- 1796. *Achirus mazatlanus* (Steindachner).** *Mexican sole; Teipalcate.*
 Pacific coast of tropical America.
Solea mazatlana Steindachner, Ichth. Notizen, IX, 1869, 23, Mazatlan,
 Mexico.
Solea pilosa Peters, Berliner Monatsber., 1869, 709, Mazatlan, Mexico.
- 1797. *Achirus comifer* Jordan and Gilbert.**
 Key West, Fla.
Achirus comifer Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884,
 31, Key West, Fla.
- 1798. *Achirus lineatus* (Linnæus).**
 Florida to Brazil.
Passer lineis transversis notatus Sloane, Jamaica, 2, 1725, 77, pl. 246,
 fig. 2, Jamaica.
Pleuronectes fuscus subrotundus glaber Browne, Jamaica, 1756, 445,
 Jamaica.
Pleuronectes lineatus Linnæus, Syst. Nat., ed. X, 1758, 268, Jamaica;
 based on Browne and Sloane; not of ed. XII, which is *Achirus
 fasciatus*.
Monochir maculipinnis Agassiz, Spix, Pisc. Brasil., 1829, 88, pl. 49,
 Brazil.
Baistoma brachialis Bean, Proc., U. S. Nat. Mus., V, 1882, 413, Appa-
 lachicola Bay and south Florida.
- 1799. *Achirus barnharti* Jordan.**
 Coast of southern California; Coronado Islands.
Achirus barnharti Jordan, Univ. Calif. Pub. Zool., XXVI, 1923, 10,
 Pl. I, Coronado Islands in about 400 feet depth of water.
- Genus 604. APIONICHTHYS Kaup.**
Apionichthys Kaup, Wiegmann's Archiv, 1858, 104 (*A. dumerili* Kaup).
Soleotalpa Günther, Cat., IV, 1862, 489 (*S. unicolor* Günther).
- 1800. *Apionichthys dumerili* Kaup.**
 West Indies.
Apionichthys dumerili Kaup, Wiegmann's Archiv, 1858, 104, no locality;
 description of species included in that of the genus.
Soleotalpa unicolor Günther, Cat., IV, 1862, 489, West Indies.
Apionichthys nebulosus Peters, Berliner Monatsb., 1869, 709, Surinam.
- Family 139. CYNOGLOSSIDÆ. Tonguefishes**
- Genus 605. SYMPHURUS Rafinesque.** *Tonguefishes.*
Sympodus Rafinesque, Indice, 1810, 13, 52 (*S. nigrescens* Rafinesque).
Bibronia Cocco, Alcuni Pesci del Mare di Messina, 1844, 15 (*B. ligulata*
 Cocco, larval form of *Sympodus*).
Plagusia Cuvier, Règne Animal, ed. 2, II, 1829, 344; based on *Plagusia*
 of Browne; name preoccupied in crustaceans, Latreille, 1806.
Plagiusa Bonaparte, Cat. Metod., 1846, 51 (*P. lactea*); name a sub-
 stitute for *Plagusia*.
Aphoristia Kaup, Archiv für Natur., 1858, 106 (*Achirus ornatus*
 Lacépède).
Glossichthys Gill, Cat. Fish. E. Coast N. Amer., 1861, 51 (*Pleuronectes
 plagusia* Linnæus).
Ammopleurops Günther, Cat., IV, 1862, 490 (*Plagusia lactea* Bonaparte).
?Bacanius Schiödte, Naturhist. Tijdskr., V, 1867, 269 (*B. tædifer*
 Schiödte; larval).
- 1801. *Sympodus piger* (Goode and Bean).**
 West Indies and Gulf of Mexico in deep water.
Aphoristia pigra Goode and Bean, Bull. Mus. Comp. Zool., XII, No. 5,
 1886, 154, St. Kitts, in about 250 fathoms.

1802. *Syphurus marginatus* (Goode and Bean.)
West Indies.
Aporistia marginata Goode and Bean, Bull. Mus. Comp. Zool., XII, No. 5, 1886, 153, off St. Vincent, West Indies.
1803. *Syphurus atramentatus* Jordan and Bollman.
Off coast of Colombia.
Syphurus atramentatus Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 177, off coast of Colombia at *Albatross* station 2795, lat. $7^{\circ} 57' N.$, long. $78^{\circ} 55' W.$, in 33 fathoms.
1804. *Syphurus fasciolaris* Gilbert.
Gulf of California.
Syphurus fasciolaris Gilbert, Proc. U. S. Nat. Mus., XIII, 1891, 506, Gulf of California.
1805. *Syphurus elongatus* (Günther).
Pacific coast of tropical America.
Aporistia ornata var. *elongata* Günther, Fishes Cent. Amer., 1869, 473, Panama.
1806. *Syphurus atricaudus* (Jordan and Gilbert). *San Diego sole*.
Lower California, north to San Diego.
Aporistia atricauda Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 23, San Diego, Calif.
1807. *Syphurus leei* Jordan and Bollman.
Off the coast of Colombia.
Syphurus leei Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 178, off coast of Colombia at *Albatross* station 2804, lat. $8^{\circ} 16' 30'' N.$, long. $79^{\circ} 37' 45'' W.$, in 47 fathoms.
1808. *Syphurus plagusia* (Bloch and Schneider). *Acedia*.
West Indies, south to Rio Janeiro.
Plagusia Browne, Jamaica, No. 1, 1756, 445, Jamaica.
Pleuronectes plagusia Bloch and Schneider, Syst. Ichth., 1801, 162, Jamaica; after Browne.
Achirus ornatus Lacépède, Hist. Nat. Poiss., IV, 1803, 659; on a specimen "presented by Holland to France."
Plagusia tessellata Quoy and Gaimard, Voyage *Uranie*, Zoology, 1824, 240, Rio Janeiro.
Plagusia brasiliensis Agassiz, in Spix, Pisc. Brasil., 1827, 89, tab. 50, Brazil.
1809. *Syphurus plagiura* (Linnæus). *Tonguefish*.
South Atlantic and Gulf coasts of United States.
Pleuronectes plagiura Linnæus, Syst. Nat., ed. XII, 1766, 455, on a specimen from Doctor Garden, probably from Charleston.
Plagusia fasciata Holbrook, in DeKay, New York Fauna: Fishes, 1842, 304, Charleston, S. C.
1810. *Syphurus pusillus* (Goode and Bean).
Off Atlantic coast of United States in deep water.
Aporistia pusilla Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 590, Gulf Stream, lat. $40^{\circ} N.$, in deep water.
1811. *Syphurus diomedeanus* (Goode and Bean).
Gulf of Mexico, off Key West, Fla.
Aporistia diomediana Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 589, Gulf of Mexico, lat. $25^{\circ} 4' 30'' N.$, long. $82^{\circ} 59' 15'' W.$
1812. *Syphurus williamsi* Jordan and Culver.
West coast of Mexico.
Syphurus williamsi Jordan and Culver, in Jordan, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 506, pl. 55, Mazatlan, Mexico.
- Genus 608. ACEDIA Jordan.
Acedia Jordan, in Jordan and Goss, Rept. U. S. Fish Comm., XIV, 1886 (1888) (*Aporistia nebulosa* Goode and Bean).
1813. *Acedia nebulosa* (Goode and Bean).
Gulf Stream.
Aporistia nebulosa Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1883, 192, Gulf Stream, off the coast of the Carolinas.

Superorder ACANTHOPTERYGII

Order LL. ZEOIDEI

Family 140. ZEIDÆ. John dories

Genus 607. ZENOPSIS Gill.

Zenopsis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 126 (*Zeus nebulosus* Temminck and Schlegel).

1814. *Zenopsis ocellata* (Storer). Ocellated dory.

Off New England coast.

Zeus ocellatus Storer, Proc. Bost. Soc. Nat. Hist., VI, 1858, 386, off Provincetown, Mass.

Genus 608. ZENION Jordan and Evermann.

Zenion Jordan and Evermann, Rept. U. S. Fish Com., XXI, 1895 (1896), 418 (*Cyttus hololepis* Goode and Bean).

1815. *Zenion hololepis* (Goode and Bean).

West Indies, in deep water.

Cyttus hololepis Goode and Bean, Oceanic Ichth., 1896, 225, figs. 233, 233a, 233b, off Yucatan and Little Bahama Bank.

Genus 609. OREOSOMA Cuvier and Valenciennes.

Oreosoma Cuvier, Règne Animal, ed. 2, II, 1829 (type not named, *Oresoma atlanticum* Cuvier).

Oreosoma Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1830, 377 (515) (*O. atlanticum* Cuvier and Valenciennes).

1816. *Oreosoma atlanticum* Cuvier and Valenciennes.

Open Atlantic; only a few specimens known.

Oreosoma atlanticum Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 377 (515), open Atlantic.

Oreosoma conferrum Cuvier and Valenciennes, loc. cit., pl. 99, same type.

Family 141. GRAMMICOLEPIDÆ

Genus 610. GRAMMICOLEPIS Poey.

Grammicolepis Poey, Anal. Soc. Esp. Hist. Nat., II, 1873 (*G. brachiusculus* Poey).

1817. *Grammicolepis brachiusculus* Poey.

Cuba; in deep water.

Grammicolepis brachiusculus Poey, Anal. Soc. Española Hist. Nat., II, 1873, Cuba.

Order MM. XENOBERYCES

Family 142. STEPHANOBERYCIDÆ

Genus 611. STEPHANOBERYX Gill.

Stephanoberyx Gill, Proc. U. S. Nat. Mus., VI, 1883, 258 (*S. monæ* Gill).

1818. *Stephanoberyx monæ* Gill.

Gulf Stream, in deep water.

Stephanoberyx monæ Gill, Proc. U. S. Nat. Mus., VI, 1883, 258, Gulf Stream at Albatross station 2077 in 1,255 fathoms.

1819. *Stephanoberyx gillii* Goode and Bean.

Gulf Stream, in deep water.

Stephanoberyx gillii Goode and Bean, Oceanic Ichth., 1895, 187, fig. 206, Gulf Stream at Albatross station 2099 in 2,949 fathoms.

Family 143. MELAMPHAIDÆ

Genus 612. CAULOLEPIS Gill.

Caulolepis Gill, Forest and Stream, XXI, Aug. 30, 1883, and Proc. U. S. Nat. Mus., VI, 1883, 258 (*C. longidens* Gill).

- 1820. Caulolepis longidens** Gill.
Gulf Stream and Pacific Ocean at Cortez Banks off San Diego, Calif.
Caulolepis longidens Gill, Proc. U. S. Nat. Mus., VI, 1883, 258, Atlantic
Ocean, lat. 39° 27' N., long. 69° 56' 20" W., in 1,346 fathoms.
- Genus 613. ANOPLOGASTER** Günther.
Anoplogaster Günther, Cat., I, 1859, 12 (*Hoplostethus cornutus* Cuvier
and Valenciennes).
- 1821. Anoplogaster cornutus** (Cuvier and Valenciennes).
Atlantic, in deep water, north and south.
Hoplostethus cornutus Cuvier and Valenciennes, Hist. Nat. Poiss.,
IX, 1833, 347 (270); from stomach of a shark, taken at 26° lat. S;
others taken by the *Albatross* 31° N., 40° W., from the stomach of
an albacore.
- Genus 614. POROMITRA** Goode and Bean.
Poromitra Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1882,
214 (*P. capito* Goode and Bean).
- 1822. Poromitra capito** Goode and Bean.
Deep water in Gulf Stream in latitude 34°.
Poromitra capito Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5,
1882, 215, Gulf Stream.
- Genus 615. PLECTROMUS** Gill.
Plectromus Gill, Proc. U. S. Nat. Mus., VI, 1883, 257 (*P. suborbitalis*
Gill).
- 1823. Plectromus suborbitalis** Gill.
Gulf Stream, in deep water.
Plectromus suborbitalis Gill, Proc. U. S. Nat. Mus., VI, 1883, 258, lat.
38° 52' N., long. 69° 24' W., at *Albatross* stations 2036, 2190, and
2535 in 1,149 to 1,800 fathoms.
- 1824. Plectromus beani** (Günther).
Gulf Stream in deep water.
Plectromus crassiceps Bean, Proc. U. S. Nat. Mus., VIII, 1885, 73, Gulf
Stream; name preoccupied.
Melamphaës beani Günther, Challenger Rept., XXII, 1887, 29, Gulf
Stream, in about latitude 40°.
- 1825. Plectromus crassiceps** (Günther).
Mid-Atlantic; Brazil.
Scopelus crassiceps Günther, Ann. Mag. Nat. Hist., II, 1878, 185,
deep water in mid-Atlantic and off Pernambuco.
- 1826. Plectromus cristiceps** (Gilbert).
Coast of Washington and Oregon and southeastern Alaska.
Melamphaës cristiceps Gilbert, Proc. U. S. Nat. Mus., XIII, 1890
(1891), 60, coasts of Oregon and Washington and *Albatross* station
3075 in 859 fathoms.
- 1827. Plectromus lugubris** (Gilbert).
Coast of California; off Bay of Panama; Bering Sea.
Melamphaës lugubris Gilbert, Proc. U. S. Nat. Mus., XIII, 1890 (1891),
59, coast of California south of Point Conception, at *Albatross* station
2923 in 822 fathoms.
- 1828. Plectromus mizolepis** (Günther).
Off Bay of Panama in deep water; south of New Guinea and Bay of
Bengal.
Scopelus mizolepis Günther, Ann. Mag. Nat. Hist., 11, 1878, 185, south
of New Guinea.
- 1829. Plectromus nycterinus** (Gilbert).
Coast of southern California, in deep water.
Melamphaës nycterinus Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915,
326, pl. 16, off Santa Catalina, 2,113 to 2,259 fathoms.
- 1830. Plectromus bispinosus** (Gilbert).
Coronado Islands, near San Diego, Calif.
Melamphaës bispinosus Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915,
325, pl. 15, fig. 5; off the Coronado Islands; near San Diego, 642 to
666 fathoms.

Order NN. BERYCOIDEI

Family 144. POLYMXIIDÆ. Barbudos

Genus 616. POLYMXIA Lowe.

Polymixia Lowe, Trans., Cambr. Phil. Soc., VI, 1838, 198 (*P. nobilis* Lowe).
Nemabrama Valenciennes, Ichth. îles Canaries, 1844, 40 (*N. webbi* Valenciennes = *Polymixia nobilis* Lowe).
Dinemus Poey, Memorias, II, 1861, 160 (*D. venustus* Poey = *Polymixia lowei* Günther).

1831. *Polymixia lowei* Günther. *Barbudo*.

Caribbean Sea; Cuba.

Polymixia lowei Günther, Cat., I, 1859, 17, Caribbean Sea.

Dinemus venustus Poey, Memorias, II, 1861, 161, Cuba.

Family 145. BERYCIDÆ

Genus 617. BERYX Cuvier. Alfonsinos.

Beryx Cuvier, Règne Anim., ed. 2, II, 1829, 151 (*B. decadactylus* Cuvier and Valenciennes).

1832. *Beryx decadactylus* Cuvier and Valenciennes. *Alfonsin a Casta Larga*. Cosmopolitan, open seas, in rather deep water; Portugal, Madeira, Japan, and Cuba.

Beryx decadactylus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 165 (222), Madeira or Portugal.

1833. *Beryx splendens* Lowe. *Alfonsin a Casta; Cumprida*.

Cosmopolitan, in the open seas, at moderate depth; Madeira, Japan, and Gulf Stream at 35° 49' 30" N., 74° 34' 45" W. in 424 fathoms.
Beryx splendens Lowe, Proc. Zool. Soc. Lond., 1833, 142, Madeira.

Family 146. ANOMALOPIDÆ

Genus 618. KRYPTOPHANARON Silvester and Fowler.

Kryptophanaron Silvester and Fowler, Proc. Acad. Nat. Sci. Phila., LXXVIII, 1926, 245 (*K. alfredi* Silvester and Fowler).

1834. *Kryptophanaron alfredi* Silvester and Fowler.

Off coast of Jamaica.

Kryptophanaron alfredi Silvester and Fowler, Proc. Acad. Nat. Sci. Phila., LXXVIII, 1926, 246, Pl. XVIII, fig. 1; Pl. XIX, figs. 2 and 4. Off Kingston Harbor, Jamaica, floating on surface.

Family 147. TRACHICHTHYIDÆ

Genus 619. HOPLOSTETHUS Cuvier and Valenciennes.

Hoplostethus Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 344 (469) (*H. mediterraneus* Cuvier and Valenciennes).

1835. *Hoplostethus mediterraneus* Cuvier and Valenciennes.

Coasts of southern Europe; Gulf Stream.

Hoplostethys mediterraneus Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 344 (469), Mediterranean Sea.

Trachichthys pretiosus Lowe, Proc. Zool. Soc. Lond. 1839, 77, Madeira.

Family 148. HOLOCENTRIDÆ. Soldierfishes; Squirrelfishes

Genus 620. OSTICHTHYS Jordan and Evermann.

Ostichthys Langsdorf, in Cuvier and Valenciennes, Hist. Nat. Poiss., III, 130 (174), 1829; name only, passing reference under *Myripristis japonicus* Cuvier.

Ostichthys Jordan and Evermann, Fish. North and Mid. Amer., 1896, 846 (*Myripristis japonicus* Cuvier and Valenciennes).

1836. *Ostichthys trachypomus* (Günther).

West Indies; Cuba.

Myripristis trachypoma Günther, Cat., I, 1859, 25, Cuba.

Myripristis fulgens Poey, Memorias, II, 1860, 160, Cuba.

Genus 621. MYRIPRISTIS Cuvier.

Myripristis Cuvier, Règne Anim., ed. 2, II, 1829, 150 (*M. jacobus* Cuvier and Valenciennes).

Rhamphoberyx Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 817 (*R. paecilopus* Gill); young.

Myripristes of authors; altered spelling.

Myriopristis of authors; altered spelling.

- 1837. Myripristis jacobus** Cuvier and Valenciennes. *Frère Jacques; Candil.* West Indies to Brazil; common.
Myripristis jacobus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 121 (162), Martinique.
Rhinoberyx chryseseus Cope, Proc. Amer. Phil. Soc., XIV, 1871, 464, St. Croix Island.

- 1838. Myripristis occidentalis** (Gill).
West coast of America, from Cape San Lucas to Panama; Galapagos; Cocos Island.
Myripristis occidentalis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 87, Cape San Lucas.
Rhamphoberyx leucopus Gill, loc. cit., 88, Cape San Lucas; immature.

- 1839. Myripristis paecilopus** (Gill).
West coast of America, from Cape San Lucas to Panama.
Rhamphoberyx paecilopus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 87, Cape San Lucas; immature example.

Genus 622. HOLOTRACHYS Günther.

Holotrichys Günther, Jour. Mus. Godeffroy, Band I, 1873, 93 (*Myripristes lima* Cuvier and Valenciennes).

- 1840. Holotrachys lima** (Cuvier and Valenciennes).
South Seas, east to Hawaii and Cocos Island.
Myripristes lima Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 493, île de France.
Myripristes humilis Kner and Steindachner, Sitz. Akad. Wiss. Wien, LIV, 1866, 357, pl. 1, fig. 1, Samoa.

Genus 623. HOLOCENTRUS Bloch. *Squirrelfishes; Soldierfishes.*

Holocentrus Gronow, Zoophyl., 1763, 65; nonbinomial.

Holocentrus Bloch, Naturgeschich. Ausländ. Fische, IV, 1790, 61 (*H. sogo* Bloch).

Holocentrum Cuvier, and of most authors; amended spelling.

Rhynchichthys Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 503 (*R. pelamidis* Cuvier and Valenciennes); young.
Rhinoberyx Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 237 (*Rhynchichthys brachyrhynchus* Bleeker); larval form.

- 1841. Holocentrus ascensionis** ¹¹ (Osbeck). *Matajuelo; Squirrelfish; Soldado; Welchman.*

Islands of the Atlantic and northward from St. Helena and Ascension Island to Florida and Cuba.

?*Percus ascensionis* Osbeck, Iter Chinensis, 1757; reprinted as Reise Ost-Ind. China, 1765, 388, Ascension Island; reprint of a pre-Linnæan paper, hence ineligible.

?*Bodianus pentacanthus* Bloch, Nat. Ausl. Fische, IV, 1790, 40, pl. 225, Brazil; after *Saguaraca* of Margrave.

?*Holocentrus sogo* Bloch, loc. cit., 61, pl. 232, Africa.

Sciaena rubra Schneider, Ed. Bloch, Syst. Ichth., 1801, 82; after Catesby.

Amphiprion matejuelo Bloch and Schneider, Syst. Ichth., 1801, 206, Cuba; after Parra.

?*Bodianus jaguar* Lacépède, Hist. Nat. Poiss., IV, 1802, 286, Brazil.

?*Holocentrum longipinne* Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 138 (185), Brazil; Martinique; Porto Rico; St. Thomas; San Domingo; Habana.

Holocentrus striatus Gronow, Cat. Fishes, 1854, 173, Antilles.

Holocentrus rostratus Gronow, loc. cit., 173, near the Equator.

¹¹ The specific name of the common mataguelo of Cuba is still wholly uncertain if the two forms (the one from Cuba, the other from Brazil) are really distinct. The earliest specific name, *ascensionis*, is apparently ineligible, occurring in reprints of a pre-Linnæan paper. The precedence of most of the other names is more or less uncertain. Here, as elsewhere where no official decision has been made by the International Commission on Zoological Nomenclature, we follow the rule of not changing current nomenclature unless, as in matters of priority, the case is reasonably clear.

1842. *Holocentrus rufus* (Walbaum).

The northern form probably varying into *H. ascensionis*, which ranges from the Bahamas to Brazil.

Perca rufa Walbaum, Artedi Pisc., 1792, 351, Bahamas; after *Perca marina rubra* of Catesby.

1843. *Holocentrus coruscus* (Poey).

Cuba; Bahama Islands; Bermudas; Tortugas.

Holocentrum coruscum Poey, Memorias, II, 1861, 158, Cuba.

Holocentrum sicciferum Cope, Trans. Amer. Phil. Soc., XIV, 1871, 465, Bahamas, New Providence.

?*Holocentrum puncticulatus* Barbour, Bull. Mus. Comp. Zool., XLVI, 1905, 117, Bermudas.

?*Holocentrus tortugae* Jordan and Thompson, Bull. U. S. Bur. Fish., XXIV, 1904 (1905), 236, fig. 1, Garden Key, Tortugas.

1844. *Holocentrus brachypterus* Poey.

Cuba.

Holocentrus brachypterus Poey, Repertorio, I, 1866, 184, Cuba.

1845. *Holocentrus vexillarius* (Poey).

Cuba; Porto Rico.

Holocentrum vexillarium Poey, Memorias, II, 1861, 158, Cuba.

?*Holocentrum riparium* Poey, Enumeratio, 1875, 37, Cuba.

1846. *Holocentrus osculus* (Poey).

Cuba.

Holocentrum osculum Poey, Memorias, II, 1861, 156, Cuba.

Holocentrum perlatum Poey, loc. cit., 157, Cuba.

1847. *Holocentrus sancti-pauli* (Günther).

St. Paul Rocks, mid-Atlantic.

Holocentrum sancti-pauli Günther, Shore Fishes, Challenger Rept., Zoology, I, part 6, 1880, 4, St. Paul Rocks, mid-Atlantic.

1848. *Holocentrus meeki* Bean.

Bermudas.

Holocentrus meeki Bean, Proc. Biol. Soc. Wash., XIX, 1906, 31, St. Davids Island, Bermudas.

Genus 624. *FLAMMEO* Jordan and Evermann.

Flammeo Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2871 (*Holocentrum Marianum* Cuvier and Valenciennes).

1849. *Flammeo Marianus* (Cuvier and Valenciennes). *Marian*.

West Indies; Cuba; Jamaica; Martinique.

Holocentrum Marianum Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 164 (219), Martinique.

Holocentrum rostratum Poey, Memorias, II, 1861, 157, Cuba.

Genus 625. *ADIORYX* Starks.

Adioryx Starks, Science, XXVIII, 1908, 614 (*Holocentrum suborbitalis* Gill).

1850. *Adioryx suborbitalis* (Gill). *Mojarra cardinalis*.

Mazatlan to Panama; Revillagigedo, Cocos, and Galapagos Islands.

Holocentrum suborbitalis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 86, Cape San Lucas, Lower California.

Genus 626. *PLECTRYPOPS* Gill.

Plectrypops Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 237 (*Holocentrum retrospinis* Guichenot).

1851. *Plectrypops retrospinis* (Guichenot).

Cuba.

Holocentrum retrospinis Guichenot, in Ramon de la Sagra, Hist. Cuba, 1853, 35, pl. 1, fig. 3, Cuba.

Order OO. THORACOSTEI

Suborder HEMIBRANCHII

Family 149. GASTEROSTEIDÆ. Sticklebacks

Genus 627. GASTEROSTEUS Linnæus. *Sticklebacks.*

Gasterosteus Linnæus, Syst. Nat., ed. X, 1758, 295 (*G. aculeatus* Linnæus).
Gasteracanthus Pallas, Zoog. Rosso.-Asiat., III, 1811 (*G. cataphractus* Pallas).

Leiurus Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 242 (*Gasterosteus aculeatus* Linnæus).

Merriamella Jordan, Univ. Calif. Publ. Geol., V, 1907, 131 (*M. dorysia* Jordan); fossil form.

1852. *Gasterosteus aculeatus* Linnæus. *Common stickleback.*

Arctic and subarctic regions, in shallow bays and estuaries, very common; also ascending streams, where it loses partially or wholly its bony plates, in Europe, Asia, and America alike. The development of these plates and in some degree of dorsal spines and caudal keel depends on the salinity of the water. Reported *G. williamsoni microcephalus* from irrigation ditch near El Rosario, Lower California, 255 miles south of U. S. border, May 21, 1925, Clinton G. Abbott, collector.

(1. *Forms completely mailed*)

Gasterosteus aculeatus Linnæus, Syst. Nat., ed. X, 1758, 489, Europe.

Gasterosteus cataphractus Pallas, Mém. Acad. Petersb., III, 1811, 325, Kamchatka.

Gasterosteus trachurus Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 481, France.

Gasterosteus teraculeatus Lacépède, Hist. Nat. Poiss., III, 1802, 296.

Gasterosteus obolarius Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 500, Kamchatka.

Gasterosteus noveboracensis Cuvier and Valenciennes, loc. cit., 502, pl. 98, fig. 3.

Gasterosteus biaculeatus Shaw, Zool., IV, 1839, 608; after Pennant.

Gasterosteus dimidiatus Reinhardt, Fauna Grænlandica, 1837, 32, 37, Greenland.

Gasterosteus loricatus Reinhardt, loc. cit., 32, 37, Greenland.

Gasterosteus bailloui Blanchard, Poiss. des eaux douces de France, 1866, 231, Abbeville.

Gasterosteus insculptus Richardson, Last Arctic Voyage, 1854, 10, Pl. XXV, figs. 1, 2, and 3, Northumberland and Puget Sounds.

Gasterosteus serratus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 47, San Francisco.

(2. *Forms armed anteriorly only*)

Gasterosteus gymnurus Cuvier, Règne Animal, ed. II, 1829, 170, waters of England and northward.

Gasterosteus leius Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 481, Seine.

Gasterosteus semiarmatus Cuvier and Valenciennes, loc. cit., 493, Havre, France.

Gasterosteus semiloricatus Cuvier and Valenciennes, loc. cit., 494, Baillon.

Gasterosteus spinulosus Jenyns and Yarrell, Man. Brit. Vert., 1835, 350, Edinburgh.

Gasterosteus ponticus Nordmann, in Demidoff, Voy. Russ. Merid., III, 1840, 357, Tauria and Black Sea.

Gasterosteus cuvieri Girard, in Storer's Fishes of Nova Scotia and Labrador, 1849, 254, p. VII, fig. 1, Bras d'Or, Red Bay, Labrador.

Gasterosteus microcephalus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 133, Four Creek (Kaweah River), a tributary to Tule Lake (Tulare Lake), San Joaquin Valley, Calif.

Gasterosteus plebeius Girard, loc. cit., 147, Presidio, Calif.

- Gasterosteus inopinatus* Girard, loc. cit., 147, Presidio, Calif.
Gasterosteus pugetti Girard, loc. cit., 1856, 135, Fort Steilacoom, Wash.
Gasterosteus neustriensis Blanchard, Poiss. des eaux douce de France, 1866, 220, Harfleur.
Gasterosteus argentissimus Blanchard, loc. cit., 232, Avignon.
Gasterosteus elegans Blanchard, loc. cit., 234, Cadillac and Langou, 6 pls.
Gasterosteus islandicus Sauvage, Revision des Épinoches, 1874, 20, Iceland.
Gasterosteus atkinsi Bean, Proc. U. S. Nat. Mus., II, 1879, 67, Schoodic Lakes, Maine.

(3. Forms naked or with one or two plates)

- Gasterosteus williamsoni* Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 133, Williamson's Pass, Calif.
Gasterosteus santa-annæ Regan, Ann. Mag. Nat. Hist., 8th ser., IV, 1909, 437, Santa Anna River, Calif.

Genus 628. **GLADIUNCULUS** Jordan and Evermann.

Gladiunculus Jordan and Evermann, Proc. Calif. Acad. Sci., ser. 4, XVI, 1927 (Apr. 27), 504 (*Gasterosteus gladiunculus* Kendall = *G. bispinosus* Walbaum).

1853. **Gladiunculus bispinosus** (Walbaum). *Common eastern stickleback.*
 Labrador to New Jersey, near the coast, in salt water; very common in New England.
Gasterosteus bispinosus Walbaum, Artedi Pisc., 1792, 450; after Pennant; said to be "New York," but more likely Hudson's Bay.
Gasterosteus wheatlandi Putnam, Proc., Essex Inst., V, 1867, 4, Nahant, Mass.
Gasterosteus gladiunculus Kendall, Proc. U. S. Nat. Mus., XVIII, 1896, off coast of Maine, near Seguin Island.

Genus 629. **EUCALIA** Jordan.

Eucalia Jordan, Man. Vert., ed. 1, 1876, 248 (*Gasterosteus inconstans* Kirtland).

1854. **Eucalia incostans** (Kirtland). *Brook stickleback.*
 New York to Kansas and northward to the Saskatchewan; Great Lakes region; central Ohio, Indiana, and Illinois.
Gasterosteus inconstans Kirtland, Bost. Jour. Nat. Hist., III, 1841, 273, brooks of Trumball County, Ohio.
Gasterosteus micropus Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 81, Fort Riley, Kans.
Gasterosteus globiceps Sauvage, Revision des Épinoches, 1874, 35, North America.
Eucalia inconstans cayuga Jordan, Man. Vert., ed. 1, 1876, 249, Cayuga Lake, Ithaca, N. Y.

1855. **Eucalia pygmæa** (Agassiz).

Lake Superior.
 A valid species if the fin rays are D. IV-I, 6; A. I, 6. In *E. inconstans* they are D. IV-I, 10; A. I, 10.
Gasterosteus pygmæus Agassiz, Lake Superior, 1850, 314, Lake Superior.

Subfamily GASTEROSTEINÆ

Genus 630. **PUNGITIUS** Costa. *Nine-spined sticklebacks.*

Pungitius Costa, Mém. Savants Etrangers, Paris, X, 1846 (*Gasterosteus pungitius* L.).

Pygosteus (Brevoort) Gill, Can. Nat., and Cat. Fish. E. Coast N. A., supplement to Proc. Acad. Nat. Sci. Phila., XIII, 1861, 39 (*Gasterosteus occidentalis* Cuvier and Valenciennes = *G. pungitius* Linnaeus).

Gasterosteus Sauvage, Révision des Épinoches, 1874, 7 (*Gasterosteus pungitius* Linnaeus).

- 1856. *Pungitius pungitius* (Linnæus).** *Nine-spined stickleback.*
Gasterosteus pungitius Linnæus, Syst. Nat., ed. X, 1758, 296, Europe;
 after *Gasterosteus aculeis* in dorso decem Arctedi.
Gasterosteus lavis Cuvier, Règne Animal, ed. 2, II, 1829, 170, streams of
 France.
Gasterosteus occidentalis Cuvier and Valenciennes, Hist. Nat. Poiss.,
 IV, 1829, 509, Newfoundland; not *G. occidentalis* Linnæus, unrec-
 ognized, probably a species of *Carangidae*.
Gasterosteus concinnus Richardson, Fauna Bor.-Amer., III, 1836, 57,
 Saskatchewan River and Great Bear Lake.
Gasterosteus nebulosus Agassiz, Lake Superior, 1850, 310, pl. 4, fig. 4,
 Lake Superior.
Gasterosteus dekayi Agassiz, Lake Superior, 1850, 311, New York.
Gasterosteus lotharingus Blanchard, Poiss. France, 1866, 244, Lorraine.
Gasterosteus breviceps Blanchard, loc. cit., 1866, 245, Caen, France.
Gasterosteus blanchardi Sauvage, Révision des Epinoches, 1874, 32, New
 York.
- 1857. *Pungitius mainensis* (Storer).**
 Kennebec River, Me.; a form or species with seven dorsal spines.
Gasterosteus mainensis Storer, Bost. Jour. Nat. Hist., I, 1837, 464,
 Kennebec County, Me.
- 1858. *Pungitius brachypoda* (Bean).**
 Baffins Bay, Greenland. A form with 10 dorsal spines and the ventral
 spines very small.
Gasterosteus pungitius-brachypoda Bean, Bull. U. S. Nat. Mus., XV,
 1879, 129, Oosooadlin Mountain, Cumberland Gulf, Greenland.

Subfamily APELTINÆ

- Genus 631. APELTES DeKay.** *Four-spined stickleback.*
Apelles DeKay, N. Y. Fauna: Fishes, 1842, 87 (*Gasterosteus quadracus*
 Mitchill).
- 1859. *Apeltes quadracus* (Mitchill).**
 Maine to New Jersey, in salt water only.
Gasterosteus quadracus Mitchill, Trans. Lit. and Phil. Soc., I, 1815,
 430, New York.
Gasterosteus apelles Cuvier and Valenciennes, Hist. Nat. Poiss., IV,
 1829, 505, no locality.
Gasterosteus millipunctatus Ayres, Bost. Jour. Nat. Hist., I, 1842, 294,
 Old Mans Harbor, Long Island.

Subfamily SPINACHIINÆ

- Genus 632: SPINACHIA Cuvier.**
Spinachia Cuvier, Règne Anim., ed I, 1817, 320 (*Gasterosteus spinachia*
 Linnæus).
Polycanthus Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 242
 (*Gasterosteus spinachia* Linnæus).
Gastræa Sauvage, Révision des Epinoches, 1874, 7 (*Gasterosteus*
spinachia Linnæus).
- 1860. *Spinachia spinachia* (Linnæus).**
 Coasts of northern Europe; recorded from Newfoundland.
Gasterosteus spinachia Linnæus, Syst. Nat., ed. X, 1758, 442.
Spinachia vulgaris Fleming, Brit. Anim., 1828, 210.
Gasterosteus marinus Gronow, Cat. Fish., 1854, 168.

Family 150. AULORHYNCHIDÆ

- Genus 633. AULORHYNCHUS Gill.**
Aulorhynchus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 169 (*A.
 flavidus* Gill).
Auliscops Peters, Berliner Monatsberichte, 1866, 570 (*A. spinescens*
 Peters).

1861. **Aulorhynchus flavidus** Gill.

Coast of California, from San Nicolas Island and Monterey northward to Sitka, Alaska.

Aulorhynchus flavidus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 169, coast of Washington.

Suborder LOPHOBRANCHII

Family 151. SYNGNATHIDÆ. Pipefishes and seahorses

Subfamily SYNGNATHINÆ

Genus 634. DERMATOSTETHUS Gill.

Dermatostethus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 283 (*D. punctipinnis* Gill).

1862. **Dermatostethus punctipinnis** Gill.

San Diego, California.

Dermatostethus punctipinnis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 283, San Diego, Calif.; not lately recognized; the genus perhaps of doubtful value.

Genus 635. SYNGNATHUS Linnæus.

Syngnathus Linnæus, Syst. Nat., ed. X, 1858, 336 (*Syngnathus acus* L., the proper type as decided by the Internat. Com. Zool. Nomencl.).
Siphostoma Rafinesque, Caratteri, 1810, 18 (*Syngnathus pelagicus* Linnæus).

1863. **Syngnathus californiensis** Storer. Great pipefish.

Santa Barbara to Monterey and northward.

Syngnathus californiensis Storer, Proc. Bost. Soc. Nat. Hist., II, 1845, 73, California.

1864. **Syngnathus griseolineatus** Ayres.

Pacific coast of the United States, from Monterey northward to Puget Sound and southeastern Alaska.

Syngnathus griseolineatus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 14, San Francisco Bay.

Syngnathus abbotti Girard, Pac. R. R. Survey, Fishes, X, 1858, 346, San Francisco.

1865. **Syngnathus leptorhynchus** Girard.

West coast of America from San Francisco to San Diego.

Syngnathus brevirostris Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 156, San Diego; name preoccupied.

Syngnathus leptorhynchus Girard, loc. cit., 156, San Diego, Calif.

Syngnathus dimidiatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 284, San Diego; substitute for *brevirostris*, preoccupied.

1866. **Syngnathus arundinaceus** Girard.

Coast of California.

Syngnathus arundinaceus Girard, Pac. R. R. Surv., X, 1858, 346, coast of California.

1867. **Syngnathus carinatus** (Gilbert).

Gulf of California.

Siphostoma carinatum Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 547, Gulf of California, lat. $31^{\circ} 31'$ N., long. $114^{\circ} 19'$ W., at *Albatross* stations 3027 and 3028.

1868. **Syngnathus auliscus** (Swain).

Southern California.

Siphostoma auliscus Swain, Proc. U. S. Nat. Mus., V, 1882, 310, Santa Barbara and San Diego, Calif.

1869. **Syngnathus barbareæ** (Swain).

California coast at Santa Barbara. Rare; not reported recently.

Siphostoma barbareæ Swain, Proc. U. S. Nat. Mus., VII, 1884, 238, Santa Barbara, Calif.

1870. *Syngnathus mackayi* (Swain and Meek).
Key West to Yucatan.
Siphonotoma mackayi Swain and Meek, Proc. U. S. Nat. Mus., VII, 1884, 239, Key West, Fla.
1871. *Syngnathus floridæ* (Jordan and Gilbert).
Virginia to Texas and Porto Rico.
Siphonotoma floridæ Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 263, Pensacola, Fla.
1872. *Syngnathus poeyi* (Jordan and Evermann).
Habana.
Syngnathus tenuis Poey, Synopsis, 1867, 444, Cuba; name preoccupied.
Siphonotoma poeyi Jordan and Evermann, Fish. North and Mid. Amer., 1896, 766, Habana, Cuba; substitute for *S. tenuis*, preoccupied.
1873. *Syngnathus elucens*⁴⁰ Poey. *Pipefish*.
Cuba; Florida Keys; Porto Rico.
Syngnathus elucens Poey, Synopsis, 1867, 443, Habana.
Syngnathus flavirostris Poey, Enumeratio, 1876, 178, Habana.
Syngnathus picturatus Poey, loc. cit., 178, Habana.
Syngnathus lineata Poey, loc. cit., 178, Habana.
Syngnathus marmoratus Poey, loc. cit., 178, Habana.
Syngnathus ascendens Poey, loc. cit., 179, Habana.
1874. *Syngnathus rousseau* Kaup.
West Indies; known from Jamaica, St. Lucia, and Martinique.
Syngnathus rousseau Kaup, Lophobranchii, 1856, 40, Martinique.
1875. *Syngnathus jonesi* Günther. *West Indian pipefish*.
Bermudas and Porto Rico.
Syngnathus jonesi Günther, Ann. Mag. Nat. Hist., ser. 4, XIV, 1874, 8, 455, Bermudas.
1876. *Syngnathus fistulatus* Peters.
Puerto Cabello, near Colon.
Syngnathus fistulatus Peters, Monatsber., Ak. Wiss. Berl., 1868, 456, Puerto Cabello, near Aspinwall.
1877. *Syngnathus pelagicus* Linnaeus.
Open sea in the Atlantic and the Mediterranean; doubtfully recorded from the West Indies; some other species, perhaps *S. fistulata*, being mistaken for it.
Syngnathus pelagicus Linnaeus, Syst. Nat., ed. I, 1758, open sea in floating seaweed.
Syngnathus ethon Risso, Eur. Mérid., III, 1826, 182, Nice, France.
1878. *Syngnathus robertsi* (Jordan and Rutter).
West Indies and Jamaica; Caribbean Sea.
Siphonotoma robertsi Jordan and Rutter, Proc. Acad. Nat. Sci. Phila., XLIX, 1897, 97, Jamaica.
1879. *Syngnathus brachycephalus* Poey.
Tortugas; Cuba.
Syngnathus brachycephalus Poey, Synopsis, 1867, 444, Habana, Cuba.
1880. *Syngnathus brevicaudus* (Meek).
Boca del Rio, Vera Cruz.
Siphonotoma brevicaudum Meek, Field Col. Mus., Zool., V, 1904, 163, Boca del Rio, Mexico.
1881. *Syngnathus crinigerum* (Bean and Dresel).
Pensacola and Key West, to Abrolhos Reef, Brazil.
Siphonotoma crinigerum Bean and Dresel, Proc. Biol. Soc. Wash., II, 1884, 99, Pensacola, Fla.
1882. *Syngnathus sinaloæ* (Jordan and Starks).
Pacific coast of Mexico.
Siphonotoma sinaloæ Jordan and Starks, Proc. Calif. Acad. Sci., 3d ser., VI, 1896, 268, Mazatlan, Mexico.

* Some of the nominal species here included may prove to be valid.

- 1883.** *Syngnathus arctus* (Jenkins and Evermann).
 Gulf of California, Guaymas.
Siphonostoma arctum Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 137, Bay of Guaymas, Sonora, Mexico.
- 1884.** *Syngnathus starksii* (Jordan and Culver).
 Estuaries about Mazatlan.
Siphonostoma starksii Jordan and Culver, Proc. Calif. Acad. Sci., 3d ser., V, 1895, 416, Pl. XXX, Rio Presidio, Mazatlan, Mexico.
- 1885.** *Syngnathus dendriticus* (Barbour).
 Ireland Island, Bermudas.
Siphonostoma dendriticum Barbour, Bull. Mus. Comp. Zool., XLVI, No. 7, 1904-1906, 115, Ireland Island, Bermudas.
- 1886.** *Syngnathus mindii* Meek and Hildebrand.
 Panama.
Syngnathus mindii Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, 261, Pl. XVIII, fig. 2; brackish creek near Mindi, Canal Zone, Panama.
- 1887.** *Syngnathus elcapitanensis* Meek and Hildebrand.
 Streams, west coast of Panama.
Syngnathus elcapitanense Meek and Hildebrand in Meek, Field Mus. Nat. Hist., Zool., X, 1914, 119, Rio Mamona at El Capitan, Panama.
- 1888.** *Syngnathus tweedliei* Meek and Hildebrand.
 Panama.
Syngnathus tweedliei Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, 259, Pl. XVIII, fig. 1, Chame Point, Panama.
- 1889.** *Syngnathus torrei* (Nichols).
 Cuba.
Siphonostoma torrei Nichols, Bull. Amer. Mus. Nat. Hist., XXXI, 1912, 183, fig. 1, Matanzas, Cuba.
- Genus 636. **SYRICTES** Jordan and Evermann.
Syrichtes Jordan and Evermann Proc. Calif. Acad. Sci., XVI, Apr. 27, 1927, 504 (*Syngnathus fuscus* Storer).
- 1890.** *Syrichtes fuscus*^a (Storer). Common pipefish; Green pipefish; Billfish.
 Atlantic coast of United States from Maine to Virginia; very common northward.
Syngnathus fuscus Storer, Rept. Fish. Mass., 1839, 162, Nahant, Mass.
Syngnathus peckianus Storer, Rept. Fish. Mass., 1839, 163, Holms Hole, Marthas Vineyard.
Syngnathus peckii Storer, Rept. Fish. Mass., 1839, 3; name only.
Syngnathus fasciatus DeKay, N. Y. Fauna: Fishes, 1842, 319, pl. 54, fig. 174, New York.
Syngnathus viridescens DeKay, loc. cit., 321, pl. 54, fig. 176, New York.
Syngnathus dekayi Duméril, Hist. Nat. Poiss., II, 1870, 569, New York.
Syngnathus milbertianus Duméril, loc. cit., 573, New York.
- 1891.** *Syrichtes affinis* (Günther).
 Gulf of Mexico, on coast of Louisiana.
Syngnathus affinis Günther, Cat., VIII, 1870, 163, Louisiana.
- 1892.** *Syrichtes scovelli* (Evermann and Kendall).
 Coast of Texas and both coasts of Florida.
Siphonostoma scovelli Evermann and Kendall, Proc. U. S. Nat. Mus., XVIII, 1895, 113, Shamrock Point, Corpus Christi, Tex.
- 1893.** *Syrichtes louisianæ* (Günther).
 Atlantic and Gulf coasts of the United States; North Carolina to Texas and south to Key West.
Syngnathus pelagicus, var., Linnaeus, Syst. Nat., ed. XII, 1766, 416, Charleston, S. C.
Syngnathus louisianæ Günther, Cat., VIII, 1870, 160, New Orleans, La.
- 1894.** *Syrichtes bairdianus* (Duméril).
 West coast of Lower California.
Syngnathus bairdianus Duméril, Hist. Nat. Poiss., II, 1870, 574, "Coast of Mexico, near California."

^a The names *fuscus* and *peckianus* are of the same date. The author who first perceived this identity chose the second name in spite of the page priority of *fuscus*.

1895. Syrichtes exilis (Osburn and Nichols).

Gulf of California.

Siphostoma exile Osborn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 153, fig. 7, west San Benito Island, Port San Bartolome.**Genus 637. HIPPICHTHYS Bleeker.***Hippichthys* Bleeker, Verh. Batav. Genootsch., XXII, 1849, 15 (*H. heptagonus* Bleeker); not *Hippocampus heptagonus* Rafinesque.*Corythoichthys* Kaup, Archiv Naturgesch., 1853, 231 (*C. albirostris* Heckel MS. Kaup).**1896. Hippichthys albirostris** (Heckel).

West Indies, north to Pensacola and Key West; south to Bahia, about coral reefs.

Corythoichthys albirostris Heckel in Kaup, Archiv Naturgesch., 1853, 231, Mexico; Bahia; type examined.*Siphostoma zatropis* Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 264, Snapper Banks, Pensacola.**1897. Hippichthys cayorum** (Evermann and Kendall).

Key West, Fla.; Porto Rico.

Corythoichthys cayorum Evermann and Kendall, Bull. U. S. Fish Com., XVII, 1897, 128, pl. 7, fig. 7, near Crawfish Bar, Key West, Fla.**1898. Hippichthys spicifer** (Rüppell).

Tropical Pacific; recorded from Tehuantepec.

Syngnathus spicifer Rüppell, Neue Wirb., Fische, about 1830, 143.**Genus 638. DORYICHTHYS Kaup.***Doryichthys* Kaup, Archiv Naturgesch., 1853, 233 (*D. bilineatus* Heckel).**1899. Doryichthys lineatus** Valenciennes.

Tropical parts of Atlantic from Cuba to Brazil.

Doryichthys lineatus Valenciennes, in Kaup, Lophobranchii, 1856, 59, Bahia, Mexico, and Guadeloupe.**Genus 639. DORYRHAMPHUS Kaup.***Doryrhamphus* Kaup, Archiv Naturgesch., 1853, 233 (*D. excisus* Kaup).
Choeroichthys Kaup, loc. cit., 1853, 233 (*C. valenciennesi* Kaup).**1900. Doryrhamphus californiensis** Gill.

Gulf of California.

Doryrhamphus californiensis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 284, Cape San Lucas, Lower California.**1901. Doryrhamphus sierra** Nichols.

Porto Rico.

Doryrhamphus sierra Nichols, Bull. Amer. Mus. Nat. Hist., XXXIV, 1915, 145, fig. 2, Porto Rico.**Genus 640. ACUS** ⁴⁴ Swainson.*Acus* Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 333 (*Syngnathus squoreus* "Yarrell").*Osphylax* Cope, Proc. Acad. Nat. Sci. Phila., XXVII, 1875, 450 (*O. pellucidus* Cope).**1902. Acus squoreus** (Linnæus). *Ocean pipefish.*

Open sea along coasts of Europe; recorded from New Orleans, some other species probably mistaken for it.

Syngnathus squoreus Linnæus, Syst. Nat., ed. X, 1758, 417, open sea.*Syngnathus sibbaldi* Walbaum, Artedi Pisc., 1792, 6, Scotland.*Nerophis maculata* Rafinesque, Indice, 1810, 57, Palermo, Italy.*Nerophis anguineus* Kaup, Lophobr., 1856, 65, Brownsey, Dorsetshire; Berwick.**1903. Acus martinicensis** (Bibron) Kaup.

West Indies.

Nerophis martinicensis (Bibron) Kaup, Lophobr., 1856, 67, Martinique.

⁴⁴ We may adopt the generic name *Acus* for the species of the old genus *Nerophis* (Rafinesque, Indice, Sicil., 1810, 57, type *Syngnathus ophidion* L.) in which a small caudal fin is present, the tail not ending in a filament, as is the case in the type of *Nerophis* (*Syngnathus ophidion* L.). The name *Acus*, variously used by authors, is not preoccupied in binomial nomenclature.

1904. *Acus heckeli* (Kaup).

Caribbean Sea.

Nerophis heckeli Kaup, Lophobr., 1856, 66, Bogota; evidently not the exact locality.1905. *Acus pellucidus* (Cope).

Open Atlantic.

Osphylax pellucidus Cope, Proc. Acad. Nat. Sci. Phila., XXVII, 1875, 450, pl. 25, figs. 1 to 4, open Atlantic Ocean; a larval form of doubtful relationships and differing from *Acus* in the very short dorsal (D. 16; about 40 in *Acus*).

Subfamily HIPPOCAMPINÆ

Genus 641. *NANNOCAMPUS* Günther.*Nannocampus* Günther, Cat., VIII, 1870, 178 (*N. subosseus* Günther).1906. *Nannocampus nanus* Rosen.

Bahama Islands.

Nannocampus nanus Rosen, Arsskr. Univ. Lund, VII, 1912, 50, Bahamas.Genus 642. *HIPPOCAMPUS* Rafinesque.*Hippocampus* Rafinesque, Caratteri, 1810, 18 (*Syngnathus hippocampus* Linnaeus).1907. *Hippocampus hudsonius* DeKay. *Common American seahorse; Horse-fish.*

Atlantic coast of United States, from Cape Cod to the Tortugas.

Hippocampus hudsonius DeKay, N. Y. Fauna: Fishes, 1842, 322, p. 53, fig. 171, New York.*Hippocampus levicaudatus* Heckel, in Kaup, Lophobranchii, 1856, 16, North America.1908. *Hippocampus ingens* Girard. *Caballito del mar.*

Pacific coast from San Diego to Mazatlan; rare.

Hippocampus ingens Girard, Pac. R. R. Surv., Fishes, X, 1858, 342, San Diego, Calif.*Hippocampus gracilis* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 282, Cape San Lucas, Lower California.1909. *Hippocampus brunneus* Bean.

Bermudas.

Hippocampus brunneus Bean, Proc. Biol. Soc. Wash., XIX, 1906, 32, Long Bird Island, Bermudas.1910. *Hippocampus punctulatus* Guichenot. *Caballito del mar; Seahorse; Horsefish.*

Tropical parts of the Atlantic; common in the West Indies, Brazil, and western Africa; Gulf Stream; Beaufort, N. C.

Hippocampus punctulatus Guichenot in Ramon de la Sagra, Hist. île. Cuba, Poiss., 1853, 174, pl. 5, fig. 2, Cuba.*Hippocampus marginalis* Heckel, in Kaup, Lophobr., 1856, 15, Mexico.*Hippocampus fasciularis* Heckel, in Kaup, loc. cit., 15, Mexico.1911. *Hippocampus stylifer* Jordan and Gilbert.

Coasts of Florida; Snapper Banks off Pensacola and Tampa; Gulf Stream.

Hippocampus stylifer Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 265, Snapper Banks, Gulf of Mexico.1912. *Hippocampus zosterae* Jordan and Gilbert.

Pensacola Bay, Fla.

Hippocampus zosterae Jordan and Gilbert, Proc. U. S. Nat. Mus., V. 1882, 265, Grand Lagoon, Pensacola, Fla.; smallest of the seahorses,

Order PP. AULOSTOMI

Family 152. AULOSTOMIDÆ. Trumpetfishes

Genus 643. *AULOSTOMUS* Lacépède.*Aulostomus* Lacépède, Hist. Nat. Poiss., V, 1803, 357 (*A. chinensis* Lacépède).*Polypterusichthys* Bleeker, Nat. Tidschr., Neder-Indie, IV, 1853, 608 (*P. valentini* Bleeker).*Aulostoma* Schlegel, change of spelling.

1913. *Aulostomus maculatus* Valenciennes. *Trompetero; Trumpetfish.*
Caribbean Sea, north to southern Florida.
?Aulostomus maculatum Valenciennes, in Cuvier, Règne Anim., III,
Poiss., 1845, pl. 92, fig. 2.
Aulostoma coloratum Müller and Troschel, in Schomburgk, History
Barbados, 1848, 673, Barbados.

1914. *Aulostomus cinereus* (Poey).
Cuba.
Aulostoma cinereum Poey, Synopsis, 1867, 386, Cuba.

Family 153. FISTULARIIDÆ. Cornetfishes

Genus 644. FISTULARIA Linnaeus.

Fistularia Linnaeus, Syst. Nat., ed. X, 1758, 312 (*F. tabacaria* Linnaeus).
Cannorhynchus Cantor, Jour. Roy. Asiatic Soc. Bengal, XVIII, 1849
(*Fistularia tabacaria* Linnaeus); substitute name, *Fistularia* having
been used in 1750 as a pre-Linnaean name for a group of polyps.
Flagellaria Gronow, Cat. Fishes, 1854 (*F. fistularis* Gronow = *Fistularia*
tabacaria Linnaeus).

Solenostomus (Gronow) Gill, Cat. Fish. East Coast. N. A., Supl. to
Proc. Acad. Nat. Sci. Phila., XIII, 1861, 38 (*Fistularia tabacaria*
Linnaeus).

1915. *Fistularia tabacaria* Linnaeus. *Trumpetfish; Trompetero; Tobacco*
trumpetfish; Unarmed trumpetfish.
West Indies and neighboring seas; northward to Carolina and Florida,
and occasionally to Long Island.
Fistularia tabacaria Linnaeus, Syst. Nat., ed. X, 1758, 312, tropical
America.
Fistularia neo-boracensis Mitchell, Trans. Lit. Phil. Soc. Phila., I,
1815, 437, New York.
Aulastome marcgravii Castelnau, Anim. Nouv. Amer. Sud, 1855, 30,
Bahia, Rio Janeiro.
Flagellaria fistularis Gronow, Cat. Fish., 1854, 146, American Ocean.

1916. *Fistularia serrata* ⁴³ Cuvier. *Flutemouth.*

Western Pacific and other warm seas; Bermuda; Cuba.
Fistularia serrata Cuvier, Règne Anim., ed. I, II, 1817, 389, 34; after Bloch.
?Fistularia immaculata Cuvier, loc. cit., 349, Sea of the Indies.
Fistularia commersonii Rüppell, Neue Wirtheitthiere, 1834, 142, Red Sea.
Fistularia villosa Klunzinger, Abh. Zool. Ges. Wien, XXI, 1871, 516,
Red Sea.

1917. *Fistularia petimba* Lacépède. *Corneta.*

East Indies; Australia; China; Panama; Lower California; Japan;
Hawaii.
Fistularia petimba Lacépède, Hist. Nat. Poiss., V, 1803, 349, New
Britain, Isle of Reunion, equatorial Pacific.
Fistularia depressa Günther, Challenger rept. Shore Fishes, Zoology, I,
pt. 6, 1880, 69, pl. 32, fig. D, Sulu Archipelago, station 200, 255
fathoms.

1918. *Fistularia corneta* Gilbert and Starks.

Pacific coast of tropical America; Panama.
Fistularia corneta Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904,
56, Pl. X, Bay of Panama; Mazatlan.

Family 154. MACRORHAMPHOSIDÆ. Snipefishes

Genus 645. MACRORHAMPHOSUS Lacépède. Snipefishes.

Macrorhamphosus Lacépède, Hist. Nat. Poiss., V, 1803, 136 (*Silurus*
cornutus Forskål = *Balistes scolopax* Linnaeus).
Centriscus Cuvier, Règne Animal, ed. I, II, 1817, 350; not *Centriscus*
Linnaeus.
Macrogynathus Gronow, Cat. Fishes, 1854, 147 (*Centriscus scolopax*
Linnaeus).

⁴³ Weber calls this species *villosa* but quotes *serrata* in synonymy. The synonymy of *serrata* and *petimba* seem hopelessly confused.

1919. *Macrorhamphosus scolopax* (Linnaeus). *Snipefish; Trumpetfish; Bellowsfish.*
Mediterranean Sea, northward to southern England, accidental on our North Atlantic Coast.
Balistes scolopax Linnaeus, Syst. Nat., ed. X, 1758, 329, Mediterranean.
Silurus cornutus Forskål, Descr. Anim., 1775, 66, Marseilles.
1920. *Macrorhamphosus hawaiiensis* Gilbert.
Deep waters about Hawaii; once taken off San Pedro, Calif. (W. F. Thompson).
Macrorhamphosus hawaiiensis Gilbert, Bull. U. S. Fish Com., XXIII, 1903 (1905), 613, Laysan Island.

Order QQ. PERCOMORPHI

Suborder PERCESOCES

- ✓ Family 155. ATERINIDÆ. Silversides; Hardheads; Pescados del Rey; Peixe Rey

Genus 646. *HEPSETIA* Bonaparte.
Hepsetia Bonaparte, Fauna Italica, Fasc. 91 (no pages), 1837 (*Atherina boyeri* Risso).
Atherinomorus Fowler, Proc. Acad. Nat. Sci. Phila., LV, 1903, 730 (*Atherina laticeps* Poey = *Atherina stipes* Müller and Troschel).

1921. *Hepsetia stipes* (Müller and Troschel). *Cabezote, Bristle herring.*
West Indies; common north to western Florida.
? *Atherina tenuiflava* Spix in Agassiz, Pisc. Bras., 1829, 135, pl. 30, fig. 2, Atlantic Ocean, off Brazil.
Atherina stipes Müller and Troschel, in Schomburgk, Hist. Barbados, 1847, 671, Barbados.
Atherina laticeps, Poey, Memorias, II, 1861, 265, Habana.
Atherina velieana Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 342, Clear Water Harbor, Fla.

1922. *Hepsetia evermanni* (Eigenmann).
Cuba; San Cristobal; Pinar del Rio.
Atherina evermanni Eigenmann, Bull. U. S. Fish Com., XXII, 1902, 228, fig. 9, San Cristobal, Cuba.

Genus 647. *ATHERINA* (Artedi) Linnaeus. *Friars.*
Atherina Artedi in Linnaeus, Syst. Nat., ed. X, 1758, 315 (*Atherina hepsetus* Linnaeus).
Aphia Risso, Europ. Mérid. 1826 (*Aphia meridionalis* Risso = apparently very young of *Atherina hepsetus* L.), but Risso describes their laying of eggs, and may refer to some other species.
Tæniomembras Ogilby, Proc. Linn. Soc. New South Wales, XXIII, 1898, 41 (*Atherina microstoma* Günther).

1923. *Atherina arxa* Jordan and Gilbert.
Gulf of Mexico at Key West, Cozumel, and south to Caribbean Sea.
Atherina arxa Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 27, Key West, Fla.

1924. *Atherina harringtonensis* Goode.
Bermudas and south to Curaçao.
Atherina harringtonensis Goode, Amer. Jour. Sci. Arts, 3d ser., XIV, No. 82, 1877, 297, Bermuda Islands.

1925. *Atherina microps* Poey.
Habana.
Atherina microps Poey, Memorias, II, 1861, 266; Habana, Cuba.

1926. *Atherina carolina* Cuvier and Valenciennes.
South Carolina (reference doubtful).
Atherina carolina Cuvier and Valenciennes, Hist. Nat. Poiss., X, 1835, 330 (445), "Carolinias."

Genus 648. MENIDIA Bonaparte.

Menidia Bonaparte, Fauna Italica, Tome III, Pesci, in description of *Atherina hepsetus* p. 91, 1836, no type, *Atherina menidia* Linnæus intended.

Ischnomembras Fowler, Proc. Acad. Nat. Sci. Phila., LV, 1903, 730, pl. 42, "Gabun River, Gabun County, West Africa," probably an error.

Phoxargyreia Fowler, loc. cit., 1903, 732, *Phoxargyreia dayi* Fowler from "India," doubtless an error.

Subgenus ISCHNOMEMBRAS Fowler.

1927. *Menidia beryllina* (Cope). *Silverside; Fresh-water silverside.*

Coastwise streams, Massachusetts to Florida and lower Mississippi, north to Tennessee.

?*Atherinichthys gracilis* Günther, Cat., III, 1861, 405, no locality given. *Christoma beryllinum* Cope, Trans. Amer. Phil. Soc., 1866, 403, Potomac River, near Washington.

Menidia audens Hay, Bull. U. S. Fish Com., II, 1882, 64, Memphis, Tenn., and Vicksburg, Miss.

Ischnomembras gabunensis Fowler, Proc. Acad. Nat. Sci. Phila., LV, 1903, 731, pl. 42, Gabon River, West Africa; probably an error.

1928. *Menidia beryllina cerea* Kendall.

Atlantic coast, Massachusetts to South Carolina.

Menidia beryllina cerea Kendall, Rept. U. S. Fish Com., XXVII, 1901, 261, Waquoit Bay, Mass.

1929. *Menidia peninsulæ* (Goode and Bean).

Florida and Gulf coast, in salt water.

Chiostoma peninsulæ Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 148, Pensacola.

1930. *Menidia peninsulæ atrimentis* Kendall.

Florida; South Lake, Lake Monroe, Salt Lake, etc.

Menidia peninsulæ atrimentis Kendall, Rept. U. S. Fish Com., XXVII, 1901, 258, South Lake, Fla.

1931. *Menidia menidia* (Linnæus).

Atlantic coast, Cape Hatteras to Florida.

Atherina menidia Linnæus, Syst. Nat., ed. XII, 1766, 519, Charleston, S. C.

Atherina boscii Cuvier and Valenciennes, Hist. Nat. Poiss., X, 1835, 345 (465), Charleston.

Menidia dentex Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 429, St. Johns River, Fla.

1932. *Menidia conchorum* Hildebrand and Ginsberg.

Key West, Fla.

Menidia conchorum Hildebrand and Ginsberg, Bull. U. S. Bur. Fish., XLII, 1926 (1927), 207, fig. 1, Boca Chica, Key West, Fla.

1933. *Menidia notata* (Mitchill). *Silversides; Friar; Whitebait.*

Atlantic coast of United States, Nova Scotia to Cape May, intergrading south with *M. menidia*.

Atherina notata Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 446, New York.

Atherina viridescens Mitchill, loc. cit., 447, New York.

Phoxargyreia dayi Fowler, Proc. Acad. Nat. Sci. Phila., LV, 1903, 732, pl. 41, "India"; no doubt an error in locality.

1934. *Menidia starksi* Meek and Hildebrand.

Panama.

Menidia starksi Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 267, Pl. XX, fig. 2, Taboga Island, Panama.

1935. *Menidia sardina* (Jenkins and Evermann). *Pez del rey.*

Gulf of California.

Atherina sardina Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 137, Guaymas, Sonora, Mexico.

1936. **Menidia clara** Evermann and Jenkins.
Gulf of California.
Menidia clara Evermann and Jenkins, Proc. U. S. Nat. Mus., XIII, 1891, 136, Guaymas, Sonora, Mexico.
- Genus 649. **ARCHOMENIDIA** Jordan and Hubbs.
Archomenidia Jordan and Hubbs, Rev. Silversides, 1919, 54 (*Atherinichthys sallae* Regan).
1937. **Archomenidia sallaei** (Regan).
Mexico.
Atherinichthys sallaei Regan, Proc. Zool. Soc. Lond., 1903, 60, Mexico.
- Genus 650. **HUBBESIA** Jordan.
Hubbesia Jordan, Proc. U. S. Nat. Mus., LV, 1920 (May 2, 1919), 310 (*Menidia gilberti* Jordan and Bollman).
1938. **Hubbesia gilberti** (Jordan and Bollman).
Panama.
Menidia gilberti Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 155, Panama.
- Genus 651. **MEMBRAS** Bonaparte.
Membras Bonaparte, Fauna Italica, fasc. 91 (not paged), 1837 (type not named directly, *Atherina martinica* Cuvier and Valenciennes being indicated).
1939. **Membras martinicus** (Cuvier and Valenciennes).
Martinique.
Atherina martinica Cuvier and Valenciennes, Hist. Nat. Poiss., X, 1835, 340 (459), Martinique.
1940. **Membras beani** (Meek and Hildebrand).
Atlantic waters of Panama.
Kirtlandia beani Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 270, Pl. XXI, Fox Bay, Colon.
- Genus 652. **KIRTLANDIA** Jordan and Evermann.
Kirtlandia Jordan and Evermann, Fishes North and Mid. Amer., 1896, 794 (*Chirostoma vagrans* Goode and Bean).
1941. **Kirtlandia vagrans** (Goode and Bean). *Rough silverside.*
Gulf of Mexico; Florida to Texas; occasionally northward.
Chirostoma vagrans Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 148, Pensacola, Fla.
1942. **Kirtlandia laciniata** (Swain). *Silverfish; Whitebait; Silversides.*
Lower Chesapeake to South Carolina, intergrading with *K. vagrans*.
Menidia vagrans laciniata Swain, in Jordan and Gilbert, Synopsis, 1882, 908, Beaufort, N. C.
- Genus 653. **THYRINA** Jordan and Culver.
Thyrina Jordan and Culver, in Jordan, Proc. Calif. Acad. Sci., 3d ser., V, 1895, 419 (*Atherinella evermanni* Jordan and Culver).
Melaniris Meek, Field Mus.-Zool., III, 1902, 117 (*Melaniris balsanus* Meek).
1943. **Thyrina crystallina** Jordan and Culver.
Rio Presidio near Mazatlan, Sinaloa.
Thyrina crystallina Jordan and Culver, in Jordan, Proc. Calif. Acad. Sci., 3d ser., V, 1895, 420, Rio Presidio, below Presidio, Sinaloa.
1944. **Thyrina evermanni** Jordan and Culver.
West coast of Mexico; Mazatlan.
Thyrina evermanni Jordan and Culver, Proc. Calif. Acad. Sci., 3d ser., V, 1895, 419, Mazatlan, Mexico.
1945. **Thyrina guija** Hildebrand. *Pepesca; Manjuda; Alfiler; Rovalete.*
Lake Guija, Rio del Desague near Lake Guija and Rio Lempa at Suchitoto and San Marcos, El Salvador.
Thyrina guija Hildebrand, Bull. U. S. Bur. Fish., XLI, 1925, 264, fig. 18, Lake Guija, El Salvador.

1946. **Thyrina balsana** (Meek).
 Mexico, Rio Balsas, Guerrero.
Melaniris balsanus Meek, Pub. Field Col. Mus., Zool., III, 1902, 117,
 Rio Balsas, at Balsas.
1947. **Thyrina guatemalensis** (Günther).
 Lakes of Huamachal.
Atherinichthys guatemalensis Günther, Proc. Zool. Soc. London, 1864,
 151, Lakes of Huamachal, Pacific coast of Guatemala.
1948. **Thyrina sardina** (Meek). *Sardina*.
 Lakes Nicaragua and Managua.
Melaniris sardina Meek, Field Col. Mus., Zool., VII, 1907, 114, Lake
 Managua, Nicaragua.
Chirosoma guatemalensis Gill, Proc. Acad. Nat. Sci. Phila., XXIX,
 1877, 187, Lake Nicaragua; not of Günther.
1949. **Thyrina chagresi** (Meek and Hildebrand).
 Streams of Atlantic slope of Panama.
Menidia chagresi Meek and Hildebrand, Field Mus. Nat. Hist., Zool.,
 X, 1914, 119, Rio Chagres at Gorgona, Zent and Parismina, Costa
 Rica.
1950. **Thyrina meeki** Miller.
 Rio Motagua, Guatemala.
Thyrina meeki Miller, Bull. Amer. Mus. Nat. Hist., XXIII, 1907,
 110, fig. 2, Rio Motagua, east coast of Guatemala.
- Genus 654. THYRINOPS Hubbs.**
Thyrinops Hubbs, Proc. Acad. Nat. Sci. Phila., LXIX, 1918, 306
 (*Atherinichthys pachylepis* Günther).
1951. **Thyrinops pachylepis** (Günther).
 Panama.
Atherinichthys pachylepis Günther, Proc. Zool. Soc. Lond., 1864, 25,
 Panama.
- Genus 655. MELANORHINUS Metzelaar.**
Melanorhinus Metzelaar, Tropisch Atl. Visschen, 1919, 38 (*M. boeki*
 Metzelaar).
1952. **Melanorhinus boeki** Metzelaar.
 West Indies.
Melanorhinus boeki Metzelaar, Trop. Atl. Visschen, 1919, 38, Oyster
 Pond, St. Martin Island.
- Genus 656. XENATHERINA Regan.**
Xenatherina Regan, Biologia Centrali Amer., Pisces, 1906, 64 (*Menidia*
lisa Meek).
1953. **Xenatherina lisa** (Meek).
 Streams of Vera Cruz, Mexico.
Menidia lisa Meek, Field Col. Mus., Pub. Zool., V, 1904, 182, Refugio,
 Vera Cruz, Basin of Rio Papaloapam, Mexico.
- Genus 657. ATHERINELLA Steindachner.**
Atherinella Steindachner, Ichth. Beitr., II, 1875, 35 (*A. panamensis*
 Steindachner).
1954. **Atherinella panamensis** Steindachner.
 Panama.
Atherinella panamensis Steindachner, Ichth. Beitr., II, 1875, 35,
 Panama.
- Genus 658. EURYSTOLE Jordan and Evermann.**
Eurystole Jordan and Evermann, Fishes North and Mid. Amer., 1896,
 802 (*Atherinella eriarcha* Jordan and Gilbert).
1955. **Eurystole eriarcha** (Jordan and Gilbert).
 West coast of Mexico; Mazatlan.
Atherinella eriarcha Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881,
 348, Mazatlan, Mexico.

Genus 659. CHIROSTOMA Swainson. *Pescados blancos.*

Chiostoma Swainson, Class. Fishes, etc., II, 1839, 243 (*Atherina humboldtiana* Cuvier and Valenciennes).

Atherinoides Bleeker, Verh. Bat. Gen., Japan, XXV, 1853, 40 (*Chiostoma vomerina* Cuvier and Valenciennes = *Atherina humboldtiana* Cuvier and Valenciennes).

Atherinichthys Bleeker, loc. cit., 40 (*Atherina humboldtiana* Cuvier and Valenciennes).

Heterognathus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 198 (*Atherina humboldtiana* Cuvier and Valenciennes).

Lethostole Jordan and Evermann, Fish. North and Mid. Amer., 1896, 792 (*Chiostoma estor* Jordan).

Eslopsarum Jordan, Rept. U. S. Fish Com., XIX, 1895, 330 (*Chiostoma jordani* Woolman).

Subgenus ESLOPSARUM Jordan.

1956. *Chiostoma jordani* Woolman.

Basin of Rio Lerma and Valley of Mexico.

Chiostoma jordani Woolman, Bull. U. S. Fish Com., XIV, 1894, 62, pl. 2, canals at Salamanca and City of Mexico.

Atherinichthys brevis Therese von Baier, Anz. Kais. Akad. Wiss. Wien, 1894, 149, Lake Cuitzeo, Mexico.

1957. *Chiostoma mezquital* Meek.

Durango.

Chiostoma mezquital Meek, Field Col. Mus., pub. 93, V, 1904, 170, Durango, Mexico.

1958. *Chiostoma labarcæ* Meek.

Rio Lerma, Mexico.

Chiostoma labarcæ Meek, Field Col. Mus., Zool., III, 1902, 113, Rio Lerma, Mexico.

1959. *Chiostoma arge* (Jordan and Snyder).

Rio Verde, Mexico.

Eslopearum arge Jordan and Snyder, Bull. U. S. Fish Com., XIV, 1899, 133, fig. 12, Rio Verde, Mexico.

Subgenus CHIROSTOMA Swainson.

1960. *Chiostoma bartoni* Jordan and Evermann.

tributary of the Rio Lerma, near Guanajuato, Mexico; Lake Patzcuaro.

Chiostoma bartoni Jordan and Evermann, Fish. North and Mid. Amer., 1896, 793, tributary of the Rio Lerma, near Guanajuato, Mexico.

Chiostoma attenuatum Meek, Field Col. Mus., Zool., III, 1902, 112, Lake Patzcuaro, Mexico.

1961. *Chiostoma zirahuen* Meek.

Lago de Zirahuen, Mexico.

Chiostoma zirahuen Meek, Field Col. Mus., Zool., III, 1902, 114, Lago de Zirahuen, Mexico.

1962. *Chiostoma patzcuaro* Meek.

Lago de Patzcuaro, Mexico.

Chiostoma patzcuaro Meek, Field Col. Mus., Zool., III, 1902, 113, Lago de Patzcuaro, Mexico.

1963. *Chiostoma humboldtianum* (Cuvier and Valenciennes). *Pescado blanco de Chalso.*

Lakes near the City of Mexico; La Laguna, Jalisco, Mexico.

Atherina humboldtiana Cuvier and Valenciennes, Hist. Nat. Poiss., X, 1835, 355 (479), Mexico.

Atherina vomerina Cuvier and Valenciennes, loc. cit., 357 (481), lake near City of Mexico.

1964. *Chiostoma regani* Jordan and Hubbs.

Valley of Mexico; Lake Patzcuaro.

Chiostoma regani Jordan and Hubbs, Leland Stanford Univ. Pub., Univ. Ser., 1919, 74, Xochimilco, Morelos, Mexico. After *Chiostoma breve* Regan; not of Steindachner.

Subgenus LETHOSTOLE Jordan and Evermann.

1965. *Chiostoma chapala* Jordan and Snyder.
Laguna de Chapala, Mexico.
Chiostoma chapala Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 135, fig. 13, Laguna de Chapala, Mexico, at Ocotlan.
1966. *Chiostoma consocium* Jordan and Hubbs.
Basin of Rio Lerma, Mexico.
Chiostoma consocium Jordan and Hubbs, Rev. Silversides, in Leland Stanford Univ. Pub. Univ. Ser., 1919, 76, Lake Chapala at La Palma, Mexico.
1967. *Chiostoma grandocule* (Therese von Baier).
Lake Patzcuaro, Mexico.
Atherinichthys grandoculis Therese von Baier, Anz. Kais. Akad. Wiss. Wien, 1894, Lake Patzcuaro, Mexico.
1968. *Chiostoma promelas* Jordan and Snyder.
Laguna de Chapala; Guadalajara, Mexico.
Chiostoma promelas Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 136, market of Guadalajara, Mexico, from Laguna de Chapala.
1969. *Chiostoma lucius* Boulenger.
Lake Chapala, Mexico.
Chiostoma lucius Boulenger, Ann. and Mag. Nat. Hist., ser. 7, V, January, 1900, 54, Lake Chapala, Mexico.
Chiostoma crystallinum Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1900), 139, fig. 16, Laguna de Chapala, Mexico, at Ocotlan.
1970. *Chiostoma diazi* Jordan and Snyder.
Lake Chapala, Mexico.
Chiostoma diazi Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1900), 137, fig. 15, Lake Chapala, Mexico.
1971. *Chiostoma sphyræna* Boulenger.
Lake Chapala, Mexico.
Chiostoma sphyræna Boulenger, Ann. Mag. Nat. Hist., ser. 7, V, 1900, 54, Lake Chapala, Mexico.
Chiostoma lerme Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1900), 142, fig. 19.
1972. *Chiostoma ocoatlane* Jordan and Snyder.
Lago de Chapala, Mexico.
Chiostoma ocoatlane Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899 (1900), 140; Lago de Chapala, Mexico.
1973. *Chiostoma estor* Jordan. *Pescado blanco de Chapala.*
Lake Chapala, Lake Patzcuaro, Lake Zirahuen, and Lake Xochimilco, Mexico.
Chiostoma estor Jordan, Proc. U. S. Nat. Mus., II, 1879, 298, Lake Chapala (an error for Lake Patzcuaro).
Atherinichthys albus Therese von Baier, Anz. Kais. Akad. Wiss. Wien, 1894, 148, Lake Patzcuaro, Mexico.
- Genus 660. **LABIDESTHES** Cope.
Labidesthes Cope, Proc. Amer. Phil. Soc., XI, 1870, 455 (*Chiostoma siculum* Cope).
1974. *Labidesthes sicculus* (Cope). *Brook silverside; Skipjack; Glassfish.*
Lake Ontario and southern Michigan to Iowa, Florida, and Texas, clear streams and lakes.
Chiostoma siculum Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 81, Grosse Isle, Detroit River, Mich.
- Genus 661. **LEURESTHES** Jordan and Gilbert. *Grunion.*
Leuresthes Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 29 (*Atherinopsis tenuis* Ayres).
1975. *Leuresthes tenuis* (Ayres). *Grunion.*
Sandy coasts of California from San Francisco to Lower California; abundant southward.
Atherinopsis tenuis Ayres, Proc. Calif. Acad. Nat. Sci., II, 1860, 75, San Francisco.
Leuresthes crameri Jordan and Evermann, Fishes North and Mid. Amer., 1896, 802, Ballenas Bay, Lower California.

- Genus 662. MUGILOPS Meek and Hildebrand.**
Mugilops Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, 271 (*M. cyanellus* Meek and Hildebrand).
- 1976. *Mugilops cyanella* Meek and Hildebrand.**
 Panama Bay.
Mugilops cyanellus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, 271, Pl. XXII, fig. 1, Balboa, Panama Bay.
- 1977. *Mugilops marina* Meek and Hildebrand.**
 Panama Bay.
Mugilops marinus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1923, 272, Pl. XXII, fig. 2, Porto Bello, Panama Bay.
- Genus 663. COLPICHTHYS Hubbs.**
Colpichthys Hubbs, Proc. Acad. Nat. Sci. Phila., LXVIX, 1917 (1918), 67 (*Atherinops regis* Jenkins and Evermann).
- 1978. *Colpichthys regis* (Jenkins and Evermann). *Pez del rey.***
 Gulf of California.
Atherinops regis Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 138, Bay of Guaymas, Sonora, Mexico.
- Genus 664. ATHERINOPS Steindachner. *Pascadillos del rey.***
Atherinops Steindachner, Ichth. Beitr., III, 1875, 61 (*Atherinopsis affinis* Ayres).
- 1979. *Atherinops affinis* (Ayres). *Little smelt; Pescadillo del rey.***
 Coast of California, and north to Oregon.
Atherinopsis affinis Ayres, Proc. Calif. Acad. Nat. Sci., II, 1860, 73, San Francisco.
Atherinops oregonia Jordan and Snyder, Proc. U. S. Nat. Mus., XLV, 1913, 575, pl. 46, Yachats River, Oreg.
- 1980. *Atherinops insularum* Gilbert.**
 San Clemente and San Nicolas Islands of the Santa Barbara Group; Guadalupe Island.
Atherinops insularum Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 549, San Clemente, San Nicolas, and Guadalupe Islands.
- 1981. *Atherinops insularum cedrosensis* Hubbs.**
 Shores of Cedros and San Benito Islands, Lower California.
Atherinops insularum cedrosensis Hubbs, Bull. Amer. Mus. Nat. Hist., XXXVIII, 1918, 434, southeast side of Cedros Island, Lower California.
- 1982. *Atherinops insularum guadalupae* Hubbs.**
 Guadalupe Island, off Lower California.
Atherinops insularum guadalupae Hubbs, Bull. Amer. Mus. Nat. Hist., XXXVIII, 1918, 435, Guadalupe Island.
- Genus 665. ATHERINOPSIS Girard. *Pescado del rey.***
Atherinopsis Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 134 (*A. californiensis* Girard).
- 1983. *Atherinopsis californiensis* Girard. *California blue 'smelt; Pescado del rey; Peize rey; Pesce rey.***
 Coast of California, from Cape Mendocino to Ballenas Bay, Lower California; abundant.
Atherinopsis californiensis Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 134, San Francisco, Calif.
- 1984. *Atherinopsis sonoreæ* Osburn and Nichols.**
 Cedros Island, Lower California.
Atherinopsis sonoreæ Osborn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 156, fig. 8, Cedros Island, Lower California.

✓ Family 156. MUGILIDÆ. Mullets

- Genus 666. MUGIL Linnaeus. *Striped mullets (soft dorsal and anal not scaled).***
Mugil Artedi, in Linnaeus, Syst. Nat., Ed. X, 1758, 316 (*M. cephalus* Linnaeus).
Arnion Gistel, Thierreich, 1848, X (*M. cephalus* L.).

- 1985.** *Mugil brasiliensis* Agassiz. *Liza; Lebrancho; Queriman; Blueback mullet; Fantail mullet.*
 Cuba to Patagonia; West Indies; Brazil; Habana, probably north to Virginia.
Mugil brasiliensis Agassiz, in Spix, Pisc. Brasil., 134 (234 by mispaging), 1829, pl. 72, Atlantic Ocean off Brazil.
Mugil liza Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 61 (83), Brazil; Porto Rico; Maracaibo; Surinam; Martinique.
Mugil lebranchus Poey, Memorias, II, 1861, 260, plate 18, fig. 3 Cuba.
Querimana gyrans Jordan and Gilbert, Proc., U. S. Nat. Mus., VII, 1884, 26, Key West; "the whirligig mullet" a little fish, with two anal spines, now regarded as the young of *Mugil*, but if so of *M. brasiliensis* as the scales are 28 to 29 (41 in *Mugil cephalus*). It is recorded from Woods Hole and Fortress Monroe; in revolving schools, perhaps mixed with the young of *M. cephalus*.
- 1986.** *Mugil cephalus* Linnaeus. *Common mullet; Striped mullet; Céfalo; Macho; Machuto; Lisa cabezuda; Callifaver mullet.*
 Cosmopolitan; coasts of southern Europe and northern Africa; Atlantic coast of America from Cape Cod to Brazil; Pacific coast of America from Monterey to Chile; Hawaii; Japan; entering rivers. Perhaps more than one species included, but we fail to detect differences. If different, the American species must be known as *Mugil albula* of the Atlantic and *Mugil mexicanus* of the Pacific.
Mugil cephalus Linnaeus, Syst. Nat., Ed. X, 1758, 316, Europe; based on Aristedi.
Mugil albula Linnaeus, Syst. Nat., Ed. XII, 1766, 520, Charleston, S. C.
? *Mugil tang* Bloch, Ichthyologia, 1794, 171, Africa.
? *Mugil plumieri* Bloch, loc. cit., 173, St. Vincent; on a drawing by Plumier.
Mugil lineatus Mitchell, in Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 71 (96), New York.
Mugil rammelsbergii Tschudi, Fauna Peruana, Ichth., 1845, 20, Peru.
Mugil berlandieri Girard, U. S. and Mex. Bound. Surv., 1859, 20, pl. 10, figs. 1 to 4, St. Josephs Island, Indianola, Brazos Santiago; Brazos; Galveston; all on coast of Texas.
Mugil guentherii Gill, Proc. Acad. Nat. Sci. Phila., XXI, 1863, 169, western coast of Central America; not of Steindachner.
Mugil mexicanus Steindachner, Ichth. Beitr., III, 1875, 58, Acapulco, Mexico.
- Genus 687. QUERIMANA** Jordan and Gilbert. *White mullets.* (Soft dorsal and anal closely scaled.)
Querimana Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 588 (*Myxus harengus* Günther, now regarded as the young of *Mugil curema*).
- 1987.** *Querimana curema* (Cuvier and Valenciennes). *White mullet; Blue-back mullet; Lisa; Lisa blanca.*
 Both coasts of America, Cape Cod to Brazil and Mazatlan to Chile.
Mugil curema Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 64 (87), Brazil; Martinique; Cuba.
Mugil petrosus Cuvier and Valenciennes, loc. cit., 65 (89), Brazil; Surinam; Gulf of Mexico; Cuba.
Myxus harengus Günther, Cat., III, 1861, 467, Pacific coast of Central America; young examples with two anal spines.
- 1988.** *Querimana incilis* (Hancock). *Trench mullet.*
 Waters of Central America; Rio Chagres to Pará and Bahia.
Mugil incilis Hancock, Quart. Jour. Sci., 1830, 127, Guiana.
Mugil guntheri Steindachner, Ichth. Notizen, I, 1864, 12, British Guiana.
- 1989.** *Querimana thoburni* (Jordan and Starks).
 Pacific coast of tropical America, from Guatemala to Galapagos Islands.
Mugil thoburni Jordan and Starks, in Jordan and Evermann, Fishes North and Mid. Amer., 1896, 812, Galapagos Islands.

1990. *Querimana hospes* (Jordan and Culver). *Lisita.*
West coast of Mexico to Panama Bay.
Mugil hospes Jordan and Culver, in Jordan, Proc. Calif. Acad. Sci., V, 1895, 422, pl. 31, Mazatlan, Mexico.
1991. *Querimana gaimardiana* (Desmarest). *Redeye mullet; Liza ojo de Perdiz.*
Florida Keys to Cuba; Key West.
Mugil gaimardianus Desmarest, Dict. Class., 1831, pl. 109, Cuba.
1992. *Querimana setosus* (Gilbert).
Clarion Island of the Revillagigedo Group.
Mugil setosus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 549, Clarion Island.
1993. *Querimana trichodon* (Poey). *Fantail mullet.*
Florida Keys to Brazil; Key West.
Mugil trichodon Poey, Ann. Lyc. Nat. Hist. N. Y., XI, 1875, 66, pl. 8, figs. 4-8, Cuba.
- Genus 668. **CHÆNOMUGIL** Gill.
Chænomugil Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 169 (*Mugil proboscideus* Günther).
1994. *Chænomugil proboscideus* (Günther).
Pacific coast of tropical America; M: zatlan; Cordova; Panama.
Mugil proboscideus Günther, Cat., III, 1861, 459, Island of Cordova (Cardon), west coast of Central America.
- Genus 669. **DAJAUS** Cuvier and Valenciennes. *Mountain mullets.*
? *Agonostomus* Bennet, Proc. Zool. Soc. Lond., I, 1830, 105 (*A. telfairii* Bennett).
Dajaus Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 122 (164) (*Mugil monticola* Bancroft).
Neomugil Vaillant, Bull. Soc. Philom., Paris, 1894, 73 (*N. digueti* Vaillant).
1995. *Dajaus monticola* (Bancroft). *Trucha; Mountain mullet.*
Fresh waters of the West Indies and the Atlantic and Pacific streams of Mexico and Central America.
Mugil monticola Bancroft, in Griffith's ed. Cuvier's Animal Kingdom, Fishes, 1836, 367, pl. 36, Jamaica.
Mugil irrelitus Gosse, Nat. Sojourn Jamaica, 1851, 84, Jamaica.
Agonostoma nasutum Günther, Cat., III, 1861, 463, Rio Geronimo, Guatemala.
Neomugil digueti Vaillant, Bull. Soc. Philom., IV, 1894, 73, Lower California.
Agonostomus salvini Regan, Ann. Mag. Nat. Hist., ser. XIX, 1907, 7, and Biol. Cent. Amer., Pisces., 1907, 68, Pl. XI, fig. 2, Nacasil, Guatemala.
1996. *Dajaus percoides* (Günther).
San Domingo.
Agonostoma percoides Günther, Cat., III, 1861, 464, San Domingo; perhaps not distinct from *Dajaus monticola*.
1997. *Dajaus microps* (Günther).
West Indies and Central America.
Agonostoma microps Günther, Cat., III, 1861, 462; probably West Indies.
- Genus 670. **JOTURUS** Poey.
Joturus Poey, Memorias, II, 1861, 263 (*J. pichardi* Poey).
Xenorhynchichthys Regan, Ann. Mag. Nat. Hist., II, 1908, 461 (*Joturus stipes* Jordan and Gilbert).
1998. *Joturus pichardi* Poey. *Joturo.*
Rivers of Cuba; at the foot of waterfalls.
Joturus pichardi Poey, Memorias, II, 1861, 263, Rio Almendares, near Habana.

1999. *Joturus globiceps* (Günther). *Bobo.*

Rivers and cataracts of eastern Mexico, from Jalapa to Costa Rica and Panama.

Agonostoma globiceps Günther, Ann. Mag. Nat. Hist., ser. 4, XIV, 1874, 370, Myzantla, Vera Cruz.

Joturus stipes Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 373, Rio Bayano, near Panama.

Family 157. SPHYRÆNIDÆ. Barracudas

Genus 671. SPHYRÆNA Röse.

Sphyraena Röse, in Artedi Piscium, 1793, 52 (*Esox sphyraena* Linnaeus).

Sphyraena Artedi, in Bloch and Schneider, Syst. Ichth., 1801, 109 (*Esox sphyraena* Linnaeus).

Agriosphyræna Fowler, Proc. Acad. Nat. Sci. Phila., LV, 1903, 749 (*Esox barracuda* Walbaum).

Subgenus SPHYRÆNA Röse.

2000. *Sphyraena sphyraena* (Linnaeus). *European barracuda; Spet; Sennet.*
Coasts of southern Europe and neighboring islands, west to the Bermudas.

Esox sphyraena Linnaeus, Syst. Nat., Ed. X, 1758, 313, Mediterranean Sea.

Esox spet Haüy, Encycl. Metod. Poiss., 1787; after Linnaeus.

Sphyraena vulgaris Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 242 (327), Mediterranean.

Sphyraena viridensis Cuvier and Valenciennes, loc. cit., 251 (339), St. Jago, Cape Verde Islands.

2001. *Sphyraena argentea* Girard. *California barracuda; Barracouta.*
Pacific coast, from San Francisco southward to Cape San Lucas.
Sphyraena argentea Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 144
San Diego, Calif.
Sphyraena lucasana Gill, loc. cit., XV, 1863, 86, Cape San Lucas.

2002. *Sphyraena ensis* Jordan and Gilbert. *Vicuda.*
Gulf of California to Panama.
Sphyraena ensis Jordan and Gilbert, Bull. U. S. Fish Com., II, 1882, 106,
Mazatlan, Mexico.

2003. *Sphyraena borealis* DeKay. *Northern barracuda; Sennet.*
Atlantic coast of the United States, from Cape Cod to North Carolina.
Sphyraena borealis DeKay, N. Y. Fauna: Fishes, 1842, 39, pl. 60,
New York.

2004. *Sphyraena guachancho* Cuvier and Valenciennes. *Guaguanche pelón;*
Guanuanche.
West Indies north to Pensacola; Gulf Stream to Woods Hole.
Sphyraena guachancho Cuvier and Valenciennes, Hist. Nat. Poiss., III,
1829, 252 (342), Habana.
Sphyraena guntheri Haly, Ann. Mag. Nat. Hist., XV, 1875, 270, Colon.

2005. *Sphyraena picudilla* Poey. *Picudilla; Baracuda.*
West Indies, on the coasts of Cuba; Bahia.
Sphyraena picudilla Poey, Memorias, II, 1861, 162; Habana.

Subgenus AGRIOSPHYRÆNA Fowler.

2006. *Sphyraena barracuda* (Walbaum). *Great barracuda; Picuda; Becuna;*
Short barracuda.
West Indies and Brazil, north to Pensacola, Charleston, and the Bermudas; accidental to Woods Hole, Mass.
Umbra minor, etc. (the barracuda) Catesby, Fishes of Carolina, etc.,
II, 1743, 1, pl. 1, Bahamas.
Esox barracuda Walbaum, Artedi Piscium, III, 1792, 94; after Catesby.
Sphyraena sphyraena var. *picuda* Bloch and Schneider, ed. Bloch, Syst.
Ichth., 1801, 110, Habana; after Parra.
Sphyraena becuna Lacépède, Hist. Nat. Poiss., V, 1803, pl. 9, fig. 3;
after drawing by Plumier at Martinique.

Suborder RHEGNOPTERI

Family 158. POLYNEMIDÆ. Threadfins

Genus 672. POLYNEMUS Linnæus. Barbudos.

Polynemus (Gronow) Linnæus, Syst. Nat., Ed. X, 1758, 317, (*P. quinquarius*, *virginicus*, and *paradiseus* Linnæus); properly restricted by Günther and by general usage to *Polynemus paradiseus* Linnæus.

Polydactylus Lacépède, Hist. Nat. Poiss., V, 1803, 419 (*P. plumieri* Lacépède = *Polynemus virginicus* Linnæus).

Trichidion Klein, in Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 274 (*Polynemus virginicus* Linnæus).

2007. *Polynemus approximans* Lay and Bennett. *Raton.*

Pacific coast of tropical America; Guaymas to Panama; Santa Catalina and San Diego.

Polynemus approximans Lay and Bennett, Beechey's Voyage, Zool., Fish., 1839, 57, Mazatlan, Mexico.

Polynemus californiensis Thominot, Bull. de la Soc. Philom. de Paris, Séance, 27 Juin, 1886, (7), X, 161, California.

2008. *Polynemus virginicus* Linnæus. *Barbudo; Barbu; Catfish.*

West Indies, north to the Florida Keys.

Polynemus virginicus Linnaeus, Syst. Nat., Ed. X, 1758, 317, America.

Polydactylus plumieri Lacépède, Hist. Nat. Poiss., V, 1803, 419, Martinique; from a drawing by Plumier.

Polynemus mango Lacépède, loc. cit., 413, America, after Linnæus.

Polynemus americanus Cuvier and Valenciennes, loc. cit., III, 1829, 291 (393), Cayenne, San Domingo, Martinique.

Polynemus oligodon Günther, Cat., II, 1860, 322, Rio Janeiro.

2009. *Polydactylus opercularis* (Gill).

Pacific coast of tropical America, from Cape San Lucas to Panama.

Trichidion opercularis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 168, Cape San Lucas.

Polynemus melanopoma Günther, Trans. Zool. Soc. Lond., VI, 1868, 421, San José de Guatemala.

2010. *Polynemus octonemus* Girard. *Threadfin; Eight-threaded threadfish.*

New York to the Rio Grande.

Polynemus octonemus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 167, Brazos Santiago; Galveston.

Trichidion octofilis Gill, loc. cit., XIII, 1861, 280, New York.

Genus 673. PENTANEMUS Günther.

Pentanemus Günther, Cat., II, 1860, 331 (*Polynemus quinquarius* Linnæus).

2011. *Pentanemus quinquarius* (Linnæus).

Atlantic Ocean; West Indies to coast of Africa; rare.

Polynemus quinquarius Linnæus, Syst. Nat., Ed. X, 1758, 317, America; after Gronow.

Polynemus artedii Bennett, Proc. Zool. Soc. Lond., 1831, 146, Africa.

Polynemus macronemus Pel, Bydrage tot de Dierk., 1851, 9, Africa.

Series SCOMBRIFORMES

Family 159. ACANTHOCYBIIDÆ

Genus 674. ACANTHOCYBIUM Gill. *Petos.*

Acanthocybium Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 125 (*Cybum sara* Bennett).

2012. *Acanthocybium petus* (Poey). *Peto.*

West Indies north to Key West; recorded from Italy.

Cybum petus Poey, Memorias, II, 1861, 234, pl. 16, fig. 1, Habana.

Cybum verany Döderlein, Giorn. di Sci. Nat., ed. Econ., VIII, 1872, Palermo.

2013. *Acanthocybium solandri* (Cuvier and Valenciennes). *Ono; Peto.*

Warmer parts of the Pacific; Hawaii; Pacific coast of Mexico; and South Seas.

Cybum solandri Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 192, open seas; exact locality unknown; after Solander.

✓ Family 160. CYBIIDÆ "4"

Genus 675. SIERRA Fowler.

Sierra Fowler, Proc. Acad. Nat. Sci. Phila., LVI, 1904 (1905), 766
(*Cybium cavalla* Cuvier).

2014. **Sierra cavalla** (Cuvier). *Sierra; Kingfish; Cavalla.*

Tropical Atlantic; Florida Keys and Charleston; Cape Cod; Africa; Brazil.

Cybium cavalla Cuvier, Régne Animal, ed. 2, II, 1829, 200; after *Guara-puca* of Marcgrave.

Cybium immaculatum Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 140 (191), no locality.

Cybium caballa Cuvier and Valenciennes, loc. cit., 187, Brazil.

Genus 676. SCOMBEROMORUS Lacépède.

Scomberomorus Lacépède, Hist. Nat. Poiss., III, 1802, 292 (*S. plumieri* Lacépède = *Scomber regalis* Bloch).

Polipturus Rafinesque, Anal. de la Nat., 1815, 89; substitute for *Scomberomorus*.

Chriomitra Lockington, Proc. Acad. Nat. Sci. Phila., XXXI, 1879, 133 (*C. concolor* Lockington).

2015. **Scomberomorus maculatus** (Mitchill). *Spanish mackerel; Spaniard; Spotted cybium; Bay mackerel; Spotted mackerel.*

East coast of America from Cape Ann to Brazil; common in Gulf of Mexico but not recorded from Cuba.

Scomber maculatus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 426, New York; not of Couch.

2016. **Scomberomorus sierra** Jordan and Starks. *Sierra.*

Pacific coast of tropical America.

Scomberomorus sierra Jordan and Starks, in Jordan, Proc. Calif. Acad. Sci., V, 1895, 428, Mazatlan, Mexico.

2017. **Scomberomorus regalis** (Bloch). *Sierra; Pintado; Cero; Sier; Siering; Searer; Spotted cero; Kingfish; Black-spotted Spanish mackerel.*

Cape Cod to Brazil; Cuba.

Scomber regalis Bloch, Naturgesch. ausl. Fische, VII, 1793, 38, Martinique; after a drawing by Plumier.

Scomberomorus plumieri Lacepédè, Hist. Nat. Poiss., III, 1802, 292, Martinique.

Cybium acervum Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 136 (180), Martinique; Cuba; St. Domingo.

2018. **Scomberomorus concolor** (Lockington). *Monterey Spanish mackerel.*

Coast of California.

Chriomitra concolor Lockington, Proc. Acad. Nat. Sci. Phila., XXXI, 1879, 133, Monterey, Calif.

✓ Family 161. SCOMBRIDÆ

Genus 677. SCOMBER Linnaeus.

Scomber Artedi, in Linnaeus, Syst. Nat., ed. X, 1758, 297 (*S. scombrus* Linnaeus).

Cordylus Gronow, Cat. Fish., 1854, 163 (*Scomber scombrus* L.).

2019. **Scomber scombrus** Linnaeus. *Common mackerel; Boston mackerel; Wauwhunnekessug; Caballa.*

North Atlantic; Norway and Labrador, south to Spain and Cape Hatteras.

Scomber scombrus Linnaeus, Syst. Nat., Ed. X, 1758, 297, Atlantic.

Scomber vernalis Mitchell, Trans. Lit. and Phil. Soc. N. Y., I, 1815, 423, Sandy Hook, N. J.

Genus 678. PNEUMATOPHORUS Jordan and Gilbert.

Pneumatophorus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 593 (*Scomber pneumatophorus* De la Roche = *Scomber colias* Gmelin).

"The family name *Cybidae* may be applied to the allies of *Scomberomorus* and *Cybium*. The name may be preferred to one derived from *Scomberomorus*, as the genus *Cybium* is also valid. *Cybium commersoni*, the Asiatic type of *Cybium* Cuvier, has but two gillrakers, while in *Scomberomorus* there are 8 to 11 below the angle. *Acanthocybium* (giant fishes with the reticulated gills of a swordfish) should constitute a separate family. The earliest of the many fossils of this type are mainly allies of *Scomberomorus*.

2020. *Pneumatophorus grex* (Mitchill). *Chub mackerel; Tinker mackerel; Easter mackerel; Thimble-eyed mackerel; Spanish mackerel of England.* The small mackerel called by the above names is very abundant on our Atlantic coast. It is very close to the similar species of Europe, *Scomber colias* Gmelin.

?*Scomber colias* Gmelin, Syst. Nat., 1788, 1329, Sardinia; after Cetti.

?*Scomber lacertus* Walbaum, Arctidi Pisc., 1792, 209, Sardinia; after Cetti.

?*Scomber pneumatophorus* De la Roche, Ann. Mus. Nat. Hist., XIII,

1809, 315, Balearic Islands.

?*Scomber macropthalmus* Rafinesque, Indice, 1810, 20 and 53, Palermo.

Scomber grex Mitchell, Trans. Lit. and Phil. Soc. N. Y., I, 1815, 422, New York.

?*Scomber maculatus* Couch, Mag. Nat. Hist., V, 1832, 22, England.

?*Scomber undulatus* Swainson, Nat. Hist. Fishes, Amphib., etc., II, 1839, 409, Sicily.

?*Scomber gracilis* Swainson, loc. cit., 410, Sicily.

Scomber dekayi Storer, Fishes Mass., 1867, 130, Massachusetts.

2021. *Pneumatophorus diego* (Ayres).

Coast of California; very abundant southward. Perhaps not separable from the Japanese species, *Scomber japonicus*; some small differences apparently permanent.

?*Scomber japonicus* Houttuyn, Verhand. Holl. Maatsch. Haarl., XX, 1782, 331, Japan.

?*Scomber auratus* Houttuyn, loc. cit., 333, Japan.

Scomber diego Ayres, Proc., Calif. Acad. Nat. Sci., I, 1856, 101, Santa Barbara, Calif.

2022. *Pneumatophorus australasicus* (Cuvier and Valenciennes).

North coast of Australia, from Moreton Bay, Queensland to Lord Howe Island, Hawaii and Socorro Island off the west coast of Mexico.

Scomber australasicus Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 49, Australia.

Family 162. KATSUWONIDÆ. The victor fishes

Genus 679. *KATSUWONUS* Kishinouye. *Oceanic bonito.*

Katsuwonus Kishinouye, Scombroïd Fish. Japan, Coll. Agric. Tokyo, VIII, 1923, 452 (*Scomber pelamis* L.).

2023. *Katsuwonus pelamis* (Linnaeus). *Oceanic bonito.*

Open Atlantic; rarely north to Cape Cod and Bermudas on the Atlantic coast.

Scomber pelamis Linnaeus, Syst. Nat., ed. X, 1758, 297, "in Pelago inter Tropicos."

Scomber pelamides Lacépède, Hist. Nat. Poiss., III, 1802, 14; after Linnaeus.

2024. *Katsuwonus vagans* (Lesson). *Aku; Pacific bonito.*

Pacific Ocean, Hawaii, southern California, and Japan; inordinately abundant in open sea.

Thynnus vagans Lesson, Voy. Coquille, Zool., II, 1828, 162, pl. 32.

Thynnus affinis Cantor, Cat. Malay Fishes, 1849, 106.

Genus 680. *EUTHYNNUS* Lütken.

Euthynnus Lütken, Jordan and Evermann, Fishes North and Mid. Amer., 1896, 868 (*Thynnus thunnina* Cuvier and Valenciennes = *Scomber alleteratus* Rafinesque).

This genus proves to be quite distinct from *Gymnosarda*, with which it has been confused. *Gymnosarda* is an Asiatic ally of *Sarda*.

2025. *Euthynnus alleteratus* (Rafinesque). *Little tunny; Bonito.*

Warm seas; north in Gulf Stream to Cape Cod; rather common at New York in summer: West Indies; Mediterranean; Japan.

Scomber alleteratus Rafinesque, Caratteri, 1810, 46, Palermo.

Thynnus leachianus Risso, Eur. Mérid., III, 1828, 414, Nice.

Scomber quadripunctatus Geoffrey St. Hilaire, Descr. Egypt. Poiss., 1827, pl. 24, fig. 3, Egypt.

Thynnus thunnina Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 76, (104), pl. 212, Mediterranean.

Thynnus brasiliensis Cuvier and Valenciennes, loc. cit., 80 (110), Brazil.

Thynnus brevipinnis Cuvier and Valenciennes, loc. cit., 81 (112), Mediterranean.

2026. *Euthynnus yaito* Kishinouye. *Kawakawa; Yaito.*
Japan; Hawaii; Pacific coast of Mexico.
Euthynnus yaito Kishinouye, Scombrid Fishes, 1923, 420, Japan.
2027. *Euthynnus lineatus* Kishinouye.
Pacific coast of Mexico.
Euthynnus lineatus Kishinouye, Sici. Gakko, No. 111, 1920, 113; and
in Scombrid Fishes, 1923, 430, fig. 2, Manzanillo, Mexico.
- Genus 681. *AUXIS* Cuvier. *Frigate mackerels.*
Auxis Cuvier, Règne Anim., ed. 2, II, 1829, 199 (*Scomber rochei* Risso =
Scomber thazard Lacépède).
2028. *Auxis rochei* (Risso). *Frigate mackerel.*
Open Atlantic, at times northward to Cape Cod; perhaps not separable
from *A. thazard* (Lacépède) of the South Seas.
Scomber rochei Risso, Ichth. Nice, 1810, 165, Nice.
Scomber bisus Rafinesque, Caratteri, 1810, 45, Palermo.

✓ Family 163. THUNNIDÆ. Tunnies and albacores

- Genus 682. *SARDA* Cuvier. *Bonitos.*
Sarda Cuvier, Règne Animal, ed. 2, II, 1829, 199 (*Scomber pelamys*
Brünich; not of Linnæus = *Scomber sarda* Bloch).
Pelamys Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 149
(*Scomber sarda* L.); name preoccupied.
Palamita Bonaparte, Ittiologia Generale, 1831, 107, substitute for
Pelamys.
Cretotroctes Gistel, Thierreich, X, 1848 (*Scomber sarda* L.); substitute
for *Pelamys*,

2029. *Sarda sarda* (Bloch). *Bonito; Bone jack; Bone-eater; Skipjack.*
Atlantic ocean, on both coasts, north to Cape Cod; very abundant.
Scomber sarda Bloch, Naturgesch. Ausl. Fische, VII, 1793, 44, Europe.
Scomber mediterraneus Bloch and Schneider, Syst. Ichth., 1801, 23,
Marseilles; after Brünich.
Scomber pelamitus Rafinesque, Caratteri, 1810, 44, pl. 2, Palermo.

2030. *Sarda lineolata*⁴⁵ (Girard). *California bonito; Skipjack.*
California, Patagonia and Japan; if *S. orientalis* is not distinct.
? *Pelamys chilensis* Cuvier and Valenciennes, Hist. Nat. Poiss., VIII,
1831, 118 (163), Valparaíso.
? *Pelamys orientalis* Temminck and Schlegel, Fauna Japon., Poiss., 1844,
99, pl. 52, Japan.
Pelamys lineolata Girard, Pac. R. R. Surv., X, 1858, 106, San Diego,
Calif.

2031. *Sarda velox* Meek and Hildebrand.
Panama.
Sarda velox Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV,
1923, 320, pl. 24, Panama market. Reported by Nichols and
Breder from Block Island, R. I.; probably an error.

- Genus 683. *THUNNUS* South. *Great tunnies.*
Thynnus Cuvier, Règne Anim., ed. 1, 1817, 313 (*Scomber thynnus* L.);
name preoccupied in Lepidoptera.
Thunnus South, Encyclop. Metropol., V, 1845, 620 (*Scomber thynnus*
Linnæus); substitute for *Thynnus*.
Albacora Jordan, Man. Vert., ed. V, 1888, 106 (*Scomber thynnus* L.).
Orycnus Cooper, Proc. Calif. Acad. Nat. Sci., III, 1863, 77 (*Scomber*
thynnus L.); not *Orycnus* Gill, 1861, the name originally a misprint
for *Orcynus*.

2032. *Thunnus saliens* Jordan and Evermann. *Leaping tuna.*
Eastern Pacific; lower California, north to mouth of Columbia River;
the type locality Santa Catalina Island.
Thunnus saliens Jordan and Evermann, Occ. Papers, Calif. Acad. Sci.,
XII, Sept. 30, 1926, 10, Santa Catalina Island, Calif.

⁴⁵ This species is well separated from *Sarda sardo* of the Atlantic. *Sarda orientalis* of Japan and Hawaii
is nearly related, and both may be inseparable from *S. chilensis* of Chile, as has been assumed by Jordan
and Evermann (1896).

2033. *Thunnus secundodorsalis* Storer.
 North Atlantic; New York to Labrador.
Thunnus secundodorsalis Storer, Hist. Fish. Mass., 1875, 163, Pl. XXII,
 fig. 1, Provincetown, Mass.
2034. *Thunnus subulatus* (Poey).
 West Indies; Cuba.
Orcynus subulatus Poey, Enumeratio, 1875, 71, Cuba; perhaps same
 as *T. coretta*.
2035. *Thunnus coretta* (Cuvier and Valenciennes). *Bonita of Martinique*.
 West Indies.
Tynnus coretta Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831,
 102, a young fish 11 inches long from Martinique.
- Genus 684. *GERMO* Jordan. *Long-finned albacores*.
Orcynus Cuvier, Règne Anim., Ed. I, 1817, 314 (*Scomber germo*
 Lacépède = *Scomber alalunga* Gmelin; not *Orcynus* Rafinesque, 1810,
 which is *Scomberoides*).
Gerмо Jordan, Proc. Acad. Nat. Sci. Phila., XL, 1888, 180 (*Scombrus*
alalonga Gmelin); substitute for *Orcynus*, preoccupied.
2036. *Germō alalunga* (Gmelin). *Albacore; Alilonghi*.
 Mediterranean; Atlantic; all warm seas.
Scomber alalunga Gmelin, Syst. Nat., I, 1788, 1330, Sardinia; mis-
 printed *Alatunga*.
Thynnus atlanticus Lesson, Voy. Coquille Zool., II, 1830, 165, Atlantic
 ocean.
Thynnus balleatus Cuvier and Valenciennes, Hist. Nat. Poiss., VIII,
 1831, 98 (136), coast of Brazil ("vis-à-vis la Trinité du Brazil, par les
 20° de latitude Australe").
2037. *Germō germo* (Lacépède).
 Pacific Ocean; California; Hawaii, Japan, and southward.
Scomber germo Lacépède, Hist. Nat. Poiss., II, 1800, 598, lat. 17° S.,
 long. 103° W.
Thynnus pacificus Cuvier and Valenciennes, op. cit., VIII, 1831, 96
 (133), lat. 26° S.; long. 103° W.
- Genus 685. *NEOTHUNNUS* Kishinouye. *Yellow-finned albacores*.
Neothunnus Kishinouye, Coll. Agric. Tokyo, VIII, 1923, 445 (*Thynnus*
macropterus Temminck and Schlegel).
2038. *Neothunnus catalinae* Jordan and Evermann. *Yellow-finned albacore*.
 Open Pacific; southern California, about Santa Catalina Island.
Neothunnus catalinae Jordan and Evermann, Occ. Paper No. 12, Calif.
 Acad. Sci., Sept. 30, 1926, 19, pl. 4, Santa Catalina Island, Calif.
2039. *Neothunnus albacores* (Bonnaterre). *West Indies long-finned albacore*.
 West Indies.
Scomber albacores Bonnaterre, Encyc. Ichth., 1788, 140, Jamaica; based
 on Sloane; not *albacora* Lowe.
Scomber sloani Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831,
 107 (148), Jamaica; after Sloane.
2040. *Neothunnus allisoni* (Mowbray). *Allison's tuna*.
 Coast of Florida.
Thunnus allisoni Mowbray, Copeia, No. 78, 1920, 9, with text fig.,
 Biscayne Bay, east coast of Florida.
- Family 164. **GEMPYLIDÆ**. Snake mackerel; Escolares
- Genus 686. *ESCOLAR* Jordan and Evermann.
Escolar Jordan and Evermann, in Goode and Bean, Oceanic Ichth.,
 1896, 519 (*Thyrsitops violaceus* Bean).
Bipinnula Jordan and Evermann, Fishes North and Mid. Amer., 1896,
 878 (*Thyrsitops violaceus* Bean); duplication of names accidental
 through noncorrection of proofs.
2041. *Escolar violaceus* (Bean).
 Atlantic coast of America in deep water; LaHave Bank.
Thyrsitops violaceus Bean, Proc. U. S. Nat. Mus., X, 1887, 513, LaHave
 Bank; perhaps the same as *Escolar niger*.

2042. *Escolar niger* (Poey).*Cuba.**Thyrsites niger* Poey, *Enumeratio*, 1875, 74, deep water off Cuba.Genus 687. *RUVENTTUS* Cocco. *Escolares.**Ruvettus* Cocco, *Giorn. Sci. Sicilia*, XLII, 1829, 2 (*R. pretiosus* Cocco).
Aplurus Lowe, *Trans. Zool. Soc. London*, II, 1841, 180 (*A. simplex* Lowe).*Acanthoderma* Cantraine, *Jour. Acad. Sci. Belles-Lettres Bruxelles*, X, 1837 (*A. temminckii*).2043. *Ruvettus pretiosus*⁴⁶ Cocco. *Escolar; Rovetto; Ruvetto; Chicolar; Oilfish; Scourfish; Plaintail.*

Mediterranean and westward to Cuba in Atlantic.

Ruvettus pretiosus Cocco, in *Giornale di Scienze per la Sicilia*, XLII, 1829, 21, Messina.*Tetragonurus simplex* Lowe, *Proc. Zool. Soc. Lond.*, 1833, 143, Madeira.
Ruvettus temminckii Cantraine, *Giorn. Sci. Ft. Litt. Pisa*, 1833, Malta; *fide* Poey.*Thyrsites acanthoderma* Lowe, *Proc. Zool. Soc. Lond.*, 1839, 78, Madeira.
Thyrsites scholaris Poey, *Memorias*, I, 1854, 372, pl. 32, fig. 1, Cuba.Genus 688. *EPINNULA* Poey.*Epinnula* Poey, *Memorias*, I, 1854, 369 (*E. magistralis* Poey).2044. *Epinnula magistralis* Poey. *Dómine.**Habana.**Epinnula magistralis* Poey, *Memorias*, I, 1854, 369, Habana.Genus 689. *NEALOTUS* Johnson.*Nealotus* Johnson, *Proc. Zool. Soc. Lond.*, 1865, 434 (*N. tripes* Johnson).2045. *Nealotus tripes* Johnson.

About Madeira in deep water; between the Bahamas and Bermudas.

Nealotus tripes Johnson, *Proc. Zool. Soc. Lond.*, 1865, 434, Madeira.Genus 690. *PROMETHICHTHYS* Gill. *Conejos.**Prometheus* Lowe, *Trans. Zool. Soc. Lond.*, II, 1841, 181, Madeira
(*Psempylus prometheus* Cuvier and Valenciennes); name preoccupied.*Promethichthys* Gill, *Mem. Nat. Acad. Sci.*, VI, 1893, 115 and 123
(*Prometheus atlanticus* Lowe = *Gempylus prometheus* Cuvier and Va-
lenciennes).2046. *Promethichthys prometheus* (Cuvier and Valenciennes). *Rabbitfish;*
Coelho; Conejo; Bermuda catfish.

Tropical Atlantic; Cuba; Bermuda.

Gempylus prometheus Cuvier and Valenciennes, *Hist. Nat. Poiss.*, VIII,
1831, 156 (213), pl. 222, St. Helena.Genus 691. *PTAX*⁴⁷ Jordan and Evermann.*Ptax* Jordan and Evermann, *Proc. Calif. Acad. Sci.*, 4th ser., XVI,
Apr. 27, 1927, 504 (*Dicrotus parvipinnis* Goode and Bean).2047. *Ptax parvipinnis* (Goode and Bean).

Western Atlantic in deep water.

Dicrotus parvipinnis Goode and Bean, *Oceanic Ichth.*, 1895, 201, fig. 212,
Gulf Stream at Albatross stations 2537, 2542, and 2601, at about
40° N., 70° W.Genus 692. *GEMPYLUS* Cuvier and Valenciennes. *Snake mackerels.**Gempylus* Cuvier and Valenciennes, *Hist. Nat. Poiss.*, VIII, 1831, 152
(207) (*G. serpens* Cuvier and Valenciennes).*Lemnisoma* Lesson, *Voyage Coquille*, Poiss., 1826 (1831), 160, the date
later than that of *Gempylus*, as the work of subsequent Cuvier and
Valenciennes is quoted by Lesson (*Lemnisoma thyrsitoides* Lesson).*Zyphothycus* Swainson, *Nat. Hist. Fish., Amphib., etc.*, II, 1839, 239
(*Gempylus coluber* Cuvier and Valenciennes).⁴⁶ A species very similar, regarded by Doctor Gudger as the same, is *Ruvettus pretiosus* Weber = *R. pacificus* Jordan and Jordan, from the tropical Pacific north to Hawaii.⁴⁷ From *rrax*, a rabbit, conejo, a Spanish name of *Promethichthys*. The genus is characterized by the number of fin rays—D. XXI, 11; A. II, 8—while in *Promethichthys* the rays are D. XVIII, 10-II; A. 10-II. *Dicrotus* has two lateral lines, the fin rays about as in *Promethichthys*.

2048. *Gempylus serpens* Cuvier and Valenciennes.

Deep seas of the Atlantic.

Gempylus serpens Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 152 (207), Martinique.

Gempylus ophidianus Poey, Memorias, II, 1861, 246, Cuba.

Family 165. TRICHIURIDÆ. Hairtails; Cutlassfishes

Genus 693. TRICHIURUS Linnaeus. Hairtails.

Trichiurus Linnaeus, Syst. Nat., Ed. X, 1758, 246 (*T. lepturus* Linnaeus); after Artedi.

Gymnogaster Gronow, Mus. Ichth., I, 1754, 17, No. 47, and Zooph., I, 1763, 136 (*Trichiurus lepturus* Linnaeus).

Enchelyopus Klein, in Bleeker, Versl. Akad. Amsterdam, XIV, 1862, 109 (*Trichiurus lepturus* Linnaeus); after Klein; not of Schneider nor of Gronow.

Lepturus (Artedi) Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 126 (*Trichiurus lepturus* Linnaeus, a revival of a pre-Linnaean name).

2049. *Trichiurus lepturus* Linnaeus. Cutlassfish; Scabbardfish; Silverfish; Sable; Savola; Hairtail; Ribbonfish; Silvery hairtail.

Western Atlantic north to New Jersey; West Indies.

Trichiurus lepturus Linnaeus, Syst. Nat., Ed. X, 1758, 246, America; after *Lepturus* of Artedi.

Trichiurus argenteus Shaw, Gen. Zool., IV, 1803, 90, pl. 12; after Linnaeus.

2050. *Trichiurus nitens* Garman.

Off Galapagos Islands. This or a similar species in the Gulf of California.

Trichiurus nitens Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 69, off Galapagos Islands, Albatross stations 3354, 3389.

Genus 694. APHANOPUS Lowe.

Aphanopus Lowe, Proc. Zool. Soc. Lond., Pt. VI, 1839, 79 (*A. carbo* Lowe).

2051. *Aphanopus minor* Collett.

East coast of Greenland.

Aphanopus minor Collett, Vidensk. Selsk. Forhandl. Christiania, No. 19, 1886, 3, east coast of Greenland at 65° N., 31° W.

Genus 695. EVOXYMETOPON Poey. Tirantes.

Evoxymetopon Poey, in Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 227 (*E. tenuatus* Poey).

2052. *Evoxymetopon tenuatus* Poey. Tirante; Tyrantfish.

Cuba.

Evoxymetopon tenuatus Poey, in Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 228, Cuba.

Genus 696. LEPIDOPUS Gouan. Frostfishes.

Lepidopus Gouan, Hist. Nat. Poiss., 1770, 107 and 185 (*gouani*).

Vandellius Shaw, Gen. Zool., IV, 1803, 199 (needless substitute for *Lepidopus* Gouan).

Scarcina Rafinesque, Caratteri, 1810, 20 (*S. argyrea* Rafinesque).

Xiphotheca Montagu, Werner. Mus., I, 1812, 82 (*Lepidopus caudatus* Euphrasen).

2053. *Lepidopus xantusi* Goode and Bean. Scabbardfish.

Avalon, Santa Catalina; Lower California north to Santa Catalina, Calif.; San José del Cabo, Cape San Lucas.

Lepidopus xantusi Goode and Bean, Oceanic Ichth., 1895, 519, Cape San Lucas.

Genus 697. BENTHODESMUS Goode and Bean.

Benthodesmus Goode and Bean, Proc. U. S. Nat. Mus., IV, 1881, 380 (*B. elongatus* Goode and Bean, not *Lepidopus elongatus* Clarke = *Benthodesmus atlanticus* Goode and Bean).

2054. *Benthodesmus atlanticus* Goode and Bean.

Western edge of Grand Banks of Newfoundland in 80 fathoms.

Benthodesmus elongatus Goode and Bean, Proc. U. S. Nat. Mus., IV, 1881, 380-383; not of Clarke.

Benthodesmus atlanticus Goode and Bean, Oceanic Ichth., 1895, 205, Grand Banks.

Series XIPHIIFORMES

Family 166. ISTIOPHORIDÆ. Sailfishes; Spearfishes

- Genus 698. **ISTIOPHORUS** Lacépède. *Sailfishes.*
Istiophorus Lacépède, Hist. Nat. Poiss., III, 1803, 374 (*I. gladifer*
Lacépède=*Scomber gladius* Broussonet).
Notistium Herrmann, Obs. Zool., 1804, 305 (*N. gladius* Herrmann).
Zanclurus Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 239
(*Histiophorus indicus* Cuvier and Valenciennes=*H. gladius*).
Histiophorus of authors, revised spelling.
2055. **Istiophorus americanus** (Cuvier and Valenciennes).
Brazil, and West Indies, north to Woods Hole.
Histiophorus americanus Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 303, Brazil; after Guebucu of Marcgrave.
2056. **Istiophorus wrighti** Jordan and Evermann.
West Indies northward to Florida.
Istiophorus wrighti Jordan and Evermann, Occ. Paper No. 12, Calif.
Acad. Sci., Sept. 30, 1926, 43, pl. 12, fig. 2, coast of Florida.
2057. **Istiophorus greyi** Jordan and Hill. *Mexican sailfish; Pacific volador.*
Pacific coast of Mexico, north to Cape San Lucas.
Istiophorus greyi Jordan and Hill, in Jordan and Evermann, Occ. Paper
No. 12, Proc. Calif. Acad. Sci., Sept. 30, 1926, 41, pl. 14, near Cape
San Lucas, Lower California.
2058. **Istiophorus volador** Jordan and Evermann. *Volador; Florida sailfish.*
About Florida Keys.
Istiophorus volador Jordan and Evermann, Occ. Paper No. 12, Calif.
Acad. Sci., Sept. 30, 1926, 46, pl. 12, fig. 1, Long Key, Fla.
Istiophorus maguirei Jordan and Evermann, loc. cit., 45, pl. 13, West
Indies; apparently young of *volador*.
- Genus 699. **MAKAIRA** Lacépède. *Marlinspike fishes; Spearfishes.*
Makaira Lacépède, Hist. Nat. Poiss., IV, 1803, 688 (*M. nigricans* Lac.).
2059. **Makaira mazara** (Jordan and Snyder). *Black marlin; Kurokajiki*
(*Black spearfish*); *Mazaara* (*real swordfish*); *Kuroka* (*black marlin*).
Pacific Ocean, from Japan to Hawaiian Islands and the Channel
Islands, Calif.
Tetrapturus mazara Jordan and Snyder, Jour. Coll. Sci. Imp. Univ.
Tokyo, XV, May 20, 1901, 4, 305, Misaki, Japan.
2060. **Makaira ampla** (Poey). *Aguja de Casta.*
West Indies.
Tetrapturus amplus Poey, Memorias, II, 1861, 243, Habana.
2061. **Makaira marlina** Jordan and Evermann. *Giant black marlin.*
Southern California, south to Cape San Lucas.
Makaira marlina Jordan and Evermann, Occ. Paper No. 12, Calif.
Acad. Sci., Sept. 30, 1926, 59, pl. 17, off Cape San Lucas, Lower
California.
2062. **Makaira mitsukurii** (Jordan and Snyder). *Marlinspike fish; Barred*
marlin; Makajiki or true spearfish of Japan.
Coasts of Japan; Hawaii; southern California; abundant off Santa
Catalina.
? *Tetrapturus indicus* Cuvier and Valenciennes, Hist. Nat. Poiss., VIII,
1831, 286, Sumatra, on a figure by Banks.
? *Histiophorus audax* Philippi.
Tetrapturus mitsukurii Jordan and Snyder, Jour., Coll. Sci. Imp. Univ.
Tokyo, XV, pt. 2, 1901, 304, Pl. XVI, fig. 5, fishwell at Misaki.
2063. **Makaira holei** Jordan and Evermann. *Striped marlin.*
Off Pacific coast of Mexico, north to Cape San Lucas.
Makaira holei Jordan and Evermann, Occ. Papers No. 12, Calif. Acad.
Sci., Sept. 30, 1926, 63, pl. 19, fig. 1, off Cape San Lucas, Lower
California.

✓ Family 167. XIPHIIDÆ. Swordfishes

Genus 700. XIPHIAS Linnæus. *Swordfishes.*

Xiphias Linnæus, Syst. Nat., Ed. X, 1758, 248 (*X. gladius* Linnæus).
Phæthonichthys Nichols, Amer. Mus. Nov., XIX, I, 1923 (*Phæthonichthys tuberculatus* Nichols; a young swordfish).

- 2064. Xiphias gladius** Linnæus. *Common swordfish; Broadbill; Espada; Espadón emperador.*

Open sea, Atlantic and Pacific Oceans, on both coasts; Cuba to Cape Breton and Newfoundland Banks; Pacific off Santa Barbara Islands, Hawaii, Japan, San Diego.

Xiphias gladius Linnæus, Syst. Nat., Ed. X, 1758, 248, Europe.
Xiphias rondeletii Leach, in Wern. Mem., II, 1818, 58, pl. 2, fig. 1, Firth of Forth.

Xiphias imperator Schneider, ed. Bloch, Syst. Ichth., 1801, 93, Pl. XXI, Mediterranean, after an erroneous figure of Duhamel.

Phæthonichthys tuberculatus Nichols, Amer. Mus. Nat. Hist. Novitates, XIX, pl. 1, Oct. 19, 1923; very young of *X. gladius* L.

Series LUVARIFORMES

Family 168. LUVARIDÆ

Genus 701. LUVARUS Rafinesque.

Luvarus Rafinesque, Caratteri, 1810, 22 (*L. imperialis* Rafinesque).

Diana Risso, Europe Méridionale, 1826, 267 (*D. semilunata* Risso).

Ausonia Risso, op. cit., 341 (*A. civieri* Risso).

Proctostegus Nardo, Diaro, Chem. et Hist. Nat. Ticino, I, 1827, 18, 42 (*P. proctostegus* Nardo).

Astrodermus (Bonelli) Cuvier, Règne Anim., ed. 2, II, 1829, about 215 (*A. guttatus* Bonelli MS).

Scafaria Gistel, Thierreich, VIII; 1848 (*Ausonia civieri* Risso).

- 2065. Luvarus imperialis** Rafinesque.

Open sea; once taken at San Pedro, Calif. (W. F. Thompson).

Luvarus imperialis Rafinesque, Caratteri, 1810, 22.

Diana semilunata Risso, Europe Mérid., 1826, 267.

Ansonia civieri Risso, loc. cit., 341.

Proctostegus proctostegus Nardo, Diaro, Chem. et Hist. Nat. Ticino, I, 1827, 18, 42.

Astrodermus guttatus Bonelli, in Cuvier, Règne Anim., ed. 2, II, 1829, 215 ca.

Scafaria Gistel, Thierreich, VIII, 1848.

Series CORYPHÆNIFORMES

Family 169. CORYPHÆNIDÆ. Dolphins; Dorados

Genus 702. CORYPHÆNA Linnæus. *Dorados.*

Coryphæna Linnæus, Syst. Nat., Ed. X, 1758, 261 (*C. hippurus* L.).

Lamprugus Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 317 (*Scomber pelagicus* L.; female).

- 2066. Coryphæna hippurus** Linnæus. *Common dolphin; Dorado; Dourade.*

Pelagic to Cape Cod and northward, South Carolina to Texas; also throughout the mid-Pacific.

Coryphæna hippurus Linnæus, Syst. Nat., ed. X, 1758, 261, open seas.

Scomber pelagicus Linnæus, loc. cit., 299, no locality.

Coryphæna fasciolata Pallas, Spicil. Zool., VIII, 1770, 23, Amboina.

Coryphæna imperialis Rafinesque, Caratteri, 1810, 33, Sicily.

Coryphæna immaculata Agassiz in Spix, Pisc. Brasil., 1829, 111, pl. 56, Atlantic, off Brazil.

Coryphæna marcgravii Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 223 (301), America.

Coryphæna suerii Cuvier and Valenciennes, loc. cit., 224 (302), Philadelphia.

Coryphæna dorado Cuvier and Valenciennes, loc. cit., 225 (303), Brazil.

Coryphæna dolfin Cuvier and Valenciennes, loc. cit., 226 (305), Antilles.

- Coryphæna virgata* Cuvier and Valenciennes, loc. cit., 228 (308), Martinique.
Coryphæna argyrurus Cuvier and Valenciennes, loc. cit., 232 (314), Sea of Coromandel.
Coryphæna virmingii Cuvier and Valenciennes, loc. cit., 233 (315), seas of India.
Lampugus sculus Cuvier and Valenciennes, loc. cit., 240 (323), Sicily.
Coryphæna scomberoides Cuvier and Valenciennes, loc. cit., 234 (315), South Sea.
Ectenias brunneus Jordan and Thompson, Mem. Carnegie Mus., VI, 1914, 241, Japan; young.
2067. *Coryphæna equisetis* Linnæus. *Small dolphin.*
 Open Atlantic and Pacific; rare in the West Indies. Recently taken in some numbers off coast of southern California.
Coryphæna equisetis Linnæus, Syst. Nat., Ed. X, 1758, 261 (misprinted *equiselis*), high seas; after Osbeck.
Coryphæna aurata Rafinesque, Caratteri, 1810, 33, Sicily.
Coryphæna lessonii Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 228 (307), India.
Lampugus punctulatus Cuvier and Valenciennes, loc. cit., 242 (429), Atlantic at the Equator.
Lampugus neapolitanus Cuvier and Valenciennes, loc. cit., 241 (325), Naples; after Bloch.
Coryphæna azorica Cuvier and Valenciennes, loc. cit., 227 (306), Azores.

Series BRAMIFORMES

✓

Family 170. BRAMIDÆ. Sea breams

- Genus 703. **BRAMA** Bloch and Schneider. *Pomfret.*
Brama Bloch and Schneider, Ed. Bloch, Syst. Ichth., 1801, 98 (*Sparus raii* Bloch).
Lepodus Rafinesque, Caratteri, 1810, 54 (*L. saragus* Rafinesque = *Sparus raii* Bloch).
2068. **Brama raii** (Bloch). *Sea bream; Pomfret; Catagnole; Rondanin.*
 Open seas; Europe; Faroe Islands; Bermuda; Grand Banks; Pacific coast from Santa Catalina to Puget Sound; Japan; reported recently from Cape Flattery.
Brama marina cauda forcipata Ray, Syn. Meth. Piscium, 1713, 115, Middelburg.
Sparus raii Bloch, Naturgesch. Ausl. Fische, V, 1791, 95; after Ray, etc.
Sparus castaneola Shaw, Gen. Zool., IV, 1803, 424; after Lacépède.
Sparus niger Turton, British Fauna, 1807, 98, Swansen.
Lepodus saragus Rafinesque, Caratteri, etc., 1810, 54, Palermo.
Brama dussumieri Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 219 (294), Indian Sea, long. 85° E.
? *Brama orcinii* Cuvier and Valenciennes, loc. cit., 220 (295), Indian Sea, long. 85° E.
? *Brama chilensis* Gay, Hist. Chile, Peces, 1843, 218, Chile.
2069. **Brama agassizii** Poey.
 Cuba; rare.
Brama agassizii Poey, Memorias, II, 1861, 204, Cuba.
2070. **Brama brevoortii** Poey.
 Cuba; rare.
Brama brevoortii Poey, Memorias, II, 1861, 206, Cuba.
- Genus 704. **TARACTES** Lowe.
Taractes Lowe, Proc. Zool. Soc. Lond., 1843, 82 (*T. asper* Lowe; probably young of *Taractes longipinnis*).
Argo Döderlein, Fische Japans, II, Denkschr. Acad. Wiss. Wien, XLVIII, 1883, 34 (*Argo steindachneri* Döderlein); name preoccupied.
2071. **Taractes saussurii** (Lunel).
 Habana.
Brama saussurii Lunel, Revue du Genre *Brama*, in Mém. Soc. Phys. Hist. Nat. Genève, XVIII, 1865, 185, tab. 2, Cuba.

Family 171. STEINEGERIIDÆ

Genus 705. STEINEGERIA Jordan and Evermann.

Steinegeria Jordan and Evermann, Proc. U. S. Nat. Mus., IX, 1886, 467 (*S. rubescens* Jordan and Evermann).

2072. Steinegeria rubescens Jordan and Evermann.

Gulf of Mexico in deep water; one specimen known.

Steinegeria rubescens Jordan and Evermann, Proc. U. S. Nat. Mus., IX, 1886, 467, snapper banks off Pensacola, Fla.

Family 172. PTERACLIDÆ

Genus 706. PTERACLIS Gronow.

Pteraclis Gronow, Acta Helvetica, VII, 1772, 43 (*Coryphæna velifera* Pallas).

Pteridium Scopoli, Introd. Hist. Nat., 1777, 454 (*Coryphæna velifera* Pallas).

Oligopodus Lacépède, Hist. Nat. Poiss., II, 1800, 511 (*Coryphæna velifera* Pallas).

Pteraclidus Rafinesque, Analyse de la Nat., 1815, 82, substitute for *Oligopodus* Lacépède.

Oligopodes Risso, in Cuvier, Règne Anim., ed. 1, 1817, 328 (*l'Oligopode noir* Risso).

Oligopus Risso, Europe Mérid., 1826, 105, substitute for *Oligopodus*.

2073. Pteraclis carolinus Cuvier and Valenciennes.

Coast of South Carolina.

Pteraclis carolinus Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 272 (368), off coast South Carolina.

?*Pteraclis trichipterus* Cuvier and Valenciennes, loc. cit., 272 (367), locality unknown.

Genus 707. CENTROPHOLIS Hilgendorf.

Centropholis Hilgendorf, Sitzber. Naturf. Freunde, Berlin, 1878, 135 (*C. petersi* Hilgendorf).

2074. Centropholis goodei Jordan.

Gulf Stream.

Pteraclis carolinus Goode and Bean, Oceanic Ichth., 212, Pl. LIX, fig. 218, Gulf Stream; not of Linnaeus.

Centropholis goodei Jordan, Ann. Carnegie Mus., IX, 1915, 333, Gulf Stream.

Series STROMATEIFORMES

Family 173. STROMATEIDÆ. Fiatolas

Genus 708. PEPRILUS Cuvier.

Rhombus Lacépède, Hist. Nat. Poiss., II, 1800, 321 (*Chætodon alepidotus* Linnaeus); preoccupied in mollusks.

Peprilus Cuvier, Règne Anim., ed. 2, II, 1829, 213 (*Stromateus longipinnis* Mitchell = *Stromateus paru* Linnaeus).

2075. Peprilus paru (Linnaeus). *Harvestfish; Poppyfish; Rudderfish.*

South Atlantic coast of United States; West Indies; Cape Cod to Jamaica and Brazil.

Stromateus paru Linnaeus, Syst. Nat., Ed. X, 1758, 248, Jamaica; based on Sloane.

Chætodon alepidotus Linnaeus, op. cit., Ed. XII, 1766, 460, Charleston, S. C.

?*Sternopyx gardenii* Bloch and Schneider, Syst. Ichth., 1801, 494, Carolinas; after Linnaeus.

Stromateus longipinnis Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 366, New York Bay.

Genus 709. PALOMETA Jordan and Evermann.

Palometa Jordan and Evermann, Fishes North and Mid. Amer., 1896, 966 (*Stromateus palometa* Jordan and Bollman).

2076. Palometa simillima (Ayres). *California pampano.*

Pacific coast of United States, Puget Sound to San Diego.

Poronotus simillimus Ayres, Proc. Calif. Acad. Nat. Sci., 1st ser., II, 1860, 84, San Francisco.

- 2077. *Palometa media* (Peters).** *Palometa*.
Pacific coast of North America, Mazatlan to Panama.
Stromateus medius Peters, Berl. Monatsb., 1869, 707, Mazatlan, Mexico.
- 2078. *Palometa palometa* (Jordan and Bollman).**
Pacific Ocean, off Colombia.
Stromateus palometa Jordan and Bollman, Proc. U. S. Nat. Mus., XII,
1889, 156, off coast of Colombia, lat. $8^{\circ} 16' 30''$ N., long. $79^{\circ} 37' 45''$ W., at Albatross station 2804.
- 2079. *Palometa snyderi* (Gilbert and Starks).**
Panama.
Peprilus snyderi Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904,
87, Panama.
- Genus 710. *PORONOTUS* Gill.**
Poronotus Gill, Cat. Fish. E. Coast N. Amer., Sup. to Proc. Acad. Nat. Sci. Phila., XIII, 1861, 35 (*Stromateus triacanthus* Peck).
- 2080. *Poronotus triacanthus* (Peck).** *Dollarfish; Harvestfish; Butterfish; Lafayette.*
Nova Scotia to Florida; common.
Stromateus triacanthus Peck, Mem. Amer. Acad., II, 1800, pt. 2, 48,
pl. 2, fig. 2, Piscataqua River, N. H.
Stromateus cryptos Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 365,
New York Bay.
- Genus 711. *EUCROTUS* Bean (B. A.).**
Eucrotus Bean, Proc. Biol. Soc. Wash., XXV, 1912, 123 (*E. ventralis* Bean).
- 2081. *Eucrotus ventralis* Bean.**
Bermudas.
Eucrotus ventralis Bean, Proc. Biol. Soc. Wash., XXV, 1912, 125,
Bermudas.

Family 174. CENTROLOPHIDÆ. Rudderfishes

- Genus 712. *CENTROLOPHUS* Lacépède.** *Black ruffs.*
Centrolophus Lacépède, Hist. Nat. Poiss., IV, 1803, 441 (*Perca nigra* Gmelin = *Coryphæna pomphilus* L.).
Acentrolophus Nardo, Prodromus. Adriat. Ichth. Giorn. Fisica di Pavia, sp. 62, 1827, (A. *maculatus* Nardo = *Centrolophus niger* Lacépède, the name *Centrolophus* deemed inappropriate).
Gymnocephalus Cocco, Nuov. Ann. Sci. Nat., II, 1838, 26 (*G. messinensis* Cocco); name preoccupied.
Pompilus Lowe, Suppl. Fish Madeira, Proc. Zool. Soc. London, VII, 1839, 81 (*Coryphæna pomphilus* L.).
- 2082. *Centrolophus niger* (Gmelin).** *Blackfish; Black ruffe; Borlase.*
Coasts of southern Europe; once taken at Dennis, Mass.
Perca nigra Gmelin, Syst. Nat., 1788, 1321, Cornwall.
Centrolophus liparis Risso, Eur. Mérid., III, 1826, 337, Nice, Italy.
Centrolophus morio Lacépède in Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 254 (342), Mediterranean.
Centrolophus pomphilus Cuvier and Valenciennes, loc. cit., 247 (334), pl. 269; not of Linnaeus.
Acentrolophus maculosus Nardo, Prodr. Ichth. Adriat., No. 62, Isis, 1827, Adriatic.
Gymnocephalus messinensis Cocco, Nuovi Ann. Sci. Nat., II, 1838, 26, Messina.
- Genus 713. *PALINURICHTHYS* Bleeker.** *Black rudderfishes.*
Palinurus DeKay, N. Y. Fauna: Fishes, 1842 (*Coryphæna perciformis* Mitchell); name preoccupied in Crustacea.
Palinurichthys Bleeker, Enum. Spec. Pisc. Arch. Ind., in Acta Soc. S. Indo-Neerl., VII, November, 1859, 22 (*Coryphæna perciformis*, Mitchell).
Palinurichthys Gill, Proc. Acad. Nat. Sci. Phila., XII, January, 1860, 20 (*Coryphæna perciformis* Mitchell).
Pammelas Günther, Cat., II, June, 1860, 485 (*Coryphæna perciformis* Mitchell).

- 2083.** *Palinurichthys perciformis* (Mitchill). *Rudderfish; Logfish; Snipnose mullet; Black rudderfish*
 Atlantic coast of North America from Cape Hatteras to Maine; once taken in Cornwall.
Coryphena perciformis Mitchill, Amer. Month. Mag., II, 1818, 244, New York Harbor.
Pimelepterus cornubiensis Cornish Zoologist, IX, 1874, 4255, Penzance, Cornwall (a stray specimen floated across in a fish box).
- Genus 714. SCHEDOPHILUS Cocco.**
Schedophilus Cocco, Giorn. Innom. Messina, Ann., III, 1834, No. 7, 57, (*S. medusophagus* Cocco).
Crius Valenciennes, Ichth. Iles Canaries, II, 1844, 43 (*Crius bertheloti Valenciennes* = *Schedophilus medusophagus* Cocco).
- 2084.** *Schedophilus heathi* Gilbert.
 Monterey Bay.
Schedophilus heathii Gilbert, Proc. Calif. Acad. Sci., ser. 3, Zool., III, 1904, 260, Pl. XXVI, Pacific Grove, Calif.
- Genus 715. ICICHTHYS Jordan and Gilbert.**
Icichthys Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 305 (*I. lockingtoni* Jordan and Gilbert).
- 2085.** *Icichthys lockingtoni* Jordan and Gilbert.
 Off California, in deep water; a surface swimmer, hovering under jelly-fishes; known from off Monterey and San Francisco.
Icichthys lockingtoni Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 305, off San Francisco.
- Family 175. NOMEIDÆ. Man-of-war fishes**
- Genus 716. NOMEUS Cuvier.**
Nomeus Cuvier, Règne Anim., ed. 1, II, 1817, 315 (*Gobius gronovii* Gmelin).
- 2086.** *Nomeus gronovii* (Gmelin). *Portuguese man-of-war fish; Harder; Pastor.*
 Tropical parts of the Atlantic and Indian Oceans; Sargasso Sea; Florida; Bermuda; Woods Hole, Mass.; rocky pools at Panama.
Gobius gronovii Gmelin, Syst. Nat., Ed. XIII, 1788, 1205, tropical America; after Gronow.
Electriss mauritii Bloch and Schneider, Syst. Ichth., 1801, 66, Brazil.
Nomeus maculosus Bennett, Proc. Comm. Zool. Soc., I, 1831, 146, East Indies.
Nomeus oxyurus Poey, Memorias, II, 1861, 236, Habana.
- Genus 717. PSENES Cuvier and Valenciennes.**
Psenes Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 193 (259), (*P. cyanophrys* Cuvier and Valenciennes).
- 2087.** *Psenes cyanophrys* Cuvier and Valenciennes.
 Open sea; Atlantic, Pacific, and Indian Oceans; Jamaica and Martinique.
Psenes cyanophrys Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 193-260, pl. 265, New Ireland.
Psenes javanicus Cuvier and Valenciennes, loc. cit., 196 (264), Batavia.
Psenes auratus Cuvier and Valenciennes, loc. cit., 197 (264), Gulf of Bengal.
Psenes fuscus Guichenot, Mem. Soc. Sci. Nat. Cherbourg, XII, 1866, 138, Madagascar.
Cubiceps multiradiatus Günther, Proc. Zool. Soc. Lond., 1871, 661, pl. 61, Manado.
- 2088.** *Psenes maculatus* Lütken.
 Open Atlantic.
Psenes maculatus Lütken, Spolia Atlantica, 1880, 110, open Atlantic, lat. 39° N., long. 25° S. and between 34° and 27° W. in 600 to 700 fathoms.
- 2089.** *Psenes regulus* Poey.
 Coasts of Cuba.
Psenes regulus Poey, Synopsis, 1868, 375, Cuba.

2090. *Psenes pellucidus* Lütken.
 Gulf Stream at 32° 24' N., 76° 55' W., in 528 fathoms.
Psenes pellucidus Lütken, Spolia Atlantica, 1880, 516 (108), with fig.,
 Strait of Surabaja.
2091. *Psenes edwardsii* Eigenmann.
 Off Rhode Island.
Psenes edwardsii Eigenmann, Bull. U. S. Fish Com., XXI, 1901 (1902),
 35, fig.; taken from under a medusa, 30 miles south of Newport, R. I.
2092. *Psenes chapmani* Fowler.
 Open Atlantic, Cape Verde Islands, northwest of the Lesser Antilles.
Psenes chapmani Fowler, Proc. Acad. Nat. Sci. Phila., LVIII, 1906,
 120; open Atlantic, lat. 24° 21' N., long. 34° 32' W.
2093. *Psenes pacificus* Meek and Hildebrand.
 Panama.
Psenes pacificus Meek and Hildebrand, Field Mus. Nat. Hist., Zool.,
 XV, 1925, 409, Chame Point, Panama.

Family 176. TETRAGONURIDÆ. Squaretails

- Genus 718. **TETRAGONURUS** Risso.
Tetragonurus Risso, Ichth. Nice, 1810, 247 (*T. cuvieri* Risso).
Ctenodax Macleay, Proc. Linn. Soc. New South Wales, 1885, 716 (*C. williamsoni* Macleay).

2094. *Tetragonurus cuvieri* Risso. *Squaretail; Escolar de natura; Courpata; Sea raven.*
 Open seas; Nice; Toulon and Marseilles; Madeira; once taken off Santa Catalina Island, Calif. (W. F. Thompson), and once at Woods Hole, Mass.; these possibly different species.
Tetragonurus cuvieri Risso, Ichth. Nice, 1810, 347, Nice.
Tetragonurus atlanticus Lowe, Fishes Madeira, pt. 5, 1860 (1843), 129, Pl. XIX, Madeira.

Family 177. ICOSTEIDÆ. Ragfishes

- Genus 719. **ICOSTEUS** Lockington.
Icosteus Lockington, Proc. U. S. Nat. Mus., III, 1880, 63 (*I. ænigmaticus* Lockington).
Schedophilopsis Steindachner, Ichth. Beitr., XI, 1881, 4 (*S. spinosus* Steindachner = *Icosteus ænigmaticus* Lockington).

2095. *Icosteus ænigmaticus* Lockington. *Ragfish.*
 Open sea off the coasts of California, Oregon, and Washington; scarce.
Icosteus ænigmaticus Lockington, Proc. U. S. Nat. Mus., III, 1880, 63, off San Francisco.
Schedophilopsis spinosus Steindachner, Ichth. Beitr., XI, 1881, 4, with figure, off San Francisco.

Family 178. ACROTIDÆ

- Genus 720. **ACROTUS** Bean.
Acrotus Bean, Proc. U. S. Nat. Mus., X, 1887, 631 (*A. willoughbyi* Bean).

2096. *Acrotus willoughbyi* Bean.
 Off coast of Washington, Whidbey Island, Puget Sound, south to San Pedro, Calif. (W. F. Thompson), in rather deep water.
Acrotus willoughbyi Bean, Proc. U. S. Nat. Mus., X, 1887, 631, Quinault Agency, Wash.

✓ Family 179. CARANGIDÆ. Cavallas; Pampanos

The *Carangidae*, as heretofore understood, may be naturally divided into two families, the *Seriolidæ* (*Seriola*, *Elagates*, *Naukrates*, etc.) differing strongly in lacking the bony arch behind the abdominal cavity, this being made up of enlarged and ankylosed hemals, its character somewhat different in different genera but developed in all *Carangidae* in the narrow sense.

- Genus 721. DECAPTERUS Bleeker.** *Mackerel scads.*
Decapterus Bleeker, *Naturk. Tijdsch.*, VIII, 1855, 417 (*Caranx kurra* Bleeker).
Eustomatodus Gill, *Proc. Acad. Nat. Sci. Phila.*, XIV, 1862, 261 (*Decapterus kurroides* Bleeker).
Gymnepiagnathus Gill, loc. cit., 261 (*Decapterus macrosoma* Bleeker).
Evepigymnus Gill, loc. cit., 261 (*Decapterus hypodus* Gill).
- 2097. Decapterus punctatus** (Agassiz). *Scad; Round robin; Cigarfish; Quiaquia; Dotted scad.*
Cape Cod to Brazil, coasts of Florida and the West Indies.
Caranx punctatus Agassiz, in Spix, *Pisc. Brasil.*, 1829, 108, pl. 56a, fig. 2, Brazil.
Caranx aureus Risso, in Cuvier and Valenciennes, *Hist. Nat. Poiss.*, IX, 1833, 25 (33), Mediterranean.
- 2098. Decapterus sanctæ-helenæ** (Cuvier and Valenciennes).
Tropical Atlantic; Cuba.
Caranx sanctæ-helenæ Cuvier and Valenciennes, *Hist. Nat. Poiss.*, IX, 1833, 28 (37), St. Helena.
- 2099. Decapterus hypodus** Gill.
Cape San Lucas, Lower California.
Decapterus hypodus Gill, *Proc. Acad. Nat. Sci. Phila.*, XIV, 1862, 262, Cape San Lucas, Lower California.
- 2100. Decapterus macarellus** (Cuvier and Valenciennes). *Mackerel scad; Antonino; Cigarfish.*
West Indies, northward to Cape Cod.
Caranx macarellus Cuvier and Valenciennes, *Hist. Nat. Poiss.*, IX, 1833, 30 (40), Martinique.
- Genus 722. TRACHURUS Rafinesque.** *Saurels.*
Trachurus Rafinesque, *Caratteri*, 1810, 41 (*Scomber trachurus* Linnaeus).
- 2101. Trachurus symmetricus** (Ayres). *Horse mackerel.*
Coast of California.
Caranx symmetricus Ayres, *Proc. Calif. Acad. Nat. Sci.*, I, 1855, 62, Bay of San Francisco.
- 2102. Trachurus trachurus** (Linnaeus). *Saurel; Gascon.*
North Atlantic, chiefly on the northern coast of Europe, south to Spain and Naples; Newport, R. I.; Pensacola.
Scomber trachurus Linnaeus, *Syst. Nat.*, Ed. X, 1758, 298, Mediterranean.
?*Caranxomorus plumieri* Lacépède, *Hist. Nat. Poiss.*, III, 64, pl. 11, scarcely identifiable, from a figure by Plumier, supposed to have been made in Martinique.
Caranx semispinosus Nilsson, *Prodr. Ichthyol. Scand.*, 1832, 84, Scandinavia; the northern form usually called *trachurus*, the valid name for the European form.
Trachurus europæus Gronow, *Cat.*, Ed. Gray, 1854, 125, seas of Europe.
Trachurus linnæi Malm, *Bohusläns Fauna*, 421, Bohuslän, Sweden.
- 2103. Trachurus japonicus** (Schlegel).
Coast of Japan, east of Lower California, the Mexican specimens few and not surely identified; may be *T. nova-zealandica* Richardson.
Caranx trachurus japonicus Schlegel, *Faun. Japon.*, Poiss., 1850, 109, pl. 59, fig. 1, Japan.
- 2104. Trachurus lathami** Nichols.
Coast of Long Island (a stray from the Gulf Stream).
Trachurus lathami Nichols, *Bull. Amer. Mus. Nat. Hist.*, XLII, 1920, 479, Long Island.
- Genus 723. TRACHUROPS** ⁴⁸ Gill.
Trachurops Gill, *Proc. Acad. Nat. Sci. Phila.*, XIV, 1862, 238, 261, 431 (*Scomber crumenophthalmus* Bloch).

⁴⁸The genus *Selar* Bleeker was based on four species, the first named being *Caranx hasselti*, which is a species of the modern genus *Atule* Jordan and Jordan. It was first restricted to *Scomber boops* Bleeker, a species of *Trachurops*; but this restriction can not stand as *boops* was not mentioned by Bleeker in connection with *Selar*. *Caranx hasselti*—affinis of the south Pacific must therefore be the type of *Selar*.

- 2105. *Trachurops crumenophthalmus* (Bloch). *Goggler; Big-eyed scad; Goggle-eye jack; Chicharro; Horse-eyed jack.***
 Atlantic coast of the United States, Central America, South America, and Africa; West Indies; probably not in the Pacific.
Scomber crumenophthalmus Bloch, Naturgeschich. Ausl. Fische, VII, 1793, 77, Acara in Guinea.
Scomber plumieri Bloch, loc. cit., 79, Antilles.
Scomber balantiophthalmus Bloch and Schneider, Syst. Ichth., 1801, 29, Guinea.
Caranx macrophtalmus Agassiz, Spix, Pisc. Brasil., 1829, 107, Brazil.
- 2106. *Trachurops brachychira* Gill. *Akule; Atule.***
 Pacific Ocean, north to Panama, and the Gulf of California and Hawaii.
Trachurops brachychirus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 261, Cape San Lucas.
- Genus 724. *HEMICARANX* Bleeker.**
Alepes Swainson, Classn. Anim., II, 1839, 248 (*Alepes melanoptera* Swainson); identification uncertain.
Hemicaranz Bleeker, Vesl. Kon. Ak. Wet., XIV, 1862, 135 (*H. marginatus* Bleeker).
Carangops Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 238 (*Caranx heteropygus* Poey).
- 2107. *Hemicaranz amblyrhynchus* (Cuvier and Valenciennes).**
 Cape Hatteras to Brazil; West Indies; Carolina and Florida.
Caranx amblyrhynchus Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 76 (100), pl. 248, Brazil.
Caranx heteropygus Poey, Memorias, II, 1861, 344, Cuba.
- 2108. *Hemicaranz falcatus* (Holbrook).**
 Charleston, S. C.
Caranx falcatus Holbrook, Ichth. South Carolina, 1855, 92, pl. 13, fig. 2, Charleston, S. C.
- 2109. *Hemicaranz atrimanus* (Jordan and Gilbert).**
 Pacific coast of tropical America; Panama.
Caranx atrimanus Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 308, Panama.
- 2110. *Hemicaranz furthii* (Steindachner).**
 Panama.
Caranx furthii Steindachner, Ichth. Beitr., IV, 1875, 12, Panama.
- 2111. *Hemicaranz leucurus* (Günther).**
 Pacific coast of tropical America; Panama.
Caranx leucurus Günther, Proc. Zool. Soc. Lond., 1864, 24, Panama.
- 2112. *Hemicaranz zelotes* Gilbert.**
 Panama.
Hemicaranz zelotes Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2845, Panama.
- 2113. *Hemicaranz rhombooides* Meek and Hildebrand.**
 Atlantic coast of Isthmus of Panama.
Hemicaranz rhombooides Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1925, 345, Fox Bay, Colon, Panama.
- Genus 725. *ELAPHOTOXON* ^a Fowler.**
Elaphotoxon Fowler, Proc. Acad. Nat. Sci. Phila., LVII, 1905, 76 (*Scomber ruber* Bloch).
- 2114. *Elaphotoxon ruber* (Bloch). *Cibi mancho; Carbonero; Cibi; Green jack.***
 West Indies and coast of the Carolinas.
Scomber ruber Bloch, Naturgeschich. Ausl. Fische, VII, 1793, 75, St. Croix.
Caranx blochii Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 52 (69), St. Croix.
Caranx iridinus Poey, Memorias, II, 1861, 226, Cuba.

^aThis genus is near *Caranx*, the body rather elongate, the teeth in few series, without canines; the arch of the lateral line moderate, more than half the straight part; the dorsal and anal fins anteriorly but little elevated, maxillary not narrowed.

- 2115. *Elaphotoxon bartholomaei* (Cuvier and Valenciennes).** *Yellow jack; Cibi amarilla.*
 West Indies, northward to Florida and North Carolina; Cuba.
Caranx bartholomaei Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 75 (100), St. Bartholomew.
Caranx cibi Poey, Memorias, II, 1861, 224, Cuba.
Caranx beani Jordan, Proc. U. S. Nat. Mus., III, 1880, 486, Beaufort, N. C.
- Genus 726. PARATRACTUS Gill.**
Parattractus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 432 (*Caranx pisquetus* Cuvier and Valenciennes = *Scomber cryos* Mitchell).
- 2116. *Parattractus cryos* (Mitchill).** *Hardtail; Runner; Blue runner; Jurel; Yellow mackerel; Crevalle; Cojinera.*
 Cape Cod to Brazil.
Scomber cryos Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 424, New York.
Trachurus squamosus Gronow, Cat. Fishes, ed. Gray, 1854, 125, Carolina.
Caranx pisquetus Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 73, Santo Domingo; Cuba; Brazil.
- 2117. *Parattractus caballus* (Günther).** *Cocinero; Cocinero dorado; Jurel.*
 Pacific coast of tropical America, San Diego to Panama, recently taken at San Pedro.
Caranx caballus Günther, Trans. Zool. Soc. Lond., VI, 1868, 431 (Nov. 27), Panama.
Trachurus boops Girard, Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 108, San Diego; not *Caranx boops* of Cuvier and Valenciennes.
Caranx girardi Steindachner, Ichthyol. Notizen, IX, 1869, 25, Mazatlan.
- Genus 727. XUREL Jordan and Evermann.**
Xurel Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, Apr. 27, 1927, 505 (*Caranx vinctus* Jordan and Gilbert).
- 2118. *Xurel vincta* (Jordan and Gilbert).** *Cocinera.*
 Pacific coast of Mexico, Mazatlan to Punta Arenas; Gulf of California to Panama.
Caranx vinctus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 349, Mazatlan, Mexico.
- 2119. *Xurel fasciatus* (Cuvier and Valenciennes).** *Segundo; Volantin.*
 Cuba; Atlantic coast of Mexico.
 ? *Caranx fasciatus* Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 53 (70), east coast of Mexico.
Caranx secundus Poey, Memorias, II, 1861, 223, Cuba.
- 2120. *Xurel lata* (Agassiz).** *Jurel; Xurel; Horse-eye jack.*
 Tropical Atlantic, north to Virginia; probably not in the Pacific, where it is represented by the near related *Caranx melampygus*.
Caranx latus Agassiz, in Spix, Pisc. Bras., 1829, 105, Brazil.
Caranx lepturus Agassiz, loc. cit., 106, Brazil.
Caranx fallax Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 71 (95), Antilles; Brazil.
Caranx richardi Holbrook, Ichthyol. South Carolina, 1855, 94, pl. 13, fig. 1, South Carolina.
Caranx aureus Poey, Enumeratio, 1875, 76, Cuba.
- 2121. *Xurel medusicola* (Jordan and Starks).**
 Pacific coast of Mexico, south to Panama.
Caranx medusicola Jordan and Starks, in Jordan, Proc. Calif. Acad. Sci., V, 1895, 430, pl. 34, Mazatlan.
- 2122. *Xurel marginatus* (Gill).**
 Pacific Ocean; Mazatlan, Panama, and Hawaii.
Caranx marginatus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 166, Panama.
- 2123. *Xurel melampygus* (Cuvier and Valenciennes).** *Ulua.*
 Tropical Pacific, east to off-shore islands of Mexico; Hawaii. A very important food fish.
Caranx melampygus Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 103, East Indies.

- 2124. Xurel stellatus** (Eydoux and Souleyet). *Omilu*.
 Pacific Ocean generally; Hawaii; Revillagigedo Islands; with the
 "Ulua" one of the staple food fishes of the Pacific islands.
Caranx punctatus Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833,
 38; name preoccupied.
Caranx stellatus Eydoux and Souleyet, Voy. Bonite, Zool., I, Poiss.,
 1840, 167, Hawaiian Islands.

- 2125. Xurel lugubris** (Poey). *Tiñosa*.
 Tropical Atlantic; West Indies.
Caranx lugubris Poey, Memorias, II, 1861, 222, Cuba.
Caranx frontalis Poey, loc. cit., 222, Cuba.

- 2126. Xurel tenebrosus** Jordan and Evermann.
 South seas; Revillagigedos; Hawaii.
Caranx lugubris Jordan and Evermann, Fishes North and Mid. Amer.,
 1896, 924, in part; not of Poey, based on examples so listed by Jordan
 and McGregor, Report U. S. Fish Com., XXIV, 1899, 227, from
 Clarion Island of the Revillagigedos Group.
Caranx tenebrosus Jordan and Evermann, Proc. Calif. Acad. Sci., 4th
 Ser., XVI, Nov. 14, 1927, 656, Clarion Island.

- Genus 728. CARANX Lacépède.**
Caranx Lacépède, Hist. Nat. Poiss., III, 1802, 57 (*Scomber carangus*
 Bloch = *Scomber hippo* L., as now usually restricted). Species with
 the breast naked except for a central patch of scales.
Tricropodus Rafinesque, Caratteri, 1810, 41 (*Scomber carangus* Bloch =
Scomber hippo Linnaeus), as restricted.
Carangichthys Bleeker, Nat. Tijdschr. Neder. Indie, III, 1852, 760 (*C.
 typus* Bleeker); immature; fish with breast naked; strong teeth
 present.

- 2127. Caranx hippo** (Linnaeus). *Crevallé*; *Toro*; *Horse crevallé*; *Cavalla*; *Jack*;
Jiguagua; *Yellow caranx*; *Yellow mackerel*.
 East coast of tropical America, north to Cape Cod; apparently repre-
 sented in the Pacific by the closely related *Caranx ignobilis*.
Scomber hippo Linnaeus, Syst. Nat., Ed. XII, 1766, 494, Charleston,
 S. C.
Scomber carangus Bloch, Naturgeschich. Ausl. Fische, VII, 1793, 69,
 Antilles.
Caranx erithrurus Lacépède, Hist. Nat. Poiss., III, 1802, 68, South
 Carolina.
Caranx daubentonii Lacépède, loc. cit., 72, Martinique.
Caranx antillarum Bennett, Whaling Voyage, II, 1840, 282, West Indies.
Caranx defensor DeKay, N. Y. Fauna: Fishes, 1842, 120, New York.
Caranx esculentus Girard, U. S. Mex. Bound. Surv., 1859, 23, pl. XI,
 figs. 1-3, Brazos Santiago, Tex.

- 2128. Caranx ignobilis** Forskål.
 Tropical Pacific, east to offshore islands of Mexico; Hawaii.
Scomber ignobilis Forskål, Descr. Anim., 1775, Red Sea.
Caranx hippocoides Jenkins, Bull. U. S. Fish Com., XXII, 1902, 443,
 fig. 15, Honolulu.

- Genus 729. GNATHANODON Bleeker.**
Gnathanodon Bleeker, Verh. Batav. Genootsch., XXIV, Makreele,
 art. 5, 1851 (1852), pp. 6, 18, 30, 72, etc. (*Scomber speciosus* Forskål).
Hypocaranx Klunzinger, Fische Rothen Meeres, 1884, 92 (*Scomber
 speciosus* Forskål).

- 2129. Gnathanodon speciosus** (Forskål). *Mojarra dorada*.
 Tropical parts of the Pacific Ocean; East Indies; Mazatlan to Panama;
 Hawaii.
Scomber speciosus Forskål, Descr. Anim., 1775, 54, Red Sea.
Scomber rim Forskål, loc. cit., 54, Djidda.
Caranx panamensis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 166,
 Panama.

- Genus 730. CITULA Cuvier.**
Citula Cuvier, Règne Anim., ed. 1, 1817, 315 (*Sciæna armata* Forskål).

- 2130.** *Citula orthogramma* (Jordan and Gilbert).
Revillagigedo Islands; Pacific Islands.
Caranx orthogrammus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 226, Sulphur Bay, Clarion Island, Revillagigedo Archipelago.
- 2131.** *Citula dorsalis* (Gill). *Pámpano.*
Pacific coast of tropical America; Mazatlan; Panama.
Carangoides dorsalis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 166, Panama.
Caranx otrynter Jordan and Gilbert, Proc. U. S. Nat. Mus., 1883, 202; new name substituted for *dorsalis* preoccupied in *Caranx*, but not in *Citula*.
- Genus 731. ALECTIS** ⁵⁰ Rafinesque. *Cobblerfishes; Cockfishes.*
*Gallus Lacépède, Hist. Nat. Poiss., IV, 1803 (*Zeus gallus* L.); preoccupied in birds.*
Alectis Rafinesque, Anal. Nature, 1815, (*Zeus gallus* L.); substitute for *Gallus*, preoccupied.
*Blepharis Cuvier, Règne Animal, 1817, 322 (*Zeus ciliaris* Bloch).*
*Gallichthys Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 264 (*Zeus gallus* L.); substitute for *Gallus*, preoccupied.*
*Bleparichthys Gill, Cat. Fish. East Coast U. S., suppl. to Proc. Acad. Nat. Sci. Phila., XIII, 1861, 36; substitute for *Blepharis*.*
- 2132.** *Alectis crinitus* (Mitchill). *Threadfish; Cobblerfish; Sunfish; Shoemaker.*
West Indies north to Cape Cod, Florida Keys and Cuba.
Zeus crinitus Mitchell, Amer. Jour. Sci. Arts, XI, 1826, 144, Shoreham, Mass.
Blepharis sutor Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 120 (161), pl. 253, Caribbean Sea.
Blepharis major Cuvier and Valenciennes, loc. cit., 121 (163), West Indies; Martinique; not *Gallichthys major*, an East Indian species.
Carangoides blepharis Bleeker, Verhand. Batav. Genootsch., XXIV, 1852, Makr. 667, East Indies.
Carangoides gallichthys Bleeker, loc. cit., 668, East Indies.
- Genus 732. SCYRIS** Cuvier.
Scyris Cuvier, Règne Anim., ed. 2, II, 1829, 209 (*S. indica* Cuvier and Valenciennes).
- 2133.** *Scyris analis* Poey.
Cuba.
Scyris analis Poey, Synopsis, 1868, 369, Cuba.
- Genus 733. HYNNIS** Cuvier and Valenciennes.
Hynnus Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 145 (195), pl. 257 (*H. goreensis* Cuvier and Valenciennes).
- 2134.** *Hynnus cubensis* Poey.
Cuba.
Hynnus cubensis Poey, Memorias, II, 1861, 235, Habana.
- 2135.** *Hynnus hopkinsi* Jordan and Starks. *Pámpano.*
Puerto Viejo, near Mazatlan, Mexico.
Hynnus hopkinsi Jordan and Starks, in Jordan, Proc. Calif. Acad. Sci., V, 1895, 435, pl. 35, Mazatlan.
- Genus 734. VOMER** Cuvier and Valenciennes.
Vomer Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 141 (189), pl. 256 (*V. brownii* Cuvier and Valenciennes = *Zeus setapinnis* Mitchell).
Platysomus Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 250 (*Vomer brownii* Cuvier and Valenciennes).

⁵⁰ The genera and species allied to *Alectis* have been much mixed from the first. While other possibilities exist, and may be defended, it seems best to adhere to current nomenclature, which, in fact, accords best with the Code.

- 2136.** *Vomer setapinnis* (Mitchill). *Jorobado; Moonfish; Horsefish; Bluntnosed shiner; Sunfish; Pug-nosed shiner.*
 Atlantic coast of America; from Maine to Florida.
Zeus setapinnis Mitchill, Trans. Lit. Phil. Soc. N. Y., 1815, 384, New York.
Vomer brownii Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 141 (189), pl. 256, New York.
Argyreiosus unimaculatus Batchelder, Proc. Bost. Soc. Nat. Hist., II, 1845, 78, Maine.
Vomer curtus Cope, Proc. Acad. Nat. Sci. Phila., XXII, 1870, 119, Rhode Island.
- 2137.** *Vomer setapinnis cubensis* Nichols.
 Coast of Cuba.
Vomer setapinnis cubensis Nichols, Bull. Amer. Mus. Nat. Hist., XXXVIII, 1918, 672, Cuba.
- 2138.** *Vomer dorsalis* Gill.
 West coast of Africa, recorded from the West Indies.
Vomer dorsalis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 436; after Günther.
Argyreiosus setipinnis var. B. Günther, Cat., II, 1860, 459, St. Vincent; Cape Verde Islands; West Indies.
Vomer senegalensis Guichenot, Ann. Soc. Linn. Maine et Loire, 1865, 35, Senegal.
Vomer goreensis Guichenot, loc. cit., 37, Gorea.
- 2139.** *Vomer spixii* (Swainson).
 West Indies south to Brazil.
Platysomus spixii Swainson, Class. Fishes, etc., II, 1839, 250 and 406, Brazil; after Spix and Agassiz, pl. 37.
Platysomus micropteryx Swainson, loc. cit., 406, Pernambuco.
Vomer sanctæ-marthæ, columbianus, martinicensis, dominicensis, novæ-boracensis, sanctæ-petri, brasiliensis, cayennensis, and cubæ Guichenot, Ann. Soc. Linn. Maine et Loire, 38 to 44, 1865, localities indicated by names.
- 2140.** *Vomer declivifrons* Meek and Hildebrand.
 Cape San Lucas to Peru.
Vomer declivifrons Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1925, 367, Pl. XXIX, fig. 2, Panama.
- 2141.** *Vomer gabonensis* Guichenot.
 West Indies and west coast of Africa.
Vomer gabonensis Guichenot, Ann. Soc. Linn. Maine et Loire, 1865, 42, Gaboon.
- Genus 735. **ARGYREIOSUS** Lacépède. *Moonfishes.*
Selene Lacépède, Hist. Nat. Poiss., IV, 1803, 560 (*S. argentea* Lacépède = *Zeus vomer* L.).
Argyreiosus Lacépède, loc. cit., IV, 1803, 566 (*Zeus vomer* Linnaeus).
- 2142.** *Argyreiosus vomer* (Linnaeus). *Moonfish; Jorobado; Lookdown; Horse-head; Hair-finned dory; Hair-finned silverfish; Dollarfish.*
 Tropical America, on both coasts; Cape Cod to Brazil; Lower California to Peru.
Zeus vomer Linnaeus, Syst. Nat., Ed. X, 1758, 266, America.
Zeus gallus Linnaeus, loc. cit., America; after Artedi.
Zeus capillaris Mitchill, Trans. Lit. and Phil. Soc., I, 383, 1815, New York.
Zeus rostratus Mitchill, loc. cit., 384, New York.
Zeus geometricus Mitchill, Amer. Month. Mag., II, 245, 1818, New York.
Argyrius triacanthus Swainson, Nat. Hist. Classn. Fish., 1839, 250, Brazil.
Argyrius filamentosus Swainson, loc. cit., 250; after Cuvier and Valenciennes.
Argyrius setifer Swainson, loc. cit., 407; after Cuvier and Valenciennes.
Argyrius mauriceii Swainson, loc. cit., 408, Brazil.
Argyrius mitchilli DeKay, N. Y. Fauna: Fishes, 1842, 126, New York.
- 2143.** *Argyreiosus ærstedii* (Lütken).
 Pacific coast, Mazatlan to Panama.
Selene ærstedii Lütken, Spolia Atlantica, 1880, 144, Punta Arenas.

- 2144. *Argyreiosus brevoorti* Gill.**
Lower California to Peru.
Argyreiosus brevoorti Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 83,
Lower California.
Argyreiosus pacificus Lockington, Proc. Calif. Acad. Sci., first ser.,
VII, 1876, 84, Magdalena Bay, Lower California.
- Genus 736. CHLOROSCOMBRUS Girard. Casabes.**
Chloroscombrus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 168
(*Micropteryx cosmopolita* Agassiz = *Scomber chrysurus* Linnaeus).
Micropteryx Günther, not of Agassiz, which is *Seriola*.
- 2145. *Chloroscombrus chrysurus* (Linnaeus).** *Casabe; Bumper; Yellowtail.*
Cape Cod to Brazil; common on our South Atlantic coast and in Cuba.
Scomber chrysurus Linnaeus, Syst. Nat., ed. XII, 1766, 494, Charleston,
S. C.
Micropteryx cosmopolita Agassiz, in Spix, Pisc. Bras., 1829, 104, Tab.
LIX, Atlantic Ocean.
Scomber latus Gronow, Cat. Fishes, ed. Gray, 1854, 127, Carolina.
Chloroscombrus caribbaeus Girard, Mex. Bound. Surv., Zool., 1859, 21,
Pl. XI, fig. 6, St. Joseph Island, Tex.
- 2146. *Chloroscombrus orqueta* Jordan and Gilbert.** *Orqueta; Xurel de Castilla.*
Pacific coast of tropical America, Magdalena Bay to Panama.
Chloroscombrus orqueta Jordan and Gilbert, Proc., U. S. Nat. Mus., V,
1862, 646, Panama.
Chloroscombrus stirurus Jordan and Gilbert, loc. cit., VI, 1883, 206,
Panama; slip for *C. orqueta* imperfectly corrected proof.
- Genus 737. TRACHINOTUS Lacépède. Pámpanos.**
Trachinotus Lacépède, Hist. Nat. Poiss., III, 1802, 79 (*Labrus falcatus*
Linnaeus).
Cæsiomorus Lacépède, loc. cit., 92 (*C. bailloni* Lacépède).
Acanthinion Lacépède, loc. cit., IV, 1803, 499 (*Chætodon rhomboides*
Linnaeus).
Baillonus Rafinesque, Analyse Nature, X, 1815 (*Trachinotus bailloni*
Lacépède); substitute for *Cæsiomorus*.
Bothroænus Holbrook, Ichth. South Carolina, 1855, 80 (*Trachinotus*
pampanus Cuvier and Valenciennes).
Doliodon Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 168 (*Gasterosteus*
carolinus Linnaeus).
- 2147. *Trachinotus palometra* Regan.** *Gafftopsail; Pámpano; Oldwife; Palometa; Banner pompano.*
Tropical America, from Virginia to the Caribbean Sea.
Chætodon glaucus Bloch, Ichth., 1787, pl. 210, Martinique; on a figure
by Plumier (name preoccupied by *Lichia glauca* L., also a *Trachinotus*).
Trachynotus palometa Regan, Ann. Mag. Nat. Hist., (7), XII, 1903,
349; substitute for *glaucus* Bloch).
- 2148. *Trachinotus rhodopus* (Gill).** *Pampanito.*
Pacific coast of tropical America; south to Panama.
Trachynotus rhodopus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 85,
Cape San Lucas.
? *Trachynotus nasutus* Gill, loc. cit., 85, Cape San Lucas.
Trachynotus fasciatus Gill, loc. cit., 86, Panama.
Trachynotus glacooides Günther, Proc. Zool. Soc. Lond., 1864, 150, San
Jose de Guatemala.
- 2149. *Trachinotus falcatus* (Linnaeus).** *Round pampano; Palometa; Permit*
(Indian River) *Spinous dory; Ovate pampano.*
East coast of United States, Cape Cod to Florida.
Labrus falcatus Linnaeus, Syst. Nat., ed. X, 1758, 284, America.
Trachinotus spinosus DeKay, N. Y. Fauna: Fishes, 1842, 117, pl. 19,
fig. 53, New York Harbor.
Trachinotus fuscus Cuvier and Valenciennes, Hist. Nat. Poiss., VIII,
1831, 302 (410), Brazil.

2150. *Trachinotus rhombooides* (Bloch). *Round pampano.*
 West Indies to Brazil.
Chætodon rhombooides Bloch, Nat. Ausl. Fische, III, 1787, 100, Martinique.
2151. *Trachinotus culveri* Jordan and Starks.
 Pacific coast of Mexico.
Trachinotus culveri Jordan and Starks, in Jordan, Proc. Calif. Acad. Sci., V, 1895, 439, pl. 36, Astillero at Mazatlan, Mexico.
2152. *Trachinotus kennedyi* Steindachner. *Palometa.*
 Magdalena Bay to Panama.
Trachinotus kennedyi Steindachner, Ichth. Beitr., III, 1875, 47, Pl. VII, Magdalena Bay.
2153. *Trachinotus goodei* Jordan and Evermann. *Permit; Palometa; Great pampano.*
 West Indies, north to southern Florida.
Trachinotus goodei Jordan and Evermann, Fishes North and Mid. Amer., 1896, 943, Key West, Fla.
2154. *Trachinotus carolinus* (Linnæus). *Common pampano; Pampano; Cobblerfish; Butterfish; Carolina pompano.*
 South Atlantic and Gulf coasts of United States; Cape Cod; West Indies; Brazil. One of our best food fishes.
Gasterosieus carolinus Linnæus, Syst. Nat., ed. XII, 1766, 490, Carolina.
?Trachinotus argenteus Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 304 (410), New York; Rio Janeiro.
?Trachinotus cupreus Cuvier and Valenciennes, loc. cit., 305 (414), Martinique.
Trachinotus pampanus Cuvier and Valenciennes, loc. cit., 305 (415), Brazil; Charleston, S. C.
2155. *Trachinotus paloma* Jordan and Starks.
 Cape San Lucas, Mazatlan, San Juan Lagoon.
Trachinotus paloma Jordan and Starks, in Jordan, Proc. Calif. Acad. Sci., V, 1895, 437, Mazatlan, Mexico.
- Genus 738. **ZALOCYS** Jordan and McGregor.
Zalocys Jordan and McGregor (R. C.), Rept. U. S. Fish. Com., XXIV, 1898 (1899), 276 (*Zalocys stilbe* Jordan and McGregor).
2156. *Zalocys stilbe* Jordan and McGregor.
 Off Pacific coast of Mexico, Revillagigedo Islands.
Zalocys stilbe Jordan and McGregor, Rept. U. S. Fish Com., XXIV, 1898 (1899), 277, pl. 5, Clarion Island.
- Genus 739. **OLIGOPLITES** Gill. *Leatherjackets.*
Oligoplites Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 166 (*O. occidentalis* Gill.).
2157. *Oligoplites saurus* (Bloch and Schneider). *Leatherjacket; Leathercoat; Runner; Zapatero; Quiebra; Sauteur.*
 Both coasts of Central America, and West Indies, north to New York and Lower California.
Scomber saurus Bloch and Schneider, ed. Bloch, Syst. Ichth., 1801, 321, Jamaica; after *Saurus argenteus* of Browne.
Centronotus argenteus Lacépède, Hist. Nat. Poiss., III, 1802, 316, equatorial America.
Lichia quiebra Quoy and Gaimard, Voy. Freycinet, Zool., 1824, 365, equatorial America.
Chorinemus saltans Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 289 (393), Martinique, Brazil, and San Domingo.
2158. *Oligoplites inornatus* Gill. *Plain leatherjacket.*
 Panama.
Oligoplites inornatus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 166 Panama.
2159. *Oligoplites saliens* (Bloch). *Sauteur.*
 West Indies.
Scomber saliens Bloch, Naturgeschich. Ausl. Fische, VII, 1793, 49, Martinique; on a figure by Plumier.
Scomberoides saltator Lacépède, Hist. Nat. Poiss., II, 1800, pl. 19, fig. 2.

2160. *Oligeplites palometa* (Cuvier and Valenciennes).
Lake Maracaibo, Venezuela.
Chorinemus palometa Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 288 (392), Lake Maracaibo, Venezuela.
2161. *Oligeplites altus* (Günther). *Jurel; Monda*.
Pacific coast of tropical America; Panama.
Chorinemus altus Günther, Trans. Zool. Soc. London, VI, 1869, 433, Panama.
2162. *Oligeplites mundus* Jordan and Starks.
West coast of Mexico; Mazatlan.
Oligeplites mundus Jordan and Starks, in Jordan and Evermann, Check List, 1896, 344, Mazatlan, Mexico (name only); Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2844, San Juan Lagoon, Mexico.
2163. *Oligeplites refulgens* Gilbert and Starks.
Panama.
Oligeplites refulgens Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 73, Panama.

✓ Family 180. SERIOLIDÆ

Genus 740. NAUCRATES Rafinesque.

Naucrates Rafinesque, Caratteri 1810, 44 (*Centronotus conductor* Lacépède = *Gasterosteus ductor* Linnaeus).
Centronotus Lacépède, Hist. Nat. Poiss., III, 1802, 311, 51 (*C. conductor* Lacépède).
Xystophorus Richardson, Voy. *Erebus and Terror*, 1844, 52, pl. 30, fig. 22 (larval form, no species named).
Nauclerus Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 247 (*N. compressus*, Cuvier and Valenciennes; young of *Naucrates*).

2164. *Naucrates ductor*. (Linnaeus). *Pilotfish; Romero*.
Atlantic coast from Cape Cod to West Indies.
Gasterosteus ductor Linnaeus, Syst. Nat., ed. X, 1758, 295, "in Pelago."
Centronotus conductor Lacépède, Hist. Nat. Poiss., III, 1802, 311, no locality.
Naucrates fangsarius Rafinesque, Caratteri, 1810, 45, Sicily.
Naucrates noveboracensis Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 239 (325), New York.
Seriola dussunieri Cuvier and Valenciennes, loc. cit., IX, 1833, 162 (217), Gulf of Bengal.
Seriola succincta Cuvier and Valenciennes, loc. cit., 163 (218), south of St. Helena.
Nauclerus compressus Cuvier and Valenciennes, loc. cit., 185 (249), pl. 263, fig. 1, Molucca.
Nauclerus abbreviatus Cuvier and Valenciennes, loc. cit., 187 (251), Atlantic Ocean.
Nauclerus brachycentrus Cuvier and Valenciennes, loc. cit., 187 (253), Indian Ocean.
Nauclerus triacanthus Cuvier and Valenciennes, loc. cit., 188 (253), Atlantic Ocean.
Naucrates annularis Cuvier and Valenciennes, loc. cit., 189 (254), south of St. Helena.
Naucrates lecurus Cuvier and Valenciennes, loc. cit., 189 (255), south of St. Helena.
Naucrates cyanophrys Swainson, Classn. Fishes, II, 1839, 412, Palermo.
Naucrates serratus Swainson, loc. cit., 413, Palermo.
Thynnus pomphilus Gronow, Cat. Fishes, 1854, 123, Mediterranean; seas of America and Spain.
Xystophorus Richardson, Voy. *Erebus and Terror*, 1844, 52, Palermo.

- 2165 *Naucrates polysarcus* Fowler.
Pacific coast of Mexico.
Naucrates polysarcus Fowler, Proc. Acad. Nat. Sci. Phila., LVII, 1905, 62, west coast of Mexico.

Genus 741. ELAGATIS Bennett.

Elagatis Bennett, Narrative of Whaling Voyage, II, 1840, 283 (*Seriola bipinnulata* Quoy and Gaimard).
Seriolichthys Bleeker, Nat. Tijdschr. Neder.-Indie, VI, 1854, 196
 (*Seriola bipinnulata* Quoy and Gaimard).
Decaptus Poey, Memorias, II, 1861, 391 (*Seriola pinnulata* Poey).

2166. *Elagatis bipinnulata* (Quoy and Gaimard). *Runner; Yellowtail; Skipjack; Shoemaker.*

Tropical seas; West Indies; Long Island; Hawaii.
Seriola bipinnulata Quoy and Gaimard, Voy. *Uranie*, Zool., I, 1824, 363, pl. 61, fig. 3, Keeling Islands.
Seriola pinnulata Poey, Memorias, II, 1861, 233, Habana, Cuba.

Genus 742. SERIOLA Cuvier.

Seriola Cuvier, Règne Animal, ed. 2, II, 1829, 205 (*Caranx dumerili* Risso).

Micropteryx Agassiz, Pisc. Brasil., 1829, 102, type (not named) *Seriola dumerili* Cuvier, substitute for *Seriola*, used in botany.
Zonichthys Swainson, Nat. Hist. Fishes, Amphib., etc., II, 1839, 248 (*Scomber fasciatus* Bloch).
Halatractus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 442 (*Seriola dumerili*, substitute for *Seriola*, used in botany).
Lepidomegas Thominot, Bull. Soc. Philom. Paris, ser. 7, IV, 1880, 173 (*L. muelleri* Thominot).

2167. *Seriola dorsalis* (Gill). *Yellowtail; White salmon.*

Pacific coast of America from Monterey southward to Mazatlan; abundant about the Santa Barbara Islands.
Halatractus dorsalis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 84, Cape San Lucas.

2168. *Seriola zonata* (Mitchill). *Rudderfish; Shark pilot; Pilofish; Banded rudderfish; Jackfish; Banded mackerel.*

Atlantic coast of the United States from Cape Cod to Cape Hatteras.
Scomber zonatus Mitchill, Trans. Lit. Phil. Soc. N. Y., 1815, 427, New York Bay.

2169. *Seriola carolinensis* Holbrook.

Gulf of Mexico, and on Atlantic coast north to Cape Hatteras.
Seriola carolinensis Holbrook, Ichthyol. South Carolina, 1855, 70, Charleston, S. C.
Seriola stearnsii Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 48, Pensacola, Fla.

2170. *Seriola lalandi* Cuvier and Valenciennes. *Great amberfish; Amber jack; Coronado; Jenny Lind.*

New Jersey to Brazil; occasionally to New York.
Seriola lalandi Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 155 (208), Brazil.
Seriola gigas Poey, Memorias, II, 1861, 227, Cuba.

2171. *Seriola dumerili* (Risso). *Amber jack; Coronado.*

Mediterranean to West Indies, north to Key West and Pensacola.
Caranx dumerili Risso, Ichthyol. Nice, 1810, 175, pl. 6, fig. 20, Nice.
Trachurus alciolus Rafinesque, Caratteri, 1810, 42, Pl. XI, fig. 2, Sicily.
Trachurus fasciatus Rafinesque, Indice, No. 108, 21 and 53, Sicily; not *Scomber fasciatus* of Bloch.
Seriola proxima Poey, Memorias, II, 1861, 229, Cuba.

Genus 743. ZONICHTHYS Swainson.

Zonichthys Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 248 (*Scomber fasciatus* Bloch).

2172. *Zonichthys mazatlanus* (Steindachner).

West coast of Mexico, Mazatlan.
Seriola mazatlana Steindachner, Ichth. Beitr., V, 1876, 8, Mazatlan, Mexico.

2173. *Zonichthys fasciatus* (Bloch). *Madregal; Bonito.*

West Indies, north to Charleston, S. C.
Scomber fasciatus Bloch, Nat. Ausl. Fische, VII, 1793, 73, pl. 341, no locality.
Seriola semicoronata Poey, Memorias, II, 1861, 232, Habana.

- 2174.** *Zonichthys rivolianus* Cuvier and Valenciennes.
Mediterranean to Brazil, West Indies, and South Carolina; Florida coast.
Seriola rivoliana Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 154 (207), the Greek Archipelago.
Seriola boscii Cuvier and Valenciennes, loc. cit., 156 (208), waters of Carolina.
Seriola bonariensis Cuvier and Valenciennes, loc. cit., 157 (211), Buenos Aires.
Seriola coronata Poey, Memorias, II, 1861, 232, Cuba.
- 2175.** *Zonichthys falcatus* (Cuvier and Valenciennes). *Madregal; Rock salmon.*
West Indies, north to Florida and the Carolinas.
Seriola falcata Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 156 (210), Gulf of Mexico.
Seriola dubia Lowe, Proc. Zool. Soc. London, VII, 1839, 80, Madeira.
Seriola declivis Poey, Memorias, II, 1861, 230, Habana.
Seriola ligulata Poey, loc. cit., 231, Cuba.

Family 181. NEMATISTIIDÆ. Peacockfishes; Papagallos

- Genus 744. NEMATISTIUS** Gill.
Nematistius Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 258 (*N. pectoralis* Gill).
Seriolophus Guichenot, Mém. Soc. Nat. Sci. Cherbourg, XIII, 1868, 90 (*Seriolophus carangooides* Guichenot).

- 2176.** *Nematistius pectoralis* Gill. *Pez de gallo; Papagallo; Roosterfish.*
Gulf of California to Panama; generally common.
Nematistius pectoralis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 259, Cape San Lucas.
Seriolophus carangooides Guichenot, Mém. Soc. Nat. Sci. Cherbourg, XIII, 1868, 90, locality unknown.

✓ Family 182. POMATOMIDÆ. Bluefishes

- Genus 745. POMATOMUS** Lacépède. *Bluefishes.*
Pomatomus Lacépède, Hist. Nat. Poiss., IV, 1802, 436 (*P. skib* Lacépède).
Lopharis Rafinesque, Caratteri, 1810, 52 (*Perca lophar* Forskål).
Gonenion Rafinesque, loc. cit., 53 (*G. serra* Rafinesque).
Temnodon Cuvier, Règne Animal, ed. 1, II, 1817, 346 (*Cheilodipterus heptacanthus* Lacépède).
Sypterus Eichwald, Fauna Caspio-Caucas, 1841 (*Perca lophar* Forskål).
Chromis Gronow, Cat. Fishes, 1854, 149 (*C. epicurorum* Gronow = *Gasterosteus saltatrix* Linnaeus); name preoccupied.
Sparactodon Rochebrune, Bull. Soc. Philom. Paris, ser. 7, IV, 1880, 162 (*S. natal* Rochebrune).

- 2177.** *Pomatomus saltatrix* (Linnaeus). *Bluefish; Snap mackerel; Skipjack; Fallback; Snapping mackerel; Tailor; Snapper; Horse mackerel; Greenfish; Skip mackerel.*
Atlantic coast of United States; eastern part of the Mediterranean and South to Australia and Madagascar; locally and irregularly abundant.

Perca saltatrix Linnaeus, Syst. Nat., ed. X, I, 1758, 293, Carolinas.
Perca lophar Forskål, Descr. Anim., 1775, 38, Constantinople.
Cheilodipterus heptacanthus Lacépède, Hist. Nat. Poiss., III, 1802, 542, Fort Dauphin, Madagascar.
Pomatomus skib Lacépède, loc. cit., IV, 1802, 436, Carolinas.
Lopharis mediterraneus Rafinesque, Indice, 1810, 17, after Forskål.
Gonenion serra Rafinesque, Caratteri, 1810, 53, Sicily.
Chromis epicurorum Gronow, Cat., ed. Gray, 1854, 149, Carolinas.
Sparactodon natal De Rochebrune, Bull. Soc. Philom. Paris, 1880, 159, Senegambia.

Family 183. RACHYCENTRIDÆ. The Sergeantfishes

Genus 746. RACHYCENTRON Kaup. *Crab-eaters.**Rachycentron* Kaup, Isis, XIX, 1826, col. 89 (*R. typus* Kaup).*Elacate* Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 328
(*E. malabarica* Cuvier and Valenciennes).*Mcladerma* Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 243
(*M. nigerrima* Swainson = *Elacate pondiceriana* Cuvier and Valen-
ciennes).2178. *Rachycentron canadus* (Linnæus). *Sergeantfish; Crab-eater; Bonito;*
*Cobia.*In all warm seas; Atlantic coast north to Cape Cod; also in Japan and
the East Indies, if all members of the genus belong to one species.*Gasterosteus canadus* Linnæus, Syst. Nat., ed. XII, 1766, 491, Carolinas.
Centronotus gardenii Lacépède, Hist. Nat. Poiss., III, 1802, 357,
Carolinas.*Centronotus spinosus* Mitchell, Trans. Lit. and Phil. Soc., I, 1815, 490,
New York.? *Elacate pondiceriana* Cuvier and Valenciennes, Hist. Nat. Poiss.,
VIII, 1831, 241 (329), Pondicherry.? *Elacate mollis* Cuvier and Valenciennes, loc. cit., 244 (332), Orixa.? *Elacate malabarica* Cuvier and Valenciennes, loc. cit., 244 (332),
Malabar.*Elacate atlantica* Cuvier and Valenciennes, loc. cit., 245 (334), Brazil.? *Elacate bivittata* Cuvier and Valenciennes, loc. cit., 248 (338), Molucca.
Elacata falcipinnis Gosse, Nat. Sojourn Jamaica, 1851, 208, Jamaica.

Family 184. BATHYCLUPEIDÆ

Genus 747. BATHYCLUPEA Alcock.

BathyCLUPEA Alcock, Ann. Mag. Nat. Hist., VIII, 1891, 130 (*B. hoskynii*
Alcock).2179. *BathyCLUPEA argentea* Goode and Bean.

West Indies, in deep water.

BathyCLUPEA argentea Goode and Bean, Oceanic Ichth., 1895, 190, off
Neyis Island, West Indies, at Blake station 37 in 365 fathoms.

✓ Family 185. PERCIDÆ. Perches

Genus 748. PERCA Linnaeus. *River perches.**Perca* Linnaeus, Syst. Nat., ed. X, 1758, 289 (*P. fluviatilis* Linnaeus).
Epitrachys Schulze, Jahrsber. Naturw. Ver. Magdeburg, 1889, 209
(*Perca fluviatilis* Linnaeus).2180. *Perca flavescens* (Mitchill). *Yellow perch; American perch; Ringed*
*perch; Raccoon perch; Yellow Ned; River perch; Lake perch.*Fresh waters of the eastern United States, chiefly northward and
eastward; abundant in the Great Lakes and in coastwise streams
from Nova Scotia to North Carolina; common in the tributaries of
the upper Mississippi, especially in lakes of Indiana, Michigan,
Wisconsin, Iowa, and Minnesota; unknown from central Ohio
southwestward; not known from the Ohio River or the lower Missouri.
Perca americana Schrank, Abh. Privatges. Oberdeutschland, I, 1792,
100, America; name preoccupied in *Morone*.*Morone flavescens* Mitchell, Rept. Fish. N. Y., 1814, 18, near New
York City.*Centropomus luteus* Rafinesque, Précis des Découvertes Somiologiques,
1814, 19, Pennsylvania.*Perca notata* Rafinesque, Amer. Month. Mag., II, 1818, 205, Lake Erie.
Perca serrato-granulata Cuvier and Valenciennes, Hist. Nat. Poiss.,
II, 1828, 34 (48), New York.*Perca granulata* Cuvier and Valenciennes, loc. cit., 35 (48), pl. 9, New
York.*Perca acuta* Cuvier and Valenciennes, loc. cit., 36 (49), pl. 10, Lake
Ontario.*Perca gracilis* Cuvier and Valenciennes, loc. cit., 36 (50), Skaneateles
Lake, N. Y.

Genus 749. STIZOSTEDION ⁵¹ Rafinesque. *American pike perches.*
Stizostedion Rafinesque, Ichth. Ohiensis, 1820, 23 (*Perca salmonea*
 Rafinesque = *Perca vitrea* Mitchell).
Stizostethium, attempted correction.

2181. *Stizostedion vitreum* (Mitchill). *Walleye; Wall-eyed pike; Pike perch;*
Dory; Glass-eye; Yellow pike; Blue pike; Jack salmon; White-eye;
Spike (small specimens about 8 inches long, at Maumee Bay,
 Sandusky Bay, Port Clinton, etc.); *Pickerel; River trout.*
 Great Lakes region, upper Mississippi, north to Assiniboina; east to
 Vermont and Pennsylvania; south to Georgia and Alabama; especially common northward.
Perca vitrea Mitchell, Am. Month. Mag., II, 1818, 247, Cayuga Lake,
 N. Y.
Lucioperca americana Cuvier and Valenciennes, Hist. Nat. Poiss., II,
 1828, 90 (122), pl. 16, New York.
Perca salmonea Rafinesque, Amer. Month. Mag. III, 1818, 354, Ohio
 River.

2182. *Stizostedion glaucum* Hubbs. *Blue pike.*
 Great Lake region.
Stizostedion glaucum Hubbs, Univ. Mich. Mus. Zool., Misc. Publ. No.
 15, July 7, 1926, 58, Pl. IV, fig. 1, Lake Erie.

- Genus 750. CYNOOPERCA Gill and Jordan.
Cynoperca Gill and Jordan, Bull. U. S. Nat. Mus., X, 1877, 44 (*Lucioperca canadensis* C. H. Smith, in Griffith's edition of Cuvier's Règne Animal).

2183. *Cynoperca grisea* (DeKay). *Sauger; Sand pike; Gray pike; Pickering.*
 Great Lakes region and westward to Montana, Tennessee, and Arkansas;
 the common species which varies northward into the next.
Lucioperca grisea DeKay, New York Fauna: Fishes, 1842, 19, New
 York.
Lucioperca borea Girard, Proc. Acad. Nat. Sci. Phila., 1857, 201, Fort
 Sarpi, Nebr.
Lucioperca pepinus Estes, in Hallock's Sportsman's Gazetteer, 1827,
 322, Lake Pepin.

2184. *Cynoperca canadensis* (Smith). *Sauger; Sand pike; Gray pike.*
 Northeastern North America, from Vermont to the Great Lakes.
Lucioperca canadensis C. H. Smith, in Griffith's edition of Cuvier's
 Règne Anim., Fishes, 1836, 275, Canada.

✓ Family 186. ETHEOSTOMIDÆ Darters

We begin this series with the species most nearly related to the *Percidæ*, from which the darters are apparently descended. *Percina* is the most primitive genus, and in most respects the highest, the other forms being progressively specialized to their life in swift brooks. From *Percina* two lines diverge in different directions; the allies of *Crystallaria* and *Vigil* to *Ammocrypta* toward a quiescent life, buried in sand; the other through *Pæciliichthys*, *Nothonotus* and *Oligocephalus* for life in weedy brooks, culminating in *Microperca* and *Alvarius*, the lowest forms.

- Genus 751. PERCINA Haldeman. *Log perches.*
Percina Haldeman, Jour. Acad. Nat. Sci. Phila., VIII, 1842, 330 (*P. nebulosa* Haldeman = *Sciæna caprodes* Rafinesque).
Pileoma DeKay, New York Fauna: Fishes, 1842, 16 (*P. semifasciata* DeKay = *Etheostoma caprodes* Rafinesque).

⁵¹ The European genus *Sander* Oken (*Lucioperca* Cuvier and Valenciennes) differs from *Stizostedion* in the close-set ventrals and the less close approximation of the anal spines.

- 2185** *Percina caprodes* (Rafinesque). *Log perch; Rockfish; Hogmolly; Hogfish.* Great Lakes and streams of the South and West from Quebec to Lake Superior, and Iowa.
Sciaena caprodes Rafinesque, Amer. Month. Mag., Vol. II, 1818, 354, Ohio River.
Perca (Percina) nebulosa Haldeman, Jour. Acad. Nat. Sci. Phila., VIII, 1842, 330, Susquehanna River.
Pileoma semifasciatum DeKay, New York Fauna: Fishes, 1842, 16, pl. 50, fig. 162, Lake Champlain, Westport, N. Y.
Percina bimaculata Haldeman, Proc. Boston Soc. Nat. Hist., 1844, 157, Susquehanna River.
- 2186.** *Percina carbonaria* (Baird and Girard).
Streams of Mississippi to the Rio Grande.
Pileoma carbonaria Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 387, Rio Salada; Rio Medina; San Pedro Creek, Tex.
- 2187.** *Percina zebra* (Agassiz). *Manitou darter.*
Lakes of Michigan, northern Indiana, Wisconsin, and northward to Lake Superior; Cayuga Lake Basin. Probably varying into *P. caprodes*.
Pileoma zebra Agassiz, Lake Superior, 1850, 308, Lake Superior.
Percina manitou Jordan, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 53, Lake Manitou, Ind.
- 2188.** *Percina rex* (Jordan and Evermann).
Roanoke River, Va. One of the largest and most imposing of the darters.
Etheostoma rex Jordan and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 357, Pl. XLV, fig. 9, Roanoke River near Roanoke, Va.
- Genus 752. *HADROPTERUS* Agassiz. *Black-sided darters.*
Hadropodus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 305 (*H. nigrofasciatus* Agassiz).
- 2189.** *Hadropodus nigrofasciatus* Agassiz. *Crawl-a-bottom.*
South Carolina to Louisiana.
Hadropodus nigrofasciatus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 305, Mobile, Ala.
Alvordius spillmani Hay, Proc. U. S. Nat. Mus., III, 1880, 491, Chickasawha River, Miss.
- Genus 753. *ALVORDIUS* Girard.
Alvordius Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 68 (*A. maculatus* Girard).
- 2190.** *Alvordius maculatus* Girard. *Black-sided darter.*
Great Lakes region to the middle Missouri and north to Minnesota, southward through Missouri, Indiana, and Kentucky to Arkansas; especially common in the Ohio Valley.
Alvordius maculatus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 68, Fort Gratiot, Mich.
Alvordius aspro Cope and Jordan, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 51, White River, Indianapolis, Ind.; substitute for *Etheostoma blennioides* of Kirtland, not of Rafinesque; perhaps distinct from *A. maculatus* which, if correctly described, has not been found later.
- 2191.** *Alvordius guntheri* (Eigenmann and Eigenmann). *Günther's darter.*
Winnipeg south to Iowa.
Etheostoma guntheri Eigenmann and Eigenmann, Amer. Nat., November, 1892, 962, Souris River, Winnipeg.
- 2192.** *Alvordius peltatus* (Stauffer). *Shielded darter.*
Southeastern Pennsylvania, southward to South Carolina in coastwise streams.
Hadropodus maculatus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 100, tributary of Potomac River, Md.; name preoccupied.
Etheostoma pellatum Stauffer, in Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 233, Conestoga Creek, near Lancaster, Pa.
Etheostoma nevisense Cope, Proc. Amer. Phil. Soc., XI, 1870, 261, falls of Neuse River, N. C.
Alvordius crassus Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 12, near Greenville, S. C.

- 2193** *Alvordius sellaris* (Radcliffe and Welch). *Maryland darter.*
 Streams of Maryland.
Hadropterus sellaris Radcliffe and Welch, Bull. U. S. Fish Com., XXXII, 1912, 31, Swan Creek, Havre de Grace, Md.
- 2194.** *Alvordius ouachitae* (Jordan and Gilbert). *Ouachita darter.*
 Southern Indiana, western Kentucky, southwest to Arkansas, not rare; known from Patoka River, Ind.; lower Green and Obion Rivers, Kentucky and Tennessee; Saline River, Ark.
Etheostoma (Hadropterus) ouachitae Jordan and Gilbert, Proc. U. S. Nat. Mus., X, 1887, 49, Saline River, Benton, Ark., a tributary of the Washita.
- 2195.** *Alvordius maxinkuckiensis* (Evermann). *Maxinkuckee darter.*
 Lake Maxinkuckee, Ind.
Hadropterus maxinkuckiensis Evermann, Rept. U. S. Fish Com., XXV, 1899 (1900), 366, pl. 17, fig. 2, Aubeenaubee Creek, an eastern tributary to Lake Maxinkuckee, Ind.
- 2196.** *Alvordius evermanni* (Moenkhaus). *Evermann's darter.*
 Lake Tippecanoe, Ind.
Hadropterus evermanni Moenkhaus, Bull. U. S. Fish Com., XXII, 1902 (1904), 397, with figure on p. 398, Tippecanoe Lake, Ind.
- 2197.** *Alvordius roanoka* (Jordan and Jenkins).
 Roanoke River, Va.
Hadropterus roanoka Jordan and Jenkins, Proc. U. S. Nat. Mus., XI, 1888, 358, Roanoke River at Roanoke, Salem, and Allegheny Springs, Va.
- 2198.** *Alvordius macrocephalus* (Cope). *Big-headed darter.*
 West slope of the Alleghenies from Pennsylvania southward in mountain streams; known from Youghiogheny River, Pa.; North Fork of Holston River, Saltville, Va.; Middle Fork of Holston River, Glade Spring, Va.; Big Sandy, upper Green, and Cumberland Rivers, Ky.; Clinch River, Tenn.
Etheostoma macrocephalum Cope, Trans. Amer. Phil. Soc., XIII, 1869, 400, Youghiogheny River, Pa.
- 2199.** *Alvordius phoxocephalus* (Nelson).
 Ohio to Iowa, south to Kentucky and Oklahoma, in sandy rivers; locally common.
Etheostoma phoxocephalum Nelson, Bull. Ill. Lab. Nat. Hist., I, 1876, 35, Illinois River and its tributaries.
- Genus 754. SERRARIA** Gilbert.
Serraria Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 205 (*Hadropterus scierus* Swain).
- 2200.** *Serraria sciera* (Swain). Northern Indiana to Tennessee and Texas.
Hadropterus scierus Swain, Proc. U. S. Nat. Mus., VI, 1883, 252, Bean Blossom Creek, Monroe County, Ind.
- 2201.** *Serraria serrula* (Jordan and Gilbert).
 From southern Arkansas southward through eastern Texas.
Hadropterus scierus serrula Jordan and Gilbert, Proc. U. S. Nat. Mus., IX, 1886, 16, Red River, Fulton, Ark.
- Genus 755. ERICOSMA** Jordan and Copeland.
Ericosma Jordan and Copeland, Bull. U. S. Nat. Mus., X, 1877, 8 (*Alvordius evides* Jordan and Copeland).
- 2202.** *Ericosma evides* (Jordan and Copeland).
 Indiana in the Wabash Basin, west to Cedar River, Iowa, and southward in Arkansas, Kentucky, and Tennessee; French Broad River and the Ozark region.
Alvordius evides Jordan and Copeland, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 51, White River near Indianapolis, Ind.
- Genus 756. HYPOHOMUS** Cope.
Hypohomus Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 449 (*Cottogaster aurantiacus* Cope).

- 2203. *Hypohomus aurantiacus* (Cope).**
Upper Tennessee Basin; North Fork of Holston, Clinch, Watauga, and French Broad Rivers.
Cottogaster aurantiacus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1869, 211, North Fork of Holston River, Saltville, Va.
- 2204. *Hypohomus cymatotenia* (Gilbert and Meek).**
Western Kentucky and southern Missouri.
Etheostoma (Hadropterus) cymatotenia Gilbert and Meek, Proc. U. S. Nat. Mus., X, 1887, 51, Niangua River and Osage Fork of the Gasconade near Marshfield, Mo.; Sac River near Greenfield, Mo.
- 2205. *Hypohomus nianguae* (Gilbert and Meek). *Niangua darter.***
Niangua River, Ozark region of Missouri.
Etheostoma (Hadropterus) nianguae Gilbert and Meek, Proc. U. S. Nat. Mus., X, 1887, 52, Niangua River near Marshfield, Mo.
- 2206. *Hypohomus spilotus* (Gilbert).**
Kentucky River.
Etheostoma nianguae spilotum Gilbert, Proc. U. S. Nat. Mus., X, 1887, 53, Sturgeon Creek, a tributary of Kentucky River near Travelers Rest, Owsley County, Ky.
- Genus 757. *SWAINIA* Jordan and Evermann.**
Swainia Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1040 (*Etheostoma (Hadropterus) squamatus* Gilbert and Swain).
- 2207. *Swainia squamata* (Gilbert and Swain).**
Upper Tennessee River Basin; known from Watauga and French Broad Rivers.
Etheostoma (Hadropterus) squamatus Gilbert and Swain, Proc. U. S. Nat. Mus., X, 1887, 50, French Broad River at mouth of Wolf Creek, Tenn.
- Genus 758. *COTTOGASTER* Putnam.**
Cottogaster Putnam, Bull. Mus. Comp. Zool., I, 1863, 4 (*Bolcosoma tessellatum* Thompson; not of DeKay = *Cottogaster putnami* Jordan and Gilbert).
Rheocrypta Jordan, Bull. U. S. Nat. Mus., X, 1877, 9 (*R. copelandi* Jordan).
- 2208. *Cottogaster copelandi* (Jordan). *Copeland's darter.***
Great Lakes region, south to the Black Warrior, westward to Missouri and through the Ozark region, where it is abundant; also about Indianapolis in clear brooks.
Rheocrypta copelandi Jordan, Bull. U. S. Nat. Mus., X, 1877, 9, White River near Indianapolis, Ind.
- 2209. *Cottogaster putnami* (Jordan and Gilbert). *Putnam's darter.***
Lake Champlain.
Cottogaster putnami Jordan and Gilbert, Synopsis, 1883, 498, Westport Brook, Lake Champlain, N. Y. Probably intergrades with *C. copelandi*.
- 2210. *Cottogaster uranidea* (Jordan and Gilbert).**
Lower Wabash Basin to southern Missouri, south through Arkansas and Alabama to the Escambia; mostly in sandy lowland streams.
Etheostoma (Cottogaster) uranidea Jordan and Gilbert, Proc. U. S. Nat. Mus., X, 1887, 48, Washita River, Arkadelphia, Ark.
- Genus 759. *IMOSTOMA* Jordan.**
Imostoma Jordan, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 49 (*Hadropterus shumardi* Girard).
- 2211. *Imostoma shumardi* (Girard). *Shumard's darter.***
Michigan, Ohio, Indiana, and Illinois southward to Kentucky and Arkansas.
Hadropterus shumardi Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 100, Arkansas River near Fort Smith, Ark.
Etheostoma wrighti McCormick, Bull. Oberlin College Lab., No. 2, 1891, 30, near Oberlin, Ohio.

- 2212.** *Imostoma gilberti* (Evermann and Thoburn). *Gilbert's darter.*
Clinch River, Tenn.
Ulocentra gilberti Evermann and Thoburn in Jordan and Evermann,
Fishes North and Mid. Amer., 1896, 1049, Clinch River at Walker's
ford near Tazewell, Tenn.
- 2213.** *Imostoma meadiæ* (Jordan and Evermann). *Meadie's darter.*
Indian Creek, Powell River basin, eastern Tennessee.
Ulocentra meadiæ Jordan and Evermann, Fishes North and Mid. Amer.,
Nov. 26, 1898, 2852, Indian Creek, Cumberland Gap, Tenn.
- 2214.** *Imostoma longimane* (Jordan).
Basin of James River, Va.
Etheostoma longimana Jordan, Proc. Acad. Nat. Sci. Phila., XL, 1888,
179, tributary of James River, Va.
- 2215.** *Imostoma cheneyi* (Evermann and Kendall). *Cheney's darter.*
Racket River near Norfolk, St. Lawrence County, N. Y.
Cottogaster cheneyi Evermann and Kendall, Bull. U. S. Fish Com.,
XVII, 1897 (Feb. 9, 1898), 129, pl. 8, fig. 8, Racket River, near
Norfolk, N. Y.
- 2216.** *Imostoma podostemone* (Jordan and Jenkins).
Roanoke River Basin, Va.
Etheostoma podostemone Jordan and Jenkins, Proc. U. S. Nat. Mus.,
XI, 1888, 359, Roanoke River, at Roanoke, Salem, and Allegheny
Springs, Va.
- Genus 760. ETHEOSTOMA Rafinesque.**
Etheostoma Rafinesque, Jour. Phys., etc., 1819, 419 (*E. blennioides*
Rafinesque, as restricted by code).
Diplosoma Rafinesque, Ichth. Ohiensis, 1820, 37 (*Etheostoma blennioides*
Rafinesque).
Hyostoma Agassiz, Amer. Jour. Sci., XVII, 1854, 305 (*H. newmani*
Agassiz = *Etheostoma blennioides* Rafinesque).
- 2217.** *Etheostoma blennioides* Rafinesque. *Green-sided darter.*
New York to South Dakota and Kansas, south to lower Alabama
Basin; a beautiful fish.
Etheostoma blennioides Rafinesque, Jour. de Physique, LXXXVIII,
1819, 419, Ohio River.
Hyostoma newmani Agassiz, Amer. Jour. Sci. and Arts, X, VII, 1854,
305, vicinity of Huntsville, Ala.
Pileoma cymatogramma Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860,
327, no locality.
Hyostoma blennioperca Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868,
215, tributaries of Kanawha and Holston Rivers.
- Genus 761. ULOCENTRA Jordan.**
Ulocentra Jordan, Man. Vertebrates, ed. 2, 1878, 223 (*Arlina atripinnis*
Jordan = *Hyostoma simoterum* Cope).
- 2218.** *Ulocentra simotera* (Cope).
West Virginia, eastern Kentucky, and Tennessee, in the basins of the
Green, Cumberland, and Tennessee Rivers, southward through
Alabama to the Escambia River.
Hyostoma simoterum Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 215,
Holston River and its tributaries, near Nashville, Tenn.
Arlina atripinnis Jordan, Bull. U. S. Nat. Mus., X, 1877, 10, tributary
of Cumberland River near Nashville, Tenn.
? *Etheostoma duryi* Henshall, Jour. Cincinnati Soc. Hist., 1889, 32,
tributary of Tennessee River.
- 2219.** *Ulocentra histrio* (Jordan and Gilbert).
Southern Indiana, southwestward to Arkansas; known from the lower
Wabash and tributaries; also from Green River, Ky., and Black,
Poteau, and Washita Rivers, Ark.
Etheostoma (Ulocentra) histrio Jordan and Gilbert, Proc. U. S. Nat.
Mus., X, 1887, 47, Poteau River near Hackett City, Ark.; Saline
River at Benton, and Washita River at Arkadelphia, Ark.

- 2220. *Ulocentra blennius* (Gilbert and Swain).**
 Northern Alabama, in streams tributary to the Tennessee River.
Etheostoma (Rhothæca) blennius Gilbert and Swain, Proc. U. S. Nat. Mus., X, 1887, 55, Cox Creek and Shoal Creek, tributary to Tennessee River, Florence, Ala.
- 2221. *Ulocentra verecunda* (Jordan and Evermann).**
 Tributaries of Holston River, Va.; rare.
Etheostoma verecundum Jordan and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 360, Pl. XLV, fig. 12, Middle Fork of the Holston River, about 5 miles south of Glade Spring, Va.
- 2222. *Ulocentra phlox* (Cope).**
 Trinity River, Tex.
Boleosoma phlox Cope, Bull. U. S. Nat. Mus., XVII, 1880, 30, Trinity River at Fort Worth, Tex.
- 2223. *Ulocentra stigmæa* (Jordan). *Speck.***
 Tennessee and Arkansas to Georgia and Louisiana.
Boleosoma stigmæum Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876 (1877), 311, small tributaries of Etowah and Oostanaula Rivers near Rome, Ga.
Pæcilichthys saxatilis Hay, Proc. U. S. Nat. Mus., III, 1880, 495, stream at Enterprise, Miss.
- Genus 762. BOLEOSOMA DeKay. *Tessellated darters.***
Boleosoma DeKay, New York Fauna: Fishes, 1842, 20 (*B. tessellatum* DeKay = *Etheostoma olmstedi* Storer).
Arlina Girard, Proc. Acad. Nat. Sci. Phila., 1859, 64 (*A. effulgens* Girard).
Estrella Girard, loc. cit., 65 (*E. atramaculata* Girard).
- 2224. *Boleosoma olmstedi* (Storer). *Tessellated darter; Grand Oranhee darter; Sand darter.***
 Lake Ontario to Massachusetts and south to North Carolina, east of the Alleghenies.
Etheostoma olmstedi Storer, Jour., Bost. Soc. Nat. Hist., IV, 1842, 61, pl. 5, fig. 2, Hartford, Conn.
Perca minima Haldeman, Jour. Acad. Nat. Sci. Phila., VIII, 1842, 330, Susquehanna River.
Boleosoma tessellatum DeKay, New York Fauna: Fishes, 1842, 20, pl. 20, fig. 57; New York fresh-water streams.
Estrella atramaculata Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 66, Potomac River, vicinity of Washington, D. C.
?Boleosoma xoporus Cope, Proc. Amer. Phil. Soc., XI, 1870, 270, Loy-also Creek, Pa.
- 2225. *Boleosoma nigrum* (Rafinesque). *Johnny darter; Blind simon.***
 Eastern United States, Ohio Valley, Great Lakes region, and upper Mississippi, west to Colorado and north to Manitoba.
Etheostoma nigra Rafinesque, Ichth. Ohiensis, 1820, 37, Green River, Ky.
Boleosoma maculatum Agassiz, Lake Superior, 1850, 305, Lake Superior.
Boleosoma olmstedi brevipinnis Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 214, Kiskiminitas River, Pa.
Pæcilichthys beani Jordan, Proc. U. S. Nat. Mus., VII, 1884, 479, Tabo Creek, Mo.
- 2226. *Boleosoma effulgens* (Girard). *Tassellated darter.***
 Maryland to North Carolina.
Arlina effulgens Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 65, brooks and streams flowing into Potomac River.
- 2227. *Boleosoma mesæum* (Cope).**
 Platte River near Fort Kearney, Nebr.
Pæcilichthys mesæus Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 232, Platte River near Fort Kearney, Nebr.
- 2228. *Boleosoma maculaticeps* Cope.**
 Upper waters of Catawba River, N. C.
Boleosoma maculaticeps Cope, Proc. Amer. Phil. Soc., XI, 1870, 269, upper waters of Catawba River, N. C.

2229. *Boleosoma vexillare* Jordan.

Tributaries of the James, Roanoke, and Rappahannock Rivers.

Boleosoma vexillare Jordan, Proc. U. S. Nat. Mus., II, 1879, 235, Rappahannock River at Warrenton, Va.

2230. *Boleosoma susanæ* Jordan and Swain. *Susan's darter.*

Basin of upper Cumberland River, Ky.

Boleosoma susanæ Jordan and Swain, Proc. U. S. Nat. Mus., VI., 1883, 249, Wolf Creek and other tributaries of the Clear Fork of Cumberland River near Pleasant View, Whitley County, Ky.

Genus 763. *VAILLANTIA* Jordan.

Vaillantia Jordan, Bull. U. S. Nat. Mus., XII, 1878, 89 (*Boleosoma camurum* Forbes).

2231. *Vaillantia camura* (Forbes).

Indiana to Iowa and Mississippi, southwest to Houston, Tex.

Boleosoma camurum Forbes, Bull. Ill. Lab. Nat. Hist., I, 1878, 40, Cache River and Clear Creek, Union County, Johnston County, and Pekin, Ill.; name preoccupied in *Nothonotus* but not in *Boleosoma* or *Vaillantia*,

Vaillantia chlorosoma Hay, Proc. U. S. Nat. Mus., III, 1880, 495, Mississippi.

Genus 764. *CRYSTALLARIA* Jordan and Gilbert.

Crystallaria Jordan and Gilbert, in Jordan, Cat. Fishes N. Amer., 1885, 78 (*Pleurolepis asprellus* Jordan).

2232. *Crystallaria asprella* (Jordan).

Southern Indiana and Illinois to Arkansas and Alabama, chiefly in the larger, clearer streams. Recorded from Ohio River at Rising Sun, Ind.; Wabash River at New Harmony, Vincennes, and Terre Haute; Green River, Ky.; Choctaw Creek, Ala.; and Washita River, Ark.; besides the original locality in Illinois.

Pleurolepis asprellus Jordan, Bull. Ill. Lab. Nat. Hist., I, 1878, No. 2, 38, 1878, rocky tributary of Mississippi River in Hancock County, Ill.

Genus 765. *VIGIL* Jordan.

Pleurolepis (Agassiz) Putnam, Bull. Mus. Comp. Zool., I, 1863, 5 (*Etheostoma pellucidum* (Baird) Agassiz; name preoccupied by Egerton for a genus of fossil fishes).

Vigil Jordan, Proc. Acad. Nat. Sci. Phila., LXX, 1918 (1919), 344 (*Pleurolepis pellucidus* Agassiz).

2233. *Vigil pellucidus* (Baird). *Sand darter.*

Lake Erie to Minnesota, Kentucky, and Texas.

Pleurolepis pellucidus Baird, in Agassiz, Bull. Mus. Comp. Zool., I, 1863, 5, no locality; the specimens from Black River, Ohio, the types of *Etheostoma pellucidum* Baird (MS., 1853).

2234. *Vigil clarus* (Jordan and Meek).

Mississippi Valley; Wabash River, west to central Iowa and Minnesota, and south to Arkansas and northern Texas.

Ammocrypta clara Jordan and Meek, Proc. U. S. Nat. Mus., VIII, 1885, 8, Des Moines River, Ottumwa, Iowa.

2235. *Vigil vivax* (Hay).

Mississippi and northwest through Arkansas and south to Texas.

Ammocrypta vivax Hay, Bull. U. S. Fish Com., II, 1882, (1883), 58, Pearl River, Jackson, Miss.

Genus 766. *IOA* Jordan and Brayton.

Ioa Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 88 (*Paeclichthys vitreus* Cope).

2236. *Ioa vitrea* (Cope).

Southeastern Virginia and western North Carolina.

Paeclichthys vitreus Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 263, Walnut Creek, a tributary of Neuse River, Wake County, N. C.

2237. *Ioa vigil* Hay.

Pearl River, Jackson, Miss.

Ioa vigil Hay, Bull. U. S. Fish Com., II, 1882 (1883), 59, Pearl River, Jackson, Miss.

- Genus 767. AMMOCRYPTA** Jordan. *Sand darters.*
Ammocrypta Jordan, Bull. U. S. Nat. Mus., X, 1877, 5 (*A. beanii* Jordan).
- 2238. Ammocrypta beanii** Jordan.
 Gulf States, Alabama to Louisiana.
Ammocrypta beanii Jordan, Bull. U. S. Nat. Mus., X, 1877, 5, Notalbany River, Tickfaw, La.
Ammocrypta gelida Hay, Proc. U. S. Nat. Mus., III, 1880, 490, Enterprise, Miss.
- Genus 768. PÆCILICHTHYS** Agassiz.
Pæcilosoma Agassiz, Lake Superior, 1850, 299 (*Etheostoma variata* Kirtland); name preoccupied.
Pæciliichthys Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 305 (*Etheostoma variata* Kirtland).
- 2239. Pæciliichthys variatus** (Kirtland).
 Ohio River Basin, from western Pennsylvania to Kentucky; scarce.
 Long confused with *Oligocephalus cæruleus*.
Etheostoma variata Kirtland, Zoology of Ohio, 1838, 168, 192, Mahoning River, Ohio.
Hadropterus tessellatus Jordan, Bull. U. S. Nat. Mus., X, 1877, 7, Allegheny River, Foxburg, Pa.
- Genus 769. NANOSTOMA** Putnam.
Nanostoma Putnam, in Jordan, Bull. U. S. Nat. Mus., X, 1877, 6 (*Pæciliichthys zonalis* Cope); not *Nannostomus* Günther.
Rhothæca Jordan, Rept. U. S. Fish Com., XIII, 1885 (1887), 868, p. 80 of reprint issued 1885; substitute for *Nanostoma* Putnam, regarded as preoccupied by *Nannostomus* Günther.
- 2240. Nanostoma zonale** (Cope). *Banded darter.*
 Mississippi Valley from Ohio and northern Indiana (Kankakee River) west through Iowa and south through Kentucky and Tennessee into Alabama, Louisiana, and Arkansas.
Pæciliichthys zonalis Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 212, Holston River, Va.
Nanostoma vinctipes Jordan, Proc. U. S. Nat. Mus., II, 1879, 236, tributary of Illinois River at Naperville, Ill.
- 2241. Nanostoma arcansanum** (Jordan and Gilbert).
 Ozark region of Missouri and Arkansas.
Etheostoma zonale arcansanum Jordan and Gilbert, Proc. U. S. Nat. Mus., IX, 1886, 5, Spring River, Carthage, Mo.; James River, Marshfield, Mo.; Poteau River, Hackett City, Ark.; Washita River, Arkadelphia, Ark., and Saline River, Benton, Ark.
- 2242. Nanostoma swannanoa** (Jordan and Evermann).
 Upper waters of the Tennessee River.
Etheostoma swannanoa Jordan and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 360, Pl. XLV, fig. 13, South Fork of Holston River, Holstein Mills, Va.; Middle Fork of Holston River, Marion, Va. and South Fork of Swannanoa River, Black Mountain, N. C.
- 2243. Nanostoma elegans** (Hay).
 Chickasawha River, Miss.
Nanostoma elegans Hay, Proc. U. S. Nat. Mus., III, 1880, 493, shallow rocky branch of Chickasawha River, Enterprise, Miss.
- 2244. Nanostoma thalassinum** (Jordan and Brayton).
 Santee River Basin in North and South Carolina.
Nothonotus thalassinus Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 13, Reedy River, Greenville, S. C.; Catawba River and tributaries in North Carolina; Ennoree River, Chick Springs, S. C.; and Saluda River, Farr's Mill, S. C.
- 2245. Nanostoma inscriptum** (Jordan and Brayton).
 Oconee River, Ga.
Nothonotus inscriptus Jordan and Brayton, Bull. U. S. Nat. Mus., XII, 1878, 34, Oconee River, Sulphur Springs, Hall County, Ga.

- 2246.** *Nanostoma rupestre* (Gilbert and Swain).
 North River, a tributary of Black Warrior River, Ala.
Etheostoma (Rhothaea) rupestre Gilbert and Swain, Proc. U. S. Nat. Mus., X, 1887, 57, North River near Tuscaloosa, Ala.
- Genus 770. NOTHONOTUS** Agassiz.
Nothonotus Agassiz, Bull. Mus. Comp. Zool., I, 1863, 3 (*Etheostoma maculata* Kirtland).
- 2247.** *Nothonotus maculatus* Kirtland.
 Ohio and Indiana, southward through Kentucky and Tennessee to northern Alabama, in tributaries of the Wabash, Ohio, Cumberland, and Tennessee Rivers.
Etheostoma maculata Kirtland, Jour. Bost. Soc. Nat. Hist., 1840, 276, Mahoning River, Ohio.
Pacilichthys sanguinolentus Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 264, headwaters of the South Fork of the Cumberland River in Tennessee.
- 2248.** *Nothonotus camurus* (Cope). *Blue-breasted darter.*
 Lake Erie to Tennessee, in clear, swift waters; not common; the most brilliantly colored of fresh-water fishes.
Pacilichthys camurus Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 265, headwaters of Cumberland River in Tennessee.
- 2249.** *Nothonotus vulneratus* (Cope).
 French Broad River.
Pacilichthys vulneratus Cope, Proc. Amer. Phil. Soc. Phila., XI, 1870, 266, tributary of French Broad River, Warm Springs, N. C.
- 2250.** *Nothonotus cinereus* (Storer).
 Tennessee and Cumberland Rivers; Tennessee River at Florence, Ala.; Obey River at Olympus, Tenn.; Rock Creek near Whitley Station, Ky.
Etheostoma cinereum Storer, Proc. Bost. Soc. Nat. Hist., II, 1845, 49, Florence, Ala.
- 2251.** *Nothonotus tessellatus* (Storer). *Tessellated darter.*
 Florence, Ala.
Etheostoma tessellata Storer, Proc., Bost. Soc. Nat. Hist., II, 1845, 48, Tennessee River at Florence, Ala. (Not *Boleosoma tessellatum* DeKay).
- 2252.** *Nothonotus rufolineatus* (Cope).
 Upper tributaries of the Tennessee, Cumberland, and Green Rivers.
Pacilichthys rufolineatus Cope, Proc. Amer. Phil. Soc., XI, 1870, 267, Warm Springs Creek, French Broad River, Madison County, N. C.
- 2253.** *Nothonotus jordani* (Gilbert). *Jordan's darter.*
 Tributaries of Coosa River in the Alabama River Basin.
Etheostoma (Nothonotus) jordani Gilbert, Bull. U. S. Fish Com., I, 1889 (1891), 156, pl. 43, fig. 3, Choccolo Creek, Oxford, Ala., and Chestnut Creek, Verbena, Ala.
- Genus 771. RAFINESQUIELLUS** Jordan and Evermann.
Aplecton Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 102 (*A. pottsii* Girard); not of Rafinesque.
Rafinesquius Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1082 (*Aplecton pottsii* Girard).
- 2254.** *Rafinesquius pottsii* (Girard).
 Streams of Chihuahua, Mexico, in Rio Grande Basin.
Aplecton pottsii (misprinted *potsii*) Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 102, tributaries of Chihuahua River, Mexico.
Etheostoma micropterus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 289, Chihuahua, Mexico.
- Genus 772. OLIGOCEPHALUS** Girard.
Oligocephalus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 67 (*Boleosoma lepida* Baird and Girard).

- 2255.** *Oligocephalus cæruleus* (Storer). *Blue darter; Rainbow darter; Soldierfish; Blue johnny.*
Upper Mississippi Valley, from Kentucky to Michigan and east to New York.
Etheostoma cærulea Storer, Proc. Bost. Soc. Nat. Hist., II, 1845, 47, Fox River, Ill.
Pacilosoma erythrogasterum Kirtland, Cleveland Annals Sci., 1854, 4, near Cleveland, Ohio.
Pacilichthys versicolor Agassiz, Amer. Jour. Sci. Arts, 1854, 304, Quincy, Ill.
Pacilichthys spectabilis Agassiz, Amer. Jour. Sci. Arts, 1854, 304, Osage River, Mo.
Pacilosoma transversum Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 326, Lake Superior.
- 2256.** *Oligocephalus lepidus* (Baird and Girard).
Streams of Arkansas, Texas, and Chihuahua; perhaps more than one species included.
Boleosoma lepida Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 388, upper tributaries of the Rio Nueces, Tex.
Oligocephalus grahami Girard, loc. cit., XI, 1859, 102, Devil River, Tex.
Oligocephalus leonensis Girard, loc. cit., 102, Leon River, Tex.
Oligocephalus pulchellus Girard, loc. cit., 102, Gypsum Creek, a tributary to the Canadian River.
- 2257.** *Oligocephalus fricksia* (Hildebrand).
Tributaries of Savannah River.
Etheostoma fricksia Hildebrand, Bull. U. S. Bur. Fish., XXIX, 1924, 7, Brook near Augusta, Ga.
- 2258.** *Oligocephalus iowæ* (Jordan and Meek).
Upper Mississippi Valley, from Iowa, Nebraska, and Colorado north to Assiniboria; abundant and widely distributed.
Etheostoma iowæ Jordan and Meek, Proc. U. S. Nat. Mus., VIII, 1885, 10, Chariton River, Chariton, Iowa.
Etheostoma quappella Eigenmann and Eigenmann, Amer. Nat., 1892, 963, Qu'Apelle River, Manitoba.
- 2259.** *Oligocephalus aubeenaubaei* (Evermann). *Aubeenaubee darter.*
Aubeenaubee Creek, an eastern inlet to Lake Maxinkuckee, Ind.
Etheostoma aubeenaubaei Evermann, Rept., U. S. Fish Com., XXV, 1899 (1900), 367, Aubeenaubee Creek, Lake Maxinkuckee, Ind.
- 2260.** *Oligocephalus hildebrandti* (Evermann and Clark).
Fletcher Lake, Fulton Co., Ind.
Etheostoma hildebrandti Evermann and Clark, Proc. Biol. Soc. Wash., XXIII, May 4, 1910, 81-88, fig. 1, Fletcher Lake, Fulton Co., Ind.
- 2261.** *Oligocephalus jessiae* (Jordan and Brayton). *Jessie's darter.*
Southern Illinois to Georgia and Mississippi.
Pacilichthys jessiae Jordan and Brayton, in Jordan, Man. Vert., ed. 2, 1878, 227; creek near Ringgold, Ga.
Pacilichthys asprigenis Forbes, Bull. Ill. Lab. Nat. Hist., I, No. 2, 1877, 41, small creek at Pekin, Ill.
Pacilichthys swaini Jordan, Proc. U. S. Nat. Mus., VII, 1884, 479, tributary of Pearl River, Monticello, Miss.
- 2262.** *Oligocephalus luteovinctus* (Gilbert and Swain).
Stone River, Tenn.
Etheostoma luteovinctum Gilbert and Swain, Proc. U. S. Nat. Mus., X, 1887, 58, Stone River near Nashville, Tenn.
- Genus 773. NIVICOLA Jordan and Evermann.**
Nivicola Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1082 (*Pacilichthys borealis* Jordan).
- 2263.** *Nivicola borealis* (Jordan). *Northern darter.*
St. Lawrence River at Montreal, Canada, and Muskoka Lakes near Lake Huron.
Pacilichthys borealis Jordan, Proc. U. S. Nat. Mus., VII, 1884, 477, Montreal, Canada.

- 2264. *Nivicola cragini* (Gilbert).** *Cragin's darter.*
 Western portion of the Arkansas River Basin, from Garden City, Kans., to Canyon City, Colo.
Etheostoma cragini Gilbert, Bull. Washburn Coll. Lab., March and April, 1885, 99, small stream connecting the "Lake" at Garden City, Kans., with Arkansas River.
- 2265. *Nivicola punctulata* (Agassiz).**
 Ozark region of southwestern Missouri.
Pacilichthys punctulatus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 304, Osage River, Mo.
- 2266. *Nivicola pagei* (Meek).**
 Tributary of Neosho River, Arkansas Basin.
Etheostoma pagei Meek, Amer. Nat., 1894, 957, Neosho, Mo.; only two specimens known.
- 2267. *Nivicola artesiae* (Hay).**
 Georgia to central Texas.
Pacilichthys artesiae Hay, Proc. U. S. Nat. Mus., III, 1880, 494, from a small branch of Catawba River, Artesia, Lowndes County, Miss.
- 2268. *Nivicola lepidogenys* (Evermann and Kendall).**
 Rio Comal, Tex.
Etheostoma lepidogenys Evermann and Kendall, Bull. U. S. Fish. Com., XII, 1892 (Feb. 6, 1894), 114, pl. 35, fig. 3, Rio Comal, New Braunfels, Tex.
- 2269. *Nivicola virgata* (Jordan).**
 Tributaries of Cumberland River in western Kentucky.
Pacilichthys virgatus Jordan, Proc. U. S. Nat. Mus., II, 1879, 236, Rock Castle River, Livingston, Ky.
- 2270. *Nivicola obeyense* (Kirsch).**
 Tributaries of the Cumberland River in Clinton County, Ky.
Etheostoma obeyense Kirsch, Bull. U. S. Fish Com., X, 1890 (1892), 292, with figure; all streams tributary to the Cumberland River in Clinton County, Ky.
- 2271. *Nivicola tippecanoe* (Jordan and Evermann).**
 Tippecanoe River, Ind.
Etheostoma tippecanoe Jordan and Evermann, Proc. U. S. Nat. Mus., XIII, 1890, 3 (with a figure of *Etheostoma camurum* by error of printer), Tippecanoe River, Marshland, Ind.
- Genus 774. CLARICOLA Jordan and Evermann.**
Claricola Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1093 (*Etheostoma juliae* Meek).
- 2272. *Claricola juliae* (Meek).** *Julia's darter.*
 Basin of White River, Mo.
Etheostoma juliae Meek, Bull. U. S. Fish Com., IX, 1889 (1891), 130, pl. 42, fig. 2, James River, near Springfield, Mo.
- 2273. *Claricola whipplei* (Girard).** *Whipple's darter.*
 Lower Arkansas Basin; locally abundant in clear tributaries of the Saline, Washita, etc.
Boleichthys whipplei Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 103, Coal Creek, Ark.
- 2274. *Claricola alabamæ* (Gilbert and Swain).** *Alabama darter.*
 Black Warrior and Big Catawba Rivers, Ala.
Etheostoma whipplei alabamæ Gilbert and Swain, Proc. U. S. Nat. Mus., X, 1887, 62, Black Warrior River near Morris and Tuscaloosa, Ala.
- 2275. *Claricola squamiceps* (Jordan).**
 Lower Wabash Valley, Ind., through western Kentucky and Tennessee to Georgia and western Florida. A peculiar species resembling *Catotomus*; perhaps not a *Claricola*.
Etheostoma squamiceps Jordan, Bull. U. S. Nat. Mus., X, 1877, 11, Russellville, Ky.
Etheostoma parvipinne Gilbert and Swain, Proc. U. S. Nat. Mus., X, 1887, 59, small spring branch, tributary to Black Warrior River, Tuscaloosa, Ala.

- Genus 775. CATONOTUS Agassiz.** *Fantail darters.*
Calonotus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 305 (*C. lineolatus* Agassiz).
Etheostoma Rafinesque, as restricted by Cope and Jordan.
Richia Coker, Bull. U. S. Bur. Fish., XLII, 1926, 105, 106; *R. brevispina* Coker, preoccupied by *Richia*, a genus of moths.
Richella Coker, Copeia, No. 162, March, 1927, 17, substitute for *Richia*, preoccupied.
- 2276. Catonotus flabellaris** (Rafinesque). *Fan-tailed darter.*
 New York to Virginia, west to Iowa and south to South Carolina and northern Alabama.
Etheostoma flabellare Rafinesque, Jour. de Physique, LXXXVIII, 1819, 419, tributaries of Ohio River.
Etheostoma flabellata Rafinesque, Ichth. Ohiensis, 1820, 36, Ohio River.
Etheostoma fontinalis Rafinesque, loc. cit., 85, Ohio River.
Etheostoma linsleyi Storer, Proc. Boston Soc. Nat. Hist., IV, 1851, 37, Wolcott, Wayne County, N. Y.
Oligocephalus humeralis Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 67, James River, Va.
Catonotus fasciatus Girard, loc. cit., 68, Madrid, N. Y., and Grass River.
Catonotus kennicotti Putnam, Bull. Mus. Comp. Zool., I, 1863, 3, southern Illinois.
Richia brevispina Coker, Bull. U. S. Bur. Fish., XLII, March, 1927, 106, fig. 1, Paddys Creek, N. C.
- 2277. Catonotus flabellare lineolatum** (Agassiz).
 Minnesota to northern Indiana and northern Missouri.
Catonotus lineolatus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 305, small creek near Quincy, Ill.
- 2278. Catonotus flabellare cumberlandicum** (Jordan and Swain). *Cumberland darter.*
 Brooks in Cumberland Mountains tributary to Cumberland River.
Etheostoma cumberlandicum Jordan and Swain, Proc. U. S. Nat. Mus., VI, 1883, 251, small streams of the Cumberland Mountains; Wolf Creek and Briar Creek near Pleasant View, Whitley County, Ky.
- Genus 776. TORRENTARIA Jordan and Evermann.**
Torrentaria Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1080 (*Etheostoma sagitta* Jordan and Swain).
- 2279. Torrentaria sagitta** (Jordan and Swain).
 Cumberland River.
Pæciliichthys sagitta Jordan and Swain, Proc. U. S. Nat. Mus., VI, 1883, 250, Wolf Creek, near Pleasant View, Whitley County, Ky.
- 2280. Torrentaria fasciata** (Girard).
 Chihuahua River, Mexico, in Rio Grande Basin.
Diplesion fasciatum Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 101, Chihuahua River, Mexico; the name preoccupied in *Etheostoma*, not in *Torrentaria*.
Etheostoma australe Jordan, Proc. U. S. Nat. Mus., XI, 1888, 362, Chihuahua River, Mexico; based on the types of *Diplesion fasciatum* Girard.
Etheostoma scovelli Woolman, Amer. Nat., March, 1892, 260, Rio de las Conchas, Chihuahua, Mexico.
- Genus 777. PSYCHROMASTER Jordan and Evermann.**
Psychromaster Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1099 (*Etheostoma tuscumbia* Gilbert and Swain).
- 2281. Psychromaster tuscumbia** (Gilbert and Swain). *Tuscumbia darter.*
 Tennessee River Basin in northern Alabama.
Etheostoma tuscumbia Gilbert and Swain, Proc. U. S. Nat. Mus., X, 1887, 63, stream flowing from the large spring at Tuscumbia, Ala.
- Genus 778. BOLEICHTHYS Girard.**
Boleichthys Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 103 (*Boleichthys exilis* Girard).

- 2282. *Boleichthys fusiformis* (Girard).** *Fusiform darter.*
 Massachusetts and south in coastwise streams.
Boleosoma fusiforme Girard, Proc. Bost. Soc. Nat. Hist., IV, 1854, 41,
 Charles River, Mass.
Hololepis erochrous Cope, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 233,
 Brown's Mill, N. J., opposite Philadelphia.
- 2283. *Boleichthys gracilis* (Girard).** *Crimson darter.*
 Southern Indiana to Georgia and Texas; perhaps more than one species
 included; perhaps varying into the preceding.
Boleosoma gracile Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 103,
 Rio Seco and Leona River, Tex.
Pæciliichthys butterianus Hay, Bull. U. S. Fish Com., II, 1882, 61, Big
 Black River near Vaughns station, Yazoo County, Miss.
Pæciliichthys palustris Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 209,
 Switz City Swamp, Greene County, Ind.
- 2284. *Boleichthys eos* (Jordan and Copeland).**
 New York to Minnesota.
Boleichthys eos Jordan and Copeland, Proc. Acad. Nat. Sci. Phila.,
 XXIX, 1877, 46, Wisconsin and Rock Rivers, Wis.; Fox River, Ill.;
 northern Indiana.
- 2285. *Boleichthys exilis* Girard.**
 Upper Missouri River Basin and Red River of the North.
Boleichthys exilis Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 103,
 Little Muddy River, a tributary of the upper Missouri.
Boleichthys warreni Girard, loc. cit., 104, Cannonball River.
- Genus 779. HOOLEPIS Agassiz.**
Hololepis Agassiz, in Putnam, Bull. Mus. Comp. Zool., I, 1863, 4
 (*Etheostoma barrati* Holbrook).
Copelandellus Jordan and Evermann, Fishes North and Mid. Amer.,
 1896, 1100 (*Pæciliichthys quiescens* Jordan).
- 2286. *Hololepis barratti* Holbrook.**
 Swamps and streams of the lowlands from Virginia to Florida; known
 from Blackwater River, Zuni, Va.; Allaphaha River, Nashville, Ga.;
 and various streams in Florida and South Carolina.
Hololepis barratti Holbrook, Jour. Acad. Nat. Sci. Phila., III, 1855, 56,
 South Carolina.
Pæciliichthys quiescens Jordan, Proc. U. S. Nat. Mus., VII, 1884, 478,
 tributary of Allaphaha River, Nashville, Ga.
- Genus 780. MICROPERCA Putnam.**
Micropurca Putnam, Bull. Mus. Comp. Zool., I, 1863, 4 (*M. punctulata*
 Putnam).
- 2287. *Micropurca punctulata* Putnam.** *Least darter.*
 Northwestern States; Indiana, Michigan, and Minnesota, south to
 Arkansas; rare outside of tributaries of the Great Lakes.
Micropurca punctulata Putnam, Bull. Mus. Comp. Zool., I, 1863, 4,
 various points in Michigan, Wisconsin, Illinois, and Alabama; those
 from Alabama probably *M. præliaris* (Hay); name preoccupied in
Etheostoma.
Etheostoma micropurca Jordan and Gilbert, in Jordan's Man. Vert.,
 ed. V, 1888, 134, lakes of Laporte County, Ind.; substitute name.
- 2288. *Micropurca præliaris* Hay.**
 Alabama and Mississippi, in lowland streams and ponds.
Micropurca præliaris Hay, Proc. U. S. Nat. Mus., III, 1880, 496, small
 branch of the Tuscmibia River at Corinth, Miss.
- Genus 781. ALVARIUS Girard.**
Alvarius Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 101 (*A. lateralis*
 Girard).
 This genus differs from *Micropurca* in having but one anal spine. The
 fins are short and low and the scales in the type are said to be small.
- 2289. *Alvarius lateralis* Girard.**
 Rio Grande about Brazos Santiago, Tex.
Alvarius lateralis Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 101,
 mouth of Rio Grande.

2290. *Alvarius fonticola* Jordan and Gilbert.

Arkansas and Texas, in clear rocky streams; known only from Washita River at Arkadelphia; San Marcos River at San Marcos, Tex.; Comal Creek at New Braunfels, Tex.

Alvarius fonticola Jordan and Gilbert, Proc. U. S. Nat. Mus., IX, 1886, 23, San Marcos Spring at San Marcos, Tex.

Family 187. APOGONIDÆ. Cardinal fishes

Genus 782. APOGON Lacépède.

Amia Gronow, Zoophyl., 1763, 80 (*Apogon moluccensis* Cuvier and Valenciennes); not *Amia* L., nonbinomial.

Apogon Lacépède, Hist. Nat. Poiss., III, 1802, 411 (*A. ruber* Lacépède = *Mullus imberbis* L.).

Ostorrhinchus Lacépède, loc. cit., IV, 1803, 23 (*O. fleurieu* Lacépède).

Dipteron Lacépède, loc. cit., 165 (*D. hexacanthus* Lacépède).

Macrolepis Rafinesque, Analyse de la Nat., 1815, 86; a bare name, no type.

Monoprion Poey, Memorias, II, 1861, 123 (*M. maculatus* Poey).

2291. *Apogon imberbis* (Linnæus). King of the mullets; Alfoncino; Fucinita. Mediterranean and neighboring waters; one from Newport, R. I., and one from the island of Fernando de Noronha.

Mullus imberbis Linnæus, Syst. Nat., ed. X, 1758, 300, Malta; after Artedi.

Apogon ruber Lacépède, Hist. Nat. Poiss., III, 1802, 411, Malta.

Perca pusilla Dc la Roche, Ann. Mus. Nat. Hist., XIII, 1809, 318, Nice.

Apogon rex-mullorum Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 105 (143), Marseilles, Nice, Genoa, Naples, Palermo.

2292. *Apogon maculatus* (Poey). Pensacola to Bahia; recorded from as far north as Woods Hole; common on the "snapper banks"; often found in the stomachs of snappers and groupers.

Monoprion maculatus Poey, Memorias, II, 1861, 123, Cuba.

2293. *Apogon pigmentarius* (Poey).

Cuba; common at Habana.

Monoprion pigmentarius Poey, Memorias, II, 1861, 123, Cuba.

2294. *Apogon sellicauda* Evermann and Marsh.

West Indies; Tortugas Archipelago; Porto Rico.

Apogon sellicauda Evermann and Marsh, Fish. Porto Rico, Bull. U. S. Fish Com., XX, 1900, 143, fig. 40, Culebra Island, Porto Rico.

2295. *Apogon americanus* Castelnau.

Brazil, north to the Tortugas.

Apogon americanum Castelnau, Anim. Amer. Sud, Poiss., 1855, 3, pl. III, fig. 2, Bahia.

2296. *Apogon binotatus* (Poey).

Cuba.

Amia binotata Poey, Repertorio, 1867, 234, Cuba.

2297. *Apogon retrosella* (Gill). Cardenal.

Pacific coast of Mexico.

Amia retrosella Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 251, Cape San Lucas.

2298. *Apogon dovii* Günther.

Mazatlan to Panama.

Apogon dovii Günther, Proc. Zool. Soc. Lond., 1861, 371, Panama.

2299. *Apogon atricaudus* (Jordan and McGregor).

Offshore islands, Pacific coast of Mexico.

Apogon atricaudus Jordan and McGregor, Rept. U. S. Fish Com., XXIV, 1898, 277, and in Jordan and Evermann, Fish. North and Mid. Amer., 1898, 2853, Socorro, Clarion, and San Benedicto islands.

2300. *Apogon atrodorsatus* Heller and Snodgrass.

Galapagos Islands.

Apogon atrodorsatus Heller and Snodgrass, Proc. Wash. Acad. Sci., V, 1903, 192, Pl. III, Charles Island.

- 2301. *Apogon guadalupensis* (Osburn and Nichols).**
Guadalupe Island, off Lower California.
Amia guadalupensis Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 160, fig. 9, Guadalupe Island.
- Genus 783. APOGONICHTHYS Bleeker.**
Apogonichthys Bleeker, Naturkundig Tijd. Nederl.-Indie, Floris, VI, 1854, 321 (*A. perdix* Bleeker).
- 2302. *Apogonichthys alutus* (Jordan and Gilbert).**
Snapper banks off Pensacola and Tampa, Fla.; and at Porto Rico.
Apogon alutus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 279, and in Synopsis, 1883, 931, snapper banks off Pensacola, Fla.
- 2303. *Apogonichthys puncticulatus* Poey.**
Cuba; Tortugas.
Apogonichthys puncticulatus Poey, Repertorio, II, 1867, 233, Cuba.
- 2304. *Apogonichthys melampodus* Blosser.**
St. Croix Island, West Indies.
Apogonichthys melampodus Blosser, Ann. Carnegie Mus., VI, 1909, No. 1, 296, St. Croix.
- Genus 784. ASTRAPOGON Fowler.**
Astrapogon Fowler, Proc. Acad. Nat. Sci. Phila., LXIII, 1906, 527 (*Apogonichthys siellatus* Cope).
- 2305. *Astrapogon stellatus* (Cope).**
Bahamas.
Apogonichthys stellatus Cope, Trans. Amer. Phil. Soc., XIII, 1869, 400, Nassau, Bahama Islands.
Apogonichthys strombi Plate, Zool. Anzeiger, XXXIII, 1908, 393, with plate, Bahamas; living symbiotic in shells of *Strombus gigas*.
- Genus 785. XYSTRAMIA Jordan.**
Xystramia Jordan, Copeia, May 24, 1917, 46 (*Glossamia pandionis* Goode and Bean).
- 2306. *Xystramia pandionis* (Goode and Bean).**
Deep water off Chesapeake Bay.
Apogon pandionis Goode and Bean, Proc. U. S. Nat. Mus., IV, 1881, 160, deep water off Chesapeake Bay.
- Genus 786. EPIGONUS Rafinesque.**
Epigonus Rafinesque, Indice, 1810, 64 (*E. macrophthalmus* Rafinesque).
Pomatomus Risso, in Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 171 (*P. telescopium* Risso); not of Lacépède.
Telescopus Bleeker, Arch. Néerl. Sci. Nat., XI, 1876, 261 (*Pomatomus telescopium* Risso); substitute for *Pomatomus* Cuvier; not of Lacépède.
Pomatominchthys Giglioli, Espos. Int. d. Pesca, Berlin, 1880, 20 (*P. constancia* Giglioli = *Pomatomus telescopium* Risso).
- 2307. *Epigonus occidentalis* Goode and Bean.**
Deep water off Barbados.
Epigonus occidentalis Goode and Bean, Oceanic Ichth., 1895, 233, off Barbados.
- Genus 787. CHEILODIPTERUS Lacépède.**
Cheilodipterus Lacépède, Hist. Nat. Poiss., II, 1802, 539; restricted by Cuvier and Valenciennes, 1828, to *C. macrodon* Lacépède.
Clodipterus Rafinesque, Analyse Nat., 1815, 86, substitute for *Cheilodipterus* Lacépède; regarded as too long.
Paramia Bleeker, Neder. Tijdschr. Dierk., I, 1863, 233 (*Cheilodipterus lineatus* Lacépède); substitute for *Cheilodipterus* (Lacépède) Cuvier.
- 2308. *Cheilodipterus affinis* Poey.**
Cuba; three specimens known.
Cheilodipterus affinis Poey, Ann. Acad. Nat. Sci. N. Y., XI, 1876, 58, Habana.

Family 188. SCOMBROPIDÆ

Genus 788. SPHYRÆNOPS Poey.

Sphyraenops Poey, Memorias, II, 1861, 249 (*S. bairdianus* Poey).2309. *Sphyraenops bairdiana* Poey.

Deep water of Cuba; one specimen known.

Sphyraenops bairdianus Poey, Memorias, II, 1861, 350, Cuba.

Genus 789. PARASPHYRÆNOPS Bean.

Parasphyraenops B. A. Bean, Proc. Biol. Soc. Wash., XXV, 1912, 194
(*P. atrimanus* B. A. Bean).2310. *Parasphyraenops atrimanus* B. A. Bean.

Bermudas.

Parasphyraenops atrimanus B. A. Bean, Proc. Biol. Soc. Wash., XXV, 1912, 194, Bermuda.

Genus 790. LATEBRUS Poey.

Latebrus Poey, Memorias, II, 1860, 168 (*L. oculatus* Poey).2311. *Latebrus oculatus* Poey. *Escalar chino*.

Coast of Cuba; in deep water.

Latebrus oculatus Poey, Memorias, II, 1861, 168, with plate, Cuba.

Genus 791. HYPOCLYDONIA Goode and Bean.

Hypoclydonia Goode and Bean, Oceanic Ichth., 1895, 265 (*H. bella* Goode and Bean).2312. *Hypoclydonia bella* Goode and Bean.

Gulf Stream.

Hypoclydonia bella Goode and Bean, Oceanic Ichth., 1895, 236, Albatross stations 2314, 2397, 2401, 2417, etc., at depths from 90 to 159 fathoms.

Genus 792. AMIICHTHYS Poey.

Amiichthys Poey, in Jordan, Proc. U. S. Nat. Mus., IX, 1886, 586 (*diapterus* Poey, 1861; only the specific name of this fish being published by Poey).2313. *Amiichthys diapterus* Poey.

Coast of Cuba; two specimens known.

diapterus Poey, Synopsis, 1861, 305, Cuba.

✓ Family 189. CENTRARCHIDÆ. Black basses; Sunfishes and pumpkinseeds

Genus 793. HURO Cuvier and Valenciennes. ~ *Largemouth black bass*.*Nemocampsis*⁵² Rafinesque, Ichth. Ohiensis, 1820, 30 (*Lepomis flexuolaris* Raf.).*Huro* Cuvier and Valenciennes, Hist. Nat., Poiss., II, 1828, 125 (*H. nigricans* Cuvier and Valenciennes = *Cichla floridana* LeSueur).2314. *Huro floridana* (LeSueur). Large-mouthed black bass; Oswego bass; Green bass; Bayou bass; Chub; Trout; Jumper; Welshman; Achigan grand bouche; Achigan noir; Big-mouthed black bass; Black bass of the Huron; Black huron; Bride perch; Cow bass; Dotted paintedtail; Freshwater trout; Gray bass; Green perch; Green trout; Growler; Huron; Lake bass; Marsh bass; Millpond chub; Moss bass; Mud bass; Paintedtail; Pale river bass; Perch; River bass; Rock bass; Salmon-formed growler; Slough bass; Southern chub; Speckled hen; Spotted bass; Straw bass; Striped bass; White bass; White salmon; White trout; Yellow bass; Yellow pondperch.Rivers of the United States from the Great Lakes and Red River of the North to Florida, Texas, and Mexico; west to the Dakotas, Nebraska and Kansas; introduced into lakes and streams of California. *Cichla floridana* LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1822, 219, east Florida.*Huro nigricans* Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 93 (124), pl. 17, Lake Huron.*Micropterus salmoides* of authors; not *Labrus salmoides* Lacépède.

⁵² Except for the phrase "Mouth large," Rafinesque gives no distinctive characters for the young black bass he calls "*Nemocampsis*"; but in *Micropterus* the mouth is decidedly large, though smaller than in *Huro*.

Genus 794. MICROPTERUS Lacépède. Black bass.

Micropterus Lacépède, Hist. Nat. Poiss., IV, 325, 1802 (*M. dolomieu* Lacépède).

Calliurus Rafinesque, Jour. de Physique, 1819, 420 (*C. punctulatus* Raf.).

Aplites Rafinesque, Ichth. Ohiensis, 1820, 30 (*Lepomis pallida* Raf.).

?*Nemocampsis* Rafinesque, loc. cit., 30 (*Lepomis flexuolaris* Raf.).

Dioplites Rafinesque, loc. cit., 32 (*Lepomis notata* Raf.).

Aplectes Rafinesque, loc. cit., 36 (*Etheostoma calliura* Raf.).

- 2315. Micropterus dolomieu** Lacépède. Small-mouthed black bass; Achigan noir; Achigan petite bouche; Bass; Bass hogfish; Black bass; Black fresh-water bass; Black perch; Bronze-backer; Brown bass; Brown trout; Dwarf bass; Gold bass; Green bass; Green perch; Green trout; Hog bass; Jumper; Little bass; Minny bass; Mountain trout; Redeye; River bass; Rock bass; Speckled hen; Spotted river bass; Streaked-checked river bass; Streaked lead; Swago bass; Suego; Trout bass; Trout perch; White trout; Yellow bass; Yellow perch.

Lake Champlain to Manitoba and southward on both sides of the mountains to South Carolina and Arkansas.

Micropterus dolomieu Lacépède, Hist. Nat. Poiss., IV, 1802, 325, type locality uncertain, probably South Carolina.

Labrus salmoides Lacépède, loc. cit., IV, 1802, 716, South Carolina.

Bodianus achigan Rafinesque, Amer. Month Mag., II, 1817, 120, New York.

Calliurus punctulatus Rafinesque, Jour. de Physique, 1819, 420, falls of the Ohio River.

Lepomis pallida Rafinesque, Ichth. Ohiensis, 1820, 30, Ohio, Miami, and Hocking Rivers.

Lepomis trifasciata Rafinesque, loc. cit., 31, Ohio River.

?*Lepomis flexuolaris* Rafinesque, loc. cit., 31, Ohio River.

Lepomis salmonneus Rafinesque, loc. cit., 32, Kentucky, Green, Licking, and Ohio Rivers.

Lepomis notata Rafinesque, loc. cit., 32, Ohio River.

Etheostoma calliura Rafinesque, loc. cit., 36, Ohio River; Salt River, Ky.

Cichla fasciata LeSueur, Jour., Acad. Nat. Sci. Phila., II, 1822, 216, Lake Erie.

Cichla ohiensis LeSueur, loc. cit., 218, Ohio River.

Cichla minima LeSueur, loc. cit., 220, Lake Erie.

Centrarchus obscurus DeKay, New York Fauna: Fishes, 1842, 30, pl. 17, fig. 48, Onondaga Creek, N. Y.

Dioplites variabilis Vaillant and Bocourt, Miss. Sci. Mex., Poissons, 1875-1883, 142, Wabash River, New Harmony, Ind.

- 2316. Micropterus pseudoplites** Hubbs. Kentucky black bass.

Southeastern United States from western West Virginia northward to southern Ohio, westward to Kansas and central Texas.

Micropterus pseudoplites Hubbs, Proc. Okla. Acad. Sci., Part I, V, Jan. 1, 1927, 137, nomen nudum.

Micropterus pseudoplites Hubbs, Occ. Papers, Mus. Zool. Univ. Mich. No. 184, June 29, 1927, 8, Pl. I, figs. 2, 3, and 5, Forbush Creek near Mill Springs, Ky.

Genus 795. APOMOTIS Rafinesque.

Apomotis Rafinesque, Jour. de Physique, LXXXVIII, 1819, 420 (*Lepomis cyanellus* Rafinesque).

Telipomis Rafinesque, Ichth. Ohiensis, 1820, 27 (*Lepomis cyanellus* Rafinesque).

Bryttus Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 454, 461 (*B. punctatus* Cuvier and Valenciennes).

- 2317. Apomotis cyanellus** (Rafinesque). Blue-spotted sunfish; Green sunfish; Little red-eye; Blackeyes; Blackeye sunfish; Blue-and-green sunfish; Blue bass; Bluefish; Blue sunfish; Bream; Buffalo sunfish; Perch; Red-eye; Sunfish; Blanco perch; Wood bass.

Great Lakes region to Mexico; very abundant from Ohio southwestward to the Rio Grande.

Lepomis cyanellus Rafinesque, Jour. de Physique, LXXXVIII, 1819, 420, Ohio River.

Icthelis melanops Rafinesque, Ichth. Ohiensis, 1820, 28, Ohio River.

- Pomotis longulus* Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 391, Otter Creek, Ark.
- Calliurus diaphanus* Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 200, Rio Blanco, Tex.
- Calliurus formosus* Girard, loc. cit., 1857, 200, Arkansas.
- Calliurus microps* Girard, loc. cit., 200, Rio Brazos, Tex.
- Calliurus murinus* Girard, loc. cit., 200, Texas.
- Bryttus signifer* Girard, loc. cit., 201, Rio Medina, Tex.
- Bryttus mineopas* Cope, loc. cit., XVII, 1865, 84, Whittlesey Lake, Minneapolis, Minn.
- 2318. *Apomotis euryorus* (McKay).**
Upper Great Lakes region. A form regarded as this occasional in the upper Mississippi.
Lepomis euryorus McKay, Proc. U. S. Nat. Mus., IV, 1881, 89, Lake Huron at Fort Gratiot, Mich. Regarded by Hubbs as a hybrid between *Apomotis cyanellus* and *Eupomotis gibbosus*.
- 2319. *Apomotis ischyurus* (Jordan and Nelson). *Big-nosed sunfish.***
Upper Mississippi Valley.
Lepiopomus ischyurus Jordan and Nelson, Bull. U. S. Nat. Mus., X, 1877, 25, Illinois River, Ill. Rare; thought by Hubbs to be a hybrid.
- 2320. *Apomotis phenax* Cope and Jordan. *Deceptive sunfish.***
Streams of New Jersey.
Apomotis phenax Cope and Jordan, Bull. U. S. Nat. Mus., X, 1877, 26, Beaseley's Point, N. J.; locality possibly erroneous; rare; thought to be a hybrid.
- 2321. *Apomotis punctatus* (Cuvier and Valenciennes). *Bream; Chinquapin perch; Spotted bream.***
South Carolina to Florida.
Bryttus punctatus Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 347 (462), no locality, probably Charleston, S. C.
Bryttus reticulatus Cuvier and Valenciennes, loc. cit., 348 (463), Charleston, S. C.
Lepomis apiatus Cope, Proc. Amer. Phil. Soc., XVII, 1877, 65, Volusia, Fla.
- 2322. *Apomotis symmetricus* (Forbes)**
Mississippi Valley; Illinois to Louisiana and Texas.
Lepomis symmetricus Forbes, in Jordan and Gilbert, Synopsis, 1883, 473, Illinois River, Ill.
- Genus 796. **LEPOMIS** Rafinesque. *Sunfishes.*
Lepomis Rafinesque, Jour. de Physique, LXXXVIII, 1819, 420 (*Labrus auritus* Linnaeus).
Pomotis Rafinesque, loc. cit., 420 (*Labrus auritus* Linnaeus).
Icthelis Rafinesque, Ichth. Ohiensis, 1820, 27 (*Labrus auritus* Linnaeus).
- 2323. *Lepomis pallidus* (Mitchill).**
New York.
Labrus palladus Mitchill, Trans. Lit. and Phil. Soc. N. Y., 1815, 407, New York. Misprint for *pallidus*.
Lepomis appendix Mitchill, Amer. Month. Mag., 1818, 247, New York. An uncertain species, perhaps based on *L. auritus*.
- 2324. *Lepomis auritus* (Linnaeus).** *Yellowbelly; Redbreast bream; Sunfish; Tobaccobox; Long-eared sunfish; Sun perch; Bream; Redbreast; Black-eared pondfish; Flatfish; Flounder; Fresh-water sunfish; Kiver; Leather-ear; Leatherwing; Perch; Quiver; Redbelly; Red-bellied bream; Red-bellied robin perch; Red-headed bream; Red perch; Red sunfish; Red-tailed bream; Red-tailed pomotis; Roach; Robin; Robin perch; Yellow perch; River sunfish; Horned-eared sunfish.*
Maine to Virginia; abundant in all streams east of the Alleghenies and south of New York.
Labrus auritus Linnaeus, Syst. Nat., Ed. X, 1758, 283, Philadelphia.
Bryttus unicolor Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 348 (464), Philadelphia and Charleston.
Pomotis rubri-cauda Storer, Jour. Boston Soc. Nat. Hist., IV, 1842, 177, Concord, N. H.
Lepomis ophthalmicus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 223, 224, Roanoke River, Va.

- 2325.** *Lepomis mystacalis* Cope.
East Florida.
Lepomis mystacalis Cope, Proc. Amer. Phil. Soc., XVII, 1877, 66, east Florida.
- 2326.** *Lepomis solis* (Cuvier and Valenciennes).
Virginia to Louisiana, in coastwise streams.
Pomotis solis Cuvier and Valenciennes, Hist. Nat. Poiss., 1831, 352 (468), Lake Pontchartrain, La.
Pomotis elongatus Holbrook, Jour. Acad. Nat. Sci. Phila., III, 1855, 47, pl. 5, fig. 1, St. Johns River, Fla.
- Genus 797. SCLEROTIS Hubbs.**
Sclerotis Hubbs, Proc. Okla. Acad. Sci., Pt. I, VI, Jan. 1, 1927, 138 (*Lepomis miniatus* Jordan).
- 2327.** *Sclerotis miniatus* (Jordan). *Red perch.*
Mississippi Valley; southern Illinois to Louisiana and Texas.
Lepiopomus miniatus Jordan, Bull. U. S. Nat. Mus., X, 1877, 26, Tangipahoa River.
Lepomis garmani Forbes, Bull. Ill. Lab. Nat. Hist., 1885, 135, Little Fox River at Phillipstown and Wabash River and Drew Pond at Carmi, Ill.
- Genus 798. ALLOTIS Hubbs.**
Allotis Hubbs, Proc. Okla. Acad. Sci., Pt. I, VI, Jan. 1, 1927, 139 (*Bryttus humilis* Girard).
- 2328.** *Allotis humilis* (Girard). *Red-spotted sunfish; Bream; Orange-spotted sunfish; Perch; Sunfish.*
Ohio and Kentucky to the Dakotas, Kansas, and Texas; locally abundant, especially in the lower Missouri Basin.
Bryttus humilis Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 201, Sugar Loaf Creek, Ark.
Bryttus oculatus Cope, loc. cit., XVII, 1865, 83, Lake Whittlesey, Minn.
Lepomis anagallinus Cope, Jour. Acad. Nat. Sci. Phila., 2d ser., VI, January, 1869, 221, Leavenworth, Kans.
- Genus 799. XENOTIS Jordan.**
Xenotis Jordan, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 76 (*Pomotis fallax* Baird and Girard).
- 2329.** *Xenotis megalotis* (Rafinesque.) *Long-eared sunfish; Big-eared sunfish; Blackears; Black-tailed sunfish; Bloody sunfish; Bream; Brilliant sunfish; Common sunfish; Eared sunfish; Long-scaled sunfish; Perch; Red-bellied bream; Redbelly; Red-eyed sunfish; Small green sunfish; Sun perch.*
Michigan to Minnesota, South Carolina, and southward to the Rio Grande.
Ichthelis megalotis Rafinesque, Ichth. Ohiensis, 1820, 29, Kentucky, Licking, and Sandy Rivers, Ky.
Pomotis nitida Kirtland, Bost. Jour. Nat. Hist., III, 1840, 472, pl. 28, fig. 1, Mahoning River, Ohio.
Pomotis breviceps Baird and Girard, Proc. Acad. Nat. Sci. Phila., VI, 1853, 390, Otter Creek, Ark.
Pomotis fallax Baird and Girard, loc. cit., VII, 1854, 24, Elm Creek, Tex.
Pomotis convexifrons Baird and Girard, loc. cit., 1854, 24, Rio Cibolo, Tex.
Pomotis nefastus Baird and Girard, loc. cit., 24, Rio Cibolo and Salado Creek, Tex.
Pomotis sanguinolentus Agassiz, Amer. Jour. Sci. and Arts, XVII, 1854, 301, Huntsville, Ala.
Pomotis inscriptus Agassiz, loc. cit., 1854, 302, Huntsville, Ala.
Pomotis popei Girard, Pacific R. R. Surv., X, 1858, 26, headwaters of Colorado River, Tex.
Lepomis peltastes Cope, Proc. Amer. Phil. Soc., 1870, 454, Huron River, Mich.
Xenotis lythrochloris Jordan, Ann. N. Y. Lyc. Nat. Hist., XI, 1876 (1877), 376, White River.
Xenotis aureolus Jordan, loc. cit., 376, White River.

2330. *Xenotis occidentalis* (Meek).

Rio Grande Basin in Mexico.

Lepomis occidentalis Meek, Field Col. Mus., Zool., III, 1902, 118, Rio Conchos, Mexico.

Genus 800. *HELIOPERCA* Jordan.

Helioperca Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876 (1877), 355
(*Labrus pallidus* Cuvier and Valenciennes; not of Mitchell = *Pomotis incisor* Cuvier and Valenciennes).

Erichzta Jordan, Klippart's Rept. Fish Com. Ohio, 1876 (1877), 48
(*Pomatis incisor* Cuvier and Valenciennes); uncorrected in proof,
reading for *Helioperca*.

2331. *Helioperca incisor* (Cuvier and Valenciennes). *Bluegill; Blue bream;*
Blue sunfish; Copper-nosed bream; Dollardee; Black-eared pondfish;
Blue Joe; Blue-mouthed sunfish; Blue perch; Bream; Copper-headed
bream; Perch; Sunfish.

Great Lakes to Florida and the Rio Grande.

Pomotis incisor Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831,
350 (466), New Orleans.

Pomotis gibbosus Cuvier and Valenciennes, loc. cit., 1831, 351 (467),
Charleston, S. C.; not of Linnaeus.

Pomotis speciosus Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII,
1854, 24, Brownsville, Tex.

Pomotis obscurus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 302,
Huntsville, Ala.; specimens almost black.

Pomotis luna Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 201, Fort
Snelling, Minn.

Lepomis longispinis Cope, loc. cit., 1865, 83, Leavenworth, Kans.

Lepomis ardesiacus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 222,
Kiskiminitas River, Pa.

Lepomis purpurascens Cope, Proc. Amer. Phil. Soc., XI, 1870, 454,
Yadkin River, N. C.

2332. *Helioperca macrochira* (Rafinesque). *Bream; Chain-sided perch; Chain-*
side; Gilded sunfish; Gold perch; Perch; Sunfish.

Ohio Valley and southwestward to Missouri and Kentucky.

Lepomis macrochirus Rafinesque, Jour. de Physique, LXXXVIII,
1819, 420, Ohio, Wabash, Green, and Licking Rivers.

2333. *Helioperca nephela* (Cope).

Western Pennsylvania.

Lepomis nephelus Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 222,
Kiskiminitas River, western Pennsylvania. Perhaps the same as
H. macrochira.

2334. *Helioperca marginata* (Holbrook).

Florida.

Pomotis marginatus Holbrook, Jour. Acad. Nat. Sci. Phila., III, 1855,
49, Pl. VI, fig. 2, St. Johns River, Fla. This and the remaining species
here referred to *Helioperca* are of uncertain value and of doubtful
relationship.

2335. *Helioperca haplognatha* (Cope).

Monterey, Nuevo Leon, Mexico.

Lepomis haplognathus Cope, Proc. Amer. Phil. Soc., XXII, 1884 (1885),
168, Monterey, Nuevo Leon, Mexico.

2336. *Helioperca bombifrons* (Agassiz).

Tennessee River.

Pomotis bombifrons Agassiz, Amer. Jour. Sci. and Arts, XVII, 1854,
303, Huntsville, Ala.

Genus 801. *XYSTROPLITES* Jordan.

Xystroplites Jordan, Bull. X, U. S. Nat. Mus., 1877, 24 (*X. gilli* Jordan =
Pomotis pallidus Agassiz).

2337. **Xystropelites pallidus** (Agassiz).

Georgia to Texas.

Pomotis pallidus Agassiz, Amer. Jour. Sci. Arts, 1854, 303 (name pre-occupied in *Lepomis* by *Labrus palladus* Mitchell, but not in *Eupomotis*).

Bryttus albulus Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 200, Rio Blanco, Tex.

Xystropelites gillii Jordan, Bull. U. S. Nat. Mus., X, 1877, 24, "Garden Key, Florida"; evidently an error, as there is no fresh-water stream on Garden Key.

Lepomis lirus McKay, Proc. U. S. Nat. Mus., IV, 1881, 89; name a substitute for *pallidus*.

Genus 802. **EUPOMOTIS** Gill and Jordan.

Eupomotis Gill and Jordan, Field and Forest, II, 1877, 190 (*Sparus aureus* Walbaum = *Perca gibbosa* Linnaeus).

2338. **Eupomotis heros** (Baird and Girard).

Southern Illinois, Indiana, and Ohio to Florida and the Rio Grande Basin.

Pomotis heros Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, March, 1854, 25, Rio Cibolo, Tex.

Pomotis notatus Agassiz, Amer. Jour. Sci. and Arts, XVII, 1854, 302, Huntsville, Ala.

2339. **Eupomotis holbrookii** (Cuvier and Valenciennes). *Yellow bream; Shellcracker; Strawberry bass; Branch perch.*

Virginia to Florida.

Pomotis holbrookii Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 350 (466), Charleston, S. C. Misprint for *holbrookii*.

Pomotis speciosus Holbrook, Jour. Acad. Nat. Sci. Phila., III, 1855, 48, pl. 5, fig. 2, St. Johns River, Fla.; not of Baird and Girard.

Pomotis microlophus Günther, Cat., I, 1859, 264, St. Johns River, Fla.

2340. **Eupomotis longimanus** (Cope).

Florida to Alabama.

Xystropelites longimanus Cope, Proc. Amer. Phil. Soc., XVII, 1877, 66, Volusia and Port Bay, Fla. Perhaps identical with *E. holbrookii*.

2341. **Eupomotis gibbosus** (Linnaeus). *Common sunfish; Bream; Pumpkinseed; Sunfish; Sunny; Crapet jaune; Female perch; Flatfish; Flounder; Harlequin roach; Kwer; Northern pomotis; Perch; Pond perch; Quiver; Redbelly; Roach; Robin; Robin perch; Ruff; Sand perch; Sun bass; Tobaccobox; Yellowbelly; Yellow perch.*

Great Lakes region to Maine and southward to Florida, east of the Alleghenies.

Perca gibbosa Linnaeus, Syst. Nat., Ed. X, 1758, 293, Carolinas; after *Perca fluviatilis gibbosa, ventre luteo*, of Catesby.

Sparus aureus Walbaum, Artedi Pisc., 1792, 290, lakes of New York.

Morone maculata Mitchell, Fishes of New York, 1814, 18, near New York City.

Pomotis vulgaris Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 67 (91), Lake Huron, New York, Virginia, and the Carolinas.

Pomotis ravenelii Cuvier and Valenciennes, loc. cit., VII, 1831, 349 (465), Philadelphia.

Pomotis catesbeii Cuvier and Valenciennes, loc. cit., 352 (469), Philadelphia.

Genus 803. **CHÆNOBRYTTUS** Gill.

Chænobrytus Gill, Amer. Jour. Sci. Arts, XXXVII, 1864, 92 (*Calliurus melanops* Girard = *Pomotis gulosus* Cuvier and Valenciennes).

Glossopelites Jordan, Man. Vert., ed. 1, 1876, 223 (*Calliurus melanops* Girard).

2342. **Chænobryttus gulosus** (Cuvier and Valenciennes). *Warmouth; Goggle-eye; Bigmouth; Big-mouthed sunfish; Black sunfish; Black warmouth; Bream; Buffalo bass; Chub; Jugmouth; Moremouth bream; Mud chub; Perch; Perchmouth bream; Redeye; Sac-a-lait; Sunfish; Sun trout; Warmouth perch; Warmouth bream; Wide-mouthed sunfish; Yawmouth perch; Indianfish.*

- Eastern United States, from the Great Lakes to Florida and Texas, west to Kansas and the Dakotas; chiefly west or south of the Alleghenies; common in South Carolina.
- Pomotis gulosus* Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 367 (498), Lake Pontchartrain and lagoons about New Orleans.
- Calliurus floridensis* Holbrook, Jour. Acad. Nat. Sci. Phila., III, 1855, 53, Pl. VI, fig. 1, St. Johns River, Fla.
- Calliurus melanops* Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 200, Texas.
- Lepomis gillii* Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 222, 225, Texas.
- Lepomis charybdis* Cope, loc. cit., 223, Texas.
- Chenobrytus antistius* McKay, Proc. U. S. Nat. Mus., IV, 1881, 88, Lake Michigan.
- Genus 804. **AMBLOPLITES** Rafinesque. *Rock basses.*
- Ambloplites* Rafinesque, Ichth. Ohiensis, 1820, 33 (*Lepomis ichtheloides* Rafinesque = *Bodianus rupestris* Rafinesque).
2343. **Ambloplites rupestris** (Rafinesque). *Common rock bass; Redeye; Goggle-eye; Redeye perch; Bream; Bronzed centrarchus; Crapet mondoux; Crapet noir; Crapet vert; Fresh-water bass; Lake bass; Red-eyed bream; Red-eyed sunfish; Sunfish; Sunfish bass; Sun perch; White bass.*
Vermont to Great Lakes region and Manitoba, south to Louisiana; very abundant west of the Alleghenies.
Bodianus rupestris Rafinesque, Amer. Month. Mag., II, 1817, 120, Lakes of New York, Vermont, and Canada.
Icthelis erythrops Rafinesque, Ichth. Ohiensis, 1820, 29, Ohio River.
Lepomis ichtheloides Rafinesque, loc. cit., 32, Ohio River.
Cichla zenez Le Sueur, Jour. Acad. Nat. Sci. Phila., II, 1822, 214, Lake Ontario.
Centrarchus pentacanthus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 65 (88), Wabash River.
2344. **Ambloplites cavifrons** Cope. *Southern rock bass.*
Roanoke River, Va.
Ambloplites cavifrons Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868, 217, Roanoke River, Va.
- Genus 805. **ARCHOPLITES** Gill.
Archoplites Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 165 (*Centrarchus interruptus* Girard).
2345. **Archoplites interruptus** (Girard). *Sacramento perch.*
Sacramento and San Joaquin Rivers, Calif.
Centrarchus interruptus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 129, San Joaquin and Sacramento Rivers, Calif.
Centrarchus maculosus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 7, Sacramento River.
- Genus 806. **ACANTHARCHUS** Gill.
Acantharchus Gill, Amer. Jour. Sci. Arts, XXXVI, 1864, 92 (*Centrarchus pomotis* Baird).
2346. **Acantharchus pomotis** (Baird). *Mud sunfish; Bass sunfish; Mud bass; Mud perch.*
Southern New York to South Carolina.
Centrarchus pomotis Baird, Ninth Smiths. Rept., 1855, 325, New Jersey, New York.
- Genus 807. **MESOGONISTIUS** Gill.
Mesogonistius Gill, Amer. Jour. Sci. Arts, XXXVII, 1864, 92 (*Pomotis chætodon* Baird).
2347. **Mesogonistius chætodon** (Baird). *Black-banded sunfish.*
New Jersey to Maryland.
Pomotis chætodon Baird, Ninth Smiths. Rept., 1855, 324, Cedar Swamp Creek, N. J.
- Genus 808. **ENNEACANTHUS** Gill.
Enneacanthus Gill, Amer. Jour. Sci. Arts, XXXVII, 1864, 92 (*Pomotis obesus* Baird).

Hemiplites Cope, Jour. Acad. Nat. Sci. Phila., 2d ser., VI, 1868, 217
 (*H. simulans* Cope = *Enneacanthus gloriosus*; abnormal example with
 A. IV, the dorsal erroneously stated to be X.
Copelandia Jordan, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 56 (*C.
 eriarcha* Jordan, an abnormal example with D. X; A. IV, of *Ennea-
 canthus gloriosus*).

- 2348.** *Enneacanthus obesus* (Girard). *Banded sunfish; Little bream; Little sunfish; Spotted sunfish; Rock sunfish.*

Charles River, Mass.; to Florida.

Pomotis obesus Girard, Proc. Bost. Soc. Nat. Hist., V, 1854, 40, waters
 about Hingham and in branch of Charles River at Holliston, Mass.

Bryttus fasciatus Holbrook, Jour. Acad. Nat. Sci. Phila., III, 1855, 51,
 Pl. V, fig. 3, St. Johns River, Fla.

Pomotis gullatus Morris, Proc. Acad. Nat. Sci. Phila., XI, 1859, 3,
 Delaware River, Philadelphia.

- 2349.** *Enneacanthus gloriosus* (Holbrook). *Blue-spotted sunfish; Little sunfish; Spotted-finned sunfish; Spotted sunfish.*

Atlantic States; New York to Florida.

Bryttus gloriosus Holbrook, Jour. Acad. Nat. Sci. Phila., III, 1855,
 52, Pl. V, fig. 4, Cooper River, S. C.

Hemiplites simulans Cope, Jour. Acad. Nat. Sci. Phila., VI, 1868,
 217, Tuckahoe Creek near Richmond, Va.; abnormal examples.

Enneacanthus pinniger Gill and Jordan, Bull. U. S. Nat. Mus., X,
 1877, 27, Tar River, Kinston, N. C.

Enneacanthus margarotis Gill and Jordan, loc. cit., 28, Beaseleys
 Point, N. J.

Copelandia eriarcha Jordan, Proc. Acad. Nat. Sci. Phila., XXIX, 1877,
 56, Menominee River, Wis.; evidently an error, an abnormal
 example; the locality for New Jersey.

Genus 809. CENTRARCHUS Cuvier and Valenciennes. *Round basses.*

Centrarchus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 62
 (84) (*Labrus irideus* Lacépède = *Labrus macropterus* Lacépède).

- 2350.** *Centrarchus macropterus* (Lacépède). *Round sunfish; Fliter; Flying perch; Large-finned bass; Long-finned sunfish; Many-spined sunfish; Millpond perch; Perch; Round bass; Sac-a-lait; Shining bass; Sunfish.*

Lowland streams from Virginia southward to Florida and Louisiana;
 northward in the Mississippi Valley to southern Illinois.

Labrus macropterus Lacépède, Hist. Nat. Poiss., III, 1802, 432, Charles-
 ton, S. C.

Labrus irideus Lacépède, loc. cit., IV, 716, Charleston, S. C.

Genus 810. POMOXIS Rafinesque. *Crappies.*

Pomoxis Rafinesque, Amer. Month. Mag., IV, 1818, 41 (*Pomoxis annularis* Rafinesque).

Hyperistius Gill, Amer. Jour. Sci. Arts, 1864, 92 (*Centrarchus hexa-
 canthus* Cuvier and Valenciennes = *Labrus sparoides* Lacépède).

Subgenus POMOXIS Rafinesque.

- 2351.** *Pomoxis annularis* Rafinesque. *Crappie; Bachelor; Sac-a-lait; New-light; Campbellite; Crapet; Bachelor perch; Bridge perch; Chinquapin perch; Goggle-eye; Goldring; John demon; Pale crappie; Papermouth; Shad; Silver perch; Speckled perch; Strawberry perch; Suckley perch; Timber crappie; Tinmouth; Tin perch; White crappie; White perch.*

Eastern United States, from the Great Lakes south to Texas and west
 to Kansas and Nebraska. Introduced into Columbia River and
 other waters on the Pacific coast.

Pomoxis annularis Rafinesque, Amer. Month. Mag., IV, 1818, 41,
 falls of the Ohio River.

Cichla storeria Kirtland, Rept., Zool. Ohio, 1838, 191, Ohio River.

Pomoxis nitidus Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 200,
 Houston River, Ky.

Pomoxys brevicauda Gill, loc. cit., XVII, 1865, 64, North Grand River,
 Mo.

Pomoxys intermedius Gill, loc. cit., 65, no locality.

Pomoxys protacanthus Gill, loc. cit., 66, Tarboro, N. C.

Subgenus HYPERISTIUS Gill.

- 2352. Pomoxis sparoides** (Lacépède). *Calico bass; Grass bass; Barfish; Strawberry bass; Bachelor; Banklick bass; Bitterhead; Calico bream; Chinquapin perch; Crapet; Crappie; Dolly Varden; Goggle-eye; Lake bass; Lake Erie bass; Millpond flier; Razorback; Roach; Rockfish; Sand perch; Silver bass; Silver perch; Speckled bass; Spotted perch; Spotted trout; Straw bass; Sun perch; Tinmouth; White perch; Black crappie.*

Great Lakes and upper Mississippi Valley to New Jersey and southward to Florida, Louisiana, and Texas. Introduced into Pacific coast streams.
Labrus sparoides Lacépède, Hist. Nat. Poiss., III, 1802, 517 and 518, South Carolina.

Cantharus nigro-maculatus LeSueur, in Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 65 (88), Wabash River.

Centrarchus hexacanthus Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 344 (459), Charleston, S. C.

Hyperistius carolinensis Gill, Amer. Jour. Sci. Arts, XXXVI, 1864, 93, South Carolina.

Family 190. ELASSOMIDÆ

Genus 811. ELASSOMA Jordan.

Elassoma Jordan, Bull. U. S. Nat. Mus., X, 1877, 50 (*E. zonata* Jordan).
Elassosoma Boulenger; attempted correction but the name is from Ελάσσωμα, a diminution.

- 2353. Elassoma zonatum** Jordan. *Pigmy sunfish.*

Southern Illinois to Texas and Alabama.

Elassoma zonata Jordan, Bull. U. S. Nat. Mus., X, 1877, 50, Little Red River, White County, Ark.

- 2354. Elassoma evergladei** Jordan. *Everglades pigmy sunfish.*

Swamps of southern Georgia and Florida; locally common in dark waters tributary to the Everglades.

Elassoma evergladei Jordan, Proc. U. S. Nat. Mus., VII, 1884, 323, St. Johns and Suwanee Rivers, Fla.

Family 191. KUHLIIDÆ. Mountain bass; Sesele

Genus 812. SAFOLE Jordan.

Safole Jordan, Proc. U. S. Nat. Mus., XLII, 1912, 655 (*Dules tæniurus* Cuvier and Valenciennes).

- 2355. Safole tæniura** (Cuvier and Valenciennes).

Galapagos and Revillagigedo archipelagoes; locally common; South Seas and Hawaii to Japan; throughout the warm parts of the Pacific, in the open sea, away from coral reefs; not in California.

Dules tæniurus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 85 (114), Java.

Kuhlia arge Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 159, Chatham Island, Galapagos.

Family 192. CENTROPOMIDÆ. Robalos

Genus 813. CENTROPOMUS Lacépède. Robalos.

Centropomus Lacépède, Hist. Nat. Poiss., IV, 1802, 248 (*lucioperca undecimalis*, etc.; restricted to *Sciæna undecimalis* Bloch by Cuvier and Valenciennes).

Oxylabrax Bleeker, Archiv Neerl., XI, 1876, 264 (same type *Centropomus* being restricted to the first species named by Lacépède).

- 2356. Centropomus undecimalis** (Bloch). *Robalo; Sergeantfish; Snook; Brochet de mer.*

Coasts of Florida and Texas, southward among the West Indies to Surinam or beyond; Bay of Panama.

Sciæna undecimalis Bloch, Naturgeschich. Ausl. Fische, VI, 1792, 60, Jamaica.

Centropomous undecim-radiatus Lacépède, Hist. Nat. Poiss., IV, 1802, 268; after Bloch.

Perca loubina Lacépède, loc. cit., 397, Cayenne.

Sphyraena aureoviridis Lacépède, loc. cit., V, 1803, 324, Martinique.

Centropomus appendiculatus Poey, Memorias, II, 1860, 119, Cuba.

2357. *Centropomus viridis* Lockington. *Robalo.*
 Pacific coast of Mexico; common from Gulf of California to Panama.
Centropomus viridis Lockington, Proc. Calif. Acad. Sci., VII, 1876
 (1877), 110, off Ascension Island, Lower California. Probably not
 separable from *C. undecimalis*.
2358. *Centropomus nigrescens* Günther. *Robalo prieto.*
 Pacific coast of tropical America; generally common from Mazatlan to
 Panama.
Centropomus nigrescens Günther, Proc. Zool. Soc. Lond., 1864 (Mar. 22),
 144, Chiapam.
2359. *Centropomus pedimacula* Poey. *Constantino de las aletas prietas;*
Robalito de las aletas prietas.
 Both coasts of tropical America; common in Cuba; Jamaica; south to
 Brazil.
Centropomus pedimacula Poey, Memorias, II, 1860, 122, Habana and
 Cienfuegos, Cuba.
Centropomus medius Günther, Proc. Zool. Soc. Lond., 1864, 144,
 Chiapam.
Centropomus cuvieri Bocourt, Ann. Sci. Nat. Paris, IX, 1868, 91, Haiti.
Centropomus grandoculatus Jenkins and Evermann, Proc. U. S. Nat.
 Mus., XI, 1888, 139, Guaymas, Sonora. Probably distinct.
- 2360 *Centropomus parallelus* Poey.
 Coasts of Cuba; San Domingo, Jamaica, Barbados, and Rio Chagres;
 Guiana, Pernambuco, and Bahia; Porto Rico.
Centropomus parallelus Poey, Memorias, II, 1861, 120, Habana and
 Cienfuegos, Cuba.
Centropomus mexicanus Bocourt, Ann. Sci. Nat. Paris, IX, 1868, 90,
 Gulf of Mexico.
2361. *Centropomus pectinatus* Poey.
 Coasts of Cuba.
Centropomus undecimalis Cuvier and Valenciennes (part; not of Bloch),
 Hist. Nat. Poiss., II, 1828, 102 (*Centropomus pectinatus* Poey, Memo-
 rias, II, 1861, 121; Habana and Cienfuegos, Cuba).
2362. *Centropomus unionensis* Bocourt.
 Pacific coast of Central America; rather common at Panama.
Centropomus unionensis Bocourt, Ann. Sci. Nat. Paris, IX, 1868, 90,
 La Union, San Salvador.
2363. *Centropomus armatus* Gill.
 Pacific coast of Central America; common from Chiapas to Panama.
Centropomus armatus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 163,
 Panama.
2364. *Centropomus robalito* Jordan and Gilbert. *Constantino; Robalito*
las aletas amarillas.
 Pacific coast of Mexico, from Mazatlan to Panama.
Centropomus robalito Jordan and Gilbert, Proc. U. S. Nat. Mus., IV,
 1881, 462, Mazatlan; Acapulco.
2365. *Centropomus ensiferus* Poey.
 West Indies; generally common from Cuba to Surinam.
Centropomus ensiferus Poey, Memorias, II, 1861, 122, Habana.
? *Centropomus brevis* Günther, Proc. Zool. Soc. Lond., 1864, 145,
 locality unknown, tropical America.
Centropomus affinis Steindachner, Ichth. Notizen, I, 1864, 1, pl. 1,
 fig. 1, Rio Janeiro and Cajutuba; Demerara, Guiana.
Centropomus scaber Bocourt, Ann. Sci. Nat., 5 ser., IX, 1868, 90, marshes
 of Belize.

✓ Family 193. MORONIDÆ. White bass

Genus 814. ROCCUS Mitchell. *Striped bass.*

Roccus Mitchell, Fishes of New York, 1814, 24 (*R. striatus* Mitchell = *Sciæna lineata* Bloch).

2366. *Roccus saxatilis* ^{as} (Walbaum). *Striped bass; Rockfish; Rock; Mis-sucke-kequock.*

Atlantic coasts of North America from New Brunswick to Escambia River, Fla.; introduced by the U. S. Fish Commission into Sacramento River, Calif.

Perca saxatilis Walbaum, Artedi Gen. Pisc., 1792, 330, New York.

Perca septentrionalis Bloch and Schneider, Systema Ichthyol., 1801, 90, New York.

Roccus striatus Mitchell, Rep. Fishes N. Y., 1814, 24, New York.

Perca mitchilli Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 413, pl. 3, fig. 4, New York.

Perca mitchilli alternata Mitchell, loc. cit., 415, New York.

Perca mitchilli interrupta Mitchell, loc. cit., 415, New York.

Genus 815. LEPIBEMA Rafinesque.

Lepibema Rafinesque, Ichth. Ohiensis, 1820, 23 (*Perca chrysops* Rafinesque).

2367. *Lepibema chrysops* (Rafinesque). *White bass; White lake bass; sometimes silver bass at Sandusky, Ohio.*

Great Lakes region, upper Mississippi and Ohio Valleys, south to Washita River.

Perca chrysops Rafinesque, Ichth. Ohiensis, 1820, 22, falls of the Ohio.

Labrax multilineatus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1830, 360 (488), Wabash River.

Labrax notatus Richardson, Fauna Bor.-Amer., III, 1836, 8, lower St. Lawrence River.

Labrax albodus DeKay, N. Y. Fauna: Fishes, 1842, 13, pl. 51, fig. 165, Buffalo, N. Y.

Labrax osculatii Filippi, Rev. et. Mag. de Zoologie, 2d series, V, 1853, 164, Canada.

Genus 816. CHRYSOPERCA Fowler.

Chrysoperca Fowler, Proc. Acad. Nat. Sci. Phila., LIX, 1907, 250 (*Morone interrupta* Gill).

2368. *Chrysoperca interrupta* (Gill). *Yellow bass.*

Lower Mississippi Valley north to Cincinnati, Terre Haute, and St. Louis.

Morone interrupta Gill, Proc. Acad. Nat. Sci. Phila., XII, 1860, 118, St. Louis; New Orleans.

Morone mississippiensis Jordan and Eigenmann, Proc. Acad. Nat. Sci. Phila., XXXIX, 1887, 295; substitute for *Morone interrupta*, preoccupied in *Roccus*, not in *Morone* nor in *Chrysoperca*.

Genus 817. MORONE Mitchell. *White perch.*

Morone Mitchell, Fishes of New York, 1814, 17 (*M. pallida* Mitchell = *Perca americana* Gmelin).

2369. *Morone americana* (Gmelin). *White perch; Silver perch; Sea perch.*

Atlantic coast of North America from Nova Scotia to South Carolina in brackish water, ascending streams, and frequently landlocked in ponds.

Perca americana Gmelin, Syst. Nat., I, III, 1789, 1308, New York; after Schöpf.

Perca immaculata Walbaum, Artedi Genera Piscium, 1792, 330, New York; after Schöpf.

Morone rufa Mitchell, Rep. Fishes N. Y., 1814, 18, New York.

Morone pallida Mitchell, loc. cit., 17, New York.

?*Centropomus albus* Rafinesque, Précis des Découvertes Somiolog., 1814, 19, Philadelphia.

^a This species is usually called *Roccus lineatus* after *Sciæna lineata* Bloch (Ausländische Fische, VI, 1792, 62); but it can not be the same. The form, serræ of the preopercle, and the stout spines of the fin, as well as the asserted locality ("Mediterranean") indicate that the species concerned is *Dicentrarchus lupus* of Europe. The only resemblance to *Roccus* is found in the striped color, but Bloch says that the stripes on the sides are yellow.

Perca mucronata Rafinesque, Amer. Month. Mag. and Crit. Rev., II, 1818, 204, Delaware, Schuylkill, and Susquehanna Rivers.
Labrax nigricans DeKay, Nat. Hist. N. Y., Fishes, 1842, 12, pl. 50, fig. 160, Long Island.

✓ Family 194. EPINEPHELIDÆ. Sea basses, groupers, garrupas, etc.

Genus 818. STEREOLEPIS Ayres. Jewfishes.

Stereolepis Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 28 (*S. gigas* Ayres).

Megaperca Hilgendorf, Sitzb. Gesell. Naturf. Freunde, Berlin, 1878, 155 (*M. ishinagi* Hilgendorf).

2370. *Stereolepis gigas* Ayres. California jewfish.

Coast of California, north to the Farallons.

Stereolepis gigas Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 28, southern California.

Stereolepis californicus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 330, California; substitute for *S. gigas*, supposed to be preoccupied in this genus.

Genus 819. POLYPRION Cuvier. Wreckfishes.

?*Polyprion* Cuvier, Règne Animal, ed. 1, II, 1817, 282 (?*Amphiprion americanus* Bloch and Schneider).

2371. *Polyprion americanus* (Bloch and Schneider). Wreckfish; Stone bass; Cernier.

Gulf stream; Grand Banks and New Jersey; also in Europe if *P. massiliensis* is the same, which is not certain.

Amphiprion americanus Bloch and Schneider, Syst. Ichth., 1801, 205, pl. 47; based on a drawing sent by Latham to Schneider representing some fish called in America "Girom"; called *Amphiprion australis* on pl. 47.

?*Scorpaena massiliensis* Risso, Ichth. Nice, 1810, 184, Marseilles.

?*Polyprion cernuum* Cuvier, Mém. du Museum, XI, 1824, 265, Pl. XVII, France.

?*Holocentrus gulo* Risso, Europe Mérid., III, 1826, 367, Nice.

?*Serranus couchii* Yarrell, British Fishes, ed. 1, I, 1836, 12, Great Britain.

Genus 820. GONIOPLECTRUS Gill. Spanish flags.

Gonioplectrus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 236, 237 (*Plectropoma hispanum* Cuvier and Valenciennes).

2372. *Gonioplectrus hispanus* (Cuvier and Valenciennes). Spanish flag; *Ouatilibi espagnol*; *Biajaiba de lo alto*.

West Indies; Cuba.

Plectropoma hispanum Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 298 (396), Martinique.

Genus 821. PETROMETOPON Gill. Enjambre.

. *Petrometopon* Gill, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 105 (*Perca guttatus* Poey = *Sparus cruentatus* Lacépède).

2373. *Petrometopon cruentatus* (Lacépède). Enjambre; Coney; Rock hind.

West Indian fauna, Brazil to Florida Keys; very common on Cuba coast.

Sparus cruentatus Lacépède, Hist. Nat. Poiss., IV, 1802, 156, Martinique; on a copy of Plumier's drawing.

?*Perca guttata* Bloch, Ichthyol., 1792, 312, Martinique, not of Linnaeus.

Saranus apiarius Poey, Memorias, II, 1861, 143, Habana.

2374. *Petrometopon cruentatus coronatus* (Cuvier and Valenciennes).

Brown hind; Petite nègre.

West Indies, north to Key West.

Serranus coronatus Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 278 (371), Martinique.

2375. *Petrometopon panamensis* (Steindachner).

Panama.

Serranus panamensis Steindachner, Ichth. Beitr., IV, 1875, 1, with plate, Panama.

Genus 822. CEPHALOPHOLIS Bloch and Schneider.

Cephalopholis Bloch and Schneider, Syst. Ichth., 1801, 311 (*C. argus* Bloch and Schneider).

Uriphaëton Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 202 (*Serranus*⁶⁵ *phaëton* Cuvier and Valenciennes = *U. microleptes* Swainson).

Enneacentrus Gill, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 105 (*Serranus ouatalibi* Cuvier and Valenciennes = *Labrus fulvus* Linnaeus). *Bodianus* Bloch, in part, restricted by Cuvier and Valenciennes to *Bodianus guttatus*, etc., later restricted to *Bodianus bodianus* = *Harpe rufa* through tautonymy.

- 2376. *Cephalopholis fulvus*⁶⁵ (Linnaeus).** *Guativere; Niggerfish; Yellowfish; Butterfish; Lemon-yellow butterfish; Guativere amarilla.*

West Indies; Bermuda and Florida Keys to Bahia.

Labrus fulvus Linnaeus, Syst. Nat., Ed. X, 1758, 287, Bahamas; after Catesby.

Guativere amarillas Parra, Descr. Dif. Piezas, Hist. Nat. Cuba, 8 lam., V, 1787, fig. 2, Cuba.

Holocentrus auratus Bloch, Naturgesch. Ausl. Fische, IV, 1790, 75, East Indies.

Bodianus guativere Bloch and Schneider, Syst. Ichth., 1801, 336 (after Parra's figure).

- 2377. *Cephalopholis fulvus ruber* (Bloch and Schneider).** *Red guativere; Ouatalibi; Rock hind.*

West Indies, etc.; very common.

Gymnocephalus ruber Bloch and Schneider, Syst. Ichth., 1801, 346, pl. 67, on *Carauna* of Marcgrave; not *Epinephelus ruber* Bloch.

Serranus ouatalibi Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 286 (381), Habana.

Serranus carauna Cuvier and Valenciennes, loc. cit., 288 (384), no locality given.

- 2378. *Cephalopholis fulvus punctatus* (Linnaeus).** *Niggerfish; Negrofish; Black guativere.*

West Indies, etc.; everywhere common.

Perca punctata Linnaeus, Syst. Nat., ed. X, 1758, 291, Bahamas; based on Catesby.

- 2379. *Cephalopholis stellatus* (Blosser).**

St. Croix.

Bodianus stellatus Blosser, Ann. Carnegie Mus., VI, 1909, 297, St. Croix.

- 2380. *Cephalopholis tæniops* (Cuvier and Valenciennes).**

West coast of Africa, straying to Florida.

Serranus tæniops Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 277 (370), Cape Verde Islands.

Genus 823. MENEPHORUS Poey.

Menephorus Poey, Ann. Lyc. Nat. Hist. N. Y., X, 1871, 50 (*Serranus dubius* Poey).

- 2381. *Menephorus dubius* (Poey).**

Cuba.

Serranus dubius Poey, Memorias, II, 1861, 142, Cuba.

- 2382. *Menephorus punctiferus* (Poey).**

Cuba.

Menephorus punctiferus Poey, Enumeratio, 1875, 21, Habana.

Genus 824. ENNEISTUS Jordan and Evermann.

Enneistus Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1147 (*Bodianus acanthistius* Gilbert).

- 2383. *Enneistus acanthistius* (Gilbert).**

Cape Lobos, eastern shore of Gulf of California.

Bodianus acanthistius Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 552, Cape Lobos, eastern shore of Gulf of California at Albatross station 3017, in 58 fathoms.

⁶⁵ A spurious species, *Cephalopolis nigrispinis* to which the tail of a *Fistularia* had been attached.

⁶⁶ The various forms here treated as subspecies seem to inhabit varying depths of water and should be regarded rather as forms of *C. fulvus*.

Genus 825. EPINEPHELUS Bloch. *Groupers.*

- Epinephelus* Bloch, Nat. Ausl. Fische, VII, 1793, 11 (*E. marginalis*).
Cerna Bonaparte, Introd. Fauna Italica, III, 1833, pt. 1 (*Perca gigas* Brünich).
Cromileptes Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 201 (*Serranus altivelis* Cuvier and Valenciennes).
Cynichthys Swainson, loc. cit., 201 (*Perca flavopurpurea* Bennett = *Holothodus flavocæruleus* Lacépède).
Hyporthodus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 98 (*Serranus niveatus* Cuvier and Valenciennes).
Labroperca Gill, loc. cit., XIV, 1862, 236 (*Serranus labriformis* Jenyns).
Schistorus Gill, loc. cit., 237 (*Serranus mystacinus* Poey).
Merus Poey, Ann. Lyc. Nat. Hist. N. Y., X, 1871, 39 (*Serranus gigas* Cuvier and Valenciennes).
Priacanthichthys Day, Proc. Zool. Soc. Lond., 1868, 193 (*P. maderaspensis* Day = *Epinephelus latefasciatus* Temminck and Schlegel).
Homalagrystes Alleyne and Macleay, Proc. Linn. Soc. New South Wales, 1877, 268 (*H. guentheri* Alleyne and Macleay).
Hypserranus Klunzinger, Fische des Rothen Meeres, 1884, 3 (*Serranus morrhua* Cuvier and Valenciennes).

Subgenus SCHISTORUS Gill.

- 2384. *Epinephelus mystacinus* (Poey).** *Cherna de lo alto.*

West Indies south to Brazil.

Serranus mystacinus Poey, Memorias, I, 1852, 52, Tab. 10, fig. 1, Cuba

Subgenus EPINEPHELUS Bloch.

- 2385. *Epinephelus analogus* Gill.** *Cabrilla pinta.*

Pacific coast of tropical America; common on coast of Mexico.

Epinephelus analogus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 163, Panama.

?*Serranus courtadei* Bocourt, Ann. Sci. Nat. Paris, X, 1868, 222, La Union, San Salvador.

- 2386. *Epinephelus adscensionis* (Osbeck).** *Rock hind; Cabra mora.*

West Indies; Florida Keys to Brazil; Ascension and St. Helena Islands; Woods Hole, Mass.

Perca tota Siba, Thesaurus, LII, 1758, Tab. 27.

Trachinus adscensionis Osbeck, Iter Chin., etc., 1757, and English ed., 1771, 96, Ascension Island.

Trachinus punctatus Bonnaterre, Tableau Encycl., 1788, 46; after Osbeck.

Perca stellio Walbaum, Artedi Pisc., 1792, 349; after Siba.

?*Perca maculata* Bloch, Nat. Ausl. Fische, 1792, 92, Martinique; on a figure by Plumier. Name preoccupied in *Epinephelus*.

Trachinus osbeck Lacépède, Hist. Nat. Poiss., II, 1800, 364; after Osbeck.

Sparus atlanticus Lacépède, loc. cit., IV, 1803, 158, pl. 5, fig. 1, Martinique; on a drawing by Plumier.

Serranus nigriculus Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 381 (375); Martinique.

Serranus pixana Cuvier and Valenciennes, loc. cit., 288 (383), after Pira-Pixanga or Gat-Vichof, Marcgrave, Hist. Brazil, 1648, 152.

Serranus aspersus Jenyns, Voy. Beagle, Fishes, 1842, 6, Porto Praya, St. Jago of the Cape Verde Islands.

Serranus impetiginosus Müller and Troschel, Schomburgk's Hist. Barb., 1848, 665, Barbados.

?*Serranus varius* Bocourt, Ann. Sci. Nat., X, 1868, 222, Gulf coast of Mexico.

- 2387. *Epinephelus guaza* (Linnæus).** *Merou; Mérou; Guasa.*

Coasts of southern Europe and western Africa, ranging north to England, southward to Cape of Good Hope, and westward to Rio Janeiro and to Guiana. The American form, *E. mentzeli* (Cuvier and Valenciennes), may prove different.

Labrus guaza Linnæus, Syst. Nat., ed. X, 1758, 285, "Habitat in Pelago."

Perca gigas Brünich, Ichth. Massiliensis, No. 81, 1768, 65, Marseilles.

Holothodus merou Lacépède, Hist. Nat. Poiss., IV, 1803, 377, Marseilles; after Brünich.

- ?*Serranus mentzelii* Cuvier and Valenciennes, loc. cit., II, 1828, 216 (291), coast of Brazil.
- Perca robusta* Couch, Mag. Nat. Hist., V, 1832, 21, fig. 7, Polperro, Cornwall.
- Serranus marginatus* Lowe, Proc. Zool. Soc. Lond., pt. 1, 1833, 142, Madeira.
- Serranus fimbriatus* Lowe, Trans. Cambr. Phil. Soc., 1836, 195, pl. 1, Madeira.
- ?*Epinephelus brachysoma* Cope, Trans. Amer. Phil. Soc., XIV, 1871, 466, Rio Janeiro.
- 2388. *Epinephelus labrifomis* (Jenyns).** Pacific coast of tropical America; Cape San Lucas to Galapagos Islands. *Serranus labrifomis* Jenyns, Voyage Beagle, Fishes, 1842, 8, pl. 3, Galapagos Islands.
- Epinephelus sellicauda* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 250, Cape San Lucas.
- Epinephelus ordinatus* Cope, Trans. Amer. Phil. Soc., XIV, 1871, 466, Panama.
- 2389. *Epinephelus flavolimbatus* Poey.** *Yellow-finned grouper.* West Indies, north to Pensacola.
- Epinephelus flavolimbatus* Poey, Repertorio, I, 1868, 183, Cuba.
- 2390. *Epinephelus niveatus* (Cuvier and Valenciennes).** *Spotted grouper;* *Snowy grouper.* West Indies to Brazil, occasionally northward in the Gulf Stream as far as Newport, R. I., and Woods Hole, Mass.
- Serranus niveatus* Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 285 (380), Brazil.
- Serranus margaritifer* Günther, Cat., I, 1859, 131, South America.
- Serranus conspersus* Poey, Memorias, II, 1860, 139, Cuba.
- Hyporthodus flavicauda* Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 98, Newport, R. I.
- 2391. *Epinephelus striatus* (Bloch).** *Nassau grouper;* *Hamlet;* *Cherna criolla;* *Grouper;* *Rockfish.* West Indian fauna; Key West to Brazil.
- Cherna Parra*, Dif. Piezas, 1787, 50, Habana.
- Anthias stratus* Bloch, Naturgesch. Ausl. Fische, VI, 1792, 125, Martinique; on figure by Plumier.
- Anthias cherna* Bloch and Schneider, Syst. Ichth., 1801, 310, Cuba; after Parra.
- Sparus chrysomelanurus* Lacépède, Hist. Nat. Poiss., IV, 1803, 160, Martinique; on a copy of Plumier.
- 2392. *Epinephelus guttatus* (Linnæus).** *Cabrilla;* *Red hind.* West Indies; Carolina to Brazil, occasionally north to Charleston, the Florida Keys, and the Bermudas.
- Perca guttata* Linnæus, Syst. Nat., ed. X, 1758, 292, Brazil; after Marckgrave, in part, but with a type specimen, examined by Lönnberg.
- Cabrilha Parra*, Dif. Piezas, Hist. Nat. Cuba, 1787, 93, pl. 36, Habana.
- Lutjanus lunulatus* Bloch and Schneider, Syst. Ichth., 1801, 329; after Parra, not *L. lunulatus* of Park.
- Serranus catus* Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 250 (373), Martinique.
- Serranus maculosus* Cuvier and Valenciennes, loc. cit., 332; verified by Boulenger.
- Serranus arara* Cuvier and Valenciennes, loc. cit., 283 (377), Habana.
- Epinephelus cubanus* Poey, Repertorio, I, 1868, 202, Cuba.
- Serranus stadhouderi* Vaillant, Miss. Sci. Mex., Poiss., 1877, 69, substitute for *maculosus* regarded as preoccupied by *maculatus*.
- 2393. *Epinephelus angustifrons* (Steindachner).** Coast of Cuba.
- Serranus angustifrons* Steindachner, Verh. Ges. Wien, XIV, 230, Pl. VII, figs. 2 and 3, 1864, Cuba.

- 2394.** *Epinephelus drummond-hayi* Goode and Bean. *Speckled hind; John paw.*
 Bermudas, south Atlantic and Gulf coast of United States; Charleston, S. C.
Epinephelus drummond-hayi Goode and Bean, Proc. U. S. Nat. Mus., I, 1878, 173, 174, Pensacola; Bermuda.
- 2395.** *Epinephelus niphobles* Gilbert and Starks.
 Magdalena Bay, Lower California.
Epinephelus niphobles Gilbert and Starks, Proc. U. S. Nat. Mus., XIX, 1896, 442, Magdalena Bay, Lower California.
- Subgenus *VIVERO*^{*} Jordan and Evermann.
- 2396.** *Epinephelus morio* (Cuvier and Valenciennes). *Red grouper; Cherna americana; Cherna de vivero; Nègre.*
 Atlantic coast of America; Woods Hole to Rio Janeiro.
Serranus morio Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 211 (285), New York and San Domingo.
Serranus erythrogaster DeKay, New York Fauna: Fishes, 1842, 21, pl. 19, fig. 52, Florida.
Serranus remotus Poey, Memorias, II, 1860, 140, Cuba.
- Genus 826. *GARRUPA* Jordan. *Black groupers.*
Garrupa Jordan, Bull. U. S. Fish Com., VIII, 1888 (1890), 353 (*Serranus nigritus* Holbrook).
- 2397.** *Garrupa nigrita* (Holbrook). *Black jewfish; Black grouper; Mero de lo alto.*
 South Atlantic and Gulf coast of the United States, Charleston and Pensacola to Cuba and Brazil; perhaps straying to Sicily.
Serranus nigritus Holbrook, Ichth. South Carolina, Ed. I, 1855, 173, pl. 25, fig. 11, Charleston, S. C.
Centropristes merus Poey, Synopsis Piscium Cubensem, 1868, 288, Cuba.
?Cerna sicana Doderlein, Rivista delle Specie del Genere *Epinephelus* o *Cerna*, 1882, 81, Palermo.
- Genus 827. *PROMICROPS* Gill. *Guasas.*
Promicrops Gill, in Poey, Synopsis, 1868, 287 (*Serranus guasa* Poey = *Itaiara* Lichtenstein).
Itaiara Vaillant and Bocourt, Miss. Sci. Mexique, 1875, 90 (*Serranus itaiara* Lichtenstein).
- 2398.** *Promicrops itaiara* (Lichtenstein). *Guasa; Spotted jewfish; Mero.*
 Atlantic coast of tropical America, from Florida to Brazil; not uncommon about rocks.
Cugupuguacu Catesby, Nat. Hist. Carolina, 1771, 14; nonbinomial.
Serranus itaiara Lichtenstein, Abhandl. Berlin Acad. Wiss., 1821 (1822), 278, Brazil; not *Itaiara* of Marcgrave.
Serranus galeus Müller and Troschel, in Schomburgk's Reise in Brit. Guiana, 1848, 621, Guiana.
Serranus guasa Poey, Memorias, II, 1861, 141; 1861, 354, Cuba.
- 2399.** *Promicrops quinquefasciata* (Bocourt).
 Pacific coast of Central America.
Serranus quinquefasciatus Bocourt, Ann. Sci. Nat., X, 1868, 223, Naguate, Pacific coast of Guatemala; very close to *P. itaiara*.
- Genus 828. *ALPHESTES* Bloch and Schneider.
Alphestes Bloch and Schneider, Syst. Ichth., 1801, 236 (*Epinephelus afer* Bloch).
Prospinus Poey, in Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 287 (*Plectropoma chloropterum* Cuvier and Valenciennes = *Alphestes afer* Bloch).
- 2400.** *Alphestes afer* (Bloch).
 West Indies; Cuba to Brazil; recorded perhaps doubtfully from Guinea, if so, the American species, *A. chloropterus*, may prove different.
Epinephelus afer Bloch, Ichthyol., 1793, pl. 327, Acará in Guinea.
Plectropoma chloropterum Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 300 (398), San Domingo, Martinique.

* The subgenus *Vivero* is based on *Epinephelus morio* (Cuv. & Val.) which differs from all other species of *Epinephelus* in the elevation of the second dorsal spine.

- Plectropoma monacanthus* Müller and Troschel, in Schomburgk's Hist. Barbados, 1847, 665, Barbados.
- ?*Serranus armatus* Osorio, Jour. Sci. Lisbon, III, 1894, 74.
- 2401. *Alpestes multiguttatus* (Günther).**
Pacific coast of tropical America; Mazatlan to Panama.
Plectropoma multiguttatum Günther, Proc. Zool. Soc. Lond., 1866, 600, Panama.
- 2402. *Alpestes lightfooti* (Fowler).**
San Domingo, West Indies.
Epinephelus lightfooti Fowler, Proc. Acad. Nat. Sci. Phila., LIX, 1907, 258, fig. 3, San Domingo.
- Genus 829. *DERMATOLEPIS* Gill.**
Dermatolepis Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 54 (*Derma-*
tolepis punctatus Gill).
Lioperca Gill, loc. cit., XIV, 1862, 237 (*Serranus inermis* Cuvier and Valenciennes).
- 2403. *Dermatolepis punctatus* Gill.**
West coast of Mexico; Cape San Lucas; the Venados; the Revillagigedos.
Dermatolepis punctatus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 54, Cape San Lucas.
- 2404. *Dermatolepis zanclus* Evermann and Kendall.**
Key West.
Dermatolepis zanclus Evermann and Kendall, Bull. U. S. Fish Com., XVII, 1897, 129, pl. 8, fig. 9, Key West, Fla.
- Subgenus *LIOPERCA* Gill.**
- 2405. *Dermatolepis inermis* (Cuvier and Valenciennes).**
West Indies.
Serranus inermis Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 322 (436), Antilles.
- Genus 830. *TRISOTROPIS* Gill.**
Trisotropis Gill, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 104 (*Johnius*
guttatus Bloch and Schneider = *Perca venenosa* Linnaeus).
Archoperca Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1171 (*Mycteroperca boulengeri* Jordan and Starks).
- Subgenus *ARCHOPERCA* Jordan and Evermann.**
- 2406. *Trisotropis boulengeri* (Jordan and Starks).** *Cabrilla de raízero; Man-*
grove grouper.
Mazatlan, Mexico.
Mycteroperca boulengeri Jordan and Starks, in Jordan, Fishes Sinaloa,
Proc. Calif. Acad. Sci., V, 1895, 445, pl. 38, taken in the Astillero at
Mazatlan, Mexico.
- Subgenus *TRISOTROPIS* Gill.**
- 2407. *Trisotropis venenosus* (Linnaeus).** *Rockfish; Yellow-finned grouper;*
Bonaci de Piedra.
Bahamas, Florida Keys, and southward.
Perca venenosa Linnaeus, Syst. Nat., Ed. X, 1758, 292, Bahamas; after
Catesby.
Serranus petrosus Poey, Memorias, II, 1861, 136, Cuba.
- 2408. *Trisotropis venenosus apua* (Bloch).** *Bonaci cardenal.*
West Indies, Florida Keys, and southward to Brazil. A red form of
T. venenosus.
Bodianus apua Bloch, Nat. Ausl. Fische, IV, 1790, 50, Brazil; after a
drawing by Prince Maurice—the same used by Marcgrave.
Pirati apia Marcgrave, Hist. Brasil., 1648, 158, Brazil.
Bonaci cardenal Parra, Piezas de Hist. Nat. Cuba, 1787, 29, Pl. XVI,
Habana.
Johnius guttatus Bloch and Schneider, Syst. Ichth., 1801, 77; after
Parra.
Bodianus marginatus Bloch and Schneider, loc. cit., 331; after Marc-
grave.
Serranus rupestris Cuvier and Valenciennes, Hist. Nat. Poiss., IX,
1833, 323 (437), San Domingo.

2409. *Trisotropis bonaci* (Poey). *Marbled rockfish; Bonaci arara; Black grouper; Aguaji; Bonaci.*
 West Indies; Pensacola to Brazil; Woods Hole, Mass.
Serranus bonaci Poey, Memorias, II, 1861, 129, Cuba.
Serranus brunneus Poey, loc. cit., 131, Cuba.
Serranus decimalis Poey, loc. cit., 138, Cuba; specimen with ten dorsal spines.
Trisotropis aguaji Poey, Repertorio, II, 1868, 229, Habana.
Serranus cyclopomatus Poey, Memorias, 1861, 353, Cuba.
Serranus latepictus Poey, loc. cit., 353, Cuba.
2410. *Trisotropis xanthostictus* (Jordan and Swain).
 Snapper Banks off Pensacola, Fla.
Mycteroperca bonaci xanthosticta Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884, 371, Snapper Banks off Pensacola, Fla.
2411. *Trisotropis jordani* (Jenkins and Evermann). *Cabrilla de astillero; Baya.*
 Pacific coast of Mexico from Gulf of California to Mazatlan.
Epinephelus jordani Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 140, Guaymas, Sonora.
2412. *Trisotropis microlepis* Goode and Bean. *Gog; Aguaji.*
 South Atlantic and Gulf coast of United States, north to Virginia, south to Pensacola and Key West.
Trisotropis microlepis Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 141, west Florida.
Trisotropis stomias Goode and Bean, loc. cit., V, 1882, 427, Pensacola; Key West.
2413. *Trisotropis interstitialis* (Poey).
 Coast of Cuba; Woods Hole, Mass.
Serranus interstitialis Poey, Memorias, II, 1861, 127, Cuba.
Trisotropis chlorostomus Poey, Repertorio, II, 1868, 231, Cuba.
2414. *Trisotropis dimidiatus* (Poey).
 Coast of Cuba.
Serranus dimidiatus Poey, Memorias, II, 1861, 129, Cuba.
2415. *Trisotropis xenarchus* (Jordan).
 Pacific coast of Mexico to the Venados Islands near Mazatlan and to the Galapagos Archipelago.
Mycteroperca xenarcha Jordan, Proc. Acad. Nat. Sci. Phila., XXXIX, 1887, 387, Galapagos Islands; Payta, Peru.
- Genus 881. **MYCTEROPTERA** Gill.
Mycteroperca Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 80 (*Serranus olfax* Jenyns).
2416. *Mycteroperca venadorum* Jordan and Starks. *Garlopa.*
 West coast of Mexico at Mazatlan.
Mycteroperca venadorum Jordan and Starks, in Jordan, Proc. Calif. Acad. Sci., V, 1895, 446, Mazatlan, Mexico.
2417. *Mycteroperca bowersi* Evermann and Marsh.
 Culebra and Vieques Islands, Porto Rico.
Mycteroperca bowersi Evermann and Marsh, Bull. Bur. Fish., XX, 1900 (1902), 158, Culebra Island.
2418. *Mycteroperca hopkinsi* Jordan and Rutter.
 Jamaica.
Mycteroperca hopkinsi Jordan and Rutter, Proc. Acad. Nat. Sci. Phila., XLIX, 1897, 105, Jamaica.
2419. *Mycteroperca olfax* (Jenyns). *Yellow grouper; Gourami.*
 James Island, Galapagos Archipelago; Panama.
Serranus olfax Jenyns, Voyage Beagle, Fishes, 1840, 9, pl. 4, Galapagos Islands.
2420. *Mycteroperca rosacea* (Streets). *Cabrilla calamaria.*
 Gulf of California.
Epinephelus rosaceus Streets, Bull. U. S. Nat. Mus., VII, 1877, 51, Angel Island, Gulf of California.

- 2421.** *Mycteroperca falcata* (Poey). *Scamp; Bacalao; Abadejo.*
West Indies, north to Bermuda.
Serranus falcatus Poey, Memorias, II, 1861, 138, Habana.
- 2422.** *Mycteroperca phenax* Jordan and Swain. *Scamp; Bacalao.*
Coasts of southern Florida; abundant about the keys.
Mycteroperca falcata phenax Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884, 363, Key West, Fla.
- 2423.** *Mycteroperca calliura* Poey.
Coast of Cuba.
Mycteroperca calliura Poey, Repertorio, I, 1867, 181, Cuba.
- 2424.** *Mycteroperca tigris* (Cuvier and Valenciennes). *Bonaci gato; Rockfish.*
West Indies, north to Bermuda.
Serranus tigris Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 325 (440), San Domingo.
Serranus felinus Poey, Memorias, II, 1861, 134, Habana.
Serranus repandus Poey, loc. cit., 135, Habana.
Trisotropis reticulatus Gill, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 105, Barbados.
- 2425.** *Mycteroperca camelopardalis* (Poey).
West Indies.
Serranus camelopardalis Poey, Memorias, II, 1861, 132, Cuba, a bright red form, otherwise similar to *M. tigris*.
- Genus 832. XYSTROPERCA** Jordan and Evermann.
Xystroperca Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1181 (*Mycteroperca pardalis* Gilbert).
- 2426.** *Xystroperca pardalis* (Gilbert). *Cabrilla pirlita.*
Gulf of California.
Mycteroperca pardalis Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 551, La Paz Bay, Lower California.
- Genus 833. PAREPINEPHELUS** Bleeker.
Parepinephelus Bleeker, Systema Percarum Revisum, in Archiv Neer. des Sci. Exet. Nat., XI, 1875, 1876, 257 (*Serranus acutirostris* Cuvier and Valenciennes = *Epinephelus ruber* Bloch).
- 2427.** *Parepinephelus acutirostris* (Cuvier and Valenciennes). *Scirenga.*
West Indies, Brazil, Mediterranean Sea, and islands of the eastern Atlantic.
?*Epinephelus ruber*⁶⁷ Bloch, Nat. Ausl. Fische, VII, 1793, 22, pl. 330, "Japan"; probably from the Mediterranean.
Serranus acutirostris Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 212 (286), Brazil.
Serranus undulosus Cuvier and Valenciennes, loc. cit., 219 (295), Brazil.
Serranus tinca Cantraine, "Nouv. Mem. Ac. Brux., XI, 1831," Naples.
Serranus fuscus Lowe, Trans. Cambr. Philos. Soc., VI, 1838, 196, Madeira.
Cerna nebulosa Cocco, "Indice Pesci. Messina, Gen. 45, sp. 2," 1844, Messina.
Cerna macrogenis Sassi, "Descr. Genovæil Genovasato, I, 139," 1846, Genoa.
Serranus emarginatus Valenciennes, Ichth. îles Canaries, 1839, 10, Canary Islands.
Epinephelus chalinus Cope, Trans. Amer. Phil. Soc., XIV, 1871, 465, St. Martins.
Cerna acutirostris lata Doderlein, loc. cit., 74, Palermo.
- Genus 834. CHORISTISTIUM** Gill.
Chorististium Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 15 (*Liopropoma rubra* Poey).
- 2428.** *Chorististium rubrum* (Poey).
Coast of Cuba.
Liopropoma rubre Poey, Memorias, II, 1861, 418, Habana.

⁶⁷ The original type of *E. ruber* has 24 gill rakers; all others examined (Athens, Brazil) have 30 to 35; for these we retain the name *acutirostris* pending investigation.

- Genus 835. LIOPROPOMA Gill.**
Liopropoma Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 52 (*Perca aberrans* Poey).
- 2429. Liopropoma aberrans (Poey).**
 Coast of Cuba.
Perca aberrans Poey, Memorias, II, 1861, 125, Cuba.
- 2430. Liopropoma longilepis Garman.**
 Eastern Pacific in deep water.
Liopropoma longilepis Garman, Mem. Mus. Comp. Zool., XXIV, 1889, 47, lat. $7^{\circ} 30' N.$, long. $78^{\circ} 34' W.$
- Genus 836. RYPTICUS Cuvier. Soapfishes.**
Rypticus Cuvier, Règne Animal, ed. 2, II, 1829, 144 (*Anthias saponaceus* Bloch and Schneider).
Smecticus Valenciennes, Ichth. Voy. Vénus, 1855, 305 (*S. bicolor* Valenciennes).
Eleutheractis Cope, Trans. Amer. Phil. Soc., XIV, 1871, 467 (*E. coriaceus* Cope).
Rhypticus authors, corrected spelling.
- 2431. Rypticus saponaceus (Bloch and Schneider).** *Soapfish; Jabon; Jaboncillo.*
 West Indies; Pensacola to west Africa and Brazil.
Jaboncillo Parra, Difer. Piezas de Hist. Nat., 1787, 51, lam. 24, fig. 2, Habana.
Anthias saponaceus Bloch and Schneider, Syst. Ichth., 1801, 310, Habana.
- 2432. Rypticus arenatus Cuvier and Valenciennes.**
 West Indies and coast of Brazil; recorded from Jamaica, Trinity, Barbados, and St. Thomas.
Rypticus arenatus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 48 (65), pl. 46, Brazil.
?Rhypticus subbifrenatus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 53, St. Thomas.
?Rhypticus nigromaculatus Steindachner, Ichth. Notiz., VI, 1867, 42, Barbados; dorsal with four spines instead of three.
- 2433. Rypticus coriaceus (Cope).** *Black soapfish.*
 West Indies; recorded from St. Martins and Jamaica.
Eleutheractis coriaceus Cope, Trans. Amer. Phil. Soc., XIV, 1870 (1871), 467, St. Martins.
- 2434. Rypticus xanti (Gill).**
 Pacific coast of Mexico, from Cape San Lucas to Colima and Mazatlan.
Rhypticus xanti Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 250, Cape San Lucas.
- 2435. Rypticus bicolor (Valenciennes).**
 Galapagos Archipelago.
Smecticus bicolor Valenciennes, Voy. Vénus, Poissons, 1855, 307, pl. 2, fig. 2, Galapagos Archipelago.
- Genus 837. PROMICROPTERUS Gill.**
Promicropterus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 53 (*Rhyp-ticus maculatus* Holbrook = *Bodianus bistrispinus* Mitchell).
- 2436. Promicropterus bistrispinus (Mitchill).** *Soapfish.*
 South Atlantic coast of United States; frequent off Charleston, Pensacola, and Key West; Porto Rico; occasional as far north as Newport, R. I.
Bodianus bistrispinus Mitchell, Amer. Month. Mag. and Crit. Review, II, February, 1818, 247, Straits of Bahama.
Rypticus maculatus Holbrook, Ichth. South Carolina, Ed. I, 1855, 39, Cape Romain, S. C.
Rhypticus pituitosus Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 341, Key West, Fla.

2437. **Promicropterus nigripinnis** (Gill).

Pacific coast of tropical America, Cape San Lucas to Panama.

Rhypticus nigripinnis Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 53,
Panama.

Rhypticus maculatus Gill, loc. cit., XIV, 1862, 251, Cape San Lucas.
Promicropterus decoratus Gill, loc. cit., XV, 1863, 164, Panama.

✓ Family 195. SERRANIDÆ. Sea bass

Genus 838. **PARALABRAX** Girard. *Cabrillas verdes*.

Paralabrax Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 131 (*Labrax nebulifer* Girard).

2438. **Paralabrax nebulifer** (Girard). *Johnny verde; Kelp bass.*

Southern California, from Monterey to Magdalena Bay.

Labrax nebulifer Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 142,
Monterey, Calif.

2439. **Paralabrax maculatofasciatus** (Steindachner). *Spotted cabrilla.*

Pacific coast of America from Lower California to San Pedro and
Mazatlan.

Serranus maculatofasciatus Steindachner, Ichth. Notizen, VII, 1868, 5,
Mazatlan, Mexico.

?*Serranus acanthophorus* Bocourt, Ann. Sci. Nat., X, 1870, 223, Pacific
coast of Mexico.

2440. **Paralabrax humeralis** (Cuvier and Valenciennes).

Pacific coast of South America; Panama to Juan Fernandez.

Serranus humeralis Cuvier and Valenciennes, Hist. Nat. Poiss., II,
1828, 183 (246), Chile.

Serranus albomaculatus Jenyns, Zool. Beagle, Fishes, 1840, 3, pl. 2,
Galapagos Archipelago.

?*Serranus semifasciatus* Guichenot, in Gay, Hist. Chile, II, 1847, 151,
pl. 1, fig. 2, Juan Fernandez.

2441. **Paralabrax clathratus** (Girard). *Cabrilla; Rock bass.*

Coast of southern California, from San Francisco to Cedros Island.

Labrax clathratus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 143,
San Diego.

Genus 839. **CRATINUS** Steindachner.

Cratinus Steindachner, Ichth. Beitr., VII, 1878, 19 (*C. agassizii* Stein-
dachner).

2442. **Cratinus agassizii** Steindachner.

Galapagos Archipelago.

Cratinus agassizii Steindachner, Ichth. Beitr., VII, 1878, 19, Galapagos
Islands.

Genus 840. **CENTROPRISTES** Cuvier and Valenciennes. *Black sea bass.*

Centropristes Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829,
27 (36) (*C. nigricans* Cuvier and Valenciennes = *Labrus striatus*
Linnaeus).

Centropristes of authors; variation in spelling.

2443. **Centropristes striatus** (Linnæus). *Black sea bass; Blackfish; Tallywag;*

Hannahill; Black Will; Black Harry; Black perch; Black bass; Bluefish.

Atlantic coast of United States, Cape Ann to northern Florida.
Labrus striatus Linnæus, Syst. Nat., ed. X, 1758, 285, "America";
description very brief, but not to be referred to any other fish.

Perca atraria Linnæus, loc. cit., ed. XII, 1766, 485, Carolina.

Perca furva Walbaum, Artedi Pisc., 1792, 336, New York; after Schöpf.
Coryphaena nigrescens Bloch and Schneider, Syst. Ichth., 1801, 297,
New York.

Lutjanus trilobatus Lacépède, Hist. Nat. Poiss., IV, 1802, 246, locality
unknown.

Perca varia Mitchell, Trans. Lit. & Phil. Soc. N. Y., I, 1815, 415, pl. 3,
fig. 6, New York.

- 2444.** *Centropristes oxyurus* (Jordan and Evermann). *Gulf sea bass.*
 Gulf of Mexico at Snapper Banks off Pensacola, Fla.
Serranus oxyurus Jordan and Evermann, Proc. U. S. Nat. Mus., IX,
 1886, 468, Snapper Banks off Pensacola, Fla.
- 2445.** *Centropristes rufus* Cuvier and Valenciennes. *Hogfish.*
 Martinique.
Centropristes rufus Cuvier and Valenciennes, Hist. Nat. Poiss., III,
 1829, 35 (47), Martinique.
- Genus 841. TRILOBURUS** Gill.
Triloburus Gill, Cat. Fish. East Coast U. S., Supplement to Proc. Acad.
 Nat. Sci. Phila., XIII, 1861, 30 (*Perca trifurca* Storer = *Perca philadelphica* Linnæus).
- 2446.** *Triloburus philadelphicus* (Linnæus). *Rock sea bass; Redfish.*
 Coast of South Carolina.
Perca philadelphica Linnæus, Syst. Nat., ed. X, 1758, 291, America.
Perca trifurca Linnæus, loc. cit., ed. XII, 1766, 489, Carolinas.
Lutjanus tridens Lacépède, Hist. Nat. Poiss., IV, 1802, 246, Carolinas.
- Genus 842. DIPLECTRUM** Holbrook. *Squirrelfishes.*
Diplectrum Holbrook, Ichth. South Carolina, ed. I, 1855, 32 (*Serranus fascicularis* Cuvier and Valenciennes = *Perca formosa* Linnæus).
Haliperca Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 236, (*Serranus bivittatus* Cuvier and Valenciennes = *Serranus radialis* Quoy and Gaimard, and other species).
Paraserranus Bleeker, Verh. Akad. Amst., XIV, 1873, 6 (*P. hasseltii* Bleeker).
- Subgenus HALIPERCA** Gill.
- 2447.** *Diplectrum radiale* (Quoy and Gaimard). *Aguavina.*
 Cuba to Brazil.
Serranus radialis Quoy and Gaimard, Voyage *Uranie*, 1824, 316, Rio Janeiro.
Serranus bivittatus Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 179 (241), Martinique.
Centropristes ayresi Steindachner, Ichth. Notizen, VII, 1868, 1, pl. 1, fig. 1, Santos, Brazil.
- 2448.** *Diplectrum pacificum* Meek and Hildebrand.
 Panama.
Diplectrum radiale Gilbert and Starks, Memoir Calif. Acad. Sci., IV, 1904, 97 (not of Quoy and Gaimard).
Diplectrum pacificum Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, Apr. 15, 1925, 477, Pl. XLVI, fig. 2, Naos Island, Panama.
- 2449.** *Diplectrum macropoma* (Günther).
 Pacific coast of tropical America.
Centropristes macropoma Günther, Proc. Zool. Soc. Lond., 1864, 145, Panama.
- 2450.** *Diplectrum sciurus* Gilbert.
 Gulf of California.
Diplectrum sciurus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 550, Gulf of California at Albatross stations Nos. 3014, 3021, 3026, and 3033, all in shallow water.
- 2451.** *Diplectrum euryplectrum* Jordan and Bollman.
 Coast of Colombia, southwest of Panama.
Diplectrum euryplectrum Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 157, Pacific Ocean, off coast of Colombia.
- Subgenus DIPLECTRUM** Holbrook.
- 2452.** *Diplectrum formosus* (Linnæus). *Squirrelfish; Sandfish; Serrano.*
 West Indies; common on the south Atlantic and Gulf coasts of the United States, from Charleston south to Montevideo.
Perca formosa Linnæus, Syst. Nat., ed. XII, 1766, 488, Carolinas.
Epinephelus striatus Bloch, Naturgesch. Ausl. Fische, VII, 1793, 20, Jamaica (not *Anthias striatus* Bloch).
Serranus radians Quoy and Gaimard, Voy. *Uranie*, Poiss., 1824, 313, pl. 58, fig. 2, Rio de Janeiro.

Serranus irriadians Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 181 (244), Montevideo.
Serranus fascicularis Cuvier and Valenciennes, loc. cit., 182 (245), pl. 30, Brazil.

Genus 843. PRIONODES Jenyns.

Prionodes Jenyns, Voy. *Beagle*, Fishes, 1842, 46 (*P. fasciatus* Jenyns).

2453. *Prionodes aequidens* (Gilbert).

Gulf of California.

Serranus aequidens Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 61, Gulf of California at Albatross station No. 2996, in 112 fathoms.

2454. *Prionodes fusculus* (Poey).

Cuba.

Centropristes fusculus Poey, Memorias, II, 1861, 342, Cuba.

2455. *Prionodes phœbe* (Poey). *Phœbe*.

West Indies, north to Pensacola, Fla.

Serranus phœbe Poey, Memorias, I, 1852, 55, pl. 2, fig. 3, Cuba.

2456. *Prionodes baldwini* Evermann and Marsh.

Porto Rico.

Prionodes baldwini Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (1900), 353, Culebra and Vieques Islands.

2457. *Prionodes fasciatus* Jenyns.

Pacific coast of Mexico, Cape San Lucas to Galapagos Islands.

Prionodes fasciatus Jenyns, Voy. *Beagle*, Fishes, 1840, 46, Chatham Island, Galapagos Archipelago.

Serranus psittacinus Valenciennes, Voy. *Vénus*, Poiss., 1855, 299, pl. 1, fig. 1, Galapagos Islands.

Serranus calopteryx Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 350, Mazatlan.

2458. *Prionodes bulleri* (Boulenger).

Coast of Jalisco, western Mexico.

Serranus bulleri Boulenger, Cat., I, 1895, 288, Las Peñas, Jalisco, Mexico.

2459. *Prionodes deweegeri* (Metzelaar).

Caribbean Sea.

Serranus (Paralabrax) deweegeri Metzelaar, Trop. Atl. Visschen, 1919, 52, fig. 20, Guanta, coast of Venezuela.

2460. *Prionodes tabacaria* (Cuvier and Valenciennes). *Jacome; Bout de tabac*.

West Indies.

Centropristes tabacarius Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 33 (44), Martinique.

Serranus jacome Poey, Memorias, I, 1852, 57, pl. 2, fig. 1, Cuba.

2461. *Prionodes tigrinus* (Bloch).

West Indies.

Holocentrus tigrinus Bloch, Naturgesch. Ausl. Fische, IV, 1790, 77, East Indies; after Seba, Thesaurus, III, Pl. XXVII, fig. 5.

Serranus præstigiator Poey, Memorias, I, 1852, 58, pl. 2, fig. 2, Habana.

?*Centropristes annularis* Günther, Shore Fishes, Challenger Rept., Zool., I, pt. 6, 1880, 6, pl. 1, fig. C, Pernambuco.

2462. *Prionodes flavescens* (Cuvier and Valenciennes).

Martinique.

Serranus flavescens Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830, 380 (506), Martinique.

Genus 844. MENTIPERCA Gill.

Mentiperca Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 236 (*Serranus lucioperca* Poey).

2463. *Mentiperca lucioperca* (Poey).

Coast of Cuba.

Serranus luciopercanus Poey, Memorias, I, 1852, 56, pl. 9, fig. 1, Cuba.

Genus 845. CENTRISTHMUS Garman.

Centrithmus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 47 (*Centrithmus signifer* Garman).

2464. *Centristhmus signifer* Garman.

Off Bay of Panama in deep water.

Centristhmus signifer Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
48, Pl. LXIX, off Bay of Panama.

Genus 846. *DULES* Cuvier.

Dules Cuvier, Règne Animal, ed. 2, II, 1829, 147 (*Dules auriga* Cuvier
and Valenciennes).

Eudulus Fowler, Proc. Acad. Nat. Sci. Phila., LIX, 1907, 264 (*Dules
auriga* Cuvier and Valenciennes; *Dulus* being a genus of birds).

2465. *Dules auriga* Cuvier and Valenciennes. *Coachman; Charioteer.*

Coasts of Brazil and Uruguay, and occasionally northward.

Dules auriga Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 83
(112), pl. 51, Brazil.

Centropristes brasiliensis Brissot de Barneville, Revue Zoologique, 1847,
131, Bahia.

Genus 847. *CALLIDULUS* Fowler.

Callidulus Fowler, Proc. Acad. Nat. Sci. Phila., LIX, 1907, 265 (*Cen-
tropristis subligarius* Cope).

**2466. *Callidulus flaviventris* Cuvier and Valenciennes. *Grassy ground rock-
fish.***

Tropical America, Charleston to Brazil.

Dules flaviventris Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829,
85, Brazil.

Centropristes dispilurus Günther, Proc. Zool. Soc. Lond., 1867, 99,
Trinidad.

Centropristes subligarius Cope, Proc. Acad. Nat. Sci. Phila., XXII,
1870, 120, Pensacola, Fla.

Genus 848. *PARANTHIAS* Guichenot.

Brachyrhinus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 286 (*Serranus
creolus* Cuvier and Valenciennes).

Paranthias Guichenot, Ann. Soc. Linn. Maine-et-Loire, X, 1868 (*Ser-
ranus furcifer* Cuvier and Valenciennes= *Serranus creolus* Cuvier and
Valenciennes).

Creolus Jordan and Gilbert, Synopsis, Bull. U. S. Nat. Mus., XVI, 1882,
XXXVI (withdrawn in the text) (*Serranus furcifer* Cuvier and
Valenciennes).

**2467. *Paranthias furcifer* (Cuvier and Valenciennes). *Rabirubia de lo alto;
Creolefish.***

Both coasts of tropical America, Cuba to Brazil, Cape San Lucas to
the Galapagos Archipelago.

Serranus furcifer Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828,
196 (264), Brazil.

Serranus creolus Cuvier and Valenciennes, loc. cit., 197 (265), Marti-
niique; San Domingo.

Corvina oxyptera DeKay, New York Fauna: Fishes, 1842, 77, Pl. XXX,
fig. 96, locality unknown.

Serranus colonus Valenciennes, Voy. Vénus, Zool., 1855, 300, pl. 2, fig. 1,
Galapagos Islands.

Genus 849. *OCYANTHIAS* Jordan and Evermann.

Ocyanthias Jordan and Evermann, Fishes North and Mid. Amer.,
1896, 1227 (*Aylopon martinicensis* Guichenot).

2468. *Ocyanthias martinicensis* (Guichenot).

West Indies.

?*Aylopon martinicensis* Guichenot, Anthiani, Ann. Linn. Soc., X, 1868,
Martinique.

Genus 850. *HYPOPLECTRUS* ⁵⁸ Gill. *Vacas.*

Hypoplectrus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 236 (*Plec-
tropoma puella* Cuvier and Valenciennes=variety of *Perca unicolor*
Walbaum).

⁵⁸ The West Indian forms correctly referred to this species fall into several groups here provisionally designated as subspecies of the form first named (*H. unicolor*), apparently differing only in color. These range from mottled olive and orange to intense indigo blue.

- 2469. *Hypoplectrus unicolor* (Walbaum). *Vaca; Petitnègre.***
 West Indies, north to Florida Keys.
Perca unicolor Walbaum, Artedi Piscium, III, 1792, 352, locality unknown.
Plectropoma ephippium Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 307, (408), locality unknown.
Hypoplectrus maculiferus Poey, Ann. Lyc. Nat. Hist. N. Y., X, 1871, 78, Habana.
- 2470. *Hypoplectrus unicolor puella* (Cuvier and Valenciennes). *Vaca.***
 Martinique.
Plectropoma puella Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 305 (405), pl. 37, Martinique.
- 2471. *Hypoplectrus unicolor vitulinus* (Poey).**
 Cuba.
Plectropoma vitulinum Poey, Memorias, I, 1852, 68, Cuba.
- 2472. *Hypoplectrus unicolor pinnivarius* (Poey).**
 Habana.
Hypoplectrus pinnivaria Poey, Synopsis, 1868, 291, Habana.
- 2473. *Hypoplectrus unicolor guttavarius* (Poey).**
 Cuba.
Plectropoma guttavarium Poey, Memorias, I, 1852, 70, Cuba.
Plectropoma melanorhina Guichenot, Poissons, in Ramon de la Sagra, Hist. Cuba, 1853, 18, pl. 1, fig. 1, Cuba.
- 2474. *Hypoplectrus unicolor gummigutta* (Poey).**
 Cuba.
Plectropoma gummigutta Poey, Memorias, I, 1852, 70, Cuba.
- 2475. *Hypoplectrus unicolor crocatus* (Cope).**
 West Indies.
Plectropoma crocota Cope, Trans. Amer. Phil. Soc., XIV, 1870 (1871), 466, St. Martins, West Indies.
- 2476. *Hypoplectrus unicolor aberrans* (Poey).**
 Cuba.
Hypoplectrus aberrans Poey, Synopsis, 1868, 291, Cuba.
- 2477. *Hypoplectrus unicolor accensus* (Poey).**
 Cuba.
Plectropoma accensum Poey, Memorias, I, 1852, 72, Cuba.
- 2478. *Hypoplectrus unicolor affinis* (Poey).**
 Cuba.
Plectropoma affine Poey, Memorias, II, 1861, 427, Cuba.
- 2479. *Hypoplectrus unicolor chlorurus* (Cuvier and Valenciennes).**
 Martinique.
Plectropoma chlorurum Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 306 (406), Martinique.
- 2480. *Hypoplectrus unicolor nigricans* (Poey).**
 Cuba.
Plectropoma nigricans Poey, Memorias, I, 1852, 71, Cuba.
- 2481. *Hypoplectrus unicolor indigo* (Poey). *Añil.***
 Cuba.
Plectropoma indigo Poey, Memorias, I, 1852, 69, pl. 3, fig. 1, Cuba.
- 2482. *Hypoplectrus unicolor bovinus* (Poey).**
 Cuba.
Plectropoma bovinum Poey, Memorias, I, 1852, 69, Cuba.
- 2483. *Hypoplectrus unicolor gemma* Goode and Bean.**
 Florida Keys; known from one specimen from Garden Key, Fla.
Hypoplectrus gemma Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 428, Garden Key, Fla.
- 2484. *Hypoplectrus lamprurus* (Jordan and Gilbert).**
 Panama.
Serranus lamprurus Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 322, Panama.

Genus 851. ANTHIAS Bloch. Barbiers.

Anthias Bloch, Nat. Ausl. Fische, VI, 1792, 97 (*Labrus anthias* L.).
Aylopon Rafinesque, Caratteri, 1810, 52, (*Labrus anthias* L.; the name
Anthias said to be preoccupied).

2485. Anthias tenuis Nichols.

Bermuda.

Anthias tenuis Nichols, Proc. Biol. Soc. Wash., XXXIII, 1920, 60,
 Bermudas.

Genus 852. HEMIANTHIAS Steindachner.

Hemianthias Steindachner, Ichth. Beitr., I, 1874, 4 (*Anthias (Hemian-*
thias) peruanus Steindachner).

2486. Hemianthias peruanus (Steindachner).

Coasts of Peru and Chile, occasionally northward; off the coast of
 Lower California.

Anthias (Hemianthias) peruanus Steindachner, Ichth. Beitr., I, 1874, 4,
 Payta; Trujillo.

2487. Hemianthias vivanus (Jordan and Swain).

Gulf of Mexico; Snapper Banks between Pensacola and Tampa, Fla.
Anthias vivanus Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884,
 544, from stomach of a red snapper taken at the Snapper Banks off
 Pensacola, Fla.

Genus 853. PRONOTOGRAMMUS Gill.

Pronotogrammus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 80 (*P.*
multifasciatus Gill).

2488. Pronotogrammus multifasciatus Gill.

Off Bay of Panama; Cape San Lucas, Lower California.

Pronotogrammus multifasciatus Gill, Proc. Acad. Nat. Sci. Phila., XV,
 1863, 81, Cape San Lucas, Lower California.

2489. Pronotogrammus eos Gilbert.

Pacific coast of tropical America.

Pronotogrammus eos Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 62,
 west coast of Mexico, at Albatross station 2996, in 112 fathoms.

Family 196. PSEUDOCHROMIDÆ**Genus 854. GRAMMA Poey.**

Gramma Poey, Synopsis, 1868, 296 (*G. loreto* Poey).

2490. Gramma loreto Poey.

Matanzas, Cuba; only the type known.

Gramma loreto Poey, Synopsis, 1868, 296, Matanzas, Cuba.

Family 197. RHEGMATIDÆ**Genus 855. RHEGMA Gilbert.**

Rhegma Gilbert, in Jordan and Evermann, Fishes North and Mid.
 Amer., 1898, 3169 (*Rhegma thaumasium* Gilbert).

2491. Rhegma thaumasium Gilbert.

Panama.

Rhegma thaumasium Gilbert, in Jordan and Evermann, Fishes North
 and Mid. Amer., 1898, 3170, Panama.

Family 198. PRIACANTHIDÆ. Big-eyes or catalufas**Genus 856. PRIACANTHUS Cuvier.**

Priacanthus Cuvier, Règne Animal, ed. 1, 1817, 281 (*P. macrophthalmus*
 Cuvier).

2492. Priacanthus cruentatus (Lacépède). Big-eye; Catalufa.

Tortugas; West Indies to St. Helena and the Canaries (*Priacanthus*
carolinus Lesson), a species very similar or identical in Hawaii and
 the western Pacific.

Labrus cruentatus Lacépède, Hist. Nat. Poiss., III, 1802, 522, Martinique;
 from a copy by Aubriet of a plate made by Plumier at Martinique.

Priacanthus cepedianus Desmarest, Prém. Dec. Ichthy., 1823, 9, pl. 1, Habana.
?Priacanthus carolinus Lesson, Voy. Coquille, 1826, 204, Caroline Islands.

2493. *Priacanthus arenatus* Cuvier and Valenciennes. *Catalufa; Big-eye.*
 West Indies, south to Brazil; occasionally northward in the Gulf Stream to Newport and Woods Hole, Mass.
Catalufa Parra, Dif. Piezas Hist. Nat., 1787, 20, Habana.
Priacanthus arenatus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 75 (101), Brazil.
Priacanthus fulgens Lowe, Trans. Zool. Soc. Lond., II, 1839, 174, Madeira.
Priacanthus catalufa Poey, Proc. Acad. Nat. Sci. Phila., XV, 1863, 182, Habana.

Genus 857. **PSEUDOPRIACANTHUS** Bleeker.
Pseudopriacanthus Bleeker, Versl. Ak. Wet. Amsterdam, 1869 (2), 241
 (*Priacanthus niphonius* Cuvier and Valenciennes).

2494. *Pseudopriacanthus altus* (Gill).
 West Indies and the Gulf Stream.
Priacanthus altus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 732, Narragansett Bay.

2495. *Pseudopriacanthus serrula* (Gilbert).
 Pacific coast of Colombia.
Priacanthus serrula Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 450, Albatross station, 2797, west coast of Colombia.

Family 199. **PEMPHERIDÆ. Deep-water catalufas**

Genus 858. **PEMPHERIS** Cuvier.
Pempheris Cuvier, Règne Anim., ed. 2, II, 1829, 195 (*Kurtus argenteus* Bloch and Schneider = *Pempheris touea* Cuvier).

2496. *Pempheris schomburgkii* Müller and Troschel.
 Barbados and Cuba.
Pempheris schomburgkii Müller and Troschel, in Schomburgk's History of Barbados, 1848, 669, Barbados.

2497. *Pempheris mülleri* Poey. *Catalufa de lo alto.*
 West Indies to Brazil.
Pempheris mülleri Poccy, Memorias, II, 1861, 203, Cuba.

2498. *Pempheris poeyi* Bean.
 Cuba.
Pempheris poeyi Bean, Proc. U. S. Nat. Mus., VIII, 1885, 229, Habana.

2499. *Pempheris mexicanus* Cuvier and Valenciennes.
 Pacific coast of Mexico, Acapulco.
Pempheris mexicana Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 230 (308), Acapulco, Mexico.

Family 200. **LOBOTIDÆ. Flashers or tripletails**

Genus 859. **LOBOTES** Cuvier.
Lobotes Cuvier, Règne Anim., ed. 2, II, 1829, 177 (*L. erate* Cuvier and Valenciennes).

2500. *Lobotes surinamensis* (Bloch). *Flasher; Tripletail; Dormeur; Black perch; Black grunt.*
 Atlantic coast from Cape Cod to Surinam; Mediterranean; Japan.
Holocentrus surinamensis Bloch, Naturgesch. Ausl. Fische, IV, 1790, 98, Surinam.

Bodianus triourus Mitchell, Trans. Lit. and Phil. Soc. N. Y., I, 1815, 418, Powells Hook, N. J.

Lobotes somnolentus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 244 (324), pl. 126, San Domingo.

Lobotes erate Cuvier and Valenciennes, loc. cit., 322, Pondicherry.

Lobotes auctorum Günther, Cat., I, 1859, 338, Cuba; Calcutta; China.

Genus 860. **VERRUGATO** Jordan.

Verrugato Jordan, Classn. Fishes, 1923, 195 (*Lobotes pacificus* Gilbert).

2501. *Verrugato pacificus* (Gilbert). *Berrugate.*

Panama.

Lobotes pacificus Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2857, Panama.

Family 201. XENICHTHYIDÆ

Genus 861. XENICHTHYS Gill.

Xenichthys Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 82 (*X. xanti* Gill.).2502. *Xenichthys xanti* Gill.

Pacific coast of tropical America, from Cape San Lucas to Panama.

Xenichthys xanti Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 82, Cape San Lucas.*Xenichthys xenops* Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 325, Panama.

Genus 862. XENISTIUS Jordan and Gilbert.

Xenistius Jordan and Gilbert, Synopsis, 1883, 920 (*Xenichthys californiensis* Steindachner).2503. *Xenistius californiensis* (Steindachner).

Pacific coast of America, from San Diego southward to Guaymas, Mexico.

Xenichthys californiensis Steindachner, Ichth. Beitr., III, 1875, 3, San Diego.

Family 202. HOPLOPAGRIDÆ

Genus 863. HOPLOPAGRUS Gill.

Hoplopagrus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 78; and 1862, 253 (*Hoplopagrus guntheri* Gill.).2504. *Hoplopagrus guntheri* Gill. *Pargo.*

Pacific coast of tropical America, from Guaymas to Panama.

Hoplopagrus guntheri Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 78, and 1862, 253, Cape San Lucas, Lower California.

✓ Family 203. LUTIANIDÆ. Snappers

Genus 864. LUTIANUS Bloch.

Lutianus Bloch, Nat. Ausl. Fische, IV, 1790, 105 (*L. lutianus* Bloch).*Diacope* Cuvier, Mém. Mus. Hist. Nat. Paris, I, 1816, 360 (*Holocentrus bengalensis* Bloch, preoccupied).*Mesopriion* Cuvier (Règne Animal, 143), Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 441 (*L. lutianus* Bloch).*Genyoroge* Cantor, Jour. Roy. Asiat. Soc. Bengal, XVIII, 1849, 12 (*Diacope sebæ* Cuvier and Valenciennes).*Neomænis* Girard, Proc. Acad. Nat. Sci. Phila., X, 1858 (1859), 167 (*Lobotes emarginatus* Baird and Girard = *Labrus griseus* Linnaeus).*Hypoplites* Gill, loc. cit., XIV, 1862, 236 (*Mesopriion retrospinis* Cuvier and Valenciennes).*Evoplites* Gill, loc. cit., 236 (*Mesopriion pomacanthus* Bleeker, young of *L. kasmira*).*Lutjanus* Bloch and later writers, change of spelling; the word from the Indian "Lutjang."

Subgenus EVOPLITES Gill.

2505. *Lutianus viridis* (Valenciennes).

Rocky islands of the eastern Pacific; known from Galapagos, Tres Marias, and Revillagigedo Islands.

Diacope viridis Valenciennes, Voy. Vénus, 1855, 303, pl. 1, fig. 2 (very bad), Galapagos Islands.

Subgenus NEOMÆNIS Girard.

2506. *Lutianus jordani* (Gilbert)

Panama; Cocos Island.

Neomænis jordani Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1252, Panama.

- 2507.** *Lutjanus novemfasciatus* (Gill). *Pargo prieto; Pargo moreno; Pargo negro.*
 Pacific coast of tropical America; Cape San Lucas; Guaymas; Mazatlan; Punta Arenas; San Blas; Panama.
Lutjanus novemfasciatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 251, Cape San Lucas, Lower California.
Mesoprion pacificus Bocourt, Ann. Sci. Nat. Paris, X, 1868, 223, Taguesco, Pacific coast of Guatemala.
Lutjanus prieto Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 353, Mazatlan.
- 2508.** *Lutjanus cyanopterus* (Cuvier and Valenciennes). *Cubera.*
 West Indies to Brazil.
Mesoprion cyanopterus Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 357 (472), Brazil.
?Mesoprion pargus Cuvier and Valenciennes, loc. cit., 358 (473), Porto Rico.
Lutjanus cubera Poey, Ann. Lyc. Nat. Hist. N. Y., X, 1871, 75, Cuba.
Lutjanus dentatus A. Duméril in Vaillant and Bocourt, Misc. Sci. Mex. Poiss., 1881, 125, Brazil.
- 2509.** *Lutjanus griseus* (Linnæus). *Gray snapper; Mangrove snapper; Cabellerote; Lawyer.*
 West Indies; ranging from New York to Brazil.
Turdus pinnis branchialibus carens Catesby, Hist. Carolinas, II, 1743, 5, pl. 9, Bahamas.
Labrus griseus Linnæus, Syst. Nat., ed. X, 1758, 283, after Catesby.
Caballerote Parra, Descr. Dif. Piezas, Hist. Nat., 1787, 52, pl. 25, fig. 1, Habana.
Sparus tetracanthus Bloch, Nat. Ausl. Fische, 1791, 116, Martinique.
Anthias caballerote Bloch and Schneider, Syst. Ichth., 1801, 310, Cuba; after Parra.
Bodianus vivanet Lacépède, Hist. Nat. Poiss., IV, 1802, pl. 4, fig. 3, Martinique.
Mesoprion griseus Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 355 (469), San Domingo, not after Linnæus.
Lobotes emarginatus Baird and Girard, 9th Smith. Rept., 1854 (1855), 332, Beaseley Point, N. J.
Lutjanus stearnsi Goode and Bean, Proc. U. S. Nat. Mus., I, 1878, 179, Pensacola, Fla.
- 2510.** *Lutjanus jocú* (Bloch and Schneider). *Dog snapper; Jocú.*
 West Indies, north to Florida Keys, south to Bahia; one young recorded from Woods Hole, Mass. (Nichols and Breder).
Anthias jocú Bloch and Schneider, Syst. Ichth., 1801, 310, Cuba; after Parra.
Jocú Parra, Descr. Dif. Piezas, Hist. Nat., I, 1787, 53, pl. 25, fig. 2, Cuba.
Mesoprion litura Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 353 (467), Cayenne; St. Thomas.
- 2511.** *Lutjanus apodus* (Walbaum). *Schoolmaster; Cajá.*
 West Indies, north to Key West, south to Bahia; young casual at Woods Hole, Mass. (Nichols and Breder).
Perca apoda Walbaum, Artedi Piscium, 1792, 351; based on the "Schoolmaster" of Catesby, which, in the very bad plate, lacks the pectoral fin.
Caxis Parra, Descr. Dif. Piezas, Hist. Nat., 1787, 14, pl. 8, fig. 2, Habana.
Sparus caxis Bloch and Schneider, Syst. Ichth., 1801, 284, Habana; after Parra.
Bodianus striatus Bloch and Schneider, loc. cit., 1801, 335, pl. 65, West Indies, misprinted *albostriatus* and *fasciatus*.
Lutjanus acutirostris Desmarest, Prem. Dec. Ichth., 1823, 12, pl. 3, Cuba.
Mesoprion cynodon Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 351 (465), Martinique, San Domingo.
Mesoprion linea Cuvier and Valenciennes, loc. cit., 354 (468), Cuba; San Domingo.
Mesoprion flavescens Cuvier and Valenciennes, loc. cit., 357 (472), Martinique.

2512. *Lutjanus lutjanoides* (Poey).

Cuba.

Ocyurus lutjanoides Poey, Ann. Lyc. Nat. Hist., IX, 1870, 319, Cuba; known from one specimen, suspected to be a hybrid between *Ocyurus chrysurus* and *Lutjanus apodus*.**2513. *Lutjanus brachypterus* (Cope).**

Bahama Islands.

One specimen known.

Lutjanus brachypterus Cope, Trans. Amer. Phil. Soc., XIV, 1871, 470, New Providence, Bahama Islands; perhaps a hybrid of *Lutjanus griseus* and *L. synagris*.**2514. *Lutjanus argentiventris* (Peters). *Pargo amarillo*.**

Pacific coast of tropical America; Galapagos and Cocos Islands.

Mesoprion argentiventris Peters, Berlin Monatsber., 1869, 704, Mazatlan.**2515. *Lutjanus hastingsi* (Bean). *Bermuda silk snapper*.**

Bermuda.

Neomaznis hastingsi Bean, Bull. Amer. Mus. Nat. Hist., X, Art. III, 1898, 45, Bermuda.**2516. *Lutjanus buccanella* (Cuvier and Valenciennes). *Sesi de lo alto; Orielle noire; Boucanello; Blackfin snapper*.**

West Indies; Habana.

Mesoprion buccanella Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 344 (455), Martinique.*Mesoprion caudanotatus* Poey, Memorias, I, 1854, pl. 3, fig. 3, 440, Cuba.**2517. *Lutjanus vivanus* (Cuvier and Valenciennes). *Pargo de lo alto; Silk snapper; West Indian snapper*.**

West Indies; Habana.

Mesoprion vivanus Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 343 (454), Martinique.*Mesoprion profundus* Poey, Memorias, II, 1861, 150, Cuba.*Lutjanus torridus* Cope, Trans. Amer. Phil. Soc., XIV, 1871, 469, St. Kitts.*Lutjanus purpureus* Poey, Enumeratio, 1875, 29.**2518. *Lutjanus campechanus*⁶⁰ (Poey). *Red snapper; Mexican snapper; Pargo guachenarigo*.**

Long Island to Yucatan and Texas; a valued food fish.

Mesoprion campechanus Poey, Memorias, II, 1861, 149, Campeche.**2519. *Lutjanus blackfordii* Goode and Bean. *Pensacola snapper*.***Lutjanus blackfordii* Goode and Bean, Proc. U. S. Nat. Mus., I, 1878, 176, Pensacola, Fla.**2520. *Lutjanus analis* (Cuvier and Valenciennes). *Muttonfish; Pargo; Pargo criollo*.**

West Indies; Pensacola to Brazil, rarely north to Woods Hole, common and valued.

Anthias quartus rondeleti (the muttonfish) Catesby, Nat. Hist. Carolina, II, 1743, 25, pl. 25, Bahamas.*Mesoprion analis* Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 341 (452), San Domingo.*Mesoprion sobra* Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 342 (453), Martinique.*Mesoprion isodon* Cuvier and Valenciennes, loc. cit., IX, 1833, 328 (443), San Domingo.*Mesoprion rosaceus* Poey, Ann. Lyc. Nat. Hist. N. Y., IX, 1870, 317, Cuba.**2521. *Lutjanus colorado* (Jordan and Gilbert). *Pargo colorado*.**

Pacific coast of Mexico, Guaymas to Panama.

Lutjanus colorado Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 338, name only, 351, 355, Mazatlan, Mexico.**2522. *Lutjanus guttatus* (Steindachner). *Flamenco*.**

Pacific coast of Mexico, from Guaymas to Panama.

Mesoprion guttatus Steindachner, Ichthyol. Notizen, IX, 1869, 18, pl. 8, Mazatlan, Mexico.

⁶⁰ The common red snapper is here admitted provisionally as distinct from the Brazilian *Lutjanus aya* from which it is said by Hildebrand to differ tangibly.

- 2523.** *Lutianus synagris* (Linnæus). *Lane snapper; Biajaiba; Redtail snapper.*
West Indies; Florida Keys to Colon and Brazil.
Sparus synagris Linnæus, Syst. Nat., Ed. X, 1758, 280, Carolinas; after
Catesby.
Salpa purpurascens variegata (Lane snapper) Catesby, Nat. Hist. Caro-
lina, II, 1743, 17, pl. 17, Bahamas.
Sparus vermicularis Bloch and Schneider, Syst. Ichth., 1801, 275,
Martinique, on a drawing by Plumier.
? *Lutjanus auberti* Desmarest, Prem. Dec. Ichth., 1823, 17, pl. 2, Cuba.
Mesopriion uninotatus Cuvier and Valenciennes, Hist. Nat. Poiss., II,
1828, 339 (449), pl. 39, San Domingo; Martinique.
- 2524.** *Lutianus ambiguus* (Poey).
Cuba.
Mesopriion ambiguus Poey, Memorias, II, 1861, 152, pl. 12, fig. 4; pl. 13,
fig. 8; Habana; two specimens known; suspected to be a hybrid of
Lutianus synagris and *Ocyurus chrysurus*.
- 2525.** *Lutianus mahogoni* (Cuvier and Valenciennes). *Ojanco; Mahogany
snapper.*
West Indies; Habana.
Mesopriion mahogoni Cuvier and Valenciennes, Hist. Nat. Poiss., II,
1828, 338 (447), Martinique.
Mesopriion ricardi Cuvier and Valenciennes, loc. cit., 339 (447), Marti-
nique? No locality given.
Mesopriion ojanco Poey, Memorias, II, 1861, 150, pl. 13, fig. 10, Cuba.
- 2526.** *Lutianus megalophthalmus* (Evermann and Marsh).
Porto Rico.
Neomænis megalophthalmus Evermann and Marsh, Bull. U. S. Fish
Com., XX, 1900 (1902), 177, Puerto Real, Porto Rico.
- Genus 865. RAIZERO** Jordan and Fesler.
Raisero Jordan and Fesler, Rept. U. S. Fish. Com., XVII, 1889 (1893),
438 (*Mesopriion aratus* Günther).
- 2527.** *Raizero aratus* (Günther). *Pargo de raizero.*
Pacific coast of tropical America.
Mesopriion aratus Günther, Proc. Zool. Soc. Lond., 1864, 145, Panama;
Chiapam.
- Genus 866. RABIRUBIA** Jordan and Fesler.
Rabirubia Jordan and Fesler, Rept. U. S. Fish Com., XVII, 1889
(1893), 438 (*Mesopriion inermis* Peters).
- 2528.** *Rabirubia inermis* (Peters).
Pacific coast of America from Mazatlan to Panama.
Mesopriion inermis Peters, Berliner Monatsber., 1869, 705, Mazatlan,
Mexico.
- Genus 867. OCYURUS** Gill. *Rabirubias.*
Ocyurus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 236 (*Sparus
chrysurus* Bloch).
- 2529.** *Ocyurus chrysurus* (Bloch). *Yellowtail; Rabirubia.*
West Indies, southern Florida to Brazil; single specimen recorded at
Woods Hole, Mass.
Sparus chrysurus Bloch, Nat. Ausl. Fische, V, 1791, 28, Brazil; after
Marcgrave.
Acara pitamba Marcgrave, Hist. Brasil., 1648, 155, Brazil.
Rabirubia Parra, Descr. Dif. Piezas, Hist. Nat., 1787, 42, pl. 20, fig. 1,
Cuba.
Anthias rabirubia Bloch and Schneider, Syst. Ichth., 1801, 309, Cuba;
after Parra.
Sparus semiluna Lacépède, Hist. Nat. Poiss., IV, 1802, 141, Martinique.
Mesopriion auroritatus Agassiz, Spix, Pisc. Brasil., 1829, 121, pl. 66,
Brazil.
Ocyurus riddersmaei Cope, Trans. Amer. Phil. Soc., XIV, 1871, 468,
fig. 4, St. Kitts.
- Genus 868. RHOMBOPLITES** Gill.
Rhomboplites Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 237 (*Cen-
trophistis aurorubens* Cuvier and Valenciennes).

2530. **Rhomboplites aurorubens** (Cuvier and Valenciennes). *Cagon de lo alto.*

West Indies, north to Charleston, south to Rio Janeiro.

Centropristes aurorubens Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 34 (45), Brazil, Martinique, and San Domingo.

Mesopriion elegans Poey, Memorias, II, 1861, 153, Cuba.

Apriom ariommus Jordan and Gilbert, Proc. U. S. Nat. Mus., VI, 1883, 142, Pensacola, Fla.

Genus 869. TROPIDINIUS Gill.

Tropidinius Gill, in Poey, Synopsis, 1868, 296 (*Mesopriion arnillo* Poey = *Apsilus dentatus* Guichenot).

2531. **Tropidinius dentatus** (Guichenot). *Arnillo.*

West Indies; Habana.

Apsilus dentatus Guichenot, in Ramon de la Sagra, Hist. Cuba, Poiss., 1853, 29, pl. 1, fig. 2, Habana.

Mesopriion arnillo Poey, Memorias, II, 1861, 154, Cuba.

Genus 870. PRISTIPOMOIDES Bleeker.

Pristipomoides Bleeker, Tijdschr. Ned. Ind., III, 1852, 574 (*Pristipomoides typus* Bleeker).

Platyinius Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 237 (*Mesopriion vorax* Poey = *Centropristes macrophthalmus* Müller and Troschel).

Bowersia Jordan and Evermann, Bull. U. S. Fish Com., XXII, 1902 (1903), 182 (*B. violescens* Jordan and Evermann).

Subgenus PLATYINIUS Gill.

2532. **Pristipomoides macrophthalmus** (Müller and Troschel). *Voraz.*

West Indies, Habana.

Centropristes macrophthalmus Müller and Troschel, in Schomburgk, Hist. Barbados, 1848, 666, Barbados.

Mesopriion vorax Poey, Memorias, II, 1861, 151, Cuba.

Genus 871. ETELIS Cuvier and Valenciennes.

Etelis Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 93 (127) (*Etelis carbunculus* Cuvier and Valenciennes, an East Indian species).

Elastoma Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 202 (*Etelis oculatus* Cuvier and Valenciennes).

Hesperanthias Lowe, Fishes of Madeira, 1843, 14 (*Serranus oculatus* Cuvier and Valenciennes).

Macrops Duméril, Mém. Acad. Sci. Paris, XXVII, 1856, 279 (*Serranus aculeatus (oculatus)* Cuvier and Valenciennes).

2533. **Etelis oculatus** [“] (Cuvier and Valenciennes). *Cachucho.*

Warm seas; West Indies to Madeira; Hawaii.

Etelis carbunculus Cuvier and Valenciennes, Hist. Nat. Poiss., II, 1828, 94 (127), pl. 18, Seychelles Archipelago.

Serranus oculatus Cuvier and Valenciennes, loc. cit., II, 198 (266), pl. 32, Martinique.

?*Etelis evirurus* Jordan and Evermann, Bull. U. S. Fish Com., XXII, 1902, 142, Hawaii.

Genus 872. ETELIDES Jordan and Thompson.

Etelides Jordan and Thompson, Bull. U. S. Bur. Fish., XXIV, 1904, 241 (*Anthias aquilonaris* Goode and Bean).

2534. **Etelides aquilonaris** (Goode and Bean).

Gulf of Mexico; Gulf Stream near Carolina coast.

Anthias aquilonaris Goode and Bean, Oceanic Ichth., 1895, 238, Gulf of Mexico.

Family 204. VERILIDÆ

Genus 873. VERILUS Poey.

Verilus Poey, Memorias, II, 1861, 124 (*V. sordidus* Poey).

2535. **Verilus sordidus** Poey. *Escolar chino.*

West Indies.

Verilus sordidus Poey, Memorias, II, 1861, 125, pl. 12, fig. 6, Cuba.

[“] This widely distributed and brilliantly colored fish seems to be of a single species throughout the warm seas. We let the American species stand, pending investigation, though we can not detect any difference from the common *evirurus*, Hawaii, and *carbunculus* of Japan.

Family 205. HÆMULIDÆ

Genus 874. HÆMULON Cuvier. *Roncos or grunts.*

Diabasis Desmarest, Prem. Décade Ichth., 1823, 34 (*D. parra* Desmarest; name preoccupied in beetles).

Hæmulon Cuvier, Règne Anim., ed. 2, II, 1829, 175 (*elegans*, etc., restricted later to *Hæmulon elegans* Cuvier = *Sparus sciurus* Shaw).

Anarmostus Scudder, in Putnam, Bull. Mus. Comp. Zool., I, 1863, 12; name preoccupied in insects.

2536. Hæmulon sexfasciatum Gill. *Mojarra almejero.*

Pacific coast of tropical America, Guaymas to Panama.

Hæmulon sexfasciatum Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 254, Cape San Lucas.

Hæmulon maculosum Peters, Berliner Monatsber., 1869, 705, Mazatlan.

2537. Hæmulon album Cuvier and Valenciennes. *Margatefish; Jallao; Margaret grunt.*

West Indies; Florida Keys to Brazil.

Perca marina gibbosa cinerea (margatefish) Catesby, Nat. Hist. Carolina, II, 1743, 2, pl. 2, Bahamas.

Perca gibbosa Walbaum, Artedi Pisc., 1792, 342, not of Linnaeus, after Catesby.

Hæmulon album Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 179 (241), St. Thomas.

Hæmulon microphthalmum Günther, Cat., I, 1859, 306, America.

2538. Hæmulon mowbrayi Jordan and Evermann.

Key West, Fla.

Hæmulon chrysopterum Mowbray, N. Y. Zool. Soc. Bull., XVIII, No. 6, 1915, 1298, with fig., Key West; not of Cuvier, which is

Bathystoma rimator Jordan and Swain.

Hæmulon mowbrayi Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, Apr. 27, 1927, 505.

2539. Hæmulon macrostomum Günther. *Gray grunt; Striped grunt.*

West Indies, north to Florida Keys.

Hæmulon macrostoma Günther, Cat., I, 1859, 308, Jamaica.

Hæmulon fremebundum Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 340, Clearwater Harbor, Fla.

2540. Hæmulon bonariense Cuvier and Valenciennes. *Black grunt; Ponce prieto.*

West Indies, south to Buenos Aires.

Hæmulon bonariense Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 174 (254), Buenos Aires.

Hæmulon canna Cuvier and Valenciennes, loc. cit., V, 173 (233), Martinique.

Hæmulon notatum Poey, Memorias, II, 1861, 179, Cuba.

Hæmulon retrocurrens Poey, Repertorio, II, 1868, 236, 461, Cuba.

Hæmulon continuum Poey, Enumeratio, 1875, 46, Cuba.

2541. Hæmulon parra (Desmarest). *Sailors' choice; Ronco blanco; Ronco prieto; Bastard margaret.*

West Indies; southern Florida to Brazil; very common at Key West and Habana.

Diabasis parra Desmarest, Prem. Décade Ichth., 1823, 30, pl. 2, fig. 2, Habana.

Hæmulon caudimacula Cuvier, Règne Anim., ed. 2, II, 1829, 176, Brazil; Habana.

Hæmulon chromis Broussonet, in Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 180 (242), Jamaica.

Hæmulon acutum Poey, Memorias, II, 1861, 180, Cuba.

Hæmulon serratum Poey, loc. cit., 181, Cuba.

Hæmulon albidum Poey, loc. cit., 181, Cuba.

Hæmulon canna Agassiz in Spix, Pisc. Brasil., 1829, 130, pl. 69, Brazil; not of Cuvier and Valenciennes.

- 2542.** *Hæmulon scudderii* Gill. *Mojarra prieta*.
 Pacific coast of tropical America, Guaymas to Panama.
Hæmulon scudderii Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 253,
 Cape San Lucas.
Hæmulon brevirostrum Günther, Trans. Zool. Soc. Lond., VI, 1868, 418
 (Nov. 27), Panama.
Hæmulon undecimale Steindachner, Ichth. Beitr., III, 1875, 11, Panama.
- 2543.** *Hæmulon carbonarium* Poey. *Ronco carbonero; Cæsar grunt*.
 West Indies and Bermudas, south to Brazil; Habana.
Hæmulon carbonarium Poey, Memorias, II, 1861, 176, Cuba.
- 2544.** *Hæmulon steindachneri* (Jordan and Gilbert). *Roncador raiado*.
 Both coasts of tropical America; Guaymas to Panama; a similar
 species (unnamed) recorded from St. Lucia and Rio Janeiro.
Diabasis steindachneri Jordan and Gilbert, Bull. U. S. Fish Com., I,
 1881 (1882), 322, Panama and Mazatlan.
- 2545.** *Hæmulon melanurum* (Linnaeus). *Jeniguana; Cæsar grunt*.
 West Indies; Habana and southward; Bush Key; Tortugas.
Perca marina cauda nigra (blacktail) Catesby, Hist. Carolina, II, 1743,
 7, pl. 7, fig. 2, Bahamas.
Perca melanura Linnaeus, Syst. Nat., ed. X, 1758, 292, and ed. XII,
 1766, 486, Bahamas; based on Catesby.
Hæmulon schrankii Agassiz in Spix, Pisc. Brasil., 1829, 131, pl. 69a,
 Brazil.
Hæmulon dorsale Poey, Memorias, II, 1861, 179, Cuba.
- 2546.** *Hæmulon sciurus* (Shaw). *Yellow grunt; Ronco amarillo; Boar
 grunt; Humpback grunt*.
 West Indies; Florida Keys to Brazil.
Anthias formosus Bloch, Nat. Ausl. Fische, VI, 1792, 122, Antilles;
 not *Perca formosa* of Linnaeus as identified.
Sparus sciurus Shaw, General Zoology, IV, 1803, 439, pl. 64, Antilles;
 based on the description and figure of Bloch.
?Diabasis obliquatus Bennett, Zool. Jour. London, V, 1837, 90, Jamaica.
?Hæmulon similis Castelnau, Anim. Nouv. Rares, II, 1855, Bahia.
Hæmulon luteum Poey, Memorias, II, 1861, 174, Cuba.
Hæmulon multilineatum Poey, loc. cit., 178, Cuba.
?Hæmulon hians Haly, Ann. Nat. Hist., XV, 1875, 268, Aspinwall.
- 2547.** *Hæmulon plumieri* (Lacépède). *Common grunt; Ronco ronco; Ronco
 arará; Boar grunt; Squirrelfish; Ronco grande*.
 West Indies; abundant from Cape Hatteras to Rio Janeiro.
Perca marina capite striato Catesby, Nat. Hist. Carolina, II, 1743, 6,
 pl. 6, Bahamas.
Labrus plumieri Lacépède, Hist. Nat. Poiss., III, 1802, 480, pl. 2, fig. 2,
 Martinique; on a copy of a drawing by Plumier, identified with this
 species by Cuvier.
Guabicoara brasiliensis Marcgrave, Hist. Brasil., 1648, 163, Brazil.
Hæmulon formosum Cuvier, Règne Anim., ed. 2, II, 1829, 175, Mar-
 tinique.
Hæmulon arcuatum Cuvier and Valenciennes, Hist. Nat. Poiss., IX,
 1833, 356 (481), Charleston, S. C.
Hæmulon arara Poey, Memorias, II, 1861, 177, Cuba.
Hæmulon subarcuatum Poey, loc. cit., 419, Cuba.
- 2548.** *Hæmulon flavolineatum* (Desmarest). *French grunt; Open-mouthed
 grunt; Ronco condenado*.
 West Indies; Florida Keys and Bermudas to Brazil.
Diabasis flavolineatus Desmarest, Prem. Décade Ichth., 1823, 35, pl. 2,
 fig. 1, Cuba.
Hæmulon heterodon Cuvier, Règne Anim., ed. 2, II, 1829, 176, Cuba,
 based on Desmarest.
Hæmulon xanthopteron Cuvier and Valenciennes, Hist. Nat. Poiss., V,
 1830, 174 (254), Martinique.
- Genus 875. BRACHYGENYS** Seudder.
Brachygenys Scudder, in Poey, Enumeratio, 1875, 47 (*Hæmulon tænia-*
tum Poey = *Hæmulon chrysargyreum* Günther).

2549. *Brachygenys chrysargyreus* (Günther).

Key West, Habana, and Trinidad.

Hæmulon chrysargyreum Günther, Cat., 1859, 314, Trinidad.

Hæmulon tæniatum Poey, Memorias, II, 1861, 182, Cuba.

Genus 876. *BATHYSTOMA* Scudder. *Tomates.*

Bathystoma Scudder, in Putnam, Bull. Mus. Comp. Zool., I, 1863, 12; no definition. (*Hæmulon jeniguano* Poey = *Hæmulon aurolineatum* Cuvier and Valenciennes); not *Bathystomus* Förster, 1862.

2550. *Bathystoma rimator* (Jordan and Swain). *Tomate; Redmouth grunt; Cesar; Young grunt.*

West Indies; Cape Hatteras to Trinidad.

Hæmulon rimator Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884, 308, Charleston; Key West; Pensacola.

2551. *Bathystoma aurolineatum* (Cuvier and Valenciennes). *Jeniguano; Tomate.*

West Indies; Florida Keys to Brazil.

Hæmulon aurolineatum Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 176 (237), Brazil; San Domingo.

Hæmulon jeniguano Poey, Memorias, II, 1861, 183, Cuba.

2552. *Bathystoma striatum* (Linnaeus). *White grunt.*

West Indies; Bermudas to Brazil.

Capeuna brasiliensis Marckgrave, Hist. Brasil., 1648, 155, Brazil (nonbinomial).

Perca striata Linnaeus, Syst. Nat., Ed. X, 1758, 293, North America. *Grammistes trivittatus* Bloch and Schneider, Syst. Ichth., 1801, 188, Brazil, after Marckgrave.

Serranus capuena Lichtenstein, Abhandl. Berlin. Akad., 1820, 288, Brazil; after Marckgrave.

Hæmulon quadrilineatum Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 177 (238), pl. 120, San Domingo.

Hæmulon quinquelineatum Poey, Memorias, II, 1861, 419, Cuba.

Genus 877. *LYTHRULON* Jordan and Swain.

Lythrulon Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884, 287 (*Hæmulon flaviguttatus* Gill).

2553. *Lythrulon flaviguttatum* (Gill). *Peixe fonda.*

Pacific coast of tropical America, Guaymas to Panama.

Hæmulon flaviguttatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 254, Cape San Lucas, Lower California.

Hæmulon margaritiferum Günther, Proc. Zool. Soc. Lond., 1864, 147, Panáma.

2554. *Lythrulon opalescens* Jordan and Starks.

Mazatlan, Mexico.

Lythrulon opalescens Jordan and Starks, in Jordan, Proc. Calif. Acad. Sci., V, 1895, 459, pl. 40, Mazatlan, Mexico.

Genus 878. *ORTHOSTÆCHUS* Gill. *Striped grunts.*

Orthostæchus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 255 (*O. maculicauda* Gill).

2555. *Orthostæchus maculicauda* Gill. *Roncador raiado.*

Pacific coast of tropical America; Guaymas to Panama.

Orthostæchus maculicauda Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 225, Cape San Lucas, Lower California.

Hæmulon mazallanum Steindachner, Ichth. Notizen, VIII, 1869, 12, Pl. VI, Mazatlan.

Genus 879. *PARACONODON* Bleeker.

Paraconodon Bleeker, Archiv Neerl., XI, 1876, 272 (*Conodon pacifici* Günther).

2556. *Paraconodon pacifici* (Günther).

Pacific coast of tropical America.

Conodon pacifici Günther, Proc. Zool. Soc. Lond., 1864, 147, Chiapam.

2557. *Paraconodon cæsius* (Jordan and Gilbert).

Pacific coast of Mexico; Mazatlan and Acapulco; Bay of Panama.

Pomadasys cæsius Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 383, Mazatlan.

- 2558. *Paraconodon dovii* (Günther).**
 Pacific coast of tropical America at Mazatlan and Panama.
Pristipoma dovii Günther, Proc. Zool. Soc. Lond., 1864, 23, Panama
- Genus 880. *ANISOTREMUS* Gill.**
Anisotremus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 105 (*Sparus virginicus* Linnaeus).
Genytremus Gill, loc. cit., XIV, 1862, 256 (*Pristipoma bilineatum* Cuvier and Valenciennes = *Anisotremus surinamensis* Bloch).
- Subgenus GENYTREMUS Gill.**
- 2559. *Anisotremus surinamensis* (Bloch). *Pompon.***
 Tropical America, from Louisiana (New Orleans market) to Brazil.
?Lutjanus surinamensis Bloch, Naturgesch. Ausl. Fische, IV, 1790, 98, Surinam.
Pristipoma bilineatum Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 202 (271), pl. 122, Martinique.
Pristipoma melanopterum Cuvier and Valenciennes, loc. cit., 203 (273), Brazil.
Hæmulon obtusum Poey, Memorias, II, 1861, 182, Habana.
Hæmulon labridum Poey, loc. cit., 1861, 419, Cuba.
- 2560. *Anisotremus interruptus* (Gill). *Mojarron.***
 Pacific coast, Magdalena Bay to Panama and the Galapagos.
Genytremus interruptus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1861, 256, Cape San Lucas, Lower California.
Pristipoma furthii Steindachner, Ichth. Beitr., V, 1876, 4, Panama.
- 2561. *Anisotremus bicolor* (Castelnau). *Maria prieta.***
 Coast of Brazil; perhaps Cuba.
Pristipoma bicolor Castelnau, Anim. Amér. du Sud, Poiss., 1855, 8, pl. 2, fig. 2, Bahia, Brazil.
?Pristipoma trilineatum Poey, Memorias, II, 1861, 343, Habana, young.
Pristipoma brasiliense Steindachner, Sitzungsb. k. Akad. Wiss. Wien, 1863, 1013, Bahia.
- Subgenus ANISOTREMUS Gill.**
- 2562. *Anisotremus scapularis* (Tschudi).**
 Coast of Peru; said to have been once taken at Mazatlan, a doubtful record.
Pristipoma scapulare Tschudi, Fauna Peruana, 12, 1845, Huacho, Peru.
?Diagramma melanospilum Kner, Sitzungsb. k. Akad. Wiss., 1807, 4, west coast of South America.
Pristipoma notatum Peters, Monatsb. Akad. Wiss. Berlin, 1869, 706, "angeblich aus Mazatlan."
- 2563. *Anisotremus davidsønii* (Steindachner). *Sargo raiado.***
 Coast of southern California.
Pristipoma davidsonii Steindachner, Ichth. Beitr., III, 1875, 6, San Diego.
- 2564. *Anisotremus spleniatus* (Poey).**
 Habana.
Pristipoma spleniatum Poey, Memorias, II, 1861, 187, Habana.
- 2565. *Anisotremus tæniatus* Gill. *Catalina; Striped grunt.***
 Pacific coast of tropical America; Magdalena Bay to Panama.
Anisotremus tæniatus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 107, Panama.
- 2566. *Anisotremus virginicus* (Linnaeus). *Porkfish; Catalineta.***
 West Indies; Florida Keys to Brazil.
Guatucupa juba Marcgrave, Hist. Brasil., 1648, 147, Brazil.
Acara pinima Marcgrave, loc. cit., 1648, 152, Brazil.
Sparus virginicus Linnaeus, Syst. Nat., ed. X, 1758, 281, South America.
?Sparus vittatus Bloch, Naturgesch. Ausl. Fische, V, 1791, 33, fig. 2, Brazil; after *Acara pinima* Marcgrave.
Perca juba Bloch, loc. cit., VI, 1792, 77, Brazil, after Marcgrave.
Grammistes mauritii Bloch and Schneider, Syst. Ichthyol., 1801, 185, after Bloch.

?*Pristipoma catharinæ* Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 201, 269, St. Catharine Island, Brazil.

Pristipoma acara-pinima Castelnau, Anim. Nouv. Rares, 1856, 8, Brazil.

Pristipoma rodo Cuvier and Valenciennes, loc. cit., 205 (274), Martinique, Brazil, Porto Rico, and San Domingo.

2567. *Anisotremus serrula* (Cuvier and Valenciennes). *Tête de roche; Petite scie.*

Martinique.

Pristipoma serrula Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 203 (272), Martinique.

Pristipoma auratum Cuvier and Valenciennes, loc. cit., 203 (272), Martinique.

Genus 881. **CONODON** Cuvier and Valenciennes.

Conodon Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 156 (*C. antillanus* Cuvier and Valenciennes = *Perca nobilis* Linnæus).

Cheloniger Plumier, in Lacépède, Hist. Nat. Poiss., V, 1803 (*Perca nobilis* Linnæus); name not accepted by Lacépède.

2568. *Conodon nobilis* (Linnæus).

West Indies; coast of Texas to Brazil.

Perca nobilis Linnæus, Syst. Nat., Ed. X, 1758, 291, North America.

Sciæna plumieri Bloch, Nat. Ausl. Fische, VI, 1792, 66, taf. 306, Martinique.

Sciæna coro Bloch, loc. cit., 1792, 70, pl. 307, fig. 2, Brazil; after *Coro coro* Marckgrave.

Conodon antillanus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 116 (156), Jamaica.

2569. *Conodon serrifer* Jordan and Gilbert.

Lower California.

Conodon serrifer Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 351, Boca Soledad, Lower California.

Genus 882. **BRACHYDEUTERUS** Gill. *Burritos.*

Brachydeuterus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 17 (*B. auritus*, an African species).

Hæmulopsis Steindachner, Ichth. Notizen, VIII, 1869, 9, Mazatlan (*Hæmulon corvinæformis* Steindachner).

2570. *Brachydeuterus nitidus* (Steindachner).

Pacific coast of tropical America; Panama, Mazatlan, and the Gulf of California.

Pristipoma (*Hæmulopsis*) *nitidum* Steindachner, Ichth. Notizen, VIII, 1869, 5, Mazatlan.

2571. *Brachydeuterus leuciscus* (Günther). *Burrito.*

Pacific coast of tropical America; not rare at Mazatlan and Panama; south to northern Peru.

Pristopoma leuciscus Günther, Proc. Zool. Soc. Lond., 1864, 147, San Jose de Nicaragua; Chiapam.

Pomadasys elongatus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 352, Mazatlan, Mexico.

2572. *Brachydeuterus axillaris* (Steindachner). *Burro blanco.*

Pacific coast of Mexico; Mazatlan and Guaymas.

Pristipoma (*Hæmulopsis*) *axillare* Steindachner, Ichth. Notizen, VIII, 1869, 7, Mazatlan.

2573. *Brachydeuterus corvinæformis* (Steindachner).

West Indies to Brazil; Jamaica.

Hæmulon corvinæforme Steindachner, Ichth. Notizen, VII, 1868, 16, Santos, Brazil.

Genus 883. **RHENCUS** Jordan and Evermann.

Rhencus Jordan and Evermann, Check List, 1895, 387 (*Pristipoma panamense* Steindachner).

2574. **Rhencus panamensis** (Steindachner).

Pacific coast of tropical America; Panama and Mazatlan.

Pristipoma panamense Steindachner, Ichth. Beitr., III, 1875, 8, Panama.

Genus 884. **POMADASYS** Lacépède. *Burros.*

Pomadasys Lacépède, Hist. Nat. Poiss., IV, 1803, 516 (*Sciæna argentea* Forskål).

Pristipomus Oken, Isis, 1817, 279 (*Lutjanus hasta* Bloch; after *les Pristipomes* Cuvier).

Pristipoma Cuvier and Valenciennes, changed spelling.

Subgenus **PRISTOPOMUS** Cuvier.2575. **Pomadasys bayanus** Jordan and Evermann.

Pacific coast of Panama, in fresh water.

Pristipoma humile Kner and Steindachner, Sitzgber. Akad. Wiss. Münch., 1863, 222, Rio Bayano, near Panama; name preoccupied by *Pristopoma humilis* Bowdich, Fishes of Madeira, 1825, 236.

Pomadasys bayanus Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1331, Rio Bayano, near Panama.

2576. **Pomadasys productus** (Poey).

Cuba.

Pristipoma productum Poey, Memorias, II, 1861, 186, Habana.

2577. **Pomadasys macracanthus** (Günther). *Burro.*

Pacific coast of tropical America; Panama; Mazatlan; Chiapas, and Punta Arenas.

Pristipoma macracanthum Günther, Proc. Zool. Soc. Lond., 1864, 146, Chiapam.

2578. **Pomadasys grandis** Meek.

Lake Nicaragua.

Pomadasys grandis Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 116, Lake Nicaragua.

2579. **Pomadasys templei** Meek.

Valles, San Luis Potosí, Mexico.

Pomadasys templei Meek, Field Col. Mus., Zool., V, 1904, 201, Valles, San Luis Potosí, Mexico.

2580. **Pomadasys boschmae** (Metzelaar).

West Indies.

Pristipoma boschmae Metzelaar, Tropisch Atl. Viessch., 1919, 83, fig. 27; probably from Curaçao.

2581. **Pomadasys starri** Meek. *Burro.*

Perez, Vera Cruz, Mexico.

Pomadasys starri Meek, Field Col. Mus., Zool., V, 1904, 200, Perez, Vera Cruz, Mexico.

Genus 885. **RHONCISCUS** Jordan and Evermann.

Rhonciscus Jordan and Evermann, Check List, 1895, 387 (*Pristipoma crocro* Cuvier and Valenciennes).

2582. **Rhonciscus crocro** (Cuvier and Valenciennes).

West Indies; Cuba to Brazil.

Pristipoma crocro Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 197 (264), Martinique.

Pristipoma cultiferum Poey, Memorias, II, 1861, 185, Habana.

Pomadasys approximans Bean and Dresel, Proc. U. S. Nat. Mus., VII, 1884, 160, Jamaica.

2583. **Rhonciscus branickii** (Steindachner). *Burrito.*

Pacific coast of tropical America; Mazatlan to Peru.

Pristipoma branickii Steindachner, Denkschr. K. Akad. Wiss. Wien, XLI, 1879, 28, Tumbez, Peru.

2584. **Rhonciscus ramosus** (Poey).

West Indies; south to Brazil.

Pristipoma ramosum Poey, Memorias, II, 1861, 186, Habana.

Pristipoma boucardi Steindachner, Ichth. Notizen, VIII, 1869, 1, Cojimar River near Habana.

- 2585. *Rhenciscus sinuosus* (Eigenmann).**
North coast of Colombia.
Pomadasys sinuosus Eigenmann, Proc. Amer. Phil. Soc., LVI, 1917, 683,
Patia, between Magna and Telembi.
- Genus 886. *ORTHOPRISTIS* Girard. *Pigfishes.***
Orthopristis Girard, U. S. Mex. Bound. Surv., Zool., 1859, 15 (*O. duplex*
Girard = *Perca chrysopterus* Linnaeus).
Pristocantharus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 256
(*Pristipoma cantharinum* Jenyns).
- 2586. *Orthopristis chalceus* (Günther).**
Pacific coast of tropical America, from Cape San Lucas to the Galapagos.
Pristipoma chalceum Günther, Proc. Zool. Soc. Lond., 1864, 146,
Panama.
Pristipoma kneri Steindachner, Ichth. Notizen, VIII, 1869, 3, Mazatlan,
Mexico.
- 2587. *Orthopristis reddingi* Jordan and Richardson.**
La Paz, Lower California.
Orthopristis reddingi Jordan and Richardson, in Jordan, Proc. Calif.
Acad. Sci., V, 1895, 509, pl. 41, La Paz, Lower California.
- 2588. *Orthopristis chrysopterus* (Linnaeus). *Pigfish; Sailors' choice, Hogfish;*
*Red-mouled grunt.***
South Atlantic and Gulf coasts of the United States; young common
as far north as New York.
Perca chrysoptera Linnaeus, Syst. Nat., Ed. XII, 1766, 485, Charleston,
S. C.
Labrus fulvo-maculatus Mitchell, Trans. Lit. and Phil. Soc. N. Y.,
1814, 406, New York.
Pristipoma fasciatum Cuvier and Valenciennes, Hist. Nat. Poiss., V,
1830, 213 (285), New York.
Orthopristis duplex Girard, U. S. Mex. Bound. Surv., Zool., Fishes, 1859,
15, pl. 9, figs. 1 to 4, Indianola and Brazos Santiago, Tex.
- 2589. *Orthopristis poeyi* Seudder.**
West Indies; Habana.
Orthopristis poeyi Seudder, in Poey, Synopsis, 1868, 312, Habana.
- Genus 887. *MICROLEPIDOTUS* Gill.**
Microlepidotus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 255 (*M.
inornatus* Gill).
Isaciella Jordan and Fessler, Rept. U. S. Fish Com., XVII, 1889-1891
(1893), 497 (*Pristipoma brevipinne* Steindachner).
- 2590. *Microlepidotus brevipinnis* (Steindachner).**
Pacific coast of Mexico.
Pristipoma (Hamulopsis) brevipinne Steindachner, Ichth. Notizen, VIII,
1869, 10, Mazatlan.
- 2591. *Microlepidotus inornatus* Gill. *Jopaton.***
Gulf of California; La Paz and Mazatlan.
Microlepidotus inornatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862,
256, Cape San Lucas, Lower California.
- Genus 888. *GENYATREMUS* Gill.**
Genyatremus Gill, Proc., Acad. Nat. Sci. Phila., XIV, 1862, 256 (*Diagramma cavifrons* Cuvier and Valenciennes = *Lutianus luteus* Bloch).
- 2592. *Genyatremus luteus* (Bloch).**
Lesser Antilles to Brazil.
Lutianus luteus Bloch, Nat. Ausl. Fische, IV, 1795, 111, Martinique;
on a drawing by Plumier.
Grammistes hepatus Bloch and Schneider, Syst. Ichth., 1801, 187,
after Bloch.
Diagramma cavifrons Cuvier and Valenciennes, Hist. Nat. Poiss., V,
1830, 217 (290), pl. 123, Rio Janeiro.

✓ Family 206. SPARIDÆ. Porgies; Pargos

Genus 889. OTRYNTER Jordan and Evermann.

Otrynter Jordan and Evermann, Check List, 1895, 388 (*Stenotomus caprinus* Bean).

2593. Otrynter caprinus (Bean).

Banks off the west coast of Florida.

Stenotomus caprinus Bean, Proc. U. S. Nat. Mus., V, 1882, 426, Snapper Banks off Pensacola, Fla.

Genus 890. STENOTOMUS Gill.

Stenotomus Gill, Canadian Nat., 1865, 266 (*Sparus argyrops* Linnæus = *Sparus chrysops* Linnæus).

Stenesthes Jordan, Copeia, Oct. 4, 1917 (same type); substitute name, there being an earlier genus *Stenotoma*.

2594. Stenotomus versicolor (Mitchill). *Common scup; Porgy; Scuppaug; Mishcuppuog; Bream; Fair maid.*
Atlantic coast of United States, from Cape Cod to North Carolina; common northward.
Labrus versicolor Mitchell, Trans. Lit. and Phil. Soc., I, 1815, 404, New York.
Sargus ambassis Günther, Cat., I, 1859, 449, New York.

2595. Stenotomus chrysops (Linnæus). *Southern porgy.*

Southern Atlantic and Gulf coast of United States, Cape Hatteras to Texas.

Sparus argyrops Linnæus, Syst. Nat., Ed. XII, 1766, 471, Charleston; young.

Sparus chrysops Linnæus, loc. cit., 1766, 471, Charleston, S. C.

Sparus xanthurus Lacépède, Hist. Nat. Poiss., IV, 1803, 120, Charleston; after *argyrops* L.

Chrysophrys aculeata Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1830, 137, Charleston, S. C.

Genus 891. CALAMUS Swainson.

Calamus Swainson, Nat. Hist. Fishes, etc., II, 1839, 171 and 221 (*Pagellus calamus* Cuvier and Valenciennes).

Grammateus Poey, Ann. Lyc. Nat. Hist. N. Y., X, 1874 (*Pagellus microps* Guichenot = *Pagellus penna* Cuvier and Valenciennes).

2596. Calamus calamus (Cuvier and Valenciennes). *Saucereye porgy; Pez de pluma.*
West Indies, north to Florida Keys.
Pagellus calamus Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830, 152 (206), pl. 152, Martinique; San Domingo.
Calamus megacephalus Swainson, Nat. Hist. Fishes, II, 1839, 222, after Cuvier and Valenciennes.
Pagellus orbitarius Poey, Memorias, II, 1861, 201, Cuba.

2597. Calamus bajonado (Bloch and Schneider). *Jolt-head porgy; Bajonado.*
West Indies, north to Florida Keys.
Sparus bajonado Bloch and Schneider, ed. Bloch, Syst. Ichth., 1801, 284, West Indies; after Parra.
Bajonado Parra, Dif. Piezas Hist. Nat. Cuba, 1787, 13, lam. 8, Habana.
Pajellus caninus Poey, Memorias, II, 1861, 199, Habana.
Calamus plumatula Guichenot, Rev. Pagels in Mém. Soc. Sci. Nat. Cherbourg, XIV, 1868, 119, Martinique.

2598. Calamus proridens Jordan and Gilbert. *Littlehead porgy; Pez de pluma.*
West Indies, north to the Florida Keys.
Calamus proridens Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 150, Key West, Fla.

2599. Calamus pennatula Guichenot. *Littlehead porgy.*
West Indies.
Calamus pennatula Guichenot, Révision des Pagels in Mém. Soc. Sci. Nat. Cherbourg, XIV, 1868, 116, Martinique.

- 2600.** *Calamus brachysomus* (Lockington). *Mojarra garabata*.
 Gulf of California and neighboring waters, locally abundant from
 Magdalena Bay to Panama.
Sparus brachysomus Lockington, Proc. U. S. Nat. Mus., III, 1880, 284,
 Magdalena Bay, Lower California.
- 2601.** *Calamus leucosteus* Jordan and Gilbert. *Whitebone porgy*.
 South Atlantic coast of United States; known only from the markets
 of Charleston, S. C.
Calamus leucosteus Jordan and Gilbert, in Jordan, Cat. Fish. N. Amer.,
 in Rept. U. S. Fish Com., XIII, 1885, 879, Charleston, S. C.
- 2602.** *Calamus macrops* Poey.
 Cuba.
Calamus macrops Poey, Ann. Lyc. Nat. Hist. N. Y., X, 1872, 181,
 Pl. VII, fig. 3, Habana.
- 2603.** *Calamus penna* (Cuvier and Valenciennes). *Littlemouth porgy; Sheep-head porgy; Speckled porgy*.
 Southern Florida to Brazil; known from southern Florida, Key West,
 Rio Janeiro, St. Thomas, Habana, Camaru, and Rio Grande do Sul.
Pagellus penna Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830,
 154 (209), Brazil.
Pagellus microps Guichenot, in Ramon de la Sagra, Hist. Ile Cuba,
 1853, 83, pl. 3, fig. 1, Habana.
Pagellus humilis Poey, Synopsis, 1868, 308, Habana.
Pagellus milneri Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 134,
 Charlotte Harbor, Fla.
- 2604.** *Calamus arctifrons* Goode and Bean. *Grass porgy; Shad porgy*.
 Gulf of Mexico, from Pensacola to Key West; West Indies; Porto Rico.
Calamus arctifrons Goode and Bean, Proc. U. S. Nat. Mus., V, 1882,
 425, Pensacola, Fla.
- 2605.** *Calamus medius* (Poey).
 West Indies; Habana.
Grammateus medius Poey, Ann. Lyc. Nat. Hist. N. Y., X, 1872, 183,
 pl. 7, fig. 4, Habana.
- 2606.** *Calamus kendalli* Evermann and Marsh. *Pluma*.
 Porto Rico.
Calamus kendalli Evermann and Marsh, Rept. U. S. Fish Com., XXV,
 1899, 354, Mayaguez, Porto Rico.
- Genus 892. **PAGRUS** Cuvier.
Pagrus Cuvier, Règne Animal, ed. 1, 1817, 272 (*Sparus argenteus* Bloch
 and Schneider = *Sparus pagrus* Linnaeus).
- 2607.** *Pagrus pagrus* (Linnaeus). *Red porgy; Besugo; Pargo colorado*.
 Southern Europe and south Atlantic and Gulf coasts of the United
 States; south to Uruguay; less common in America.
Sparus pagrus Linnaeus, Syst. Nat., Ed. X, 1758, 279, southern Europe.
Sparus argenteus Bloch and Schneider, Syst. Ichth., 1801, 271, Europe.
Pagrus vulgaris Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830,
 104 (142), pl. 148, southern Europe.
- Genus 893. **LAGODON** Holbrook. *Chopa spina; Sailors' choice*.
Lagodon Holbrook, Ichth. South Carolina, 1855, 56 (*Sparus rhombooides*
 Linnaeus).
- 2608.** *Lagodon rhomboides* (Linnaeus). *Pinfish; Bream; Sailors' choice;*
Chopa spina; Sargo.
 Atlantic and Gulf coasts of United States; Cape Cod to Cuba.
Sparus rhomboides Linnaeus, Syst. Nat., Ed. XII, 1766, 470, Charleston,
 S. C.
- Genus 894. **SALEMA** Jordan and Evermann.
*Salema*¹ Jordan and Evermann, Check List, 1895, 390 (*Perca uni-*
maculata Bloch).

¹ Not *Seleima* Bowdich 1825 = *Kyphosus*.

2609. *Salema rhomboidalis* (Linnæus). *Salema*.

West Indies, north to Key West, south to Rio Janeiro.
Perca rhomboidalis Linneus, Syst. Nat., Ed. X, 1758, Jamaica; after
 Bream, Browne, Jamaica, 1756, 446.
Perca unimaculata Bloch, Nat. Ausl. Fische, VI, 1792, 75, Brazil.
Sparus salin Lacépède, Hist. Nat. Poiss., IV, 1802, 136; based on
 Bloch.
Sargus humerimaculatus Quoy and Gaimard, Voy. Freycinet, Zool.,
 1824, 297, Rio Janeiro.
Sargus flavolineatus Cuvier and Valenciennes, Hist. Nat. Poiss., VI,
 1830, 44 (60), Cuba.
Cynædus brama Gronow, Cat. Fishes, ed. Gray, 1854, 56, South Carolina.
Sargus caribæus Poey, Memorias, II, 1861, 197, Cuba.

2610. *Salema tridens* (Poey).

Cuba.
Sargus tridens Poey, Enumeratio, 1875, 57, Cuba.

Genus 895. *ARCHOSARGUS* Gill. *Sheepsheads*.

Archosargus Gill, Canadian Nat., 1865, 266 (*Sparus probatocephalus* Walbaum).

2611. *Archosargus probatocephalus* (Walbaum). *Sheepshead; Sargo raiado*.
 Atlantic and Gulf coasts of the United States; Cape Cod to Florida Keys and Texas.

Sparus probatocephalus Walbaum, Artedi Pisc., 1792, 295, New York;
 based on *Sparus* of Schöpf.
Sparus oviccephalus Bloch and Schneider, Syst. Ichth., 1801, 280, New York.
Sargus ovis Mitchell, Trans. Lit. and Phil. Soc. N. Y., I, 1814, 392, pl. 2, fig. 5, New York.

2612. *Archosargus aries* (Cuvier and Valenciennes).

Honduras to Brazil; Rio Janeiro; Maracaibo; Belize.

Sargus aries Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830, 42 (58), Rio Janeiro; Maracaibo.

Genus 896. *DIPLODUS* Rafinesque.

Diplodus Rafinesque, Indice, 1810, 54 (*Sparus annularis* L.).
Sargus Cuvier, Règne Anim., I, 1817, 272 (*Sparus sargus* Linnæus); preoccupied in insects.

2613. *Diplodus holbrookii* (Bean).

South Atlantic and Gulf coasts of the United States; Cape Hatteras to Cedar Keys.

Sargus holbrookii Bean, Forest and Stream, June 13, 1878, 359, Charleston, S. C.

2614. *Diplodus argenteus* (Cuvier and Valenciennes). *Sargo*.

West Indies; Florida and Bermudas, south to Argentina.

Sargus argenteus Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830, 44 (60), Brazil.

Sargus caudimacula Poey, Memorias, II, 1861, 198, Cuba.

2615. *Diplodus sargus* (Linnæus). *Sargo*.

Coasts of southern Europe; Bermudas.

Sargus sargus Linnæus, Syst. Nat., Ed. X, 1758, 278, Mediterranean.

Sargus variegatus Lacépède, Hist. Nat. Poiss., IV, 1803, 207, Mediterranean.

Sargus raucus Geoffrey St. Hilaire, Descr. de l'Egypte, Poiss., 1813, Pl. XVIII, fig. 1, coast of Egypt.

Sargus rondeletii Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830, 10 (14), pl. 141, Mediterranean.

Sargus vitula Cuvier and Valenciennes, loc. cit., 35 (48), Naples.

Family 207. *GIRELLIDÆ*Genus 897. *GIRELLA* ² Gray.

Girella Gray, Illus. Indian Zool., II, 1833, 98 (*G. punctata* Gray).

Camarina Ayres, Proc. Calif. Acad. Nat. Sci., II, 1860 (1863), 81 (*C. nigricans* Ayres).

² *Melanichthys* Schlegel, 1844 = *Inciadens* Gill, 1862; differs from *Girella* in having two to four rows of incisors and is probably a valid genus.

- 2616.** *Girella nigricans* (Ayres). *Greenfish; California bluefish; White spot.*
Coast of southern California from Monterey to Cape San Lucas.
Camarina nigricans Ayres, Proc. Calif. Acad. Nat. Sci., II, 1860, 81, fig. 22, California.
Girella dorsomacula Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 244, Cape San Lucas.

- Genus 898. INCISIDENS** Gill.
Incisidens Gill, Proc. Acad. Sci. Phila., XIV, 1862, 244 (*Crenidens simplex** Richardson).
Aplodon Duméril, in Thominot, Bull. Soc. Philom. Paris, 1883, 141 (*Aplodon margaritiferum* Duméril); name preoccupied.

- 2617.** *Incisidens simplicidens* (Osburn and Nichols.)
Gulf of California.
Girella simplicidens Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 165, fig. 10, San Francisquito.

Family 208. KYPHOSIDÆ. Pilotfishes

Genus 899. KYPHOSUS Lacépède.

- Kyphosus* Lacépède, Hist. Nat. Poiss., III, 1802, 114 (*K. bigibbus* Lacépède = *fucus* Lacépède).
Pimelepterus Lacépède, Hist. Nat. Poiss., IV, 1803, 429 (*P. boscii* Lacépède).
Dorsuarius Lacépède, loc. cit., V, 1803, 482 (*D. nigrescens* Lacépède.).
Xyster Lacépède, loc. cit., X, 484 (*X. fuscus* Lacépède).
Xysterus Rafinesque, Anal. Nature, 1815, 95; substitute for *Xyster*.
Selaima Bowdich, Fishes Madeira, 1825, 238 (*S. aurata* Bowdich).
Opisthistius Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 245 (*Sciæna tahmel* Forskål).
Cyphosus of authors, corrected spelling.

- 2618.** *Kyphosus analogus* (Gill). *Salema.*
Pacific coast of tropical America, Gulf of California to Panama.
Pimelepterus analogus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 245, Cape San Lucas.

- 2619.** *Kyphosus incisor* (Cuvier and Valenciennes). *Chopa amarilla.*
West Indies to Brazil and the Canaries; Tortugas.
Pimelepterus incisor Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 198 (266), Brazil.
? *Saleima aurata* Bowdich, Excursion Madeira, 1825, 238, Bona Vista Island.
Pimelepterus flavo-lineatus Poey, Repertorio, I, 1866, 319, Habana.

- 2620.** *Kyphosus elegans* (Peters). *Chopa.*
Pacific coast of tropical America, from Guaymas to Mazatlan; a species apparently the same in Hawaii.
Pimelepterus elegans Peters, Berlin. Monatsb., 1869, 707, Mazatlan, Mexico.
Pimelepterus sandvicensis Sauvage, Bull. Soc. Philom., July, 1880 (7), IV, 221, Sandwich Islands.

- 2621.** *Kyphosus metzelaari* Jordan and Evermann.
Caribbean Sea.
Cyphosus elegans Metzelaar, Trop. Atl. Vissch., 1919, 44, Curaçao; Venezuela; not *Pimelepterus elegans* Peters.
This species, according to Metzelaar's account, is extremely close to *Kyphosus elegans* of the Pacific, but the scales are larger; 52 in the lateral line (67 in *elegans*), the teeth are also fewer, "15 in upper jaw"; in *elegans* they are "86 in each jaw."
Kyphosus metzelaari Jordan and Evermann, Proc., Calif. Acad. Sci., 4th Ser., XVI, Apr. 27, 1927, 506, Curaçao, Venezuela.

* This species with entire teeth is said to be the female of *Girella tricuspidata*. If this is true the genus *Incisidens* may not be tenable.

2622. *Kyphosus sectatrix* (Linnæus). Rudderfish; Bermuda chub; Chub; *Chopa blanca*.

West Indies, ranging from Cape Cod to Brazil, crossing the ocean to the Canary Islands; accidental in the Mediterranean; one taken at Palermo.

Perca marina sectatrix Catesby, Nat. Hist. Carolina, II, 1743, 8, Carolinas.

Perca sectatrix Linnæus, Syst. Nat., Ed. XII, 1766, 486 (correction), misprinted for *salatatrix*, after Catesby.

Chatodon cyprinaceus Gmelin, Syst. Nat., I, 1789, 1269, tropical America.

Pimelepterus bosquii Lacépède, Hist. Nat. Poiss., IV, 1803, 429, South Carolina.

Pimelepterus oblongior Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 197 (264), locality unknown.

Pimelepterus bosci sicula Doderlein, Nat. Siciliano, Giornale Scienze Nat., III, 1883, 83, Gulf of Palermo.

2623. *Kyphosus lutescens* (Jordan and Gilbert).

Revillagigedo Archipelago.

Pimelepterus lutescens Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 229, Braithwaite Bay, Socorro Island.

Genus 900. SECTATOR Jordan and Fesler.

Sectator Jordan and Fesler, Rept. U. S. Fish Com., XVII, 1889-91, 534 (*Pimelepterus oxyurus* Jordan and Gilbert).

2624. *Sectator oxyurus* (Jordan and Gilbert).

Panama.

Pimelepterus oxyurus Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 327, 328, Bay of Panama.

Genus 901. HERMOSILLA Jenkins and Evermann.

Hermosilla Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 144 (*H. azurea* Jenkins and Evermann).

2625. *Hermosilla azurea* Jenkins and Evermann.

Gulf of California.

Hermosilla azurea Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 144, Guaymas, Mexico.

2626. *Hermosilla robusta* Osburn and Nichols.

Gulf of California.

Hermosilla robusta Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 166, fig. 11, Tiburon Island.

Family 209. INERMIIDÆ

Genus 902. INERMIA Poey.

Inermia Poey, Memorias, II, 1861, 193 (*I. vittata* Poey).

2627. *Inermia vittata* (Poey). Boga.

Caribbean Sea, north to Cuba; often in large schools, off Venezuela.

Inermia vittata Poey, Memorias, II, 1861, 193, Habana.

Family 210. MÆNIDÆ

Genus 903. SPICARA Rafinesque.

Spicara Rafinesque, Caratteri, 1810, 51 (*S. flexuosa* Rafinesque = *Sparus smaris* Linnæus).

Smaris Cuvier, Règne Animal, ed. 1, 1817, 269 (*Sparus smaris* Linnæus).

2628. *Spicara martinica* (Cuvier and Valenciennes).

West Indies; probably an error.

Smaris martinicus Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830, 319 (424), Martinique; most likely an error, type probably from the Mediterranean.

Family 211. GERRIDÆ. Mojarras; Silver perch

Genus 904. EUCINOSTOMUS Baird and Girard. Mojarritas.

Eucinostomus Baird and Girard, Ninth Smithson. Rept., 1854, 344 (*E. argenteus* Baird and Girard = *Gerres gula* Cuvier and Valenciennes).

- 2629.** *Eucinostomus gula* (Cuvier and Valenciennes). *Silver jenny; Mojarra de ley; Petite gueule; Shad.*
Gerres gula Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830, 349
(464), Martinique.
Eucinostomus argenteus Baird and Girard, Ninth Smith. Rept., 1854,
335, Bessley Point, N. J.
Eucinostomus gulula Poey, Enumeratio, 1875, 54, pl. 2, Habana.
Diapterus homonymus Goode and Bean, Proc. U. S. Nat. Mus., II, 1879,
340, Clearwater Harbor, Fla.
- 2630.** *Eucinostomus pseudogula* Poey.
Tortugas; West Indies to Brazil; Bermudas; St. Lucia; Bahia.
Eucinostomus pseudo-gula Poey, Enumeratio, 1875, 53, pl. 1, Cuba.
Gerres jonesii Günther, Ann. Mag. Nat. Hist., 1879, 150, Bermudas.
Eucinostomus harenigulus Goode and Bean, Proc. U. S. Nat. Mus., II,
1879, 132, west Florida.
- 2631.** *Eucinostomus californiensis* (Gill). *Mojarra cantilena.*
Pacific coast of Mexico; Guaymas and Cape San Lucas to Panama.
Diapterus californiensis Gill, Proc. Acad. Nat. Sci., Phila., XIV, 1862,
245, Cape San Lucas, Lower California.
Diapterus gracilis Gill, loc. cit., 245, Cape San Lucas.
Diapterus downii Gill, loc. cit., XV, 1863, 162, Panama.
- 2632.** *Eucinostomus elongatus* Meek and Hildebrand.
Panama.
Eucinostomus californiensis elongatus Meek and Hildebrand, Field Mus.
Nat. Hist., Zool., XV, 1925, 586, Sobago Island, Panama.
- Genus 905. ULÆMA Jordan and Evermann.**
Ulæma Jordan and Evermann, Proc. Calif. Acad. Sci., V, 1895, 471
(*Diapterus lefroyi* Goode).
- 2633.** *Ulæma lefroyi* (Goode).
West Indies, on sandy shores north to Cedar Keys; Bermudas; Cuba;
Key West; recorded north at Beaufort, N. C.
Diapterus lefroyi Goode, Amer. Jour. Sci. Arts, 3d ser., VIII, 1874, 123,
Bermudas.
- 2634.** *Ulæma producta* (Poey).
Cuba.
Eucinostomus productus Poey, Enumeratio, 1875, 55, Habana.
- 2635.** *Ulæma meeki* (Eigenmann).
Rio San Juan, in western Cuba.
Eucinostomus meeki Eigenmann, Bull. U. S. Fish Com., XXII, 1902,
229, fig. 10, San Juan River, just below its first ford, west Cuba;
apparently a *Ulæma* having but two anal spines, the second haemal
spine not described.
- Genus 906. GERRES (Cuvier) Quoy and Gaimard. Mojarras blancas.**
Gerres Cuvier in Quoy and Gaimard, Voyage *Uranie*, 1824, 293,
restricted by Jordan 1919 to *Gerres vaigiensis* Quoy and Gaimard =
Sciæna argyrea Forskål, an ally of *Mugil cinereus* Walbaum. The
only other species named in this earlier paper is *Gerres gula*, belonging
to *Eucinostomus*. This use of the name *Gerres* from Cuvier's MS.
antedates that in the *Règne Animal*, 1829, and unfortunately must
stand.
Podager Gistel, Thierreich, IX, 1848, a substitute for *Gerres* regarded
as preoccupied by *Gerris*; *Podager* is actually so, however.
Catocænum Cantor, Malayan Fishes, 1849, 55, same type, being a
substitute for *Gerres*, deemed preoccupied by *Gerris*, a genus of insects.
Xystæma Jordan and Evermann, Proc. Calif. Acad. Sci., V, 1895, 471
(*Mugil cinereus* Walbaum).
- 2636.** *Gerres cinereus* (Walbaum). *Mojarra de Casta; Mojarra blanca;*
Broad shad.
Coasts of tropical America; West Indies, north to Florida; Habana;
Jamaica; Martinique; Bahamas; Barbados; Florida Keys.
Turdus cinereus Peltius Catesby, Nat. Hist. Carolinas, II, 1743, 11,
Bahamas.
Mugil cinereus Walbaum, Artedi Pisc., 1792, 228, Bahamas; after
Catesby.

- Gerres aprion* Cuvier, Règne Animal, ed. 2, II, 1829, 188; based on
Catesby.
- Gerres zebra* Müller and Troschel, in Schomburgk's Hist. Barbados,
1848, 668, Barbados.
- Gerres squamipinnis* Günther, Cat., I, 1859, 349, Jamaica; Guatemala.
- 2637. *Gerres simillimus* Regan.**
Pacific coast of tropical America, Mazatlan to Panama.
Gerres simillimus Regan, Biol. Centrali-Amer., Pisces, 1906, 38, Panama.
- 2638. *Gerres havana* (Nichols).**
Miami, Florida; Cuba.
Xystema havana Nichols, Bull. Amer. Mus. Nat. Hist., XXXI, 1912,
189, fig. 2, Marianas Beach, Habana.
- Genus 907. MOHARRA Poey.**
Moharra Poey, Enumeratio, 1875, 50 (*Gerres rhombeus* Cuvier and
Valenciennes).
- 2639. *Moharra rhombea* (Cuvier and Valenciennes).**
West Indies and Atlantic coast of tropical America; Jamaica; San
Domingo; Martinique; Puerto Cabello; Habana; Aspinwall; Rio
Magdalena; Santa Lucia; Bahia.
Gerres rhombeus Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830,
345 (459), Martinique; San Domingo.
- 2640. *Moharra evermanni* (Meek and Hildebrand).**
Atlantic waters of Panama.
Diapterus evermanni Meek and Hildebrand, Field Mus. Nat. Hist.,
Zool., XV, 1925, 594, Pl. LXII, Mindi River near Mindi, Panama
Canal Zone.
- Genus 908. DIAPTERUS Ranzani.**
Diapterus Ranzani, Nov. Comment. Bonon., V, 1842, 340 (*Diapterus*
auratus Ranzani, a species near *D. olisthostomus*).
- 2641. *Diapterus aureolus* (Jordan and Gilbert).**
Panama; only the type known.
Gerres aureolus Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881,
328, Panama.
- 2642. *Diapterus axillaris* (Günther).**
Pacific coast of Mexico.
Gerres axillaris Günther, Proc. Zool. Soc. Lond., 1864, 152, Chiapas.
- 2643. *Diapterus peruvianus* (Cuvier and Valenciennes). *Mojarra de las*
aletas amarillas.**
West coast of tropical America; Mazatlan; Salina Cruz; Chiapas;
Panama; Peru.
Gerres peruvianus Cuvier and Valenciennes, Hist. Nat. Poiss., VI,
1830, 351 (467), Payta, northern Peru.
Gerres brevirostris Sauvage, Bull. Soc. Philom. Paris, III, 1879, 208,
Rio Guayas, Ecuador.
- 2644. *Diapterus olisthostomus* (Goode and Bean). *Irish pompano; Mutton-*
fish.
West Indies, north to southern Florida.
Gerres olistostoma Goode and Bean, Proc. U. S. Nat. Mus., V, 1882,
423, Indian River, Fla.**
- Genus 909. EUGERRES Jordan and Evermann.**
Eugerres Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI,
Apr. 27, 1927, 506 (*Gerres plumieri* Cuvier and Valenciennes).
- 2645. *Eugerres lineatus* (Humboldt). *Mojarra china.*
Pacific coast of Mexico; Acapulco; Mazatlan; San Blas; Chiapas.
Smaris lineatus Humboldt, Observ. Zool., II, 1811, 185, pl. 46, Acapulco.**
- 2646. *Eugerres brasiliensis* (Cuvier and Valenciennes). *Pataó.*
West Indies to Brazil.
Gerres brasiliensis Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830,
344, 458, Brazil.
Gerres patao Poey, Synopsis, VI, 1868, 320, Habana; scarcely dis-
tinguishable from *G. lineatus* of the Pacific coast.
Gerres embryx Jordan and Starks, in Jordan and Evermann, Fishes
North and Mid. Amer., 1898, 1379, Charleston, S. C.; large example.**

2647. *Eugerres brevimanus* (Günther).
 Pacific coast of tropical America and Panama Bay.
Gerres brevimanus Günther, Proc. Zool. Soc. Lond., 1864, 152, Chiapam.
2648. *Eugerres plumieri* (Cuvier and Valenciennes). *Mojarra*.
 Atlantic coast of tropical America and West Indies; Habana; Porto Rico; San Domingo; Jamaica; Pernambuco; Bahia; Aspinwall; Guatemala.
Gerres plumieri Cuvier and Valenciennes, Hist. Nat. Poiss., VI, 1830, 340 (452), pl. 467, Antilles; Porto Rico.
2649. *Eugerres mexicanus* (Steindachner). *Mexican mojarra*.
 Rio Teapa, Mexico.
Gerres mexicanus Steindachner, Ueber eine Neue Gerres-Art aus Mexico, Verh. K. K. Ges. Wien, XIII, 1863, 383, Rio Teapa, Mexico.

✓ Family 212. MULLIDÆ. Surmulletts

Genus 910. **MULLUS** Linnaeus. *Surmulletts*.
Mullus Linnaeus, Syst. Nat., Ed. X, 1758, 299 (*M. barbatus* Linnaeus).

2650. *Mullus auratus* Jordan and Gilbert. *Goatfish; Red mullet*.
 Eastern coast of North America, Cape Cod to Pensacola; rare.
Mullus barbatus auratus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 280, Pensacola, Fla.

Genus 911. **UPENEUS** Cuvier. *Goatfishes*.
Upeneus Cuvier, Règne Animal, Ed. II, 1829, 157 (*vittatus*; *russelii*; *bifasciatus*; *trifasciatus*; first restricted by Bleeker to *bifasciatus*, an ally to *U. maculatus*).
Pseudupeneus Bleeker, Versl. Akad. Amsterdam, XIV, 1862, 134 (*P. prayenensis* Bleeker).
Parupeneus Bleeker, Neder. Tijdschr. Dierk., I, 1863, 342 (*Mullus bifasciatus* Lacépède).
Mullhypeneus Poey, Repertorio, II, 1867, 160 (*Upeneus flavovittatus* Poey).
Brachymullus Bleeker, Arch. Neerl. Sci. Nat., XI, 1876, 333 (*Upeneus tetraspis* Günther).
Hypeneus Agassiz, corrected spelling.

2651. *Upeneus maculatus* (Bloch). *Red goatfish; Salmonete*.
 West Indies and Brazil, Key West to Rio Janeiro; generally common.
Mullus maculatus Bloch, Naturgesch. Ausl. Fische, VII, 1893, 95, Brazil.
Upeneus punctatus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 355, 482, Martinique.

2652. *Upeneus dentatus* Gill.
 Pacific coast of Mexico, Cape San Lucas, La Paz, and the Tres Marias Islands.
Upeneus dentatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 256, Cape San Lucas.

2653. *Upeneus parvus* Poey.
 Cuba.
Upeneus parvus Poey, Memorias, I, 1853, 226, Cuba.

2654. *Upeneus martinicus* Cuvier and Valenciennes. *Yellow goatfish; Salmonete amarilla; King mullet*.
 West Indies, north to Key West.
Upeneus martinicus Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 356 (483), Martinique.
Upeneus balteatus Cuvier and Valenciennes, loc. cit., 356 (483), Cuba.
Upeneus flavovittatus Poey, Memorias, I, 1853, 224, Cuba.

2655. *Upeneus xanthogrammus* Gilbert.
 Coast of Lower California.
Upeneus xanthogrammus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 553, La Paz, Lower California.

2656. *Upeneus grandisquamis* Gill. *Chivo.*

Pacific coast of Mexico and Central America, Guaymas to Panama.
Upeneus grandisquamis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863,
 168, western coast of Central America.
Upeneus tetraspis Günther, Proc. Zool. Soc. Lond., 1864, 148, Panama.

Genus 912. **MULLOIDES** Bleeker.

Mulloides Bleeker, Verh. Bat. Gen., XXII, 1849, 12 (*Mullus flavolineatus* Lacépède).

2657. *Mulloides rathbuni* (Evermann and Jenkins).

Gulf of California.

Upeneus rathbuni Evermann and Jenkins, Proc. U. S. Nat. Mus., XIV,
 1891, 158, pl. 2, fig. 4, Bay of Guaymas, Sonora.

✓ Family 213. **SCIÆNIDÆ. Croakers; Roncadores**Genus 913. **LARIMUS** Cuvier and Valenciennes.

Larimus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 108 (145),
 (*L. breviceps* Cuvier and Valenciennes).

Amblyscion Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 165 (*A. argenteus* Gill).

Monosira Poey, Anales de l'Hist. Nat. Esp., 1881, 326 (*M. stahli* Poey).

Subgenus **LARIMUS.**2658. *Larimus effulgens* Gilbert.

Pacific coast of Mexico, Sonora to Panama.

Larimus effulgens Gilbert, in Jordan and Evermann, Fishes North and
 Mid. Amer., 1898, 1421, San Juan Lagoon, Sonora.

Subgenus **AMBLYSCION** Gill.2659. *Larimus argenteus* (Gill).

Panama.

Amblyscion argenteus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 165,
 western coast of Central America, the extreme form of the series.

2660. *Larimus acclivis* Jordan and Bristol.

Pacific coast of Mexico and Central America, from Sonora to Panama.

Larimus acclivis Jordan and Bristol (S. B.), in Jordan and Evermann,
 Fishes North and Mid. Amer., 1898, 1422, San Juan Lagoon, Sonora.

2661. *Larimus breviceps* Cuvier and Valenciennes. *Cabezon.*

West Indies, south to Brazil.

Larimus breviceps Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830,
 108 (145), pl. 140, Brazil; San Domingo.

Monosira stahli Poey, Anal. Soc. Española Hist. Nat., X, 1881, 326,
 Pl. VI, Porto Rico.

2662. *Larimus pacificus* Jordan and Bollman.

Pacific Ocean, off coast of Colombia.

Larimus pacificus Jordan and Bollman, Proc. U. S. Nat. Mus., XII,
 1889, 161, Albatross station 2802, 8° 38' N., 79° 31' 30" W., between
 Galapagos Islands and Panama.

2663. *Larimus fasciatus* Holbrook. *Banded larimus.*

South Atlantic coast of the United States, from Chesapeake Bay to
 Galveston, Tex.; not common.

Larimus fasciatus Holbrook, Ichth. South Carolina, 1855, 153, pl. 22,
 fig. 1, Charleston, S. C.

Genus 914. **ODONTOSCION** Gill.

Odontoscion Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 18 (*Corvina*
dentex Cuvier and Valenciennes).

2664. *Odontoscion dentex* (Cuvier and Valenciennes). *Corvina.*

West Indies, north to the Tortugas.

Corvina dentex Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830,
 103 (139), pl. 109, San Domingo.

2665. *Odontoscion xanthops* Gilbert.

Panama.

Odontoscion xanthops Gilbert in Jordan and Evermann, Check List,
 1895, 396, Panama.

Genus 915. CORVULA Jordan and Eigenmann.

Corvula Jordan and Eigenmann, Review Scienidæ Europe and America, in Rept. U. S. Fish Com., XIV, 1886, 377 (35), (*Johnius batabanus* Poey).

2666. *Corvula batabana* (Poey).

Cuba and Porto Rico.

Johnius batabanus Poey, Memorias, II, 1861, 184, Batabano, south coast of Cuba.

Genus 916. VACUOQUA Jordan and Evermann.

Vacuoqua Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, Apr. 27, 1927, 506 (*Corvina macrops* Steindachner).

This genus differs from the aberrant *Corvula batabana* in having the deeper, symmetrical body and the silvery coloration of *Bairdiella* and related species.

2667. *Vacuoqua macrops* (Steindachner). *Vacuoqua*.

Pacific coast of tropical America; Mazatlan to Panama.

Corvina macrops Steindachner, Ichth. Beitr., III, 1875, 24, fig. 2. Panama.

2668. *Vacuoqua sialis* (Jordan and Eigenmann).

Florida Keys.

Corvula sialis Jordan and Eigenmann, Rept. U. S. Fish Com., XIV, 1886, 379, Key West, Fla.

2669. *Vacuoqua subæqualis* (Poey).

West Indies.

Corvina subæqualis Poey, Ann. Lyc. Nat. Hist. N. Y., XI, 1875, 58, Cuba.

2670. *Vacuoqua sanctæ-luciae* (Jordan).

West Indies.

Corvula sanctæ-luciae Jordan, Proc. U. S. Nat. Mus., XII, 1889, 649, Port Castries, St. Lucia.

Genus 917. ELATTARCHUS Jordan and Evermann.

Elattarchus Jordan and Evermann, Check List, 1895, 397 (*Odontoscion archidium* Jordan and Gilbert).

2671. *Elattarchus archidium* (Jordan and Gilbert).

Panama.

Odontoscion archidium Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 317, Panama.

Genus 918. BAIRDIELLA Gill. *Mademoiselles*.

Bairdiella Gill, Cat. Fish. E. Coast N. Amer., Suppl. to Proc. Acad. Nat. Sci. Phila., XIII, 1861, 33 (*Bodianus argyroleucus* Mitchell = *Diplerodon chrysurus* Lacépède).

2672. *Bairdiella chrysura* (Lacépède). *Mademoiselle; Yellowtail; Silver perch.*

South Atlantic and Gulf coasts of United States, north to New York; very abundant on our sandy shores from Long Island to Texas.

Caranx chrysurus Lacépède, Hist. Nat. Poiss., III, 1802, 64, South Carolina; after Linnaeus.

Perca punctata Linnaeus, Syst. Nat., Ed. XII, 1766, 482, in part, South Carolina.

Bodianus argyro-leucus et *exiguus* Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 417, pl. 6, fig. 9, New York.

Bodianus pallidus Mitchell, loc. cit., 420, New York.

Homopriion subtruncatus Gill, Cat. Fish. E. Coast N. Amer., Suppl. to Proc. Acad. Nat. Sci. Phila., XIII, 1861, 33, South Carolina; after Holbrook.

2673. *Bairdiella ensifera* (Jordan and Gilbert).

Panama; common.

Sciaena ensifera Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 313, Bay of Panama; Punta Arenas.

Corvina fulgens Vaillant, Miss. Sci. au Mexique, 1883, 164, Pacific coast of Mexico.

- 2674.** *Bairdiella icistia* (Jordan and Gilbert). *Corbineta*.
Pacific coast of Mexico; rather common about Mazatlan.
Sciæna icistia Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881,
356, Mazatlan.
- 2675.** *Bairdiella ronchus* (Cuvier and Valenciennes). *Ronco; Ground drummer*.
Atlantic coasts of tropical America; generally common in the West
Indies and along the coast of Brazil.
Corvina ronchus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830,
79 (107), Maracaibo; Surinam.
- 2676.** *Bairdiella bedoti* (Regan).
Cuba.
Sciæna (Bairdiella) bedotis Regan, Rev. Suisse Zool., XIII, 1905, 391,
Pl. VI, Cuba.
- Genus 919. NECTOR** Jordan and Evermann.
Nector Jordan and Evermann, Fishes North and Mid. Amer., 1898,
1432 (*Corvina chrysoleuca* Günther).
- 2677.** *Nector armatus* (Gill).
Both coasts of tropical America; not uncommon on the Pacific coast
about Panama, and equally abundant on the Atlantic coast.
Bairdiella armata Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 164,
west coast of Central America.
Corvina acutirostris Steindachner, Ichth. Beitr., III, 1875, 28, pl. 4,
Panama.
- 2678.** *Nector veræ-crucis* (Jordan and Dickerson).
Gulf of Mexico, Vera Cruz.
Bairdiella veræ-crucis Jordan and Dickerson, Proc. U. S. Nat. Mus.,
XXXIV, 1908, 16, fig. 1, Vera Cruz, Mexico.
- 2679.** *Nector alatus* (Jordan and Gilbert).
Pacific coast of Central America.
Sciæna aluta Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 232,
La Union, Salvador.
- 2680.** *Nector chrysoleucus* (Günther).
Panama.
Corvina chrysoleuca Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov.
27), 387, 427, pl. 67, fig. 1, Panama.
- Genus 920. STELLIFER** (Cuvier) Oken.
Les Stellifères Cuvier, Règne Anim., Ed. I, 1817, 283 (*Bodianus stellifer*
Bloch).
Stellifer (Cuvier) Oken, Isis, I, 1817, 1182 (*Bodianus stellifer* Bloch).
Stelliferus Stark, Elem. Nat. Hist., I, 1828, 459 (*Bodianus stellifer*
Bloch).
Homopriion Holbrook, Ichth. South Car., 1st ed. (mostly destroyed by
fire), 1855, 168 (*Homopriion lanceolatus* Holbrook).
- 2681.** *Stellifer stellifer* (Bloch).
Coasts of Guiana and Brazil.
Bodianus stellifer Bloch, Naturgesch. Ausl. Fische, IV, 1790, 55, Cape
of Good Hope.
Corvina trispinosa Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830,
80 (109), Brazil; Cayenne.
- 2682.** *Stellifer lanceolatus* (Holbrook).
South Atlantic and Gulf coasts of United States, Charleston to Texas.
Homopriion lanceolatus Holbrook, Ichth. South Carolina, Ed. I, 1855,
168, pl. 23, Port Royal Sound, S. C.
- 2683.** *Stellifer ericymba* (Jordan and Gilbert).
Panama.
Sciæna ericymba Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881,
311, Bay of Panama.
- 2684.** *Stellifer microps* (Steindachner).
Coast of Brazil and Guiana.
Corvina microps Steindachner, Ichth. Notizen, I, 1864, 6, pl. 2, fig. 1,
Guiana.
Corvina stellifera Günther, Cat., II, 1860, 299, West Indies (not *Bodianus*
stellifer Bloch).

2685. *Stellifer mindii* Meek and Hildebrand.
 Atlantic coast of Panama.
Stellifer mindii Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1925, 626, Pl. LXVI, fig. 2, Mindi Reef, near Colon.
- Genus 921. *ZESTIS* Gilbert.
Zestis Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1439 (*Stellifer oscitans* Jordan and Gilbert).
2686. *Zestis oscitans* (Jordan and Gilbert).
 Panama.
Sciena oscitans Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 312, Bay of Panama.
2687. *Zestis furthii* (Steindachner).
 Panama.
Corvina (Homopriion) fürthii Steindachner, Ichth. Beitr., III, 1875, 26, fig. 3, Panama.
2688. *Zestis melanochir* (Eigenmann).
 North coast of Colombia.
Stellifer melanochir Eigenmann, Proc. Am. Phil. Soc., LVI, 1917, 682, Tumneo.
- Genus 922. *ZESTIDIUM* Gilbert.
Zestidium Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1439 (*Stellifer illecebrosus* Gilbert).
2689. *Zestidium illecebrosum* (Gilbert).
 Panama.
Stelliferus illecebrosus Gilbert, in Jordan and Evermann, Check List, 1895, 398, and Fishes North and Mid. Amer., 1896, 1443, Panama.
2690. *Zestidium colonense* (Meek and Hildebrand).
 Coral reefs near Colon, Atlantic side of Isthmus of Panama.
Stellifer colonensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1925, 623, Mindi Reef, Colon.
- Genus 923. *STELLICARENS* Gilbert.
Stellicarens Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1445 (*Stellifer zestocarus* Gilbert).
2691. *Stellicarens zestocarus* (Gilbert).
 Panama.
Stelliferus zestocarus Gilbert, in Jordan and Evermann, Check List, 1895, 398, and Fishes North and Mid. Amer., 1896, 1445, Panama.
- Genus 924. *OPHIOSCION* Gill.
Ophioscion Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 164 (*Ophioson typicus* Gill).
2692. *Ophioscion adustus* (Agassiz).
 West Indies to coast of Brazil.
Corvina adusta Agassiz, in Spix, Pisc. Brasil., 1829, 126, pl. 70, Brazil.
2693. *Ophioscion punctatissimus* Meek and Hildebrand.
 Panama.
Ophioscion punctatissimus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1925, pt. 2, 644, Pl. LXVIII, Panama Canal at Cristobal.
2694. *Ophioscion typicus* Gill.
 Panama.
Ophioscion typicus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 165, west coast Central America.
Corvina ophioscion Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 387, 428, Panama.
2695. *Ophioscion strabo* Gilbert.
 Panama.
Ophioscion strabo Gilbert, Proc. U. S. Nat. Mus., XIX, 1896, 444, San Juan Lagoon, south of Guaymas, Mexico.

2696. *Ophioscion simulus* Gilbert.*Panama.**Ophioscion simulus* Gilbert, in Jordan and Evermann, Check List, 1895, 398, and Fishes North and Mid. Amer., 1896, 1449, Panama.2697. *Ophioscion imiceps* (Jordan and Gilbert).*Panama.**Sciæna imiceps* Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 309, Bay of Panama.2698. *Ophioscion scierus* (Jordan and Gilbert).

Pacific coast of tropical America, from Mazatlan to Panama.

Sciæna sciera Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 480, Panama.Genus 925. *SIGMURUS* Gilbert.*Sigmurus* Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1446 (*Corvina vermicularis* Günther).2699. *Sigmurus vermicularis* (Günther).*Panama.**Corvina vermicularis* Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 387, 427, pl. 67, fig. 2, Panama.Genus 926. *SCLENOPNS* Gill. *Red drums.**Sciænops* Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 30 (*Perca ocellata* Linnaeus).2700. *Sciænops ocellata* (Linnaeus). *Red drum; Channel bass; Redfish; Pescado colorado; Bull redfish; Bass; Red bass; Sea bass; Spotted bass; Spottail.*
South Atlantic and Gulf coasts of the United States, New York to Texas; very abundant south.*Perca ocellata* Linnaeus, Syst. Nat., Ed. XII, 1766, 483, South Carolina.
Lutjanus triangulum Lacépède, Hist. Nat. Poiss., IV, 1802, 181, pl. 24, fig. 3, Sumatra.*Sciæna imberbis* Mitchell, Trans. Lit. and Phil. Soc. New York, 1815, 411, New York.Genus 927. *CALLAUS* Jordan.*Callaus* Jordan, Rept. U. S. Fish Com., XIV, 1886, 401, 406 (59) (*Corvina delicosa* Tschudi).2701. *Callaus delicosa* (Tschudi).

Pacific coast of South America, from Panama to Peru.

Corvina delicosa Tschudi, Fauna Peruana, Ichthyol., 1845, 8, Peru.Genus 928. *RHINOSCIION* * Gill. *Red roncador.**Rhinoscion* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 17 (*Amblodon saturnus* Girard).2702. *Rhinoscion saturnus* (Girard). *Red roncador; Black croaker.*Coast of southern California, from Santa Barbara to Cedros Island.
Amblodon saturnus Girard, U. S. Pac. R. R. Surv., 1858, 98, San Diego, Calif.*Corvina (Johnius) jacobi* Steindachner, Ichth. Beitr., VIII, 1879, 3, San Diego; young.Genus 929. *RONCADOR* Jordan and Gilbert.*Roncador* Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 28 (*Corvina stearnsi* Steindachner).2703. *Roncador stearnsi* (Steindachner). *Roncador.*

Coast of southern California, north to Santa Barbara.

Corvina stearnsi Steindachner, Ichth. Beitr., III, 1875, 22, San Diego, Calif.Genus 930. *LEIOSTOMUS* Lacépède. *Goodies.**Leiostomus* Lacépède, Hist. Nat. Poiss., IV, 1803, 439 (*L. xanthurus* Lacépède).*Liostomus* authors, revised spelling.

* This genus is very close to the European genus *Cortina* on the one hand and to the Peruvian *Chelotrema* on the other, but probably deserves to stand. There is no genuine species in America of *Sciæna* as now restricted.

2704. *Leiostomus xanthurus* Lacépède. *Spot; Goody; Postcroaker; Oldwife; Lafayette; Cape May goody; Porgy; Yellowtail.*
 South Atlantic and Gulf coasts of United States; Cape Cod to Texas; once doubtfully recorded from Martinique.
Leiostomus xanthurus Lacépède, Hist. Nat. Poiss., IV, 1802, 439, pl. 10, fig. 1, Carolina.
Mugil obliquus Mitchell, Trans. Lit. Phil. Soc. New York, 1815, 405, New York.
Sciæna multofasciata LeSueur, Jour. Acad. Nat. Sci. Phila., II, 1821, 255, east Florida.
Leiostomus humeralis Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 105 (141), pl. 110, New York.
- Genus 931. **PACHYPOPS** Gill.
Pachypops Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 87 (*Micro-*
pogon trifilis Müller and Troschel).
2705. *Pachypops furcræa* (Lacépède).
 Rivers and estuaries of Guiana.
Perca furcræa Lacépède, Hist. Nat. Poiss., IV, 1802, 398, 424, Surinam.
Corvina biloba Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 83 (112), habitat unknown.
- Genus 932. **GENYONEMUS** Gill.
Genyonemus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 87 (*Leiostomus lineatus* Ayres).
2706. *Genyonemus lineatus* (Ayres). *Little roncador; Kingfish; Croaker.*
 Coast of southern California, San Francisco to Cedros Island.
Leiostomus lineatus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 24, San Francisco.
- Genus 933. **MICROPOGON** Cuvier and Valenciennes. *Croaker.*
Micropogon Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 158 (213) (*M. lineatus* Cuvier and Valenciennes = *Perca undulata* Linnaeus).
Micropogonias Bonaparte, Ittiologia Generale, 1831, 104 (*Perca undulata* Linnaeus; substitute for *Micropogon* said to be preoccupied; but the genus of birds so named dates from 1838).
2707. *Micropogon undulatus* (Linnaeus). *Croaker; Roncodina; Corvina; Hardhead.*
 South Atlantic and Gulf coasts of United States; Cape Cod to Texas.
Perca undulata Linnaeus, Syst. Nat., Ed. XII, 1766, 483, South Carolina.
Sciæna croker Lacépède, Hist. Nat. Poiss., IV, 1802, 309, Carolinas.
Bodianus costatus Mitchell, Trans. Lit. Phil. Soc. New York, 1815, 417, New York.
Micropogon lineatus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 160 (215), pl. 119, New York.
2708. *Micropogon furnieri* (Desmarest). *Verrugato; Whitemouth drummer.*
 West Indies and coasts of South America.
Umbrina furnieri Desmarest, Première Décade Ichth., 22, pl. 2, fig. 3, 1823, 182, Habana.
? *Micropogon argenteus* Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 162 (218), Surinam.
2709. *Micropogon megalops* Gilbert.
 Gulf of California.
Micropogon megalops Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 64, Gulf of California in 14 fathoms at Albatross station 3021.
2710. *Micropogon ectenes* Jordan and Gilbert. *Verrugate.*
 Pacific coast of Mexico, Mazatlan.
Micropogon ectenes Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 355, Mazatlan.
2711. *Micropogon altipinnis* Günther.
 Panama.
Micropogon altipinnis Günther, Proc. Zool. Soc. Lond., 1864, 149, San Jose; Panama; Chiapam.

Genus 934. UMBRINA Cuvier.

Umbrina Cuvier, Règne Anim., Ed. I, II, 1817, 297 (*Sciæna cirrhosa* Linnæus).

Attitus Gistel, Naturgesch. d. Thierreichs, 1848, 109 (*Sciæna cirrhosa* Linnæus).

2712. *Umbrina broussonnetii* Cuvier and Valenciennes.

West Indies; Florida to Brazil.

Umbrina broussonnetii Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 139 (187), Jamaica.

2713. *Umbrina coroides* Cuvier and Valenciennes.

Southern Florida to the West Indies and Brazil.

Umbrina coroides Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 139 (187), Brazil.

2714. *Umbrina gracilicirrhus* Metzelaar.

Coast of Venezuela.

Umbrina gracilicirrhus Metzelaar, Tropisch Atlantisch Vissch., 1919, 72, coast of Venezuela.

2715. *Umbrina roncador* Jordan and Gilbert. *Yellow-finned roncador*;

Yellow-tailed croaker.

Coast of southern California, from Point Conception to Guaymas.

Umbrina roncador Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 277, Pequena Bay, west coast of Lower California.

2716. *Umbrina xanti* Gill. *Codorniz*.

Pacific coast of tropical America, Cape San Lucas to Panama.

Umbrina xanti Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 257, Cape San Lucas.

Umbrina analis Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 387, 426. Panama.

2717. *Umbrina sinaloæ* Scofield.

Pacific coast of Mexico; Mazatlan.

Umbrina sinaloæ Scofield, Proc. Calif. Acad. Sci., VI, 1896, 220; Pl. XXV, Mazatlan, Mexico.

2718. *Umbrina dorsalis* Gill.

Pacific coast of Mexico.

Umbrina dorsalis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 257, Cape San Lucas.

Genus 935. MENTICIRRHUS Gill. *Kingfishes*.

Menticirrhus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 86 (*Perca alburnus* Linnæus = *Cyprinus americanus* Linnæus).

2719. *Menticirrhus americanus* (Linnæus). *Carolina whiting*; *Sand whiting*;

Kingfish; *Sea smelt*; *Sea mullet*.

South Atlantic and Gulf coasts of United States; Chesapeake Bay to Texas.

Cyprinus americanus Linnaeus, Syst. Nat., Ed. X, 1758, 321, Carolinas; (based on *Alburnus americanus* the whiting of Catesby, nonbinomial; not *Cyprinus americanus* of 12th edition of *Systema Naturæ*, which is a cyprinoid, *Abramis boscii* Cuvier and Valenciennes).

Perca alburnus Linnæus, Syst. Nat., Ed. XII, 1766, 482. Charleston, S. C.

Umbrina phalæna Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 167, Indianola and Brazos Santiago, Tex.

2720. *Menticirrhus saxatilis* (Bloch and Schneider). *Kingfish*; *Sea mink*;

Northern whiting; *Hake*; *Barb*.

Atlantic and Gulf coasts of the United States; Cape Ann to Key West and Pensacola; most common northward.

Johnius saxatilis Schneider, Ed. Bloch, Syst. Ichth., 1801, 75, New York.

2721. *Menticirrhus undulatus* (Girard). *California whiting*; *Sand sucker*.

Southern California, north to Santa Barbara.

Umbrina undulata Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 148, San Diego, Calif.

- 2722. *Menticirrhus martinicensis* (Cuvier and Valenciennes). Jewsharp drummer.**
 West Indies; Martinique; Jamaica to Patagonia; common in Brazil.
Umbrina martinicensis Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 186, Martinique.
Umbrina gracilis Cuvier and Valenciennes, loc. cit., 1830, 186, Brazil.
Umbrina arenata Cuvier and Valenciennes, loc. cit., 141 (190), Brazil.
Umbrina januaria Steindachner, Ichth. Beitr., V, 1876, 122, Rio Janeiro.
- 2723. *Menticirrhus nasus* (Günther).**
 Panama.
Umbrina nasus Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 387, 426, Panama.
Menticirrhus simus Jordan and Eigenmann, Review Sciænidæ, Rept. U. S. Fish Com., XIV, 1886, 427, Mazatlan and Panama.
- 2724. *Menticirrhus panamensis* (Steindachner).**
 Pacific coast of tropical America; Mazatlan to Panama.
Umbrina panamensis Steindachner, Ichth. Beitr., IV, 1875, 9, Panama.
- Genus 936. *UMBRULA* Jordan and Eigenmann.**
Umbrula Jordan and Eigenmann, Review Sciænidæ, Rept. U. S. Fish Com., XIV, 1886, 424 (*Umbrina littoralis* Holbrook).
- 2725. *Umbrula littoralis* (Holbrook).** Surf whiting; Silver whiting; Shore whiting.
 South Atlantic and Gulf coasts of United States, North Carolina to Texas.
Umbrina littoralis Holbrook, Ichth. South Carolina, Ed. I, 1855, 142, pl. 20, fig. 1, South Carolina.
- 2726. *Umbrula elongata* (Günther).** Verrugata.
 Pacific coast of tropical America, Mazatlan to Panama.
Umbrina elongata Günther, Proc. Zool. Soc. Lond., 1864, 148, Chiapam.
- Genus 937. *PARALONCHURUS* Bocourt.**
Paralonchurus Bocourt, Nouv. Arch. Mus. d'Hist. Nat., IV, 1869, 21 (*P. petersi* Bocourt).
- 2727. *Paralonchurus petersi* Bocourt.**
 Coast of Central America; rare at Panama.
Paralonchurus petersi Bocourt, Nouv. Archiv du Muséum, IV, 1869, 22, La Union, San Salvador.
- Genus 938. *POLYCLEMUS* Berg.** Corvalos.
Polycirrus Bocourt, Nouv. Arch. Mus. Paris, IV, 1869, 22 (*P. dumerili* Bocourt); name preoccupied in worms.
Polycomus Berg, Ann. Mus. Nac. Buenos Aires, IV, 1895, 52 (*Polycirrus dumerili* Bocourt).
- 2728. *Polycomus dumerili* (Bocourt).**
 Panama.
Polycirrus dumerili Bocourt, Nouv. Arch. Mus. d'Hist. Nat., IV, 1869, 23, La Union, San Salvador.
Genyonemus fasciatus Steindachner, Ichth. Beitr., II, 1875, 31, Panama.
- Genus 939. *ZONOSCIUS* Jordan and Evermann.**
Zonoscion Jordan and Evermann, Check List, 1895, 401 (*Polycirrus rathbuni* Jordan and Bollman).
- 2729. *Zonoscion rathbuni* (Jordan and Bollman).**
 Panama.
Polycirrus rathbuni Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 162, Panama.
- Genus 940. *ZACLEMUS* Gilbert.**
Zaclemus Gilbert, in Jordan and Evermann, Check List, 1895, 401 (*Paralonchurus goodei* Gilbert).
- 2730. *Zaclemus goodei* (Gilbert).**
 Panama.
Paralonchurus goodei Gilbert, in Jordan and Evermann, Check List, 1895, 401, and Fishes North and Mid. Amer., 1896, 1480, Panama.

Genus 941. LONCHIURUS Bloch.

Lonchiurus Bloch, Naturgesch. Ausl. Fische, VII, 1793, 144 (*Lonchiurus*

barbata Bloch = *Perca lanceolata* Bloch).

Lonchurus authors, preferred spelling.

2731. *Lonchiurus lanceolatus* (Bloch).

West Indies to Guiana.

Perca lanceolata Bloch, Nov. Act. Sc., Copenh., III, 1788, 383, India.

Lonchiurus depressus Bloch, Naturgesch. Ausl. Fische, VII, 1793, 144, Surinam.

Genus 942. POGONIAS Lacépède. Sea drums.

Pogonias Lacépède, Hist. Nat. Poiss., III, 1802, 138 (*P. fasciatus*

Lacépède = *Labrus cromis* Linnæus).

Pogonathus Lacépède, loc. cit., V, 1803, 121 (*P. courbina* Lacépède).

Pogonathus Bonaparte, Ittiologia Generale, 1831 (*Pogonias fasciatus* Lacépède).

2732. *Pogonias cromis* (Linnæus). Gray drum; Drum; Black drum; Drumfish; Banded drum; Big drum.

Atlantic coast of America; Long Island to mouth of Rio Grande.

Labrus cromis Linnæus, Syst. Nat., Ed. XII, 1766, 479, Carolinas.

Pogonias fasciatus Lacépède, Hist. Nat. Poiss., III, 1802, 137.

Mugil grunniens Mitchell, Rept. in part, Fishes New York, 1814, 15, New York.

Sciæna fusca Mitchell, Trans. Lit. Phil. Soc. N. Y., 1815, 409, New York.

Sciæna gigas Mitchell, loc. cit., 412, New York.

2733. *Pogonias courbina* (Lacépède).

Guiana to Uruguay; rather common on coast of Brazil.

Pogonathus courbina Lacépède, Hist. Nat. Poiss., V, 1803, 121, Rio de la Plata.

Genus 943. APLODINOTUS Rafinesque. River drums; Thunderpumpers.

Aploidonotus Rafinesque, Jour. de Phys., LXXXVIII, 1819, 418 (*A. grunniens* Rafinesque).

Amblodon Rafinesque, Jour. de Phys., 1819, 421, based on teeth supposed to belong to *Ictiobus*, later restricted by Rafinesque to *Amblodon grunniens*.

Eutychelithus Jordan, Man. Vert., Ed. I, 1876, 242 (*Corvina richardsoni* Cuvier and Valenciennes, an injured or deformed example of *A. grunniens*).

Haploidonotus Gill, revised spelling.

2734. *Aploidonotus grunniens* Rafinesque. Fresh-water drum; Gaspergou; Thunderpumper; Lake sheepshead; Croaker; Bubbler; White perch.

Great Lakes to Texas; abundant in lakes and large streams between the Alleghenies and the Great Plains.

Aploidonotus grunniens Rafinesque, Jour. de Phys., 1819, 88, Ohio River.

Sciæna oscula LeSueur, Jour. Acad. Nat. Sci. Phila., 1822, 252, pl. 13, Lake Ontario.

Sciæna grisea LeSueur, loc. cit., 254, Ohio River at Pittsburgh, Pa.

Corvina richardsoni Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 100, Lake Huron, based on an abnormal specimen.

Amblodon concinnus Agassiz, Amer. Jour. Sci. Arts, XVII, 1854, 307, Tennessee River.

Amblodon lineatus Agassiz, loc. cit., 307, Osage River, Mo.

2735. *Aploidonotus neglectus* (Girard).

Streams of eastern Mexico from Rio Grande to Rio Usumacinta.

Amblodon neglectus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 167, mouth of the Rio Grande; Rio Usumacinta. This Mexican form apparently distinguished by larger scales (43 instead of 55 to 60).

Genus 944. EQUETUS Rafinesque.

Eques Bloch, Nat. Ausl. Fische, VII, 1793, 91 (*E. americanus* Bloch =

Chælodon lanceolatus Linnæus); preoccupied.

Equetus Rafinesque, Anal. Nat., 1815, 89; substitute for *Eques* Bloch.

- 2736.** *Equetus lanceolatus* (Linnaeus). *Ribbonfish; Guapena; Serrana.*
 West Indies, ranging northward to Pensacola.
Ribband fish, Edwards, Gleanings, pl. 210, 1758, Guadeloupe; Carolina.
Chætodon lanceolatus Linnaeus, Syst. Nat., Ed. X, 1758, 277, "Caraibes
 Islands"; based on Edwards, pl. 210.
?Serrana Parra, Piezas, d'Hist. de Cuba, 1787, pl. 2, Cuba.
Eques americanus Bloch, Nat. Ausl. Fische, VII, 1793, West Indies.
Eques balteatus Cuvier, Règne Anim., ed. 2, II, 1829, 175, pl. 29, fig. 2,
 Martinique, after Edwards.
Sciæna edwardsi Gronow, Cat. Fish., ed. Gray, 1854, 53, after Edwards.
- Genus 945. PAREQUES Gill.**
Pareques Gill in Goode, Bull. U. S. Nat. Mus., V, 1875, 50 (*Grammistes
 acuminatus* Bloch and Schneider).
- 2737. Pareques viola** (Gilbert).
 Panama.
Eques viola Gilbert in Jordan and Evermann, Check List, 1895, 402,
 Panama.
- 2738. Pareques acuminatus** (Bloch and Schneider).
 West Indies; South Carolina to Brazil.
Grammistes acuminatus Bloch and Schneider, ed. Bloch, Syst. Ichth.,
 1801, 184, no locality; after Seba.
Eques lineatus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830,
 126 (169), Brazil.
- 2739. Pareques umbrosus** (Jordan and Eigenmann).
 Southeast coast of the United States, Charleston to Pensacola.
Eques acuminatus umbrosus Jordan and Eigenmann, Review Sciænidæ,
 Rept. U. S. Fish Com., XIV, 1886, 440, Charleston and Pensacola.
- 2740. Pareques punctatus** (Bloch and Schneider). *Serrana hispana.*
 West Indies.
Serrana hispanis Parra, Piezas, Hist. Nat. Cuba, 1787, 2, pl. 2, lower
 figure, Cuba.
Eques punctatus Bloch and Schneider, Syst. Ichth., 1801, 106, Cuba;
 based on Parra, 2, pl. 2, fig. 2.
- 2741. Pareques pulcher** (Steindachner).
 Tortugas Archipelago; Barbados.
Eques pulcher Steindachner, Ichth. Notizen, VI, 1867, 43, Barbados.
- ✓ **Family 214. OTOLITHIDÆ. Weakfishes; Queenfishes**
- Genus 946. SERIPHUS Ayres.**
Seriphus Ayres, Proc. Calif. Acad. Nat. Sci., II, 1858-1862 (1863;
 read Nov. 5, 1860), 80 (*S. politus* Ayres).
- 2742. Seriphus politus** Ayres. *Queenfish; White croaker.*
 Coast of southern California from Point Conception to Cedros Island;
 common on sandy shores.
Seriphus politus Ayres, Proc. Calif. Acad. Nat. Sci., II, 1858-1862 (1863;
 read Nov. 5, 1860), 80, fig. 21, about San Diego.
- Genus 947. ISOPISTHUS Gill.**
Isopisthus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 18 (*Ancylodon
 parvipinnis* Cuvier and Valenciennes).
- 2743. Isopisthus parvipinnis** (Cuvier and Valenciennes).
 Coast of Brazil, north to Cayenne and Guiana.
Ancylodon parvipinnis Cuvier and Valenciennes, Hist. Nat. Poiss., V,
 1830, 62 (84), pl. 105, Cayenne.
Isopisthus affinis Steindachner, Denksch. Mat. Nat. Kais. Acad. Wiss.,
 XLI, Bd. I, Abth., 1879, 43, pl. 2, fig. 2, Porto Alegre, Brazil.
- 2744. Isopisthus remifer** Jordan and Gilbert.
 Panama, on sandy shores, rather common.
Isopisthus remifer Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881,
 320, Panama.

2745. *Isopisthus harroweri* Fowler.

Atlantic shore of Central America.

Isopisthus harroweri Fowler, Proc. Acad. Nat. Sci. Phila., LXVIII, 1916, 402, Colon.

Genus 948. *NEBRIS* Cuvier and Valenciennes.

Nebris Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 111 (149) (*N. microps* Cuvier and Valenciennes).

2746. *Nebris microps* Cuvier and Valenciennes.

Atlantic coast of northern South America; sandy shores.

Nebris microps Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 111 (149), pl. 112, Surinam.

2747. *Nebris occidentalis* Vaillant. *Guavina*.

Pacific coast of tropical America.

Nebris occidentalis Vaillant, Bull. Mus. Paris, 1897, 124, Pacific coast of Central America.

Nebris zetus Jordan and Starks, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1417, Panama.

Genus 949. *MACRODON* Schinz.

Ancylodon Cuvier, in Oken, Isis, 1817, 1182 (*Lonchurus ancylodon* Schneider); name preoccupied in mammals.

Macrodon Schinz, Das Thierreich, II, 1822, 482 (*Lonchurus ancylodon* Schneider).

Nomalus Gistel, Thierreich, X, 1848, substitute for *Ancylodon*.

Sagenichthys Carlos Berg, Ann. Mus. Nat. Buenos Aires, 1895, 42 (*Lonchurus ancylodon* Schneider); substitute for *Ancylodon*.

2748. *Macrodon ancylodon* (Bloch and Schneider). *Pescadillo del rey*.

West Indies; Guiana; Brazil; and Uruguay; rather scarce.

Lonchurus ancylodon Bloch and Schneider, ed. Bloch, Syst. Ichth., 1801, 102, pl. 25, Surinam.

Ancylodon jaculidens Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 60 (81), Cayenne.

2749. *Macrodon mordax* (Gilbert and Starks).

Panama.

Sagenichthys mordax Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 121, Pl. XVI, Panama.

Genus 950. *BUCCONE* Jordan and Evermann.

Buccone Jordan and Evermann, Check List, 1895, 394 (*Cestreus prædatorius* Jordan and Gilbert).

2750. *Buccone prædatoria* (Jordan and Gilbert). *Bocone*.

Coast of Panama; scarce.

Cestreus prædatorius Jordan and Gilbert, Rept. U. S. Fish Com., XIV, 1886, 363, Panama.

Genus 951. *CYNOSCIION* Gill.

Cestreus Gronow, Cat. Fish., ed. Gray, 1854, 49 (*Cestreus carolinensis* Gronow = *Otolithus nebulosus* Cuvier and Valenciennes); preoccupied in gobies.

Cynoscion Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 81 (*Johnius regalis* Bloch and Schneider).

Apseudobranchus Gill, loc. cit., XIV, 1862, 18 (*Otolithus tæ-ræ* Cuvier and Valenciennes = *Cheilodipterus acoupa* Lacépède).

Subgenus *CYNOSCIION* Gill.2751. *Cynoscion acoupa* (Lacépède). *Acoupa; Toeroe*.

Atlantic coast of South America. Brazil north to Venezuela; generally common.

Cheilodipterus acoupa Lacépède, Hist. Nat. Poiss., III, 1802, 546, Cayenne.

Lutjanus cayanensis Lacépède, loc. cit., IV, 1802, 196, Cayenne.

Otolithus rhomboidalis Cuvier, Règne Anim., ed. 2, II, 1829, 173, Cayenne.

Otolithus toe-roe Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 54 (72), pl. 103, Cayenne.

2752. *Cynoscion squamipinnis* (Günther).
 Pacific coast of tropical America; known from a few specimens taken at La Union and Panama.
Otolithus squamipinnis Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 387, 429, Panama.
2753. *Cynoscion othonopterus* Jordan and Gilbert.
 Gulf of California.
Cynoscion othonopterum Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 274, Punta San Felipe, Mexico.
2754. *Cynoscion obliquatus* (Cuvier and Valenciennes).
 Martinique.
Otolithus obliquatus Cuvier and Valenciennes, in Sauvage, Bull. Soc. Philom. Paris, III, 1879, 209, Martinique.
2755. *Cynoscion jamaicensis* (Vaillant and Bocourt). *Mongolar drummer*.
 Jamaica and Porto Rico.
Otolithus jamaicensis Vaillant and Bocourt, Miss. Sci. au Mexique, Poissons, 1874, 156, Jamaica.
2756. *Cynoscion regalis* (Bloch and Schneider). *Common weakfish; Squeaque; Sea trout; Squit; Shecuts; Chickwick; Salt-water trout; Gray trout*.
 Atlantic and Gulf coasts of United States, Cape Cod southward to Mobile.
Johnius regalis Bloch and Schneider, Syst. Ichth., 1801, 75, New York.
Roccus comes Mitchell, Report, in part, Fishes New York, 1814, 25, New York.
Labrus squeaque Mitchell, Trans. Lit. Phil. Soc. New York, 1815, 396, pl. 2, fig. 1, New York.
2757. *Cynoscion nothus* (Holbrook). *Bastard weakfish*.
 South Atlantic and Gulf coasts of United States; rather rare at Charleston. This unspotted fish is now generally regarded as a pale variant of *Cynoscion regalis*.
Otolithus nothus Holbrook, Ichth. South Carolina, 1855, 134, pl. 10, fig. 1, South Carolina.
2758. *Cynoscion thalassinus* (Holbrook).
 Pensacola; Pass Christian, Miss.; Hampton Roads, Va.
Otolithus thalassinus Holbrook, Ichth. South Carolina, 1855, 132, pl. 18, fig. 2, Charleston, S. C.
2759. *Cynoscion leiaarchus* (Cuvier and Valenciennes).
 Coasts of Brazil and Guiana.
Otolithus leiaarchus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 58 (78), Brazil; Cayenne.
2760. *Cynoscion microlepidotus* (Cuvier and Valenciennes).
 Coast of Brazil and Guiana.
Otolithus microlepidotus Cuvier and Valenciennes, Hist. Nat. Poiss.; V, 1830, 59 (79), Surinam.
- Genus 952. *ERISCIUS* Jordan and Evermann.
Eriscion Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, Apr. 27, 1927, 506 (*Otolithus nebulosus* Cuvier and Valenciennes).
2761. *Eriscion reticulatus* (Günther). *Corvina*.
 Pacific coast of tropical America, Mazatlan to Panama.
Otolithus reticulatus Günther, Proc. Zool. Soc. Lond., 1864, 149, San Jose de Guatemala; Chiapam.
2762. *Eriscion nebulosus* (Cuvier and Valenciennes). *Spotted weakfish; Spotted sea trout; Spotted Squeaque*.
 South Atlantic and Gulf coasts of the United States; New York to Texas; everywhere common on our southern coast; rare north of Virginia.
Labrus squeaque maculatus Mitchell, Trans. Lit. Phil. Soc., 1815, 396, New York; not *Labrus maculatus* Bloch.
Otolithus nebulosus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 59 (79), locality unknown.
Otolithus carolinensis Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 351, (475), South Carolina.
Otolithus drummondi Richardson, Fauna Bor.-Amer., Fish., 1836, 70, New Orleans.

- 2763. *Eriscion parvipinnis* (Ayres).** *California bluefish.*
 Coast of Lower California; Santa Barbara Islands to Guaymas; common along the coasts of southern California as far north as San Pedro.
Cynoscion parvipinnis Ayres, Proc. Calif. Acad. Nat. Sci., II, 1858-1862 (read Dec. 1, 1861, published Sept., 1862), 156, fig. 48, coast of Lower California.
Otolithus magdalena Steindachner, Ichth. Beit., III, 1875, 34, Magdalena Bay, Lower California.
- 2764. *Eriscion xanthulus* (Jordan and Gilbert).** *Corvina de las aletas amarillas; Yellowmouth.*
 Pacific coast of Mexico; not rare about Mazatlan.
Cynoscion xanthulum Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 460, Mazatlan.
- 2765. *Eriscion albus* (Günther).**
 Pacific coast of tropical America.
Otolithus albus Günther, Proc. Zool. Soc. Lond., 1864, 149, Chiapam; Panama.
Otolithus cayennensis Vaillant and Bocourt, Miss. Sci. Mexique, Poiss., 1874, 156, La Union, Salvador.
- 2766. *Eriscion stolzmanni* (Steindachner).**
 Pacific coast of tropical America, from Panama to Peru.
Otolithus stolzmanni Steindachner, Denkschr. Mat. Nat. Kais. Acad. Wiss., XLI, Bd. I, Abth. 35, 1879, pl. 2, fig. 1, Tumbez, Peru.
- 2767. *Eriscion virescens* (Cuvier and Valenciennes).**
 Coasts of Guiana and Brazil.
Otolithus virescens Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 54 (72), Surinam.
Otolithus microps Steindachner, Denkschr. Mat. Nat. Kais. Acad. Wiss., XLI, Bd. I, Abth. 38, 1879, pl. 8, fig. 2, Porto Alegre, Brazil.
- 2768. *Eriscion macdonaldi* (Gilbert).** *Totuava.*
 Gulf of California.
Cynoscion macdonaldi Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 64, head of Gulf of California.
- Genus 953. *ATRACTOSCION* Gill.**
Atractoscion Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 18 (*Otolithus squidens* Cuvier and Valenciennes).
- 2769. *Atractoscion nobilis* (Ayres).** *White sea bass of California.*
 Coast of California, north to San Francisco, occasionally straying as far north as Boca de Quadra Inlet, Alaska.
Johnius nobilis Ayres, Proc. Calif. Acad. Nat. Sci., II, 1860, 77, San Francisco.
Otolithus californiensis Steindachner, Ichth. Beit., III, 1875, 31, San Diego, Calif., and Magdalena Bay, Lower California.
- 2770. *Atractoscion phoxocephalus* (Jordan and Gilbert).**
 Pacific coast of tropical America; Panama.
Cynoscion phoxocephalum Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 318, Panama.
- Family 215. MALACANTHIDÆ. Matajuelos**
- Genus 954. *MALACANTHUS* Cuvier.**
Malacanthus Cuvier, Règne Animal, ed. 2, II, 1829, 264 (*Coryphæna plumieri* Bloch).
- 2771. *Malacanthus plumieri* (Bloch).** *Matajuelo blanco.*
 West Indies, generally common.
Coryphæna plumieri Bloch, Ichth., V, 1787, 119, pl. 175, Martinique; from a drawing by Plumier.

Family 216. BRANCHIOSTEGIDÆ.⁶⁵ Blanquillos

Genus 955. CAULOLATILUS Gill.

Caulolatilus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 240 (?*Latilus chrysops* Cuvier and Valenciennes).
Dekaya Cooper, Proc. Calif. Acad. Nat. Sci., III, pt. 1, 1863 (Nov.), 70, fig. 17 (*D. anomala* Cooper).

2772. *Caulolatilus chrysops* (Cuvier and Valenciennes). *Blanquito*.
 Gulf of Mexico and southward.
Latilus chrysops Cuvier and Valenciennes, Hist. Nat. Poiss., IX, 1833, 496, Brazil.
2773. *Caulolatilus microps* Goode and Bean.
 Gulf of Mexico.
Caulolatilus microps Goode and Bean, Proc. U. S. Nat. Mus., I, 1878, 43, off Pensacola, Fla.
2774. *Caulolatilus princeps* (Jenyns). *Blanquito*.
 Cape San Lucas to the Galapagos.
Latilus princeps Jenyns, Zool. Voy. *Beagle*, Fishes, 1842, 52, Chatham Island, Galapagos Archipelago.
Caulolatilus affinis Gill, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 68, Cape San Lucas.
2775. *Caulolatilus anomalus* (Cooper). *Blanquito; Sea whitefish*.
 Coast of southern California, about rocky islands from Point Conception southward; common.
Dekaya anomala Cooper, Proc. Calif. Acad. Nat. Sci., III, pt. 1, 1863 (Nov.), 70, fig. 17, Santa Catalina Island off southern California.

2776. *Caulolatilus cyanops* Poey. *Blanquito*.
 Coasts of Cuba and Porto Rico.
Caulolatilus cyanops Poey, Repertorio, I, 1866, 311-312, Cuba.
- Genus 956. LOPHOLATILUS Goode and Bean. *Tilefish*.
Lopholatilus Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 205 (*L. chamæleonticeps* Goode and Bean).

- 2777-2778. *Lopholatilus chamæleonticeps* Goode and Bean. *Tilefish*.
 Deep waters of the western Atlantic; abundant at intervals but subject to calamities; a valued food fish.
Lopholatilus chamæleonticeps Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 205; 80 miles S. by E. of No-man's Land, 40° N., 70° W., in 84 fathoms.

Family 217. TRICHODONTIDÆ

Genus 957. TRICHODON Steller.

Trichodon Steller, in Tilesius, Mém. Acad. St. Petersb., IV, 1811, 468 (*Trachinus trichodon* Telesius).

2779. *Trichodon trichodon* (Tilesius). *Sandfish*.
 Coasts of Kamchatka and eastward; Herendeen Bay and elsewhere in Bering Sea, and south to coast of Oregon.
Trachinus trichodon Tilesius, Mém. Acad. St. Petersb., IV, 1811, fig. 8, 73, Kamchatka.
Trachinus gasteropelecus Tilesius, loc. cit., 1811, 466, Kamchatka.
Trichodon lineatus Ayres, Proc. Calif. Acad. Nat. Sci., II, 1860, 60, San Francisco.

Genus 958. ARCTOSCOPUS Jordan and Evermann.

Arctoscopus Jordan and Evermann, Check List, 1895, 464 (*Trichodon japonicus* Steindachner).

2780. *Arctoscopus japonicus* (Steindachner).
 Sea of Japan, eastward to Sitka, Alaska.
Trichodon japonicus Steindachner, Ichth. Beitr., X, 1881, 4 Strietok and Sitka.

⁶⁵ The name *Branchistegus* Rafinesque (1815) has priority over *Latilus* Cuvier and Valenciennes (1830).

Family 218. CIRRHITIDÆ

Genus 959. CIRRHITUS Lacépède.

Cirrhitus Lacépède, Hist. Nat. Poiss., V, 1803, 3 (*C. maculatus* Lacépède).
Cirrhitichthys Günther, Cat., II, 1860, 73; not of Bleeker.
Cirrhites of authors; revised spelling.

2781. *Cirrhitus rivulatus* (Valenciennes).

Cape San Lucas to the Revillagigedos and Galapagos Islands.
Cirrhites rivulatus Valenciennes, Voy. *Vénus*, Poiss., 1855, 309, pl. 3,
fig. 1, Galapagos Islands.
Cirrhitus betaurus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 259,
Cape San Lucas; young.

Family 219. ANTIGONIIDÆ

Genus 960. ANTIGONIA Lowe.

Antigonia Lowe, Proc. Zool. Soc. Lond., 1843, 85 (*A. capros* Lowe).
Caprophanus Müller and Troschel, Horæ Ichth., III, 1845, 28, pl. 5,
fig. 1 (*C. aurora* Müller and Troschel).
Hypsinotus Temminck and Schlegel, Faun. Japon., Poiss., 1844, 84
(*H. rubescens* Günther).

2782. *Antigonia capros* Lowe.

Tropical Atlantic, in rather deep water.
Antigonia capros Lowe, Proc. Zool. Soc. Lond., 1843, 85, Madeira.
Caprophanus aurora Müller and Troschel, Horæ Ichthyologicæ, III,
1845, p. 28, pl. 5, fig. 1, Barbados.

Family 220. SCORPIDÆ. Halfmoon fishes

Genus 961. MEDIALUNA Jordan and Fesler.

Medialuna Jordan and Fesler, Rept. U. S. Fish Com., XVII, 1889-1891,
536 (*Scorpis californiensis* Steindachner).

2783. *Medialuna californiensis* (Steindachner). *Medialuna*; *Halfmoon*.

Coast of southern California from Point Conception southward to
Cedros Island.
Scorpis californiensis Steindachner, Ichth. Beitr., III, 1875, 19, San
Diego.

Suborder SQUAMIPENNES

Family 221. EPHIPPIDÆ

Genus 962. CHÆTOIDIPTERUS Lacépède.

Chætodipterus Lacépède, Hist. Nat. Poiss., IV, 1803, 503 (*Chætodon*
plumieri Bloch = *Chætodon faber* Broussonet).
Parehippus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 165 (*Ephip-*
pus zonatus Girard).

2784. *Chætodipterus faber* (Broussonet). *Angelfish*; *Spadefish*; *Moonfish*;
Tripletail; *Three-banded sheephead*; *Three-tailed porgy*.

West Indies, Cape Cod to Rio Janeiro, south Atlantic coast.
Faber marinus fere quadratus (the pilotfish) Sloane, Jamaica, II, 1793,
290, pl. 251, Jamaica.

Chætodon faber Broussonet, Ichth., V, Dec. 1, 1782, pl. 4, Jamaica.
Zeus quadratus Gmelin, Syst. Nat., I, 1788, 1225, Jamaica; after
Sloane.

Chætodon plumieri Bloch, Naturgesch. Ausl. Fische, III, 1787, 104,
Martinique.

Selene quadrangularis Lacépède, Hist. Nat. Poiss., IV, 1803, 564,
Jamaica; after Sloane.

Chætodon ovisformis Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 247,
pl. 5, fig. 4, New York.

Ephippus gigas Cuvier, Règne Anim., ed. 2, II, 1829, 191, America.

2785. *Chætodipterus zonatus* (Girard).

Pacific coast of America, from San Diego to Panama; common south-
ward.

Ephippus zonatus Girard, Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 110,
San Diego, Calif.

Genus 963. PARAPSETTUS Steindachner.

Parapsettus Steindachner, Ichth. Beitr., III, 1875, 50 (*P. panamensis* Steindachner).

2786. *Parapsettus panamensis* Steindachner.

Panama.

Parapsettus panamensis Steindachner, Ichth. Beitr., III, 1875, 51,
Panama.

Family 222. CHÆTODONTIDÆ. Butterflyfishes; Coralfishes

Genus 964. PROGNATHODES Gill.

Prognathodes Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 238 (*Chelmo pelta* Günther = *Chelmon aculeatus* Poey).

2787. *Prognathodes aculeatus* (Poey).

West Indies.

Chelmon aculeatus Poey, Memorias, II, 1861, 202, Cuba.

Chelmo pelta Günther, Cat., II, 1860, 38, locality unknown.

Genus 965. FORCIPIGER Jordan and McGregor.

Forcipiger Jordan and McGregor, Rept. U. S. Fish Com., XXIV, 1898, 279 (*Chætodon longirostris* Broussonet).

2788. *Forcipiger longirostris* (Broussonet).

South Seas east to Revillagigedo Islands.

Chætodon longirostris Broussonet, Desc. Ichth., I, 1782, 23, pl. 7, Society and Sandwich Islands.

Forcipiger flavissimus Jordan and McGregor, Rept. U. S. Fish Com., XXIV, 1898, 279, Clarion Island, Revillagigedo Archipelago.

Genus 966. CHÆTODON ⁶⁶ Linnæus. Butterflyfishes.

Chætodon Linnæus, Syst. Nat., Ed. X, 1758, 272 (*C. capistratus* Linnæus). *Rabdophorus* Swainson, Classn. Fishes, II, 1839, 21 (*Chætodon ephippium* Cuvier and Valenciennes).

Citraræsus Kaup, Wiegmann's Archiv, XXVI, 1860, 136 (*Chætodon ornatissimus* Solander).

Linophora Kaup, loc. cit., 137 (*Chætodon auriga* Forskål).

Sarothrodus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 99 (*Chætodon capistratus* Gill).

Tholichthys Günther, Ann. Mag. Nat. Hist., I, 4th ser., 1868, 457 (*T. osseus* Günther; a larval form).

Chætodontops Bleeker, Arch. Neerl. Soc. Nat., XI, 1876, 304 (*Chætodon collaris* Bloch).

Tetragonopterus Klein, in Bleeker, loc. cit., 305 (*Chætodon striatus* L.); not *Tetragonopterus* of Cuvier.

Lepidochætodon Bleeker, loc. cit., 305 (*Chætodon unimaculatus* Bloch).

Gonochætodon Bleeker, loc. cit., 306 (*Chætodon triangulum* Kuhl and Van Hasselt).

Orychætodon Bleeker, loc. cit., 306 (*Chætodon lineolatus* Quoy and Gaimard).

Hemichætodon Bleeker, Versl. Ak. Amsterdam, XII, 1876, 38 (*Chætodon capistratus* Linnæus).

Anisochætodon Klunzinger, Fische Rothen Meeres, I, 1884, 3 (*Chætodon auriga* Cuvier and Valenciennes).

Subgenus CHÆTODON Linnæus.

2789. *Chætodon capistratus* Linnæus. Parché.

West Indies, Habana, north to Woods Hole.

Chætodon capistratus Linnæus, Syst. Nat., Ed. X, 1758, 275, India.

Chætodon bricei Smith, Bull. U. S. Fish Com., XVII, 1897, 102, with text fig., Woods Hole, Mass.; young.

2790. *Chætodon striatus* Linnæus. Portuguese butterfly.

West Indies.

Chætodon striatus Linnæus, Syst. Nat., Ed. X, 1758, 275, India.

⁶⁶ These nominal genera, here included under *Chætodon*, are mostly imperfectly defined. The following may be valid: *Chætodon*, lines of scales running upward and backward from the lateral line, no produced rays; *Linophora*, similar but one dorsal ray much produced; *Orychætodon*, scales of sides enlarged, snout sharp; *Chætodontops*, similar, but snout blunt, etc.; *Lepidochætodon*, scales of sides much enlarged, in horizontal rows, etc.; *Rabdophorus*, scales horizontal, much reduced above lateral line (see Jordan and Jordan, Mem. Carnegie Mus. X, 68, 1922).

Subgenus CHÆTODONTOPS Bleeker.

- 2791. *Chætodon bimaculatus* Bloch.⁶⁷ *Parché; Isabelita de la alto.***
 West Indies, north in Gulf Stream to New Jersey and Rhode Island.
Chætodon ocellatus Bloch, Naturgesch. Ausl. Fische, III, pl. 211, 1787,
 105, "also from the East Indies."
Chætodon bimaculatus Bloch, loc. cit., 1790, pl. 219, fig. 1.
Sarothrodus maculo-cinctus Gill, Proc. Acad. Nat. Sci. Phila., XIII,
 1861, 99, Newport, R. I.
Sarothrodus amplexicollis Poey, Enumeratio, 1875, 63, Cuba.
- 2792. *Chætodon nigrirostris* (Gill).**
 Cape San Lucas, Lower California; Panama Bay.
Sarothrodus nigrirostris Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862,
 243, Cape San Lucas, Lower California.
- 2793. *Chætodon humeralis* Günther. *Muñeca; Dollfish.***
 Pacific coast of America, from Guaymas to Panama.
Chætodon humeralis Günther, Cat., II, 1860, 19, "Sandwich Islands,"
 by error for Central America.
- 2794. *Chætodon sedentarius* Poey.**
 West Indian fauna.
Chætodon sedentarius Poey, Memorias, II, 1860, 203, Cuba.
- 2795. *Chætodon gracilis* Günther.**
 West Indies.
Chætodon gracilis Günther, Cat., II, 1860, 20, Caribbean Sea, West
 Indies.
- 2796. *Chætodon aya* Jordan.**
 Gulf of Mexico.
Chætodon aya Jordan, Proc. U. S. Nat. Mus., IX, 1886, 225, Snapper
 Banks, south of Pensacola, Fla.
- 2797. *Chætodon atæniatus* (Poey).**
 Habana Cuba.
Sarothrodus atæniatus Poey, Synopsis, 1868, 353, Cuba.
? *Chætodon unicolor* Sauvage, Bull. Soc. Philom., IV, 1880, 222, Mar-
tinique.
- Genus 967. POMACANTHUS Lacépède. *Chirivitas.***
Pomacanthus Lacépède, Hist. Nat. Poiss., IV, 1803, 517 (*Chætodon*
arcuatus Linnaeus, as restricted by Cuvier).
- 2798. *Pomacanthus arcuatus* (Linnaeus). *Black angel; Chirivita; Portugais.***
 West Indies, north to New Jersey, south to Bahia.
Chætodon arcuatus Linnaeus, Syst. Nat., Ed. X, 1758, 273, "India."
Chætodon aureus Bloch, Naturgesch. Ausl. Fische, III, 1787, 49, Mar-
tinique; on drawing by Plumier.
Chætodon lutescens Bonnaterre, Encycl. Meth., 1788, 82, Jamaica; after
Browne.
Chætodon littoricola Poey, Synopsis, 1868, 351, Cuba.
Pomacanthus balteatus Cuvier and Valenciennes, Hist. Nat. Poiss., VII,
1831, 156 (208), Porto Rico.
Pomacanthus cinctulus Cuvier and Valenciennes, loc. cit., 156 (209),
pl. 185, West Indies (probably Porto Rico).
Pomacanthus quinquecinctus Cuvier and Valenciennes, loc. cit., 157
(210), West Indies (probably Porto Rico).
- 2799. *Pomacanthus paru* (Bloch). *Paru; Indianfish; Flatfish; Chirivita.***
 West Indies, south to Bahia; Tortugas.
Chætodon paru Bloch, Naturgesch. Ausl. Fische, III, 1787, 57, Brazil;
on a drawing.
- Genus 968. POMACANTHOIDES Gill.**
Pomacanthodes Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 244 (*P.*
zonipictus Gill).

⁶⁷ Poey seems to have been the first to recognize the identity of *ocellatus* (which has page priority) with *bimaculatus*. He chose to use the name *bimaculatus*, which should therefore stand.

- 2800. Pomacanthodes zonipectus** Gill. *Mojarra de las piedras.*
 West coast of tropical America, from Mazatlán to Panama.
Pomacanthodes zonipectus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862,
 244, San Salvador.
Pomacanthus crescentalis Jordan and Gilbert, Proc. U. S. Nat. Mus.,
 IV, 1881, 358, Mazatlán.
- Genus 969. CENTROPYGE** Kaup.
Centropyge Kaup, Wiegmann's Archiv, XXVI, 1860, 140 (*Holacanthus*
tibicen Cuvier and Valenciennes).
- 2801. Centropyge passer** (Valenciennes).
 West coast of tropical America; Cape San Lucas to Galapagos Islands.
Holacanthus passer Valenciennes, Voy. Vénus, 1846, 327, pl. 6, Galapagos
 Archipelago.
Holacanthus strigatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 243,
 Cape San Lucas.
- 2802. Centropyge clarionensis** (Gilbert).
 Clarion, Socorro, and San Benedicto Islands of the Revillagigedo group.
Holacanthus clarionensis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890,
 72, Revillagigedo Islands; Clarion, Socorro, and San Benedicto.
- 2803. Centropyge lunata** (Blosser).
 St. Croix.
Holacanthus lunatus Blosser, Ann. Carnegie Mus., VI, 1909, 299, St.
 Croix.
 This small species with low fins and rounded caudal if fairly mature
 can not be placed in *Holacanthus*. It seems to belong to the genus
Centropyge, a genus thus far known only from the Pacific.
- Genus 970. HOLACANTHUS** Lacépède. *Catalinetas.*
Holacanthus Lacépède, Hist. Nat. Poiss., IV, 1803, 525 (*Chætodon*
tricolor Bloch).
- 2804. Holacanthus tricolor** (Bloch). *Rock beauty; Catalineta; Vaqueta de los*
colores.
 West Indies and Bermudas; south to Bahia.
Chætodon tricolor Bloch, Naturgesch. Ausl. Fische, IX, 1795, 103, Cuba.
Catalineta Parra, Descr. Dif. Piez., Hist. Nat. Cuba, 1787, 12, Pl. VII,
 fig. 2, Cuba.
- Genus 971. ANGELICHTHYS** Jordan and Evermann.
Angelichthys Jordan and Evermann, Check List, 1895, 420 (*Chætodon*
ciliaris Linnaeus).
- 2805. Angelichthys ciliaris** (Linnaeus). *Angelfish; Isabelita; Yellow angel;*
Golden angel; Blue angel.
 West Indies north to Key West.
Chætodon ciliaris Linnaeus, Syst. Nat., Ed. X, 1758, 276, Indies, in part.
Isabelita Parra, Dif. Piezas, Hist. Nat. Cuba, 1787, 11, pl. 7, fig. 1, Cuba.
Chætodon squamulosus Shaw, Naturalist's Miscellany, VIII, 1796, 275;
 after the angelfish of Catesby.
Chætodon parræ Bloch and Schneider, Syst. Ichth., 1801, 235, Cuba;
 after *Isabelita* of Parra.
Holacanthus cornutus Desmarest, Décade Ichthyologique, 1823, 44,
 pl. 3, fig. 3, Cuba.
Holacanthus formosus Castelnau, Anim. Amer. Sud, Pois., 1855, 19,
 Pl. XI, fig. 2, Bahia.
- 2806. Angelichthys isabelita** Jordan and Rutter. *Angelfish.*
 West Indies; north to Key West, Fla.
Angelichthys isabelita Jordan and Rutter, in Jordan and Evermann,
 Fishes North and Mid. Amer., 1898, 1685, Key West.
- 2807. Angelichthys townsendi** Nichols and Mowbray.
 Florida.
Angelichthys townsendi Nichols and Mowbray, Bull. Amer. Mus. Nat.
 Hist., XXXIII, 1914, 581, Key West, Fla.

Family 223. ZANCLIDÆ. Moorish idols

Genus 972. ZANCLUS Cuvier and Valenciennes

Zanclus Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 70 (102)*(Chætodon cornutus* Linnaeus).*Gonopterus* Gronow, Cat. Fish., ed. Gray, 1854, 77 (*G. mærens* Gronow = *Chætodon cornutus* Linnaeus).*Gnathocentrum* Guichenot, Ann. Maine et Loire, IX, 1866, 4 (*Zanclus centrognathus* Cuvier and Valenciennes); young.

- 2808. Zanclus cornutus** (Linnaeus). *Moorish idol; Besan; Piquier; Porte enseigne.*

Warm seas of the Pacific; Revillagigedo, Cocos, and Galapagos islands; Hawaii; Samoa; East Indies and islands of Polynesia; common and widely distributed throughout the Pacific.

Chætodon cornutus Linnaeus, Syst. Nat., Ed. X, 1758, 273, Indies; after Aristedi.

Zanclus centrognathos Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 397 (528), near Equator, 75° E.

Gonopterus mærens Gronow, Cat. Fishes, ed. Gray, 1854, 77, India.

Family 224. ACANTHURIDÆ. Surgeonfishes; Tangs

Genus 973. ACANTHURUS Forskål. Surgeonfishes.

Acanthurus Forskål, Deser. Anim., 1775, 59 (*Chætodon unicornis*), etc.*Harpurus* (Forster) Gmelin, Enchiridion, 1778, 84 (*H. fasciatus* Forster).*Aspisurus* Lacépède, Hist. Nat. Poiss., IV, 1803, 556 (*Chætodon sohar* Forskål),*Acronurus* Gronow, Cat. Fishes, 1854, 190 (*Teuthis hepatus* Linnaeus).*Rhombotides* (Klein) Bleeker, Neder. Tijdschr. Dierk., I (*Chætodon triostegus* Linnaeus).

Subgenus HARPURUS Forster.

- 2809. Acanthurus triostegus** (Linnaeus).

South Pacific, from New Zealand and Australia to the Revillagigedos.

Chætodon triostegus Linnaeus, Syst. Nat., Ed. X, 1758, 274, India.

Harpurus fasciatus Forster, Enchiridion, 1778, 84, South Seas.

Acanthurus zebra Lacépède, Hist. Nat. Poiss., III, 1802, 548, fig. 3, no locality.

Chætodon couaga Lacépède, loc. cit. IV, 1802, 727, no locality.

Teuthis australis Gray, in King, Surv. Australia, II, 1826, 435.

Acanthurus hirudo Bennett, Ceylon Fishes, 1851, 11, pl. 11, Ceylon.

Acanthurus subarmatus Bennett, Whaling Voyage, II, 1840, 278, Raiatea, Society Islands.

Subgenus TEUTHIS.

- 2810. Acanthurus cæruleus** Bloch and Schneider. *Barbero; Blue surgeon; Blue tang.*

West Indies, Key West, and Bermuda to Bahia, north to Woods Hole, Mass.

Acanthurus cæruleus Bloch and Schneider, Syst. Ichth., 1801, 214, Carolinas; Habana; Jamaica; after Catsby, Parra, and Browne.

Acanthurus broussonetii Desmarest, Prem. Dec. Ichth., 1823, 26, Cuba.

Acanthurus brevis Poey, Memorias, II, 1861, 207, Antilles and Habana.

Acronurus cæruleatus Poey, Enumeratio, 1875, 69, Cuba; young.

- 2811. Acanthurus hepatus** (Linnaeus). *Common surgeon; Doctor fish; Lancelet fish; Tang; Tangbarbero; Saigneur; Barbero.*

West Indian fauna; Florida to Bahia; Key West and Cuba; Charleston; north to Woods Hole, Mass.

Teuthis hepatus Linnaeus, Syst. Nat., Ed. XII, 1766, 507, Carolinas.

Chætodon chirurgus Bloch, Naturgesch. Ausl. Fische, III, 1787, 99, Martinique; on drawing by Plumier.

Acanthurus phlebotomus Cuvier and Valenciennes, Hist. Nat. Poiss., X, 1835, 129 (176), pl. 287 Martinique, Brazil, Habana, New York.

Acronurus fuscus Gronow, Fish., ed. Gray, 119, 1854, West Indies; young.

Acronurus carneus Poey, Memorias, II, 1860, 207, Cuba.

2812. *Acanthurus bahianus* Castelnau. *Barbero; Ocean tang.*
 West Indies to Brazil.
Acanthurus bahianus Castelnau, Anim. Amer. Sud, Poiss., 1855, 24,
 Pl. XI, fig. 1, Bahia.
Acanthurus tractus Poey, Memorias, II, 1861, 208, Cuba.
Acronurus nigriculus Poey, Enumeratio, 1875, 69, Cuba.
2813. *Acanthurus crestonis* (Jordan and Starks). *Barbero negro.*
 West coast of Mexico and Central America, Mazatlan to Panama.
Teuthis crestonis Jordan and Starks, Proc. Calif. Acad. Sci., ser. 2,
 1895, V, 485, pl. 47, Mazatlan.
2814. *Acanthurus aliala* Lesson. *Philosophe.*
 East Indies to the west coast of Mexico; abundant at Socorro and
 Clarion Islands.
Acanthurus aliala Lesson, Voy. Coquille, Zool., II, 1830, 150, Oualan.
Acanthurus glaucopareius Cuvier and Valenciennes, Hist. Nat. Poiss.,
 X, 1835, 139 (190), Oualan; Molucca.
- Genus 974. *XESURUS* Jordan and Evermann.
Xesurus Jordan and Evermann, Check List, 1895, 421 (*Prionurus*
punctatus Gill).
2815. *Xesurus punctatus* (Gill). *Cochinito.*
 Cape San Lucas; Mazatlan; Creston Island.
Prionurus punctatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 242,
 Cape San Lucas.
2816. *Xesurus clarionis* Gilbert and Starks.
 Galapagos Islands.
Xesurus clarionis Gilbert and Starks, Proc. U. S. Nat. Mus., XIX,
 1896, 437, 445, Clarion Island, Galapagos Archipelago.
- Genus 975. *CTENOCHÆTUS* Gill.
Ctenodon Bonaparte, in Klunzinger, Fische Rothen Meeres, 1884,
 I Theil, 54 (*Acanthurus strigosus* Bennett); name preoccupied.
Ctenochætus Gill, Proc. U. S. Nat. Mus., VII, 1884, 277 (*Acanthurus*
strigosus Bennett).
2817. *Ctenochætus striatus* (Quoy and Gaimard). *Kale.*
 South seas, recorded from Cocos Island.
Acanthurus striatus Quoy and Gaimard, Voyage l'Uranie, Zool., 1824,
 373, pl. 63, fig. 3, Guam.
Acanthurus strigosus Bennett, Zool. Jour., IV, XIII, Art. III, 1828, 41,
 Honolulu.
Acanthurus ctenodon Cuvier and Valenciennes, Hist. Nat. Poiss., X,
 1835, 178 (241), pl. 289, the Carolines and New Guinea.

Order RR. CATAPHRACTI

Family 225. SCORPÆNIIDÆ. Rockfishes and scorpionfishes

- Genus 976. *SEBASTES* Cuvier. *Rosefishes.*
Sebastes Cuvier, Règne Anim., ed. 2, II, 1829, 166 (*Perca norvegica*
Ascanius = *Perca marina* Linnaeus).
Eusebastes Sauvage, Nouv. Archives Mus. Paris, 1878, (2), 1, 1421
 (*Perca norvegica* Ascanius).
2818. *Sebastes marinus* (Linnaeus). *Rosefish; Redfish; Snapper; Hemdurgan;*
Red seafish; Norway haddock.
 North Atlantic, both coasts, south to Faroe Islands; Norway; Maine;
 in deep water off coast of middle New Jersey.
Perca marina Linnaeus, Syst. Nat., Ed. X, 1758, 290, Norway.
Perca norvegica Ascanius, Icones Rev. Nat., II, 1772, 7, Norway.
Holocentrus sanguineus Faber, Natur. der Fische Islands, 1829, 126,
 Iceland.
Sebastes fasciatus Storer, Proc. Bost. Soc. Nat. Hist., V, 1854, 31,
 Provincetown, Mass.
Sebastes regulus Ekstrom, Skand. Fish., Ed. I, 197, plate only.

- 2819. *Sebastes viviparus* Krøyer.**
 North Atlantic, in shallow bays, more southerly not certainly recorded from America.
Sebastes viviparus Krøyer, Naturh. Tidskr., I, 1845, 275, Norway, in shallow water.
- Genus 977. SEASTOLOBUS Gill.**
Sebastolobus Gill, Rept. Smithson. Inst., 1880, 375 (*Sebastes macrochir* Günther).
- 2820. *Sebastolobus altivelis* Gilbert.**
 Aleutian Islands, Alaska, and in deep water off the coast of California.
Sebastolobus altivelis Gilbert, Rept. U. S. Fish Com., XIX, 1893, 410, pl. 23, south of Alaska Peninsula, at *Albatross* station 3338, in 625 fathoms.
- 2821. *Sebastolobus alascanus* Bean.**
 Bering Sea; Pacific coast of Alaska, Washington, Oregon, and California, in deep water; recently taken in San Pedro.
Sebastolobus alascanus Bean, Proc. U. S. Nat. Mus., XIII, 1890, 44, off Trinity Islands, Alaska, at *Albatross* station 2853, 56° N., 154° W., in 159 fathoms.
- Genus 978. EMMELAS Jordan and Evermann.**
Emmelas Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1773 (*Sebastes glaucus* Hilgendorf).
- 2822. *Emmelas glaucus* (Hilgendorf).**
 Hokkaido, north to Bering Island.
Sebastes glaucus Hilgendorf, Sitzber. Ges. Naturf. Freunde, Berlin, 1880, 172, Hokkaido, north Japan; no description given.
- Genus 979. SEASTODES^{**} Gill. Rockfishes.**
Sebastodes Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 165 (*Sebastes paucispinis* Ayres).
- 2823. *Sebastodes paucispinis* (Ayres). *Bocaccio; Mérou; Jack.***
 San Diego to Barclay Sound, British Columbia; common south. A valued food fish.
Sebastes paucispinis Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 6, San Francisco.
- 2824. *Sebastodes jordani* Gilbert.**
 Coast of California in deep water.
Sebastodes jordani Gilbert, Rept. U. S. Fish Com., XIX, 1893, 466, coast of California, at *Albatross* station 2935, 3103, and 3114, in 64 to 124 fathoms.
- 2825. *Sebastodes goodei* Eigenmann and Eigenmann.**
 San Diego to San Francisco; not common.
Sebastodes goodei Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., III, 1890, 12, San Diego, Calif.
- Genus 980. SEASTOSOMUS Gill.**
Sebastosomus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 59 (*Sebastes melanops* Girard).
Primospina Eigenmann and Beeson, Am. Nat., XXVII, 669 (*Sebastichthys mystinus* Jordan and Gilbert).
- Subgenus PRIMOSPINA Eigenmann and Beeson.**
- 2826. *Sebastosomus flavidus* (Ayres). *Yellowtail rockfish.***
 San Diego to San Francisco; common northward.
Sebastodes flavidus Ayres, Proc. Calif. Acad. Nat. Sci., ser. I, II, 1862, 209, fig. 64, San Francisco.

^{**} The 13-spined allies of *Sebastodes* constitute a most striking feature of the fauna of the north Pacific, the species being very numerous on the rocky coast of California, Alaska, and Japan; a few also in Chile. The species are in general well defined, showing little individual variation and no sex changes. It is extremely difficult, however, to divide them into genera, the various traits being intertangled to a remarkable degree. Yet such species as *paucispinis*, *rosaceus*, and *nigroinctus* could hardly stand in one genus in a natural system. It may be noted that the group called *Sebastodes* is the simplest in organization and also the oldest, four well-marked species related to *Sebastodes paucispinis* occurring in the upper Miocene of California.

- 2827. *Sebastosomus serranoides* (Eigenmann and Eigenmann).**
Cortez Banks, off San Diego.
Sebastodes serranoides Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., III, 1890, 36, Cortez Banks.
- 2828. *Sebastosomus melanops* (Girard).**
Pacific coast of America, from Monterey to Sitka.
Sebastodes melanops Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 135, Astoria, Oreg.
Sebastosomus simulans Gill, loc. cit., XVI, 1864, 147, Cape Flattery.
- 2829. *Sebastosomus mystinus* (Jordan and Gilbert).** *Pêche prêtre; Priestfish; Black rockfish.*
Pacific coast of America, from Aleutian Islands to San Diego.
Sebastichthys mystinus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 455, San Francisco.
- 2830. *Sebastosomus ciliatus* (Tilesius).**
Coast of Alaska.
Epinephelus ciliatus Tilesius, Mem. Acad. Sci. St. Petersb., IV, 1810, 474, Aleutian Islands.
Perca variabilis Pallas, Zoogr. Rosso-Asiat., III, 1811 (1831), 241, Aleutian Islands; including *S. aleuticus*.
- Genus 981. *ROSCOLA* Jordan and Evermann.**
Rosicola Jordan and Evermann, Check List, 1895, 429 (*Sebastosomus pinniger* Gill).
- 2831. *Rosicola babcocki* (Thompson).**
Coast of Alaska and British Columbia, common.
Sebastodes babcocki W. F. Thompson, Fisheries British Columbia, 1914, Middleton Island, Alaska.
- 2832. *Rosicola pinniger* (Gill).** *Flioma; Orange rockfish.*
Pacific coast of America, from San Diego to Puget Sound.
Sebastosomus pinniger Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 147, San Francisco.
- 2833. *Rosicola miniatus* (Jordan and Gilbert).** *Rasciera; Rasher.*
Pacific coast of America, from San Francisco to San Diego.
Sebastichthys miniatus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 70, Monterey, Calif.
- 2834. *Rosicola alexandri* (Evermann and Goldsborough).**
Monterey Bay.
Sebastodes alexandri Evermann and Goldsborough, Proc. U. S. Nat. Mus., XXXI, 1907, 651, Monterey Bay in 65 fathoms.
- Genus 982. *ZALOPYR* Jordan and Evermann.**
Zalopyr Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1795 (*Sebastodes aleutianus* Jordan and Evermann).
- 2835. *Zalopyr aleutianus* (Jordan and Evermann).**
Alaska; Aleutian Islands.
Sebastodes aleutianus Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1795, Shelikof Straits, Alaska.
- 2836. *Zalopyr atrorubens* (Gilbert).**
Monterey, Calif.
Sebastodes atrorubens Gilbert, in Jordan and Evermann, Check List, 1895, 429, and Fishes North and Mid. Amer., 1898, 1797, San Francisco.
- 2837. *Zalopyr atrovirens* (Jordan and Gilbert).** *Garrupa.*
Coast of California, from San Diego, to San Francisco.
Sebastichthys atrovirens Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 289, Monterey, Calif.
- Genus 983. *ACUTOMENTUM* " Eigenmann and Beeson.**
Acutomentum Eigenmann and Beeson, Amer. Nat., XXVII, 1893, 669 (*Sebastes ovalis* Ayres).

" As here arranged, this is far from a natural assemblage.

- 2838. *Acutomentum ovale* (Ayres).** *Viwa; Widowfish.*
 Pacific coast of United States from San Diego to San Francisco.
Sebastodes ovalis Ayres, Proc. Calif. Acad. Nat. Sci., II, 1862, 209-212,
 fig. 65, San Francisco.
- 2839. *Acutomentum prorigerum* (Jordan and Gilbert).**
 San Diego to San Francisco.
Sebastichthys proriger Jordan and Gilbert, Proc. U. S. Nat. Mus., III,
 1880, 327, Monterey, Calif.
- 2840. *Acutomentum brevispine* (Bean).**
 Coast of Alaska.
Sebastichthys proriger var. *brevispinis* Bean, Proc. U. S. Nat. Mus.,
 VI, 1883, 359, Hassler Harbor, Alaska.
- 2841. *Acutomentum clavilatum* (Starks).**
 Puget Sound.
Sebastodes clavilatus Starks, Ann. Carnegie Mus., VII, 1911, 181, San
 Juan Islands, Puget Sound.
- 2842. *Acutomentum entomelas* (Jordan and Gilbert).**
 Pacific coast of America, from Port Harford to Monterey Bay.
Sebastichthys entomelas Jordan and Gilbert, Proc. U. S. Nat. Mus.,
 III, 1880, 142, Monterey.
- 2843. *Acutomentum wilsoni* (Gilbert).**
 Monterey Bay.
Sebastodes wilsoni Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 333,
 pl. 16, fig. 8, Monterey Bay.
- 2844. *Acutomentum rufum* (Eigenmann and Eigenmann).**
 Pacific coast of America, near San Diego; Cortez Banks.
Sebastodes rufus Eigenmann and Eigenmann, Proc. Calif. Acad. Sci.,
 2d ser. III, 1890, 13, Point Loma; Cortez Banks.
- 2845. *Acutomentum macdonaldi* Eigenmann and Beeson.**
 San Diego, Calif.
Acutomentum macdonaldi Eigenmann and Beeson, Am. Nat., XXVII,
 1893, 669, San Diego, Calif.
- 2846. *Acutomentum eigenmanni* (Cramer).**
 Coast of California.
Sebastodes eigenmanni Cramer, Proc. Calif. Acad. Sci., VI, 1896, 239,
 Pl. XLII, Monterey, Calif.
- 2847. *Acutomentum hopkinsi* (Cramer).**
 Monterey Bay, Calif.
Sebastodes hopkinsi Cramer, Proc. Calif. Acad. Sci., V, 1895, 594, with
 plate, Monterey Bay, Calif.
- 2848. *Acutomentum alutum* (Gilbert).**
 Pacific coast of America, from Bering Sea to Santa Barbara.
Sebastichthys alutus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 76,
 Santa Barbara Islands, Calif.
- Genus 984. EOSEBASTES Jordan and Evermann.**
Eosebastes Jordan and Evermann, Check List, 1895 430 (*Sebastichthys*
aurora Gilbert).
- 2849. *Eosebastes aurora* (Gilbert).**
 Santa Barbara Islands, in deep water.
Sebastichthys aurora Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 80,
 Santa Barbara Islands, in 266 fathoms.
- 2850. *Eosebastes melanostomus* (Eigenmann and Eigenmann).**
 Coast of California to Alaska.
Sebastodes melanostomus Eigenmann and Eigenmann, Proc. Calif.
 Acad. Sci., 2d ser., III, 1890, 17, Point Loma, near San Diego, Calif.
- 2851. *Eosebastes introniger* (Gilbert).**
 Santa Barbara to Bering Sea.
Sebastichthys introniger Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 81,
 Santa Barbara Islands.

- 2852. *Eosebastes saxicola* (Gilbert).**
 Southern California to southeast Alaska in moderate depth.
Sebastichthys saxicola Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 78,
 Santa Barbara Islands at *Albatross* stations 2893, 2907, and others,
 in 44 to 155 fathoms.
- 2853. *Eosebastes deani* (Starks).**
 Puget Sound.
Sebastodes deani Starks, Ann. Carnegie Mus., VII, 1911, 178, Puget
 Sound.
- 2854. *Eosebastes crameri* (Jordan).**
 Coast of Oregon.
Sebastodes crameri Jordan, Proc. U. S. Nat. Mus., XIX, 1896, 451,
 coast of Oregon, at *Albatross* station 3091.
- 2855. *Eosebastes swifti* (Evermann and Goldsborough).**
 Southern Alaska.
Sebastodes swifti Evermann and Goldsborough, Bull. U. S. Bur. Fish.,
 XXVI, 1906, 285, Yes Bay and Kasaan Bay, Alaska.
- 2856. *Eosebastes semicinctus* (Gilbert).**
 Coast of California.
Sebastodes semicinctus Gilbert, Proc. U. S. Nat. Mus., XIX, 1896, 449,
 Santa Barbara Channel, Calif.
- 2857. *Eosebastes diploproa* (Gilbert).**
 Pacific coast from the Coronado Islands northward to Nanaimo,
 British Columbia.
Sebastichthys diploproa Gilbert, Proc. U. S. Nat. Mus., XIII, 1890,
 79, Coronado Islands at *Albatross* station 2935, in 124 fathoms.
- 2858. *Eosebastes emphæus* (Starks).**
 Puget Sound, in deep water.
Sebastodes emphæus Starks, Ann. Carnegie Mus., VII, 1911, 183, about
 San Juan Islands, Puget Sound.
- Genus 985. SEBASTOMUS Gill.**
Sebastomus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 147 (*Sebastes*
rosaceus Girard).
- 2859. *Sebastomus constellatus* (Jordan and Gilbert).** *Spotted rockfish.*
 Coast of California, from San Diego to San Francisco.
Sebastichthys constellatus Jordan and Gilbert, Proc. U. S. Nat. Mus.,
 III, 1880, 295, Santa Barbara Channel; San Francisco.
- 2860. *Sebastomus rosaceus* (Girard).** *Corsair.*
 Pacific coast from San Diego to Puget Sound.
Sebastes rosaceus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1854, 146,
 San Diego, San Francisco.
Sebastes helvomaculatus Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859,
 26, San Francisco.
- 2861. *Sebastomus rhodochloris* (Jordan and Gilbert).** *Flyfish.*
 Off Monterey and San Francisco.
Sebastichthys rhodochloris Jordan and Gilbert, Proc. U. S. Nat. Mus.,
 III, 1880, 144, Monterey, Calif.
- 2862. *Sebastomus chlorostictus* (Jordan and Gilbert).** *Pesca vermiciglia.*
 San Diego; Monterey; San Francisco.
Sebastichthys chlorostictus Jordan and Gilbert, Proc. U. S. Nat. Mus.,
 III, 1880, 294, Monterey, Calif.
- 2863. *Sebastomus umbrosus* (Jordan and Gilbert).**
 Coast of California, Point Conception to Coronado Islands.
Sebastichthys umbrosus Jordan and Gilbert, Proc. U. S. Nat. Mus., V,
 1882, 410, Santa Barbara.
Sebastodes aereus Eigenmann and Eigenmann, Proc. Calif. Acad. Sci.,
 2d ser., III, 1890, 20, Coronado Islands.
- 2864. *Sebastomus ayresii* (Gilbert and Cramer).**
 Coast of southern California at Cortez Banks, near San Diego.
Sebastodes ayresii Gilbert and Cramer, Proc. U. S. Nat. Mus., XIX,
 1896, 450, Cortez Banks, near San Diego, Calif.

- 2865. *Sebastomus eos* (Eigenmann and Eigenmann).**
 Coast of California at Point Loma, near San Diego.
Sebastodes eos Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., III, 1890, 18, Point Loma, near San Diego, Calif.
- 2866. *Sebastomus gillii* (Eigenmann and Eigenmann).**
 Coast of California at Point Loma, near San Diego.
Sebastodes gillii Eigenmann and Eigenmann, Amer. Nat., XXV, February, 1891, 154, Point Loma, near San Diego, Calif.
- 2867. *Sebastomus rupestris* (Gilbert).**
 Santa Barbara Islands northward to southeast Alaska.
Sebastichthys rupestris Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 76, Santa Barbara Islands, at *Albatross* station 2946, in 150 fathoms.
- Genus 986. SEBASTOPYR⁷⁰ Jordan and Evermann.**
Sebastopyr Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, Apr. 27, 1927, 506 (*Sebastodes ruberrimus* Cramer).
- 2868. *Sebastopyr ruberrimus* (Cramer).** *Red rockfish; Tambor.*
 Pacific coast from San Diego to Puget Sound and southeastern Alaska.
Sebastes ruber of early writers, not of Ayres.
Sebastodes ruberrimus Cramer, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 597, pls. 62, 63, and 69, Monterey, Calif.
- Genus 987. HISPANISCUS Jordan and Evermann.**
Hispanicus Jordan and Evermann, Check List, 1895, 431 (*Sebastichthys rubrivinctus* Jordan and Gilbert).
- 2869. *Hispaniscus rubrivinctus* (Jordan and Gilbert).** *Spanish flag.*
 California coast from San Diego to Monterey; not rare south. A most brilliantly colored fish.
Sebastichthys rubrivinctus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 291, Santa Barbara Channel.
- 2870. *Hispaniscus levis* (Eigenmann and Eigenmann).**
 Coast of California, from San Diego to Monterey.
Sebastichthys levis Eigenmann and Eigenmann, West American Scientist, Nov. 9, 1889, 128, San Diego, Calif.
- 2871. *Hispaniscus zacentrus* (Gilbert).**
 Coast of California.
Sebastichthys zacentrus Gilbert. Proc. U. S. Nat. Mus., XIII, 1890, 77, Santa Barbara Islands, at *Albatross* station 2893 and 2946, in 145 and 150 fathoms.
- 2872. *Hispaniscus elongatus* (Ayres).** *Reina.*
 California coast, from San Diego to San Francisco.
Sebastes elongatus Ayres, Proc. Calif. Acad. Nat. Sci., II, 1858-1862 (Oct. 17, 1859), 26, San Francisco.
- 2873. *Hispaniscus sinensis* (Gilbert).**
 Gulf of California.
Sebastichthys sinensis Gilbert, Proc. U. S. Nat. Mus., XLIII, 1890, 81, Gulf of California, at *Albatross* station 3015 in 145 fathoms.
- Genus 988. AUCTOSPINA Eigenmann and Beeson.**
Auctospina Eigenmann and Beeson, Amer. Nat., XXVII, 1893, 670 (*Sebastes auriculatus* Girard).
- 2874. *Auctospina auriculata* (Girard).** *Brown rockfish.*
 Pacific coast of America, from Cape Mendocino to Cedros Island; common.
Sebastes auriculatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 131, 146, San Francisco.
Sebastes ruber Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854 (Sept. 11), 7, San Francisco.
Sebastes ruber parvus Ayres, loc. cit., I, 1854 (Sept. 11), 7, San Francisco.

⁷⁰ This group is allied to *Sebastomus*, but of much coarser build, the cranial spines in the adult rough with blunt spinules.

- 2875. *Auctospina dallii* (Eigenmann and Beeson).**
 Pacific coast of America, from San Francisco to Puget Sound; perhaps a variation of *A. auriculata*.
Pteropodus dallii Eigenmann and Beeson, Amer. Nat., XXVIII, January, 1894, 66, San Francisco.
- Genus 989. PTEROPODUS Eigenmann and Beeson.**
Pteropodus Eigenmann and Beeson, Amer. Nat., XXVII, 1893, 670 (*Sebastichthys maliger* Jordan and Gilbert); not *Pteropus*, a genus of birds.
- 2876. *Pteropodus rastrelliger* (Jordan and Gilbert).** *Grass rockfish.*
 California coast, from San Diego to San Francisco.
Sebastichthys rastrelliger Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 296, Monterey, Calif.
- 2877. *Pteropodus caurinus* (Richardson).**
 Pacific coast of America from Puget Sound to Sitka.
Sebastes caurinus Richardson, Voy. *Sulphur*, Ichthyology, 1845, 77, pl. 41, fig. 1, Sitka.
- 2878. *Pteropodus vexillaris* (Jordan and Gilbert).**
 California coast, from San Diego to Cape Mendocino, replacing *caurinus* southward.
Sebastichthys vexillaris Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 292, Santa Barbara Channel; San Francisco.
- 2879. *Pteropodus maliger* (Jordan and Gilbert).** *Yellow-backed rock cod;*
Brown rock cod.
 Pacific coast of America, from Monterey to Sitka; not rare.
Sebastichthys maliger Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 322, San Francisco.
- 2880. *Pteropodus gilberti* (Cramer).**
 Coast of California.
Sebastodes gilberti Cramer, Proc. Calif. Acad. Sci., VI, 1896, 241, Pl. XLIII, San Francisco.
- 2881. *Pteropodus carnatus* (Jordan and Gilbert).** *Flesh-colored rock cod.*
 California coast, from San Diego to Point Reyes, replacing *P. chrysomelas* in shallow water.
Sebastichthys carnatus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 73, Monterey, Calif.
- 2882. *Pteropodus chrysomelas* (Jordan and Gilbert).** *Black and yellow rock cod.*
 Pacific coast of America, from Puget Sound to San Diego; common in moderate depths.
Sebastichthys chrysomelas Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 455, 465, Monterey, Calif.
- 2883. *Pteropodus nebulosus* (Ayres).** *Yellow-spotted rock cod.*
 Pacific coast; southeastern Alaska; Puget Sound region, Vancouver to Point Conception; common.
Sebastes nebulosus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854 (Sept. 11), 5, San Francisco.
Sebastes fasciatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 146, Presidio, California; not of Storer.
Sebastichthys fasciolaris Lockington, Proc. U. S. Nat. Mus., III, 1880, 297, San Francisco.
- Genus 990. SEBASTOCARUS¹¹ Jordan and Evermann.**
Sebastocarbus Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, Apr. 27, 1927, 507 (*Sebastichthys serriceps* Jordan and Gilbert).
- 2884. *Sebastocarbus serriceps* (Jordan and Gilbert).** *Treefish.*
 Coast of California, from Point Reyes to Cedros Island; common south.
Sebastichthys serriceps Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 38, Santa Barbara and Santa Catalina Islands, Calif.

¹¹ This genus is nearly allied to *Sebastichthys* differing in the more compressed body and especially in the strict and high ridges on the head, all of them free from serrations or accessory tubercles.

Genus 991. **SEBASTICHTHYS** Gill.

Sebastichthys Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 278, 329
(*Sebastes nigrocinctus* Ayres).

2885. **Sebastichthys nigrocinctus** (Ayres). *Black-banded rock cod.*

Monterey Bay to Naha Bay, Alaska.

Sebastes nigrocinctus Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 25,
fig. 6, San Francisco.

Genus 992. **METZELAARIA** Jordan.

Metzelaaria Jordan, Class. Fishes, 1923, 209 (*Scorpæna tridecimspinosa*
Metzelaar).

2886. **Metzelaaria tridecimspinosa** (Metzelaar).

Caribbean Sea.

Scorpæna tridecimspinosa Metzelaar, Tropisch Atlantische Visschen,
1919, 146, fig. 44, Bonaire, Aruba.

Genus 993. **SCORPÆNODES** Bleeker.

Scorpænodes Bleeker, Nat. Hist. Ned. Ind., XIII, 1857, 371 (*Scorpæna*
polylepis Bleeker).

Sebastopsis Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 278 (*Scorpæna*
polylepis Bleeker).

Sebastodes Sauvage, Ann. Sci. Nat. Zool., V, 1873, 1 (*Sebastes minutus*
Cuvier); not *Sebastodes* Gill, 1861.

2887. **Scorpænodes xyris** (Jordan and Gilbert).

West coast of Mexico.

Sebastopsis xyris Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882,
369, Cape San Lucas, Lower California.

Genus 994. **HELICOLENUS** Goode and Bean.

Helicolenus Goode and Bean, Oceanic Ichth., 1896, 248 (*Scorpæna*
dactylopterus De la Roche).

2888. **Helicolenus dactylopterus** (De la Roche). *Seran imperial; Fanegal;*
Cardonniera; Scorfaniudi; Funal; Crabra; Redfish.

Deep waters of Atlantic off both coasts from Narragansett and Chesapeake Bays, and to coasts of Italy and Spain.

Scorpæna dactyloptera De la Roche, Ann. Mus., XIII, 1809, pl. 22,
fig. 2, Ivica, Barcelona.

Sebastes imperialis Cuvier and Valenciennes, Hist. Nat. Poiss., IV,
1829, 246 (336), Nice.

Helicolenus maderensis Goode and Bean, Oceanic Ichth., 1895, 250,
Madeira, and in the Gulf Stream.

Genus 995. **SCORPÆNA** (Artedi) Linnæus.

Scorpæna (Artedi) Linnæus, Syst. Nat., ed. X, 1758 266 (*S. porcus*
Linnæus).

Parascorpæna Bleeker, Versl. Ak. Amsterdam., IX, pt. 3, 1876, 295
(*Scorpæna picta* Kuhl and van Hasselt in Cuvier and Valenciennes).

Pseudosebastes Sauvage, Nouv. Arch. Mus. Hist. Nat. Paris, II, 1878
(*Sebastes bougainvillae* Cuvier and Valenciennes).

2889. **Scorpæna atlantica** Nichols and Breder.

Coast of Texas.

Scorpæna russula atlantica Nichols and Breder, Proc. Biol. Soc. Wash.,
XXXVII, 1924 (Feb. 21), 21, Pl. VII, off Galveston, Tex.

2890. **Scorpæna colesi** Nichols.

Coast of North Carolina.

Scorpæna colesi Nichols, Bull. Amer. Mus. Nat. Hist., XXXIII, 1914,
537, fig. 1, Cape Lookout, N. C.

2891. **Scorpæna plumieri** Bloch. *Rescacio; Poison grouper; Plumier's*
pigfoot.

West Indies and Brazil, north to Florida; rarely northward to Woods Hole.

Scorpæna plumieri Bloch, Nya. Handl. Stockh., X, 1789, 234, Martinique.

Scorpæna bufo Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829,
224 (306), Martinique.

Scorpæna rascacio Poey, Memorias, II, 1861, 169, Cuba.

Apistes exul Gosse, Nat. Sojourn Jamaica, 1851, 207, Jamaica.

2892. *Scorpæna mystes* Jordan and Starks. *Lapon.*
Coast of Mexico.
Scorpæna mystes Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser. V, 1895, 491, pl. 52, Mazatlan, Mexico.
2893. *Scorpæna grandicornis* Cuvier and Valenciennes. *Lionfish; Poison grouper; Sculpin.*
Florida Keys to Brazil, in the Gulf Stream to Woods Hole, Mass.
Scorpæna grandicornis Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 227 (309), pl. 86, Martinique, Porto Rico, Habana, San Domingo.
2894. *Scorpæna brasiliensis* Cuvier and Valenciennes.
Atlantic coast, from Charleston to Rio Janeiro.
Scorpæna brasiliensis Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 224 (305), Brazil.
Scorpæna stearnsi Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 421, Pensacola, Fla.
2895. *Scorpæna inermis* Cuvier and Valenciennes.
West Indies, north to Florida.
Scorpæna inermis Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 228 (311), Martinique.
Scorpæna occipitalis Poey, Memorias, II, 1861, 171, Cuba.
Scorpæna calcarata Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 422, Clearwater Harbor, Fla.
2896. *Scorpæna sonoræ* Jenkins and Evermann.
Guaymas, Gulf of California, and Santa Margarita Island.
Scorpæna sonoræ Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 150, Guaymas, Sonora.
2897. *Scorpæna guttata* Girard. *Scorpene; Scorpion; Sculpin.*
Coast of California, from Monterey southward to Cedros Island.
Scorpæna guttata Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 145, Monterey.
2898. *Scorpæna russula* Jordan and Bollman.
Pacific coast of Colombia.
Scorpæna russula Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 165, Pacific coast of Colombia.
2899. *Scorpæna histrio* Jenyns.
Pacific coast of America, Panama to Juan Fernandez; Galapagos Islands.
Scorpæna histrio Jenyns, Voy. *Beagle*, Fishes, 1842, 35, pl. 8, Chatham Island, Galapagos Archipelago.
2900. *Scorpæna pannosa* Cramer.
Panama.
Scorpæna pannosa Cramer, Proc. U. S. Nat. Mus., XIX, 1896, 446, with plate, Panama.
2901. *Scorpæna albofasciata* Metzelaar.
Caribbean Sea.
Scorpæna albofasciata Metzelaar, Tropisch Atlantische Vissch., 1919, 145, fig. 43; Bonaire, Arubi.
2902. *Scorpæna albifimbria* Evermann and Marsh.
Porto Rico.
Scorpæna albifimbria Evermann and Marsh, Bull. U. S. Fish Com., XX, 1900, 275, fig. 82, Culebra Island; Porto Rico.
2903. *Scorpæna nuttingi* Evermann and Seale.
West Indies (Barbados).
Scorpæna nuttingi Evermann and Seale, University of Iowa Studies Nat. Hist., X, 1924, 14, 39, Carlisle Bay, Barbados.
2904. *Scorpæna agassizii* Goode and Bean.
Mid-ocean east of Cuba.
Scorpæna agassizii Goode and Bean, Oceanic Ichth., 1896, 247, fig. 243, latitude 32° 13' N., longitude 39° 10' W.

- 2905. *Scorpæna cristulata* Goode and Bean.**
 Atlantic Ocean, off coast of Georgia; perhaps also from northern Spain.
Scorpæna cristulata Goode and Bean, Oceanic Ichth., 1896, 247, fig. 242,
 off Georgia.
Scorpæna echinata Koehler, Ann. Univ. Lyons, XXVI, 1896, 474, pl. 27,
 figs. 4 and 6, Bay of Biscay, in 700 fathoms.
- Genus 996. MERINTHE Snyder.**
Merinthe Snyder, Bull. U. S. Fish Com., XXII, 1902, 536 (*Sebastes*
macrocephalus Sauvage).
- 2906. *Merinthe nematophthalma* (Günther).**
 West Indies, exact locality uncertain.
Sebastes nematophthalmus Günther, Cat., II, 1861, 99, West Indies.
- Genus 997. PONTINUS Poey.**
Pontinus Poey, Memorias, II, 1861, 172 (*P. castor* Poey).
- 2907. *Pontinus castor* Poey.**
 Cuba.
Pontinus castor Poey, Memorias, II, 1861, 173, Habana, Cuba.
- 2908. *Pontinus furcirhinus* Garman.**
 Eastern Pacific in deep water.
Pontinus furcirhinus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
 51, Pl. VII, numerous Albatross stations off Panama.
- 2909. *Pontinus pollux* Poey.**
 Cuba.
Pontinus pollux Poey, Memorias, II, 1861, 174, Habana, Cuba.
- 2910. *Pontinus rathbuni* Goode and Bean.**
 Coast of North Carolina, probably in the Gulf Stream.
Pontinus rathbuni Goode and Bean, Oceanic Ichth., 1895, 255, fig. 245,
 off Cape Hatteras, N. C.
- 2911. *Pontinus macrolepis* Goode and Bean.**
 Yucatan.
Pontinus macrolepis Goode and Bean, Oceanic Ichth., 1895, 257, fig. 247,
 off Yucatan.
- 2912. *Pontinus longispinis* Goode and Bean.**
 Coast of western Florida.
Pontinus longispinis Goode and Bean, Oceanic Ichth., 1895, 258, fig. 246,
 west coast of Florida, lat. $28^{\circ} 36' N.$, long. $85^{\circ} 33' 30'' W.$, in 111
 fathoms.
- 2913. *Pontinus beanorum* Evermann and Marsh.**
 Porto Rico.
Pontinus beanorum Evermann and Marsh, Bull. U. S. Fish Com., XX,
 pt. 1, 1900, 279, fig. 85, San Juan Harbor, Porto Rico.
- 2914. *Pontinus sierra* (Gilbert).**
 Gulf of California.
Scorpæna (Sebastoplus) sierra Gilbert, Proc. U. S. Nat. Mus., XIII,
 1890, 82, Gulf of California, at Albatross stations 2996 and 3011 in
 112 and 71 fathoms.
- 2915. *Pontinus bergii* (Evermann and Marsh).**
 Porto Rico.
Scorpæna bergii Evermann and Marsh, Bull. U. S. Fish Com., XX, pt. 1,
 1900, 276, fig. 83, Mayaguez, Culebra; perhaps not a *Pontinus*.
- Genus 998. SETARCHEΣ Johnson.**
Setarches Johnson, Proc. Zool. Soc. Lond., 1862, 177 (*S. guntheri*
 Johnson).
- 2916. *Setarches parmatus* Goode.**
 Coasts of Rhode Island, North Carolina, and western Florida, in deep
 water.
Setarches parmatus Goode, Proc. U. S. Nat. Mus., III, 1880, 480, off
 coast of Rhode Island.

Family 226. ANOPOLOPOMIDÆ. Skilfishes; Sablefishes

Genus 999. ANOPOLOPOMA Ayres.

Anoplopoma Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 27 (*A. merlangus* Ayres = *Gadus fimbria* Pallas).

Scombrococtonus Peters, Monatsber. Akad. Wiss. Berlin, 1872, 569 (*S. salmoneus* Peters = *Gadus fimbria* Pallas).

2917. **Anoplopoma fimbria** (Pallas). *Beshow; Calfish; Skil; Black cod; Sable-fish.*

Unalaska to San Diego, common northward; a valued food fish in Alaska when it becomes fat.

Gadus fimbria Pallas, Zoogr. Rosso-Asiat., III, 1811, 200, probably Aleutian Islands.

Anoplopoma merlangus Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 27, San Francisco market.

Scombrococtonus salmoneus Peters, Monatsber. Akad. Wiss. Berlin, 1872, 569, Vancouver Island.

Family 227. ERILEPIDIDÆ. Fat priests

Genus 1000. ERILEPIS Gill.

Myriolepis Lockington, Proc. U. S. Nat. Mus., III, 1880, 248 (*Myriolepis zonifer* Lockington); name preoccupied in fossil fishes.

Erilepis Gill, Science, Jan. 26, 1894, 54 (*Myriolepis zonifer* Lockington). *Ebisus* Jordan and Snyder, Jour. Coll. Sci. Tokyo, XV, 1901, 308 (*E. sagamius* Jordan and Snyder).

2918. **Erilepis zonifer** (Lockington). *Aburabozu; Fat priest.*

Coast of Japan to Alaska; once taken at Monterey Bay, Calif.

Myriolepis zonifer Lockington, Proc. U. S. Nat. Mus., III, 1880, 248, Monterey.

Ebisus sagamius Jordan and Snyder, Jour. Coll. Sci., Tokyo, XV, 1901, 308, Misaki, Japan.

Family 228. HEXAGRAMMIDÆ. Greenlings

Genus 1001. HEXAGRAMMOS Steller.

Hexagrammos Steller, in Tilesius, Act. Ac. Petrop., II, 1810, 335 (*H. stelleri* Tilesius = *H. asper* Tilesius).

Labrax Pallas, Mém. Acad. Sci. Petersb., II, 1810, 382 (*L. lagocephalus* Pallas), not of Klein nor of Cuvier.

Lebus (Steller MS.) Pallas, Zoog. Rosso-Asiat., III, 1811, 279 (*Labrax superciliatus* Pallas).

Chirus Steller, in Pallas, loc. cit., 279 (*Labrax superciliatus* Pallas).

Octogrammus Bleeker, Versl. Akad. Amsterdam (2), VIII, 1874, 370 ("*Labrax octogrammus* Schlegel" = *Octogrammus pallasi* Bleeker).

Acantholebius Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 166 (*Chiropterus nebulosus* Girard = *Hexagrammos stelleri* Tilesius); specimen with broken dorsal spines.

Grammatopleurus Gill, loc. cit., XIII, 1861, 166 (*Labrax lagocephalus*. Pallas).

2919. **Hexagrammos decagrammus** (Pallas). *Rock trout; Boregal; Bodieron* North Pacific, from Point Conception to southern Alaska; common south.

Labrax decagrammus Pallas, Mém. Acad. Petersb., II, 1810, 386, pl. 22, fig. 2, Elias Bay.

Chirus guttatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 132, San Francisco.

Gryastes lineatus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 9, Bay of San Francisco.

Chiropsis constellatus Girard, U. S. Pac. R. R. Surv., X, Pt. 4, Fishes, 1858, 42, San Francisco.

Chirus maculo-serialis Lockington, Proc. U. S. Nat. Mus., III, 1880, 55, San Francisco.

- 2920.** *Hexagrammos octogrammus* (Pallas). *Alaska greenfish.*
 Aleutian Islands, westward to Petropaulski and Robben Island.
Labrax octogrammus Pallas, Zoogr. Rosso-Asiat., III, 1811, 283, Kamchatka and Aleutian Islands; doubtful species.
Chirus ordinatus Cope, Proc. Amer. Phil. Soc., XIII, 1873, 28, Unalaska.
Octogrammus pallasi Bleeker, Versl. Akad. Amsterdām, VI, 1870; after Pallas.
- 2921.** *Hexagrammos stelleri* Tilesius. *Greenling.*
 Kamchatka and Unalaska to San Francisco; abundant in Puget Sound and northward.
Hexagrammos stelleri Tilcsius, Mém. Acad. St. Petersb., II, 1809, 335, Kamchatka.
Hexagrammos asper Steller in Tilesius, loc. cit., 340, pl. 15, 1810, Kamchatka.
Labrus hexagrammus Pallas, loc. cit., 395, pl. 23, fig. 3, 1810, Kamchatka.
Chiropsis nebulosus Girard, U. S. Pac. R. R. Surv., X, Fishes, 1858, 45, Puget Sound and at Fort Steilacoom.
Chirus trigrammus Cope, Proc. Amer. Phil. Soc., XIII, 1873, 29, Unalaska.
- 2922.** *Hexagrammos superciliatus* (Pallas). *Red rock trout.*
 Bering Sea to Monterey Bay.
Labrax superciliatus Pallas, Mém. Acad. St. Petersb., II, 1810, 388, Unalaska.
Chirus pictus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 132, San Francisco; Humboldt Bay.
Chirus balias Cope, Proc. Amer. Phil. Soc., XIII, 1873, 28, Captains Harbor, Unalaska.
Hexagrammus scaber Bean, Proc. U. S. Nat. Mus., IV, 1881, 154, Amchitka, Unalaska.
- 2923.** *Hexagrammos lagocephalus* (Pallas).
 Western Bering Sea, Kuril Islands; Alaska; Attu.
Labrax lagocephalus Pallas, Mém. Acad. St. Petersb., II, 1810, 384, Kuril Islands.
- Genus 1002. **PLEUROGRAMMUS** Gill. *Atka mackerel.*
Pleurogrammus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 166 (*Labrax monopterygius* Pallas).
- 2924.** *Pleurogrammus monopterygius* (Pallas). *Atka mackerel; Atka fish; Yellowfish.*
 Northern Pacific, common about Atka and other Aleutian Islands east to Unalaska.
Labrax monopterygius Pallas, Mém. Acad. Sci. St. Petersburg, II, 1810, 391, Unalaska.

Family 229. OPHIODONTIDÆ. Cultus cods

Genus 1003. **OPHIODON** Girard.
Ophiodon Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 133 (*Ophiodon elongatus* Girard).
Oplopoma Girard, loc. cit., VIII, 1856, 135 (*Oplopoma pantherina* Girard = *Ophiodon elongatus* Girard; young).

- 2925.** *Ophiodon elongatus* Girard. *Cultus cod; Blue cod; Buffalo cod; Ling.*
 Pacific coast from Alaska to Santa Barbara.
Ophiodon elongatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 133, San Francisco.
Oplopoma pantherina Girard, loc. cit., VIII, 1856, 135, Cape Flattery, Wash.

Family 230. OXYLEBIIDÆ

Genus 1004. **OXYLEBIUS** Gill.
Oxylebius Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 277 (*O. pictus* Gill).

- 2926.** *Oxylebius pictus* Gill.
 From Monterey northward to Puget Sound; not rare.
Oxylebius pictus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 278, California; no definite locality given.

Family 231. ZANIOLEPIDÆ

Genus 1005. ZANIOLEPIS Girard.

Zaniolepis Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 202 (*Z. latipinnis* Girard).

2927. *Zaniolepis latipinnis* Girard.

Coast of California, from San Francisco northward.

Zaniolepis latipinnis Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 202, Fort Steilacoom, Puget Sound.

Genus 1006. XANTOCLES Jordan.

Xantocles Jordan, Copeia, Oct. 4, 1917, 88 (*Zaniolepis frenatus* Eigenmann).

2928. *Xantocles frenatus* (Eigenmann).

Shore banks of southern California; rare.

Zaniolepis frenatus Eigenmann, West American Scientist, VI, Nov. 9, 1889, 147, Cortez Banks off San Diego.

Family 232. JORDANIIDÆ

Genus 1007. JORDANIA Starks.

Jordania Starks, Proc. Acad. Nat. Sci. Phila., XLVII, 1895, 410 (*J. zonope* Starks).

2929. *Jordania zonope* Starks.

Puget Sound.

Jordania zonope Starks, Proc. Acad. Nat. Sci. Phila., XLVII, 1895, 410, Port Orchard, Puget Sound.

Genus 1008. PARICELINUS Eigenmann and Eigenmann.

Paricelinus Eigenmann and Eigenmann, West Amer. Sci., VI, 1889, 131 (*P. hopliticus* Eigenmann and Eigenmann).

2930. *Paricelinus hopliticus* Eigenmann and Eigenmann.

Cortez Banks, off San Diego.

Paricelinus hopliticus Eigenmann and Eigenmann, West Amer. Sci., VI, Nov. 9, 1889, 131, Cortez Banks.

Genus 1009. ALCIDEA Jordan and Evermann.

Alcidea Jordan and Evermann, Fishes North and Mid. Amer., 1886, 1898 (*Paricelinus thoburni* Gilbert).

2931. *Alcidea thoburni* (Gilbert).

Coast of Oregon.

Paricelinus thoburni Gilbert, Rept. U. S. Fish Com., XIX, 1893, 432, pl. 30, coast of Oregon, in 75 fathoms.

Family 233. ICELIDÆ. Scaled sculpins

Genus 1010. CHITONOTUS Lockington.

Chitonotus Lockington, Proc. U. S. Nat. Mus., IV, 1881, 141 (*C. megacephalus* Lockington = *Artedius pugetensis* Steindachner).

2932. *Chitonotus pugetensis* (Steindachner).

San Francisco to Puget Sound.

Artedius pugetensis Steindachner, Ichth. Beitr., V, 1876, 133, pl. 14, fig. 2, Puget Sound.

Chitonotus megacephalus Lockington, Proc. U. S. Nat. Mus., IV, 1881, 142, off San Francisco.

Genus 1011. TARANDICHTHYS Jordan and Evermann.

Tarandichthys Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 225 (*Icelinus filamentosus* Gilbert).

2933. *Tarandichthys cavifrons* (Gilbert).

Coast of southern California.

Icelinus cavifrons Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 83, off Santa Barbara Islands, at Albatross stations 2907 and 2945, in 44 and 30 fathoms.

- 2934. *Tarandichthys filamentosus* (Gilbert).**
 Coast of southern California; Alaska, off St. Mary's Mission; Hood Canal, Puget Sound.
Icelinus filamentosus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 85, off Santa Barbara Islands, in 145 and 55 fathoms.
- 2935. *Tarandichthys tenuis* (Gilbert).**
 Coast of southern California, Alaska, Fort Rupert and Gulf of Georgia.
Icelinus tenuis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 86, off Santa Barbara Islands, at *Albatross* stations 2893, 2946, and others, in 45 to 150 fathoms.
- Genus 1012. *ICELINUS* Jordan.**
Icelinus Jordan, Cat. Fish. N. Amer., 1885, 110 (*Artediush quadriseriatus* Lockington).
- 2936. *Icelinus fimbriatus* Gilbert.**
 Southern California.
Icelinus fimbriatus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 87, off Santa Barbara Islands, at *Albatross* stations 2893 and 2975, in 145 and 36 fathoms.
- 2937. *Icelinus oculatus* Gilbert.**
 Coast of southern California.
Icelinus oculatus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 87, off Santa Barbara Islands, at *Albatross* station 2935, in 124 fathoms.
- 2938. *Icelinus burchami* Evermann and Goldsborough.**
 Loring and Behm Canal, Alaska.
Icelinus burchami Evermann and Goldsborough, Bull. U. S. Bur. Fish., XXVI, 1906, 297, fig. 48, off Loring, Alaska.
- 2939. *Icelinus borealis* Gilbert.**
 Coast of Alaska, north and south of Aleutian Islands and in Bristol Bay.
Icelinus borealis Gilbert, Rept. U. S. Fish Com., XIX, 1893, 415, pl. 25, Aleutian Islands and Bristol Bay at *Albatross* stations 3213, 3214, and others, in 11 to 121 fathoms.
- 2940. *Icelinus quadriseriatus* (Lockington).**
 Off San Francisco, between Point Reyes and Golden Gate.
Artediush quadriseriatus Lockington, Proc. U. S. Nat. Mus., II, 1879, 330, off San Francisco.
- 2941. *Icelinus strabo* Starks.**
 Puget Sound at Port Ludlow, Wash.
Icelinus strabo Starks, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 551, Port Ludlow, Wash.
- 2942. *Icelinus fuscescens* Gilbert.**
 Southern California, Santa Barbara Islands to Monterey.
Icelinus fuscescens Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 340, pl. 17, fig. 9, Santa Barbara Islands, depth 260 to 310 fathoms.
- Genus 1013. *ASTROLYTES* Jordan and Starks.**
Astrolytes Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 807 (*Artediush fenestralis*, Jordan and Gilbert).
- 2943. *Astrolytes fenestralis* (Jordan and Gilbert).**
 Puget Sound to Unalaska.
Artediush fenestralis Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 577, Puget Sound.
Artediush asperulus Starks, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 553, Port Ludlow, Wash.
- Genus 1014. *PARASTROLYTES* Hubbs.**
Parastrolytes Hubbs, Occ. Papers, Mus. Zool. Univ. Mich., No. 170, Feb. 20, 1926, 2 (*Artediush notospilotus* Girard).
- 2944. *Parastrolytes notospilotus* (Girard).**
 Coast of California from Straits of Fuca southward; abundant at Santa Barbara and Puget Sound.
Artediush notospilotus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 134, Tomales Bay, Calif.

Genus 1015. AXYRIAS Starks.

Axyrius Starks, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 554 (*Axyrius harringtoni* Starks).

2945. *Axyrius harringtoni* Starks.

Puget Sound in the vicinity of Port Ludlow, Wash.

Axyrius harringtoni Starks, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 554, pl. 74, Port Ludlow, Wash.

Genus 1016. ARTEDIUS Girard.

Artedius Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 134 (*Scorpænichthys lateralis* Girard).

2946. *Artedius lateralis* (Girard).

Pacific coast of North America, from Puget Sound to Port Harford, Calif.

Scorpænichthys lateralis Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 145, San Luis Obispo (Port Harford), Calif.

Genus 1017. PARARTEDIUS Hubbs.

Parartedius Hubbs, Occ. Papers, Mus. Zool. Univ. Mich., No. 170, Feb. 20, 1926, 3 (*P. hankinsoni* Hubbs).

2947. *Parartedius hankinsoni* Hubbs.

Tidepool, near end of Point Loma, California.

Parartedius hankinsoni Hubbs, Occ. Papers, Mus. Zool. Univ. Mich., No. 171, Feb. 20, 1926, 4, Tidepool near end of Point Loma, Calif.

Genus 1018. ALLARTEDIUS Hubbs.

Allartedius Hubbs, Occ. Papers, Mus. Zool. Univ. Mich., No. 170, Feb. 20, 1926, 8 (*A. corallinus* Hubbs).

2948. *Allartedius corallinus* Hubbs.

Tidepool on Point Lobos, Monterey Co., Calif. Only one specimen known.

Allartedius corallinus Hubbs, Occ. Papers, Mus. Zool. Univ. Mich., No. 170, Feb. 20, 1926, 8, deep low pool on Point Lobos, Monterey Co., Calif.

Genus 1019. ARCHAULUS Gilbert and Burke.

Archaulus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 36 (*A. biseriatus* Gilbert and Burke).

2949. *Archaulus biseriatus* Gilbert and Burke.

West Bering Sea.

Archaulus biseriatus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 36, Petrel Bank, Bering Sea.

Genus 1020. ARTEDIELLUS Jordan.

Artediellus Jordan, Cat. Fish. N. Amer., 1885, 110 (*Cottus uncinatus* Reinhardt).

2950. *Artediellus uncinatus* (Reinhardt).

Greenland to Labrador.

Cottus uncinatus Reinhardt, K. Dansk. Vid. Selsk. Natur. Math. Afh., VI, 1833; XLIV, 1837, Greenland.

Artediellus atlanticus Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1906, Massachusetts Bay.

2951. *Artediellus pacificus* Gilbert.

Coast of Alaska; in Bristol Bay, south of Sannak Island, and north of Unalaska.

Artediellus pacificus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 416, Unalaska, Sannak, Bristol Bay, at Albatross station 3216, south of Sannak Island, many other stations in Bristol Bay, and station 3323 north of Unalaska, in 8 to 61 fathoms.

2952. *Artediellus clarkii* (Evermann and Goldsborough).

Bering Sea.

Blennicottus clarkii Evermann and Goldsborough, Bull. U. S. Fish Com., XXVI, 1906, 323, fig. 85; Bering Sea. Young, perhaps of *Artediellus pacificus*.

- 2953. Artediellus miacanthus** Gilbert and Burke.
 Western Bering Sea.
Artediellus miacanthus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 47, fig. 7, east coast of Kamchatka.
- 2954. Artediellus camchaticus** Gilbert and Burke.
 Western Bering Sea.
Artediellus camchaticus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 46, fig. 6, Avatcha Bay, east coast of Kamchatka.
- Genus 1021. RUSCARIUS** Jordan and Starks.
Ruscarius Jordan and Starks, Proc. Calif. Acad. Sci., 2d series, V, 1895, 805 (*Ruscarius meanyi* Jordan and Starks).
- 2955. Ruscarius meanyi** Jordan and Starks.
 Puget Sound, at Port Orchard near Seattle.
Ruscarius meanyi Jordan and Starks, Proc. Calif. Acad. Sci., 2d series, V, 1895, 805, pl. 80, Port Orchard, Puget Sound.
- Genus 1022. RUSCARIOPS** Hubbs.
Ruscariops Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 170, Feb. 20, 1926, 12 (*R. creaseri* Hubbs).
- 2956. Ruscariops creaseri** Hubbs.
 Low rock pool at Bird Rock, San Diego County, Calif.
Ruscariops creaseri Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 170, Feb. 20, 1926, 12, very low rock pool at Bird Rock, San Diego Co., Calif.
- Genus 1023. RASTRINUS** Jordan and Evermann.
Rastrinus Jordan and Evermann, Check List, 1895, 437 (*Icelus scutiger* Bean).
- 2957. Rastrinus scutiger** (Bean).
 South of the Alaskan Peninsula.
Icelus scutiger Bean, Proc. U. S. Nat. Mus., XIII, 1890, 41, Trinity Islands, Alaska, at *Albatross* station 2853, in 159 fathoms.
- Genus 1024. ICELUS** Krøyer.
Icelus Krøyer, Natur. Hist. Tidsskr., I, 1845, 253 (*Icelus hamatus* Krøyer= *Cottus bicornis* Reinhardt).
- 2958. Icelus bicornis** (Reinhardt).
 Arctic seas, circumpolar; Spitzbergen to northern Russia, Finland, Alaska, Greenland, Labrador, and Cape Cod.
Cottus bicornis Reinhardt, Vid. Selsk. Natur. og Math. Afh., VIII, 1841, LXXV, Greenland.
Icelus hamatus Krøyer, Nat. Hist. Tidsskr., I, 1845, 253, Belsund in Spitzbergen.
Icelus furciger Malm, Forhandl. Skand. Naturf., 1865, 9 Mote, 410, Bohuslan, Sweden.
- 2959. Icelus spiniger** Gilbert.
 Coast of Alaska in the vicinity of Unalaska Island and in Bristol Bay.
Icelus spiniger Gilbert, Rept. U. S. Fish Com., XIX, 1893, 412, pl. 24, fig. 1, Bristol Bay and Unalaska, at *Albatross* stations 3216, 3223, and many others, in 17 to 121 fathoms.
- 2960. Icelus euryops** Bean.
 Bering Sea, off Trinity Islands, Alaska.
Icelus euryops Bean, Proc. U. S. Nat. Mus., XIII, 1890, 41, off Trinity Islands, at *Albatross* station 2853, in 159 fathoms.
- 2961. Icelus vicinalis** Gilbert.
 Bristol Bay, Alaska.
Icelus vicinalis Gilbert, Rept. U. S. Fish Com., XIX, 1893, 413, Bristol Bay at *Albatross* stations 3324, 3330, 3331, and 3332, in 109 to 406 fathoms.
- 2962. Icelus canaliculatus** Gilbert.
 Bering Sea north of Unalaska.
Icelus canaliculatus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 412, pl. 24, fig. 2, off Unalaska, at *Albatross* station 3329, in 399 fathoms.

2963. *Icelus uncinalis* Gilbert and Burke.
 Coast Bering Sea, off Commander Islands.
Icelus uncinalis Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910,
 39, fig. 2, Petrel Bank, near Commander Islands.
2964. *Icelus spatula* Gilbert and Burke.
 East coast of Kamchatka.
Icelus spatula Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910,
 41, fig. 3, Avatcha Bay.
2965. *Icelus australis* (Eigenmann and Eigenmann).
 Cortez Banks off San Diego.
Icelinus australis Eigenmann and Eigenmann, West Amer. Sci., 1889,
 131, Cortez Banks, Calif.
- Genus 1025. **STELGIDONOTUS** Gilbert and Thompson.
Stelgidonotus Gilbert and Thompson, Proc. U. S. Nat. Mus., XXVIII,
 1905, 977 (*S. latifrons* Gilbert and Thompson).
2966. *Stelgidonotus latifrons* Gilbert and Thompson.
 Puget Sound region, Admiralty Head, Friday Harbor.
Stelgidonotus latifrons Gilbert and Thompson, Proc. U. S. Nat. Mus.,
 XXVIII, 1905, 977, Friday Harbor, Wash.
- Genus 1026. **RADULINUS** Gilbert.
Radulinus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 88 (*R. asprellus*
 Gilbert).
2967. *Radulinus boleoides* Gilbert.
 Off Santa Catalina Island in shallow water.
Radulinus boleoides Gilbert, Report U. S. Fish Com., XXIV, 1898, 26,
 pl. 1, and in Jordan and Evermann, Fishes North and Mid. Amer.,
 II, 1898, 1919, Santa Catalina Island, Calif.
2968. *Radulinus asprellus* Gilbert.
 Coast of Oregon and Washington, and in Puget Sound.
Radulinus asprellus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 88, off
 Oregon and Washington, at Albatross stations 3046, 3057, and others,
 in 43 to 77 fathoms.
- Genus 1027. **STELGISTROPS** Hubbs.
Stelgistrops Hubbs, Occ. Papers, Mus. Zool. Univ. Mich. No. 170,
 Feb. 20, 1926, 8 (*Stelgistrum beringianum* Gilbert and Burke).
2969. *Stelgistrops beringiana* (Gilbert and Burke).
 South Bering Sea.
Stelgistrum beringianum Gilbert and Burke, Bull. U. S. Bur. Fish.,
 XXX, 1910, 52, fig. 9, Petrel Bank, Aleutian Islands.
- Genus 1028. **THYRISCUS** Gilbert and Burke.
Thyriscus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 43
 (*T. anoplus* Gilbert and Burke).
2970. *Thyriscus anoplus* Gilbert and Burke.
 Bering Sea.
Thyriscus anoplus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX,
 1910, 43, fig. 4, Attu Island.
- Genus 1029. **TRIGLOPS** Reinhardt.
Triglops Reinhardt, Vid. Selsk. Natur. Math. Afh., V, 1832, LII (*T.
 pingeli* Reinhardt).
2971. *Triglops pingeli* Reinhardt.
 North Atlantic, from Spitzbergen and western Norway to Greenland.
Triglops pingeli Reinhardt, Vid. Selsk. Natur., V, 1832, LII, Greenland.
Triglops pleurostictus Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865,
 81, Godhaven, Greenland.
2972. *Triglops ommatistius* Gilbert.
 North Atlantic coast of North America, Newfoundland to Cape Ann.
Triglops ommatistius Gilbert, Proc. U. S. Nat. Mus., XLIV, 1913, 465,
 Gloucester, Mass.; heretofore wrongly called *Triglops pingeli*, which
 is mainly a European species.

2973. *Triglops terræ-novæ* Gilbert.
Coast of Newfoundland.
Triglops ommatistius terræ-novæ Gilbert, Proc. U. S. Nat. Mus., XLIV, 1913, 467, off Newfoundland, in 39 fathoms.
2974. *Triglops metopias* Gilbert and Burke.
West Bering Sea.
Triglops metopias Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 50, fig. 8, Petrel Bank, Bering Sea.
2975. *Triglops beani* Gilbert.
Alaska, both north and south of the Aleutian Islands and in Bristol Bay.
Triglops beani Gilbert, Rept. U. S. Fish Com., XIX, 1893, 426, pl. 28, upper fig., Aleutian Islands and Bristol Bay, at *Albatross* stations 3214, 3217, and many others, in 7½ to 42 fathoms.
2976. *Triglops scepticus* Gilbert.
Aleutian Islands, south of Sannak and north of Unalaska Island.
Triglops scepticus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 428, pl. 28, lower fig., Aleutian Islands, at *Albatross* stations 3215, 3222, and others, in 43 to 138 fathoms.
- Genus 1030. **PRIONISTIUS** Bean.
Prionistius Bean, Proc. U. S. Nat. Mus., VI, 1883, 355 (*P. macellus* Bean).
2977. *Prionistius macellus* Bean.
British Columbia, in Carter Bay.
Prionistius macellus Bean, Proc. U. S. Nat. Mus., VI, 1883, 355, Carter Bay, British Columbia.
- Genus 1031. **ELANURA** Gilbert.
Elanura Gilbert, Rept. U. S. Fish Com., XIX, 1893, 429 (*E. forficata* Gilbert).
2978. *Elanura forficata* Gilbert.
Aleutian Islands, south of Sannak and north of Uimak Island.
Elanura forficata Gilbert, Rept. U. S. Fish Com., XIX, 1893, 430, pl. 30, fig. 1, Aleutian Islands, at *Albatross* stations 3213, 3214, and 3222, in 38 to 50 fathoms.
- Genus 1032. **STERNIAS** Jordan and Evermann.
Sternias Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1926 (*Triglops xenostethus* Gilbert).
2979. *Sternias xenostethus* (Gilbert).
Bering Sea, north to Unalaska.
Triglops xenostethus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 429, pl. 29, two figures, north of Unalaska, at *Albatross* station 3230, in 34 fathoms.
- Genus 1033. **PTERYGIOCOTTUS** Bean and Weed.
Pterygiocottus Bean and Weed, Trans. Roy. Soc. Canada, XII, 1920, 73 (*Pterygiocottus macouni* Bean and Weed).
2980. *Pterygiocottus macouni* Bean and Weed.
Vancouver Island.
Pterygiocottus macouni Bean and Weed, Trans. Roy. Soc. Canada, XIII, 1920, 73, Vancouver Island.
- Genus 1034. **MELLETES** Bean.
Melletes Bean, Proc. U. S. Nat. Mus., II, 1879, 354 (*M. papilio* Bean).
2981. *Melletes papilio* Bean.
Aleutian Islands.
Melletes papilio Bean, Proc. U. S. Nat. Mus., II, 1879, 354, St. Paul Island, Alaska.
- Genus 1035. **HEMILEPIDOTUS** Cuvier.
Hemilepidotus Cuvier, Règne Animal, ed. 2, II, 1829, 165 (*Cottus hemilepidotus* Tilesius).
Tennista Richardson, Fauna Bor.-Amer., III, 1831, 59 (*Blepsias ventricosus* Eschscholtz = *Cottus hemilepidotus* Tilesius).

- 2982. *Hemilepidotus hemilepidotus* (Tilesius). *Red sculpin.***
 Kamchatka to San Francisco; not common in Bering Sea; abundant from Sitka to Puget Sound.
Cottus hemi'epidotus Tilesius, Mém. Acad. Petersb., III, 1810, 262, Sea of Okhotsk.
Cottus trachurus Pallas, Zoogr. Ross.-Asiat., III, 1811, 138, Kuril Islands.
Hemilepidotus tilesii Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 203 (276), pl. 85, after Tilesius and Pallas.
?Blepsias ventricosus Eschscholtz, Zool. Atlas, 3d Heft, 1829, 14, t. 13, Norfolk Sound and Sitka.
Hemilepidotus gibbsii Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 13, no definite locality; Northwest boundary survey.
- 2983. *Hemilepidotus jordani* Bean. *Irish Lord; Yellow sculpin.***
 Alaska; Unalaska.
Hemilepidotus jordani Bean, Proc. U. S. Nat. Mus., IV, 1881, 153, Unalaska.
- 2984. *Hemilepidotus zapus* Gilbert and Burke.**
 Bering Sea.
Hemilepidotus zapus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 54, fig. 10, Attu Island.
- Genus 1036. CALYCILEPIDOTUS Ayres.**
Calycilepidotus Ayres, Proc. Calif. Acad. Nat. Sci., ed. 2, I, 1855, 81 (76) (*C. spinosus* Ayres).
- 2985. *Calycilepidotus spinosus* Ayres.**
 Coast of California; known only from about Monterey and San Francisco.
Calycilepidotus spinosus Ayres, Proc. Calif. Acad. Nat. Sci., ed. 2, I, 1855, 81 (76), San Francisco.
- Genus 1037. ENOPHRYS Swainson.**
Enophrys Swainson, Classn. Fishes, II, 1839, 271 (*Cottus claviger* Cuvier and Valenciennes).
- 2986. *Enophrys claviger* (Cuvier and Valenciennes).**
 Bering Sea.
Cottus claviger Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 143 (195), pl. 79, Kamchatka.
Cottus e'egans Gray, in Cuvier and Valenciennes, loc. cit., 143 (195), same type.
- Genus 1038. ASPICOTTUS Girard.**
Aspicottus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 130 (*A. bison* Girard).
Clypeocottus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 12 (*C. robustus* Ayres = *Aspicottus bison* Girard).
- 2987. *Aspicottus bison* Girard. *Buffalo sculpin.***
 San Francisco to Alaska; abundant about Puget Sound.
Aspicottus bison Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 130, San Francisco.
Clypeocottus robustus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 11, San Francisco.
- 2988. *Aspicottus taurinus* (Gilbert).**
 Monterey Bay.
Enophrys taurinus Gilbert, Proc. U. S. Nat. Mus., XLVII, 1915, 135, Monterey Bay.
- Genus 1039. CERATOCOTTUS Gill.**
Ceratocottus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 165 (*Cottus diceraus* Pallas).
- 2989. *Ceratocottus diceraus* (Pallas).**
 Alaska and Kamchatka.
Cottus diceraus Pallas, Nov. Act. Petropol., 1783, 354, pl. 10, fig. 7, Petropaulski, Kamchatka.
Synanceia cervus Tilesius, Mém. Acad. Petersb., III, 1811, 278, pl. 13, Petropaulski.
Cottus stelleri Bloch and Schneider, Syst. Ichth., 1801, 63, after Steller.

2990. Ceratocottus lucasi Jordan and Gilbert.

Bering Sea.

Ceratocottus lucasi Jordan and Gilbert, in Jördan and Evermann, Fishes North and Mid. Amer., 1898, 1940, St. Paul Island, Bering Sea.

Family 234. BLEPSIIDÆ

Genus 1040. HISTIOCOTTUS Gill.

Peropos Lay and Bennett, Beechey's Voy., Zool., Fish., 1839, 50 (*Blepsias bilobus* Cuvier and Valenciennes); name preoccupied.

Histiocottus Gill, Proc. U. S. Nat. Mus., XI, 1888, 573 (*Blepsias bilobus* Cuvier and Valenciennes).

2991. Histiocottus bilobus (Cuvier and Valenciennes).

Coast of Alaska and Kamchatka.

Blepsias bilobus Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 279 (379), no locality, somewhere off Kamchatka.

Genus 1041. BLEPSIAS Cuvier.

Blepsias Cuvier, Règne Anim., ed. 2, II, 1829, 167 (*Trachinus cirrhosus* Pallas).

2992. Blepsias cirrhosus (Pallas). *Nee-kaio-kaiung.*

Alaska to San Francisco; common at Unalaska, but not often south of Puget Sound.

Trachinus cirrhosus Pallas, Zoogr. Rosso-Asiat., III, 1811, 237, Bering Sea.

Blepsia trilobus Cuvier and Valenciennes; Hist. Nat. Poiss., IV, 1829, 276 (375); same types.

Genus 1042. NAUTICHTHYS Girard.

Nautichthys Girard, Pac. R. R. Sury., X, pt. 4, Fishes, 1858, 74 (*Blepsias oculofasciatus* Girard).

Nautiscus[?] Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2019 (*Nautichthys pribilovius* Jordan and Gilbert).

Subgenus NAUTICHTHYS Girard.

2993. Nautichthys oculofasciatus (Girard).

Pacific coast, Alaska to Monterey, chiefly northward; common in Bristol Bay and about Unalaska.

Blepsias oculofasciatus Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 202, Fort Steilacoom, Wash.

Subgenus NAUTISCUS Jordan and Gilbert.

2994. Nautichthys pribilovius (Jordan and Gilbert).

Bering Sea, Alaska.

Nautiscus pribilovius Jordan and Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2020; off St. George Island, Bering Sea.

Family 235. SCORPÆNICHTHYIDÆ. Cabezones

Genus 1043. SCORPÆNICHTHYS Girard.

Scorpenichthys Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 131 (*Hemitripterus marmoratus* Ayres).

2995 Scorpænichthys marmoratus (Ayres). *Cabezon.*

Pacific coast of America from Puget Sound to San Diego.

Hemitripterus marmoratus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 4, San Francisco.

* The genus *Nautiscus* is now regarded as scarcely distinct from *Nautichthys*.

Family 236. COTTIDÆ. Scaleless sculpins

Genus 1044. COTTUS ¹² Linnaeus. Miller's thumbs.

Cottus Linnaeus, Syst. Nat., Ed. X, 1758, 264 (*C. gobio* Linnaeus).

Pegedictis Rafinesque, Ichth. Ohiensis, 1820, 85 (*P. ictalops* Rafinesque = ?*Cottus bairdi* Girard, apparently also confused with the darter *Catotomus flabellaris*).

Uranidea DeKay, New York Fauna: Fishes, 1842, 61 (*U. quiescens* DeKay = *Cottus gracilis* Heckel).

Cottopsis Girard, Proc. Bost. Soc. Nat. Hist., III, 1850, 303 (*Cottus asper* Richardson).

Potamocottus Gill, loc. cit., VIII, 1861, 40 (*P. carolinæ* Gill).

Tauridea Jordan and Rice, in Jordan, Man. Vert., Ed. II, 1878, 255 (*Cottopsis ricei* Nelson).

2996. *Cottus gulosus* (Girard).

Streams of the Coast Ranges in California; Columbia, Klamath, and Sacramento River regions; Alaska; Loring and Boca de Quadra.

Cottopsis gulosus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 120, San Mateo Creek and San Joaquin River, Calif.

Cottopsis parvus Girard, loc. cit., 1854, 144, Monterey, Presidio, Fort Reading, Petaluma.

Uranidea semiscabra centropleura Eigenmann and Eigenmann, West Amer. Scientist, VI, No. 49, 1889, 149, Allen Springs, Lake Co., Calif.

Subgenus COTTOPSIS Girard.

2997. *Cottus asper* Richardson. Prickly bullhead.

Streams of the Cascade Range, southeastern Alaska to Oregon; south to Sacramento River.

Cottus asper Richardson, Fauna Bor.-Amer., Fish., 1836, 295, Columbia River at Fort Vancouver.

Trachidermis richardsoni Heckel, Ann. Wiener Mus., II, 1840, 162, Columbia River.

2998. *Cottus evermanni* Gilbert.

Lost River, Oreg.

Cottus evermanni Gilbert, Bull. U. S. Fish Com., XVII, 1897, 11, with figure, Lost River, Lostine, Oreg.

2999. *Cottus rhothea* (Rosa Smith).

Hangman Creek, Tekoa, Washington, and elsewhere in Columbia River Basin.

Uranidea rhothea Rosa Smith, Proc. U. S. Nat. Mus., V, 1882, 347, Spokane Falls, Wash.

3000. *Cottus shasta* Jordan and Starks.

Upper Sacramento Basin, about Mount Shasta.

Cottus shasta Jordan and Starks, Proc. Calif. Acad. Sci. 2d ser., VI, 1896, 224, McCloud River, Baird, Shasta County, Calif.

3001. *Cottus punctulatus* (Gill).

Headwaters of Green River, Wyo.; Silver Creek, Harney County, Oreg.

Potamocottus punctulatus Gill, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 40, Bridger Pass, Wyo.

¹² In every cold stream of northern Europe, northern Asia, and northern America species of this genus abound, lurking on the bottom and devouring the eggs of trout which inhabit the same waters. The species are ill differentiated and like those of trout subject to much individual variation. Those of the United States and Canada are not yet well known and the number of species has probably been unduly expanded. Efforts to subdivide the group have not been very successful and we place all the American forms in a single genus, *Cottus*. Western species, allies of *Cottus asper*, termed *Cottopsis*, are in general larger in size, more prickly, with well-developed palatine teeth and the ventral rays 1, 4. Those of the middle west allied to *Cottus bairdi*, called *Potamocottus*, show similar characters but are for the most part smaller and smoother, the colors more mottled. Rafinesque's *Pegedictis ictalops*, the "Cat's-eye Spring fish," seems to us a *Cottus* described carelessly or more likely from memory. In New England and south to Pennsylvania the allies of *Cottus gracilis* called *Uranidea* are small, smooth, without palatine teeth and the ventral rays 1, 3. But all these traits are subject to intergrading. A few species in deeper waters in Lake Michigan and Canada, allies of *Cottus ricei* called *Tauridea*, are more or less rough and have the preopercular spine larger, hooked and somewhat spiral, otherwise much as in *Cottopsis*.

- 3002. *Cottus semiscaber* (Cope).** *Rocky Mountain bullhead.*
 Rocky Mountain region in clear streams on both slopes; abundant in Colorado, northern New Mexico, Wyoming, Montana, Idaho, Utah, eastern Washington, and Oregon.
Cottopsis semiscaber Cope, Hayden Surv. Mont., 1871, 476, Fort Hall, Idaho.
Uranidea wheeleri Cope, Proc. Amer. Phil. Soc. Phila., XIV, 1874, 138, Beaver River, Utah.
- 3003. *Cottus asperrimus* Rutter.**
 California; Sacramento-San Joaquin Basin; Fall River.
Cottus asperrima Rutter, Bull. U. S. Bur. Fish., XXVII, 1907, 144, fig. 3, Fall River, Calif.
- 3004. *Cottus beldingii* Eigenmann and Eigenmann.**
 Streams of Columbia River Basin, south to Lake Lahontan; abundant east of the Cascades and in Lake Tahoe; fossil in Lahontan Basin, in Pleistocene deposits.
Cottus beldingii Eigenmann and Eigenmann, Amer. Nat., XXV, 1891, 1132, Lake Tahoe.
- 3005. *Cottus annæ* Jordan and Starks.**
 Eagle River, a tributary of Grand River, Colo., Colorado River Basin.
Cottus annæ Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 223, Pl. XXVII, Eagle River at Gypsum, Colo.
- 3006. *Cottus philonips* Eigenmann and Eigenmann.**
 Columbia River Basin at Field, British Columbia, and in Idaho.
Cottus philonips Eigenmann and Eigenmann, Amer. Nat., XXVI, 1892, 963, Kicking Horse River, Field, British Columbia.
- 3007. *Cottus klamathensis* Gilbert.**
 Klamath Lakes, Oreg.
Cottus klamathensis Gilbert, Bull. U. S. Fish. Com., XVII, 1897, 10, figure, Upper Klamath Lake, Oreg.
- 3008. *Cottus macrops* Rutter.**
 Fall River, Calif.
Cottus macrops Rutter, Bull. U. S. Bur. Fish., XXVII, 1907, 146, fig. 4, Fall River, Calif.
- 3009. *Cottus perplexus* Gilbert and Evermann.**
 Skookumchuck and Newaukum Rivers, near Chehalis, Wash.
Cottus perplexus Gilbert and Evermann, Bull. U. S. Fish Com., XIV, 1894, 202, pl. 20, fig. 1, Skookumchuck River, near Chehalis, Wash.
- 3010. *Cottus princeps* Gilbert.**
 Klamath Lakes, Oreg.
Cottus princeps Gilbert, Bull. U. S. Fish Com., XVII, 1897, 12, figure, Upper Klamath Lake, Oreg.
- 3011. *Cottus bendirei* (Bean).**
 Walla Walla, Wash.; Rattlesnake Creek near Camp Harney, Oreg., and Goose Creek near Meadows, Idaho.
 • *Potamocottus bendirei* Bean, Proc. U. S. Nat. Mus., IV, 1881, 27, Rattlesnake Creek near Camp Harney, Oreg.
- 3012. *Cottus greenei* (Gilbert and Culver).**
 Snake River Basin, Idaho.
Uranidea greenei Gilbert and Culver, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1965, Thousand Springs, Snake River, Idaho.
- 3013. *Cottus marginatus* (Bean).**
 Tributaries of Walla Walla River, Wash.
Uranidea marginata Bean, Proc. U. S. Nat. Mus., IV, 1881, 26, Walla Walla, Wash.
- 3014. *Cottus tenuis* (Evermann and Meek).**
 Klamath Lakes, Oreg.
Uranidea tenuis Evermann and Meek, Bull. U. S. Fish Com., XVII, 1897, 83, fig. 6, Upper Klamath Lake, Klamath Falls, Oreg.

3015. *Cottus leiopomus* Gilbert and Evermann.
Wood River, Shoshone, Idaho.
Cottus leiopomus Gilbert and Evermann, Bull. U. S. Fish Com., XIV, 1894, 203, pl. 20, fig. 2, Wood River, Shoshone, Idaho.
3016. *Cottus aleuticus* Gilbert.
Unalaska; Departure Bay, Vancouver Island; southward to Monterey in the Coast Range.
Uranidea microstoma Lockington, Proc. U. S. Nat. Mus., III, 1880, 58, St. Paul, Kodiak Island; name preoccupied in *Cottus*.
Cottus aleuticus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 418, streams at Unalaska; also in Departure Bay, Vancouver Island.
3017. *Cottus chamberlaini* Evermann and Goldsborough.
Southeastern Alaska.
Cottus chamberlaini Evermann and Goldsborough, Bull. U. S. Bur. Fish., XXVI, 1906, 308, fig. 66, Loring, Alaska.
- Subgenus PEGEDICTIS Rafinesque.
3018. *Cottus ictalops* (Rafinesque.) *Miller's thumb; Mufflejaws; Springfish; Blob; Bulthead.*
Middle and northern States, east of the Dakotas and Nebraska; Lake Superior to the Ohio and Potomac Rivers.
Pegedictus ictalops Rafinesque, Ichth. Ohiensis, 85, supplement, 1820. Falls of the Ohio; perhaps confused with *Catotomus flabellaris*.
Cottus richardsoni Agassiz, Lake Superior, 1850, 300, northern shore of Lake Superior; not *Trachidermis richardsoni* Heckel, also a species of *Cottus*.
Cottus bairdii Girard, Proc. Amer. Assoc. Adv. Sci., II, 1850, 410, Mahoning River, Poland, Ohio.
Cottus wilsoni Girard, Monogr. Cottoids, in Smithson. Contrib. to Knowledge, 1850, 42, Pittsburgh, Pa.
Cottus alvordii Girard, loc. cit., 46, Fort Gratiot, Lake Huron.
Uranidea spilota Cope, Proc. Acad. Nat. Sci. Phila., XVII, 1865, 82, Grand River at Grand Rapids, Mich.
3019. *Cottus meridionalis* Girard. *Southern miller's thumb; Blob; Mufflejaw*
Cottus meridionalis Girard, Proc. Am. Assn. Adv. Sci., II, 1850, 410. James River, Va.
Potamocottus carolinæ Gill, Proc. Bost. Soc. Nat. Hist., 1861, 40, Carolinas.
Potamocottus zopherus Jordan, Ann. Lyc. Nat. Hist. N. Y., XI, 1876, 320, small tributaries of the Etowah and Oostanaula Rivers near Rome, Ga.
- Subgenus URANIDEA DeKay
3020. *Cottus franklini* Agassiz.
Lake Superior and Lake Michigan in deep water.
Cottus franklini Agassiz, Lake Superior, 1850, 303, north and east shores of Lake Superior.
Uranidea hoyi Putnam, in Nelson, Bull. Ill. Lab. Nat. Hist., I, No. 1, 1876, 41, Lake Michigan, in deep water.
Uranidea kumlieni Hoy, in Nelson, loc. cit., 41, Lake Michigan, in deep water.
3021. *Cottus gracilis*¹⁴ Heckel. *Miller's thumb; Rock cusk; Little stargazer.*
Streams of New England, Maine to New York; recorded from tributaries of the Connecticut, Lake Champlain, Hudson, Delaware, and Susquehanna.
Cottus gracilis Heckel, Ann. Wien Mus., II, 1837, 148, New York.
Cottus viscosus Haldeman, Suppl. Mongr. Linnaea, 1840, 3. Susquehanna River.
Uranidea quiescens DeKay, New York Fauna: Fishes, 1842, 61, Lake Pleasant, Adirondack region.
Cottus gobiooides Girard, Proc. Amer. Assoc. Adv. Sci., II, 1850, 411, Lamville River, Johnson, Vt.
Cottus boleoides Girard, loc. cit., II, 411, Windsor, Vt.

¹⁴ May not be distinct from the northern *C. cognatus*.

- Cottus copei* Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 326,
probably New Jersey.
- Cottus gracilis cayuga* Meek, Ann. New York Acad. Sci., IV, 1888, 315,
Cayuga Lake, N. Y.
- 3022. *Cottus formosus* Girard.**
Lake Ontario, in deep water.
Cottus formosus Girard, Monogr. Cott., in Smithson. Contrib., III,
1850, 58, Lake Ontario, off Oswego, in stomach of *Lota maculosa*.
- Subgenus TAURIDEA Jordan and Rice.**
- 3023. *Cottus ricei* (Nelson).**
Lake Michigan; Lake Ontario.
Cottopsis ricei Nelson, Bull. Ill. Lab. Nat. Hist., I, 1876, 40, Lake
Michigan off Evanston.
Uranidea pollicaris Jordan and Gilbert, Proc. U.S. Nat. Mus., V,
1882, 222, Lake Michigan, off Racine, Wis.
- 3024. *Cottus onychus* Eigenmann and Eigenmann.**
Saskatchewan Basin.
Cottus onychus Eigenmann and Eigenmann, Am. Nat., XXVI, No-
vember, 1892, 963, Bow River at Calgary, a tributary of the South
Saskatchewan, Alberta Territory.
- 3025. *Cottus cognatus* Richardson.** *Northern miller's thumb.*
Northern United States and northward, through Canada.
Cottus cognatus Richardson, Fauna Bor.-Amer., III, 1836, 40, Great
Bear Lake.
- Genus 1045. ACANTHOCOTTUS Girard.** *Daddy sculpins.*
Acanthocottus Girard, Proc. Bost. Soc. Nat. Hist., III, 1849, 185 (*Cottus*
grænlandicus Cuvier and Valenciennes).
- 3026. *Acanthocottus mitchilli* ¹⁵ (DeKay).**
Coast of southern New England and New York.
Cottus mitchilli DeKay, in Cuvier and Valenciennes, Hist. Nat. Poiss.,
IV, 1829, 188, New York; after *Cottus scorpio* Mitchell.
- 3027. *Acanthocottus bubalis* (Euphrasen).** *Fatherlasher; Lucky proach.*
Barbados.
Cottus bubalis Euphrasen, König. Vet. Nya Abhandl., VII, 1786, 65,
taf. 3, fig. 8, 1883, Barbados.
Cottus maculatus Fischer, Jahrb. Wissenschaft. Aust. Hamburg, II, 1885,
78, taf. 2, fig. 8, Barbados.
- 3028. *Acanthocottus æneus* (Mitchill).** *Grubby; Brassy sculpin; Pigmy*
sculpin.
Coast of southern New England and New York.
Cottus æneus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 380, New
York.
Cottus anceps Sauvage, Nouv. Archiv Mus. Paris, I, 1878, 145, pl. 1,
fig. 13, New York.
? *Cottus nigricans* LeSueur, in Vaillant, Bull. Soc. Philom., serie 8,
tome VIII, No. 1, 1895, 12, no locality.
- 3029. *Acanthocottus octodecemspinosis* (Mitchill).** *Long-spined sculpin;*
Common sculpin; Bullhead; Eighteen-spined sculpin; Hacklehead.
Atlantic coast south to Virginia; common northward.
Cottus octodecem-spinosus Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815,
380, New York.
- 3030. *Acanthocottus scorpius* (Linnæus).** *European sculpin; Ulke.*
Norther Europe and Arctic regions, not common on our coasts; recorded
by Doctor Lütken from the Baltic, Finland, Spitzbergen, Nova
Zembla, coasts of England and northern Asia; also recorded from
Eastport, Me.
Cottus scorpius Linnæus, Syst. Nat., Ed. X, 1758, 265, in Atlantic
Ocean off Europe.
- 3031. *Acanthocottus scorpioides* (Fabricius).** *Arctic sculpin.*
Arctic regions of America, Greenland to Siberia.
Cottus scorpioides Fabricius, Fauna Greenl., 1780, 157, Greenland.
Cottus pachypus Günther, Cat., II, 1860, 161, Port Leopold.

¹⁵ Differentiated by Nichols, Copeia, No. 31, 40, 1910.

- 3032. *Acanthocottus grønlandicus* (Cuvier and Valenciennes). *Daddy sculpin.***
 New York to Greenland; according to Fabricius it is abundant in all the bays and inlets of Greenland.
Cottus grønlandicus Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 133 (156), no locality given, probably Greenland; after Fabricius.
Cottus porosus Cuvier and Valenciennes, loc. cit., VIII, 1831, 367 (498), Baffins Bay.
Acanthocottus mucosus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854 (Oct. 9), 12, coast of New England.
Acanthocottus ocellatus Storer, Boston Jour. Nat. Hist., 1850, 253, Labrador.
Cottus glacialis Richardson, in Belcher, Last Arctic Voy. H. M. S. Assistant, App. 3, 1855, 349, pl. 24, Northumberland Sound.
- Genus 1046. *MYOXOCEPHALUS* (Steller) Tilesius.**
Myoxocephalus Tilesius, Mém. Acad. Sci. Petersb., IV, 1811, 273 (*Myoxocephalus stelleri* Tilesius).
Boreocottus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 166 (*B. axillaris* Gill).
- 3033. *Myoxocephalus polyacanthocephalus* (Pallas). *Great sculpin; Kalog.***
 Alaska to Kamchatka; abundant, Bering Sea to Puget Sound; tide pools of Karluk.
Cottus polyacanthocephalus Pallas, Zoogr. Rosso.-Asiat., III, 1811, 133, Aleutian Islands.
- 3034. *Myoxocephalus jack* (Cuvier and Valenciennes).**
 Kamchatka.
Cottus jack Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 126 (172), Kamchatka.
Cottus humilis Bean, Proc. U. S. Nat. Mus., IV, 1881, 149, Chamisso Island, Bering Straits.
- 3035. *Myoxocephalus verrucosus* (Bean).**
 Plover Bay, Siberia, near Bering Straits; Unalaska; Sitka; Bristol Bay; Kings Island; Grantley Harbor, Port Clarence; Litnik.
Cottus verrucosus Bean, Proc. U. S. Nat. Mus., IV, 1881, 152, Plover Bay, Siberia.
- 3036. *Myoxocephalus axillaris* (Gill).**
 Bering Straits.
Boreocottus axillaris Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 166, Bering Straits.
- 3037. *Myoxocephalus stelleri* Tilesius.**
 Western shore of Bering Sea and Okhotsk Sea.
Myoxocephalus stelleri Tilesius, Mém. Acad. Petersb., IV, 1811, 273, with plate, Petropaulski.
Cottus decastrensis Kner, Denk. Kais. Acad. Wissen., XXIV, 1865, 2, taf. 2, figs. 1, 1a, Decastris Bay, near mouth of the Amur.
?Cottus marmoratus Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 366 (497), Petropaulski.
- 3038. *Myoxocephalus mednius* Bean.**
 Medni Island, Bering Sea.
Myoxocephalus mednius Bean, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1983, Medni Island, Bering Sea.
- 3039. *Myoxocephalus parvulus* Gilbert and Burke.**
 Medni Island, Commander Group.
Myoxocephalus parvulus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 59, Preobazensko Bar, Medni (Copper) Island.
- 3040. *Myoxocephalus niger* (Bean). *Black sculpin.***
 Pribilof Islands.
Cottus niger Bean, Proc. U. S. Nat. Mus., IV, 1881, 151, St. Paul Island, Bering Sea.
?Cottus mertensi Cuvier and Valenciennes, Hist. Nat. Poiss., VIII, 1831, 366 (496), Petropaulski.
- Genus 1047. *MEGALOCOTTUS* Gill.**
Megalocottus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 166 (*Cottus platycephalus* Pallas).

3041. *Megalocottus platycephalus* (Pallas).
 Kamchatka.
Cottus platycephalus Pallas, Zoogr. Rosso.-Asiat., III, 1811, 135, Kamchatka; after Steller.
Cottus tenuiopterus Kner, Sitzungsber. Ak. Wiss. Wien, LXIII, 1868, 28, Taf. 14, fig. 10, Decastris Bay, near mouth of Amur River.
3042. *Megalocottus laticeps* (Gilbert).
 Eastern shore of Bering Sea.
Acanthocottus laticeps Gilbert, Rept. U. S. Fish. Com., XIX, 1893, 422, pls. 26 and 27, Nushagak River and Herendeen Bay, Alaska.
 Genus 1048. **POROCOTTUS** Gill.
Porocottus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 166 (*P. quadrifilis* Gill).
3043. *Porocottus sellaris* (Gilbert).
 Bristol Bay, Alaska.
Acanthocottus sellaris Gilbert, Rept. U. S. Fish. Com. XIX, 1893, 419, Bristol Bay, at Albatross stations 3229, 3231, and others, in 5 to 17 fathoms.
3044. *Porocottus quadratus* Bean.
 Bering Island.
Porocottus quadratus Bean, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1998, Bering Island.
3045. *Porocottus polaris* (Sabine).
 East side of peninsula of Boothia.
Cottus polaris Sabine, App. Parry's First Voyage, CCXIII, 1824, north Georgia.
3046. *Porocottus quadrifilis* Gill.
 Bering Straits.
Porocottus quadrifilis Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 166, Bering Straits.
3047. *Porocottus bradfordi* Rutter.
 Alaska; rock pools at Karluk.
Porocottus bradfordi Rutter, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2862, Karluk.
- Genus 1049. **MALACOCOTTUS** Bean.
Malacocottus Bean, Proc. U. S. Nat. Mus., XIII, 1890, 42 (*M. zonurus* Bean).
3048. *Malacocottus zonurus* Bean.
 Coast of Alaska off Trinity Islands.
Malacocottus zonurus Bean, Proc. U. S. Nat. Mus., XIII, 1890, 43, off Trinity Islands, Alaska, 56° N., 154° W., at Albatross station 2853, in 159 fathoms.
3049. *Malacocottus kincaidi* Gilbert and Thompson.
 Puget Sound.
Malacocottus kincaidi Gilbert and Thompson, Proc. U. S. Nat. Mus., XXVIII, 1905, 979, fig. 2, Brinnon, Hood's Canal, Wash.
- Genus 1050. **ONCOCOTTUS** Gill.
Oncocottus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 13 (*Cottus quadricornis* Linnaeus).
3050. *Oncocottus quadricornis* (Linnaeus).
 Arctic America to Baltic Sea, chiefly northward; abundant in eastern Baltic and in Lakes Ladoga and Onega; north to White Sea and Nova Zembla; rare in England and eastern Greenland; unknown in western Greenland.
Cottus quadricornis Linnaeus, Syst. Nat., ed. X, 1758, 264, Baltic Sea.
3051. *Oncocottus hexacornis* (Richardson).
 Arctic Sea about Hershel Island.
Cottus hexacornis Richardson, Franklin's Journal, 1823, 726, mouth of Tree River near Coppermine River.
Acanthocottus labradoricus Girard, Bost. Jour. Nat. Hist., VI, 1850, 247, pl. 7, fig. 3, coast of Labrador,

3052. *Oncocottus giberti* Fowler.
 Point Barrow, Alaska; Arctic Sea.
Oncocottus hexacornis giberti Fowler, Proc. Acad. Nat. Sci. Phila., LVII, 1905, 366, figs. 2, 3, 4, and 5, Point Barrow, Alaska.
- Genus 1051. **TRIGLOPSIS** Girard.
Triglopsis Girard, Proc. Bost. Soc. Nat. Hist., IV, 1851, 16 (*T. thompsoni* Girard).
3053. *Triglopsis thompsoni* Girard. *Lake sculpin*.
 Deep waters of Lake Ontario.
Triglopsis thompsoni Girard, Proc. Bost. Soc. Nat. Hist., IV, 1851, 19, Lake Ontario off Oswego, N. Y.
3054. *Triglopsis stimpsoni* Hoy.
 Lake Michigan in deep water.
Triglopsis stimpsoni Hoy, Trans. Wis. Acad. Sci., 1872, 99, Lake Michigan, off Chicago.
3055. *Triglopsis ontariensis* Jordan and Thompson.
 Deep waters of Lake Ontario.
Triglopsis ontariensis Jordan and Thompson, Proc. U. S. Nat. Mus., XXXVIII, 1911, 75, Lake Ontario, off Toronto.
- Genus 1052. **GYMNOCANTHUS** Swainson.
Gymnacanthus Swainson, Class. Fishes, Amphib., etc., II, 1839, 271 (*Cottus ventralis* Cuvier and Valenciennes = *Cottus pistilliger* Pallas).
Phobetor Krøyer, Naturh. Tidskr., I, 1844, 263 (*Cottus tricuspidatus* Reinhardt).
Elaphocottus Sauvage, Nouv. Arch. Mus. Paris (2), 1, 1878, 142 (*Cottus pistilliger* Pallas).
Gymnacanthus of many authors. Corrected spelling.
3056. *Gymnacanthus pistilliger* (Pallas).
 Coasts of Alaska; taken abundantly in Bristol Bay.
Cottus pistilliger Pallas, Zoogr. Rosso-Asiat., III, 1811, 143, Unalaska Island.
Cottus ventralis Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 142 (194), Kamchatka.
Cottus cephaloides Gray, in Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 142 (194), Kamchatka.
3057. *Gymnacanthus tricuspidatus* (Reinhardt).
 Arctic seas, south to Norway and Labrador; not very common on our coasts.
Cottus tricuspidatus Reinhardt, Vidensk. Selsk. Nat. Math., V, 1832, LII, Greenland.
Acanthocottus patris Storer, Bost. Jour. Nat. Hist., VI, 1857, 250, Labrador.
3058. *Gymnacanthus galeatus* (Bean).
 Aleutian Islands.
Gymnacanthus galeatus Bean, Proc. U. S. Nat. Mus., IV, 1881, 153, Unalaska.
- Genus 1053. **CROSSIAS** Jordan and Starks.
Crossias Jordan and Starks, Proc. U. S. Nat. Mus., XXVII, 1904, 296 (*C. allisi* Jordan and Starks).
3059. *Crossias albomaculatus* Schmidt.
 Bering Sea.
Crossias albomaculatus Schmidt, Ann. Mus. Zool. Russ., XX, 1923, 614, fig. 2, Bering Sea.
3060. *Crossias beringi* Soldatov.
 Bering Sea.
Crossias beringi Soldatov, Ann. Mus. Zool. Russ., XXI, 1924, 218, Bering Sea.
3061. *Crossias camtschaticus* Schmidt.
 Kamchatka.
Crossias camtschaticus Schmidt, Ann. Mus. Zool. Russ., XX, 1923, 17, Kamchatka.

Genus 1054. LEILOCOTTUS Girard.

Leiocottus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 133 (*L. hirundo* Girard).

3062. *Leiocottus hirundo* Girard.

Coast of southern California, chiefly about the Santa Barbara Islands; extremely local.

Leiocottus hirundo Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 133, San Miguel Island near Santa Barbara, Calif.

Genus 1055. LEPTOCOTTUS Girard.

Leptocottus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 130 (*L. armatus* Girard).

3063. *Leptocottus armatus* Girard. *Smooth cabezon.*

Pacific coast, from Kodiak to San Diego; abundant northward.

Leptocottus armatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 131, Cape Flattery, Fort Steilacoom, Shoalwater Bay, Humboldt Bay, San Francisco, Monterey, San Pedro, Fort Point, San Diego, and Tomales Bay.

Genus 1056. ASEMATICHTHYS Gilbert.

Asematichthys Gilbert (*Asematichthys taylori* Gilbert).

3064. *Asematichthys taylori* Gilbert.

Vancouver Island.

Asematichthys taylori Gilbert, Departure Bay, Nanaimo, Vancouver Island.

Genus 1057. CLINOCOTTUS Gill.

Clinocottus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 166 (*Oligocottus analis* Girard).

Rusulus Starks and Mann, Univ. Calif. Pub. Zool., VIII, 1911, 14 (*Clinocottus sabarrensis* Starks and Mann; young).

3065. *Clinocottus analis* (Girard).

Coast of California from Point Conception northward.

Oligocottus analis Girard, Proc. Acad. Nat. Sci. Phila., IX, 1857, 201, Monterey.

Cottus criniger Günther, Cat., II, 1860, 522, Monterey, Calif.

Rusulus sabarrensis Starks and Mann, Univ. Calif. Pub. Zool., VIII, 1911, 14, Ballast Point, off mouth of San Diego Bay; young.

3066. *Clinocottus australis* Hubbs.

Southern California from Los Angeles County and Santa Catalina Island southward to Todos Santos, San Martin, San Benito, and Guadalupe Islands. Probably varies into *C. analis*, of which it is the southern form.

Clinocottus analis australis Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 171, Feb. 20, 1926, 11, Point Loma, San Diego, Calif.

Genus 1058. OLIGOCOTTUS Girard.

Oligocottus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 132 (*O. maculosus* Girard).

3067. *Oligocottus maculosus* Girard. *Johnny.*

San Francisco to central Alaska.

Oligocottus maculosus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 133, Tomales Bay, San Francisco.

Oligocottus borealis Jordan and Snyder, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 225, Neah Bay, Wash.

Genus 1059. GREELEYA⁷⁶ Jordan.

Eximia Greeley, Bull. U. S. Fish Com., XIX, 1899, 18 (*Eximia rubellio* Greeley); name preoccupied.

Greeleya Jordan, Genera of Fishes, IV, 1920, 493 (*Eximia rubellio* Greeley).

⁷⁶ The separation of this genus from *Oligocottus* has been questioned by Starks and approved by Hubbs.

- 3068.** *Greeleya rubellio* (Greeley).
 Monterey Bay, in deep tide pools.
Exinia rubellio Greeley, Bull. U. S. Fish Com., XIX, 1899, 18, fig. 5,
 Pacific Grove, Calif.
- Genus 1060.** *DIALARCHUS* Greeley.
Dialarchus Greeley, Bull. U. S. Bur. Fish., XIX, 1899, 14 (*Oligocottus snyderi* Greeley).
- 3069.** *Dialarchus snyderi* (Greeley).
 Coast of California, Crescent City to San Luis Obispo, in tide pools.
Oligocottus snyderi Greeley, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2871, Pacific Grove, Calif.
- Genus 1061.** *SIGMISTES* Rutter.
Sigmistes Rutter, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2863 (*S. caulias* Rutter).
- 3070.** *Sigmistes caulias* Rutter.
 Rock pools at Kodiak Island, Alaska.
Sigmistes caulias Rutter, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2863, rock pools at Karluk, Kodiak Island, Alaska.
- Genus 1062.** *OXYCOTTUS* Jordan and Evermann.
Oxycoetus Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2015 (*Oligocottus acuticeps* Gilbert).
- 3071.** *Oxycottus acuticeps* (Gilbert).
 Alaska south to Cape Flattery, Departure Bay, Vancouver Island, Kadiak.
Oligocottus acuticeps Gilbert, Rept. U. S. Fish Com., XIX, 1893, 432, Unalaska.
- Genus 1063.** *ALLOCOTTUS* Hubbs.
Allocottus Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 171, 1926, 14 (*Oligocottus embryum* Jordan and Starks).
- 3072.** *Allocottus embryum* (Jordan and Starks).
 Washington (Neah Bay), Point Lobos, Calif., Karluk and Sitka, Alaska.
Oligocottus embryum Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 808, pl. 82, Neah Bay, Wash.
- Genus 1064.** *RUSCICULUS* Greeley.
Rusciculus Greeley, Bull. U. S. Fish Com., XIX, 1899, 13 (*R. rimensis* Greeley).
- 3073.** *Rusciculus rimensis* Greeley.
 Monterey Bay in tide pools.
Rusciculus rimensis Greeley, Bull. U. S. Fish Com., XIX, 1899, 13, fig. 3, Point Lobos, near Carmel, Calif.
- Genus 1065.** *BLENNICOTTUS* Gill.
Blennicottus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 166 (*Oligocottus globiceps* Girard).
- 3074.** *Blennicottus globiceps* (Girard).
 Pacific coast from Coronado Islands northward to Cape Mendocino.
Oligocottus globiceps Girard, Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 58, South Farallons.
Blennicottus globiceps bryosus Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 808, Neah Bay, Wash.
- Genus 1066.** *MONTEREYA* Hubbs.
Montereya Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 171, 1926, 16 (*Blenicottus recalvus* Greeley).
- 3075.** *Montereya recalva* (Greeley).
 Monterey Bay to San Diego, in tide pools.
Blennicottus recalvus Greeley, Bull. U. S. Fish Com., XIX, 1899 (Dec. 13), 9, fig. 1, Pacific Grove, Calif.
- Genus 1067.** *ORTHONOPIAS* Starks and Mann.
Orthonopias Starks and Mann, Univ. Cal. Publ. Zool., VIII, 1911, 11, (*O. triacis* Starks and Mann).

- 3076.** *Orthonopias triacis* Starks and Mann.
 Coast of California, Monterey to San Diego.
Orthonopias triacis Starks and Mann, Univ. Cal. Publ. Zool., VIII, 1911, 11, fig. 1, Cortez Banks off San Diego.
- Genus 1068. DASYCOTTUS Bean.**
Dasycocttus Bean, Proc. U. S. Nat. Mus., XIII, 1890, 42 (*D. setiger* Bean).
- 3077.** *Dasycottus setiger* Bean.
 Puget Sound to Bering Sea.
Dasycocttus setiger Bean, Proc. U. S. Nat. Mus., XIII, 1890, 42, off Sitkalidak Island, Alaska.
- Genus 1069. ZESTICELUS Jordan and Evermann.**
Zesticelus Jordan and Evermann, Check List, 1895, 443 (*Acanthocottus profundorum* Gilbert).
- 3078.** *Zesticelus profundorum* (Gilbert).
 Bering Sea, north of Unalaska.
Acanthocottus profundorum Gilbert, Rept. U. S. Fish Com., XIX, 1893, 423, pl. 27, upper fig., Bering Sea, north of Unalaska, at *Albatross* station 3328, in 399 fathoms.
- Genus 1070. THECOPTERUS Smith.**
Thecopterus Smith (H. M.), Proc. Biol. Soc. Wash., XVII, 1904, 163 (*Thecopterus aleuticus* Smith).
- 3079.** *Thecopterus aleuticus* Smith.
 Bering Sea.
Thecopterus aleuticus Smith, Proc. Biol. Soc. Wash., XVII, 1904, 163, near Rat Islands, in 270 fathoms.
- Genus 1071. COTTUNCULUS Collett.**
Cottunculus Collett, Norges Fiske, 1875, 20 (*C. microps* Collett).
- 3080.** *Cottunculus microps* Collett.
 Deep waters off coasts of Norway and Rhode Island.
Cottunculus microps Collett, Norges Fiske, 1875, 20, pl. I, figs. 1-3, Hasvig near Hammerfest, Norway, in 200 fathoms.
- 3081.** *Cottunculus thomsonii* (Günther).
 North Atlantic, in deep water.
Cottus thomsonii Günther, Proc. Royal Soc. Edinburgh, XI, 1882, 679, Faroe Channel, in 535 fathoms.
Cottunculus torvus Goode, Proc. U. S. Nat. Mus., III, 1880, 479, Gulf Stream off Rhode Island. *A nomen nudum.*
- Genus 1072. ASCELICHTHYS Jordan and Gilbert.**
Ascelichthys Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 264 (*A. rhodorus* Jordan and Gilbert).
- 3082.** *Ascelichthys rhodorus* Jordan and Gilbert.
 Pacific coast from Sitka to Cape Mendocino; abundant in Neah Bay, Wash.
Ascelichthys rhodorus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 264, Waadda Island, Neah Bay, near Cape Flattery.
- Genus 1073. PSYCHROLUTES Günther.**
Psychrolutes Günther, Cat., III, 1861, 516 (*P. paradoxus* Günther).
- 3083.** *Psychrolutes paradoxus* Günther.
 Puget Sound to Kodiak Island.
Psychrolutes paradoxus Günther, Cat., III, 1861, 516, Gulf of Georgia.
Psychrolutes zebra Bean, Proc. U. S. Nat. Mus., XIII, 1890, 43, Aleutian Islands, between Unga and Nagai Islands, in 110 fathoms.
- Genus 1074. GILBERTIDIA Berg.**
Gilbertina Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 811 (*Gilbertina sigalutes* Jordan and Starks); name preoccupied.
Gilbertidia Berg, Comun. del Museo Nac. de Buenos Aires, I, 1898, 42 (*Gilbertina sigalutes* Jordan and Starks).

3084. *Gilbertidia sigalutes* (Jordan and Starks).

Puget Sound.

Gilbertina sigalutes Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 811, pl. 86, Port Orchard, near Seattle.

Family 237. SYNCHIRIDÆ

Genus 1075. SYNCHIRUS Bean.

Synchirus Bean, Proc. U. S. Nat. Mus., XII, 1889, 641 (*S. gilli* Bean).

3085. *Synchirus gilli* Bean.

Puget Sound; Barclay Sound, Port Townsend; rare.

Synchirus gilli Bean, Proc. U. S. Nat. Mus., XII, 1889, 642, Barclay Sound, British Columbia.

Family 238. RHAMPHOCOTTIDÆ

Genus 1076. RHAMPHOCOTTUS Günther.

Rhamphocottus Günther, Ann. Mag. Nat. Hist., Series 4, XIV, 1874, 369 (*R. richardsoni* Günther).

3086. *Rhamphocottus richardsoni* Günther.

North Pacific, Alaska to Monterey; not infrequent in Puget Sound.

Rhamphocottus richardsoni Günther, Ann. Mag. Nat. Hist., XIV, 1874, 369, Fort Rupert, British America.

Genus 1077. ULCA Jordan and Evermann.

Ulca Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 227 (*Hemitripterus marmoratus* Bean).

3087. *Ulca marmorata* (Bean).

Alaska, Bering Sea and off Sitkalidak Island.

Hemitripterus marmoratus Bean, Proc. U. S. Nat. Mus., XIII, 1890, 43, off Sitkalidak Island.

Family 239. HEMITRIPTERIDÆ. Sea ravens

Genus 1078. HEMITRIPTERUS Cuvier.

Hemitripterus Cuvier, Règne Anim., ed. 2, II, 1829, 164 (*Scorpaena americanus* Gmelin).

3088. *Hemitripterus americanus* (Gmelin). *Sea raven; Yellow sculpin.*

Atlantic coast of America from Tortugas to Labrador, chiefly northward.

Scorpaena americana Gmelin, Syst. Nat., ed. XIV, 1788, 1220, after "Diable en Crapaud de l'Amérique" of Duhamel de Moreau.

Cottus acadianus Walbaum, Artedi Fisc., 1792, 392, 1784, Nova Scotia; after Pennant.

Cottus hispidus Bloch and Schneider, Syst. Ichth., 1801, 63, New York.

Cottus tripterygius Bloch and Schneider, Syst. Ichth., 1801, 63, Nova Scotia; after Pennant.

3089. *Hemitripterus villosus* (Pallas).

Coast of Alaska.

Cottus villosus Pallas, Zoogr. Rosso-Asiat., III, 1811, 129, Cape Kronok, Kamchatka, Itscha River.

Hemitripterus cauifrons Lockington, Proc. Acad. Nat. Sci. Phila., XXXII, 1880, 233, Kodiak Island, Alaska.

Family 240. AGONIDÆ. Sea poachers; Alligatorfishes

Genus 1079. HYPSEGONUS Gill.

Hypsagonus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 167 (*Aspidophorus quadricornis* Cuvier and Valenciennes).

Cheiragonus Herzenstein, Bull. Acad. Sci. St. Petersb., XIII, 1890, 116 (*Hypsagonus gradiens* Hertzenstein = *Aspidophorus quadricornis* Cuvier and Valenciennes).

3090. *Hypsagonus quadricornis* (Cuvier and Valenciennes).

North Pacific; Kamchatka; Bering Sea and coasts of Alaska south to Bristol Bay and Puget Sound.

Aspidophorus quadricornis Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829, 162 (221), Kamchatka.

Hypsagonus gradiens Herzenstein, Bull. Acad. Int. Sci. St. Petersb., XIII, 1890, 116, Kamchatka, Sinus Awatcha Bay, Petropaulski.

Genus 1080. STELLERINA Cramer.

Stellerina Cramer, in Jordan and Evermann, Check List, 1895, 447
 (*Brachyopsis xyosternus* Jordan and Gilbert).

3091. *Stellerina xyosterna* (Jordan and Gilbert).

Coast of California and Oregon.

Brachyopsis xyosternus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, July 2, 1880, 152, Santa Cruz, Calif.

Agonus annæ Steindachner, Ichth. Beitr., IX, 1880, 254, pl. 6, figs. 1, 1a, Sitzb. K. Acad. Wiss., LXXXII, San Francisco, Calif.

Genus 1081. OCCA Jordan and Evermann.

Occa Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2043
 (*Brachyopsis verrucosus* Lockington).

3092. *OCCA verrucosa* (Lockington).

Coast of California, south to San Francisco.

Brachyopsis verrucosus Lockington, Proc. U. S. Nat. Mus., III, May 6, 1880, 60, Drake Bay, near San Francisco.

Agonus barkani Steindachner, Ichth. Beitr., IX, 1880, 253, pl. 5, Sitzb. K. Akad. Wiss., LXXXII, San Francisco, Calif.

Genus 1082. OCCELLA Jordan and Hubbs.

Ocella Jordan and Hubbs, Mem. Carnegie Mus., X, No. 2, 1925, 291
 (*Agonus dodecaëdron* Tilesius).

3093. *Occella dodecaëdron* (Tilesius).

North Pacific, Kamchatka, Bristol Bay, coast of Alaska.

Agonus dodecaëdrus Tilesius, Mém. Acad. Petersb., IV, 1810, pl. 13, Kamchatka.

Phalangistes loricatus Pallas, Zoog. Rosso-Asiat., III, 1811, 114, tab. 19, Kamchatka.

Genus 1083. PALLASINA Cramer.

Pallasina Cramer, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 815 (*Siphagonus barbata* Steindachner).

3094. *Pallasina barbata* (Steindachner).

Bering Sea on both coasts to Hokkaidō, northern Japan.

Siphagonus barbatus Steindachner, Ichth. Beitr., V, July, 1876, 140, pl. 5, Sitzb. der K. Akad. Wiss., LXXIV, Hakodate, Japan; the locality of Nagasaki also mentioned is impossible.

3095. *Pallasina aix* Starks.

Puget Sound to the Aleutian Islands.

Pallasina aix Starks, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 558, pl. 75, Puget Sound near Port Ludlow, Wash.

Genus 1084. LEPTAGONUS Gill.

Leptagonus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 167, 259
 (*Aspidophorus spinosissimus* Krøyer = *Aspidophorus decagonus* Cuvier and Valenciennes).

3096. *Leptagonus decagonus* (Bloch and Schneider).

Arctic Ocean, south to Newfoundland and Norway.

Agonus decagonus Bloch and Schneider, Syst. Ichth., I, 1801, 105, pl. 27, erroneously recorded from the East Indies; the type from Greenland.
Aspidophorus spinosissimus Krøyer, Naturhist. Tidskr., I, 1844, 250, Greenland.

Genus 1085. PODOTHECUS Gill.

Podothecus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 77, 259 (*P. peristethus* Gill = *Agonus acipenserinus* Tilesius).

Paragonus Gill, loc. cit., 167, 259 (*Agonus acipenserinus* Tilesius).

3097. *Podothecus acipenserinus* (Tilesius). Common alligatorfish.

Kamchatka to Puget Sound.

Agonus acipenserinus Tilesius, Mém. Acad. Petersb., IV, 1811, 422, pl. 11, fig. 163, Unalaska.

Podothecus peristethus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 259, Simeahmoo, Washington Territory.

3098. *Podothecus gilberti* Collett.

Kamchatka.

Podolthecus gilberti Collett, Proc. Zool. Soc. London, 1894, 67, pl. 45,
Kamchatka.

Genus 1086. AGONUS Bloch and Schneider:

Agonus Bloch and Schneider, Syst. Ichth., 1801, 104 (*Cottus cataphractus*
Linnæus).*Aspidophorus* Lacépède, Hist. Nat. Poiss., III, 1802, 221 (*A. armatus*
Lacépède = *Cottus cataphractus* Linnæus).*Phalangistes* Pallas, Zoog. Rosso-Asiat., III, 1811, 113 (*Cottus cata-*
phractus L.).3099. *Agonus cataphractus* (Linnæus). Sea poacher; Pogge.

Northern Europe to western Greenland.

Cottus cataphractus Linnæus, Syst. Nat., Ed. X, 1758, 264; after Artedi.*Cottus brodami* Bonnaterre, Encycl. Meth., 1788, 67, northern seas.
Aspidophorus armatus Lacépède, Hist. Nat. Poiss., III, 1802, 222

Atlantic Ocean.

Calaphractus schoneveldii Fleming, British Animals, 1842, 216, Great
Britain.*Aspidophorus europaeus* Cuvier and Valenciennes, Hist. Nat. Poiss.,
IV, 1829, 147 (201), sandy shores of northern Europe.

Genus 1087. STELGIS Cramer.

Stelgis Cramer, in Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser.,
V, 1895, 821 (*Agonus vulsus* Jordan and Gilbert).3100. *Stelgis vulsus* (Jordan and Gilbert).

Off San Francisco.

Agonus vulsus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880,
330, Point Reyes, near San Francisco.

Genus 1088. AVERRUNCUS Jordan and Starks.

Avveruncus Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895,
821, 824 (*A. emmelane* Jordan and Starks).3101. *A verruncus emmelane* Jordan and Starks.

Puget Sound, near Port Orchard.

Avveruncus emmelane Jordan and Starks, Proc. Calif. Acad. Sci., 2d
ser., V, 821, pl. 91, Port Orchard, Puget Sound.3102. *A verruncus sterletus* Gilbert.

Coast of southern California.

Avveruncus sterletus Gilbert, in Jordan and Evermann, Fishes North
and Mid. Amer., 1898, 2072, off Coronado Island.

Genus 1089. XYSTES Jordan and Starks.

Xystes Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 824
(*X. axinophrys* Jordan and Starks).3103. *Xystes axinophrys* Jordan and Starks.

Puget Sound to Monterey.

Xystes axinophrys Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser.,
V, 1895, 824, pl. 92, Port Orchard, Admiralty Inlet.

Genus 1090. SARRITOR Cramer.

Sarritor Cramer, in Jordan and Evermann, Check List, 1895, 448, and
Fishes of North and Mid. Amer., 1898, 2072 (*Odontopyxix frenatus*
Gilbert).3104. *Sarritor frenatus* (Gilbert).

Coast of Alaska.

Odontopyxix frenatus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 435,
pl. 30, fig. 3, Alaska and Aleutian Islands, at Albatross stations 3219,
3225 and many others, in 16 to 351 fathoms.3105. *Sarritor leptorhynchus* (Gilbert).

Coast of Alaska.

Odontopyxix leptorhynchus Gilbert, Rept. U. S. Fish Com., XIX, 1893,
437, north and south of the Alaska peninsula, at Albatross stations
3215, 3219, and others in 32 to 59 fathoms.

Genus 1091. BATHYAGONUS Gilbert.

Bathyagonus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 89 (*B. nigri-*
pinnis Gilbert).

3106. *Bathyagonus nigripinnis* Gilbert.

North Pacific; known from the Aleutian Islands, Bering Sea, south to the coast of Washington.

Bathyagonus nigripinnis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 89, coast of Washington, at *Albatross* station 3073, in 477 fathoms.

Genus 1092. *XENERETMUS* Gilbert.

Xenochirus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 90 (*X. triacanthus* Gilbert); name preoccupied.

Xeneretmus Gilbert in Jordan, Amer. Nat., 1903, 360 (*Xenochirus triacanthus* Gilbert).

3107. *Xeneretmus triacanthus* (Gilbert).

Pacific coast, from Santa Barbara to Washington.

Xenochirus triacanthus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 91, off coast of California, Oregon, and Washington, at *Albatross* stations 2893, 2973, and 3059, in 145, 68, and 77 fathoms.

Genus 1093. *ASTEROTHECA* Gilbert.

Asterotheca Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 343 (*Xenochirus pentacanthus* Gilbert).

3108. *Asterotheca pentacantha* (Gilbert).

Coast of Washington.

Xenochirus pentacanthus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 91, coast of Washington, at *Albatross* station 3076, in 178 fathoms.

3109. *Asterotheca alascanus* (Gilbert).

Puget Sound to Bering Sea; coast of Washington.

Xenochirus alascanus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 438, Unimak Pass, Aleutian Islands, at *Albatross* stations 3216, 3219, and others, in 35 to 138 fathoms.

3110. *Asterotheca infraspinata* (Gilbert).

Off coast of Washington.

Xeneretmus infraspinatus Gilbert, Proc. Calif. Acad. Sci., 3d ser., III, 1904, 262, pl. 27, off Cape Flattery, Wash., in 77 fathoms.

Genus 1094. *XENOPYXIS* Gilbert.

Xenopyxis Gilbert, Proc. U. S. Nat. Mus., XLIII, 1915, 345 (*Xeneretmus latifrons* Gilbert).

3111. *Xenopyxis latifrons* (Gilbert).

North Pacific coast from Oregon to San Diego.

Xenochirus latifrons Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 92, off coast of Oregon and San Diego, at *Albatross* stations 2898, 2935, etc., in 61 and 158 fathoms.

3112. *Xenopyxis leiops* (Gilbert).

Off southern California.

Xeneretmus leiops Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 348, pl. 17, fig. 10, off Santa Catalina Island.

3113. *Xenopyxis ritteri* (Gilbert).

Off San Diego, Calif.

Xeneretmus ritteri Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 350, pl. 17, fig. 11, off San Diego, in 176 to 181 fathoms.

Genus 1095. *ODONTOPYXIS* Lockington.

Odontopyxis Lockington, Proc. U. S. Nat. Mus., II, 1879, 328 (*O. trispinosus* Lockington).

3114. *Odontopyxis trispinosus* Lockington.

Pacific coast, Alaska to Santa Barbara.

Odontopyxis trispinosus Lockington, Proc. U. S. Nat. Mus., II, 1879, 328, San Francisco.

Genus 1096. *BOTHRAGONUS* Gill.

Bothragonus Gill, in Jordan and Gilbert, Synopsis, 1883, 728 (*Hypsagonus swanii* Steindachner).

3115. *Bothragonus swanii* (Steindachner).

Puget Sound.

Hypsagonus swanii Steindachner, Ichth. Beitr., V, July, 1876, 144, pl. 4, Sitzb. Akad. Wiss. Wien, LXXIV, Port Townsend, Wash.

Family 241. ASPIDOPHOROIDIDÆ

Genus 1097. ULCINA Cramer.

Ulcina Cramer, in Jordan and Evermann, Check List, 1895, 449, 1896
(*Aspidophoroides olriki* Lütken).

3116. *Ulcina olriki* (Lütken).

Arctic Ocean; known from west coast of Greenland, Davis Strait, Kara Sea, Barrors Bay, and Nova Zembla.

Aspidophoroides olriki Lütken, Forelob. Meddel. om Nord Ulkefisiche Vidensk. Meddel. Naturhist. Foren. Kjøb., 1876, 386, 3 figs., Greenland.

3117. *Ulcina güntheri* (Bean).

Coast of Alaska.

Aspidophoroides güntheri Bean, Proc. U. S. Nat. Mus., VIII, 1885, 74, Alaska.

Genus 1098. ASPIDOPHOROIDES Lacépède.

Aspidophoroides Lacépède, Hist. Nat. Poiss., III, 1802, 228 (*A. tranquebar* Lacépède = *Cottus monopterygius* Bloch).

Canthirynchus Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 272 (*A. monopterygius* Bloch).

3118. *Aspidophoroides monopterygius* (Bloch). *Sea poacher*.

Greenland to Cape Cod; accidental to New York.

Cottus monopterygius Bloch, Ichth., II, 1786, 156, pl. 178, figs. 1, 2, Tranquebar; an error.

Cottus indicus Bonnaterre, Tableau Encycl. Méth., 1788, 68, pl. 87, fig. 367, India; after Bloch.

Aspidophoroides tranquebar Lacépède, Hist. Nat. Poiss., III, 1802, 228, Tranquebar.

Aspidophoroides grænlandicus Valenciennes, in Cuvier, Règne Anim., ed. 2, II, 1829, pl. 21, fig. 3, Greenland.

3119. *Aspidophoroides bartoni* Gilbert.

Aleutian Islands.

Aspidophoroides bartoni Gilbert, Rept. U. S. Fish Com., XIX, 1893, 434, Aleutian Islands and Bristol Bay, Alaska, at Albatross stations 3213, 3223, and many others, in 11½ to 121 fathoms.

Genus 1099. ANOPLAGONUS Gill.

Anoplagonus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 259 (*Aspidophoroides inermis* Günther).

3120. *Anoplagonus inermis* (Günther).

Coasts of Alaska, south to Vancouver Island.

Aspidophoroides inermis Günther, Cat., II, 1860, 524, Vancouver Island.

Family 242. CYCLOPTERIDÆ

Genus 1100. CYCLOPTERUS Linnæus.

Cyclopterus Linnæus, Syst. Nat., Ed. X, 1758, 260 (*C. lumpus* Linnæus). *Lumpus* Oken, Isis, 1817, 1782; based on "les Lumps" Cuvier (*Cyclopterus lumpus* Linnæus).

3121. *Cyclopterus lumpus* Linnæus. *Lumpfish; Cock and hen paddle; Sea owl; Paddle; Licorne de mer; Lumpsucker*.

North Atlantic; rocky shores of both coasts, south to Cape Cod and France.

Cyclopterus lumpus Linnæus, Syst. Nat., Ed. X, 1758, 260, Baltic and North Sea.

Cyclopterus minutus Pallas, Spicilegia, Zool., VII, 1769, 12, Atlantic Ocean.

Cyclopterus pavoninus Shaw, Nat. Hist., IX, 1797, pl. 310.

Cyclopterus pyramidatus Shaw, Gen. Zool., V, 1804, 390.

Cyclopterus caruleus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 480, New York Harbor.

Cyclopterus coronatus Couch, Nat. Hist. Cornwall, in Trans. Linn. Soc. London, XIV, 1823, 47, Cornwall.

Lumpus vulgaris McMURTRIE, Eng. ed. Cuvier, Anim. King., 1831; after
C. lumpus LINNÆUS.

Lumpus anglorum DeKay, N. Y. Fauna: Fishes, IV, 1842, 305, after
 LINNÆUS.

Genus 1101. EUMICRO TREMUS Gill.

Eumicrotremus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 190
 (*Cyclopterus spinosus* Müller).

3122. *Eumicrotremus spinosus* (Müller). *Spinous lumpfish.*

North Atlantic and Arctic Oceans, south to Maine and Denmark; and
 Eastport, Me.

Cyclopterus spinosus Müller, Prodr. Zool. Dan., IX, 1777, Denmark.

3123. *Eumicrotremus orbis* (Günther).

Aleutian Islands, south to Vancouver Island.

Cyclopterus orbis Günther, Cat., III, 1861, 158, Vancouver Island.

3124. *Eumicrotremus phrynoïdes* Burke.

Bering Sea.

Eumicrotremus phrynoïdes Burke, Bull. U. S. Bur. Fish., XXX, 1910,
 69, Petrel Bank, Alaska.

Genus 1102. LETHOTREMUS Gilbert.

Lethotremus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 449 (*L. muticus*
 Gilbert).

3125. *Lethotremus muticus* Gilbert.

Aleutian Islands.

Lethotremus muticus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 449,
 pl. 31, Unimak Pass, Alaska, at Albatross stations 3223 and 3258,
 in 56 to 70 fathoms.

3126. *Lethotremus vinolentus* Jordan and Starks.

Puget Sound, Wash.

Lethotremus vinolentus Jordan and Starks, Proc. Calif. Acad. Sci., 2d
 ser., V, 1895, 827, pl. 94, Puget Sound, near Seattle, Wash.

Genus 1103. CYCLOPTEROIDES Garman.

Cyclopterooides Garman, Mem. Mus. Comp. Zool., XIV, 1892, 37 (*C.
 gyrinops* Garman).

3127. *Cyclopterooides gyrinops* Garman.

Pribilof Islands, Alaska.

Cyclopterooides gyrinops Garman, Mem. Mus. Comp. Zool., XIV, 1892,
 37, St. Paul Island.

Family 243. LIPAROPIDÆ

Genus 1104. APTOCYCLUS De la Pylaie.

Aptocyclus De la Pylaie, Congrès Scientifique Potiers, 1834, 529 (*Cy-
 clopterus ventricosus* Pallas; misprinted *Cyclogasterus*).

Cycloptericthys Steindachner, Ichth., X, 1881, 14, 192 (*C.
 glaber* Steindachner = *Cyclopterus ventricosus* Pallas).

3128. *Aptocyclus ventricosus* (Pallas).

Kamchatka; Sea of Okhotsk.

Cyclopterus ventricosus Pallas, Spicilegia Zool., VII, 1770, 15, t. 2; no
 locality given, somewhere about Kamchatka.

Cycloptericthys glaber Steindachner, Ichth. Beitr., X, 1881, 14, pl. 8,
 Sea of Okhotsk.

Genus 1105. LIPAROPS Garman.

Liparops Garman, Mem. Mus. Comp. Zool., XV, 1892, 42 (*Cyclopterus
 stelleri* Pallas).

3129. *Liparops stelleri* (Pallas).

Peter and Paul Harbor, Kamchatka.

Cyclopterus stelleri Pallas, Zoogr. Rossj-Asiatica, III, 1811, 73, Peter
 and Paul Harbor, Kamchatka.

Family 244. LIPARIDÆ. Sea snails

Genus 1106. POLYPERA Burke.

Polypera Burke, Proc. U. S. Nat. Mus., XLIII, 1913, 567 (*Liparis greeni* Jordan and Starks).

3130. *Polypera greeni* (Jordan and Starks).

Puget Sound at Victoria, British Columbia.

Neoliparis greeni Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 829, pl. 96, figs. 2 and 3, Victoria, British Columbia.

Genus 1107. NEOLIPARIS Steindachner.

Neoliparis Steindachner, Ichth. Beitr., III, 1875, 54 (*Liparis mucosus* Ayres).

3131. *Neoliparis atlanticus* Jordan and Evermann. *Sea snail; Lumpsucker.*

Common along rocky shores from Newfoundland to Cape Cod.

Neoliparis atlanticus Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2107, Godbout, Quebec.

Liparis montagui of authors, not of Donovan, the European cognate.

3132. *Neoliparis mucosus* (Ayres).

Pacific coast from Puget Sound to San Francisco; rare.

Liparis mucosus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 22, San Francisco.

Neoliparis floræ Jordan and Starks, Proc. Calif. Acad. Sci., 2d. ser., V, 1895, 830, pl. 96, fig. 1, Waadda Island, Straits of Juan de Fuca.

3133. *Neoliparis fissuratus* Starks.

Puget Sound.

Neoliparis fissuratus Starks, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 560, Port Ludlow, Puget Sound.

3134. *Neoliparis rutteri* Gilbert and Snyder.

Kodiak Island, Alaska.

Neoliparis rutteri Gilbert and Snyder, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2108, Ugah Bay, Kodiak Island.

3135. *Neoliparis bristolensis* (Burke).

Bristol Bay, Bering Sea.

Cyclogaster bristolensis Burke, Proc. U. S. Nat. Mus., XLIII, 1913, 567, Bristol Bay in about 20 fathoms.

3136. *Neoliparis callyodon* (Pallas).

Coasts of Alaska, north to Kamchatka and Bering Sea; recorded from St. Paul, Kamchatka, Plover Bay, Siberia, Unalaska, and Kodiak.

Cyclopterus callyodon Pallas, Zoogr. Rosso-Asiat., III, 1811, 75, Kamchatka and Aleutian Islands.

3137. *Neoliparis grebnitzkii* Schmidt.

Commander Islands.

Neoliparis grebnitzkii Schmidt, Fauna Mare Orientale, 1875, 61, Bering Island.

3138. *Neoliparis beringianus* (Gilbert and Burke).

Bering Island.

Cyclogaster (*Neoliparis*) *beringianus* Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 72; Nikolski, Bering Island.

3139. *Neoliparis micraspidophorus* (Burke).

Bering Island.

Cyclogaster (*Neoliparis*) *micraspidophorus* Burke, Bull. U. S. Bur. Fish., XXX, 1910, 71, Nikolski, Bering Island.

Genus 1108. LIPARIS Scopoli.

Cyclogaster " Gronow, Act. Helvet., IV, 1760, 265 (*Cyclopterus liparis* Linnaeus); nonbinomial.

Cyclogasterus Gronow, Zoophyl., No. 198, 1763; nonbinomial.

Liparis (Artedi) Scopoli, Introd. Hist. Nat., 1777, 453 (*Cyclopterus liparis* Linnaeus).

Cyclogaster Gronow, Cat. Fishes, ed. Gray, 1854, 40 (*Cyclopterus liparis* L.).

ⁿ We do not find that Gronow used the name *Cyclogaster* as binomial. Although later than Linnaeus (1758), he had not then adopted, most likely had not heard of, the Linnaean system.

3140. *Liparis liparis* (Linnæus). *Sea snail; Striped sea snail.*

North Atlantic, on both shores, south to Connecticut and France, common north; very variable, a dozen color varieties having received names.

Cyclopterus liparis Linnæus, Syst. Nat., Ed. XII, I, 1766, 414, northern ocean; after Artedi and Gronow.

Cyclopterus lineatus Lepechin, Nov. Comm. Petropol., XVIII, 1774, 520, White Sea.

Cyclopterus liparis minor Fabricius, Fauna Grœnländica, 1780, 135, Greenland.

Cyclopterus musculus Lacépède, Hist. Nat. Poiss., IV, 1802, 683, near Havre.

3141. *Liparis osborni* Townsend and Nichols.

Off southern California in 440 fathoms.

Liparis osborni Townsend and Nichols, Bull. Amer. Mus. Nat. Hist., LII, 1925, 14, fig. 4, off southern California.

3142. *Liparis cyclopus* Günther.

Aleutian Islands to Puget Sound.

Liparis cyclopus Günther, Cat., III, 1861, 162, Esquimault Harbor, Vancouver Island.

3143. *Liparis megacephalus* (Burke).

Southeastern Bering Sea.

Cyclogaster megacephalus Burke, Proc. U. S. Nat. Mus., XLIII, 1913, 569, southeastern Bering Sea at 37 fathoms.

3144. *Liparis fucensis* Gilbert.

Straits of Fuca.

Liparis fucensis Gilbert, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 837, and Rept. U. S. Fish Com., XIX, 1893, 447, Port Angeles, Straits of Juan de Fuca.

Genus 1109. *CARELIPARIS* Garman.

Careliparis Garman, Mem. Mus. Comp. Zool., XIV, 1892, 62 (*Liparis agassizii* Putnam).

3145. *Careliparis tunicatus* (Reinhardt).

Greenland.

Liparis tunicata Reinhardt, Overs. Kong. Danske Vidensk. Selsk., VI, CXI, 1836, Greenland; after Fabricius.

Liparis fabricii Krøyer, Naturh. Tidsskr., II, 1847, 274, Greenland; D. XIII, 27; A. II, 30; P. 34; C. 9; Vert. 46.

Liparis arctica Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 191, Port Foulke, Greenland.

3146. *Careliparis agassizii* (Putnam).

Bering Sea; Sakhalin Island.

Liparis agassizii Putnam, Proc. Amer. Assoc. Adv. Sci., 1874, 339, Sakhalin Channel of Tartary.

Liparis gibbus Bean, Proc. U. S. Nat. Mus., IV, 1881, 148, Unalaska, St. Paul Island, Indian Point, Cape Chaplin, and Plover Bay, Siberia.

3147. *Careliparis herschelinus* (Scofield).

Arctic Ocean.

Liparis herschelinus Scofield, in Jordan and Evermann, Fishes North and Mid Amer., 1898, 2123, Herschel Island, Arctic Ocean, near mouth of the Mackenzie.

3148. *Careliparis dennysi* (Jordan and Starks).

North Pacific, from Aleutian Islands to Puget Sound.

Liparis dennysi Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 835, pl. 98, Admiralty Inlet near Seattle.

Genus 1110. *LYOLIPARIS* Jordan and Evermann.

Lyoliparis Jordan and Evermann, Check List, 1895, 451 (*Liparis pulchellus* Ayres).

3149. *Lyoliparis pulchellus* (Ayres).

Point Reyes, Calif.; northern Pacific, Alaska to Monterey.

Liparis pulchellus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 22, San Francisco.

Genus 1111. ACTINOCHIR Gill.

Actinochir Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 190, 193 (A. major Gill).

3150. *Actinochir major* Gill.

Coasts of Greenland.

Actinochir major Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 193, Greenland.

Cyclopterus liparis major Fabricius, Fauna Grœnland., 1780, 136, Greenland; major is here simply an adjective.

Genus 1112. CRYSTALLICHTHYS Jordan and Gilbert.

Crystallichthys Jordan and Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2864 (*C. mirabilis* Jordan and Gilbert).

3151. *Crystallichthys mirabilis* Jordan and Gilbert.

Bering Sea.

Crystallichthys mirabilis Jordan and Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2865, Provostmaya, Kamchatka.

3152. *Crystallichthys cyclospilus* Gilbert and Burke.

Bering Sea, Petrel Bank, St. Paul Island.

Crystallichthys cyclospilus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 74, Petrel Bank.

3153. *Crystallichthys cyclostigma* (Gilbert).

Bristol Bay, Alaska.

Liparis cyclostigma Gilbert, Rept. U. S. Fish Com., XIX, 1893, 446, Bristol Bay, Alaska, at Albatross station 3252, in 29½ fathoms.

Genus 1113. BATHYPHASMA Gilbert.

Bathypasma Gilbert, Rept. U. S. Fish Com., XIX, 1893, 447 (*B. ovigerum* Gilbert).

3154. *Bathypasma ovigerum* Gilbert.

Off Queen Charlotte Island, British Columbia.

Bathypasma ovigerum Gilbert, Rept. U. S. Fish Com., XIX, 1893, 448, Queen Charlotte Islands, at Albatross station 3342 in 1,588 fathoms.

Genus 1114. CAREMITRA Jordan and Evermann.

Caremitra Jordan and Evermann, Check List, 1895, 452 (*Careproctus simus* Gilbert).

3155. *Caremitra sima* (Gilbert).

Off Unalaska Island.

Careproctus simus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 444, off Unalaska, at Albatross station 3331 in 350 fathoms.

Genus 1115. LIPARISCUS Gilbert.

Lipariscus Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 358 (*L. nanus* Gilbert).

3156. *Lipariscus nanus* Gilbert.

Monterey Bay, Calif.

Lipariscus nanus Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 358, pl. 19, Monterey Bay in 32 to 309 fathoms.

Genus 1116. EURYMEM Gilbert and Burke.

Eurymen Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 64 (*E. gyrinus* Gilbert and Burke).

3157. *Eurymen gyrinus* Gilbert and Burke.

Bering Sea.

Eurymen gyrinus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 64, Avatcha Bay, east coast of Kamchatka.

Genus 1117. ACANTHOLIPARIS Gilbert and Burke.

Acantholiparis Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 83 (*A. opercularis* Gilbert and Burke).

3158. *Acantholiparis opercularis* Gilbert and Burke.

East coast of Kamchatka.

Acantholiparis opercularis Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 83, Starichkof Island.

Genus 1118. **NECTOLIPARIS** Gilbert and Burke.

Nectoliparis Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 82 (*N. pelagicus* Gilbert and Burke).

3159. **Nectoliparis pelagicus** Gilbert and Burke.

Bering Sea to Hokkaido, Japan; pelagic.

Nectoliparis pelagicus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 82, fig. 27, Bering Sea.

Genus 1119. **ECLASSODISCUS** Gilbert and Burke.

Elassodiscus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 81 (*E. tremebundus* Gilbert and Burke).

3160. **Elassodiscus tremebundus** Gilbert and Burke.

Coast of Kamchatka.

Elassodiscus tremebundus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 81, Avatcha Bay.

Genus 1120. **CAREPROCTUS** Krøyer.

Careproctus Krøyer, Naturh. Tidskr., I, 1862, 253-257 (*Liparis reinhardi* Krøyer).

Allochir Jordan and Evermann, Check List, 1895, 452 (*Careproctus melanurus* Gilbert).

Allurus Jordan and Evermann, loc. cit., 452 (*Careproctus ectenes* Gilbert).

Alinectes Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2866 (*Careproctus ectenes* Gilbert, substitute for *Allurus*, preoccupied).

Subgenus **CAREPROCTUS** Krøyer.

3161. **Careproctus reinhardi** (Krøyer).

Greenland, Jan Mayen, and Beeren Island.

Liparis reinhardi Krøyer, Naturh. Tidskr., I, 1862, 252, Greenland.

3162. **Careproctus colletti** Gilbert.

South of Alaska Peninsula to Sea of Okhotsk.

Careproctus colletti Gilbert, Rept. U. S. Fish Com., XIX, 1893, 442, south of Alaska Peninsula, at *Albatross* station 3338 in 625 fathoms.

3163. **Careproctus phasma** Gilbert.

Bristol Bay, Alaska.

Careproctus phasma Gilbert, Rept. U. S. Fish Com., XIX, 1893, 443, Bristol Bay, Alaska, at *Albatross* stations 3254 and 3256 in 46 and 49 fathoms.

3164. **Careproctus spectrum** Bean.

Between Unga and Nagai Islands, Alaska.

Careproctus spectrum Bean, Proc. U. S. Nat. Mus., XIII, 1890, 40, between Unga and Nagai Islands, 160° 18' W., 55° 10' N., at *Albatross* station 2848 in 110 fathoms.

3165. **Careproctus ranulus** (Goode and Bean).

Off Halifax Harbor.

Liparis ranula Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 46, Halifax.

3166. **Careproctus gelatinosus** (Pallas).

Peter and Paul Harbor, Kamchatka.

Cyclopterus gelatinosus Pallas, Spicilegia, VII, 1769, 19, Peter and Paul Harbor, Kamchatka.

3167. **Careproctus ostentum** Gilbert.

Unalaska Island.

Careproctus ostentum Gilbert, Rept. U. S. Fish Com., XIX, 1893, 444, north of Unalaska Island, at *Albatross* stations 3324 and 3331 in 109 and 350 fathoms.

3168. **Careproctus furcellus** Gilbert and Burke.

Bering Sea.

Careproctus furcellus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 80, between Petrel Bank and Agattu Island, Bering Sea.

3169. **Careproctus gibberti** Burke.

Southeastern Alaska and British Columbia.

Careproctus gibberti Burke, Proc. U. S. Nat. Mus., XLIII, 1913, 570, Kodiak Island and neighborhood in 102 to 482 fathoms.

3170. *Careproctus candidus* Gilbert and Burke.
Attu Island, Aleutian Chain.
Careproctus candidus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 77, fig. 22, East Cape, Attu Island, Alaska.
3171. *Careproctus mollis* Gilbert and Burke.
Attu Island, Aleutian Chain.
Careproctus mollis Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 77, fig. 21, East Cape, Attu Island, Alaska.
3172. *Careproctus bowersianus* Gilbert and Burke.
Bering Sea.
Careproctus bowersianus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 76, fig. 20, Bowers Bank, Bering Sea.
3173. *Careproctus opisthotremus* Gilbert and Burke.
Bering Sea.
Careproctus opisthotremus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 78, fig. 23, Petrel Bank, between Attu and Agattu Islands, Alaska.
3174. *Careproctus longifilis* Garman.
Off Galapagos Islands and Panama in deep water.
Careproctus longifilis Garman, Mem. Mus. Comp. Zool., XIV, 1892, 9, southwest Panama.
3175. *Careproctus longipinnis* Burke.
Arctic Ocean; Faroe Islands to Beeren Island.
Careproctus longipinnis Burke, Ann. Mag. Nat. Hist., 8th ser., IX, 1912, 510, Ingolf station 139, north of the Faroe Islands, depth 702 Danish fathoms.
3176. *Careproctus melanurus* Gilbert.
Off coast of California and Oregon.
Careproctus melanurus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 560, off coasts of California and Oregon, at Albatross stations 2840, 2891, and others in 178 to 339 fathoms.
3177. *Careproctus attenuatus* Gilbert and Burke.
Bering Sea.
Careproctus attenuatus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 79, Petrel Bank and Agattu Island, Alaska in 482 fathoms.
- Subgenus ALLINECTES Jordan and Evermann.
3178. *Careproctus ectenes* Gilbert.
Unalaska Island.
Careproctus ectenes Gilbert, Rept. U. S. Fish Com., XIX, 1893, 442, north of Unalaska, at Albatross station 3331 in 350 fathoms.
- Genus 1121. *PROGNURUS* Jordan and Evermann.
Prognurus Jordan and Evermann, Fishes North and Mid. Amer., 1896, 2866 (*P. cypselurus* Jordan and Gilbert).
3179. *Prognurus cypselurus* Jordan and Gilbert.
Bering Sea and north Pacific.
Prognurus cypselurus Jordan and Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2866, Bogoslof Island, Bering Sea.
- Genus 1122. *GYRINICHTHYS* Gilbert.
Gyrinichthys Gilbert, Rept. U. S. Fish Com., XIX, 1893, 444 (*G. minytremus* Gilbert).
3180. *Gyrinichthys minytremus* Gilbert.
Off Unalaska Island, Alaska.
Gyrinichthys minytremus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 444, north of Unalaska at Albatross station 3331 in 350 fathoms.

Genus 1123. PARALIPARIS⁷⁸ Collett.

Paraliparis Collett, Vid. Selsk. Forsk. Christiania, No. 14, 1878, 32
(*Paraliparis bathybius* Collett); species allied to *P. cephalus* with narrow gill openings.

Amitra Goode, Proc. U. S. Nat. Mus., III, 1880, 478 (*A. liparina* Goode).

Monomitra Goode, loc. cit., VI, 1883, 109 (*Amitra liparina* Goode); substitute for *Amitra* regarded as preoccupied by *Amitrus* Schönherr, a beetle.

Hilgendorfia Goode and Bean, Oceanic Ichthyol., 1895, 280 (*Paraliparis membranaceus* Günther).

Amitrichthys Jordan and Evermann, Check List, 1895, 453 (*Paraliparis cephalus* Gilbert).

3181. *Paraliparis holomelas* Gilbert.

North of Unalaska Island, Alaska.

Paraliparis holomelas Gilbert, Rept. U. S. Fish Com., XIX, 1893, 441, north of Unalaska, at *Albatross* stations 3308 and 3332 in 406 and 1,625 fathoms.

3182. *Paraliparis cephalus* Gilbert.

Alaska to California.

Paraliparis cephalus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 561, off California and Oregon, at *Albatross* station 2919 in 984 fathoms.

3183. *Paraliparis mento* Gilbert.

Coast of Oregon.

Paraliparis mento Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 562, off coast of Oregon, at *Albatross* station 3071 in 685 fathoms.

3184. *Paraliparis albescens* Gilbert.

Off Monterey Bay, Calif.

Paraliparis albescens Gilbert, Proc. U. S. Nat. Mus., XLVIII, 355, pl. 18, fig. 13, off Monterey Bay.

3185. *Paraliparis caudatus* Gilbert.

Monterey Bay, Calif.

Paraliparis caudatus Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 356, pl. 18, fig. 14, Monterey Bay.

3186. *Paraliparis deani* Burke.

Bering Sea; once taken in Monterey Bay.

Paraliparis deani Burke, Proc. U. S. Nat. Mus., XLIII, 1912, 571, Stevens Passage, Alaska.

3187. *Paraliparis dactylosus* Gilbert.

Off Santa Cruz, Calif.

Paraliparis dactylosus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 469, pl. 34, fig. 2, off Santa Cruz, Calif., at *Albatross* station 3112, in 296 fathoms.

3188. *Paraliparis grandiceps* Garman.

Off Lower California in deep water.

Paraliparis grandiceps Garman, Mem. Mus. Comp. Zool., XXIV, 1899, off Lower California.

3189. *Paraliparis rosaceus* Gilbert.

Pacific coast of Mexico.

Paraliparis rosaceus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 93, Pacific coast of Mexico, at *Albatross* stations 2898, 2935, and others in 61 to 158 fathoms.

3190. *Paraliparis fimbriatus* Garman.

Off Colombia and Galapagos Islands in deep water.

Paraliparis fimbriatus Garman, in Mem. Mus. Comp. Zool., XIV, 1892, 9, southwest of Panama.

3191. *Paraliparis attenuatus* Garman.

Off Bay of Panama in deep water.

Paraliparis attenuatus Garman, Deep Sea Fishes, Mem. Mus. Comp. Zool., XXIV, 1899, 118, pl. XXVII, off Bay of Panama.

⁷⁸This group of degenerate liparids is excessively abundant in all northern sea bottoms at considerable depth. It should, no doubt, be subdivided, but the present knowledge of the group hardly permits this.

3192. *Paraliparis angustifrons* Garman.
Off Bay of Panama in deep water.
Paraliparis angustifrons Garman, Mem. Mus. Comp. Zool. XXIV, 1899, 119, Pl. XXVII, *Albatross* station 3394, off Bay of Panama.
3193. *Paraliparis latifrons* Garman.
Off Bay of Panama in deep water.
Paraliparis latifrons Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 120, Pl. XXVII, off Bay of Panama.
3194. *Paraliparis garmani* Burke
Off coast of New England.
Paraliparis garmani Burke, Proc. U. S. Nat. Mus., XLIII, 1912, 572, off coast of New England in 328 to 542 fathoms.
3195. *Paraliparis copei* Goode and Bean.
Gulf Stream.
Paraliparis copei Goode and Bean, Oceanic Ichth., 1896, 279, fig. 353, Gulf Stream, at *Albatross* station 2232, 39° 12' 17" N., 72° 9' 30" W., in 520 fathoms.
Subgenus AMITRA Goode and Bean.
3196. *Paraliparis liparinus* (Goode).
Atlantic Ocean, off Rhode Island.
Amitra liparina Goode, Proc. U. S. Nat. Mus., III, 1880, 478, off Rhode Island.
Subgenus HILGENDORFIA Goode and Bean.
3197. *Paraliparis ulochir* Gilbert.
Gulf of California to Unalaska.
Paraliparis ulochir Gilbert, Rept. U. S. Fish Com., XIX, 1893, 441, Gulf of California, at *Albatross* station 3010 in 1005 fathoms.
- Genus 1124. RHINOLIPARIS Gilbert.
Rhinoliparis Gilbert, Rept. U. S. Fish Com., XIX, 1893, 445 (*R. barbulifer* Gilbert).
3198. *Rhinoliparis barbulifer* Gilbert.
Bering Sea.
Rhinoliparis barbulifer Gilbert, Rept. U. S. Fish Com., XIX, 1893, 445, north of Unalaska, Alaska, at *Albatross* stations 3237, 3325, and others, in 225 to 576 fathoms.
3199. *Rhinoliparis attenuatus* Burke.
Bering Sea.
Rhinoliparis attenuatus Burke, Proc. U. S. Nat. Mus., XLIII, 1912, 573, Bering Sea in 576 fathoms.

Family 245. PERISTEDIIDÆ

Genus 1125. PERISTEDION Lacépède.
Peristedion Lacépède, Hist. Nat. Poiss., III, 1802, 368 (*P. malarmat* Lacépède = *Trigla catachrycta* L.).
Peristethus Kaup, amended spelling.

3200. *Peristedion brevirostre* (Günther).
Caribbean Sea.
Peristethus brevirostre Günther, Cat., II, 1860, 217, West Indies.
3201. *Peristedion miniatum* Goode.
Gulf Stream.
Peristedion miniatum Goode, Proc. U. S. Nat. Mus., III, 1880, 349, Gulf Stream, off Rhode Island
3202. *Peristedion longispathum* (Goode and Bean).
West Indies, off Cuba and Barbados.
Peristedium longispatha Goode and Bean, Bull. Mus. Comp. Zool., XII, 1886, No. 5, 166, off Santa Cruz Island, West Indies, and off Barbados.
3203. *Peristedion gracile* Goode and Bean.
Gulf of Mexico and Porto Rico.
Peristedion gracile Goode and Bean, Oceanic Ichth., 1896, 473, Gulf of Mexico.

3204. *Peristedion platycephalum* (Goode and Bean).

West Indies; Barbados.

Peristedium platycephalum Goode and Bean, Bull. Mus. Comp. Zool., XII, 1886, No. 5, 167, off Barbados.

3205. *Peristedion barbiger* (Garman).

Off Bay of Panama in deep water.

Peristedium barbiger Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 110, off Bay of Panama.

3206. *Peristedion crustosum* (Garman).

Off Bay of Panama in deep water.

Peristedium crustosum Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 112, Pl. A, off Bay of Panama.

Genus 1126. *VULSICULUS* Jordan and Evermann.

Vulsiculus Jordan and Evermann, Check List, 1895, 489 (*Peristedion imberbe* Poey).

3207. *Vulsiculus imberbis* (Poey).

Gulf of Mexico.

Pterystedion imberbe Poey, Memorias, II, 1861, 367, Cuba; apparently never described.

Peristedion micronemus Poey, Ann. Lyc. Nat. Hist. N. Y., IX, 1870, 321, Cuba; substitute name on discovery of the minute barbels.

Family 246. TRIGLIDÆ

Genus 1127. *COLOTRIGLA* Gill.

Colotrigla Gill, Proc. U. S. Nat. Mus., XXVII, 1905, 240 (*Prionotus stearnsi* Jordan and Swain).

3208. *Colotrigla stearnsi* (Jordan and Swain).

Gulf of Mexico.

Prionotus stearnsi Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884, 541, Pensacola, Fla.

Genus 1128. *GURNARDUS* Jordan and Evermann.

Gurnardus Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2152 (*Prionotus gymnostethus* Gilbert).

3209. *Gurnardus birostratus* (Richardson).

Gulf of Fonseca, Pacific coast of Central America.

Prionotus birostratus Richardson, Voy. *Sulphur*, Ichth., pt. II, 1845, 81, April, Gulf of Fonseca, Pacific coast of Central America.

3210. *Gurnardus gymnostethus* (Gilbert).

Gulf of California.

Prionotus gymnostethus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 559, Gulf of California, in shallow water.

3211. *Gurnardus xenisma* (Jordan and Bollman).

Pacific Ocean, off coast of Colombia.

Prionotus xenisma Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 169, Pacific coast of Colombia.

3212. *Gurnardus loxias* Jordan.

Pacific coast of Central America.

Prionotus loxias Jordan, Proc. U. S. Nat. Mus., XIX, 1896, 452, Bay of Panama.

3213. *Gurnardus ruscarius* (Gilbert and Starks).

Panama Bay; Mazatlan; Magdalena Bay.

Prionotus ruscarius Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 165, Pl. XXVII, Panama Bay.

Genus 1129. *MERULINUS* Jordan and Evermann.

Merulinus Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2148 (*Trigla carolina* Linnaeus).

- 3214.** *Merulinus carolinus* (Linnæus). *Common gurnard; Red-winged sea robin; Pigfish; Rock pigfish; Flyingfish; Spinous gurnard; Small-scaled sea robin; Web-fingered sea robin; Carolina robin; Carolina sea robin.* Cape Ann to South Carolina, chiefly northward.
Trigla carolina Linnæus, Mantissa, II, 1771, 528, Carolina.
Trigla palmipes Mitchell, Trans. Lit. Phil. Soc. New York, 1815, 431, pl. 4, fig. 5, New York Harbor.
Prionotus pleatus Storer, Proc. Bost. Soc. Nat. Hist., II, 1845, 77, Massachusetts Bay.
- 3215.** *Merulinus scitulus* (Jordan and Gilbert).
 South Atlantic coast of United States, Beaufort to St. Augustine.
Prionotus scitulus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 288, Beaufort, N. C.
- 3216.** *Merulinus roseus* (Jordan and Evermann).
 Gulf of Mexico; Tampa Bay; Pensacola; Tortugas.
Prionotus roseus Jordan and Evermann, Proc. U. S. Nat. Mus., IX, 1886, 470, off Tampa Bay, Fla.
- 3217.** *Merulinus salmonicolor* Fowler.
 New River, Dade County, Fla.
Merulinus salmonicolor Fowler, Proc. Acad. Nat. Sci. Phila., LV, 1903, 333, New River, Fla.
- Genus 1180. PRIONOTUS Lacépède.** *Gurnards; Sea robins.*
Prionotus Lacépède, Hist. Nat. Poiss., III, 1802, 337 (*Trigla evolans* Linnæus).
Oreichthys Swainson, Classn. Fishes, Amphib., etc., 1839, 262 (*Trigla punctata* Bloch).
Chriolax Jordan and Gilbert, Proc. U. S. Nat. Mus., I, 1878, 374 (*Trigla evolans* Linnæus).
- 3218.** *Prionotus evolans* (Linnæus). *Southern striped gurnard; Lined sea robin; Flyingfish; Striped sea robin.*
 South Atlantic coast of United States.
Trigla evolans Linnæus, Syst. Nat., Ed. XII, 1766, 498, Carolinas.
Prionotus sarrtor Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 615, Charleston; Beaufort, N. C.
- 3219.** *Prionotus strigatus* (Cuvier). *Northern striped gurnard; Brown-winged sea robin; Sea robin; Sand pigfish; Pigfish; Flyingfish; Red-winged gurnard.*
 Atlantic coast of northern States, Cape Cod to Virginia; replacing *P. evolans* northward.
Trigla strigata Cuvier, Règne Anim., ed. 2, II, 1829, 161, New York; after *lineata* of Mitchell.
- 3220.** *Prionotus rubio* Jordan. *Rubio volador.*
 West Indies.
Prionotus rubio Jordan, Proc. U. S. Nat. Mus., IX, 1886, 50, Habana.
Rubio volador Parra, Descr. Dif. Piezas de Hist. Nat., 1787, 98, lam. 38, Habana.
- 3221.** *Prionotus punctatus* Bloch. *Spotted sea robin; Rubio volador.*
 West Indies and Caribbean Sea; Cameron, La.
Prionotus punctatus Bloch, Ichth., 1793, pl. 353, Martinique; on a drawing by Plumier.
- 3222.** *Prionotus ophryas* Jordan and Swain.
 Gulf of Mexico; Snapper Banks off Pensacola, Fla.
Prionotus ophryas Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884, 542, Snapper Banks off Pensacola, Fla.
- 3223.** *Prionotus pectoralis* (Nichols and Breder).
 Coast of Texas.
Prionotus miles pectoralis Nichols and Breder, Proc. Biol. Soc. Wash., XXXVII, 1924, 22, Gulf of Mexico, off Aransas Pass, Tex.
- 3224.** *Prionotus stephanophrys* Lockington.
 Deep water off San Francisco, Point Reyes, and Monterey; very rare.
Prionotus stephanophrys Lockington, Proc. U. S. Nat. Mus., III, 1880, 529, Point Reyes, near San Francisco.

3225. *Prionotus quiescens* Jordan and Bollman.
 Pacific Ocean off the coast of Colombia; Gulf of California.
Prionotus quiescens Jordan and Bollman, Proc. U. S. Nat. Mus., XII,
 1889, 166, off Pacific coast of Colombia.
3226. *Prionotus frontalis* Garman.
 Off Bay of Panama in deep water.
Prionotus frontalis Garman, Deep Sea Fishes, in Mem. Mus. Comp.
 Zool., XXIV, 1899, 108, off Bay of Panama.
3227. *Prionotus albirostris* Jordan and Bollman.
 Pacific Ocean off the coast of tropical America.
Prionotus albirostris Jordan and Bollman, Proc. U. S. Nat. Mus., XII,
 1889, 168, Pacific Ocean off the coast of Colombia.
3228. *Prionotus beanii* Goode.
 Off Trinidad.
Prionotus beanii Goode, in Goode and Bean, Oceanic Ichth., 1896, 468,
 off Trinidad.
3229. *Prionotus tribulus* Cuvier and Valenciennes. *Big-headed gurnard; Sea robin.*
 South Atlantic coast from Long Island to Brazos, Tex.
Prionotus tribulus Cuvier and Valenciennes, Hist. Nat. Poiss., IV, 1829,
 72 (98), pl. 74, New York; Carolinas.
? *Trigla carolina* Bloch, Ichth., 1793, 352, Carolinas; not of Linnaeus.
3230. *Prionotus horrens* Richardson.
 Pacific coast of tropical America.
Prionotus horrens Richardson, Voy. *Sulphur*, Ichth., 1843, 79, pl. 42,
 figs. 1-3, Gulf of Fonseca.
- Genus 1131. *FISSALA* Gill.
Fissala Gill, Proc. U. S. Nat. Mus., XXVIII, 1905, 342 (*Prionotus alatus* Goode and Bean).
3231. *Fissala alata* (Goode and Bean).
 Off Charleston, S. C.
Prionotus alatus Goode and Bean, Bull. Mus. Comp. Zool., X, 1883,
 210, deep sea off Charleston, S. C.
- Genus 1132. *BELLATOR* Jordan and Evermann.
Bellator Jordan and Evermann, Check. List, 1895, 488 (*Prionotus militaris* Goode and Bean).
3232. *Bellator militaris* (Goode and Bean).
 Off Cape Catoche, Yucatan; Gulf of Mexico.
Prionotus militaris Goode and Bean, Oceanic Ichth., 1896, 464, figs. 380
 and 384, off Cape Catoche, Yucatan.
3233. *Bellator egretta* (Goode and Bean).
 Barbados.
Prionotus egretta Goode and Bean, Oceanic Ichth., 1896, 465, fig. 381, off
 Barbados, at *Blake* station 64.

Family 247. DACTYLOPTERIDÆ

- Genus 1133. *DACTYLOPTERUS* Lacépède. *Flying gurnards.*
Cephalacanthus Lacépède, Hist. Nat. Poiss., III, 1802, 323 (*C. spinarella*
 Lacépède); young examples.
Dactylopterus Lacépède, loc. cit., 325 (*D. pirapeda* Lacépède = *Trigla volitans* Linnaeus); adult form.
Cephalacanthus Rafinesque, Analyse Nature, 1815, 85; substitute for
Cephalacanthus; too long a name.
Gonocephalus Gronow, Cat. Fish., ed. Gray, 1854, 106 (*G. macrocephalus*
 Gray = *Trigla volitans* Linnaeus).

- 3234. *Dactylopterus volitans* (Linnaeus).** *Flying robin; Batfish; Volador; Flying gurnard; Murcielago; Civetta de mer.*
 Atlantic Ocean on both coasts; also Gulf of Mexico.
Trigla volitans Linnaeus, Syst. Nat., Ed. X, I, 1758, 302; after Artedi.
Hirundo Catesby, Nat. Hist. Carol., II, 1771, Tab. 8, p. 8, Bahamas.
Trigla tentabunda Walbaum, Artedi Piscium, III, 1792, 362; after Klein.
Trigla fasciata Bloch and Schneider, Syst. Ichth., 1801, 16, tab. 3, fig. 1; after Klein.
Dactylopterus pirapeda Lacépède, Hist. Nat. Poiss., III, 1802, 326, Mediterranean and almost all warm seas.
Polygnemus securidatus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, pl. 4, fig. 10, New York.
Calionymus pelagicus Rafinesque, Amer. Month. Mag., 1818, 205, Atlantic Ocean.
Gonocephalus macrocephalus Gronow, Cat. Fish., ed. Gray, 1854, 106; pelagic.

Order SS. HOLCONOTI

Family 248. EMBIOTOCIDÆ. Surf-fishes; Viviparous perchés

Genus 1134. MICROMETRUS Gibbons.

Micrometrus Gibbons, Proc. Calif. Acad. Nat. Sci., in Daily Placer Times and Transcript, May 30, 1854 (*M. minimus* Gibbons).
Abeona Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 322 (*A. trowbridgii* Girard).

3235. *Micrometrus minimus* (Gibbons).

San Francisco to San Diego.

Cymatogaster minimus Gibbons, Proc. Acad. Nat. Sci. Phila., VII, 1854, 106, San Francisco Bay.
Holoconotus trowbridgei Girard, loc. cit., 152, no locality given.

Genus 1135. AMPHIGONOPTERUS Hubbs.

Amphigonopterus Hubbs, Proc. Biol. Soc. Wash., XXXI, May 16, 1918, 13 (*Abeona aurora* Jordan and Gilbert).

3236. *Amphigonopterus aurora* (Jordan and Gilbert).

Monterey Bay, Calif.

Abeona aurora Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 299, Monterey Bay, Calif.

Genus 1136. CYMATOGASTER Gibbons.

Cymatogaster Gibbons, Proc. Calif. Acad. Nat. Sci., in Daily Placer Times and Transcript, May 18, 1854 (*C. aggregatus* Gibbons).

Metrogaster Agassiz, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 128 (*Cymatogaster aggregatus* Gibbons).

Sema Jordan, Bull. U. S. Geol. Surv., IV, 1878, 399 (*S. signifer* Jordan; larvæ with very high vertical fins and closely compressed body).

3237. *Cymatogaster aggregatus* Gibbons. *Sparada; Viviparous perch; Perch.* Pacific coast from Port Wrangel, Alaska, to Todos Santos Bay.

Cymatogaster aggregatus Gibbons, Proc. Calif. Acad. Nat. Sci., in Daily Placer Times and Transcript, May 18, 1854, San Francisco.

Sema signifer Jordan, Bull. Hayden's Geol. Surv., IV, 1878, 399, Rio Grande, Tex.; error for San Francisco.

Metrogaster lineolatus Agassiz, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 129, no locality.

Genus 1137. BRACHYISTIUS Gill.

Brachyistius Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 275 (*B. frenatus* Gill).

3238. *Brachyistius frenatus* Gill.

Vancouver Island to San Diego.

Brachyistius frenatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 275, California coast.

Ditrema brevipinne Günther, Cat., IV, 1862, 248, Esquimault Harbor, Vancouver Island.

Genus 1138. ZALEMBIUS Jordan and Evermann.

Zalembius Jordan and Evermann, Fishes North and Mid. Amer., 1896, 1499 (*Cymatogaster rosaceus* Jordan and Gilbert).

3239. *Zalembius rosaceus* (Jordan and Gilbert).

San Francisco, Calif.

Cymatogaster rosaceus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 303, off San Francisco in rather deep water; the other species of this family live on sandy shores.

Genus 1139. HYPOCRITICHTHYS Gill.

Hypocritichthys Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 275 (*Hyperprosopon analis* Alexander Agassiz).

3240. *Hypocritichthys analis* (Agassiz).

San Francisco to Point Conception.

Hyperprosopon analis Alexander Agassiz, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 133, San Francisco.

Genus 1140. TOCICHTHYS Hubbs.

Tocichthys Hubbs, Proc. Biol. Soc. Wash., XXXI, May 16, 1918, 12 (*Hyperprosopon agassizii* Gill).

3241. *Tocichthys agassizii* (Gill).

Coast of California, San Francisco to Santa Barbara; most common along San Luis Obispo County.

Hyperprosopon agassizii Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 276, California.

Genus 1141. HYPERPROSOPON Gibbons.

Hyperprosopon Gibbons, Proc. Calif. Acad. Nat. Sci., in Daily Placer Times and Transcript, May 18, 1854 (*H. argenteum* Gibbons).

Eunichthys Girard, Proc. Acad. Nat. Sci. Phila., VII, 1855, 322 (*E. megalops* Girard).

Bramopsis Alexander Agassiz, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 132 (*B. mento* Agassiz = *Hyperprosopon argenteus* Girard).

3242. *Hyperprosopon argenteus* Gibbons. Wall-eyed surfish; White perch.

Coast of California, Cape Disappointment to Todos Santos Bay.

Hyperprosopon argenteum Gibbons, Proc. Acad. Nat. Sci. Phila., VII, 1854, 105, San Francisco.

Hyperprosopon argenteum punctatum Gibbons, loc. cit., 106, San Francisco.

Hyperprosopon arcuatum Gibbons, loc. cit., 125, San Francisco.

Holconotus megalops Girard, loc. cit., Presidio, Humboldt Bay; Astoria.

Bramopsis mento Agassiz, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 133,

no locality.

Genus 1142. HOLCONOTUS Agassiz.

Holconotus Agassiz, Amer. Jour. Sci. Arts., XVII, May, 1854, 367 (*H. rhodoterus* Agassiz).

Cymatogaster Gibbons, Proc. Calif. Acad. Nat. Sci., in Daily Placer Times and Transcript, June 21, 1854 (*C. pulchellus* Gibbons); not of Gibbons, May 18, 1854.

3243. *Holconotus rhodoterus* Agassiz..

Coast of California, San Francisco to San Diego.

Holconotus rhodoterus Agassiz, Amer. Jour. Sci. Arts., XVII, May, 1854, 368, San Francisco.

Cymatogaster pulchellus Gibbons, Proc. Acad. Nat. Sci. Phila., VII, July, 1854, 123, San Francisco.

Cymatogaster larkinsi Gibbons, loc. cit., 123, San Francisco.

Cymatogaster ellipticus Gibbons, loc. cit., 124, San Francisco.

Genus 1143. AMPHISTICHUS Agassiz.

Amphistichus Agassiz, Amer. Jour. Sci. Arts., May, 1854, 367 (*A. argenteus* Agassiz).

Mytilophagus Gibbons, Proc. Acad. Nat. Sci. Phila., VI, July, 1854, 125 (*M. fasciatus* Gibbons = *A. argenteus* Agassiz).

3244. *Amphistichus argenteus* Agassiz.

Pacific coast, from Cape Flattery to San Diego; very abundant on sandy shores.

Amphistichus argenteus Agassiz, Amer. Jour. Sci. Arts, XVII, May, 1854, 367, San Francisco.

Mytilophagus fasciatus Gibbons, Proc. Acad. Nat. Sci. Phila., VII, July, 1854, 125, San Francisco.

Amphistichus heermannii Girard, loc. cit., 135, Cape Flattery, San Francisco.

Amphistichus similis Girard, loc. cit., 153, San Francisco.

Genus 1144. *EMBIOTOMA* Agassiz.

Embiotoma Agassiz, Amer. Jour. Sci. Arts, XVI, 1853 (November), 388 (*E. jacksoni* Agassiz).

3245. *Embiotoma jacksoni* Agassiz. *Black surfish.*

Vancouver Island to Todos Santos Bay; abundant and variable.

Embiotoma jacksoni Agassiz, Amer. Jour. Sci. Arts, XVII, 1853, 387, and XVIII, 1854, 366, San Francisco.

Holconotus fuliginosus Gibbons, Proc. Acad. Nat. Sci. Phila., VII, 1854, 123, San Francisco.

Embiotoma cassidyi Girard, loc. cit., 151, San Diego.

?*Embiotoma webbii* Girard, loc. cit., VIII, 1855, 320, San Diego.

Genus 1145. *TÆNIOTOMA* Alexander Agassiz.

Holconotus Gibbons, Proc. Acad. Nat. Sci. Phila., VII, 1854 (July), 122 (*H. agassizi* Gibbons = *Embiotoma lateralis* Agassiz); not *Holconotus* Agassiz.

Tæniotoma Alexander Agassiz, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 133 (*Embiotoma lateralis* Agassiz).

3246. *Tæniotoma lateralis* (Agassiz). *Striped surfish.*

Vancouver Island to Todos Santos Bay; abundant southward.

Embiotoma lateralis Agassiz, Amer. Jour. Sci. Arts., XVIII, 1854 (May), 356, San Francisco.

Holconotus agassizii Gibbons, Proc. Acad. Nat. Sci. Phila., VII, 1854 (July), 122, San Francisco.

Embiotoma lineata Girard, loc. cit., (Aug.), 134, San Francisco, Tomales Bay, Presidio.

Embiotoma ornata Girard, loc. cit., VII, 1855, 321, San Diego, Calif.

Embiotoma perspicabilis Girard, loc. cit., 321, Puget Sound.

Genus 1146. *PHANERODON* Girard.

Phanerodon Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 153 (*P. furcatus* Girard).

3247. *Phanerodon furcatus* Girard. *White surfish.*

Pacific coast from Vancouver Island to Todos Santos Bay; abundant southward.

Phanerodon furcatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 153, Presidio, San Francisco, Calif.

3248. *Phanerodon atripes* (Jordan and Gilbert).

Monterey to San Diego; scarce.

Ditrema atripes Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 320, Monterey Bay, Calif.

Ditrema orthonotus Eigenmann and Eigenmann, West American Scientist, October, 1889, 127, Cortez Banks, off San Diego.

Genus 1147. *RHACOCHILUS* Agassiz.

Rhacochilus Agassiz, Amer. Jour. Sci. Arts, XVIII, 1854 (May), 367 (*R. toxotes* Agassiz).

Pachylabrus Gibbons, Proc. Calif. Acad. Sci., in Daily Placer Times and Transcript, June 21, 1854 (*P. variegatus* Gibbons = *R. toxotes* Agassiz).

3249. *Rhacochilus toxotes* Agassiz. *Bigmouth surfish; Alfione.*

Rhacochilus toxotes Agassiz, Amer. Jour. Sci. Arts, XVIII, 1854 (May), 367, San Francisco.

Pachylabrus variegatus Gibbons, Proc. Acad. Nat. Sci. Phila., VII, 1854 (July), 126, San Francisco.

Genus 1148. HYPSURUS Alexander Agassiz.

Hypsurus Alexander Agassiz, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 133 (*Embiotoca caryi* Agassiz).

3250. *Hypsurus caryi* (Agassiz). *Bugara*.

Cape Mendocino to San Diego; common northward.

Embiotoca caryi Agassiz, Amer. Jour. Sci. Arts, XVI, 1853, 389, and XVII, 1854, 366, San Francisco.

Holconotus gibbonsi (Gibbons MS., Proc. Calif. Acad. Sci., 1854), Gibbons, Proc. Acad. Nat. Sci. Phila., VII, 1854, 12, San Francisco

Genus 1149. DAMALICHTHYS Girard.

Damalichthys Girard, Proc. Acad. Nat. Sci. Phila., VII, 1855, 321 (*D. vacca* Girard = *D. argyrosoma* Girard).

3251. *Damalichthys argyrosomus* Girard.

Vancouver Island to Todos Santos Bay; very abundant northward.

Damalichthys argyrosoma Girard, Proc. Acad. Nat. Sci. Phila., VII, 1855, 136, San Francisco; young individual in bad condition.

Damalichthys vacca Girard, loc. cit., 321, Puget Sound.

Genus 1150. HYSTEROCARPUS Gibbons.

Hysteroxarpus Gibbons, Proc. Calif. Acad. Nat. Sci., in Daily Placer Times and Transcript, May 18, 1854, and in Proc. Acad. Nat. Sci. Phila., VII, 1854, 124 (*H. traski* Gibbons).

Sargosomus Agassiz, in Alexander Agassiz, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 128 (*S. fluvialis* Agassiz = *Hysteroxarpus traski* Gibbons).

Dacentrus Jordan, Bull. U. S. Geol. Surv., IV, 1878, 667 (*D. lucens* Jordan; very young).

3252. *Hysteroxarpus traski* Gibbons.

Rivers of central California, chiefly in the Sacramento-San Joaquin Valley from Lake County to Santa Clara County.

Hysteroxarpus traski Gibbons, Proc. Acad. Nat. Sci., Phila., VII, 1854, 105, lower Sacramento River.

Sargosomus fluvialis Agassiz, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 130, Sacramento.

Decentrus lucens Jordan, Bull. U. S. Geol. Surv., IV, 1878, 667, "Rio Grande, Tex." locality erroneous, should be San Francisco.

Order TT. CHROMIDES

Family 249. POMACENTRIDÆ

Genus 1151. AZURINA Jordan and McGregor.

Azurina Jordan and McGregor, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1544 (*A. hirundo* Jordan and McGregor).

3253. *Azurina hirundo* Jordan and McGregor.

Pacific coast of Mexico.

Azurina hirundo Jordan and McGregor, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1544, Guacalupe Island, off Lower California.

Genus 1152. FURCRIA Poey.

Furcaria Poey, Memorias, II, 1860, 194 (*F. puncta* Poey = *Heliases multilineatus* Guichenot in Ramon de la Sagra, Hist. fisica Cuba, IV).

3254. *Furcaria atrilobata* (Gill).

Pacific coast of America, from Cape San Lucas southward.

Chromis (*Furcaria*) *atrilobata* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 149, Cape San Lucas, Lower California.

3255. *Furcaria cyanea* Poey.

Cuba.

Furcaria cyanea Poey, Memorias, II, 1860, 196, Cuba.

3256. *Furcaria multilineata* (Guichenot).

Cuba.

Heliases multilineatus Guichenot, in Ramon de la Sagra, Poiss. Cuba, 1853, 76, pl. 2, fig. 2, Habana.

Furcaria puncta Poey, Memorias, II, 1860, 195, Cuba.

3257. *Furcaria bermudæ* (Nichols).
Bermuda.
Chromis bermudæ Nichols, Proc. Biol. Soc. Wash., XXXIII, 1920, 60,
Bermuda.
- Genus 1153. *HELIASES* Cuvier and Valenciennes.
Heliases Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 369 (495)
(H. insolatus Cuvier and Valenciennes).
Heliases Günther, revised spelling.
3258. *Heliases insolatus* Cuvier and Valenciennes. *Chauffe soleil.*
West Indies, north to Pensacola.
Heliases insolatus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830,
 370 (494), pl. 137, Martinique.
3259. *Heliases enchyrsurus* (Jordan and Gilbert).
Snapper Banks off Pensacola and Tampa, Fla.
Chromis enchyrsurus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882,
 286, Pensacola, Fla.
3260. *Heliases marginatus* Castelnau.
Brazil; St. Croix Island, West Indies.
Heliases marginatus Castelnau, Anim. Amer. Sud, Poiss., 1855, 9, Pl.
 III, fig. 1, Bahia.
- Genus 1154. *AYRESIA*[®] Cooper.
Ayresia Cooper, Proc. Calif. Acad. Nat. Sci., 1st ser., III, 1803, 73 (*A. punctipinnis* Cooper).
3261. *Ayresia punctipinnis* Cooper. *Blacksmith.*
Coast of California, from Point Conception to Cedros Island.
Ayresia punctipinnis Cooper, Proc. Calif. Acad. Nat. Sci., 1st ser., III,
 1863, 73, San Diego Bay, San Pedro, and Santa Barbara.
- Genus 1155. *EUPOMACENTRUS*[®] Bleeker.
Eupomacentrus Bleeker, Nat. Verh. Holl. Maats., Wetensk., II, 1877,
 73 (*Pomacentrus lividus* Bleeker); teeth in one row, truncate; snout
 scaly; lower jaw naked; dorsal not notched.
Amblypomacentrus Bleeker, loc. cit., 68 (*Pomacentrus breviceps*); similar,
 snout naked.
3262. *Eupomacentrus fuscus* (Cuvier and Valenciennes). *Maria molle.*
West Indies and coast of Brazil north to Key West.
Pomacentrus fuscus Cuvier and Valenciennes, Hist. Nat. Poiss., V,
 1830, 324 (432), Brazil.
Pomacentrus variabilis Castelnau, Anim. Nouv. ou Rares, Poiss., 1855,
 9, pl. 3, fig. 3, Bahia.
Pomacentrus atro-cyanus Poey, Memorias, II, 1861, 190, Cuba.
3263. *Eupomacentrus diencaeus* Jordan and Rutter.
Jamaica.
Eupomacentrus diencaeus Jordan and Rutter, Proc. Acad. Nat. Sci.
 Phila., XLIX, 1897, 116, Jamaica.
3264. *Eupomacentrus leucostictus* (Müller and Troschel). *Cockeye pilot;*
Beau Gregoire; Black pilot.
*West Indies to Snapper Banks, west Florida; accidental north to Woods
 Hole, Mass.*
Pomacentrus leucostictus Müller and Troschel, in Schomburgk's Hist.
 Barbados, 1848, 674, Barbados.
Pomacentrus xanthurus Poey, Memorias, II, 1861, 190, Cuba.
Pomacentrus caudalis Poey, Synopsis, 1867, 328, Cuba.
Pomacentrus dorso-punicans Poey, loc. cit., 328, Cuba.
3265. *Eupomacentrus analis* (Poey).
Cuba, Key West, and Porto Rico.
Pomacentrus analis Poey, Synopsis, 1867, 327, Cuba.

[®] This genus is near the Mediterranean genus *Chromis* Cuvier, based on *Sparus chromis* Linnaeus, a species with 14 dorsal spines, the head high and narrow, the tail short, the caudal not deeply forked. *Furcaria* and *Heliases* are also closely related and usually the four groups have been united into one as *Chromis* or "Hellastes."

[®] In the closely related old world genus *Pomacentrus* the teeth are in two rows. Several of the species here accepted are of very doubtful validity, and the genera of Bleeker derived from the subdivisions of *Pomacentrus* need further study.

3266. *Eupomacentrus otophorus* (Poey).
Cuba.
Pomacentrus otophorus Poey, Memorias, II, 1861, 188, Cuba.
3267. *Eupomacentrus chrysus* Bean.
Bermudas.
Eupomacentrus chrysus Bean, Proc. Biol. Soc. Wash., XIX, 1906, 32, Whites Flat Channel, Bermudas.
3268. *Eupomacentrus flaviventer* (Troschel).
Atlantic coast of Mexico.
Pomacentrus flaviventer Troschel, in J. W. von Müller's Reisen in den Vereinigten Staaten, Canada, und Mexico, III, 1865, 633, Atlantic Ocean.
3269. *Eupomacentrus gillii* (Gilbert and Starks).
Panama.
Pomacentrus gillii Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 141, pl. XXII, fig. 44, Panama.
3270. *Eupomacentrus partitus* (Poey).
Cuba.
Pomacentrus partitus Poey, Synopsis, 1867, 327, Cuba.
3271. *Eupomacentrus planifrons* (Cuvier and Valenciennes). *Petite jaquette*.
Tortugas; West Indies, Jamaica to Martinique.
Pomacentrus planifrons Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 323 (431), Martinique.
3272. *Eupomacentrus leucorus* (Gilbert).
Socorro Island, one of the Revillagigedo group, off Pacific coast of Mexico.
Pomacentrus leucorus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 554, Socorro Island.
3273. *Eupomacentrus adustus* (Troschel).
Cuba.
Pomacentrus adustus Troschel, in J. W. von Müller's Reisen in den Vereinigten Staaten, Canada und Mexico, III, 1865, 633, Mexico.
Pomacentrus obscuratus Poey, Enumeratio, 1876, 101, Habana.
3274. *Eupomacentrus rectifrenum* (Gill). *Pescado azul de dos colores*.
Pacific coast of America from Cape San Lucas to Panama; common.
Pomacentrus rectifrenum Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 148, Cape San Lucas, Lower California.
Pomacentrus analigutta Günther, Cat., IV, 1862, 27, Cape San Lucas.
3275. *Eupomacentrus flavilatus* (Gill). *Pescado azul de dos colores*.
Pacific coast of America, from Cape San Lucas to Mazatlan and beyond.
Pomacentrus flavilatus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 148, Cape San Lucas, Lower California.
3276. *Eupomacentrus nepenthe* Nichols.
Bahamas.
Eupomacentrus nepenthe Nichols, Amer. Mus. Novitates, XXVI, 1921, 1, fig. 1, Bahamas.
- Genus 1156. **NEXILARIUS** Gilbert.
Nexilarius Gilbert, in Jordan and Evermann, Check List, 1895, 512 (*Euschistodus concolor* Gill); misprinted *Nexilaris*.
3277. *Nexilarius concolor* (Gill).
Panama.
Euschistodus concolor Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 145, Panama.
- Genus 1157. **ABUDEFDUF**⁸¹ Forskål. *Pintanos*.
Abudefduf Forskål, Deser. Anim., etc., 1775, 59 (*Chætodon sordidus* Forskål).
Glyphtodon Lacépède, Hist. Nat. Poiss., IV, 1803, 542 (*Chætodon saxatilis* Linnæus).
Euschistodus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 145 (*E. declivifrons* Gill).

⁸¹ The small brilliantly colored species of coral reefs in the South Seas constituting the genus *Chrysiptera* Swainson are probably separable from *Abudefduf*.

- 3278. Abudedefduf marginatus** (Bloch). *Pintano; Cow pilot; Jaqueta; Mojarra raiado; Demoiselle.*
 Atlantic coast of tropical America, West Indies, to Brazil; once reported from Newport, R. I.
Jaguacanguare Marçgrave, Hist. Brazil, 1648, 156, Brazil.
Chætodon marginatus Bloch, Ichthyol., III, 1787, 98, Martinique; after Plumier.
Chætodon sargoides Lacépède, Hist. Nat. Poiss., IV, 1803, 452, Martinique.
Glyphidodon moucharra Lacépède, loc. cit., 542, Brazil.
Glyphidodon saxatilis of authors, not of Linnaeus, an Asiatic species.
?Chætodon mauritii Bloch, Naturgesch. Ausl. Fische, III, 1787, 109, Brazil, on a drawing by Prince Maurice. A doubtful species, apparently the same as *A. marginatus*.
- 3279. Abudedefduf troschelii** (Gill).
 Pacific coast of Mexico; Cape San Lucas; Socorro Island; Cocos Islands; Galapagos.
Glyphidodon troschelii Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 150, Cape San Lucas.
- 3280. Abudedefduf declivifrons** (Gill).
 Pacific coast of tropical America, from Cape San Lucas southward.
Euschiistodus declivifrons Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 145, 146, Cape San Lucas.
- 3281. Abudedefduf analogus** (Gill).
 Atlantic coast of Central America.
Euschiistodus analogus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 220, Aspinwall.
- 3282. Abudedefduf taurus** (Müller and Troschel). *Dovetail fish.*
 Tortugas; Barbados.
Glyphidodon taurus Müller and Troschel, in Schomburgk, Hist. Barbados, 1848, 674, Barbados.
- Genus 1158. CENTROCHROMIS** Norman.
Centrochromis Norman, Ann. Mag. Nat. Hist., ser. 9, IX, 1922, 534 (*Glyphidodon rufus* Poey.)
- 3283. Centrochromis rufus** (Poey).
 Cuba.
Glyphidodon rufus Poey, Memorias, II, 1861, 191, Cuba.
- Genus 1159. HYPSEPOPS** Gill. *Garibaldi.*
Hypsopops Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 165 (*Glyphidodon rubicundus* Girard).
- 3284. Hypsopops rubicundus** (Girard). *Garibaldi; Sea goldfish;* a magnificent and hardy aquarium fish.
 Coast of California south of Monterey; abundant about Santa Catalina Island.
Glyphidodon rubicundus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 148, Monterey, Calif.
- Genus 1160. STEGASTES** Jenyns.
Stegastes Jenyns, Voy. Beagle, Fishes, 1842, 63 (*S. imbricatus* Jenyns).
Microspathodon Günther, Cat., IV, 1862, 35 (*Glyphidodon chrysurus* Cuvier and Valenciennes).
Pomataprion Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 214, 216 (*Hypsopope dorsalis* Gill).
- 3285. Stegastes chrysurus** (Cuvier and Valenciennes).
 Tortugas; West Indies; Cuba and St. Thomas.
Glyphidodon chrysurus Cuvier and Valenciennes, Hist. Nat. Poiss., V, 1830, 356 (476), St. Thomas.
Pomacentrus denegatus Poey, Memorias, II, 1861, 190, Cuba.
- 3286. Stegastes bairdii** (Gill).
 Pacific coast of tropical America, from Cape San Lucas to Panama.
Pomacentrus bairdii Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 149, Cape San Lucas.

3287. *Stegastes niveatus* (Poey).*Cuba.**Pomacentrus niveatus* Poey, *Enumeratio*, 1876, 102, Habana.3288. *Stegastes dorsalis* (Gill).

Pacific coast of America, from Cape San Lucas to Mazatlan.

Hypsypops dorsalis Gill, *Proc. Acad. Nat. Sci. Phila.* XIV, 1862, 147, Cape San Lucas; young.*Pomacentrus quadriguttata* Gill, *loc. cit.*, 149, Cape San Lucas.*Microspathodon cinereus* Gilbert, *Proc. U. S. Nat. Mus.*, XIII, 1890, 71, Clarion and Socorro Islands.*Microspathodon cururissimus* Jordan and Starks, in *Jordan*, *Proc. Calif. Acad. Sci.*, 2d ser., V, 1895, 478, pl. 44, Venados Islands, near Mazatlan; adult.Family 250. CICHLIDÆ.⁸² Mojarras de rioGenus 1161. *CICHLAURUS* Swainson.*Cichlaurus* Swainson, *Nat. Hist. Classn. Fishes, etc.*, II, 1839, 173, definition, but no type named; this is given on p. 230, where, without assigned reason, the name is changed to *Cichlasoma*.*Cichlasoma* Swainson, *Nat. Hist., etc.*, *loc. cit.*, 230 (*Labrus punctatus* Bloch).*Heros* Heckel, *Natterers Reise*, II, 1840, 362 (*Heros severus* Heckel).3289. *Cichlaurus bifasciatus* (Steindachner).*Mexico.**Heros bifasciatus* Steindachner, *Denks. Ak. Wiss. Wien.*, XXIII, 1864, 60, pl. II, Mexico.3290. *Cichlaurus angulifer* (Günther).*East slope of Guatemala.**Heros angulifer* Günther, *Cat.*, IV, 1862, 298, River de Santa Isabel, Guatemala.3291. *Cichlaurus microphthalmus* (Günther).*Rio Motagua, Guatemala.**Heros microphthalmus* Günther, *Cat.*, IV, 1862, 295, Rio Motagua, Guatemala.3292. *Cichlaurus guttulatus* (Günther).*Pacific drainage of Guatemala.**Heros guttulatus* Günther, *Proc. Zool. Soc. Lond.*, 1864, 152, Lake Amatitlan, Guatemala.3293. *Cichlaurus meeki* (Hildebrand). *Mojarra; Mojarra negra; Mojarra plateada.**Lakes and rivers of El Salvador, Pacific slope of Central America.**Cichlasoma meeki* Hildebrand, *Bull. U. S. Bur. Fish.*, XLI, 1925, 275, fig. 20, Lake Guiga, El Salvador.3294. *Cichlaurus melanurus* (Günther).*Lake Peten, Guatemala.**Heros melanurus* Günther, *Cat.*, IV, 1862, 288, Lake Peten, Guatemala.3295. *Cichlaurus fenestratum* (Günther).*Atlantic slope of southern Mexico.**Chromis fenestrata* Günther, *Proc. Zool. Soc. Lond.*, 1860, 318, Mexico.3296. *Cichlaurus godmanni* (Günther).*Atlantic slope of Guatemala.**Heros godmanni* Günther, *Cat.*, IV, 1862, 296, Rio Cahabon, Guatemala.3297. *Cichlaurus intermedius* (Günther).*Lake Peten, Guatemala.**Heros intermedius* Günther, *Cat.*, IV, 1862, 298, Lake Peten, Guatemala.3298. *Cichlaurus deppii* (Heckel).*Mexico.**Heros deppii* Heckel, *Ann. Wiener Mus.*, II, 1840, 382, Mexico.

⁸² The many species of this genus in the Mexican region are chiefly the overflow from the vast Brazilian fauna. The larger genera as here recognized are very closely related and were recently regarded as subgenera of *Cichlasoma* (*Cichlaurus*) = *Heros* Heckel; several of the species are only provisionally located and a few (*godmani*, *hedricki*, *lentigonosus*) may represent new groups not yet defined.

3299. *Cichlaurus calobrensis* (Meek and Hildebrand).
 Streams of Panama.
Cichlasoma calobrense Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 90, Rio Calobra, Panama.
3300. *Cichlaurus umbrifer* (Meek and Hildebrand).
 Rivers of Panama.
Cichlasoma umbriferum Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1913, 88, Rio Cupe, Cituro, Panama.
- Genus 1162. *THERAPS* Günther.
Theraps Günther, Cat., IV, 1862, 284 (*T. irregularis* Günther).
? *Baiodon* Agassiz, Proc. Bost. Soc. Nat. Hist., VI, 1858, 401 (*B. fasciatus* Agassiz); scanty description; said to be slender as in the Sparoid genus *Boops* or *Box*.
3301. *Theraps irregularis* Günther.
 Atlantic slope of Guatemala and southern Mexico.
Theraps irregularis Günther, Cat., IV, 1862, 284, Guatemala.
3302. *Theraps mojarra* (Meek).
 Pacific slope of Isthmus of Tehuantepec.
Cichlasoma mojarra Meek, Field Mus., Zool., V, 1904, 93, 217, San Geronimo, Oaxaca, Mexico.
3303. *Theraps nebulifer* (Günther).
 Atlantic lowland streams of the Isthmus of Tehuantepec.
Heros nebulifer Günther, Proc. Zool. Soc. Lond., 1860, 318, Mexico.
3304. *Theraps gibbiceps* (Steindachner).
 Rivers of Tabasco, Mexico.
Heros gibbiceps Steindachner, Denks. Ak. Wiss. Wien, XXIII, 1864, 66, Pl. V, figs. 1 and 2, Rio Teapa, Tabasco, Mexico.
3305. *Theraps zonatus* (Meek).
 Pacific slope of Isthmus of Tehuantepec, Mexico.
Cichlasoma zonatum Meek, Proc. Biol. Soc. Wash., XVIII, 1905, 245, Niltepec, Oaxaca, Mexico.
Baiodon fasciatus Agassiz, Proc. Bost. Soc. Nat. Hist., VI, 1858, 408, Lake Nicaragua.
3306. *Theraps eigenmanni* (Meek).
 Rio Tehuacan, Mexico.
Cichlasoma eigenmanni Meek, Field Col. Mus., Zool., III, 1902, 119, Rio Tehuacan, Mexico.
3307. *Theraps teapæ* (Evermann and Goldsborough).
 Rio Teapa, Mexico.
Cichlasoma teapæ Evermann and Goldsborough, Bull. U. S. Fish Com., XXI, 1901, 156, fig. 6, Rio Teapa, Mexico.
3308. *Theraps frontalis* (Meek).
 Turubales, Costa Rica.
Cichlasoma frontale Meek, Field Col. Mus., Zool., VIII, 1909, 210, Turubales, Costa Rica.
3309. *Theraps punctatum* (Meek).
 Buenos Aires de Terraba, Costa Rica.
Cichlasoma punctatum Meek, Field Mus. Nat. Hist., Zool., VII, 1909, 210, Buenos Aires de Terraba, Costa Rica; name preoccupied in *Cichlasoma* but not in *Theraps*.
Theraps terrabæ Jordan and Evermann, Proc. Calif. Acad. Sci., 4th ser., XVI, Apr. 27, 1927, 507, substitute for *Cichlasoma punctatum* Meek, preoccupied.
3310. *Theraps gadovii* (Regan).
 Atlantic slope of southern Mexico.
Cichlosoma gadovii Regan, Ann. Mag. Nat. Hist., 1905, 232, Motozongo, Mexico.
3311. *Theraps milleri* (Meek).
 El Rancho, Guatemala.
Cichlasoma milleri Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 142, El Rancho, Guatemala.

- 3312.** *Theraps balteatus* (Gill and Bransford).
 Great Lakes of Nicaragua.
Heros balteatus Gill and Bransford, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 184, Lake Nicaragua.
- 3313.** *Theraps nicaraguensis* (Günther).
 Great Lakes of Nicaragua.
Heros nicaraguensis Günther, Proc. Zool. Soc. Lond., 1864, 153, Lake Nicaragua.
- 3314.** *Theraps nigritus* (Meek).
 Lake Nicaragua.
Cichlasoma nigratum Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 128, Lake Nicaragua.
- 3315.** *Theraps lentiginosus* (Steindachner).
 Rivers of Mexico.
Heros lentiginosus Steindachner, Denks. Akad. Wiss. Wien, XXIII, 1864, 62, Pl. III, fig. 1, Mexico.
- Genus 1163. PARAPETENIA Regan.**
Hypsophrys Agassiz, Proc., Bost. Soc. Nat. Hist., VI, 1858, 401 (*H. unimaculatus* Agassiz); no description except that it resembles *Chrysophrys* (*Sparus*) and has a large round black spot on the shoulder; lakes of Nicaragua.
?Parachromis Agassiz, Proc., Bost. Soc. Nat. Hist., VI, 1858, 401 (*P. gulosus* Agassiz); not described but said to resemble *Dentex*; lakes of Nicaragua.
Parapetenia Regan, Ann. Mag. Nat. Hist., XVI, 7th ser., 1905, 324 (*Acara adspersum* Günther).
- 3316.** *Parapetenia adspersa* (Günther).
 Barbados.
Acara adspersum Günther, Cat., IV, 1862, 282, Barbados.
- 3317.** *Parapetenia dovii* (Günther).
 Great Lakes of Nicaragua.
Heros dovii Günther, Proc. Zool. Soc. Lond., 1864, 154, Lake Nicaragua.
- 3318.** *Parapetenia managuensis* (Günther).
 Great Lake of Nicaragua.
Heros managuensis Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 392, 463, pl. 77, fig. 3, Lake Managua, Nicaragua.
- 3319.** *Parapetenia motaguensis* (Günther).
 Atlantic slope of Guatemala.
Heros motaguensis Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 391, 462, pl. 77, fig. 2, Lake Motagua, Guatemala.
- 3320.** *Parapetenia friedrichstahli* (Heckel).
 Lake Peten, Guatemala.
Heros friedrichstahli Heckel, Ann. Mus. Wien, II, 1840, 381, Lake Peten, Guatemala.
- 3321.** *Parapetenia multifasciata* (Regan).
 Guatemala; British Honduras.
Cichlasoma multifasciatum Regan, Ann. Mag. Nat. Hist., 7th ser., XVI, 1905, 335, Lake Peten, Guatemala.
- 3322.** *Parapetenia salvini* (Günther).
 Atlantic slope of southern Mexico to Guatemala.
Heros salvini Günther, Cat., IV, 1862, 294, Rio de Santa Isabel, Guatemala.
Heros triogramma Steindachner, Denkschr. Akad. Wiss. Wien, XXIII, 1864, 70, Taf. 3, fig. 2, central Mexico.
- 3323.** *Parapetenia urophthalma* (Günther).
 Atlantic slope of southern Mexico to Guatemala; British Honduras and Yucatan.
Heros urophthalmus Günther, Cat., IV, 1862, 291, Lake Peten.
- 3324.** *Parapetenia istlana* (Jordan and Snyder).
 Rio Balsas, Mexico.
Heros istlanus Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 144, fig. 21, Rio Ixtla at Puente de Ixtla, Mexico.

3325. *Parapetenia tetracantha* (Cuvier and Valenciennes). *Viajaca*.
 Rivers of Cuba.
Centrarchus tetracanthus Cuvier and Valenciennes, Hist. Nat. Poiss., VII, 1831, 346 (460), Habana.
Chromis fusco-maculatus Guichenot, in Ramon de la Sagra, Hist. Ile Cuba, Poiss., 1853, 78, pl. 2, fig. 3, Habana.
3326. *Parapetenia tetracantha torralbasi* (Eigenmann).
 Cuba; Calabazar, Almendares River; San Cristobal; Los Palacios.
Heros tetracanthus torralbasi Eigenmann, Bull. U. S. Fish Com., XXII, 1902, 230, fig. 11, Calabaza, Cuba.
3327. *Parapetenia tetracantha grisea* (Eigenmann).
 San Antonio, Cuba.
Heros tetracanthus griseus Eigenmann, Bull. U. S. Fish Com., XXII, 1902, 233, San Antonio, Cuba.
3328. *Parapetenia tetracantha lata* (Eigenmann).
 San Juan, Cuba.
Heros tetracanthus latus Eigenmann, Bull. U. S. Fish Com., XXII, 1902, 234, San Juan, Cuba.
3329. *Parapetenia tetracantha cincta* (Eigenmann).
 Paso Real, Cuba.
Heros tetracanthus cinctus Eigenmann, Bull. U. S. Fish Com., XXII, 1902, 234, fig. 16, Paso Real, Cuba.
3330. *Parapetenia parma* (Günther).
 Mexico and Guatemala.
Heros parma Günther, Cat., IV, 1862, 285, Mexico and Guatemala.
3331. *Parapetenia nigricans* (Eigenmann).
 Pinar del Rio, Cuba.
Heros nigricans Eigenmann, Bull. U. S. Fish Com., XXII, 1902, 235, fig. 17, Pinar del Rio, Cuba.
3332. *Parapetenia steindachneri* (Jordan and Snyder).
 Lower portion of the basin of the Rio Panuco and Rio Verde, Mexico.
Cichlasoma steindachneri Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 143, fig. 20, Rio Verde, Mexico.
3333. *Parapetenia bartoni* (Bean).
 Upland portion of the Rio Verde in San Luis Potosi, Mexico.
Acaro bartoni Bean, Proc. U. S. Nat. Mus., XV, 1892, 286, Pl. XLIV, fig. 3, Huasteca Potosina, San Luis Potosi, Mexico.
3334. *Parapetenia labridens* (Pellegrin).
 Upland portion of the Rio Verde in San Luis Potosi, Mexico.
Heros labridens Pellegrin, Bull. Mus. Hist. Nat. Paris, 1903, 122, Huasteca, Potosi, Mexico.
3335. *Parapetenia beani* (Jordan). *Mojarra verde*.
 Lowland streams of Sinaloa and Jalisco.
Heros beani Jordan, Proc. U. S. Nat. Mus., XI, 1888, 332, Rio Presidio, near Mazatlan, Mexico.
3336. *Parapetenia tenuis* (Meek).
 Atlantic slope of the isthmus of Tehuantepec; Achatol; Vera Cruz, Mexico.
Cichlasoma tenue Meek, Field Mus. Nat. Hist., Zool., VII, 1906, 94, Achatal, Vera Cruz, Mexico.
3337. *Parapetenia centralis* (Meek).
 Pacific slope of Guatemala; Caballo Blanco.
Cichlasoma centrale Meek, Field Mus. Nat. Hist., Zool., VII, 1906, 94, Caballo Blanco, Guatemala.
3338. *Parapetenia alfari* (Meek).
 Atlantic slope of Costa Rica; Turrialba.
Cichlasoma alfari Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 148, Turrialba, Costa Rica.
3339. *Parapetenia mento* (Vaillant and Pellegrin).
 Southern Mexico.
Heros mento Vaillant and Pellegrin, Bull. Mus. Hist. Nat. Paris, 1902, 88, Rio Negro, southern Mexico.

- 3340. Parapetenia trimaculata** (Günther).
 Pacific coast streams of southern Mexico and Guatemala.
Heros trimaculatum Günther, Trans. Zool. Soc. Lond., VI, 1868, 461,
 pl. 76 (Nov. 27), Chiapam and Huamuchal, Mexico.
- Genus 1164. ARCHOCENTRUS** Gill.
Archocentrus Gill, Proc. Acad. Nat. Sci. Phila., XXIX, 1877, 186
 (*Heros centrarchus* Gill and Bransford).
- 3341. Archocentrus centrarchus** (Gill and Bransford).
 Great Lakes of Nicaragua.
Heros centrarchus Gill and Bransford, Proc. Acad. Nat. Sci. Phila.,
 XXIX, 1877, 185, Lake Nicaragua.
- 3342. Archocentrus nigrofasciatus** (Günther).
 Lakes Amatitlan and Atitlan, Guatemala.
Heros nigrofasciatus Günther, Trans. Zool. Soc. Lond., VI, 1868, 391,
 452, pl. 74, fig. 3, Lakes Amatitlan and Atitlan.
- 3343. Archocentrus spilurus** (Günther).
 Rio Motagua and Lake Isabel, Guatemala.
Heros spilurus Günther, Cat., IV, 1862, 289, Lake Yzabal, Guatemala.
- 3344. Archocentrus spinosissimus** (Vaillant and Pellegrin).
 Rio Polochic, Guatemala.
Heros (Cichlasoma) spinosissimus Vaillant and Pellegrin, Bull. Mus.
 Hist. Nat. Paris, 1902, 87, Rio Polochic, Guatemala.
- 3345. Archocentrus immaculatus** (Pellegrin).
 Rio Polochic, Guatemala.
Cichlasoma spinosissimum immaculatum Pellegrin, Mém. Soc. Zool.
 France, XVI, 1903, 225, Rio Polochic, Guatemala.
- Genus 1165. ASTATHEROS**³³ Pellegrin.
Astatheros Pellegrin, Mém. Soc. Zool. France, 1833, 205 (*Heros heterodontus* Vaillant and Pellegrin).
Erythrichthys Meek, Field Mus. Nat. Hist., Zool., VII, 1907, 118, 121
 (*Heros citrinellus* Günther); name twice preoccupied as *Erythrichthys*.
- 3346. Astatheros heterodontus** (Vaillant and Pellegrin).
 Isthmus of Tehuantepec, Mexico.
Heros heterodontus Vaillant and Pellegrin, Bull. Mus. Nat. Hist., Paris,
 1902, 86, Isthmus of Tehuantepec.
- 3347. Astatheros evermanni** (Meek).
 Pacific slope of Isthmus of Tehuantepec.
Cichlasoma evermanni Meek, Field Col. Mus. Zool., V, 1904, 214, fig.
 70, Rio Tehuantepec, Tehuantepec, Mexico.
- 3348. Astatheros robertsoni** (Regan).
 British Honduras.
Cichlasoma robertsoni Regan, Ann. Mag. Nat. Hist., ser. 7, XVI,
 1905, 239, Stann Creek, British Honduras.
- 3349. Astatheros lethrinus** (Regan).
 Rio Iroquois, Costa Rica.
Cichlasoma lethrinus Regan, Ann. Mag. Nat. Hist., ser. 8, II, 1908, 462,
 Rio Iroquois, Costa Rica.
- 3350. Astatheros dorsatus** (Meek).
 Lake Managua, Nicaragua.
Cichlasoma dorsatum Meek, Field Col. Mus. Nat. Hist., Zool., VII,
 1907, 123, Lake Managua.
- 3351. Astatheros longimanus** (Günther).
 Lake Nicaragua.
Heros longimanus Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27),
 391, 453, pl. 72, fig. 2, Lake Nicaragua.
- 3352. Astatheros altifrons** (Kner and Steindachner).
 Panama and Colombia.
Heros altifrons Kner and Steindachner, Sitz. Bayer. Akad., 1863, 223,
 Colombia.

³³This group is usually united with the two large-lipped species *labiatum* and *lobochilus*, to which the name *Amphilophus* was given earlier. *Citrinellus* and *Erythruus* seem intermediate.

3353. *Astatheros lauræ* (Regan).
 Tampico, Mexico.
Cichlasoma lauræ Regan, Ann. Mag. Nat. Hist., 8th ser., 2, 1908, 223,
 Tampico, Mexico.
3354. *Astatheros margaritifer* (Günther).
 Lake Peten, Guatemala.
Heros margaritifer Günther, Cat., IV, 1862, 287, Lake Peten.
Heros melanopogon Steindachner, Denks. Akad. Wiss. Wien, XXIII,
 1864, 72, pl. 1, fig. 3, Central America.
3355. *Astatheros sexfasciatus* (Regan).
 Guapote, Mexico.
Cichlosoma sexfasciatum Regan, Ann. Mag. Nat. Hist., (7), XVI, 1905,
 230, Guapote, Mexico.
3356. *Astatheros septemfasciatus* (Regan).
 Costa Rica.
Cichlosoma septemfasciatum Regan, Ann. Mag. Nat. Hist., II, ser. 8,
 1908, 461, Rio Iroquois, Costa Rica.
3357. *Astatheros basilaris* (Gill and Bransford).
 Lake Nicaragua.
Heros basilaris Gill and Bransford, Proc. Acad. Nat. Sci. Phila., XXIX,
 1877, 182, Lake Nicaragua.
3358. *Astatheros rectangularis* (Steindachner).
 Mexico.
Acara rectangularis Steindachner, Chromiden Mejicos, 1864, pl. 1, fig. 1,
 1, Mexico.
3359. *Astatheros maculicauda* (Regan).
 Atlantic slope of Guatemala and Panama.
Cichlasoma maculicauda Regan, Ann. Mag. Nat. Hist., XVI, ser. 7,
 1905, 227, Lake Yzabal, Rio Motagua, and Rio Chagres.
3360. *Astatheros tuyrensis* (Meek and Hildebrand).
 Rivers of Panama.
Cichlasoma tuyrensis Meek and Hildebrand, Field Mus. Nat. Hist.,
 Zool., X, 1913, 89, Rio Tuyra, Boca de Cupa, Panama.
3361. *Astatheros oblongus* (Günther).
 Rio Motagua; Guatemala.
Heros oblongus Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27),
 392, 464, Rio Motagua, Guatemala
3362. *Astatheros montezuma* (Heckel).
 Mexico.
Heros montezuma Heckel, Ann. Wiener Mus., II, 1840, 382, Mexico.
3363. *Astatheros globosum* (Miller).
 Puerto Barrios, Guatemala.
Cichlasoma globosum Miller, Bull. Amer. Mus. Nat. Hist., XXIII,
 1907, 114, fig. 4, Puerto Barrios, Guatemala.
3364. *Astatheros rostratum* (Gill and Bransford).
 Great Lakes of Nicaragua.
Heros rostratus Gill and Bransford, Proc. Acad. Nat. Sci. Phila., XXIX,
 1877, 181, Lake Nicaragua.
3365. *Astatheros hedricki* (Meek).
 Large Atlantic slope rivers of Mexico, south of Vera Cruz.
Cichlasoma hedricki Meek, Field Col. Mus., Zool., V, 1904, 208, fig. 66,
 Obispo, Vera Cruz, Mexico.
3366. *Astatheros octofasciatus* (Regan).
 Atlantic slope, Isthmus of Tehuantepec to British Honduras.
Heros octofasciatus Regan, Ann. Mus. Hist. Nat. Geneva, Rev. Suisse,
 Zool., 1903, 417, pl. 13, fig. 1, Mexico.
3367. *Antatheros granadense* (Meek).
 Lake Nicaragua.
Cichlasoma granadense Meek, Field Mus. Nat. Hist., Zool., VII, 1907,
 121, Lake Nicaragua, Nicaragua, Granada.

3368. *Astatheros citrinellus* (Günther).

Great Lakes of Nicaragua.

Heros citrinellus Günther, Proc. Zool. Soc. Lond., 1864, 153, Lake Nicaragua.

3369. *Astatheros erythræum* (Günther).

Great lakes of Nicaragua.

Heros erythræus Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 391, 457, pl. 75, fig. 2, Lake Managua.

Genus 1166. *AMPHILOPHUS* Agassiz.

Amphilophus Agassiz, Proc. Bost. Soc. Nat. Hist., VI, 1858, 401 (*A. frebeli* Agassiz); a large-lipped species from Lake Nicaragua not otherwise described, probably *labiatus*; these species, *labiatus* and *lobochilus*, have been placed in *Astatheros*, but the name *Amphilophus* has priority.

3370. *Amphilophus labiatus* (Günther).

Great lakes of Nicaragua.

Heros labiatum Günther, Proc. Zool. Soc. Lond., 1864, 27, Pl. IV, fig. 1, Lake Nicaragua.

3371. *Amphilophus lobochilus* (Günther).

Lake Managua, Nicaragua.

Heros lobochilus Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 391, 457, pl. 75, fig. 1, Lake Managua, Nicaragua.

Genus 1167. *THORICHTHYS* Meek.

Thorichthys Meek, Field Col. Mus., Zool., V, 1904, 222 (*T. ellioti* Meek).

3372. *Thorichthys ellioti* Meek.

Atlantic slope of Vera Cruz to Guatemala.

Thorichthys ellioti Meek, Field Col. Mus., Zool., V, 1904, 223, Motzorongo, Vera Cruz, Mexico.

3373. *Thorichthys callolepis* (Regan).

Atlantic slope of Isthmus of Tehuantepec, Mexico.

Heros callolepis Regan, Ann. Mag. Nat. Hist., ser. 7, XIII, 1904, 258, San Domingo de Guzman, Mexico.

3374. *Thorichthys aureus* (Günther).

Atlantic slope, Vera Cruz, Mexico, to Guatemala.

Heros aureus Günther, Cat., IV, 1862, 292, Lake Yzabal, Guatemala.

3375. *Thorichthys affinis* (Günther).

Lake Peten, Guatemala.

Heros affinis Günther, Cat., IV, 1862, 292, Lake Peten.

3376. *Thorichthys acutus* (Miller).

Small tributaries of the Rio Motagua, Guatemala.

Cichlasoma acutum Miller, Bull. Amer. Mus. Nat. Hist., XXIII, 1907, 117, fig. 6, Rio Tenedores, Guatemala.

3377. *Thorichthys helleri* (Steindachner).

Lowland streams of Mexico emptying into the Gulf south of Vera Cruz.

Heros helleri Steindachner, Denks. Akad. Wiss. Wien, XXIII, 1864, 64.

Pl. IV, fig. 1, Rio Teapa, Tabasco, Mexico.

Heros maculipinnis Steindachner, loc. cit., 69, Pl. IV, fig. 2, Rio Zanopa, Mexico.

Genus 1168. *HERICHTHYS* Baird and Girard.

Herichthys Baird and Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854,

25 (*H. cyanoguttatus* Baird and Girard).

3378. *Herichthys cyanoguttatus* Baird and Girard.

East slope of Mexico from Rio Grande to Rio Panuco.

Herichthys cyanoguttatus Baird and Girard, Proc. Acad. Nat. Sci. Phila.,

VII, 1854, 25, Rio Grande, Brownsville, Tex.

Heros leporatus Fowler, loc. cit., LV, 1903, 321, Victoria, Tamaulipas.

3379. *Herichthys pavonaceus* (Garman).

Spring near Monclova, Coahuila, Mexico.

Heros pavonaceus Garman, Bull. Mus. Comp. Zool., VIII, 1881, 93, spring near Monclova, Coahuila, Mexico.

3380. *Herichthys bocourtii* (Vaillant and Pellegrin).
Lake Isabel, Guatemala.
Neetroplus bocourtii Vaillant and Pellegrin, Bull. Mus. Hist. Nat., Paris, 1902, 85, Lake Isabel, Guatemala.
3381. *Herichthys carpintis* (Jordan and Snyder).
Rio Panuco, Mexico, and its lowland tributaries.
Neetroplus carpintis Jordan and Snyder, Bull. U. S. Fish Com., XIX, 1899, 146, fig. 22, Laguna de Carpinte, Tampico, Mexico.
3382. *Herichthys troscheli* (Steindachner).
Mexico.
Heros troscheli Steindachner, Ichthyologische Notizen, IV, 1867, 12, Mexico.
- Genus 1169. **TOMOCICHLA** Regan.
Tomocichla Regan, Ann. Mag. Nat. Hist., II, ser. 8, 1908, 463 (*Herichthys underwoodi* Regan).
3383. *Tomocichla underwoodi* (Regan).
Costa Rica.
Herichthys underwoodi Regan, Biol. Central Amer., Pisces, 1908, 30, Costa Rica.
3384. *Tomocichla geddesi* (Regan).
Southern Mexico.
Herichthys geddesi Regan, Ann. Mag. Hist., XVI, ser. 7, 1905, 436, southern Mexico.
- Genus 1170. **PETENIA** Günther.
Petenia Günther, Cat., IV, 1862, 301 (*P. splendida* Günther).
3385. *Petenia splendida* Günther.
Lake Peten, Guatemala; Montecristo; Frontera.
Petenia splendida Günther, Cat., IV, 1862, 301, Lake Peten, Guatemala.
- Genus 1171. **PARANEETROPLUS** Regan.
Paraneetroplus Regan, Ann. Mag. Nat. Hist., XVI, ser. 7, 1905, 436 (*P. bulleri* Regan).
3386. *Paraneetroplus bulleri* Regan.
Rio de Sarabia, Mexico.
Paraneetroplus bulleri Regan, Ann. Mag. Nat. Hist., XVI, ser. 7, 1905, 437, Rio de Sarabia, Mexico.
- Genus 1172. **NEETROPLUS** Günther.
Neetroplus Günther, Trans. Zool. Soc. Lond. VI, 1866, 469 (*N. nematopus* Günther).
3387. *Neetroplus sieboldii* (Kner and Steindachner).
New Granada, and from the Pacific slope of Panama.
Heros sieboldii Kner and Steindachner, Abhandl. Bayer Akad. Wiss., X, 1864, 13, pl. 2, fig. 2, Panama, Colombia.
3388. *Neetroplus nematopus* Günther.
Great lakes of Nicaragua.
Neetroplus nematopus Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 392, 470, pl. 74, fig. 4, Lake Managua.
- Genus 1173. **HEROTILAPIA** Regan.
Herotilapia Regan, Ann. Mag. Nat. Hist., V, 38 (*Tilapia buettikoferi* Boulenger).
3389. *Herotilapia multispinosa* (Günther).
Great lakes of Nicaragua.
Heros multispinosus Günther, Trans. Zool. Soc. Lond., VI, 1868 (Nov. 27), 391, 453, pl. LXXIV, fig. 2, Lake Managua.
- Genus 1174. **ÆQUIDENS** Eigenmann and Bray.
Æquidens Eigenmann and Bray, Ann. N. Y. Acad. Sci., 1894, 616 (*Acara tetramerus* Heckel).
3390. *Æquidens caeruleopunctata* (Kner and Steindachner).
Panama to northwest Ecuador.
Acara caeruleopunctata Kner and Steindachner, Sitz. Bayer. Akad., 1863, 222, northwest Ecuador.

Genus 1175. **GEOPHAGUS** Heckel.

Geophagus Heckel, Ann. Mus. Wien, II, 1840, 383 (*G. magasema* Heckel = *Sparus surinamensis* Bloch).

Mesops Günther, Cat., IV, 1862, 311 (*Geophagus cupido* Heckel).
Satanoperca Günther, loc. cit., 312 (*Geophagus demon* Heckel).

Subgenus **SATANOPERCA** Günther.

3391. **Geophagus crassilabris** Steindachner.

Panama.

Geophagus (*Satanoperca*) *crassilabris* Steindachner, Sitz. Akad. Wiss. Wien, 1876, 74, 65, Pl. VII, Panama.

Order UU. **PHARYNGOGNATHI**Family 251. **LABRIDÆ**Genus 1176. **CENTROLABRUS** Günther. *Rock cooks.*

Centrolabrus Günther, Cat., IV, 1862, 92 (*Labrus exoletus* Linnæus).

3392. **Centrolabrus exoletus** (Linnæus). *Rock cook.*

Coast of northern Europe south to Cornwall; Greenland.

Labrus exoletus Linnæus, Syst. Nat., Ed. X, 1758, 287, Atlantic Ocean.

Labrus pentacanthus Lacépède, Hist. Nat. Poiss., III, 1803, 503; after Linnæus.

Crenilabrus microstoma Couch, in Thompson, Proc. Zool. Soc. London, 1837, 55, Cornwall.

Genus 1177. **TAUTOGOLABRUS** Günther. *Cunners.*

Tautogolabrus Günther, Cat., IV, 1862, 89 (*Ctenolabrus bergall* Günther = *Labrus adspersus* Walbaum).

3393. **Tautogolabrus adspersus** (Walbaum). *Cunner; Chogset; Blue perch; Bergall; Berg-gylt; Nibbler; Conner; Gall Bengal.*

Atlantic coasts of North America from Labrador to Sandy Hook.

Labrus adspersus Walbaum, Artedi Piscium, 1792, 254, no locality; after *Bergall* of Schöpf.

Tautoga niger Mitchell, Report, in part, in Fishes of New York, 1814, 23, New York.

Tautoga cærulea Mitchell, Report, loc. cit., 1814, 24, New York.

Labrus chogset Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 402, pl. 3, fig. 2, New York.

Labrus chogset fulva Mitchell, loc. cit., 1815, 403, New York.

Ctenolabrus uninotatus Cuvier and Valenciennes, Hist. Nat. Poiss., XIII, 1839, 174 (239), New York.

Ctenolabrus bergall Günther, Cat., IV, 1862, 90, Canada.

Genus 1178. **TAUTOGA** Mitchell. *Tautogs.*

Hiatula Lacépède, Hist. Nat. Poiss., II, 1800, 522; name preoccupied.

Tautoga Mitchell, Rept. Fishes New York, 1814, 23 (*T. niger* Mitchell = *Labrus tautoga* Mitchell = *Labrus onitis* Linnæus).

3394. **Tautoga onitis** (Linnæus). *Tautog; Blackfish; Oysterfish; Tautoga; Smooth blackfish; Cub; Sea dog.*

Atlantic coast of United States; Cape Ann to Charleston.

Labrus onitis Linnæus, Syst. Nat., Ed. X, 1758, 286, locality unknown.

Labrus hiatula Linneaus, Syst. Nat., Ed. XII, 1766, 475, Carolinas.

Labrus carolinus Bonnaterre, Tableau Encyclopéd. et Méthod. Ichthyologie, 1788, 113, Carolinas.

Labrus subfucus Walbaum, Artedi Pisc., 1792, 254; after Schöpf.

?*Labrus tessellatus* Bloch, Naturgesch. Ausl. Fische, VI, 1792, pl. 8, Norway.

Hiatula gardeniana Lacépède, Hist. Nat. Poiss., II, 1800, 522, Carolinas.

Labrus americanus Bloch and Schneider, Syst. Ichth., 1801, 261; after Schöpf.

Labrus tautoga Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 398, Long Island; Rhode Island; Cape Cod; Sandy Circle.

Tautoga fusca Mitchell, Rept. Fish. N. Y., 1814, 23, New York.

Tautoga rubens Mitchell, loc. cit., 23, New York.

Tautoga alia Mitchell, loc. cit., 23, New York.

Tautoga niger Mitchell, loc. cit., 23, New York.

Genus 1179. ACHNOLAIMUS Cuvier and Valenciennes. *Capitaines.*
Lachnolaimus Cuvier and Valenciennes, Hist. Nat. Poiss., XIII, 1839,
 200 (274) (*L. aigula* Cuvier and Valenciennes = *Labrus maximus*
 Walbaum).
Lachnolaimus Günther; corrected spelling.

- 3395. *Lachnolaimus maximus* (Walbaum).** *Capitaine; Perro perro; Hogfish.*
 West Indies, north to Key West and the Bermudas.
Suillus Catesby, Nat. Hist. Carol., II, 1743, 15, pl. 15; no locality;
 probably Bahamas.
Labrus maximus Walbaum, Arredi Pisc., 1792, 261, no locality; after
 Catesby.
Lachnolaimus suillus Cuvier, Règne Animal, ed. 2, II, 1829, 257; after
 Catesby.
Lachnolaimus aigula Cuvier and Valenciennes, Hist. Nat. Poiss., XIII,
 1839, 202 (277), pl. 378, St. Bartholomew.
Lachnolaimus dux Cuvier and Valenciennes, loc. cit., 208 (285), Mar-
 tinique.
Lachnolaimus caninus Cuvier and Valenciennes, loc. cit., 210 (288),
 St. Thomas; San Domingo.
Lachnolaimus psittacus Cuvier and Valenciennes, loc. cit., 212 (291);
 Porto Rico.

Genus 1180. BODIANUS Bloch.
Bodianus Bloch, Nat. Ausl. Fische, IV, 1790, 31, 33 (*B. bodianus*
 Bloch = *Labrus rufus* Linnaeus); in part; type by tautonomy; not by
 first restriction to species of *Cephalopholis*.
Harpe Lacépède, Hist. Nat. Poiss., IV, 1803, 426 (*H. cæruleo-aureum*
 Lacépède = *Labrus rufus* Linnaeus).
Cossyphus Cuvier and Valenciennes, loc. cit., XIII, 1839, 102 (*Bodianus*
bodianus Bloch); name twice preoccupied.
Crenilabrus Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839 (*Lutjanus*
verres Bloch = *Labrus rufus* Linnaeus); name preoccupied.

- 3396. *Bodianus rufus* (Linnaeus).** *Ladyfish; Spanish ladyfish; Spanish hogfish;*
Pudiano; Perro colorado.
 West Indies, north to Key West, south to Rio de Janeiro.
Labrus rufus Linnaeus, Syst. Nat., Ed. X, 1758, 284, America; after
 Catesby.
Pudiano vermelho Marcgrave, Hist. Bras., 1648, 145, Brazil.
Turdus flavus Catesby, Nat. Hist. Carolina, II, 1743, 11, pl. 11, fig. 1,
 Bahamas.
Perro colorado Parra, Descr. Dif. Piez. Hist. Nat. Cuba, 1787, 3, lam. 3,
 fig. 1, Habana.
?Bodianus bodianus Bloch, Naturgesch. Ausl. Fische, IV, 1790, 33,
 Brazil.
?Lutjanus verres Bloch, loc. cit., V, 1791, 9, locality uncertain.
?Sparus falcatus Bloch, loc. cit., 20, Martinique; after Plumier.
Labrus semiruber Lacépède, Hist. Nat. Poiss., III, 1802, 428, Rio
 Janeiro.
Bodianus blockii Lacépède, loc. cit., IV, 1803, 279; after Bloch.
Harpe coeruleo-aureus Lacépède, loc. cit., 426, pl. 8, fig. 2, Martinique.

- 3397. *Bodianus diplostenia* (Gill).**
 Pacific coast of tropical America; Cape San Lucas; Panama; Revilla-
 gigedo Islands; Mazatlan.
Harpe diplostenia Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 140,
 Cape San Lucas; female.
Harpe pectoralis Gill, loc. cit., 141, Cape San Lucas; male.

- 3398. *Bodianus pulchellus* (Poey).**
 Cuba.
Cossyphus pulchellus Poey, Memorias, II, 1860, 208, Habana.

Genus 1181. DECODON Günther.
Decodon Günther, Cat., IV, 1862, 101 (*Cossyphus puellaris* Poey).

- 3399. *Decodon puellaris* (Poey).**
 West Indies; Cuba; Barbados; Snapper Banks off Pensacola, Fla.
Cossyphus puellaris Poey, Memorias, II, 186, 2101, Habana.

Genus 1182. PIMELOMETOPON Gill. *Fatheads.*

Pime洛metopon Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 58 (*Labrus pulcher* Ayres).

3400. *Pime洛metopon pulcher* (Ayres). *California redfish; Fathead; Sheep-head.*

Southern California, from Point Conception to Ascension Island, Lower California.

Labrus pulcher Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 3, San Diego, Calif.

Genus 1183. CLEPTICUS Cuvier.

Clepticus Cuvier, Règne Anim., ed. 2, II, 1829, 261 (*Rabirubia genizara Parra*=*Brama parræ* Bloch and Schneider).

3401. *Clepticus parræ* (Bloch and Schneider). *Genizara; Janissary; Creole.*

West Indies.

Rabirubia genizara Parra, Dif. Piezas de Hist. Nat. Cuba, 1787, 44, pl. 21, fig. 1, Habana; author not binomial.

Brama parræ Bloch and Schneider, Syst. Ichth., 1801, 100, Habana.

Family 252. CORIDÆ

Genus 1184. IRIDIO⁴ Jordan and Evermann. *Doncellas.*

Halichæres Rüpell, Neue Wirbelthiere, Fische, 1835, 17 (*bimaculatus*); not *Halichærus* Nillson, 1820, a genus of seals.

Hermilus Swainson, Nat. Hist. Classn. Fishes, II, 1839 228, (*Labrus guttatus* Bloch); as restricted by Bonaparte (1841); restricted by Swain (1882) to *Cheilio auratus* Quoy and Gaimard.

Parajulis Bleeker, Ned. Tijdschr. Dierk., II, 1865, 251 (*Julis pacilopterus* Temminck and Schlegel).

Iridio Jordan and Evermann, Check List, 1895, 412 (*Julis dimidiatus* Agassiz= *Labrus cyanocephalus* Bloch).

3402. *Iridio radiata* (Linnæus). *Puddingwife; Doncella; Pudiano verde.*

West Indies, Brazil north to Florida Keys, Bermuda, and St. Paul Rocks.

Labrus radiatus Linnæus, Syst. Nat., Ed. X, 1758, 288, Bahamas; based on Catesby.

Pudiano verde Marcgrave, Hist. Pisc. Brasil., 1648, 146, Brazil.

Turdus oculo-radiato Catesby, Nat. Hist. Carolina, II, 1743, 12, pl. 12, fig. 1, Bahamas.

Doncella Parra, Desc. Dif. Piez. Hist. Nat. Cuba, 1787, 95, lam. 37, fig. 1, Habana.

Labrus brasiliensis Bloch, Naturgesch. Ausl. Fische, V, 1791, 125, Brazil.

Julis crotaphus Cuvier, Règne Anim., ed. 2, II, 1829, 258; after Parra.

Julis cyanostigma Cuvier and Valenciennes, Hist. Nat. Poiss., XIII, 1839, 286 (391), Martinique.

Julis opalina Cuvier and Valenciennes, loc cit., 287 (392), Martinique.

Julis patatus Cuvier and Valenciennes, loc. cit., 291 (398), Martinique; Cuba.

Julis principis Cuvier and Valenciennes, loc. cit., 294 (402), Bahia.

3403. *Iridio bivittata*⁴ (Bloch). *Slippery dick; Doncella.*

West Indies, north to Beaufort, N. C., and south to Brazil.

Labrus bivittatus Bloch, Naturgesch. Ausl. Fische, V, 1791, 133, from a painting by Plumier made at Martinique.

Sparus radialatus Linnaeus. Syst. Nat., ed. XII, 1766, 472, Carolinas.

Labrus psittacus Lacépède, Hist. Nat. Poiss., III, 1800, 452, Martinique; after Plumier.

Julis humeralis Poey, Memorias, II, 1860, 212, Cuba.

Chærojulis grandisquamis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 206, Beaufort, N. C.

Chærojulis arangoi Poey, Enumeratio, 1876, 109, Habana.

Platyglossus florealis Jordan and Gilbert, Proc. U. S. Nat. Mus., V,

1882, 287, Pensacola, Fla.

⁴ This genus is very close to the Old World genus *Halichæres*.

⁴ Regarded by some authors as the young of *Iridio radiata*.

- 3404.** *Iridio cyanoccephalus* (Bloch).
 West Indies, south to Brazil.
Labrus cyanoccephalus Bloch, Naturgesch. Ausl. Fische, V, 1791, 139,
 location unknown; Museum of Link.
Julis dimidiatus Agassiz, in Spix, Pisc. Bras., 1829, 96, pl. 53, Brazil.
Julis internasalis Poey, Memorias, II, 1861, 421, Cuba.
- 3405.** *Iridio maculipinna* (Müller and Troschel).
 West Indies to Beaufort, N. C.
Julis maculipinna Müller and Troschel, in Schomburgk, Hist. Barbados,
 1848, 674, Barbados.
- 3406.** *Iridio garnoti* (Cuvier and Valenciennes).
 West Indies, Cuba, Martinique, St. Croix.
Julis garnoti Cuvier and Valenciennes, Hist. Nat. Poiss., XIII, 1839,
 285 (390), Martinique.
Julis cinctus Poey, Memorias, II, 1860, 211, Cuba.
Julis ruptus Poey, loc. cit., 212, Cuba.
- 3407.** *Iridio kirschii* Jordan and Evermann.
 Cuba to Brazil.
Julis crotophus Cuvier and Valenciennes, Hist. Nat. Poiss., XIII,
 1839, 289 (395), Bahia; not of Cuvier.
Iridio kirschii Jordan and Evermann, Check List, 1895, 413, Bahia,
 Brazil.
- 3408.** *Iridio poeyi* (Steindachner).
 Surinam.
Platyglossus poeyi Steindachner, Ichth. Notizen, VI, 1867, 49, Surinam.
- 3409.** *Iridio caudalis* (Poey).
 Cuba.
Julis caudalis Poey, Memorias, II, 1860, 213, Habana.
- 3410.** *Iridio pictus* (Poey).
 West Indies, north to the Snapper Banks off Pensacola, Fla.
Julis pictus Poey, Memorias, II, 1860, 214, Cuba.
- 3411.** *Iridio elegans* Bean.
 Bermudas to Curaçao.
Iridio elegans Bean, Field Mus. Nat. Hist., Zool., VII, 1906, 65, fig. 6,
 Nonesuch Island, Bermudas.
- 3412.** *Iridio similis* Nichols.
 Bermudas.
Iridio similis Nichols, Proc. Biol. Soc. Wash., XXXIII, 1920, 61,
 Bermudas.
- 3413.** *Iridio frenatus* Nichols.
 Bermuda.
Iridio frenatus Nichols, Proc. Biol. Soc. Wash., XXXIII, 1920, 61,
 Bermuda.
- 3414.** *Iridio decoratus* Bean.
 Bermudas.
Iridio decoratus Bean, Proc. Biol. Soc. Wash., XIX, 1906, 29, Nonesuch
 Island, Bermudas.
- 3415.** *Iridio meyeri* Bean.
 Bermudas.
Iridio meyeri Bean, Proc. Biol. Soc. Wash., XIX, 1906, 29, Nonesuch
 Island, Bermudas.
- 3416.** *Iridio microstomus* Bean.
 Bermudas.
Iridio microstomus Bean, Proc. Biol. Soc. Wash., XIX, 1906, 30, None-
 such Island, Bermudas.
- 3417.** *Iridio semicinctus* (Ayres). *Kelpfish; Señorita.*
 Southern California, Santa Barbara Islands to Cedros Island.
Julis semicinctus Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 32,
 Cerros [Cedros] Island, off coast of Lower California.
- 3418.** *Iridio dispilus* (Günther).
 Pacific coast of tropical America, Mazatlan to Panama.
Platyglossus dispilus Günther, Proc. Zool. Soc. Lond., 1864, 25, Panama.

3419. *Iridio nicholsi* (Jordan and Gilbert).
 Revillagigedo and Galapagos Islands.
Platyglossus nicholsi Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 231, Braithwaite Bay, Socorro Island.
3420. *Iridio sellifer* (Gilbert).
 Revillagigedo Archipelago.
Halichæres sellifer Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 67, Clarion Island.
- Genus 1185. *OXYJULIS* Gill. *Señoritas.*
Oxyjulis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 330 (*Julis modestus* Girard).
3421. *Oxyjulis californicus* (Günther). *Señorita.*
 Coast of southern California; common from Monterey to Guadalupe Island.
Julis modestus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 151, San Diego, Monterey, San Miguel; preoccupied.
Halichæres californicus Günther, Ann. Mag. Nat. Hist., ser. 3, VIII, 1861, 386, name only; substitute for *Julis modestus*, preoccupied.
- Genus 1186. *PSEUDOJULIS* Bleeker.
Pseudojulis Bleeker, Proc. Zool. Soc. Lond., 1861, 412 (*P. girardi* Bleeker).
- Subgenus **PSEUDOJULIS.**
3422. *Pseudojulis inornatus* Gilbert.
 Pacific coast of Mexico; only the type known.
Pseudojulis inornatus Gilbert, Proc. U. S. Nat. Mus., XI II, 1890, 67, Pacific coast of Mexico.
3423. *Pseudojulis melanotis* Gilbert.
 Pacific coast of Mexico; only the type known.
Pseudojulis melanotis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 66, Pacific coast of Mexico.
- Genus 1187. *EMMEEKIA* Jordan and Evermann.
Emmekia Jordan and Evermann, Check List, 1895, 413 (*Pseudojulis venustus* Jenkins and Evermann).
3424. *Emmekia venusta* (Jenkins and Evermann).
 Gulf of California.
Pseudojulis venustus Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 145, Guaymas, Sonora, Mexico.
- Genus 1188. *JULIDIO* Jordan and Evermann.
Julidio Jordan and Evermann, Check List, 1895, 413 (*Pseudojulis adustus* Gilbert).
3425. *Julidio adustus* (Gilbert).
 Revillagigedo Islands.
Pseudojulis adustus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 66, Socorro Island.
3426. *Julidio notospilus* (Günther).
 Pacific coast of Mexico, Panama, and Clarion Island of Revillagigedo Archipelago.
Pseudojulis notospilus Günther, Proc. Zool. Soc. Lond., 1864, 26, Panama.
- Genus 1189. *THALASSOMA* Swainson:
Thalassoma Swainson, Nat. Hist. Class. Fishes, II, 1839, 224 (*Labrus purpureus* Rüppell).
Chlorichthys Swainson, loc. cit., 232 (*Labrus bifasciatus* Bloch).
Julis Günther, not of Cuvier and Valenciennes.

- 3427. *Thalassoma bifasciatum* (Bloch).**
 West Indies; Tortugas; Jamaica; Cuba; San Domingo; and Martinique.
?Labrus bifasciatus Bloch, Naturgesch. Ausl. Fische, V, 1791, 131,
 West Indies.
Labrus bifasciatus torquatus Bloch and Schneider, Syst. Ichth., 1801,
 243, Antilles; after Gronow.
Julis detensor Cuvier and Valenciennes, Hist. Nat. Poiss., XIII, 1839,
 298 (408), San Domingo.
Labrus ornatus Gronow, Syst. Nat., ed. Gray, 1854, 83, Antilles.
Julis gillianus Poey, Memorias, II, 1860, 214, Cuba.
?Julis nitida Günther, Cat., IV, 1862, 190, Jamaica
- 3428. *Thalassoma socorroensis* Gilbert.**
 Revillagigedo Archipelago.
Thalassoma socorroense Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 69,
 Socorro Island.
- 3429. *Thalassoma nitidum* (Günther).**
 West Indies; Jamaica; Tortugas.
Julis nitida Günther, Cat., IV, 1862, 190, Jamaica.
Julis nitidissima Goode, Am. Jour. Sci. Arts, XIV, 1877, 293, Bermuda.
- 3430. *Thalassoma grammaticum* Gilbert.**
 Revillagigedo Archipelago.
Thalassoma grammaticum Gilbert, Proc. U. S. Nat. Mus., XIII, 1890,
 68, Socorro Island; Clarion Island.
- 3431. *Thalassoma virens* Gilbert.**
 Revillagigedo Archipelago.
Thalassoma virens Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 68,
 Socorro Island.
- 3432. *Thalassoma lucasanum* (Gill).**
 Gulf of California, Cape San Lucas to Mazatlan and Tres Marias
 Islands; Panama Bay.
Julis lucasanus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 142,
 Cape San Lucas.
- 3433. *Thalassoma steindachneri* (Jordan).**
 Acapulco, Pacific coast of Mexico.
Thalassoma steindachneri Jordan, Rept. U. S. Fish Com., XV, 1887,
 654, Acapulco; after Steindachner.
Julis melanochir Steindachner, Ichth. Beitr., III, 1875, 63, Acapulco;
 not of Bleeker.
- Genus 1190. *DORATONOTUS* Günther.**
Doratonotus Günther, Cat., IV, 1862, 124 (*D. megalepis* Günther).
- 3434. *Doratonotus megalepis* Günther.**
 West Indies, north to Key West.
Doratonotus megalepis Günther, Cat., IV, 1862, 125, St. Christopher.
Doratonotus thalassinus Jordan and Gilbert, Proc. U. S. Nat. Mus.,
 VII, 1884, 28, Key West, Fla.
- 3435. *Doratonotus decoris* Evermann and Marsh.**
 Porto Rico.
Doratonotus decoris Evermann and Marsh, Rept. U. S. Fish Com.,
 XXV, 1899, 354, Ponce, Porto Rico.
- 3436. *Doratonotus boekei* Metzelaar.**
 Caribbean Sea.
Doratonotus boekei Metzelaar, Trop. Atlant. Visschen, 1919, 107, fig. 31,
 Curaçao, Bonaire.
- Genus 1191. *XYRULA* Jordan.**
Xyrula Jordan, Rept. U. S. Fish Com., XV, 1887, 656 (*Xyrichtys*
jessiae Jordan).
- 3437. *Xyrula jessiae* (Jordan).**
 Snapper Banks off Tampa, Fla.
Xyrichtys jessiae Jordan, Proc. U. S. Nat. Mus., X, 1887, 698, off
 Tampa Bay, Fla.

Genus 1192. NOVACULICHTHYS Bleeker.

Novacutichthys Bleeker, Proc. Zool. Soc. Lond., 1861, 414 (*Labrus tenuirostris* Bloch).

Dimalacocentrus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 223 (*Novacula kallasoma* Bleeker).

3438. Novaculichthys martinicensis (Cuvier and Valenciennes).

Martinique.

Xyrichthys martinicensis Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 37 (49), Martinique.

Xyrichthys vitta Cuvier and Valenciennes, loc. cit., 38 (51), no locality.

3439. Novaculichthys infirmus (Bean).

Cozumel, Yucatan.

Xyrichthys infirmus Bean, Bull. U. S. Fish Com., VIII, 1888, 199, pl. 29, fig. 2, Cozumel.

3440. Novaculichthys rosipes (Jordan and Gilbert).

Key West; Florida; Tortugas.

Xyrichthys rosipes Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884, 27, Key West, Tortugas.

3441. Novaculichthys ventralis (Bean).

Cozumel Island, Yucatan.

Xyrichthys ventralis Bean, Bull. U. S. Fish Com., VIII, 1888, 198, pl. 29, fig. 1, Cozumel, Yucatan.

Genus 1193. XYRICHTHYS Cuvier. *Razorfishes.*

Xyrichthys Cuvier, Mém. Mus. d'Hist. Nat., I, 1815, 324, 329 (*X. novacula* Cuvier).

Novacula Cuvier, Règne Anim., Ed. I, 1817, 265 (*Coryphæna novacula* Linnæus).

Amorphocephalus Bowdich, Exc. Madeira, 1825, 238 (*A. granulatus* Bowdich).

3442. Xyrichthys psittacus (Linnæus). *Razorfish.*

West Indies, north to Pensacola and Charleston.

Coryphæna psittacus Linnæus, Syst. Nat., Ed. XII, 1766, 448, Charleston, S. C.

Coryphæna lineata Gmelin, Syst. Nat., 1788, 1195, Charleston, S. C.

Xyrichthys venustus Poey, Enumeratio, 1876, 110, Martinique.

Xyrichthys vermiculatus Poey, Memorias, II, 1860, 215, Cuba.

3443. Xyrichthys mundiceps Gill.

Cape San Lucas, Lower California.

Xyrichthys mundiceps Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 143, Cape San Lucas.

3444. Xyrichthys modestus Poey.

Cuba.

Xyrichthys modestus Poey, Repertorio, II, 1867, 238, Cuba.

3445. Xyrichthys binghami Mowbray.

Bahama Islands.

Xyrichthys binghami Mowbray, Marine Life, I, Apr. 2, 1925, 1, Cay Sal Bank, Bahamas.

Genus 1194. INIISTIUS Gill.

Iniistius Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 143 (*Xyrichthys pavo* Cuvier and Valenciennes).

3446. Iniistius pavoninus (Cuvier and Valenciennes).

Hawaii and Pacific coast of Mexico.

Xyrichthys pavoninus Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 63, Hawaii.

Iniistius mundicorpus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 145, Cape San Lucas.

Genus 1195. BERMUDICHTHYS Nichols.

Bermudichthys Nichols, Proc. Biol. Soc. Wash., XXXIII, 1920, 62 (*B. subfurcatus* Nichols).

3447. Bermudichthys subfurcatus Nichols.

Bermuda.

Bermudichthys subfurcatus Nichols, Proc. Biol. Soc. Wash., XXXIII, 1920, 62, Bermuda.

Family 253: SPARISOMIDÆ

Genus 1196. CRYPTOTOMUS Cope.

Cryptotomus Cope, Trans. Amer. Philos. Soc., XIV, 1870, 462 (*C. roseus* Cope).

Nicholsina Fowler, Copeia, No. 14, 1915, 3 (*Cryptotomus beryllinus* Jordan and Swain).

Subgenus CRYPTOTOMUS Cope.

3448. *Cryptotomus roseus* Cope.

West Indies, south to Brazil.

Cryptotomus roseus Cope, Trans. Amer. Phil. Soc., XIV, 1871, 462, St. Martins, West Indies.

3449. *Cryptotomus crassiceps* Bean.

Bermudas.

Cryptotomus crassiceps Bean, Proc. Biol. Soc. Wash., XIX, 1906, 32, Coopers Island, Bermudas.

Subgenus NICHOLSINA Fowler.

3450. *Cryptotomus beryllinus* Jordan and Swain. *Parrotfish*.

Florida Keys, north to New Jersey, south to Rio Janeiro.

Cryptotomus beryllinus Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884, 101, Habana.

Sparisoma sp. Bean, Bull. U. S. Fish Com., VII, 1887, 137, Somers Point, N. J.

3451. *Cryptotomus dentiens* (Poey).

Cuba.

Scarus dentiens Poey, Memorias, II, 1861, 422, Cuba.

3452. *Cryptotomus retractus* (Poey).

West Indies, north to Pensacola.

Callidodon retractus Poey, Synopsis, 1868, 245, Habana.

3453. *Cryptotomus ustus* (Cuvier and Valenciennes).

West Indies, north to Charleston, south to Bahia.

Callyodon ustus Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 212 (286), pl. 405, Brazil.

3454. *Cryptotomus auropunctatus* (Cuvier and Valencieunes).

San Domingo.

Callyodon auro-punctatus Cuvier and Valencieunes, Hist. Nat. Poiss., XIV, 1839, 215 (290), Santo Domingo.

Genus 1197. LEPTOSCARUS Swainson.

Leptoscarus Swainson, Nat. Hist. Classn. Fishes, 1839, 11 (*Scarus vaigiensis* Quoy and Gaimard = *Scarus spinidens* Cuvier and Valenciennes).

Calotomus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 70 (*C. xenodon* Gilbert).

3455. *Leptoscarus xenodon* (Gilbert).

Socorro Island, Revillagigedo Archipelago.

Calotomus xenodon Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 70, Socorro Island, Revillagigedo Islands.

Genus 1198. SPARISOMA Swainson. Viejas.

Sparisoma Swainson, Nat. Hist. Classn. Fishes, etc., II, 1839, 227 (*Scarus abildgardi* Bloch).

Euscarus Jordan and Evermann, Check List, 1895, 416 (*Labrus cretensis* Linnaeus).

Subgenus SPARISOMA Swainson.

3456. *Sparisoma xystrodon* Jordan and Swain.

West Indies, north to Key West.

Sparisoma xystrodon Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884, 99, Key West; Habana.

3457. *Sparisoma atomarium* (Poey).

Cuba.

Scarus atomarius Poey, Memorias, II, 1861, 423, Habana.

- 3458.** *Sparisoma radians* (Cuvier and Valenciennes).
 West Indies, south to Brazil; Tortugas.
Scarus radians Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839,
 153 (206), Brazil.
Scarus lacrimosus Poey, Memorias, II, 1861, 422, Cuba.
- 3459.** *Sparisoma hoplomystax* (Cope).
 West Indies, from Key West to Bahia.
Scarus hoplomystax Cope, Trans. Amer. Phil. Soc., XIV, 1871, 462,
 St. Martins, West Indies.
Sparisoma cyanolene Jordan and Swain, Proc. U. S. Nat. Mus., VII,
 1884, 98, Key West, Fla.
- 3460.** *Sparisoma niphobles* Jordan and Bollman.
 Bahamas; Tortugas; Cape Florida; Key West; Porto Rico; abundant
 about Porto Rico.
Sparisoma niphobles Jordan and Bollman, Proc. U. S. Nat. Mus., XI,
 1888, 551, Green Turtle Cay, Bahamas.
- 3461.** *Sparisoma aurofrenatum* (Cuvier and Valenciennes).
 West Indies.
Scarus aurofrenatus Cuvier and Valenciennes, Hist. Nat. Poiss., XIV,
 1839, 142 (191), San Domingo.
Scarus miniofrenatus Poey, Memorias, II, 1860, 279, Cuba.
- 3462.** *Sparisoma oxybrachium* (Poey).
 Cuba.
Scarus oxybrachius Poey, Synopsis, 1868, 342, Habana.
- 3463.** *Sparisoma abildgaardi* (Bloch). *Red parrotfish; Roseback parrot.*
 West Indies south to Brazil; Tortugas; generally common.
Sparus abildgaardi Bloch, Nat. Ausl. Fische, V, 1791, 22, pl. 259,
 America.
Vieja Parra, Descr. Dif. Piezas Hist. Nat., 1787, 58, pl. 28, fig. 2, Cuba.
Scarus coccineus Bloch and Schneider, Syst. Ichth., 1801, 289, Cuba;
 after Parra.
Sparus aureo-ruber Lacépède, Hist. Nat. Poiss., IV, 1803, 55, 163,
 Martinique; on a drawing by Plumier.
Scarus amplus Ranzani, Nov. Comm. Ac. Scient. Inst. Bonon., V,
 1842, 324, pl. 25, Brazil (fide Guichenot).
Scarus erythrinoides Guichenot, Mem. Soc. Sci. Nat. Cherbourg, XI,
 1865, 10, San Domingo.
- 3464.** *Sparisoma distinctum* (Poey).
 Cuba.
Scarus distinctus Poey, Memorias, II, 1861, 423, Habana.
- 3465.** *Sparisoma chrysopterum* (Bloch and Schneider). *Blue parrotfish;*
Vieja.
 West Indies, south to Bahia.
Scarus chrysopterus Bloch and Schneider, Syst. Ichth., 1801, 286, pl. 57,
 American seas.
Vieja Parra, Descr. Dif. Piezas Hist. Nat., 1787, 59, pl. 28, fig. 4, Cuba.
Scarus chloris Bloch and Schneider, Syst. Ichth., 1801, 289, Cuba.
Scarus lateralis Poey, Memorias, II, 1860, 219, Cuba.
Scarus spinidens Guichenot, Mem. Soc. Sci. Nat. Cherbourg, XI,
 1865, 15, Bahia.
- 3466.** *Sparisoma lorito* Jordan and Swain.
 West Indies.
Sparisoma lorito Jordan and Swain, Proc. U. S. Nat. Mus., VII, 1884,
 95, Habana.
- 3467.** *Sparisoma viride* (Bonnaterre). *Green parrotfish.*
 Tortugas to West Indies.
Scarus viridis Bonnaterre, Enc. Meth., 1788, 96, Bahamas; after
 Catesby.
Piscis viridis bahamensis Catesby, Nat. Hist. Car., II, 1743, 29, pl. 29,
 Bahamas.
Scarus melanotis Bleeker, Verslagen Koning. Akad. Wetenschappen,
 XIV, 1862, 126, St. Croix.
Scarus (Catesby) Lacépède, Hist. Nat. Poiss., IV, 1803, 16; after
 Catesby.

- 3468. *Sparisoma flavescens* (Bloch and Schneider). *Mud parrot; Vieja colorada; Vieja muger; Parrotfish.***
 West Indies; Key West to Rio Janeiro; Woods Hole, Mass.
Scarus flavescens Bloch and Schneider, Syst. Ichth., 1801, 290, Cuba; after Parra.
Vieja Parra, Descr. Dif. Piezas Hist. Nat., 1787, 59, pl. 28, fig. 4, Cuba.
Scarus squamidus Poey, Memorias, II, 1860, 218, Cuba.
- 3469. *Sparisoma rubripinne* (Cuvier and Valenciennes).**
 West Indies.
Scarus rubripinnis Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 147 (199), San Domingo.
Scarus virens Cuvier and Valenciennes, loc. cit., 1839, 150 (203), Porto Rico; Martinique.
Scarus circumnotatus Poey, Memorias, II, 1861, 423, Cuba.
Scarus truncatus Poey, Synopsis, 1868, 339, Cuba.
?Scarus emarginatum Poey, Synopsis, 1868, 340, Cuba.
- 3470. *Sparisoma strigatum* (Günther).**
Scarus strigatus Günther, Cat., IV, 1862, 212, locality unknown, but as the genus *Sparisoma* is chiefly confined to American waters, this species is probably American.
- 3471. *Sparisoma frondosum* (Cuvier).**
 West Indies, south to Brazil.
Scarus frondosus Cuvier, in Agassiz, Spix, Pisc. Brasil., 1829, 98, Bahia.
Scarus aracanga Günther, Cat., IV, 1862, 209, Jamaica.
- 3472. *Sparisoma brachiale* (Poey).**
 Cuba.
Scarus brachialis Poey, Memorias, II, 1861, 345, Cuba.
?Scarus humeralis Poey, loc. cit., 422, Habana.
- 3473. *Sparisoma maschalespilos* (Bleeker).**
 Surinam.
Scarus maschalespilos Bleeker, Verslagen König. Akad. Vet., XIV, 1862, 127, Surinam.

Family 254. SCARIDÆ

Genus 1199. SCARUS Forskål.

Callyodon Gronow, Zoophyl., I, 1763, 72 (*Scarus croicensis* Bloch); work not binomial.
Scarus Forskål, Descr. Anim., 1775, 25 (*Scarus croicensis* Bloch); not of Gronow.
Hemistoma Swainson, Classn. Fish., etc., II, 1839, 226 (*Scarus pepo* Bennett = *H. reticulata* Swainson).
Petronason Swainson, loc. cit., 226 (*Coryphæna psittacus* Linnæus).
Erychthys Swainson, loc. cit., 226 (*Labrus croicensis* Bloch).
Chlorurus Swainson, loc. cit., 227 (*Scarus gibbus* Rüppell).

- 3474. *Scarus bollmani* Jordan and Evermann.**
 Gulf of Mexico in moderately deep water.
Scarus bollmani Jordan and Evermann, Proc. U. S. Nat. Mus., IX, 1886, 470, snapper banks off Tampa Bay, Fla.
- 3475. *Scarus punctulatus* Cuvier and Valenciennes.**
 West Indies; Tortugas.
Scarus punctulatus Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 144 (195), Martinique.
Scarus diadema Cuvier and Valenciennes, loc. cit., 146 (196), Martinique.
Pseudoscarus tæniopterus Günther, Cat., IV, 1862, 226, Trinidad.
- 3476. *Scarus tæniopterus* Desmarest.**
 West Indies.
Scarus tæniopterus Desmarest, Dict. Classique, XV, 1831, 244, pl. 12, Cuba.
Scarus vetula Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 143 (193), St. Thomas; not of Bloch and Schneider.
Scarus virginalis Jordan and Swainson, Proc. U. S. Nat. Mus., VII, 1884, 88, Habana.

3477. ***Scarus trispinosus*** Cuvier and Valenciennes.*Brazil.**Scarus trispinosus* Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 135 (182), *Brazil.**Scarus quadrispinosus* Cuvier and Valenciennes, loc. cit., 146 (197), *Martinique.*3478. ***Scarus aracanga*** Günther.*West Indies.**Scarus aracanga* Günther, Cat., IV, 1862, 209, *Jamaica.**?Pseudoscarus aracanga* Günther, loc. cit., 227, *Jamaica.*3479. ***Scarus cuzamilæ*** Bean.*Cozumel Island, Yucatan.**Scarus cuzamilæ* Bean, Bull. U. S. Fish. Com., VIII, 1888, 196, *Cozumel, Yucatan.*3480. ***Scarus vetula*** Bloch and Schneider. *Mudfish; Vieja; Oldwife.**Tortugas; West Indies.**Vieja Parra*. Dif. Piezas, etc., 1787, 58, pl. 28, fig. 1, *Habana.**Scarus vetula* Bloch and Schneider, Ichth., 1801, 289, *Cuba.**Scarus superbus* Poey, Memorias, II, 1860, 218, *Cuba.*3481. ***Scarus gnathodus*** (Poey).*Cuba.**Pseudoscarus gnathodus* Poey, Repertorio, II, 1867, 240, *Habana.*Subgenus **CALLYODON** Gronow.3482. ***Scarus croicensis*** Bloch. *Bullion.**West Indies to Key West and Woods Hole, Mass.**Scarus croicensis* Bloch, Nat. Ausl. Fische, IV, 1790, 27, *St. Croix.**Scarus insulæ-santaæ-crucis* Bloch and Schneider, Syst. Ichth., 1801, 285, *St. Croix.**Scarus alternans* Cuvier and Valenciennes, Hist. Nat. Poiss., 1839, 148 (200), *Martinique.**Pseudoscarus lineolatus* Poey, Repertorio, II, 1867, 239, *Cuba.*3483. ***Scarus evermanni*** Jordan.*Gulf of Mexico, in moderately deep water.**Scarus evermanni* Jordan, Proc. U. S. Nat. Mus., IX, 1886, 469, *Snapper Banks off Tampa Bay, in moderately deep water.*3484. ***Scarus flavomarginatus*** Cuvier and Valenciennes.*Martinique.**Scarus flavo-marginalis* Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 150 (202), *Martinique.*3485. ***Scarus acutus*** Poey.*Cuba.**Scarus acutus* Poey, Memorias, II, 1861, 216, *Habana.*3486. ***Scarus cæruleus*** (Bloch). *Blue parrotfish; Loro; Clamagore.**West Indies; Chesapeake Bay.**Coryphæna cærulea* Bloch, Naturgesch. Ausl. Fische, II, 1786, 120 (148), pl. 176, probably *Martinique*; in part; after Catesby.*Loro Parra*, Descr. Dif. Piezas Hist. Nat., 1787, 57, pl. 27, fig. 1, *Cuba.**Trompa Parra*, loc. cit., fig. 2, *Cuba.**Scarus trilobatus* Lacépède, Hist. Nat. Poiss., IV, 1803, 21, *Martinique.**?Scarus holocyaneos* Lacépède, loc. cit., 45, *Martinique.**Scarus obtusus* Poey, Memorias, II, 1861, 217, *Cuba.**Scarus nuchalis* Poey, loc. cit., 220, *Cuba.*3487. ***Scarus compressus*** (Osburn and Nichols).*Gulf of California.**Callyodon compressus* Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 170, fig. 12, *Concepcion Bay, Gulf of California.*3488. ***Scarus noyesi*** Heller and Snodgrass.*Gulf of California to Galapagos Islands.**Scarus noyesi* Heller and Snodgrass, Proc. Wash. Acad. Sci., V, 1903, 206, *Tagus Cove; Albemarle Island.*

- 3489. *Scarus emblematicus* Jordan and Rutter.**
Jamaica.
Scarus emblematicus Jordan and Rutter, Proc. Acad. Nat. Sci. Phila., XLIX, 1897, 122, Jamaica.
- Genus 1200. PSEUDOSCARUS Bleeker.**
Pseudoscarus Bleeker, Versl. Akad. Vet. Amsterdam, Scaroiden, XII, 1861, 230 (*P. microrrhinos* Bleeker).
Loro Jordan and Evermann, Check List, 1895, 418 (*Scarus guacamaia* Cuvier).
- Subgenus PSEUDOSCARUS Bleeker.**
- 3490. *Pseudoscarus caelestinus* (Cuvier and Valenciennes).** *Loro.*
West Indies.
Scarus caelestinus Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 134 (180), St. Thomas.
- 3491. *Pseudoscarus microps* (Osburn and Nichols).**
Gulf of California.
Callichodon microps Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 170, fig. 12, Santa Catalina Island, Gulf of California.
- 3492. *Pseudoscarus simplex* Poey.**
Cuba.
Pseudoscarus simplex Poey, Repertorio, I, 1867, 185, Habana.
- 3493. *Pseudoscarus pleianus* (Poey).**
St. Thomas; only the type known.
Scarus pleianus Poey, Memorias, II, 1861, 393, St. Thomas, West Indies.
- 3494. *Pseudoscarus guacamaia* (Cuvier).** *Guacamaia; Green parrotfish.*
West Indies, north to Florida Keys, south to Rio de Janeiro; straying to Argentine.
Guacamaya Parra, Descr. Dif. Piezas Hist. Nat., 1787, p. 54, pl. 26, Cuba.
Scarus guacamaia Cuvier, Règne Anim., ed. 2, II, 1829, 265; after Parra.
Scarus turchesius Cuvier and Valenciennes, Hist. Nat. Poiss., XIV, 1839, 134 (181), Porto Rico.
Scarus rostratus Poey, Memorias, II, 1861, 221, Cuba.
- 3495. *Pseudoscarus perrico* (Jordan and Gilbert).** *Perrico.*
Pacific coast of Mexico, from La Paz to Mazatlan.
Scarus perrico Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 357, Mazatlan.
- 3496. *Pseudoscarus plumbeus* Bean.**
Bermudas.
Pseudoscarus plumbeus Bean, Proc. Biol. Soc. Wash., XXV, 1912, 125, Bermudas.

Order VV. GOBOIDEA

Family 255. ELEOTRIDÆ. Sleepers

- Genus 1201. GOBIOMORUS Lacépède.**
Gobiomorus Lacépède, Hist. Nat. Poiss., II, 1800, 583 (*G. dormitor* Lacépède); first restriction to this species.
Philypnus Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 255 (*Platycephalus dormitator* Bloch).
Lembus Günther, Cat., I, 1859, 505 (*L. maculatus* Günther).
- 3497. *Gobiomorus dormitor* Lacépède.** *Sleeper; Guavina.*
Streams of the West Indies and the Atlantic shores of Central America, Mexico and Surinam.
Gobiomorus dormitor Lacépède, Hist. Nat. Poiss., II, 1800, 583, Martinique; from a drawing by Plumier.
Guavina Parra, Descr. Dif. Piezas Hist. Nat. Cuba, 1787, 105, lam. 39, fig. 1, Habana.
Platycephalus dormitator Bloch and Schneider, Ichth., 1801, 60, Martinique.
Eleotris longiceps Günther, Proc. Zool. Soc. Lond., 1864, 151, Nicaragua.
Eleotris dormitatrix Cuvier, Règne Anim., ed. 2, II, 1829, 246, Antilles.

- 3498. *Gobiomorus lateralis* (Gill). *Aboma de mar.***
 Streams of Pacific coast of Mexico and Central America; from Sonora to Panama.
Philypnus lateralis Gill, Proc. Acad. Nat. Sci. Phila., XII, 1860, 123, Cape San Lucas.
- Genus 1202. DORMITATOR Gill.**
Prochilus Cuvier, Règne Anim., ed. 1, II, 1817, 294 (*Sciæna maculata* Bloch); name preoccupied.
Dormitator Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 240 (*D. gundlachi* Poey).
- 3499. *Dormitator maculatus* (Bloch). *Guavina; Mapo; Paneca; Mapiro; Striped sleeper.***
 Both coasts of tropical America, ranging from South Carolina through the West Indies to Para, Cape San Lucas, and Panama.
Sciæna maculata Bloch, Ichth., 1785, pl. 299, fig. 2, West Indies.
Eleotris mugiloides Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 170 (226), Martinique; Surinam.
Eleotris sima Cuvier and Valenciennes, loc. cit., 174 (232), Vera Cruz.
Eleotris latifrons Richardson, Voy. *Sulphur*, Fishes, 1844, 57, pl. 35, figs. 4 and 5, locality unknown, supposed to be from Pacific coast of Central America.
Eleotris grandisquama Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 173 (229), America.
Eleotris somnolentus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 169, near mouth of Rio Grande.
Eleotris omocyanus Poey, Memorias, II, 1861, 269, Habana.
Dormitator microphthalmus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 170, Panama.
Dormitator gundlachi Poey, Synopsis, 1868, 396, Cuba.
Dormitator lineatus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 271, Savannah.
- Genus 1203. GUAVINA Bleeker.**
Guavina Bleeker, Esquisse d'un Syst. Nat. Gobiod., in Archiv Neerl. Sci., IX, 1874 (*Eleotris guavina* Cuvier and Valenciennes).
- 3500. *Guavina guavina* (Cuvier and Valenciennes). *Guarubaco; Guavina.***
 East coast of tropical America, Cuba to Rio Janeiro.
Eleotris guavina Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 168 (223) Martinique.
- Genus 1204. LEPTOPHILYPNUS Meek and Hildebrand.**
Leptophilypnus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 361 (*L. fluviatilis* Meek and Hildebrand).
- 3501. *Leptophilypnus fluviatilis* Meek and Hildebrand.**
 Fresh waters of Panama.
Leptophilypnus fluviatilis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 361, Mindi, Canal Zone, Panama.
- Genus 1205. ELEOTRIS Schneider.**
Eleotris Schneider, Syst. Ichth., ed. Bloch, 1801, 65 (*Gobius pisonis*, etc., Gmelin).
Culius Bleeker, Esquisse, Gobioides, 1874, 304 (*C. fuscus* Bleeker).
- 3502. *Eleotris pisonis* (Gmelin). *Guavina; Tetard; Sleeper.***
 Streams of the West Indies; southern Florida to Brazil.
Gobius pisonis Gmelin, Syst. Nat., 1789, 1206, Rio Almendares, Cuba; based on *Eleotris* of Gronow.
Amore pixuma Marcgrave, Hist. Brasil., IV, 1648, 166, Brazil.
Eleotris capite plagioplateo Gronow, Mus. Ichth., II, 1757, 168; after Marcgrave.
Gobius amoreo Walbaum, Artedi Pisc., III, 1792, 205; based on *Eleotris* of Gronow.
Eleotris gyriinus Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 166 (220), pl. 356, Martinique; San Domingo; Surinam.
Eleotris (Culius) belizianus Sauvage, Bull. Soc. Philom. Paris, ser. 7, IV, 1879, 55, Belize, Cayenne.

3503. **Eleotris perniger** (Cope).
 West Indies, south to Rio Janeiro.
Culius perniger Cope, Trans. Amer. Phil. Soc., XIV, 1870, 473, St. Martins, West Indies.
3504. **Eleotris amblyopsis** (Cope).
 Atlantic coast of America, from Charleston, S. C., to Surinam; streams of West Indies.
Culius amblyopsis Cope, Trans. Amer. Phil. Soc., XIV, 1870, 473, Surinam.
3505. **Eleotris abacurus** Jordan and Gilbert.
 Coast of South Carolina.
Eleotris abacurus Jordan and Gilbert, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 228, Charleston, S. C.
3506. **Eleotris pictus** Kner and Steindachner. *Guavina*.
 Streams about the Gulf of California, from Sonora south to Panama; Rio Presidio, Mazatlan.
Eleotris pictus Kner and Steindachner, Abh. Akad. Wiss. Wien, X, 1864, 18, pl. 3, fig. 1, Rio Bayano, near Panama.
Culius xquidens Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 461, Rio Presidio, near Mazatlan.
3507. **Eleotris tubularis** Heller and Snodgrass.
 Cocos Island.
Eleotris tubularis Heller and Snodgrass, Proc. Wash. Acad. Sci., V, 1903, 210, Pl. X, Cocos Island.
- Genus 1206. **ALEXURUS** Jordan.
Alexurus Jordan, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 511 (*A. armiger* Jordan and Richardson); perhaps inseparable from *Eleotris*.
3508. **Alexurus armiger** Jordan.
 La Paz, Lower California, to Panama.
Alexurus armiger Jordan, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 511, pl. 48, La Paz, Lower California.
- Genus 1207. **HEMIELEOTRIS** Meek and Hildebrand.
Hemieleotris Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 364 (*Eleotris latifasciatus* Meek and Hildebrand).
3509. **Hemieleotris latifasciatus** (Meek and Hildebrand).
 Streams of Panama.
Eleotris latifasciatus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1912, 68, Rio Cardenais, Panama.
- Genus 1208. **EROTELIS** Poey. *Esmeraldas de Mar*.
Erotelis Poey, Memorias, II, 1861, 273 (*E. valenciennesi* Poey = *Eleotris smaragdus* Cuvier and Valenciennes).
3510. **Erotelis smaragdus** (Cuvier and Valenciennes). *Esmralda negra*; *Esmralda de mar*.
 Key West and Cuba; coral shores among algae.
Eleotris smaragdus Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 173 (231), Cuba.
Erotelis valenciennesi Poey, Memorias, II, 1861, 273, Cuba.
- Genus 1209. **MICROELEOTRIS** Meek and Hildebrand.
Microeleotris Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 362 (*M. panamensis* Meek and Hildebrand).
3511. **Microeleotris panamensis** Meek and Hildebrand.
 Pacific slope of central Panama.
Microeleotris panamensis Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 363, Rio Juan Diaz, Panama.
3512. **Microeleotris mindii** Meek and Hildebrand.
 Mindi, near Atlantic coast of Canal Zone.
Microeleotris mindii Meek and Hildebrand, Field Mus. Nat. Hist., Zool., X, 1916, 364, brackish creek at Mindi, Canal Zone.

Genus 1210. EVIOTA Jenkins.

Eviota Jenkins, Bull. U. S. Fish Com., XXII, 1902, 501 (*Eviota epiphanes* Jenkins).

3513. *Eviota personata* Jordan and Thompson.

Coral reefs about the Tortugas, Fla.; Garden Key.

Eviota personata Jordan and Thompson, Bull. U. S. Bur. Fish., XXIV, 1904, 251, fig. 3, Garden Key, Fla.

Genus 1211. IOGLOSSUS Bean.

Ioglossus Bean, in Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 297 (*I. calliurus* Bean).

3514. *Ioglossus calliurus* Bean.

Gulf of Mexico; Snapper Banks off Pensacola, Fla.

Ioglossus calliurus Bean, in Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 297, Pensacola, Fla.

Genus 1212. GYMNELEOTRIS Bleeker.

Gymneleotris Bleeker, Esquisse d'un Syst. Nat. des Gobioid., in Archiv Nér. Sci., IX, 1874, 304 (*Eleotris seminuda* Günther).

3515. *Gymneleotris seminuda* (Günther).

Panama.

Eleotris seminudus Günther, Proc. Zool. Soc. Lond., 1864, 24, pl. 4, figs. 2, 2a, Panama.

Genus 1213. PYCNOMMA Rutter.

Pycnomma Rutter, Proc. Calif. Acad. Sci., 3d ser., III, 1904, 252 (*P. semisquamatum* Rutter).

3516. *Pycnomma semisquamatum* Rutter.

Gulf of California.

Pycnomma semisquamatum Rutter, Proc. Calif. Acad. Sci., 3d ser., III, 1904, 252, Pl. XXIV, Gulf of California.

Genus 1214. CHRIOLEPIS Gilbert.

Chriolepis Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 557 (*C. minutillus* Gilbert).

3517. *Chriolepis minutillus* Gilbert.

Gulf of California.

Chriolepis minutillus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 558, Gulf of California, at Albatross station 2825.

Family 256. GOBIIDÆ. Gobies

Genus 1215. LOPHOGOBIUS Gill. Crested gobies.

Lophogobius Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 240 (*Gobius cristagalli* Valenciennes, in Cuvier and Valenciennes = *Gobius cyprinoides* Pallas).

3518. *Lophogobius cyprinoides* (Pallas).

West Indies; Cuba.

Gobius cyprinoides Pallas, Spic., Zool., VIII, 1770, 17, pl. 1, fig. 5, "Amboina."

Gobius crista-galli Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 98 (130), Habana.

Genus 1216. EVORTHODUS Gill.

Evorthodus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 195 (*E. breviceps* Gill).

3519. *Evorthodus breviceps* Gill.

Trinidad and Surinam.

Evorthodus breviceps Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 195, Trinidad.

Genus 1217. CTENOGOBIUS Gill.

Ctenogobius Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 374, 430 (*C. fasciatus* Gill).

3520. *Ctenogobius fasciatus* Gill.

West Indies.

Ctenogobius fasciatus Gill, Ann. Lyc. Nat. Hist. N. Y., VI, 1858, 376, Trinidad.

Genus 1218. RHINOGOBius ³⁶ Gill.

Rhinogobius Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 145 (*R. similis* Gill).

Coryphopterus Gill, loc. cit., XV, 1863, 263 (*C. glaucofrænum* Gill).

3521. *Rhinogobius glaucofrænum* (Gill).

Florida Keys; Tortugas.

Coryphopterus glaucofrænum Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 263, coast of the State of Washington; evidently an error.

3522. *Rhinogobius manglicola* (Jordan and Starks).

Pacific coast of Mexico at Mazatlan.

Gobius manglicola Jordan and Starks, in Jordan, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 495, Mazatlan, Mexico.

3523. *Rhinogobius quadriporus* (Cuvier and Valenciennes).

Surinam.

Gobius quadriporus Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 66 (87), Surinam.

3524. *Rhinogobius stigmaturus* (Goode and Bean).

Florida Keys.

Gobius stigmaturus Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 418, no type locality given, but specimens probably from Florida Keys.

3525. *Rhinogobius shufeldti* (Jordan and Eigenmann).

Gulf coast of United States.

Gobius shufeldti Jordan and Eigenmann, Proc. U. S. Nat. Mus., IX, 1886, 495, New Orleans.

3526. *Rhinogobius boleosoma* (Jordan and Gilbert).

Gulf of Mexico, Pensacola to Key West.

Gobius boleosoma Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 295, Laguna Grande, Pensacola, Fla.

3527. *Rhinogobius encæomus* (Jordan and Gilbert).

South Carolina to Key West.

Gobius encæomus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 611, Charleston, S. C.

3528. *Rhinogobius eigenmanni* (Garman).

Off Key West.

Gobius eigenmanni Garman, Bull. Lab. Nat. Sci. State Univ. Iowa, IV, No. 1, 1896, 88, pl. 3, fig. 1, off Key West in 60 fathoms.

3529. *Rhinogobius curaçao* (Metzelaar).

Caribbean Sea.

Gobius curaçao Metzelaar, Trop. Atl. Viss., 1919, 136, fig. 38, Antilles.

3530. *Rhinogobius mowbrayi* Bean.

Bermudas to Caribbean Sea.

Rhinogobius mowbrayi Bean, Field Mus. Nat. Hist., Zool., VII, 1906, 81, Challenger Bank, 28 fathoms, rock cavity, Bermudas.

3531. *Rhinogobius tortugæ* (Jordan).

Dry Tortugas, Garden Key, West Indies.

Ctenogobius tortugæ Jordan, Bull. U. S. Fish Com., XXII, 1902, 541, pl. 1, Garden Key, Tortugas.

3532. *Rhinogobius parvus* (Meek).

Rio San Francisco, La Antigua, Mexico.

Gobius parvus Meek, Field Col. Mus., Zool., III, 1902, 121, La Antigua.

3533. *Rhinogobius claytonii* (Meek).

Rio San Francisco, La Antigua, Mexico.

Gobius claytonii Meek, Field Col. Mus., Zool., III, 1902, 121, La Antigua.

3534. *Rhinogobius garmani* (Eigenmann and Eigenmann).

Dominica; Fort de France; Martinique; St. Kitts.

Gobius garmani Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888, 61, Dominica; Fort de France, Martinique, and St. Kitts.

* Some of the species named below may belong to *Ctenogobius*, from which this genus differs but slightly.

Genus 1219. GOBIUS Linnæus.

Gobius Linnæus, Syst. Nat., Ed. X, I, 1758, 263 (*G. niger* Linnæus).

3535. *Gobius translucens* Nichols.

Porto Rico.

Gobius translucens Nichols, Bull. Amer. Mus. Nat. Hist., XXXIV, 1915, 145, fig. 2, near San Antonio Bridge, San Juan, Porto Rico.

3536. *Gobius hipoliti* Metzelaar.

West Indies.

Gobius hipoliti Metzelaar, Bijdr. Dierk. Amsterdam, XXII, 1922, 138, with figure, West Indies.

Genus 1220. BATHYGOBIUS Bleeker.

*Bathygobius*⁷ Bleeker, Arch. Neerl. Acad. Sci., XIII, 1878, 54 (*Gobius nebulos-punctatus* Rüppell=*Gobius fuscus* Rüppell).

Mapo Smitt, Oefv. Kong. Vet. Akad. Stockholm, 1899, 542 (*Gobius soporator* Cuvier and Valenciennes).

3537. *Bathygobius soporator* (Cuvier and Valenciennes). *Sleeper; Mapo; Caiman; Oopo.*

Barbados, Pará, Itapuana, Cuba, Bahamas, Orange Key, Bahia, Pernambuco, St. Thomas, Charleston, Tortugas, Florida Keys, Martinique, São Mathias, Curaçao, Rio de Janeiro, Rio Doce.

Gobius soporator Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 42 (56), Martinique.

?*Gobius albopunctatus* Cuvier and Valenciennes, loc. cit., 43 (57), Isle of France.

?*Gobius padangensis* Bleeker, Nat. Tijds. Ned. Ind., I, 1850, 249, Padang.

Gobius catulus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 169, St. Joseph Island, Tex.

?*Gobius breviceps* Blyth, Proc. Asia. Soc. Beng., 1858, 271, Great Andaman Island (in Jour., XXVII, 1858).

Gobius mapo Poey, Memorias, II, 1861, 277, Cuba.

Gobius lacertus Poey, loc. cit., 278, Cuba.

Gobius carolinensis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 268, Charleston, S. C.

Gobius brunneus Poey, Synopsis, 1868, 393, Habana.

3538. *Bathygobius fuscus* (Rüppell).

Red Sea to Galapagos Islands.

'*Gobius fuscus* Rüppell, Atl. Reis. Nord. Af. Fish., 1828, 137, Red Sea.

Gobius punctillatus Rüppell, loc. cit., 138, Red Sea.

Gobius nebulos-punctatus Rüppell, Neue Wirbelthiere, Fische, 1835, 139, Red Sea.

Gobius lineatus Jenyns, Zool. Voy. Beagle, 1842, 95, pl. 19, fig. 2, Galapagos Islands.

Gobius homocyanus Vaillant and Sauvage, Rev. Mag. Zool., III, 1875, 280, Sandwich [Hawaiian] Islands.

Gobius sandwicensis Günther, Rept. Shore Fishes, Challenger, I, 1880, pt. 6, 60, Honolulu.

Genus 1221. RHINOGOBIOPS Hubbs.

Rhinogobiops Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 169, Feb. 17, 1926, p. 1 (*Gobius nicholsii* Bean).

3539. *Rhinogobiops nicholsii* (Bean).

Coast of British Columbia, south to southern California.

Gobius nicholsii Bean, Proc. U. S. Nat. Mus., IV, 1881, 469, Departure Bay, British Columbia.

Genus 1222. GOBIONELLUS Girard.

Gobionellus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 168 (*G. hastatus* Girard).

Smaragdus Poey, Memorias, II, 1860, 280 (*Gobius smaragdus* Cuvier and Valenciennes).

⁷ An ill-chosen name for these shallow-water gobies.

- 3540.** *Gobionellus oceanicus* (Pallas). *Esmeralda; Endormi; Emerande; Bacalhao sabara.*
 Gulf coast of United States, southward through the West Indies.
Gobius oceanicus Pallas, Spic., VIII, 1770, 4, locality unknown; after Gronow.
Gobius lanceolatus Bloch, Fische Deutschlands, II, 1783, 8, pl. 38, fig. 1, Martinique.
Gobius bacalaus Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 90 (119), Surinam; Cayenne.
- 3541.** *Gobionellus hastatus* Girard. *Emerald fish; Sharp-tailed goby;*
 Gulf of Mexico.
Gobionellus hastatus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 168, St. Joseph Island, Tex.
- 3542.** *Gobionellus smaragdus* (Cuvier and Valenciennes). *Esmeralda.*
 West Indies, south to Rio Janeiro, north to St. Augustine, Fla., and to Charleston, N. C.
Gobius smaragdus Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 91 (120), Cuba.
Smaragdus valencienensis Poey, Memorias, II, 1861, 280, Cuba; after Valenciennes.
- 3543.** *Gobionellus sagittula* (Günther).
 Gulf of California, south to Panama.
Euctenogobius sagittula Günther, Proc. Zool. Soc. Lond., 1861, pt. 3, 371, 372, Pacific coast of Central America.
Gobius longicaudus Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 146, Guaymas.
- 3544.** *Gobionellus strigatus* (O'Shaughnessy).
 Coast of Surinam.
Euctenogobius strigatus O'Shaughnessy, Ann. Mag. Nat. Hist., ser. IV, XV, 1875, 145, Surinam.
Gobius kraussi Steindachner, Ichth. Beitr., VIII, 1879, 16, Surinam.
- 3545.** *Gobionellus bayamonensis* (Evermann and Marsh).
 Porto Rico.
Gobius bayamonensis Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 355, San Juan Market, Porto Rico.
- 3546.** *Gobionellus lyricus* (Girard).
 Gulf of Mexico, from Galveston to Cuba and the Lesser Antilles.
Gobius lyricus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 169, Brazos Santiago, Tex.
Gobius wurdemanni Girard, loc. cit., 169, Brazos Santiago.
Smaragdus costalesii Poey, Memorias, II, 1861, 280, Habana.
- 3547.** *Gobionellus stigmaticus* (Poey).
 Coast of North Carolina, Florida Keys, West Indies, southward to Rio Janeiro; common at Habana.
Smaragdus stigmaticus Poey, Memorias, II, 1861, 281, Cuba.
- 3548.** *Gobionellus microdon* (Gilbert).
 San Juan Lagoon, west coast of Mexico.
Gobius microdon Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 554, San Juan Lagoon, north of Rio Ahomé, Mexico.
- Genus 1223. **ZONOGOBIUS** Bleeker.
Zonogobius Bleeker, Esquisse Syst. Nat. Gobioiden, 1874, 323 (*Gobius semifasciatus* Kner).
- 3549.** *Zonogobius zebra* (Gilbert).
 Pacific coast of Mexico and southern California.
Gobius zebra Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 73, *Albatross* station 2989, Pacific coast of Mexico.
- Genus 1224. **EUCTENOGOBIUS** Gill
Euctenogobius Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1859, VII, 45 (*E. badius* Gill).

3550. *Euctenogobius badius* Gill.

Mouth of Amazon; recorded also from Martinique.
Euctenogobius badius Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1859, 45,
 Amazon.

Gobius boscii Sauvage, Bull. Soc. Philom. Paris, IV, 1880, 44, Mar-
 tinique; not *Gobius boscii* Lacépède.

3551. *Euctenogobius poeyi* (Steindachner).

Barbados.
Gobius poeyi Steindachner, Ichth. Notizen, VI, 1867, 44, Barbados.

Genus 1225. *LYTHRYPNUS* Jordan and Evermann.

Lythrypnus Jordan and Evermann, Check List, 1895, 458 (*Gobius dalli*
 Gilbert).

3552. *Lythrypnus dalli* (Gilbert).

Catalina Harbor, Calif.
Gobius dalli Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 73, Avalon,
 Catalina Harbor, Calif., in 35 fathoms.

Genus 1226. *XENOGOBIUS* Metzelaar.

Xenogobius Metzelaar, Trop. Atlant. Vissch., 1919, 140 (*X. weberi* Metze-
 laar).

3553. *Xenogobius weberi* Metzelaar.

Caribbean Sea.
Xenogobius weberi Metzelaar, Trop. Atlant. Vissch., 1919, 140, fig. 40,
 Curaçao.

Genus 1227. *ELACATINUS* Jordan.

Elecatinus Jordan, Bull. U. S. Fish Com., XXII, 1902, 542 (*E. oceanops*
 Jordan).

3554. *Elacatinus oceanops* Jordan.

Garden Key, Tortugas.
Elacatinus oceanops Jordan, Bull. U. S. Fish Com., XXII, 1902, 542,
 pl. 2, Garden Key, Fla.

Genus 1228. *CHONOPHORUS* Poey.

Chonophorus Poey, Memorias, II, 1860, 274 (*C. bucculentus* Poey =
Gobius banana Cuvier and Valenciennes; published in July, 1860, and
 therefore presumably older in date than *Awaous*).

Awaous Steindachner, Ver. Mat. Phrs., XLII, 1860, 289 (*Gobius ocellaris*
 Cuvier and Valenciennes; in an article presented on July 12, 1860).

3555. *Chonophorus banana* (Cuvier and Valenciennes). *Guavina hoyera*;
Aboma de rio.

West Indies; both coasts of Mexico, south to Brazil; Cuba; Sinaloa;
 common; Lower California to Panama.

?*Gobius tajasicus* Lichtenstein, Abh. Akad. Wiss. Berlin, 1822, 273,
 Brazil; not *Tajaasicus* Marckgrave.

Gobius banana Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837,
 78 (103), San Domingo.

Gobius martinicus Cuvier and Valenciennes, loc. cit., 79 (105), Marti-
 nique.

Chonophorus bucculentus Poey, Memorias, II, 1860, 275, Cuba.

Chonophorus contractus Poey, loc. cit.; Cuba.

Gobius dolichocephalus Cope, Trans. Amer. Phil. Soc. Phila., XIII,
 1869, 403, near Orizaba, Mexico.

Euctenogobius latus O'Shaughnessy, Ann. Mag. Nat. Hist., ser. 4, XV,
 1875, 146, Bahia.

3556. *Chonophorus mexicanus* (Günther).

Fresh-water streams of Atlantic slope of Mexico.

Gobius mexicanus Günther, Cat., III, 1861, 61, Mexico.

3557. *Chonophorus nelsoni* (Evermann).

Streams of Sinaloa.

Awaous nelsoni Evermann, Proc. Biol. Soc. Wash., XII, 1898, 3, fresh-
 water pools at Rosario, Sinaloa, Mexico.

Genus 1229. *GNATHOLEPIS* Bleeker.

Gnatholepis Bleeker, Esquisse Syst. Nat. Gobioiden, 1874, 318 (*Gobius*
anjerensis Bleeker).

3558. *Gnatholepis thompsoni* Jordan.
Bush Key, Tortugas.
Gnatholepis thompsoni Jordan, Bull. U. S. Fish Com., XXII, 1902, 541,
pl. 1, Bush Key, Tortugas.
- Genus 1230. **BOLLMANNIA** Jordan.
Bollmannia Jordan, Proc. U. S. Nat. Mus., XII, 1889, 164 (*B. chlamydes*
Jordan).
3559. *Bollmannia chlamydes* Jordan.
Pacific Ocean off coast of Colombia.
Bollmannia chlamydes Jordan, Proc. U. S. Nat. Mus., XII, 1889, 164,
Pacific Ocean off coast of Colombia.
3560. *Bollmannia ocellata* Gilbert.
Gulf of California.
Bollmannia ocellata Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 555,
northern part of Gulf of California at *Albatross* stations 3031 and 3035
in 30 and 33 fathoms.
3561. *Bollmannia macropoma* Gilbert.
Gulf of California.
Bollmannia macropoma Gilbert, Proc. U. S. Nat. Mus., XIV, 1891,
556, near La Paz, Gulf of California, at *Albatross* station 2996 in
112 fathoms.
3562. *Bollmannia stigmatura* Gilbert.
Northern part of the Gulf of California.
Bollmannia stigmatura Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 556,
Gulf of California at *Albatross* stations 3016 and 3017 in 76 and 58
fathoms.
3563. *Bollmannia boqueronensis* Evermann and Marsh.
Porto Rico.
Bollmannia boqueronensis Evermann and Marsh, Rept. U. S. Fish Com.,
XXV, 1899 (Dec. 19), 356, Ensenada del Boqueron, Porto Rico.
- Genus 1231. **ABOMA** Jordan and Starks.
Aboma Jordan and Starks, Fishes of Sinaloa, Proc. Calif. Acad. Sci.,
2d ser., V, 1895, 497 (*A. ethostoma* Jordan and Starks).
3564. *Aboma ethostoma* Jordan and Starks.
Pacific coast of Mexico.
Aboma ethostoma Jordan and Starks, Fishes of Sinaloa, Proc. Calif.
Acad. Sci., 2d ser., V, 1895, 498, pl. 50, Mazatlan, Mexico.
3565. *Aboma lucretiae* (Eigenmann and Eigenmann).
Pearl Island, Gulf of Panama.
Gobius lucretiae Eigenmann and Eigenmann, Proc. Calif. Acad. Sci.,
2d ser., I, 1888, 57, Pearl Island, Gulf of Panama.
3566. *Aboma chiquita* (Jenkins and Evermann).
Gulf of California.
Gobius chiquita Jenkins and Evermann, Proc. U. S. Nat. Mus., XI,
1888, 146, Guaymas, Sonora, Mexico.
- Genus 1232. **MICROGOBIUS** Poey.
Microgobius Poey, Enumeratio, 126, 1876 (*M. signatus* Poey).
Zalypnus Jordan and Evermann, Check List, 1895, 459 (*Gobius emblematicus* Jordan and Gilbert). .
3567. *Microgobius emblematicus* (Jordan and Gilbert).
Lower California to Panama.
Gobius emblematicus Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881,
330, Bay of Panama.
Microgobius cyclolepis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 74,
Albatross station 3020, Lower California.
3568. *Microgobius signatus* Poey.
West Indies; common about Cuba.
Microgobius signatus Poey, Enumeratio, 1876, 127, pl. 5, fig. 3, Cuba.
3569. *Microgobius gulosus* (Girard).
Coast of Florida to Texas; common north to Indian River, Fla.
Gobius gulosus Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 169,
Indianola, Tex.

- 3570. *Microgobius meeki* Evermann and Marsh.**
Culebra and Vieques Islands, Porto Rico.
Microgobius meeki Evermann and Marsh, Rept. U. S. Fish Com., XXV,
 1899 (Dec. 19), 356, between Vieques and Culebra Islands east of
 Porto Rico.
- 3571. *Microgobius thalassinus* (Jordan and Gilbert).**
South Atlantic coast, Virginia to South Carolina.
Gobius thalassinus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882,
 612, Charleston Harbor, S. C.
Microgobius eulipis Eigenmann and Eigenmann, Proc. Calif. Acad. Sci.,
 2d ser., I, 1888, 69, Fortress Monroe, Va.
- 3572. *Microgobius miraflorensis* Gilbert and Starks.**
Bay of Panama.
Microgobius miraflorensis Gilbert and Starks, Mem. Calif. Acad. Sci.,
 IV, 1904, 176, Pl. XXIX, Bay of Panama.
- Genus 1233. *EUCYCLOGOBIUS* Gill.**
Eucyclogobius Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 279 (*Gobius newberryi* Girard).
- 3573. *Eucyclogobius newberryi* (Girard).**
Streams of California, in small brooks near the sea; locally common in San Luis Obispo Creek.
Gobius newberryi Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 136, Tomales Bay, Calif.
- Genus 1234. *LEPIDOGOBIUS* Gill.**
Lepidogobius Gill, Ann. Lyc. Nat. Hist. N. Y., VII, 1859, 14 (*Gobius lepidus* Girard).
Cyclogobius Steindachner, Sitzb. Kais. Akad. Wiss. Wien, XLII, 1860, 284 (*Gobius lepidus* Girard).
- 3574. *Lepidogobius lepidus* (Girard).**
Pacific coast of North America, from Vancouver Island to Lower California, in rather deep water off San Francisco Bay.
Gobius lepidus Girard, Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 127, 25a, figs. 5 and 6; substitute for *Gobius gracilis*, preoccupied.
Gobius gracilis Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 134, San Francisco.
- Genus 1235. *GILLICHTHYS* Cooper.**
Gillichthys Cooper, Proc. Calif. Acad. Nat. Sci., III, 1863, 109 (*G. mirabilis* Cooper).
Gillia Günther, Zool. Rec., 1864, 137 (*Gillichthys mirabilis* Cooper); substitute name preoccupied.
Aprolepis Hubbs, Occ. Papers, Univ. Mich. Mus. Zool., I, 1921, 99 (*A. barbaræ* Hubbs); young of *Gillichthys mirabilis*.
- 3575. *Gillichthys mirabilis* Cooper. *Long-jawed goby.***
Pacific coast of North America, from San Francisco to Guaymas.
Gillichthys mirabilis Cooper, Proc. Calif. Acad. Nat. Sci., III, 1863, 109, San Diego Bay.
Gobius townsendi Eigenmann and Eigenmann, Proc. U. S. Nat. Mus., XI, 1888, 463, San Diego, Calif.
Aprolepis barbaræ Hubbs, Occ. Papers Univ. Mich. Mus. Zool., I, 1921, 99, young.
- 3576. *Gillichthys detrusus* Gilbert and Scofield.**
Horseshoe Bend, near mouth of Colorado River, in Mexico.
Gillichthys detrusus Gilbert and Scofield, Proc. U. S. Nat. Mus., XX, 1896, 498, near mouth of Colorado River.
- Genus 1236. *QUIETULA* Jordan and Evermann.**
Quietula Jordan and Evermann, in Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 839 (*Gillichthys y-cauda* Jenkins and Evermann).
- 3577. *Quietula y-cauda* (Jenkins and Evermann).**
Pacific coast of North America from Guaymas to Vancouver Island.
Gillichthys y-cauda Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 147, Guaymas, Sonora.
Gillichthys guayamasæ Jenkins and Evermann, loc. cit., 148, Guaymas, Sonora.

Genus 1237. CLEVELANDIA Eigenmann and Eigenmann.

Clevelandia Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888, 73 (*C. longipinnis* Eigenmann and Eigenmann = *Clevelandia rosæ* Jordan and Evermann).

3578. *Clevelandia ios* (Jordan and Gilbert).

Puget Sound and neighboring waters.

Gobiosoma ios Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 437, Sanianch Arm, Vancouver Island.

Clevelandia longipinnis Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888, 73; not of Steindachner.

Clevelandia rosæ Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 229, San Diego, Calif.

Genus 1238. ILYPNUS Jordan and Evermann.

Ilypnus Jordan and Evermann, Check List, 1895, 460 (*Lepidogobius giberti* Eigenmann and Eigenmann).

3579. *Ilypnus giberti* (Eigenmann and Eigenmann).

San Diego Bay and southward; abundant at Magdalena Bay, Conception Bay, and St. George's Bay in the Gulf of California.

Lepidogobius giberti Eigenmann and Eigenmann, Proc. U. S. Nat. Mus., XI, 1888, 464, San Diego Bay, Calif.

Genus 1239. EVERMANNIA Jordan.

Evermannia Jordan, Proc. Calif. Acad. Sci., 2d ser., IV, 1895, 592 (*Gobiosoma zosterurum* Jordan and Gilbert).

3580. *Evermannia zosterura* (Jordan and Gilbert).

Pacific coast of Mexico at Mazatlan, Mexico.

Gobiosoma zosterurum Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 361, Mazatlan.

3581. *Evermannia longipinnis* (Steindachner).

Gulf of California.

Gobiosoma longipinne Steindachner, Ichth. Beitr., VIII, 1879, 27, Las Animas Island, Gulf of California.

3582. *Evermannia panamensis* Gilbert and Starks.

Panama Bay.

Evermannia panamensis Gilbert and Starks, Mem. Calif. Acad. Sci., VI, 1904, 179, Pl. XXX, Panama Bay.

Genus 1240. GARMANNIA Jordan and Evermann. *Half-naked gobies.*

Garmannia Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 496, pl. 49 (*Gobius paradoxus* Günther).

3583. *Garmannia paradoxus* (Günther).

Pacific coast of Mexico, Mazatlan to Panama.

Gobius paradoxus Günther, Cat., III, 1861, 549, Pacific coast of Central America.

3584. *Garmannia hemigymna* (Eigenmann and Eigenmann).

West Indies.

Gobius hemigymnus Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser., I, 1888, 66, dredged in the West Indies.

3585. *Garmannia rubra* Rosen.

Bahamas.

Garmannia rubra Rosen, Arsek. Univ. Lund, VII, 1912, 50, Bâhamas.

Genus 1241. ENYPNIAS Jordan and Evermann.

Enypnias Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2233 (*Gobius seminudus* Günther).

3586. *Enypnias seminudus* (Günther).

Pacific coast of Central America.

Gobius seminudus Günther, Cat., III, 1861, 554, Pacific coast of Central America.

Genus 1242. GOBIOSOMA Girard.

Gobiosoma Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 169 (*Gobius alepidotus* Bloch and Schneider = *Gobius boscii* Lacépède).

- 3587. *Gobiosoma boscii* (Lacépède).** *Naked goby; Clinging goby; Goby; Variegated goby; Mud-creeper; Oysterfish.*
 Atlantic coast of United States, Cape Cod to Florida.
Gobius boscii Lacépède, Hist. Nat. Poiss., II, 1798, 555, pl. 16, fig. 1.
 Charleston, S. C.
Gobius alepidotus Bloch and Schneider, Syst. Ichth., 1801, 547; after
 Lacépède.
Gobiosoma viridipallidus Mitchell, Trans. Lit. and Phil. Soc. N. Y., I, 1814,
 379, pl. 1, fig. 8, New York.
- 3588. *Gobiosoma molestum* Girard.**
 Gulf coast of the United States from Key West to Texas, and south
 to Bahia.
Gobiosoma molestum Girard, Proc. Acad. Nat. Sci. Phila., X, 1858,
 169, Indianola, Tex.
Gobiosoma alepidotum Jordan and Gilbert, Proc. U. S. Nat. Mus., V,
 1882, 297, Laguna Grande, Pensacola, Fla.
- 3589. *Gobiosoma histrio* Jordan.**
 Known only from the Gulf of California at Guaymas.
Gobiosoma histrio Jordan, Proc. U. S. Nat. Mus., VII, 1884, 260, Guay-
 mas, Mexico.
- 3590. *Gobiosoma crescentale* Gilbert.**
 Gulf of California.
Gobiosoma crescentalis Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 557,
 Gulf of California, at Albatross station 2825.
- 3591. *Gobiosoma multifasciatum* Steindachner.**
 West Indies; known from Cuba, St. Thomas, and the Lesser Antilles.
Gobiosoma multifasciatum Steindachner, Ichth. Beitr., V, 1876, 183,
 Lesser Antilles.
Gobius lineatus Poey, Memorias, II, 1861, 424, Cuba; not of Jenyns.
- 3592. *Gobiosoma horsti* Metzelaar.**
 Curaçao, Caribbean Sea.
Gobiosoma horsti Metzelaar, Marine Fishes Lesser Antilles, 1923,
 Curaçao.
- 3593. *Gobiosoma longum* Nichols.**
 West Indies, north to Florida.
Gobiosoma longum Nichols, Bull. Amer. Mus. Nat. Hist., XXXIII,
 1914, 143, fig. 1, Florida.
- Genus 1243. *BARBULIFER* Eigenmann and Eigenmann.**
Barbulifer Eigenmann and Eigenmann, Proc. Calif. Acad. Sci., 2d ser.
 I, 1888, 70 (*B. papillosus* Eigenmann and Eigenmann = *Gobiosoma cauthæcum* Jordan and Gilbert).
- 3594. *Barbulifer cauthæcus* (Jordan and Gilbert).**
 About Key West, Fla.
Gobiosoma cauthæcum Jordan and Gilbert, Proc. U. S. Nat. Mus., VII,
 1884, 29, Key West, Fla.
Barbulifer papillosus Eigenmann and Eigenmann, Proc. Calif. Acad.
 Sci., 2d ser., I, 1888, 70, Key West, Fla.
- Genus 1244. *EVERMANNICHTHYS* Metzelaar.**
Evermannichthys Metzelaar, Trop. Atl. Vissch., 1919, 139 (*E. spongicola*
 Metzelaar).
Radcliffea Hubbs, Occ. Papers, Mus. Zool. Univ. Mich., No. 99, 1921,
 2 (*Garmannia spongicola* Radcliffe).
- 3595. *Evermannichthys spongicola* (Radcliffe).**
 Off shore banks of North Carolina and Florida in cavities of sponges.
Garmannia spongicola Radcliffe, Proc. U. S. Nat. Mus., LII, 1917,
 423, off Cape Hatteras, N. C., and near Tarpon Springs, west coast
 of Florida.

3596. *Evermannichthys metzelaari* Hubbs.

Caribbean sea, living in sponges.

Evermannichthys spongicola Metzelaar, Trop. Atl. Vissch., 1919, 189, figs. 39 and 40, 139, Curaçao; name preoccupied by *Evermannichthys spongicola* (Radcliffe).

Evermannichthys metzelaari Hubbs, Occ. Papers Mus. Zool. Univ. Mich., No. 144, Dec. 21, 1923, pl. 1, Curaçao; substitute for *E. spongicola* Metzelaar, preoccupied.

Genus 1245. *LETHOPS* Hubbs.

Lethops Hubbs, Occ. Papers Univ. Mich. Mus. Zool., 1925 (*Lethops connectens* Hubbs).

3597. *Lethops connectens* Hubbs.

Coast of California in tide pools.

Lethops connectens Hubbs, Occ. Papers Mus. Zool. Univ. Mich., 1925, Carmel Bay; Point Firmian, near Los Angeles.

Genus 1246. *TYPHLOGOBIUS* Steindachner. *Blind gobies.*

Typhlogobius Steindachner, Ichth. Beitr., VIII, 1879, 24 (*T. californiensis* Steindachner).

Othonops Rosa Smith (Mrs. Eigenmann), Proc. U. S. Nat. Mus., II, 1881, 19 (*O. eos* Smith = *Typhlogobius californiensis* Steindachner).

3598. *Typhlogobius californiensis* Steindachner. *Blind goby of Point Loma; Pinkfish.*

Coast of Lower California, from San Diego southward to Cedros Island; especially common at Point Loma.

Typhlogobius californiensis Steindachner, Ichth. Beitr., VIII, 1879, 24, False Bay, San Diego, Calif.

Othonops eos Rosa Smith, Proc. U. S. Nat. Mus., II, 1881, 19, Point Loma, San Diego, Calif.

Genus 1247. *SICYDIUM* Cuvier and Valenciennes.

Sicydium Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 126 (168) (*Gobius plumieri* Bloch).

3599. *Sicydium plumieri* (Bloch). *Sirago.*

Fresh waters of the West Indies.

Gobius plumieri Bloch, Ichth., 1786, 125, pl. 178, fig. 3, Martinique; on a drawing by Plumier.

Sicydium siragus Poey, Memorias, II, 1861, 278, Santiago de Cuba.

3600. *Sicydium caguiae* Evermann and Marsh.

Porto Rico.

Sicydium caguiae Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 355, Rio de Caguas at Caguas, Porto Rico.

3601. *Sicydium vincente* Jordan and Evermann.

Kingston, St. Vincent Island; Haiti.

Sicydium vincente Jordan and Evermann, Check List, 1895, 45, St. Vincent Island.

3602. *Sicydium antillarum* Ogilvie-Grant.

Barbados.

Sicydium antillarum Ogilvie-Grant, Proc. Zool. Soc. Lond., 1884, 157, Barbados.

3603. *Sicydium punctatum* Perugia.

Martinique.

Sicydium punctatum Perugia, Annali del Museo Civico di Storia Naturale di Genova, ser. 2, XVI, 1896, 18, Martinique.

3604. *Sicydium buscki* Evermann and Clark.

Santo Domingo.

Sicydium buscki Evermann and Clark, Proc. U. S. Nat. Mus., XXX, 1906, 854, fig. 3, San Francisco Mountain, Santo Domingo.

Genus 1248. *COTYLOPUS* Guichenot.

Cotylopus Guichenot, in Maillard, Notes sur l'Isle de la Réunion, II, 1862, annex b., 9 (*C. acutipinnis*).

Sicya Jordan and Evermann, Check List, 1895, 456, (*Sicydium gymnogaster* Ogilvie-Grant).

3605. *Cotylopus salvini* (Ogilvie-Grant).
 Streams near Panama.
Sicydium salvini Ogilvie-Grant, Proc. Zool. Soc. Lond., 1884, 159, pl. 12, fig. 2, Panama.
3606. *Cotylopus cocoensis* Heller and Snodgrass.
 Chatham Bay, Cocos Island, northwest of Panama.
Cotylopus cocoensis Heller and Snodgrass, Proc. Wash. Acad. Sci., V, 1903, 211, Pl. XI, Cocos Island.
3607. *Cotylopus punctatus* Regan.
 Tequesixtlán, Mexico.
Cotylopus punctatus Regan, Ann. Mag. Nat. Hist., ser. 7, XVI, 1905, 362, Tequesixtlán, Mexico.

Family 257. GOBIOOIDIDÆ

Genus 1249. GOBIOIDES Lacépède.
Gobiooides Lacépède, Hist. Nat. Poiss., II, 1800, 580 (*G. broussonnetii* Lacépède).

3608. *Gobiooides broussonnetii* Lacépède.
 West Indies to Brazil; once taken near New Orleans.
Gobiooides broussonnetii Lacépède, Hist. Nat. Poiss., II, 1800, 580, probably from Surinam; "given by Holland to France."
Gobius brasiliensis Bloch and Schneider, Syst. Ichth., 1801, 69, Brazil.
Gobius oblongus Bloch and Schneider, loc. cit., 548; based on Lacépède.
Gobiooides barreto Poey, Memorias, II, 1861, 282, Cuba.
Amblyopsis mexicanus O'Shaughnessy, Ann. Mag. Nat. Hist., ser. 4, XV, 1875, 147, Mexico.

Genus 1250. TYNTLASTES Günther.
Tyntlastes Günther, Proc. Zool. Soc. Lond., 1862, 193 (*Amblyopus sagitta* Günther).

3609. *Tyntlastes sagitta* (Günther).
 Coast of Lower California.
Amblyopus sagitta Günther, Proc. Zool. Soc. Lond., 1862, 193, "California"; probably from Lower California.

3610. *Tyntlastes brevis* (Günther).
 Panama.
Amblyopus brevis Günther, Proc. Zool. Soc. Lond., 1864, 151, Panama.

Genus 1251. CAYENNIA Sauvage.
Cayennia Sauvage, Bull. Soc. Philom., ser. 7, IV, 1880, 57 (*C. guichenoti* Sauvage).

3611. *Cayennia guichenoti* Sauvage.
 Cayenne, French Guiana.
Cayennia guichenoti Sauvage, Bull. Soc. Philom., ser. 7, IV, 1880, 57, Cayenne.

Order WW. DISCOCEPHALI

Family 258. ECHENEIDÆ. Remoras

Genus 1252. ECHENEIS Linnæus.
Echeneis Linnæus, Syst. Nat., Ed. X, 1758, 261 (*E. naucrates* Linnæus, as first restricted by Gill, 1862).
Leptecheneis Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 60 (*Echeneis naucrates* Linnæus).

3612. *Echeneis naucrates* Linnæus. *Sharksucker; Pega; Pegador; Suckingfish; White-tailed remora; Indian remora.*
 Warm seas, universally distributed; common north to Cape Cod and San Francisco.
Echeneis neucrates (misprint for *naucrates*) Linnæus, Syst. Nat., Ed. X, 1758, 261, Pelago Indico.
Echeneis albacauda Mitchell, Amer. Month. Mag., II, 1818, 244, New York.
Echeneis lunata Bancroft, Proc. Comm. Zool. Soc., I, 1830, 174, Kingston, Jamaica.

- Echeneis vittata* Lowe, Proc. Zool. Soc. Lond., 1839, 89, Madeira.
Echeneis guacan Poey, Memorias, II, 1861, 248, Cuba.
Echeneis verticalis Poey, loc. cit., 253, Cuba.
Echeneis metallica Poey, loc. cit., 252, Cuba.
Echeneis fusca Gronow, Cat. Fish., ed. Gray, 1854, 92; after Linnæus.
Echeneis neucratoides Zuieuw, Nova Acta Acad. Sci. Imp. Petropol., IV, 1789, 279, no locality.
Echeneis lineata Holbrook, Ichth. South Carolina, 1855, 101, Charleston, S. C.; not of Menzies.
Echeneis holbrooki Günther, Cat., II, 1860, 382, Jamaica.

Genus 1253. REMILEGIA Gill.

- Remilegia* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 239 (*Echeneis scutata* Günther).

3613. Remilegia australis (Bennett).

- Tropical seas; recorded by Doctor Lütken from 10° N., 39° W.
Echeneis australis Bennett, Nar. Whaling Voyage, II, 1840, 273, pls. 24–26.
Echeneis scutata Günther, Ann. Mag. Nat. Hist., 1860, 401, pl. 10, fig. B, Ceylon.

Genus 1254. REMORA Forster.

- Remora* Forster, Cat. Anim. N. Amer., 1770, 20 (*Echeneis remora* Linnæus).

3614. Remora remora (Linnæus). *Remora; Sucker; Pega.*

- Warm seas, north to New York and San Francisco.
Echeneis remora Linnæus, Syst. Nat., Ed. X, 1758, 260, Pelago Indico.
Echeneis squalipeta Daldorf, Skrft. Naturrh. Selsk., II, 1793, 157, Atlantic Ocean between the tropics.
Echeneis jacobæa Lowe, Proc. Zool. Soc. Lond., 1839, 89, Madeira.
Echeneis remoroides Bleeker, Natur. Tijd. Ned. Ind. (Batæ, IX, 1855, 70, Batæ).
Echeneis parva Gronow, Cat. Fish., ed. Gray, 1854, 92, no locality; after Linnæus.
Echeneis postica Poey, Memorias, II, 1861, 255, Habana.

Genus 1255. REMORINA Jordan and Evermann.

- Remorina* Jordan and Evermann, Check List, 1895, 490 (*Echeneis albescens* Temminck and Schlegel).

3615. Remorina albescens (Temminck and Schlegel).

- Tropical Pacific, northward to San Francisco and Japan; common at Galapagos Islands.
Echeneis albescens Temminck and Schlegel, Fauna Japonica, Poiss., 1845, 272, pl. 120, fig. 4, Nagasaki, Japan.

Genus 1256. REMOROPSIS Gill.

- Remoropsis* Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 60 (*Echeneis brachyptera* Lowe).

3616. Remoropsis brachyptera (Lowe). *Swordfish sucker.*

- Warm seas; occasionally north to Cape Cod.
Echeneis brachyptera Lowe, Proc. Zool. Soc. Lond., 1839, 89, Madeira.
Echeneis sexdecim-lamellata Eydoux and Gervais, Voy. Favorite, V, 1839, 77, pl. 31, Indian Ocean.
Echeneis quatuordecimlaminatus Storer, Rept. Fishes Mass., 1839, 155, Holmes Hole.
Echeneis pallida Temminck and Schlegel, Fauna Japonica, Poiss., 1845, 271, pl. 120, fig. 2, 3, Japan.
Echeneis nieuhofii Bleeker, Sumatra, Naturk. Tijd. Neder. Indie, IV, 1853, 279, Priaman.

Genus 1257. PHTHEIRICHTHYS Gill.

- Phtheirichthys* Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 239 (*Echeneis lineata* Menzies).

3617. Phtheirichthys tropicus (Euphrasen).

- Tropical seas, ranging north to South Carolina and Pensacola.
Echeneis tropica Euphrasen, Nya. Handl., XII, 1791, 315, Atlantic between the tropics.
Echeneis apicalis Poey, Memorias, II, 1861, 254, Cuba.
? *Echeneis sphyraenarum* Poey, loc. cit., 255, Cuba, on barracudas.

Genus 1258. RHOMBOCHIRUS Gill.

Rhombochirus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 88 (*Echeneis osteochir* Cuvier).

3618. *Rhombochirus osteochir* (Cuvier). *Spearfish sucker.*

West Indies, north to Cape Cod.

Echeneis osteochir Cuvier, Règne Animal, ed. 2, II, 1829, 348, no locality given.

Echeneis terapturorum Poey, Memorias, II, 1861, 256, Cuba.

Order XX. JUGULARES**Family 259. DRACONETTIDÆ****Genus 1259. DRACONETTA Jordan and Fowler.**

Draconetta Jordan and Fowler, Proc. U. S. Nat. Mus., XXV, 1903, 939 (*D. xenica* Jordan and Fowler).

3619. *Draconetta acanthopoma* Regan.

Depths of north Atlantic.

Draconetta acanthopoma Regan, Ann. Mag. Nat. Hist., May, 1904; north Atlantic, in deep water.

Family 260. CALLIONYMIDÆ. Dragonets**Genus 1260. CALLIONYMUS Linnæus.**

Callionymus Linnæus, Syst. Nat., Ed. X, 1758, 249, (*C. lyra* Linnæus).

3620. *Callionymus bairdi* Jordan.

Snapper Banks, Gulf of Mexico, between Pensacola and Tampa, Fla. *Callionymus bairdi* Jordan, Proc., U. S. Nat. Mus., X, 1887, 501, Snapper Banks off Pensacola, Fla.

3621. *Callionymus agassizii* Goode and Bean.

West Indies off Barbados and Santa Cruz

Callionymus agassizii Goode and Bean, Three Cruises of the *Blake*, II, 1888, 29, fig., Gulf of Mexico.

Callionymus himantophorus Goode and Bean, Oceanic Ichth., 1895, 296, figs. 268 A and B, off Barbados in 209 fathoms.

3622. *Callionymus calliurus* Eigenmann and Eigenmann.

Key West, Fla.

Callionymus calliurus Eigenmann and Eigenmann, Proc., Calif. Acad. Sci., 2d ser., I, 1888, 76, South Beach, Key West, Fla.

3623. *Callionymus pauciradiatus* Gill.

Matanzas, Cuba.

Callionymus pauciradiatus Gill, Ann. Lyc. Nat. Hist. N. Y., VIII, 1867, 143, Matanzas, Cuba.

3624. *Callionymus bermudarum* Barbour.

Castle Island, Bermudas.

Callionymus bermudarum Barbour, Bull. Mus. Comp. Zool., XLVI, 1905, No. 7, 129, Castle Island, Bermudas.

3625. *Callionymus bækei* Metzelaar.

Caribbean Sea.

Callionymus bækei Metzelaar, Trop. Atl. Vissch., 1916, 149, fig. 47, Curaçao.

3626. *Callionymus sancti-eustatii* Metzelaar.

Callionymus sancti-eustatii Metzelaar, Trop. Atl. Vissch., 1916, 150, fig. 47, St. Eustatius.

3627. *Callionymus atrilabiatus* Garman.

Off Bay of Panama in deep water.

Callionymus atrilabiatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 122, off Bay of Panama.

Family 261. PTEROPSARIDÆ

Genus 1261. HYPSICOMETES Goode.

Hypsicometes Goode, Proc. U. S. Nat. Mus., III, 1880, 347 (*H. gobiooides* Goode).3628. *Hypsicometes gobiooides* Goode.

Gulf Stream.

Hypsicometes gobiooides Goode, Proc. U. S. Nat. Mus., III, 1880, 348, Gulf Stream at Albatross station 871, $40^{\circ} 2' 54''$ N., $70^{\circ} 23' 40''$ W., in 115 fathoms.

Family 262. AMMODYTIIDÆ. Sand lances or sand launces

Genus 1262. AMMODYTES Linnaeus. *Sand launces; Sand lances.**Ammodytes* Linnaeus, Syst. Nat., Ed. X, 1758, 247 (*A. tobianus* Linnaeus). *Argyrotenia* Gill, Fishes East Coast U. S., Suppl. to Proc. Acad. Nat. Sci. Phila., XIII, 1861, 40 (*Ammodytes vittatus* DeKay; probably a deformed example).3629. *Ammodytes americanus* DeKay. *Sand lance; Sand launce; Sand eel; Lant.*

Newfoundland to Cape May and Cape Hatteras; abundant on sandy shores.

Ammodytes americanus DeKay, New York Fauna: Fishes, 1842, 317, Stratford, Conn.*Ammodytes vittatus* DeKay, loc. cit., 318, pl. 60, fig. 197, New York.3630. *Ammodytes dubius* Reinhardt.

Greenland to Cape Cod.

Ammodytes dubius Reinhardt, Dansk. Vidensk. Selsk. Afhandl., VII, 1838, 131, Greenland.3631. *Ammodytes personatus* Girard. *Sand lance; Sand launce.*

Japan to Alaska, south to Monterey, Calif.

Ammodytes personatus Girard, Proc. Acad. Nat. Sci. Phila., VIII, 1856, 137, Cape Flattery, Wash.3632. *Ammodytes alascanus* Cope. *Alaska sand lance*

North Pacific coast of North America, Sitka to Aleutian Islands.

Ammodytes alascanus Cope, Proc. Am. Phil. Soc., XIII, 1873, 30, Sitka, Alaska.

Family 263. BLEEKERIIDÆ

Genus 1263. RHYNCHIAS Gill.

Rhynchias Gill, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2841 (*Ammodytes septipinnis* Pallas); species and genus doubtful, not seen by recent authors.3633. *Rhynchias septipinnis* (Pallas).

Kamchatka.

Ammodytes septipinnis Pallas, Rosso-Asiat., III, 1811, 227, Tab. XLVIII, fig. 3, Kamchatka.

Family 264. BATHYMASTERIDÆ. Ronquils

Genus 1264. BATHYMASTER Cope.

Bathymaster Cope, Proc. Amer. Phil. Soc., XIII, 1873, 31 (*B. signatus* Cope).3634. *Bathymaster signatus* Cope.

Shores of Alaska.

Bathymaster signatus Cope, Proc. Amer. Phil. Soc. XIII, 1873, 31, near Sitka, Alaska.3635. *Bathymaster cæruleofasciatus* Gilbert and Burke.

Southern Bering Sea.

Bathymaster cæruleofasciatus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 84, fig. 29, Agattu Island, Alaska.

Genus 1265. RONQUILUS Jordan and Starks.
Ronquilus Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 838 (*Bathymaster jordani* Gilbert).

- 3636. *Ronquilus jordani*** (Gilbert). *Ronquil*.
Alaska to Puget Sound.
Rathymaster jordani Gilbert, Proc. U. S. Nat. Mus., XI, 1888, 554, Elliott Bay at Seattle; Alaska.

Genus 1266. RATHBUNELLA Jordan and Evermann.
Rathbunella Jordan and Evermann, Check List, 1895, 463 (*Bathymaster hypoplectus* Gilbert).

- 3637. *Rathbunella hypoplecta*** (Gilbert).
Coast of California about the Santa Barbara Islands.
Bathymaster hypoplectus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 97, coast of California, south of Point Conception, at *Albatross* station 2944.

- 3638. *Rathbunella allenii*** Gilbert.
Monterey Bay, Calif.
Rathbunella allenii Gilbert, Proc. Calif. Acad. Sci., 3d ser., III, 1904, 267, Monterey Bay, Calif.

Family 265. ZAPRORIDÆ

Genus 1267. ZAPRORA Jordan.
Zaprora Jordan, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 202 (*Z. silenus* Jordan).

- 3639. *Zaprora silenus*** Jordan.
East coast of Vancouver Island; only four specimens known.
Zaprora silenus Jordan, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 203, pl. 20, Nanaimo, Vancouver Island.

Family 266. CHIASMODONTIDÆ. Black swallows.

Genus 1268. CHIASMODON Johnson.
Chiasmodon Johnson, Proc. Zool. Soc. Lond., 1863, 408 (*C. niger* Johnson).
Chiasmodus Günther, altered spelling.

- 3640. *Chiasmodon niger*** Johnson. *Black swallow*.
Deep waters of the Atlantic; taken at Madeira and off Massachusetts coast.
Chiasmodon niger Johnson, Proc. Zool. Soc. Lond., 1863, 408, Madeira.

- 3641. *Chiasmodon subniger*** Garman.
Pacific coast of Mexico, off Lower California, in deep water.
Chiasmodon subniger Garman, Mem., Mus. Comp. Zool., XXIV, 1899, 73, off Gulf of California at *Albatross* station 3429.

Genus 1269. PSEUDOSCOPELUS Lütken.
Pseudoscopelus Lütken, Spolia Atlantica, Scopelini, 64, in Vid. Selsk. Skr. (6), VII, 1892, 285 (*P. scriptus* Lütken).

- 3642. *Pseudoscopelus scriptus*** Lütken.
Old Bahama Straits, in deep water.
Pseudoscopelus scriptus Lütken, Spolia Atlantica, Scopelini, 64, in Vid. Selsk. Skr. (6), VII, 1892, 285, Old Bahama Straits.

Family 267. OPISTHOGNATHIDÆ. Jawfishes

Genus 1270. OPISTHOGNATHUS Cuvier.
Opistognathus Cuvier, Règne Anim., Ed. I, 1817, 252 (*O. sonnerati* Cuvier).

- 3643. *Opistognathus macrognathum*** Poey.
Florida Keys to Cuba.
Opistognathus macrognathus Poey, Memorias, II, 1861, 284, Cuba.
Opistognathus megastoma Günther, Cat., I, 1860, 255, Gulf of Mexico.
Opistognathus scaphiurus Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 417, Garden Key, Fla.

3644. *Opistognathus lonchurum* Jordan and Gilbert.
 Gulf of Mexico, in deep water; two specimens known, taken from the stomach of a red snapper at Pensacola, Fla.
Opistognathus lonchurus Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 290, Snapper Banks off Pensacola, Fla.
3645. *Opistognathus punctatum* Peters.
 West coast of Mexico; Mazatlan; Magdalena Bay.
Opistognathus punctatus Peters, Berl. Monatsber., 1869, 708, Mazatlan, Mexico.
3646. *Opistognathus ommatum* Jenkins and Evermann.
 Bay of Guaymas, Gulf of California.
Opistognathus ommata Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 153, Guaymas, Sonora.
- Genus 1271. **GNATHYPOPS**^{ss} Gill.
Gnathypops Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 241 (*Opistognathus maxillosus* Poey).
3647. *Gnathypops maxillosa* (Poey).
 Cuba, north to Garden Key, Fla.
Opistognathus maxillosus Poey, Memorias, II, 1861, 286, Cuba.
3648. *Gnathypops macrops* (Poey).
 Coast of Cuba.
Opistognathus macrops Poey, Memorias, II, 1861, 287, coast of Cuba.
3649. *Gnathypops cuvieri* (Cuvier and Valenciennes).
 Bahia.
Opistognathus cuvieri Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 504, Bahia (may be identical with *macrops*).
3650. *Gnathypops rhomalea* (Jordan and Gilbert).
 Gulf of California.
Opistognathus rhomaleus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 276, Santa Maria Cove, Lower California.
3651. *Gnathypops scops* Jenkins and Evermann.
 Gulf of California at Guaymas.
Gnathypops scops Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 152, Guaymas, Sonora, Mexico.
3652. *Gnathypops snyderi* Jordan and Evermann.
 Gulf of California.
Gnathypops snyderi Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2285, San Luis Gonzales Bay, Gulf of California.
3653. *Gnathypops mystacina* Jordan.
 Gulf of Mexico; known from the Pensacola "Snapper Banks," from the stomach of a red snapper (*Neomexicanus blackfordi*).
Gnathypops mystacinus Jordan, Proc. U. S. Nat. Mus., VII, 1884, 37, Snapper Banks off Pensacola, Fla.
3654. *Gnathypops aurifrons* Jordan and Thompson.
 Garden Key, Fla.
Gnathypops aurifrons Jordan and Thompson, Bull. U. S. Bur. Fish., XXIV, 1904, 252, fig. 4, Garden Key, Fla.
- Genus 1272. **LONCHOPISTHUS** Gill.
Lonchopisthus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 241 (*Opistognathus micrognathus* Poey).
3655. *Lonchopisthus micrognathus* (Poey).
 Cuba.
Opistognathus micrognathus Poey, Memorias, II, 1861, 287, Cuba.

* It is possible that some or all of the short-jawed forms are the females of species of *Opistognathus*.

Family 268. URANOSCOPIDÆ. Stargazers

Genus 1273. ASTROSCOPUS Brevoort. *Stargazers.*

Astroscopus Brevoort, in Gill, Proc. Acad. Nat. Sci. Phila., XII, 1860, 20 (*Uranoscopus anoplos* Cuvier and Valenciennes = *Uranoscopus y-græcum* Cuvier and Valenciennes).

3656. *Astroscopus y-græcum* (Cuvier and Valenciennes). *Stargazer.*

South Atlantic coast, from Cape Hatteras to the Caribbean Sea.
Uranoscopus y-græcum Cuvier and Valenciennes, Hist. Nat. Poiss., III, 1829, 229 (308), locality unknown.
Uranoscopus anoplos Cuvier and Valenciennes, loc. cit., VIII, 1831, 363 (493), Atlantic coast of North America.

3657. *Astroscopus guttatus* Abbot. *Spotted stargazer.*

Atlantic coast of the United States, from Long Island to Virginia; not known south of Cape Hatteras.
Astroscopus guttatus Abbott, Proc. Acad. Nat. Sci. Phila., XII, 1860, 365, pl. 7, Cape May, N. J.

3658. *Astroscopus zephyreus* Gilbert and Starks.

Magdalena Bay, Lower California.
Astroscopus zephyreus Jordan and Starks, Proc. U. S. Nat. Mus., XIX, 1896, 453, Magdalena Bay, Lower California.

Genus 1274. KATHETOSTOMA Günther.

Kathetostoma Günther, Cat., II, 1860, 231 (*Uranoscopus loris* Bloch and Schneider).

3659. *Kathetostoma averruncus* Jordan and Bollman.

Pacific Ocean off coast of Colombia, and Bay of Panama, in deep water.
Kathetostoma averruncus Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 163, off coast of Colombia, 8° 57' N., 79° 31' 30" W., at Albatross station 2800 in 7 fathoms.

3660. *Kathetostoma alboguttata* (Bean).

Gulf of Mexico.
Cathetostoma alboguttata Bean, Proc. U. S. Nat. Mus., XVIII, 1892, 121, Gulf of Mexico, at Albatross station 2403, 28° 42' 30" N., 85° 29' W.

Genus 1275. EXCESTIDES Jordan and Thompson.

Execestides Jordan and Thompson, Bull. U. S. Bur. Fish., XXIV, 1904, 253, (*Execestides egregius* Jordan and Thompson).

3661. *Execestides egregius* Jordan and Thompson.

Garden Key, Fla.
Execestides egregius Jordan and Thompson, Bull. U. S. Bur. Fish., XXIV, 1904, 253, fig. 5, Garden Key, Fla.

Family 269. DACTYLOSCOPIDÆ. Sand stargazers

Genus 1276. DACTYLOSCOPUS Gill.

Dactyloscopus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 132 (*D. tridigitatus* Gill).
Congrammus Fowler, loc. cit., LVIII, 1906, 105 (*C. moorei* Fowler).

3662. *Dactyloscopus tridigitatus* Gill. *Sand conger.*

West Indies, north to Key West.
Dactyloscopus tridigitatus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 132, Barbados.
Congrammus moorei Fowler, loc. cit., LVIII, 1906, 105, Hailer's Rock, Florida Keys.

3663. *Dactyloscopus pectoralis* Gill.

Cape San Lucas, Lower California.
Dactyloscopus pectoralis Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 267, Cape San Lucas, Lower California.

3664. *Dactyloscopus cinctus* Osburn and Nichols.

Gulf of California.
Dactyloscopus cinctus Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 176, fig. 14, Cape San Lucas.

3665. *Dactyloscopus poeyi* Gill.
 Cuba.
Dactyloscopus poeyi Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 266,
 Cuba.
3666. *Dactyloscopus lunaticus* Gilbert.
 Gulf of California.
Dactyloscopus lunaticus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 99,
 Gulf of California.
- Genus 1277. *GILLELLUS* Gilbert.
Gillellus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 98 (*G. semicinctus*
 Gilbert).
3667. *Gillellus semicinctus* Gilbert.
 Gulf of California; also off the Florida coast.
Gillellus semicinctus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 98,
 near north end of Gulf of California, at Albatross stations 2827, 24°
 11' 45" N., 109° 55' W., and 2829, 22° 52' N., 109° 55' W., in 10
 and 31 fathoms.
3668. *Gillellus arenicola* Gilbert.
 Cape San Lucas, Lower California.
Gillellus arenicola Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 99, Cape
 San Lucas.
3669. *Gillellus ornatus* Gilbert.
 Gulf of California.
Gillellus ornatus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 558, Gulf
 of California.
- Genus 1278. *ESLOSCOPUS* Jordan and Evermann.
Esloscopus Jordan and Evermann, Check List, 1895, 465 (*Dactyloscopus
 zelotes* Jordan and Gilbert).
3670. *Esloscopus zelotes* (Jordan and Gilbert).
 Panama.
Dactyloscopus zelotes Jordan and Gilbert, in Jordan and Evermann,
 Check List, 1895, 464, Panama.
- Genus 1279. *DACTYLAGNUS* Gill.
Dactylagnus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 505 (*D.
 mundus* Gill).
3671. *Dactylagnus mundus* Gill.
 Carmen Island, Gulf of California.
Dactylagnus mundus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 505,
 Carmen Island, Gulf of California.
- Genus 1280. *MYXODAGNUS* Gill.
Myzodagnus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 269 (*M.
 opercularis* Gill).
3672. *Myzodagnus opercularis* Gill.
 Cape San Lucas, Lower California.
Myzodagnus opercularis Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861,
 270, Cape San Lucas, Lower California.

Family 270. CLINIDÆ

Genus 1281. *MNIERPES* Jordan and Evermann.
Mnierpes Jordan and Evermann, Check List, 1895, 468 (*Clinus macrocephalus* Günther).

3673. *Mnierpes macrocephalus* (Günther).
 Pacific coast of Central America; Panama.
Clinus macrocephalus Günther, Cat., III, 1861, 266, Pacific coast of
 Central America.

Genus 1282. *GOBIOCLINUS* Gill.
Gobioclinus Gill, Proc. Acad. Nat. Sci. Phila., XII, 1860, 103 (*Clinus gobia* Cuvier and Valenciennes).

- 3674.** *Gobioclinus gobio* (Cuvier and Valenciennes).
 Lesser Antilles.
Clinus gobio Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 292 (395), Lesser Antilles.
- Genus 1283. **HISTIOCLINUS** Metzelaar.
Histioclinus Metzelaar, Trop. Atl. Vissch., 1919, 157 (*H. veliger* Metzelaar).
- 3675.** *Histioclinus veliger* Metzelaar.
 Caribbean Sea.
Histioclinus veliger Metzelaar, Trop. Atl. Vissch., 1919, 157, Bonaire.
- Genus 1284. **STARKSIA** Jordan and Evermann.
Starksia Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 231, (*Labrosomus cremnobates* Gilbert).
- 3676.** *Starksia cremnobates* (Gilbert).
 Gulf of California.
Labrosomus cremnobates Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 100, Gulf of California.
- Genus 1285. **ALLOCLINUS** Hubbs.
Allocinus Hubbs, Papers, Mich. Acad. Sci. Arts and Letters, VII, 1926 (1927), 351 (*Starksia holderi* Lauderbach).
- 3677.** *Allocinus holderi* (Lauderbach).
 Santa Catalina, Calif.
Starksia holderi Lauderbach, Proc. U. S. Nat. Mus., XXXII, 1907, 74, fig. 6, Santa Catalina Island.
- Genus 1286. **CRYPTOTREMA** Gilbert.
Cryptotrema Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 101 (*C. corallinum* Gilbert).
- 3678.** *Chryptotrema corallinum* Gilbert.
 Coast of California, south of Point Conception.
Cryptotrema corallinum Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 101, coast of California, south of Point Conception.
- Genus 1287. **EXERPES** Jordan and Evermann.
Exerpes Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 232 (*Auchenopterus asper* Jenkins and Evermann).
- 3679.** *Exerpes asper* (Jenkins and Evermann).
 Guaymas, Mexico.
Auchenopterus asper Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 154, Guaymas, Mexico.
- Genus 1288. **CORALLICOLA** Jordan and Evermann.
Corallicola Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2369 (*Cremnobates marmoratus* Steindachner).
- 3680.** *Corallicola marmoratum* (Steindachner). *Rockfish*.
 Florida Keys to Cuba; Key West.
Cremnobates marmoratus Steindachner, Ichth. Beitr., V, 1876, 174, pl. 12, fig. 6, near Cuba.
- 3681.** *Corallicola nigripinne* (Steindachner).
 Barbados.
Clinus nigripinnis Steindachner, Ichth. Notizen, VI, 1867, 45, Barbados.
- 3682.** *Corallicola altivelis* (Lockington).
 La Paz, Lower California.
Cremnobates altivelis Lockington, Proc. Acad. Nat. Sci. Phila., XXXIII, 1881, 116, La Paz, Lower California.
- Genus 1289. **AUCHENOPTERUS** Günther.
Auchenopterus Günther, Cat., III, 1861, 275 (*A. monophthalmus* Günther; not *Auchenipterus* Cuvier and Valenciennes).
Cremnobates Günther, Proc. Zool. Soc. Lond., 1861, 374 (same type; substitute name).
Tekla Nichols, Copeia, No. 110, 1922, 69 (*Cremnobates fasciatus* Steindachner).

3683. *Auchenopterus monophthalmus* Günther.
 Gulf of California to Panama.
Auchenopterus monophthalmus Günther, Cat., III, 1861, 275, Panama.
3684. *Auchenopterus integripinnis* (Rosa Smith).
 Coast of California and southward to Todos Santos, Lower California.
Cremnobates integripinnis Rosa Smith, Proc. U. S. Nat. Mus., III, 1880,
 147, La Jolla, near San Diego.
3685. *Auchenopterus mexicanus* Gilbert.
 La Paz to Mazatlan, Mexico.
Auchenopterus mexicanus Gilbert, Proc. Calif. Acad. Sci., 3d ser., III,
 1904, 269, Pl. XXIX, La Paz, Lower California.
3686. *Auchenopterus affinis* (Steindachner).
 West Indies; Key West; St. Thomas.
Cremnobates affinis Steindachner, Ichth. Beitr., V, 1876, 178, St. Thomas.
3687. *Auchenopterus fasciatus* (Steindachner).
 Florida Straits; north to Key West; Porto Rico.
Cremnobates fasciatus Steindachner, Ichth. Beitr., V, 1876, 176, Florida
 Straits.
3688. *Auchenopterus nox* (Jordan and Gilbert).
 Key West, Fla.
Cremnobates nox Jordan and Gilbert, Proc. U. S. Nat. Mus., VII, 1884,
 30, Key West.
3689. *Auchenopterus albicaudus* Evermann and Marsh.
 Porto Rico.
Auchenopterus albicaudus Evermann and Marsh, Rept. U. S. Fish
 Com., XXV, 1899 (Dec. 19), 360, Arroyo, Porto Rico.
3690. *Auchenopterus fajardo* Evermann and Marsh.
 Porto Rico.
Auchenopterus fajardo Evermann and Marsh, Rept. U. S. Fish Com.,
 XXV, 1899 (Dec. 19), 361, Fajardo, Porto Rico.
3691. *Auchenopterus rubescens* Evermann and Marsh.
 Porto Rico.
Auchenopterus rubescens Evermann and Marsh, Rept. U. S. Fish Com.,
 XXV, 1899 (Dec. 19), 360, Puerto Real, Porto Rico.
3692. *Auchenopterus cingulatus* Evermann and Marsh.
 Porto Rico.
Auchenopterus cingulatus Evermann and Marsh, Rept. U. S. Fish Com.,
 XXV, 1899 (Dec. 19), 361, Ponce, Porto Rico.
3693. *Auchenopterus grandicomus* Rosen.
 Bahamas.
Auchenopterus grandicomus Rosen, Fauna Bahamas, 1911, 69, Andros
 Island.
- Genus 1290. **AUCHENISTIUS** Evermann and Marsh.
Auchenistius Evermann and Marsh, Rept. U. S. Fish Com., XXV,
 1899 (Dec. 19), 359 (*A. stahli* Evermann and Marsh).
3694. *Auchenistius stahli* Evermann and Marsh.
 Porto Rico.
Auchenistius stahli Evermann and Marsh, Rept. U. S. Fish Com.,
 XXV, 1899 (Dec. 19), 359, Ponce, Porto Rico.
- Genus 1291. **PARACLINUS** Mocquard.
Acanthoclinus Mocquard, Bull. Soc. Philom. Paris, X, 1885, 18 (*A.
 chaperi* Mocquard); name preoccupied.
Paraclinus Mocquard, loc. cit., I, 1888, 41 (*Acanthoclinus chaperi*
 Mocquard).
3695. *Paraclinus chaperi* (Mocquard).
 Bay of Guanta, near Barcelona, in Venezuela.
Acanthoclinus chaperi Mocquard, Bull. Soc. Philom. Paris, X, 1885, 18,
 Bay of Guanta, Venezuela.

Genus 1292. ENNEAPTERYGIUS Rüppell.

Enneapterygius Rüppell, Neue Wirbelthiere, Fische, 1835, 2 (*E. pusillus* Rüppell).
Enneanectes Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 501 (*Tripterygium carmineum* Jordan and Gilbert).

3696. *Enneapterygius carminalis* (Jordan and Gilbert).

Mazatlan, Mexico, and Caribbean Sea.
Tripterygium carmineum Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 362, Mazatlan, Mexico.

Genus 1293. GILLIAS Evermann and Marsh.

Gillias Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 357 (*G. jordani* Evermann and Marsh).

3697. *Gillias jordani* Evermann and Marsh.

Porto Rico.
Gillias jordani Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899, (Dec. 19), 357, Cardona Lighthouse Reef, Ponce, Porto Rico.

Genus 1294. GIBBONSIA Cooper.

Gibbonsia Cooper, Proc. Calif. Acad. Nat. Sci., III, 1863, 109 (*Myxodes elegans* Cooper).
Blakea Steindachner, Ichth. Beitr., V, 1876, 148 (*Myxodes elegans* Cooper).

3698. *Gibbonsia elegans* (Cooper). *Spotted kelpfish; Señorita.*

Coast of southern California; Point Conception to Todos Santos.
Myxodes elegans Cooper, Proc. Calif. Acad. Nat. Sci., III, 1863, 109, San Diego; Santa Barbara.
Clinus ocellifer Mocquard, Bull. Soc. Philom. Paris, 1886, 44, California.
Clinus evides Jordan and Gilbert, Synopsis, 1883, 763, Monterey, Calif.

3699. *Gibbonsia elegans montereyensis* Hubbs.

Coast of southern California.
Gibbonsia elegans montereyensis Hubbs, Papers Mich. Acad. Sci. Arts and Letters, VII, 1926, 354, reef at Pacific Grove, Calif.

3700. *Gibbonsia metzi* Hubbs.

California coast, chiefly in tidepools.
Gibbonsia metzi Hubbs, Papers Mich. Acad. Sci. Arts and Letters, VII, 1926, 360 Pacific Grove, Calif.

Genus 1295. HETEROSTICHUS Girard.

Heterostichus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 143 (*H. rostratus* Girard).

3701. *Heterostichus rostratus* Girard. *Kelpfish.*

Coast of California from San Francisco to San Diego.
Heterostichus rostratus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 143, San Diego, Calif.

Genus 1296. NEOCLINUS Girard.

Neoclinus Girard, U. S. Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 114 (*N. blanchardi* Girard).

3702. *Neoclinus blanchardi* Girard.

Coast of California, Monterey to San Diego.
Neoclinus blanchardi Girard, U. S. Pac. R. R. Surv., X, pt. 4, Fishes, 1858, 114, off San Diego, Calif.

Genus 1297. PTEROGNATHUS Jordan and Evermann.

Pterognathus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 57 (*Neoclinus satiricus* Girard); name only suggested but not adopted.
Pterognathus Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2354 (*Neoclinus satiricus* Girard).

3703. *Pterognathus satiricus* (Girard).

Coast of California, from Monterey to Santa Barbara.
Neoclinus satiricus Girard, Proc. Acad. Nat. Sci. Phila., XI, 1859, 57, Monterey Bay.

Genus 1298. LABRISOMUS Swainson.

Labrisomus Swainson, Nat. Hist. Classn. Fishes, II, 1839, 277 (*Clinus pectinifer* Cuvier and Valenciennes = *Clinus nuchipinnis* Quoy and Gaimard).

Lepisoma DeKay, N. Y. Fauna: Fishes, 1842, 41, *L. cirrhosum* DeKay = *Clinus nuchipinnis* Quoy and Gaimard).

3704. *Labrisomus nuchipinnis* (Quoy and Gaimard).

West Indies; South Atlantic coast.

Clinus nuchipinnis Quoy and Gaimard, Voy. *Uranié*, Zool., 1824, 255, Bay of Rio de Janeiro.

Clinus pectinifer Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 276 (374), Bahia.

Clinus capillatus Cuvier and Valenciennes, loc. cit., 278 (377), Martinique.

Clinus canariensis Valenciennes, in Webb and Berthelot, Poiss. Iles Canaries, 1839, 60, Pl. XVII, fig. 3, Canary Islands.

Lepisoma cirrhosum DeKay, N. Y. Fauna: Fishes, 41, 1842, Florida.

3705. *Labrisomus heilneri* Nichols.

Bahamas.

Labrisomus heilneri Nichols, Amer. Mus. Novitates, XXVI, 1921, p. 2, fig. 2, Bahamas.

3706. *Labrisomus xanti* Gill.

Pacific coast of Mexico.

Labrisomus xanti Gill, Proc. Acad. Nat. Sci. Phila., XII, 1860, 107, Cerro Blanco, Cape San Lucas.

3707. *Labrisomus herminieri* (LeSueur).

St. Bartholomew, West Indies.

Blennius herminieri LeSueur, Jour. Acad. Nat. Sci. Phila., IV, 1824, 361, St. Bartholomew, West Indies.

3708. *Labrisomus bucciferus* Poey.

Cuba.

Labrosomus bucciferus Poey, Synopsis, 1868, 399, Cuba.

3709. *Labrisomus lentiginosus* Bean.

Paget Parish, Bermudas.

Labrisomus lentiginosus Bean, Proc. Biol. Soc. Wash., XIX, 1906, 31, tide pool at Does Rock, Paget Parish, Bermudas.

- 3710-3711. *Labrisomus microlepidotus* Poey.

Cuba.

Labrosomus microlepidotus Poey, Anal. Soc. Espaniola Hist. Nat., XIX, 1880, 246, Pl. VIII, fig. 2, Cuba.

Genus 1299-1300. MALACOCTENUS Gill.

Malacoctenus Gill, Proc. Acad. Nat. Sci. Phila., XII, 1860, 103 (*Clinus delalandi* Cuvier and Valenciennes).

3712. *Malacoctenus delalandi* (Cuvier and Valenciennes).

Coast of Brazil and the Pacific coast of Mexico.

Clinus delalandii Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 279 (378), Brazil.

Clinus zonifer Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 361, Mazatlan.

3713. *Malacoctenus ocellatus* (Steindachner).

Bahama Islands.

Clinus ocellatus Steindachner, Ichth. Beitr., V, 1876, 182, Bahama Islands.

3714. *Malacoctenus varius* (Poey).

Cuba.

Myxodes varius Poey, Enumeratio, 1876, 132, pl. 5, fig. 2, Habana.

3715. *Malacoctenus macropus* (Poey).
Cuba.
Myzodes macropus Poey, Synopsis, III, 1868, 399, Habana.
3716. *Malacoctenus lugubris* (Poey).
Cuba.
Myzodes lugubris Poey, Enumeratio, 1876, 131, Cuba.
3717. *Malacoctenus gilli* (Steindachner).
Barbados.
Clinus gilli Steindachner, Ichth. Notizen, VI, 1867, 46, Barbados.
3718. *Malacoctenus bimaculatus* (Steindachner).
Small rocky islands north coast of Cuba.
Clinus bimaculatus Steindachner, Ichth. Beitr., V, 1876, 180, islands off Cuba.
3719. *Malacoctenus puertoricensis* Evermann and Marsh.
Porto Rico.
Malacoctenus puertoricensis Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 358, Hucares, Porto Rico.
3720. *Malacoctenus versicolor* (Poey).
Cuba.
Myzodes versicolor Poey, Enumeratio, 1876, 131, pl. 5, fig. 1, Cuba.
3721. *Malacoctenus biguttatus* (Cope).
New Providence, Bahamas.
Labrisomus biguttatus Cope, Trans. Amer. Phil. Soc. Phila., XIV, 1871, 473, New Providence, Bahamas.
- Genus 1301. ERICTEIS Jordan.
Ericteis Jordan, Bull. U. S. Fish Com., XXII, 1902, 543 (*E. kalisheræ* Jordan).
3722. *Ericteis kalisheræ* Jordan.
Bush Key, Tortugas.
Ericteis kalisheræ Jordan, Bull. U. S. Fish Com., XXII, 1902, 543, pl. 2, Bush Key, Fla.
- Genus 1302. ACTEIS Jordan.
Acteis Jordan, Bull. U. S. Fish Com., XXII, 1902, 543 (*Malacoctenus moorei* Evermann and Marsh).
3723. *Acteis moorei* (Evermann and Marsh).
Porto Rico.
Malacoctenus moorei Evermann and Marsh, Fishes of Porto Rico, in Bull. U. S. Fish Com., XX, 1900, 309, fig. 97, Culebra Island, Porto Rico.
3724. *Acteis culebrae* (Evermann and Marsh).
Porto Rico.
Malacoctenus culebrae Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 357, Culebra Island.

Family 271. BLENNIIDÆ

Genus 1303. BLENNIUS Linnaeus. *Blennies*.
Blennius Linnaeus, Syst. Nat., ed. X, 1758, 256 (*B. galerita* Linnaeus).
Salaria Forskål, Descr. Anim., 1777, 22 (*Blennius galerita* Linnaeus).
Adonis Gronow, Cat. Fish., Ed. Gray, 1854, 93 (*A. pavoninus* Gronow = *Blennius ocellaris* Linnaeus).

3725. *Blennius fabbri* Nichols.
Florida.
Blennius fabbri Nichols, Bull. Amer. Mus. Nat. Hist., XXVIII, 1910, 161, Miami, Fla.; perhaps young of *Chasmodes bosquianus*.
3726. *Blennius stearnsi* Jordan and Gilbert.
Gulf of Mexico.
Blennius stearnsi Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 300, Pensacola snapper banks.

3727. **Blennius fucorum** Cuvier and Valenciennes. *Seaweed blenny.*
 Off the Azores; also off New York.
Blennius fucorum Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 194 (263), pl. 324, 240 miles south of the Azores.
Blennius oceanicus Cuvier and Valenciennes, loc. cit., 195 (265), open sea.
3728. **Blennius cristatus** Linnaeus. *Rockfish.*
 Tropical Atlantic.
Blennius cristatus Linnaeus, Syst. Nat., Ed. X, 1758, 256, Indies; after Gronow.
Blennius crinitus Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 175 (237), La Rochelle, France.
Blennius nuchiifilis Cuvier and Valenciennes, loc. cit., 186 (253), Isle of Ascension.
Blennius asterias Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 416, Garden Key, Fla.
3729. **Blennius favosus** Goode and Bean.
 Gulf of Mexico; Garden Key, Fla.
Blennius favosus Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 416, Garden Key, Fla.
3730. **Blennius pilicornis** Cuvier.
 Coast of Brazil and north to the West Indies; recorded from Rio Janeiro, Bahia, the Tortugas, and off the coast of Florida.
Blennius pilicornis Cuvier, Règne Anim., ed. 2, II, 1829, 237, coasts of Europe, Rio Janeiro.
3731. **Blennius marmoratus** Poey.
 Cuba.
Blennius marmoratus Poey, Enumeratio, 1870, 130, Cuba.
3732. **Blennius truncatus** Poey.
 Cuba.
Blennius truncatus Poey, Memorias, II, 1861, 424, Cuba.
3733. **Blennius vinctus** Poey.
 Cuba.
Blennius vinctus Poey, Repertorio, II, 1867, 243, Habana.
- Genus 1304. **LIPOPHRYS** Gill.
Pholis Fleming, Brit. Anim., 1828, 207 (*Blennius lavis* = *Blennius pholis* Linnaeus); name preoccupied.
Lipophrys Gill, Amer. Nat., XXX, June, 1896, 498 (*Blennius pholis* Linnaeus).
3734. **Lipophrys carolinus** (Cuvier and Valenciennes).
 South Carolina.
Pholis carolinus Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 204 (276), Carolina.
- Genus 1305. **SCARTELLA** Jordan.
Scartella Jordan, Proc. U. S. Nat. Mus., IX, 1886, 50 (*Blennius microstomus* Poey).
Blenniolus Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2386 (*Blennius brevipinnis* Günther).
3735. **Scartella microstoma** (Poey).
 Cuba.
Blennius microstomus Poey, Memorias, II, 1861, 288, Cuba.
3736. **Scartella brevipinnis** (Günther).
 Pacific coast of Mexico, from Mazatlan to Panama.
Blennius brevipinnis Günther, Cat., III, 1861, 226, Pacific coast of Central America.
- Genus 1306. **HYPELUROCHILUS** Gill.
Hypeurochilus Gill, Cat. Fishes E. Coast of N. Amer., Suppl. Proc. Acad. Nat. Sci. Phila., XIII, 1861, 44 (*Blennius multifilis* Girard = *Blennius geminatus* Wood).

- 3737.** *Hypseurochilus geminatus* (Wood). *Blenny.*
 South Atlantic and Gulf coasts of United States.
Blennius geminatus Wood, Jour. Acad. Nat. Sci. Phila., IV, 1824, 278,
 Charleston, S. C.
Blennius multifilis Girard, Proc. Acad. Nat. Sci. Phila., X, 1858, 169,
 St. Joseph's Island, Tex.
- Genus 1307. HYPSOBLENNIUS Gill.**
Hypsoblennius Gill, Cat. Fishes E. Coast N. Amer., Suppl. Proc. Acad.
 Nat. Sci. Phila., XIII, 1861, 44 (*Blennius hentz* LeSueur).
Isesthes Jordan and Gilbert, Synopsis, 1883, 757 (*Blennius gentilis*
 Girard).
- 3738.** *Hypsoblennius hentz* (LeSueur).
 Coasts of North and South Carolina, south of Indian River, Fla.;
 locally common.
Blennius hentz LeSueur, Jour. Acad. Nat. Sci. Phila., 1st ser., IV, 1825,
 363, Charleston, S. C.
Blennius punctatus Wood, loc. cit., 278, Charleston, S. C.; not of
 Fabricius, 1780.
- 3739.** *Hypsoblennius giberti* (Jordan).
 California, from Point Conception southward to Todos Santos; Santa
 Barbara Islands, among rocks.
Isesthes giberti Jordan, Proc. U. S. Nat. Mus., V, 1882, 349, Santa
 Barbara, Calif.
- 3740.** *Hypsoblennius gentilis* (Girard).
 Coast of California from Monterey to Cape San Lucas.
Blennius gentilis Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 149, .
 Monterey, Calif.
- 3741.** *Hypsoblennius piersoni* Gilbert and Starks.
 Bay of Panama.
Hypsoblennius piersoni Gilbert and Starks, Mem. Calif. Acad. Sci., IV,
 1904, 191, Pl. XXXII, Bay of Panama.
- 3742.** *Hypsoblennius striatus* (Steindachner).
 Panama.
Blennius striatus Steindachner, Ichth. Beitr., V, 1876, 15, Panama.
- 3743.** *Hypsoblennius ionthas* (Jordan and Gilbert).
 Pensacola, Fla.
Isesthes ionthas Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882,
 299, Pensacola.
Isesthes scrutator Jordan and Gilbert, loc. cit., 300, Pensacola; Galveston.
- Genus 1308. CHASMODES Cuvier and Valenciennes.**
Chasmodes Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 218
 (295) (*Blennius bosquianus* Lacépède).
Blenitracus Swainson, Classn. Fish., II, 1839, 78, 274 (*Pholis quadri-
 faciatus* Wood).
- 3744.** *Chasmodes bosquianus* (Lacépède). *Banded blenny; Bosc's shanny;*
Four-banded blenny.
 New York to Florida; common southward in shallow water.
Blennius bosquianus Lacépède, Hist. Nat. Poiss., II, 1800, 493, South
 Carolina; female.
- 3745.** *Chasmodes quadrifasciatus* (Wood).
 Probably south Atlantic coast of United States.
Pholis quadrifasciatus Wood, Jour. Acad. Nat. Sci. Phila., IV, 1824,
 282, locality unknown; probably South Carolina.
- 3746.** *Chasmodes saburrae* Jordan and Gilbert.
 Pensacola Bay, Fla.
Chasmodes saburrae Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882,
 298, Pensacola, Fla.
- 3747.** *Chasmodes novemlineatus* (Wood). *Nine-lined shanny.*
 South Atlantic coast of the United States, South Carolina to Florida.
Pholis novemlineatus Wood, Jour. Acad. Nat. Sci. Phila., IV, 1825, 280,
 Charleston Harbor, S. C.

3748. *Chasmodes jenkinsi* Jordan and Evermann.
Guaymas, Sonora, Mexico.
Chasmodes jenkinsi Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., VI, 1896, 232, pl. 39, Guaymas, Mexico.
- Genus 1309. **HOMESTHES** Gilbert.
Homesthes Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2394 (*H. caulopus* Gilbert).
3749. *Homesthes caulopus* Gilbert.
Panama Bay.
Homesthes caulopus Gilbert, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2394, Panama.
- Genus 1310. **SCARTICHTHYS** Jordan and Evermann.
Scartes Jordan and Evermann, Check List, 1895, 471 (*Salarias salarias rubropunctatus* Cuvier and Valenciennes); name preoccupied.
Scartichthys Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2395 (*Salarias rubro-punctatus* Cuvier and Valenciennes).
3750. *Scartichthys rubropunctatus* (Cuvier and Valenciennes).
Coast of Peru and Chile, north to Panama.
Salarias rubro-punctatus Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 257 (348), Juan Fernandez, off Chile.
- Genus 1311. **RUPISCARTES** Swainson.
Alticus Commerson, in Lacépède, Hist. Nat. Poiss., II, 1800, 458 (*A. saltatorius* Commerson); not binomial and not accepted by Lacépède.
Alticus Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 337; casual notice; the name not accepted as that of a genus (*Blennius saliens*⁸⁹=*Salarias alticus* Cuvier and Valenciennes).
Rupiscartes Swainson, Class. Fish., II, 1839, 275 (*Salarias atlanticus* Cuvier and Valenciennes).
3751. *Rupiscartes atlanticus* (Cuvier and Valenciennes).
Tropical America on both coasts, north to West Indies and to Todos Santos, Lower California; abundant.
Salarias atlanticus Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 238 (321), Atlantic Ocean.
- Genus 1312. **ENTOMACRODUS** Gill.
Entomacrodus Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 168 (*E. nigricans* Gill).
3752. *Entomacrodus chiostücktus* (Jordan and Gilbert).
Pacific coast of Mexico.
Salarias chiostücktus Jordan and Gilbert, Synopsis, 1883, 363, Mazatlan, Mexico.
3753. *Entomacrodus margaritaceus* (Poey).
Cuba.
Salarias margaritaceus Poey, Memorias, II, 1861, 289, Cuba.
3754. *Entomacrodus decoratus* Poey.
Cuba.
Entomacrodus decoratus Poey, Synopsis, 1868, 398, Cuba.
3755. *Entomacrodus nigricans* Gill.
West Indies.
Entomacrodus nigricans Gill, Proc. Acad. Nat. Sci. Phila., XI, 1859, 168, Barbados.
3756. *Entomacrodus cruentatus* Garman.
Off Cocos Island in deep water.
Entomacrodus cruentatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 123, Pl. L; off Cocos Island, Panama.
- Genus 1313. **SALARICHTHYS** Guichenot.
Salarichthys Guichenot, Mém., Soc. Sci. Nat. Cherbourg, XIII, 1868, 96 (*Salarias vomerinus* Cuvier and Valenciennes= *Salarias textile* Cuvier and Valenciennes).

⁸⁹ "Un petit *salarias* qui nous paraît être celui, la même pour laquel Commerson a établi son genre *Alticus*."

- 3757. *Salarichthys textilis* Quoy and Gaimard.**
 West Indies, from Bermudas to Brazil.
Salarias textilis Quoy and Gaimard, in Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 227 (307), Ascension Island.
Salarias vomerinus Cuvier and Valenciennes, loc. cit., 258 (349), Bahia.
- Genus 1314. *BRANNERELLA* Gilbert.**
Brannerella Gilbert, Proc. Wash. Acad. Sci., II, 1900, 180 (*B. brasiliensis* Gilbert).
- 3758. *Brannerella brasiliensis* Gilbert.**
 Brazil, north to Curaçao, in the Caribbean Sea.
Brannereila brasiliensis Gilbert, Proc. Wash. Acad. Sci., II, 1900, 180, Maceio, Brazil.
- 3759. *Brannerella sluiteri* Metzelaar.**
 Caribbean Sea.
Brannerella sluiteri Metzelaar, Trop. Atl. Vissch., 1919, 155, Bonaire.
- Genus 1315. *OPHIOLENNIUS* Gill.**
Blennophis Valenciennes, in Webb and Barthelot, îles Canaries, Poiss., 1844, 60 (*B. webbii* Valenciennes), name preoccupied.
Ophioblennius Gill, Proc. Acad. Nat. Sci. Phila., XII, 1860, 103 (*B. webbii* Valenciennes); substitute for *Blennophis* Valenciennes, not *Blennophis* Swainson, a genus of Clinidae.
- 3760. *Ophioblennius webbii* (Valenciennes).**
 West Indies and Canary Islands.
Blennophis webbii Valenciennes, in Webb and Barthelot, îles Canaries, Poiss., 1839, 60, pl. 20, fig. 1, Fort Aventura, Canary Islands.
- 3761. *Ophioblennius steindachneri* Jordan and Evermann.**
 Pacific coast of Mexico.
Ophioblennius steindachneri Jordan and Evermann, Fishes North and Mid. Amer., 1898, 2401, Tres Marias Islands; after Steindachner.
- Genus 1316. *CORALLIOZETUS* Evermann and Marsh.**
Coralliozetus Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 362 (*C. cardonæ* Evermann and Marsh).
- 3762. *Coralliozetus cardonæ* Evermann and Marsh.**
 Porto Rico.
Coralliozetus cardonæ Evermann and Marsh, Rept. U. S. Fish Com., XXV, 1899 (Dec. 19), 362, reef at Cardona Light House, Playa de Ponce, Porto Rico.
- Family 272. EMBLEMARIIDÆ**
- Genus 1317. *EMBLEMARIA* Jordan and Gilbert.**
Emblemaria Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 627 (*E. nivipes* Jordan and Gilbert).
- 3763. *Emblemaria atlantica* Jordan and Evermann.**
 Snapper Banks off Pensacola; Gulf of Mexico.
Emblemaria atlantica Jordan and Evermann, Fishes North and Mid. Amer., 1908, 2402, snapper banks off Pensacola, Fla.
- 3764. *Emblemaria nivipes* Jordan and Gilbert.**
 Pearl Island near Panama.
Emblemaria nivipes Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 627, Pearl Island, near Panama.
- 3765. *Emblemaria oculocirris* Jordan.**
 Gulf of California.
Emblemaria oculocirris Jordan, in Gilbert, Proc. U. S. Nat. Mus., XIX, 1896, 456, La Paz.
- 3766. *Emblemaria pandionis* Evermann and Marsh.**
 Porto Rico.
Emblemaria pandionis Evermann and Marsh, Bull. U. S. Fish Com., XX, 1900, 318, between Vieques and Culebra Islands.

3767. *Emblemaria marki* Mowbray.

Bermudas.

Emblemaria marki Mowbray, in Bean, Proc. Biol. Soc. Wash., XXV, 1912, 126, Hamilton Harbor, Bermudas.

Genus 1318. ACANTHEMBLEMARIA Metzelaar.

Acanthemblemaria Metzelaar, Trop. Atl. Vissch., 1919, 159 (*A. spinosa* Metzelaar).

3768. *Acanthemblemaria spinosa* Metzelaar.

Caribbean Sea.

Acanthemblemaria spinosa Metzelaar, Trop. Atl. Vissch., 1919, 159, fig. 53, Curaçao.

Family 273. ATOPOCLINIDÆ

Genus 1319. ATOPOCLINIS Vaillant.

Atopoclinus Vaillant, Bull. Soc. Philom. Paris, ser. 8, VI, 1894, 73 (*A. ringens* Vaillant).

3769. *Atopoclinus ringens* Vaillant.

Gulf of California.

Atopoclinus ringens Vaillant, Bull. Soc. Philom., ser. 8, VI, 1894, 74, Gulf of California.

Family 274. CHÆNOPSISIDÆ

Genus 1320. CHÆNOPSIS Gill.

Chænopsis Gill, Ann. Lyc. Nat. Hist. N. Y., VIII, 1863, 141, Pl. 3, fig. 3 (*C. ocellatus* Poey).

Lucioblemnius Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 103 (*L. alepidotus* Gilbert).

3770. *Chænopsis ocellata* Poey.

Matanzas, Cuba.

Chænopsis ocellatus Poey, in Gill, Ann. Lyc. Nat. Hist. N. Y., VIII, 1863, 143, Matanzas, Cuba.

3771. *Chænopsis alepidota* (Gilbert).

Gulf of California; Avalon Bay, Calif.

Lucioblemnius alepidotus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 103, Lower California.

3772. *Chænopsis lucius* (Osburn and Nichols).

Gulf of California.

Lucioblemnius lucius Osburn and Nichols, Bull. Amer. Mus. Nat. Hist., XXXV, 1916, 179, fig. 15, San Jose Island, Gulf of California.

Genus 1321. PHOLIDICHTHYS Bleeker.

Pholidichthys Bleeker, Boerœ, in Natur. Tijd. Neder. Indie, XI, 1856, 406 (*P. leucotænia* Bleeker).

3773. *Pholidichthys anguilliformis* Lockington.

Gulf of California.

Pholidichthys anguilliformis Lockington, Proc. Acad. Nat. Sci. Phila., XXXIII, 1881, 118, San Jose Island, Lower California.

Genus 1322. PSEDNOBLENNIUS Jenkins and Evermann.

Psednoblennius Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 156 (*P. hypacanthus* Jenkins and Evermann).

3774. *Psednoblennius hypacanthus* Jenkins and Evermann.

Gulf of California at Guaymas, Sonora, Mexico.

Psednoblennius hypacanthus Jenkins and Evermann, Proc. U. S. Nat. Mus., XI, 1888, 156, Guaymas, Mexico.

Genus 1323. STATHMONOTUS Bean.

Stathmonotus Bean, Proc. U. S. Nat. Mus., VIII, 1885, 191 (*S. hemphillii* Bean).

3775. *Stathmonotus hemphillii* Bean.

Key West, Fla.

Stathmonotus hemphillii Bean, Proc. U. S. Nat. Mus., VIII, 1885, 191, pl. 13, Key West, Fla.

3776. *Stathmonotus tekla* Nichols.

Key West, Fla.

Stathmonotus tekla Nichols, Bull. Amer. Mus. Nat. Hist., XXVIII, 1910, 161, fig. 1c. XXX, Pl. XI, fig. 2, Key West, Fla.

Family 275. CEBIDICHTHYIDÆ

Genus 1324. CEBIDICHTHYS Ayres.

Cebidichthys Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 59 (*Apodichthys violaceus* Girard).

3777. *Cebidichthys violaceus* (Girard).

Coast of California from Crescent City to Carpenteria.

Apodichthys violaceus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 150, Monterey, Calif.

Cebidichthys cristagalli Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 58, San Francisco.

Family 276. CHIROLOPHIDÆ

Genus 1325. BRYOSTEMMA Jordan and Starks.

Bryostemma Jordan and Starks, Proc. Calif. Acad. Sci. 2d ser., V, 1895, 841 (*Blennius polyactocephalus* Pallas).

3778. *Bryostemma polyactocephalum* (Pallas).

Northern Japan and Kamchatka to Puget Sound.

Blennius polyactocephalus Pallas, Zool. Rosso-Asiat., III, 1811, 179, Kamchatka.

Chirolophus japonicus Herzenstein, Mélanges Biologiques Soc. Sci. Petersb., XIII, 1890, 123, Yezo, Japan.

3779. *Bryostemma nugator* Jordan and Williams.

Puget Sound.

Bryostemma nugator Jordan and Williams, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 843, pl. 101, Seattle, Wash.

3780. *Bryostemma decoratum* Jordan and Snyder.

Point Orchard, Wash.

Bryostemma decoratum Jordan and Snyder, Proc. U. S. Nat. Mus., XXV, 1903, 615, Point Orchard, near Seattle, Wash.

3781. *Bryostemma tarsodes* Jordan and Snyder.

Unalaska.

Bryostemma tarsodes Jordan and Snyder, Proc. U. S. Nat. Mus., XXV, 1903, 614, Unalaska.

Genus 1326. BRYOLOPHUS Jordan and Snyder.

Bryolophus Jordan and Snyder, Proc. U. S. Nat. Mus., XXV, 1903, 617 (*B. lysimus* Jordan and Snyder).

3782. *Bryolophus lysimus* Jordan and Snyder.

Unalaska.

Bryolophus lysimus Jordan and Snyder, Proc. U. S. Nat. Mus., XXV, 1903, 617, Unalaska.

Genus 1327. GYMNOCLINUS Gilbert and Burke.

Gymnoclinus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 86 (*G. cristulatus* Gilbert and Burke).

3783. *Gymnoclinus cristulatus* Gilbert and Burke.

Bering Island.

Gymnoclinus cristulatus Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 86, fig. 30, Nikolski, Bering Island.

Family 277. PHOLIDÆ

Genus 1328. APODICHTHYS Girard.

Apodichthys Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 150 (*A. flavidus* Girard).

3784. *Apodichthys flavidus* Girard.

Point Conception to Vancouver Island.

Apodichthys flavidus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 150, Presidio (San Francisco), Calif.

Apodichthys virescens Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 55 (54), San Francisco.

Apodichthys sanguineus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 279, California.

Apodichthys inornatus Gill, loc. cit., 279, Puget Sound.

3785. *Apodichthys univittatus* Lockington.

Lower California.

Apodichthys univittatus Lockington, Proc. Acad. Nat. Sci. Phila., XXXIII, 1881, 119, Lower California, probably from the Gulf.

Genus 1329. *XERERPES* Jordan and Gilbert.

Xererpes Jordan and Gilbert, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 846 (*Apodichthys fucorum* Jordan and Gilbert).

3786. *Xererpes fucorum* (Jordan and Gilbert).

Coast of California.

Apodichthys fucorum Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 139, Point Pinos, near Monterey, Calif.

Genus 1330. *ULVICOLA* Gilbert and Starks.

Ulvicola Gilbert and Starks, Proc. U. S. Nat. Mus., XIX, 1896, 455 (*U. sanctæ-rosæ* Gilbert and Starks).

3787. *Ulvicola sanctæ-rosæ* Gilbert and Starks.

Santa Rosa Island, coast of California.

Ulvicola sanctæ-rosæ Gilbert and Starks, Proc. U. S. Nat. Mus., XIX, 1896, 453, Santa Rosa Island, Calif.

Genus 1331. *PHOLIS* Gronow.

Pholis Gronow, Acta Helvet., IV, 1760, 259 (*Blennius gunnellus* Linnaeus).

Pholis Scopoli, Introd. Hist. Nat., 1777, 456 (*Blennius gunnellus* Linnaeus).

Murænoides Lacépède, Hist. Nat. Poiss., II, 1800, 324 (*Murænoides sujef* Lacépède).

Centronotus Schneider, Systema, ed. Bloch, 1801, 165 (*Centronotus fasciatus* Schneider).

Dactyleptus Rafinesque, Anal. Nat., 1815, 82; substitute for *Murænoides*. *Gunnelus* Fleming, Brit. Anim., 1828, 207 (*G. vulgaris* Fleming = *Blennius gunnellus* Linnaeus).

Centronotus Cuvier, Règne Anim., ed. 2, II, 1829, 239 (*Blennius gunnellus* Linnaeus).

Ophisomus Swainson, Classn. Anim., II, 1839, 279 (*Blennius gunnellus* Linnaeus).

Urocentrus Kner, Sitzber. K. Akad. Wiss. Wien, LVIII, 1868, 51 (*U. pictus* Kner).

3788. *Pholis gunnellus* (Linnaeus). *Gunnel*; *Butterfish*; *Rockfish*.

North Atlantic, south to Cape Cod.

Blennius gunnellus Linnaeus, Syst. Nat., Ed. X, 1758, 257, Atlantic Ocean.

Ophidion imberbe Linnaeus, loc. cit., 259, Europe.

Blennius europæus Olafsen, Reise Island, I, 1772, 81, Iceland.

Blennius murænoides Sujeff, Act. Petrop., II, 1779, 195, no locality, probably Baltic.

Ophidium mucronatum Mitchell, Trans. Lit. and Phil. Soc. N. Y., I, 1815, 361, Pl. I, fig. 1, New York.

Gunnellus vulgaris Fleming, British Anim., 1842, 207, England.

- 3789. *Pholis fasciatus* (Bloch and Schneider).**
 Greenland to Alaska and Japan.
Centronotus fasciatus Bloch and Schneider, Syst. Ichth., 1801, 165, pl. 37, fig. 1, Tranquebar.
Gunnellus grænlandicus Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 326 (442), pl. 340, Greenland; after Bloch and Schneider.
?Gunnellus murænoïdes Valenciennes, in Cuvier, Règne Anim., Poiss., ed. 2, 1829, pl. 78, fig. 2, 916; after Bloch and Schneider.
Blennius tænia Pallas, Zoogr. Rosso-Asiat., III, 1811, 178, Kuril Islands.
Murænoïdes maxillaris Bean, Proc. U. S. Nat. Mus., IV, 1881, 147, St. Paul Island, Alaska.
- 3790. *Pholis ornatus* (Girard).**
 San Francisco to Alaska.
Gunnellus ornatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 149, Presidio (San Francisco), Calif.
Centronotus latus Cope, Proc. Amer. Phil. Soc., XIII, 1873, 27, Sitka or Unalaska.
- 3791. *Pholis gilli* Evermann and Goldsborough.**
 Bering Sea.
Pholis gilli Evermann and Goldsborough, Bull. U. S. Bur. Fish., XXVI, 1906, 337, fig. 110, Bering Sea.
- Genus 1332. *RHODYMENICHTHYS* Jordan and Evermann.**
Rhodymenichthys Jordan and Evermann, Check List, 1895, 474 (*Gunnellus ruberrimus* Cuvier and Valenciennes = *Blennius dolichogaster* Pallas).
- 3792. *Rhodymenichthys dolichogaster* (Pallas). *Butterfish*.**
 Okhotsk Sea to San Francisco.
Blennius dolichogaster Pallas, Zoogr. Rosso-Asiat., III, 1811, 175, Kamchatka.
Gunnellus ruberrimus Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836, 324 (440), Kuril Islands; after Pallas.
- Genus 1333. *ASTERNOPTERYX* Rüppell.**
Asternopteryx Rüppell, in Günther, Cat., III, 1861, 288 (*A. gunelliformis* Rüppell).
- 3793. *Asternopteryx gunelliformis* Rüppell.**
 Greenland.
Asternopteryx gunelliformis Rüppell, in Günther, Cat., III, 1861, 288, Greenland; type in Senckenberg Museum, Frankfurt; a second specimen from Greenland examined by us.
- Genus 1334. *ANOPLARCHUS* Gill.**
Anoplarchus Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 261 (*A. purpureascens* Gill).
- 3794. *Anoplarchus cristagalli* (Günther).**
 Alaska to San Francisco.
Centronotus cristagalli Günther, Cat., III, July, 1861, 289, Vancouver Island.
Anoplarchus purpureascens Gill, Proc. Acad. Nat. Sci. Phila., XIII, Sept., 1861, 262, Washington Territory.
- 3795. *Anoplarchus cristigalli archolepis* Hubbs.**
 Coast of central California.
Anoplarchus purpureascens archolepis Hubbs, Papers Mich. Acad. Sci., Arts and Letters, VII, 1926, 375, Reef at Carmel, Calif.
- 3796. *Anoplarchus insignis* Gilbert and Burke.**
 Attu Island.
Anoplarchus insignis Gilbert and Burke, Bull. U. S. Bur. Fish., XXX, 1910, 88, fig. 32, Attu Island, Bering Sea.
- Genus 1335. *PLECTOBRANCHUS* Gilbert.**
Plectobranchus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 102 (*P. evides* Gilbert).

3797. **Plectobranchus evides** Gilbert.

Coasts of Oregon and Washington.

Plectobranchus evides Gilbert, Proc. U. S. Nat. Mus., XIII, 1898, 102,
coasts of Oregon and Washington.

Family 278. XIPHISTERIDÆ

Genus 1336. **XIPHISTER** Jordan and Gilbert.

Xiphidion Girard, Pac. R. R. Surv., X, Pt. 4, Fishes, 1858, 119 (*X. mucosum* Girard); name preoccupied in *Orthoptera*.

Xiphister Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 135
(*Xiphidion mucosum* Girard).

3798. **Xiphister mucosus** (Girard).

Monterey to Alaska.

Xiphidion mucosum Girard, Pac. R. R. Surv., X, pt. 4, Fishes, 1858,
119, South Farallon Island, off California.

Xiphidium crureum Cope, Proc. Amer. Phil. Soc., XIII, 1873, 27,
Sitka, Alaska.

Genus 1337. **EPIGEICHTHYS** Hubbs.

Epineichthys Hubbs, Papers Mich. Acad. Sci. Arts and Letters, VII,
1926 (*Xiphister rupestris* Jordan and Gilbert).

3799. **Epineichthys atropurpureus** (Kittlitz).

Pacific coast of America from Alaska to Monterey.

Ophidium atropurpureum Kittlitz, Denkwürd, einer Reise Russ.-Amer.,
I, 1858, 225, Alaska.

Xiphister rupestris Jordan and Gilbert, Proc. U. S. Nat. Mus., III,
1880, 137, Monterey, Calif.

Genus 1338. **PHYTICHTHYS** Hubbs.

Xiphistes Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895,
846 (*Xiphistes chirus* Jordan and Gilbert); name preoccupied in
insects.

Phytichthys Hubbs, in Jordan, Class. Fishes, 1923, 234 (*Xiphister chirus*
Jordan and Gilbert).

3800. **Phytichthys chirus** (Jordan and Gilbert).

Monterey to Alaska.

Xiphister chirus Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880,
135, Point Pinos, near Monterey, Calif.

Xiphistes ulvae Jordan and Starks, Fishes Puget Sound, Proc. Calif.
Acad. Sci., 2d ser., V, 1895, 847, pl. 102, Waadda Island, Neah Bay.

3801. **Phytichthys versicolor** (Gilbert and Burke).

Bering Sea.

Xiphistes versicolor Gilbert and Burke, Bull. U. S. Bur. Fish., XXX,
1910, 88, fig. 33; Attu; Agattu, Bering Sea.

Family 279. STICHAEIDÆ

Genus 1339. **STICHAEUS** Reinhardt.

Stichæus Reinhardt, K. Dansk. Vidensk. Selsk. Skrift. Natur. og
Math. Afhandl., VI, 1837, CX (*Blennius punctatus* Fabricius).

Notogrammus Bean, Proc. U. S. Nat. Mus., IV, 1881, 147 (*N. rothrocki*
Bean); young example.

3802. **Stichæus punctatus** (Fabricius). *Spotted snake blenny.*

Arctic seas from Greenland to northern Siberia, south to Bristol Bay
and Newfoundland.

Blennius punctatus Fabricius, Fauna Greenlandica, 1780, 153, Greenland.

Notogrammus rothrocki Bean, Proc. U. S. Nat. Mus., IV, 1881, 146,
Plover Bay and Cape Lisburne, Siberia.

Genus 1340. **PLAGIOPRAGMUS** Bean.

Plagiogrammus Bean, Proc. U. S. Nat. Mus., XVI, 1893, 699 (*P. hopkinsi*
Bean).

3803. **Plagiogrammus hopkinsi** Bean.

Monterey Bay, Calif.

Plagiogrammus hopkinsi Bean, Proc. U. S. Nat. Mus., XVI, 1893, 700,
with fig. on p. 699; Monterey, Calif.

Genus 1341. ULVARIA Jordan and Evermann.
Ulvaria Jordan and Evermann, Check List, 1895, 475 (*Pholis subbifurcatus* Storer).

3804. Ulvaria subbifurcata (Storer). *Radiated shanny.*
 North Atlantic, south of Cape Cod.
Pholis subbifurcatus Storer, Rept. Fish. Mass., 1839, 63, Nahant, Mass.

Genus 1342. EUMESOGRAMMUS Gill.
Eumesogrammus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 210
 (*Clinus præcisus* Krøyer).

3805. Eumesogrammus præcisus (Krøyer).
 Greenland.
?Clinus præcisus Krøyer, Naturh. Tidsskr., I, 1837, 25, Greenland.
Clinus unimaculatus Reinhardt, Dansk. Vidensk. Selsk., VII, 1838,
 114, 121, Greenland.

Family 280. LUMPENIDÆ

Genus 1343. LEPTOCLINUS Gill.
Ctenodon Nilsson, Scand. Fauna, IV, 1853, 190 (*Clinus maculatus* Fries); name preoccupied.
Leptoclinus Gill, Cat. Fish. E. coast N. Amer., Suppl., Proc. Acad. Nat. Sci. Phila., XIII, 1861, 45 (*Clinus aculeatus* Reinhardt = *Clinus maculatus* Fries).

3806. Leptoclinus maculatus (Fries). *Langbarn.*
 North Atlantic coasts of Europe; this or some closely similar species is recorded from Bering Sea, Robben Island, Aleutian Islands, and Hood Canal, Puget Sound.
Clinus maculatus Fries, Kgl. Vet. Ak. Handl., 1837, 51, Bohüslaan, Sweden.
Clinus aculeatus Reinhardt, K. Dansk. Vid. Selsk. Natur. Afh., VII, 1838, 114, 122, Spitzbergen.

Genus 1344. POROCLINUS Bean.
Poroclinus Bean, Proc. U. S. Nat. Mus., XIII, 1890, 40 (*P. rothrocki* Bean).

3807. Poroclinus rothrocki Bean.
 Between Nagai and Big Koniushi Islands, Alaska.
Poroclinus rothrocki Bean, Proc. U. S. Nat. Mus., XIII, 1890, 40, Albatross station 2952, lat. 55° 15' N., long. 159° 37' W., in 58 fathoms.

Genus 1345. ANISARCHUS Gill.
Anisarchus Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 209 (*Clinus medius* Reinhardt).

3808. Anisarchus medius (Reinhardt).
 Greenland to Norway and Spitzbergen; this or a similar species on coasts of southern and central Alaska.
Clinus medius Reinhardt, K. Dansk. Vidensk. Selsk. Skrift. Afh., VII, 1838, 114, 121, Greenland.

Genus 1346. LUMPENUS Reinhardt.
Lumpenus Reinhardt, Dansk. Vidensk. Selsk. Natur., VI, 1837, CX (*Blennius lumpenus* Fabricius = *Gunnellus fabricii* Cuvier and Valenciennes).
Leptogunnellus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 26 (*L. gracilis* Ayres = *Blennius anguillaris* Pallas).
Centroblennius Gill, Proc. Acad. Nat. Sci. Phila., XVI, 1864, 209 (*Lumpenus nubilis* Richardson = *Gunnellus fabricii* Cuvier and Valenciennes).
Leptoblennus Gill, loc. cit., 1864, 209 (*Blennius serpentinus* Storer = *Blennius lampetraformis* Walbaum).

3809. Lumpenus gracilis (Ayres).
 San Francisco to Alaska.
Blennius anguillaris Pallas, Zoogr. Rosso-Asiat., III, 1811, 176, about Kamchatka; no definite locality; not of Peck.
Leptogunnellus gracilis Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 26, San Francisco.

- 3810.** *Lumpenus mackayi* Gilbert.
Nushagak River, Alaska.
Lumpenus mackayi Gilbert, Rept. U. S. Fish Com., XIX, 1893, 450,
 pl. 32, Nushagak River, Alaska.
- 3811.** *Lumpenus fabricii* (Cuvier and Valenciennes).
 Arctic seas; Greenland, Gulf of St. Lawrence, Wellington Sound,
 Bristol Bay, and other localities in Bering Sea.
Gunellus fabricii Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836,
 317 (431), Greenland.
Lumpenus nubilus Richardson, Last Arctic Voyage, Fishes, 1855, 259,
 pl. 28, Wellington Sound.
- 3812.** *Lumpenus lampetraeformis* (Walbaum). *Snake blenny; Tangbrosme;*
Eel blenny; Snakefish.
 North Atlantic and Arctic, on both shores; south to Sweden and Green-
 land; rarely to Cape Cod.
Blennius lampetraeformis Walbaum, Artedi Pisc., III, 1792, 184, Iceland.
Centronotus islandicus Schneider, ed. Bloch, Syst. Ichth., 1801, 167,
 Iceland.
? *Clinus nebulosus* Fries, Vet. Akad. Hahdl., 1837, 55, Bohuslän, Sweden.
? *Clinus mohri* Krøyer, Naturrh. Tidsskr., I, 1837, R, 32, Iceland.
Blennius gracilis Stuvitz, Nye Mag. for Naturvid., I, 1838, 406, west
 coast of Norway.
Blennius serpentinus Storer, Proc. Bost. Soc. Nat. Hist., III, 1848, 30,
 Massachusetts Bay.
- Genus 1347.** *LUMPENELLA* Hubbs.
Lumpenella Hubbs, Papers Mich. Acad. Sci. Arts and Letters, VII,
 1926, 378 (*Lumpenus longirostris* Evermann and Goldsborough).
- 3813.** *Lumpenella longirostris* (Evermann and Goldsborough).
 Southeastern Alaska.
Lumpenus longirostris Evermann and Goldsborough, Bull. U. S. Bur.
 Fish., XXVI, 1906, 340, fig. 115, Lynn Canal, Alaska.
- Family 281. PTILICHTHYIDÆ**
- Genus 1348.** *PTILICHTHYS* Bean.
Ptilichthys Bean, Proc. U. S. Nat. Mus., IV, 1881, 157 (*P. goodei* Bean).
- 3814.** *Ptilichthys goodei* Bean.
 Aleutian Islands.
Ptilichthys goodei Bean, Proc. U. S. Nat. Mus., IV, 1881, 157, Port
 Levasheff, Unalaska.
- Family 282. CRYPTACANTHODIDÆ**
- Genus 1349.** *CRYPTACANTHODES* Storer.
Cryptacanthodes Storer, Rept. Fish. Mass., 1839, 27 (*C. maculatus*
 Storer).
- 3815.** *Cryptacanthodes maculatus* Storer. *Wrymouth; Ghostfish.*
 North Atlantic, south to Cape Cod.
Cryptacanthodes maculatus Storer, Rept. Fish. Mass., 1839, 28, Massa-
 chusetts.
Fierasfer borealis DeKay, New York Fauna: Fishes, 1842, 316, New
 York; after Peck.
Cryptacanthodes inornatus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863,
 332, coast of Massachusetts.
- Genus 1350.** *LYCONNECTES* Gilbert.
Lyconectes Gilbert, Rept. U. S. Fish Com., XIX, 1893, 452 (*L. aleutensis*
 Gilbert).
- 3816.** *Lyconectes aleutensis* Gilbert.
 Aleutian Islands.
Lyconectes aleutensis Gilbert, Rept. U. S. Fish Com., XIX, 1893, 452,
 pl. 34, near Unalaska.
- Genus 1351.** *DELOLEPIST* Bean.
Delolepis Bean, Proc. U. S. Nat. Mus., IV, 1881, 465 (*D. virgatus* Bean).

3817. ***Delolepis giganteus*** Kittlitz.

Coasts of British Columbia and southern Alaska, south to Puget Sound.
Delolepis giganteus Kittlitz, Denk. Reise Russ. Amer., 226, 1858,
 Siberia.

Delolepis virgatus Bean, Proc. U. S. Nat. Mus., IV, 1881, 466, King-
 combe Inlet, British Columbia, and Port Wrangel, Alaska.

Family 283. ANARHICHADIDÆ

Genus 1352. **ANARHICHAS** Linnæus.

Anarhichas Linnæus, Syst. Nat., Ed. X, 1758, 247 (*A. lupus* Linnæus).

3818. ***Anarhichas lupus*** Linnæus. *Wolfish.*

North Atlantic, south to Cape Cod and France; rather common in
 America and Europe.

Anarhichas lupus Linnæus, Syst. Nat., Ed. X, 1758, 247, no definite
 locality.

Anarhichas strigosus Gmelin, Syst. Nat., Tome I, Pt. III, 1789, 1144,
 British seas.

Anarrhichas vomerinus Agassiz, in Storer, Hist. Fish. Mass., 1867,
 265, pl. 18, fig. 1, Cusk Rocks between Boston and Cape Ann.

3819. ***Anarhichas orientalis*** Pallas.

Alaska.

Anarhichas orientalis Pallas, Zoogr. Rosso-Asiat., III, 1811, 77,
 Kamchatka.

Anarrhichas lepturus Bean, Proc. U. S. Nat. Mus., II, 1879, 212, St.
 Michaels, Alaska.

Genus 1353. **LYCICHTHYS** Gill.

Lycichthys Gill, Ann. Rec. Indust. & Sci., 1876, CLXVII (*Anarrhichas latifrons* Steenstrup and Hallgrímsson).

3820. ***Lycichthys latifrons*** (Steenstrup and Hallgrímsson).

North Atlantic, on both coasts; chiefly north of the Arctic Circle.

Anarrhichas latifrons Steenstrup and Hallgrímsson, Förh. Skand. Naturf., 3 die Möte, 1842, 647; name only, no description; Viden-skabelige Meddeleker Natur Forening Kjobenhavn, 1877-1878, 109-114.

3821. ***Lycichthys minor*** (Olafsen).

North Atlantic, on both coasts, chiefly north of the Arctic Circle.

Anarrhichas minor Olafsen, Reise i. Island, sec. 682b, 1774, tab. 42,
 Iceland.

Anarrhichas karrak Bonnaterre, Encycl. Ichth., 1788, 38, Iceland,
 after Olafsen.

Anarrhichas maculatus Bloch and Schneider, Syst. Ichth., 1801, 496,
 Iceland; after Olafsen.

Anarrhichas leopardus Agassiz, in Spix, Pisc. Brasil., 1829, 92, tab. 51,
 "Atlantic Ocean."

Family 284. ANARRHICHTHYIDÆ

Genus 1354. **ANARRHICHTHYS** Ayres.

Anarrhichthys Ayres, Proc. Calif. Acad. Nat. Sci., I, 1855, 31 (*A. ocellatus* Ayres = *A. felis* Girard).

3822. ***Anarrhichthys felis*** (Girard). *Wolf eel.*

Pacific coast, from Monterey north to Puget Sound; common.

Anarrhichthys felis Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 150,
 San Francisco; brief notice.

Anarrhichthys ocellatus Ayres, Proc. Calif. Acad. Nat. Sc., I, 1855, 31,
 San Francisco.

Family 285. CERDALIDÆ

Genus 1355. **MICRODESMUS** Günther.

Microdesmus Günther, Proc. Zool. Soc. Lond., 1864, 26 (*M. dipus*
 Günther).

3823. *Microdesmus dipus* Günther.
Pacific coast of tropical America.
Microdesmus dipus Günther, Proc. Zool. Soc. Lond., Jan. 26, 1864,
26, pl. 3, fig. 2, Central America.
3824. *Microdesmus retropinnis* Jordan and Gilbert.
Panama.
Microdesmus retropinnis Jordan and Gilbert, Bull. U. S. Fish Com., I,
1881, 331, Panama.
- Genus 1356. **CERDALE** Jordan and Gilbert.
Cerdale Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881, 332 (*C.
ionthas* Jordan and Gilbert).
3825. *Cerdale ionthas* Jordan and Gilbert.
Panama.
Cerdale ionthas Jordan and Gilbert, Bull. U. S. Fish Com., I, 1881,
332, Panama.
- Genus 1357. **LEPTOCERDALE** Weymouth.
Leptocerdale Weymouth, Proc. U. S. Nat. Mus., XXXVIII, 1910,
142 (*L. longipinnis* Weymouth).
3826. *Leptocerdale longipinnis* Weymouth.
Coast of Louisiana.
Leptocerdale longipinnis Weymouth, Proc. U. S. Nat. Mus., XXXVIII,
1910, 142, Cameron, La.

Family 286. SCYTALINIDÆ

- Genus 1358. **SCYTALINA** Jordan and Gilbert.
Scytalina Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880, 266
(*S. cerdale* Jordan and Gilbert).
Scitellus Jordan and Gilbert, Proc. U. S. Nat. Mus., VI, 1883, III;
same type, there being already a genus *Scytalinus*.

3827. *Scytalina cerdale* Jordan and Gilbert.
Straits of Juan de Fuca, Waadda Island, near Cape Flattery; also in
Aleutian Islands burrowing deep in gravel and sand.
Scytalina cerdale Jordan and Gilbert, Proc. U. S. Nat. Mus., III, 1880,
267, Waadda Island, Neah Bay.

Family 287. ZOARCIDÆ

- Genus 1359. **MACROZOARCES** Gill.
Macrozoarces Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 258 (*Blennius
anguillaris* Peck).

3828. *Macrozoarces americanus* (Bloch and Schneider). *Eelpout; Muttonfish;*
Mother-of-eels; Ling; Lamper eel; Conger eel.
Delaware to Labrador; rather common northward.
? *Enchelyopus* Gronow, Zoophyl., No. 266, 1763, 77, America.
? *Blennius americanus* Bloch and Schneider, Syst. Ichth., 1801, 171,
America; after Gronow.
Blennius anguillaris Peck, Mem. Amer. Acad. Sci., II, 1804, 46, fig. 3,
New Hampshire.
? *Blennius fimbriatus* Mitchell, Trans. Lit. and Phil. Soc. N. Y., 1815,
374, pl. 1, fig. 6, New York.
Blennius ciliatus Mitchell, loc. cit., 374, pl. 1, fig. 6, New York.
Zoarces labrosus Cuvier, Règne Anim., ed. 2, II, 1829, 240, America.
Zoarces gronovii Cuvier and Valenciennes, Hist. Nat. Poiss., XI, 1836,
246 (469); after Gronow.

- Genus 1360. **EMBRYX** Jordan and Evermann.
Embryx Jordan and Evermann, Fishes North and Mid. Amer., 1898,
2458 (*Lycodopsis crotalinus* Gilbert).

3829. *Embryx crotalinus* (Gilbert).
Northern Pacific and off Santa Barbara Islands.
Lycodopsis crotalinus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 105,
off Santa Barbara Islands at Albatross station 2980 in 603 fathoms.

- 3830.** *Embryx crassilabris* (Gilbert).
 Coast of southern California.
Lycodopsis crassilabris Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 106,
 off southern California at *Albatross* station 2839.
- 3831.** *Embryx parallelus* Gilbert.
 Monterey Bay.
Embryx parallelus Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 360,
 Pl. X, IX, fig. 16, Monterey Bay, 394 to 406 fathoms.
- Genus 1361. **LYCODOPSIS** Collett.
Lycodopsis Collett, Proc. Zool. Soc. Lond., 1879, 381 (*Lycodes pacificus* Collett).
Leurynnis Lockington, Proc. U. S. Nat. Mus., II, 1879, 326 (*L. paucidens* Lockington = *Lycodes pacificus* Collett).
- 3832.** *Lycodopsis pacificus* (Collett).
 San Francisco to central Alaska; Afognak Bay.
Lycodes (*Lycodopsis*) *pacificus* Collett, Proc. Zool. Soc. Lond., 1879, 381; said to be from Japan; doubtless an error.
Leurynnis paucidens Lockington, Proc. U. S. Nat. Mus., II, 1879, 326, off San Francisco, Calif.
- 3833.** *Lycodopsis scaurus* Garman.
 Off Bay of Panama in deep water,
Lycodopsis scaurus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 132, Pl. XXXII, off Bay of Panama.
- Genus 1362. **APRODON** Gilbert.
Aprodon Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 106 (*A. cortezianna* Gilbert).
- 3834.** *Aprodon cortezianna* Gilbert.
 Cortez Banks, near San Diego, Calif.
Aprodon cortezianna Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 107, Cortez Banks, off San Diego, at *Albatross* stations 2925 and 2948 in 339 and 266 fathoms.
- Genus 1363. **LYCODES** Reinhardt.
Lycodes Reinhardt, Königl. Danske Vidensk. Selsk. Naturv. Math. Afh., VII, 1838, 153 (*L. vahli* Reinhardt).
- 3835.** *Lycodes vahlii* Reinhardt. *Vahl's lydodes*; *Misarkornak*.
 Coast of Greenland.
Lycodes vahlii Reinhardt, Königl. Danske Vidensk. Selsk. Nat. Math. Afh., VII, 1838, 153, pl. 5, Greenland.
- 3836.** *Lycodes frigidus* Collett.
 North Atlantic, off New England coast; Spitzbergen.
Lycodes frigidus Collett, Forh. Vid. Selsk. Christ., 1878, Nos. 14, 45, near Spitzbergen.
- 3837.** *Lycodes terræ-novæ* Collett.
 Newfoundland Banks.
Lycodes terræ-novæ Collett, Campagnes Scientifiques *L'Hirondelle*, Poiss., 1896, 55, Newfoundland Banks in 25 fathoms.
- 3838.** *Lycodes reticulatus* Reinhardt. *Akulliaqitsok*; *Spotted lycodes*.
 North Atlantic from Greenland south to Narragansett Bay; also northern Europe.
Lycodes reticulatus Reinhardt, Königl. Danske Vid. Selsk. Afh., VII, 1838, 167, Greenland.
Lycodes perspicillum Krøyer, Dansk. Vidensk. Selsk. Afhandl., XI, 1845, p. cxxxiii, Newfoundland.
- 3839.** *Lycodes esmarkii* Collett.
 North Atlantic; Gulf Stream, in about latitude 40°, to Finmark and Spitzbergen.
Lycodes esmarkii Collett, Norges Fiske, 1875, 95; American specimens from the Gulf Stream in about latitude 40°.
- 3840.** *Lycodes jordani* Evermann and Goldsborough.
 Puget Sound and southeastern Alaska.
Lycodes jordani Evermann and Goldsborough, Bull. U. S. Bur. Fish., XXVI, 1906, 343, fig. 120, Puget Sound.

3841. *Lycodes gracilis* Sars.
 Atlantic Ocean off Nova Scotia.
Lycodes gracilis Sars, Christ. Vid. Selsk. Forh., 1866, 40-45, Dröbak,
 Norway.
Lycodes zoarchus Goode and Bean, Oceanic Ichth., 1896, 308, off Nova
 Scotia.
3842. *Lycodes concolor* Gill and Townsend.
 Bering Sea.
Lycodes concolor Gill and Townsend, Proc. Biol. Soc. Wash., XI, 1897,
 233, Bering Sea in 276 fathoms.
3843. *Lycodes digitatus* Gill and Townsend.
 Bering Sea.
Lycodes digitatus Gill and Townsend, Proc. Biol. Soc. Wash., XI,
 1897, 232, Bering Sea in 49 fathoms.
3844. *Lycodes palearis* Gilbert.
 Bristol Bay, Alaska.
Lycodes palearis Gilbert, Rept. U. S. Fish Com., XIX, 1893, 454,
 Bristol Bay, Alaska, at *Albatross* stations 3253 and 3254 in 36 and
 46 fathoms.
3845. *Lycodes brevipes* Bean.
 Aleutian Islands to Puget Sound.
Lycodes brevipes Bean, Proc. U. S. Nat. Mus., XIII, 1890, 38, between
 Unga and Nagai islands in 110 fathoms.
3846. *Lycodes anguis* Garman.
 Off Lower California and Panama in deep water.
Lycodes anguis Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 133,
 off Bay of Panama.
3847. *Lycodes serpens* Garman.
 Off Lower California in deep water.
Lycodes serpens Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 134,
 off Gulf of California.
3848. *Lycodes incisus* Garman.
 Off Bay of Panama in deep water.
Lycodes incisus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 135,
 Pl. XXX, off Bay of Panama.
3849. *Lycodes cicatrifer* Garman.
 Off Bay of Panama in deep water.
Lycodes cicatrifer Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 136,
 Pl. XXXI, off Bay of Panama.
- Genus 1864. **LYCIAS** Jordan and Evermann.
Lycias Jordan and Evermann, Fishes North and Mid. Amer., 1898,
 2461 (*Lycodes seminudus* Reinhardt).
3850. *Lycias nebulosus* (Krøyer). *Cloudy lycodes*.
 Greenland.
Lycodes nebulosus Krøyer, Køgl. Danske Vidensk. Selsk. Afhand., XI,
 1845, p. ccxxii, Greenland.
3851. *Lycias seminudus* (Reinhardt). *Half-naked lycodes*.
 North Atlantic, from Greenland to Spitzbergen.
Lycodes seminudus Reinhardt, Køgl. Danske Selsk. Afhand., VII,
 1838, 221, Omenak, Greenland.
- Genus 1865. **LYCENCHELYS** Gill.
Lycenchelys Gill, Proc. Acad. Nat. Sci. Phila., XXXVI, 1884, 180
 (*Lycodes murana* Collett).
3852. *Lycenchelys verrillii* (Goode and Bean).
 Coast of Massachusetts and northward.
Lycodes verrillii Goode and Bean, Amer. Jour. Sci. Arts, XIV, 1877,
 474; 27 miles south and west of Chebuoto Head near Halifax.

3853. *Lycenchelys paxillus* (Goode and Bean).

Gulf Stream.

Lycodes paxillus Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 44,
between La Have and Sable Island Banks.

Lycodes paxilloides Goode and Bean, Bull. Mus. Comp. Zool., X, No.
5, 1883, 207, off Newfoundland.

3854. *Lycenchelys porifer* (Gilbert).

Off Lower California.

Lycodes porifer Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 104, off
Lower California at *Albatross* station 3009 in 857 fathoms.

Genus 1366. LYCODALEPIS Bleeker.

Lycodalepis Bleeker, Versl. Akad. Amst., ed. 2, VIII, 1874, 369 (*Lycodes*
mu osus Richardson).

3855. *Lycodalepis polaris* (Sabine). *Arctic lycodes*.

Gulf of Georgia and Alaska.

Blennius polaris Sabine, Parry's Jour., Voyage 1819-20, App. CCXII,
north Georgia.

Lycodes turneri Bean, Proc. U. S. Nat. Mus., I, 1878, 463, St. Michaels,
Alaska.

Lycodes coccineus Bean, Proc. U. S. Nat. Mus., IV, 1881, 144, Big
Diomede Island, Bering Strait.

3856. *Lycodalepis mucosus* (Richardson).

Arctic seas.

Lycodes mucosus Richardson, Last Arctic Voyage, 1855, 362, pl. 26,
Northumberland Sound.

Genus 1367. FURCIMANUS Jordan and Evermann.

Furcella Jordan and Evermann, Fishes North and Mid. Amer., 1896,
480 (*Lycodes diapterus* Gilbert); name preoccupied.

Furcimanus Jordan and Evermann, Fishes North and Mid. Amer.,
1898, 2472 (*Lycodes diapterus* Gilbert).

3857. *Furcimanus diapterus* (Gilbert).

Coasts of California and Oregon.

Lycodes diapterus Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 564, off
coast of Oregon at *Albatross* stations 2892, 2896, and others in 82
to 376 fathoms.

Genus 1368. LYCODONUS Goode and Bean.

Lycodonus Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1883,
208 (*L. mirabilis* Goode and Bean).

3858. *Lycodonus mirabilis* Goode and Bean.

Off the New England coast.

Lycodonus mirabilis Goode and Bean, Bull. Mus. Comp. Zool., X,
No. 5, 1883, 208, lat. $38^{\circ} 20' 8''$ N., long. $73^{\circ} 23' 20''$ W., in 740
fathoms.

Genus 1369. LYCONEMA Gilbert.

Lyconema Gilbert, Rept. U. S. Fish Com., XIX, 1893, 471 (*L. barbatum*
Gilbert).

3859. *Lyconema barbatum* Gilbert.

Coast of Alaska.

Lyconema barbatum Gilbert, Rept. U. S. Fish Com., XIX, 1893, 471,
pl. 35, coast of Alaska, at *Albatross* station 3129 in 204 fathoms.

Genus 1370. LYCOGRAMMA Gilbert.

Lycogramma Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 364,
(*Maynea brunnea* Bean).

3860. *Lycogramma brunnea* (Bean).

Coast of California to Alaska.

Maynea brunnea Bean, Proc. U. S. Nat. Mus., XIII, 1890, 38, off
Queen Charlotte Island in 876 fathoms.

Genus 1371. MAYNEA Cunningham.

Maynea Cunningham, Trans. Linn. Soc. Lond., XXVII, 1871, 471
(*M. patagonica* Cunningham).

3861. *Maynea californica* Gilbert.
Santa Barbara Islands.
Maynea californica Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 362,
pl. 19, fig. 17, San Nicolas Island off southern California.
3862. *Maynea bulbiceps* Garman.
Off Bay of Panama in deep water.
Maynea bulbiceps Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 140,
Pl. E, off Bay of Panama.
- Genus 1372. **BOTHROCARA** Bean.
Bothrocara Bean, Proc. U. S. Nat. Mus., XIII, 1890, 38 (*B. mollis* Bean).
3863. *Bothrocara pusilla* (Bean).
Coast of Alaska north to Unalaska.
Maynea pusilla Bean, Proc. U. S. Nat. Mus., XIII, 1890, 39, Aleutian
Islands, lat. $55^{\circ} 10' N.$, long. $160^{\circ} 18' W.$, in 110 fathoms.
3864. *Bothrocara mollis* Bean.
North Pacific to southern California.
Bothrocara mollis Bean, Proc. U. S. Nat. Mus., XIII, 1890, 39, off
Queen Charlotte Islands in 876 fathoms.
3865. *Bothrocara remigera* Gilbert.
Monterey Bay.
Bothrocara remigera Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 366,
pl. 20, fig. 19, Monterey Bay in 718 to 756 fathoms.
- Genus 1373. **BOTHROCAROPSIS** Garman.
Bothrocaropsis Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 127
(*B. alalonga* Garman).
3866. *Bothrocaropsis alalonga* Garman.
Off Gulf of California in deep water.
Bothrocaropsis alalonga Garman, Mem. Mus. Comp. Zool., XXIV,
1899, 127, XXXII, off Gulf of California.
3867. *Bothrocaropsis rictolata* Garman.
Off Bay of Panama in deep water.
Bothrocaropsis rictolata Garman, Mem. Mus. Comp. Zool., XXIV,
1899, 128, Pl. XXIII, off Bay of Panama.
3868. *Bothrocaropsis elongata* Garman.
Off Bay of Panama in deep water.
Bothrocaropsis elongata Garman, Mem. Comp. Zool., XXIV, 1899, 129,
Pl. XXXIII, off Bay of Panama.
- Genus 1374. **GYMNELIS** Reinhardt.
Gymnelis Reinhardt, Dansk. Vidensk. Selsk. Afhandl., VII, 1838, 116,
131 (*Ophidium viridis* Fabricius).
Cepolopolis Kaup, Archiv für Natur., 1856, 96 (*Ophidium viridis*
Fabricius).
3869. *Gymnelis viridis* (Fabricius). *Unernak*.
Arctic seas south to Alaska and Nova Scotia.
Ophidium viridis Fabricius, Fauna Græn., 1780, 141, Greenland.
Ophidium unernal Lacépède, Hist. Nat. Poiss., II, 1800, 280, Greenland.
? *Gymnelus pictus* Günther, Cat., IV, 1862, 324, no locality.
3870. *Gymnelis stigma* (Lay and Bennett).
Kotzebue Sound, Alaska.
Ophidium stigma Lay and Bennett, Zool. Beechey's Voy., 1839, 67,
pl. 20, fig. 1, Kotzebue Sound.
Gymnelis viridis unimaculatus Richardson, Last Arctic Voyage, 1855,
371, Northumberland Sound.
3871. *Gymnelis conorhynchus* Garman.
Off Bay of Panama in deep water.
Gymnelis conorhynchus Garman, Mem. Mus. Comp. Zool., XXIV, 1899,
131, Pl. XXXI, off Bay of Panama.

Genus 1375. LYCOCARA Gill.

Uronectes Günther, Cat., IV, 1862, 325 (*Ophidium parrii* Ross); name preoccupied.

Lycocara Gill, Proc. Acad. Nat. Sci. Phila., XXXVI, 1884, 180 (*Ophidium parrii* Ross).

3872. *Lycocara parrii* (Ross).

Baffins Bay.

Ophidium parrii Ross, in Parry's Third Voyage, App., 1826, 109, Baffins Bay.

Genus 1376. MELANOSTIGMA Günther.

Melanostigma Günther, Proc. Zool. Soc. Lond., 1881, 20 (*M. gelatinosum* Günther).

3873. *Melanostigma gelatinosum* Günther.

Deep waters of the western Atlantic; originally known from the Straits of Magellan, but since obtained in various localities from Cape Cod to the West Indies.

Melanostigma gelatinosum Günther, Proc. Zool. Soc. Lond., 1881, 21, pl. 2, fig. A, Straits of Magellan.

3874. *Melanostigma pammelas* Gilbert.

Coast of Alaska.

Melanostigma pammelas Gilbert, Rept. U. S. Fish Com., XIX, 1893, 472, pl. 35, coast of Alaska at *Albatross* station 3202, lat. 36° 46' 10" N., long. 121° 58' 45" W., in 382 fathoms.

Genus 1377. PHYCOCETES Jenyns.

Phycocetes Jenyns, Zool. Beagle, 1842, 168 (*P. latitans* Jenyns).

3875. *Phycocetes suspectus* Garman.

Off Gulf of California in deep water.

Phycocetes suspectus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 137, Pl. XXX, off Gulf of California.

Family 288. LYCODAPODIDÆ**Genus 1378. LYCODAPUS Gilbert.**

Lycodapus Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 107 (*L. ferasfer* Gilbert).

3876. *Lycodapus ferasfer* Gilbert.

Pacific coast of America, Lower California to southeastern Alaska.

Lycodapus ferasfer Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 108, off Lower California and Washington at *Albatross* stations 2980, 3010, and 3072 in 610 to 1005 fathoms.

3877. *Lycodapus dermatinus* Gilbert.

Coast of California in deep water.

Lycodapus dermatinus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 471, pl. 35, coast of California at *Albatross* station 3162, 37° 54' 10" N., 123° 30' W., in 552 fathoms.

3878. *Lycodapus parviceps* Gilbert.

Bering Sea, north of Unalaska Island.

Lycodapus parviceps Gilbert, Rept. U. S. Fish Com., XIX, 1893, 455, Bering Sea north of Unalaska Island at *Albatross* station 3324 in 109 fathoms.

3879. *Lycodapus extensus* Gilbert.

Bering Sea north of Unalaska Island.

Lycodapus extensus Gilbert, Rept. U. S. Fish Com., XIX, 1893, 455, Bering Sea north of Unalaska Island at *Albatross* station 3324 in 109 fathoms.

3880. *Lycodapus lycodon* Gilbert.

Monterey Bay.

Lycodapus lycodon Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 371, Monterey Bay, 152 to 268 fathoms.

3881. *Lycodapus grossidens* Gilbert.
Bering Sea.
Lycodapus grossidens Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915,
373, Funter Bay, southeast Alaska, in 350 fathoms.
3882. *Lycodapus attenuatus* Gilbert.
Monterey Bay.
Lycodapus attenuatus Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915,
372, pl. 21, Monterey Bay, in 285 to 357 fathoms.
3883. *Lycodapus mandibularis* Gilbert.
Monterey Bay.
Lycodapus mandibularis Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915,
369, pl. 20, fig. 20, Monterey Bay, in 144 to 293 fathoms.

Family 289. DEREPODICHTHYIDÆ

Genus 1379. DEREPODICHTHYS Gilbert.
Derepodichthys Gilbert, Rept. U. S. Fish Com., XIX, 1893, 456 (*D. alepidotus* Gilbert).

3884. *Derepodichthys alepidotus* Gilbert.
Coast of British Columbia, off Queen Charlotte Island.
Derepodichthys alepidotus Gilbert, Rept. U. S. Fish Com., XIX, 1893,
456, Queen Charlotte Island at Albatross station 3342 in 1,588 fathoms.

Family 290. BROTULIDÆ

Genus 1380. BROTULA Cuvier.
Brotula Cuvier, Règne Anim., ed. 2, II, 1829, 335 (*Enchelyopus barbatus* Bloch and Schneider).

3885. *Brotula barbata* (Bloch and Schneider). *Brotula*.
West Indies.
Enchelyopus barbatus Bloch and Schneider, Syst. Ichth., 1801, 52,
Habana; after Parra.
Brotula Parra, Dif. Piezas Hist. Nat., 1787, 70, lam. 31, fig. 2, Habana.

Genus 1381. STYGICOLA Gill.
Stygicola Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 252 (*Lucifuga dentatus* Poey).

3886. *Stygicola dentatus* (Poey).
Caves in southern Cuba.
Lucifuga dentatus Poey, Memorias, II, 1861, 102, cave of Cajio, Cuba.
- Genus 1382. LUCIFUGA** Poey.
Lucifuga Poey, Memorias, II, 1860, 95 (*L. subterraneus* Poey).

3887. *Lucifuga subterranea* Poey. *Pez ciego*.
Cuba, in caves.
Lucifuga subterraneus Poey, Memorias, II, 1861, 96, caves of San Antonio in southern Cuba.

Genus 1383. BROSMOPHYCIS Gill.
Brosmophycis Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 168
(*Brosmius marginatus* Ayres).
Halismus Ayres, Proc. Calif. Acad. Nat. Sci., II, 1861, 52 (*Brosmius marginatus* Ayres).

3888. *Brosmophycis marginatus* (Ayres).
San Francisco to Puget Sound; scarce, recently taken at San Pedro,
Calif.
Brosnius marginatus Ayres, Proc. Calif. Acad. Nat. Sci., I, 1854, 13,
Bay of San Francisco.

Genus 1384. OGILBIA Jordan and Evermann.
Ogilbia Jordan and Evermann, in Evermann and Kendall, Bull. U. S.
Fish Com., XVII, 1897, 132 (*O. cayorum* Evermann and Kendall).

- 3889.** *Ogilbia cayorum* Evermann and Kendall.
Key West.
Ogilbia cayorum Evermann and Kendall, Bull. U. S. Fish Com., XVII, 1897, 132, pl. 9, fig. 14, Key West, Fla.
- 3890.** *Ogilbia verrillii* (Garman).
Bermudas.
Brosmophycis verrillii Garman, Trans. Conn. Acad. Sci., X, 1900, 511, Bailey Bay, Bermuda Islands.
- 3891.** *Ogilbia ventralis* (Gill).
Lower California, about Cape San Lucas.
Brosmophycis ventralis Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 253, Cape San Lucas.
- Genus 1385. BYTHITES Reinhardt.**
Bythites Reinhardt, Königl. Dansk. Vidensk. Selsk. Skrift. Afhandl., VII, 1837, 178 (*B. fuscus* Reinhardt).
- 3892.** *Bythites fuscus* Reinhardt.
The only specimen known, now in the Museum at Copenhagen, was obtained in Greenland nearly a century ago.
Bythites fuscus Reinhardt, Königl. Dansk. Vidensk. Selsk. Skrift. Afhandl., VII, 1838, 179, Greenland.
- Genus 1386. CATÆTYX Günther.**
Catætyx Günther, Challenger Rept., XXII, 1887, 104 (*Sirembo messieri* Günther).
- 3893.** *Catætyx rubrirostris* Gilbert.
Coast of California south of Point Conception, in deep water.
Catætyx rubrirostris Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 111, coast of California, at Albatross stations 2909, 2925, and 2936 in 205 and 356 fathoms.
- 3894.** *Catætyx simus* Garman.
Off Galapagos Islands to Bay of Panama, in deep water.
Catætyx simus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, Pl. E 1, 168, off Bay of Panama.
- Genus 1387. DICROMITA Goode and Bean.**
Dicromita Goode and Bean, Oceanic Ichth., 1895, 319 (*D. agassizii* Goode and Bean).
- 3895.** *Dicromita agassizii* Goode and Bean.
Granada and Gulf of Mexico.
Dicromita agassizii Goode and Bean, Oceanic Ichth., 1896, 319, fig. 285, Gulf of Mexico in 26 fathoms.
- Genus 1388. MONOMITOPUS Alcock.**
Monomitopus Alcock, Ann. Mag. Nat. Hist., ser. 6, VI, 1890, 297 (*Sirembo nigripinnis* Alcock).
- 3896.** *Monomitopus torvus* Garman.
Off Bay of Panama in deep water.
Monomitopus torvus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 157, Pl. XL, off Bay of Panama.
- Genus 1389. MONOMEROPUS Garman.**
Monomeropus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 157 (*Monomeropus malispinosus* Garman).
- 3897.** *Monomeropus malispinosus* Garman.
Off Bay of Panama in deep water.
Monomeropus malispinosus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 158, Pl. XL, off Bay of Panama.
- Genus 1390. BASSOZETUS Gill.**
Bathynectes Günther, Ann. Nat. Hist., II, 1878, 20 (*B. compressus* Günther); name preoccupied.
Bassozetus Gill, Proc. U. S. Nat. Mus., VI, 1883, 259 (*B. normalis* Gill).
Bathyonus Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1886, 603 (*Bassozetus catena* Goode and Bean).

- 3898. Bassozetus compressus** (Günther).
 Gulf of Mexico and elsewhere in deep water.
Bathynectes compressus Günther, Ann. Mag. Nat. Hist., II, 1878, 20,
 southeast of New Guinea in 1,400 fathoms.
- 3899. Bassozetus normalis** Gill.
 West Indies, off Dominica, to 40° north latitude, in region of Gulf Stream.
Bassozetus normalis Gill, Proc. U. S. Nat. Mus., VI, 1883, 259, Gulf Stream, at Albatross station 2042, 39° 33' N., 68° 26' 45" W., in 1,555 fathoms.
- 3900. Bassozetus catena** (Goode and Bean).
 Gulf of Mexico in deep water.
Bathyonus catena Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 603, at Albatross station 2379, 28° 00' 15" N., 87° 42' W., in 1,467 fathoms.
- 3901. Bassozetus tænia** (Günther).
 Mid-Atlantic.
Bathyonus tænia Günther, Challenger Rept., XXII, 1887, 110, pl. 23, fig. A, Mid-Atlantic.
- 3902. Bassozetus nasus** Garman.
 Off Gulf of Panama and Pacific coast of southern Mexico in deep water.
Bassozetus nasus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 159, Pl. LXXVII, off Pacific coast of southern Mexico.
- Genus 1391. ERETMICHTHYS Garman.
Eretmichthys Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 164 (*E. pinnatus* Garman).
- 3903. Eretmichthys pinnatus** Garman.
 Galapagos Islands to Bay of Panama in deep water.
Eretmichthys pinnatus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 165, Pl. XXXV, off Bay of Panama.
- 3904. Eretmichthys ocella** Garman.
 Off Bay of Panama in deep water.
Eretmichthys ocella Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 166, Pl. XXVII, off Bay of Panama.
- Genus 1392. MÆBIA Goode and Bean.
Mæbia Goode and Bean, Oceanic Ichth., 1895, 331 (*Bathynectes gracilis* Günther).
- 3905. Mæbia promelas** (Gilbert).
 Gulf of California.
Porogadus promelas Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 546, Gulf of California in 1,005 fathoms.
- Genus 1393. NEOBYTHITES Goode and Bean.
Neobythites Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 600 (*N. gillii* Goode and Bean).
- 3906. Neobythites gillii** Goode and Bean.
 Gulf of Mexico to Brazil, in deep water.
Neobythites gillii Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 601, Gulf of Mexico, at Albatross station 2402, 28° 36' N., 85° 33' W., in 111 fathoms.
Neobythites ocellatus Günther, Challenger Rept., XXII, 1887, 103, pl. 21, fig. B, off Pernambuco in 350 fathoms.
- 3907. Neobythites marginatus** Goode and Bean.
 Off Barbados.
Neobythites marginatus Goode and Bean, Bull. Mus. Comp. Zool., XII, 1886, off Barbados in 209 fathoms.
- Genus 1394. HOLCOMYCTERONUS Garman.
Holcomycteronus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 162 (*H. digitatus* Garman).

- 3908. Holomycteronus digittatus Garman.**
Gulf of California to Colombia in deep water.
Holomycteronus digittatus Garman, Mem. Mus. Comp. Zool., XXXVI, 1899, 163, Gulf of California.
- Genus 1395. BENTHOCOMETES Goode and Bean.**
Benthocometes Goode and Bean, Oceanic Ichth., 1895, 327 (*Neobythites robustus* Goode and Bean).
- 3909. Benthocometes robustus** (Goode and Bean).
West Indies to lat. 29° N. in Gulf Stream.
Neobythites robustus Goode and Bean, Bull. Mus. Comp. Zool., XII, 1883, 161, off Morro Castle, Cuba.
- Genus 1396. BASSOGIGAS Gill.**
Bassogigas Gill, in Goode and Bean, Oceanic Ichth., 1895, 328 (*B. gillii* Goode and Bean).
- 3910. Bassogigas gillii** Goode and Bean.
Off Cape Henlopen, Del.
Bassogigas gillii Goode and Bean, Oceanic Ichth., 1895, 328, fig. 291, off Cape Henlopen, Del., at Albatross station 2684 in 1,106 fathoms.
- 3911. Bassogigas stelliferoides** (Gilbert).
Off Bay of Panama in deep water, and Gulf of California.
Neobythites stelliferoides Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 112, off Lower California at Albatross station 2996 in 112 fathoms.
- Genus 1397. BARATHRODEMUS Goode and Bean.**
Barathrodemus Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1882, 200 (*B. manatinus* Goode and Bean).
- 3912. Barathrodemus manatinus** Goode and Bean.
Gulf Stream, north to Bermudas.
Barathrodemus manatinus Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1882, 200, lat. 33° 51' 20" N., long. 76° W., in 647 fathoms.
- Genus 1398. NEMATONUS Günther.**
Nematonus Günther, Challenger Rept., XXII, 1887, 114 (*Bathyonus pectoralis* Goode and Bean).
- 3913. Nematonus pectoralis** (Goode and Bean).
Gulf of Mexico and off Dominica.
Bathyonus pectoralis Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 604, Albatross station 2380, lat. 28° 02' 30" N., long. 87° 43' 45" W., in 1,430 fathoms.
- Genus 1399. POROGADUS Goode and Bean.**
Porogadus Goode and Bean Proc. U. S. Nat. Mus., VIII, 1885, 602 (*P. miles* Goode and Bean).
- 3914. Porogadus miles** Goode and Bean.
Atlantic coast, off Delaware Bay, in deep water.
Porogadus miles Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 602, at Albatross station 2230, lat. 38° 27' N., long. 73° 2' W., in 1,168 fathoms.
- 3915. Porogadus longiceps** Garman.
Off Cocos Islands to Bay of Panama in deep water.
Porogadus longiceps Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 153, Pl. F, off Bay of Panama.
- 3916. Porogadus atripectus** Garman.
Off Bay of Panama and Pacific coast of Mexico in deep water.
Porogadus atripectus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 154, Pl. XXXVII, off Bay of Panama.
- 3917. Porogadus breviceps** Garman.
Gulf of California in deep water.
Porogadus breviceps Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 155, Gulf of California.
- Genus 1400. PENOPUS Goode and Bean.**
Penopus Goode and Bean, Oceanic Ichth., 1895, 335 (*P. macdonaldi* Goode and Bean).

3918. *Penopus macdonaldi* Goode and Bean.

Gulf Stream.

Penopus macdonaldi Goode and Bean, Oceanic Ichth., 1895, 336, fig. 293, Gulf Stream at Albatross station 2716, lat. $38^{\circ} 29' 30''$, long. $70^{\circ} 57'$ W., in 1,631 fathoms.

Genus 1401. *DICROLENE* Goode and Bean.

Dicrolene Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1882, 202 (*D. introniger* Goode and Bean).

3919. *Dicrolene intronigra* Goode and Bean.

Gulf Stream, from lat. 15° to 40° N.

Dicrolene introniger Goode and Bean, Bull. Mus. Comp. Zool., X, No. 5, 1882, 202, Gulf Stream, lat. 34° N.

3920. *Dicrolene pullata* Garman.

Off Gulf of Panama in deep water.

Dicrolene pullata Garman, Mem. Comp. Zool., XXIV, 1899, 152, off Gulf of Panama.

3921. *Dicrolene nigra* Garman.

Off Galapagos Islands to Bay of Panama in deep water.

Dicrolene nigra Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 150, Pl. XXXVII, off Galapagos Islands.

3922. *Dicrolene filamentosa* Garman.

Off Bay of Panama and Pacific coast of Mexico in deep water.

Dicrolene filamentosa Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 149, Pl. F, off Bay of Panama.

Genus 1402. *MIXONUS* Günther.

Mixonus Günther, Challenger Rept., XXII, 1887, 108 (*Bathynectes laticeps* Günther).

3923. *Mixonus laticeps* (Günther).

Mid-Atlantic in deep water.

Bathynectes laticeps Günther, Ann. Mag. Nat. Hist., II, 1878, 20, Mid-Atlantic at Challenger station 104 in 2,500 fathoms.

Sirembo guentheri Vaillant, Expd. Sci. Travailleur et Talisman, 1888, 268, Pl. XXIV, fig. 5, off the Cape Verde Islands at a depth of 3,200 meters.

3924. *Mixonus caudalis* Garman.

Between Galapagos and Cocos Islands in deep water.

Mixonus caudalis Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 148, Pl. XXXVI, off Cocos Islands.

Genus 1403. *ACANTHONUS* Günther.

Acanthonus Günther, Ann. Mag. Nat. Hist., II, 1878, 22 (*A. armatus* Günther).

3925. *Acanthonus spinifer* Garman.

Off Bay of Panama in deep water.

Acanthonus spinifer Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 170, Pl. F, off Bay of Panama.

Genus 1404. *LEUCICORUS* Garman.

Leucicorus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 146 (*L. lusciosus* Garman).

3926. *Leucicorus lusciosus* Garman.

Off Pacific coast of Mexico in deep water.

Leucicorus lusciosus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 146, Pl. XXXVIII, off Pacific coast of Mexico.

Genus 1405. *SCIADONUS* Garman.

Sciadonus Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 171 (*S. pedicellaris* Garman).

3927. *Sciadonus pedicellaris* Garman.

Off Cocos Island in deep water.

Sciadonus pedicellaris Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 172, Pl. F, off Cocos Island.

Genus 1406. LAMPROGRAMMUS Alcock.

Lamprogrammus Alcock, Ann. Mag. Nat. Hist., VIII, 1891, 32 (*L. niger* Alcock).

3928. Lamprogrammus illustris Garman.

Off Bay of Panama.

Lamprogrammus illustris Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 174, Pl. XXXIV, off Bay of Panama.

Genus 1407. BARATHRONUS Goode and Bean.

Barathronus Goode and Bean, Mem. Mus. Comp. Zool., XII, 1886, 164 (*B. bicolor* Goode and Bean).

3929. Barathronus bicolor Goode and Bean.

Off Guadalupe Island in deep water.

Barathronus bicolor Goode and Bean, Bull. Mus. Comp. Zool., XII, 1886, 164, off Guadalupe at *Blake* station 71 in 769 fathoms.

Genus 1408. APHYONUS Günther.

Aphyonus Günther, Ann. Mag. Nat. Hist., II, 1878, 22 (*A. gelatinosus* Günther).

3930. Aphyonus mollis Goode and Bean.

Gulf of Mexico in deep water.

Aphyonus mollis Goode and Bean, Bull. Mus. Comp. Zool., XII, 1886, 163, Gulf of Mexico at *Blake* station 221, lat. 24° 36' N., long. 84° 5' W., in 955 fathoms.

Genus 1409. ALLECTOR Heller and Snodgrass.

Alector Heller and Snodgrass, Proc. Wash. Acad. Sci., V, 1903, 228 (*A. cheloniae* Heller and Snodgrass).

3931. Alector cheloniae Heller and Snodgrass.

Between Clipperton Island and Galapagos Archipelago.

Alector cheloniae Heller and Snodgrass, Proc. Wash. Acad. Sci., V, 1903, 228, fig. 1, between Clipperton Island and the Galapagos.

Family 291. OPHIDIIDÆ**Genus 1410. LEPOPHIDIUM Gill.**

Lepophidium Gill, Am. Nat., XXIX, 1895, 16; substitute for *Leptophidium* Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 210 (*L. profundorum* Gill); preoccupied in serpents by *Leptophidium* Hallowell, 1860.

3932. Lepophidium marmoratum (Goode and Bean).

Gulf Stream.

Leptophidium marmoratum Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 423, lat. 23° 10' 39" N., long. 82° 20' 21" W., in 213 fathoms.

3933. Lepophidium profundorum (Gill).

Gulf Stream, off coast of Florida.

Leptophidium profundorum Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 211, Gulf Stream, off coast of Florida.

3934. Lepophidium cervinum (Goode and Bean).

Gulf Stream.

Leptophidium cervinum Goode and Bean, Proc. U. S. Nat. Mus., VIII, 1885, 422, lat. 40° 01' N., long. 69° 56' W., depth 76 fathoms.

3935. Lepophidium brevibarbe (Cuvier).

West Indies and Brazil.

Ophidium brevibarbe Cuvier, Règne Anim., ed. 2, II, 1829, 359, Brazil.

3936. Lepophidium prorates (Jordan and Bollman).

Panama and south of Panama.

Leptophidium prorates Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 172, Panama and south of Panama.

3937. Lepophidium emmelas (Gilbert).

Coast of Lower California.

Leptophidium emmelas Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 110, coast of Lower California at *Albatross* stations 3007 and 3008 in 362 and 306 fathoms.

3938. *Lepophidium pardale* (Gilbert).
Lower California.
Lepophidium pardale Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 108,
off Lower California.
3939. *Lepophidium microlepis* (Gilbert).
Off Lower California.
Lepophidium microlepis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890,
109, off Lower California.
3940. *Lepophidium stigmatistium* (Gilbert).
Coast of Lower California.
Lepophidium stigmatistium Gilbert, Proc. U. S. Nat. Mus., XIII, 1890,
109, off Lower California at Albatross station 2996 in 112 fathoms.
- Genus 1411. *OPHIDION* Linnaeus. *Cusk eels.*
Ophidion Linnaeus, Syst. Nat., Ed. X, 1758, 259 (*Ophidion barbatum*
Linnaeus).
Ophidion of most authors; revised spelling.
3941. *Ophidion graëlli* Poey.
Cuba.
Ophidion graëlli Poey, Memorias, II, 1861, 425, Habana.
3942. *Ophidion beani* Jordan and Gilbert.
Gulf of Mexico; Snapper Banks off Pensacola, Fla.
Ophidion beani Jordan and Gilbert, Proc. U. S. Nat. Mus., VI, 1883, 143,
Snapper Banks off Pensacola.
3943. *Ophidion holbrooki* (Putnam).
Key West, Fla.; Gulf of Mexico.
Ophidion holbrooki Putnam, Proc. Bost. Soc. Nat. Hist., XVI, 1874,
342, Key West, Fla.
- Genus 1412. *RIS SOLA* Jordan and Evermann.
Risso Jordan and Evermann, Check List, 1895, 483 (*Ophidium marginatum* DeKay).
3944. *Rissola marginata* (DeKay). *Sand eel; Slippery dick.*
South Atlantic coast of United States from New York south to Pensacola.
Ophidium marginatum DeKay, New York Fauna: Fishes, 1842, 315,
New York Harbor.
? *Ophidium josephi* Girard, U. S. and Mex. Bound. Surv., Zool., 1859,
29, Saint Joseph Island, Tex.
- Genus 1413. *OTOPHIDIUM* Gill.
Otopholidium Gill, in Jordan, Cat. Fish. N. Amer., 1885, 914 (*Genypterus omostigma* Jordan and Gilbert).
Chilaria Jordan and Evermann, Check List, 1895, 482 (*Ophidium taylori* Girard).
3945. *Otophidiump taylori* (Girard).
Coast of California, from Monterey to San Diego.
Ophidium taylori Girard, Pac. R. R. Surv., X, Pt. 4, Fishes, 1858, 138,
Monterey, Calif.
3946. *Otophidiump omostigma* (Jordan and Gilbert).
Gulf of Mexico, Pensacola snapper banks.
Genypterus omostigma Jordan and Gilbert, Proc. U. S. Nat. Mus., V,
1882, 301, Pensacola snapper banks.
3947. *Otophidiump scrippsi* Hubbs.
Long Beach to Cedros Island, Calif.
Otophidiump scrippsi Hubbs, Univ. Calif. Publ. Zool., XVI, 1916, 166,
pl. 20, Cedros Island, off coast of Lower California.
3948. *Otophidiump galeoides* (Gilbert).
Coast of Lower California.
Otophidiump galeoides Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 110,
off Lower California.

3949. *Otophidium welshi* Nichols and Breder.

Coast of Texas.

Otophidium welshi Nichols and Breder, Proc. Biol. Soc. Wash., XXXV, 1922, 15, fig. 3, Texas, *Grampus* station 10478, southwest from Brayes Light.

Family 292. CARAPIDÆ

Genus 1414. *CARAPUS* Rafinesque.

Carapus Rafinesque, Indice, 1810, 57, restricted by the International Commission of Nomenclature to *Ophidium imberbe* ⁹⁰ Linnaeus.

Fierasfer Cuvier, Règne Anim., ed. 1, II, 1817, 239 (*Ophidium imberbe* Linnaeus).

Echiodon Thompson, Proc. Zool. Soc. Lond., 1837, 55 (*E. drummondii* Thompson).

Diaphasia Lowe, loc. cit., 1843, 92 (*Gymnotus acus* Brünich).

Oxybeles Richardson, Voy. *Erebus* and *Terror*, Fishes, 1844-48, 73 (*O. homeri* Richardson).

Porobronchus Kaup, Ann. Mag. Nat. Hist., 1860, 272 (*P. linearis* Kaup; larva of *Fierasfer acus*).

Vexillifer Gasco, Bull. Assoc. Nat. Med. Napoli, 1870, 59 (*V. dephilippi* Gasco; larva of *Fierasfer acutus*).

Lefroyia Jones, Zoologist, IX, 1874, 3838 (*L. bermudensis* Jones).

3950. *Carapus affinis* (Günther). *Pearlfish*.

Lower California to Panama; abundant about the Pearl Islands off Panama, north to Point Loma, San Diego, Calif.

Fierasfer affinis Günther, Cat., IV, 1862, 381, no locality given.

Fierasfer dubius Putnam, Proc. Bost. Soc. Nat. Hist., XVI, 1874, 344, Pearl Islands.

3951. *Carapus arenicola* (Jordan and Gilbert).

Pacific coast of Mexico.

Fierasfer arenicola Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 363, Mazatlan, Mexico.

3952. *Carapus bermudensis* (Jones).

West Indies; recorded from Bermuda, Habana, Key West, and St. Thomas.

Lefroyia bermudensis Jones, Zoologist, IX, 1874, 3837, Bermuda.

Genus 1415. LEPTOFIERASFER Meek and Hildebrand.

Leptofierasfer Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1928, 964 (*L. macrurus* Meek and Hildebrand).

3953. *Leptofierasfer macrurus* Meek and Hildebrand.

Leptofierasfer macrurus Meek and Hildebrand, Field Mus. Nat. Hist., Zool., XV, 1928, 964, Pl. XCIX, fig. 1, Chame Point, Panama.

Suborder HAPLODOCÍ

Family 293. BATRACHOIDIDÆ

Genus 1416. BATRACHOIDES Lacépède.

Batrachoides Lacépède, Hist. Nat. Poiss., II, 1800, 451 (*B. tau* Lacépède); not *Gadus tau* Linnaeus = *Batrachus surinamensis* Schneider.

Batrachus Schneider, ed. Bloch, Syst. Ichth., 1801, 42 (*B. surinamensis* Schneider); substitute for *Batrachoides*.

Batrictius Rafinesque, Anal. Nat., 1815, 82; substitute for *Batrachoides*.

3954. *Batrachoides surinamensis* (Schneider). *Sapo*.

Coast of Guiana and Brazil.

Batrachus surinamensis Schneider, ed. Bloch, Syst. Ichth., 1801, 43, Surinam; from a specimen in the museum of Paris.

⁹⁰ With this decision we do not willingly concur, as the name *Carapus* is clearly cognate with *Gymnotus carapo*, a species of Gymnotidae included but not named in the original account.

3955. **Batrachoides goldmani** Evermann and Goldsborough.
Rio Usumacinta at Montecristo, Tabasco, Mexico.
Batrachoides goldmani Evermann and Goldsborough, Bull. U. S. Fish Com., XXI, 1901, 159, fig. 8, Rio Usumacinta, Mexico.
3956. **Batrachoides boulengeri** Gilbert and Starks.
Bay of Panama.
Batrachoides boulengeri Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 182, Pl. XXXI, Bay of Panama.
3957. **Batrachoides pacifici** (Günther).
Panama.
Batrachus pacifici Günther, Cat., III, 1861, 173, Panama.
- Genus 1417. **OPSANUS** Rafinesque. *Toadfishes.*
Opsanus Rafinesque, Amer. Month. Mag., II, 1818, 203 (*O. cerapalus*).
Rafinesque=Gadus tau Linnaeus.
Batrachus of authors, not of Schneider.
3958. **Opsanus tau** (Linnaeus). *Toadfish; Sapo; Slimer; Oysterfish; Scorpion.*
Atlantic coast, Massachusetts to the West Indies.
Gadus tau Linnaeus, Syst. Nat., ed. XII, 1766, 439, Charleston, S. C.
Cottus glaber Schöpf, Schrift. Naturf. Freunde, VII, 1788, 146, Long Island.
Cottus chætodon Bloch and Schneider, Syst. Ichth., 1801, 62, New York; after Schöpf.
Lophius bufo Mitchell, Trans. Lit. Phil. Soc., I, 1815, 463, New York.
Opsanus cerapalus Rafinesque, Amer. Month. Mag., II, 1818, 204, south coast of Long Island.
Batrachoides vernellas LeSueur, Mém. Mus. Hist. Nat., V, 1819, 157, pl. 17, coast of Rhode Island.
Batrachoides variegata LeSueur, Jour. Acad. Nat. Sci. Phila., III, pt. 2, 1824, 398, Egg Harbor, N. J.
Batrachus celatus DeKay, New York Fauna: Fishes, 1842, 170, pl. 50, fig. 161, New York.
3959. **Opsanus hildebrandi** Breder.
Colon Market, Panama.
Opsanus hildebrandi Breder, Amer. Mus. Novit., No. 188, Sept. 28, 1925, I, Colon Market, Panama; place of collection unknown.
3960. **Opsanus pardus** (Goode and Bean). *Sapo.*
Gulf of Mexico in deep water.
Batrachus tau pardus Goode and Bean, Proc. U. S. Nat. Mus., II, 1879, 336, Pensacola snapper banks.
- Genus 1418. **PORICHTHYS** Girard. *Midshipmen.*
Porichthys Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 141 (*P. notatus* Girard).
3961. **Porichthys notatus** Girard. *Singingfish; Midshipman; Cabezon; Sapo.*
Pacific coast from Puget Sound to Panama.
Porichthys notatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854, 141, San Francisco.
3962. **Porichthys marginatus** (Richardson).
Pacific coast of tropical America.
Batrachus marginatus Richardson, Voy. Sulphur, Fishes, 1844-1843, 67, Pacific coast of Central America.
Porichthys naupopædium Jordan and Bollman, Proc. U. S. Nat. Mus., XII, 1889, 171, Pacific Ocean off coast of Colombia in 16 fathoms.
3963. **Porichthys greenei** Gilbert and Starks.
Panama Bay.
Porichthys greenei Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 184, Panama Bay.
- Genus 1419. **NAUTOPÆDIUM** Jordan.
Nautopædium Jordan, Proc. Acad. Nat. Sci. Phila., 1919, 342 (*Batrachus porosissimus* Cuvier and Valenciennes).

- 3964.** *Nautopædium porosissimum* (Cuvier and Valenciennes). *Bagre sapo.*
South Carolina to Montevideo; Pensacola.
Batrachus porosissimus Cuvier and Valenciennes, Hist. Nat. Poiss., XII,
1837, 373 (501), Surinam, Cayenne, Rio Janeiro, and St. Catherine.
Porichthys plectrodon Jordan and Gilbert, Proc. U. S. Nat. Mus., V,
1882, 291, Galveston, Tex.
- Genus 1420. THALASSOPHYRE** Günther. *Poison toadfishes.*
Thalassophryne Günther, Cat., III, 1861, 175 (*T. maculosa* Günther).
Dæctor Jordan and Evermann, Fishes North and Mid. Amer., 1898,
2325 (*Thalassophryne dowi* Jordan and Gilbert).
- Subgenus THALASSOPHYRE.**
- 3965.** *Thalassophryne maculosa* Günther.
Caribbean Sea.
Thalassophryne maculosa Günther, Cat., III, 1861, 175, Puerto Cabello
near Aspinwall.
- 3966.** *Thalassophryne reticulata* Günther.
Panama.
Thalassophryne reticulata Günther, Proc. Zool. Soc. Lond., 1864, 150,
155, Panama.
- Subgenus DÆCTOR** Jordan and Evermann.
- 3967.** *Thalassophryne dowi* (Jordan and Gilbert).
Pacific coast of North America from Punta Arenas to Panama.
Thalassophryne dowi Jordan and Gilbert, Proc. U. S. Nat. Mus., X,
1887, 388, Punta Arenas.
- Order YY. XENOPTERYGII**
- Family 294. GOBIESOCIDÆ**
- Genus 1421. CAULARCHUS** Gill.
Caularchus Gill, Proc. Acad. Nat. Sci. Phila., XIV, 1862, 330 (*Lepado-*
gaster mæandricus Girard = *Lepadogaster reticulatus* Girard); name
preoccupied.
- 3968.** *Caularchus mæandricus* (Girard). *Suckfish.*
Pacific coast of United States, from Vancouver Island to Monterey.
Lepadogaster reticulatus Girard, Proc. Acad. Nat. Sci. Phila., VII, 1854,
155, San Luis Obispo, Calif.; name preoccupied.
Lepadogaster mæandricus Girard, Pac. R. R. Surv., X, Pt. 4, Fishes,
1858, 130, San Luis Obispo, Calif., and the Farallons.
- Genus 1422. BRYSETÆRES** Jordan and Evermann.
Brysetæres Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., VI,
1896, 230 (*Gobiesox pinniger* Gilbert).
- 3969.** *Brysetæres pinniger* (Gilbert).
Gulf of California; known from Puerto Refugio (Angel Island), San
Luis Gonzales Bay, and La Paz.
Gobiesox pinniger Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 94, Puerto
Refugio, Gulf of California.
- Genus 1423. CAULISTIUS** Jordan and Evermann.
Caulistius Jordan and Gilbert, Check List, 1895, 491 (*Gobiesox papillifer*
Gilbert).
Bryssophilus Jordan and Evermann, Fishes North and Mid. Amer.,
1898, 2329 (*Gobiesox papillifer* Gilbert); accidental duplication over-
looked in proof reading.
- 3970.** *Caulistius papillifer* (Gilbert).
Magdalena Bay, Lower California.
Gobiesox papillifer Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 96,
Magdalena Bay, Lower California.
- Genus 1424. GOBIESOX** Lacépède. *Clingfishes.*
Gobiesox Lacépède, Hist. Nat. Poiss., II, 1800, 595 (*G. cephalus*
Lacépède).
Megaphalus Rafinesque, Anal. Nat., 1815, 86 (substitute for *Gobiesox*
Lacépède).
Sicyogaster Brisout de Barneville, Rev. Zool., 1846, 144 (*Gobiesox*
marmoratus Jenyns).

3971. **Gobiesox nigripinnis** (Peters).
 Puerto Cabello, near Aspinwall, Colon.
Cotylis nigripinnis Peters, Monatsber. Akad. Wiss. Berlin, 1859, 412,
 Puerto Cabello, near Aspinwall, Colon.
3972. **Gobiesox macrophthalmus** Günther.
 Probably West Indies.
Gobiesox macrophthalmus Günther, Cat., III, 1861, 502, locality
 unknown.
3973. **Gobiesox androsiensis** Rosen.
 Bahamas.
Gobiesox androsiensis Rosen, Arsskr. Univ. Lund., VII, 1912, 50,
 Bahamas.
3974. **Gobiesox costaricensis** Meek.
 Costa Rica.
Gobiesox costaricensis Meek, Field Mus. Nat. Hist., Zool., X, 1912, 74,
 Zent, Costa Rica.
3975. **Gobiesox gyrinus** Jordan and Evermann.
 West Indies.
Gobiesox gyrinus Jordan and Evermann, Fishes North and Mid. Amer.,
 1898, 2331, Cordova; name only in Check List, 1895, 491.
3976. **Gobiesox cephalus** Lacépède. *Tétard; Testar.*
 West Indies.
Gobiesox cephalus Lacépède, Hist. Nat. Poiss., II, 1800, 595, Mar-
 tinique.
Lepadogaster testar Bloch and Schneider, Syst. Ichth., 1801, 4, Mar-
 tinique; after Plumier.
3977. **Gobiesox tudes** Richardson.
 Probably West Indies.
Gobiesox tudes Richardson, Voy. *Sulphur*, Fishes, 1845, 103, pl. 46,
 figs. 1-3; habitat unknown, erroneously supposed to be China.
3978. **Gobiesox strumosus** Cope.
 Maryland to Florida.
Gobiesox strumosus Cope, Proc. Acad. Nat. Sci. Phila., XXII, 1870,
 121, Hilton Head, S. C.
3979. **Gobiesox virgatulus** Jordan and Gilbert.
 From Pensacola Bay north to Charleston, S. C.
Gobiesox virgatulus Jordan and Gilbert, Proc. U. S. Nat. Mus., V,
 1882, 293, Pensacola, Fla.
3980. **Gobiesox adustus** Jordan and Gilbert.
 Pacific coast of Mexico.
Gobiesox adustus Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881,
 360, Mazatlan, Mexico.
3981. **Gobiesox funebris** Gilbert.
 Gulf of California; Puerto Refugio (Angel Island) and La Paz.
Gobiesox funebris Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 95, Puerto
 Refugio, Gulf of California.
3982. **Gobiesox rhodospilus** Günther.
 Panama.
Gobiesox rhodospilus Günther, Proc. Zool. Soc. Lond., 1864, 25, Panama.
3983. **Gobiesox cerasinus** Cope.
 St. Martins, West Indies.
?Cyclopterus nudus Linnæus, Syst. Nat., ed. X, 1758, 260, "India";
 not identifiable.
Gobiesox cerasinus Cope, Trans. Amer. Phil. Soc., XIV, 1871, 473, St.
 Martins, West Indies.
3984. **Gobiesox vittatus** Metzelaar.
 Caribbean Sea.
Gobiesox vittatus Metzelaar, Marine Fish. Lesser Antilles, Bijdr.
 Dierk., 1922, 140, with fig., Curaçao.

3985. *Gobiesox sancti-martini* Metzelaar.
 West Indies.
Gobiesox sancti-martini Metzelaar, Trop. Atl. Vissch., 1919, 154, fig. 48,
 St. Martin's Island.
3986. *Gobiesox yuma* Nichols.
 Coast of Florida.
Gobiesox yuma Nichols, Bull. Amer. Mus. Nat. Hist., XXXVII, Dec.
 29, 1917, 876, fig. 1, south of Sanibel Light, Florida west coast.
- Genus 1425. **SICYASES** Müller and Troschel.
Sicyases Müller and Troschel, Wiegmann's Archiv, 1843, 298, *S. sanguineus*.
Tomicodon Brisout de Barneville, Rev. Zool., 1846, 144 (*T. chilensis* Barneville).
3987. *Sicyases erythrops* (Jordan and Gilbert).
 Mazatlan; also Tres Marias Islands.
Gobiesox erythrops Jordan and Gilbert, Proc. U. S. Nat. Mus., IV,
 1881, 360, Mazatlan, Mexico.
3988. *Sicyases rubiginosus* Poey.
 Matanzas, Cuba.
Sicyases rubiginosus Poey, Synop., 1868, 391, wharves of Palmasola,
 Matanzas, Cuba.
3989. *Sicyases carneus* Poey.
 Matanzas, Cuba.
Sicyases carneus Poey, Synop., 1868, 392, wharves of Palmasola,
 Matanzas, Cuba.
3990. *Sicyases haeres* (Jordan and Bollman).
 Green Turtle Cay, Bahamas.
Gobiesox haeres Jordan and Bollman, Proc. U. S. Nat. Mus., XI, 1888,
 552, Green Turtle Cay, Bahamas.
3991. *Sicyases punctulatus* Poey.
 Cuba.
Sicyases punctulatus Poey, Enumeratio, 1876, 124, Habana.
3992. *Sicyases fasciatus* Peters.
 Puerto Cabello near Aspinwall.
Sicyases fasciatus Peters, Monatsber. Akad. Wiss. Berlin, 1859, 412,
 Puerto Cabello.
- Genus 1426. **RIMICOLA** Jordan and Evermann.
Rimicola Jordan and Evermann, Proc. Calif. Acad. Sci., 2d ser., VI,
 1896, 231 (*Gobiesox muscarum* Meek and Pierson).
3993. *Rimicola eigenmanni* (Gilbert).
 Point Loma, near San Diego, to Monterey Bay, Calif.
Gobiesox eigenmanni Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 96,
 Point Loma near San Diego, Calif.
Gobiesox muscarum Meek and Pierson, Proc. Calif. Acad. Sci., 2d ser.,
 V, 1895, 571, pl. 71, Monterey Bay, Calif.
3994. *Rimicola beryllinus* (Hildebrand and Ginsberg).
 Key West, Fla.
Gobiesox (*Rimicola*) *beryllinus* Hildebrand and Ginsberg, Bull. U. S.
 Bur. Fish., XLII, 1926, 213, fig. 5, Boca China, Key West, Fla.
- Genus 1427. **ARBACIOSA** Jordan and Evermann.
Arbaciosa Jordan and Everman, Proc. Calif. Acad. Sci., 2d ser., VI,
 1896, 230 (*Gobiesox humeralis* Gilbert).
3995. *Abraciosa rhessodon* (Rosa Smith).
 San Diego to the northern part of the Gulf of California.
Gobiesox rhessodon Rosa Smith (Mrs. Eigenmann), Proc. U. S. Nat.
 Mus., IV, 1881, 140, San Diego, Calif.
3996. *Abraciosa humeralis* (Gilbert).
 Gulf of California; Puerto Refugio, Angel Island; La Paz.
Gobiesox humeralis Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 95,
 Puerto Refugio, Gulf of California.

3997. **Arbaciosa rupestris** (Poey).
 Coral reefs of Cuba.
Gobiesox rupestris Poey, Memorias, II, 1861, 283, Cuba.
3998. **Arbaciosa zebra** (Jordan and Gilbert).
 Mazatlan, Mexico.
Gobiesox zebra Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 359, Mazatlan, Mexico.
3999. **Arbaciosa eos** (Jordan and Gilbert).
 Pacific coast of Mexico.
Gobiesox eos Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 360, Mazatlan, Mexico.

Order ZZ. PLECTOGNATHI

Suborder SCLERODERMI

Family 295. TRIACANTHIDÆ

- Genus 1428. HOLLARDIA** Poey.
Hollardia Poey, Memorias, II, 1861, 348 (*H. hollardi* Poey).
4000. **Hollardia hollardi** Poey.
 West Indies.
Hollardia hollardi Poey, Memorias, II, 1861, 348, pl. 18, fig. 11, Cuba

Family 296. BALISTIDÆ

- Genus 1429. BALISTES** Linnæus.
Balistes Linnæus, Syst. Nat., ed. X, 1758, 327 (*B. vetula* Linnæus).
Capriscus Rafinesque, Indice, 1810, 41, 58 (*C. porcus* Rafinesque = *Balistes capriscus* Gmelin).
4001. **Balistes naufragium** Jordan and Starks. *Pez puerco de piedra*.
 Mazatlan; Panama Bay.
Balistes naufragium Jordan and Starks, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 488, Mazatlan.
4002. **Balistes capriscus** Gmelin. *Leatherjacket; Cucuyo; Turbot; Triggerfish; Filefish*.
 Tropical parts of the Atlantic; Gulf Stream; Mediterranean.
Balistes carolinensis Gmelin, Syst. Nat., I, 1788, 1468, Carolinas.
Balistes capriscus Gmelin, loc. cit., 1789, 1471, Indian and American oceans; after Gronow (this name preferred by first reviewer).
Balistes buniva Lacépède, Hist. Nat. Poiss., V, 1798, 669, pl. 21, fig. 1, Nice.
Balistes caprinus Valenciennes, Ichth. îles Canaries, 1836, 94, pl. 16, Canaries.
Balistes fuliginosus DeKay, N. Y. Fauna: Fishes, 1842, 339, pl. 57, fig. 188, New York.
Balistes tenuipterus Poey, Memorias, II, 1861, 326, Cuba.
4003. **Balistes vetula** Linnæus. *Oldwife; Old wench; Cochino; Blue-striped triggerfish; Bessy cerka*.
 Tropical parts of the Atlantic; West Indies; Gulf Stream to Woods Hole.
Guapervia Maregrave, Hist. Bras., 1648, 163, Brazil.
Balistes vetula Linnæus, Syst. Nat., Ed. X, 1758, 329, Ascension Island; after *Balistes vetula* of Osbeck, Iter Chinensis, 294, 1757.
Balistes bellus Walbaum, Artedi Pisc., III, 1792, 467, West Indies; after Kröyer.
Balistes equestris Gronow, Cat. Fishes, ed. Gray, 1854, 31, American seas.

4004. **Balistes forcipatus** Gmelin.

West coast of Africa to Rhode Island and Brazil.

Balistes forcipatus Gmelin, Syst. Nat., I, 1789, 1472, Brazil; after *Guaperva lata* of Lista.

Balistes punctatus Gmelin, loc. cit., 1472; after Stipvisch of Willoughby.

Balistes spiloptopterygius Walbaum, Artedi Pisc., III, 1792, 455, Brazil.

Balistes guttatus Walbaum, loc. cit., 467, St. Vincent.

Balistes ciliaris Bloch and Schneider, Syst. Ichth., 1801, 471, Brazil.

Balistes liberensis Steindachner. Ichth. Notizen, IV, 1867, 9, Monrovia in Liberia.

?*Balistes powelli* Cope, Proc. Acad. Nat. Sci. Phila., XXII, 1870, 120, Newport, R. I.

Balistes moribundus Cope, Trans. Amer. Phil. Soc., XIV, 1870, 478, St. Martins.

Genus 1430. **SUFFLAMEN** Jordan.

Pachynathus Swainson, Classn. Fishes, II, 1839, 326 (*Balistes triangularis* = *Balistes capistratus* Shaw); not *Pachygnatha* nor *Pachynathus*; both these names earlier used for genera of spiders, the name *Pachygnathus* regarded as a misprint and not valid according to decision of the International Commission of Nomenclature.

Sufflamen Jordan, Copeia, April 12, 1916, 27 (*Balistes capistratus* Shaw).

4005. **Sufflamen verres** (Gilbert and Starks). *Pez puerco*.

Pacific coast of tropical America, from Magdalena Bay to the Galapagos.

Balistes capistratus of authors; not of Shaw, Gen. Zool., V, 1804, 417; after *Balistes bridé* Lacépède, a species of the South Seas and East Indies.

Balistes verres Gilbert and Starks, Mem. Calif. Acad. Sci., V, 1904, 153, Pl. XXVI, fig. 49, Panama and Mazatlan.

Genus 1431. **VERRUNCULUS** Jordan.

Verrunculus Jordan, Copeia, Sept., 1924, 82 (*Balistes polylepis* Steindachner).

4006. **Verrunculus polylepis** (Steindachner). *Pez puerco*.

San Diego to Panama; common southward.

Balistes polylepis Steindachner, Ichth. Beitr., V, 1876, 21, Magdalena Bay; Mazatlan; Acapulco.

Genus 1432. **CANTHIDERMIS** Swainson. *Sobacos*.

Canthidermis Swainson, Nat. Hist. Classn. Anim., II, 1839, 325, (*Balistes angulosus* Quoy and Gaimard).

4007. **Canthidermis angulosus** (Quoy and Gaimard).

Hawaii to Cocos Island; rare.

Balistes angulosus Quoy and Gaimard, Voy. *Uranie*, Zool., 1824, 210, Sandwich Islands (Hawaiian Islands).

4008. **Canthidermis sobaco** (Poey). *Sobaco*.

West Indies.

Balistes sobaco Poey, Memorias, II, 1861, 324, Habana.

Balistes asperrimus Cope, Trans. Amer. Phil. Soc., XIV, 1870, 478, St. Martins.

4009. **Canthidermis sufflamen** (Mitchill). *Sobaco*.

West Indies; Habana.

Sobaco Parra, Dif. Piezas, Hist. Cuba, 1787, 17, fig. 10, Habana.

Balistes sufflamen Mitchill, Trans. Lit. Phil. Soc. N. Y., I, 1815, 467, locality unknown, said by DeKay to be from the south Atlantic.

Balistes macrops Poey, Memorias, II, 1861, 326, Habana.

4010. *Canthidermis maculatus* (Bloch). *Ocean turbot.*

West Indies.

Balistes maculatus Bloch, Nat. Ausl. Fisch., II, 1786, 25, West Indies.*Balistes americanus* Gmelin, Syst. Nat., I, 1789, 1473, America; after Gronow.*Balistes macropterus* Walbaum, Artedi Pisc., III, 1792, 465, America.*Balistes rufus* Gronow, Cat. Fishes, ed. Gray, 1854, 36, American oceans.*Balistes longus* Gronow, loc. cit., 1854, American oceans.*Balistes melanopterus* Cope, Trans. Amer. Phil. Soc., XIV, 1870, 478, Darien.4011. *Canthidermis willughbeii* (Lay and Bennett).

Pacific coast of Mexico.

Balistes willughbeii Lay and Bennett, Zoology Beechey's Voyage, 1839, 68, pl. 21, fig. 2, Acapulco, Mexico.Genus 1433. *XANTHICHTHYS* Kaup.*Xanthichthys* Kaup, in Richardson, Enc. Brit., ed. XII, 1856, 313 (*Balistes curassavicus* Gmelin).4012. *Xanthichthys ringens* (Linnaeus). *Cocuyo.*

West Indies and southward, recorded from Mauritius.

Balistes ringens Linnaeus, Syst. Nat., Ed. X, 1758, 329, no locality.*Cucuyo Parra*, Dif. Piezas, etc., Cuba, 1787, 19, Habana; after Linnaeus.*Balistes curassavicus* Gmelin, Syst. Nat., I, 1788, 1472, Curaçao.*Balistes notatus* Gronow, Cat. Fishes, ed. Gray, 1854, 36, both Indies.*Balistes nitidus* Gronow, loc. cit., 36, American oceans.*Balistes cicatricosus* Poey, Memorias, II, 1861, 327, Cuba.*Balistes heckeli* von Müller, Reisen in Vereinigten Staaten, Canada und Mexico, I, 1864, 182, Mexico.4013. *Xanthichthys lineopunctatus* (Hollard).

Panama; Pacific coast of Mexico to East Indies; taken at San Diego.

Balistes lineopunctatus Hollard, Ann. Sci. Nat., ser. 4, I, 1854, 65, Bourbon Island.*Balistes mento* Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 228 Clarion Island of the Revillagigedo group.Genus 1434. *MELICHTHYS* Swainson.*Melichthys* Swainson, Class. Anim., II, 1839, 325 (*Balistes ringens* Bloch = *Balistes buniva* Lacépède).4014. *Melichthys piceus* (Poey). *Galafate; Black oldwife; Calafate.*

West Indies and southward.

Balistes nigra Osbeck, Iter Chinensis, 1757, 295, Ascension Island; pre-Linnaean.*Galafate Parra*, Dif. Piezas, etc., 1787, 18, Habana.*Balistes piceus* Poey, Proc. Acad. Nat. Sci. Phila., XV, 1863, 180, Cuba.4015. *Melichthys radula* (Solander).

East Indies, Clarion and Socorro Islands; Cocos Island.

Balistes radula Solander, in Richardson, Voy. H. M. S. *Samarang*, Fishes, 1848, 21.*Melichthys bispinosus* Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 125, Clarion and Socorro Islands of the Revillagigedo group.

Family 297. MONACANTHIDÆ. Filefishes

Genus 1435. *CANTHERINES* Swainson.*Cantherines* Swainson, Classn. Fishes, II, 1839, 327 (*Cantherines nasutus* Swainson = *Balistes sandwichensis* Quoy and Gaimard).*Pseudomonacanthus* Bleeker, loc. cit., 12 (*Monacanthus macrurus* Bleeker).*Liomonacanthus* Bleeker, Ned. Tijdskr., III, 1866, 13 (*Monacanthus pardalis* Rüppell).

- 4016. *Cantherines pullus* (Ranzani). *Lija colorado*.**
 West Indies and coast of Brazil; southern Florida.
Lija colorado Parra, Dif. Piezas, etc., 1787, pl. 23, p. 49, Cuba.
Monacanthus pullus Ranzani, Nov. Comm. Act. Sci. Inst. Bonon., V, 1842, 4, pl. 1, Brazil.
Monacanthus macrocerus Holland, Ann. Sci. Nat., II, 1854, 327, pl. 12, fig. 1, Bahia.
Monacanthus ruppelii Castelnau, Anim. Nouv. Amer. Sud., Poiss., 1855, 97, pl. 47, fig. 2, Bahia.
Monacanthus irroratus Poey, Memorias, II, 1861, 330, Cuba.
Monacanthus striatus Poey, loc. cit., 329, Cuba.
Monacanthus parraianus Poey, Proc. Acad. Nat. Sci. Phila., XV, 1863, 185, Cuba; after Parra.
Monacanthus punctatus Poey, Synopsis, 1868, 437, Cuba.
- 4017. *Cantherines carolæ* (Jordan and McGregor).**
 Islands off the Pacific coast of Mexico; Revillagigedo Islands.
? *Balistes sandwichensis* Quoy and Gaimard, Voy. *l'Uranie*, Zool., 1824, 214, Sandwich [Hawaiian] Islands.
? *Cantherines nasutus* Swainson, Nat. Hist. Class. Fishes, II, 1839, 327; substitute for *B. sandwichensis*.
Cantherines carolæ Jordan and McGregor, in Jordan and Evermann, Fishes North and Mid. Amer., 1898, 1713 (Clarion Island by error), Socorro Island.
- 4018. *Cantherines amphioxys* (Cope).**
 St. Martins Island, West Indies.
Monacanthus amphioxys Cope, Trans. Amer. Phil. Soc., XIV, 1870, 477, St. Martins Island.
- Subgenus MONACANTHUS.**
- Genus 1436. MONACANTHUS Cuvier.**
Monacanthus Cuvier, Règne Anim., Ed. I, 1817, 152 (*Balistes chinensis* Bloch).
Trichoderma Swainson, Nat. Hist. Fish., Amphib., etc., II, 1839, 328 (*Balistes scapus* Lacépède).
Stephanolepis Gill, Proc. Acad. Nat. Sci. Phila., XIII, 1861, 78 (*Monacanthus setifer* Bennett).
- 4019. *Monacanthus ciliatus* (Mitchill). *Leatherfish; Lija; Flapmingo*.**
 West Indies and Florida; Florida Keys.
Balistes ciliatus Mitchill, Amer. Month. Mag. and Crit. Rev., II, March, 1818, 326, Bahama Straits.
Monacanthus piracaca Kner, Reise Fregatte Novara um Erde, Zool., Fische, 1867, 396, Rio Janeiro.
Monacanthus occidentalis Günther, Cat., VIII, 1870, 237, Puerto Caballo.
Monacanthus davidsonii Cope, Trans. Amer. Phil. Soc. Phila., XIV, 1870 (1871), 476, Florida Reef.
- Subgenus STEPHANOLEPIS Gill.**
- 4020. *Monacanthus hispidus* (Linnaeus). *Foolfish; Filefish; Leatherfish; Horny cony; Lija; Thread filefish; Mingo*.**
 Cape Cod to Cuba; south Atlantic coast and Florida Keys; West Indies to Brazil.
Balistes hispidus Linnaeus, Syst. Nat., Ed. XII, 1766, 405, Carolinas.
Balistes broccus Mitchell, Trans. Lit. Phil. Soc., I, 1815, 467, New York.
Monacanthus filamentosus Valenciennes, Iles Canaries Ichth. Atl., 1836, 95, pl. 16, fig. 1, Canaries.
Monacanthus gallinula Valenciennes, loc. cit., 95, Canaries.
Monacanthus varius Ranzani, Novi Comment Ac. Sci. Inst. Bonon., V, 1842, 6, Pl. II, Brazil.
Monacanthus massachusettsensis DeKay, N. Y. Fauna: Fishes, 1842, 336, pl. 57, fig. 187, Massachusetts Bay.
Monacanthus setifer DeKay, loc. cit., 337, pl. 59, fig. 194, New York.
Monacanthus signifer Storer, Synop. Fish. N. Amer., 1846, 497, Massachusetts.
Monacanthus auriga Lowe, Proc. Zool. Soc. Lond., 1850, 253, Madeira.

4021. **Monacanthus oppositus** Poey.

Cuba.

Monacanthus oppositus Poey, Memorias, II, 1861, 331, Habana.4022. **Monacanthus spilonotus** Cope.

Gulf of Mexico.

Monacanthus spilonotus Cope, Trans. Amer. Phil. Soc. Phila., XIV, 1870, 476, Gulf of Mexico.4023. **Monacanthus tuckeri** Bean.

West Indies, north to Bermudas.

Monacanthus tuckeri Bean, Field Mus. Nat. Hist., Zool., VII, 1906, 78, fig. 11, Bermudas.Genus 1437. **CERATACANTHUS** Gill.*Ceratacanthus* Gill, Fish. E. Coast N. Amer., Proc. Acad. Nat. Sci. Phila., XIII, 1861, 57 (*Balistes aurantiacus* Mitchell = *Balistes schœpfi* Walbaum).4024. **Ceratacanthus schœpfi** (Walbaum). *Filefish; Orange filefish; Lija; Foolfish; Unicornfish; Sunfish.*

Cape Cod to the Carolinas and Florida.

Balistes schœpfi Walbaum, Artedi Pisc., 1792, 461, Long Island; after Schöpf, Berlin Ges. Naturf., VIII, 1788, 186.*Balistes aurantiacus* Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 468, New York.*Aluterus cuspicauda* DeKay, New York Fauna: Fishes, 1842, 338, New York.*Aluterus holbrooki* Hollard, Ann. Sci. Nat., IV, 1855, 7, North America.*Aluterus cultrifrons* Hollard, Ann. Sci. Nat., ser. 4, IV, 1855, 8, pl. 1, fig. 2, New York and Bahia.Genus 1438. **DAVIDIA** Miranda Ribeiro.*Davidia* Miranda Ribeiro, Archiv Mus. Nac. Rio Janeiro, XVII, 1915, 9 (*Alutera punctata* Agassiz).4025. **Davidia punctata** (Agassiz). *Long mingo.*

Jamaica to Bahia.

Alutera punctata Agassiz, Pisc. Brasil., 1829, 134, pl. 76, Brazil.Genus 1439. **OSBECKIA** Jordan and Evermann.*Osbeckia* Jordan and Evermann, Check List, 1895, 424 (*Balistes scripta* Osbeck).4026. **Osbeckia scripta**⁹¹ (Osbeck). *Unicornfish; Lija trompa; Olilepa; Ohua.*

Tropical seas; West Indies; North Carolina; off west coast of Mexico; Clarion Island, Revillagigedo Islands.

Balistes scriptus Osbeck, Iter Chinensis, I, 1757, 144, Canaries.*Unicornis piscis bahamensis* Catesby, Nat. Hist. Carolina, etc., II, 1743, 19, pl. 19, Bahamas.*Balistes lœvis* Bloch, Naturgesch. Ausl. Fische, IV, 1795, 82, pl. 414, Morocco.*Monacanthum proboscideum* Ranzani, Nov. Comm. Ac. Sc. Inst. Bonon., V, 1842, 8, Brazil.*Aluterus venosus* Hollard, Ann. Sc. Nat., ser. 4, IV, 1855, 14, pl. 1, fig. 3, New Ireland.*Alutera picturata* Poey, Proc. Acad. Nat. Sci. Phila., XV, 1863, 183, Cuba.Genus 1440. **ALUTERA** Cuvier.*Les Alutères* Cuvier, Règne Anim., ed. 1, 1817, 153 (*Balistes monoceros* Osbeck).*Alutera* Oken, Isis, 1817, 153 ("Les Alutères" Cuvier; *Balistes monoceros* Linnæus).

⁹¹ Whether *Osbeckia venosa* Hollard, East Indies to Hawaii and the Revillagigedos, is distinct from *O. scripta* of the Atlantic is a matter not yet determined. The distinction of *Osbeckia* and *Ceratacanthus* from *Alutera* is questionable.

- 4027. *Alutera monoceros* ⁹² Linnæus.** *Laja barbuda; Loulu.*
 Pacific Ocean; East Indies; Japan; Hawaii.
Capriscus minima etc., Klein, Miasus, II, 1742, 25 (no locality; figure very bad).
Balistes monoceros Osbeck, Iter Chinensis, 1757, 110, Asia; Linnæus, Syst. Nat., ed. X, 1758, 327, Asia and America.
Laja barbuda Parra, Dif. Piezas Hist. Nat., 1787, 48, pl. 22, fig. 2, Habana.
Balistes kleinii Gmelin, Syst. Nat., 1788, 1472, Indian seas; after Klein.
Balistes barbatus Walbaum, Artedi Pisc., III, 1792, 464; after Klein.
Balistes monoceros unicolor Bloch and Schneider, Syst. Ichth., 1801, 463, after Gmelin.
Aluterus berardi Lesson, Voy. Coquille, Zool., 1830, 108, pl. 7, New Guinea.
Aluterus cinerea Temminck and Schlegel, Fauna Japon., Poiss., 1850, 292, pl. 131, fig. 1, Japan.
Alutarius macracanthus Bleeker, Verh. Bat. Gen., Balistini, XXIV, Art. 11, 1852, 22, pl. 3, fig. 7, East Indies.
Alutarius amphacanthoides Bleeker, loc. cit., 23, pl. 2, fig. 5, East Indies.
Alutarius obliteratus Cantor, Malayan Fishes, 1850, 352, Pinang.
Balistes linguatula Gronow, Cat., ed. Gray, 1854, 35, Indian seas.
Aluterus anginosus Hollard, Ann. Sci. Nat., IV, 1855, 11, East Indies.
Balistes unicornu Basilewsky, Nouv. Mém. Soc. Nat. Moscow, X, 1855, 263, China.
- 4028. *Alutera guntheriana* Poey.** *Liza barbuda.*
 West Indies; once taken at Woods Hole (1898).
Alutera guntheriana Poey, Proc. Acad. Nat. Sci. Phila., XV, 1863, 184, Habana.

Suborder OSTRACODERMI

Family 298. OSTRACIIDÆ. Trunkfishes

- Genus 1441. OSTRACION** Linnæus.
Ostracion Linnæus, Syst. Nat., Ed. X, 1758, 330 (*O. cubicus* Linnæus).
Cibotion Kaup, Wiegmann's Archiv, XXI, 1855, 215 (*Ostracion cubicus* Linnæus).
Ostracium of authors; improved orthography.
- 4029. *Ostracion punctatum*** Bloch and Schneider.
 Tropical Pacific in general; Clipperton Island, Galapagos Islands, Hawaiian Islands.
Ostracion lentiginosum Bloch and Schneider, Syst. Ichth., 1801, 501, Ile de France.
Ostracion punctatus Bloch and Schneider, loc. cit., 501, Honolulu.
- 4030. *Ostracion clippertonense*** Snodgrass and Heller.
 Clipperton Island.
Ostracion clippertonense Snodgrass and Heller, Shore Fishes, Galapagos Islands, in Proc. Wash. Acad. Sci., VI, 1904, 410, Clipperton Island.
- Genus 1442. RHINESOMUS** Swainson.
Rhinesomus Swainson, Nat. Hist. Classn. Fishes, II, 1839, 194, 324 (*Ostracion triqueter* Linnæus).
- 4031. *Rhinesomus triqueter*** (Linnæus). *Trunkfish; Rock shellfish; Drunken-fish; Chapin; Platefish.*
 West Indies, north to the Bermudas; Key West and Pensacola; Woods Hole, Mass.
Ostracion triqueter Linnæus, Syst. Nat. Ed. X, 1758, 330, "India."
Ostracion concatenatus Bloch, Naturgesch. Ausl. Fische, I, 1785, 101, Martinique.
- Genus 1443. CHAPINUS** Jordan and Evermann.
Chapinus Jordan and Evermann, Check List, 1895, 424 (*Ostracion bicaudalis* Linnæus).

⁹² Whether the West Indian *Alutera guntheriana* is distinguishable from *Alutera monoceros* of the Pacific is uncertain.

- 4032. *Chapinus bicaudalis* (Linnæus).** *Chapin; Spotted trunkfish; Pesce toro; Bullfish.*
West Indies, from Cuba to Ascension Island; Tortugas.
Ostracion bicaudalis Linnæus, Syst. Nat., Ed. X, 1758, 330, "India"; after Artedi.

- Genus 1444. *LACTOPHRYNS* Swainson.** *Three-angled trunkfishes.*
Lactophrys Swainson, Nat. Hist. Classn. Fishes, II, 1839, 194, 324 (*Ostracion trigonus* Linnæus, etc.).

- 4033. *Lactophrys trigonus* (Linnæus).** *Common trunkfish; Chapin; Shellfish; Cuckold; Boxfish.*
West Indies, north to Bermuda and Key West; occasionally northward in Gulf Stream to Holmes Hole, Mass., and Chesapeake Bay.
Ostracion trigonus Linnæus, Syst. Nat., Ed. X, 1758, 330, India.
Ostracion yalei Storer, Bost. Jour. Nat. Hist., I, 1836, 353, Holmes Hole and Marthas Vineyard.
Ostracion undulatus Poey, Synopsis, 1868, 441, Habana.
Ostracium expansum Cope, Trans. Amer. Phil. Soc., XIV, 1870, 474, figs. 9-10, locality unknown.

- Genus 1445. *ACANTHOSTRACION* Bleeker.**
Acanthostracion Bleeker, Atlas Ichth., V, 1865, 27, (*Ostracion quadricornis* Linnæus = *Ostracion tricornis* Linnæus).

- 4034. *Acanthostracion quadricornis* (Linnæus).** *Cuckold; Toro; Cowfish; Horned trunkfish.*
Tropical seas; common from Carolina to Brazil; Marquesas Islands; Florida Keys; occasionally at Pensacola and Galveston and north in the Gulf Stream to Charleston and Chesapeake Bay.
Ostracion tricornis Linnæus, Syst. Nat., Ed. X, 1758, 331, "India"; after Artedi.
Ostracion quadricornis Linnæus, loc. cit., 331, "India"; after Artedi, (this name preferred to the preceding by the first reviser).
Ostracion lister Lacépède, Hist. Nat. Poiss., I, 1798, 468; after Willoughby.
Ostracion sexcornutus Mitchell, Amer. Month. Mag., II, 1818, 328, mouth of Mississippi River.
Ostracion guineensis Bleeker, Ned Tydskr. Dierk., II, 1865?, 298, Guinea.
Ostracion gronovii Bleeker, loc. cit., II, 298, Guinea.
Acanthostracion polygonius Poey, Enumeratio, 1876, 175, Cuba.

Suborder GYMNODONTES

Family 299. TETRAODONTIDÆ. Puffers

- Genus 1446. *LAGOCEPHALUS* ⁹³ Swainson.**
Lagocephalus Swainson, Nat. Hist. Classn. Fishes, II, 1839, 194, 328 (*Tetraodon stellatus* Donovan).

- 4035. *Lagocephalus lavigatus* (Linnæus).** *Smooth puffer; Rabbitfish; Smooth swellfish; Tambor.*
Cape Cod to Brazil; rare north of Cape Hatteras.
Tetrondon lavigatus Linnæus, Syst. Nat., Ed. XII, 1766, 411, Charleston, S. C.
Tetrodon curvus Mitchell, Trans. Lit. Phil. Soc., I, 1815, 472, New York.
Tetrodon mathematicus Mitchell, loc. cit., I, 1815, 474, New York.
Holocanthus melanotus Gronow, Syst. Nat., ed. Gray, 1854, 24, Carolinas.
Tetrodon lineolatus Poey, Synopsis, 1868, 432, Cuba.

- 4036. *Lagocephalus pachycephalus* (Ranzani).** *Jugfish.*
Jamaica to Brazil.
Tetrondon pachycephalus Ranzani, Nov. Comm. Ac. Sci. Inst. Bonon., IV, 1840, 73, Pl. X, fig. 2, Brazil.

⁹³ The American species of *Lagocephalus* are very well separated from *Sphoeroides* but various Japanese species of the former genus grade off into the latter.

Genus 1447. SPHÉROÏDES Lacépède. *Swellfishes.*

Les Sphéroïdes Lacépède, Hist. Nat. Poiss., II, 1798, 22, Pl. I, fig. 1
(La Sphéroïde tuberculée, a front view of *Tetraodon spengleri* Bloch).

Sphéroïdes, Allegem. Lit. Zeit. Berlin, Sept. 24, 1798, unsigned review,
adopts this spelling as a technical name for the genus.

Sphéroïdes Duméril, Zool. Anal., 1806, 342; after Lacépède.

Orbodus Rafinesque, Analyse de la Nature, 1815, 90 (substitute for *Les Sphéroïdes* Lacépède).

Crayracion Klein, Gesellschaft Schauplatz, IV, 1777, 788 (*Tetraodon spengleri* Bloch; not binomial); after Klein, 1742.

Holocanthus Gronow, Cat. Fish., 1854 (*H. leionothus* Gronow).

Anchisomus (Kaup) Richardson, Zool. H. M. S. Herald, 1854, 156
(*Tetraodon spengleri* Bloch).

Crayracion Bleeker, Atlas Ichth., 1865, 65 (*Tetraodon spengleri* Bloch).

Stenometopus Bibron, Rev. Mag. Zool., VIII, 1855, 279 (*Tetraodon testudineus* Linnæus).

1

4037. **Sphéroïdes maculatus** (Bloch and Schneider). *Puffer; Swelltoad; Blower; Swellbelly; Swellfish; Toadfish.*

Atlantic coast of the United States from Cape Ann to St. Johns River and Biscayne Bay, Fla.

Tetraodon hispidus var. *maculatus* Bloch and Schneider, Syst. Ichth., 1801, 504, Rhode Island; after Schöpf.

Tetraodon turgidus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 473, pl. 5, fig. 5, New York.

4038. **Sphéroïdes spengleri** (Bloch). *Southern puffer; Swelltoad; Tambor.*

West Indies; coasts of Texas and Florida, south to Rio Janeiro and to the Madeiras and Canaries; rarely north to Woods Hole.

Tetraodon spengleri Bloch, Ichth., I, 1782, 135, pl. 144, East Indies.

Le *Tetraodon* Plumier Hist. Nat. Poiss., I, 1797, 504, Martinique.

Le *Sphéroïde tubercule* Lacépède, Hist. Nat. Poiss., II, 1798, 22, Martinique.

Sphéroïdes tuberculatus Pillot edition of Lacépède, VI, 1831, 279, Martinique.

4039. **Sphéroïdes nephelus** (Goode and Bean). *Puffer; Swellingfish.*

Florida.

Tetraodon nephelus Goode and Bean, Proc. U. S. Nat. Mus., V, 1882, 412, Indian River and Pensacola, Fla.

4040. **Sphéroïdes marmoratus** (Ranzani). *Spinyback blowfish; Swellfish.*

West Indies to Brazil; Jamaica.

Tetraodon marmoratus Ranzani, Nov. Comm. Ac. Sci. Inst. Bonon., IV, 1840, 72, pl. 10, fig. 1, Brazil.

4041. **Sphéroïdes harperi** Nichols.

West Indies (St. Eustatius) north to Florida.

Sphéroïdes harperi Nichols, Bull. Am. Mus. Nat. Hist., XXXIII, 1914, 81, Cape Sable, Fla.

4042. **Sphéroïdes asterias** (Blosser).

St. Croix.

Sphéroïdes asterias Blosser, Ann. Carnegie Mus., VI, 300, 1909, St. Croix.

4043. **Sphéroïdes eulepidotus** (Metzelaar).

Caribbean Sca.

Tetraodon (Sphéroïdes) eulepidotus Metzelaar, Trop. Atl. Visschen, 1919, 170, Curaçao.

4044. **Sphéroïdes lobatus** (Steindachner). *Bolete.*

Gulf of California to Galapagos Islands; Mazatlan; Panama.

Canthogaster? lobatus Steindachner, Ichth. Notizen, X, 1870, 18, pl. 5, fig. 3, Altata.

4045. **Sphéroïdes angusticeps** (Jenyns).

Pacific coast Lower California (La Paz), to the Galapagos Islands.

Tetraodon angusticeps Jenyns, Voyage Beagle, Fishes, 1842, 154, 28, Galapagos Islands.

Genus 1448. CHEILICHTHYS Müller.

Cheilichthys Müller, Abhandl. Akad. Wiss. Berl., 1839, 252 (*Tetraodon testudineus* Linnæus).
Apsicephalus Hollard, Ann. Sci. Nat., Zool., VIII, 1857, 275 (*Tetrodon fluviatilis* Hamilton).

4046. **Cheilichthys testudineus** (Linnæus). *Tambor; Glovefish; Blowfish; Swellfish; Puffer.*
 West Indies; Gulf Stream, occasionally to Newport, R. I.
Orbis lœvis variegatus Catesby, Nat. Hist. Carolinas, II, 1743, 28, pl. 28, Virginia.
Tetraodon testudineus Linnæus, Syst. Nat., Ed. X, 1758, 332, India; based on Balk and Artedi.
?Tetrodon punctatus Bloch and Schneider, Syst. Ichth., 1801, 506, Brazil.
Tetronotus geometricus Bloch and Schneider, loc. cit., 508, Virginia; after Catesby.
Tetraodon ammocryptus Gosse, Nat. Sojourn Jamaica, 1851, 287, Jamaica.
Holocanthus leionothos Gronow, Cat. Fishes, ed. Gray, 1854, 24, American oceans.

4047. **Cheilichthys annulatus** (Jenyns).

Pacific coast of tropical America; Galapagos Islands.
Tetraodon annulatus Jenyns, Zool. Beagle, 1842, 153, Galapagos Islands.
Anchisomus geometricus (Kaup) Richardson, Voyage Herald, 1854, 156, pl. 30, Galapagos Islands.
Tetronotus heraldi Günther, Cat., VIII, 1870, 283, Galapagos Islands and Panama.

4048. **Cheilichthys politus** (Girard).

Pacific coast of tropical America, San Diego to Mazatlan.
Tetraodon politus Girard, Pacific R. R. Surv., X, 1858, 340, San Diego, Calif.

4049. **Cheilichthys fürthi** (Steindachner).

Panama.
Tetraodon fürthii Steindachner, Ichth. Beitr., V, 1876, 22, Panama.

4050. **Cheilichthys trichocephalus** (Cope). *Hairy blowfish.*

Gulf Stream off Newport, R. I.; only the type known.
Tetraodon trichocephalus Cope, Proc. Acad. Nat. Sci. Phila., XXII, 1870, 120, Gulf Stream off Newport, R. I.

4051. **Cheilichthys pachygaster** (Müller and Troschel).

Barbados.
Tetraodon (*Cheilichthys*) *pachygaster* Müller and Troschel, in Schomburgk's Hist. Barbados, 1848, 677, Barbados.

Genus 1449. TETRAODON Linnæus.

Tetraodon Linnæus, Syst. Nat., Ed. X, 1758, 333 (*T. lineatus* Linnæus).
Les Ovoïdes Lacépède, Hist. Nat. Poiss., I, 1797, 256 (fascié, face view).
Ovoïdes Cuvier, Leçons Anat. Comp., I, 1800, tab. 1 (*Ovoïdes fasciatus* Lacépède = *Tetraodon stellatus* Linnæus).
Ovum Bloch and Schneider, Syst. Ichth., 1801, 530 (*O. commersoni* Bloch and Schneider = "l'Ovoïde Fascié" Lacépède, 1797); preoccupied in mollusks.
Oonidus Rafinesque, Anal. Nat., 1815, 90, for "les Ovoïdes" of Lacépède.
Arothron Müller, Abh. Akad. Wiss. Berlin, 1839, 252 (*A. testudinatus* Müller = *Tetraodon reticulatus* Bloch and Schneider).
Dilibomycter Bibron, Rev. Mag. Zool., VIII, 1855, 279 (*Tetraodon reticulatus* Bloch and Schneider).
Dichotomycter Bibron, loc. cit., 279 (*Tetraodon pluvialis* Hamilton).

- 4052.** *Tetraodon hispidus* Linnæus. *Maki-maki; Kete.*
 Panama and neighboring islands; Hawaii; Samoa; to China and the Red Sea.
Tetraodon hispidus Linnæus, Syst. Nat., Ed. X, 1758, 333, India.
Tetraodon perspicillaris Rüppell, Atlas, Reise Nord Africa, 1828, 63, Red Sea.
Tetrodon implusus Jenyns, Voy. *Beagle*, Fish., 1842, 152, Keeling Island.
Tetrodon laterna Richardson, Voy. *Sulphur*, Zool., 1844, 124, pl. 61, fig. 2, Canton, China.
Arothron erethizon^{**} Jordan and Gilbert, Proc. U. S. Nat. Mus., V, 1882, 631, Panama.

Subgenus OVOIDES Cuvier.

- 4053.** *Tetraodon setosus* Smith.
 West coast of Mexico; Revillagigedo, Clipperton, Cocos, and Galapagos Islands.
Tetraodon setosus Smith, Bull. Calif. Acad. Sci., 2d ser., VI, Nov. 13, 1886, Mexico.

Genus 1450. GUENTHERIDIA Gilbert and Starks.

Guentheridia Gilbert and Starks, Mem. Calif. Acad. Sci., IV, 1904, 158 (*Tetrodon formosus* Günther).

- 4054.** *Guentheridia formosa* (Günther).

Panama.
Tetrodon formosus Günther, Cat., VIII, 1870, 283, "South America."

Genus 1451. COLOMESUS Gill.

Colomesus Gill, Proc. U. S. Nat. Mus., VII, 1884, 422 (*Tetrodon psittacus* Bloch and Schneider).

- 4055.** *Colomesus psittacus* (Bloch and Schneider).

River mouths, Guiana and northern Brazil; Rio Branco, Essequibo, Marañon, and Capin at Para; West Indies in fresh waters only.
Tetrodon psittacus Bloch and Schneider, Syst. Ichth., 1801, 505, "Malabar."
Cheilichthys asellus Müller and Troschel, in Schomburgk, British Guiana, III, 1848, 641, Guiana.

Family 300. CANTHIGASTERIDÆ. Sharp-nosed puffers

Genus 1452. CANTHIGASTER Swainson.

Canthigaster Swainson, Nat. Hist. Fish., Amphib., etc., II, 1835, 194 (*Tetraodon rostratus* Bloch, called *Psilonotus* later in the text).
Psilonotus Swainson, loc. cit., 1835, 328, same type.
Prilonotus (Kaup) Müller, Richardson, Voyage *Herald*, 1854, 162 (*Tetraodon rostratus* Bloch).
Tropidichthys Bleeker, Nat. Tijdschr. Neder-Indie, VI, 501 (*T. marginatus* Bleeker).
Anosmius Peters, Monatsber. Akad. Wiss. Berlin, XXI, 1855, 274 (*T. tenuatus* Peters).
Eumycterias Jenkins, Bull. U. S. Fish Com., XIX, 1899, 399 (*E. biteniatus* Jenkins).

- 4056. Canthigaster rostratus** (Bloch).

West Indian fauna; Pensacola; Madeiras and Bermudas.
Tetrodon rostratus Bloch, Ichth., I, 1782, pl. 146, India.
Tetrodon capistratus Lowe, Proc. Zool. Soc. Lond., 1839, 90, Madeira.
Prilonotus (Anchisomus) caudacinctus Richardson, Voyage *Herald*, 1854, 162, pl. 30, figs. 1-3, locality unknown.
Tetrodon ornatus Poey, Synopsis, 1868, 433, Cuba.

- 4057. Canthigaster punctatissimus** (Günther).

Pacific coast of America, Gulf of California to Panama.
Tetrodon punctatissimus Günther, Cat., VIII, 1870, 302, Panama.
Tetrodon oxyrhynchus Lockington, Proc. Acad. Nat. Sci. Phila., XXXIII, 1881, 116, Gulf of California.

^{**} This American form shows no distinction from the widespread *Tetraodon hispidus*.

Family 301. DIODONTIDÆ

Genus 1453. DIODON Linnaeus. *Porcupinefishes.*

Diodon Linnaeus, Syst. Nat., Ed. X, 1758, 335 (*D. hystrix* Linnaeus).
Paradiodon Bleeker, Neder. Tijdschr. Dierk., II, 1865, 271 (*Diodon hystrix* Linnaeus).

- 4058.** *Diodon hystrix* Linnaeus. *Porcupinefish; Erizo; Puerco espino; Soursap.*
 Tropical seas, north to Lower California, Hawaii, and Florida.

Diodon hystrix Linnaeus, Syst. Nat., Ed. X, 1758, 335, India; after
 Artedi.

Guamajacu guara Marcgrave, Hist. Nat. Bras., 1648, 159, Brazil.

Erizo Parra, Desc. Dif. Piezas Hist. Nat. Cuba, 1787, 60, pl. 29, fig. 1,
 Habana.

Le Diodon plumier Lacépède, Hist. Nat. Poiss., II, 1798, 1, pl. 3, fig. 3,
 Martinique.

Diodon brachialis Bloch and Schneider, Syst. Ichth., 1801, 513, Cuba.

Diodon punctatus Cuvier, Mém. du Mus. d'Hist. Nat., IV, 1818, 132,
 no locality.

Diodon echinus Rafinesque, in Bonaparte, Cat. Met. Pisc. Eur., 1846,
 87, Mediterranean Sea.

- 4059.** *Diodon holacanthus* Linnaeus. *Balloonfish.*

All warm seas; north to Florida Keys, Lower California, and Hawaiian
 Islands.

Diodon holocanthus Linnaeus, Syst. Nat., ed. X, 1758, 335, India; based
 on Artedi (misprint).

Erizo guanabana Parra, Desc. Dif. Piezas Hist. Nat. Cuba, 1787, 62,
 pl. 29, fig. 2, Habana.

Le Diodon tacheté Lacépède, Hist. Nat. Poiss., II, 1798, 13, New Cy-
 therea.

Diodon liturosus Shaw, Gen. Zool., V, 1804, 436; after Lacépède.

Diodon spinosissimus Cuvier, Mém. Mus. Hist. Nat., IV, 1818, 134,
 pl. 7, no locality.

Diodon novem-maculatus Cuvier, loc. cit., 136, pl. 6, no locality.

Diodon sex-maculatus Cuvier, loc. cit., 136, pl. 7, no locality.

Diodon multimaculatus Cuvier, loc. cit., 137, no locality.

Diodon 4-maculatus Cuvier, loc. cit., 137, pl. 6, Otaiti.

Diodon maculatus var. a, Günther, Cat., VIII, 1870, 307, St. Croix;
 Jamaica; Panama; South America, etc.; after Lacépède.

Diodon melanopsis Kaup, Wiegmann's Archiv, 1855, 228.

- 4060.** *Diodon maculifer* Kaup.

Cape of Good Hope; recorded doubtfully from Cuba.

Diodon maculifer Kaup, Wiegmann's Archiv, 1855, 229, Cape of Good
 Hope.

Genus 1454. TRICHODIODON Bleeker.

Trichodiodon Bleeker, Atlas Ichth., Gymnodontes, V, 1867, 49 (*Diodon pilosus* Mitchell).

Trichocyclus Günther, Cat., VIII, 316 (*T. erinaceus* Günther).

- 4061.** *Trichodiodon pilosus* (Mitchill). *Balloonfish; Hairy porcupinefish;*
Hairy boxfish.

New York.

Diodon pilosus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 471;
 supposed to be from New York Harbor.

Diodon asper Cuvier, Mém. Mus. Hist. Nat., IV, 1818, 138; no locality.
 ?*Trichocyclus erinaceus* Günther, Cat., VII, 1870, 316; no locality.

Genus 1455. CYCLICHTHYS Kaup. *Swelltoads.*

Cyclichthys Kaup, Wiegmann's Archiv, XXI, 1855, 231 (*Diodon orbicu-*
latus Bloch).

- 4062.** *Cyclichthys schœpfi* (Walbaum). *Burrfish; Cucumberfish; Spotted balloonfish; Balloonfish; Rabbitfish; Swelltoad; Spiny boxfish.*
Cape Cod to West Indies and occasionally northward to Woods Hole,
Mass.
Diodon schœpfi Walbaum, Artedi Pisc., 1792, 601, Long Island; after
Schöpf.
?*Diodon meulenii* Walbaum, loc. cit., 602; no locality.
Diodon geometricus lineatus Bloch and Schneider, Syst. Ichth., 1801,
513, New York.
Diodon maculato striatus Mitchell, Fish. N. Y., 470, pl. 56, fig. 3, 814,
New York.
Diodon rivulatus Cuvier, Mém. Mus. Hist. Nat., IV, 1818, 129, pl. 6,
New York.
Diodon nigrolineatus Ayres, Proc. Bost. Soc. Nat. Hist., I, 1842, 68,
Brookhaven, Long Island.
Diodon verrucosus DeKay, N. Y. Fauna: Fishes, 324, 1842, pl. 56,
fig. 184, New York.
- 4063.** *Cyclichthys spinosus* (Linnæus).
West Indies and coast of Brazil.
Guamaicu atinga Marcgrave, Hist. Brasil., 1648, 168, Brazil.
Diodon spinosus Linnæus, Syst. Nat., Ed. X, 1758, 335, India.
Le Diodon orbe Lacépède, Hist. Nat. Poiss., II, 1798, 16, Rio Janeiro.
Diodon geometricus Bloch and Schneider, Ichth., 1801, 513, pl. 96,
America.
Cyclichthys cornutus Kaup, Wiegmann's Archiv, 1855, 231, Bahia.
- 4064.** *Cyclichthys fuliginosus* (DeKay). *Black burrfish; Unspotted balloon-fish.*
Atlantic coast from Cape Cod southward; not common.
Diodon fuliginosus DeKay, N. Y. Fauna: Fishes, 1842, 324, pl. 55,
fig. 181, New York Harbor.
- 4065.** *Cyclichthys antillarum* (Jordan and Rutter).
Cuba and Jamaica.
Chilomycterus antillarum Jordan and Rutter, Proc. Acad. Nat. Sci.
Phila., XLIX, 1897, 131, Kingston, Jamaica.
- 4066.** *Cyclichthys briareos* (Metzelaar).
Lesser Antilles.
Chilomycterus briareos Metzelaar, Trop. Atl. Vissch., 1919, 173, fig. 55,
St. Eustatius.
- 4067.** *Cyclichthys antennatus* (Cuvier).
West Indies and southward; St. Croix; Jamaica; Porto Rico; Cape of
Good Hope.
Diodon antennatus Cuvier, Mém. Mus. Hist. Nat., IV, 1818, 131; no
locality.
?*Chilomycterus puncticulatus* Poey, Anal. Soc. Española Hist. Nat.,
X, 1881, 346, Porto Rico.
- Genus 1456. CHILOMYCTERUS Bibron.** *Burrfishes.*
Chilomycterus Bibron, in Barneville, Rev. Zool., 1846, 40 (*Diodon reticulatus* Linnæus = *Diodon atinga* Linnæus).
Cyanichthys Kaup, Wiegmann's Archiv, IV, 1855, 231 (*C. cærulæus*
Kaup; not of Quoy and Gaimard).
Diodon Bleeker, Atlas Ind. Gymnod., 1865, 55 (*D. atinga*); not of
authors, who, after Kaup, restrict *Diodon* to *D. hystrix*.
- Subgenus CHILOMYCTERUS Bibron.**
- 4068.** *Chilomycterus atinga* (Linnæus). *Atinga.*
West Indies to Bermuda and Florida Keys.
Diodon atinga (misprinted *atinga*) Linnæus, Syst. Nat., Ed. X, 1758,
334, India.
Diodon reticulatus Linnæus, loc. cit., 334, India; after Artedi.
- 4069.** *Chilomycterus affinis* Günther.
Galapagos; coast of southern California; Hawaii, Japan; scarce.
Chilomycterus affinis Günther, Cat., VIII, 1870, 314, no locality; California?
Chilomycterus californiensis Eigenmann, Amer. Nat., V, 1891, 1133,
San Pedro, Calif.

Genus 1457. LYOSPHÆRA Evermann and Kendall.

Lyosphæra Evermann and Kendall, Bull. U. S. Fish Com., XVII, 1897, 131 (*L. globosa* Evermann and Kendall).

4070. **Lyosphæra globosa** Evermann and Kendall. *Marblefish.*

Atlantic coast of the United States; known from the mouth of the Rappahannock River and from Biscayne Bay, Fla.

Lyosphæra globosa Evermann and Kendall, Bull. U. S. Fish Com., XVII, 1897, 131, pl. 9, figs. 11 and 12, Rappahannock River near mouth of Windmill Creek, Va.

Family 302. MOLIDÆ. Headfishes**Genus 1458. MOLA Kœlreuter. Headfishes.**

Mola Kœlreuter, Nova Comm. Act. Petropol., VII, 1770, 337 (*Diodon mola* Kœlreuter = *Diodon mola* Linnaeus).

Mola Cuvier, Tableau Elem. Hist. Nat. Anim., 1798, 323 (*M. rotunda* Cuvier = *Tetrodon mola* Linnaeus).

Orthogoriscus Schneider, ed. Bloch, Syst. Ichth., 1801, 510 (*Diodon mola* Bloch).

Cephalus Shaw, Gen. Zool., V, 1804, 432 (*Diodon mola* Linnaeus).

Orthragus Rafinesque, Caratteri, 1810, 17 (*Tetrodon mola* Linnaeus).

Diplanchias Rafinesque, loc. cit., 17 (*D. nasus* Rafinesque).

Typanomomium Ranzani, Novi Comm. Acad. Sci. Inst. Bonon., III, 1839, 81 (*T. planci* Ranzani).

Ozadura Ranzani, loc. cit., 81 (*O. orsini* Ranzani = *Mola mola* Linnaeus).

Trematopsis Ranzani, loc. cit., 81 (*T. willughbei* Ranzani).

Pedalion Swainson, Nat. Hist. Fish. Amphib. etc., I, 1838, 199, II, 195, 329 (*Tetrodon mola* Linnaeus).

Molacanthus Swainson, loc. cit., II, 1839, 329 (*M. pallasi* Swainson = *Diodon mola* Pallas, young).

Pallasio Nardo, Ann. Sci. Lombardo-Veneto, X, 1839, V, 10, 112 (*P. pallasi* Nardo).

Acanthosoma DeKay, Nat. Hist. N. Y., 1842, 350 (*A. carinatum* DeKay).

Centaurus Kaup, Wiegemann's Archiv Naturg., XXI, 1855, 221 (*Ostracion boops* Richardson; larval).

Aledon Castelnau, Mém. Poiss., Afr. Aust., VII, 1861, 75 (*A. storeri* Castelnau).

4071. **Mola mola** (Linnaeus). *Ocean sunfish; Globefish; Sunfish; Headfish; Mola; Pez luna; Pez mola.*

Temperate and tropical seas, northward to England, Cape Cod, and San Francisco; rare in the West Indies.

Tetrodon mola Linnaeus, Syst. Nat., Ed. X, 1758, 334, 412, Mediterranean.

Diodon carinatus Mitchell, Ann. Lyc. Nat. Hist. New York, II, 1815, 264, pl. 5, fig. 1, New York.

Ozadura orsini Ranzani, Nov. Comm. Ac. Sci. Inst. Bonon., III, 1839, 82, Mediterranean Sea.

Typanomomium planci Ranzani, loc. cit., table, Adriatic Sea.

Diplanchias nasus Ranzani, loc. cit., table, "in marei siculo" (Sicily).

Trematopsis willughbei Ranzani, loc. cit., table, "in oceano."

Orthragoriscus retzii Ranzani, loc. cit., table, no locality.

Orthragoriscus ghini Ranzani, loc. cit., table, Mediterranean Sea.

Orthragoriscus rondeletii Ranzani, loc. cit., Mediterranean Sea.

Orthragoriscus blochii Ranzani, loc. cit., "in mari oceano."

Orthragoriscus alexandrini Ranzani, loc. cit., Adriatic Sea.

Orthragoriscus redi Ranzani, loc. cit., Mediterranean Sea.

Orthragoriscus oculeatus Ranzani, loc. cit., no locality.

Orthragoriscus elegans Ranzani, loc. cit., Atlantic Ocean.

Orthragoriscus battaræ Ranzani, loc. cit., Adriatic Sea.

Orthragoriscus lunaris Gronow, Cat. Fishes, ed. Gray, 1854, 165, Mediterranean.

Orthragoriscus solaris Gronow, loc. cit., 165, Mediterranean.

Orthragoriscus analis Ayres, Proc. Calif. Acad. Nat. Sci., II, 1859, 31, fig. 54, San Francisco.

Ostracion boops Richardson, Voy, Erebus and Terror, Ichth., 1844, 52, South Atlantic (larva).

Genus 1459. RANZANIA Nardo. *King of the mackerels.*
Ranzania Nardo, Ann. Sci. Regn. Lombard. Venet., X, 1840, 105
 (*Tetrodon truncatus* Retzius).

4072. *Ranzania truncata* (Retzius).
 Pelagic; occasionally on the Atlantic coast; Bermudas; Martinique.
Tetrodon truncatus Retzius, Vet. Ak. Nya. Handl., VI, 1785, 2, 121,
 Brazil.

4073. *Ranzania makua* Jenkins.
 Open Pacific, Hawaii to Japan; very rare, once found on Pismo Beach
 near Oceano, San Luis Obispo County, Calif.
Ranzania makua Jenkins, Proc. Calif. Acad. Sci., 2d ser., V, 1895, 779,
 with colored plate (frontispiece), Honolulu; probably not different
 from *Ranzania truncata*; see Snyder, Proc. U. S. Nat. Mus., XLIV,
 1913, 455.

Order AAA. PEDICULATI

Family 303. LOPHIIDÆ. Fishing frogs

Genus 1460. LOPHIUS Linnæus.

Lophius Linnæus Syst. Nat., Ed. X, 1758, 236 (*Lophius piscatorius*
 Linnæus); after Artedi.

4074. *Lophius piscatorius* Linnæus. *Fishing frog; Monkfish; Goosefish;*
Angler; Allmouth; Bellowsfish.
 North Atlantic, on both coasts.
Lophius piscatorius Linnæus, Syst. Nat., Ed. XII, 1766, 402, Seas of
 Europe.
Lophius americanus Cuvier and Valenciennes, Hist. Nat. Poiss., XII,
 1837, 380, Philadelphia.

Genus 1461. LOPHIOMUS Gill.

Lophiomus Gill, Proc. U. S. Nat. Mus., V, 1882, 552 (*Lophius setigerus*
 Vahl.)

4075. *Lophiomus setigerus* (Vahl).
 Pacific Ocean; not uncommon in rather deep water off coasts of China
 and Japan; one specimen from *Albatross* station 2805, southwest of
 Panama.
Lophius setigerus Vahl, Skr. Naturh., IV, 1797, 214, tab. 3, figs. 5
 and 6, China Sea.
Lophius viviparus Bloch and Schneider, Syst. Ichth., 142, 1801, tab.
 32, China Sea; after Vahl.

Family 304. ANTENNARIDÆ. Frogfishes

Genus 1462. HISTRIO Fischer.

Histrio[“] Fischer, Zoog., Tab. Syn. Illust., Ed. III, I, 1813 (*Lophius*
histrio Linnæus).

Pterophryne Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 90 (*Chironectes*
lævigatus Cuvier; name preoccupied).

Pterophrynidoides Gill, Proc. U. S. Nat. Mus., I, 1878, 216 (*Chironectes*
lævigatus Cuvier); substitute for *Pterophryne*.

4076. *Histro pictus* (Cuvier and Valenciennes). *Mousefish; Sargassumfish.*
 Tropical parts of the Atlantic; abundant on our Gulf coast and occa-
 sional to Cape Hatteras and beyond.
Lophius tumidus Osbeck, Iter Chinensis, 1757, 400, open sea; pre-
 Linnaean.
Chironectes pictus Cuvier and Valenciennes, Hist. Nat. Poiss., XII,
 1837, 393, Surinam.
Chironectes tumidus Cuvier and Valenciennes, loc. cit., 397, Sargasso Sea.
Chironectes arcticus Dübén and Koren, K. Vet. Akad. Stockholm, 1844,
 72, Vadsö, Norway, carried northward on the Gulf Stream.

[“] Name questionable on account of error in the published diagnosis, the description interchanged with
 that of *Lophius*.

4077. *Histrio gibba* (Mitchill).

West Indies, north to Key West and the Tortugas.

Lophius gibbus Mitchell, Trans. Lit. Phil. Soc. N. Y., I, 1815, 1, pl. 4, fig. 9, off St. Croix, lat. 22° N., long. 64° W.
Chironectes levigatus Cuvier, Mém. Mus., III, 1817, 423, pl. 16, fig. 1; South Carolina.

4078. *Histrio jagua* Nichols.

Bermuda.

Histrio jagua Nichols, Proc. Biol. Soc. Wash., XXXIII, 1920, 62, Bermuda.

Genus 1463. ANTENNARIUS ^{**} Lacépède.

Antennarius Lacépède, Hist. Nat. Poiss., I, 1798, 421 (*Antennarius biverplex totus ater puncto mediorum, laterum albo* Commerson (in foot-note) = *Lophius commersonianus* Lacépède).

Chironectes Cuvier, Mém. Mus., III, 1817, 418 (*Lophius commersonianus* Lacépède).

Capellaria Gistel, Thierreich, VIII, 1848 (*Lophius commersonianus* Lacépède).

4079. *Antennarius scaber* (Cuvier).

Caribbean Sea; Port Castries; St. Lucia.

Chironectes scaber Cuvier, Mém. Mus., III, 1817, 425, pl. 16, fig. 2, Martinique.

Lophius spectrum Gronow, Cat. Fish., ed. Gray, 1854, 49, Antilles; after *Lophius acute scabra* Gronow, Zoophyl., 1781, 210.

4080. *Antennarius inops* Poey.

Porto Rico.

Antennarius inops Poey, Anal. Soc. Esp. Hist. Nat., X, 1881, 340 Porto Rico.

4081. *Antennarius principis* (Cuvier and Valenciennes).

West Indies to Brazil.

Chironectes principis Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 416, Brazil; on a drawing by Prince Maurice.

?*Chironectes mentzeli* Cuvier and Valenciennes, loc. cit., 416, Brazil; on a drawing.

4082. *Antennarius tenebrosus* (Poey).

Cuba.

Chironectes tenebrosus Poey, Memorias, I, 1853, 219, pl. 17, fig. 1, Cuba.

4083. *Antennarius reticularis* Gilbert.

Gulf of California; only the type known.

Antennarius reticularis Gilbert, Proc. U. S. Nat. Mus., XIV, 1891, 566; Gulf of California at Albatross station 2825.

4084. *Antennarius strigatus* Gill.

Cape San Lucas; Panama.

Antennarius strigatus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 92; Cape San Lucas.

Antennarius tenuifilis Günther, Fish. Cent. Amer., 1869, 440, Panama; young.

4085. *Antennarius sanguineus* Gill.

Pacific coast of tropical America, Cape San Lucas to Panama; scarce. *Antennarius sanguineus* Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 91; Cape San Lucas.

Antennarius leopardinus Günther, Proc. Zool. Soc. Lond., 1864, 151; Panama.

4086. *Antennarius ocellatus* (Bloch and Schneider).

West Indies, north to Florida.

Pescador Parra, Dif. Piezas, Hist. Nat., 1780, 1, tab. 1, Cuba.

Lophius histrio, var. *ocellatus* Bloch and Schneider, Syst. Ichth., 142, 180; after Parra.

Antennarius pleurophthalmus Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 92; Key West.

^{**} These fishes vary considerably in color and more or less in form. It is likely that the number of species should be reduced.

4087. **Antennarius tigris** (Poey).
Cuba.
Chironectes tigris Poey, *Memorias*, I, 1853, 217, pl. 17, fig. 2; Cuba.
4088. **Antennarius nuttingii** Garman.
Great Bahama Banks.
? *Chironectes mentzelii* Cuvier and Valenciennes, *Hist. Nat. Poiss.*, XII, 1837, 417; Brazil; on a drawing by Prince Maurice.
Antennarius nuttingii Garman, *Bull. Iowa Lab. Nat. Hist.*, 1896, 83, Pl. II; Great Bahama Banks.
4089. **Antennarius multiocellatus** (Cuvier and Valenciennes). *Martin pescador*.
Garden Key, Fla., to Martinique.
Chironectes multiocellatus Cuvier and Valenciennes, *Hist. Nat. Poiss.*, XI, 1837, 422, Martinique.
Antennarius annulatus Gill, *Proc. Acad. Nat. Sci. Phila.*, XV, 1863, 91, Garden Key, Fla.
Antennarius corallinus Poey, *Repertorio*, I, 1865, 188, Cuba.
4090. **Antennarius radiosus** Garman.
Key West.
Antennarius radiosus Garman, *Bull. Lab. Nat. Hist. Iowa Univ.*, 1896, 85, pl. 1, off Key West in 50 fathoms.
4091. **Antennarius sonntagii** (Baron Müller).
Gulf of Mexico.
Chironectes sonntagii Baron J. W. von Müller, *Reisen in den Verein. Staat., etc.*, Band I, 1864, 180, in floating seaweed, no exact locality stated.
4092. **Antennarius avalonis** Jordan and Starks.
Santa Barbara Islands.
Antennarius avalonis Jordan and Starks, *Proc. U. S. Nat. Mus.*, XXXII, 1907, 77, fig. 8, Santa Catalina Island at Avalon, Calif.
4093. **Antennarius verrucosus** Bean.
Bermudas.
Antennarius verrucosus Bean, *Proc. Biol. Soc. Wash.*, XIX, 1906, 31, St. George Island, Bermuda Islands.
4094. **Antennarius astroscopus** Nichols.
Barbados.
Antennarius astroscopus Nichols, *Bull. Amer. Mus. Nat. Hist.*, XXXI, 1912, 109, Barbados.

Family 305. CHAUNACIDÆ

- Genus 1464. CHAUNAX** Lowe.
Chaunax Lowe, *Proc. Zool. Soc. Lond.*, XIV, 1846, 339 (*C. pictus* Lowe).
4095. **Chaunax pictus** Lowe.
Madeira; taken off the coast of Rhode Island in the Gulf Stream.
Chaunax pictus Lowe, *Trans. Zool. Soc. Lond.*, 1846, 339, *Camera de Lobos*, Madeira.
4096. **Chaunax nuttingi** Garman.
Florida Keys.
Chaunax nuttingi Garman, *Bull. Nat. Hist. Iowa Univ.*, 1896, 86, Sand Key Light, Fla.

Family 306. OGCOCEPHALIDÆ. Batfishes

- Genus 1465. OGCOCEPHALUS** Fischer. *Seabats*.
Ogcocephalus Fischer, *Zoognosia*, ed. 3, I, 1813, 70, 78 (*Lophius vespertilio* Linnaeus).
Oncocephalus Gill, corrected spelling.
Malthe Cuvier, *Règne Anim.*, ed. 1, II, 1817, 311 (*Lophius vespertilio* Linnaeus).
Malthæa of authors, modified spelling.

4097. *Ogcocephalus vespertilio* (Linnaeus). *Batfish; Diablo.*
 West Indies, north to Carolina.
Lophius vespertilio Linnaeus, Syst. Nat., Ed. X, I, 1758, 236, American seas; after *Lophius fronti unicorni* of Artedi.
4098. *Ogcocephalus nasutus* (Cuvier and Valenciennes).
 West Indies.
Malthea nasuta Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 452, Martinique.
Malthea notata Cuvier and Valenciennes, loc. cit., 453, Surinam; snout 15 in length; body spotted.
4099. *Ogcocephalus radiatus* (Mitchill).
 Coast of Florida and neighboring waters; very common in shallow bays among weeds, especially about the Florida keys.
Lophius radiatus Mitchill, Amer. Month. Mag., March, 1818, 326, Straits of the Bahamas.
Malthe cubifrons Richardson, Fauna Bor.-Amer., III, 1836, 103; said to be from Labrador, which is certainly an error.
? *Malthea truncata* Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 454, America; snout wholly obsolete; perhaps a species of *Zalieutes*.
- Genus 1466. **ZALIEUTES** Jordan and Evermann.
Zalieutes Jordan and Evermann, Check List, 1895, 511 (*Malthe elater* Jordan and Gilbert).
4100. *Zalieutes elater* (Jordan and Gilbert).
 Pacific coast of Mexico, south to Panama, in water of moderate depth; very rare near the shore but obtained in abundance by the *Albatross* at stations 2794 and 2795 near Panama.
Malthe elater Jordan and Gilbert, Proc. U. S. Nat. Mus., IV, 1881, 365, Mazatlan.
- Genus 1467. **MALTHOPSIS** Alcock.
Malthopsis Alcock, Deep Sea Fish. Invest., Ann. Mag. Nat. Hist., VIII, 1891, 26 (*M. luteus* Alcock).
4101. *Malthopsis spinosa* Garman.
 Off Bay of Panama in deep water.
Malthopsis spinosa Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 104, Pl. XXII, *Albatross* stations 3392 and 3393, off Panama, in 1,020 to 1,270 fathoms.
4102. *Malthopsis spinulosa* Garman.
 Off Bay of Panama, in deep water.
Malthopsis spinulosa Garman, Mem. Mus. Comp. Zool., XXIV, 1899, 106, Pls. XXI, XXVI, *Albatross* station 3394, lat. $7^{\circ} 21' N.$, long. $79^{\circ} 35' W.$ in 511 fathoms.
- Genus 1468. **HALIEUTICHTHYS** Poey.
Halieutichthys Poey, Proc. Acad. Nat. Sci. Phila., XV, 1863, 90 (*H. reticulatus* Poey = *Lophius aculeatus* Mitchill).
4103. *Halieutichthys aculeatus* (Mitchill).
 Cuba to southern Florida.
Lophius aculeatus Mitchill, Amer. Month. Mag., II, 1818, 325, Straits of Bahamas.
Halieutichthys reticulatus Poey, in Gill, Proc. Acad. Nat. Sci. Phila., XV, 1863, 91, Cuba.
4104. *Halieutichthys caribbaeus* Garman.
 Jamaica to Barbados, in depths of 70 to 150 fathoms or more.
Halieutichthys caribbaeus Garman, Bull. Lab. Nat. Hist. Iowa Univ., 1896, 87, pl. 4, fig. 2, Jamaica to Barbados.
4105. *Halieutichthys smithii* Evermann and Marsh.
 Porto Rico.
Halieutichthys smithii Evermann and Marsh, Bull. U. S. Fish Com., XX, 1900, 339, fig. 112, Mayaguez Harbor, Porto Rico.
- Genus 1469. **HALIEUTÆA** Cuvier and Valenciennes.
Halieutæa Cuvier and Valenciennes, Hist. Nat. Poiss., XII, 1837, 455 (*H. stellata* Cuvier and Valenciennes).

4106. ***Halieutæa spongiosa*** Gilbert.

Pacific Ocean, off Lower California.

Halieutæa spongiosa Gilbert, Proc. U. S. Nat. Mus., XIII, 1890, 124,
west of Revillagigedo Islands, at Albatross station 2992, lat. $18^{\circ} 17' 30''$ N., long. $114^{\circ} 43' 15''$ W., in 460 fathoms.

Genus 1470. **HALIEUTELLA** Goode and Bean.

Halieutella Goode and Bean, Proc. Biol. Soc. Wash., II, 1885, 88 (*H. lappa* Goode and Bean).

4107. ***Halieutella lappa*** Goode and Bean.

Gulf Stream.

Halieutella lappa Goode and Bean, Proc. Biol. Soc. Wash., II, 1882, 88,
Gulf Stream at Fish Hawk station 1151, lat. $39^{\circ} 58' 30''$ N., long. $70^{\circ} 37'$ W., in 125 fathoms.

Genus 1471. **DIBRANCHUS** Peters.

Dibranchus Peters, Monatsber. Akad. Wiss. Berlin, 1875, 736 (*D. atlanticus* Peters).

4108. ***Dibranchus atlanticus*** Peters.

Gulf Stream.

Dibranchus atlanticus Peters, Monatsber. K. Akad. Wiss. Berlin, 1875,
736, with plate; West Africa, lat. 10° N. long. 17° W., in 360 fathoms.

Family 307. CERATHIDÆ. Sea devils

Genus 1472. **CERATIAS** Krøyer.

Ceratias Krøyer, Naturhist. Tidsskr., I, 1845, 639 (*C. hollboelli* Krøyer).

4109. ***Ceratias hollboelli*** Krøyer.

Greenland to Nova Scotia; four specimens known.

Ceratias hollboelli Krøyer, Naturh. Tidsskr., 1844, 639, Greenland.

Genus 1473. **MANCALIAS** Gill.

Mancalias Gill, Proc. U. S. Nat. Mus., I, 1878, 227 (*Ceratias uranoscopus* Murray).

Typhlosaras Gill, Forest and Stream, 1883, Nov. 8, 284 (*T. shufeldti* Gill).

4110. ***Mancalias uranoscopus*** (Murray).

Deep seas; taken at Madeira and off the coast of southern New England.

Ceratius uranoscopus Murray, in Wyville Thompson, The Atlantic, II, 1878, 67, off Madeira.

4111. ***Mancalias shufeldti*** (Gill).

Western Atlantic.

Typhlosaras shufeldti Gill, Forest and Stream, November 8, 1883, 284, western Atlantic.

Genus 1474. **CRYPTOSARAS** Gill.

Cryptosaras Gill, Forest and Stream, Nov. 8, 1883 (*C. couesi* Gill).

4112. ***Cryptosaras couesii*** Gill.

Gulf Stream.

Cryptosaras couesii Gill, Forest and Stream, Nov. 8, 1883, 284, Gulf Stream off New England.

Genus 1475. **EDRIOLYCHNUS** Regan.

Edriolychnus Regan, Proc. Roy. Soc. Lond., ser. B, XCVII, 1925, 398 (*E. schmidti* Regan).

4113. ***Edriolychnus schmidti*** Regan.

Deep waters of the western Atlantic.

Edriolychnus schmidti Regan, Proc. Royal Soc. Lond., ser. B, XCVII, 1925, 398, western Atlantic; *Dana* expedition.

Genus 1476. **MONOCERATIAS** Gilbert.

Monoceratias Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 379 (*M. acanthias* Gilbert).

4114. ***Monoceratias acanthias*** Gilbert.

Off Santa Barbara Islands, Calif.

Monoceratias acanthias Gilbert, Proc. U. S. Nat. Mus., XLVIII, 1915, 379, pl. 22, fig. 24, off Santa Cruz Island, 764 to 891 fathoms.

Genus 1477. **BOROPHRYNE** Regan.
Borophryne Regan, Ann. Mag. Nat. Hist., ser. 9, XV, 1925, 564 (*B. apogon* Regan).

4115. **Borophryne apogon** Regan.
 Gulf of Panama.
Borophryne apogon Regan, Ann. Mag. Nat. Hist., ser. 9., XV, 1925, 564, Gulf of Panama.

Genus 1478. **PHOTOCORYNUS** Regan.
Photocorynus Regan, Proc. Roy. Soc. Lond., ser. B, XCVII, 1925, 393
 (*P. spiniceps* Regan).

4116. **Photocorynus spiniceps** Regan.
 Deep waters of Panama.
Photocorynus spiniceps Regan, Proc. Roy. Soc. Lond., ser. B, XCVII, 1925, 393, figs. 6-7, deep waters off Panama.

Family 308. HIMANTOLOPHIDÆ

Genus 1479. **ONEIRODES** Lütken.
Oneirodes Lütken, Overs. Dansk. Vid. Selsk. Kjøbenhavn, 1871, 56
 (*O. eschrichtii* Lütken).

4117. **Oneirodes eschrichtii** Lütken.
 Deep sea off Greenland.
Oneirodes eschrichtii Lütken, Overs. Dansk. Vid. Selsk. Forhandl., 1871, 56, 9-18, pl. 2, deep sea off Greenland.

Genus 1480. **HIMANTOLOPHUS** Reinhardt.
Himantolophus Reinhardt, Dansk. Vid. Selsk. Afh. Kjøbenhavn, VII, 1838, 74 (*H. grænlandicus* Reinhardt).

4118. **Himantolophus grænlandicus** Reinhardt.
 Greenland.
Himantolophus grænlandicus Reinhardt, Dansk. Vid. Selsk. Nat. Math. Afh., 1837, 74, Greenland.

Genus 1481. **LOPHODOLUS** Lloyd.
Lophodolus Lloyd, Mem. Indian Mus., II, 1909, 167 (*L. indicus* Lloyd).

4119. **Lophodolus acanthognathus** Regan.
 North Atlantic.
Lophodolus acanthognathus Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 563, north Atlantic.

Genus 1482. **CORYNOLOPHUS** Gill.
Corynolophus Gill, Proc. U. S. Nat. Mus., I, 1878, 219 (*Himantolophus reinhardtii* Lütken).

4120. **Corynolophus reinhardtii** (Lütken).
 Greenland; one specimen known.
Himantolophus reinhardtii Lütken, König. Dansk. Vid. Selsk., 1878, 321, Greenland. Perhaps a species of *Himantolophus*.

Genus 1483. **DOLOPICHTHYS** Garman.
Dolopichthys Garman in Mem. Mus. Comp. Zool., XXIV, 1899, 81,
 (*D. allector* Garman).

4121. **Dolopichthys gracilispinis** Regan.
 North Atlantic.
Dolopichthys gracilispinis Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 563, north Atlantic.

4122. **Dolopichthys microlophus** Regan.
 North Atlantic.
Dolopichthys microlophus Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 563, north Atlantic.

4123. **Dolopichthys anisacanthus** Regan.
 North Atlantic.
Dolopichthys anisacanthus Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May 1925, 562, north Atlantic.

4124. **Dolopichthys luetkeni** Regan.
Gulf of Panama.
Dolopichthys luetkeni Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 562, Gulf of Panama.
4125. **Dolopichthys heteracanthus** Regan.
Gulf of Panama.
Dolopichthys heteracanthus Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, Gulf of Panama.
- Genus 1484. LASIOGNATHUS** Regan.
Lasiognathus Regan, Ann. Mag. Nat. Hist., 9 ser., XV, 1925, 563 (*L. saccostoma* Regan).
4126. **Lasiognathus saccostoma** Regan.
Caribbean Sea.
Lasiognathus saccostoma Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 563, Caribbean Sea.
- Genus 1485. LINOPHYRYNE** Collett.
Linophryne Collett, Proc. Zool. Soc. Lond., 1886, 138 (*L. lucifer* Collett).
4127. **Linophryne lucifer** Collett.
Open Atlantic.
Linophryne lucifer Collett, Proc. Zool. Soc. Lond., 1886, 138, pl. 15, off Madeira, lat. 36° N., long. 20° W.
4128. **Linophryne polypogon** Regan.
North Atlantic.
Linophryne polypogon Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 565, north Atlantic.
4129. **Linophryne arborifer** Regan.
North Atlantic.
Linophryne arborifer Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 564, north Atlantic.
4130. **Linophryne macrodon** Regan.
Gulf of Panama.
Linophryne macrodon Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1895, 564, Gulf of Panama.
- Genus 1486. CHÆNOPHYRYNE** Regan.
Chænophryne Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 564 (*C. longiceps* Regan).
4131. **Chænophryne longiceps** Regan.
North Atlantic and Gulf of Panama.
Chænophryne longiceps Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 564, north Atlantic.

Family 309. MELANOCETIDÆ

- Genus 1487. MELANOCECUS** Günther.
Melanocetus Günther, Proc. Zool. Soc. Lond., 1864, 301 (*M. johnsoni* Günther).
4132. **Melanocetus polyactis** Regan.
Gulf of Panama.
Melanocetus polyactis Regan, Ann. Mag. Nat. Hist., 9 ser., XV, May, 1925, 565, Gulf of Panama.
4133. **Melanocetus niger** Regan.
Gulf of Panama.
Melanocetus niger Regan, Ann. Mag. Nat. Hist., ser. 9, XV, May, 1925, 565, Gulf of Panama.
- Genus 1488. LYOCETUS** Günther.
Lyocetus Günther, Challenger Rept., XXII, 1887, 57 (*L. murrayi* Günther).

4134. *Lyocetus murrayi* (Günther).

Mid-Atlantic at *Challenger* stations 106 and 348.

Melanocetus (Liocetus) murrayi Günther, *Challenger Rept.*, XXII, 1887,
57, pl. 11, fig. A, mid-Atlantic.

Family 310. CAULOPHRYNIDÆ**Genus 1489. CAULOPHRYNE Goode and Bean.**

Caulophryne Goode and Bean, *Oceanic Ichth.*, 1895, 496 (*Caulophryne jordani* Goode and Bean).

4135. *Caulophryne jordani* Goode and Bean.

Gulf Stream.

Caulophryne jordani Goode and Bean, *Oceanic Ichth.*, 1895, 496, pl. 21,
fig. 409, Gulf Stream off Carolinas in 1,276 fathoms.

Family 311. ACERATHIIDÆ**Genus 1490. RHYNCHOCERATIAS Regan.**

Rhynchoceratias Regan, *Ann. Mag. Nat. Hist.*, 9 ser., XV, May, 1925,
566 (*R. brevirostris* Regan).

4136. *Rhynchoceratias leucorhinus* Regan.

North Atlantic and Gulf of Panama.

Rhynchoceratias leucorhinus Regan, *Ann. Mag. Nat. Hist.*, 9 ser., XV,
May, 1925, 566, north Atlantic.

4137. *Rhynchoceratias rostratus* Regan.

Gulf of Panama.

Rhynchoceratias rostratus Regan, *Ann. Mag. Nat. Hist.*, 9 ser., XV,
May, 1925, 567, Gulf of Panama.

Blank page retained for pagination

INDEX

LIST OF COMMON NAMES OF FISHES OCCURRING IN CHECK LIST

Page	Page	Page	
abadejo.....	315	angled.....	40
aboma de mar.....	436	angel, black.....	360
aboma de rio.....	442	angelfish.....	22, 358, 361
aburabozu.....	373	angel shark, California.....	22
acedia.....	231	angel sharks.....	22
achigan grand bouche.....	297	angler.....	504
achigan noir.....	297, 298	ahil.....	321
achigan petite bouche.....	298	antonino.....	270
acoupa.....	354	aquai.....	314
aguaji.....	314	arctic charr.....	61
aguavina.....	318	arctic flounder.....	227
aguja de casta.....	106, 263	arctic grayling.....	66
agujon.....	196, 197	arctic lake herring.....	62
aku.....	258	arctic lamprey.....	10
akule.....	271	arctic lycodes.....	476
akulliaqitsok.....	474	arctic salmon, Ross's.....	60
Alabama darter.....	292	arctic sculpin.....	386
Alabama shad.....	43	argentine, sheppy.....	73
Alaska blackfish.....	172	argentines.....	68
Alaska codfish.....	210	arnillo.....	328
Alaska greenfish.....	374	arrow-toothed halibut.....	221
Alaska pollack.....	209	Athabaska herring.....	62
Alaska sand lance.....	451	Athabaska minnow.....	115
albacore.....	200	attinga.....	502
West Indies long-finned.....	260	atkaifish.....	374
yellow-flinned.....	270	atka mackerel.....	374
albacores.....	259	atka mackerels.....	374
long-finned.....	260	Atlantic salmon, common.....	66
yellow-flinned.....	260	atule.....	271
ajkey trout.....	212	Aubeenabooe darter.....	291
alewife.....	42	aurora trout.....	60
alewives.....	41	awa.....	39, 40
alfilor.....	248		
alfione.....	411	bacalao.....	315
alfonsino.....	295	bacalhao sabare.....	441
alfonsin a casta.....	234	bachelor.....	304, 305
alfonsin a casta larga.....	234	bachelor perch.....	304
alfonsinos.....	234	bagre.....	151, 157
allonghi.....	260	bagre sapo.....	488
alligatorfish, common.....	394	bagres de rio.....	156
alligatorfishes.....	393	Bahama amphioxus.....	7
alligator gar, Cuban.....	38	Bahama lancelet.....	7
Mississippi.....	38	abajonado.....	336
alligator gars.....	38	balao.....	199
Allison's tuna.....	266	blue-tailed.....	199
allmouth.....	504	red-tailed.....	199
alma seca.....	98	balloonfish.....	501, 502
amberfish, great.....	279	spotted.....	502
amberjack.....	279	unspotted.....	502
American anchovies.....	47	bananafish.....	39
American carp.....	103	banded blenny.....	462
American eel.....	77	banded darter.....	280
American hako.....	214	banded drum.....	352
American perch.....	281	banded larimus.....	344
American pike.....	173	banded mackerel.....	279
American pike porches.....	282	banded ruddersfish.....	279
American saibling.....	60	banded sunfish.....	304
American seahorse, common.....	244	bang.....	42
American shad.....	42	banklick bass.....	305
American smelt.....	67	banner pompano.....	276
American sole.....	229	baracuda.....	355
amphioxus, Bahama.....	7	barb.....	350
California.....	7	barbero.....	362, 363
anchoa pelada.....	45	barbero negro.....	363
anchovies.....	47	barbiers.....	322
American.....	47	barbu.....	256
anchovy, bay.....	47	barbudo.....	157, 234, 256
California.....	47	barbudos.....	234, 254
fry.....	48	barfish.....	306
Porto Rican.....	51	barndoor skate.....	25
striped.....	48	barracouta.....	265

List of common names of fishes occurring in check list—Continued

	Page		Page		Page
bass, trout.....	298	black cod.....	373	blueback mullet.....	253
white.....	297, 303, 307	black crappie.....	305	blueback salmon.....	55
white lake.....	307	black croaker.....	348	blueback trout.....	61
white sea, of California.....	356	black drum.....	352	of Lake Crescent.....	57
wood.....	298	black-eared pondfish.....	299, 301	blue bass.....	298
yellow.....	297, 298, 307	blackear.....	300	blue bream.....	301
bass hogfish.....	298	blackeyes.....	299	blue-breasted darter.....	290
bass killy.....	176	blackeye sunfish.....	298	blue cat.....	152
bass sunfish.....	303	blackfin.....	126	blue cod.....	374
basses, black.....	297	blackfin of Lake Michigan.....	63	blue darter.....	291
rock.....	303	blackfin snapper.....	326	blue dog.....	13
round.....	301	blackfish.....	112, 267, 317, 423	bluefish.....	230, 293, 317
sea.....	303	Alaska.....	172	California.....	339, 356
bastard halibut.....	223	smooth.....	424	bluefishes.....	280
bastard halibuts.....	222	black fresh-water bass.....	203	bluegill.....	301
bastard margaret.....	329	black grouper.....	312, 314	bluehead sucker.....	104
bastard weakfish.....	355	black groupers.....	312	blue herring.....	42
batfish.....	31, 403, 507	black grunt.....	323, 329	blue Joe.....	301
batfishes.....	506	black guativera.....	309	blue Johnny.....	291
bay-a.....	314	black Harry.....	317	blue-lined minnow.....	132
bay anchovy.....	47	blackhead minnow.....	145	blue-mouthed sunfish.....	301
bay mackerel.....	257	blackhorse.....	102	blue mullet.....	110
bay mare.....	105	black huron.....	297	blue parrotfish.....	432, 434
bay shark.....	15	black jewfish.....	312	blue perch.....	301, 424
bayou bass.....	297	black lamprey, small.....	9	blue pike.....	282
Bear Lake herring, great.....	62	black marlin.....	263	blue runner.....	272
Bear Lake whitefish.....	65	Giant.....	283	blue shark, great.....	17
beau gregoire.....	413	black minnow.....	139	small.....	16
becuna.....	255	black moray.....	91, 92	blue smelt, California.....	252
bellowsfish.....	246, 504	black mullet.....	105	blue-spotted sunfish.....	208, 304
bellyfish.....	191	black-nosed dace.....	139, 140	blue-striped triggerfish.....	491
bergall.....	424	black oldwife.....	493	blue sucker.....	148
berg-gylt.....	424	black perch.....	298, 317, 323	blue sunfish.....	298, 301
Bermuda catfish.....	261	black pilot.....	413	blue surgeon.....	302
Bermuda chub.....	340	black rockfish.....	365	blue-tailed balao.....	199
Bermuda lancelet.....	7	black ruddersfish.....	268	blue tang.....	362
Bermuda silk snapper.....	326	black rudderfishes.....	267	blunthead.....	176
berrugate.....	324	black ruffe.....	267	bluntnosed minnow.....	145
besan.....	362	black ruffs.....	267	bluntnosed river carp.....	103
beshow.....	373	black sculpin.....	387	bluntnosed shiner.....	275
bessy cerka.....	491	black sea bass.....	317	boar grunt.....	330
besugo.....	337	black-sided darters.....	283	bobo.....	255
blajalba.....	327	blacksmith.....	413	bobtailed skate.....	25
blajalba de lo alto.....	303	black soapfish.....	316	bocaccio.....	364
bleaya ryba.....	54	black spearfish.....	283	bocce dulce.....	13
big drum.....	352	black-spotted Spanish mackerel.....	257	bocon.....	48, 51
big-eared sunfish.....	300	black sucker.....	108, 107	bocone.....	354
big-eye.....	322, 323	black sunfish.....	302	borderon.....	373
big-eyed herring.....	38, 42	black surfish.....	411	boga.....	340
big-eyed scad.....	271	black swallowe.....	452	bonac.....	314
bigeye minnow.....	134	black swallows.....	455	bonaci arara.....	314
big-eyes.....	322	black-tailed sunfish.....	300	bonaci cardinal.....	313
big-headed darter.....	284	black warmouth.....	302	bonaci de Piedra.....	313
big-headed gurnard.....	403	black will.....	317	bonaci gato.....	315
biglip sucker.....	106	black winter sucker.....	109	bone-dog.....	19, 21
bigmouth.....	302	blanco perch.....	298	bone-eater.....	259
bigmouth buffalo.....	102	blanketfish.....	32	bonefish.....	39
big-mouthed black bass.....	297	blanquillo.....	357	bone-jack.....	259
big-mouthed buffalo fish.....	102	blanquillos.....	357	bone shark.....	20
big-mouthed cat.....	153	blennies.....	400	bonita of Martinique.....	260
big-mouthed sunfish.....	302	blenny.....	462	bonito.....	258, 259, 279, 281
bigmouth sunfish.....	411	banded.....	462	California.....	259
big-nose sunfish.....	200	eel.....	471	oceanic.....	268
big skate of California.....	20	four-banded.....	462	Pacific.....	258
big-toothed redhorse.....	111	seaweed.....	461	bonitos.....	259
billfish.....	30, 196, 197, 242	snake.....	471	bonnethead shark.....	17
billy gar.....	36	spotted snake.....	469	Bonneville whitefish.....	65
bishop ray.....	30	bloater.....	154	bonyfish.....	39, 44
Bissell's herring.....	63	catfishes.....	154	bonyfishes.....	39
bitterhead.....	115, 305	blindfishes, cave.....	194	bony-tail.....	114
black and yellow rock cod.....	369	blindfish, Mammoth Cave.....	195	bootleg sucker.....	107
black angel.....	360	blindfish, small.....	195	boregat.....	373
black-banded rock cod.....	370	blind gobies.....	447	borer.....	8
black-banded sunfish.....	303	blind goby of Point Loma.....	447	borers.....	8
black bass.....	293, 317	blind simon.....	287	borlase.....	267
big-mouthed.....	297	blob.....	63	Bosc's shanny.....	462
Kentucky.....	298	of Lake Michigan.....	63	Boston mackerel.....	257
largemouth.....	297	blob.....	388	botete.....	498
large-mouthed.....	297	bloody sunfish.....	300	boucanelo.....	328
of the Huron.....	297	blower.....	498	bout de tabac.....	319
small-mouthed.....	298	bowfin.....	499	bowfin.....	38
black basses.....	297	hairy.....	499	bowfins.....	38
black-belly.....	42	spinyback.....	498	boxfish.....	497
black buffalo.....	102	blue-and-green sunfish.....	298	hairy.....	501
black bullhead.....	154	blue angel.....	361	spiny.....	502
black burrfish.....	502	blueback.....	42, 62	bramble shark.....	22

List of common names of fishes occurring in check list—Continued

Page	Page	Page			
bramble sharks.....	22	bullhead shark.....	10	capitaines.....	426
branch herring.....	42	little.....	22	caranx, yellow.....	273
branch perch.....	302	bullhead sharks.....	10	carbonero.....	271
brassy sculpin.....	388	bullock.....	434	caronetta.....	16
bream.....	115, 298-303, 336, 337	bulldredfish.....	348	cardenal.....	295
blue.....	301	bull trout.....	61	cardinafishes.....	295
calico.....	305	bumper.....	276	cardoniaria.....	370
copper-headed.....	301	burbot.....	212	Carolina pompano.....	277
copper-nosed.....	301	burbots.....	212	Carolina robin.....	407
little.....	304	burrfish.....	502	Carolina sea robin.....	407
morelmouth.....	302	black.....	502	Carolina whiting.....	350
perchmouth.....	302	burrfishes.....	502	carp.....	103
red-bellied.....	300	burrito.....	333, 334	American.....	103
redbreast.....	299	burritos.....	333	blunt-nosed river.....	103
red-eyed.....	303	burro.....	334	common river.....	103
red-headed.....	299	burro blanco.....	333	lake.....	103
red-sided.....	118	burros.....	333	silver.....	103
red-tailed.....	200	butter chub.....	146	carp sucker.....	103
sea.....	285	butterfish.....	267, 277, 309, 467, 483	carp suckers.....	103
spotted.....	299	lemon-yellow.....	309	casabe.....	276
warmouth.....	302	butterflyfishes.....	355	casabes.....	276
yellow.....	302	butterfly ray.....	29	catagnole.....	266
breams, sea.....	265	caballina.....	257	catalina.....	332
bride perch.....	297	caballito del mar.....	244	catalineta.....	332, 361
bridge perch.....	304	caballeroete.....	325	catalinetas.....	361
bridled minnow.....	135	cabezon.....	344, 382, 487	catalufa.....	322, 323
brier ray.....	25	smooth.....	390	catalufas.....	322
brilliant sunfish.....	300	cabezones.....	382	deep-water.....	323
bristle herring.....	246	cabezoete.....	246	cat, big-mouthed.....	153
broadbill.....	264	cobra mora.....	310	blue.....	152
broadhead, grubber.....	47, 50	cabrilla.....	311, 317	brown.....	154
broad shad.....	241	spotted.....	317	channel.....	163
broad soles.....	220	cabrilla calamaria.....	314	of the Potomac.....	163
brochet de mer.....	305	cabrilla de astillero.....	314	duckbill.....	36
bronze-backer.....	298	cabrilla de raizero.....	313	eel.....	152
bronzed centrarchus.....	303	cabrilla pinta.....	310	Fulton.....	152
bronze minnow.....	121	cabrilla piritita.....	315	great fork-tailed.....	152
brook lamprey.....	9, 10	cabrillas verdes.....	317	long-tailed.....	212
brook lampreys.....	10	cachuccho.....	328	Mississippi.....	162
brook minnow.....	140	Cesar.....	331	mud.....	154, 165
brook silverside.....	251	Cesar grunt.....	330	Russian.....	165
brook stickleback.....	238	caffa bota.....	11	Sacramento.....	164
brook sucker.....	105	engojo de lo alto.....	328	Schuylkill.....	164
brook trout.....	60	cailliu.....	43	sea.....	149
eastern.....	60	caillen tassart.....	44	spoonbill.....	36
of western Oregon to southern Alaska.....	58	caliman.....	440	stone.....	155
brotula.....	479	coj.....	325	white.....	152, 153
brown bass.....	298	crafate.....	493	willow.....	152
brown buffalo.....	102	calice bass.....	305	yellow.....	163, 165
brown cat.....	164	calico bream.....	305	catfish.....	176, 256
brown hind.....	308	calico salmon.....	54	sea.....	148, 149
brown rock cod.....	369	California amphioxus.....	7	small.....	164
brown rockfish.....	368	California anchovy.....	47	catfishes, blind.....	154
brown trout.....	298	California angel shark.....	22	cave.....	150
brown-winged sea robin.....	407	California barracuda.....	285	gafftopsail.....	147
bubbler.....	352	California bluefish.....	239, 386	mailed.....	160
buffalo.....	103	California blue smelt.....	27	sea.....	147
bigmouth.....	102	California bonito.....	259	South American.....	166
black.....	102	California common skate.....	26	cats, channel.....	151
brown.....	102	California dogfish.....	21	stone.....	155
gourdsseed.....	102	California flounder.....	226	tadpole.....	165
mongrel.....	102	California garfish.....	198	cat shark.....	14
quillback.....	102	California hagfish.....	8	cat sharks.....	11
razor-backed.....	102	California herring.....	41	cavalla.....	257, 273
round.....	102	California jewfish.....	308	cavallas.....	269
smallmouth.....	102	California lancelet.....	7	cave blindfishes.....	194
suckermouth.....	102	California pampano.....	266	cave catfishes.....	159
buffalo bass.....	302	California redfish.....	420	cavefish.....	194
buffalo cod.....	374	California sardine.....	43	Missouri.....	105
buffalofish, big-mouthed.....	102	California sole.....	22	cazon de playa shark.....	16
common.....	102	California stingray.....	31	céfalo.....	263
redmouth.....	102	California torpedo.....	27	centrarchus, bronzed.....	303
buffalofishes.....	102	California whiting.....	350	cernier.....	308
buffalo sculpin.....	381	Californiar mullet.....	304	cero.....	257
buffalo sunfish.....	298	campbellite.....	304	spotted.....	257
bugara.....	412	candil.....	235	chagrin.....	18
bugfish.....	44	candlefish.....	66	chainside.....	301
bulifish.....	497	cane de mare.....	19	chain-sided perch.....	301
bulldhead.....	385, 386	capelin.....	66	chanl.....	40
black.....	154	capelina.....	66	channel bass.....	348
common.....	154	Capo May goody.....	349	channel cat.....	152
prickly.....	383	capidoll.....	20	of the Potomac.....	163
Rocky Mountain.....	384	capitalne.....	425	channel cats.....	153
bullhead minnow.....	145			chapin.....	496, 497

List of common names of fishes occurring in check list—Continued

Page	Page	Page			
chappaul.....	114	coachman.....	320	common weakfish.....	355
characins.....	95	coalfish.....	208, 277	common whitefish.....	65
charioteer.....	320	coast-range trout.....	58	compressed redfin.....	120
charnia ryba.....	172	cobbler.....	175	conejo.....	261
charr, Arctic Greenland.....	61	common.....	176	coney.....	308
Oregon.....	61	cobblerfish.....	274, 277	conger, sand.....	454
charrs.....	60	cobblerfishes.....	274	conger eel.....	54, 79, 473
chauffe soleil.....	413	Cohessicantic smelt.....	67	of California.....	91
Cheney's darter.....	286	cobia.....	281	conger eels.....	70
cherna americana.....	312	cochinito.....	363	Connecticut River shad.....	42
cherna criolla.....	311	cochino.....	401	conner.....	424
cherna de lo alto.....	310	cocinera.....	272	constantino.....	300
cherna de vivero.....	312	cocinero.....	272	constantino de las aletas prietas.....	300
chibcharo.....	271	cocinero dorado.....	272	Copeland's darter.....	285
chickwick.....	355	cock and hen paddle.....	397	copper-headed bream.....	301
chicolar.....	261	cockeye pilot.....	413	copper-nosed bream.....	301
chigh.....	113	cockfishes.....	493	coralfishes.....	359
chimeras.....	10, 32	cod, black.....	373	corbineta.....	246
chimeroids.....	32	blue.....	374	cornetfishes.....	245
chinook salmon.....	55	buffalo.....	374	coronado.....	279
chinquapin perch.....	290, 304, 305	cultus.....	374	corporal.....	117
chirivita.....	360	green.....	209	corsair.....	367
chirivitas.....	360	Pacific.....	210	corvalos.....	351
chiselmouth.....	112	wachna.....	210	corvina.....	344, 349, 365
chiselmouth jack.....	66	codfish, Alaska.....	210	corvina de las aletas am. arillas.....	356
chivin.....	112	common.....	210	couchu.....	181
chivo.....	117	Greenland.....	210	coudia.....	108
chogset.....	344	codfishes.....	209, 210	Coulter's whitefish.....	65
chopa.....	424	codfish shark.....	211	courpata.....	269
chopa amarilla.....	339	codling.....	213	cow bass.....	297
chopa blanca.....	340	spotted.....	213	cowfish.....	31, 407
chopa spina.....	337	codlings.....	212	cow-nosed ray.....	31
chopper.....	102	codorniz.....	350	cow pilot.....	415
choupique.....	38	cods, cultus.....	374	cow shark.....	11
chub.....	63, 64, 109, 115, 117, 137, 297, 302, 340	cod shark.....	21	cow sharks.....	11
Bermuda.....	340	coelho.....	261	crab-eater.....	281
butter.....	146	cojinera.....	272	crab-eaters.....	281
Columbia River.....	112	Colorado River squawfish.....	114	crabra.....	370
creek.....	117	Colorado River trout.....	57	Craigin's darter.....	292
flat-headed.....	136	Columbia River chub.....	112	Craig fluke.....	228
greased.....	146	Columbia River salmon.....	55	crampfish.....	27
Indian.....	137	Columbia River sucker.....	106	crapet.....	304, 305
Kaweah.....	113	Columbia River trout.....	56	carpet jaune.....	302
lake.....	136	common alligatorfish.....	394	carpet mondoux.....	303
millpond.....	297	common American seahorse.....	244	carpet noir.....	303
mud.....	117, 302	common Atlantic salmon.....	56	carpet vert.....	303
nigger.....	146	common buffalofish.....	102	crappie.....	304, 305
of the Rio Grande.....	120	common bullhead.....	154	cravo.....	216
river.....	137	common cobbler.....	176	crawl-a-bottom.....	107, 111, 283
Sacramento.....	120	common codfish.....	210	creek chub.....	117
silver.....	117, 137	common dolphin.....	264	creekfish.....	109
southern.....	297	common eastern stickleback.....	238	creek sucker.....	109
steel-backed.....	146	common flatfish.....	227	croole.....	428
Storer's.....	137	common gar pike.....	36	creolefish.....	320
chub mackerel.....	258	common gray shark.....	15	crested gobies.....	438
chub minnow.....	136	common grunt.....	330	crevallé.....	272, 273
chub sucker.....	108, 109	common gurnard.....	407	croaker.....	349, 352
chuchin.....	158	common halfbeak.....	198	black.....	348
chum.....	54	common halibut.....	221	white.....	353
chuss.....	213	common herring.....	41	yellow-tailed.....	350
cibi.....	271	common lake herring.....	175	croakers.....	344
cibi amarilla.....	272	common mackerel.....	257	cuatros ojos.....	194
cibi mancho.....	271	common pampano.....	277	cub.....	424
cigarfish.....	270	common pipefish.....	205	Cuban alligator gar.....	38
cirstostomes.....	7	common rat-tail.....	205	cubera.....	325
cisco, Lake Michigan.....	63	common redhorse.....	110	Cub shark.....	15
mooneye.....	64	common river carp.....	113	cuckold.....	497
of Green Lake.....	64	common rock bass.....	303	cucumberfish.....	502
of Lake Bonneville.....	63	common sculpin.....	386	cucuyoco.....	491
of Lake Michigan.....	64	common seup.....	336	cultus cod.....	374
of Lake Ontario.....	63	common shad.....	42	cultus cods.....	374
of Lake Superior.....	63	common skate.....	25, 26	Cumberland darter.....	293
of Lake Tippecanoe.....	62	common sucker.....	105	cumprida.....	234
ciscos.....	61	common sunfish.....	300, 302	cunner.....	424
civette de Mer.....	409	common spotted moray.....	91		
clamagore.....	434	common stickleback.....	237		
clam-cracker.....	28	common sturgeon.....	34		
clear-nosed skate.....	25	common sunfish.....	300, 302		
clear ray.....	25	common surgeon.....	362		
clingfishes.....	458	common swordfish.....	264		
clingy goby.....	446	common trunkfish.....	497		
cloudy lycodes.....	475				

List of common names of fishes occurring in check list—Continued

Page	Page	Page	
cunners.....	424	dogfish.....	13, 10, 21, 174, 176
cusk.....	213, 214	California.....	21
European.....	214	horned.....	19
rock.....	385	picked.....	19, 21
cusk eels.....	485	smooth.....	13
cutlassfish.....	262	spiny.....	21
cutlassfishes.....	262	dogfishes.....	21
cut-lips.....	112, 146	dog salmon.....	54
cut-throat trout.....	56	dog shark.....	13
of Utah Lake.....	56	dog snapper.....	325
cybium, spotted.....	257	dollarfish.....	267, 275
dab.....	222	dolphin.....	300
sand.....	222	Dolly Varden.....	305
smear.....	228	Dolly Varden trout.....	61
dabs, mud.....	220	dolphin, common.....	204
smear.....	228	small.....	265
dace.....	115, 117, 128, 140	dolphins.....	264
black-nosed.....	139, 140	dômîne.....	261
horned.....	117	doncella.....	426
long-nosed.....	130	doucellas.....	426
red-bellied.....	113	dorado.....	264
rosy.....	121	dorados.....	264
striped.....	140	dormeur.....	323
daces.....	112	dory.....	282
daddy sculpin.....	387	hair-flinned.....	275
daddy sculpins.....	386	ocellated.....	232
darter, Alabama.....	292	spinous.....	276
Aubeenaubee.....	291	dotted paintedtail.....	297
banded.....	289	dotted scad.....	270
big-headed.....	284	double-bearded flyingfish.....	201
black-sided.....	283	doughbelly.....	140
blue.....	291	dourade.....	264
blue-breasted.....	290	dovetailfish.....	415
Cheney's.....	286	dragoneets.....	450
Copeland's.....	285	drum.....	103, 352
Cragin's.....	292	banded.....	352
crimson.....	294	black.....	352
Cumberland.....	293	fresh-water.....	352
Evermann's.....	284	gray.....	352
fan-tailed.....	293	red.....	348
fusiform.....	294	drumfish.....	352
Gilbert's.....	286	drummer, Jewsharp.....	351
Grand Oranche.....	287	mongolar.....	355
green-sided.....	286	whitemouth.....	349
Günther's.....	283	drums, red.....	348
Jessie's.....	291	river.....	352
Johnny.....	287	sea.....	352
Jordan's.....	290	drunkenfish.....	496
Julia's.....	292	Dublin Pond trout.....	60
least.....	294	duckbill cat.....	36
manitou.....	283	duckbill eels.....	82
Maryland.....	284	dusky shark.....	16
Maxinkuckee.....	284	dwarf bass.....	298
Meadie's.....	286	Eagle Lake trout.....	58
Niangua.....	285	eagle rays.....	30, 31
northern.....	291	seer sunfish.....	300
ouachite.....	284	Easter mackerel.....	258
Putnam's.....	285	eastern brook trout.....	60
rainbow.....	291	eastern mud minnow.....	174
sand.....	287, 288	eastern pickerel.....	173
shielded.....	283	eastern redhorse.....	111
Shumard's.....	285	eastern stickleback, com-mon.....	238
Susan's.....	288	East Indian basking shark.....	18
tasselled.....	287	eel.....	91
tessellated.....	290	American.....	54, 70, 473
Tuscumbia.....	293	conger.....	54, 70, 473
Whipple's.....	292	of California.....	91
darters.....	282	fresh-water.....	77
black-sided.....	283	lamprey.....	473
fantail.....	293	moray.....	8
sand.....	289	sand.....	451, 485
tessellated.....	287	sea.....	88
devilfish.....	216	snipe.....	83
deceptive sunfish.....	299	whipsnake.....	88
deep-water catalufis.....	323	wolf.....	472
demoiselle.....	415	eel-back flounder.....	227
dentuda.....	19	eel blenny.....	471
devilfish.....	28, 32	eel cat.....	152
small.....	32	eel pout.....	212, 473
diablo.....	507	eels.....	54, 77
diamond flounder.....	225	conger.....	79
doctorfish.....	362	cusk.....	485
dog, blue.....	13	duckbill.....	82
		long-necked.....	77
		coze.....	78
		snake.....	84
		snipe.....	83
		snub-nosed.....	78
		spiny.....	75
		symbranchoid.....	76
		thread.....	83
		true.....	77
		worm.....	82
		eight-spined sculpin.....	386
		eight-threaded threadfish.....	256
		electricfish.....	27
		electric rays.....	27
		elephantfish.....	33
		elephant fishes.....	32
		elephant shark.....	20
		ellwife.....	42
		emeraldfish.....	441
		emerald minnow.....	123
		emerald trout.....	58
		emerando.....	441
		endormi.....	441
		enjambre.....	308
		ontemedor.....	27
		Erie great herring.....	63
		Erie herring.....	63
		erizo.....	501
		escalar.....	261
		escolar chino.....	297, 328
		escolar de natura.....	269
		escribano.....	200, 261
		esmeralda.....	441
		esmeralda de mar.....	437
		esmeralda negra.....	437
		espada.....	204
		espadón, emperador.....	264
		elachon.....	60
		European barracuda.....	255
		European cusk.....	214
		European lake.....	214
		European sculpin.....	386
		Everglades pygmy sunfish.....	305
		Evermann's darter.....	284
		Evermann's trout.....	59
		faquito.....	191
		fair maid.....	336
		fallfish.....	117
		red.....	127
		fallfishes.....	117
		fall herring.....	42
		fanogal.....	370
		fan-tail darters.....	203
		fan-tailed darters.....	203
		fan-tail mullet.....	253, 254
		fatback.....	44, 280
		fathead.....	428
		fathead minnow.....	145
		fatheads.....	145, 428
		fatherlasher.....	386
		fat priest.....	373
		fat priests.....	373
		female perch.....	302
		flatolas.....	266
		fiddlefish.....	22
		fiddler.....	152
		fiddlerfish.....	23
		fleefish.....	401, 404, 495
		orange.....	495
		thread.....	404
		filefishes.....	493
		fine-scaled mullet.....	105
		fine-scaled sucker.....	105
		fishes, bony.....	39
		soft-rayed.....	39
		true.....	33
		fishing-frog.....	504
		fishing-frogs.....	504

List of common names of fishes occurring in check list—Continued

Page	Page	Page			
fish of Dismal Swamp.....	194	gabilian.....	31	golet.....	61
fish of the rice ditches of South Atlantic States.....	194	gaftopsail.....	148, 276	goodies.....	348
fish shark.....	18	gaftopsail catfishes.....	147	goody.....	349
five-bearded rockling.....	208	galafate.....	493	Cape May.....	349
five-bearded rocklings.....	208	gall bengal.....	424	goosefish.....	504
flamenco.....	326	gallwasp.....	163	gorbuscha.....	54
flannel-mouthed sucker.....	107	gar.....	195	goujon.....	155
flamingo.....	494	billy.....	36	gourami.....	314
flasher.....	323	Cuban alligator.....	38	gourdseed buffalo.....	102
flashers.....	323	green.....	196	gourdseed sucker.....	102
flatfish.....	299, 302, 360	long-nosed.....	36	grande écaille.....	39
common.....	227	Mississippi alligator.....	38	Grand Oranhee darter.....	287
flatfishes.....	217	Ohio.....	36	grass bass.....	303
flat-headed chub.....	136	salt-water.....	196	grass pike.....	173
flesh-colored rock cod.....	369	silver.....	196	grass porgy.....	337
dier.....	304	garfish.....	196	grass rockfish.....	369
floma.....	365	California.....	196	grassy ground rockfish.....	323
Florida lancelet.....	7	Ohio.....	36	grayback.....	63
Florida sailfish.....	283	garibaldi.....	415	gray bass.....	297
flounder.....	299, 302	garibaldis.....	415	gray drum.....	352
arctic.....	227	garlopa.....	314	gray grunt.....	329
California.....	226	gar pike, common.....	36	grayling, arctic.....	66
diamond.....	225	gar pikes.....	36	Michigan.....	68
eel-back.....	227	garrupa.....	365	Montana.....	68
four-spotted.....	224	garrupas.....	308	graylings.....	65
gulf.....	224	gars, alligator.....	37	gray piko.....	282
lemon-yellow.....	227	short-nosed.....	37	gray shark.....	15
peacock.....	217	gascon.....	270	common.....	15
pole.....	223	gaspreau.....	42	gray sharks.....	13
southern.....	224	gaspergou.....	352	gray snapper.....	325
summer.....	224	gata.....	12	gray sole.....	228
winter.....	227	gato.....	13	gray sucker.....	106
flounders.....	217	genizara.....	426	gray trout.....	355
great.....	228	Georgian Bay herring.....	62	greased chub.....	148
starry.....	228	ghostfish.....	471	great amberfish.....	279
winter.....	227	ghost smelt.....	68	great barracuda.....	255
fluke.....	224	giant black marlin.....	263	great Bear Lake herring.....	62
Craig.....	223	Gila trout.....	114	great blue shark.....	17
flutemouth.....	245	Gilbert's darter.....	286	great flounders.....	228
flyfish.....	387	gilded sunfish.....	301	great flyingfish.....	202
flyingfish.....	211, 407	gizzard shad.....	46	great fork-tailed cat.....	152
double-bearded.....	201	gizzard shads.....	216	great herring, Erie.....	63
sharp-finned.....	202	glancefish.....	282	Great Lakes trout.....	59
single-bearded.....	201	glass-eye.....	251	great northern pike.....	173
flyingfishes.....	100, 200	glassfish.....	503	great pampano.....	277
flying gurnard.....	409	globefish.....	499	great pike.....	172
flying gurnards.....	403	glovefish.....	42	great pipefish.....	240
flying perch.....	304	glut herring.....	343	great sculpin.....	387
flying robin.....	409	goatfish.....	343	great sea lamprey.....	8
foolfish.....	404, 405	red.....	343	great torpedo.....	27
fork-beard.....	213	yellow.....	343	great trout of Lake Tahoe.....	57
four-banded blenny.....	402	goatfishes.....	343	great tunnies.....	259
four-bearded rockling.....	209	gobies.....	438	great white sharks.....	20
four-bearded rocklings.....	209	blind.....	447	greenback trout.....	57
four-eyed fish.....	184	crested.....	447	green bass.....	297, 298
four-eyed fishes.....	194	half-naked.....	445	green cod.....	209
four-spined stickleback.....	239	goby.....	446	greenfish.....	280, 339
four-spotted flounder.....	224	blind, of Point Loma.....	447	Alaska.....	374
fox shark.....	18	cling.....	448	green gar.....	196
Frasier River salmon.....	55	long-jawed.....	448	green jack.....	271
French grunt.....	330	naked.....	444	green lamprey.....	8
frère jacques.....	235	sharp-tailed.....	446	Greenland charr.....	60
fresh-water bass.....	303	variegated.....	446	Greenland codfish.....	210
black.....	293	gog.....	314	Greenland halibut.....	221
fresh-water drum.....	352	goggle-eye.....	302-305	greenling.....	374
fresh-water eel.....	77	goggle-eye jack.....	271	greenlings.....	373
fresh-water shark.....	16	goggler.....	271	green parrotfish.....	432, 435
fresh-water silverside.....	247	gold bass.....	298	green perch.....	297, 298
fresh-water sunfish.....	299	golden angel.....	361	green pike.....	173
fresh-water trout.....	297	golden trout of south fork of Kern River.....	59	green pipefish.....	242
friar.....	247	golden trout of Sunapee Lake.....	60	green-sided darter.....	286
friars.....	246	golden trout of Volcano Creek.....	59	green sturgeon.....	34
frigate mackerel.....	259	golden trout of western tributaries of Kern River.....	59	green sunfish.....	298
frogfishes.....	504	goldeye.....	41	small.....	300
frostfish.....	210	goldeyes.....	41	green trout.....	27, 298
frostfishes.....	262	goldfish, sea.....	415	grenadier.....	202
fry, anchovy.....	48	gold perch.....	301	grenadiers.....	202
hogmouth.....	48	goldring.....	304	ground drummer.....	346
tarpon.....	51	goldthread shiner.....	116	ground rockfish, grassy.....	320
fucinids.....	295			ground spearling.....	163
Fulton cat.....	162			groupers.....	311
funal.....	370			black.....	312, 314
fusiform darter.....	294			Nassau.....	311
				poison.....	370, 371
				red.....	312

CHECK LIST OF FISHES

519

List of common names of fishes occurring in check list—Continued

	Page		Page	Page	
grouper, snowy-----	311	hairy boxfish-----	501	herrings, lake-----	41, 61
spotted-----	311	hairy porcupinefish-----	501	round-----	45
yellow-----	314	hake-----	213, 350	hickory jack-----	42
yellow-finned-----	311, 313	American-----	214	hickory shad-----	42, 46
groupers-----	308, 310	European-----	214	hind, brown-----	308
black-----	312	New England-----	214	red-----	312
growler-----	297	Old English-----	213	rock-----	308, 310
salmon-formed-----	297	Old England-----	214	speckled-----	312
grubber broadhead-----	47, 50	red-----	213	bitch-----	113
grubby-----	386	silver-----	214	hoe-----	19
grunion-----	251	spotted-----	213	hoe-mother-----	20
grunt, black-----	323, 329	squirrel-----	213	hog bass-----	238
boar-----	330	white-----	213	hogchoker-----	229
Casar-----	330	hakes-----	214	hogfish-----	283, 318, 335, 425
common-----	330	halfbeak, common-----	198	bass-----	208
French-----	330	slender-----	198	Spanish-----	425
gray-----	329	halfbeaks-----	198	bogmolly-----	107, 283
humpback-----	330	halimoon-----	358	hogmouth fry-----	48
margaret-----	329	halfmoonfishes-----	368	hog sucker-----	107
open-mouthed-----	330	half-naked gobies-----	445	Holbrook's top minnow-----	185
redmouth-----	331	half-naked lycodes-----	475	holia-----	54
red-mouthed-----	335	halibut-----	221	hoopid salmon-----	54
striped-----	329, 332	arrow-toothed-----	221	hoosier-----	155
white-----	331	bastard-----	223	horned dace-----	117
yellow-----	330	common-----	221	horned dogfish-----	19
young-----	331	Greenland-----	221	horned-eared sunfish-----	299
grunts-----	329	Monterey-----	223	horned pout-----	153, 154
striped-----	331	halibuts-----	221	horned pouts-----	151
guabín, small-----	180	bastard-----	222	horned trunkfish-----	497
guacamala-----	433	hamlet-----	91, 311	horny cony-----	494
guaguanche pelón-----	255	hammerhead-----	18, 107	hornyhead-----	137
guajacon-----	183	hammerhead shark-----	18	horse crevallé-----	273
guajacones-----	184	hammerheaded sharks-----	17	horse-eyed jack-----	271
guajica-----	191	handsawfish-----	171	horse-eye jack-----	272
guanuanche-----	255	handsawfishes-----	171	horsefish-----	244, 275
guapena-----	353	annahill-----	317	horsehead-----	275
guardfish-----	146	harder-----	268	horse mackerel-----	270, 280
guardubaco-----	436	hardhead-----	109, 349	hound, smooth-----	13
guasa-----	310, 312	hardheads-----	190, 246	houndfish-----	190, 197
guasas-----	312	hardmouth-----	112	houndfishes-----	195
guativera-----	309	hard-mouthed suckers-----	105	humpback grunt-----	330
black-----	309	hardtail-----	272	humpback salmon-----	54
red-----	309	harelip sucker-----	112	humpback sucker-----	108
guativera amarilla-----	309	harlequin roach-----	302	humpback whitefish-----	65
guavina-----	354, 435, 430, 437	harvestfish-----	264, 267	huron-----	297
guavina hoyera-----	442	harvest pike-----	196	black-----	297
gudgeon-----	115, 132, 143	hayko-----	54	leefish-----	67
Niagara-----	130	headfish-----	503	inconnu-----	61
gudlax-----	216	bechudo-----	60	Indian chub-----	137
guitarfish-----	24	hedgheog ray-----	25	Indianfish-----	302, 360
guitarfishes-----	23	hemdurgan-----	363	Indian remora-----	448
guitarro-----	23, 24	herring, arctic lake-----	62	Irish lord-----	381
gulf flounder-----	224	Athabasca-----	62	Irish pompano-----	342
gulf menhaden-----	44	big-eyed-----	63	isabelita-----	361
gulf sea bass-----	318	Bissell's-----	63	isabelita de la alto-----	360
gulpers-----	96	blue-----	42	iverkoak-----	212
gunnel-----	467	branch-----	42	jabon-----	316
Günther's darter-----	283	bristle-----	246	Jaboncillo-----	316
guppy-----	191	California-----	41	Jack-----	173, 273, 364
gurnard, big-headed-----	408	common-----	41	chiselmouth-----	66
common-----	407	common lake-----	63	goggle-eye-----	271
flying-----	409	Erie-----	63	green-----	271
northern striped-----	407	Erie great-----	63	hickory-----	42
red-winged-----	407	Georgian Bay-----	42	horse-eye-----	272
southern striped-----	407	glut-----	62	horse-eyed-----	271
spinous-----	407	great Bear Lake-----	62	yellow-----	272
gurnards-----	407	jumbo-----	63	jackfish-----	279
flying-----	408	king of the-----	63	jack salmon-----	282
hackleback-----	36	lake-----	63	jacome-----	319
hacklehead-----	386	Lake Huron-----	62	jalao-----	320
haddo-----	64	least lake-----	62	janissary-----	428
haddock-----	211	Mackenzie River-----	62	jaqueta-----	415
Jerusalem-----	216	Michigan-----	62	jawfishes-----	452
Norway-----	303	rainbow-----	67	Jeniguana-----	330
hagfish-----	8	Rawson Lake-----	63	Jeniquano-----	331
California-----	8	Seneca Lake-----	63	Jenny Lind-----	279
hagfishes-----	8	southern toothed-----	61	jerker-----	137
hair-finney dory-----	275	summer-----	62	Jerusalem haddock-----	216
hair-finney silverfish-----	275	tailor-----	62	Jessie's darter-----	291
hairtail-----	262	thread-----	44	jewfish, black-----	312
silvery-----	262	toothed-----	40	California-----	308
hairtails-----	262	wall-eyed-----	42	spotted-----	312
hairy blowfish-----	499			jawfishes-----	308
hairy blowfish-----	499			wallsharp drummer-----	351

List of common names of fishes occurring in check list—Continued

	Page		Page		Page
Jiguagna.....	273	lake herrings.....	61	lisa blanca.....	253
jocú.....	325	Lake Huron herring.....	62	lisa cahezunda.....	253
John A. Grindle.....	38	lake lamprey.....	9	lisa francesca.....	39
john demon.....	304	lake lawyer.....	212	lisita.....	254
john dories.....	232	Lake Michigan cisco.....	63	little bass.....	208
John Mariggle.....	39	lake minnow.....	123, 137	little bream.....	304
john paw.....	312	lake perch.....	281	little bullhead shark.....	22
johnny.....	390	lake sculpin.....	380	little head porgy.....	236
johnny darter.....	287	lake sheepshead.....	352	little killifish.....	175
johnny verde.....	317	lake sturgeon.....	35	little mouth porgy.....	337
jolt-head porgy.....	336	Lake Superior whitefish.....	64	little pickerel.....	173
jopatan.....	335	Lake Tahoe trout.....	57	little red-eye.....	298
Jordan's darter.....	290	lamia.....	15	little rocadador.....	349
Jordan's trout.....	57	lampre eel.....	473	little skate.....	25
Jorobado.....	275	lampnerina.....	8	little smelt.....	252
toturo.....	254	lamprey, arctic.....	10	little stargazer.....	385
jugfish.....	497	brook.....	9, 10	little sunfish.....	304
jugmouth.....	302	great sea.....	8	little tunny.....	258
Julia's darter.....	292	green.....	8	liza.....	253
jumbo herring.....	63	lake.....	9	liza ojo de perdiz.....	254
jumper.....	297, 298	Mexican.....	9	lizardfish.....	163
jumping mullet.....	111	mud.....	9	lizardfishes.....	163
jmprock.....	111	river.....	8	lodde.....	66
June sucker.....	108	shad.....	8	logfish.....	268
jurel.....	272, 278	silver.....	9	log perch.....	283
kale.....	363	small black.....	9	log perches.....	282
kálog.....	387	three-toothed.....	10	long-eared sunfish.....	209, 300
Kamchatka salmon trout.....	56	western.....	10	long-finned albacore, West Indies.....	260
Kamloops trout.....	58	lampreys.....	8	long-finned albacores.....	260
kawakawa.....	259	brook.....	10	long-finned sole.....	228
kaweah chub.....	113	river.....	9	long-finned sunfish.....	304
kelp bass.....	317	lancelet, Bahama.....	7	long-headed trout of Crescent Lake.....	57
kelpfish.....	427, 458	Bermuda.....	7	longjaw.....	196
spotted.....	458	California.....	7	Ontario.....	63
Kennicott's whitefish.....	65	Florida.....	7	of Lake Superior.....	64
Kentucky black bass.....	298	Virginia.....	7	long-jawed goby.....	444
Kern River trout.....	58	West Indian.....	7	longjaws.....	195
kete.....	500	lancelets.....	7	longningo.....	495
killie.....	176	lanctetfish.....	171, 312	long-mouthed minnow.....	121
killifish.....	176, 181	landlocked salmon.....	56	long-necked eels.....	77
common.....	175	Lane snapper.....	327	long-nosed dace.....	139
little.....	175	langburn.....	470	long-nosed gar.....	36
killifishes.....	174	lanternfishes.....	106, 169	long-nosed sucker.....	108
killy.....	176	lapon.....	371	long-scaled sunfish.....	100
bass.....	176	la queche.....	41	long-spined sculpin.....	786
kingfish.....	257, 349, 350	large-finned bass.....	304	long-tailed cat.....	212
kingfishes.....	350	largemouth black bass.....	287	long-tailed stingray.....	31
king mullet.....	343	large-mouthed black bass.....	297	longtail shark.....	18
king of the herring.....	216	large-scaled sucker.....	110	longdoe.....	59
King of the mackerels.....	501	larimus, banded.....	344	longdown.....	275
king of the mullets.....	295	Laretta whitefish.....	12	loro.....	434, 435
king of the salmon.....	216	lavarets.....	64	Lost River sucker.....	108
king salmon.....	55	lawyer.....	28, 25	loulu.....	496
kingston.....	22	lake.....	212	Lower California trout.....	59
kisutch.....	54	leaping tuna.....	259	lucky broach.....	286
kit.....	28	least darter.....	294	lumpfish.....	297
kiver.....	200, 302	least lake berring.....	62	spinous.....	308
kyi.....	64	leathercoat.....	277	lumpsucker.....	307, 399
Kodiak smelt.....	67	leather-ear.....	299	lycodes, arctic.....	476
krashnaya ryba.....	55	leatherfish.....	494	cloudy.....	475
kuroko.....	263	leatherjacket.....	277, 491	half-naked.....	475
kurokajiki.....	263	plain.....	277	spotted.....	474
kweewee.....	108	leatherjackets.....	277	Vahl's.....	474
kyack.....	42	leather-sided minnow.....	119		
Labrador whitefish.....	64	leatherwing.....	299	macabí.....	30
Lac de Mubre trout.....	61	lebisa.....	28	macho.....	153
lacefin.....	129	lebrancho.....	253	machuelo.....	44
lacerto.....	163	le kai salmon.....	54	machuto.....	253
ladyfish.....	30, 425	lemonfum.....	129	Mackenzie River herring.....	62
Spanish.....	425	lemon sole.....	228	mackerel, Atka.....	374
ladyfishes.....	39	lemon-yellow butterfish.....	300	banded.....	279
lady ray.....	30	lemon-yellow flounder.....	227	bay.....	257
lafayette.....	267, 349	leopard shark.....	14	black-spotted Spanish.....	257
lagarto.....	163	licorne de Mer.....	397	Boston.....	257
lake bass.....	297, 303, 305	lijá.....	494, 495	chub.....	258
white.....	307	lijá barbuda.....	496	common.....	257
lake carp.....	103	lijá colorada.....	494	Faster.....	258
lake chub.....	136	lijá trompa.....	495	frigate.....	259
Lake Erie bass.....	305	lined sea robin.....	407	horse.....	280
Lake Erie whitefish.....	65	ling.....	212, 213, 374, 473	Monterey Spanish.....	257
Lake herring.....	63	thimble-eyed.....	213	skip.....	280
arctic.....	63	lings.....	212	snake.....	260
common.....	63	lionfish.....	371	snap.....	280
least.....	62	lise.....	253	snapping.....	280

List of common names of fishes occurring in check list—Continued

Page	Page	Page			
mackerel, Spanish.....	257	menhaden.....	44	Mississippi cat.....	152
of England.....	258	gulf.....	44	Missouri cavefish.....	195
spotted.....	257	short-tailed.....	44	Missouri sucker.....	102
thimble-eyed.....	258	menhadens.....	44	missuckeke-kequock.....	307
tinker.....	258	Menominee whitefish.....	65	mojarra.....	343, 416
yellow.....	272, 273	mero.....	312	Mexican.....	343
mackerel midge.....	208	merco de lo alto.....	310, 304	mojarra almejero.....	329
mackerel porbeagle.....	19	merou.....	103	mojarra blanca.....	341
mackerel scad.....	270	metalote.....	103	mojarra cantilena.....	341
mackerel scads.....	270	Mexican lamprey.....	9	mojarra cardinal.....	236
mackerel shark.....	19	Mexican mojarra.....	343	mojarra china.....	342
mackerel sharks.....	19	Mexican sailfish.....	263	mojarra de Casta.....	341
mackerels, Atka.....	374	Mexican snapper.....	326	mojarra de las aletas anarill- las.....	342
king of the.....	504	Mexican sole.....	230	mojarra de las piedras.....	381
snake.....	261	Michigan grayling.....	18	mojarra de ley.....	341
Mackinaw trout.....	59	midshipman.....	62	mojarra dorada.....	273
mademoiselle.....	345	midshipmen.....	487	mojarra garabata.....	337
mademoiselles.....	345	milkfish.....	40	mojarra negra.....	416
madregal.....	279, 280	milkfishes.....	40	mojarra plateada.....	416
mad tom.....	158	miller's thumb.....	385	mojarra prieta.....	330
mad toms.....	155	northern.....	386	mojarra raiado.....	415
mahogany snapper.....	327	southern.....	385	mojarra verde.....	419
maid.....	29	minnie.....	133	mojarras.....	340
mailed catfishes.....	160	minnow.....	116, 121, 133,	mojarras blancas.....	341
maire d'Amplora.....	166	Athabaska.....	175	mojarras de rio.....	416
makajiki.....	263	bigeye.....	134	mojarras de río.....	340
maki-maki.....	500	black.....	139	mojarras negras.....	329
Malma trout.....	61	blackhead.....	145	mojarras plateadas.....	332
malolo.....	199	blue-lined.....	145	mojarron.....	305
Mammoth Cave blindish.....	195	blunt-nosed.....	135	mola.....	503
mammy.....	146	bridled.....	121	monda.....	278
man-eaters.....	19	bronze.....	140	mongolian drummer.....	355
man-eater sharks.....	20	brook.....	145	mongrel buffalo.....	102
mangrove minnow.....	175	bullhead.....	145	monkeyfish.....	22
mangrove snapper.....	313, 325	chub.....	136	monkfish.....	22, 504
Manitoba whitefish.....	64	eastern mud.....	174	Montana grayling.....	66
Manitou darter.....	283	emerald.....	123	Monterey halibut.....	223
Manitoulin tullibee.....	64	fathead.....	145	Monterey roach.....	122
manjuai.....	48	Holbrook's top.....	185	Monterey Spanish mack- erel.....	257
manjuari.....	38	leather-sided.....	123, 137	moo-gad-ee.....	107
manjuda.....	248	long-mouthed.....	121	mooneye.....	40, 64
man-of-war fish, Portuguese.....	268	mangrove.....	175	southern.....	41
man-of-war fishes.....	268	Moose Lake.....	136	mooneye cisco.....	64
manta.....	32	mud.....	174	mooneyes.....	40
manta raja.....	29	northern straw-colored.....	134	moonfish.....	275
manta rala colorado.....	29	pearl.....	116	Moorish idol.....	362
many-spined sunfish.....	304	pike.....	175	Moorish idols.....	362
mapro.....	436	pin.....	109	Moose Lake minnow.....	136
mapo.....	440	plumbeus.....	136	moray, black.....	91, 92
marbled rockfish.....	314	pursy.....	181	common spotted.....	91
marblefish.....	603	river.....	176	spotted.....	93
margaret grunt.....	329	rosy.....	123	moray eel.....	93
margatfish.....	329	rosy-faced.....	124	morays.....	90, 91
maria molle.....	413	rotgut.....	146	more mouth bream.....	302
marian.....	236	spotted.....	145	morena pinta.....	90
maria prieta.....	332	spotted-tail.....	129	morena pintita.....	92
mariposa.....	216	star-headed.....	178	morena verde.....	92
marlin, barred.....	263	steel-colored.....	129	mosquitofish.....	185
black.....	263	straw-colored.....	132, 143	moss bass.....	297
giant black.....	263	smelt.....	143	mossbunker.....	44
striped.....	263	spot-tailed.....	132	mother-of-oels.....	473
marlinspikefish.....	263	spotted.....	145	mountain bass.....	305
marlinspikefishes.....	263	sucker-mouthed.....	142	mountain mullet.....	254
marsh bass.....	297	top.....	177, 178, 186	mountain mullets.....	254
Marston trout.....	81	variegated.....	181	mountain sucker.....	104
Martin pescador.....	506	minnows.....	112	mountain suckers.....	104, 105
Maryland darter.....	284	mud.....	174	mountain trout.....	298
masamacush.....	59	silvery.....	123	mousefish.....	504
maskinongy.....	172	top.....	184	mud bass.....	297, 303
matajuelo.....	235	variegated.....	181	mud blower.....	212
matajuelo blanco.....	356	minnows.....	112	mud cat.....	154, 155
matajuelo real.....	39	mud.....	174	mud chub.....	117, 302
matajuelos.....	356	silvery.....	123	mud creeper.....	446
mattowacca.....	42	top.....	184	muddabbler.....	175
Maxinkuckee darter.....	284	minny bass.....	298	mud dabs.....	226
Mayfish.....	176	misarkornak.....	474	mudfish.....	38, 192, 434
May sucker.....	112	mischuppaug.....	836	mud lamprey.....	9
mazaara.....	263	Mississippi alligator gar.....	38	mud minnow.....	174, 175
McCloud River rainbow trout.....	58			striped.....	174
Meidle's darter.....	286			mud minnows.....	174
medialuna.....	358			mud parrot.....	433
medium red salmon.....	54			mud perch.....	215, 303
				mud sucker.....	107

List of common names of fishes occurring in check list—Continued

Page	Page	Page			
mud sunfish.....	303	obispo.....	30	parrotfish.....	431, 433, 434
mufflejaw.....	385	oceanic bonito.....	258	blue.....	432
mufflejaws.....	385	ocean pipefish.....	243	green.....	432, 435
multichi.....	58	ocean sunfish.....	243	red.....	432
mullet.....	109, 110, 111	ocean tang.....	363	peru.....	360
black.....	105	ocean turbot.....	492	pascadillo del rey.....	252
blue.....	110	ocellated dory.....	232	pastor.....	268
blueback.....	253	ogac.....	210	patao.....	342
Califaver.....	253	ohia.....	36	peacockfishes.....	260
common.....	253	ohio garfish.....	36	peacock flounder.....	217
fantail.....	253, 254	ohio shad.....	42	pearlip sucker.....	112
fine-scaled.....	105	ohio sturgeon.....	35	pearlfish.....	486
jumping.....	111	ohua.....	495	pearl minnow.....	116
king.....	343	oilfish.....	20, 261	pearlsides.....	73
mountain.....	254	oil shark.....	14	peche-peche.....	99
of Utah Lake.....	105	ojanco.....	327	peche pretre.....	365
red.....	111, 343	old England hake.....	214	peck-nosed skate.....	25
redeye.....	254	old English hake.....	213	pega.....	448, 449
sea.....	350	old wench.....	491	pegador.....	448
snipnose.....	268	oldwife.....	276, 349, 434, 491	peixe agulha.....	195
striped.....	253	black.....	493	peixe cargo.....	12
trench.....	253	ollepa.....	495	peixe fonda.....	331
whirligig.....	253	omitu.....	273	peixe rey.....	246, 252
white.....	100, 253	ono.....	256	pelérin.....	20
mullets.....	252	Ontario longjaw.....	63	Pensacola snapper.....	328
king of the.....	205	oolachan.....	66	pespesca.....	248
mountain.....	254	oopu.....	440	perch.....	297-302, 304, 400
striped.....	252	ooro eels.....	78	American.....	281
white.....	253	opah.....	216	backelor.....	304
murumichog.....	175, 176, 181	open-mouthed grunt.....	330	black.....	298, 317, 323
muñeca.....	380	Oquassa trout.....	81	blanco.....	298
murcelago.....	409	orango filefish.....	495	blue.....	301, 424
muscalonge.....	172	orange rockfish.....	365	branch.....	302
muskallunge.....	172	orange-spotted sunfish.....	300	bride.....	297
muskelunge.....	172	Oregon charr.....	61	bridge.....	304
Musquau River whitefish.....	64	Oregon sturgeon.....	34	chain-sided.....	301
muttonfish.....	328, 342, 473	oreille noire.....	326	chinquapin.....	299, 304, 305
mykiss.....	56	orqueta.....	276	female.....	302
naccaysh.....	41	Oswego bass.....	297	flying.....	304
nailrod.....	42	ouachita darter.....	284	gold.....	301
naked goby.....	440	ouananiche.....	56	green.....	297, 298
namaycush.....	59	outatlibi.....	303	lake.....	281
namaycush salmon.....	59	outatlibi espagnol.....	308	log.....	283
Nassau grouper.....	311	ovate pampano.....	276	millpond.....	304
Navarro roach.....	121	oysterfish.....	424, 446, 487	mud.....	215, 303
needlefish.....	105, 198	Pacific bonito.....	268	pike.....	282
needlefishes.....	195	Pacific cod.....	210	pirate.....	215
ng-e-kaio-kalung.....	382	Pacific salmon.....	54	pond.....	302
nègre.....	312	Pacific volador.....	263	raccoon.....	281
negrofish.....	309	paddle.....	397	red.....	299, 300
nejorpallujak.....	214	paddlefish.....	36	red-bellied robin.....	299
Nelson's trout.....	50	paddlefishes.....	36	redeye.....	303
Nelson's whitefish.....	65	paintedtail.....	297	ringed.....	281
nerka.....	55	dotted.....	297	river.....	281
New England hake.....	214	pajaro.....	108	robin.....	302
Newfoundland shark.....	14	pale crappie.....	304	Sacramento.....	303
newlight.....	304	pale river bass.....	297	sand.....	302, 305
Niagara gudgeon.....	139	palometta.....	267, 276, 277	sea.....	307
Niangua darter.....	285	pampamito.....	276	silver.....	304, 305, 307, 340, 345
nibbler.....	424	pampamito.....	276	speckled.....	304
nigger chub.....	146	pampano.....	274, 276, 277	spotted.....	305
nigger dick.....	148	California.....	266	strawberry.....	304
niggerfish.....	146, 309	common.....	277	Suckley.....	304
niggerhead.....	107	great.....	277	sun.....	299, 300, 303, 305
nigrome.....	18	round.....	276, 277	tin.....	304
nine-eyes.....	8	panampos.....	269, 278	trout.....	214, 215, 298
nine-lined shanny.....	402	paneca.....	438	viviparous.....	409
nine-spined sticklebacks.....	238, 239	panagallo.....	280	warmouth.....	302
nissiue trout.....	58	papagallos.....	280	white.....	304, 305, 307, 352, 410
northern barracuda.....	255	papermouth.....	304	yawmouth.....	302
northern darter.....	291	parende.....	350, 360	yellow.....	281, 298, 299
northern miller's thumb.....	386	pargo.....	324, 328	perchmouth bream.....	302
northern pomois.....	302	pargo amarillo.....	326, 337	perches.....	281
northern roach.....	122	pargo colorado.....	328	American pike.....	282
northern straw-colored minnow.....	134	pargo criollo.....	328	log.....	282
northern striped gurnard.....	407	pargo de lo alto.....	328	river.....	281
northern sucker.....	106	pargo de raízero.....	327	viviparous.....	409
northern whiting.....	350	pargo guachenargo.....	328		
North River shad.....	42	pargo moreno.....	325	permilt.....	276, 277
Norway haddock.....	363	pargo negro.....	325	perrico.....	435
no-shee trout.....	58	pargo prieto.....	325	perro colorado.....	425
numbfish.....	27	pargos.....	326	perro perro.....	425
nurse shark.....	12, 22	parrot, mud.....	433	pesca blanca.....	137
nurse sharks.....	12	roseback.....	432	pesca pavone.....	18
				pesca vermiglia.....	367
				pescadillo del rey.....	252, 254
				pescadito.....	120

List of common names of fishes occurring in check list—Continued

Page	Page	Page	
pescado azul de dos colores.....	414	pipefish, ocean.....	243
pescado blanco de chapala.....	261	West Indian.....	241
pescado blanco de chalso.....	250	pipefishes.....	240
pescado colorado.....	348	piper.....	199
pescado del rey.....	252	pipon.....	98
pescados blancos.....	250	piquier.....	362
pescados del rey.....	246	pirate.....	215
pesce rey.....	252	pirate perch.....	215
pesca tondo.....	19	pirate perches.....	215
pesca toro.....	497	plaice.....	226
petite gueule.....	341	plain leatherjacket.....	277
petite jaquette.....	414	plaintall.....	261
petite scie.....	333	plateado.....	98
petitnègre.....	321	platefish.....	496
peto.....	256	pluma.....	337
petos.....	256	plumbeus minnow.....	136
pez clego.....	479	Plumier's pigfoot.....	370
pez de espada.....	23	pogge.....	395
pez de gallo.....	280	po-he-wa.....	110
pez del rey.....	247, 252	poison grouper.....	370, 371
pez de pluma.....	336	poison toadfishes.....	488
pez luna.....	503	poisson bleu.....	65, 152
pez mola.....	503	poisson de marais.....	38
pez puerco.....	492	poisson lune.....	216
pez puerco de piedra.....	491	poissons crappie.....	304
pez sierra.....	23	polo flounder.....	228
pez zorro.....	18	pollack.....	209
philosophie.....	363	pom.fret.....	265
phoebe.....	319	pomotis, northern.....	302
plicconou.....	111	red-tailed.....	299
picked dogfish.....	19, 21	pompano, Carolina.....	277
pickerel.....	282	Irish.....	342
barred.....	173	pollacks.....	209
eastern.....	173	pom.fret.....	265
little.....	173	pomotis, northern.....	302
pickerling.....	282	red-tailed.....	299
picuda.....	255	pompano, Carolina.....	277
picudilla.....	255	yellow.....	297
pigeon stingray.....	28	pont smelt.....	68
pigfish.....	335, 407	popeye.....	205
rock.....	407	poppypfish.....	206
sand.....	307	ponch, black-eared.....	299, 301
pigfishes.....	335	pond perch.....	302
plumbeus minnow.....	28	pond smelt.....	68
pigmy sculpin.....	386	porbeagle.....	19
pigmy sunfish.....	305	mackerel.....	19
Everglades.....	305	porbeagles.....	19
pike, American.....	173	porcupinefish.....	501
blue.....	282	hairy.....	501
common gar.....	36	porcupinedfishes.....	501
grass.....	173	porkfish.....	332
gray.....	282	porches.....	336
great.....	172	porgy.....	336, 349
great northern.....	173	grass.....	337
green.....	173	lolt-head.....	336
harvest.....	106	littlehead.....	337
Sacramento.....	114	littlemouth.....	337
sand.....	282	red.....	337
sea.....	198	saucereye.....	336
wall-eyed.....	282	shad.....	337
yellow.....	282	sheepshead.....	337
pike minnow.....	175	southern.....	337
pike perch.....	282	speckled.....	337
pike perches, American.....	282	three-tailed.....	358
pike shiner.....	121	whitebone.....	337
pikes.....	172	porte enseigne.....	361
gar.....	36	Porte Rican anchovy.....	51
pilot, black.....	413	portugals.....	360
cockeye.....	413	Portuguese butterfly.....	359
cow.....	415	Portuguese man-of-war fish.....	288
shark.....	270	possum ray.....	28
plotifish.....	65, 278, 279	postcroaker.....	349
plotifishes.....	339	Potomac shad.....	42
pinceo.....	43	pout, horned.....	153, 184
pinfish.....	337	pouts, horned.....	151
pinkfish.....	447	prickly fish.....	168
pink salmon.....	54	prickly bullhead.....	383
pin minnow.....	109	prickly skate.....	26
pinous dory.....	276	pristefish.....	365
pintado.....	257	puddingwife.....	421
pintano.....	415	pudiano.....	425
pintanos.....	414	pudiano verde.....	428
common.....	242	puerco espino.....	501
great.....	240		
green.....	242		

Page	Page	Page	
puffer.....	498, 499	puffer.....	498, 499
smooth.....	497	smooth.....	497
southern.....	498	southern.....	498
puffers.....	497	puffers.....	497
sharp-nosed.....	500	sharp-nosed.....	500
Fuget Sound pollack.....	209	Fuget Sound pollack.....	209
pug-nosed shiner.....	275	pug-nosed shiner.....	275
pumpkinseed.....	302	pumpkinseed.....	302
pumpkinseeds.....	297	pumpkinseeds.....	297
puraque.....	23	puraque.....	23
pursey minnow.....	181	pursey minnow.....	181
Putnam's darter.....	285	Putnam's darter.....	285
Pyramid Lake trout.....	57	Pyramid Lake trout.....	57
quasky.....	61	quasky.....	61
queenfish.....	353	queenfish.....	353
querimen.....	253	querimen.....	253
quiaquila.....	270	quiaquila.....	270
quiabira.....	277	quiabira.....	277
quiabirubas.....	320	quiabirubas.....	320
rabbitfish.....	261, 497, 502	rabbitfish.....	261, 497, 502
rabbitmouth sucker.....	112	rabbitmouth sucker.....	112
rabitubia.....	327	rabitubia.....	327
rabitubrias.....	327	rabitubrias.....	327
raccoon perch.....	281	raccoon perch.....	281
radiated shanny.....	470	radiated shanny.....	470
ragfish.....	289	ragfish.....	289
rainfishes.....	266	rainfishes.....	266
rainbow darter.....	291	rainbow darter.....	291
rainbowfish.....	191	rainbowfish.....	191
rainbow herring.....	67	rainbow herring.....	67
rainbow trout.....	58	rainbow trout.....	58
McCloud River.....	58	McCloud River.....	58
of fish-culturists.....	58	of fish-culturists.....	58
rainfish.....	180	rainfish.....	180
rainwaterfish.....	175	rainwaterfish.....	175
rasciera.....	365	rasciera.....	365
rasher.....	365	rasher.....	365
ratfish.....	33	ratfish.....	33
raton.....	256	raton.....	256
rat-tail, common.....	205	rat-tail, common.....	205
rat-tails.....	202	rat-tails.....	202
raven, sea.....	393	raven, sea.....	393
Rawson Lake herring.....	63	Rawson Lake herring.....	63
ray, brier.....	30	ray, brier.....	30
butterfly.....	25	butterfly.....	25
clear.....	25	clear.....	25
cow-nosed.....	31	cow-nosed.....	31
hedgehog.....	25	hedgehog.....	25
lady.....	30	lady.....	30
possum.....	25	possum.....	25
shark.....	22	shark.....	22
sharn.-nosed.....	31	sharn.-nosed.....	31
small electric.....	27	small electric.....	27
rays.....	10, 23	rays.....	10, 23
eagle.....	30, 31	eagle.....	30, 31
electric.....	27	electric.....	27
thick-tailed.....	28	thick-tailed.....	28
whip-tailed.....	28	whip-tailed.....	28
razorback.....	305	razorback.....	305
razor-backed buffalo.....	102	razor-backed buffalo.....	102
razorback sucker.....	108	razorback sucker.....	108
razorback suckers.....	108	razorback suckers.....	108
razorfish.....	430	razorfish.....	430
razorfishes.....	430	razorfishes.....	430
real swordfish.....	263	real swordfish.....	263
red bass.....	348	red bass.....	348
red-bellied bream.....	299, 300	red-bellied bream.....	299, 300
red-bellied dace.....	113	red-bellied dace.....	113
red-bellied robin perch.....	299	red-bellied robin perch.....	299
redbelly.....	299, 300, 302	redbelly.....	299, 300, 302
redbreast.....	299	redbreast.....	299
redbreast bream.....	299	redbreast bream.....	299
red drum.....	348	red drum.....	348
red drums.....	348	red drums.....	348
redeye.....	298, 302, 303	redeye.....	298, 302, 303
little.....	298	little.....	298

List of common names of fishes occurring in check list—Continued

Page	Page	Page			
red-eyed bream.....	303	robabilo de las aletas amarillas.....	308	rose trout.....	59
red-eyed sunfish.....	300, 303	robabilo de las aletas prietas.....	306	Ross's arctic salmon.....	60
redeye mullet.....	254	robalo.....	305	rosy dace.....	121
redeye perch.....	303	robalo prieto.....	306	rosy-faced minnow.....	124
red fallfish.....	127	robalos.....	305	rosy minnow.....	123
redfin.....	128, 131	robin.....	299, 302	rotgut minnow.....	146
compressed.....	128	Carolina flying.....	407	roughback.....	227
redfish.....	55, 318, 348, 363, 370	robin perch.....	209, 302	rough silverside.....	248
bull.....	348	red-bellied.....	299	round bass.....	304
California.....	426	rock.....	307	round buffalo.....	102
red goatfish.....	343	rock bass.....	297, 298	round herring.....	46
red grouper.....	312	common.....	303	round herrings.....	45
red guativera.....	309	southern.....	303	round pampano.....	276, 277
red hake.....	213	rock basses.....	303	round robin.....	270
red-headed bream.....	299	rock beauty.....	361	round stingray.....	29
red hind.....	311	rock cod, black and yellow.....	360	round sunfish.....	304
redhorse.....	109	black-banded.....	370	roundtail.....	114
big-toothed.....	111	brown.....	369	round whitefish.....	65
common.....	110	flesh-colored.....	369	roussettes.....	11
eastern.....	111	yellow-backed.....	369	rovaleto.....	248
short-headed.....	111	yellow-spotted.....	369	rovetto.....	261
Texas.....	110	rock cook.....	424	royal silver trout.....	56
redmouth buffalo fish.....	102	rock cooks.....	424	rubio volador.....	407
redmouth grunt.....	331	rock cusk.....	388	rudderfish.....	279
red-mouthed grunt.....	335	rockfish.....	176, 283, 305,	banded.....	268
red mullet.....	111, 343	black.....	367	black.....	267
red parrotfish.....	432	rockfishes.....	363, 364	ruff.....	302
red perch.....	299, 300	rock hind.....	308, 309, 310	runner.....	272, 277, 279
red porgy.....	337	rockling.....	208	blue.....	272
red rockfish.....	368	five-bearded.....	208	Russian cat.....	165
red rock trout.....	374	four-bearded.....	208	ruvetto.....	261
red roncador.....	348	silvery.....	208	sabalito.....	99
red roncadores.....	348	three-bearded.....	208	sablio.....	39, 40
red salmon, medium.....	54	rocklings.....	208	sabina.....	28
red sculpin.....	381	five-bearded.....	208	sable.....	262
red seafish.....	363	four-bearded.....	208	sablefish.....	373
red-sided bream.....	118	three-bearded.....	208	sablefishes.....	373
red-sided shiner.....	120	rocklings.....	208	sac-a-lait.....	177, 302, 304
red snapper.....	328	five-bearded.....	208	Sacramento cat.....	154
red-spotted sunfish.....	300	four-bearded.....	208	Sacramento chub.....	120
red-spotted trout.....	61	three-bearded.....	208	Sacramento perch.....	303
red sturgeon.....	35	rockling.....	208	Sacramento pike.....	114
red sucker.....	106	five-bearded.....	208	Sacramento River salmon.....	55
red sunfish.....	299	four-bearded.....	208	Sacramento squawfish.....	114
red-tailed balaos.....	199	three-bearded.....	208	Sacramento sturgeon.....	34
red-tailed bream.....	209	rock pigfish.....	407	Sacramento sucker.....	108
red-tailed pomotis.....	299	rock salmon.....	280	Saginaw Bay herring.....	62
redtail snapper.....	327	rock sea bass.....	318	sailfin, American.....	60
red-winged gurnard.....	407	rock shellfish.....	496	sailfinnur.....	362
red-winged sea robin.....	407	rock trout.....	304	sailfish.....	20, 103
reina.....	368	red.....	374	Florida.....	263
remora.....	449	Rocky Mountain bullhead.....	384	Mexican.....	263
Indian.....	448	Rocky Mountain whitefish.....	385	sailfishes.....	263
white-tailed.....	448	roncador.....	278	sailing sucker.....	103
remoras.....	448	little.....	349	sailorfish.....	103
requiem.....	15	red.....	349	sailors' choice.....	328, 335, 337
requim.....	15	yellow-finned.....	350	salerna.....	338, 339
rescacio.....	370	roncador ralado.....	330, 331	salmichi.....	132
rex sole.....	228	roncodores.....	344	salmon.....	54, 55
ribbonfish.....	282, 353	red.....	348	blueback.....	55
ringed perch.....	281	ronco.....	330	calico.....	54
Rio Grande trout.....	57	ronco amarillo.....	330	chinook.....	55
river bass.....	297, 298	ronco arara.....	330	coho.....	54
pale.....	297	ronco blanco.....	329	Columbia River.....	55
spotted.....	298	ronco carbonero.....	330	common Atlantic.....	56
streaked-cheeked.....	298	ronco condenado.....	330	dog.....	54
river carp, blunt-nosed.....	103	roncodina.....	349	Fraser River.....	55
river chub.....	103	ronco grande.....	330	hoopid.....	54
river drums.....	352	ronco prieto.....	329	humpback.....	54
river lamprey.....	8	ronco ronco.....	330	jack.....	282
river lampreys.....	9	roncos.....	329	klog.....	55
river minnow.....	176	rondanin.....	265	king of the.....	216
river perch.....	281	ronquil.....	452	landlocked.....	56
river perches.....	281	ronquils.....	451	le kal.....	54
river sucker.....	111	Roosevelt trout.....	59	medium red.....	54
river sunfish.....	299	roosterfish.....	290	Namaycush.....	59
river trout.....	282	rooter.....	102	Pacific.....	54
rosach.....	116, 299, 302, 305	roseback parrot.....	432	pink.....	54
harlequin.....	302	rosefish.....	363	quinnat.....	55
Monterey.....	122	rosefishes.....	363	rock.....	280
Navarro.....	121				
northern.....	122				
venus.....	122				
robabilo.....	306				

List of common names of fishes occurring in check list—Continued

Page	Page	Page			
salmon, Ross's arctic.....	60	scamp.....	316	segundo.....	272
Sacramento River.....	55	schoolmaster.....	325	Seneca Lake herring.....	62
sau-qui.....	65	Schuylkill cat.....	154	Seneca Lake smelt.....	62
sebago.....	66	scirenga.....	315	sennet.....	255
silver.....	54	scorpanudl.....	370	señorita.....	427, 428, 458
sockeye.....	55	scopene.....	371	señoritas.....	428
spring.....	55	scorpion.....	371, 487	serat imperial.....	370
sukkegh.....	55	scorpionfishes.....	363	sergeantfish.....	281, 305
tyee.....	55	scourfish.....	291	sergeantfishes.....	281
white.....	270, 297	sculpin.....	371	serra.....	171
of the Colorado.....	114	arctic.....	386	serrana.....	353
salmonete.....	343	black.....	387	serrana hispania.....	353
salmonete amarilla.....	343	brassy.....	386	serrano.....	318
salmon-formed growler.....	297	buffalo.....	381	sesele.....	305
salmon shark.....	19	common.....	386	sesi do lo alto.....	326
salmon trout.....	58	daddy.....	387	shad.....	42, 103, 304, 341
Kamchatka.....	56	eighteen-spined.....	388	Alabama.....	43
of Lake Southerland.....	57	European.....	386	American.....	42
salt-water gar.....	196	great.....	387	broad.....	341
salt-water trout.....	355	lake.....	389	common.....	42
San Bernardino trout.....	59	long-spined.....	386	Connecticut River.....	42
sand conger.....	454	pigmy.....	386	gizzard.....	46
sand dab.....	222	red.....	381	hickory.....	42, 46
sand darter.....	287, 288	yellow.....	381, 393	North River.....	42
sand darters.....	289	sculpins, scaled.....	375	of Lake Champlain.....	64
San Diego sole.....	231	scaleless.....	383	Ohio.....	42
sand diver.....	164	daddy.....	386	Potomac.....	42
sand eel.....	451, 485	scup, common.....	336	young.....	115
sandfish.....	318, 357	scuppaug.....	336	shad lamprey.....	8
sand lance.....	451	sea ape.....	18	shad porgy.....	337
Alaska.....	451	sea bass.....	317, 348	shadwaiter.....	65
sand lances.....	451	black.....	317	shads, gizzard.....	46
sand launce.....	451	gulf.....	318	shanny, Bos'e's.....	462
sand launces.....	451	of California, white.....	366	nine-lined.....	402
sand perch.....	302, 305	rock.....	318	radiated.....	470
sand pigfish.....	407	sea basses.....	308	shark, basking.....	20
sand pike.....	282	seabats.....	506	bay.....	15
sand roller.....	215	sea breams.....	265	bone.....	20
sand rollers.....	214	sea creams.....	265	bonnethead.....	17
sand shark.....	19	sea cat.....	149	bramble.....	22
sand sharks.....	19	sea catfish.....	148, 149	bullhead.....	10
sand stargazers.....	454	sea catchises.....	147	California angel.....	22
sand sturgeon.....	35	sea devil.....	32	cat.....	14
sand sucker.....	109, 350	sea devils.....	31, 508	cazon de playa.....	16
sandunche.....	39	sea dog.....	424	cod.....	21
sand whiting.....	350	sea drums.....	352	codfish.....	21
San Pedro fish.....	216	sea eel.....	88	common gray.....	15
sapo.....	486, 487	sea fish, red.....	363	cow.....	11
sardina.....	48, 98, 249	sea fox.....	18	cub.....	15
sardina blanca.....	100	sea goldfish.....	415	dog.....	13
sardina bocona.....	60	seahorse.....	244	dusky.....	16
sardina de españa.....	43	common American.....	244	East Indian basking.....	18
sardina de loy.....	43	seahorses.....	240	elephant.....	20
sardina escamuda.....	43	sea lamprey, great.....	8	fish.....	18
sardina machete.....	44	sea mink.....	350	fox.....	18
sardine, California.....	43	sea mullet.....	350	fresh-water.....	16
smooth.....	43	sea owl.....	397	gray.....	15
sardines, scaled.....	43	sea perch.....	307	great blue.....	17
smooth-headed.....	43	sea pike.....	196	hammerhead.....	18
true.....	43	sea poacher.....	395, 397	leopard.....	14
sargassumfish.....	504	sea poachers.....	393	little bullhead.....	22
sargo.....	337, 338	sea possum.....	25	longtail.....	18
sargo raiado.....	332, 338	sea raven.....	260, 303	mackerel.....	19
satinfin.....	129	sea ravens.....	393	Newfoundland.....	14
saucereye porgy.....	336	searer.....	257	nurse.....	12, 22
sauger.....	282	sea robin.....	407, 408	oil.....	14
sau-qui salmon.....	55	brown-winged.....	407	salmon.....	19
saurrels.....	270	Carolina.....	407	sand.....	19
sauries.....	197	lined.....	407	sharp-nosed.....	14
saury.....	197	red-winged.....	407	shovelhead.....	17
sauterie.....	277	small-scaled.....	407	shovelnose.....	18, 19
savannia.....	39	spotted.....	407	shovel-nosed.....	11
savola.....	262	striped.....	407	sleeper.....	22
sawbelly.....	42	web-fingered.....	407	small blue.....	16
sawfish, common.....	23	sea robins.....	407	soup-fin.....	14
sawfishes.....	23	sea serpent.....	88	spotted.....	14
scabbardfish.....	262	sea smelt.....	350	swell.....	12
scad.....	270	sea snail.....	399, 400	thornback.....	21
big-eyed.....	271	striped.....	400	tiger.....	14
dotted.....	270	sea snails.....	399	whale.....	18
mackerel.....	270	sea trout.....	355	wharf.....	16
scaled sardines.....	43	spotted.....	365	white.....	16
scaled sculpins.....	375	sea weed blenny.....	401	shark pilot.....	279
scaleless sculpins.....	383	sea whitefish.....	367	shark ray.....	22
		sebago salmon.....	56		

List of common names of fishes occurring in check list—Continued

	Page		Page	Page	
sharks.....	10	silverside minnow.....	119	snake blenny.....	471
angel.....	22	silversides.....	246, 247, 248	spotted.....	469
basking.....	20	silver trout.....	57	snake eels.....	84
bramble.....	22	royal.....	56	snakefish.....	163, 471
bullhead.....	10	silver whiting.....	351	snake mackerel.....	260
cat.....	11	silvery gar.....	196	snake mackerels.....	261
cow.....	11	silvery hairtail.....	262	snap mackerel.....	280
gray.....	12	silvery minnow.....	132, 143	snapper.....	280, 363
great white.....	20	silvery minnows.....	123	Bermuda silk.....	326
hammer-headed.....	17	silvery rocking.....	208	blackfin.....	326
mackerel.....	19	singingfish.....	487	dog.....	325
man-eater.....	20	single-bearded flyingfish.....	201	gray.....	325
nurse.....	12	sirago.....	447	Lane.....	327
sand.....	19	siscowet.....	60	mahogany.....	327
sleepers.....	21	skate.....	25	mangrove.....	313, 325
tectospondylous.....	21	barndoor.....	25	Mexican.....	326
thresher.....	18	bottailed.....	25	Pensacola.....	326
true.....	11	California big.....	26	red.....	326
whale.....	18	California common.....	26	redtail.....	327
sharksucker.....	448	clear-nosed.....	25	silk.....	328
sharp-finned flyingfish.....	200	common.....	25, 26	West Indian.....	328
sharp-nosed flyingfish.....	199	little.....	25	snappers.....	324
sharp-nosed puffers.....	500	peck-nosed.....	25	snapping mackerel.....	280
sharp-nosed ray.....	31	prickly.....	25	snipe eel.....	83
sharp-nosed shark.....	14	summer.....	25	snipe eels.....	83
sharp-tailed goby.....	441	winter.....	25	snipefish.....	246
sheatfish.....	355	skates.....	10, 24	snipefishes.....	245
sheepshead.....	338, 426	skil.....	373	snipnose mullet.....	268
lake.....	352	skilfishes.....	373	snook.....	305
three-banded.....	358	skimback.....	103	snowy grouper.....	311
sheepshead minnow.....	181	skipjack.....	42, 197, 251, 259, 279, 280	snub-nosed eels.....	78
sheepshead porgy.....	337	skip mackerel.....	280	soapfish.....	163, 316
sheepsheads.....	338	skipper.....	197	black.....	316
shellcracker.....	302	skipjack.....	196	soapfishes.....	316
shellfish.....	497	skittle-dog.....	19, 21	sobaco.....	492
rock.....	496	skowitz.....	54	sobacos.....	492
sheppy argentine.....	73	skooper.....	435, 436, 440	sockeye salmon.....	55
shielded darter.....	283	striped.....	436	soft-rayed fishes.....	39
shiner.....	115, 123, 128, 129,	sleepers.....	435	soldado.....	235
blunt-nosed.....	132	sleeper shark.....	22	soldierfish.....	291
goldthread.....	275	sleeper sharks.....	21	soldierfishes.....	234, 235
pike.....	116	slender halfbeak.....	198	sole, American.....	229
pug-nosed.....	275	slimefish.....	8	California.....	222
red-sided.....	120	slimer.....	487	gray.....	228
smelt.....	132	slippery dick.....	426, 485	lemon.....	228
spotted.....	138	slough bass.....	297	long-finned.....	228
swallowminnow.....	133	small black lamprey.....	9	Mexican.....	230
shining bass.....	304	small blindfish.....	106	rex.....	228
shoemaker.....	274, 279	small blue shark.....	16	San Diego.....	231
shore whiting.....	351	small catfish.....	154	somka.....	56
short barracuda.....	255	small devilfish.....	32	sorcerers.....	82
short-finned roach.....	122	small dolphin.....	205	soup-fin shark.....	14
short-headed redhorse.....	111	small electric ray.....	27	sourasap.....	601
short-nosed gars.....	37	small green sunfish.....	300	South American catfishes.....	168
short-nosed sturgeon.....	34	small guabin.....	180	southern chub.....	277
short-tailed menhaden.....	44	smallmouth buffalo.....	102	southern flounder.....	224
shovelhead shark.....	17	small-mouthed black bass.....	298	southern miller's thumb.....	385
shovel-nosed shark.....	11	small-scaled sea robin.....	407	southern mooneye.....	41
shovelnose shark.....	18, 19	smear dab.....	228	southern porgy.....	336
shovelnose sturgeon.....	35	smear dabs.....	222	southern puffer.....	498
white.....	35	smelt.....	132, 143	southern rock bass.....	303
shovelnose sturgeons.....	35	American.....	67	southern stingray.....	28
Shumard's darter.....	285	California blue.....	252	southern striped gurnard.....	407
sier.....	257	Cobessicontic.....	67	southern toothed herring.....	41
sterling.....	257	ghost.....	68	spadefish.....	38, 368
sierra.....	257	Kodiak.....	67	Spaniard.....	257
sierrita.....	196	little.....	252	Spanish flag.....	338, 363
silk snapper.....	328	pond.....	68	Spanish flags.....	308
Bermuda.....	326	sea.....	350	Spanish hogfish.....	425
silver bass.....	305, 307	Seneca Lake.....	62	Spanish ladyfish.....	425
silver carp.....	103	surf.....	67	Spanish mackerel.....	257
silver chub.....	117, 137	Wilton.....	67	black-spotted.....	257
silverfin.....	129, 132	smelt minnow.....	143	Monterey.....	257
hair-finned.....	275	smelt shiner.....	132	of England.....	258
silver gar.....	198	smelts.....	60, 67	sparada.....	409
silver hake.....	214	ghost.....	68	spawneater.....	132
silver jenny.....	341	surf.....	67	spearfish.....	103
silver lamprey.....	9	smooth blackfish.....	424	black.....	263
silver-mouthed minnow.....	142	smooth cabezon.....	300	true, of Japan.....	263
silver perch.....	304, 305, 307, 340, 345	smooth dogfish.....	13	spearfishes.....	263
silver salmon.....	54	smooth-headed sardines.....	43	spearfish sucker.....	450
silverside.....	247	smooth hound.....	13	spearing, ground.....	103
brook.....	251	smooth puffer.....	497	speck.....	287
fresh-water.....	247	smooth sardine.....	43	speckled bass.....	305
rough.....	248	smooth swellfish.....	497	speckled hen.....	297, 298
				speckled hind.....	312

List of common names of fishes occurring in check list—Continued

Page	Page
speckled perch.....	304
speckled porgy.....	337
speckled trout.....	60
of Lake Crescent.....	57
spet.....	255
spike.....	282
spinous dory.....	276
spinous gurnard.....	407
spinous lumpfish.....	398
spinyback blowfish.....	498
spinyboxfish.....	302
spiny dogfish.....	21
spiny eels.....	75
split-mouth sucker.....	112
spittail.....	116
spoonbill cat.....	36
spot.....	349
white.....	339
spottail.....	348
spot-tailed minnow.....	502
spotted balloonfish.....	297, 348
spotted bass.....	299
spotted bream.....	317
spotted cabrilla.....	257
spotted codling.....	213
spotted cybium.....	257
spotted-dinned sunfish.....	304
spotted grouper.....	311
spotted hake.....	213
spotted jewfish.....	312
spotted kelpfish.....	458
spotted lycodes.....	474
spotted mackerel.....	257
spotted minnow.....	145
common.....	91
spotted moray.....	93
spotted perch.....	305
spotted river bass.....	298
spotted rockfish.....	367
spotted sea robin.....	407
spotted sea trout.....	355
spotted shark.....	14
spotted shiner.....	138
spotted snake blenny.....	469
spotted squireague.....	355
spotted stargazer.....	454
spotted stingray.....	30
spotted sucker.....	107, 109
spotted suckers.....	109
spotted sunfish.....	304
spotted-tail minnow.....	120
spotted trout.....	305
of Lake Southerland.....	57
spotted trunkfish.....	497
spotted weakfish.....	355
sprat.....	44, 51
springfish.....	385
spring salmon.....	55
squaremouth.....	112
squaretail.....	269
squaretails.....	269
squato.....	22
squawfish.....	113, 114
Colorado River.....	114
Sacramento.....	114
squireague.....	355
spotted.....	355
squirrelfish.....	235, 318, 330
squirrelfishes.....	234, 235, 318
squirrel hake.....	213
squit.....	365
Stanley's whitefish.....	65
stargazer.....	454
little.....	385
spotted.....	454
stargazers.....	454
sand.....	454
star-headed minnow.....	178
starry flounders.....	226
steel-backed chub.....	146
steel-colored minnow.....	129
steelhead.....	68
Stewart White trout.....	59
stickleback, brook.....	238
common eastern.....	238
stickleback, four-spined.....	239
nine-spined.....	239
sticklebacks.....	237
nine-spined.....	238
stingaree.....	28, 31
stingray.....	28
California.....	31
long-tailed.....	31
round.....	29
southern.....	28
spotted.....	30
whip.....	28
stingrays.....	28
stite-tse.....	58
stockfish.....	214
stone bass.....	308
stone cat.....	155
stone cats.....	155
stonelugger.....	107
stoneroller.....	107, 146
stonerollers.....	148
Storer's chub.....	137
straw bass.....	297, 305
strawberry bass.....	302, 305
strawberry perch.....	304
straw-colored minnow.....	133, 134
streaked-cheeked river bass.....	298
streaked lead.....	298
striped anchovy.....	48
striped bass.....	297, 307
striped dace.....	140
striped grunt.....	329, 332
striped grunts.....	331
striped gurnard, northern.....	407
southern.....	407
striped marlin.....	263
striped mud minnow.....	174
striped mullet.....	253
striped mullets.....	232
striped sea robin.....	407
striped sea snail.....	400
striped sleeper.....	436
striped sucker.....	109
striped surfshl.....	411
studfish.....	179
sturgeon, common.....	34
green.....	34
lake.....	35
Ohio.....	35
Oregon.....	34
red.....	35
Sacramento.....	34
sand.....	35
short-nosed.....	34
shovelnose.....	35
white.....	34, 35
white shovelnose.....	35
sturgeons.....	33
shovelnose.....	35
sturgeon sucker.....	106
subotka.....	171
sucker.....	8, 103, 110, 449
biglip.....	106
black.....	106, 107
black winter.....	109
blue.....	146
bluehead.....	104
bootleg.....	107
brook.....	105
carp.....	103
chub.....	108, 109
Columbia River.....	108
common.....	105
creek.....	109
fine-scaled.....	105
flannel-mouthed.....	107
gourdseed.....	102
gray.....	106
harelip.....	112
hog.....	107
humpback.....	108
June.....	108
large-scaled.....	110
long-nosed.....	106
Lost River.....	108
sucker, May.....	112
Missouri.....	102
mountain.....	104
mud.....	107
northern.....	106
pearlip.....	112
rabbitmouth.....	113
razorback.....	108
red.....	106
river.....	111
Sacramento.....	106
sailing.....	103
sand.....	109, 350
spearfish.....	450
split-mouth.....	112
spotted.....	107, 109
striped.....	109
sturgeon.....	106
sweet.....	109
swordfish.....	449
Tahoe.....	107
Webug.....	108
white.....	105, 110, 111
white-nosed.....	109
winter.....	109
suckrel.....	102
sucker-mouthed buffalo.....	102
sucker-mouthed minnow.....	142
suckers.....	102
carp.....	103
hard-mouthed.....	105
mountain.....	104, 105
razorback.....	108
spotted.....	109
suckfish.....	488
suckingfish.....	448
Suckley perch.....	304
sukkoghe salmon.....	55
summer flounder.....	224
summer herring.....	42
summer skate.....	25
Sunapee trout.....	60
sunfish.....	20,
115, 274, 275, 298-304, 495, 503	
banded.....	304
big-eared.....	300
big-mouthed.....	302
big-nosed.....	299
black.....	302
black-banded.....	303
blackeye.....	288
black-tailed.....	300
bloody.....	300
blue.....	298, 301
blue-and-green.....	298
blue-mouthed.....	301
blue-spotted.....	298, 304
brilliant.....	300
buffalo.....	298
common.....	300, 302
deceptive.....	299
eared.....	300
Everglades pigmy.....	305
fresh-water.....	299
gilded.....	301
green.....	288
horned-eared.....	299
little.....	304
long-eared.....	299, 300
long-finned.....	304
long-scaled.....	300
many-splined.....	304
mud.....	303
ocean.....	503
orange-spotted.....	300
pigmy.....	305
red.....	299
red-eyed.....	300, 303
red-spotted.....	300
river.....	299
rock.....	304
round.....	304
small green.....	300
spotted.....	304
spotted-fin.....	304

List of common names of fishes occurring in check list—Continued

Page	Page
sunfish, wide-mouthed.....	30:
sunfish bass.....	303
sunfishes.....	297, 299
sunny.....	302
sun bass.....	302
sun perch.....	299, 300, 303, 305
sun trout.....	302
surfshifl, bigmouth.....	411
black.....	411
striped.....	411
wall-eyed.....	410
white.....	411
surfshifl.....	409
surf smelt.....	67
surf smelts.....	67
surf whiting.....	351
surgeon, blue.....	362
common.....	362
surgeonfishes.....	362
surmullet.....	343
Susan's darter.....	288
swago bass.....	298
swallow, black.....	452
swallow minnow shiner.....	133
swallows.....	95
black.....	452
sweet sucker.....	109
swego.....	298
swellbelly.....	498
swellfish.....	498, 499
smooth.....	497
swellfishes.....	498
swellingfish.....	498
swell shark.....	12
swelltoad.....	498, 502
swingtail.....	18
switchtail.....	35
swordfish.....	196
common.....	264
swordfishes.....	264
swordfish sucker.....	449
swordtail.....	103
sympbranchoid eels.....	76
tadpole cats.....	155
Tahoe sucker.....	107
tailor.....	280
tailor herring.....	42
tallywag.....	317
tambor.....	308, 407, 408, 499
tang.....	362
blue.....	362
tanbarhero.....	362
tangbrosme.....	471
tangs.....	362
tarentola.....	163
tarpon.....	39
tarpon fry.....	51
tarpons.....	39
tarpum.....	39
tasselled darter.....	287
tautog.....	424
tautoga.....	424
tautogs.....	424
tchaviche.....	55
tectospondylous sharks.....	21
tejpalcate.....	230
tenpounder.....	39
tenpounders.....	39
tessellated darter.....	290
tessellated darters.....	287
testar.....	489
tetard.....	436, 489
tête de roche.....	333
Texas redhorse.....	110
thick-tailed rays.....	23
thimble-eyed ling.....	213
thimble-eyed mackerel.....	258
thornback shark.....	21
thread eels.....	83
thread fleshfis.....	494
threadfin.....	266
threadfins.....	258
eight-threaded.....	256
three-angled trunkfishes.....	44
three-banded sheephead.....	358
three-bearded rockling.....	208
three-bearded rocklings.....	208
three-tailed porgy.....	358
thresher.....	18
thresher sharks.....	18
thunderpumper.....	352
thunderpumpers.....	352
tiburón.....	17
tiburón ballena.....	18
ticky-ticky.....	178, 185
tiger shark.....	14
tigrone.....	14
tilefish.....	357
timber crappie.....	304
timucu.....	195
tinker mackerel.....	258
timouth.....	304, 305
tifosa.....	273
tin perch.....	204
tinoreva.....	18
tiranete.....	262
tiranetes.....	262
tiru.....	163
toadfish.....	487, 498
toadfishes.....	487
poison.....	498
tobaccobox.....	25, 260, 302
tobacco trumpetfish.....	245
toeroe.....	354
togue.....	50
tomcod.....	210
California.....	210
tomcods.....	210
tomitate.....	331
tomtates.....	331
tonguefish.....	231
tonguefishes.....	230
toothed herring.....	40
southern.....	41
top minnow.....	177, 178, 185
Halibook's.....	185
top minnows.....	184
topo.....	273, 497
torpedo.....	27
California.....	27
great.....	27
torpedofish.....	27
toro.....	214
toter.....	107
totuava.....	356
treefish.....	369
trembler.....	27
trinch mullet.....	253
triggerfish.....	491
tripletail.....	323, 358
tripletails.....	323
trompetero.....	245
trout.....	54, 55, 297
alekey.....	212
aurora.....	60
blueback.....	61
of Lake Crescent.....	57
brook.....	60
of western Oregon, etc.....	58
brown.....	298
bull.....	61
coast-range.....	57
Colorado River.....	57
Columbia River.....	57
cut-throat.....	58
Dolly Varden.....	61
Dublin Pond.....	60
Eagle Lake.....	58
eastern brook.....	60
emerald.....	58
Evermann's.....	59
fresh-water.....	297
Gila.....	114
trout, golden, of south fork.....	59
of Kern River.....	59
of Sunapee Lake.....	60
of Volcano Creek.....	59
of western tributaries.....	59
of Kern River.....	59
gray.....	355
Great Lakes.....	59
great, of Lake Tahoe.....	57
greenback.....	57
green.....	297, 298
Jordan's.....	57
Kamchatka salmon.....	56
Kamloops.....	58
Kern River.....	58
Lac de Mabre.....	61
Lake Tahoe.....	57
long-headed, of Crescent	
Lake.....	57
Lower California.....	59
Mackinaw.....	59
Malma.....	61
Marston.....	61
McCloud River rainbow.....	58
mountain.....	298
Nelson's.....	59
nissues.....	58
no-shoe.....	58
Oquassa.....	61
Pyramid Lake.....	57
rainbow.....	58
of fish-culturists.....	58
red rock.....	374
red-spotted.....	61
Rio Grande.....	57
river.....	282
rock.....	373
Roosevelt.....	59
rose.....	59
royal silver.....	56
salmon.....	58
of Lake Southerland.....	57
salt-water.....	355
San Bernardino.....	59
sea.....	355
spotted.....	355
silver.....	57
silver.....	60
of Lake Crescent.....	57
spotted.....	305
of Lake Southerland.....	57
Stewart White.....	59
Sunapee.....	60
Truckee.....	57
Verde.....	114
Volcano Creek.....	59
Waha Lake.....	56
white.....	297, 298
yellowfin.....	57
Yellowstone.....	56
trout bass.....	298
trout perch.....	214, 215, 298
trucha.....	264
Truckee trout.....	57
true eels.....	77
true fishes.....	33
true sardines.....	43
true sharks.....	11
true spearfish of Japan.....	263
trumpetfish.....	245, 246
tobacco.....	245
unarmed.....	245
trumpetfishes.....	244
trunkfish.....	406
common.....	497
horned.....	497
spotted.....	497
trunkfishes.....	498
three-angled.....	497
tschaviche.....	54
tschawytscha.....	55
tswam.....	108
tulipi.....	64

List of common names of fishes occurring in check list—Continued

Page	Page	Page			
tullibee.....	64	web-fingered sea robin.....	407	whiting.....	214
Manitoulin.....	64	webug sucker.....	108	California.....	350
tuna, Allison's.....	260	welchman.....	235	Carolina.....	350
leaping.....	250	welshman.....	297	northern.....	350
tunnies.....	250	western lamprey.....	10	of Lake Winnipesaukee.....	64
great.....	259	West Indian lancelet.....	7	sand.....	350
tunny, little.....	258	West Indian pipefish.....	241	shore.....	361
turbot.....	491	West Indian snapper.....	328	silver.....	351
ocean.....	492	West Indies long-finned albacore.....	260	surf.....	361
Tuscmibia darter.....	293	whale shark.....	18	wide-mouthed sunfish.....	302
tusk.....	214	whale sharks.....	18	widowfish.....	366
tyee salmon.....	55	wharf shark.....	16	willow cat.....	152
tyrantfish.....	262	whiffs.....	218	Wilton smelt.....	67
ulke.....	388	Whipple's darter.....	292	windfish.....	115, 117
ulmina.....	98	whipsnake eel.....	88	windowpane.....	217
ulua.....	272	whip stingray.....	28	windowpanes.....	217
unarmed trumpetfish.....	245	whip-tailed rays.....	28	winninish.....	56
unernak.....	477	whirligig mullet.....	253	winter flounder.....	227
unicornfish.....	495	whitebait.....	41, 247, 248	winter flounders.....	227
unspotted balloonfish.....	502	white bass.....	297, 303, 307	winter skate.....	25
vaagmør.....	216	white-bill.....	43	winter sucker.....	109
vaca.....	321	whitebone porgy.....	337	black.....	109
vacas.....	320	white cat.....	152, 153	wolf eel.....	472
vacuocua.....	345	white crappie.....	304	wolfish.....	171, 472
Vahl's lycodes.....	474	white croaker.....	353	wood bass.....	298
vaqueta de los colores.....	361	white-eye.....	282	worm eels.....	82
variegated goby.....	446	whitefish.....	44, 64, 114	wreckfish.....	308
variegated minnow.....	181	Bear Lake.....	65	wreckfishes.....	308
Venus roach.....	122	Bonneville.....	65	wrymouth.....	471
Verde trout.....	114	common.....	65	xurel.....	272
verrugata.....	351	Coulter's.....	65	xurel de castilla.....	276
verrugate.....	349	humpback.....	65	yaito.....	259
verrugato.....	349	Kennicott's.....	65	yawmouth perch.....	302
viajúca.....	419	Labrador.....	64	yellow angel.....	361
victorfishes.....	258	Lake Erie.....	65	yellow-backed rock cod.....	369
vicudá.....	255	Lake Superior.....	65	yellow bass.....	207, 298, 307
vieja.....	432, 434	Lauretta.....	62	yellowbelly.....	299, 302
vieja colorada.....	433	Manitoba.....	64	yellow bream.....	302
vieja muger.....	433	Monominee.....	65	yellow caraux.....	273
viejas.....	431	Musquau River.....	64	yellow cat.....	153, 155
Virginia lancelet.....	7	Nelson's.....	65	yellow-finned albacore.....	260
viuya.....	366	Rocky Mountain.....	65	yellow-finned albacores.....	260
viviparous perch.....	409	round.....	65	yellow-finned grouper.....	311, 313
viviparous perches.....	409	see.....	357	yellow-finned roncador.....	350
volador.....	202, 203, 400	Stanley's.....	65	yellowfin trout.....	57
Pacific.....	263	Yellowstone.....	65	yellowfish.....	309, 374
volantín.....	272	whitefishes.....	61, 64	yellow goatfish.....	343
Volcano Creek trout.....	59	white grunt.....	331	yellow grouper.....	314
voraz.....	328	white hake.....	213	yellow grunt.....	330
Wachna cod.....	210	white lake bass.....	307	yellow jack.....	272
Waha Lake trout.....	56	white mouth drummer.....	340	yellow mackerel.....	27, 273
walleye.....	282	white mullet.....	109, 253	yellow mouth.....	356
wall-eyed herring.....	42	white mullets.....	253	yellow Ned.....	281
wall-eyed pike.....	282	white-nosed sucker.....	109	yellow perch.....	281, 298, 299, 302
wall-eyed pollack.....	209	white perch.....	304, 305, 307, 352, 410	yellow pike.....	282
wall-eyed surfish.....	410	white salmon.....	279, 297	yellow pondporch.....	297
warmouth.....	302	of the Colorado.....	114	yellow sculpin.....	381, 393
black.....	302	white sea bass of California.....	356	yellow-spotted rock cod.....	369
warmouth bream.....	302	white shark.....	16	Yellowstone trout.....	56
warmouth perch.....	302	white sharks, great.....	20	Yellowstone whitefish.....	65
waterbelly.....	64	white shovelnose sturgeon.....	35	yellowtail.....	278, 279, 327, 345, 349
wawwhunnekesuog.....	257	white spot.....	339	yellow-tailed croaker.....	350
weakfish, bastard.....	355	white sturgeon.....	34, 35	yellowtail rockfish.....	364
common.....	355	white sucker.....	105, 110, 111	young grunt.....	331
spotted.....	355	white surfish.....	411	young shad.....	115
weakfishes.....	353	white-tailed remora.....	448	zapatero.....	277

LIST OF SCIENTIFIC NAMES

	Page		Page
<i>abacurus</i> , <i>Eleotris</i>	437	<i>Acanthocottus patris</i>	389
<i>aboti</i> , <i>Syngnathus</i>	240	<i>profundorum</i>	392
<i>abotti</i> , <i>Anchovia</i>	60	<i>scorpoides</i>	386
<i>Anchoviella</i>	50	<i>scorpius</i>	386
<i>Hydrophox</i>	127	<i>sellaris</i>	388
<i>Notropis chalybeus</i>	127	<i>Acanthocybidae</i>	256
<i>Osmerus</i>	67	<i>Acanthocybium</i>	256
<i>mordax</i>	67	<i>petus</i>	256
<i>abbreviata</i> , <i>Chimæra</i>	33	<i>solandri</i>	256
<i>abbreviatus</i> , <i>Bathyalopex</i>	33	<i>Acanthoderma</i>	261
<i>Nauclerus</i>	278	<i>temmincki</i>	261
<i>Abesona</i>	409	<i>acanthoderma</i> , <i>Thyrsites</i>	261
<i>aurora</i>	409	<i>acanthognathus</i> , <i>Lophodolus</i>	509
<i>trowbridgii</i>	409	<i>Acantholebius</i>	373
<i>aberrans</i> , <i>Hypoplectrus</i>	321	<i>Acantholepis</i>	68
<i>Hypoplectrus unicolor</i>	321	<i>Acantholiparis</i>	401
<i>Liopropoma</i>	316	<i>opercularis</i>	401
<i>Perca</i>	316	<i>Acanthonotus</i>	75
<i>abilgaardii</i> , <i>Sparisoma</i>	432	<i>nasus</i>	75
<i>Sparus</i>	432	<i>quadricornis</i>	497
<i>abilgaardii</i> , <i>Scarus</i>	431	<i>Acanthonus</i>	483
<i>Ablennes</i>	197	<i>armatus</i>	483
<i>hiangs</i>	197	<i>spinifer</i>	483
<i>Aboma</i>	443	<i>Acanthophacelus</i>	191
<i>chiquita</i>	443	<i>melenzonus</i>	191
<i>etheostoma</i>	443	<i>acanthophorus</i> , <i>Serranus</i>	317
<i>luctuosa</i>	443	<i>acanthopoma</i> , <i>Draconetta</i>	450
(<i>Abramis</i>) <i>baileatus</i> , <i>Cyprinus</i>	117	<i>Acanthopteryx</i>	232
<i>smithii</i> , <i>Cyprinus</i>	41	<i>Acanthosoma</i>	503
<i>Abramis boscii</i>	350	<i>carinatum</i>	503
<i>versicolor</i>	115	<i>Acanthostracion</i>	497
(<i>Abramis</i>) <i>baileatus</i> , <i>Cyprinus</i>	118	<i>polygonius</i>	497
<i>Abudeidu</i>	414	<i>Acanthuridae</i>	362
<i>analogus</i>	415	<i>Acanthurus</i>	392
<i>declivifrons</i>	415	<i>allua</i>	363
<i>marginatus</i>	415	<i>bahianus</i>	363
<i>taurus</i>	415	<i>brevis</i>	362
<i>troeschelli</i>	415	<i>broussoneti</i>	362
<i>abyssicola</i> , <i>Coregonus</i>	66	<i>cæruleus</i>	362
<i>Prosopium</i>	66	<i>crestonis</i>	363
<i>Raja</i>	26	<i>ctenodon</i>	363
<i>abyssorum</i> , <i>Nematonurus</i>	204	<i>glaucopareius</i>	363
<i>acadian</i> , <i>Cottus</i>	393	<i>hepatus</i>	362
<i>acadianus</i> , <i>Glyptocephalus</i>	228	<i>hirudo</i>	362
<i>Acantharchus</i>	303	<i>phlebotomus</i>	362
<i>pomotis</i>	303	<i>striatus</i>	363
<i>Acanthemblemaria</i>	465	<i>strigosus</i>	363
<i>spinosa</i>	465	<i>subarmatus</i>	362
<i>Acanthenchelys</i>	89	<i>tractus</i>	363
<i>Acanthenchelys</i>	89	<i>triostegus</i>	362
<i>spinicauda</i>	89	<i>zebra</i>	362
<i>Acanthias</i>	21	<i>Acaris adpersum</i>	418
<i>americanus</i>	21	<i>bartoni</i>	419
<i>suckleyi</i>	21	<i>ceruleopunctata</i>	423
<i>vulgaris</i>	21	<i>pinima</i>	332
<i>acanthias</i> , <i>Monoceratias</i>	508	<i>pitamba</i>	327
<i>Squalus</i>	21	<i>rectangularis</i>	421
<i>Acanthicus canensis</i>	161	<i>tetramerus</i>	423
<i>Acanthidium</i>	21	<i>acara-pinima</i> , <i>Pristipoma</i>	333
<i>calceus</i>	21	<i>Acathorhinus</i>	21
<i>pusillum</i>	21	<i>accensus</i> , <i>Plectropoma</i>	321
<i>Acanthinion</i>	276	<i>accensus</i> , <i>Hypoplectrus unicolor</i>	321
<i>acanthistius</i> , <i>Bodianus</i>	309	<i>Accipenser macrostomus</i>	35
<i>Enneistius</i>	309	<i>obtiensis</i>	35
<i>Acanthoclinus</i>	457	<i>serotinus</i>	35
<i>chaperi</i>	457	<i>acclinidens</i> , <i>Cyclothona</i>	72
<i>Acanthocottus</i>	386	<i>acclivis</i> , <i>Larimus</i>	344
<i>aneus</i>	386	<i>Acedia</i>	231
<i>bubalis</i>	386	<i>nebulosa</i>	231
<i>greenlandicus</i>	387	<i>Acentrolophus</i>	267
<i>labradoricus</i>	388	<i>maculatus</i>	267
<i>laticeps</i>	388	<i>maculosus</i>	267
<i>mitchilli</i>	386	<i>Aceratilidae</i>	511
<i>mucosus</i>	387	<i>acervum</i> , <i>Cyblum</i>	267
<i>ocellatus</i>	387	<i>Acharina</i>	229
<i>octodecemspinosis</i>	388	<i>achigan</i> , <i>Bodianus</i>	298

List of scientific names—Continued

	Page		Page
Achiridae.....	229	Acomus.....	105
Achiroplichthys typus.....	89	(Acomus) generosus, Catostomus.....	104
Achirus.....	229	griseus, Catostomus.....	107
achirus.....	229	guzmaniensis, Catostomus.....	104
barnharti.....	230	lactarius, Catostomus.....	107
comifer.....	230	acoupa, Chelodipterus.....	354
fasciatus.....	229, 230	Cynoscion.....	354
fonsecensis.....	230	Acrochelius.....	112
inscriptus.....	229	alutaceus.....	112
klunzingeri.....	229	acrolepis, Macrurus.....	203
lineatus.....	229, 230	Nematonurus.....	203
mazatlanus.....	230	acronotus, Carcharhas.....	16
ornatus.....	231	Squalus.....	16
scutum.....	229	Acropocilia.....	190
achirus, Achirus.....	229	(Acropocilia) tridens, pocilia.....	190
Pleuronectes.....	229	Acronurus.....	362
Achrophichthys.....	89	caruleatus.....	362
typus.....	89	carneus.....	362
Acipenser.....	33	fuscus.....	362
acutirostris.....	34	nigriculus.....	363
aleutensis.....	34	Acrotidae.....	269
(Antaeus) agassizii.....	34	Acrotus.....	269
alexandri.....	34	willoughbyi.....	269
buffalo.....	35	Acteis.....	460
cinchonni.....	35	culebrae.....	460
oligolepis.....	34	moorei.....	460
ayresi.....	34	Actinochir.....	401
bairdi.....	34	major.....	401
brachyrhynchus.....	34	Actinopterl.....	33
brevirostris.....	34	aculeata, Chrysophrys.....	336
brevirostrum.....	34	aculeatus, Chelmon.....	359
carbonarius.....	35	Clinus.....	470
caryi.....	34	Etomopterus.....	21
dekayi.....	34	Gastrostoeus.....	237
fulvescens.....	33, 35	Haleichthys.....	507
girardi.....	34	Lophius.....	507
glabra.....	34	Prognathodes.....	859
halops.....	34	aculeatus (oculatus), Serranus.....	328
helops.....	34	aculeatus, Gasterosteus.....	239
holbrookii.....	33	acuminata, Halecola.....	46
huso.....	35	Muraena.....	85
(Huso) anasimos.....	35	acuminatus, Etrumeus.....	46
anthracinus.....	35	Grammistes.....	353
atelaaspis.....	36	Myrichthys.....	85
copel.....	35	Pareques.....	353
kirtlandi.....	37	acuminatus umbrosus, Eques.....	353
lamarilli.....	35	Acus.....	243
megalaspis.....	35	squoroenus.....	243
nertinianus.....	35	heckeli.....	244
paranaesimus.....	35	martinicensis.....	243
rafinesquii.....	35	maxima squamosa viridis.....	36
richardsoni.....	35	pellucidus.....	244
rosarium.....	35	acus, Atopichthys.....	34
kennicotti.....	34	Gymnotus.....	101, 486
levis.....	35	Sphyraena.....	196, 197
lagenarius.....	36	Syngnathus.....	240
leontel.....	34	Tylosurus.....	197
lesueuri.....	34	acuta, Loricaria.....	161
lopeletsi.....	35	Myliobatis.....	31
macrorhinus.....	34	Ophismoma.....	79
macrostomus.....	35	Perca.....	281
maculosus.....	35	acute scabra, Lophius.....	505
medirostris.....	34	acuticeps, Oligocottus.....	391
microrhynchus.....	34	Oxycottus.....	391
milberti.....	34	acutirostris, Acipenser.....	34
mitchilli.....	34	Corvina.....	346
ohiensis.....	35	Ichthypodus.....	84
oxyrhinchus.....	34	Lutjanus.....	326
oxythynchus.....	34	Parepinophelus.....	315
platyryynchus.....	35	Pristis.....	23
putnami.....	34	Serranus.....	315
rhynchosurus.....	35	acutirostris late, Cerna.....	315
rubricundus.....	35	acutiventris, Alfaro.....	198
rupertianus.....	35	Acutomentum.....	365
ruthenus.....	34	alatum.....	366
schypa.....	34	brevispine.....	366
stellatus.....	34	clavilatum.....	366
storeri.....	34	eigenmanni.....	366
sturio.....	34	entomelas.....	366
transmontanus.....	33, 34	hopkinsi.....	366
vultharius.....	34	macdonaldi.....	366
Acipenseridae.....	33	ovale.....	366
acipenserinus, Agonus.....	394	prorigerum.....	366
Podothecus.....	394	rufum.....	366
ackleyi, Raja.....	25	wilsoni.....	366

List of scientific names—Continued

	Page		Page
acutum, <i>Cichlasoma</i>	422	affine, <i>Myctophum</i>	169
<i>Hæmulon</i>	329	<i>Plectropoma</i>	321
scutus, <i>Exocetus</i>	199	<i>Scopelus</i>	169
<i>Fodiator</i>	199	affinis, <i>Algancea</i>	116
<i>Scarus</i>	434	<i>Argyropelecus</i>	74
Synodus.....	164	<i>Atherinops</i>	252
<i>Thorichthys</i>	422	<i>Atherinopsis</i>	252
adamantinus, <i>Litholepis</i>	38	<i>Auchenipterus</i>	457
Adinia.....	180	<i>Carapus</i>	486
<i>multifasciata</i>	180	<i>Caulophilus</i>	357
adinia, <i>Fundulus</i>	179	<i>Centroponus</i>	306
Adioryx.....	236	<i>Cheilodipterus</i>	296
<i>suborbitalis</i>	236	<i>Chilomycterus</i>	502
adirondacus, <i>Salmo</i>	59	<i>Chimaera</i>	33
adobe, <i>Agosia</i>	141	<i>Clinostomus</i>	121
<i>Apococe</i>	141	<i>Cremnochites</i>	457
Adonis.....	460	<i>Exocoetus</i>	201
<i>pavoninus</i>	400	<i>Exonotus</i>	201
adscensionis, <i>Trachinus</i>	310	<i>Fierasfer</i>	486
adscensionis, <i>Epinephelus</i>	310	<i>Gila</i>	114
adspersa, <i>Parapetenia</i>	418	<i>Heros</i>	422
<i>Rhinoptera</i>	31	<i>Heterandria</i>	185
adspersum, <i>Acaris</i>	418	<i>Hypoplectrus unicolor</i>	321
adspersus, <i>Labrus</i>	424	<i>Isopisthus</i>	353
<i>Tautogolabrus</i>	424	<i>Lucania</i>	175
adusta, <i>Corvina</i>	347	<i>Pimeledous</i>	162
adustus, <i>Couesius</i>	137	<i>Pseudochelodon</i>	96
<i>Eupomacentrus</i>	414	<i>Psychichthys</i>	33
<i>Gobiesox</i>	489	<i>Stomias</i>	69
<i>Julidio</i>	428	<i>Syngnathus</i>	242
<i>Ophiocion</i>	347	<i>Syrietes</i>	242
<i>Pomacentrus</i>	414	<i>Thorichthys</i>	422
<i>Pseudojulis</i>	428	<i>Thynnus</i>	258
Æglefinus.....	210	agassizii, <i>Careliopsis</i>	400
<i>Illoai</i>	210, 211	<i>Iplops</i>	171
æglefinus, <i>Gadus</i>	210, 211	agassizii, <i>Apincenser (Anticeus)</i>	34
<i>Melanogrammus</i>	211	<i>Alepocephalus</i>	51
Ælurichthys.....	147	<i>Bathysaurus</i>	164
<i>isthmenis</i>	148	<i>Brama</i>	265
<i>longispinis</i>	148	<i>Callionymus</i>	450
<i>panamensis</i>	148	<i>Careliopsis</i>	400
<i>scutatus</i>	148	<i>Chlorophthalmus</i>	165
ænea, <i>Cichla</i>	303	<i>Chologaster</i>	194
æneolus, <i>Notropis</i>	143	<i>Cratinus</i>	317
æneus, <i>Acanthocottus</i>	386	<i>Cylindrosteus</i>	37
<i>Astyanax</i>	96	<i>Dicromita</i>	480
<i>Cottus</i>	388	<i>Forbesella</i>	194
<i>Pimeodus</i>	155	<i>Holconotus</i>	411
<i>Tetragonopterus</i>	96	<i>Hyperprosopon</i>	410
ænigmaticus, <i>Icosteus</i>	269	<i>Iplops</i>	171
æpyptera, <i>Ammocetes</i>	10	<i>Leptochilichthys</i>	53
æquatoris, <i>Talismania</i>	53	<i>Liparis</i>	400
Æquidens.....	423	<i>Pimephales</i>	145
<i>ceruleopunctata</i>	433	<i>Salmo</i>	60
æquidens, <i>Cilius</i>	437	<i>Salvelinus</i>	60
<i>Otolithus</i>	356	<i>Scorpena</i>	371
<i>Prionodes</i>	319	<i>Tocichthys</i>	410
<i>Serranus</i>	319	<i>Agenelosus</i>	169
æquoreus, <i>Acus</i>	243	<i>brevifilis</i>	159
<i>Syngnathus</i>	243	<i>caucanus</i>	169
æreus, <i>Sebastodes</i>	367	<i>aggregatus, Cymatogaster</i>	400
æsculapius, <i>Alepidosaurus</i>	171	<i>agilis, Gadus</i>	209
<i>Plagiodon</i>	171	<i>Agonidae</i>	393
æsopus, <i>Boleosoma</i>	287	<i>Agonostoma globiceps</i>	255
æstivalis, <i>Clupea</i>	42	<i>micros</i>	254
<i>Gobio</i>	138	<i>nasutum</i>	254
<i>Macryhypsos</i>	138	<i>percoides</i>	254
<i>Pomolobus</i>	42	<i>Agonostomus</i>	254
æstivalis marconis, <i>Hybopsis</i>	138	<i>salvinii</i>	254
æstuarium, <i>Paralichthys</i>	138	<i>telfairii</i>	254
æthalion, <i>Citharichthys</i>	224	<i>Agonus</i>	395
æthalurus, <i>Carcharias</i>	17	<i>acipenserinus</i>	394
Æthopora.....	168	<i>anne</i>	394
<i>effulgens</i>	168	<i>barkani</i>	394
<i>lucida</i>	168	<i>cataphractus</i>	395
Ætobatidae.....	30	<i>decagonus</i>	394
Ætobatis.....	30	<i>dodecaedron</i>	394
Ætobatus.....	30, 31	<i>dodecaedrus</i>	394
<i>californicus</i>	31	<i>vulsus</i>	395
<i>frémivillii</i>	31	<i>Agosia</i>	140
<i>goodiei</i>	31	<i>adobe</i>	141
<i>laticeps</i>	30	<i>chrysogaster</i>	140
afer, <i>Alphestes</i>	312	<i>falcata</i>	141
<i>Epinephelus</i>	312	<i>klamathensis</i>	141
<i>Gymnothorax</i>	91	<i>metallica</i>	140

List of scientific names—Continued

	Page		Page
<i>Agosia novemradiata</i>	141	<i>Albula</i>	39
<i>robusta</i>	140	<i>conorhynchus</i>	30, 40
<i>umatilla</i>	141	<i>erythrocheilus</i>	39
<i>yarwoi</i>	141	<i>forskali</i>	39
<i>Ariosphyraena</i>	255	<i>forsteri</i>	40
<i>aguia-bonita</i> , <i>Salmo</i>	50	<i>goreensis</i>	40
<i>Salmo mykiss</i>	50	<i>neoguinaica</i>	40
<i>aguil</i> , <i>Trisoptropis</i>	314	<i>parae</i>	40
<i>aguilha</i> , <i>Peixe</i>	185	<i>plumieri</i>	40
<i>Anila</i>	82	<i>rostrata</i>	40
<i>egmontis</i>	82	<i>seminuda</i>	40
<i>Ahyannodontophis</i>	94	<i>vulpes</i>	39
<i>stigmatodus</i>	94	<i>albula</i> , <i>Mugil</i>	253
<i>algula</i> , <i>Lachnolaimus</i>	425	<i>Albulidae</i>	39
<i>Allurichthys</i>	147	<i>Albuloides</i>	39
<i>pinnimaculatus</i>	148	<i>albus</i> , <i>Bryttus</i>	302
<i>allurus</i> , <i>Pimeledous</i>	153	<i>album</i> , <i>Hæmulon</i>	329
<i>almara</i> , <i>Macrodon</i>	99	<i>Moxostoma</i>	110
<i>aix</i> , <i>Pallasina</i>	394	<i>Alburnellus</i>	123
<i>alabame</i> , <i>Alosa</i>	43	<i>altipinnis</i>	127
<i>Claricola</i>	202	<i>amoenus</i>	123
<i>Etheostoma whipplei</i>	292	<i>argo</i>	124
<i>Notropis</i>	126	<i>dilectus</i>	123
<i>alalonga</i> , <i>Bothrocampsis</i>	477	<i>jaculus</i>	123
<i>Scomber</i>	200	<i>jemezanus</i>	125
<i>Scombrus</i>	260	<i>matutinus</i>	126
<i>alalunga</i> , <i>Germo</i>	200	<i>micropteryx</i>	124
<i>Scomber</i>	260	<i>percobromus</i>	124
<i>alascana</i> , <i>Asterotheca</i>	306	<i>simus</i>	135
<i>alascanus</i> , <i>Ammodytes</i>	451	<i>Alburnops</i>	133
<i>Argyrosomus</i>	62	<i>blennius</i>	133
<i>Leucichthys</i>	62	<i>heterodon</i>	130
<i>Sebastolobus</i>	364	<i>illicebrous</i>	134
<i>Xenochirius</i>	306	<i>longirostris</i>	135
<i>alata</i> , <i>Fissala</i>	403	<i>nubilis</i>	144
<i>alatus</i> , <i>Arius</i>	148	<i>plumbeolus</i>	128
<i>Lampanyctus</i>	167	<i>saludanus</i>	132
<i>Nemipteraster</i>	167	<i>shumardi</i>	134
<i>Prionotus</i>	408	<i>taurocephalus</i>	145
<i>Alausa</i>	42	<i>Alburnus armabilis</i>	121
<i>Alausella</i>	41	<i>americanus</i>	350
<i>Alazon</i>	190	<i>dilectus</i>	125
<i>alba</i> , <i>Rogenia</i>	41	<i>formosus</i>	129
<i>albacauda</i> , <i>Echeneis</i>	448	<i>lepidulus</i>	125
<i>Albacora</i>	259	<i>lineolatus</i>	135
<i>albacores</i> , <i>Nothunnus</i>	260	<i>megalops</i>	125
<i>Scomber</i>	260	<i>nitidus</i>	123
<i>Albatrossia</i>	204	<i>oligaspis</i>	125
<i>pectoralis</i>	204	<i>rubellus</i>	123
<i>albatrossis</i> , <i>Osmerus</i>	67	<i>rubrifrons</i>	124
<i>Thaumelichthys</i>	67	<i>socius</i>	124
<i>albatrossus</i> , <i>Macrourus</i>	206	<i>umbratilis</i>	125
<i>albeoelus</i> , <i>Chirlope</i>	136	<i>zonatus</i>	123
<i>Luxilus</i>	120	<i>alburnus</i> , <i>Perca</i>	350
<i>Notropis</i>	136	<i>albus</i> , <i>Atherinichthys</i>	251
<i>Notropis megalops</i>	129	<i>Centropomus</i>	307
<i>albescons</i> , <i>Echenesis</i>	449	<i>Coregonus</i>	62, 66
<i>Paraliparis</i>	404	<i>Coregonus clupeaformis</i>	65
<i>Remorina</i>	449	<i>Gymnotus</i>	101
<i>albicaudus</i> , <i>Auchenipterus</i>	457	<i>Eriscion</i>	356
<i>albidactylus</i> , <i>Excetrus</i>	202	<i>Lepisosteus</i>	37
<i>albidum</i> , <i>Hemulon</i>	329	<i>Otolithus</i>	356
<i>albidus</i> , <i>Gadus</i>	214	<i>Parascaphirhynchus</i>	35
<i>Labrax</i>	307	<i>Ptychostomus</i>	110
<i>Ostmerus</i>	163	<i>Alcidea</i>	375
<i>Pimeledous</i>	150, 153	<i>thoburni</i>	375
<i>Ptychostomus</i>	110	<i>Aldrovandia</i>	75
<i>albifimbria</i> , <i>Scorpaena</i>	371	<i>goodei</i>	75
<i>albifrons</i> , <i>Gymnotus</i>	101	<i>gracilis</i>	75
<i>albigutta</i> , <i>Cathetostoma</i>	454	<i>macrochir</i>	75
<i>Kathestostoma</i>	454	<i>palilda</i>	75
<i>Paraleichthys</i>	224	<i>rostrata</i>	75
<i>albiguttus</i> , <i>Paraleichthys</i>	224	<i>Alectis</i>	274
<i>albimentis</i> , <i>Gymnothorax</i>	93	<i>crinitus</i>	274
<i>Lycodontis</i>	93	<i>Aledon</i>	503
<i>albipinnis</i> , <i>Lucifer</i>	70	<i>storeri</i>	503
<i>albirostris</i> , <i>Corythoichthys</i>	243	<i>Alepes</i>	271
<i>Hippocampus</i>	243	<i>melanoptera</i>	271
<i>Prionotus</i>	243	<i>Alepidosaurus</i> <i>esculapius</i>	171
<i>albofasciata</i> , <i>Scorpana</i>	371	<i>borealis</i>	171
<i>albolineatus</i> , <i>Fundulus</i>	177	<i>(caulopus) poeyi</i>	171
<i>albonaculatus</i> , <i>Crossias</i>	389	<i>serra</i>	171
<i>Serranus</i>	317	<i>alepidota</i> , <i>Chenopsis</i>	465
<i>albopunctatus</i> , <i>Gobius</i>	440	<i>alepidotum</i> , <i>Gobiosoma</i>	446

List of scientific names—Continued

	Page		Page
<i>alepidotus</i> , <i>Chætodon</i>	265	<i>Allocottus</i>	391
<i>Derepodichthys</i>	479	<i>embryum</i>	391
<i>Gobius</i>	446, 446	<i>Allogamibusia</i>	188
<i>Lucioblennius</i>	465	<i>cana</i>	188
<i>Alepisauridæ</i>	171	<i>tridentiger</i>	188
<i>Alepisaurus</i>	171	<i>Allosmerus</i>	188
<i>asculapius</i>	171	<i>attenuatus</i>	67
<i>altivelis</i>	171	<i>Allosomus</i>	61, 64
<i>azureus</i>	171	<i>Allotis</i>	300
<i>ferox</i>	171	<i>humilis</i>	300
<i>Alepocephalidæ</i>	51, 54	<i>Allotriognathus</i>	215
<i>Alepocephalus</i>	51	<i>Allurus</i>	402
<i>agassizii</i>	51	<i>almeida</i> , <i>Belone</i>	195
<i>asperifrons</i>	52	<i>Alopecias</i>	18
<i>bairdii</i>	52	<i>vulpes</i>	18
<i>convexifrons</i>	52	<i>Alopas</i>	18
<i>fundulus</i>	52	<i>macrourus</i>	18
<i>macropterus</i>	53	<i>vulpinus</i>	18
<i>productus</i>	51	<i>Alopidae</i>	18
<i>rostratus</i>	51	<i>Alosa</i>	42
<i>tenebrosus</i>	52	<i>alabamae</i>	43
<i>Aleposomus</i>	53	<i>apicalis</i>	43
<i>copel</i>	53	<i>bishopi</i>	43
<i>aletes</i> , <i>Hudsonius</i>	132	<i>clupea</i>	42
<i>aleutensis</i> , <i>Acipenser</i>	34	<i>cyanonotum</i>	42
<i>Lyconectes</i>	471	<i>lineata</i>	42
<i>aleutianus</i> , <i>Sebastodes</i>	365	<i>menhaden</i>	44
<i>Zalopyr</i>	365	<i>ohiensis</i>	42
<i>aleutica</i> , <i>Raja</i>	26	<i>præstabilis</i>	42
<i>aleuticus</i> , <i>Cottus</i>	385	<i>sapidissima</i>	42
<i>Sebastosomus</i>	365	<i>teres</i>	40
<i>Thecopterus</i>	392	<i>alosoides</i> , <i>Amphiodon</i>	41
<i>alexandri</i> , <i>Acipenser (Antaceus)</i>	34	<i>alpeneæ</i> , <i>Leucichthys</i>	63
<i>Rosicola</i>	365	<i>Alpharaia</i>	24
<i>Sebastodes</i>	305	<i>Alphesthes</i>	312
<i>alexandrini</i> , <i>Orthagoriscus</i>	503	<i>afer</i>	312
<i>Alexurus</i>	437	<i>chloropterus</i>	312
<i>armiger</i>	437	<i>lightfooti</i>	313
<i>alfari</i> , <i>Cichlasoma</i>	419	<i>multiguttatus</i>	313
<i>Parapetenia</i>	419	<i>Alpismaris</i>	163
<i>Alfaro</i>	193	<i>rizzo</i>	163
<i>acutiventralis</i>	193	<i>alta</i> , <i>Yuriria</i>	137
<i>acutiventralis</i>	193	<i>altavelia</i> , <i>Pteroplatea</i>	29
<i>culturatus</i>	193	<i>Raja</i>	29
<i>alfredi</i> , <i>Kryptophanaron</i>	234	<i>alternata</i> , <i>Scarus</i>	434
<i>Algansæa</i>	115	<i>alternata</i> , <i>Perca mitchilli</i>	307
<i>affinis</i>	116	<i>alticulus</i> , <i>Catostomus</i>	105
<i>antica</i>	122	<i>Alticus</i>	463
<i>bicolor</i>	122	<i>saluatorius</i>	463
<i>dugesii</i>	116	<i>alticus</i> , <i>Salaris</i>	463
<i>formosa</i>	122	<i>altifrons</i> , <i>Astatheros</i>	420
<i>lacustris</i>	116	<i>Heros</i>	420
<i>obesa</i>	122	<i>altipinna</i> , <i>Belone</i>	197
<i>rubescens</i>	116	<i>altipinnis</i> , <i>Alburnellus</i>	127
<i>sallæi</i>	116	<i>Hydrophlox</i>	127
<i>stigmatura</i>	116	<i>Loricaria</i>	162
<i>tarascorum</i>	116	<i>Micropogon</i>	349
<i>tineella</i>	116	<i>altivelis</i> , <i>Alepisaurus</i>	171
<i>Algoma</i>	144	<i>Corallicola</i>	456
<i>amaræ</i>	144	<i>Cremnobates</i>	466
<i>alia</i> , <i>Tautoga</i>	424	<i>Sebastolobus</i>	384
<i>aliala</i> , <i>Acanthurus</i>	363	<i>Serranus</i>	310
<i>alicicæ</i> , <i>Squalus</i>	119	<i>altus</i> , <i>Bubalichthys</i>	102
<i>alictolus</i> , <i>Trachurus</i>	279	<i>Chorinemus</i>	278
<i>Alinectes</i>	402	<i>Hudsonius</i>	137
<i>alipes</i> , <i>Salmo</i>	60	<i>Oligoplites</i>	278
<i>alisthostomus</i> , <i>Dipterus</i>	342	<i>Priacanthus</i>	323
<i>Allartediæ</i>	377	<i>Pseudopriacanthus</i>	323
<i>corallinus</i>	377	<i>aluta</i> , <i>Scæna</i>	344
<i>Allector</i>	484	<i>alutaceus</i> , <i>Acrocheilus</i>	112
<i>chelonæ</i>	484	<i>Alutarius amphacanthoides</i>	496
<i>allector</i> , <i>Dolopichthys</i>	509	<i>macracanthus</i>	496
<i>alleghaniensis</i> , <i>Moxostoma</i>	111	<i>obliteratus</i>	496
<i>Salmo</i>	60	<i>Alutera</i>	495
<i>allenii</i> , <i>Rathbunella</i>	452	<i>cineræ</i>	496
<i>alleteratus</i> , <i>Euthynnus</i>	258	<i>guntheriana</i>	496
<i>Scomer</i>	258	<i>monoceros</i>	496
<i>Allinectes</i>	403	<i>picturata</i>	496
<i>allisi</i> , <i>Crossias</i>	389	<i>punctata</i>	496
<i>allisoni</i> , <i>Neothunnus</i>	260	<i>Aluteres berardi</i>	496
<i>Thunnus</i>	260	<i>cuspicauda</i>	496
<i>Allocirr.</i>	402	<i>Aluterus anginosus</i>	496
<i>Alloclinus</i>	456	<i>cultrifrons</i>	496
<i>holderi</i>	456	<i>holbrookii</i>	496

List of scientific names—Continued

	Page		Page
<i>Aluterus venosus</i>	495	<i>Ameiurus nebulosus catulus</i>	154
<i>alutum, Acutomentum</i>	366	<i>platycephalus</i>	154
<i>alutus, Apogon</i>	298	<i>ponderosus</i>	152
<i>Apogonichthys</i>	298	<i>prosthectis</i>	154
<i>Nector</i>	346	<i>vulgaris</i>	153
<i>Sebastichthys</i>	368	<i>americana, Lucioperca</i>	282
<i>Alvarius</i>	282, 294	<i>Manta</i>	32
<i>fonticola</i>	295	<i>Morone</i>	307
<i>lateralis</i>	294	<i>Morrhua</i>	210
<i>alveatus, Trigonorhina</i>	24	<i>Perca</i>	281, 307
<i>alveatus, Bathytroctes</i>	52	<i>Rala</i>	25
<i>alvifrons, Bathytroctes</i>	52	<i>Scorpaena</i>	393
<i>alvordii, Cottus</i>	385	<i>americanum, Apogon</i>	295
<i>Alvordius</i>	283	<i>americanus, Acanthias</i>	21
<i>aspro</i>	283	<i>Alburnus</i>	350
<i>crassus</i>	283	<i>Ammodytes</i>	451
<i>evermanni</i>	284	<i>Amphiprion</i>	308
<i>evides</i>	284	<i>Apogon</i>	295
<i>günterii</i>	283	<i>Ballistes</i>	403
<i>macrocephalus</i>	284	<i>Blennius</i>	473
<i>maculatus</i>	283	<i>Cyprinus</i>	115, 350
<i>maximkuckiensis</i>	284	<i>Enchelyopus</i>	213
<i>ouachites</i>	284	<i>Eques</i>	352, 353
<i>peltatus</i>	283	<i>Esox</i>	172, 173
<i>phoxocephalus</i>	284	<i>Esox lucius</i>	173
<i>roanoka</i>	284	<i>Hemipterus</i>	393
<i>sellaris</i>	284	<i>Histiophorus</i>	263
<i>spilimani</i>	283	<i>Istiphorus</i>	263
<i>Alysia</i>	168	<i>Labrus</i>	424
<i>loricata</i>	108, 169	<i>Lophius</i>	504
<i>ambassis, Alburnus</i>	124	<i>Macrozoarces</i>	473
<i>Notropis</i>	124	<i>Menticirrhus</i>	330
<i>amarra, Algona</i>	144	<i>Petromyzon</i>	8
<i>Dionda</i>	144	<i>Pleuronectes</i>	227
<i>amarillas, Gustivere</i>	309	<i>Polynemus</i>	256
<i>amarus, Hudsonius</i>	132	<i>Polyprion</i>	308
<i>amates, Phallichthys</i>	190	<i>Pseudopleuronectes</i>	227
<i>Poecilia</i>	190	<i>Scorpaena</i>	393
<i>ambassis, Sargus</i>	336	<i>Squalus</i>	19
<i>ambiguous, Lutjanus</i>	327	<i>amethystinus, Maurolicus</i>	73
<i>Merluccius</i>	214	<i>amethystinus, Salmo</i>	59
<i>Mesopriion</i>	327	<i>Amla</i>	38, 295
<i>Amblodon</i>	352	<i>binotata</i>	295
<i>bubalis</i>	102	<i>calva</i>	38
<i>concinnus</i>	352	<i>canina</i>	38
<i>grunniens</i>	352	<i>cinerra</i>	38
<i>lineatus</i>	352	<i>guadalupensis</i>	296
<i>neglectus</i>	352	<i>immaculata</i>	40
<i>saturnus</i>	348	<i>lentiginosa</i>	38
<i>Ambloplites</i>	303	<i>marmorata</i>	38
<i>cavifrons</i>	303	<i>occidentalis</i>	38
<i>rupestris</i>	303	<i>ocellata</i>	38
<i>amblops, Felichthys</i>	159	<i>ornata</i>	38
<i>Rutilus</i>	133	<i>piquotii</i>	38
<i>Trachoryctes</i>	159	<i>reticulata</i>	38
<i>Amblyopsidae</i>	194	<i>retrosella</i>	295
<i>Amblyopsis</i>	104, 195	<i>subcærulea</i>	38
<i>mexicanus</i>	448	<i>thompsoni</i>	38
<i>spelaeus</i>	195	<i>viridis</i>	38
<i>Amblyopsis, Cithlus</i>	437	<i>Amia</i>	38
<i>Electris</i>	437	<i>calva</i>	38
<i>Amblyopus brevis</i>	448	<i>Amiichthys</i>	297
<i>sagitta</i>	448	<i>dipaterus</i>	297
<i>Amblyomacentrus</i>	413	<i>Amidae</i>	38
<i>Amblyraja</i>	24, 25	<i>Amitra</i>	404, 405
<i>amblyrhynchus, Caranx</i>	271	<i>liparina</i>	404, 405
<i>Hemicarax</i>	271	<i>Amiichthys</i>	404
<i>Amblyscion</i>	344	<i>Amitrus</i>	404
<i>argenteus</i>	344	<i>Amiurus</i>	153
<i>Ameriurus</i>	161	<i>brachyacanthus</i>	154
<i>bolini</i>	153	<i>brunneus</i>	154
<i>brachyacanthus</i>	153	<i>cragini</i>	154
<i>catus</i>	153	<i>lophius</i>	153
<i>dugeei</i>	153	<i>meridionalis</i>	152
<i>erebennus</i>	153	<i>niveliventris</i>	153
<i>lacustris okeechobeensis</i>	152	<i>obesus</i>	154
<i>marmoratus</i>	154	<i>prosthectis</i>	154
<i>melas</i>	154	<i>Ammocotes</i>	8
<i>misphilensis</i>	154	<i>aepyptera</i>	10
<i>natalis</i>	153	<i>aureus</i>	10
<i>natalis analis</i>	153	<i>bicolor</i>	8
<i>nebulosus</i>	154	<i>borealis</i>	9, 10
		<i>cibarius</i>	10
		<i>concolor</i>	9

List of scientific names—Continued

	Page		Page
<i>Ammocetes unicolor</i>	9	<i>analogus</i> , <i>Abudeodus</i>	415
<i>Ammocetus</i>	8	<i>Epinephelus</i>	310
<i>Ammocrypta</i>	282, 289	<i>Euchistodus</i>	415
beani	289	<i>Kyphosus</i>	339
clara	288	<i>Pimelepterus</i>	339
gelida	289	<i>analostoma</i> , <i>Cyprinella</i>	129
vivax	288	<i>Erogala</i>	129
<i>ammocryptus</i> , <i>Tetraodon</i>	499	<i>Anarhichadidae</i>	472
<i>Ammodytes</i>	451	<i>Anarhichas</i>	472
alascanus	451	<i>lupus</i>	472
americanus	451	<i>orientalis</i>	472
dubius	451	<i>Anarmostus</i>	329
personatus	451	<i>Anarrhichthas karrak</i>	472
septiplinis	451	<i>latifrons</i>	472
tobianus	451	<i>leopardus</i>	472
vittatus	451	<i>lepturus</i>	472
<i>Ammodytidae</i>	451	<i>maculatus</i>	472
<i>Ammolerops</i>	230	<i>minor</i>	472
<i>ameenus</i> , <i>Alburnellus</i>	123	<i>strigosus</i>	472
Notropis	123	<i>Anarrhichthyidae</i>	422
<i>Amore pixuma</i>	436	<i>Anarrhichthys</i>	472
<i>amoren</i> , <i>Gobius</i>	436	<i>felis</i>	472
<i>Amorphacephalus</i>	430	<i>ocellatus</i>	472
granulatus	430	<i>anastos</i> , <i>Acipenser</i> (<i>Iuso</i>).....	35
<i>amphacanthoides</i> , <i>Alutarius</i>	496	<i>Ananchias canescens</i>	161
<i>Amphigonopterus</i>	409	<i>anceps</i> , <i>Cottus</i>	386
aurora	409	<i>Anchisomus</i>	409
<i>Amphilophus</i>	420, 422	<i>geometricus</i>	409
frebelii	422	<i>(Anchisomus) caudacinctus</i> , <i>Prionotus</i>	500
labiatus	422	<i>Anchoa</i>	47, 50
lobochilus	422	<i>Anchovia</i>	50
<i>Amphiodon</i>	41	<i>abbotti</i>	50
alooides	41	<i>arenicola</i>	49
<i>Amphioxides</i>	7	<i>brevirostrum</i>	51
<i>Amphioxus</i>	7	<i>cherostoma cayorum</i>	49
<i>amphioxys</i> , <i>Cantherines</i>	491	<i>clupeoides</i>	50
Monocanthus	494	<i>eigenmanni</i>	50
<i>Amphiprion americanus</i>	308	<i>elongata</i>	49
australis	308	<i>filifera</i>	50
matejuelo	235	<i>lepidentostole</i>	49
<i>Amphistichus</i>	410	<i>macrolepidota</i>	50
argenteus	410, 411	<i>parva</i>	49
heermanni	411	<i>platyargyreia</i>	49
similis	411	<i>scitula</i>	49
<i>Amphotistius</i>	28	<i>trinitatis</i>	50
dipterurus	29	<i>anchovia</i> , <i>Sardinella</i>	43
sabinius	29	<i>Sardinia</i>	43
say	28	<i>Anchoviella</i>	47
<i>ampla</i> , <i>Makaira</i>	263	<i>abbotti</i>	50
<i>amplexicolis</i> , <i>Sbrothrodus</i>	360	<i>arenicola</i>	49
<i>Ampliova</i>	50, 51	<i>argyrophana</i>	48
halboe	50	<i>astilbe</i>	49
<i>amplus</i> , <i>Searus</i>	432	<i>branchiomelas</i>	49
<i>Tetrapturus</i>	263	<i>cayorum</i>	49
<i>ampullaceus</i> , <i>Ophiongnathus</i>	95	<i>cherostoma</i>	48
<i>Saccopharynx</i>	95	<i>compressa</i>	50
<i>Anablepsidæ</i>	104	<i>cubana</i>	48
<i>Anableps</i>	104	<i>culttrata</i>	48
anableps	104	<i>curta</i>	48
dowei	104	<i>delicatissima</i>	48
tetraphthalmus	104	<i>eigenmanni</i>	50
<i>anableps</i> , <i>Anableps</i>	104	<i>elongata</i>	49
<i>Anacanthini</i>	202	<i>epsetus</i>	48
<i>Anacyrtus guatemalensis</i>	98	<i>exigua</i>	47
<i>anagallinus</i> , <i>Lepomis</i>	300	<i>filliera</i>	50
<i>Anago</i>	80	<i>guianensis</i>	49
macrurus	80	<i>helleri</i>	48
nitens	80	<i>ischana</i>	48
<i>anago</i> , <i>Conger</i>	80	<i>lepidentostole</i>	49
<i>analigutta</i> , <i>Pomacentrus</i>	414	<i>lucida</i>	48
<i>analis</i> , <i>Ameirus natalis</i>	153	<i>lyolepis</i>	49
Clinocottus	390	<i>miarchus</i>	47
<i>Conger</i>	80	<i>mitchilli</i>	48
<i>Eupomacentrus</i>	413	<i>mundcola</i>	50
<i>Hyperprosopon</i>	410	<i>naso</i>	49
<i>Hyporcritichthys</i>	410	<i>panamensis</i>	50
<i>Lutjanus</i>	326	<i>parva</i>	49
<i>Mesopriion</i>	326	<i>perfasciata</i>	47
<i>Notacanthus</i>	76	<i>perthecata</i>	47
<i>Oligocottus</i>	390	<i>platyargyreia</i>	49
<i>Orthagoriscus</i>	503	<i>rastralis</i>	50
<i>Pomacentrus</i>	413	<i>robertsi</i>	49
<i>Scyris</i>	274	<i>scitula</i>	49
<i>Umbrina</i>	350		

List of scientific names—Continued

	Page		Page
<i>Anchoviella</i> <i>scofieldi</i>	49	<i>Anisotremus</i> <i>scapularis</i>	332
<i>spinifera</i>	50	<i>serrula</i>	333
<i>trinitatis</i>	50	<i>splenatus</i>	332
<i>Ancistrus</i>	161	<i>surinamensis</i>	332
<i>aspidolepis</i>	161	<i>taeniatus</i>	332
<i>chagresi</i>	160	<i>virginicus</i>	332
<i>guacharote</i>	161	<i>anisurum</i> , <i>Moxostoma</i>	109
<i>mediana</i>	161	<i>anisurus</i> , <i>Catostomus</i>	109
<i>medians</i>	161	<i>anjerensis</i> , <i>Gobius</i>	442
<i>planiceps</i>	161	<i>anna-carolina</i> , <i>Mugilomorus</i>	39
<i>spinosa</i>	160	<i>annæ</i> , <i>Agonus</i>	394
<i>Ancyloodon</i>	354	<i>Cottus</i>	384
<i>jaculidens</i>	354	<i>annectans</i> , <i>Gambusia</i>	187
<i>parvipinnis</i>	353	<i>annectens</i> , <i>Coccolus</i>	217
<i>ancyloodon</i> , <i>Lonchurus</i>	354	<i>Gambusia</i>	187
<i>Macrodon</i>	354	<i>Priapichthys</i>	187
<i>Ancylopsetta</i>	224	<i>annectens hisporis</i> , <i>Priapichthys</i>	187
<i>dendritica</i>	224	<i>annularis</i> , <i>Centropristes</i>	319
<i>quadrocincta</i>	224	<i>Nauclerus</i>	278
<i>andreae</i> , <i>Rhinoscopelus</i>	168	<i>Pomoxis</i>	304
<i>Scopelus</i>	168	<i>Sparus</i>	338
<i>androsiensis</i> , <i>Gobiesox</i>	489	<i>annulata</i> , <i>Melanura</i>	174
<i>Angelichthys</i>	361	<i>annulatum</i> , <i>Exoglossum</i>	146, 174
<i>ciliaris</i>	361	<i>annulatus</i> , <i>Antennarius</i>	608
<i>isabelita</i>	361	<i>Chelidonichthys</i>	499
<i>townsendi</i>	361	<i>Tetradon</i>	499
<i>angelus</i> , <i>Squatina</i>	22	<i>anogenus</i> , <i>Hybopsis</i>	136
<i>anginosus</i> , <i>Aluterus</i>	496	<i>Notropis</i>	136
<i>anglerum</i> , <i>Lumpus</i>	398	<i>anolis</i> , <i>Saurus</i>	164
<i>anguiformis</i> , <i>Ophichthys</i> (<i>Sphagebranchus</i>).....	84	<i>anomala</i> , <i>Dekaya</i>	357
<i>Sphagebranchus</i>	84, 91	<i>Anomalopidae</i>	234
<i>Anguilla</i>	77	<i>anomalium</i> , <i>Campostoma</i>	148
<i>anguilla</i>	77	<i>anomalus</i> , <i>Caulolatitus</i>	357
<i>aferrima</i>	77	<i>Rutilus</i>	146
<i>australis</i>	77	<i>Anoplagonus</i>	397
<i>blephura</i>	77	<i>Inermis</i>	398
<i>chrysypa</i>	77	<i>Anoplarchus</i>	468
<i>cubana</i>	77	<i>eristigalli</i> , <i>archolepis</i>	468
<i>laticauda</i>	77	<i>insignis</i>	468
<i>lutea</i>	77	<i>purpurescens</i>	468
<i>novæorleanensis</i>	77	<i>Anoplogaster</i>	233
<i>novæterreæ</i>	77	<i>cornutus</i>	233
<i>oceânica</i>	77	<i>Anopllopionta</i>	373
<i>punctatissima</i>	79	<i>fimbria</i>	373
<i>rostrata</i>	77	<i>merlangus</i>	373
<i>tenuirostris</i>	77	<i>Anoplopomidae</i>	373
<i>texana</i>	77	<i>anoplos</i> , <i>Uranoscopus</i>	454
<i>tyrannus</i>	77	<i>anoplus</i> , <i>Thyrsicus</i>	379
<i>wabashensis</i>	77	<i>Anopopus</i>	8
<i>xanthomelas</i>	77	<i>Anopsmius</i>	600
<i>anguilla</i> , <i>Anguilla</i>	77	<i>Anostomidae</i>	100
<i>Ictalurus</i>	152	<i>Ansonia</i> , <i>cuvieri</i>	264
<i>Muraena</i>	77	<i>ansonii</i> , <i>Salmo</i>	55
<i>anguillaris</i> , <i>Bleennius</i>	470, 473	<i>Antaceidae</i>	10
<i>Tribranchus</i>	77	<i>Antaceus</i>	34
<i>Anguillidae</i>	77	<i>Antelochimera</i>	33
<i>anguillifer</i> , <i>Pholidichthys</i>	465	<i>chætrirhampus</i>	33
<i>Anguilloidei</i>	77	<i>Antennaria</i> , <i>avalonis</i>	506
<i>anguina</i> , <i>Muraena</i>	90	<i>Antennariidae</i>	504
<i>anguineus</i> , <i>Nerophis</i>	243	<i>Antennarius</i>	505
<i>anguis</i> , <i>Lycodes</i>	475	<i>annulatus</i>	506
<i>anguiceps</i> , <i>Macrourus</i>	200	<i>astroscopus</i>	506
<i>Macrurus</i>	200	<i>avalonis</i>	506
<i>anguilifer</i> , <i>Clethlaurus</i>	416	<i>corallinus</i>	506
<i>Heros</i>	416	<i>inops</i>	505
<i>angulosus</i> , <i>Balistes</i>	492	<i>leopardinus</i>	505
<i>Canthidermis</i>	492	<i>multilocellatus</i>	506
<i>angusticeps</i> , <i>Belone</i>	196	<i>nuttingii</i>	506
<i>Coregonus</i>	136	<i>ocellatus</i>	505
<i>Sphaeroides</i>	408	<i>pleuroptalmus</i>	505
<i>Strongylura</i>	196	<i>principis</i>	505
<i>Tetraodon</i>	498	<i>radiosus</i>	506
<i>angustidens</i> , <i>Macrostoma</i>	166	<i>reticularis</i>	505
<i>angustifrons</i> , <i>Epinephelus</i>	311	<i>sanguineus</i>	505
<i>Paraliparis</i>	405	<i>scaber</i>	505
<i>Serranus</i>	311	<i>sonnagasti</i>	506
<i>angustimanus</i> , <i>Merluccius</i>	214	<i>striatus</i>	505
<i>anisacanthus</i> , <i>Dolopichthys</i>	509	<i>tenebrosus</i>	505
<i>Ansarchus</i>	470	<i>tenuifilis</i>	505
<i>medius</i>	470	<i>tgiris</i>	506
<i>Anisochætodon</i>	359	<i>verrucosus</i>	506
<i>Anisotremus</i>	332	<i>antennatus</i> , <i>Cyclichthys</i>	502
<i>bicolor</i>	332	<i>Diodon</i>	502
<i>davidsonii</i>	332	<i>Anthias</i>	322
<i>interruptus</i>	332		

List of scientific names—Continued

	Page		Page
<i>Anthins aquilonarias</i>	328	<i>Apocope adobe</i>	141
caballerote	325	<i>carringtonii</i>	140, 141
cherna	311	<i>cousii</i>	141
formosus	330	<i>falcata</i>	141
(<i>Hemianthias</i>) peruanus	322	<i>henshavii</i>	140
jocu	325	<i>klamathensis</i>	141
quartus rondeleti	326	<i>nevadensis</i>	141
rabilubia	327	<i>nubila</i>	140
saponaceus	318	<i>oscula</i>	141
striatus	311	<i>robusta</i>	140
tenuis	322	<i>umatilla</i>	141
vivanus	322	<i>vallifera</i>	141
<i>anthias</i> , <i>Labrus</i>	322	<i>vulnerata</i>	141
<i>anthracinus</i> , <i>Acipenser</i> (<i>Huso</i>)	35	<i>yarrowi</i>	141
<i>antica</i> , <i>Algassea</i>	122	(<i>Apocope</i>) <i>nevadensis</i> , <i>Rhinichthys</i>	141
<i>anticus</i> , <i>Hesperoleucus</i>	122	<i>velifera</i> , <i>Rhinichthys</i>	141
<i>Antigona</i>	358	<i>spoda</i> , <i>Perca</i>	325
capros	368	Pleuronectes	229
<i>Antigonidae</i>	368	<i>Apodes</i>	77
<i>antillanus</i> , <i>Conodon</i>	333	apodes, <i>Lutianus</i>	325
<i>antillarum</i> , <i>Caranx</i>	273	<i>Apodichthys</i>	466
Chilomycterus	502	flavidus	466, 467
Cylichthys	502	fucorum	467
Fundulus	175	inornatus	467
Sicydium	447	sanguineus	467
Talismans	63	univittatus	467
<i>Antimora</i>	211	violaceus	466
microlepis	211	virescens	467
rhina	211	<i>spodus</i> , <i>Lutianus</i>	328
rostrata	211	<i>Apogon</i>	38, 295
viola	211	alutus	296
<i>antistius</i> , <i>Chænobryttus</i>	303	americanum	295
antoniensis	153	americanus	295
antrostomus	70	atricaudus	295
anuli	146	atrodorsatus	295
<i>Apareiodon</i>	100	binotatus	295
compressus	100	dovii	295
dariensis	100	guadalupensis	296
<i>Apeltes</i>	239	imbuebis	295
quadracus	239	maculatus	295
<i>apeletes</i> , <i>Gasterosteus</i>	239	mokuicensis	295
<i>Apeltinæ</i>	239	pandionis	296
<i>Aphla</i>	246	pigmentarius	295
meridionalis	246	retrocella	295
<i>Aphenopnus</i>	262	rex-mullorum	295
carbo	262	ruber	295
minor	262	sellaeauda	295
<i>Aphododerus</i>	215	<i>apogon</i> , <i>Borophryne</i>	509
cocklanus	215	<i>Apogonichthys</i>	298
<i>Aphorista</i>	230	aiutus	296
atricauda	231	melampodus	296
diomediana	231	perdix	296
marginalis	231	puncticulatus	296
nebulosa	231	stellatus	296
ornata elongata	231	strombi	296
pigra	230	<i>Apogonidæ</i>	295
pusilla	231	<i>Apomotis</i>	298
<i>Aphredoderidæ</i>	215	cyanellus	298
<i>Aphredoderus</i>	215	euryurus	299
gibbosus	215	ischyrus	299
sayanus	215	phonax	299
<i>Aphyonus</i>	484	punctatus	299
gelatinosus	484	symmetricus	299
mollis	484	appendiculatus, <i>Centroponius</i>	305
<i>apia</i> , <i>Pirati</i>	313	<i>Exocetus</i>	201
<i>apiarius</i> , <i>Saranus</i>	308	appendix, <i>Lepomis</i>	299
<i>apiatus</i> , <i>Lepomis</i>	299	Lethenteron	9
apicalis, <i>Alosa</i>	43	Petromyzon	9
Echenesis	449	approximans, <i>Polynemus</i>	256
<i>Apionichthys</i>	230	Pomadasys	334
dumerili	230	<i>Aprion</i>	15
nebulosus	230	arlongus	328
<i>Apistes exul</i>	370	aprion, <i>Gerres</i>	342
<i>Aplension</i>	290, 298	Aprionodon	15
pottsville	290	isodon	15
<i>Aptites</i>	298	punctatus	15
<i>Aploiodon</i>	352	<i>Apristurus</i>	12
grunnienus	352	brunneus	12
neglectus	352	profundorum	12
<i>Aplodon</i>	339	<i>Aprodon</i>	474
margaritiferum	339	corteziana	474
<i>Aplurus</i>	261	cortezianus	474
simplex	261	<i>Aprolepis</i>	444
<i>Apocope</i>	140	barbaræ	444

List of scientific names—Continued

	Page		Page
<i>Apseudobranchus</i>	354	<i>arctifrons</i> , <i>Calamus</i>	337
<i>Apsicephalus</i>	499	<i>Citharichthys</i>	219
<i>Apsilus dentatus</i>	328	<i>Arctoscopus</i>	357
<i>Apterichthys</i>	84	<i>japonicus</i>	357
<i>selachops</i>	84	<i>Arctozenus</i>	165
<i>Apterontotus</i>	101	borealis	165
<i>passan</i>	101	<i>coruscans</i>	165
<i>rostratus</i>	101	<i>arctum</i> , <i>Siphostoma</i>	242
<i>Apterurus</i>	31	<i>arcturus</i> , <i>Salmo</i>	61
<i>fabroni</i>	31	<i>Salvelinus</i>	61
<i>Apthalmichthys</i>	89	<i>actrus</i> , <i>Syngnathus</i>	242
<i>caribbeus</i>	89	<i>arcutatum</i> , <i>Haemulon</i>	330
<i>javanicus</i>	89	<i>Hyperprosopon</i>	410
<i>Aptocyclus</i>	398	<i>arcuatus</i> , <i>Bathygadus</i>	202
<i>ventricosus</i>	308	<i>Chætodon</i>	360
<i>apua</i> , <i>Bodianus</i>	313	<i>Gadomus</i>	202
<i>Trisotropis venenosus</i>	313	<i>Pomacanthus</i>	360
<i>apus</i> , <i>Platypterus</i>	53	<i>ardens</i> , <i>Catostomus</i>	105
<i>aquedulcis</i> , <i>Gymnothorax</i>	93	<i>Hypsilepis</i>	126
<i>aqua-dulcis</i> , <i>Muraena</i>	91, 93	<i>Lythrurus</i>	126
<i>Rubula</i>	91	<i>ardeola</i> , <i>Belone</i>	195
<i>aquila</i> , <i>Raja</i>	31	<i>ardesiaca</i> , <i>Gla</i>	118
<i>Raja</i>	31	<i>ardesiacus</i> , <i>Lepomis</i>	301
<i>Sciene</i>	61	<i>arenaceus</i> , <i>Citharichthys</i>	219
<i>auquilarum</i> , <i>Salmo</i>	58	<i>arenata</i> , <i>Umbrina</i>	351
<i>auquilonaris</i> , <i>Anthias</i>	328	<i>arenatus</i> , <i>Arius</i>	150
<i>Etelides</i>	328	<i>Priacanthus</i>	322
<i>aquosus</i> , <i>Pleuronectes</i>	217	<i>Rhinichthys</i>	140
<i>arabatch</i> , <i>Salmo</i>	55	<i>Rypticus</i>	316
<i>arabicus</i> , <i>Chanos</i>	40	<i>arenicola</i> , <i>Anchovia</i>	49
<i>aracanga</i> , <i>Pseudoscarus</i>	434	<i>Anchoviella</i>	49
<i>Scarus</i>	433, 434	<i>Carapus</i>	486
<i>aracanensis</i> , <i>Notropis</i>	124	<i>Flerasfer</i>	486
<i>Notropis telecopus</i>	124	<i>Gillellus</i>	455
<i>area</i> , <i>Pristiurus</i>	12	<i>arenosus</i> , <i>Gadus</i>	210
<i>area</i> , <i>Atherina</i>	246	<i>argalus</i> , <i>Belone</i>	196
<i>areopodus</i> , <i>Catostomus</i>	108	<i>Strongylura</i>	196
<i>Aramaca</i>	218	<i>arge</i> , <i>Alburnellus</i>	124
<i>aramaca</i> , <i>Heimirhombus</i>	218	<i>Chiromystoma</i>	250
<i>Pleuronectes</i>	218	<i>Esloroparum</i>	250
<i>Rhombus</i>	218	<i>Kuhlia</i>	305
<i>arangoi</i> , <i>Cherojulis</i>	426	<i>Notropis</i>	124
<i>arara</i> , <i>Haemulon</i>	330	<i>argentata</i> , <i>Ciliata</i>	208
<i>Serranus</i>	311	<i>Motella</i>	208
<i>aratus</i> , <i>Mesoprion</i>	327	<i>argentatus</i> , <i>Astyanax</i>	96
<i>Raizero</i>	327	<i>Plagyrus</i>	128
<i>Arbaciosa</i>	490	<i>argentea</i> , <i>Bathyclupea</i>	281
<i>eos</i>	491	<i>Muraena</i>	77
<i>humeralis</i>	490	<i>Sciene</i>	334
<i>rbessodon</i>	490	<i>Sélène</i>	275
<i>rupestris</i>	491	<i>Sphyraena</i>	265
<i>zebra</i>	491	<i>Steindachneria</i>	203
<i>arboreifer</i> , <i>Limophryne</i>	510	<i>argenteolus</i> , <i>Lycophopis</i>	72, 73
<i>arcansanum</i> , <i>Etheostoma zonale</i>	289	<i>argenteum</i> , <i>Hyperprosopon</i>	410
<i>Nanostoma</i>	289	<i>punctatum</i> , <i>Hyperprosopon</i>	410
<i>arcansanus</i> , <i>Notropis</i>	124	<i>argenteus</i>	9
<i>Notropis telecopus</i>	124	<i>Amblyscion</i>	344
<i>Archaulius</i>	377	<i>Amphistichus</i>	411
<i>biseriatus</i>	377	<i>Brycon</i>	98
<i>archidium</i> , <i>Elattarchus</i>	345	<i>Centronotus</i>	277
<i>Odontoscion</i>	345	<i>Diplodus</i>	338
<i>Archocentrus</i>	420	<i>Eucinostomus</i>	340
<i>centrarchus</i>	420	<i>Fario</i>	55
<i>immaculatus</i>	420	<i>Hyperprosopon</i>	410
<i>nigrofasciatus</i>	420	<i>Kurtus</i>	323
<i>splirurus</i>	420	<i>Larimus</i>	344
<i>spinosissimus</i>	420	<i>Leuciscus</i>	117
<i>archolepis</i> , <i>Anoplarchus cristigall</i>	468	<i>Microponer</i>	349
<i>Anoplarchus purpureus</i>	468	<i>Petromyzon</i>	9
<i>Archomenidia</i>	248	<i>Sarchirus</i>	36
<i>sallei</i>	248	<i>Sargus</i>	338
<i>Archoperca</i>	313	<i>Saurus</i>	277
<i>Archoplites</i>	303	<i>Sparus</i>	337
<i>interruptus</i>	303	<i>Synodus</i>	40
<i>Archosargus</i>	338	<i>Trachinotus</i>	277
<i>aries</i>	338	<i>Trichiurus</i>	262
<i>probatocephalus</i>	338	<i>Argentina</i>	68
<i>artica</i> , <i>Liparis</i>	400	<i>carolina</i>	39
<i>articum</i> , <i>Benthosema</i>	170	<i>glossodonta</i>	40
<i>arcticus</i> , <i>Chironectes</i>	504	<i>pennanti</i>	73
<i>Galeocerdo</i>	14	<i>pretirosa</i>	67
<i>Gymnogaster</i>	216	<i>stali</i>	68
<i>Salmo</i>	68	<i>silus</i>	68
<i>Scopelus</i>	170	<i>sphyraena</i>	68
<i>Squalus</i>	14	<i>striata</i>	68
<i>Trachipterus</i>	216	<i>syrtensium</i>	68

List of scientific names—Continued

Page	Page
Argentinidae.....	68
argentinus, <i>Pimelodus</i>	152
argentissimus, <i>Gasterosteus</i>	238
<i>Plagopterus</i>	147
argiventris, <i>Lutjanus</i>	328
<i>Mesopriion</i>	326
argentosa, <i>Dionda</i>	144
Arges.....	159
<i>sabalo</i>	159
Argo.....	265
<i>steindachneri</i>	265
argus, <i>Cephalopholis</i>	309
<i>Gymnothorax (Limamurana)</i>	90
<i>Muraena</i>	90
<i>Pleuronectes</i>	217
<i>Squalus</i>	12
<i>argyrea, Scarcina</i>	262
<i>Scienea</i>	341
Argyrellosus.....	275
<i>brevortii</i>	276
<i>cerstedi</i>	275
<i>pacificus</i>	276
<i>setiphinis</i>	275
<i>unimaculatus</i>	275
<i>vomer</i>	275
Argyrellosus, <i>Pogonichthys</i>	116
Argyreus.....	139
<i>duleci</i>	139
<i>notabilis</i>	141
<i>nubilus</i>	140
<i>osculus</i>	141
<i>rubripinnis</i>	128
<i>argyreus, Fario</i>	55
Argyrius brevoorti.....	278
<i>filamentosus</i>	275
<i>mauricei</i>	275
<i>mitchilli</i>	275
<i>seifer</i>	275
<i>triacanthus</i>	275
argyrinus, <i>Hybognathus</i>	144
argyroleucus, <i>Bodianus</i>	345
argyro-leucus et exiguum, <i>Bodianus</i>	345
Argyropelecus.....	74
<i>affinis</i>	74
<i>durvillii</i>	74
<i>hemigymnus</i>	74
<i>lychnus</i>	74
<i>olfersi</i>	74
argyrophana, <i>Anchoviella</i>	48
argyrophanus, <i>Engraulis</i>	48
argyrops, <i>Sparus</i>	336
argyrosoma, <i>Damalichthys</i>	412
Argyrosomus.....	61
<i>alascanus</i>	62
<i>erleni</i>	63
<i>hoyi</i>	64
<i>huronius</i>	62
<i>johannae</i>	63
<i>nigriponis</i>	62, 63
<i>sisco</i>	62
<i>tullibee</i>	64
<i>zenithicus</i>	64
argyrosomus, <i>Damalichthys</i>	412
Argyrotrema.....	451
argyrurus, <i>Coryphena</i>	268
aryrus, <i>Pimelodus</i>	152
aries, <i>Archosargus</i>	338
<i>Sargus</i>	338
Ariidae.....	147
Ariodes.....	150
arliomus, <i>Aprión</i>	328
<i>Paranotropis</i>	125
<i>Photogenus</i>	125
Ariopsis.....	149
Ariosoma.....	79
<i>caudale</i>	80
<i>gilberti</i>	80
<i>opisthopthalmia</i>	80
Arius.....	150
<i>alatus</i>	148
<i>arenarius</i>	150
<i>arenatus</i>	150
<i>arius</i>	150
<i>assimilis</i>	150
<i>brandti</i>	148
<i>ceruleoceanus</i>	160
Arius commersonii.....	149
<i>dasycephalus</i>	150
<i>elatturus</i>	149
<i>emmeline</i>	151
<i>equestris</i>	161
<i>evermanni</i>	161
<i>furthi</i>	161
<i>grandicassis</i>	149
<i>guatemalensis</i>	150
<i>hypothalmus</i>	151
<i>inscriptus</i>	149
<i>kessleri</i>	149
<i>laticeps</i>	150
<i>llropus</i>	151
<i>melanopus</i>	150
<i>milberti</i>	149
<i>multiradiatus</i>	151
<i>osculus</i>	149
<i>planiceps</i>	149
<i>platypogon</i>	149
<i>seemanni</i>	150
<i>spixii</i>	150
<i>steindachneri</i>	151
<i>taylori</i>	151
<i>truncatus</i>	149
Arizona, <i>Pantosteus</i>	104
Arlina.....	287
<i>atripinnis</i>	286
<i>effulgens</i>	287
arlingtonia, <i>Gambusia</i>	177
armata, <i>Baldriella</i>	346
<i>Sciæna</i>	273
armatus, <i>Aspidophorus</i>	395
<i>Centropomus</i>	306
<i>Dactylobatus</i>	27
<i>Leptocottus</i>	390
<i>Nector</i>	346
<i>Nematonorus</i>	203
<i>Serranus</i>	313
armiger, <i>Alexurus</i>	437
arnillo, <i>Mesoprion</i>	328
Arnon.....	252
Arnoldi, <i>Limia</i>	191
Arothron.....	499
<i>erethizon</i>	500
<i>testudinatus</i>	499
Artedi.....	8
artedi, <i>Coregonus</i>	63
<i>Leucichthys</i>	63
<i>Rhamphichthys</i>	101
artedi bisselli, <i>Leucichthys</i>	63
Artediellus.....	377
<i>atlanticus</i>	377
<i>camchaticus</i>	378
<i>clarki</i>	377
<i>miacanthus</i>	378
<i>pacificus</i>	377
<i>unicinatus</i>	377
artedi, <i>Polynemus</i>	266
Artedius.....	377
<i>asperulus</i>	376
<i>fenestrals</i>	376
<i>lateralis</i>	377
<i>notospilotus</i>	378
<i>pugentensis</i>	375
<i>quadriseriatus</i>	376
artesia, <i>Nivicola</i>	292
<i>Poccilichthys</i>	292
Arthrophalius.....	184
arubensis, <i>Poccilis vanderpolli</i>	193
arundinaceus, <i>Syngnathus</i>	240
ascani, <i>Silus</i>	68
Ascelichthys.....	392
<i>rhodorus</i>	392
ascendens, <i>Syngnathus</i>	241
ascensionis, <i>Holocentrus</i>	235
<i>Perca</i>	235
ascita, <i>Pimelodus</i>	158
asellus, <i>Chelichthys</i>	500
Asematichthys.....	390
<i>taylori</i>	390
asper, <i>Auchenipterus</i>	456
<i>Cottus</i>	383
<i>Diodon</i>	501

List of scientific names—Continued

	Page		Page
asper, <i>Exerpes</i>	456	Asthenurus.....	208
<i>Hexagrammos</i>	373	<i>atripinnis</i>	208
<i>Macrurus</i>	203	<i>astilbe</i> , <i>Anchoviella</i>	49
<i>Pleuronectes</i>	227	<i>Stolephorus</i>	49
<i>Taractes</i>	265	<i>astillero</i>	49
aspera, <i>Limanda</i>	227	<i>astori</i> , <i>Petromyzon</i>	10
asperifrons, <i>Alepocephalus</i>	52	<i>Astrapogon</i>	296
asperima, <i>Cottus</i>	384	<i>stellatus</i>	296
asperinus, <i>Balistes</i>	492	<i>Astrobleplis</i>	159
<i>Cottus</i>	384	<i>Astroblepus</i>	159
<i>Myliobatis</i>	30	<i>grixovii</i>	159
<i>Pteromylaeus</i>	30	<i>pirrensis</i>	160
aspersus, <i>Serranus</i>	310	<i>Astrodermus</i>	264
asperulus, <i>Artedius</i>	376	<i>guttatus</i>	264
Aspicottus.....	381	<i>Astrolytes</i>	376
<i>bison</i>	381	<i>fenestralis</i>	376
<i>taurinus</i>	381	<i>Astronesthes</i>	71
aspidolepis, <i>Ancistrus</i>	161	<i>gemmifer</i>	71
<i>Chetostomus</i>	161	<i>niger</i>	71
Aspidophoroides.....	397	<i>Astronesthes nigra</i>	71
<i>bartoni</i>	397	<i>richardsoni</i>	71
<i>greenlandicus</i>	397	<i>Astronesthidae</i>	71
<i>güntheri</i>	397	<i>Astroscopus</i>	454
<i>inermis</i>	397	<i>guttatus</i>	454
<i>monopterygius</i>	397	<i>y-gracum</i>	454
<i>olrikii</i>	397	<i>zephyreus</i>	454
<i>tranquebar</i>	397	<i>astroscopus</i> , <i>Antennarius</i>	506
Aspidophoroidæ.....	397	<i>Astyana</i>	98
Aspidophorus.....	395	<i>eneus</i>	98
<i>armatus</i>	395	<i>argentatus</i>	98
<i>decagonus</i>	395	brevirostris	97
<i>europeus</i>	395	<i>emperador</i>	97
<i>quadricornis</i>	393	<i>fasciatus</i>	97
<i>spinosissimus</i>	394	<i>macrophthalmus</i>	97
aspidurus, <i>Urolophus</i>	30	<i>mexicanus</i>	96
<i>Urotrygon</i>	30	<i>nasutus</i>	97
Aspisurus.....	362	<i>petenensis</i>	97
asprella, <i>Crystallaria</i>	288	<i>ruberinus</i>	97
asprellus, <i>Pleurolepis</i>	288	<i>scabripinnis</i>	97
<i>Radiulinus</i>	379	<i>Asymmetron</i>	7
asprigens, <i>Pseuclichthys</i>	291	<i>lucayanum</i>	7
aspro, <i>Alvoridius</i>	283	<i>atenuatus</i> , <i>Chetodon</i>	380
assimilis, <i>Arius</i>	150	<i>Sarotherodon</i>	380
<i>Galeichthys</i>	150	<i>atchafalaya</i> , <i>Signalosa</i>	46
Astatheros.....	420, 422	<i>vanhyningi</i> , <i>Signalosa</i>	47
<i>altifrons</i>	420	<i>atelaspis</i> , <i>Acipenser</i> (<i>Huso</i>).....	35
<i>basilaris</i>	421	<i>Atelaxia</i>	216
<i>citrinellus</i>	422	<i>Atelobrachium</i>	207
<i>dorsatus</i>	420	<i>pterotum</i>	207
<i>erythraeum</i>	422	<i>aterrima</i> , <i>Anguilla</i>	77
<i>evermanni</i>	420	<i>athabasca</i> , <i>Leucichthys</i>	62
<i>globosum</i>	421	<i>Atheresthes</i>	221
<i>granadense</i>	421	<i>stomias</i>	221
<i>hedricki</i>	421	<i>Atherinæ</i>	246
<i>heterodontus</i>	420	<i>area</i>	246
<i>laurea</i>	421	<i>boscii</i>	247
<i>lethrinus</i>	420	<i>boyeri</i>	246
<i>longimanus</i>	420	<i>brownii</i>	48
<i>maculicauda</i>	421	<i>carolina</i>	246
<i>margaritifer</i>	421	<i>evermanni</i>	246
<i>montezuma</i>	421	<i>harringtonensis</i>	246
<i>oblongus</i>	421	<i>hepsetus</i>	246, 247
<i>octofasciatus</i>	421	<i>humboldtianus</i>	250
<i>rectangularis</i>	421	<i>laticeps</i>	246
<i>robertsoni</i>	420	<i>martinica</i>	248
<i>rostratum</i>	421	<i>menidia</i>	247
<i>septemfasciatus</i>	421	<i>microps</i>	246
<i>serfasciatus</i>	421	<i>microstoma</i>	246
<i>tuyensis</i>	421	<i>mordax</i>	67
asterias, <i>Blennius</i>	461	<i>notata</i>	247
<i>Galens</i>	13	<i>sardina</i>	247
<i>Mustelus</i>	13	<i>stipes</i>	246
<i>Spheroides</i>	498	<i>tæniata</i>	246
<i>Sphaeroides</i>	498	<i>vellicans</i>	246
<i>Urolophus</i>	30	<i>viridescens</i>	247
asterius, <i>Blennius</i>	461	<i>vomerina</i>	250
Asternopteryx.....	488	<i>Atherinella</i>	249
<i>gunelliformis</i>	488	<i>eriarcha</i>	249
Asternotremis.....	215	<i>evermanni</i>	248
<i>mesotrema</i>	215	<i>panamensis</i>	249
Asterrothea.....	396	<i>Atherinichthys</i>	250
<i>alsacana</i>	396	<i>albus</i>	251
<i>infraspinata</i>	396	<i>brevis</i>	250
<i>pentacantha</i>	396	<i>gracilis</i>	247
		<i>grandoculis</i>	251

List of scientific names—Continued

	Page		Page
<i>Atherinichthys guatemalensis</i>	249	<i>atricaudus</i> , <i>Apogon</i>	295
<i>pachylepis</i>	249	<i>Syphurus</i>	231
<i>sallei</i>	248	<i>atrilabiatus</i> , <i>Callionymus</i>	450
<i>Atherinidae</i>	246	<i>atrilitus</i> , <i>Zygonectes</i>	185
<i>Atherinoides</i>	260	<i>atrilobata</i> , <i>Chromis</i> (<i>Furcaria</i>).....	412
<i>atherinoides</i> , <i>Chlorodorus</i>	199	<i>Furcaria</i>	412
<i>Clupea</i>	51	<i>atrimana</i> , <i>Monolete</i>	220
<i>Notropis</i>	123	<i>atrimanus</i> , <i>Caranx</i>	271
<i>Pterengraulis</i>	51	<i>Hemicarax</i>	271
<i>Atherinomorus</i>	246	<i>Parasphyraenops</i>	297
<i>Atherinops</i>	252	<i>atrimensis</i> , <i>Menidia peninsulae</i>	247
<i>affinis</i>	252	<i>atripectus</i> , <i>Porogadus</i>	482
<i>insularum</i>	262	<i>atripes</i> , <i>Ditrema</i>	411
<i>cedrosensis</i>	252	<i>Lythrurus</i>	126
<i>guadalupae</i>	252	<i>Phanerodon</i>	411
<i>oregona</i>	252	<i>atripinnis</i> , <i>Arlina</i>	286
<i>regis</i>	252	<i>Asthenurus</i>	208
<i>Atherinopsis</i>	252	<i>Goodea</i>	183
<i>affinis</i>	252	<i>atriventer</i> , <i>Stomias</i>	69
<i>californiensis</i>	252	<i>atrocaudalis</i> , <i>Hybopsis</i>	133
<i>sonorensis</i>	252	<i>Notropis cayuga</i>	133
<i>tenuis</i>	251	<i>atrocaudus</i> , <i>Apogon</i>	295
<i>Athlennes</i>	197	<i>atrocyanus</i> , <i>Pomacentrus</i>	413
<i>Atima</i> , <i>Chilomycterus</i>	502	<i>atrodorsatus</i> , <i>Apogon</i>	295
<i>Diodon</i>	502	<i>atromaculata</i> , <i>Estrella</i>	287
<i>Guamaiulus</i>	502	<i>atromaculatus</i> , <i>Cyprinus</i>	117
<i>atkinsi</i> , <i>Gasterosteus</i>	238	<i>SEMOTILLIS</i>	117
<i>atlantica</i> , <i>Elacate</i>	281	<i>atromaculatus thorestanus</i> , <i>Serranillus</i>	117
<i>Emblemaria</i>	464	<i>atranus</i> , <i>Cyprinus</i>	139, 140
<i>Scorpaena</i>	370	<i>Rhinichthys</i>	140
<i>russula</i>	370	<i>atropurpureum</i> , <i>Ophidium</i>	489
<i>atlanticum</i> , <i>Oreosoma</i>	232	<i>atropurpureus</i> , <i>Epiileichthys</i>	469
<i>atlanticus</i> , <i>Arteediellus</i>	377	<i>atrorubens</i> , <i>Sebastodes</i>	365
<i>Benthodesmus</i>	292	<i>Zalopyr</i>	365
<i>Bregmaceros</i>	208	<i>atrovirens</i> , <i>Sebastichthys</i>	365
<i>Callorhynchus</i>	32	<i>Zalopyr</i>	365
<i>Dibranchus</i>	508	<i>attavilla</i> , <i>Dasyatis</i>	29
<i>Megalops</i>	39	<i>attenuatus</i> , <i>Cyclichthys</i>	502
<i>Neoliparis</i>	390	<i>attenuatus</i> , <i>Vinciguerria</i>	73
<i>Prometheus</i>	261	<i>attenuatum</i> , <i>Chiostoma</i>	250
<i>Rupisarcetes</i>	463	<i>attenuatus</i> , <i>Allosmerus</i>	67
<i>Salarias</i>	463	<i>Careproctus</i>	403
<i>Sparus</i>	310	<i>Halosaurus</i>	75
<i>Tarpon</i>	39	<i>Lycodapus</i>	479
<i>Tetragonurus</i>	269	<i>Maurolicus</i>	73
<i>Thynnus</i>	280	<i>Osmerus</i>	67
<i>atomarium</i> , <i>Sparisoma</i>	431	<i>Paraliparis</i>	404
<i>atomarius</i> , <i>Scarus</i>	431	<i>Rhinoliparis</i>	405
<i>Atopichthys</i>	54	<i>Atticus</i>	350
<i>acus</i>	54	<i>Atule</i>	270
<i>cinctetus</i>	54	<i>atwoodi</i> , <i>Carcharias</i>	20
<i>cingulus</i>	54	<i>aubeaubaibel</i> , <i>Etheostoma</i>	291
<i>dentatus</i>	54	<i>Oligocephalus</i>	291
<i>ensunculus</i>	54	<i>aubrieti</i> , <i>Lutjanus</i>	327
<i>falcidens</i>	54	<i>Auchenipterus</i>	456
<i>longidens</i>	54	<i>Auchenistius</i>	457
<i>lychnus</i>	54	<i>stainii</i>	457
<i>obtusus</i>	54	<i>Auchenopterus</i>	456
<i>opichthys</i>	54	<i>affinis</i>	457
<i>sclarus</i>	54	<i>albicaudus</i>	457
<i>Atopoclinidae</i>	465	<i>asper</i>	456
<i>Atopoclinus</i>	465	<i>cingulatus</i>	457
<i>ringens</i>	465	<i>fajardo</i>	457
<i>Atractoscion</i>	356	<i>fuscatus</i>	457
<i>nobilis</i>	356	<i>grandicormis</i>	457
<i>phoxocephalus</i>	356	<i>integripinnis</i>	457
<i>Atractosteus</i>	38	<i>mexicanus</i>	457
<i>lucius</i>	38	<i>monophthalmus</i>	456, 457
<i>tristochus</i>	38	<i>nox</i>	457
<i>tropicus</i>	38	<i>rubescens</i>	457
(<i>Atractosteus</i>) <i>berlandieri</i> , <i>Lepidosteus</i>	38	<i>auctorum</i> , <i>Lobotes</i>	323
<i>ferox</i> , <i>Lopisosteus</i>	38	<i>Auctospina</i>	368
<i>atramentatus</i> , <i>Syphurus</i>	231	<i>auriculata</i>	368
<i>straria</i> , <i>Perca</i>	231	<i>dalli</i>	369
<i>Siboma</i>	317	<i>audax</i> , <i>Histiophorus</i>	263
<i>Tigoma</i>	119	<i>audens</i> , <i>Menidia</i>	247
<i>atratius</i> , <i>Pimelodus</i>	119	<i>Aulastome marcgravii</i>	245
<i>Xenomystax</i>	154	<i>Auliscops</i>	239
<i>stratum</i> , <i>Myctophum</i>	169	<i>spiniceps</i>	239
<i>atricauda</i> , <i>Aporistia</i>	231	<i>aulicus</i> , <i>Siphonotoma</i>	240
<i>Hydrangia</i>	174	<i>Syngnathus</i>	240
<i>atricaudata</i> , <i>Deuterodon</i>	96	<i>Aulophalus</i>	190
<i>atricaudatus</i> , <i>Gephyrocharax</i>	96	<i>elongatus</i>	190

List of scientific names—Continued

	Page		Page
<i>Aulorhynchus</i>	239	<i>australis</i> , <i>Esox</i>	173
<i>flavidus</i>	239, 240	<i>Remilegia</i>	449
<i>Aulostoma</i>	244	<i>Tautis</i>	362
<i>cinerum</i>	245	<i>australis</i> , <i>Icelus</i>	379
<i>coloratum</i>	245	<i>austriana</i> , <i>Myoxostoma</i>	110
<i>Aulostomus marcgravii</i>	245	<i>austrinum</i> , <i>Myoxostoma</i>	110
<i>Aulostomi</i>	244	<i>Auris</i>	259
<i>Aulostomidae</i>	244	<i>rochei</i>	259
<i>Aulostomus chinensis</i>	244	<i>thazard</i>	259
<i>cinerous</i>	245	<i>avalonis</i> , <i>Antennaria</i>	506
<i>maculatum</i>	245	<i>Averruncus</i>	295
<i>surantiacus</i> , <i>Balistes</i>	495	<i>emmeline</i>	395
<i>Cottogaster</i>	254, 255	<i>sterletus</i>	396
<i>auratus</i> , <i>Chelio</i>	428	<i>averruncus</i> , <i>Kathetostoma</i>	454
<i>auriga</i> , <i>Monosanthus</i>	494	<i>avitus</i> , <i>Chologaster</i>	104
<i>surantiacus</i> , <i>Hypophomus</i>	285	<i>avocetta</i> , <i>Nemichthys</i>	83
<i>aurata</i> , <i>Coryphaena</i>	265	<i>Avocettina</i>	83
<i>Moniana</i>	132	<i>elongata</i>	83
<i>Saleima</i>	339	<i>gilli</i>	83
<i>auratum</i> , <i>Pristipoma</i>	333	<i>infans</i>	83
<i>auratus</i> , <i>Dipterus</i>	342	<i>Awaous</i>	442
<i>Gadus</i>	210	<i>nelsoni</i>	442
<i>Holocentrus</i>	309	<i>axillare</i> , <i>Pristipoma (Hæmulopsis)</i>	333
<i>Mullus</i>	343	<i>axillaris</i> , <i>Boreocottus</i>	387
<i>barbatus</i>	343	<i>Brachydeuterus</i>	333
<i>Notenigonus</i>	115	<i>Dipterus</i>	342
<i>Psenes</i>	268	<i>Gerres</i>	342
<i>Scomber</i>	258	<i>Myoxocephalus</i>	387
<i>aurea</i> , <i>Loricaria</i>	102	<i>axinophrys</i> , <i>Xystes</i>	395
<i>aureolum</i> , <i>Myoxostoma</i>	110	<i>Axyris</i>	377
<i>aureodus</i> , <i>Catostomus</i>	109, 110	<i>harringtoni</i>	377
<i>Dipterus</i>	342	<i>ayxa</i> , <i>Chætodon</i>	380
<i>Gerres</i>	342	<i>Lutjanus</i>	326
<i>Salvelinus</i>	60	<i>Neomens</i>	453
<i>Xenotis</i>	300	<i>Aylopon</i>	322
<i>aureo-ruber</i> , <i>Sparus</i>	432	<i>martinicensis</i>	320
<i>aureoviridis</i> , <i>Sphyraena</i>	305	<i>ayresi</i> , <i>Acipenser</i>	34
<i>aureus</i> , <i>Ammocetes</i>	10	<i>Centropristes</i>	318
<i>Caranx</i>	272	<i>Lepidotestes</i>	37
<i>Caranx</i>	270	<i>Parophrys</i>	225
<i>Chaetodon</i>	360	<i>Ayresia</i>	413
<i>Fundulus</i>	177	<i>punctipinnis</i>	413
<i>Heros</i>	422	<i>ayresii</i> , <i>Lampetra</i>	10
<i>Sparus</i>	302	<i>Petromyzon</i>	10
<i>Thorichthys</i>	422	<i>Sebastodes</i>	367
<i>auriculata</i> , <i>Auctospina</i>	368	<i>Sebastomus</i>	367
<i>Mobula</i>	31	<i>Azevia</i>	221
<i>auriculatus</i> , <i>Sebastes</i>	368	<i>panamensis</i>	221
<i>aurifrons</i> , <i>Gnathopogon</i>	453	<i>querina</i>	221
<i>auriga</i> , <i>Chætodon</i>	359	<i>azorica</i> , <i>Coryphaena</i>	265
<i>Dules</i>	330	<i>Azteca</i>	143
<i>aurita</i> , <i>Sardiniella</i>	43	<i>Aztecula</i>	143
<i>Selima</i>	339	<i>lermæ</i>	143
<i>auritus</i> , <i>Brachydeuterus</i>	333	<i>mexicana</i>	143
<i>Labrus</i>	299	<i>vittata</i>	143
<i>Lepomis</i>	299	<i>aztecus</i> , <i>Notropis</i>	143
<i>Macrodon</i>	99	<i>azurea</i> , <i>Hermosilla</i>	340
<i>aurofrenatum</i> , <i>Sparisoma</i>	432	<i>azureus</i> , <i>Alepisaurus</i>	171
<i>aurofrenatus</i> , <i>Scarus</i>	432	<i>Carcharias</i>	16
<i>auroguttatus</i> , <i>Zygonectes</i>	177	<i>Galeichthys</i>	160
<i>aurolateratum</i> , <i>Myctophum</i>	170	<i>Azurina</i>	412
<i>aurolineatum</i> , <i>Bathystoma</i>	331	<i>hirundo</i>	412
<i>Hæmulon</i>	831	<i>azurissimus</i> , <i>Microspathodon</i>	416
<i>auro-punctatus</i> , <i>Callyodon</i>	431		
<i>aupropunctatus</i> , <i>Cryptotomus</i>	431	 <i>babcocki</i> , <i>Rosicola</i>	365
<i>aurora</i> , <i>Abiesa</i>	409	<i>Sebastodes</i>	365
<i>Amphigonopterus</i>	409	<i>bacalau</i> , <i>Gobius</i>	441
<i>Csprophonus</i>	358	<i>badius</i> , <i>Euctenogobius</i>	441, 442
<i>Catostomus</i>	106	<i>fundulus heteroclitus</i>	176
<i>Eusebastes</i>	366	<i>Rhinichthys</i>	140
<i>Fario</i>	55	<i>Bagre</i>	147
<i>Sebastichthys</i>	366	<i>bahiensis</i>	148
<i>aurorubens</i> , <i>Centropristes</i>	327	<i>eydouxii</i>	148
<i>Rhomboplites</i>	323	<i>isthmeensis</i>	148
<i>aurovittatus</i> , <i>Mesopriion</i>	327	<i>marina</i>	148
<i>Ausonia</i>	264	<i>panamensis</i>	148
<i>cuvieri</i>	264	<i>pinnimaculatus</i>	148
<i>australasicus</i> , <i>Pneumatophorus</i>	268	<i>scutatus</i>	148
<i>Scomber</i>	258	<i>secundæ</i> , etc.	153
<i>australe</i> , <i>Etheostoma</i>	203	<i>bagre</i> , <i>Silurus</i>	147
<i>Zophendum</i>	116	<i>Bagrus propus</i>	148
<i>australis</i> , <i>Amphilophion</i>	308	<i>sundiacus</i>	149
<i>Clinocottus</i>	390	<i>(Sciaes) emphysetus</i>	148
<i>analis</i>	390	<i>bahamensis</i> , <i>Pisces viridis</i>	432
<i>Echeneis</i>	449	<i>Unicornis piscis</i>	496

List of scientific names—Continued

	Page		Page
bahamensis, <i>Vulpes</i>	39	Balistes kleinii.....	496
bahianus, <i>Acanthurus</i>	363	laevis.....	496
Rhompus.....	217	liberiensis.....	492
bahiensis, <i>Bagre</i>	148	lineopunctatus.....	493
Cypselurus.....	202	linguatula.....	496
Engraulis.....	40	longus.....	493
Exocetus.....	202	macrops.....	492
Galeichthys.....	148	macropterus.....	449
baileyi, <i>Cyprinodon</i>	182	maculatus.....	493
Cyprinodon macularius.....	182	melanopterus.....	493
ballioni, <i>Casiomorus</i>	276	mento.....	493
Trachinotus.....	276	monoceros.....	495, 496
Ballonotus.....	276	monoceros unicolor.....	496
balloui, <i>Gasterosteus</i>	237	moribundus.....	492
Baiondon.....	417	nausfragium.....	491
fasciatus.....	417	nigra.....	493
Baione.....	60	nitidus.....	493
fontinalis.....	60	notatus.....	493
Balostoma.....	221	piceus.....	493
brachialis.....	230	polylepis.....	492
bairdi, <i>Acipenser</i>	34	pwelli.....	492
Callionymus.....	430	punctatus.....	492
Cottus.....	383	radula.....	493
Macrourus.....	205	ringens.....	493
bairdiana, <i>Sphyraenops</i>	297	rufus.....	493
bairdianus, <i>Sphyraenops</i>	297	sandwichensis.....	493, 494
Syngnathus.....	242	sandwichiensis.....	494
Syrichtes.....	242	scapus.....	494
Bairdiella.....	345	schepfi.....	495
armata.....	348	scolopax.....	246
bedoti.....	348	scripta.....	495
chrysura.....	345	scriptus.....	495
ensifera.....	345	sobaco.....	492
icistia.....	348	spilotopterygius.....	492
ronchus.....	346	sufflamen.....	492
verso-crucis.....	346	tenlopodus.....	491
bairdii, <i>Alepocephalus</i>	52	triangularis.....	492
Bathymyzon.....	8	unicornu.....	490
Cottus.....	385	verres.....	492
Gasterostomus.....	95	vetula.....	491
Mitchillina.....	52	wilburgi.....	493
Petromyzon (Bathymyzon).....	8	Balistidae.....	491
Pomacentrus.....	415	Balistichthys.....	183
Salmo.....	61	balsana, <i>Thyrina</i>	249
Stegastes.....	415	balsanus, <i>Istalarius</i>	161
Bajacalifornia.....	53	Melaniris.....	248
burragei.....	53	balsas, <i>Pecilolopsis</i>	189
Bajonado.....	338	balteatus, <i>Cyprinus (Abramis)</i>	118
bajonado, <i>Calamus</i>	336	Eques.....	353
Sparus.....	336	Heros.....	418
balantophthalmus, <i>Scomber</i>	271	Pomacanthus.....	360
balo, <i>Hemiramphus</i>	199	Richardsonius.....	118
balboæ, <i>Amplova</i>	50	Theraps.....	418
Fundulus.....	179	Thynnus.....	260
Oxygymnocytes.....	179	Upeneus.....	343
baldwini, <i>Prionodes</i>	319	banana, <i>Butyrinus</i>	30, 40
balearica, <i>Muraena</i>	79	Chonophorus.....	442
balearicum, <i>Ophisoma</i>	80	Gobius.....	442
balearus, <i>Cyprinus (Abramis)</i>	117	bancrofti, <i>Torpedo</i>	27
balias, <i>Chirurus</i>	374	Barathrodermus.....	482
Balistes.....	491	manatinus.....	482
americanus.....	493	Barathronus.....	484
angulosus.....	492	bicolor.....	484
asperimus.....	492	barbare, <i>Aprolepis</i>	444
aurantiacus.....	495	Siphonotoma.....	240
barbatus.....	496	Syngnathus.....	240
bellus.....	491	barbata, <i>Brotula</i>	479
bridé.....	492	Lonchirurus.....	352
broccus.....	494	Loricaria.....	101, 162
buniva.....	491, 493	Pallasina.....	394
capistratus.....	492	Rhamdia.....	158
caprinus.....	491	barbatula, <i>Leomonema</i>	213
capriscus.....	491	barbatulum, <i>Leomonema</i>	213
carolinensis.....	491	barbatum, <i>Echiostoma</i>	70
chinensis.....	494	Lyconema.....	476
cicatricosus.....	493	Ophidion.....	485
ciliaris.....	492	barbatus, <i>Balistes</i>	496
ciliatus.....	494	Chauliodus.....	71
curassavicus.....	493	Enchelyopus.....	479
equestris.....	491	barbatus, <i>Mullus</i>	343
forcipatus.....	492	Siphagonus.....	394
fuliginosus.....	491	barbatus suratus, <i>Mullus</i>	343
guttatus.....	492	barbiger, <i>Macrourus</i>	205
heckeli.....	493	Macrurus.....	205
hispidus.....	494		

List of scientific names—Continued

	Page		Page
barbiger, <i>Peristedion</i> .	406	Bathylagus <i>milleri</i> .	69
<i>Peristedium</i> .	406	<i>pacificus</i> .	69
barbuda, <i>Liza</i> .	496	Bathymaster.	451
Barbulifer		<i>caeruleofasciatus</i> .	451
<i>cauthocæus</i> .	446	<i>hypoplectus</i> .	452
<i>papillosus</i> .	446	<i>jordani</i> .	452
barbulifer, <i>Rhinoliparis</i> .	405	<i>signatus</i> .	451
bardus, <i>Minomus</i> .	104	bathymaster, <i>Bregmaceros</i> .	208
barkani, <i>Agonus</i> .	394	Bathymasteridae.	451
barnharti, <i>Achirus</i> .	230	Bathymyzon.	8
baroni-s-müller, <i>Pimelodus</i> .	157	<i>bairdii</i> .	8
<i>Rhamdia</i> .	157	(<i>Bathymyzon</i>) <i>bairdii</i> , <i>Petromyzon</i> .	8
barracuda, <i>Esox</i> .	255	Bathynectes.	480
<i>Sphyraena</i> .	255	<i>compressus</i> .	480, 481
barracudina, <i>Paralepis</i> .	165	<i>gracilis</i> .	481
barratii, <i>Etheostoma</i> .	294	<i>laticeps</i> .	483
<i>Holeolepis</i> .	294	Bathyponus.	480
barreto, <i>Gobloides</i> .	448	<i>catena</i> .	481
bartholomaei, <i>Caranx</i> .	272	<i>pectoralis</i> .	482
<i>Elephothon</i> .	272	<i>tania</i> .	481
bartoni, <i>Acara</i> .	419	Bathyophis.	70
<i>Aspidophoroïdes</i> .	397	<i>ferox</i> .	70
<i>Chirostoma</i> .	250	Bathyphasma.	401
<i>Cylindrostomus</i> .	37	<i>ovigerum</i> .	401
<i>Parapetenia</i> .	37	bathyphila, <i>Cyclothone</i> .	72
Bascanichthys.		bathyphilum, <i>Neostoma</i> .	72
<i>bascanium</i> .	86	Bathypteroideæ.	166
<i>bascanoïdes</i> .	86	Bathypterois.	166
<i>cylindricus</i> .	86	<i>longifilis</i> .	166
<i>panamensis</i> .	86	<i>pectoralis</i> .	166
<i>peninsularis</i> .	86	<i>quadririnis</i> .	166
<i>scuticaris</i> .	86	<i>ventralis</i> .	166
<i>bascanium</i> , <i>Bascanichthys</i> .	86	Bathyaurus.	164
<i>Cœcula</i> .	86	<i>agassizii</i> .	164
Bascanius.		<i>ferox</i> .	164
<i>tædifer</i> .	230	Bathypterus.	166
<i>bascanoïdes</i> , <i>Bascanichthys</i> .	86	<i>longifilis</i> .	166
basilaris, <i>Astatheros</i> .		<i>pectoralis</i> .	166
<i>Heros</i> .	421	<i>quadririnis</i> .	166
Bassogigas.		<i>ventralis</i> .	166
<i>gilli</i> .	482	Bathyraurus.	164
<i>stelliferoides</i> .	482	<i>agassizii</i> .	164
Bassozetus.		<i>ferox</i> .	164
<i>catena</i> .	480, 481	Bathyrauridae.	164
<i>compressus</i> .	481	<i>molitis</i> .	164
<i>nasus</i> .	481	Bathystoma.	331
<i>normalis</i> .	480, 481	<i>aurolineatum</i> .	331
<i>tænia</i> .	481	<i>rimator</i> .	329, 331
batabana, <i>Corvula</i> .	345	<i>striatum</i> .	331
batabanus, <i>Johnius</i> .	345	Bathystomus.	331
bathicector, <i>Salmo</i> .	57	Bathytroctes.	52
Bathyagonus.		<i>alveatus</i> .	52
<i>nigripinnis</i> .	395, 396	<i>alvifrons</i> .	52
Bathyalopex.		<i>homopterus</i> .	53
<i>abbreviatus</i> .	33	<i>inspector</i> .	52
<i>plumbeus</i> .	33	<i>microlepis</i> .	52
bathybiß, <i>Paraliparis</i> .	404	<i>stomias</i> .	52
bathybiß, <i>Cynoglossus</i> .		Batis.	24
<i>Embastichthys</i> .	228	<i>radula</i> .	24
<i>Histiobranchus</i> .	78	Batis, <i>Raja</i> .	24, 25
<i>Synaphobranchus</i> .	78	Batoidæ.	23
Bathyclupea.		Batrachoides.	486
<i>argentea</i> .	281	<i>bouleengeri</i> .	487
<i>hoskynii</i> .	281	<i>goldmani</i> .	487
Bathyclupeidae.		<i>pacifici</i> .	487
Bathygadus.		<i>surinamensis</i> .	486
<i>arcuatus</i> .	203	<i>tau</i> .	486
<i>cavernosus</i> .	207	<i>variegata</i> .	487
<i>cottoides</i> .	203	<i>vernillas</i> .	487
<i>favosus</i> .	203	Batrachoididae.	486
<i>longidorsus</i> .	202	Batrachus.	487
<i>macrorops</i> .	203	<i>celatus</i> .	487
<i>melanobranchus</i> .	203	<i>margaritatus</i> .	487
Bathygobius.		<i>pacifici</i> .	487
<i>fuscus</i> .	440	<i>porosissimus</i> .	487, 488
<i>soporator</i> .	440	<i>surinamensis</i> .	486
Bathylacea.		<i>tau pardus</i> .	487
<i>nigricans</i> .	164	Batrictius.	486
Bathylagidae.		battaræ, <i>Orthragoriscus</i> .	503
Bathylagus.		bayamonensis, <i>Gobionellus</i> .	441
<i>antarcticus</i> .	68	<i>Gobius</i> .	441
<i>benedicti</i> .	69	bayanas, <i>Pomadasys</i> .	334
<i>borealis</i> .	69	<i>Pomadasys</i> .	334
<i>euryops</i> .	69	bdellium, <i>Petromyzon</i> .	9
		Bdellostomidae.	8
		<i>bdellium</i> , <i>stoutii</i> .	8
		beadlei, <i>Synechoglanis</i> .	151, 152
		beani, <i>Belonocharax</i> .	100
		<i>Caranx</i> .	272
		<i>Ctenolucius</i> .	100
		<i>Heros</i> .	419
		<i>Kirtlandia</i> .	248
		<i>Melamphæs</i> .	233
		<i>Membras</i> .	248

List of scientific names—Continued

	Page		Page
beani, <i>Ophidion</i>	486	beringi, <i>Crossias</i>	389
<i>Parapetenia</i>	419	beringiana, <i>Stegistrops</i>	379
<i>Plectromus</i>	233	beringianum, <i>Stegistrum</i>	379
<i>Pecilichthys</i>	287	beringianus, <i>Cyclogaster</i> (<i>Neoliparis</i>)	399
<i>Triglops</i>	380	<i>Neoliparis</i>	399
beanii, <i>Ammocrypta</i>	289	berlandieri, <i>Lepidosteus</i> (<i>Attractosteus</i>)	38
<i>Limanda</i>	227	<i>Mugil</i>	253
<i>Prionotus</i>	408	bermude, <i>Branchiostoma</i>	7
<i>Serrivomer</i>	84	<i>Chromis</i>	413
beanorum, <i>Pontinus</i>	372	<i>Fundulus</i>	175
beardslei, <i>Salmo</i>	57	<i>heteroclitus</i>	175
<i>Salmo gairdneri</i>	57	<i>Furcaria</i>	413
beckwithi, <i>Cyprinella</i>	130	bermudarum, <i>Callionymus</i>	450
becuna, <i>Sphyraena</i>	255	bermudensis, <i>Carapus</i>	486
bedoti, <i>Bairdiella</i>	246	<i>Lefroyia</i>	483
bedotis, <i>Sciaena</i> (<i>Bairdiella</i>)	246	bermudichthys	430
beldingii, <i>Cottus</i>	384	<i>subfurcatus</i>	430
belis, <i>Silurus</i>	149	bernardini, <i>Catostomus</i>	101
belizanus, <i>Belonesox</i>	184	berthelotii, <i>Cirrus</i>	203
belizianus, <i>Electris</i> (<i>Cultris</i>)	486	Berycidae	234
bella, <i>Hypoclydonia</i>	197	berycoides	234
Bellator	408	beryllina, <i>Menidia</i>	247
<i>egretta</i>	408	beryllina cerea, <i>Menidia</i>	247
<i>militaris</i>	408	beryllinum, <i>Christoma</i>	247
bellicus, <i>Nocomis</i>	187	beryllinus, <i>Cryptotomus</i>	431
bellus, <i>Balistes</i>	491	<i>Gobiesox</i> (<i>Rimicola</i>)	490
<i>Lythyrurus</i>	120	<i>Rimicola</i>	490
<i>Minnlius</i>	126	Beryx	234
Belone almeida	195	<i>decadactylus</i>	234
<i>altipinna</i>	197	<i>splendens</i>	234
<i>angusticeps</i>	198	Betarala	25
<i>ardeola</i>	195	<i>betarus</i> , <i>Cirrhitus</i>	358
<i>argalus</i>	198	<i>baculeatus</i> , <i>Gasterosteus</i>	237
<i>caribbea</i>	197	Bibronia	230
<i>cigonella</i>	195	<i>ilgulata</i>	230
<i>crassa</i>	196	bicaudalis, <i>Chapinus</i>	497
<i>depressa</i>	195	<i>Ostracion</i>	498, 497
<i>diplostenia</i>	198	<i>bicolor</i> , <i>Algansea</i>	122
<i>exilis</i>	198	<i>Ammocetes</i>	8
<i>galeata</i>	197	<i>Anisotremus</i>	332
<i>geranla</i>	196	<i>Barathronus</i>	484
<i>hilans</i>	197	<i>Chiopis</i>	82
<i>jonesi</i>	197	<i>Exocetus</i>	201
<i>latimanus</i>	197	<i>Grammiconotus</i>	197
<i>maculata</i>	197	<i>Pristipoma</i>	332
<i>melanochira</i>	198	<i>Rondeletia</i>	171
<i>microps</i>	196	<i>Rypticus</i>	316
<i>notata</i>	195	<i>Siphateles</i>	122
<i>pacificus</i>	197	<i>Smecticus</i>	316
<i>raphidoma</i>	196	<i>Tigoma</i>	120
<i>scutator</i>	196	bicornis, <i>Cottus</i>	378
<i>stolzmanni</i>	196	<i>Icelus</i>	378
<i>strongylura</i>	195	bifasciatum, <i>Thalassoma</i>	429
<i>subtruncata</i>	195	bifasciatus, <i>Cichlaurus</i>	416
<i>timucu</i>	195	<i>Heros</i>	416
<i>truncata</i>	196	<i>Labrus</i>	428, 429
Belonesox	184	<i>Mullus</i>	343
<i>belizanus</i>	184	<i>Upeneus</i>	343
Belonidae	195	bifasciatus torquatus, <i>Labius</i>	429
Belonocharax beanii	100	bifrenata, <i>Hybopsis</i>	135
Belonopsis	83	bifrenatus, <i>Hybopsis</i>	133, 135
<i>leuchtenbergii</i>	83	<i>bibbus</i> , <i>Kyphosus</i>	339
bellus, <i>Minnlius</i>	126	<i>biguttatus</i> , <i>Cochleognathus</i>	146
bendirei, <i>Cottus</i>	384	<i>Labrisomus</i>	460
<i>Potomacottus</i>	384	<i>Malacoctenus</i>	460
benedicti, <i>Bathylagus</i>	69	<i>Semotilus</i>	137
bengalensis, <i>Holocentrus</i>	324	bilinearis, <i>Merluccius</i>	214
benoiti, <i>Myctophum</i>	169	<i>Stomodon</i>	214
<i>Scopelus</i>	169	bilineata, <i>Lepidopsetta</i>	228
Benthocometes	482	<i>Platesa</i>	228
<i>robustus</i>	482	<i>Skiffia</i>	183
Benthodesmus	262	<i>Umbrina pygmaea</i>	174
<i>atlanticus</i>	262	bilineatum, <i>Pristipoma</i>	332
<i>elongatus</i>	262	bilineatus, <i>Charodon</i>	183
Benthosauridae	166	<i>Doryichthys</i>	243
Benthosaurus	166	billingsiana, <i>Cyprinella</i>	131
<i>grallator</i>	166	biloba, <i>Corvina</i>	349
Benthosema	170	<i>bilobus</i> , <i>Blepsias</i>	382
<i>arcticum</i>	170	<i>Histiophotus</i>	382
<i>mulleri</i>	170	<i>bimaculata</i> , <i>Percina</i>	281
berardi, <i>Aluterus</i>	496	<i>bimaculatus</i> , <i>Chatodon</i>	360
bergalli, <i>Ctenolabrus</i>	424	<i>Clinus</i>	460
bergii, <i>Pontinus</i>	372	<i>Halichoeres</i>	426
<i>Scorpaen</i>	372	<i>Malacoctenus</i>	460
bergla, <i>Macrourus</i>	205	<i>Poecilodiodes</i>	187
		<i>Pseudophorophorus</i>	187

List of scientific names—Continued

	Page		Page
bimaculatus Sayris.	197	Blennius gunnellus.	471
Xiphophorus.	187	hentz.	462
bimaculatus Jonesi. Pseudoxiphophorus.	188	herminieri.	459
bimaculatus tenuistius. Pseudoxiphophorus.	188	lævis.	461
binghami. Xyrichthys.	430	lampetraformis.	470, 471
binocula, Raja.	26	lumpenus.	470
binotata, Amia.	295	marmoratus.	461
binotatus, Apogon.	295	microstomus.	461
Bipinnula.	260	multifilis.	461, 462
bipinnulata, Seriota.	279	murænoides.	467
bipinnulata, Elagatis.	279	nuchifilus.	461
birgei, Leucichthys.	64	oceanicus.	461
birostratus, Gurnardus.	406	ocellaris.	460
Prionotus.	406	opholis.	461
birostris, Manta.	22	pilicornis.	461
Raja.	32	polaris.	476
biserialis, Ophichthys.	88	polyactecephalus.	466
Ophichthus.	88	punctatus.	462, 469
biseriatus, Archaulus.	377	regius.	213
bishop, Aloa.	43	salfensis.	463
bison, Aspicottus.	381	serpentinus.	470, 471
Carpoides.	103	stearnsii.	460
Lepisosteus.	37	striatus.	462
bispinacis, Gasterosteus.	233	tænia.	468
Gladiunculus.	233	truncatus.	461
Melanophæs.	233	vinctus.	461
Melichthys.	492	blennius, Alburnops.	133
Myloobatidae.	31	Etheostoma (Rhothocæa).	287
Plectromus.	233	Hybopsis.	133
bisselli, Coregonus tullibee.	63	Ulocentra.	287
Leucichthys artedi.	63	Blennophis.	404
bis-trispinus, Bodianus.	316	webbi.	464
bistrispinus, Bodianus.	316	Bleparichthys.	274
Promicropterus.	316	Blepharis.	274
bisus, Scomber.	250	major.	274
bitemnatus, Eumycteris.	500	sutor.	274
Bivittata, Elacate.	281	blepharis, Carangoides.	274
Iridio.	426	blephure, Anguilla.	77
bivittatus, Hybopsis.	119	Blepsis.	382
Labrus.	426	bilobus.	382
Serranus.	318	cirrhosus.	382
blackfordi, Lutjanus.	320	oculofasciatus.	382
Lutjanus.	326	trilobus.	382
Neomenis.	453	ventricosus.	380, 381
Yarrella.	72	Blepsidae.	382
blackfordii, Lutjanus.	326	blochii, Bodianus.	425
Lutjanus.	326	Caranx.	271
Blakea.	458	Orthragoriscus.	503
blanchardi, Gasterosteidae.	238	Piramutana.	159
Neoclinus.	468	boa, Esox.	69
bleekeriana, Chirocentrodon.	46	bocourti, Herichthys.	423
Pellona.	46	Neotropius.	423
Bleekeridae.	451	Bodianus.	300, 425
Blennitrichus.	462	acanthistius.	300
Blenniocottus.	391	achigan.	298
clarki.	377	albostriatus.	326
globiceps.	301	apua.	313
bryosus.	391	argyroleucos.	345
recalvus.	391	argyro-leucos et exiguis.	345
Bleniidae.	460	bistrispinus.	316
blennioides, Etheostoma.	283, 286	bis-trispinus.	316
Blenniolus.	461	blochii.	425
blennioperca, Hyostoma.	286	bodianus.	309, 425
Blennius.	460	costatus.	349
americanus.	478	diplostenia.	425
anguillaris.	470, 478	fasciatus.	325
asterias.	461	guativera.	309
asterius.	461	guttatus.	309
bosquianus.	462	jaguar.	235
brevipinnis.	461	marginatus.	313
chiaso.	212	pallidus.	345
chuss.	213	pentacanthus.	235
ciliatus.	473	pulchellus.	425
crinitus.	461	rufus.	425
cristatus.	461	rupestris.	303
dolichogaster.	468	stellatus.	309
euroopeus.	467	stelleri.	346
fabri.	460	striatus.	325
favosus.	461	triourus.	323
fimbriatus.	473	vivanei.	325
fucorum.	461	bodianus, Bodianus.	309, 425
galerita.	460	beckei, Callionymus.	450
geminatus.	461, 462	Doratonotus.	429
gentilis.	462	Melanorhinus.	249
gracilis.	471	Moringua.	59

List of scientific names—Continued

	Page		Page
<i>Bogmarus</i>	218	<i>borealis</i> , <i>Notorhynchus</i>	11
<i>islandicus</i>	218	<i>Oligocottus</i>	390
<i>bogmarus</i> , <i>Trachypterus</i>	218	<i>Opsopoeodus</i>	116
<i>Bogoslovius</i>	204	<i>Paralepis</i>	165
<i>clarki</i>	204	<i>Petromyzon</i>	10
<i>fimisquamis</i>	204	<i>Pimelodus</i>	152
<i>Boleichthys</i>	293	<i>Poecilichthys</i>	291
<i>eos</i>	294	<i>Sphyraena</i>	255
<i>exilis</i>	293, 294	<i>Squalus</i>	21, 22
<i>fusiformis</i>	294	<i>Boreocottus</i>	387
<i>gracilis</i>	294	<i>axillaris</i>	387
<i>wareni</i>	294	<i>Boreogadus</i>	209
<i>whippili</i>	292	<i>salda</i>	209
<i>boleoides</i> , <i>Cottus</i>	385	<i>Boreogaleus</i>	14
<i>Radulinus</i>	370	<i>boreus</i> , <i>Esox</i>	173
<i>Boleosoma</i>	287, 288	<i>Borphryne</i>	509
<i>aesopus</i>	287	<i>Borborys</i>	509
<i>camurum</i>	288	<i>boschma</i> , <i>Pomadasys</i>	334
<i>effulgens</i>	287	<i>Pristipoma</i>	334
<i>fusiforme</i>	294	<i>boscii</i> , <i>Abramsi</i>	350
<i>gracile</i>	294	<i>Gobiosoma</i>	446
<i>leptida</i>	290	<i>Gobius</i>	442, 445, 446
<i>maculaticeps</i>	287	<i>Notemigonus</i>	446
<i>maculatum</i>	287	<i>Pimeleopterus</i>	339
<i>meseum</i>	287	<i>sicula</i> , <i>Pimeleopterus</i>	340
<i>nigrum</i>	287	<i>boscii</i> , <i>Atherina</i>	247
<i>olmstedi</i>	287	<i>Gobius</i>	442
<i>brevipinnis</i>	287	<i>Leuciscus</i>	115
<i>phlox</i>	287	<i>Seriola</i>	280
<i>stigmatum</i>	287	<i>bosquianus</i> , <i>Biennius</i>	462
<i>susane</i>	288	<i>Chasmodes</i>	462
<i>tassellatum</i>	287	<i>bosquianus</i> , <i>Chasmodes</i>	460
<i>tessellatum</i>	285, 287	<i>bosquii</i> , <i>Pimeleopterus</i>	340
<i>verillare</i>	288	<i>bostoniensis</i> , <i>Catostomus</i>	105
<i>bolemomn</i> , <i>Gobius</i>	439	<i>Muraena</i>	77
<i>Rhinogobius</i>	430	<i>Bothidæ</i>	217
<i>bolli</i> , <i>Ameiurus</i>	163	<i>Bothragonus</i>	396
<i>bolmani</i> , <i>Hippoglossina</i>	223	<i>swanii</i>	396
<i>Opsoprea</i>	115	<i>Bothrocænæ</i>	477
<i>Opsopœodus</i>	115	<i>mollis</i>	477
<i>Scarus</i>	433	<i>pusilla</i>	477
<i>Bollmannia</i>	443	<i>remigera</i>	477
<i>boqueronensis</i>	443	<i>Bothrocæropsis</i>	477
<i>chlamydes</i>	443	<i>alalonga</i>	477
<i>macropoma</i>	443	<i>elongata</i>	477
<i>ocellata</i>	443	<i>rictolata</i>	477
<i>stigmatura</i>	443	<i>Bothrolæmus</i>	276
<i>bombifrons</i> , <i>Heloperca</i>	301	<i>Bothrus</i>	217
<i>Pomotis</i>	301	<i>boucardi</i> , <i>Leuciscus</i>	132
<i>Bonaci cardenal</i>	313	<i>Moniana</i>	132
<i>bonaci</i> , <i>Serranus</i>	314	<i>Poecilia</i>	192
<i>Trisotropis</i>	314	<i>Pristipoma</i>	334
<i>xanthosticta</i> , <i>Mycteroperca</i>	314	<i>bougainvillie</i> , <i>Sebastæ</i>	370
<i>bonapartei</i> , <i>Lampanyctus</i>	167	<i>boulengeri</i> , <i>Batrachoides</i>	487
<i>Poecilosephalus</i>	87	<i>Mycteroperca</i>	313
<i>Bonapartia</i>	72	<i>Plecostomus</i>	160
<i>pedaliota</i>	72	<i>Trisotropis</i>	313
<i>bonariense</i> , <i>Haemulon</i>	329	<i>bouvieri</i> , <i>Salmo</i>	56
<i>bonariensis</i> , <i>Seriola</i>	280	<i>Salmo purpuratus</i>	56
<i>bonasus</i> , <i>Bulbilichthys</i>	102	<i>bovinum</i> , <i>Plectropoma</i>	321
<i>Rhinoptera</i>	31	<i>bovinus</i> , <i>Cyprinodon</i>	181
<i>bonita</i> , <i>Gambusia</i>	188	<i>Hoploplectrus unicolor</i>	321
<i>Præpella</i>	188	<i>rubrofutialis</i> , <i>Cyprinodon</i>	181
<i>Boops</i>	417	<i>bowersi</i> , <i>Labichthys</i>	84
<i>boops</i> , <i>Ceranx</i>	272	<i>Mycteroperca</i>	314
<i>Hybopsis</i>	134	<i>Rhinichthys</i>	139
<i>Macrourus</i>	205	<i>Bowersi</i>	328
<i>Macrurus</i>	205	<i>violescens</i>	328
<i>Myctophum</i>	169	<i>bowersianus</i> , <i>Careproctus</i>	403
<i>Notropis</i>	134	<i>bowmani</i> , <i>Plargyrus</i>	128
<i>Ostracion</i>	503	<i>Box</i>	417
<i>Scomber</i>	270	<i>boyeri</i> , <i>Atherina</i>	246
<i>Trachurus</i>	272	<i>brachiale</i> , <i>Sparisoma</i>	433
<i>boqueronensis</i> , <i>Bollmannia</i>	272	<i>brachialis</i> , <i>Baiostoma</i>	230
<i>borea</i> , <i>Lutiperca</i>	443	<i>Scarus</i>	433
<i>borealis</i> , <i>Alepidosaurus</i>	282	<i>brachiatius</i> , <i>Diodon</i>	501
<i>Ammocetes</i>	9	<i>Brachioptilon</i>	32
<i>Arctozenus</i>	165	<i>hamiltoni</i>	32
<i>Bathylagus</i>	89	<i>brachiusculus</i> , <i>Grammicolepis</i>	232
<i>Caulopus</i>	171	<i>brachyacanthus</i> , <i>Amelurus</i>	154
<i>Flerasfer</i>	471	<i>Amilurus</i>	154
<i>Icelinus</i>	376	<i>brachycentrus</i> , <i>Nauclerus</i>	278
<i>Lampetra</i>	10	<i>brachycephala</i> , <i>Rhamdia</i>	157
<i>Molvella</i>	208		
<i>Nivicolæ</i>	291		

List of scientific names—Continued

	Page		Page
brachycephalus, <i>Exocetus</i> .	200	Brachyrhaphichthys.	101
Pimelodus.	157	Brachyrhaphis.	186
Syngnathus.	241	cascajalensis.	187
Uranichthys.	87	episcopi.	187
Bramidae.	265	olomius.	187
Bramiformes.	265	parismina.	187
Bramocharax.	69	rhabdophora.	188
bransfordi.	99	terrabensis.	188
elongatus.	99	Brachyrhinus.	320
Bramops.	410	brachyrhynchus, <i>Acipenser</i> .	34
mento.	410	Rhynchichtys.	235
branchialis, <i>Ammocetus</i> .	8	brachysoma, <i>Ephippophelus</i> .	311
branchiomelas, <i>Anchoviella</i> .	49	Brachysomophis.	89
Stelephorus.	49	crocodilinus.	89
Branchiostegidae.	357	horridus.	89
Branchiostoma.	7	brachysomus, <i>Calamus</i> .	337
bermude.	7	Sparus.	337
californense.	7	brachyurus, <i>Oxydorlichthys</i> .	88
caribaeum.	7	bradfordi, <i>Porocottus</i> .	388
floridæ.	7	Brama.	265
lanceolatum.	7	agassizii.	265
lubricum.	7	brevirostris.	265
pelagicum.	7	chilensis.	265
plate.	7	dussumieri.	265
virginie.	7	marina cauda forcipata.	265
Branchiostomidae.	7	orcinus.	265
Branchistegus.	357	parra.	428
Branchyconger.	78	raii.	265
brandieriana, <i>Cæcilia</i> .	84	saussurii.	265
Branderup.	84	brahma, <i>Cynedus</i> .	338
braadtii, <i>Arius</i> .	148	brasiliensis, <i>Torpido</i> .	27
bralecki, <i>Pristipoma</i> .	334	Vomer.	276
Rhönchicus.	334	brasiliensis corallina, <i>Narcine</i> .	27
Branneroella.	404	braytoni, <i>Notropis</i> .	135
brasiliensis.	404	Bregmaceros.	208
sluiteri.	464	atlanticus.	208
bransfordi, Bramocharax.	99	bathyraja.	208
Loricaria.	102	longipes.	208
Rhamdia.	157	macrolellandii.	208
brasilianus, <i>Eugerres</i> .	342	Bregmacerotidae.	208
Geres.	342	breve, <i>Chirostoma</i> .	250
brasiliense, <i>Pristipoma</i> .	332	brevibarbe, <i>Lepophidium</i> .	484
brasiliensis, <i>Caepuna</i> .	331	Ophidium.	484
Guabiocoara.	330	brevibarbis, <i>Chalinura</i> .	204
brasiliensis, <i>Brannerella</i> .	404	brevicauda, <i>Pomoxys</i> .	304
Centropristis.	320	Salmo.	56
Clupea.	40	brevicaudata, <i>Brevoortia tyrannus</i> .	44
Conger.	78	brevicaudatum, <i>Siphonstoma</i> .	241
Erythrinus.	99	brevicaudus, <i>Diplanichthys</i> .	79
Esox.	109	Syngnathus.	241
Gobius.	448	Breviceps.	147
Hemiramphus.	199	filamentosus.	147
Hippoglossus.	224	breviceps, <i>Evorthodus</i> .	438
Hypostomus.	160	Gasterosteus.	239
Lebris.	426	Gobius.	440
Mugil.	223	Larinus.	344
Narcine.	27	Moxostoma.	111
Paralichthys.	224	Pomacentrus.	413
Plagiusa.	231	Pomotis.	300
Scorpaena.	371	Porogadus.	482
Thynnus.	258	Ptychostomus.	111
brachychira, <i>Trachurops</i> .	271	brevifilis, <i>Agenelosus</i> .	169
brachychirus, <i>Trachurops</i> .	271	brevimanus, <i>Eugerres</i> .	343
Brachydeuterus.	333	Geres.	343
auritus.	333	Tetragonopterus.	96
axillaris.	333	brevipes, <i>Lycodes</i> .	475
corviniformis.	333	brevipinnis, <i>Somniosus</i> .	21, 22
leuciscus.	333	Ditrema.	409
nitidus.	333	Pristipoma.	335
Brachygenys.	330	(Hæmulopsis).	336
chrysargyreus.	331	brevipinnis, <i>Bleennius</i> .	461
Brachylipus.	409	Microlepidotus.	335
frenatus.	409	Thynxus.	258
Brachymimus.	343	brevirostre, <i>Peristedion</i> .	405
Brachyopstis verrucosus.	394	Peristethus.	405
xyosternus.	394	Pristipoma.	34
brachypoda, <i>Gasterosteus pungitius</i> .	239	Chasmistes.	108
Pungitius.	239	Colobabis.	198
Brachyprosopon.	228	Engraulis.	51
brachyptera, <i>Echeneis</i> .	449	Geres.	342
Remoraopsis.	449	Hypopomus.	101
brachypterus, <i>Holocentrus</i> .	236	Hypoprion.	17
Lutjanus.	326	Macrognathus.	199
Lutjanus.	326	Rhamphichthys.	101
Pimelodus.	157	Rhynchoceratias.	511
Zygonectes (<i>Microstius</i>)	185	Saurus.	163

List of scientific names—Continued

Page	Page
brevirostris, <i>Scombrasox</i>	198
<i>Syngnathus</i>	240
brevirostrum, <i>Acipenser</i>	34
<i>Anchovia</i>	51
<i>Haemulon</i>	330
<i>Stolephorus</i>	50
brevis, <i>Acanthurus</i>	362
<i>Amblyopus</i>	448
<i>Atherinichthys</i>	250
<i>Campostoma</i>	146
<i>Centropomus</i>	306
<i>Dasyatis</i>	29
<i>Engraulis</i>	51
<i>Fundulus parvipinnis</i>	176
<i>Tyntastes</i>	448
<i>Xiphophorus</i>	104
<i>helleri</i>	194
brevispina, <i>Ricchia</i>	293
<i>Richella</i>	293
brevispine, <i>Acutomentum</i>	366
brevispinis, <i>Sebastichthys proriger</i> var.....	366
brevortii, <i>Argyrelosus</i>	276
<i>Argyrius</i>	276
<i>Brama</i>	265
<i>Euleptorhamphus</i>	198
<i>Poecilurichthys</i>	96
brevortii, <i>Poecilurichthys</i>	97
Brevoortia.....	44
<i>patronus</i>	44
<i>tyrannus</i>	44
<i>breviceaudata</i>	44
brevortii, <i>Astyanax</i>	97
<i>Euleptorhamphus</i>	198
briareos, <i>Chilomycterus</i>	502
<i>Cyclichthys</i>	602
bricel, <i>Chetodon</i>	359
brimleyi, <i>Coccotis</i>	127
<i>Notropis</i>	127
bristolensis, <i>Cyclogaster</i>	399
neoliparis, <i>Neoliparis</i>	399
broccus, <i>Balistes</i>	494
broadmus, <i>Cottus</i>	395
Brontes.....	150
<i>prenadilla</i>	159
Brosme.....	213
<i>brosme</i>	214
brosme, <i>Brosme</i>	214
<i>Gadus</i>	213, 214
Brosmerus.....	213
<i>flavesny</i>	213, 214
brosmane, <i>Lota</i>	212
Brosnius.....	213
<i>marginatus</i>	479
Bromophycis.....	479
<i>marginatus</i>	479
<i>ventralis</i>	480
<i>verrillii</i>	480
Brotula.....	479
<i>barbata</i>	479
Brotulidae.....	479
broussoneti, <i>Acanthurus</i>	362
broussonetti, <i>Gobiooides</i>	448
<i>Umbrina</i>	360
browni, <i>Lucania</i>	181
<i>Solea</i>	229
brownii, <i>Atherina</i>	48
<i>Hemiramphus</i>	199
Vomer.....	274, 275
brucus, <i>Echinorhinus</i>	22
Squalus.....	22
brunnea, <i>Lycogramma</i>	476
<i>Maynei</i>	476
brunneus, <i>Amiurus</i>	154
<i>Apristurus</i>	12
Catulus.....	12
Ectenias.....	285
Gobius.....	440
Gymnothorax.....	92
Hippocampus.....	244
Ilyophis.....	78
Rivulus.....	180
Serranus.....	314
Brycon.....	98
<i>argenteus</i>	98
<i>cephalus</i>	98
<i>dentex</i>	98
<i>falcatus</i>	98
<i>guatemalensis</i>	98
<i>oligolepis</i>	99
<i>petrosus</i>	98
<i>striatus</i>	98
Bryconamericus.....	97
<i>cascajalensis</i>	97
<i>emperador</i>	97
<i>exodon</i>	97
Bryopholus.....	466
<i>lysinus</i>	466
bryoporus, <i>Spatelloides</i>	41
Bryostemma.....	466
<i>decoratum</i>	466
<i>nugator</i>	466
<i>polyactocephalum</i>	466
<i>tarsodes</i>	466
Bryssateres.....	488
<i>pinniger</i>	488
Bryssophilus.....	488
Brytus.....	298
<i>albus</i>	302
<i>fuscatus</i>	304
<i>gloriosus</i>	304
<i>humilis</i>	300
<i>mineopas</i>	299
<i>oculatus</i>	300
<i>punctatus</i>	298, 299
<i>reticulatus</i>	299
<i>signifer</i>	299
<i>unicolor</i>	299
Bunalfichthys.....	102
<i>altus</i>	102
<i>bonasus</i>	102
<i>bubalinus</i>	102
<i>niger</i>	102
<i>urus</i>	102
bubalina, <i>Cyprinella</i>	130
bubalinus, <i>Bubalichthys</i>	102
<i>Leuciscus</i>	130
<i>bubalis</i> , <i>Acanthocottus</i>	386
<i>Amblodon</i>	102
<i>Cottus</i>	386
<i>Ichthyobus</i>	102
<i>Ictiobus</i>	326
<i>buccanella</i> , <i>Lutjanus</i>	326
<i>Mesoprlion</i>	326
<i>buccata</i> , <i>Ericymba</i>	141, 142
<i>bucciferus</i> , <i>Labrisomus</i>	459
<i>Labrosomus</i>	459
<i>bucco</i> , <i>Maxostoma</i>	110
<i>Ptychostomus</i>	110
<i>Buccone</i>	354
<i>praedatoria</i>	354
<i>bucculentus</i> , <i>Chonophorus</i>	442
<i>bucephalus</i> , <i>Macrourus</i>	205
<i>Macrurus</i>	205
<i>buchananii</i> , <i>Hybopsis</i>	134
<i>Notropis</i>	134
<i>buettikoferi</i> , <i>Tilapia</i>	423
<i>buffalo</i> , <i>Acipenser</i> (<i>Antaeus</i>).....	35
<i>bufo</i> , <i>Lophius</i>	487
<i>Scorpaena</i>	370
<i>bulbiceps</i> , <i>Macrourus</i>	205
<i>Macrurus</i>	205
<i>Maynei</i>	477
<i>bullaris</i> , <i>Cyprinus</i>	117
<i>bulleri</i> , <i>Paraneetroplus</i>	423
<i>Prionodes</i>	319
<i>Serranus</i>	319
<i>buniva</i> , <i>Ballistes</i>	491, 493
<i>burchami</i> , <i>Icelinus</i>	376
<i>burgali</i> , <i>Ctenolabrus</i>	424
<i>burragei</i> , <i>Baja California</i>	53
<i>burtonianus</i> , <i>Leuciscus</i>	121
<i>buscki</i> , <i>Sicydium</i>	447
<i>butleri</i> , <i>Poecilia</i>	192
<i>butlerianus</i> , <i>Poecilichthys</i>	294

List of scientific names—Continued

	Page		Page
Butyrinus.....	39	californiensis, <i>Dipterus</i>	341
banana.....	39, 40	<i>Coryphampus</i>	243
Bythites.....	480	<i>elongatus</i> , <i>Eucinostomus</i>	341
fuscus.....	480	<i>Eucinostomus</i>	341
caballia, <i>Cybium</i>	257	<i>Medialuna</i>	358
Caballeroite.....	325	<i>Ophisurus</i>	88
caballeroite, <i>Anthias</i>	325	<i>Otolithus</i>	356
caballus, <i>Caranx</i>	272	<i>Poly nemus</i>	256
<i>Paratractus</i>	272	<i>Scorpi</i>	358
caboverdianus, <i>Ginglymostoma</i>	12	<i>Syngnathus</i>	240
cabrerae, <i>Rhamdia</i>	167	<i>Typhlogobius</i>	447
Cabilla.....	311	<i>Xenichthys</i>	324
Cæcilia.....	84	<i>Xenistius</i>	324
Cæcorhamphia.....	188	callarias, <i>Gadus</i>	210
urichi.....	188	<i>Salmo</i>	61
cærulea, <i>Cheonoda</i>	120	Callaus.....	348
<i>Clupea</i>	41	<i>delleciosa</i>	348
<i>Coryphena</i>	434	<i>Callechelys</i>	86
<i>Erogala</i>	130	<i>guichenoti</i>	86
<i>Etheostoma</i>	291	<i>muraena</i>	86
<i>Melletta</i>	43	<i>peninsulae</i>	88
<i>Sardinia</i>	43	Callichthys.....	160
cæruleatus, <i>Acronurus</i>	362	<i>Callichthys (Holosternum) thoracatus</i>	160
cæruleoauereum, <i>Harpie</i>	425	<i>levigatus</i>	160
cæruleofasciatus, <i>Bathymaster</i>	451	<i>littoratus</i>	160
cæruleoceans, <i>Arius</i>	160	Callidulus.....	320
<i>Galeichthys</i>	150	<i>flaviventris</i>	320
cæruleus, <i>Acanthurus</i>	362	Callionodon retractus.....	431
<i>Carcharias</i>	16	Callionymidae.....	450
<i>Oligocephalus</i>	289, 291	<i>Callionymus</i>	450
<i>Photogenis</i>	130	<i>agassizii</i>	450
<i>Scarus</i>	434	<i>atrilabiatus</i>	450
<i>Squalus</i>	17	<i>bairdi</i>	450
Cæsiomorus.....	276	<i>bermudarium</i>	450
<i>balloni</i>	276	<i>boekeli</i>	450
cæsius, <i>Paraconodon</i>	331	<i>callurus</i>	450
<i>Pomadasys</i>	331	<i>himantophorus</i>	450
caguita, <i>Scydium</i>	447	<i>lyra</i>	450
cablepta, <i>Dermatocheil</i>	97	<i>pauciradiatus</i>	450
Calamus.....	236	<i>pelagicus</i>	450
<i>arcifrons</i>	237	<i>sanci-eustati</i>	450
<i>baionado</i>	236	callipteryx, <i>Campostoma</i>	146
<i>brachysomus</i>	237	callisema, <i>Episema</i>	132
<i>calamus</i>	236	<i>Moniana</i>	132
<i>kendalli</i>	237	callisoma, <i>Herpetolichthys</i>	88
<i>leucosteus</i>	237	callistia, <i>Erogala</i>	130
<i>macrops</i>	237	callistius, <i>Photogenis</i>	130
<i>medius</i>	237	calliura, <i>Cyprinella</i>	129
<i>megacephalus</i>	236	<i>Etheostoma</i>	298
<i>penna</i>	237	<i>Mycteroperca</i>	315
<i>peniculata</i>	238	Callurus.....	298
<i>plumatus</i>	236	<i>disparitus</i>	299
<i>proridens</i>	236	<i>floridensis</i>	303
calamus, <i>Calamus</i>	237	<i>formosus</i>	299
<i>Pagellus</i>	236	<i>melanops</i>	302, 303
calcarata, <i>Scorpæna</i>	271	<i>micros</i>	299
calcea, <i>Acanthidium</i>	21	<i>murinus</i>	298
caliente, <i>Xenendum</i>	184	<i>punctulatus</i>	298
caliente, <i>Goodea</i>	184	callurus, <i>Callionymus</i>	450
<i>Hybopsis</i>	134	<i>loglossus</i>	438
<i>Notropis</i>	134	callotopis, <i>Harengula</i>	43
california, <i>Alausa</i>	43	<i>Heros</i>	422
<i>Mayneia</i>	477	<i>Thorichthys</i>	422
<i>Morrhus</i>	210	callopterus, <i>Cypselurus</i>	202
<i>Squatina</i>	22	<i>Exocetus</i>	202
<i>Tetranarce</i>	27	Calloptilum.....	208
<i>Torpedo</i>	27	<i>mirrum</i>	208
californicus, <i>Arotatus</i>	31	callorhini, <i>Therobromus</i>	68
<i>Cyprinurus</i>	202	Callorhynchus atlanticus.....	32
<i>Exocetus</i>	202	<i>centrina</i>	32
<i>Halichoeres</i>	428	Callyodon.....	433, 434
<i>Hippoglossus</i>	223	<i>auro-punctatus</i>	431
<i>Mustelus</i>	13	<i>compressus</i>	434
<i>Myliobatis</i>	31	<i>micros</i>	435
<i>Myliobatus</i>	31	<i>ustus</i>	431
<i>Oryzius</i>	428	callyodon, <i>Cyclopterus</i>	390
<i>Paralichthys</i>	223	<i>Neoliparis</i>	390
<i>Stereolepis</i>	308	calobrese, <i>Cichlasoma</i>	417
californiense, <i>Branchiostoma</i>	7	<i>calobrese</i> , <i>Cichlaurus</i>	417
<i>Myotophum</i>	169	calopteryx, <i>Serranus</i>	319
californiensis, <i>Atherinopsis</i>	252	Calotomus.....	431
<i>Chilomycterus</i>	602	<i>xenodon</i>	431
<i>Cyprinodon</i>	181		

List of scientific names—Continued

	Page		Page
calva, <i>Amia</i>	38	canus, <i>Macrurus</i>	206
<i>Amiatus</i>	38	<i>Macrurus</i>	206
<i>Calyclepidotus</i>	381	<i>Capellaria</i>	505
<i>spinosus</i>	381	<i>capellei</i> , <i>Lophotes</i>	215
<i>Camarina</i>	338	<i>capense</i> , <i>Scyllium</i>	11
<i>nigricans</i>	338, 339	<i>capensis</i> , <i>Carcharodon</i>	20
<i>camchaticus</i> , <i>Artediellus</i>	378	<i>capensis</i> , <i>Loricaria</i>	162
<i>camelopardalis</i> , <i>Mycteroperca</i>	315	<i>Capeuna brasiliensis</i>	331
<i>Serranus</i>	315	<i>capeuna</i> , <i>Serranus</i>	331
<i>campechanus</i> , <i>Lutjanus</i>	328	<i>capillaris</i> , <i>Zeus</i>	275
<i>Mesopriion</i>	328	<i>capillatus</i> , <i>Clinus</i>	459
<i>campbelli</i> , <i>Moxostoma</i>	109	<i>capistratus</i> , <i>Balistes</i>	492
<i>Salmo</i>	61	<i>Chaetodon</i>	359
<i>camperi</i> , <i>Scomberesox</i>	197	<i>Tetradon</i>	500
<i>camperi</i> , <i>Scomberesox</i>	197	<i>capite plagioplateo</i> , <i>Eleotris</i>	436
<i>campi</i> , <i>Signalosa mexicana</i>	47	<i>striato</i> , <i>Perca marina</i>	330
<i>Campostoma</i>	146	<i>capito</i> , <i>Macrurus</i>	205
<i>anomalum</i>	146	<i>Macrurus</i>	205
<i>brevis</i>	146	<i>Poromitra</i>	223
<i>callipteryx</i>	146	<i>caprinus</i> , <i>Balistes</i>	491
<i>formulosum</i>	147	<i>Otrynter</i>	336
<i>gobionum</i>	146	<i>Stenotomus</i>	336
<i>hippos</i>	146	<i>Caprichthys</i>	491
<i>mormyrus</i>	146	<i>minima</i>	496
<i>nasutum</i>	146	<i>porcus</i>	491
<i>ornatum</i>	146	<i>caprichthys</i> , <i>Balistes</i>	491
<i>princei</i>	147	<i>caprodes</i> , <i>Etheostoma</i>	282
<i>Campylodon</i>	75	<i>Percina</i>	283
<i>fabrieli</i>	75	<i>Sciæna</i>	282, 283
<i>camtharinum</i> , <i>Pristipoma</i>	335	<i>Caphrophorus</i>	358
<i>cantschaticus</i> , <i>Crossias</i>	389	<i>aurora</i>	358
<i>camura</i> , <i>Clœla</i>	120	<i>capros</i> , <i>Antigonia</i>	358
<i>Erogala</i>	120	<i>captiva</i> , <i>Goodea</i>	184
<i>Vallantia</i>	288	<i>caracasesis</i> , <i>Hemiloricaria</i>	161
<i>camurum</i> , <i>Bolesoma</i>	288	<i>caramuru</i> , <i>Muraenophis</i>	91
<i>Etheostoma</i>	292	<i>Carangidae</i>	239, 269
<i>camurus</i> , <i>Nothonotus</i>	290	<i>Carangichthys</i>	273
<i>Pœciliichthys</i>	290	<i>typus</i>	273
<i>cana</i> , <i>Allogamibia</i>	188	<i>Carangooides blepharis</i>	274
<i>Gambusia</i>	188	<i>dorsalis</i>	274
<i>canadensis</i> , <i>Cynoperca</i>	282	<i>gallichthys</i>	274
<i>Salmo</i>	60	<i>carangooides</i> , <i>Seriolophus</i>	280
<i>canadensis</i> , <i>Lucoperca</i>	282	<i>Carangops</i>	271
<i>canadus</i> , <i>Gasterosteus</i>	281	<i>Carangus aureus</i>	272
<i>Rachycentron</i>	281	<i>esculentus</i>	273
<i>canalculatus</i> , <i>Icelus</i>	378	<i>hippolyoides</i>	273
<i>canariensis</i> , <i>Clinus</i>	459	<i>carangus</i> , <i>Scomber</i>	273
<i>Pteraplatea</i>	29	<i>Caranx</i>	271, 273
<i>cancrivorus</i> , <i>Pisodonophis</i>	85, 86	<i>amblyrhynchus</i>	271
<i>candalerius</i> , <i>Profundulus</i>	174	<i>antillarum</i>	273
<i>candidissimus</i> , <i>Leptocephalus</i>	79	<i>atrimanus</i>	271
<i>candidus</i> , <i>Careproctus</i>	403	<i>aureus</i>	270, 272
<i>canensis</i> , <i>Acanthicus</i>	161	<i>bartholomaei</i>	272
<i>Leptoancistrus</i>	161	<i>beani</i>	272
<i>canescens</i> , <i>Ancanthias</i>	161	<i>biocellii</i>	271
<i>canicula</i> , <i>Squalus</i>	11	<i>caballus</i>	272
<i>canina</i> , <i>Amia</i>	38	<i>chrysurus</i>	345
<i>caninianus</i> , <i>Scopelinus</i>	169	<i>cibi</i>	272
<i>caninus</i> , <i>Lechnolaimus</i>	425	<i>daubentonii</i>	273
<i>Pajellus</i>	336	<i>defensor</i>	273
<i>canis</i> , <i>Cynias</i>	13	<i>dumerili</i>	279
<i>Salmo</i>	54	<i>erithrurus</i>	273
<i>Squalus</i>	13	<i>esculentus</i>	273
<i>canna</i> , <i>Hæmulon</i>	329	<i>falciatus</i>	271
<i>Cannorhynchus</i>	245	<i>fallax</i>	272
<i>Cantharus nigro-maculatus</i>	305	<i>fasciatus</i>	272
<i>Cantherines</i>	493	<i>frontalis</i>	273
<i>amphioxys</i>	494	<i>fürthii</i>	271
<i>carolæ</i>	494	<i>girardi</i>	272
<i>nasutus</i>	493, 494	<i>hasselti</i>	270
<i>pullus</i>	494	<i>heterophygus</i>	271
<i>Cantidermis</i>	492	<i>hippopodus</i>	273
<i>angulosus</i>	492	<i>hippos</i>	273
<i>maculatus</i>	493	<i>ignobilis</i>	273
<i>sabaco</i>	492	<i>ridibulus</i>	271
<i>sufflamen</i>	492	<i>kurra</i>	270
<i>wilughbali</i>	493	<i>latus</i>	272
<i>Canthigaster</i>	500	<i>lepturus</i>	272
<i>punctatissimus</i>	500	<i>leucurus</i>	271
<i>rostratus</i>	500	<i>lugubris</i>	273
<i>Canthigasteridae</i>	500	<i>macarellus</i>	270
<i>Canthirhynchus</i>	397	<i>macrophthalmus</i>	271
<i>Canthogaster lobatus</i>	498	<i>marginatus</i>	272
<i>cantrainei</i> , <i>Tylosurus</i>	197	<i>medusicola</i>	272
<i>cantrini</i> , <i>Tylosurus</i>	196	<i>melampygus</i>	272

List of scientific names—Continued

	Page		Page
<i>Caranx orthogrammus</i>	274	<i>Carcharius isodon</i>	15
otrynter	274	punctatus	15
<i>panamensis</i>	273	<i>carcharius</i> , <i>Carcharodon</i>	15
<i>pisquetos</i>	272	<i>Carcharodon</i>	20
<i>pisquetus</i>	272	capensis	20
<i>punctatus</i>	270, 273	carcharis	15, 20
richardi	272	rondeleti	20
<i>santae-helenae</i>	270	verus	20
<i>secundus</i>	272	<i>carcinognathus</i> , <i>Gymnothorax</i>	93
<i>semispinosus</i>	270	<i>cardenai</i> , <i>Bonaci</i>	313
<i>stellatus</i>	273	<i>cardonei</i> , <i>Coralliozetus</i>	464
<i>symmetricus</i>	270	<i>Carelliparis</i>	400
<i>tenebrosus</i>	273	agassizii	400
<i>trachurus japonicus</i>	270	dennysi	400
<i>vinetus</i>	272	herschelinius	400
<i>Caranxomorus plumieriiranus</i>	270	tunclatus	400
<i>Carapidae</i>	480	<i>Caremira</i>	401
<i>carapinus</i> , <i>Coryphaenoides</i>	205	sima	401
<i>Carapo</i>	101	<i>Carencheli</i>	77
<i>carapo</i> , <i>Gymnotus</i>	101, 488	<i>Careproctus</i>	402
<i>Carapus</i>	101, 488	attenuatus	403
affinis	488	bowerbanki	403
arenicola	488	candidus	403
bermudensis	488	colletti	402
<i>Carauana</i>	309	ectenes	402, 403
<i>carauana</i> , <i>Serranus</i>	309	furcellus	402
<i>carbo</i> , <i>Aphanopus</i>	262	galeatus	402
<i>carbonaria</i> , <i>Percina</i>	283	gilberti	402
Pileoma	283	longifilis	403
<i>carbonarium</i> , <i>Haemulon</i>	330	longipinnis	403
<i>carbonarius</i> , <i>Acipenser</i>	35	melanurus	402, 403
<i>Gadus</i>	269	mollis	403
<i>carbunculus</i> , <i>Etelis</i>	328	opisthotremus	403
<i>Carcharhinus</i>	15	ostentum	402
cerdale	16	phasma	402
menisorrah	16	ranulus	402
natator	17	reinhardti	402
velox	16	sinus	401
<i>Carcharias</i>	15	spectrum	402
acronotus	16	<i>carlbeum</i> , <i>Branchiostoma</i>	7
ethalorus	17	<i>caribbeus</i> , <i>Sargus</i>	338
atwoodi	20	<i>caribbicus</i> , <i>Belone</i>	197
azureus	16	Ilisha	45
cæruleus	16	<i>caribbus</i> , <i>Chloroscombrus</i>	276
cerdale	16	<i>Celorhynchus</i>	207
commersonii	15	<i>Haleichthys</i>	507
falciformis	16	<i>Macrurus</i>	207
falcipinnis	16	<i>Tylosurus</i>	197
fronto	17	<i>caribbeus</i> , <i>Aphthalmichthys</i>	89
griseus	19	<i>carinatum</i> , <i>Acanthosoma</i>	503
hemiodon	16	Siphonotoma	240
henlei	17	<i>carinatus</i> , <i>Diodon</i>	503
isodon	15	<i>Labichthys</i>	84
lalandi	15	<i>Placopharynx</i>	111
lamia	15, 20	Salmo	56
lamella	15	Syngnathus	240
limbatus	16, 17	<i>carinicauda</i> , <i>Halosaurichthys</i>	74
longurio	15	<i>carletoni</i> , <i>Leuciscus</i>	116
maclovi	17	<i>carinatale</i> , <i>Tripterygium</i>	458
menisorrah	17	<i>carinalis</i> , <i>Enneapterygius</i>	458
microps	16	<i>carinatus</i> , <i>Celorhynchus</i>	207
milberti	16, 17	Macrurus	207
natator	17	carinifer, Macrourus	206
nicaraguensis	16	Macrurus	206
obesus	14	<i>carinatus</i> , <i>Pteropodus</i>	369
obscurus	16	Sebastichthys	369
oxyrhynchus	15, 16	carneus, Acronurus	362
platyodon	15	Sicyases	490
platyrhynchus	15	carolae, Cantherines	494
porosus	16	carolina, Argentina	39
(<i>Prionodon</i>) <i>limbatus</i>	16	Atherina	246
milleri	16	Trigla	406-408
mulleri	16	<i>carolinensis</i> , <i>Potamocottus</i>	383, 385
punctatus	15	Balistes	401
remotus	16	Cestreas	854
sorrah	15	Clupea	44
taurus	15	Gobius	440
temminckii	15, 19	Hyperistius	305
tigris	15	Otolithus	355
velox	16	Seriola	279
<i>carcharias</i> , <i>Carcharodon</i>	15, 20	<i>carolinus</i> , <i>Gasterosteus</i>	276, 277
Squalus	15, 20	Labrus	424
(<i>Carcharias</i>) <i>terrae-nova</i> , <i>Squalus</i>	14	Lipophrys	461
<i>Carcharidae</i>	13	Merulius	407
		Pholis	461

List of scientific names—Continued

	Page
<i>carolinus</i> , <i>Priacanthus</i>	322, 323
<i>Pteraclis</i>	286
<i>Trachinotus</i>	277
<i>carpintis</i> , <i>Herichthys</i>	423
<i>Neotropius</i>	423
<i>carpio</i> , <i>Carpoides</i>	103
<i>Catostomus</i>	103, 109
<i>Cyprinodon</i>	180
<i>Floridichthys</i>	180
<i>Carpoides</i>	103
<i>bison</i>	103
<i>carpio</i>	103
<i>cultanserinus</i>	103
<i>cypinus</i>	103
<i>damalis</i>	103
<i>diformis</i>	103
<i>elongatus</i>	104
<i>grayi</i>	103
<i>labiosus</i>	103
<i>meridionalis</i>	104
<i>microstomus</i>	104
<i>niger</i>	102
<i>nummifer</i>	103
<i>selene</i>	103
<i>taurus</i>	102
<i>thompsoni</i>	103
<i>tumidus</i>	103
<i>urus</i>	102
<i>vacca</i>	103
<i>velifer</i>	103
<i>vitulus</i>	102
<i>carringtoni</i> , <i>Apocope</i>	140
<i>carringtonii</i> , <i>Apocope</i>	141
<i>caryi</i> , <i>Acipenser</i>	34
<i>Embiotoca</i>	412
<i>Hypsurus</i>	412
<i>cascalensis</i> , <i>Brachyrhaphis</i>	187
<i>Bryconamericus</i>	97
<i>Gambusia</i>	187
<i>cassidyl</i> , <i>Embiotoca</i>	411
<i>castanea</i> , <i>Sidera</i>	92
<i>castaneola</i> , <i>Sparus</i>	265
<i>castaneum</i> , <i>Macrostoma</i>	167
<i>castaneus</i> , <i>Gymnothorax</i>	92
<i>Ichthiomyzon</i>	9
<i>Notoscopelus</i>	167
<i>castelnauii</i> , <i>Cylindrosteus</i>	37
<i>castor</i> , <i>Pontinus</i>	372
<i>Catablemella</i>	166
<i>brachychir</i>	166
<i>Catetyx</i>	480
<i>rubrifrons</i>	480
<i>simus</i>	480
<i>catalinae</i> , <i>Neothunnus</i>	260
<i>Catalineta</i>	361
<i>Catalufa</i>	323
<i>catalufa</i> , <i>Priacanthus</i>	323
<i>Cataphracti</i>	363
<i>Cataphractus schoneveldii</i>	395
<i>cataphractus</i> , <i>Agonus</i>	395
<i>Cottus</i>	395
<i>Gasteracanthus</i>	237
<i>Gasterosteus</i>	237
<i>cataractæ</i> , <i>Gobio</i>	139
<i>Rhinichthys</i>	139
<i>cataractus</i> , <i>Leucosomus</i>	117
<i>catena</i> , <i>Bassozetus</i>	480, 481
<i>Bathyonus</i>	481
<i>catenata</i> , <i>Echidna</i>	94
<i>Poecilia</i>	179
<i>catenatum</i> , <i>Xenisma</i>	179
<i>catenatus</i> , <i>Gymnothorax</i>	94
<i>catenula</i> , <i>Muraenophis</i>	94
<i>catesbeii</i> , <i>Pomotis</i>	302
<i>catharinæ</i> , <i>Pristipoma</i>	333
<i>Cathetostoma alboguttata</i>	454
<i>Cathorops</i>	151
<i>gulosa</i>	151
<i>hypothalma</i>	151
<i>Cathochænum</i>	341
<i>Catotodus</i>	292, 293
<i>fuscatus</i>	293
<i>flabellare cumberlandicum</i>	293
<i>lineolatum</i>	293
<i>Catostomus</i>	105, 106
<i>(Acomus) generosus</i>	104
<i>griseus</i>	107
<i>guizmaniensis</i>	104
<i>lactarius</i>	107
<i>alticulus</i>	105
<i>anisurus</i>	109
<i>arceopus</i>	106
<i>ardens</i>	105
<i>aureolus</i>	109, 110
<i>aurora</i>	108
<i>bernardini</i>	108
<i>bostoniensis</i>	105
<i>bubalus</i>	102
<i>carpio</i>	103, 109
<i>catostomus</i>	106
<i>chamberlaini</i>	107
<i>chloropteron</i>	105
<i>clarkii</i>	104
<i>commersonii</i>	105
<i>communis</i>	105
<i>conchos</i>	107
<i>congestus</i>	110
<i>cypho</i>	108
<i>cyprinus</i>	103
<i>discobolus</i>	107
<i>duquesnii</i>	110
<i>elongatus</i>	101
<i>erythrus</i>	110
<i>fuscatus</i>	109
<i>fecundus</i>	108
<i>flexuosus</i>	105
<i>forsterianus</i>	103, 106
<i>gibbosus</i>	109
<i>gila</i>	106
<i>gracilis</i>	105
<i>griseus</i>	107
<i>hudsonius</i>	106
<i>insignis</i>	105, 108
<i>labialis</i>	106
<i>latipinnis</i>	107
<i>lesueuri</i>	111
<i>longirostrum</i>	106
<i>macrocheilus</i>	106
<i>macrolepidotus</i>	111
<i>maculosus</i>	107
<i>megastomus</i>	107
<i>melanops</i>	109
<i>melanotus</i>	137
<i>microps</i>	106
<i>nanomyzon</i>	106
<i>nebuliferus</i>	104
<i>nigricans</i>	107
<i>etowanus</i>	107
<i>occidentalis</i>	106
<i>oneida</i>	110
<i>pallidus</i>	105
<i>planiceps</i>	107
<i>plebeius</i>	104
<i>pocatillo</i>	107
<i>reticulatus</i>	105
<i>retropinnis</i>	107
<i>rex</i>	108
<i>rhotocetus</i>	107, 108
<i>richardsoni</i>	106
<i>snyderi</i>	106
<i>sonoriensis</i>	106
<i>suckili</i>	105
<i>tahoensis</i>	107
<i>texanus</i>	108
<i>tsiticoensis</i>	106
<i>tuberculatus</i>	109
<i>utawana</i>	105
<i>velifer</i>	103
<i>vittatus</i>	109
<i>xanthopus</i>	107
<i>catostomus</i> , <i>Catostomus</i>	106
<i>Cyprinus</i>	105, 106
<i>Phenacobius</i>	142

List of scientific names—Continued

	Page		Page
Catulus.....	11, 17	cayanus, <i>Pristigaster</i>	45
brunneus.....	12	cayennensis, <i>Citharichthys</i>	218, 219
cephalus.....	11	Otolithus.....	368
saxatilis.....	11	Vomer.....	275
uter.....	12	Cayennia.....	448
xanthurus.....	12	guichenoti.....	448
catulus, <i>Ameiurus nebulosus</i>	164	caymanensis, <i>Gambusia</i>	186
(<i>caniculus</i>), <i>Scylliorhinus</i>	13	cayorum, <i>Anchovia cheirostoma</i>	49
Galeus.....	13	Anchoviella.....	49
Gobius.....	440	Corythoichthys.....	243
<i>Pimelodus</i>	164	Hippichthys.....	479, 480
catus, <i>Haustor</i>	163	Ogilbia.....	388
<i>Serranus</i>	311	Eucallichthys.....	238
<i>Silurus</i>	163	Hybopsis.....	133
caukanus, <i>Agenesiosus</i>	159	<i>Notropis</i>	133
cauda forcipata, <i>Brama marina</i>	265	<i>cayuge atrocaudalis</i> , <i>Notropis</i>	133
cauda nigra, <i>Perca marina</i>	320	<i>Cebidichthys</i>	466
caudacinctus, <i>Prionotus (Anchisomus)</i>	500	<i>Cebidichthys</i>	466
caudacuta, <i>Motella</i>	209	<i>cristagalli</i>	466
caudafurcatus, <i>Pimelodus</i>	152	<i>violaceus</i>	466
caudale, <i>Ariosa</i>	80	cedrosensis, <i>Atherinops insularum</i>	262
caudalis, <i>Congermuræna</i>	80	celatus, <i>Batrachus</i>	487
<i>Iridio</i>	427	Centaurus.....	503
<i>Julis</i>	427	centrale, <i>Cichlasoma</i>	419
<i>Leptocephalus</i>	81	centralis, <i>Parapetenia</i>	419
<i>Leptocephalus (Microconger)</i>	81	Centrarchidæ.....	297
<i>Microconger</i>	81	Centrarchus.....	304
<i>Mixonus</i>	483	<i>hexacanthus</i>	304, 305
<i>Pomacentrus</i>	413	<i>interruptus</i>	303
caudanotatus, <i>Mesopriion</i>	326	<i>macropodus</i>	304
caudata, <i>Lamna</i>	16	<i>maculosus</i>	303
<i>Motella</i>	209	<i>obscurus</i>	298
caudatus, <i>Lepidopus</i>	282	<i>pentacanthus</i>	303
<i>Paraliparis</i>	404	<i>pomotis</i>	303
caudicula, <i>Conger</i>	79	<i>tetraacanthus</i>	419
caudillimbatus, <i>Conger</i>	79	centrarchus, <i>Archocentrus</i>	420
<i>Echelus</i>	79	<i>Heros</i>	420
caudimacula, <i>Hæmulon</i>	329	centrina, <i>Calorhynchus</i>	32
<i>Sargus</i>	338	Centriscus.....	245
audimaculata, <i>Strongylura</i>	195	<i>scolopax</i>	245
caudimaculatus, <i>Hemilexocottus</i>	199	Centrismus.....	319
caudofasciata, <i>Limia</i>	191	<i>signifer</i>	319
Caularchus.....	488	Centrobranchiellus.....	470
<i>meandricus</i>	488	Centrochromis.....	415
caulias, <i>Sigmatistes</i>	391	<i>rudis</i>	415
Caulistius.....	488	centrognathos, <i>Zanclus</i>	362
<i>papillifer</i>	488	centrognathus, <i>Zanclus</i>	362
Caulolatilus.....	357	Centrolabrus.....	424
<i>affinis</i>	357	<i>exoletus</i>	424
<i>anomalus</i>	357	Centrolophidae.....	267
<i>chrysops</i>	357	Centrolophus.....	267
<i>cyanops</i>	357	<i>liparis</i>	267
<i>microps</i>	357	<i>morio</i>	267
<i>princeps</i>	357	<i>niger</i>	267
Caulolepis.....	232	<i>pomphilus</i>	267
<i>longidens</i>	232	Centronotus.....	278, 467
Caulophryne.....	611	<i>argenteus</i>	277
<i>jordani</i>	511	<i>conductor</i>	278
Caulophrynidæ.....	511	<i>cristagalli</i>	468
Caulopus.....	171	<i>fuscatus</i>	467
<i>altivelis</i>	171	<i>gardenii</i>	281
<i>borealis</i>	171	<i>islandicus</i>	471
(Caulopus) <i>poeyi</i> , <i>Alepidosaurus</i>	171	<i>letus</i>	468
caulopus, <i>Homestes</i>	468	<i>spinosus</i>	281
caurina, <i>Clarkina</i>	118	Centropholis.....	266
caurinus, <i>Cyprinus</i>	118	<i>goodei</i>	266
<i>Cyprinus (Leuciscus)</i>	118	<i>pettersi</i>	266
<i>Leuciscus</i>	112, 118	centropleura, <i>Uranidea semiscabra</i>	383
<i>Mylochelus</i>	118	Centropomidæ.....	305
<i>Pteropodus</i>	369	Centropomus.....	305
<i>Sebastes</i>	369	<i>affinis</i>	306
cauthosum, <i>Gobiosoma</i>	446	<i>albus</i>	307
cauthosus, <i>Barbillifer</i>	446	<i>armatus</i>	308
cavalla, <i>Cyblum</i>	257	<i>appendiculatus</i>	305
<i>Sierra</i>	257	<i>brevis</i>	306
cavernosus, <i>Bathygadus</i>	207	<i>cuvieri</i>	306
<i>Hymenocephalus</i>	207	<i>ensiferus</i>	306
cavifrons, <i>Ambloplites</i>	303	<i>grandoculatus</i>	306
<i>Diagramma</i>	335	<i>lucioperca undecimalis</i>	306
<i>Hemitripterus</i>	393	<i>luteus</i>	281
<i>Icelinus</i>	375	<i>medius</i>	306
<i>Tarandichthys</i>	375	<i>mexicanus</i>	306
Caxis.....	325	<i>nigrescens</i>	306
caxis, <i>Sparus</i>	325	<i>parallelus</i>	306
cayanensis, <i>Lutjanus</i>	384		

List of scientific names—Continued

	Page		Page
<i>Centropomus pectinatus</i>	306	<i>Ceratias, holboelli</i>	508
pedimacula	306	<i>Ceraticthys</i>	137
robalito	306	<i>physignathus</i>	136
scaber	306	<i>prosthemius</i>	136
undecimalis	305, 306	<i>smithi</i>	145
undecim-radiatus	305	<i>squamiferus</i>	136
unionensis	306	<i>vigilax</i>	137, 144, 145
viridis	306	<i>Ceratidae</i>	508
<i>Centropristes</i>	317	<i>Ceratobatis</i>	32
ayresi	318	<i>robertsi</i>	32
fuscus	319	<i>Ceratocottus</i>	381
merus	312	<i>diceraus</i>	381
nigricans	317	<i>lucasii</i>	382
ocyurus	318	<i>Ceratoptera</i>	32
rufus	318	<i>Ceratoscopelus</i>	167
striatus	317	<i>maderensis</i>	167
subligarius	321	<i>Ceraticthys</i>	137, 144
tabacarius	319	<i>cumingii</i>	138
<i>Centropristes</i>	317	<i>cyclotis</i>	137
annularis	319	<i>halinus</i>	137
aurorubens	327, 323	<i>hypsinotus</i>	138
brasiliensis	320	<i>labrosus</i>	138
displururus	320	<i>leptocephalus</i>	137
macrophthalmus	328	<i>lucens</i>	137
macropoma	318	<i>micropogon</i>	137
subligarius	320	<i>monacus</i>	138
<i>Centropyge</i>	361	<i>physignathus</i>	136
clarionensis	361	<i>prosthemius</i>	136
lunata	361	<i>sallei</i>	145
passer	361	<i>smithi</i>	145
<i>Centrosyllium</i>	21	<i>squamiferus</i>	138
fabricii	21	<i>Sterletidae</i>	139
<i>Centroscymnus</i>	21	<i>stigmaticus</i>	137
cocolepis	21	<i>ventricosus</i>	141
centaura, <i>Raja</i>	28	<i>vigilax</i>	137, 144, 145
centrourus, <i>Pastinachus</i>	28	<i>zanemus</i>	138
<i>Centrurophis</i>	28	<i>Ceratius uranoscopus</i>	508
spadiceus	87	<i>Ceratoscopelus</i>	167
cepediania, <i>Megalops</i>	46	<i>maderensis</i>	167
cepedianum, <i>Dorosoma</i>	46	<i>cercostigma, Cyprinella</i>	130
cepedianum exile, <i>Dorosoma</i>	46	<i>Erogia</i>	130
cepedianus, <i>Lophotes</i>	215	<i>Cerdale</i>	473
<i>Friacanthus</i>	323	<i>ionthas</i>	473
<i>Cephalacanthus</i>	408	<i>cerdale, Carcharlas</i>	10
<i>Cephalacanthus</i>	408	<i>Carcharhinus</i>	16
spindrella	408	<i>Scyatlina</i>	473
<i>Cephalocassis</i>	149	<i>Cerdalidae</i>	472
<i>cephaloides, Cottus</i>	339	<i>Cerdo</i>	210
<i>Cephalopholis</i>	309, 425	<i>cerea, Menidia beryllina</i>	247
argus	309	<i>Cerna</i>	310
fulvus	309	<i>acutirostris lata</i>	315
fulvus punctatus	309	<i>macrogenis</i>	315
ruber	309	<i>nebulosa</i>	315
stellatus	309	<i>cernum, Polypriion</i>	308
taeniolus	309	<i>cervinum, Lepophidium</i>	484
<i>Cephalopeltis nigripinnis</i>	309	<i>Leptophidium</i>	484
<i>Cephaloptera jobni</i>	32	<i>ervinus, Ptychostomus</i>	111
manta	32	<i>Scartozymzon</i>	111
vampyrus	32	<i>Teretulus</i>	111
<i>Cephalopterus</i>	31	<i>cervus, Synanceia</i>	381
gloria	32	<i>Cestracion</i>	18
hypotomus	32	<i>francisci</i>	10
manta	32	<i>pantherinus</i>	10
olfersi	32	<i>quoyi</i>	10
vampyrus	32	<i>Cestraciontes</i>	10
<i>Cephaloscyllium</i>	11	<i>Cestreus</i>	354
uter	12	<i>carolinensis</i>	354
<i>Cephalus</i>	503	<i>predatorius</i>	354
<i>cephalus, Brycon</i>	98	<i>Castrorhinus</i>	18
Catulus	11	<i>cetaceus, Squalus</i>	20
Gobiesox	488, 489	<i>Cetengraulis</i>	51
Mugil	252, 253	<i>edentulus</i>	51
Paraliparis	404	<i>garmani</i>	51
Scyliorhinus	11	<i>gilberti</i>	51
Semotilus	117	<i>mysticetus</i>	51
<i>Cephola trachiptera</i>	216	<i>Cetomimidæ</i>	172
<i>Cephus</i>	210	<i>Cetomimus</i>	172
<i>Cepolopsis</i>	477	<i>gilli</i>	172
<i>cerapalus, Opsanus</i>	487	<i>storeri</i>	172
<i>cerasinus, Gobiesox</i>	499	<i>Cetorhinidae</i>	20
<i>Hypsilepis cornutus</i>	128	<i>Cetorhinus</i>	20
<i>Luxilus</i>	128	<i>maximus</i>	20
<i>Ceratacanthus</i>	495	<i>shavlanus</i>	20
<i>schoepfi</i>	495	<i>Cetunculi</i>	171
<i>Ceratias</i>	508		

List of scientific names—Continued

	Page		Page
<i>Chænobrytus</i>	302	<i>chagresi</i> , <i>Pimelodus</i>	188
<i>antistius</i>	303	<i>Thyrina</i>	249
<i>gulosus</i>	302	<i>Xenocara</i>	160
<i>Chænomugil</i>	254	<i>chalceum</i> , <i>Pristipoma</i>	335
<i>proboscideus</i>	254	<i>chalceus</i> , <i>Orthopristis</i>	335
<i>Chænophryne</i>	510	<i>Chalcinopsis</i>	98
<i>longiceps</i>	510	<i>chagresensis</i>	98
<i>Chænopsetta</i>	223	<i>striatulus</i>	98
<i>Chænopsidea</i>	405	<i>chalcogramma</i> , <i>Theragra</i>	209
<i>Chænopsis</i>	405	<i>chalcogrammus</i> , <i>Gadus</i>	209
<i>alepidota</i>	405	<i>chalogrammus fucensis</i> , <i>Pollachius</i>	209
<i>lucus</i>	405	<i>chalinus</i> , <i>Epinephelus</i>	315
<i>ocellata</i>	405	<i>Chalinura</i>	204
<i>ocellatus</i>	405	<i>brevibarbis</i>	204
<i>chætirhamphus</i> , <i>Anteliochimæra</i>	33	<i>dilera</i>	204
<i>Chætodipterus</i>	358	<i>occidentalis</i>	204
<i>faber</i>	358	<i>serrula</i>	204
<i>zonatus</i>	358	<i>simula</i>	204
<i>Chætodon</i>	359	<i>spinulosa</i>	204
<i>alepidotus</i>	268	<i>Chalinurus</i>	204
<i>arcuatus</i>	360	<i>challengeri</i> , <i>Notacanthus</i>	76
<i>atæniatus</i>	360	<i>Polyacanthonus</i>	76
<i>aureus</i>	360	<i>chalybeus</i> , <i>Hypsopus</i>	127
<i>auriga</i>	359	<i>Hydrophox</i>	127
<i>aya</i>	360	<i>chalybeus abbotti</i> , <i>Notropis</i>	127
<i>bimaculatus</i>	360	<i>chalybeus</i> , <i>Chlorophthalmus</i>	165
<i>bricei</i>	359	<i>Hyphalonedrus</i>	165
<i>capistratus</i>	359	<i>chameleonticeps</i> , <i>Lopholatilus</i>	357
<i>chirurgus</i>	362	<i>chamberlaini</i> , <i>Catostomus</i>	107
<i>ciliaris</i>	361	<i>Chasmistes</i>	107
<i>collaris</i>	369	<i>Cottus</i>	385
<i>cornutus</i>	362	<i>Hudsonius</i>	132
<i>couaga</i>	362	<i>Notropis</i>	132
<i>cyprinaceus</i>	340	<i>chamensis</i> , <i>Ophichththus</i>	88
<i>ephippium</i>	359	<i>Chanideæ</i>	40
<i>faber</i>	358	<i>Channomuraena</i>	95
<i>glaucus</i>	378	<i>cubensis</i>	95
<i>gracilis</i>	360	<i>vittata</i>	95
<i>humeralis</i>	360	<i>Channo-Muraena</i>	95
<i>lanceolatus</i>	352, 353	<i>Chanos</i>	40
<i>lineolatus</i>	359	<i>arabicus</i>	40
<i>littoralis</i>	360	<i>cyprinella</i>	40
<i>longirostris</i>	359	<i>chanos</i> , <i>Mugil</i>	40
<i>lutescens</i>	360	<i>chanteray</i> , <i>Raja</i>	25
<i>marginitatus</i>	415	<i>chapalite</i> , <i>Chirostoma</i>	261
<i>mauritii</i>	415	<i>Falcula</i>	142
<i>nigrirostris</i>	360	<i>Falcularius</i>	142
<i>ocellatus</i>	360	<i>Chapalichthys</i>	184
<i>ornatissimus</i>	359	<i>encaustus</i>	184
<i>oviformis</i>	358	<i>chaperi</i> , <i>Acanthoclinus</i>	457
<i>parræ</i>	361	<i>Paraclinus</i>	457
<i>pardi</i>	360	<i>Chapinus</i>	496
<i>plumieri</i>	358	<i>bicaudalis</i>	497
<i>rhomboides</i>	276, 277	<i>chapmani</i> , <i>Psenes</i>	269
<i>sargoides</i>	415	<i>Tænloconger</i>	82
<i>saratilis</i>	414	<i>Characidium</i>	100
<i>sedentarius</i>	360	<i>fasclatum</i>	100
<i>sohar</i>	362	<i>marshi</i>	100
<i>sordidus</i>	414	<i>Characiniðæ</i>	95
<i>squamulosus</i>	361	<i>Characodon</i>	182
<i>striatus</i>	359	<i>billneatus</i>	183
<i>triangulum</i>	359	<i>eiseni</i>	183
<i>tricolor</i>	361	<i>encaustus</i>	184
<i>tristegus</i>	362	<i>ferrugineus</i>	183
<i>unicornis</i>	362	<i>furcidens</i>	183
<i>unicolor</i>	360	<i>garmani</i>	183
<i>unimaculatus</i>	359	<i>geddesi</i>	182
<i>chætodon</i> , <i>Cottus</i>	487	<i>lateralis</i>	182, 183
<i>Mesogoniistius</i>	303	<i>luitpoldi</i>	183, 184
<i>Pomotis</i>	303	<i>multiradiatus</i>	183
<i>Chætodontidae</i>	359	<i>variatus</i>	182
<i>Chætodontops</i>	359, 360	<i>Charax maxilla superiore longiore</i>	100
<i>Chætostoma</i>		<i>chartes</i> , <i>Platotichthys</i>	221
<i> fischeri</i>	160	<i>charybdis</i> , <i>Lepomis</i>	303
<i> loborhynchus</i>	160	<i>Chasmistes</i>	108
<i>Chætostomus</i>	160	<i>brevirostris</i>	108
<i> aspidolepis</i>	161	<i>chamberlaini</i>	107
<i> fischeri</i>	161	<i>copei</i>	108
<i> heteracanthus</i>	161	<i>cujus</i>	108
<i> latifrons</i>	160	<i>fecundus</i>	108
<i>chagresi</i> , <i>Chalcinopsis</i>	161	<i>llorus</i>	108
<i>chagresi</i> , <i>Ancistrus</i>	160	<i>luxatus</i>	108
<i>Menidia</i>	249	<i>stomias</i>	108
<i>Pimeledella</i>	168	<i>tahoensis</i>	107

List of scientific names—Continued

	Page		Page
<i>Chasmodes</i>	462	<i>Chilomycterus californiensis</i>	502
bosquianus	462	<i>puncticulatus</i>	502
Jenkinsi	463	<i>Chilarhinus</i>	502
<i>novenlineatus</i>	462	<i>suezensii</i>	502
<i>quadrifasciatus</i>	462	<i>Chimæra</i>	502
<i>saburrae</i>	462	<i>abbreviata</i>	502
<i>Chatoëtus</i>	46	<i>affinis</i>	502
<i>ellipticus</i>	40	<i>collici</i>	502
<i>eumorphus</i>	44	<i>cristata</i>	502
<i>mexicanus</i>	46, 47	<i>mediterranea</i>	502
<i>signifer</i>	44	<i>mirabilis</i>	502
<i>Chauliodontidae</i>	71	<i>monstrosa</i>	502
<i>Chauliodus</i>	71	<i>plumbea</i>	502
<i>barbatus</i>	71	<i>Chimæridæ</i>	502
<i>macouni</i>	71	<i>Chimæroides</i>	502
<i>richardsoni</i>	71	<i>chinensis</i> , <i>Aulostomus</i>	502
<i>sloanii</i>	71	<i>Balistes</i>	502
<i>Chaunacidae</i>	506	<i>Tachysurus</i>	502
<i>Chaunax</i>	506	<i>chlonostigma</i> , <i>Echidna</i>	502
<i>nuttingi</i>	506	<i>chiostictus</i> , <i>Entomacrodus</i>	502
<i>plexus</i>	506	<i>Salaria</i>	502
<i>Cheilichthys</i>	499	<i>chiquita</i> , <i>Aboma</i>	502
<i>annulatus</i>	499	<i>Gobius</i>	502
<i>asellus</i>	500	<i>Chiocentrodon</i>	502
<i>furthii</i>	499	<i>bleekeriana</i>	502
<i>türthii</i>	499	<i>temnatus</i>	502
<i>pachygaster</i>	499	<i>Chirolophidae</i>	502
<i>politius</i>	499	<i>Chirolophus japonicus</i>	502
<i>testudineus</i>	499	<i>Chironectes</i>	502
<i>trichocephalus</i>	499	<i>arcticus</i>	502
(<i>Cheilichthys</i>) <i>pachygaster</i> , <i>Tetronodon</i>	499	<i>lævigatus</i>	502
<i>Cheilo</i> <i>auratus</i>	423	<i>mentzelli</i>	502
<i>Cheilodipterus</i>	296	<i>mentzelli</i>	502
<i>acoupa</i>	354	<i>multicellatus</i>	502
<i>affinis</i>	296	<i>pictus</i>	502
<i>heptacanthus</i>	280	<i>principis</i>	502
<i>lineatus</i>	296	<i>scaber</i>	502
<i>macrodon</i>	296	<i>sonnagili</i>	502
<i>Cheilonemus</i>	117	<i>tenebrosus</i>	502
<i>Cheilotrema</i>	348	<i>tigris</i>	502
<i>Cheiragonus</i>	393	<i>tumidus</i>	502
<i>Cheirodon</i> <i>gorgone</i>	96	<i>Chiropais constellatus</i>	502
<i>insignis</i>	96	<i>nebulosus</i>	502
<i>Cheimo</i> <i>pelta</i>	359	<i>Chiostoma</i>	502
<i>Cheimon</i> <i>aculeatus</i>	359	<i>arge</i>	502
<i>chelonias</i> , <i>Allector</i>	484	<i>attenuatum</i>	502
<i>Cheoniger</i>	333	<i>bartoni</i>	502
<i>chemnitzi</i> , <i>Notacanthus</i>	75	<i>beryllinum</i>	502
<i>cheneyi</i> , <i>Cottogaster</i>	288	<i>breve</i>	502
<i>Imostoma</i>	288	<i>chapala</i>	502
<i>Cheoanda</i>	117, 118	<i>conoscium</i>	502
<i>carules</i>	120	<i>crystallinum</i>	502
<i>cooperi</i>	118	<i>diaz</i>	502
<i>copel</i>	119	<i>estor</i>	502
<i>egregia</i>	118	<i>grandocule</i>	502
<i>evermanni</i>	118	<i>guatemalensis</i>	502
<i>humboldtii</i>	119	<i>humboldtianum</i>	502
<i>hydrophlox</i>	119	<i>jordani</i>	502
<i>microdon</i>	118	<i>labarces</i>	502
<i>modesta</i>	120	<i>lermes</i>	502
<i>montana</i>	119	<i>lucius</i>	502
<i>Cherna</i>	311	<i>mezquital</i>	502
<i>cherno</i> , <i>Anthias</i>	311	<i>oocathame</i>	502
<i>chesteri</i> , <i>Phycis</i>	213	<i>patzcuaro</i>	502
<i>Urophycis</i>	213	<i>promelas</i>	502
<i>Chiasmodon</i>	452	<i>regani</i>	502
<i>niger</i>	452	<i>siccum</i>	502
<i>subniger</i>	452	<i>sphyrena</i>	502
<i>Chiasmodontidae</i>	452	<i>vagrans</i>	502
<i>Chiasmodus</i>	452	<i>vomerina</i>	502
<i>chickasavensis</i> , <i>Luxilus</i>	130	<i>tzirahuen</i>	502
<i>chiuahua</i> , <i>Monilaea</i>	132	<i>chirurgus</i> , <i>Chætodon</i>	502
<i>Notropis</i>	132	<i>Chirus</i>	502
<i>Chilaria</i>	485	<i>ballas</i>	502
<i>chilensis</i> , <i>Brama</i>	265	<i>guttatus</i>	502
<i>Sarda</i>	259	<i>maculo-seriatus</i>	502
<i>chilensis</i> , <i>Exocoetus</i>	200	<i>ordinatus</i>	502
<i>Pelamys</i>	259	<i> pictus</i>	502
<i>chilisticus</i> , <i>Hybopsis</i>	127	<i>trigrammus</i>	502
<i>Hydroplox</i>	127	<i>chirus</i> , <i>Phytichthys</i>	502
<i>Chilomycterus</i>	502	<i>Xiphister</i>	502
<i>affinis</i>	502	<i>chisoensis</i> , <i>Procilia</i>	502
<i>antillarum</i>	502	<i>chiso</i> , <i>Blennius</i>	502
<i>atinga</i>	502		
<i>briareos</i>	502		

List of scientific names—Continued

	Page		Page
<i>Chitonotus</i>	375	<i>Chromis</i>	280
<i>megacephalus</i>	375	<i>epicurorum</i>	280
<i>pugentensis</i>	375	<i>fenestrata</i>	416
<i>chittendeni</i> , <i>Cyclopsetta</i>	221	<i>(Furcaria) atrilobata</i>	412
<i>chlamydes</i> , <i>Bolimannia</i>	443	<i>fusco-maculatus</i>	419
<i>chlevastes</i> , <i>Gymnothorax</i>	93	<i>chromis</i> , <i>Hæmulon</i>	329
<i>Sidera</i>	93	<i>Pogonius</i>	352
<i>Chlopsis</i>	82	<i>Sparus</i>	413
<i>bicolor</i>	82	<i>dakotensis</i>	113
<i>equatorialis</i>	82	<i>eos</i>	113
<i>gilberti</i>	82	<i>erythrogaster</i>	113
<i>chlora</i> , <i>Chloa</i>	135	<i>oreas</i>	113
<i>Chlorichthys</i>	428	<i>pyrrhogaster</i>	113
<i>chloris</i> , <i>Scarus</i>	432	<i>chrosomus</i> , <i>Hybopsis</i>	127
<i>chloristia</i> , <i>Codoma</i>	129	<i>Hydrophox</i>	127
<i>Ergala</i>	129	<i>chrysargyreum</i> , <i>Hæmulon</i>	330, 331
<i>chlorocephalus</i> , <i>Hybopsis</i>	127	<i>chrysargyreus</i> , <i>Brachygenys</i>	331
<i>Hydrophox</i>	127	<i>chrysargyreus</i> , <i>Rhinoberyx</i>	235
<i>chloropteron</i> , <i>Catostomus</i>	105	<i>Chrysiptera</i>	414
<i>chloropterum</i> , <i>Plectropoma</i>	312	<i>Chrysitis</i> , <i>Dionda</i>	144
<i>chloropterus</i> , <i>Alphistes</i>	312	<i>chrysocephalus</i> , <i>Luxilus</i>	128
<i>Chlorophthalmidae</i>	165	<i>Luxilus cornutus</i>	128
<i>Chlorophthalmus</i>	165	<i>chrysochilus</i> , <i>Pomolobus</i>	41, 42
<i>agassizii</i>	165	<i>chrysogaster</i> , <i>Agosia</i>	140
<i>chalibatus</i>	165	<i>chrysoleuca</i> , <i>Corvina</i>	346
<i>mento</i>	165	<i>chrysoleuca</i> , <i>Leucosomus</i>	117
<i>trunculentus</i>	165	<i>Notemigonus</i>	117
<i>Chloroscombrus</i>	276	<i>chrysoleucus</i> , <i>Nector</i>	346
<i>caribbeus</i>	276	<i>chrysomelanurus</i> , <i>Sparus</i>	311
<i>chrysurus</i>	276	<i>chrysomelas</i> , <i>Pteropodus</i>	369
<i>orquesus</i>	276	<i>Sebastichthys</i>	369
<i>stirurus</i>	276	<i>Chrysopera</i>	307
<i>chlorosoma</i> , <i>Vallantia</i>	288	<i>interrupta</i>	307
<i>chlorostictus</i> , <i>Sebastichthys</i>	367	<i>Chrysophrys aculeata</i>	336
<i>Sebastomus</i>	367	<i>chrysops</i> , <i>Caulolatilis</i>	357
<i>chlorostomus</i> , <i>Trisoptropis</i>	314	<i>Latilis</i>	357
<i>chlorurum</i> , <i>Plectropoma</i>	321	<i>Lepidema</i>	307
<i>Chlorurus</i>	433	<i>Ophidurus</i>	88
<i>chlorurus</i> , <i>Hypoplectrus unicolor</i>	96	<i>Perca</i>	307
<i>choensis</i> , <i>Gyrocharax</i>	243	<i>Sparus</i>	336
<i>Chœrolchthys</i>	243	<i>Stenotomus</i>	338
<i>valenciennesi</i>	243	<i>chrysopsis</i> , <i>Hydon</i>	41
<i>Chœrojulis arangoi</i>	426	<i>chrysoptera</i> , <i>Perca</i>	335
<i>grandisquamis</i>	426	<i>chrysopterum</i> , <i>Hæmulon</i>	329
<i>cherostoma</i> , <i>Anchoviella</i>	48	<i>Sparisoma</i>	432
<i>cherostomus</i> , <i>Engraulis</i>	48	<i>chrysopterus</i> , <i>Leuciscus</i>	117
<i>chogset</i> , <i>Labrus</i>	424	<i>Orthopristis</i>	335
<i>chogset fulva</i> , <i>Labrus</i>	424	<i>Perca</i>	335
<i>Chilogaster</i>	194	<i>Scarus</i>	432
<i>agassizii</i>	194	<i>chrysos</i> , <i>Scomber</i>	272
<i>avitus</i>	194	<i>Chrysotus</i>	216
<i>cornutus</i>	194	<i>Fundulus</i>	177
<i>papilliferus</i>	194	<i>Zygocetes</i>	177
<i>Chondrostoma pullum</i>	146	<i>chrysura</i> , <i>Balduilla</i>	345
<i>Chondropterygi</i>	10	<i>chrysurus</i> , <i>Caranx</i>	345
<i>Chonophorus</i>	442	<i>Chloroscombrus</i>	276
<i>banana</i>	442	<i>Dipterodon</i>	345
<i>bucculentus</i>	442	<i>Glyphidodon</i>	415
<i>contractus</i>	442	<i>Glyphtisodon</i>	415
<i>mexicanus</i>	442	<i>Ocyurus</i>	827
<i>nelsoni</i>	442	<i>Scomber</i>	276
<i>chordatus</i> , <i>Saccopharynx</i>	05	<i>Sparus</i>	327
<i>Choregon</i>	06	<i>Stegastes</i>	415
<i>Chorinemus altus</i>	278	<i>chrysus</i> , <i>Eupomacentrus</i>	414
<i>palometra</i>	278	<i>chrysypa</i> , <i>Anguilla</i>	77
<i>saltans</i>	277	<i>chucunaque</i> , <i>Rivulus</i>	180
<i>Chlorististium</i>	315	<i>chucunaque subculti</i> , <i>Rivulus</i>	180
<i>ruberum</i>	315	<i>chuss</i> , <i>Blennius</i>	213
<i>Chriodorus</i>	199	<i>Urophycis</i>	213
<i>atherinoides</i>	199	<i>cibarius</i> , <i>Ammocetes</i>	10
<i>Chriolax</i>	407	<i>Cibi amarilla</i>	272
<i>Chriolepis</i>	438	<i>cibi</i> , <i>Caranx</i>	272
<i>minutillus</i>	438	<i>Clobion</i>	496
<i>Chromintra</i>	257	<i>cloacricosus</i> , <i>Balistes</i>	493
<i>concolor</i>	257	<i>Pleuronectes</i>	227
<i>Chriope</i>	133, 135, 136	<i>cleatirifer</i> , <i>Lycodes</i>	475
<i>alboeolus</i>	136	<i>Clinus senes</i>	303
<i>Chriopeops</i>	175	<i>fasciata</i>	298
<i>goodei</i>	175	<i>floridana</i>	297
<i>Christoma erythrinum</i>	247	<i>minima</i>	298
<i>peninsulae</i>	247	<i>ohiensis</i>	298
<i>Chromides</i>	412	<i>storera</i>	304
<i>Chromis</i>	280, 413	<i>Cichlasoma</i>	416
<i>bermudæ</i>	413	<i>acutum</i>	422
<i>enchyrsurus</i>	413	<i>alfari</i>	419

List of scientific names—Continued

Page	Page
Cichlasoma calobrense.....	417
centrale.....	419
dorsatum.....	420
eigenmanni.....	417
evermanni.....	420
frontale.....	417
gadovii.....	417
globosum.....	421
granadense.....	421
hedricki.....	421
laure.....	421
lethrinus.....	420
maculicauda.....	421
meeki.....	416
milleri.....	417
mojarra.....	417
multifasciatum.....	418
nigrum.....	418
punctatum.....	417
robertsoni.....	420
septemfasciatum.....	421
sexfasciatum.....	421
spinosisstimum innacutum.....	420
steindachneri.....	419
teape.....	417
tenue.....	419
tuyensis.....	421
umbiferum.....	417
zonatum.....	417
(Cichlasoma) spinosissimus, Heros.....	420
Cichlurus.....	416
angulifer.....	416
bifasciatus.....	416
calobrensis.....	417
deppii.....	416
fenestratum.....	416
godmani.....	416
guttulatus.....	416
intermedius.....	416
meeki.....	416
microphthalmus.....	416
melanurus.....	416
umbriifer.....	417
Cichlidæ.....	418
cigonella, Belone.....	195
ciliaris, Angelichthys.....	361
Balistes.....	492
Chætodon.....	361
Zeus.....	274
Ciliata.....	208
argentata.....	238
glauca.....	208
ciliatus, Ballistes.....	494
Blennius.....	473
Entosphenus.....	10
Epinephelus.....	365
Monacanthus.....	494
Petromyzon.....	10
Sebastosomus.....	365
cmbrilus, Enchelyopus.....	209
Gadus.....	209
cincinnati, Acpenser (Antaeus).....	35
cincta, Parapetenia tetricantha.....	419
cinctus, Atopichthys.....	54
Dactyloscopus.....	454
Heros tetraacanthus.....	419
Julis.....	427
cineræa, Alutera.....	496
Amia.....	38
Etheostoma.....	290
Perca marina gibbosa.....	329
cinereum, Aulostoma.....	245
cinereus, Aulostomus.....	245
Gerres.....	341
Macrourus.....	205
Macrurus.....	205
Microspathodon.....	416
Mugil.....	341
Nothonotus.....	290
peleatus, Turdus.....	341
cingulatus, Auchenopterus.....	457
Fundulus.....	177
Pomacanthus.....	360
Zygonectes.....	177
cingulus, Atopichthys.....	54
circularis, Raia.....	24
Raja.....	24
circumnotatus, Scarus.....	433
cirratum, Gingymostoma.....	12
cirratus, Physcis.....	212
Squalus.....	12
Cirrhites.....	358
rivulatus.....	358
Cirrhitichthys.....	358
Cirrhitidæ.....	368
Cirrhitus.....	368
betaurus.....	368
maculatus.....	368
rivulatus.....	358
cirrhosum, Lepisoma.....	459
cirrhosus, Blepsias.....	282
Hypostomus.....	160
Trachinus.....	382
Cirrostomi.....	7
Cisco.....	62, 63
cismontanus, Coregonus williamsoni.....	65
Prosopium williamsoni.....	65
Citharichthys.....	218, 219
æthallon.....	218
artifrons.....	219
arenaceus.....	219
cayennensis.....	218, 219
dinoceros.....	219
fragilis.....	218
gilberti.....	219
guatemalensis.....	219
latifrons.....	218
macrops.....	219
micros.....	219
microstomus.....	219
pætulus.....	218
panamensis.....	221
platophrys.....	219
sordidus.....	218
spilopterus.....	219
stigmæus.....	219
sumichrasti.....	219
uhleri.....	219
unicornis.....	219
ventralis.....	220
xanthostigma.....	218
Citharus.....	222
Citraceodus.....	349
Citrinellus.....	420
citrinellus, Astatheros.....	422
Heros.....	422
Citula.....	278, 274
dorsals.....	274
orthogramma.....	274
citurensis, Oxylophorcaria.....	162
civilis, Hybognathus.....	144
clara, Menidia.....	248
claris, Pimelodus.....	159
Silurus.....	158, 159
Claricola.....	292
alabamæ.....	292
julie.....	292
squamiceps.....	292
whippiti.....	292
clarionensis, Centropyge.....	361
Holacanthus.....	361
clarionis, Xesurus.....	363
clarlus, Pimelodus.....	159
clarki, Artedielius.....	377
Blennicotitus.....	377
Bogoslovius.....	204
clarkii, Catostomus.....	104
Notolepidomyzon.....	104
Salmo.....	56
clarkii declivifrons, Salmo.....	57
clarkii jordani, Salmo.....	57
clarkii tahoensis, Salmo.....	57
Clarkina.....	118
caurina.....	118
clarus, Vigil.....	288
clathratus, Labrax.....	317
Paralabrax.....	317
clavata, Raia.....	25
claviformis, Moxostoma.....	109

List of scientific names—Continued

	Page		Page
claviger Cottus.....	381	Olupea macrophthalmia.....	43
Enophrys	381	mattowacca	42
clavilatum , Acutomentum	380	mediocris	41, 42
claviflatus , Sebastodes	380	megalops	42
claytoni , Gobius	439	membras	41
Rhinogobius	439	menhaden	44
leptosydra , Muraena	90	micropus	46
Clepticus	426	minima	41
parre	426	mirabilis	41
clevelandi , Phoxinus	118	neglecta	44
Clevelandia	445	pallasii	41
los	445	parvula	41, 42
longipinnis	445	pilchardus	42
rosea	445	pseudoharengus	42
Clinidae	455, 404	pusilla	42
Clinocottus	390	sadina	46
analis	390	sapidissima	42
australis	390	sternica	99
australis	390	stolifera	44
sabarræ	390	thrissa	44
Clinostomus	120	thrissina	44
affinis	121	tyrannus	44
elongatus	120	vernalis	42
funduloides	121	villosa	66
hydrophlox	119	virescens	42
margarita	116	vittata	41
montanus	119	clupeiformis , Coregonus	62, 64, 66
pandora	120	Salmo	64
phlegothontis	121	clupeiformis albus , Coregonus	65
tenia	119	Clupeidae	41
vandoisulus	121	clupeiformis , Argyrosomus	61
Clintoni , Lepidosteus	37	Coregonus	62
Clinus aculeatus	470	Clupeoidei	40
binaculatus	400	clupeoides , Anchovia	50
canariensis	450	Engraulis	50
capillatus	459	clupeola , Harengula	43
delalandii	459	Clypeocottus	381
evides	458	robustus	381
gilli	460	Cobitis heteroclitia	175
gobio	455, 456	killifish	176
macrocephalus	455	macrolepidota	176
maculatus	470	majalis	176
medius	470	tiaroga	143
mohri	471	coccineus , Lycodes	476
nebulosus	471	Scarus	432
nigripinnis	456	Coccogenia	127
nuchipinnis	459	coccogenis , Coccotis	127
ocellatus	459	Hypsilepis	127
ocellifer	468	coccoi , Rhinoscopelus	109
pectinifer	459	Scopelus	108, 169
procisus	470	Coccolus	127
unimaculatus	470	annectens	217
zonifer	469	Coccots	127
Cliola	144	brimleyi	127
camura	129	coccogenis	127
chlora	135	macdonaldi	127
montirogi	131	zonatus	128
smithi	145	zonistius	128
velox	145	Cochlognathus	146
vivax	145	biguttatus	146
clippertonense , Ostracion	490	ornatus	146
clodalis , Hiodon	40	cocoensis , Cotylopus	448
Clodipterus	296	cocosa , Echidna	94
Clupanodon	44	Codoma	143
Clupea	41	chlorista	129
festivals	42	eurystoma	130
africana	44	ornata	143
alosa	42	topeka	143
anchovia	43	trichroistia	130
atherinoides	51	vittata	143
brasiliensis	40	coca , Murina	84
carulea	41	Cocilia	84
carolinensis	44	brandtiana	84
elongata	41	Coclophilis	87
encrasichlius	47	compar	87
fasciata	42	Coccula bascanium	86
haleo	41	coccus , Gastrobranchus	8
harengus	41	cocolepis , Centroscymnus	21
var.	41	coelestinus , Pseudoscarus	436
heterura	40	Scarus	435
hudsonia	132	Colorhynchus	207
indigena	42	caribbeus	207
lamproteria	46	carminatus	207
leachi	41	laville	207
lineolata	41	occa	207
macrocephala	40		

List of scientific names—Continued

	Page		Page
<i>Cælorhynchus scaphopsis</i>	207	<i>Compsura</i>	95
(<i>Cælorhynchus</i>) <i>scaphopsis</i> , <i>Macrurus</i>	207	<i>gorgona</i>	95
<i>cænicala</i> , <i>Precilla</i>	175	<i>heterura</i>	95
<i>cænous</i> , <i>Pimeodus</i>	153	<i>concatenatus</i> , <i>Ostracion</i>	496
<i>Pleuronichthys</i>	225	<i>concentricus</i> , <i>Urobatis</i>	30
<i>cerulea</i> , <i>Tautoga</i>	424	<i>Conchognathus</i>	78
<i>ceruleopunctata</i> , <i>Acara</i>	423	<i>grimaldii</i>	78
<i>Aequidens</i>	423	<i>conchorum</i> , <i>Menidia</i>	247
<i>ceruleus</i> , <i>Cyclopterus</i>	397	<i>conchos</i> , <i>Catostomus</i>	107
<i>cerulo-aureus</i> , <i>Hari</i> c.	425	<i>cincinnus</i> , <i>Amblodon</i>	362
<i>cerulopunctata</i> , <i>Acara</i>	423	<i>Gasterosteus</i>	239
<i>cognatus</i> , <i>Cottus</i>	385, 386	<i>coerulea</i> , <i>Ammocetes</i>	9
<i>Cogrus</i>	87	<i>Chlormitra</i>	257
<i>maculatus</i>	87	<i>Euschistodus</i>	414
<i>Coleoptera</i>	86	<i>Gymnoterax</i>	93
<i>colei</i> , <i>Scorpæna</i>	370	<i>Ichthyomyzon</i>	9
<i>collas</i> , <i>Scomber</i>	258	<i>Lycodes</i>	475
<i>colinus</i> , <i>Gadus</i>	209	<i>Nebrius</i>	94
<i>Coliscus</i>	145	<i>Nexilarius</i>	414
<i>parletalis</i>	145	<i>Scomberomorus</i>	267
<i>collapsum</i> , <i>Moxostoma</i>	110	<i>Thyreoidæa</i>	92, 93
<i>collapse</i> , <i>Ptychostomus</i>	110	<i>conductor</i> , <i>Centronotus</i>	278
<i>collaris</i> , <i>Chestodon</i>	359	<i>conferrum</i> , <i>Oreosoma</i>	232
<i>Collettia</i>	188	<i>confertus</i> , <i>Hyborhynchus</i>	146
<i>nocturna</i>	188	<i>Plimphales</i>	146
<i>rufinesquii</i>	188	<i>confinis</i> , <i>Pimeodus</i>	154
<i>colletti</i> , <i>Careproctus</i>	402	<i>Salmo</i>	59
<i>colliet</i> , <i>Chimæra</i>	33	<i>confluentus</i> , <i>Fundulus</i>	177
<i>Hydrolagus</i>	33	<i>Salmo</i>	55
<i>Colocephalæ</i>	90	<i>conformis</i> , <i>Lavinia</i>	120
<i>Colobolabis</i>	108	<i>Siboma</i>	120
<i>brevirostris</i>	198	<i>Conger</i>	79
<i>saira</i>	198	<i>anago</i>	80
<i>colombianus</i> , <i>Heterandria</i>	189	<i>analis</i>	80
<i>Colomesus</i>	500	<i>brasiliensis</i>	78
<i>pistacæus</i>	500	<i>caudicula</i>	79
<i>colonense</i> , <i>Zestidium</i>	347	<i>caudilimbatus</i>	79
<i>colonensis</i> , <i>Stellifer</i>	347	<i>conger</i>	79
<i>colonus</i> , <i>Serranus</i>	320	<i>esculentus</i>	79
<i>colorado</i> , <i>Lija</i>	494	<i>impressus</i>	80
<i>Lutjanus</i>	328	<i>limbatus</i>	78
<i>Lutjanus</i>	326	<i>macrops</i>	79
<i>Perro</i>	423	<i>microstomus</i>	80
<i>coloratum</i> , <i>Aulostoma</i>	245	<i>mordax</i>	88
<i>Colorigla</i>	406	<i>obtusa</i>	79
<i>stearnsi</i>	406	<i>opisthopthalmus</i>	80
<i>Colpichthys</i>	252	<i>orbignyanus</i>	79
<i>regis</i>	252	<i>rubescens</i>	79
<i>coluber</i> , <i>Gempylus</i>	261	<i>verreauxi</i>	79
<i>colubrinus</i> , <i>Stomias</i>	69	<i>verus</i>	79
<i>Columbia</i>	215	<i>vulgaris</i>	79
<i>transmontana</i>	215	<i>conger</i> , <i>Conger</i>	79
<i>columbianus</i> , <i>Pantosteus</i>	105	<i>Muraena</i>	79
<i>Rutilus</i>	122	<i>conger brasiliensis</i> , <i>Muraena seu</i>	94
<i>Siphateles</i>	122	<i>Congermuraena</i>	79
<i>Vomer</i>	275	<i>caudalis</i>	80
<i>columns</i> , <i>Rhinobatos</i>	23	<i>flava</i>	80
<i>comatus</i> , <i>Eroccetus</i>	201	<i>guppil</i>	80
<i>comes</i> , <i>Roccus</i>	355	<i>congestum</i> , <i>Moxostoma</i>	110
<i>comifer</i> , <i>Achirus</i>	230	<i>congestus</i> , <i>Catostomus</i>	110
<i>commersonianus</i> , <i>Lophius</i>	505	<i>Congrammus</i>	454
<i>commersoni</i> , <i>Ovum</i>	499	<i>moorei</i>	454
<i>commersoni</i> , <i>Arius</i>	149	<i>Congrellus</i>	79
<i>Carcharias</i>	15	<i>gilberti</i>	80
<i>Catostomus</i>	105	<i>Congridæ</i>	79
<i>Cyprinus</i>	105	<i>Congrosoma</i>	80
<i>Fistularia</i>	246	<i>evermanni</i>	80
<i>Squalus</i>	15	<i>Congrus</i>	79
<i>communis</i> , <i>Catostomus</i>	105	<i>curridens</i>	78
<i>Dasybatus</i>	24	<i>coniceps</i> , <i>Muraenesox</i>	79
<i>Pogonichthys</i>	138	<i>conirostris</i> , <i>Pimelodus</i>	156
<i>compar</i> , <i>Cædilophis</i>	87	<i>connectens</i> , <i>Lethops</i>	447
<i>complanata</i> , <i>Monias</i>	131	<i>Conocara</i>	53
<i>compressa</i> , <i>Anchoviella</i>	50	<i>macdonaldi</i>	53
<i>compressus</i> , <i>Apareiodon</i>	100	<i>macroptera</i>	53
<i>Bassozetus</i>	481	<i>conocephalia</i> , <i>Gila</i>	112
<i>Bathyneutes</i>	480, 481	<i>conocephalus</i> , <i>Mylopharodon</i>	112
<i>Callyodon</i>	434	<i>Conodon</i>	333
<i>Engraulis</i>	47, 50	<i>antillanus</i>	333
<i>Gadus</i>	212	<i>nobilis</i>	333
<i>Hemigrammus</i>	97	<i>serifer</i>	333
<i>Nauclerus</i>	278	<i>Conorhynchichthys</i>	156
<i>Rutilus</i>	128	<i>Conorhynchos</i>	156
<i>Scarus</i>	484	<i>nelsoni</i>	156
		<i>Conorhynchus</i>	39, 156

List of scientific names—Continued

	Page		Page
<i>conorhynchus</i> <i>Albula</i>	39, 40	<i>Coregonus</i> <i>spilonotus</i>	65
<i>Gymnella</i>	47	<i>stanleyi</i>	65
<i>consocium</i> , <i>Chirostoma</i>	251	<i>thymalloides</i>	66
<i>conspersa</i> , <i>Gla</i>	120	<i>thymallus</i>	64
<i>conspersus</i> , <i>Gymnothorax</i>	92	<i>tullibee</i>	61
<i>Serranus</i>	311	<i>bisselli</i>	63
<i>constancie</i> , <i>Pomatomichthys</i>	298	<i>williamsoni</i>	65
<i>constellatus</i> , <i>Chiropsis</i>	373	<i>cismontanus</i>	65
<i>Platophrys</i>	217	(<i>Coregonus</i>) <i>harangus</i> , <i>Salmo</i>	62
<i>Sebastichthys</i>	367	<i>lucidus</i> , <i>Salmo</i>	62
<i>Sebastomus</i>	367	<i>sapidissimus</i> , <i>Salmo</i>	64
<i>consuetus</i> , <i>Salmo</i>	54	<i>tullibee</i> , <i>Salmo</i>	64
<i>continuum</i> , <i>Hæmulon</i>	329	<i>coregonus</i> , <i>Moxostoma</i>	110
<i>contractus</i> , <i>Chomophorus</i>	442	<i>Ptychostomus</i>	110
<i>conus</i> , <i>Moreostoma</i>	111	<i>coretta</i> , <i>Thunnus</i>	260
<i>Psychostomus</i>	111	<i>Tynnus</i>	260
<i>convergens</i> , <i>Macrourus</i>	206	<i>coriaceus</i> , <i>Eleutheracis</i>	316
<i>Macrurus</i>	206	<i>Rypticus</i>	316
<i>convexus</i> , <i>Alepocephalus</i>	52	<i>Coridae</i>	426
<i>Pomotis</i>	300	<i>Corinus</i> , <i>Hæxanchus</i>	11
<i>cockianus</i> , <i>Aphododerus</i>	215	<i>cornuta</i> , <i>Fistularia</i>	245
<i>cooperi</i> , <i>Cheonota</i>	118	<i>cornubiensis</i> , <i>Pimelepterus</i>	268
<i>Metanops</i>	218	<i>Rhombus</i> <i>levius</i>	228
<i>Metapomops</i>	218	<i>cornubicus</i> , <i>Squalus</i>	19
<i>Raija</i>	26	<i>cornutus</i> , <i>Anoplogaster</i>	233
<i>Salmo</i>	55	<i>Chetodon</i>	362
<i>copel</i> , <i>Acipenser</i> (<i>Huso</i>).....	35	<i>Chilogaster</i>	194
<i>Aleposomus</i>	53	<i>Cyclichthys</i>	502
<i>Chasmistes</i>	108	<i>Cyprinus</i>	128
<i>Cheonda</i>	119	<i>Holacanthus</i>	361
<i>Cottus</i>	388	<i>Hoplostethus</i>	233
<i>Paraliparis</i>	406	<i>Luxilus</i>	123, 128
<i>Lepidosteus</i>	37	<i>Pleuronectes</i>	225
<i>Leuciscus</i>	119	<i>Silurus</i>	245, 246
<i>Squalius</i>	119	<i>Zanclus</i>	362
<i>Xenodermichthys</i>	53	<i>cornutus</i> <i>cerasinus</i> , <i>Hypsilepis</i>	128
<i>Copelandellus</i>	294	<i>chryscephalus</i> , <i>Luxilis</i>	128
<i>copelandi</i> , <i>Cottogaster</i>	285	<i>cyanus</i> , <i>Hypsilepis</i>	128
<i>Rheocrypta</i>	285	<i>gibbus</i> , <i>Hypsilepis</i>	128
<i>Copelandia</i>	304	<i>Isolepis</i> , <i>Luxilis</i>	128
<i>eriaroha</i>	304	<i>Notropis</i>	128
<i>topili</i> , <i>Leuciscus</i>	123	<i>Coro</i> <i>coro</i>	333
<i>Corallicola</i>	456	<i>coro</i> , <i>Coro</i>	333
<i>altivelis</i>	456	<i>Sciaena</i>	333
<i>marmoratum</i>	456	<i>coroides</i> , <i>Umbrina</i>	340
<i>nigripinnis</i>	456	<i>coronata</i> , <i>Seriola</i>	280
<i>coralina</i> , <i>Narcine brasiliensis</i>	27	<i>coronatus</i> , <i>Cylopterus</i>	397
<i>corallinum</i> , <i>Cryptotrema</i>	466	<i>Petrometopon orientatus</i>	308
<i>corallinus</i> , <i>Allardius</i>	377	<i>Serranus</i>	308
<i>Antennarius</i>	506	<i>corporalis</i> , <i>Cyprinus</i>	117
<i>Corallizetus</i>	464	<i>Leucosomus</i>	117
<i>cardone</i>	464	<i>corpore variegata</i> , <i>Trutta</i>	55
<i>Cordylus</i>	257	<i>cortezianus</i> , <i>Aprodon</i>	474
<i>Coregoni</i>	64	<i>coruscans</i> , <i>Arctozenus</i>	165
<i>Coregonidae</i>	61	<i>Paralepis</i>	165
<i>coregonoides</i> , <i>Paralepis</i>	165	<i>coruscum</i> , <i>Holocentrum</i>	230
<i>Coregonus</i>	64	<i>coruscus</i> , <i>Holocentrus</i>	236
<i>abyssicola</i>	66	<i>Corvina</i>	348
<i>albus</i>	65	<i>acutirostris</i>	346
<i>angusticeps</i>	136	<i>adusta</i>	347
<i>artedi</i>	62, 63	<i>biloba</i>	349
<i>clupeiformis</i>	62, 64, 65	<i>chrysoleuca</i>	346
<i>albus</i>	65	<i>delticosa</i>	348
<i>clupeiformis</i>	62	<i>dentex</i>	344
<i>couchii</i>	65	<i>fulgens</i>	345
<i>coulteri</i>	65	(<i>Homopriion</i>) <i>furthii</i>	347
<i>hoyi</i>	62	(<i>Johnius</i>) <i>jacobi</i> !.....	348
<i>kennicotti</i>	65	<i>macrops</i>	345
<i>labradoricus</i>	64	<i>microps</i>	346
<i>javarensis</i>	64	<i>ophioscion</i>	347
<i>laurette</i>	62	<i>oxyptera</i>	320
<i>nelsonii</i>	65	<i>richardsoni</i>	362
<i>neohantonus</i>	64	<i>ronchus</i>	346
<i>nigripinnis</i>	63	<i>stearnsi</i>	348
<i>nova-angliae</i>	65	<i>stellifera</i>	346
<i>oregonius</i>	66	<i>subsequalis</i>	345
<i>osmeriformis</i>	62	<i>trispinosa</i>	346
<i>paralepis</i>	165	<i>vermicularis</i>	348
<i>preblei</i>	65	<i>corvinæformis</i> , <i>Hæmulon</i>	333
<i>prognathus</i>	63	<i>corvinæformis</i> , <i>Brachydeuterus</i>	333
<i>pusillus</i>	62	<i>Hæmulon</i>	333
<i>quadrilateralis</i>	65	<i>Corvula</i>	345
<i>richardsoni</i> !.....	64	<i>batabana</i>	345
<i>ruber</i>	163	<i>sanctaë-luciae</i>	345
<i>signifer</i>	66		

List of scientific names—Continued

	Page		Page
<i>Corvula stialis</i>	345	<i>Cottus bubalis</i>	386
<i>Corynolophus</i>	509	<i>cataphractus</i>	395
<i>reinhardti</i>	509	<i>cephaloïdes</i>	389
<i>Coryphena</i>	264	<i>chætodon</i>	487
<i>argyrurus</i>	265	<i>chamberlaini</i>	385
<i>aurata</i>	265	<i>claviger</i>	381
<i>azorica</i>	265	<i>cognatus</i>	385, 386
<i>cærulea</i>	434	<i>copel</i>	386
<i>dolfin</i>	264	<i>criniger</i>	390
<i>dorado</i>	264	<i>decastrensis</i>	387
<i>equisetis</i>	265	<i>diceraus</i>	381
<i>fuscilata</i>	264	<i>elegans</i>	381
<i>hippurus</i>	264	<i>evermanni</i>	383
<i>immaculata</i>	264	<i>formosus</i>	386
<i>imperialis</i>	264	<i>franklini</i>	385
<i>lessonii</i>	265	<i>glaber</i>	487
<i>lineata</i>	430	<i>glacialis</i>	387
<i>marcgravii</i>	264	<i>gobio</i>	383
<i>nigrescens</i>	317	<i>gobioïdes</i>	385
<i>novacula</i>	430	<i>gracilis</i>	383, 385
<i>perciformis</i>	267, 268	<i>cayuga</i>	386
<i>plumieri</i>	356	<i>greeni</i>	384
<i>pomphilus</i>	267	<i>grønlandicus</i>	386, 387
<i>psittacus</i>	430, 433	<i>gulosus</i>	383
<i>rupestris</i>	205	<i>hemilepidotus</i>	380, 381
<i>scomberoides</i>	265	<i>hexacornis</i>	388
<i>suerli</i>	284	<i>hispidus</i>	393
<i>velifera</i>	266	<i>humilis</i>	387
<i>virgata</i>	265	<i>ictalops</i>	385
<i>viamingii</i>	265	<i>indicus</i>	397
<i>Coryphaenidae</i>	204	<i>jaok</i>	387
<i>Coryphaeniformes</i>	204	<i>klamathensis</i>	384
<i>Coryphaenoides</i>	204	<i>leiopomus</i>	385
<i>carapinus</i>	205	<i>macrops</i>	384
<i>filicauda</i>	207	<i>maculatus</i>	386
<i>rupestris</i>	204, 205	<i>margnatus</i>	384
<i>sulcatus</i>	207	<i>marmoratus</i>	387
<i>Coryphaenoididae</i>	202	<i>meridionalis</i>	385
<i>Coryphopterus</i>	439	<i>mertensi</i>	387
<i>glaucofrenum</i>	439	<i>mitchilli</i>	386
<i>Corythoichthys</i>	243	<i>monopterygius</i>	397
<i>albirostris</i>	243	<i>niger</i>	387
<i>cayorum</i>	243	<i>nigricans</i>	386
<i>cosmopolita</i> , <i>Micropteryx</i>	276	<i>octodecem-spinosus</i>	386
<i>Cossyphus</i>	425	<i>onychus</i>	386
<i>puellaris</i>	425	<i>pachypus</i>	386
<i>pulchellus</i>	425	<i>plexipus</i>	384
<i>costalesi</i> , <i>Smaragdus</i>	441	<i>philomips</i>	384
<i>costaricensis</i> , <i>Gobiesox</i>	499	<i>pistilliger</i>	389
<i>costatus</i> , <i>Bodianus</i>	340	<i>platycephalus</i>	387, 388
<i>Cottidae</i>	383	<i>polaris</i>	388
<i>Cottogaster</i>	285	<i>porosus</i>	387
<i>aurantiacus</i>	234, 285	<i>polyacanthocephalus</i>	387
<i>cheneyi</i>	288	<i>principis</i>	384
<i>copelandi</i>	285	<i>punctulatus</i>	383
<i>putnami</i>	285	<i>quadricornis</i>	388
<i>uranidea</i>	285	<i>rhothea</i>	383
(<i>Cottogaster</i>) <i>uranidea</i> , <i>Etheostoma</i>	285	<i>ricei</i>	383, 386
<i>cottoides</i> , <i>Bathygadus</i>	203	<i>richardsoni</i>	385
<i>Cottopsis</i>	393	<i>scorpioides</i>	386
<i>gulosus</i>	393	<i>scorpius</i>	386
<i>parvus</i>	393	<i>semiscaber</i>	384
<i>ricei</i>	383, 386	<i>shasta</i>	383
<i>semiscaber</i>	384	<i>stelleri</i>	381
<i>Cottunculus</i>	392	<i>tælopopterus</i>	388
<i>micros</i>	392	<i>tenuis</i>	384
<i>thomsonii</i>	392	<i>thomsonii</i>	392
<i>torvus</i>	392	<i>trachurus</i>	381
<i>Cottus</i>	393	<i>tricuspidis</i>	389
<i>acadian</i>	393	<i>triptygyius</i>	393
<i>seneus</i>	396	<i>uncinatus</i>	377
<i>aleuticus</i>	395	<i>ventrils</i>	389
<i>alvodii</i>	395	<i>verrucosus</i>	387
<i>anceps</i>	396	<i>villosus</i>	393
<i>annæ</i>	394	<i>viscosus</i>	385
<i>asper</i>	393	<i>wilsoni</i>	385
<i>asperima</i>	394	<i>Cotylopus nigripinnis</i>	489
<i>asperimus</i>	394	<i>Cotylopus</i>	447
<i>bairdii</i>	393	<i>acutipinnis</i>	447
<i>beldingii</i>	394	<i>cocoonsis</i>	448
<i>bendirei</i>	394	<i>punctatus</i>	448
<i>bicornis</i>	378	<i>salvinii</i>	448
<i>boleoides</i>	395	<i>cousaga</i> , <i>Chætodon</i>	362
<i>brodamius</i>	395	<i>couchi</i> , <i>Dionda</i>	144
		<i>Montana</i>	181

List of scientific names—Continued

Page	Page
Couchia.....	208
minor.....	208
couchiana, <i>Limna</i>	193
couchianus, <i>Platypterus</i>	193
couchii, <i>Serranus</i>	303
couesi, <i>Coregonus</i>	66
<i>Prosopium</i>	65
couesi, <i>Apocope</i>	141
<i>Cryptosarasa</i>	508
Couesius.....	136
<i>adustus</i>	137
<i>dissimilis</i>	137
<i>greeni</i>	137
<i>plumbeus</i>	136
<i>rubrilateralis</i>	118
Couessius <i>squamilentus</i>	136
coulteri, <i>Coregonus</i>	65
<i>Prosopium</i>	65
courbina, <i>Pogonathus</i>	352
Pogonias.....	352
courtadei, <i>Serranus</i>	310
cragini, <i>Amiurus</i>	154
Etheostoma.....	291
<i>Nivicola</i>	291
crameri, <i>Etheostoma</i>	367
<i>Hybopsis</i>	138
<i>Leuresthes</i>	251
<i>Oregonichthys</i>	138
<i>Sebastodes</i>	307
<i>Umbra</i>	174
crasse, <i>Belone</i>	196
<i>Tigoma</i>	120
<i>crassicauda, Lavinia</i>	120
<i>Siboma</i>	120
<i>crassiceps, Cryptotomus</i>	431
<i>Plectromus</i>	233
<i>Scopelus</i>	233
<i>crassilabre, Morostoma</i>	111
<i>crassilabris, Embryx</i>	474
<i>Geophagus</i>	424
<i>(Satanoperca)</i>	424
<i>Lycodopsis</i>	474
<i>Ptychostomus</i>	111
<i>crassus, Alvordius</i>	283
<i>Esox</i>	173
<i>Lepidosteus</i>	38
<i>craticula, Zygogetes</i>	178
<i>Cratinus</i>	317
<i>agassizii</i>	317
<i>Crayacion</i>	498
<i>Creagrus</i>	98
<i>notropoides</i>	98
<i>simus</i>	98
<i>creaseri, Ruscarlopa</i>	378
<i>crebripunctata, Pteroplates</i>	29
<i>Cremnobates</i>	450
<i>affinis</i>	457
<i>altivelis</i>	456
<i>fuscatus</i>	456
<i>integripinnis</i>	457
<i>marmoratus</i>	456
<i>nox</i>	457
<i>cremnobates, Labrosomus</i>	456
<i>Starksia</i>	456
<i>Crenidens simplex</i>	339
<i>Crenilabrus</i>	425
<i>microstoma</i>	424
<i>crenulate, Myctophum</i>	170
<i>crenularis, Tarletonbeania</i>	170
<i>Creolus</i>	320
<i>creolus, Girardinus</i>	189
<i>Serranus</i>	320
<i>Creotroctes</i>	259
<i>crecentale, Gobiosoma</i>	446
<i>crecentalis, Gobiosoma</i>	446
<i>Pomacanthus</i>	381
<i>crecentis, Salmo</i>	57
<i>Salmo gairdneri</i>	57
<i>crestonis, Acanthurus</i>	363
<i>Teuthis</i>	363
<i>cretensis, Labrus</i>	431
<i>criniger, Cottus</i>	390
<i>crinigerum, Siphostoma</i>	241
<i>Syngnathus</i>	241
<i>crinitus, Alectis</i>	274
<i>Blennius</i>	461
<i>Zeus</i>	274
<i>cristagalli, Anoplarchus</i>	468
<i>archolepis, A. noplarchus</i>	468
<i>Cebedichthys</i>	466
<i>Centronotus</i>	468
<i>cristata, Chimera</i>	32
<i>cristatus, Biennius</i>	461
<i>Pinneodus</i>	158
<i>cristiceps, Melanophæs</i>	233
<i>Plectromus</i>	233
<i>cristigall, Gobius</i>	438
<i>cristi-gall, Gobius</i>	438
<i>Cristivomer</i>	59
<i>namaycush</i>	59
<i>siscowet</i>	60
<i>cristulata, Scorpæna</i>	372
<i>cristulatus, Gymnocephalus</i>	466
<i>Crirus</i>	268
<i>bertheloti</i>	140
<i>croceus, Leuciscus</i>	140
<i>Rhinichthys</i>	140
<i>crocodilinus, Brachysomophis</i>	89
<i>Ophisaurus</i>	89
<i>crocota, Plectropoma</i>	321
<i>crocutus, Hypoplectrus unicolor</i>	321
<i>crocro, Pristipoma</i>	334
<i>Rhondiscus</i>	334
<i>croicensis, Scarus</i>	433, 434
<i>croker, Schæna</i>	349
<i>Cromileptes</i>	310
<i>cromis, Labrus</i>	352
<i>Pogonias</i>	352
<i>Crossias</i>	389
<i>albomaculatus</i>	389
<i>alisl</i>	389
<i>beringi</i>	389
<i>camtschaticus</i>	389
<i>crossetus, Etropus</i>	219, 220
<i>crotalinus, Embryx</i>	473
<i>Lycodopsis</i>	473
<i>Crotalopsis punitifer</i>	88
<i>crotaphus, Juilis</i>	426, 427
<i>Crotolopsis</i>	88
<i>quintififer</i>	88
<i>crudellis, Eurymyctera</i>	91
<i>cruentatus, Entomacrodus</i>	463
<i>Labrus</i>	322
<i>Petrometopon</i>	308
<i>Priacanthus</i>	322
<i>Sparus</i>	308
<i>cruentatus coronatus, Petrometopon</i>	308
<i>cruentifer, Omochelys</i>	88
<i>Pisoedenophis</i>	86
<i>crumenophthalmus, Trachurops</i>	271
<i>crumenophthalmus, Scomber</i>	270, 271
<i>cruroeum, Xiphidium</i>	469
<i>crurorous, Squalius</i>	119
<i>crustosum, Peristedion</i>	406
<i>Peristedium</i>	406
<i>Cryptacanthodes</i>	471
<i>Inornatus</i>	471
<i>maculatus</i>	471
<i>Cryptacanthodidae</i>	471
<i>Cryptops</i>	101
<i>Cryptopterenchelys</i>	85
<i>frontalis</i>	85
<i>puncticeps</i>	85
<i>(Cryptopterus) frontalis, Ophichthys</i>	85
<i>Cryptopterus</i>	85
<i>punctatus</i>	85
<i>puncticeps</i>	85
<i>Cryptosarasa</i>	508
<i>couesi</i>	508
<i>cryptosus, Stromateus</i>	267
<i>Cryptotomus</i>	431
<i>europunctatus</i>	431
<i>beryllinus</i>	431
<i>crassiceps</i>	431
<i>dentiens</i>	431
<i>retractus</i>	431
<i>roseus</i>	431
<i>ustus</i>	431

List of scientific names—Continued

	Page		Page
Cryptotrema	456	Curimata	100
corallinum	456	magdalenae	100
crysoleucas, <i>Cyprinus</i>	114, 115	Curimatella	100
<i>Notemigonus</i>	115	lepidurus	100
crysos, <i>Paratractus</i>	272	Curimates, <i>Les.</i>	100
<i>Scomber</i>	272	Curimatidae	100
Crystallaria	282, 288	Curimatus	100
<i>asprella</i>	288	<i>magdalenae</i>	100
Crystallichthys	401	<i>spilurus</i>	100
<i>cyclospilus</i>	401	curta, <i>Anchoviella</i>	48
<i>cyclostigma</i>	401	<i>curtiss-jamesi</i> , <i>Harriotta</i>	33
<i>mirabilis</i>	401	<i>Pollistotrema</i>	8
crystallina, <i>Thyridina</i>	248	curtus, <i>Stolephorus</i>	48
crystallinum, <i>Chiostoma</i>	251	<i>Vomer</i>	275
Ctenochætus	363	curvidens, <i>Congrus</i>	78
<i>striatus</i>	363	curvilineata, <i>Muraenophis</i>	91
Ctenodax	269	curvus, <i>Tetronodon</i>	497
<i>williamsi</i>	269	cuspicauda, <i>Aluterus</i>	495
Ctenodon	363, 470	cuspidatus, <i>Macrourus</i>	206
ctenodon, <i>Acanthurus</i>	363	<i>Macrurus</i>	206
Ctenogobius	438	cutisanserinus, <i>Carploides</i>	103
<i>fuscatus</i>	438	cuvieri, <i>Ausonia</i>	244
<i>tortugae</i>	439	<i>Centropomus</i>	306
Ctenolabrus bergall	424	<i>Gasterosteus</i>	237
<i>uninotatus</i>	424	<i>Gnathopops</i>	453
Ctenolucius	100	<i>Opistognathus</i>	453
<i>beani</i>	100	<i>Tetragonurus</i>	269
<i>insculptus</i>	100	cuzamilæ, <i>Scarus</i>	434
cubæ, <i>Vomer</i>	275	cyanea, <i>Furcaria</i>	412
cubana, <i>Anchoviella</i>	48	cyanella, <i>Mugiloglossa</i>	252
<i>Anguilla</i>	77	cyanellus, <i>Apomotis</i>	298, 299
Cubanichthys	174	<i>Ichthyobus</i>	102
<i>cubensis</i>	174	<i>Lepomis</i>	298
cubanus, <i>Engraulis</i>	48	<i>Mugilops</i>	252
<i>Epinephelus</i>	311	cyaneus, <i>Hypsilepis cornutus</i>	128
cubensis, <i>Channomuraena</i>	95	Cyanichthys	502
<i>Cubanichthys</i>	174	<i>cæruleus</i>	502
<i>Fundulus</i>	174	<i>cyanoccephalus</i> , <i>Iridio</i>	427
<i>Heterandria</i>	188	<i>Labrus</i>	426, 427
<i>Hymnis</i>	274	<i>Lythyrurus</i>	128
<i>Limla</i>	191	<i>cyanoguttatus</i> , <i>Herichthys</i>	422
<i>Toxus</i>	188	<i>cyanoleue, Sparisoma</i>	432
<i>Vomer septapinnis</i>	275	<i>cyanonotus</i> , <i>Alosa</i>	42
cubera, <i>Lutjanus</i>	325	<i>cyanophrys</i> , <i>Naucrates</i>	278
Cubiceps multiradiatus	298	<i>Psenes</i>	288
cubicus, <i>Ostracion</i>	496	<i>cyanops</i> , <i>Caulolatius</i>	357
cubifrons, <i>Malthe</i>	507	<i>cyanopterus</i> , <i>Cypselurus</i>	202
cucurl, <i>Prionodon</i>	16	<i>Exocetus</i>	202
Cucuy	493	<i>Leucichthys</i>	63
Cugupuguacu	312	<i>Lutjanus</i>	325
cujus, <i>Chasmistes</i>	108	<i>Mesopriion</i>	325
cuebræ, <i>Acteis</i>	460	<i>cyanostigma</i> , <i>Julis</i>	428
<i>Malacotenus</i>	460	Cybidae	257
Cutilus	456	<i>Cybium</i>	257
<i>equidens</i>	437	<i>aeratum</i>	257
<i>amblyopsis</i>	437	<i>caballa</i>	277
<i>fusca</i>	436	<i>cavalla</i>	257
<i>perniger</i>	437	<i>commersoni</i>	257
(Cutilus) <i>brevianus</i> , <i>Electris</i>	438	<i>immaculatum</i>	257
cultiferum, <i>Pristipoma</i>	334	<i>petus</i>	256
culturata, <i>Anchoviella</i>	48	<i>sara</i>	256
culturatum, <i>Petalosoma</i>	193	<i>solandri</i>	256
culturatus, <i>Alfaro</i>	193	<i>verany</i>	256
<i>Stolephorus</i>	48	Cycleptus	101
culturifrons, <i>Aluterus</i>	495	<i>elongatus</i>	101
culveri, <i>Trachinotus</i>	277	<i>nigrescens</i>	101
cumberlandicum, <i>Catnotus fimbriatus</i>	293	Cyclichthys	501
<i>Testostoma</i>	293	<i>antillarum</i>	502
cumingii, <i>Ceratichthys</i>	138	<i>attenuatus</i>	502
<i>Oregonichthys</i>	138	<i>briareos</i>	502
cummingsi, <i>Hybopsis</i>	133	<i>cornutus</i>	502
<i>Notropis</i>	133	<i>fuliginosus</i>	502
cuneata, <i>Mollisenesia sphenocephala</i>	192	<i>schopfi</i>	502
<i>Poecilia</i>	193	<i>spinosus</i>	502
cupido, <i>Geophagus</i>	424	Cyclogaster	399
cupresoides, <i>Pimelodus</i>	163	<i>bristoliensis</i>	399
cypreus, <i>Pimelodus</i>	163	<i>megacephalus</i>	400
<i>Silurus</i>	163	<i>(Neoliparis) beringianus</i>	399
<i>Trachinotus</i>	277	<i>micasplendorhus</i>	399
curacao, <i>Gobius</i>	439	Cyclogasterus	398, 399
<i>Rhinogobius</i>	439	Cyclogobius	444
curassavicus, <i>Balistes</i>	493	cyclolepis, <i>Dolloa</i>	203
<i>Mugil</i>	253	<i>Microgobius</i>	443
<i>Querimana</i>	263	<i>Nematoturus</i>	203
curlitus, <i>Salmo</i>	61	Cyclonarce	27

List of scientific names—Continued

	Page		Page
<i>Cycloplum</i>	159	<i>Cynoperca</i>	282
<i>humboldti</i>	159	<i>canadensis</i>	282
<i>pirrense</i>	160	<i>grisea</i>	282
<i>cyclopomatus</i> , <i>Serranus</i>	314	<i>Cynoscion</i>	354
<i>Cyclopsetta</i>	220	<i>acoupa</i>	354
<i>chittendeni</i>	221	<i>jamaiensis</i>	355
<i>fimbriata</i>	220	<i>lefarchus</i>	355
<i>quernea</i>	221	<i>macdonaldi</i>	356
<i>Cyclopterichthys</i>	398	<i>microlepidotus</i>	355
<i>glaber</i>	398	<i>nothus</i>	365
<i>Cyclopteridae</i>	397	<i>obliquatus</i>	355
<i>Cyclopteroidea</i>	398	<i>othonoropterum</i>	355
<i>gyrinos</i>	398	<i>othonoropterus</i>	355
<i>Cyclopterus</i>	397	<i>parvipinnis</i>	356
<i>callyodon</i>	399	<i>phoxocephalum</i>	356
<i>ceruleus</i>	397	<i>regalis</i>	355
<i>corenatus</i>	397	<i>squamipinnis</i>	365
<i>gelatinosus</i>	402	<i>thalassinus</i>	355
<i>lineatus</i>	400	<i>xanthulum</i>	356
<i>liparis</i>	399, 400	<i>cypho</i> , <i>Catostomus</i>	108
<i>major</i>	401	<i>Esox</i>	173
<i>minor</i>	400	<i>Cyphodus</i>	339
<i>lumpus</i>	397	<i>elegans</i>	330
<i>minutus</i>	397	<i>Cyprinaceus</i> , <i>Chetodon</i>	340
<i>musculus</i>	400	<i>Cyprinella</i>	130, 131
<i>nudus</i>	489	<i>analcostana</i>	129
<i>orbis</i>	397	<i>beckwithi</i>	130
<i>pavoninus</i>	397	<i>billingsiana</i>	131
<i>pyramidatus</i>	397	<i>bubalina</i>	130
<i>spinosus</i>	397	<i>callura</i>	129
<i>stelleri</i>	398	<i>cercostigma</i>	130
<i>ventricosus</i>	398	<i>forbesi</i>	131
<i>Cycloptus lumpus</i>	397	<i>forlonensis</i>	131
<i>cyclopum</i> , <i>Pime odus</i>	159	<i>gunnisoni</i>	130
<i>cyclopum</i> , <i>Liparis</i>	400	<i>leptida</i>	130
<i>cyclopilus</i> , <i>Crystallichthys</i>	401	<i>ludibunda</i>	130
<i>cyclostigma</i> , <i>Crystallichthys</i>	401	<i>lugubris</i>	131
<i>Liparis</i>	401	<i>luxiloides</i>	131
<i>Cyclostomi</i>	8	<i>macrostoma</i>	131
<i>Cyclothona</i>	72	<i>notata</i>	130
<i>accinidens</i>	72	<i>rubripinna</i>	131
<i>bathypilla</i>	72	<i>suavis</i>	131
<i>elongata</i>	72	<i>texana</i>	130
<i>lusca</i>	72	<i>umbroosa</i>	130
<i>microdon</i>	72	<i>venusta</i>	130
<i>signata</i>	72	<i>whippili</i>	129
<i>cyclotis</i> , <i>Ceratichthys</i>	137	<i>cyprinella</i> , <i>Chanos</i>	40
<i>cylindraceus</i> , <i>Rivulus</i>	179	<i>Megastomatobus</i>	102
<i>cylindricus</i> , <i>Bascanichthys</i>	86	<i>Scherogynathus</i>	102, 103
<i>Cylindrosteus</i>	37	<i>Cyprinidae</i>	112
<i>agassizii</i>	37	<i>Cyprinodon</i>	180
<i>bartoni</i>	37	<i>baileyi</i>	182
<i>castelnaudii</i>	37	<i>bovinus</i>	181
<i>megalops</i>	37	<i>rubrofluviatilis</i>	181
<i>platostomus</i>	37	<i>californiensis</i>	181
<i>productus</i>	37	<i>carpio</i>	180
<i>rafinesquii</i>	37	<i>dearborni</i>	181
<i>scabriceps</i>	37	<i>elegans</i>	181
<i>zadockii</i>	37	<i>eximius</i>	181
(<i>Cylindrosteus</i>) <i>oculatus</i> , <i>Lepisosteus</i>	37	<i>felicianus</i>	181
<i>Cymatogaster</i>	409, 410	<i>gibbosus</i>	181
<i>aggregatus</i>	409	<i>latifasciatus</i>	181
<i>ellipticus</i>	410	<i>macularius</i>	181
<i>larkinsi</i>	410	<i>baileyi</i>	182
<i>minimus</i>	409	<i>martei</i>	181
<i>puichellus</i>	410	<i>mydrus</i>	180
<i>rosaceus</i>	410	<i>nevadensis</i>	181
<i>cymatogramma</i> , <i>Pleoma</i>	286	<i>parvus</i>	175
<i>cymatotenia</i> , <i>Etheostoma</i> (<i>Iladropterus</i>).....	286	<i>riverendii</i>	181
<i>Hypophomus</i>	286	<i>rubrofluviatilis</i>	181
<i>Cynodus brama</i>	338	<i>variegatus</i>	180, 181
<i>Cynias</i>	13	<i>Cyprinodontes</i>	174
<i>canis</i>	13	<i>Cyprinodontidae</i>	174
<i>lunulatus</i>	13	<i>cyprinoides</i> , <i>Gobius</i>	488
<i>Cynichthys</i>	310	<i>Lophogobius</i>	438
<i>Cynicoglossus</i>	228	<i>Cyprinus</i> (<i>Abramus</i>) <i>balteatus</i>	117, 118
<i>bathybius</i>	228	(<i>Abramus?</i>) <i>smithii</i>	141
<i>Cynocephalus</i>	17	<i>americanus</i>	116, 350
<i>cynodon</i> , <i>Mesopriion</i>	825	<i>atromaculatus</i>	117
<i>Cynodonichthys</i>	179	<i>atronasus</i>	130, 140
<i>tenuis</i>	179, 180	<i>bullaris</i>	117
<i>Cynoglossa</i>	228	<i>catostomus</i>	105, 106
<i>Cynoglossidae</i>	230	<i>caurinus</i>	118
<i>cynoglossus</i> , <i>Glyptocephalus</i>	228	<i>commersoni</i>	105
<i>Pleuronectes</i>	221, 228	<i>cornutus</i>	128

List of scientific names—Continued

	Page		Page
<i>Cyprinus corporalis</i>	117	<i>dariensis</i> , <i>Apareiodon</i>	100
<i>crysoleucus</i>	114, 115	<i>Hemibrycon</i>	97
<i>hemiplus</i>	115	<i>Oxyloricaria</i>	162
<i>leuciscus</i>	117	<i>Parodon</i>	100
(<i>Leuciscus</i>) <i>caurinus</i>	118	<i>Sternopygus</i>	101
<i>gracilis</i>	130	<i>Dasibatis</i>	28
<i>oregonensis</i>	113, 114	<i>daspiota</i> , <i>Pisodonophis</i>	88
<i>maxilingua</i>	146	<i>Dasyatidæ</i>	28
<i>megalops</i>	128	<i>Dasyatis</i>	28
<i>melanurus</i>	128	<i>attavilla</i>	29
<i>oblongus</i>	109	<i>guttatus</i>	28
<i>sucetta</i>	109	<i>hastatus</i>	28
<i>teres</i>	105	<i>longus</i>	28
<i>vittatus</i>	140	<i>schmardae</i>	28
<i>cypinus</i> , <i>Carpoides</i>	103	<i>torreli</i>	28
<i>Catostomus</i>	103	<i>Dasybatis dipterurus</i>	29
<i>Sclerognathus</i>	102	<i>Dasybatis</i>	28
<i>Cypselurus</i>	201	<i>dasycephalus</i> , <i>Arius</i>	150
<i>babiensis</i>	202	<i>Galeichthys</i>	150
<i>californicus</i>	202	<i>Dasycottus</i>	392
<i>callopterus</i>	202	<i>scutiger</i>	392
<i>cyanopterus</i>	202	<i>daubentonii</i> , <i>Caranx</i>	273
<i>fucorum</i>	202	<i>Davidia</i>	495
<i>fureatus</i>	201	<i>punctata</i>	495
<i>gibbifrons</i>	202	<i>davidsonii</i> , <i>Anisotremus</i>	332
<i>heterurus</i>	201	<i>Monacanthus</i>	494
<i>holubi</i>	202	<i>Pristipoma</i>	332
<i>lineatus</i>	202	<i>davisi</i> , <i>Rabula</i>	91
<i>lutkeni</i>	201	<i>dayi</i> , <i>Phoxargyreus</i>	247
<i>nigricans</i>	201	<i>deani</i> , <i>Eosebastes</i>	367
<i>spilonotopterus</i>	202	<i>Paraliparis</i>	404
<i>xenopterus</i>	202	<i>Pollistotrema</i>	9
<i>cypselurus</i> , <i>Prognurus</i>	403	<i>Sebastodes</i>	367
<i>Cypsilurus</i>	201	<i>dearborni</i> , <i>Cyprinodon</i>	181
(<i>Cypsilurus</i>) <i>procne</i> , <i>Exocetus</i>	201	<i>Decaetulus</i>	105
<i>Cytus hololepis</i>	232	<i>decadactylus</i> , <i>Beryx</i>	234
<i>Dacentrus</i>	412	<i>decagonus</i> , <i>Agonus</i>	394
<i>lucens</i>	412	<i>Leptagonus</i>	394
<i>Dactylagnus</i>	455	<i>decagrammus</i> , <i>Hexagrammos</i>	373
<i>mundus</i>	455	<i>Labrax</i>	373
<i>Dactyleptus</i>	467	<i>Decapterus</i>	270
<i>Dactylobatus</i>	27	<i>hypodus</i>	270
<i>armatus</i>	27	<i>macarellus</i>	270
<i>dactyloptera</i> , <i>Scorpaena</i>	370	<i>punctatus</i>	270
<i>Dactylopterus</i>	408	<i>sanctaëhelene</i>	270
<i>Dactylopterus</i>	408	<i>Decapturus</i>	279
<i>pirapeda</i>	409	<i>decastrensis</i> , <i>Cottus</i>	387
<i>voltans</i>	409	<i>Decentrus lucens</i>	412
<i>dactylopterus</i> , <i>Helcolenus</i>	370	<i>decimalis</i> , <i>Serranus</i>	314
<i>Dactyloscopidae</i>	464	<i>declyfrons</i> , <i>Abudedefduf</i>	415
<i>Dactyloscopus</i>	464	<i>Euschistodus</i>	415
<i>cluctus</i>	454	<i>Salmo</i>	57
<i>lunaticus</i>	455	<i>clarkii</i>	57
<i>pectoralis</i>	454	<i>Vomer</i>	275
<i>poeyi</i>	455	<i>decivis</i> , <i>Seriola</i>	280
<i>tridigitatus</i>	454	<i>Decodon</i>	425
<i>zelotes</i>	455	<i>puellaris</i>	425
<i>Dactylostomias</i>	69	<i>decoratum</i> , <i>Bryostemma</i>	466
<i>filifer</i>	69, 70	<i>decoratus</i> , <i>Entomacrodus</i>	463
<i>dactylosus</i> , <i>Paraliparis</i>	404	<i>Iridio</i>	427
<i>Dæctor</i>	488	<i>Promicropterus</i>	317
<i>dækayi</i> , <i>Oxyrhina</i>	19	<i>decoris</i> , <i>Doratonotus</i>	429
<i>Dajaus</i>	254	<i>decurrens</i> , <i>Pleuronichthys</i>	225
<i>microps</i>	254	<i>defensor</i> , <i>Caranx</i>	273
<i>monticola</i>	254	<i>Dekaya</i>	357
<i>percoides</i>	254	<i>anomala</i>	357
<i>dakotensis</i> , <i>Chrosomus</i>	113	<i>dekayi</i> , <i>Acipenser</i>	34
<i>dalli</i> , <i>Gobius</i>	442	<i>Gasterosteus</i>	239
<i>Lythrypnus</i>	442	<i>Isuropolis</i>	19
<i>Dallia</i>	172	<i>Phycis</i>	213
<i>pectoralis</i>	172	<i>Pimelodus</i>	153
<i>dallii</i> , <i>Auctospina</i>	309	<i>Scomber</i>	258
<i>Pteropodus</i>	309	<i>Syngnathus</i>	242
<i>Dalliidæ</i>	172	<i>delalandii</i> , <i>Maiacocetus</i>	459
<i>Damalichthys</i>	412	<i>delalandii</i> , <i>Clinus</i>	459
<i>argyrosomus</i>	412	<i>Umbra</i>	48
<i>vacca</i>	412	<i>delicatissimus</i> , <i>Engraulis</i>	48
<i>damalis</i> , <i>Carpoides</i>	103	<i>deliclosa</i> , <i>Corvina</i>	348
<i>daneæ</i> , <i>Xenophthalmichthys</i>	69	<i>Hybopsis</i>	134
<i>darienensis</i> , <i>Darienichthys</i>	189	<i>Monias</i>	134
<i>Gambusia</i>	189	<i>Deloepis</i>	471
<i>Darienichthys</i>	189	<i>giganteus</i>	472
<i>darienensis</i>	189	<i>virgatus</i>	472
		<i>Delothyris</i>	220

List of scientific names—Continued

	Page		Page
<i>delphinus</i> , <i>Minomus</i>	104	<i>Diaphus</i>	168
<i>Pantosteus</i>	104	<i>theta</i>	168
<i>Deltaria</i>	25	<i>Dipterus</i>	297, 342
<i>Deltistes</i>	108	<i>aureolus</i>	342
<i>lucatus</i>	108	<i>axillaris</i>	342
<i>dendrica</i> , <i>Ramularia</i>	224	<i>californiensis</i>	341
<i>dendritica</i> , <i>Ancylospetta</i>	224	<i>dowii</i>	341
<i>dendriticum</i> , <i>Siphostoma</i>	242	<i>evermanni</i>	342
<i>dendriticus</i> , <i>Syngnathus</i>	242	<i>gracilis</i>	341
<i>denegatus</i> , <i>Pomacentrus</i>	415	<i>homonymus</i>	341
<i>dennyi</i> , <i>Carelliparis</i>	400	<i>lefroyi</i>	341
<i>Liparis</i>	400	<i>olistostomus</i>	342
<i>dentatus</i> , <i>Aspislus</i>	328	<i>peruvianus</i>	342
<i>Grammatostomias</i>	70	<i>dapterus</i> , <i>Amictichthys</i>	297
<i>Lucifuga</i>	479	<i>Furcimanus</i>	476
<i>Lutjanus</i>	325	<i>Lycodes</i>	476
<i>Paralichthys</i>	224	<i>diazzi</i> , <i>Chirostoma</i>	251
<i>Pleuronectes</i>	224	<i>Zoogoneticus</i>	182
<i>Stygicola</i>	479	<i>Dibranchus</i>	508
<i>Tropidinthus</i>	328	<i>atlanticus</i>	508
<i>Upeneus</i>	343	<i>diceraus</i> , <i>Ceratocottus</i>	381
<i>dentex</i> , <i>Brycon</i>	98	<i>Cottus</i>	381
<i>Corvina</i>	344	<i>Dicerobatus</i>	31
<i>Engraulis</i>	51	<i>Dicholomycter</i>	499
<i>Menidia</i>	247	<i>Dicrolene</i>	483
<i>Odontoscion</i>	344	<i>filamentosa</i>	483
<i>Osmerus</i>	67	<i>intriongra</i>	483
<i>denticulatus</i> , <i>Girardinus</i>	189	<i>nigra</i>	483
<i>dentiens</i> , <i>Cryptotomus</i>	431	<i>pullata</i>	483
<i>Scarus</i>	431	<i>Dicrotus</i> , <i>parvipinnis</i>	261
<i>denudata</i> , <i>Gonostoma</i>	72	<i>diego</i> , <i>Pneumatophorus</i>	258
<i>denudatum</i> , <i>Gonostoma</i>	72	<i>Scomber</i>	258
<i>deppii</i> , <i>Cichlaurus</i>	416	<i>diencephalus</i> , <i>Eupomacentrus</i>	413
<i>Heros</i>	416	<i>disformis</i> , <i>Carpoides</i>	103
<i>depraudus</i> , <i>Esox</i>	173	<i>digitatus</i> , <i>Lycodes</i>	475
<i>depressa</i> , <i>Belone</i>	196	<i>digitatus</i> , <i>Holomycteronus</i>	482
<i>Fistularia</i>	245	<i>dignabilis</i> , <i>Pseudopleuronectes</i>	227
<i>Rhamdia</i>	158	<i>digrammus</i> , <i>Pleuronectes</i>	226
<i>depressus</i> , <i>Lonchurus</i>	352	<i>digueti</i> , <i>Neomugil</i>	254
<i>Derepodichthysidae</i>	479	<i>Tæniconger</i>	82
<i>Derepodichthys</i>	479	<i>directive</i> , <i>Notosema</i>	225
<i>alepidotus</i>	479	<i>directive</i> , <i>Notosema</i>	225
<i>Derichthyidae</i>	77	<i>directive</i> , <i>Alburnus</i>	125
<i>Derichthys</i>	77	<i>Notropsis</i>	126
<i>serpentinus</i>	77	<i>Dilibomycter</i>	499
<i>dermatinus</i> , <i>Lycodapus</i>	478	<i>Dimalacobentrus</i>	430
<i>Salmo</i>	54	<i>dimidiatus</i> , <i>Gasterosteus</i>	237
<i>Dermatochaefer</i>	97	<i>Julis</i>	427
<i>Dermatolepis</i>	313	<i>Leucus</i>	122
<i>inermis</i>	313	<i>Serranus</i>	314
<i>punctatus</i>	313	<i>Syngnathus</i>	240
<i>zanculus</i>	313	<i>Trisoptropis</i>	314
<i>Dermatostethus</i>	240	<i>Directus truncatus</i>	35
<i>punctipinnis</i>	240	<i>Dinemus</i>	234
<i>desmarestia</i> , <i>Rala</i>	25	<i>venustus</i>	234
<i>deterior</i> , <i>Julis</i>	429	<i>dinemus</i> , <i>Minnius</i>	123
<i>detrusus</i> , <i>Gillichthys</i>	444	<i>dinoceros</i> , <i>Citharichthys</i>	219
<i>Deuterodon</i> , <i>atricaudata</i>	98	<i>Diodon</i>	501, 502
<i>deweegeri</i> , <i>Prionodes</i>	319	<i>antennatus</i>	502
<i>Serranus</i> (<i>Paralabrax</i>).....	319	<i>asper</i>	501
<i>Diabasis</i>	329	<i>atinga</i>	502
<i>flavolineatus</i>	330	<i>brachylatus</i>	501
<i>obliquatus</i>	330	<i>carinatus</i>	503
<i>parra</i>	329	<i>echinus</i>	501
<i>steindachneri</i>	330	<i>4-maculatus</i>	501
<i>Diabolichthys</i>	82	<i>fuliginosus</i>	502
<i>elliotti</i>	32	<i>geometricus</i>	502
<i>Diacope</i>	324	<i>lineatus</i>	502
<i>viridis</i>	324	<i>holacanthus</i>	501
<i>diadema</i> , <i>Scarus</i>	433	<i>holocanthus</i>	501
<i>Diagramma cavifrons</i>	335	<i>hystric</i>	501
<i>melanospilum</i>	332	<i>liturosus</i>	501
<i>Dialarchus</i>	381	<i>maculato-striatus</i>	502
<i>synderi</i>	391	<i>maculatus</i> var.	501
<i>Diana</i>	204	<i>maculifer</i>	501
<i>semilunata</i>	264	<i>melanopsis</i>	501
<i>diaphana</i> , <i>Hydrargyrus</i>	178	<i>meuleini</i>	502
<i>Sternoptix</i>	74	<i>mola</i>	503
<i>diaphanes</i> , <i>Raja</i>	25	<i>multimaculatus</i>	501
<i>Raja</i>	25	<i>nigrolineatus</i>	502
<i>Diaphanichthys</i>	79	<i>novem-maculatus</i>	501
<i>diaphanus</i> , <i>Callurus</i>	299	<i>pilosus</i>	501
<i>Zygoectes</i>	178	<i>punctatus</i>	501
<i>Diaphasia</i>	486	<i>reticulatus</i>	502

List of scientific names—Continued

	Page		Page
<i>Diodon rivulatus</i>	502	<i>Dolopichthys</i>	509
<i>schoepfi</i>	502	<i>anisacanthus</i>	509
<i>sex-maculatus</i>	501	<i>gracilispinis</i>	509
<i>spinosissimus</i>	501	<i>heteracanthus</i>	510
<i>spinulos</i>	502	<i>luetkeni</i>	510
<i>verrucosus</i>	502	<i>microlophus</i>	509
<i>Diodon plumier</i> , le.....	502	<i>dombeyi</i> , <i>Gastrobranchus</i>	8
<i>Diodontidae</i>	501	<i>dominicensis</i> , <i>Gambusia</i>	186
<i>diomediana</i> , <i>Aphoristia</i>	231	<i>Limia</i>	191
<i>diomedianus</i> , <i>Hoplunnis</i>	81	<i>Platyceclius</i>	191
<i>Sympnurus</i>	231	<i>Poecilia</i>	191
<i>Dionda</i>	141	<i>dominicus</i> , <i>Protoporus</i>	119
<i>amara</i>	144	<i>Doncella</i>	426
<i>argentosa</i>	144	<i>dorado</i> , <i>Coryphaen</i>	284
<i>chrysitis</i>	144	<i>Doratonotus</i>	429
<i>conchi</i>	144	<i>boekeli</i>	429
<i>episcopa</i>	144	<i>decorus</i>	429
<i>grisea</i>	144	<i>megalepis</i>	429
<i>melanops</i>	144	<i>thalassinus</i>	429
<i>nubila</i>	144	<i>Dormitator</i>	436
<i>papalis</i>	144	<i>guindachii</i>	436
<i>plumbea</i>	144	<i>lineatus</i>	436
<i>serena</i>	144	<i>maculatus</i>	436
<i>spadicea</i>	144	<i>microphthalmus</i>	436
<i>texensis</i>	144	<i>dormitor</i> , <i>Platyccephalus</i>	435
<i>Diplolites</i>	298	<i>dormitatrix</i> , <i>Eleotris</i>	435
<i>variabilis</i>	298	<i>dormitor</i> , <i>Gobionorus</i>	435
<i>Diplanchias</i>	503	<i>Dorosoma</i>	46
<i>nsus</i>	503	<i>cepedianum</i>	46
<i>Diplectrum</i>	318	<i>exile</i>	46
<i>euryleptrum</i>	318	<i>insoclabills</i>	46
<i>formosum</i>	318	<i>notata</i>	46
<i>macropoma</i>	318	<i>Dorosomidae</i>	46
<i>pacificum</i>	318	<i>dorsale</i> , <i>Hæmulon</i>	330
<i>radiale</i>	318	<i>dorsalis</i> , <i>Carangoides</i>	274
<i>scirurus</i>	318	<i>Citula</i>	274
<i>diplemia</i> , <i>Semotilus</i>	117	<i>Halactractus</i>	279
<i>Diplestion</i>	286	<i>Hybopsis</i>	134
<i>fasciatus</i>	293	<i>Hypsypops</i>	416
<i>Diplodus</i>	338	<i>Macrurus</i>	207
<i>argenteus</i>	338	<i>Mustelus</i>	13
<i>holbrookii</i>	338	<i>Semotilus</i>	117
<i>sargus</i>	338	<i>Seriola</i>	279
<i>diplopros</i> , <i>Eosebastes</i>	367	<i>Stegastes</i>	416
<i>Sebastichthys</i>	367	<i>Umbrina</i>	350
<i>Diplopomidae</i>	11	<i>Vomer</i>	275
<i>diploaenia</i> , <i>Belone</i>	198	<i>dorsatum</i> , <i>Cichlasoma</i>	420
<i>Bodianus</i>	425	<i>dorsatus</i> , <i>Astatheros</i>	420
<i>Harpe</i>	425	<i>Petromyzon marinus</i>	9
<i>Strongylura</i>	196	<i>dorsoniaculata</i> , <i>Girella</i>	339
<i>Dipterodon</i>	295	<i>dorsopunicea</i> , <i>Pomacentrus</i>	413
<i>dipterus</i> , <i>Amphotistius</i>	29	<i>Dorsarius</i>	339
<i>Dasyatis</i>	29	<i>Dorylchthys</i>	248
<i>Dipturus</i>	24	<i>lineatus</i>	243
<i>dipus</i> , <i>Microdesmus</i>	473	<i>Doryrhamphus</i>	243
<i>discobolus</i> , <i>Catostomus</i>	107	<i>californiensis</i>	243
<i>Discocephalidae</i>	448	<i>sierra</i>	243
<i>Discopyge</i>	27	<i>dovii</i> , <i>Apogon</i>	295
<i>ommatum</i>	27	<i>Gambusia</i>	186
<i>dispar</i> , <i>Scopelengys</i>	170	<i>Gymnothorax</i>	92
<i>Zygonectes</i>	178	<i>Haplochilus</i>	179
<i>dispus</i> , <i>Iridio</i>	427	<i>Heros</i>	418
<i>Platyglossus</i>	427	<i>Muræna</i>	92
<i>displurus</i> , <i>Centropristes</i>	320	<i>Opisthophterus</i>	45
<i>disinclus</i> , <i>Couesius</i>	137	<i>Oxyzygonectes</i>	179
<i>Erimystax</i>	138	<i>Paraconodon</i>	332
<i>Leucosomus</i>	137	<i>Parapetenia</i>	418
<i>Luxilus</i>	138	<i>Poecilia</i>	192
<i>distinctum</i> , <i>Sparisoma</i>	432	<i>Pristigaster</i>	45
<i>distinctus</i> , <i>Scarus</i>	432	<i>Pristipoma</i>	332
<i>Distrema atripes</i>	411	<i>dowei</i> , <i>Anableps</i>	194
<i>brevipinne</i>	409	<i>dowii</i> , <i>Leptarius</i>	148
<i>orthonotus</i>	411	<i>Thalassophryne</i>	488
<i>Dixoniina</i>	40	<i>dowii</i> , <i>Diapterus</i>	341
<i>nemoptera</i>	40	<i>Exocetus</i>	201
<i>dodecaedron</i> , <i>Ocellia</i>	394	<i>Selenaspis</i>	148
<i>dodecaëdrus</i> , <i>Agonus</i>	394	<i>Draconetidae</i>	450
<i>dolifyn</i> , <i>Coryphæna</i>	284	<i>acanthopoma</i>	450
<i>dolichocephalus</i> , <i>Gobius</i>	442	<i>Draconetidæ</i>	450
<i>dolichogaster</i> , <i>Blennius</i>	488	<i>Drepanopsetta</i>	222
<i>Rhodymenichthys</i>	488	<i>Dropsaridae</i>	208
<i>Dollodon</i>	276	<i>drummond-hayi</i> , <i>Epinephelus</i>	312
<i>Dollosa</i>	203	<i>drummondii</i> , <i>Otolithus</i>	365
<i>cyclolepis</i>	203	<i>dubia</i> , <i>Seriola</i>	280
<i>dolomieu</i> , <i>Micropterus</i>	298	<i>dubium</i> , <i>Exoglossum</i>	146

List of scientific names—Continued

	Page		Page
dubius, <i>Ammodytes</i>	401	Echinorhinus.....	22
<i>Flerasfer</i>	486	<i>brucus</i>	22
<i>Menepterus</i>	309	<i>obesus</i>	22
<i>Serranus</i>	309	<i>echinus</i> , <i>Diodon</i>	501
ductor, <i>Gasterosteus</i>	278	Echilodon.....	486
<i>Nauirates</i>	278	<i>drummondii</i>	486
dugesi, <i>Algancea</i>	116	Echiopsis.....	88
<i>Amelurus</i>	153	Echistoma.....	70
<i>Haustor</i>	153	<i>barbatum</i>	70
dugesii, <i>Fundulus</i>	182	<i>margarita</i>	70
<i>Zoogeneticus</i>	182	<i>margaritum</i>	70
duclis, <i>Argyreus</i>	139	etenes, <i>Careproctus</i>	403
<i>Rhinichthys</i>	139	<i>Microponodon</i>	349
Dules.....	320	Ectenias, <i>brunneus</i>	265
<i>auriga</i>	320	edentula, <i>Platirostra</i>	38
<i>flaviventris</i>	320	edentulus, <i>Cetenograulis</i>	51
<i>tenuirurus</i>	305	Engraulis.....	51
dumerili, <i>Squatina</i>	22	Edriolichthys.....	508
dumerili, <i>Aplonichthys</i>	230	schmidti.....	508
<i>Caranx</i>	279	edwardsi, <i>Sciaena</i>	353
<i>Polyclirrus</i>	851	edwardsi, <i>Psenes</i>	269
<i>Polyclemus</i>	351	Stilibicus.....	89
<i>Seriola</i>	279	eeltenkee, <i>Myliobatis</i>	30
duodecim, <i>Engraulis</i>	48	<i>fulgens</i> , <i>Zithopora</i>	168
duplex, <i>Orthopristis</i>	335	<i>Artilia</i>	287
duquesnii, <i>Catostomus</i>	110	<i>Boleosoma</i>	287
durvillii, <i>Argyropelecus</i>	74	<i>Larimus</i>	344
duryi, <i>Etheostoma</i>	286	eglanteria, <i>Raja</i>	25
dussumieri, <i>Brama</i>	265	<i>Raja</i>	25
<i>Seriola</i>	278	egmontis, <i>Ahlia</i>	82
Dussumieriella, <i>stolifera</i>	46	Myrophis.....	82
Dussumieriidae.....	45	egregia, <i>Cheonda</i>	118
dux, <i>Lachnolaimus</i>	425	Tigoma.....	118
dviniensis, <i>Platesa</i>	227	egregius, <i>Exocetoides</i>	454
earlii, <i>Phycis</i>	212	egretta, <i>Bellator</i>	408
Ebisus.....	373	Prionatus.....	408
<i>sagamius</i>	373	eigenmanni, <i>Acutomentum</i>	366
Echelidae.....	82	Cichlasoma.....	417
Echelus caudillimbatus.....	79	<i>Evara</i>	142
Echemythes.....	216	<i>Gobiesox</i>	490
Echeneidae.....	448	<i>Gobius</i>	439
Echenes.....	443	<i>Rhinogobius</i>	439
<i>albacauda</i>	448	<i>Rimicola</i>	490
<i>albescens</i>	449	<i>Sebastodes</i>	366
<i>apicalis</i>	449	<i>Theraps</i>	417
<i>australis</i>	449	Eigenmannia.....	101
<i>brachypтера</i>	449	<i>virescens</i>	101
<i>fusca</i>	449	eigenmanni, <i>Anchovia</i>	60
<i>guianan</i>	449	<i>Anchoviella</i>	60
<i>holbrookii</i>	449	eliseni, <i>Characodon</i>	183
<i>jacobae</i>	449	elaborata, <i>Muraena</i>	93
<i>lineata</i>	449	elaboratus, <i>Gymnothorax</i>	93
<i>lunata</i>	448	Elacate.....	281
<i>metallicus</i>	449	<i>atlantica</i>	281
<i>naucrates</i>	448	<i>bivittata</i>	281
<i>neurbrates</i>	448	<i>falcipinnis</i>	281
<i>neuratoidea</i>	449	<i>malabarica</i>	281
<i>nieuhoffi</i>	449	<i>motta</i>	281
<i>osteochir</i>	450	<i>pondiceriana</i>	281
<i>pallida</i>	449	Elacatinus.....	442
<i>parva</i>	449	<i>oceanoops</i>	442
<i>postica</i>	449	Elagatis.....	279
<i>quatuordecimlaminatus</i>	449	<i>bipinnulatus</i>	279
<i>remora</i>	449	Elanura.....	380
<i>remoroides</i>	449	<i>forficata</i>	380
<i>scutata</i>	449	Elaphocottus.....	389
<i>sexdecimlamellata</i>	449	<i>Elaphotoxon</i>	271
<i>sphrenarum</i>	449	<i>bartholomaei</i>	272
<i>squalipeta</i>	449	<i>ruber</i>	271
<i>tetrapturonum</i>	449	Elapsophis.....	87
<i>tropica</i>	450	Elasmobranchii.....	10
<i>verticalis</i>	449	<i>elassochir</i> , <i>Noturus</i>	156
<i>vittata</i>	449	Elassodiscus.....	402
Echidna.....	94	<i>tremebundus</i>	402
<i>catenata</i>	94	<i>elassodon</i> , <i>Hippoglossoides</i>	222
<i>chionostigma</i>	94	Elassomea.....	305
<i>cocosa</i>	94	<i>evergladei</i>	305
<i>flavoscripta</i>	94	<i>zonata</i>	305
<i>fuscomaculata</i>	94	<i>tonatum</i>	305
<i>nebulosa</i>	94	Elassomidae.....	305
<i>nocturna</i>	94	Elastoma.....	328
<i>scabra</i>	94	elater, <i>Malthe</i>	507
echinatum, <i>Leiodon</i>	21	<i>Zaleutes</i>	507
echinita, <i>Scorpaena</i>	372	Elattonistius.....	41
Echinorhinidae.....	23	Elattarchus.....	345
		<i>archidium</i>	345

List of scientific names—Continued

	Page		Page
<i>elattura, Netuma</i>	149	<i>elongatus, Luxilius</i>	120
<i>elatturus, Arius</i>	149	<i>Megalops</i>	39
<i>elcapitanensis, Syngnathus</i>	242	<i>Ophiodon</i>	374
<i>elcapitanensis, Syngnathus</i>	242	<i>Osmerus</i>	67
<i>electricus, Rhinobatos</i>	23	<i>Pomadasys</i>	333
<i>elegans, Cottus</i>	381	<i>Pomotis</i>	300
<i>Cyphosus</i>	339	<i>Sebastes</i>	368
<i>Cyprinodon</i>	181	<i>Syphurus</i>	231
<i>Gasterosteus</i>	238	<i>Elopidae</i>	39
<i>Gibbonsia</i>	458	<i>Elopoides</i>	39
<i>Gila</i>	114	<i>Elops</i>	39
<i>Hæmulon</i>	329	<i>hawaiiensis</i>	39
<i>Iridio</i>	427	<i>inermis</i>	39
<i>Kyphosus</i>	339	<i>saurus</i>	39
<i>Labeo</i>	109	<i>elucaen, Syngnathus</i>	241
<i>Mesopion</i>	328	<i>emarginatum, Scarus</i>	433
<i>montereyensis, Gibbonsia</i>	458	<i>emarginatus, Lobotes</i>	325
<i>Myxodes</i>	458	<i>Serranus</i>	315
<i>Nanostoma</i>	289	<i>Embassichthys</i>	228
<i>Orthragoriscus</i>	503	<i>bathybius</i>	228
<i>Pimalepterus</i>	339	<i>Embiotoca</i>	411
<i>Eleginus</i>	209	<i>caryi</i>	412
<i>nawaga</i>	210	<i>cassidyi</i>	411
<i>Electrides</i>	435	<i>jacksoni</i>	411
<i>Eleotris</i>	436	<i>lateralis</i>	411
<i>abacuris</i>	437	<i>lineata</i>	411
<i>amblyopsis</i>	437	<i>ornata</i>	411
<i>capite plagioplateo</i>	436	<i>perspicabilis</i>	411
<i>(Culius) belizianus</i>	436	<i>webbi</i>	411
<i>dormitatrix</i>	435	<i>Embiotocidae</i>	409
<i>grandisquama</i>	436	<i>Emblemaria</i>	464
<i>guavina</i>	436	<i>atlantica</i>	464
<i>gyrinus</i>	436	<i>marki</i>	464
<i>latifrons</i>	436	<i>nivipes</i>	464
<i>latifasciatus</i>	437	<i>oculocirrís</i>	464
<i>longiceps</i>	435	<i>pandionis</i>	464
<i>mauritii</i>	268	<i>Emblemariidæ</i>	464
<i>mugiloides</i>	436	<i>emblematicus, Gobius</i>	443
<i>omocyanus</i>	436	<i>Microgobius</i>	443
<i>perniger</i>	437	<i>Scarus</i>	436
<i>pictus</i>	437	<i>embryum, Allocottus</i>	391
<i>pisonis</i>	436	<i>Oligocottus</i>	391
<i>semifasciata</i>	438	<i>Embryx</i>	473
<i>semifasciatus</i>	438	<i>crassilabris</i>	473
<i>sima</i>	436	<i>crotalinius</i>	473
<i>smaragdus</i>	437	<i>parallelus</i>	473
<i>somnolentus</i>	436	<i>embryx, Gerres</i>	342
<i>tubularis</i>	437	<i>emiliae, Opsopœodus</i>	115
<i>Elephas, Squinus</i>	20	<i>Eumeekia</i>	428
<i>eleuthera, Rabida</i>	158	<i>venusta</i>	428
<i>Eleutheractis</i>	316	<i>emmeline, Arius</i>	151
<i>cornuta</i>	316	<i>A verruncus</i>	395
<i>eleutherus, Noturus</i>	156	<i>Tachysurus</i>	151
<i>elizabeth, Lepidosteus</i>	37	<i>Emmelas</i>	364
<i>Elllops</i>	151	<i>glaucus</i>	354
<i>ellioti, Thorichthys</i>	422	<i>emmelas, Lepophidium</i>	484
<i>ellotti, Diabolichthys</i>	32	<i>Leptophidium</i>	484
<i>ellipsoides, Lebias</i>	181	<i>emoryi, Gila</i>	114
<i>ellipticus, Chatoctus</i>	46	<i>emperor, Astyanax</i>	97
<i>Cynatogaster</i>	410	<i>Bryconamericus</i>	97
<i>Platophrys</i>	217	<i>Empetrichthys</i>	182
<i>Pleuronectes</i>	217	<i>Empetrichthys</i>	182
<i>Ellops</i>	34	<i>merriami</i>	182
<i>elongata, Anchovia</i>	49	<i>emphesus, Ecsobastes</i>	367
<i>Anchoviella</i>	49	<i>Sebastodes</i>	367
<i>Aporistia ornata var.</i>	231	<i>Emphyicus</i>	213
<i>Avocettina</i>	83	<i>emphysetus, Bagrus (Sciades)</i>	148
<i>Bothrocarpopsis</i>	477	<i>Sciadiceithys</i>	148
<i>Clupea</i>	41	<i>encionomus, Gobius</i>	439
<i>Cyclothona</i>	72	<i>Rhinogobius</i>	439
<i>Fossilia</i>	190	<i>encaustus, Chapalichthys</i>	184
<i>Umbrina</i>	351	<i>Characodon</i>	184
<i>Umbrula</i>	351	<i>encenadæ, Rhinoptera</i>	31
<i>elongatum, Gonostoma</i>	72	<i>Enchelycephali</i>	77
<i>elongatus, Aulophallus</i>	190	<i>Enchelycore</i>	90
<i>Bramocharax</i>	99	<i>euryrhina</i>	90
<i>Carpoides</i>	104	<i>nigricans</i>	90
<i>Catostomus</i>	102	<i>nigrocastaneus</i>	90
<i>Clinostomus</i>	120	<i>Enchelyopus</i>	209, 202, 473
<i>Cyclostomus</i>	102	<i>americanus</i>	218
<i>Eucinostomus</i>	341	<i>barbatus</i>	479
<i>californiensis</i>	341	<i>cimbrius</i>	209
<i>Hispanicus</i>	368	<i>mediterraneus</i>	208
<i>Labeo</i>	109	<i>regalis</i>	213
<i>Labichthys</i>	83		

List of scientific names—Continued

	Page		Page
<i>enchrurus</i> , <i>Chromis</i> .	413	<i>Eosebastes deani</i> .	367
<i>Heliases</i> .	413	<i>diplopros</i> .	367
<i>Engrasicholus</i> .	47	<i>emphaeus</i> .	367
<i>Encrotos</i> .	180	<i>intruder</i> .	368
<i>Engraulidae</i> .	47	<i>melanostomus</i> .	368
<i>engraulinus</i> , <i>Photogenis leucops</i> .	123	<i>saxicola</i> .	367
<i>Engraulis</i> .	47	<i>semicinctus</i> .	367
<i>argyrophanus</i> .	48	<i>swifti</i> .	367
<i>brevis</i> .	51	<i>Eperlanus</i> .	67
<i>brevirostris</i> .	51	<i>Ephippidae</i> .	358
<i>chærostomus</i> .	48	<i>ephippium</i> , <i>Chaetodon</i> .	359
<i>clupeoides</i> .	50	<i>Plectropoma</i> .	321
<i>compressus</i> .	47-50	<i>Ephippus gigas</i> .	358
<i>cubanus</i> .	48	<i>zonatus</i> .	358
<i>delicatissimus</i> .	48	<i>Epipodus</i> .	214
<i>denter</i> .	51	<i>epleurorum</i> , <i>Chromis</i> .	280
<i>duodecim</i> .	48	<i>Epigieichthys</i> .	409
<i>edentulus</i> .	51	<i>atropurpureus</i> .	469
<i>grossidens</i> .	51	<i>Epigonichthysidæ</i> .	7
<i>janeiro</i> .	51	<i>Epigonus</i> .	206
<i>lemniscatus</i> .	48	<i>macrophthalmus</i> .	296
<i>louisiana</i> .	48	<i>occidentalis</i> .	296
<i>macrolepidotus</i> .	50	<i>Epilichodon</i> , <i>Petromyzon</i> .	10
<i>mitchilli</i> .	48	<i>Epinephelus afer</i> .	312
<i>mordax</i> .	47	<i>Epinepheloidæ</i> .	308
<i>mysticetus</i> .	51	<i>Epinephelus</i> .	310
<i>nanus</i> .	47	<i>adscensionis</i> .	310
<i>panamensis</i> .	50	<i>analogus</i> .	310
<i>perfasciatus</i> .	47	<i>angustifrons</i> .	311
<i>pooyl</i> .	51	<i>brachysoma</i> .	311
<i>productus</i> .	50	<i>chalinius</i> .	315
<i>spinifer</i> .	50	<i>ciliatus</i> .	365
<i>tricolor</i> .	48	<i>cubanus</i> .	311
<i>engymen</i> , <i>Ctenograulis</i> .	51	<i>drummondhayi</i> .	312
<i>Engyophrys</i> .	220	<i>flavolimbatus</i> .	311
<i>sancti-laurentii</i> .	220	<i>guaza</i> .	310
<i>Enneacanthus</i> .	303	<i>guttatus</i> .	311
<i>gloriosus</i> .	304	<i>jordani</i> .	314
<i>margarotis</i> .	304	<i>labriformis</i> .	311
<i>obesus</i> .	304	<i>latefasciatus</i> .	310
<i>pinniger</i> .	304	<i>lightfooti</i> .	313
<i>Enneacentrus</i> .	309	<i>marginalis</i> .	310
<i>Enneapterectes</i> .	458	<i>mentzeli</i> .	310
<i>Enneapterygius</i> .	458	<i>morio</i> .	311, 312
<i>carminalis</i> .	458	<i>mystacinus</i> .	310
<i>pusillus</i> .	458	<i>niphobles</i> .	312
<i>Enneistius</i> .	309	<i>niveatus</i> .	311
<i>acanthistius</i> .	309	<i>ordinatus</i> .	311
<i>Enophryss</i> .	381	<i>rosaceus</i> .	314
<i>claviger</i> .	381	<i>ruber</i> .	309, 315
<i>taurinus</i> .	381	<i>selicauda</i> .	311
<i>ensifera</i> , <i>Bairdiella</i> .	345	<i>striatus</i> .	311, 318
<i>Sciæna</i> .	345	<i>Epinnula</i> .	201
<i>ensiferus</i> , <i>Centropomus</i> .	306	<i>magistralis</i> .	261
<i>ensis</i> , <i>Gaidropsarus</i> .	208	<i>episcola</i> , <i>Dionda</i> .	144
<i>Motella</i> .	208	<i>episcoli</i> , <i>Brachyrhaphis</i> .	187
<i>Sphyraena</i> .	255	<i>Gambusa</i> .	187
<i>entemedor</i> , <i>Narcine</i> .	27	<i>Episema</i> .	125
<i>Entomacrodus</i> .	463	<i>callisema</i> .	132
<i>chloستicus</i> .	463	<i>jejuna</i> .	125
<i>cruentatus</i> .	463	<i>Epitrachys</i> .	281
<i>decoratus</i> .	463	<i>epsetus</i> , <i>Anchoviella</i> .	48
<i>margaritaceus</i> .	463	<i>Esox</i> .	48
<i>nigricans</i> .	463	<i>Eptatretidae</i> .	8
<i>antomelas</i> , <i>Acutomentum</i> .	366	<i>equatorialis</i> , <i>Chiopsis</i> .	82
<i>Sebastichthys</i> .	366	<i>Raja</i> .	26
<i>entomophagus</i> , <i>Leucichthys</i> .	63	<i>Eques</i> .	352
<i>Entosphenus</i> .	9	<i>acuminatus umbrosus</i> .	353
<i>ciliatus</i> .	10	<i>americanus</i> .	352, 353
<i>tridentatus</i> .	9	<i>balteatus</i> .	353
<i>Enypnias</i> .	445	<i>lanceolatus</i> .	353
<i>seminudus</i> .	445	<i>lineatus</i> .	353
<i>Eopsetta</i> .	222	<i>pulcher</i> .	353
<i>jordanii</i> .	222	<i>punctatus</i> .	353
<i>eos</i> , <i>Arbaciosa</i> .	491	<i>viola</i> .	353
<i>Boleichtthys</i> .	294	<i>equestris</i> , <i>Arius</i> .	149
<i>Chrosomus</i> .	113	<i>Balistes</i> .	491
<i>Gobiesox</i> .	491	<i>Equetus</i> .	352
<i>Othonops</i> .	447	<i>equirostrum</i> , <i>Scomberesox</i> .	197
<i>Pronotogrammus</i> .	322	<i>equisetis</i> , <i>Coryphaena</i> .	265
<i>Sebastodes</i> .	368	<i>erate</i> , <i>Lobotes</i> .	323
<i>Sebastomus</i> .	368	<i>erebennus</i> , <i>Ameiurus</i> .	153
<i>Eosebastes</i> .	366	<i>ebrebus</i> , <i>Muraena</i> .	92, 93
<i>aurora</i> .	366	<i>eremogenes</i> , <i>Salmo</i> .	58
<i>crameri</i> .	367		

List of scientific names—Continued

	Page		Page
<i>erethizon</i> , <i>Arothron</i>	600	<i>Erythrinus brasiliensis</i>	99
<i>Eretmichthys</i>	481	<i>macrodon</i>	99
<i>ocella</i>	481	<i>microcephalus</i>	99
<i>pinnatus</i>	481	<i>trahira</i>	99
<i>erfarcha</i> , <i>Atherinella</i>	249	<i>erythrocillus</i> , <i>Albula</i>	40
<i>Copelandia</i>	304	<i>erythrogaster</i> , <i>Chrosomus</i>	113
<i>Eurystole</i>	249	<i>Luxilus</i>	113
<i>Ericara</i>	53	<i>Serranus</i>	312
<i>salmonea</i>	53, 54	<i>erythrogasterum</i> , <i>Pocilosoma</i>	291
<i>Erichthys</i>	301	<i>erythrops</i> , <i>Gobiesox</i>	490
<i>Ericosma</i>	284	<i>Ichthelis</i>	303
<i>evides</i>	284	<i>Sicyas</i>	400
<i>Erieteis</i>	460	<i>erythroptera</i> , <i>Pimelodus</i>	152
<i>kalishore</i>	460	<i>erythrorhynchos</i> , <i>Salmo</i>	61
<i>Ericymba</i>	141	<i>erythrus</i> , <i>Catostomus</i>	110
<i>buccata</i>	142	<i>escamblie</i> , <i>Zygoectes</i>	178
<i>ericymba</i> , <i>Sciema</i>	246	<i>eschrichtii</i> , <i>Oneirodes</i>	504
<i>Stellifer</i>	246	<i>Escolar</i>	260
<i>erlensis</i> , <i>Argyrosomus</i>	63	<i>niger</i>	260, 281
<i>Leucichthys</i>	63	<i>violaceus</i>	260
<i>Erilepidæ</i>	373	<i>esculentus</i> , <i>Carangus</i>	273
<i>Erilepis</i>	373	<i>Conger</i>	79
<i>zonifer</i>	373	<i>Merluccius</i>	214
<i>Erimonax</i>	138	<i>Eslopsarum</i>	250
<i>monacus</i>	138	<i>arge</i>	250
<i>Erimystax</i>	138	<i>Esloscopus</i>	455
<i>dissimilis</i>	138	<i>zelotes</i>	455
<i>watauga</i>	138	<i>esmarkii</i> , <i>Lycodes</i>	474
<i>Erimyzon</i>	108	<i>Esocidae</i>	172
<i>gooden</i>	109	<i>esopus</i> , <i>Labeo</i>	109
<i>oblongus</i>	109	<i>estor</i> , <i>Gila</i>	120
<i>sucetta</i>	109	<i>Esox</i>	172, 173
<i>erinacea</i> , <i>Rain</i>	25	<i>americanus</i>	172
<i>Raja</i>	25	<i>australis</i>	173
<i>erinaceus</i> , <i>Trichocyclus</i>	501	<i>barracuda</i>	255
<i>Erineimus</i>	137	<i>bozeus</i>	173
<i>hyalinus</i>	137	<i>brusiliensis</i>	199
<i>hypsinotus</i>	138	<i>crassus</i>	173
<i>labrosus</i>	138	<i>cypho</i>	173
<i>ruberfrons</i>	138	<i>depraudus</i>	173
<i>storerianus</i>	137	<i>epsetus</i>	48
<i>Erision</i>	355	<i>estor</i>	173
<i>albus</i>	356	<i>fuscatus</i>	173
<i>mackdonaldi</i>	356	<i>flavulus</i>	176
<i>nebulosus</i>	355	<i>immaculatus</i>	173
<i>parvipinnis</i>	355	<i>imperialis</i>	197
<i>reticulatus</i>	355	<i>lineatus</i>	173
<i>stolzmanni</i>	356	<i>longirostris</i>	106
<i>virespons</i>	356	<i>lucius americanus</i>	173
<i>xanthulus</i>	356	<i>lucoides</i>	173
<i>erithrus</i> , <i>Caraux</i>	273	<i>lugubris</i>	173
<i>Eritrema</i>	140	<i>malabaricus</i>	99
<i>Erizo</i>	501	<i>marinus</i>	106
<i>guanabana</i>	501	<i>masquinony</i>	172
<i>erochrous</i> , <i>Hololepis</i>	294	<i>immaculatus</i>	173
<i>Erogala</i>	129	<i>ohiensis</i>	172
<i>analostana</i>	129	<i>niger</i>	173
<i>cerulea</i>	130	<i>nobilior</i>	172
<i>callista</i>	130	<i>nobilis</i>	172
<i>camura</i>	129	<i>ohiensis</i>	172
<i>cercostigma</i>	130	<i>ornatus</i>	173
<i>chloristia</i>	129	<i>osseus</i>	36
<i>eurystoma</i>	130	<i>ovinus</i>	181
<i>fornosa</i>	129	<i>phaleratus</i>	173
<i>glaetatura</i>	129	<i>pisculentus</i>	176
<i>nivea</i>	129	<i>pisciculus</i>	176
<i>notata</i>	130	<i>porosus</i>	173
<i>pyrhomelas</i>	129	<i>raveneli</i>	173
<i>stigmatura</i>	129	<i>reticulatus</i>	173
<i>texana</i>	130	<i>salmoneus</i>	103, 172
<i>trichroistia</i>	130	<i>saurus</i>	197
<i>venusta</i>	130	<i>scomberius</i>	173
<i>whippii</i>	129	<i>spet</i>	255
<i>xenura</i>	130	<i>sphyrema</i>	255
<i>Erotelis</i>	437	<i>synodus</i>	163
<i>smaragdus</i>	437	<i>timucu</i>	195
<i>valenciennesi</i>	437	<i>tridecemlineatus</i>	173
<i>Error</i>	228	<i>tristechus</i>	38
<i>zachirus</i>	228	<i>umbrosus</i>	173
<i>Erychthys</i>	433	<i>vermiculatus</i>	173
<i>erythrum</i> , <i>Astatheros</i>	422	<i>vittatus</i>	173
<i>Erythreas</i>	420	<i>vulpes</i>	39
<i>erythraeus</i> , <i>Heros</i>	422	<i>zonatus</i>	176
<i>Erythrichthys</i>	420	<i>estor</i> , <i>Chiromysa</i>	250, 251
<i>erythrinolodes</i> , <i>Scarus</i>	432	<i>Esox</i>	172

List of scientific names—Continued

Page	Page
estor, <i>Gila</i>	121
Estrella.....	287
<i>atromaculata</i>	287
esunculus, <i>Atopichthys</i>	54
Etelides.....	328
<i>aqulonaris</i>	328
Eteis.....	328
<i>carbunculus</i>	328
<i>evirurus</i>	328
<i>oculatus</i>	328
Etheostoma.....	236, 293, 294
<i>artesiae</i>	292
<i>aubeenaubeli</i>	291
<i>australe</i>	293
<i>barrati</i>	294
<i>blennioides</i>	283, 286
<i>cerulea</i>	291
<i>calliura</i>	298
<i>camurum</i>	292
<i>caprodes</i>	282
<i>cinernea</i>	290
(<i>Cottogaster</i>) <i>uranidea</i>	285
<i>cragini</i>	292
<i>cumberlandicum</i>	293
<i>duryi</i>	286
<i>flabellaria</i>	293
<i>flabellata</i>	293
<i>fontinalis</i>	293
<i>fricksia</i>	291
<i>güntheri</i>	283
(<i>Hadropterus</i>) <i>cymatotenia</i>	285
<i>ninigue</i>	285
<i>ouachitae</i>	284
<i>squamatus</i>	285
<i>hildebrandti</i>	291
<i>lowei</i>	291
<i>julie</i>	292
<i>lepidogenys</i>	292
<i>linsleyi</i>	293
<i>longimanus</i>	286
<i>luteovinctum</i>	291
<i>macrocephalum</i>	284
<i>maculata</i>	290
<i>microperca</i>	294
<i>micropterus</i>	290
<i>nevilsense</i>	283
<i>nianguae spilotum</i>	285
<i>nigra</i>	287
(<i>Nothonotus</i>) <i>jordani</i>	290
<i>obeyense</i>	292
<i>olmstedi</i>	287
<i>pagei</i>	292
<i>parvipinne</i>	292
<i>pellucidum</i>	288
<i>petatum</i>	283
<i>phoxocephalum</i>	284
<i>podostemone</i>	286
<i>quapilla</i>	291
<i>rex</i>	283
(<i>Rhothcea</i>) <i>biennius</i>	287
<i>rupstre</i>	290
<i>sagitta</i>	293
<i>scovelli</i>	293
<i>squamiceps</i>	293
<i>swannanoa</i>	289
<i>tessellata</i>	290
<i>tippecanoe</i>	292
<i>tuscumbia</i>	293
(<i>Ulocentra</i>) <i>listrio</i>	286
<i>varifata</i>	289
<i>vereocundum</i>	287
<i>whipplei alabamae</i>	292
<i>wrighti</i>	285
<i>zonale arcansanum</i>	290
ethestoma, <i>Abona</i>	443
Etheostomidae.....	282
ethon, <i>Syngnathus</i>	241
Etmopterus.....	21
<i>hilliurus</i>	21
etowanus, <i>Catostomus nigricans</i>	107
Etropus.....	219
<i>crossotus</i>	219, 220
<i>micros</i>	219
<i>microstomus</i>	219
<i>rimosus</i>	220
Etrumeus.....	46
<i>acuminatus</i>	46
<i>micropus</i>	46
<i>sadina</i>	46
Eucilia.....	238
<i>inconstans</i>	238
<i>cayuga</i>	238
<i>pygmaea</i>	238
Euchalarodus.....	227
<i>putnami</i>	227
Eucinostomus.....	340, 341
<i>argenteus</i>	340, 341
<i>californiensis</i>	341
<i>elongatus</i>	341
<i>elongatus</i>	341
<i>gula</i>	341
<i>gulula</i>	341
<i>harengulus</i>	341
<i>meeki</i>	341
<i>productus</i>	341
<i>pseudogula</i>	341
<i>pseudo-gula</i>	341
Eucrotus.....	267
<i>ventralis</i>	267
Euctenogobius.....	441
<i>badius</i>	441, 442
<i>latuus</i>	442
<i>poeyi</i>	442
<i>sagittula</i>	441
<i>strigatus</i>	441
Eucyclogobius.....	444
<i>newberryi</i>	444
Eudulus.....	320
Eugaleus.....	14
Eugerres.....	342
<i>brasilianus</i>	342
<i>brevimanus</i>	343
<i>lineatus</i>	342
<i>mexicanus</i>	343
<i>plumieri</i>	343
Eulamia.....	15
<i>nicaraguensis</i>	16
<i>platyrhynchus</i>	15
eulepidotus, <i>Spherooides</i>	498
<i>Tetronotus (Spherooides)</i>	408
<i>eulepis</i> , <i>Microgobius</i>	444
Euleptorhamphus.....	198
<i>brevoorti</i>	198
<i>brevoortii</i>	198
<i>longirostris</i>	198
<i>velox</i>	198
Eumesogrammus.....	470
<i>praeclus</i>	470
Eunichtromus.....	398
<i>orbis</i>	398
<i>phrynooides</i>	398
<i>spiniferus</i>	398
<i>eumorphaeus</i> , <i>Chatoëssus</i>	44
Eumycteris.....	600
<i>blauvatus</i>	600
Eunichthys.....	410
<i>megalops</i>	410
Eupomacentrus.....	413
<i>adustus</i>	414
<i>analis</i>	413
<i>chrysus</i>	414
<i>dienceus</i>	413
<i>flavilatus</i>	414
<i>flaviventer</i>	413
<i>fuscus</i>	414
<i>gilli</i>	414
<i>leucorus</i>	414
<i>leucostictus</i>	413
<i>neopomacentrus</i>	414
<i>otophorus</i>	414
<i>partitus</i>	414
<i>planifrons</i>	414
<i>rectifranum</i>	414
Eupomotis.....	302
<i>gibbosus</i>	299, 302
<i>heros</i>	302
<i>holbrookii</i>	302
<i>holbrookii</i>	302
<i>longimanus</i>	302
<i>europaeus</i> , <i>Trachurus</i>	270

List of scientific names—Continued

	Page		Page
europaeus, <i>Aspidophorus</i> .	395	Evorhodus..	438
<i>Blennius</i>	467	breviceps	438
Eurymen.....	401	Evoxymetopon.....	262
<i>gyrinus</i>	401	<i>tæniatus</i>	263
Eurymyctera.....	91	exasperata, <i>Platyrhina</i>	24
eurypaus, <i>Hudsonius</i>	132	exasperatus, <i>Zapteryx</i>	24
euryops, <i>Bathylagus</i>	69	excisus, <i>Doryrhamphus</i>	243
<i>Icelus</i>	378	Execastides.....	454
<i>Myxostoma</i>	110	<i>egregius</i>	454
<i>Strongylura</i>	196	Exerpes.....	456
<i>Tylosurus</i>	196	<i>asper</i>	456
euryorus, <i>A pomotis</i>	209	exigua, <i>Anchoviella</i>	47
<i>Lepomis</i>	299	exiguus, <i>Bodianus argyro-leucos</i> et al.	345
Eurypharyngidae.....	95	<i>Stolephorus</i>	47
euryplectrum, <i>Diplectrum</i>	318	exile, <i>Dorosoma cepedianum</i>	48
euryhina, <i>Enchelycore</i>	90	<i>Siphonostoma</i>	243
Eurystole.....	249	exilicauda, <i>Lavinia</i>	113
<i>erinacea</i>	249	exilis, <i>Belone</i>	106
eyrstole, <i>Stolephorus</i>	48	<i>Boleichtthys</i>	204
eurystoma, <i>Erogala</i>	130	<i>Hippoglossoides</i>	222
Eurystomus.....	107	<i>Lyopsetta</i>	222
eurystomus, <i>Photogenis</i>	130	<i>Noturus</i>	156
Euscarus.....	431	<i>Rabida</i>	156
Euschistodus.....	414	<i>Strongydura</i>	198
<i>analogus</i>	416	<i>Syrichtes</i>	243
<i>concolor</i>	414	Eximia.....	390
<i>declivifrons</i>	414, 415	<i>rubellio</i>	390, 391
Eusebastes.....	363	eximus, <i>Cyprinodon</i>	181
Euseelachii.....	11	Exocetidae.....	199
Eusphyra.....	18	Exocetus.....	200
Eustomatodus.....	270	<i>acutus</i>	199
Euthynnus.....	258	<i>affinis</i>	201
<i>alleteratus</i>	258	<i>albidactylus</i>	202
<i>lineatus</i>	259	<i>appendiculatus</i>	201
<i>yaito</i>	259	<i>bahiensis</i>	202
Eutychelithus.....	352	<i>bicolor</i>	201
evansi, <i>Hybognathus</i>	143	<i>brachycephalus</i>	200
Evara.....	142	<i>californicus</i>	202
<i>eigenmanni</i>	142	<i>callipterus</i>	202
<i>tlahuacensis</i>	142	<i>chiliensis</i>	200
Eventognathi.....	102	<i>comatus</i>	201
Evepigymnus.....	270	<i>cyanopterus</i>	202
evergladei, <i>Elassoma</i>	305	<i>(Cypsilurus) procne</i>	201
evermanni, <i>Alvordius</i>	284	<i>dowii</i>	201
<i>Arius</i>	151	<i>evolans</i>	200
<i>Astatheros</i>	420	<i>exsiliens</i>	200
<i>Atherina</i>	246	<i>fasciatus</i>	200
<i>Atherinella</i>	248	<i>fucorum</i>	202
<i>Cheonda</i>	118	<i>furcatus</i>	201
<i>Cichlasoma</i>	420	<i>georgianus</i>	200
<i>Congrosoma</i>	80	<i>gibbifrons</i>	202
<i>Cottus</i>	383	<i>gryllus</i>	200
<i>Diapterus</i>	342	<i>heterurus</i>	201
<i>Hadropterus</i>	294	<i>hillianus</i>	200
<i>Hepsetia</i>	240	<i>holubi</i>	202
<i>Leuciscus</i>	118	<i>lamolifer</i>	200
<i>Moharra</i>	342	<i>lineatus</i>	202
<i>Salmo</i>	59	<i>lutkeni</i>	201
<i>Scarus</i>	434	<i>maculipinnis</i>	201
<i>Synodus</i>	164	<i>melanurus</i>	201
<i>Tachysurus</i>	151	<i>mento</i>	200
<i>Thyrsina</i>	248	<i>mesogaster</i>	200
Evermannia.....	445	<i>micropterus</i>	109
<i>longipinnis</i>	445	<i>nigricans</i>	201
<i>panamensis</i>	445	<i>noveboracensis</i>	201
<i>zosterura</i>	445	<i>nuttalli</i>	201
Evermannichthys.....	446	<i>obtusirostris</i>	200
<i>metzelniki</i>	447	<i>orbignianus</i>	200
<i>spongicola</i>	446, 447	<i>parra</i>	202
evides, <i>Alvordius</i>	284	<i>quadriremus</i>	201
<i>Clinus</i>	453	<i>roberti</i>	201
<i>Ericosma</i>	284	<i>rondeletii</i>	201
<i>Plectobranchus</i>	408	<i>rubescens</i>	201
evionthas, <i>Ophichthus</i>	87	<i>rufipinnis</i>	201
<i>Quassiremus</i>	87	<i>scylla</i>	201
Eviota.....	438	<i>speculiger</i>	201
<i>epiphanes</i>	438	<i>spilonotopterus</i>	202
<i>personata</i>	438	<i>spilopodus</i>	201
eviirus, <i>Eels</i>	328	<i>splendens</i>	200
evolans, <i>Exocetus</i>	200	<i>vermiculatus</i>	202
<i>Prionotus</i>	407	<i>vincigueriae</i>	200
<i>Trigla</i>	407	<i>volador</i>	200
Evolantia.....	199	<i>volitans</i>	200, 201
<i>microptera</i>	199	<i>volitens</i>	200
Evopites.....	324	<i>xenopterus</i>	202

List of scientific names—Continued

	Page	Page	
Exoglossum	140	fasciatus, Achirus.....	229
annulatum.....	140	Astyanax.....	97
dubium.....	140	Auchenopterus.....	457
lesurianum.....	140	Baionodon.....	417
maxillingua.....	140	Bodianus.....	325
mirabile.....	142	Bryttus.....	304
nigrescens.....	146	Caranx.....	272
spinicephalum.....	146	Catotomus.....	293
eroleatus, Labrus	424	Catostomus.....	109
Centrolabrus.....	424	Centronotus.....	467, 468
Exonantes	200	Crenichthys.....	456, 457
affinis.....	201	Ctenogobius.....	438
exsilis.....	200	Diplesion.....	203
rondeletti.....	200	Esox.....	203
ribescens.....	201	Exocetus.....	200
rufipinnis.....	201	Genyonemus.....	351
speculiger.....	201	Gymnachirus.....	229
vinciguerra.....	200	Gymnotus.....	101
expansum, Ostracion.....	497	Harpurus.....	362
exsiliens, Exocetus.....	200	Hemirhamphus.....	198
Exonantes.....	200	Larimus.....	344
extensus, Fundulus.....	179	Mytilophagus.....	410, 411
Lycodapus.....	478	Ovoides.....	299
Zygoletes.....	179	Pholis.....	468
Extrarius	139	Prinephales.....	145
tetranemus.....	139	Pogonias.....	352
exul, Apistos.....	370	Prionodes.....	319
eydouxii, Bagre.....	148	Scomber.....	279
Galeichthys.....	148	Schistes.....	369
fabbri, Blennius.....	460	Sicyas.....	400
Faber marinus fere quadratus.....	358	Syngnathus.....	242
faber, Chætodipterus.....	358	Trachichthys.....	279
Chaetodon.....	358	Trachynotus.....	278
fabricii, Campylodon.....	75	Xurel.....	272
Centroscyllium.....	21	Zonichthys.....	279
Gadus.....	209	fascicularis, Serranus.....	318, 319
Gunnellus.....	470, 471	fasciolaris, Idiacanthus.....	70
Liparis.....	400	fasciolaris, Catostomus.....	109
Lumpenus.....	471	Lythrurus.....	126
Spinax.....	21	Notropis umbratilis.....	126
fahronii, Apterurus.....	31	Sebastichthys.....	369
fajardo, Auchenopterus.....	457	Syphurus.....	231
falcata, Agosin.....	141	fasciolata, Coryphaena.....	264
Apococe.....	141	favosus, Bathygadus.....	203
Mycteroperca.....	315	Blennius.....	461
phenax, Mycteroperca.....	315	fecundus, Catostomus.....	108
Seriola.....	280	Chasmistes.....	108
falcatus, Brycon.....	98	Felichthys.....	147
Caranx.....	271	amblops.....	159
Gleridichthys.....	188	felicanus, Cyprinodon.....	181
Hemicaranx.....	271	Trifrcatus.....	181
Labrus.....	276	felinus, Pimelodus.....	153
Serranus.....	315	Serranus.....	315
Sparus.....	425	felis, Anarrhichthys.....	472
Trachinotus.....	270	Galeichthys.....	140
Zonichthys.....	280	Mustelus.....	14
falcidens, Atopichthys.....	54	Pimelodus.....	154
falciformis, Carcharias.....	16	Silurus.....	149
falcipinnis, Carcharias.....	16	fenestralis, Artedius.....	376
Elacatinus.....	281	Astrotyle.....	376
Falcula.....	142	fenestra, Chromis.....	416
chapala.....	142	fenestratum, Cichlaurus.....	416
Falculorius.....	142	ferox, Alepisaurus.....	171
chapala.....	142	Bathyophis.....	70
fallax, Caranx.....	272	Idiancanthus.....	70
Pomotis.....	300	Lepisosteus.....	38
Ianfarus, Naucrates.....	278	(Atractosteus).....	38
Fario.....	55	Macrodon.....	99
argenteus.....	55	Squalus.....	19
argyreus.....	55	Stomias.....	69
newberryi.....	58	ferruginea, Limanda.....	227
stellatus.....	56	Platessa.....	226, 227
tsuppitch.....	57	ferrugineus, Characodon.....	183
Fario, Salmo.....	55	festae, Piahuicina.....	99
fasciatus, Hippocampus.....	224	feuille, Polyodon.....	36
fasciatus, Cichla.....	208	fibulatus, Spinicophalus.....	145
Clupea.....	42	Fierasfer.....	101, 486
Ganibus.....	189	acutus.....	486
Molinessa.....	192	affinis.....	486
Plagiusa.....	231	arenicola.....	486
Pocilloopsis.....	189	borealis.....	471
Torrentaria.....	293	dubius.....	486
Trigla.....	409	fierasfer, Lycodapus.....	478
fuscatum, Characidium.....	100	filamentosa, Dicrolene.....	483
Pristipoma.....	335		

List of scientific names—Continued

	Page		Page
filamentosus, <i>Argyriosus</i>	275	flaviventer, <i>Eupomacentrus</i>	414
<i>Breviceps</i>	147	<i>Pomacentrus</i>	414
<i>Hemiramphus</i>	199	flaviventris, <i>Callidulus</i>	320
<i>Icelinus</i>	375, 376	<i>Dules</i>	320
<i>Monacanthus</i>	494	flavomaculatus, <i>Halocentrus</i>	310
<i>Tarandichthys</i>	376	flavolimbatus, <i>Epinephelus</i>	311
filicauda, <i>Coryphaenoides</i>	207	flavolineatum, <i>Hemulon</i>	330
filifer, <i>Dactylostomias</i>	69, 70	flavolineatus, <i>Diabasis</i>	330
filifera, <i>Anchovia</i>	50	<i>Mullus</i>	344
<i>Anchoviella</i>	50	<i>Sargus</i>	333
<i>Chalinura</i>	204	flavo-lineatus, <i>Pimelepterus</i>	339
fimbra, <i>Anoplopoma</i>	373	flavopurpurea, <i>Perca</i>	310
<i>Gadus</i>	373	flavoscripta, <i>Echidna</i>	94
fimbriata, <i>Cyclopsetta</i>	220	flavoscriptus, <i>Gymnothorax</i>	91
<i>Loricaria</i>	162	flavovittatus, <i>Upeneus</i>	343
<i>Raiia</i>	32	flavulus, <i>Esox</i>	170
<i>Squatina</i>	22	flavus, <i>Turdus</i>	423
fimbriatus, <i>Blennius</i>	473	<i>Noturus</i>	155
<i>Hemirhombus</i>	220	flleurieu, <i>Ostorbinius</i>	295
<i>Icelinus</i>	376	flexularis, <i>Lepomis</i>	298
<i>Paraliparis</i>	404	flexiosus, <i>Catostomus</i>	105
<i>Serranus</i>	311	flore, <i>Neoliparis</i>	399
Fimbriotorpedo.....	27	florealis, <i>Platyglossus</i>	420
fimisquamis, <i>Bogoslovicus</i>	204	floridae, <i>Branchiostoma</i>	7
<i>Macrurus</i>	204	<i>Jordanelle</i>	180
fischeri, <i>Chatostomus</i>	161	<i>Siphonotoma</i>	241
<i>Chatostomus</i>	161	<i>Syngnathus</i>	241
<i>Solea</i>	230	floridana, <i>Cichla</i>	297
<i>Tetragonopterus</i>	97	<i>Huro</i>	297
Fissala.....	408	floridanus, <i>Phycis</i>	213
<i>alata</i>	408	<i>Urophycis</i>	213
fissuratus, <i>Neoliparis</i>	399	floridensis, <i>Calliturus</i>	303
Fistularia.....	245	<i>Fundulus</i>	170
<i>cominnersonii</i>	245	floridichthys.....	180
<i>corneta</i>	245	<i>carpio</i>	180
<i>depressa</i>	245	<i>floripinnis, Haplochilus</i>	177
<i>immaculata</i>	245	<i>fluviatilis gibbosa, Perca</i>	302
<i>neo-boraccensis</i>	245	<i>Hudsonius</i>	132
<i>petimba</i>	245	<i>Leptophilypnus</i>	430
<i>serrata</i>	245	<i>Sargosomus</i>	412
<i>tabacaria</i>	245	<i>Tetronodon</i>	499
<i>villosa</i>	245	Fodiator.....	199
Fistulariidae.....	245	<i>acutus</i>	199
fistularis, <i>Flagellaria</i>	245	fodiator, <i>Tylosurus</i>	196
fistulata, <i>Syngnathus</i>	241	fontens, <i>Salmo</i>	163, 172
fistulatus, <i>Syngnathus</i>	241	folium, <i>Polyodon</i>	36
flabellare, <i>cumberlandicum</i> , <i>Catonotus</i>	293	fonsecaensis, <i>Achirus</i>	230
<i>lineolatum</i> , <i>Catonotus</i>	293	<i>Solea</i>	230
flabellaris, <i>Catonotus</i>	293, 383, 385	Fonticola.....	178
<i>Etheostoma</i>	293	fonticola, <i>Alvarius</i>	295
flabellata, <i>Etheostoma</i>	293	<i>Fundulus</i>	175
<i>flabellicauda</i> , <i>Rivulus</i>	180	<i>Fundulus heteroclitus</i>	175
Flagellaria.....	245	fontinalis, <i>Etheostoma</i>	293
<i>fistularis</i>	245	<i>Salmo</i>	60
flagellum, <i>Goniobatis</i>	30	<i>Salvelinus</i>	60
<i>Raiia</i>	30	forbesi, <i>Cyprinella</i>	131
<i>Saccopharynx</i>	95	Forbesella.....	194
Flammeo.....	236	<i>agenzzii</i>	194
<i>mariannus</i>	236	<i>papillifera</i>	194
flammeus, <i>Phoxinus</i>	121	Forcipatus, <i>Balistes</i>	492
flava, <i>Congerimurena</i>	80	Forcipiger.....	359
<i>Hildebrandia</i>	80	<i>flavissimus</i>	359
<i>Loricaria</i>	100	<i>longirostris</i>	359
flavescens, <i>Mesoprion</i>	325	forficata, <i>Elanura</i>	380
<i>Morone</i>	281	forlonensis, <i>Cyprinella</i>	131
<i>Perca</i>	281	<i>Nototroplus</i>	131
<i>Prionodes</i>	319	formosa, <i>Algansea</i>	122
<i>Scarus</i>	433	<i>Ergaula</i>	129
<i>Serranus</i>	319	<i>Heterandria</i>	188
<i>Sparisoma</i>	433	<i>Limia</i>	192
flavesny, <i>Brosmerus</i>	213, 214	<i>Mollieuscia</i>	192
flavicauda, <i>Hyporthodus</i>	311	<i>Moniana</i>	120, 131
flavidus, <i>Apodichthys</i>	466, 467	<i>Perca</i>	318
<i>Aulorhynchus</i>	240	formosulum, <i>Campostoma</i>	147
<i>Sebastodes</i>	364	formosum, <i>Hæmulon</i>	330
<i>Sebastosomus</i>	364	formosus, <i>Albumnus</i>	120
flaviguttatum, <i>Lythryalon</i>	331	<i>Anthias</i>	330
flaviguttatus, <i>Haemulon</i>	331	<i>Calliturus</i>	299
flavilatus, <i>Eupomacentrus</i>	414	<i>Cottus</i>	386
<i>Pomacentrus</i>	414	<i>Diplectrum</i>	318
<i>Hybognathus</i>	144	<i>Holacanthus</i>	361
<i>Pristigaster</i>	44	<i>Siphateles</i>	122
flavirostris, <i>Syngnathus</i>	241	<i>Tetrodon</i>	500
<i>flavissimus, Forcipiger</i>	359		

List of scientific names—Continued

	Page		Page
formulosum, <i>Campostoma</i>	147	Fundulus bermudæ.....	175
forskæll, <i>Albula</i>	39	chrysotus.....	177
Glossodus.....	40	cingulatus.....	177
forsteri, <i>Albula</i>	40	confuentus.....	177
Scomberesox.....	197	cubensis.....	174
forsterianus, <i>Catostomus</i>	105, 106	dugesii.....	182
fosser, <i>Ichthyomyzon</i>	9	extensus.....	170
fosteri, <i>Priapichthys</i>	189	floridensis.....	176
4-maculatus, <i>Diodon</i>	501	fonticola.....	175
fragilis, <i>Citharichthys</i>	218	fuscus.....	174
Macrourus.....	206	grandis.....	176
Macrurus.....	206	guatemalensis.....	174
francisci, <i>Cestracion</i>	10	heteroclitus.....	175
Gyroleurodus.....	10	badius.....	176
franklini, <i>Cottus</i>	385	bermudæ.....	175
Pleuronectes.....	227	fonticola.....	175
fraterculus, <i>Mylocheilus</i>	112	grandis.....	176
freniebundum, <i>Hæmulon</i>	329	macrolepidotus.....	176
freminvillii, <i>Aetobatus</i>	31	kansæ.....	170
Mylobatris.....	31	labialis.....	174
frenata, <i>Loricaria</i>	102	limi.....	178
frenatus, <i>Brachystius</i>	400	majalis.....	175, 176
Iridio.....	427	meeki.....	178
Odontopyxis.....	305	menopis.....	178
Sarritor.....	395	mudfish.....	175
Xantocles.....	375	nisorius.....	175
Zanolepis.....	375	oaxacae.....	174
fratensis, <i>Hybopsis</i>	120	ocellaris.....	176
fricksia, <i>Etheostoma</i>	291	pachycephalus.....	174
Oligocephalus.....	291	pallidus.....	176
friedrichsthalii, <i>Heron</i>	418	parvipinnis.....	176
Parapetenia.....	418	<i>brevis</i>	176
frigida, <i>Moniana</i>	131	punctatus.....	174
frigidus, <i>Lycodes</i>	474	rathbuni.....	175, 176
freebelli, <i>Amphilophus</i>	422	rhizophorus.....	175
frondosum, <i>Sparsisoma</i>	433	scartes.....	177
frondosus, <i>Scarus</i>	433	sciadicus.....	177
frontale, <i>Clethasoma</i>	417	seminolis.....	178
frontalis, <i>Caranx</i>	273	similis.....	177
Cryptopterenchelys.....	85	tenellus.....	177
Gastropsetta.....	225	vinctus.....	176
Leuciscus.....	128	viridescens.....	176
Ophichthys (<i>Cryptopterus</i>).....	85	xenicus.....	180
Prionotus.....	408	zebra.....	176
Theraps.....	417	zebrinus.....	179
fronti unicorni, <i>Lophius</i>	507	zonatus.....	178
fronto, <i>Carcharhinus</i>	17	Fundulus, <i>Alceocephalus</i>	52
Nemichthys.....	83	funebris, <i>Gobiesox</i>	489
frost, <i>Gadus</i>	210	Gymnothorax.....	92
fucensis, <i>Liparis</i>	400	Noturus.....	156
Pollachius <i>chalcopterus</i>	209	Rabida.....	156
Theragra.....	209	Furcaria.....	412
fucorum, <i>Apodichthys</i>	467	<i>atrilobata</i>	412
Bleennius.....	461	<i>bermudæ</i>	413
Cypsolurus.....	202	<i>cyanea</i>	412
Exocetus.....	202	<i>multilineata</i>	412
Xeropetra.....	407	<i>puncta</i>	412
fulgens, <i>Corvina</i>	345	(<i>Furcaria</i>) <i>atrilobata</i> , <i>Chromis</i>	412
Mylopristis.....	234	<i>furcatus</i> , <i>Cypselurus</i>	201
Placanthus.....	323	<i>letalus</i>	151, 152
Tetragonopterus.....	96	<i>Exocetus</i>	201
fulgida, <i>Meda</i>	147	<i>Phanerodon</i>	411
fulginosus, <i>Balistes</i>	401	<i>Pimelodus</i>	152
Cyclichthys.....	502	Furcella.....	476
Diodon.....	502	<i>furcellus</i> , <i>Careproctus</i>	402
Synbranchus.....	70	<i>furcidens</i> , <i>Characodon</i>	183
fuliginosus, <i>Holconotus</i>	411	<i>furcifer</i> , <i>Paranthias</i>	320
fulva, <i>Labrus chogoset</i>	424	<i>Pimelodus</i>	152
fulvescens, <i>Acipenser</i>	33, 35	<i>Serranus</i>	320
fulvo-maculatus, <i>Labrus</i>	335	<i>furciger</i> , <i>Icelus</i>	378
fulvum, <i>Glynglymostomus</i>	12	<i>Furcimanus</i>	476
fulvus, <i>Cephalopholis</i>	309	<i>diapterus</i>	476
<i>Labrus</i>	309	<i>furcifrinus</i> , <i>Pontinus</i>	372
<i>Physcilius</i>	212	<i>furcata</i> , <i>Pachhypops</i>	349
<i>punctatus</i> , <i>Cephalopholis</i>	309	<i>Perca</i>	349
<i>ruber</i> , <i>Cephalopholis</i>	309	Furlosa, <i>Rabida</i>	150
fumeus, <i>Notropis</i>	125	furlous, <i>Noturus</i>	156
funduloides, <i>Clinostomus</i>	121	furnieri, <i>Micropogon</i>	349
Zygonectes.....	176	<i>Umbrina</i>	349
Fundulus.....	175, 180	furthi, <i>Arius</i>	151
<i>adinala</i>	179	<i>Chelichthys</i>	499
<i>albolineatus</i>	177	furtheri, <i>Arius</i>	151
<i>antillarum</i>	175	<i>Caranx</i>	271
<i>aurous</i>	177	<i>Corvina</i> (<i>Homopriion</i>).....	347
<i>balboæ</i>	179		

List of scientific names—Continued

	Page		Page
<i>furthii</i> , <i>Hemicaranx</i>	271	<i>Gadus punctatus</i>	213
<i>Pristipoma</i>	333	<i>purpureus</i>	209
<i>Tetradon</i>	499	<i>pygmæus</i>	210
<i>Zestis</i>	347	<i>ruber</i>	214
<i>furva</i> , <i>Perca</i>	317	<i>rupestris</i>	210
<i>fusca</i> , <i>Echeneis</i>	449	<i>ssida</i>	209
<i>Hydrargyra</i>	174	<i>tau</i>	486, 487
<i>Sciaena</i>	342	<i>tenuis</i>	213
<i>Tautoga</i>	424	<i>tomecod</i>	210
<i>fuscescens</i> , <i>Icelinus</i>	376	<i>tomeodus fuscus</i>	210
<i>fuscomaculata</i> , <i>Echidna</i>	94	<i>tomeodus luteus</i>	210
<i>fusco-maculatus</i> , <i>Chromis</i>	419	<i>mixtus</i>	210
<i>fusculus</i> , <i>Centropristes</i>	319	<i>pruinosus</i>	210
<i>Prionodes</i>	159	<i>torsk</i>	214
<i>fuseum</i> , <i>Pygidium</i>	362	<i>vertagus</i>	210
<i>fuscus</i> , <i>Acronurus</i>	362	<i>virens</i>	209
<i>Bathygobius</i>	440	<i>wachna</i>	210
<i>Bythites</i>	480	<i>Gaidropsaridae</i>	209
<i>Culius</i>	436	<i>Gaidropsaruss</i>	208
<i>Eupomacentrus</i>	413	<i>ensis</i>	208
<i>Fundulus</i>	174	<i>mustelurus</i>	208
<i>Gadus tomcodus</i>	210	<i>septentrionalis</i>	208
<i>Gobius</i>	440	<i>gaimardiana</i> , <i>Querimanna</i>	254
<i>Hemirhambus</i>	219	<i>gaimardianus</i> , <i>Mugil</i>	254
<i>Kyphosus</i>	339	<i>gaiderneri beardslei</i> , <i>Salmo</i>	57
<i>Pomacentrus</i>	413	<i>crescentis</i> , <i>Salmo</i>	57
<i>Psenes</i>	268	<i>gilberti</i> , <i>Salmo</i>	58
<i>Serranus</i>	315	<i>shasta</i> , <i>Salmo</i>	58
<i>Silurus</i>	153	<i>stonei</i> , <i>Salmo</i>	58
<i>Syngnathus</i>	242	<i>gaiderneri</i> , <i>Salmo</i>	58
<i>Syrichtes</i>	242	<i>glaucula</i> , <i>Erogala</i>	120
<i>Trachinotus</i>	276	<i>glaucus</i> , <i>Hypsilepis</i>	129
<i>Xyster</i>	339	<i>Galaate</i>	493
<i>fusiforme</i> , <i>Boleosoma</i>	294	<i>Galasaceus</i>	175
<i>fusiformis</i> , <i>Boleichthys</i>	204	<i>galeata</i> , <i>Belone</i>	177
<i>tylæ</i> , <i>Raja</i>	25	<i>galeatus</i> , <i>Gymnacanthus</i>	389
<i>gabonensis</i> , <i>Vomer</i>	275	<i>Gymnocaanthus</i>	389
<i>gabunensis</i> , <i>Ischnomembras</i>	217	<i>Silurus</i>	159
<i>Gadella</i>	211	<i>Tylosurus</i>	197
<i>gracilis</i>	211	<i>Galei</i>	11
<i>maraldi</i>	211	<i>Galeichthys</i>	149
<i>Gadidae</i>	209	<i>assimilis</i>	150
<i>Gadomus</i>	202	<i>azurceus</i>	150
<i>arcutus</i>	202	<i>boehmiensis</i>	148
<i>longithilis</i>	202	<i>carulescens</i>	150
<i>gadovii</i> , <i>Cichlosoma</i>	417	<i>dasycephalus</i>	150
<i>Theraps</i>	417	<i>cydouxit</i>	148
<i>Gadus</i>	210	<i>feliceps</i>	149
<i>agilinus</i>	210, 211	<i>felis</i>	148, 149
<i>agilis</i>	209	<i>gilberti</i>	150
<i>albidus</i>	214	<i>guatemalensis</i>	150
<i>arenosus</i>	210	<i>guentheri</i>	149
<i>auratus</i>	210	<i>jordani</i>	150
<i>brosme</i>	213	<i>letigiosus</i>	149
<i>callarias</i>	210	<i>longicephalus</i>	150
<i>carbonarius</i>	209	<i>peruvianus</i>	149
<i>chalocrammus</i>	209	<i>sewanianus</i>	150
<i>cimbrius</i>	208	<i>surinamensis</i>	150
<i>colinus</i>	209	<i>xenauchen</i>	150
<i>compressus</i>	212	<i>Galeocerdo</i>	14
<i>fabricii</i>	209	<i>arcticus</i>	14
<i>flimbriis</i>	373	<i>tigrinus</i>	14
<i>frost</i>	210	<i>galeoides</i> , <i>Otophidium</i>	485
<i>gracilis</i>	210, 211	<i>Galeorhinus</i>	14
<i>heteroglossus</i>	210	<i>zyopterus</i>	14
<i>leucostriis</i>	152, 212	<i>galerita</i> , <i>Bleennius</i>	400
<i>lepidion</i>	211	<i>Galeus</i>	13, 17
<i>longipes</i>	213	<i>asterias</i>	13
<i>lotia</i>	212	<i>catulus</i>	13
<i>macrocephalus</i>	210	<i>glauces</i>	17
<i>maculosus</i>	212	<i>maculatus</i>	14
<i>marilli</i>	211	<i>mechistostomus</i>	12
<i>merluccius</i>	214	<i>instinctus</i>	13
<i>merlus</i>	214	<i>vulpecula</i>	13
<i>molva</i>	212	<i>galeus</i> , <i>Serranus</i>	312
<i>morrhua</i>	210	<i>Squalus</i>	13, 14
<i>mustela</i>	208	<i>Galichthys</i>	274
<i>narus</i>	210	<i>gallichthys</i> , <i>Carangooides</i>	274
<i>nawaga</i>	209, 210	<i>gallinula</i> , <i>Monacanthus</i>	404
<i>ogac</i>	210	<i>Gallus</i>	274
<i>periscopius</i>	209	<i>gallus</i> , <i>Zeus</i>	274, 275
<i>pollachius</i>	209	<i>galtié</i> , <i>Squalius</i>	118
<i>polymorphus</i>	210		
<i>proximus</i>	210		

List of scientific names—Continued

	Page		Page
Gambusia.....	184, 185	Gasterosteus aculeatus.....	239
annectans.....	177	apeletes.....	239
arlingtonia.....	188	argentissimus.....	238
bonita.....	188	atkinsi.....	238
cana.....	188	bailoul.....	237
cascajalensis.....	187	biaculeatus.....	237
caymanensis.....	186	bispinosus.....	238
darienensis.....	189	blanchardi.....	239
dominicensis.....	189	breviceps.....	239
dovii.....	186	cannatus.....	281
episcopi.....	187	carolinus.....	276, 277
fusciata.....	189	cataphractus.....	237
gracilior.....	186	concininus.....	239
gracilis.....	185	cuvieri.....	237
heckeli.....	185	dehayi.....	239
holbrookii.....	184	dimidiatus.....	237
humilis.....	185	duktor.....	278
infans.....	189	elegans.....	238
latipunctata.....	187	gladiunculus.....	238
manni.....	186	globiceps.....	238
menieri.....	186	gymnurus.....	237
metapleura.....	185	inconstans.....	238
modesta.....	185, 192	inopinatus.....	238
myersi.....	185	insculptus.....	237
nicaraguensis.....	184, 186	islandicus.....	238
nobilis.....	185	levius.....	239
oligosticta.....	186	leucus.....	237
panuco.....	186	loricatus.....	237
parismima.....	187	lotharingus.....	239
patruelis.....	185	mainensis.....	239
picturata.....	185	marinus.....	239
plumbea.....	192	microcephalus.....	237
punctata.....	184, 185	micropus.....	238
punctulata.....	185	millipunctatus.....	239
puncticulata.....	185	nebulosus.....	239
regani.....	184, 186	neustrianus.....	238
rhabdophorus.....	186	noveboracensis.....	237
senilis.....	185	obolarius.....	237
speciosa.....	185	occidentalis.....	238, 239
terrabensis.....	186	plebeius.....	237
tridentiger.....	188	ponticus.....	237
turrubarensis.....	189	pugniti.....	238
umbratilis.....	190	brachypoda.....	239
vittata.....	186	pygmoeus.....	238
wrayi.....	186	quadracus.....	239
yucatana.....	180	saltatrix.....	280
Gambusinus.....	175, 179	santa-annae.....	238
Ganunaraia.....	25	semilarvatus.....	237
gardeniana, Hiatula.....	424	semiloricatus.....	237
gardenii, Contronotus.....	281	serratus.....	237
Sternopyx.....	206	spinachia.....	239
gardoneus, Leuciscus.....	115	spinulosus.....	237
Notemigonus.....	115	terracanteus.....	237
garmani, Cetengraulis.....	51	trachurua.....	237
Characodon.....	183	wheatlandi.....	238
Girardinus.....	189	williamsoni.....	238
Gobius.....	439	microcephalus.....	237
Lepomis.....	300	Gastrea.....	239
Notropis.....	131	Gastrobranchus.....	8
Paraliparis.....	405	cæcus.....	8
Rhinogobius.....	430	domboi.....	8
Stolephorus.....	51	Gastropsetta.....	225
Garmannia.....	445	frontalis.....	225
hemigymna.....	445	Gastrostomus.....	95
paradoxa.....	445	baIRDII.....	95
rubra.....	445	pacificus.....	95
spongicola.....	446	gausa, Serranus.....	312
garnoti, Iridio.....	427	gauza, Epinephelus.....	310
Julis.....	427	Labrus.....	310
Garrupa.....	312	gavialis, Lepisosteus.....	36
nigrita.....	312	guyi, Merlus.....	214
Gasteracanthus.....	237	geddesi, Characodon.....	182
cataphractus.....	237	HORICHThYs.....	423
Gasteropelecidae.....	90	Tomocichla.....	423
Gasteropelecanus.....	90	gelatinosum, Melanostigma.....	478
humboldtii.....	169	gelatinosus, Aphyonus.....	484
maculatus.....	99	Careproctus.....	402
sterniflu.....	99	Cyclopterus.....	402
gasteropelecanus, Trachinus.....	357	gelida, Anniomyrypta.....	280
Gasterosteidae.....	238	Macrhybopsis.....	139
Gasterosteidae.....	237	gelidus, Gobio.....	138, 139
Gasterosteidae.....	238	genimatus, Blennius.....	461, 462
Gasterosteus.....	237	Hypoleurochilus.....	462
aculeatus.....	237		

List of scientific names—Continued

	Page		Page
gemina, <i>Hypoplectrus</i> .	321	Gibbonsia metzi	458
<i>Hypoplectrus unicolor</i> .	321	gibbosa cinerea, <i>Perca marina</i>	329
geminifer, <i>Astronesthes</i> .	71	Gila	119
<i>Lampanyctus</i> .	167	Montana	131
<i>Leucichthys</i> .	63	Perca	302, 329
Gempylidae	260	<i>fluviatilis</i>	302
Gempylus	261	Tigoma	119
<i>cobuber</i>	261	glbbosus, <i>Aphredoderus</i>	215
<i>ophidianus</i>	262	Catostomus	109
<i>prometheus</i>	261	Cyprinodon	181
<i>serpens</i>	261, 262	Eupomotis	302
Generosus, <i>Catostomus</i> (<i>Acomus</i>)	104	Pomotis	301
<i>Notolepidomyzon</i>	104	gibbsii, <i>Homilepidotus</i>	381
genizara, <i>Kabirubia</i>	420	Salmo	57
gentilis, <i>Bleennius</i>	462	gibbus, <i>Hypsilepis cornutus</i>	128
<i>Hypsoblennius</i>	462	Liparis	400
Genyatremus	335	Lophius	505
<i>luteus</i>	335	Scarus	433
Genyonemus	349	gigas <i>Ephippus</i>	358
<i>fasciatus</i>	351	<i>Hippoglossus</i>	221
<i>lineatus</i>	349	Perca	310
Genyorge	324	Sciæna	352
Genypterus omostigma	485	Seriola	279
Genytremus	332	Stereolepis	308
<i>interruptus</i>	332	giganteus, <i>Dololepis</i>	472
geometricus, <i>Anchisomus</i>	499	Gila	114
<i>Diodon</i>	502	<i>affinis</i>	114
<i>lineatus</i> , <i>Diodon</i>	502	<i>ardesiac</i>	118
<i>Tetronodon</i>	499	<i>concephala</i>	113
<i>Zeus</i>	275	<i>conspersa</i>	120
Geophagus	424	<i>elegans</i>	114
<i>crassilabris</i>	424	<i>emoryi</i>	114
<i>cupido</i>	424	<i>estor</i>	121
<i>demon</i>	424	<i>gibbosa</i>	119
<i>magasoma</i>	424	<i>gracilis</i>	114
(<i>Satanoperca</i>) <i>crassilabris</i>	424	<i>grahami</i>	114
georgianus, <i>Exocetus</i>	200	<i>grandis</i>	114
Gephyrocharax	96	<i>gula</i>	120
<i>atricaudatu</i>	96	<i>microlepido</i>	112
<i>atricaudatus</i>	96	<i>minace</i>	114
<i>choconensis</i>	96	<i>nacre</i>	114
<i>intermedius</i>	96	<i>nigra</i>	119
gerania, <i>Belone</i>	196	<i>pulchella</i>	119, 120
germanus, <i>Notropis</i>	136	<i>robusta</i>	114
Germe	260	<i>seminuda</i>	114
<i>alalunga</i>	260	gilla, <i>Catostomus</i>	106
<i>gerino</i>	260	gilberti, <i>Ariosoma</i>	80
germo, <i>Germo</i>	260	<i>Careproctus</i>	402
<i>Scomber</i>	260	<i>Ctenograulis</i>	51
germon, <i>Scomber</i>	280	Chlopsis	82
Gerres	341	Citharichthys	219
<i>apiron</i>	342	Congrellus	80
<i>aureolus</i>	342	Galeichthys	150
<i>axillaris</i>	342	Hubbesia	248
<i>brasilianus</i>	342	Hybopsis	134
<i>brevimanus</i>	343	Hypsoblemnius	462
<i>brevirostris</i>	342	Ilyonus	445
<i>cinerereus</i>	341	Imostoma	286
<i>embryx</i>	342	Isesthas	462
<i>gula</i>	340, 341	Lepidogobius	445
<i>havana</i>	342	Notropis	134
<i>jonesii</i>	341	Noturus	156
<i>lineatus</i>	342	Oncocottus	389
<i>mexicanus</i>	343	<i>hexacanthus</i>	389
<i>olifistostoma</i>	342	Podothecus	305
<i>patao</i>	342	Pteropodus	369
<i>peruvianus</i>	342	Rabida	158
<i>plumieri</i>	342, 343	Salmo	68
<i>rhombeus</i>	342	<i>gaideri</i>	58
<i>simillimus</i>	342	Sebastodes	309
<i>squamipinnis</i>	342	Stolephorus	51
<i>vaigiensis</i>	341	Ulocentra	286
<i>zebra</i>	342	Gilbertida	392
Gerrido	340	<i>sigulatus</i>	393
Gerris	341	Gilbertina	392
ghini, <i>Orthragoriscus</i>	503	<i>sigulatus</i>	392, 393
gibba, <i>Ilistrio</i>	505	Gillellus	455
gibber, <i>Salmo</i>	54	<i>arenicola</i>	455
gibbiceps, <i>Heros</i>	417	<i>ornatus</i>	455
<i>Theraps</i>	417	<i>semicinctus</i>	455
gibblifrons, <i>Cypselurus</i>	202	gilli, <i>Avocettina</i>	83
<i>Exocattus</i>	202	Clinus	460
gibbonsi, <i>Holconotus</i>	412	Hyperhamphus	198
gibbonsi	458	Labichthys	83
<i>elegans</i>	458	Leuciscus	118
<i>montereyensis</i>	458		

List of scientific names—Continued

	Page		Page
<i>gilli</i> , <i>Malacoctenus</i> .	460	<i>glaucæ</i> , <i>Ciliata</i> .	208
Pholts.	468	<i>Læchia</i> .	276
<i>Pleuronectes</i> .	228	<i>Prionace</i> .	17
<i>Synchirus</i> .	393	<i>glaucofrænum</i> , <i>Coryphopterus</i> .	439
<i>Gillia</i> .	444	<i>Rhinogobius</i> .	439
<i>gilliæ</i> , <i>Julis</i> .	420	<i>glaukopareiæus</i> , <i>Acanthurus</i> .	363
<i>Gillias</i> .	458	<i>Glaucostegus</i> .	23
<i>jordani</i> .	458	<i>glaucostictus</i> , <i>Rhinobatus</i> .	23
<i>Gillichthys</i> .	444	<i>glaucostigma</i> , <i>Rhinobatos</i> .	24
<i>detrusus</i> .	444	<i>Rhinobatus</i> .	24
<i>guayanensis</i> .	444	<i>glaucæ</i> , <i>Sitzistedion</i> .	282
<i>marmoratus</i> .	444	<i>glaucus</i> , <i>Chirodon</i> .	276
<i>y-cauda</i> .	444	<i>Emmelas</i> .	364
<i>gillii</i> , <i>Bassogigas</i> .	482	<i>Galeus</i> .	17
<i>Cetomimus</i> .	172	<i>Sebastes</i> .	364
<i>Eupomacentrus</i> .	414	<i>Squalus</i> .	17
<i>Lepomis</i> .	303	<i>Trachynotus</i> .	276
<i>Lipogramys</i> .	75	<i>globiceps</i> , <i>Agonostoma</i> .	255
<i>Neobrythites</i> .	481	<i>Blennicottus</i> .	391
<i>Pomacentrus</i> .	414	<i>bryosus</i> , <i>Blennicottus</i> .	391
<i>Sebastodes</i> .	368	<i>Gasterosteus</i> .	238
<i>Sebastomus</i> .	368	<i>Joturus</i> .	255
<i>Stephanoberyx</i> .	232	<i>Oligocottus</i> .	391
<i>Xiphophorus</i> .	192	<i>globosa</i> , <i>Lyosphæra</i> .	503
<i>Xystropeltes</i> .	301, 302	<i>globosum</i> , <i>Astatheros</i> .	421
<i>Ginglymostomiæ</i> .	36	<i>Cichlasoma</i> .	421
<i>Ginglymostoma</i> .	12	<i>gloriosus</i> , <i>Brytus</i> .	304
<i>caboverdianus</i> .	12	<i>Enneacanthus</i> .	304
<i>cirratum</i> .	12	<i>Glossamia</i> , <i>pandionis</i> .	206
<i>fulvum</i> .	12	<i>Glossichthys</i> .	230
<i>Ginglymostomidae</i> .	12	<i>Glossodon</i> .	40
<i>giorna</i> , <i>Cephalopterus</i> .	32	<i>harengoides</i> .	40
<i>girardi</i> , <i>Acipenser</i> .	34	<i>hetorurus</i> .	40
<i>Caranx</i> .	272	<i>glossodonta</i> , <i>Argentina</i> .	40
<i>Pseudojulis</i> .	428	<i>Glossodus</i> .	39
<i>Girardinichthys</i> .	182	<i>forskali</i> .	40
<i>innominatus</i> .	182	<i>Glossoplites</i> .	302
<i>limnurgus</i> .	183	<i>gloveri</i> , <i>Salmo</i> .	56
<i>Girardinus</i> .	189	<i>glutinosa</i> , <i>Myxine</i> .	8
<i>creolus</i> .	189	<i>Glyphtodon</i> <i>chrysurus</i> .	415
<i>denticulatus</i> .	189	<i>rudis</i> .	415
<i>garnmani</i> .	189	<i>saxatilis</i> .	415
<i>guppii</i> .	191	<i>taurus</i> .	415
<i>metallicus</i> .	189	<i>troeschellii</i> .	415
<i>pleurospilus</i> .	190	<i>Glyphtodon</i> .	414
<i>sonoriensis</i> .	189	<i>chrysurus</i> .	415
<i>uninotatus</i> .	188	<i>moucharra</i> .	415
<i>versicolor</i> .	191	<i>rubicundus</i> .	415
<i>Girella</i> .	338	<i>Glyptocephalus</i> .	228
<i>dorsoinacula</i> .	339	<i>acadianus</i> .	228
<i>nigricans</i> .	339	<i>cynoglossus</i> .	228
<i>punctata</i> .	338	<i>pacificus</i> .	228
<i>simplicidens</i> .	339	<i>zachirus</i> .	228
<i>Girellidae</i> .	338	<i>Gnathanodon</i> .	273
<i>Giton</i> .	101	<i>Gnathobolus</i> .	45
<i>glaber</i> , <i>Cottus</i> .	487	<i>Gnathocentrum</i> .	302
<i>Cylopterichthys</i> .	398	<i>Gnathanodon</i> .	273
<i>Pleuronectes fuscus subrotundus</i> .	230	<i>speciosus</i> .	273
<i>glabra</i> , <i>Acipenser</i> .	34	<i>gnathodus</i> , <i>Pseudoscarus</i> .	434
<i>Liopsetta</i> .	227	<i>Scarus</i> .	434
<i>Platessa</i> .	227	<i>Gnatholepis</i> .	442
<i>Pleuronectes satis parva et</i> .	217	<i>thompsoni</i> .	443
<i>glacialis</i> , <i>Cottus</i> .	387	<i>Gnathophis</i> .	80
<i>Liopsetta</i> .	227	<i>guppii</i> .	80
<i>Pleuronectes</i> .	227	<i>heterogenanthus</i> .	80
<i>Scopeus</i> .	170	<i>proriger</i> .	80
<i>Squalus</i> .	22	<i>Gnathoprops</i> .	453
<i>glacoides</i> , <i>Trachynotus</i> .	276	<i>aurifrons</i> .	453
<i>gladiifer</i> , <i>Istrophorus</i> .	263	<i>cuvieri</i> .	453
<i>Gladiunculus</i> .	238	<i>macrops</i> .	453
<i>bispinosus</i> .	238	<i>maxillosa</i> .	453
<i>gladiunculus</i> , <i>Gasterosteus</i> .	238	<i>mystacina</i> .	453
<i>gladius</i> , <i>Histiophorus</i> .	263	<i>mystacinus</i> .	453
<i>Notistium</i> .	263	<i>rhomalea</i> .	453
<i>Scomer</i> .	263	<i>scops</i> .	453
<i>Tylosurus</i> .	196	<i>snyderi</i> .	453
<i>Xiphæs</i> .	264	<i>Goblescodium</i> .	488
<i>Glanidostomi</i> .	33	<i>Gobiasox</i> .	488
<i>Glaridichthys</i> .	188	<i>adustus</i> .	489
<i>falcatus</i> .	188	<i>androsensis</i> .	489
<i>torralbasi</i> .	188	<i>cephalus</i> .	488, 489
<i>uninotatus</i> .	188	<i>cerasinus</i> .	489
<i>Glaridodon</i> .	188	<i>costaricensis</i> .	489
<i>latidens</i> .	188, 189	<i>eigenmanni</i> .	490
		<i>eos</i> .	491

List of scientific names—Continued

	Page		Page
<i>Gobiesox</i> , <i>erythrops</i> .	490	<i>Gobius</i> , <i>claytonii</i> .	439
<i>funebris</i> .	489	<i>cristagalli</i> .	438
<i>gyrinus</i> .	489	<i>crista-galli</i> .	438
<i>laevis</i> .	490	<i>curaçao</i> .	439
<i>humeralis</i> .	490	<i>cyprioides</i> .	438
<i>macrophthalmus</i> .	489	<i>dalli</i> .	442
<i>marmoratus</i> .	488	<i>dolichocephalus</i> .	442
<i>muscarum</i> .	490	<i>eigenmanni</i> .	430
<i>nigripinnis</i> .	489	<i>emblematicus</i> .	443
<i>papillifer</i> .	488	<i>encaezomus</i> .	439
<i>piniger</i> .	488	<i>fucus</i> .	440
<i>rheosodon</i> .	490	<i>garmani</i> .	439
<i>rhodospilus</i> .	489	<i>gracilis</i> .	441
(<i>Rimicola</i>) <i>beryllinus</i> .	490	<i>gronovii</i> .	268
<i>rupestris</i> .	491	<i>gulosus</i> .	443
<i>sancti-martini</i> .	490	<i>hemigymnus</i> .	445
<i>strunmosus</i> .	489	<i>hipoliti</i> .	440
<i>tudes</i> .	489	<i>homocyanus</i> .	440
<i>virgatulus</i> .	489	<i>kraussi</i> .	441
<i>vittatus</i> .	489	<i>lacerus</i> .	440
<i>yuma</i> .	490	<i>lanceolatus</i> .	441
<i>zebra</i> .	491	<i>lepidus</i> .	444
<i>Gobiidae</i> .	438	<i>lineatus</i> .	440, 446
<i>Gobio</i> , <i>estivalis</i> .	138	<i>longicaudus</i> .	441
<i>cataractae</i> .	139	<i>lucretiae</i> .	443
<i>gelidus</i> .	138, 139	<i>lyricus</i> .	441
<i>plumbeus</i> .	136	<i>manglicola</i> .	439
<i>vernalis</i> .	137	<i>mapo</i> .	440
<i>gobio</i> , <i>Clinus</i> .	455, 456	<i>martinicus</i> .	442
<i>Cottus</i> .	383	<i>mexicanus</i> .	442
<i>Gobioclinus</i> .	456	<i>microdon</i> .	441
<i>Gobioclinus</i> .	455	<i>nebulopunctatus</i> .	440
<i>gobio</i> .	456	<i>nebulo-punctatus</i> .	440
<i>Gobioides</i> .	448	<i>newberryi</i> .	444
<i>barreto</i> .	448	<i>nicholsii</i> .	440
<i>broussonnetii</i> .	448	<i>niger</i> .	440
<i>gobioides</i> , <i>Cottus</i> .	385	<i>oblongus</i> .	448
<i>Hypsicometes</i> .	451	<i>oceanicus</i> .	441
<i>Gobioididae</i> .	448	<i>ocellaris</i> .	442
<i>Gobiomorus</i> .	435	<i>padangensis</i> .	440
<i>dormitor</i> .	435	<i>paradoxus</i> .	445
<i>lateralis</i> .	436	<i>parvus</i> .	439
<i>Gobionellus</i> .	440	<i>pisonis</i> .	436
<i>bayamonensis</i> .	441	<i>plumieri</i> .	447
<i>bastatus</i> .	440, 441	<i>poeyi</i> .	442
<i>lyricus</i> .	441	<i>punetillus</i> .	440
<i>microdon</i> .	441	<i>quadriporus</i> .	439
<i>oceanicus</i> .	441	<i>sandwicensis</i> .	440
<i>sagittula</i> .	441	<i>semifasciatus</i> .	441
<i>sinuargus</i> .	441	<i>seminudus</i> .	445
<i>strigatus</i> .	441	<i>shufeldti</i> .	439
<i>stigmatus</i> .	441	<i>smaragdus</i> .	440, 441
<i>gobionum</i> , <i>Campostoma</i> .	146	<i>soporator</i> .	440
<i>Gobiosoma</i> .	445	<i>stigmaturus</i> .	439
<i>alepidotum</i> .	446	<i>taenifica</i> .	442
<i>bosci</i> .	446	<i>thalassinus</i> .	444
<i>cauthoecum</i> .	446	<i>townsendi</i> .	444
<i>crescentiale</i> .	446	<i>translucens</i> .	440
<i>crescentalis</i> .	446	<i>viridi pallidus</i> .	446
<i>histrio</i> .	446	<i>wurdemani</i> .	441
<i>horsti</i> .	446	<i>zebra</i> .	441
<i>ios</i> .	445	<i>Goboidea</i> .	435
<i>longipinne</i> .	445	<i>godmani</i> , <i>Pimeledodus</i> .	157
<i>longum</i> .	446	<i>Rhamdia</i> .	157
<i>moestum</i> .	446	<i>godmanni</i> , <i>Cichlaurus</i> .	416
<i>multifasciatum</i> .	446	<i>Heros</i> .	416
<i>zosterurum</i> .	445	<i>Rivulus</i> .	180
<i>Gobius</i> .	440	<i>goldmani</i> , <i>Batrachoides</i> .	487
<i>albopunctatus</i> .	440	<i>gomesii</i> , <i>Ophichthus</i> .	88
<i>alepidotus</i> .	445, 446	<i>Ophisurus</i> .	87, 88
<i>amoreo</i> .	436	<i>gomphodon</i> , <i>Oxyrhina</i> .	19
<i>anjerensis</i> .	442	<i>Gonionemion</i> .	280
<i>bacalaus</i> .	441	<i>serra</i> .	280
<i>banana</i> .	442	<i>Goniobatis</i> .	30
<i>bayamonensis</i> .	441	<i>macroptera</i> .	30
<i>boleosoma</i> .	439	<i>Goniodus</i> .	22
<i>bosci</i> .	445, 446	<i>Gonionarce</i> .	27
<i>boscii</i> .	442	<i>Gonioplectrus</i> .	308
<i>brasiliensis</i> .	448	<i>hispanus</i> .	308
<i>breviceps</i> .	440	<i>Gonocephalus</i> .	408
<i>brunneus</i> .	440	<i>macrocephalus</i> .	408
<i>carolinensis</i> .	440	<i>Gonocheilodon</i> .	359
<i>catulus</i> .	440	<i>Gonopterus</i> .	362
<i>chiquita</i> .	443	<i>microrens</i> .	362
		<i>microreus</i> .	362

List of scientific names—Continued

	Page		Page
Gonostoma...	72	gracilis, <i>Scopelus</i> ...	169
denudata...	72	Tigoma...	119
denudatum...	72	Umbrina...	351
elongatum...	72	Xiphophorus...	185
microdon...	72	gracilispinis, <i>Dolopichthys</i> ...	509
Gonostomidae...	72	gracilipes, <i>Læmonema</i> ...	213
Goodea...	183	graciosus, <i>Pimelodus</i> ...	152
atripinnis...	183	gradiens, <i>Hypsagonus</i> ...	393
caliantis...	184	gratilis, <i>Ophidion</i> ...	485
captiva...	184	grahami, <i>Gila</i> ...	114
luitpoldi...	184	Oligocephalus...	291
toweri...	184	grallator, <i>Benthosaurus</i> ...	166
whitei...	183	Gramma...	322
goodei, <i>Aldrobitus</i> ...	31	ioreto...	372
Aldrovandia...	75	Grammateus...	336
Centropholis...	266	medius...	337
Chiropeops...	175	grammaticum, <i>Thalassoma</i> ...	429
Erimyzon...	109	Grammatopleurus...	373
Halosaurus...	75	Grammatostoma...	70
Lucania...	175	dentatus...	70
Myliobatis...	31	Grammichthys...	229
Paralonchurus...	351	Grammicoelopidae...	232
Ptilichthys...	471	Grammicolepis...	232
Sabastodes...	304	brachiusculus...	232
Spinivomer...	84	Grammicotonotus...	197
Trachinotus...	277	bicolor...	197
Urolophus...	30	Grammistes acuminatus...	353
Urotrygon...	30	hepatus...	335
Zaclemus...	351	mauritii...	322
Goodeidae...	182	trivittatus...	331
goodii, <i>Macrurus</i> ...	203	granadensis, <i>Astatheros</i> ...	421
Nematorurus...	203	Cichlasoma...	421
gorbuscha, <i>Oncorhynchus</i> ...	54	grandicassis, <i>Arius</i> ...	149
Salmo...	54	grandiceps, <i>Microlepidium</i> ...	211
Gordichthys...	89	Paraliparis...	404
irretitus...	89	grandicomis, <i>Auchenipterus</i> ...	457
goreensis, <i>Albulia</i> ...	40	grandicorvus, <i>Scorpaena</i> ...	371
Hymnus...	274	grandipinnatus, <i>Photogenis</i> ...	129
Vomer...	275	grandis, <i>Fundulus</i> ...	176
Gorgasia...	77	Fundulus heteroclitus...	176
punctata...	77	Gila...	114
gorgonea, <i>Cheirodon</i> ...	96	Pomadasys...	334
Compsura...	06	Pomadasys...	334
goreensis, Vomer...	275	Ptychocheilus...	114
gouani, <i>Lepidopus</i> ...	262	grandisquamis, <i>Electris</i> ...	436
Trachipterus...	216	grandisquamis, <i>Chirodus</i> ...	426
gracile, <i>Bolesoma</i> ...	294	Upeneus...	344
Myctophum...	169	grandoculatus, <i>Centropomus</i> ...	306
Peristedion...	405	grandoculus, <i>Chiostoma</i> ...	251
gracilicauda, <i>Macrourus</i> ...	206	grandoculus, <i>Atherinichthys</i> ...	251
Macrurus...	206	granulatus, <i>Perca</i> ...	281
gracilicirrhus, <i>Umbrina</i> ...	350	Raia...	25
graciliors, <i>Gambusia</i> ...	186	Raja...	25
gracilis, <i>Aldrovanda</i> ...	75	granulatus, <i>Amphocephalus</i> ...	430
Atherinichthys...	247	granulosa, <i>Pristis</i> ...	23
Bathynectes...	481	Graodus...	133
Blennius...	471	nigrotentatus...	132, 133
Boleichterus...	294	grayi, <i>Carioides</i> ...	103
Catostomus...	105	Lepidotosteus...	37
cayuga, <i>Cottus</i> ...	389	grebnitzkii, <i>Neoliparis</i> ...	399
Chætodon...	300	Greeleya...	390
Cottus...	383, 385	rubellio...	391
Cyprinus (Leuciscus)...	136	greenoi, <i>Cottus</i> ...	384
Diapterus...	341	Porichthys...	487
Gadella...	211	Uranidea...	334
Gadus...	210, 211	greeni, <i>Ctenoides</i> ...	137
Gambusia...	185	Liparis...	399
Gila...	114	Neoliparis...	399
Gobius...	444	Polypora...	390
Hippocampus...	244	rex, <i>Pneumatophorus</i> ...	258
Hybopsis...	133	Scomber...	258
Leptocephalus...	70	greyi, <i>Istiophorus</i> ...	263
Leptogunnellus...	470	grimaldii, <i>Conchognathus</i> ...	78
Leuciscus...	128	grisea, <i>Cynopercula</i> ...	252
Lumpenus...	470	Dionda...	144
Lycodes...	475	Lucioperca...	282
Mollienesia...	192	Parapetenia tetracantha...	419
Molinessa...	192	Sciæna...	352
Monopterus...	131	griseolineatus, <i>Syngnathus</i> ...	240
Perca...	281	griseus, <i>Carcharias</i> ...	19
Photonectes...	70	Catostomus...	107
Pimelodus...	152	Catostomus (Acomus)...	107
Platygobio...	136	Heros tetracanthus...	419
Ptychocheilus...	113, 114	Hexanchus...	11
Scomber...	258	Lebrus...	324

List of scientific names—Continued

	Page		Page
<i>griseus</i> , <i>Lutianus</i>	325, 326	<i>Eula</i> , <i>Eucinostomus</i>	341
<i>Mesoprión</i>	325	<i>Gerres</i>	340, 341
<i>Saurus</i>	163	<i>Gila</i>	120
<i>Squalus</i>	11	<i>gulo</i> , <i>Holocentrus</i>	308
<i>grizalvii</i> , <i>Astroblepus</i>	159	<i>gulonellus</i> , <i>Pogonichthys (Platygobio)</i>	136
<i>greenlandicus</i> , <i>Microstomus</i>	68	<i>gułosia</i> , <i>Cathorops</i>	151
<i>Nansenii</i>	68	<i>gulosus</i> , <i>Chænobrytus</i>	302
<i>greenlandicus</i> , <i>Acanthocottus</i>	387	<i>Cottopsis</i>	383
<i>Aspidophoroides</i>	397	<i>Cottus</i>	383
<i>Cottus</i>	386, 387	<i>Gobius</i>	443
<i>Gunnellus</i>	468	<i>Microgobius</i>	443
<i>Himantolophus</i>	509	<i>Parachromis</i>	418
<i>Hippoglossus</i>	221	<i>Pomotis</i>	302, 303
<i>Salmo</i>	66	<i>Tachisurus</i>	151
<i>Gronias</i>	154	<i>gulula</i> , <i>Eucinostomus</i>	341
<i>nigritabilis</i>	154	<i>gummigutta</i> , <i>Hypoplectrus unicolor</i>	321
<i>gronovii</i> , <i>Gobius</i>	268	<i>Plectropoma</i>	321
<i>Nameus</i>	268	<i>gundlachi</i> , <i>Dormitor</i>	436
<i>Ostracion</i>	497	<i>gunelliformis</i> , <i>Asternopteryx</i>	468
<i>Solea</i>	220	<i>Gunnellus</i>	467
<i>Zoarcis</i>	473	<i>fabricii</i>	470, 471
<i>grossidens</i> , <i>Engraulis</i>	51	<i>greenlandicus</i>	468
<i>Lycengraulis</i>	51	<i>murenoides</i>	468
<i>Lycodapus</i>	479	<i>ornatus</i>	468
<i>grunniens</i> , <i>Amblodon</i>	352	<i>ruberinus</i>	468
<i>Aplodonutus</i>	352	<i>vulgaris</i>	467
<i>Mugil</i>	352	<i>gunnellus</i> , <i>Bleennius</i>	467
<i>gryllus</i> , <i>Exocetetus</i>	200	<i>Pholis</i>	467
<i>Gryistes lineatus</i>	373	<i>gunneri</i> , <i>Scomber</i>	217
<i>Guebicoara brasiliensis</i>	330	<i>Squalus</i>	20
<i>guacamala</i> , <i>Pseudoscarus</i>	435	<i>gunnisoni</i> , <i>Cyprinella</i>	130
<i>Scarus</i>	435	<i>güntheri</i> , <i>Alvordius</i>	283
<i>Guacamaya</i>	435	<i>Aspidophoroides</i>	387
<i>guacari</i> , <i>Hypostomus</i>	160	<i>Halosaurus</i>	74
<i>guachanco</i> , <i>Sphyraena</i>	255	<i>Hoplopagrus</i>	824
<i>guacharote</i> , <i>Anicistrus</i>	161	<i>Lampancyctus</i>	167
<i>Hypostomus</i>	161	<i>Mugil</i>	253
<i>guadalupae</i> , <i>Atherinops insularum</i>	252	<i>Sphyraena</i>	255
<i>guadalupensis</i> , <i>Amin</i>	206	<i>Ulcinus</i>	397
<i>Apono</i>	296	<i>Xiphophorus</i>	103
<i>gualcan</i> , <i>Echeneis</i>	449	<i>Xiphophorus helleri</i>	103
<i>Guamaícu atinga</i>	502	<i>guntheriana</i> , <i>Aluterata</i>	496
<i>Guamájacu</i>	501	<i>guppil</i> , <i>Congromurena</i>	80
<i>guanabana</i> , <i>Erizo</i>	501	<i>Girardinus</i>	191
<i>Guapervia</i>	491	<i>Gnathophis</i>	80
<i>lata</i>	492	<i>Gurnardus</i>	406
<i>guara</i> , <i>Guamájaci</i>	501	<i>birostratus</i>	406
<i>Guatapuca</i>	257	<i>gymnostethus</i>	406
<i>guasa</i> , <i>Serranus</i>	312	<i>loxias</i>	406
<i>guatemalensis</i> , <i>Anacyrtus</i>	98	<i>russarius</i>	406
<i>Arius</i>	150	<i>xenisma</i>	406
<i>Brycon</i>	98	<i>guttata</i> , <i>Liinamurana</i>	90
<i>Chirostoma</i>	249	<i>Perca</i>	311
<i>Citharichthys</i>	219	<i>Raja</i>	28
<i>Fundulus</i>	174	<i>Scorpaena</i>	371
<i>Galeichthys</i>	150	<i>guttatus</i> , <i>Astrodermus</i>	264
<i>Pimelodus</i>	157	<i>Astrocopus</i>	454
<i>Rhamdia</i>	167	<i>Balistes</i>	492
<i>Rhaboldes</i>	98	<i>Bodianus</i>	300
<i>Thyrina</i>	249	<i>Chirus</i>	373
<i>Guativere amarillas</i>	309	<i>Dasyatis</i>	28
<i>guativere</i> , <i>Bodianus</i>	309	<i>Epinephelus</i>	311
<i>Guatucupa juba</i>	332	<i>Johnius</i>	313
<i>Guavina</i>	435, 436	<i>Labrus</i>	426
<i>guavina</i>	436	<i>Lutianus</i>	326
<i>guavina</i> , <i>Electris</i>	436	<i>Mesoprión</i>	326
<i>Guavina</i>	436	<i>Ophisurus</i>	87
<i>guayamasie</i> , <i>Gillichthys</i>	444	<i>Perca</i>	308
<i>guaza</i> , <i>Epinephelus</i>	310	<i>Percopis</i>	214, 215
<i>Labrus</i>	310	<i>Pomotis</i>	304
<i>guentheri</i> , <i>Galeichthys</i>	149	<i>Zeus</i>	216
<i>Homaloglystes</i>	310	<i>Zygogenes</i>	178
<i>Mugil</i>	253	<i>guttavarium</i> , <i>Plectropoma</i>	321
<i>Strembo</i>	483	<i>guttavarius</i> , <i>Hypoplectrus unicolor</i>	321
<i>Guentheridia</i>	500	<i>guttifer</i> , <i>Ophichthus</i>	87
<i>formosa</i>	500	<i>guttifrons</i> , <i>Ophichthys</i>	87
<i>guianensis</i> , <i>Anchoviella</i>	49	<i>guttulata</i> , <i>Hypsopsetta</i>	225
<i>Stolephorus</i>	49	<i>guttulatus</i> , <i>Cichlaurus</i>	416
<i>Strongylura</i>	196	<i>Heros</i>	416
<i>Tylosurus</i>	196	<i>Pisodonophis</i>	85
<i>guichenoti</i> , <i>Callechelys</i>	86	<i>Pleuronichthys</i>	225
<i>Cayenna</i>	448	<i>gymnacanthus</i> , <i>Catostomus (Acomus)</i>	104
<i>gulia</i> , <i>Thyrina</i>	248	<i>gymnacanthus</i>	389
<i>guineensis</i> , <i>Ostracion</i>	407	<i>galeatus</i>	389
<i>Guiritings</i>	149	<i>Gymnachirinae</i>	229

List of scientific names—Continued

	Page		Page
<i>Gymnachirus</i>	220	<i>gymnurus</i> , <i>Gasterosteus</i>	237
<i>fuscatus</i>	220	<i>gyrans</i> , <i>Querimana</i>	253
<i>nudus</i>	220	<i>Gyrinichthys</i>	403
<i>Gymnepigynathus</i>	270	<i>minytrema</i>	403
<i>Gymnoleotris</i>	438	<i>gyrinops</i> , <i>Cyclopterooides</i>	398
<i>seminuda</i>	438	<i>gyrinus</i> , <i>Fleotris</i>	436
<i>Gymnolis</i>	477	<i>Eurytemen</i>	401
<i>conorhynchus</i>	477	<i>Gobiesox</i>	489
<i>pictus</i>	477	<i>Schilbeoides</i>	165
<i>stigma</i>	477	<i>Silurus</i>	165
<i>viridis</i>	477	<i>Gyropleurodon</i>	10
<i>unimaculatus</i>	477	<i>francisi</i>	10
<i>Gymnelus pictus</i>	477		
<i>Gymnoanthus</i>	389	Hadropterus	283
<i>galeatus</i>	389	<i>evermanni</i>	284
<i>pistilliger</i>	389	<i>maculatus</i>	283
<i>tricuspidis</i>	389	<i>maxinkuckiensis</i>	284
<i>Gymnocephalus</i>	267	<i>nigrofasciatus</i>	283
<i>messiniensis</i>	267	<i>roanoka</i>	284
<i>ruber</i>	309	<i>scierus</i>	284
<i>Gymnolinus</i>	466	<i>serrula</i>	284
<i>eristulatus</i>	466	<i>sellaris</i>	284
<i>Gymnodontes</i>	497	<i>schumardi</i>	285
<i>Gymnogaster</i>	262	<i>tessellatus</i>	289
<i>arcticus</i>	216	(<i>Hydropterus</i>) <i>cymatotenia</i> , <i>Etheostoma</i>	285
<i>gymnogaster</i> , <i>Sicydium</i>	447	<i>langue</i> , <i>Etheostoma</i>	285
<i>Gymnomuraena</i>	94	<i>ouachitae</i> , <i>Etheostoma</i>	284
<i>doliata</i>	94	<i>squamatus</i> , <i>Etheostoma</i>	283
<i>nectaria</i>	95	<i>haematurus</i> , <i>Hypopsis</i>	145
<i>Gymnonotus</i>	101	Hæmulidae	329
<i>Gymnotus acus</i>	486	Hæmulon	329
<i>carapo</i>	486	<i>acutum</i>	329
<i>Gymnopsis</i>	94	<i>albidum</i>	329
<i>Gymnosarda</i>	258	<i>album</i>	329
<i>gymnostethus</i> , <i>Gurnardus</i>	406	<i>arara</i>	330
<i>Prionotus</i>	406	<i>arcuatum</i>	330
<i>Gymnothorax</i>	91	<i>aurolineatum</i>	331
<i>afer</i>	91	<i>bonariense</i>	329
<i>albimentis</i>	93	<i>brevirostrum</i>	330
<i>aque dulcis</i>	93	<i>canna</i>	329
<i>brunneus</i>	92	<i>carbonarium</i>	330
<i>carcino gnathus</i>	93	<i>caudimacula</i>	329
<i>castaneus</i>	92	<i>chromis</i>	329
<i>catenatus</i>	94	<i>chrysargyreum</i>	330, 331
<i>chile vestes</i>	93	<i>chrysopterum</i>	329
<i>concolor</i>	93	<i>continuum</i>	329
<i>conversus</i>	92	<i>corviniforme</i>	333
<i>dovii</i>	92	<i>corviniformis</i>	333
<i>elaboratus</i>	93	<i>dorsale</i>	330
<i>flavoscriptus</i>	91	<i>elegans</i>	329
<i>funebris</i>	92	<i>flaviguttatus</i>	331
<i>jordanii</i>	92	<i>flavolineatum</i>	330
<i>(Limamurina) argus</i>	90	<i>formosum</i>	330
<i>miliaris</i>	92	<i>fremebundum</i>	329
<i>mordax</i>	91	<i>heterodon</i>	330
<i>moringa</i>	91	<i>hians</i>	330
<i>murena</i>	91	<i>joniquano</i>	331
<i>nigrocastaneus</i>	90	<i>labridum</i>	332
<i>obscuratus</i>	93	<i>luteum</i>	330
<i>ocellatus</i>	91, 93	<i>macrostoma</i>	329
<i>nigromarginatus</i>	93	<i>macrostomum</i>	329
<i>sexcola</i>	93	<i>maculosum</i>	329
<i>picturatus</i>	91	<i>margariferum</i>	331
<i>polygonus</i>	92	<i>mazatlanum</i>	331
<i>punctarenas</i>	92	<i>melanurum</i>	330
<i>punctatus</i>	92	<i>microphthalmum</i>	329
<i>rostratus</i>	91	<i>mowbrayi</i>	329
<i>sancti-helenae</i>	92	<i>multilineatum</i>	330
<i>scriptus</i>	92	<i>notatum</i>	329
<i>umbrosus</i>	90	<i>obtusum</i>	332
<i>verilli</i>	93	<i>parra</i>	329
<i>versipunctatus</i>	92	<i>plumieri</i>	330
<i>vicinus</i>	92	<i>quadrilineatum</i>	331
<i>virescens</i>	92	<i>quinqueleatum</i>	331
		<i>retrocurrons</i>	329
Gymnotidae	101	<i>rimator</i>	331
Gymnotorpedo	27	<i>schrankii</i>	330
<i>Gymnotus</i>	101	<i>selurus</i>	330
<i>acus</i>	101	<i>scudderii</i>	330
<i>albifrons</i>	101	<i>serratum</i>	329
<i>albus</i>	101	<i>sexfasciatum</i>	329
<i>carapo</i>	101	<i>soxfasciatum</i>	329
<i>fuscatus</i>	101	<i>similis</i>	330
<i>macrourus</i>	101	<i>steindachneri</i>	330
<i>maercurius</i>	101	<i>subarcuatum</i>	330
<i>putao</i>	101	<i>taeniatum</i>	330, 331
<i>gymnura</i> , <i>Trygon</i>	28		

List of scientific names—Continued

	Page		Page
Hæmulon undecimale.....	330	Harpe cœruleo-aureus.....	425
xanthopteron.....	330	diplotenia.....	425
Hæmulops.....	333	pectoralis.....	425
(Hæmulopsis) axillare, <i>Pristipoma</i>	333	rufa.....	309
brevipinne, <i>Pristipoma</i>	335	harperi, <i>Spheroïdes</i>	498
nitidum, <i>Pristipoma</i>	333	Sphœroïdes.....	498
hares, <i>Gobiesox</i>	490	Harpurus.....	362
Sloiates.....	490	fasciatus.....	362
Halactractus.....	279	harringtoni, <i>Atherina</i>	246
dorsalis.....	279	harringtoni, <i>Axyrias</i>	377
halec, <i>Clupea</i>	41	Harriotta.....	33
Halecomorphi.....	38	curtiss-jamesi.....	33
Halecula.....	46	raleighana.....	33
acuminata.....	46	harroweri, <i>Isopisthus</i>	354
Hallas.....	479	harti, <i>Haplochilus</i>	180
Halichthres.....	426	Rivulus.....	180
californicus.....	428	hasselti, <i>Caranx</i>	270
sellifer.....	428	Paraserranus.....	318
Halleutæa.....	507	hasta, <i>Lutjanus</i>	334
spongiosa.....	508	hastata, <i>Pastinaca</i>	28
stellata.....	507	hastatus, <i>Dasyatis</i>	28
Halleutella.....	508	Gobionellus.....	440, 441
lappa.....	508	hastingsi, <i>Lutjanus</i>	326
Halleutichthys.....	507	Neomenis.....	326
aculeatus.....	507	Haustor.....	152
caribbeus.....	507	catus.....	153
reticulatus.....	507	dugesi.....	153
smithii.....	507	lacustris.....	152
Hallperca.....	318	lophius.....	153
halleri, <i>Urobatis</i>	29, 30	lupus.....	153
Crotalus.....	29	okeechobeensis.....	152
Haloporphyrus viola.....	211	prael.....	153
Haloporphyrus.....	211	havana, <i>Gerres</i>	342
halops, <i>Acipenser</i>	34	Xystéma.....	342
Halosurichthys.....	74	havannensis, <i>Muraena</i>	87
Halosauridæ.....	74	hawaiensis, <i>Elops</i>	39
Halosauropsis.....	75	hawaliensis, <i>Macrorhamphosus</i>	246
Halesaurus.....	74	haydeni, <i>Ptychostomus</i>	109
attenuatus.....	75	hayi, <i>Hybognathus</i>	144
goodei.....	75	hearnel, <i>Salmo</i>	60
günterii.....	74	heathi, <i>Schedophilus</i>	268
macrochir.....	75	heckeli, <i>Acus</i>	244
oweni.....	74	Balistes.....	403
ovenii.....	74	Gambusia.....	185
radiatus.....	74	Nerophis.....	244
rostrata.....	75	hedricki, <i>Astatheros</i>	421
rostratus.....	75	Cithlasoma.....	421
hamatus, <i>Icterus</i>	378	höermannii, <i>Amphistichus</i>	411
hamiltoni, <i>Brachiptilon</i>	32	heinleri, <i>Labrisomus</i>	459
Hippoglossoides.....	222	Heiropætra.....	25
Manta.....	32	helena, <i>Muraena</i>	90
hammondi, <i>Pimelodus</i>	152	Murænophis.....	90
hammondi, <i>Percopsis</i>	215	Hæliastes.....	413
Semotilus.....	117	enchyrsurus.....	413
hankinsoni, <i>Parapterius</i>	377	insolatus.....	413
Haplochilus dovii.....	179	marginatus.....	413
floripinnis.....	177	multilineatus.....	412
harti.....	180	Hæliastes.....	413
melanops.....	185	Helicolenus.....	370
Haplodoci.....	486	dactylopterus.....	370
haplognatha, <i>Helioperca</i>	301	madereusis.....	370
haplognathus, <i>Lepomis</i>	301	Hælioperca.....	301
Haplodonotus.....	352	bombifrons.....	301
Haplomi.....	172	haplognatha.....	301
hardwicki, <i>Ratlabouri</i>	89	incisor.....	301
harengoides, <i>Glossodon</i>	40	macrochira.....	301
Harengula.....	43	marginata.....	301
callolepis.....	43	nephela.....	301
clupeola.....	43	helleri, <i>Anchoviella</i>	48
humeralis.....	43	brevis, <i>Xiphophorus</i>	104
latula.....	43	guntheri, <i>Xiphophorus</i>	103
macrothymus.....	43	Heros.....	422
maculosa.....	43	ranchovii, <i>Xiphophorus</i>	103
pensacolae.....	43	strigatus, <i>Xiphophorus</i>	103
sardina.....	43	Thorichthys.....	422
thrissina.....	43	Xiphophorus.....	193
barengulus, <i>Eucinostomus</i>	341	helolepis, <i>Trachyrincus</i>	203
barengus, <i>Clupea</i>	41	Hæmichthys.....	79
Lavinia.....	113	diaphanus.....	79
Leucichthys.....	52	Hæmictis.....	79
Myxus.....	253	heloops, <i>Trachyrhynchus</i>	203
Salmo (Coregonus).....	62	Trachyrhynchus.....	203
harfordi, <i>Ptychocheilus</i>	114	Helops.....	34
bariani, <i>Lepidostous</i>	37	helops, <i>Acipenser</i>	34
Härpe.....	425	helomaculatus, <i>Sebastes</i>	367
cœruleoaurorum.....	425	Hæmiancistrus.....	161

List of scientific names—Continued

Page	Page
<i>Hemianthias</i>	322
<i>peruanus</i>	322
<i>vivanus</i>	322
(<i>Hemianthias</i>) <i>peruanus</i> , <i>Anthias</i>	322
<i>Hemiarius</i>	149
<i>Hemibranchii</i>	237
<i>Hembrycon</i>	97
<i>dariensis</i>	97
(<i>Hembrycon</i>) <i>polyodon</i> , <i>Tetragonopterus</i>	97
<i>Hemicarax</i>	271
<i>amblyrhynchus</i>	271
<i>atrimanus</i>	271
<i>falcatus</i>	271
<i>furthii</i>	271
<i>leucus</i>	271
<i>marginatus</i>	271
<i>rhomboides</i>	271
<i>zelotes</i>	271
<i>Hemicatodon</i>	359
<i>Hemieleotris</i>	437
<i>latifasciatus</i>	437
<i>Hemieroxocetus</i>	199
<i>caudimaculatus</i>	199
<i>Hemigalea</i>	14
<i>Hemigaleus</i>	14
<i>microstoma</i>	14
<i>pectoralis</i>	14
<i>Hemigrammus</i>	98
<i>compressus</i>	97
<i>minutus</i>	97
<i>unilineatus</i>	98
<i>hemigymna</i> , <i>Garmannia</i>	445
<i>hemigymnus</i> , <i>Argyropelecus</i>	74
<i>Gobius</i>	445
<i>Hemilepidotus</i>	380
<i>gibbsi</i>	381
<i>hemilepidotus</i>	381
<i>jordani</i>	381
<i>tileisi</i>	381
<i>zapus</i>	381
<i>hemilepidotus</i> , <i>Cottus</i>	380, 381
<i>llenilepidotus</i>	381
<i>Hemiodon</i>	161
<i>depressus</i>	161
<i>platycephalus</i>	161
<i>hemiodon</i> , <i>Carcharias</i>	17
<i>Hemiodontidae</i>	100
<i>Hemioleorcaria</i>	161
<i>caracasenses</i>	161
<i>Hemioptilites</i>	304
<i>simulans</i>	304
<i>Hemiplus</i>	114
<i>lacustris</i>	114, 115
<i>hemiplus</i> , <i>Cyprinus</i>	115
<i>Hemiramphidae</i>	198
<i>Hemiramphus</i>	199
<i>balao</i>	199
<i>brasiliensis</i>	199
<i>brownii</i>	199
<i>fuscatus</i>	198
<i>filamentosus</i>	199
<i>longirostris</i>	198
<i>macrochirius</i>	199
<i>picarti</i>	198
<i>pleii</i>	199
<i>richardi</i>	198
<i>roberti</i>	198
<i>rosei</i>	198
<i>saltator</i>	199
<i>unifasciatus</i>	198
<i>Hemirhombus</i>	218
<i>aramaca</i>	218
<i>fimbriatus</i>	220
<i>fucus</i>	210
<i>guineensis</i>	218
<i>ovalis</i>	218
<i>Hemistoma</i>	433
<i>reticulatum</i>	433
<i>Hemitremia</i>	121
<i>maculata</i>	135
<i>milneriana</i>	121
<i>vittata</i>	121
<i>Hemitriptericidae</i>	393
<i>Hemitripterus</i>	393
<i>americanus</i>	393
<i>cavifrons</i>	393
<i>marmoratus</i>	382, 393
<i>villosum</i>	393
<i>Hemixiphophorus</i>	184, 185
<i>hemphilli</i> , <i>Stathmonotus</i>	465
<i>henlei</i> , <i>Carcharias</i>	16
<i>Rhinotrincis</i>	13, 14
<i>henshalli</i> , <i>Zygogenes</i>	177
<i>henshawi</i> , <i>Rhinichthys</i>	141
<i>henshawi</i> , <i>Salmo</i>	57
<i>hentz</i> , <i>Blennius</i>	402
<i>Hypsoblennius</i>	462
<i>hepatus</i> , <i>Acanthurus</i>	362
<i>Grammistes</i>	335
<i>Teuthis</i>	302
<i>Hepsetia</i>	240
<i>evermanni</i>	240
<i>stipes</i>	246
<i>Hepsetus</i> , <i>Atherina</i>	246, 247
<i>heptacanthus</i> , <i>Chelodipterus</i>	280
<i>heraldii</i> , <i>Tetronotus</i>	499
<i>herichthys</i>	422
<i>bocourti</i>	423
<i>carpintis</i>	423
<i>cyanoguttatus</i>	422
<i>geddesi</i>	423
<i>pavonaceus</i>	422
<i>troscheli</i>	423
<i>underwoodi</i>	423
<i>herminieri</i> , <i>Blennius</i>	459
<i>Labrisomus</i>	459
<i>Herinopsis</i>	426
<i>Hermosilla</i>	340
<i>azurae</i>	340
<i>robusta</i>	340
<i>Heros</i>	416
<i>affinis</i>	422
<i>altifrons</i>	420
<i>angulifer</i>	410
<i>aureus</i>	422
<i>balticus</i>	418
<i>basilaris</i>	421
<i>beani</i>	419
<i>bifuscatus</i>	416
<i>callolepis</i>	422
<i>contrarius</i>	420
(<i>Clinusoma</i>) <i>spinosissimus</i>	420
<i>citrinellus</i>	420, 422
<i>deppii</i>	416
<i>dovii</i>	418
<i>erythreus</i>	422
<i>friedrichstahli</i>	418
<i>geddesi</i>	423
<i>gibbiceps</i>	417
<i>godmani</i>	416
<i>guttulatus</i>	416
<i>helleri</i>	422
<i>heterodontus</i>	420
<i>intermedius</i>	416
<i>istianus</i>	418
<i>inabilitum</i>	422
<i>labridens</i>	419
<i>lentiginosus</i>	418
<i>lobochilus</i>	422
<i>longimanus</i>	420
<i>maculipinnis</i>	422
<i>managuensis</i>	418
<i>margaritifer</i>	421
<i>melanopogon</i>	421
<i>mento</i>	419
<i>microphthalmus</i>	410
<i>montezuma</i>	421
<i>motaguensis</i>	418
<i>multispinosus</i>	423
<i>nebulifer</i>	417
<i>nicaraguensis</i>	418
<i>nigricans</i>	419
<i>nigrofasciatus</i>	420
<i>oblongus</i>	421
<i>octofasciatus</i>	421

List of scientific names—Continued

	Page		Page
<i>Heros parma</i>	419	<i>heterurus, Cypselurus</i>	201
<i>pavonaceus</i>	422	<i>Erotactus</i>	201
<i>rostratus</i>	421	<i>Glossodon</i>	40
<i>salvini</i>	418	<i>hexacanthus, Centrarchus</i>	304, 305
<i>severus</i>	410	<i>Dipteron</i>	295
<i>sieboldii</i>	423	<i>hexacornis, Cottus</i>	388
<i>spilurus</i>	420	<i>Oncocottus</i>	388
<i>teporatus</i>	422	<i>hexagonatus, Stomias</i>	69
<i>tetraacanthus cinctus</i>	419	<i>Hexagrammidae</i>	373
<i>griseus</i>	419	<i>Hexagrammos</i>	373
<i>latus</i>	419	<i>aspor</i>	373, 374
<i>torralbasi</i>	419	<i>deagrammus</i>	373
<i>triogramma</i>	418	<i>lagocephalus</i>	374
<i>trimaculatum</i>	420	<i>octogrammus</i>	374
<i>troscellum</i>	423	<i>stellori</i>	373, 374
<i>urophthalmus</i>	418	<i>superchilios</i>	374
<i>heros, Eupomotis</i>	302	<i>Hexagrammus scaber</i>	374
<i>Pomotis</i>	302	<i>hexagrammus, Labrus</i>	374
<i>Herotilapia</i>	423	<i>Hexanchidae</i>	11
<i>multispinosa</i>	423	<i>Hexanchus</i>	11
<i>Herpetoichthys</i>	87	<i>corinus</i>	11
<i>callisoma</i>	88	<i>griseus</i>	11
<i>ornatissimus</i>	87	<i>Hexanematichthys</i>	149
<i>sulcatus</i>	87	<i>surinamensis</i>	150
<i>herschelinus, Carelliparis</i>	400	<i>heyel, Rutilus</i>	180
<i>Liparis</i>	400	<i>hians, Abelones</i>	197
<i>hertzbergi, Silurus</i>	148	<i>Belone</i>	197
<i>Hesperianthias</i>	328	<i>Hæmulon</i>	330
<i>herperis, Priapichthys annectens</i>	187	<i>Myctophum</i>	169
<i>Hesporoleucus</i>	121	<i>Sayris</i>	197
<i>anticus</i>	122	<i>Hiatula</i>	424
<i>mearnsi</i>	122	<i>gardeniana</i>	424
<i>mitrulus</i>	122	<i>Hiatula, Labrus</i>	424
<i>navarrensis</i>	121	<i>hieroglyphicus, Zygonectes</i>	178
<i>parvipinnis</i>	122	<i>Hiroptera</i>	24
<i>subditus</i>	122	<i>abredonensis</i>	24
<i>symmetricus</i>	121	<i>Hildebrandia</i>	80
<i>venustus</i>	122	<i>flava</i>	80
<i>heteracantha, Rhamdia</i>	158	<i>hildebrandi, Opsanus</i>	487
<i>heteracanthus, Dolopichthys</i>	510	<i>hildebrandti, Etheostoma</i>	291
<i>Heterandria</i>	188	<i>Oligocephalus</i>	201
<i>affinis</i>	185	<i>hilianus, Etmopterus</i>	404, 405
<i>colombianus</i>	189	<i>Exocetus</i>	21
<i>cubensis</i>	188	<i>Spinax</i>	21
<i>formosa</i>	188	<i>Himantolophidae</i>	509
<i>formosus</i>	188	<i>Himantolophus</i>	509
<i>holbrookii</i>	185, 188	<i>greenlandicus</i>	509
<i>lutzii</i>	189, 190	<i>reinhardtii</i>	509
<i>nobilis</i>	185	<i>Himantophorus, Callionymus</i>	450
<i>occidentalis</i>	189	<i>Hiodon</i>	40
<i>ommata</i>	174	<i>clodalis</i>	40
<i>patruelis</i>	184, 185	<i>solenops</i>	41
<i>zonata</i>	191	<i>tergisus</i>	40
<i>heterandria, Limia</i>	191	<i>Hiodontidae</i>	40
<i>heterocita, Cobitis</i>	175	<i>Hipolliti, Gobius</i>	440
<i>heteroclitus badius, Fundulus</i>	176	<i>Hippichthys</i>	243
<i>heteroclitus bermudæ, Fundulus</i>	175	<i>albirostris</i>	243
<i>heteroclitus fonticola, Fundulus</i>	175	<i>cayorum</i>	243
<i>heteroclitus, Fundulus</i>	176	<i>heptagonus</i>	243
<i>heteroclitus grandis, Fundulus</i>	176	<i>spicifer</i>	243
<i>heteroclitus macrolepidotus, Fundulus</i>	176	<i>Hippocampinae</i>	244
<i>heterodon, Alburnops</i>	136	<i>Hippocampus</i>	244
<i>Hæmulon</i>	136	<i>brunneus</i>	244
<i>Hybopsis</i>	136	<i>fasciularis</i>	244
<i>richardsoni, Hybopsis</i>	136	<i>gracilis</i>	244
<i>Heterodontidae</i>	10	<i>heptagonus</i>	243
<i>heterodontus, Astatheros</i>	420	<i>hudsonius</i>	244
<i>Heros</i>	420	<i>ingens</i>	244
<i>heteroglossus, Gadus</i>	210	<i>lavicaudatus</i>	244
<i>Heterognathidae</i>	95	<i>marginalis</i>	244
<i>Heterognathus</i>	250	<i>punctulatus</i>	244
<i>heterognathus, Ophichthidae</i>	80	<i>styliifer</i>	244
<i>heterolepis, Hybopsis</i>	134	<i>zosteræ</i>	244
<i>Notropis</i>	134	<i>Hippoglossidae</i>	221
<i>Heteromii</i>	75	<i>Hippoglossina</i>	222
<i>Heterophallina</i>	184, 186	<i>bolmani</i>	223
<i>Heterophallus</i>	184	<i>macrops</i>	222, 223
<i>rachovii</i>	184	<i>sabanensis</i>	223
<i>Heterosomatidae</i>	217	<i>stomata</i>	223
<i>Heteropropsonopon</i>	225	<i>Hippoglossoides</i>	222
<i>heteropygus, Caranx</i>	271	<i>elassodon</i>	222
<i>Heterostichus</i>	458	<i>exilis</i>	222
<i>rostratus</i>	458	<i>hamiltoni</i>	222
<i>heterura, Clupea</i>	46	<i>jordani</i>	222
<i>Compsura</i>	95		

List of scientific names—Continued

	Page		Page
Hippoglossoides platessoides.	222	holbrookii, Sargus.	338
robustus.....	222	Holcomycteronus.	481
hippoglossoides, Pleuronectes.	221	digitatus.....	481, 482
Reinhardtius.	221	Holconoti.	409
Hippoglossus.	221	Holconotus.....	410, 411
brasiliensis.....	224	agassizii.....	411
californicus.....	223	fuliginosus.....	411
gigas.....	221	gibbonsi.....	412
greenlandicus.....	221	inegalops.....	410
hippoglossus.....	221	rhodoterus.....	410
maximus.....	221	trowbridgei.....	409
ocellatus.....	218	holderi, Allocinclus.....	458
ponticus.....	221	Starckia.....	456
hippoglossus, Hippoglossus.	221	holoi, Makaira.....	263
Pleuronectes.....	221	Hollardia.	491
hippoides, Caranx.	273	hollardi.....	491
hippops, <i>Capostriona</i>	148	hollardi, Hollardia.....	491
hippos, <i>Caranx</i>	273	hollbolli, Ceratias.....	508
Acomber.....	273	Holocanthus.	498
hippurus, <i>Coryphina</i>	264	leionothos.....	498, 499
hirudo, <i>Acanthurus</i>	362	lunatus.....	361
Ichthiomyzon.....	9	melanophos.....	497
hirundinaceus, <i>Squalus</i>	17	holocanthus, <i>Diodon</i>	501
Hirundo.	409	Holocentridae.	234
hirundo, <i>Azurina</i>	412	Holocentrum.	235
Lelocottus.....	390	coruscum.....	236
Hispanisicus.	388	longipinne.....	235
elongatus.....	368	marijanum.....	236
levius.....	368	osculum.....	236
rubrifinctus.....	368	perlatum.....	236
sinensis.....	368	puncticulatum.....	236
zacontrus.....	368	retrospinis.....	236
hispanis, Serrana.	353	riparium.....	236
hispanum, Plectropoma.	308	rostratum.....	236
hispanus, Gonioplectrus.	308	sancti-pauli.....	236
Serrana.....	353	sicciferum.....	236
hispidus, <i>Balistes</i>	494	suborbitalis.....	236
Cottus.....	393	vexillarium.....	236
Monacanthus.....	494	Holocentrus.	235
var. maculatus, <i>Tetronotus</i>	498	auratus.....	309
Histichthys.	215	bengalensis.....	324
Histiobranchus.	78	brachypterus.....	236
bathybius.....	78	coruscus.....	236
infernalis.....	78	flavocæruleus.....	310
Histioclinius.	456	gulo.....	308
veiliger.....	456	meeki.....	236
Histiocottus.	382	morou.....	310
bilobus.....	382	osculus.....	236
Histiophorus.	263	rostratus.....	235
americanus.....	263	rufus.....	236
audax.....	263	sancti-pauli.....	236
gladius.....	263	sanguineus.....	363
indicus.....	263	sogo.....	235
Histro.	504	striatus.....	235
gibba.....	505	surinamensis.....	323
jagua.....	505	tigrinus.....	319
pictus.....	504	tortugae.....	236
histrio, Ethostoma (Ulocentra).	286	vexillarius.....	236
Gobiosoma.....	446	Holocephalus.	32
Lophius.....	504, 505	holocyaneos, <i>Scarus</i>	434
Scorpaena.....	371	Hololepis.	294
Ulocentra.....	286	barratti.....	294
hiulcus, Stolephorus.	48	erochrous.....	294
Holancanthus.	361	hololepis, <i>Cytinus</i>	232
clarionensis.....	361	zenion.....	232
cornutus.....	361	holomelas, Paraliparis.	404
formosus.....	361	Holorhinus.	31
lunatus.....	361	Holosteus.	38
passer.....	361	Holostomi.	76
strigatus.....	361	Holotrachys.	235
tibicen.....	361	lima.....	235
tricolor.....	361	holubi, <i>Cypselurus</i>	202
holanthus, Diodon.	501	Exocetus.....	202
Neopocilla.....	190	Homalogrystes.	310
holbrookii, <i>Acipenser</i>	34	guentheri.....	310
Aluterus.....	495	Homalopomus trowbridgi.	214
Echeneis.....	449	Homesthes.	463
Gambusia.....	184	caulopus.....	463
Heterandria.....	185, 188	homianus, <i>Squalus</i>	20
Ophidion.....	485	homocyanus, <i>Gobius</i>	440
Ophidium.....	485	homonymus, <i>Diapterus</i>	341
Pomotis.....	302	Homopiron.	346
holbrookii, <i>Diplodus</i>	338	lanceolatus.....	346
Eupomotis.....	302	subtruncatus.....	345
Gamhusia.....	185	(Homopiron) firthii, <i>Corvina</i>	347
Pomotis.....	302	homopterus, <i>Bathytyctes</i>	53

List of scientific names—Continued

	Page		Page
hopkinsi, <i>Acutomentum</i>	366	humills, <i>Pristipoma</i>	334
Hynnus.....	274	Tetragonopterus.....	96
Mycteroperca.....	314	humilis, <i>Pagellus</i>	337
Plagiogrammus.....	469	huntnia, <i>Molva</i>	212
Sebastodes.....	366	Huro.....	297
Hopladelus.....	155	floridana.....	297
Hoplias.....	99	nigricans.....	297
malabaricus.....	99	huronensis, <i>Lepisosteus</i>	36
microlepis.....	99	huronius, <i>Argyrosomus</i>	62
Hopliticus, <i>Paricelinus</i>	375	Leucichthys sisco.....	62
Hoplomystax, <i>Scarus</i>	432	(<i>Huso</i>) anasimios, <i>Acipenser</i>	35
Sparisoma.....	432	anthracinus, <i>Acipenser</i>	35
Hoplopagridæ.....	324	atelaspis, <i>Acipenser</i>	35
Hoplopagrus.....	324	copel, <i>Acipenser</i>	35
guntheri.....	324	kirtlandi, <i>Acipenser</i>	35
Hoplosternum.....	160	lamarili, <i>Acipenser</i>	35
punctatum.....	160	megalepis, <i>Acipenser</i>	35
punctatus.....	160	nertinianus, <i>Acipenser</i>	35
(<i>Hoplosternum</i>) thoracatus, <i>Callichthys</i>	160	paranasmios, <i>Acipenser</i>	35
Hoplostethus.....	234	platyrhinus, <i>Acipenser</i>	35
cornutus.....	233	rathesquii, <i>Acipenser</i>	35
mediterraneus.....	234	rauchi, <i>Acipenser</i>	35
Hoplunnis.....	81	richardsoni, <i>Acipenser</i>	35
diomedianus.....	81	rosarium, <i>Acipenser</i>	35
schmidti.....	81	hyalina, <i>Sudis</i>	165
horatii, <i>Hybopsis</i>	134	hyalinus, <i>Ceratichthys</i>	137
Lepidosteus.....	345	Erinemus.....	137
Notropis.....	134	hyalocephalus, <i>Squalius</i>	117
horkeli, <i>Rhinobatos</i>	23	Hybognathus.....	143
horrens, <i>Prionotus</i>	408	argyritis.....	144
horridus, <i>Brachysomophis</i>	89	civilis.....	144
horstii, <i>Gobiosoma</i>	446	(<i>Dionda</i>) punctifer.....	144
hoskynii, <i>Bathyclupea</i>	281	evansi.....	143
hospes, <i>Mugil</i>	254	flavipinnis.....	144
Querimana.....	254	hai.....	144
houghi, <i>Pimeodus</i>	152	nigrotentata.....	144
hoysi, <i>Argyrosomus</i>	64	nuchalis.....	143, 144
Coregonus.....	62	osmerinus.....	143
Leucichthys.....	64	placitus.....	143
Pimeodus.....	154	procne.....	133
Uranidea.....	386	regius.....	143
hubbardii, <i>Parophrys</i>	226	siderius.....	140
Hubbesia.....	218	stramineus.....	133
gilberti.....	248	volucellus.....	133
huberi, <i>Priapichthys</i>	157	(<i>Hybognathus</i>) siderius, <i>Hyborhynchus</i>	140
hudsonia, <i>Clupea</i>	132	Hybopsis.....	117, 133
hudsonicus, <i>Salmo</i>	69	festivalis marconis.....	138
Hudsonius.....	132	anogenus.....	136
aletes.....	132	atrocaudalis.....	133
altus.....	137	bifrenata.....	135
amarus.....	132	bifrenatus.....	133, 135
chamberlaini.....	132	bivittatus.....	119
euryopa.....	132	blennius.....	133
fluvialis.....	132	boops.....	134
hudsonius.....	132	buchananii.....	134
saladanus.....	132	calentis.....	134
hudsonius, <i>Catostomus</i>	106	cayuga.....	133
Hippocampus.....	241	chalibaeus.....	127
Hudsonius.....	132	chilictus.....	127
humboldti, <i>Cheonda</i>	119	chlorocephalus.....	127
Cyclopium.....	159	chromosomus.....	127
Myctophum.....	160	crameri.....	138
Sternopygus.....	101	cummingsi.....	133
Stygogenes.....	159	deliciosa.....	134
Tigoma.....	119	straminea.....	134
humboldtiana, <i>Atherina</i>	250	deliciosa.....	134
humboldtianum, <i>Chiostoma</i>	250	dorsalis.....	134
humboldti, <i>Gasteropelecus</i>	169	fretensis.....	129
humeralis, <i>Arbacia</i>	490	gilberti.....	134
Chætodon.....	300	gracilis.....	133
Gobiesox.....	490	haematurus.....	145
Harengula.....	43	heterodon.....	136
Julis.....	426	heterolepis.....	184
Leiostomus.....	349	horatii.....	134
Oligocephalus.....	293	jordani.....	136
Paralabrax.....	317	kanawha.....	134
Scarus.....	433	keimi.....	133
Serranus.....	317	kendalli.....	135
humeralmaculatus, <i>Sargus</i>	338	lacertosus.....	129
humile, <i>Pristipoma</i>	334	longiceps.....	133
humilis, <i>Allotis</i>	300	longirostris.....	135
Bryttus.....	300	maculata.....	135
Cottus.....	397	meeki.....	139
Gambusia.....	185	missuriensis.....	133
Myripristes.....	235		

List of scientific names—Continued

	Page		Page
<i>Hybopsis montanus</i>	139	<i>hyostomus, Nocomis</i>	139.
<i>uitida</i>	135	<i>hypacanthus, Psednoblennius</i>	405.
<i>niveus</i>	129	<i>Hypargyrus</i>	144
<i>nocomis</i>	135	<i>Hypeneus</i>	343
<i>nux</i>	135	<i>Hypentelium</i>	107
<i>ortenburgeri</i>	135	<i>macroplectum</i>	107
<i>ozarkanus</i>	135	<i>macropterum</i>	107
<i>phatima</i>	132	<i>nigricans</i>	107
<i>phenacobius</i>	134	<i>Hyperchorisitus</i>	70
<i>piptolepis</i>	134	<i>tanueri</i>	70
<i>procne</i>	133	<i>Hyperistius</i>	304, 305
<i>rubriocroceus</i>	125, 127	<i>carolinensis</i>	305
<i>sabinæ</i>	135	<i>Hyperoartia</i>	8
<i>scylla</i>	135	<i>Hyperotreta</i>	8
<i>shumardi</i>	134	<i>Hyperprosopon</i>	410
<i>simus</i>	135	<i>agassizii</i>	410
<i>spectrunculus</i>	134	<i>analis</i>	410
<i>stramineus</i>	134	<i>arcuatum</i>	410
<i>tetranemus</i>	139	<i>argenteum</i>	410
<i>timpanogensis</i>	119	<i>punctatum</i>	410
<i>tuditanus</i>	144, 145	<i>argenteus</i>	410
<i>volucella</i>	133	<i>Hyphalonedrus</i>	165
<i>watauga</i>	138	<i>chalybeatus</i>	165
<i>winchelli</i>	133	<i>Hypnessoxycon</i>	97
<i>xenoccephalus</i>	127	<i>pnamensis</i>	97
<i>Hyborhynchus</i>	145	<i>Hypeurochilus</i>	461
<i>confertus</i>	145	<i>geminatus</i>	462
<i>(Hybognathus) siderius</i>	140	<i>Hypocaranx</i>	273
<i>nigrellus</i>	145	<i>Hypoclydonia</i>	297
<i>notatus</i>	145	<i>bella</i>	297
<i>perspicuus</i>	145	<i>Hypocritichthys</i>	410
<i>punicus</i>	145	<i>analis</i>	410
<i>superciliosus</i>	145	<i>Hypodus</i> , <i>Dactypterus</i>	270
<i>tonellus</i>	145	<i>Hypophomus</i>	284
<i>Hydargira atricauda</i>	174	<i>aurantiacus</i>	285
<i>lima</i>	174	<i>cymatotaenia</i>	285
<i>Hydrargyreus</i>	175	<i>nianqua</i>	285
<i>diaphana</i>	178	<i>spilotus</i>	285
<i>fusca</i>	174	<i>Hypolophus</i>	28
<i>lima</i>	174	<i>Hypomesus</i>	67
<i>lucie</i>	178	<i>olidus</i>	68
<i>multifaciata</i>	178	<i>pretiosus</i>	67
<i>nigrofasciata</i>	176	<i>hypolecta, Rathbunella</i>	452
<i>ornata</i>	176	<i>Hypolepturus</i>	320
<i>similis</i>	175, 177	<i>aberrans</i>	321
<i>swampina</i>	175	<i>genima</i>	321
<i>trifasciata</i>	176	<i>lamprurus</i>	321
<i>vernalis</i>	176	<i>maculiferus</i>	321
<i>zebra</i>	179	<i>pinnivarius</i>	321
<i>Hydrolagus</i>	33	<i>unicolor</i>	320, 321
<i>colliei</i>	33	<i>aberrans</i>	321
<i>waitei</i>	33	<i>accensus</i>	321
<i>Hydrophlox</i>	126	<i>afinis</i>	321
<i>abbotti</i>	127	<i>bovinus</i>	321
<i>altipinnis</i>	127	<i>chlorurus</i>	321
<i>chalybeus</i>	127	<i>crocotus</i>	321
<i>chiliticus</i>	127	<i>gemma</i>	321
<i>chlorocephalus</i>	127	<i>gummigutta</i>	321
<i>chromosus</i>	127	<i>guttavarius</i>	321
<i>lutipinnis</i>	127	<i>indigo</i>	321
<i>roseus</i>	127	<i>nigricans</i>	321
<i>rubriocroceus</i>	127	<i>pinnivarius</i>	321
<i>stonei</i>	127	<i>puelia</i>	321
<i>xenoccephalus</i>	127	<i>vittulinus</i>	321
<i>hydrophlox, Cheonda</i>	119	<i>hypolectus, Bathymaster</i>	452
<i>Clinostomus</i>	119	<i>Hyppolites</i>	324
<i>hygomi, Myctophum</i>	169	<i>Hypopomus</i>	101
<i>Scopelus</i>	169	<i>brevirostris</i>	101
<i>Hylomyzon</i>	107	<i>occidentalis</i>	101
<i>Hymenocephalus</i>	207	<i>hypothalmus, Arius</i>	151
<i>cavernosus</i>	207	<i>Hypopriion</i>	17
<i>italicus</i>	207	<i>brevirostris</i>	17
<i>Hynnus</i>	274	<i>longirostris</i>	17
<i>cubensis</i>	274	<i>signatus</i>	17
<i>goreensis</i>	274	<i>Hypoprionodon</i>	17
<i>hopkinsi</i>	274	<i>hypothalmus, Cathorops</i>	151
<i>Hyodon</i>	40	<i>hypothalmus, Arius</i>	151
<i>amphiodon</i>	41	<i>Hyporhamphus</i>	198
<i>chrysopsis</i>	41	<i>gilli</i>	198
<i>seleneops</i>	41	<i>roberti</i>	198
<i>vernalis</i>	40	<i>rose</i>	198
<i>Hyostoma</i>	286	<i>snyderi</i>	199
<i>blennioperca</i>	286	<i>tricuspidatus</i>	198
<i>newmani</i>	286	<i>unifasciatus</i>	198
<i>sinoturum</i>	286	<i>Hyorthodus</i>	310
<i>hyostoma, Macrhybopsis</i>	139	<i>flavicauda</i>	311

List of scientific names—Continued

	Page		Page
<i>Hypserranus</i>	310	<i>Icelus hamatus</i>	378
<i>hyposticta</i> , <i>Propterygia</i>	24	<i>scutiger</i>	378
<i>hypostoma</i> , <i>Mobula</i>	32	<i>spatula</i>	379
<i>Hypostomus</i>	160	<i>spiniger</i>	378
<i>brasiliensis</i>	160	<i>uncinialis</i>	379
<i>cirrhosus</i>	160	<i>vicinalis</i>	378
<i>guacari</i>	160	<i>Ichthocolla</i>	33
<i>guacharote</i>	161	<i>Ichthyapus</i>	84
<i>plecostomus</i>	160	<i>ichthyomyzon</i>	9
<i>robinii</i>	160	<i>castaneus</i>	9
<i>hypostomus</i> , <i>Cephalopterus</i>	32	<i>concolor</i>	9
<i>Hypsagonus</i> <i>gradiens</i>	393	<i>fossor</i>	9
<i>swani</i>	396	<i>hirudo</i>	9
<i>hypselosoma</i> , <i>Leptocephalichthys</i>	79	<i>Ichthys</i>	268
<i>hypselurus</i> , <i>Rhamdia</i>	157	<i>lockingtoni</i>	268
<i>hypselurus</i> , <i>Pimelodus</i>	157	<i>Icista</i> , <i>Bairdiella</i>	346
<i>hypselopterus</i> , <i>Leuciscus</i>	129	<i>Scienea</i>	346
<i>Hypsicometes</i>	451	<i>Icosteidae</i>	289
<i>gobioides</i>	451	<i>Icosteus</i>	289
<i>Hypsifario</i>	55	<i>enigmaticus</i>	289
<i>Hypsilepis</i>	123	<i>ictalops</i> , <i>Cottus</i>	385
<i>ardens</i>	126	<i>Pegedictis</i>	385
<i>coccognis</i>	127	<i>Italurus</i>	151
<i>cornutus</i>	128	<i>anguilla</i>	152
<i>cyanus</i>	128	<i>furcatus</i>	151, 152
<i>gibbus</i>	128	<i>kevinski</i>	153
<i>galacturus</i>	129	<i>macrasyei</i>	153
<i>iris</i>	131	<i>meridionalis</i>	152
<i>Hypsinotus</i>	358	<i>oceanoboeensis</i>	152
<i>rubescens</i>	358	<i>punctatus</i>	152
<i>hypsinotus</i> , <i>Ceratichthys</i>	138	<i>robustus</i>	152
<i>Erinimus</i>	138	<i>simpsoni</i>	152
<i>Hypselepis</i>	128	<i>Icteliis</i>	299
<i>Hynophrys</i>	418	<i>erythrops</i>	303
<i>unimaculatus</i>	418	<i>megalotis</i>	300
<i>Hypsopsetta</i>	225	<i>melanops</i>	298
<i>guttulata</i>	225	<i>ichtyloides</i> , <i>Lepomis</i>	303
<i>hypostoma</i> , <i>Mobula</i>	32	<i>Ichthyelurus</i>	151
<i>Hypostomus</i>	160	<i>Ichthyapus</i>	84
<i>plectostomus</i>	160	<i>acutirostris</i>	84
<i>hypostoma</i> , <i>Cephalopterus</i>	32	<i>Ichthyobius</i>	102
<i>Hypsagonus</i>	393	<i>cyanellus</i>	102
<i>gradiens</i>	393	<i>rauchi</i>	102
<i>quadricornis</i>	393	<i>Ichthyobiphs</i>	94
<i>swani</i>	396	<i>vittatus</i>	95
<i>hypselura</i> , <i>Rhamdia</i>	157	<i>Ictiobus</i>	102, 352
<i>Hypsicometes</i>	451	<i>bubalis</i>	102
<i>gobioides</i>	451	<i>uris</i>	102
<i>Hypsifario</i>	54	<i>Idiacanthidae</i>	70
<i>hypsinotus</i> , <i>Erinimus</i>	138	<i>Idiacanthus</i>	70
<i>Hypsoblennius</i>	462	<i>antrostomus</i>	70
<i>gentilis</i>	462	<i>ferox</i>	70
<i>gilberti</i>	462	<i>ignobilis</i> , <i>Caranx</i>	273
<i>hentz</i>	462	<i>Scomber</i>	273
<i>ionthas</i>	462	<i>Ilictis</i>	155
<i>piersoni</i>	462	<i>Ilisha</i>	44
<i>striatus</i>	462	<i>abnormis</i>	44
<i>Hypsopsetta</i>	225	<i>caribbea</i>	45
<i>guttulata</i>	225	<i>Illecebrosus</i> , <i>Zestidium</i>	347
<i>Hypsufus</i>	412	<i>Illecebrosus</i> , <i>Albunrops</i>	344
<i>caryl</i>	412	<i>Stelliferus</i>	347
<i>Hypsypops</i>	415	<i>illistris</i> , <i>Lampogrammus</i>	484
<i>dorsalis</i>	415, 416	<i>Ilyophidae</i>	78
<i>rubicundus</i>	415	<i>Ilyophis</i>	78
<i>Hysteroxarpus</i>	412	<i>brunneus</i>	78
<i>traski</i>	412	<i>Ilypnus</i>	445
<i>hystrix</i> , <i>Diodon</i>	501, 502	<i>gilberti</i>	445
<i>Icelidae</i>	375	<i>imberbe</i> , <i>Ophiodon</i>	467
<i>Icelinus</i>	376	<i>Peristedion</i>	406
<i>borealis</i>	376	<i>imberbis</i> , <i>Apogon</i>	295
<i>burchami</i>	376	<i>Mullus</i>	295
<i>cavifrons</i>	375	<i>Sciena</i>	348
<i>filamentosus</i>	375	<i>Vulsculus</i>	406
<i>fimbriatus</i>	378	<i>imiceps</i> , <i>Ophioscion</i>	348
<i>fuscescens</i>	378	<i>Sciena</i>	348
<i>oculatus</i>	376	<i>immaculata</i> , <i>Amia</i>	40
<i>quadriseriatus</i>	376	<i>Coryphana</i>	284
<i>strabo</i>	376	<i>Fistularia</i>	245
<i>tenuis</i>	376	<i>Perca</i>	307
<i>Icelus</i>	378	<i>immaculatum</i> , <i>Cichlasoma spinosissimum</i>	420
<i>australis</i>	379	<i>Cybium</i>	257
<i>bicornis</i>	378	<i>immaculatus</i> , <i>Archocentrus</i>	420
<i>canaliculatus</i>	378	<i>Esor</i>	173
<i>euryops</i>	378	<i>masquinony</i>	173
<i>furciger</i>	378	<i>Platypterus</i> , <i>maculatus</i>	193
		<i>Salmo</i>	60
		<i>Synbranchus</i>	76

List of scientific names—Continued

	Page		Page
<i>Imostoma</i>	285	<i>insigne</i> , <i>Pimelodus</i>	155
<i>cheneyi</i>	286	<i>insignis</i> , <i>Anoplarchus</i>	488
<i>gilberti</i>	286	<i>Catostomus</i>	105, 106
<i>longimanus</i>	286	<i>Cheirodon</i>	96
<i>meadi</i>	286	<i>Pimelodus</i>	156
<i>podostomone</i>	286	<i>Rabida</i>	156
<i>shumardi</i>	285	<i>insociabilis</i> , <i>Dorosoma</i>	46
<i>imperator</i> , <i>Xiphias</i>	284	<i>insolatus</i> , <i>Hellas</i>	413
<i>imperialis</i> , <i>Coryphaea</i>	284	<i>inspector</i> , <i>Bathypteroctes</i>	52
<i>Esox</i>	197	<i>insulæ-santa-crucis</i> , <i>Scarus</i>	424
<i>Luvaris</i>	264	<i>insularum</i> , <i>Atherinops</i>	232
<i>Sebastes</i>	370	<i>Cedrosensis</i>	252
<i>Zeus</i>	217	<i>Muraena</i>	90
<i>impeditinus</i> , <i>Serranus</i>	310	<i>integripinnis</i> , <i>Auchenipterus</i>	457
<i>implutus</i> , <i>Tetronotus</i>	500	<i>Cremonobates</i>	457
<i>impressus</i> , <i>Conger</i>	80	<i>intermedia</i> , <i>Tigonia</i>	119
<i>inæquilobus</i> , <i>Pogonichthys</i>	116	<i>intermedius</i> , <i>Cichlaurus</i>	416
<i>incili</i> , <i>Querimana</i>	253	<i>Gephyrocharax</i>	96
<i>Incisidens</i>	239	<i>Heros</i>	416
<i>simplicidens</i>	339	<i>Paralepis</i>	165
<i>inceror</i> , <i>Helloperca</i>	301	<i>Pomoxys</i>	304
<i>Kyphosus</i>	339	<i>Saurus</i>	104
<i>Pimelepterus</i>	339	<i>Sudis</i>	165
<i>Pomotis</i>	301	<i>Synodus</i>	164
<i>incisus</i> , <i>Lycodes</i>	475	<i>internasalis</i> , <i>Julis</i>	427
<i>inconstans</i> , <i>Eucalia</i>	238	<i>interrupta</i> , <i>Chrysopera</i>	307
<i>Gasterosteus</i>	238	<i>Morone</i>	307
<i>inconstans cayuga</i> , <i>Eucalia</i>	238	<i>Perca mitchilli</i>	307
<i>incrassatus</i> , <i>Leucosomus</i>	117	<i>Raia</i>	26
<i>indica</i> , <i>Lutodeira</i>	40	<i>Raja</i>	26
<i>indicus</i> , <i>Cottus</i>	397	<i>interruptus</i> , <i>Anisotremus</i>	332
<i>Tetrapterurus</i>	263	<i>Archoplites</i>	303
<i>indigena</i> , <i>Clupea</i>	42	<i>Centrarchus</i>	303
<i>indigo</i> , <i>Hypoplectrus unicolor</i>	321	<i>Gentremus</i>	332
<i>Plectropoma</i>	321	<i>Luxilus</i>	128
<i>Inermia</i>	340	<i>intertinctus</i> , <i>Mystriophus</i>	88
<i>vittata</i>	340	<i>Ophisurus</i>	88
<i>Inermidae</i>	340	<i>interstitialis</i> , <i>Serranus</i>	314
<i>inermis</i> , <i>Anoplagonus</i>	397	<i>Trisopteris</i>	314
<i>Aspidophoroides</i>	397	<i>introneriger</i> , <i>Dicrolene</i>	483
<i>Dermatolepis</i>	313	<i>Eosebastes</i>	366
<i>Elops</i>	39	<i>Sebastichthys</i>	366
<i>Mesopion</i>	327	<i>intronigra</i> , <i>Dicrolene</i>	483
<i>Robirubia</i>	327	<i>inurus</i> , <i>Zygonectes</i>	185
<i>Raia ornata</i>	26	<i>Ioa</i>	288
<i>Scorpena</i>	371	<i>vigil</i>	288
<i>Serranus</i>	313	<i>vitreus</i>	288
<i>infans</i> , <i>Avocettina</i>	83	<i>Ioglossus</i>	438
<i>Gambusia</i>	189	<i>callurus</i>	438
<i>Nemichthys</i>	83	<i>ionthas</i> , <i>Cerdale</i>	473
<i>Pacilopsis</i>	189	<i>Hypsoblennius</i>	462
<i>internalis</i> , <i>Histiobranchus</i>	78	<i>Isasthes</i>	462
<i>Muraena</i>	92	<i>ios</i> , <i>Clevelandia</i>	445
<i>infirmis</i> , <i>Novaculichthys</i>	430	<i>Gobiosoma</i>	445
<i>Xyrichthys</i>	430	<i>iotlichthys</i>	121
<i>infraspinata</i> , <i>Asterotheca</i>	396	<i>phlegethoni</i>	121
<i>infraspinatus</i> , <i>Xeneretmus</i>	396	<i>iowa</i> , <i>Etheostoma</i>	291
<i>ingens</i> , <i>Hippocampus</i>	224	<i>Oligocephalus</i>	291
<i>ingolfi</i> , <i>Macrourus</i>	206	<i>Ipnopidae</i>	170
<i>inlistus</i>	430	<i>Ipnops</i>	170
<i>mundicorpus</i>	430	<i>ugassizi</i>	171
<i>pavonius</i>	430	<i>murnayi</i>	170
<i>Inloml</i>	103	<i>irideus</i> , <i>Labrus</i>	304
<i>innominado</i>	87	<i>Salmo</i>	58
<i>innominatus</i> , <i>Girardichthys</i>	182	<i>iridio</i> , <i>Salmo</i>	58
<i>inopinatus</i> , <i>Gasterosteus</i>	238	<i>caranx</i>	271
<i>inops</i> , <i>Antennarius</i>	605	<i>Iridio</i>	426
<i>inopsetta</i>	226	<i>bivittata</i>	426
<i>ischya</i>	226	<i>caudalis</i>	427
<i>inornata</i> , <i>Lota</i>	212	<i>cyanoccephalus</i>	427
<i>Raja</i>	26	<i>decoratus</i>	427
<i>inornatus</i> , <i>Apodichthys</i>	467	<i>displiis</i>	427
<i>Cryptocanthodes</i>	471	<i>elegans</i>	427
<i>Microlepidotus</i>	335	<i>frenatus</i>	427
<i>Oligolophites</i>	277	<i>garnoti</i>	427
<i>Pseudojulis</i>	428	<i>kirschii</i>	427
<i>inscriptum</i> , <i>Nanostoma</i>	289	<i>maculipinna</i>	427
<i>inscriptus</i> , <i>Achirus</i>	229	<i>meyeri</i>	427
<i>Nothonotus</i>	289	<i>microstomus</i>	427
<i>Pomotis</i>	300	<i>nicholsi</i>	428
<i>insculpta</i> , <i>Netuma</i>	149	<i>pictus</i>	427
<i>insculptus</i> , <i>Arius</i>	149	<i>poeyi</i>	427
<i>Ctenolucius</i>	100	<i>radiata</i>	428
<i>Gasterosteus</i>	237	<i>sellifer</i>	428
<i>Luciobarbus</i>	100	<i>semicinctus</i>	427
<i>insignarius</i> , <i>Pimelodus</i>	155, 156	<i>similis</i>	427

List of scientific names—Continued

	Page		Page
Irrillion.	66	jagua, <i>Histro</i> .	505
oregonius.	66	Jaguacanguare.	415
iris, <i>Hypsilepis</i> .	131	jaguar, <i>Bodianus</i> .	235
Leuciscus.	117	jalape, <i>Xiphophorus</i> .	103
irradians, <i>Serranus</i> .	319	jamaicensis, <i>Cynoscion</i> .	355
irregularis, <i>Theraps</i> .		Otolithus.	355
irretitus, <i>Gordiichthys</i> .	417	Raja.	29
Mugil.	89	janeiro, <i>Engraulis</i> .	51
irroratus, <i>Monacanthus</i> .	254	januaria, <i>Umbrina</i> .	351
irroratus, <i>Monacanthus</i> .	494	jaok, <i>Cottus</i> .	387
isabelita.	361	Myoxocephalus.	387
isabelita, <i>Angelichthys</i> .	361	japonicus, <i>Arotocopus</i> .	357
Isaciella.	335	Caranx trachurus.	270
ischana, <i>Anchoviella</i> .	48	Chiropodus.	466
ischanus, <i>Notemigonus</i> .	115	Salmo.	54
Stolephorus.	48	Scomber.	258
Ischnomembras.	247	Trachurus.	270
gabonensis.	247	Trichodon.	357
gabunensis.	247	jarrovii, <i>Lepidomedra</i> .	147
ischanus, <i>Notemigonus</i> .	115	Minomus.	104
Ischnomembras.	247	Javanicus, <i>Psenes</i> .	268
ischyra, <i>Inopsetta</i> .	226	jejunn, <i>Episeima</i> .	125
ischyra, <i>Apomotis</i> .	299	jejunus, <i>Paranotropis</i> .	125
Lepiopomus.	299	jemenezanus, <i>Alburnellus</i> .	125
Parophrys.	223	jeniquano, <i>He-mulon</i> .	331
Isesthes.	462	Jenkinsi, <i>Chasmodes</i> .	403
gilberti.	462	Synodus.	163
ionthas.	462	Zygonectes.	178
scrutator.	462	Jenkinsia.	45
ishinagi, <i>Megapercra</i> .	308	Iamprotinia.	46
islandicus, <i>Bogmarus</i> .	216	stolifera.	46
Centronotus.	471	jessiae, <i>Oligocephalus</i> .	291
Gasterosteus.	238	Poeciliichthys.	291
Isodon, <i>Aprionodon</i> .	15	Xyrichtys.	429
Carcharius.	15	Xyrula.	429
Mesoprion.	320	Jouc, <i>Anthias</i> .	325
isodus, <i>Squalus</i> .	20	Lutianus.	325
Isogomphodon.	15	Johannae, <i>Argyrosomus</i> .	63
maculipinnis.	16	Leucichthys.	63
isolepis, <i>Isopsetta</i> .	226	johni, <i>Cephaloptra</i> .	32
Lepidopsetta.	226	Johnius batabanus.	345
Luxilus cornutus.	128	guttatus.	313
Notropis cornutus.	128	nobilis.	356
Sternotremia.	215	regalis.	356
Isopisthus.	353	saxatilis.	350
affinis.	353	jonesi, <i>Belone</i> .	107
barroweri.	354	Mollienisia.	188
parvipinnis.	353	Syngnathus.	241
remifer.	353	jonesii, <i>Gerres</i> .	341
Isoplagonodon.	15	Pseudoxiphophorus bimaculatus.	188
Isopsetta.	226	Jordanella.	180
isolepis.	225	floridei.	180
Isoppondyli.	39	Jordanini, <i>Bathymaster</i> .	452
Isthmennis, <i>Muraenichthys</i> .		Caulophryne.	511
Bagre.	148	Chiromystoma.	250
Preciliopsis.	148	Eopsetta.	222
Rivulus.	190	Epinephelus.	314
Istiophoridae.		Etheostoma (<i>Nothonotus</i>)	290
Istiophorus.	263	Galeichthys.	150
americanus.	263	Oillas.	458
greyi.	263	Gymnothorax.	92
maguirei.	263	Hemilepidotus.	381
volador.	263	Hippoglossoides.	222
wrighti.	263	Hybopsis.	136
istiana, <i>Parapetenia</i> .	418	Lampanyctus.	107
istianus, <i>Heros</i> .	418	Lutianus.	324
Istiarus.	151	Lycodes.	474
balsanus.	151	Lyconodontis.	92
Isuropsis.	19	Neonemis.	324
dekayi.	19	Nothonotus.	290
Isurus.	19	Notropis.	136
oxyrinchus.	19	Pantosteus.	104
punctatus.	19	Rain.	26
spallanzani.	19	Ronquilus.	452
tigris.	19	Salmo.	57
Italara.	312	clarkii.	57
italara, <i>Promicrops</i> .	312	Sebastodes.	364
Serranus.	312	Tachisurus.	150
Jaboncillo.	316	Trisoptropis.	314
jacksoni, <i>Embiotoca</i> .	411	Jordanina.	375
jacobsæ, <i>Echeneis</i> .	449	zonope.	375
jacobi, <i>Corvina (Johnius)</i> .	348	Jordanidae.	375
jacobus, <i>Myripristes</i> .	235	josephi, <i>Ophidium</i> .	485
jacome, <i>Serranus</i> .	319	Joturus.	254
jaculidens, <i>Acanthodon</i> .	354	globiceps.	255
jaculus, <i>Alburnellus</i> .	123	pichardi.	254
		stipes.	255

List of scientific names—Continued

	Page		Page
<i>juba</i> , <i>Guatucupa</i>	332	<i>kunzingeri</i> , <i>Achirus</i>	220
<i>Perca</i>	332	<i>Solea</i>	220
<i>jugalis</i> , <i>Moniana</i>	131	<i>kneri</i> , <i>Pristipoma</i>	335
<i>Jugulares</i>	450	<i>kraussi</i> , <i>Gobius</i>	441
<i>juliiæ</i> , <i>Claricola</i>	292	<i>Kryptophaneron</i>	234
<i>Julidio</i>	428	<i>alfredi</i>	234
<i>adustus</i>	428	<i>Kuhliidae</i>	305
<i>notospilus</i>	428	<i>Kuhlia arga</i>	305
<i>Julis</i>	428	<i>kumlienii</i> , <i>Uranidea</i>	385
<i>caudalis</i>	427	<i>Kyphosidae</i>	339
<i>cinctus</i>	427	<i>Kyphosus</i>	339
<i>crotaphus</i>	426, 427	<i>analogus</i>	339
<i>cyahostigma</i>	420	<i>elegans</i>	339
<i>detersor</i>	420	<i>incisor</i>	339
<i>dimidiatus</i>	427	<i>lutescens</i>	340
<i>garnoti</i>	427	<i>metzelaari</i>	339
<i>gillianus</i>	420	<i>secatrix</i>	340
<i>humeralis</i>	426	<i>labarcae</i> , <i>Chirostoma</i>	250
<i>internasalis</i>	427	<i>Labeo elegans</i>	109
<i>lucasanus</i>	429	<i>labialis</i> , <i>Fundulus</i>	174
<i>maculipinna</i>	427	<i>Profundulus</i>	174
<i>melanochir</i>	427	<i>labiatum</i> , <i>Heros</i>	422
<i>modestus</i>	428	<i>labiatus</i> , <i>Amphilophus</i>	422
<i>nitida</i>	429	<i>Catostomus</i>	106
<i>nitidissima</i>	429	<i>Labichthys</i>	84
<i>opalina</i>	426	<i>bowersi</i>	84
<i>patatus</i>	426	<i>carinatus</i>	84
<i>pictus</i>	427	<i>elongatus</i>	83
<i>principis</i>	426	<i>gilli</i>	83
<i>ruptus</i>	427	<i>Labidesthes</i>	251
<i>semininctus</i>	427	<i>siccarius</i>	251
<i>kalisheræ</i> , <i>Eretis</i>	460	<i>lablosus</i> , <i>Carpoides</i>	103
<i>kanloops</i> , <i>Oncorhynchus</i>	58	<i>labradoricus</i> , <i>Acanthocottus</i>	388
<i>kanawha</i> , <i>Hybopsis</i>	134	<i>Coregonus</i>	64
<i>Notropis</i>	134	<i>Labrax</i>	373
<i>kanseæ</i> , <i>Fundulus</i>	170	<i>albidus</i>	307
<i>Plancterus</i>	170	<i>clathratus</i>	317
<i>karrak</i> , <i>Anarrhichas</i>	472	<i>decaenamus</i>	373
<i>Kathetostoma</i>	454	<i>lagocephalus</i>	374
<i>albigutta</i>	454	<i>monopterygius</i>	374
<i>averruncus</i>	454	<i>multilineatus</i>	307
<i>Katsuwondæ</i>	238	<i>nebulifer</i>	317
<i>Katsuwonus</i>	258	<i>nigricans</i>	308
<i>pelamis</i>	258	<i>notatus</i>	307
<i>vagans</i>	258	<i>octogrammus</i>	373, 374
<i>kaupi</i> , <i>Physiculus</i>	211	<i>osculatus</i>	307
<i>kaupi</i> , <i>Synaphobranchus</i>	78	<i>superciliosus</i>	374
<i>kayko</i> , <i>Salmo keta vol.</i>	54	<i>Labridæ</i>	424
<i>kefml</i> , <i>Hybopsis</i>	133	<i>labridens</i> , <i>Heros</i>	410
<i>Notropis</i>	133	<i>Parapetenia</i>	419
<i>kendallii</i> , <i>Calamus</i>	337	<i>labridum</i> , <i>Hæmulum</i>	332
<i>Hybopsis</i>	135	<i>labriformis</i> , <i>Epinephelus</i>	311
<i>Sphagebranchus</i>	84, 85	<i>Serranus</i>	311
<i>Vermæ</i>	85	<i>Labrisomus</i>	450
<i>kennedyi</i> , <i>Trachinotus</i>	277	<i>biguttatus</i>	460
<i>kenneryi</i> , <i>Moxostoma</i>	109	<i>bucciferus</i>	459
<i>Oncorhynchus</i>	55	<i>heilneri</i>	459
<i>Salmo</i>	55	<i>herminier</i>	459
<i>kennicotti</i> , <i>Acipenser</i>	34	<i>lentiginosus</i>	459
<i>Catotonus</i>	203	<i>microlepidotus</i>	459
<i>Coregonus</i>	65	<i>nuchipinnis</i>	459
<i>Prosopium</i>	65	<i>xanti</i>	459
<i>Kenoza</i>	173	<i>Labroperca</i>	310
<i>kentuckiensis</i> , <i>Luxilis</i>	137	<i>Labrosomus bucciferus</i>	459
<i>Nocomis</i>	137	<i>cremnobates</i>	456
<i>kessleri</i> , <i>Artus</i>	140	<i>microlepidotus</i>	459
<i>Netuma</i>	149	<i>labrosus</i> , <i>Ceratichthys</i>	138
<i>keta</i> , <i>Oncorhynchus</i>	54	<i>Erinemus</i>	138
<i>kevinskii</i> , <i>Ictalurus</i>	153	<i>Zoarces</i>	473
<i>killifish</i> , <i>Cobitis</i>	176	<i>Labrus adspersus</i>	424
<i>kincaidi</i> , <i>Malacocottus</i>	388	<i>americanus</i>	424
<i>kirshii</i> , <i>Iridio</i>	427	<i>auritus</i>	299
<i>kirtlandi</i> , <i>Acipenser</i> (<i>Huso</i>).....	35, 37	<i>biuncinatus</i>	429
<i>laciniata</i>	248	<i>torquatus</i>	420
<i>vagrana</i>	248	<i>bivittatus</i>	420
<i>klutwich</i> , <i>Oncorhynchus</i>	54	<i>brasiliensis</i>	426
<i>Salmo</i>	54, 55	<i>carolinus</i>	424
<i>kitt</i> , <i>Micromystomus</i>	228	<i>chogoset</i>	424
<i>Pleuronectes</i>	228	<i>fulva</i>	424
<i>klyli</i> , <i>Leucichthys</i>	64	<i>cromis</i>	352
<i>klamathensis</i> , <i>Agosia</i>	141	<i>cruentatus</i>	322
<i>Apocope</i>	141	<i>cyanocephalus</i>	427
<i>Cottus</i>	384	<i>exoletus</i>	424
<i>kleinii</i> , <i>Balistes</i>	490	<i>falcatus</i>	276
		<i>fulvo-maculatus</i>	335

List of scientific names—Continued

	Page		Page
<i>Labrus fulvus</i> .	309	<i>Lagochila lacera</i> .	112
<i>gauza</i> .	310	<i>Lagodon</i> .	337
<i>griseus</i> .	325	<i>rhomboides</i> .	337
<i>hexagrammus</i> .	374	<i>lahontan</i> , <i>Pantosteus</i> .	105
<i>hiatula</i> .	424	<i>lalandi</i> , <i>Carcharias</i> .	15
<i>irideus</i> .	304	<i>Scollodon</i> .	15
<i>macropterus</i> .	304	<i>Seriola</i> .	279
<i>maximus</i> .	424	<i>lamellifer</i> , <i>Exocetus</i> .	200
<i>ornatus</i> .	424	<i>Lamia</i> .	20
<i>pallidus</i> .	299	<i>lamia</i> , <i>Carcharias</i> .	15
<i>pentacanthus</i> .	424	<i>lamarii</i> , <i>Acipenser</i> .	35
<i>plumieri</i> .	330	<i>Lepidosteus</i> .	37
<i>psittacus</i> .	426	<i>lamella</i> , <i>Carcharias</i> .	15
<i>pulcher</i> .	426	<i>Lamiopsis</i> .	15
<i>radiatus</i> .	426	<i>Lamna</i> .	19
<i>rufus</i> .	425	<i>caudata</i> .	16
<i>salmoides</i> .	298	<i>nasus</i> .	19
<i>semiruber</i> .	425	<i>punctata</i> .	19
<i>sparoides</i> .	305	<i>Lamnidae</i> .	19
<i>squamiceps</i> .	355	<i>lamotteti</i> , <i>Okkelbergia</i> .	10
<i>maculatus</i> .	355	<i>Petromyzon</i> .	10
<i>striatus</i> .	317	<i>Lampadena</i> .	167
<i>subfuscus</i> .	424	<i>speculigera</i> .	187
<i>tassellatus</i> .	424	<i>Lamppanyctus</i> .	187
<i>tautoga</i> .	424	<i>alatus</i> .	187
<i>versicolor</i> .	336	<i>gemmifer</i> .	187
<i>lacera</i> , <i>Lagochila</i> .	112	<i>günterii</i> .	187
<i>lacerta</i> , <i>Lamppanyctus</i> .	167	<i>lacerta</i> .	187
<i>Synodus</i> .	163	<i>ritteri</i> .	187
<i>lacertinus</i> , <i>Synodus</i> .	164	<i>Lampetra</i> .	10
<i>lacertosus</i> , <i>Ilyobops</i> .	129	<i>ayresi</i> .	10
<i>Luxilus</i> .	129	<i>borealis</i> .	10
<i>lacertus</i> , <i>Gobius</i> .	440	<i>spadicea</i> .	9
<i>Scomber</i> .	258	<i>wilderi</i> .	9
<i>Lachnoltemus</i> .	425	<i>Lampetraformis</i> , <i>Blennius</i> .	470, 471
<i>Lachnolaimus</i> .	425	<i>Lumpenus</i> .	471
<i>aigula</i> .	425	<i>Lampreda</i> .	10
<i>caninus</i> .	425	<i>Lampridae</i> .	216
<i>dux</i> .	425	<i>Lampris</i> .	216
<i>maximus</i> .	425	<i>lauta</i> .	217
<i>psittacus</i> .	425	<i>regius</i> .	216
<i>sullus</i> .	425	<i>Lamprogrammus</i> .	484
<i>lachrymalis</i> , <i>Ptychostomus</i> .	111	<i>illistris</i> .	484
<i>lacinata</i> , <i>Merluca vulgaris</i> .	248	<i>lamproteenia</i> , <i>Clupea</i> .	46
<i>laciniata</i> , <i>Kirtlandia</i> .	248	<i>Jenkinsia</i> .	46
<i>lacrimosus</i> , <i>Scarus</i> .	432	<i>lamprurus</i> , <i>Hypoplectrus</i> .	321
<i>lactarius</i> , <i>Catostomus</i> (<i>Aconius</i>).	107	<i>Serranus</i> .	321
<i>Lactophrys</i> .	497	<i>Lampugus</i> .	284
<i>trigonus</i> .	497	<i>neopolitanus</i> .	265
<i>lacustris</i> , <i>Algansea</i> .	116	<i>punctulatus</i> .	265
<i>Gadus</i> .	152	<i>sculus</i> .	265
<i>Haustor</i> .	212	<i>lanatus</i> , <i>Merluccius</i> .	214
<i>Hemiplus</i> .	152	<i>lanceolata</i> , <i>Perca</i> .	352
<i>okeechobeensis</i> , <i>Amelurus</i> .	115	<i>lanceolatus</i> , <i>Chætodon</i> .	352, 353
<i>Pomolobus pseudoharengus</i> .	152	<i>Equetus</i> .	353
<i>Leuciscus</i> .	21	<i>Gobius</i> .	441
<i>Læmonema</i> .	213	<i>Homopriion</i> .	346
<i>barbatula</i> .	213	<i>Lonchirurus</i> .	352
<i>barbatulum</i> .	213	<i>Stellifer</i> .	346
<i>gracillipes</i> .	213	<i>lappa</i> , <i>Italiutella</i> .	508
<i>melanurum</i> .	213	<i>Laproporca</i> .	310
<i>Latrabialis</i> , <i>Moniana</i> .	131	<i>Larinus</i> .	344
<i>latetus</i> , <i>Centronotus</i> .	468	<i>acclivis</i> .	344
<i>lavicaudatus</i> , <i>Hippocampus</i> .	244	<i>argenteus</i> .	344
<i>levigatus</i> , <i>Callichthys</i> .	160	<i>breviceps</i> .	344
<i>Chitronectes</i> .	505	<i>aufulgens</i> .	344
<i>Lagocephalus</i> .	497	<i>fasciatus</i> .	344
<i>Salmo</i> .	61	<i>pacificus</i> .	344
<i>Tetraodon</i> .	407	<i>larkinsi</i> , <i>Cymatogaster</i> .	410
<i>Leviraja</i> .	24	<i>Lasiancistrus</i> .	161
<i>levis</i> , <i>Acipenser</i> .	35	<i>planiceps</i> .	161
<i>Balistes</i> .	495	<i>Lasionognathus</i> .	510
<i>Gasterosteus</i> .	239	<i>saccostoma</i> .	510
<i>Mustelus</i> .	13	<i>lata</i> , <i>Parapetenia tetricantha</i> .	419
<i>Raja</i> .	25	<i>Xurel</i> .	272
<i>Squatina</i> .	22	<i>Latebrus</i> .	297
<i>variegatus</i> , <i>Orbis</i> .	499	<i>oculatus</i> .	297
<i>lagenarius</i> , <i>Acipenser</i> .	36	<i>lateplicatus</i> , <i>Serranus</i> .	314
<i>Lagocephalus</i> .	407	<i>lateralis</i> , <i>Alvarius</i> .	294
<i>levigatus</i> .	497	<i>Artedius</i> .	377
<i>pachycephalus</i> .	497	<i>Characodon</i> .	183
<i>lagocephalus</i> , <i>Hexagrammos</i> .	374	<i>Embletoca</i> .	411
<i>Labrax</i> .	374	<i>Gobiomorus</i> .	436
<i>Salmo</i> .	54	<i>Mylochelus</i> .	112
<i>Lagochila</i> .	112	<i>Philypnus</i> .	436
		<i>Richardsonius</i> .	118

List of scientific names—Continued

	Page		Page
<i>lateralis</i> , <i>Scarus</i>	432	<i>lelopomus</i> , <i>Cottus</i>	385
<i>Scorpaenichthys</i>	377	<i>lelops</i> , <i>Xeneretmus</i>	398
<i>Tæniotoca</i>	411	<i>Xenopyxis</i>	398
<i>Zygoneutes</i>	177	<i>Leiostomus</i>	348
<i>laterna</i> , <i>Tetodon</i>	500	<i>humeralis</i>	349
<i>laternatum</i> , <i>Myctophum</i>	170	<i>lineatus</i>	349
<i>laternatus</i> , <i>Polyipnus</i>	74	<i>nanthurus</i>	349
<i>laterum albo</i> , <i>Antennarius</i>	505	<i>Leiurus</i>	237
<i>lathamii</i> , <i>Trachurus</i>	270	<i>leiurus</i> , <i>Gasterosteus</i>	237
<i>laticauda</i> , <i>Anguilla</i>	77	<i>omarii</i> , <i>Acipenser (Huso)</i>	35
<i>Rhamphus</i>	157	<i>Lombus</i>	435
<i>aticeaudus</i> , <i>Pimelodus</i>	157	<i>maculatus</i>	435
<i>Scoliodon</i>	14	<i>lemonni</i> , <i>Squalius</i>	119
<i>laticeps</i> , <i>Acanthocottus</i>	388	<i>lemniscatus</i> , <i>Engraulis</i>	48
<i>Ætobatus</i>	30	<i>Osmerus</i>	163
<i>Arrius</i>	150	<i>Lemnismora</i>	261
<i>Atherina</i>	246	<i>Lencoraja</i>	24
<i>Bathynectes</i>	483	<i>lentiginosa</i> , <i>Muraena</i>	90
<i>Megalocottus</i>	388	<i>lentiginosum</i> , <i>Ostracion</i>	496
<i>Mixonius</i>	483	<i>lentiginosus</i> , <i>Ania</i>	38
<i>latidens</i> , <i>Gliridodon</i>	189	<i>Galeichthys</i>	149
<i>Microstomus</i>	228	<i>Heros</i>	418
<i>Poecilopsis</i>	189	<i>Labrisomus</i>	459
<i>latifasciatus</i> , <i>Cyprinodon</i>	181	<i>Rhinobatos</i>	23
<i>Electris</i>	437	<i>Rhinobatoides</i>	23
<i>Hemicleotris</i>	437	<i>Tachisurus</i>	149
<i>latifrons</i> , <i>Anarrhichthys</i>	472	<i>Theraps</i>	418
<i>Citharichthys</i>	213	<i>leoniensis</i> , <i>Oligocephalus</i>	291
<i>Electris</i>	436	<i>leonia</i> , <i>Mormina</i>	131
<i>Lycichthys</i>	472	<i>leopardinus</i> , <i>Antennarius</i>	705
<i>Paraliparis</i>	405	<i>Platophrys</i>	218
<i>Stegidionotus</i>	379	<i>Rhomboidichthys</i>	218
<i>Syacium</i>	218	<i>leopardus</i> , <i>Anarrhichthys</i>	472
<i>Xenochirus</i>	396	<i>Lepadogaster mæandricus</i>	488
<i>Xenopylax</i>	396	<i>reticulatus</i>	488
<i>Latilus chrysops</i>	357	<i>testar</i>	489
<i>princeps</i>	357	<i>Lophodolus indicus</i>	509
<i>latimaculatus</i> , <i>Ophisurus</i>	85	<i>Lepibrama</i>	307
<i>Latimana</i> , <i>Belone</i>	197	<i>chrysops</i>	307
<i>latimasutus</i> , <i>Macrourus</i>	206	<i>epida</i> , <i>Boleosoma</i>	291
<i>Macrurus</i>	206	<i>Cyprinella</i>	130
<i>lator</i> , <i>Salmo</i>	64	<i>lepidostole</i> , <i>Anchovia</i>	49
<i>latipinna</i> , <i>Mollinessa</i>	192	<i>Anchoviella</i>	49
<i>latipinnis</i> , <i>Catostomus</i>	107	<i>Lepidia</i>	211
<i>Zanolepis</i>	375	<i>Lepidion</i>	211
<i>latipunctata</i> , <i>Gambusia</i>	187	<i>vereendum</i>	211
<i>latirostratus</i> , <i>Macrourus</i>	206	<i>Lepidochætodon</i>	359
<i>Macrurus</i>	206	<i>lepidogenys</i> , <i>Etheostoma</i>	292
<i>laitura</i> , <i>Loricaria</i>	162	<i>Nivicolia</i>	292
<i>Loricaria filamentosa</i>	162	<i>Lepidogobius</i>	444
<i>atus</i> , <i>Caranx</i>	272	<i>giberti</i>	445
<i>Euctenogobius</i>	442	<i>lepidus</i>	444
<i>Heros tetricanthus</i>	419	<i>Lepidolepus trachyrincus</i>	203
<i>Scomber</i>	276	<i>Lepidolepus norvegicus</i>	205
<i>laurea</i> , <i>Astheros</i>	421	<i>Lepidomedra</i>	147
<i>Clinlusoma</i>	421	<i>arrovi</i>	147
<i>laurettæ</i> , <i>Coregonus</i>	62	<i>vittata</i>	147
<i>Leucichthys</i>	62	<i>Lepidomeganas</i>	279
<i>lauta</i> , <i>Lampris</i>	217	<i>Lepidopsetta</i>	226
<i>Lavinia</i>	113	<i>billinata</i>	226
<i>conformis</i>	120	<i>isolepis</i>	226
<i>crassicauda</i>	120	<i>Lepidopus</i>	262
<i>exilicauda</i>	113	<i>caudatus</i>	262
<i>harengus</i>	113	<i>elongatus</i>	262
<i>leachi</i> , <i>Clupea</i>	41	<i>gouani</i>	262
<i>leachianus</i> , <i>Thynnus</i>	258	<i>xantusi</i>	262
<i>Lebias ellipsoidea</i>	181	<i>Lepidosteus</i>	36
<i>rhomboidalis</i>	181	(Atractosteus) <i>berlandieri</i>	38
<i>Lebistes</i>	191	<i>ayresi</i>	37
<i>paciloides</i>	191	<i>clintonii</i>	37
<i>reticulatus</i>	191	<i>copel</i>	37
<i>Lebius</i>	374	<i>crassus</i>	39
<i>lebranchus</i> , <i>Mugil</i>	253	<i>elizabeth</i>	37
<i>lecontii</i> , <i>Acipenser</i>	34	<i>grayi</i>	37
<i>lecurus</i> , <i>Naucrates</i>	278	<i>harlanii</i>	37
<i>lei</i> , <i>Sympodus</i>	291	<i>horatii</i>	37
<i>lesroyi</i> , <i>Diplapterus</i>	341	<i>lamarli</i>	37
<i>Uliemata</i>	341	<i>latirostris</i>	37
<i>Lesroyia</i>	488	<i>leptorhynchus</i>	37
<i>bermudensis</i>	488	<i>lesuerii</i>	37
<i>lelarchus</i> , <i>Cynoscion</i>	355	<i>louisianensis</i>	38
<i>Otolithus</i>	355	<i>manjuei</i>	38
<i>Lelobatus</i> , <i>sloani</i>	20	<i>milberti</i>	37
<i>Lelocottus</i>	390	<i>otarius</i>	37
<i>hirundo</i>	390	<i>piquitanus</i>	37
<i>Lelodon</i>	21	<i>semiradiatus</i>	36
<i>lelonothos</i> , <i>Holocanthus</i>	499	<i>smithi</i>	37

List of scientific names—Continued

	Page		Page
<i>Lepidosteus thompsoni</i>	37	<i>Leptocephalus spallanzani</i>	79
troosti	37	<i>leptocephalus</i> , <i>Ceratichthys</i>	137
<i>lepidulus</i> , <i>Alburnus</i>	125	Merlangus	209
<i>lepidus</i> , <i>Gobius</i>	444	<i>Leptocerdale</i>	473
<i>Lepidogobius</i>	444	longipinnis	473
<i>Oligocephalus</i>	291	<i>Leptochilichthys</i>	53
<i>Lepiopomus ischyurus</i>	299	ngassizii	53
<i>miniatius</i>	300	<i>Leptoclinus</i>	470
<i>Lepisoma</i>	459	maculatus	470
cirrhosum	459	<i>Leptoconger</i>	82
<i>Lepisosteidae</i>	36	pediongus	82
<i>Lepisosteus</i>	36	<i>Leptocottus</i>	390
<i>nibus</i>	37	armatus	390
(<i>Actrostosteus</i>) <i>ferox</i>	38	<i>Leptodes</i>	71
<i>bison</i>	37	<i>Leptoherasfer</i>	486
(<i>Cylindrostoeus</i>) <i>oculatus</i>	37	macrurus	486
<i>gavialis</i>	36	<i>Leptogunnellus</i>	470
<i>huronensis</i>	36	gracilis	470
<i>lineatus</i>	37	<i>Leptolucania</i>	174
<i>longirostris</i>	36	ommata	175
<i>osseus</i>	36	<i>Leptophidium cervinum</i>	484
<i>oxyurus</i>	37	emmelas	484
<i>platostomus</i>	37	· marmoratum	484
<i>semiradiatus</i>	36	microlepis	485
<i>spatula</i>	33	pardale	485
<i>Lepodus</i>	265	profundorum	484
<i>suragis</i>	265	prorates	484
<i>Lepomis</i>	299	<i>Leptostomias</i>	485
<i>anagallinus</i>	300	<i>Leptophidium</i>	436
<i>apiatus</i>	299	fluvialis	436
<i>appendix</i>	299	<i>Leptops</i>	155
<i>ardescacus</i>	301	<i>Leptopus</i>	215
<i>auritus</i>	299	peregrinus	215
<i>charybdis</i>	303	<i>Leptorhaphis</i>	189
<i>cyanellus</i>	293	<i>Leptorhinophis</i>	87
<i>euryorus</i>	299	<i>Leptorhynchus</i>	83
<i>flexuolaris</i>	298	leuchtenbergii	83
<i>garmani</i>	300	<i>leptorhynchus</i> , <i>Lepidosteus</i>	37
<i>gillii</i>	303	<i>Odontopyxis</i>	395
<i>haplognathus</i>	301	<i>Sarrtorus</i>	395
<i>icthoeloides</i>	303	<i>Syngnathus</i>	240
<i>llrus</i>	302	<i>Leptoscarus</i>	431
<i>longispinis</i>	301	xenodon	431
<i>macrochirus</i>	301	<i>leptosomus</i> , <i>Luxilus</i>	115
<i>miniatius</i>	300	<i>Lepturus</i>	262
<i>mystacalis</i>	300	<i>lepturus</i> , <i>Anarrhichas</i>	472
<i>nephelus</i>	301	Caranx	272
<i>notata</i>	298	Macrourus	207
<i>occidentalis</i>	301	Trichiurus	202
<i>ophthalmicus</i>	299	<i>ierme</i> , <i>Aztecula</i>	143
<i>pallida</i>	298	Chiostoma	251
<i>paillidus</i>	299	Notropis	143
<i>peltastes</i>	300	Skiffia	183
<i>purpurascens</i>	301	<i>Lermilichthys</i>	183
<i>salmonina</i>	298	multiradiatus	183
<i>solis</i>	300	les Alutères	495
<i>symmetrius</i>	299	Curimates	100
<i>trifasciata</i>	298	Lottes	212
<i>Lepophidium</i>	484	Musteles	208
<i>brevibarbe</i>	484	Ovoides	499
<i>cervinum</i>	484	Spheroïdes	498
<i>emmelas</i>	484	Stellifères	340
<i>marmoratum</i>	484	lessonii, <i>Coryphæna</i>	265
<i>microlepis</i>	485	Lestidiops	166
<i>pardale</i>	485	sphyraenops	166
<i>profundorum</i>	484	lesueuri, <i>Acipenser</i>	34
<i>prorates</i>	484	Catostomus	111
<i>stigmatistium</i>	485	Moxostoma	111
<i>leptacantha</i> , <i>Rabida</i>	155	lesueurianum, <i>Exoglossum</i>	146
<i>leptacanthus</i> , <i>Noturus</i>	155	lesueuri, <i>Lepidosteus</i>	37
<i>Leptagonus</i>	394	Lotharchus	85
<i>decagonus</i>	394	pacificus	85
<i>Leptarius</i>	148	velifer	86
<i>dowii</i>	148	Lethenteron	9
<i>Leptecheneis</i>	448	appendix	447
<i>Leptoancistrus</i>	161	connectens	447
<i>canescens</i>	161	lethostigma, <i>Paralichthys</i>	224
<i>Leptoblemnus</i>	470	lethostigmus, <i>Paralichthys</i>	224
<i>Leptocardii</i>	7	Lethostole	250
<i>Leptocephalichthys</i>	79	Lethotremus	398
<i>Leptocephalida</i>	79	muticus	398
<i>Leptocephalus</i>	54, 79	vinolentus	398
<i>candidissimus</i>	79	lethrinus, <i>Astatheros</i>	420
<i>gracilis</i>	79	Clethlasoma	420
(<i>Microconger</i>) <i>caudalis</i>	81	letonal, <i>Priapichthys</i>	199
<i>morrisi</i>	54, 79		

List of scientific names—Continued

	Page		Page
leuchtenbergii, Belonopsis	83	leucopsarum, Myctophum (Stenobrachius)	168
Leptorhynchus	83	Nanobrachium	168
Leucichthys	61	leucopus, Photogenis	130
alascanus	62	Rhamphoberyx	235
alpeno	63	leucorhinus, Rhynchoseratus	511
artedi	63	leucorhynchus, Rhinobatos	24
bissell	63	Rhinobatus	24
athabascae	62	leucorus, Eupomacentrus	414
birgel	64	Pomacentrus	414
cyanopterus	63	Leucosomus	116, 117
entomophagus	63	cataractus	117
eriensis	63	corporalis	117
gemmifer	63	dissimilis	137
harengus	62	incrassatus	117
hoyi	64	occidentalis	113
johannae	63	pallidus	117
klyli	63	rhotheus	117
laurettæ	62	leucosteus, Calamus	337
lucidus	62	leucostictus, Eupomacentrus	413
macrognathus	62	Pomacentrus	413
manitoulinus	64	leucotænia, Pholidichthys	465
nigripinnis	63	leucurus, Caranx	271
nipigon	64	Heunicarax	271
optaricensis	62	Leucus dimidiatus	122
osmeriformis	62	olivaceus	122
prognathus	63	Leuresthes	251
pusillus	62	crameri	251
reighardi	63	tonus	251
sisco	62	Leuroglossus	68
huronius	62	stilbius	68
supernas	63	Leuryrhnis	474
tullibee	64	paucidens	474
zenithicus	64	levius, Hispanicus	368
Leucicorvus	483	Sebastichthys	368
lusciosus	483	lewini, Zygonem	18
Leucidius	123	lewis, Squatina	22
pectinifer	123	lewisi, Salar	55, 56
leucodus, Notropis	124	Salmo	66
Photogenis	124	iberiensis, Balistes	492
Leuciscus	117, 119	libertate, Meletta	44
argenteus	117	Opisthonema	44
boscii	115	Lichia quiebra	277
boucardi	132	lightfooti, Alphestes	313
bubalinus	130	Epinephelus	313
burtonianus	121	ligulata, Seriola	280
carletoni	116	Liza barbuda	496
caurinus	112, 118	colorado	494
chrysopterus	117	Lile	44
copii	123	stolifera	44
croceus	140	lima, Holothrichys	235
evermanni	118	Hydrargira	174
frontalis	128	Loricaria	162
gardoneus	115	Myripristic	235
gilli	118	Rineloricaria	162
gracilis	129	Lima-murena	90
hypselopterus	120	melanotis	90
iris	117	Limanda	226
longirostris	120	aspera	227
lutrensis	131	beanii	227
macrolepidotus	116	ferruginea	227
nachtriebeli	116	proboscidea	227
nasutus	139	limanda, Pleuronectes	222
nitidus	117	Himantoides, Pleuronectes	222
obesus	128	limbatus, Carcharias	16
productus	120	Carcharias (Prionodon)	16
prolixus	146	Conger	78
pulchelloides	117	Oxydontaichthys	88
pulchellus	117	limi, Fundulus	178
pygmæus	174	Umbra	174
rotengulus	117	Zygonectes	178
siuslawi	119	Limia	191
splirilingulius	128	arnoldi	191
storeri	117	caudofasciata	191
tincalla	115	couchiana	193
vandoisulus	121	cubensis	191
vittatus	128	dominicensis	191
zunnensis	114	formosa	192
(Leuciscus) caurinus, Cyprinus	118	heterandra	191
oregonensis, Cyprinus	114	matamorensis	192
leuciscus, Brachydeuterus	333	melanonotata	191
Pristipoma	333	nigrofasciata	191
leucophætus, Macrourus	205	ornata	191
Macrurus	205	pavonina	191
leucops, Photogenis	123	perugiae	191
engraulinus, Photogenis	123	pœcioides	192

List of scientific names—Continued

	Page	Page	
<i>Limia venusta</i>	175	<i>Liopsetta obscura</i>	228
<i>versicolor</i>	191	<i>putnami</i>	227
<i>vittata</i>	191	<i>Ilorus, Chasmistes</i>	108
<i>Limnurus</i>	182	<i>Ilosternus, Phenacobius teretulus</i>	142
<i>variegatus</i>	182	<i>Liostomus</i>	348
<i>limnurus, Girardinichthys</i>	183	<i>Liparidae</i>	399
<i>limosa, Myxine</i>	8	<i>liparina, Amitra</i>	405
<i>Pilodictis</i>	155	<i>liparitus, Paraliparis</i>	405
<i>limosus, Pylodictis</i>	155	<i>Liparis</i>	399
<i>linea, Mesopriion</i>	325	<i>agassizii</i>	400
<i>Syngnathus</i>	241	<i>arctica</i>	400
<i>lineata, Alosa</i>	42	<i>cyclopus</i>	400
<i>Coryphena</i>	430	<i>cyclostigma</i>	401
<i>Echenels</i>	449	<i>denneyi</i>	400
<i>Embletoca</i>	411	<i>fabricii</i>	400
<i>Scienea</i>	307	<i>fuconensis</i>	400
<i>Tigoma</i>	119	<i>gibbus</i>	400
<i>lineatus, Achirus</i>	230	<i>herschelianus</i>	400
<i>Amblyodon</i>	352	<i>liparis</i>	400
<i>Cyclopterus</i>	400	<i>megacephalus</i>	400
<i>Cypselurus</i>	202	<i>montagui</i>	399
<i>Diodon geometricus</i>	502	<i>mucosus</i>	399
<i>Dormitor</i>	436	<i>osborni</i>	400
<i>Doryichthys</i>	243	<i>pulchellus</i>	400
<i>Eques</i>	353	<i>ranula</i>	402
<i>Esox</i>	173	<i>reinhardti</i>	402
<i>Eugerres</i>	342	<i>tunicata</i>	400
<i>Euthynnus</i>	259	<i>liparis, Centrolophus</i>	267
<i>Exocetus</i>	202	<i>Cyclopterus</i>	399, 400
<i>Genyonemus</i>	349	<i>Liparis</i>	400
<i>Gobius</i>	440, 446	<i>major, Cyclopterus</i>	401
<i>Grystes</i>	374	<i>minor, Cyclopterus</i>	400
<i>Leiostomus</i>	349	<i>Lipariscus</i>	401
<i>Lepisosteus</i>	37	<i>nanus</i>	401
<i>Micropogon</i>	349	<i>Liparopidae</i>	398
<i>Mugil</i>	253	<i>Liparops</i>	398
<i>Pleuronectes</i>	230	<i>stelleri</i>	398
<i>Smaris</i>	342	<i>Lipogenyidae</i>	75
<i>Sternopygus</i>	101	<i>Lipogenys</i>	75
<i>Trichodon</i>	357	<i>gilli</i>	75
<i>Zygonectes</i>	177	<i>Lipomyzon</i>	109
<i>lineis transversis, Passer</i>	230	<i>Lipophrys</i>	461
<i>lineolata, Clupea</i>	41	<i>carolinus</i>	461
<i>Pelamys</i>	259	<i>liropus, Arius</i>	151
<i>Poecilia</i>	192	<i>Tachysurus</i>	151
<i>Sarda</i>	259	<i>lirus, Lepomis</i>	302
<i>lineolatum, Catonotus flabellare</i>	293	<i>Lythurus</i>	126
<i>lineolatus, Albinus</i>	135	<i>Nototroplus</i>	126
<i>Catonotus</i>	293	<i>lisa, Menidia</i>	249
<i>Metrogaster</i>	409	<i>Xenatherina</i>	249
<i>Pseudoscarus</i>	474	<i>lister, Ostracion</i>	407
<i>Tetronodon</i>	497	<i>lita, Muræna</i>	93
<i>Zygonectes</i>	178	<i>Litholepis</i>	38
<i>lineoplatus, Muræna</i>	92	<i>littoralis, Odontaspis</i>	19
<i>lineopunctatus, Balistes</i>	493	<i>Squalus</i>	10
<i>Xanthichthys</i>	493	<i>Umbrina</i>	251
<i>linguistula, Balistes</i>	496	<i>Umbrula</i>	351
<i>Pleuronectes</i>	222	<i>littoralis, Chaetodon</i>	360
<i>linnei, Eglienus</i>	211	<i>littorosus, Diodon</i>	501
<i>Trachurus</i>	270	<i>lividus, Petromyzon</i>	10
<i>Linophora</i>	350	<i>Silurus</i>	153
<i>Linophryne</i>	510	<i>livree, Pimelode</i>	156
<i>arborifer</i>	510	<i>liza, Mugil</i>	253
<i>lucifer</i>	510	<i>lobatus, Canthigaster</i>	498
<i>macrodon</i>	510	<i>Sphaeroides</i>	498
<i>polygong</i>	510	<i>lobochilus, Amphilius</i>	422
<i>linsleyi, Etheostoma</i>	293	<i>Heros</i>	422
<i>lioglossina</i>	223	<i>lobotes</i>	323
<i>tetraphthalma</i>	223	<i>auctorum</i>	323
<i>tetraphthalmus</i>	223	<i>emarginatus</i>	325
<i>holepis, Lionurus</i>	207	<i>erata</i>	323
<i>Macrusus (Lionurus)</i>	207	<i>pacificus</i>	324
<i>Xystreurus</i>	223	<i>somnolentus</i>	323
<i>liomonacanthus</i>	403	<i>surinamensis</i>	323
<i>Lioniscus</i>	34	<i>lobotidae</i>	323
<i>Lionurus</i>	207	<i>lockingtoni, Ichthys</i>	208
<i>holepis</i>	207	<i>lonchurus</i>	352
<i>(Lionurus) holepis, Macrurus</i>	207	<i>depressus</i>	352
<i>liopeiris, Acipenser</i>	35	<i>lanceolatus</i>	352
<i>Liperca</i>	313	<i>lorchophilus</i>	453
<i>liopropoma</i>	316	<i>micrognathus</i>	453
<i>aberrans</i>	316	<i>lonchurum, Opistognathus</i>	453
<i>longilepis</i>	316	<i>lonchurus</i>	352
<i>rubra</i>	315	<i>ancyodon</i>	354
<i>liopsetta</i>	227	<i>lonchurus, Opistognathus</i>	453
<i>glacialis</i>	227	<i>longa, Trygon</i>	28

List of scientific names—Continued

	Page		Page
<i>longicauda</i> , <i>Muraena</i> .	91	<i>Lophopsetta</i> .	217
<i>Rubula</i> .	91	<i>maculata</i> .	217
<i>longicaudus</i> , <i>Gobius</i> .	441	<i>Lophotes</i> .	215
<i>longicephalus</i> , <i>Galeichthys</i> .	150	<i>capellei</i> .	215
<i>Tachisurus</i> .	150	<i>cephedianus</i> .	215
<i>longiceps</i> , <i>Chenophryne</i> .	510	<i>Lophotidote</i> .	215
<i>Eleotris</i> .	435	<i>lordii</i> , <i>Salmo</i> .	61
<i>Hybopsis</i> .	133	<i>loreti</i> , <i>Gramma</i> .	322
<i>Porogadus</i> .	482	<i>Loricaria</i> .	161
<i>longidens</i> , <i>Caulolepis</i> .	233	<i>acuta</i> .	161
<i>longifilis</i> , <i>Gadomus</i> .	202	<i>altipinnis</i> .	162
<i>longifilus</i> , <i>Bathygadus</i> .	202	<i>aurea</i> .	162
<i>Careproctus</i> .	403	<i>barbata</i> .	161
<i>longilepis</i> , <i>Ilopropoma</i> .	316	<i>bransfordi</i> .	162
<i>longimana</i> , <i>Etheostoma</i> .	286	<i>capetensis</i> .	162
<i>longimane</i> , <i>Limostoma</i> .	286	<i>filamentosa latiura</i> .	162
<i>longimanus</i> , <i>Astatheros</i> .	420	<i>fimbriata</i> .	162
<i>Eupomotis</i> .	302	<i>flava</i> .	160
<i>Heros</i> .	420	<i>frenata</i> .	162
<i>Squalus</i> .	15	<i>latiura</i> .	162
<i>Xystroptiles</i> .	302	<i>lima</i> .	162
<i>longipes</i> , <i>Bregmaceros</i> .	208	<i>panamensis</i> .	162
<i>Gadus</i> .	213	<i>plecostomus</i> .	160
<i>Physcetus</i> .	212	<i>rostrata</i> .	161, 162
<i>longipinne</i> , <i>Gobiosoma</i> .	445	<i>strigilata</i> .	162
<i>Holocentrum</i> .	235	<i>tuyrensis</i> .	162
<i>longipinnis</i> , <i>Careproctus</i> .	403	<i>uraeanthra</i> .	162
<i>Clevelandia</i> .	445	<i>variegata</i> .	161
<i>Evermannia</i> .	445	<i>Loricariichthys</i> .	161
<i>Leptocerdale</i> .	473	<i>Loricariidae</i> .	160
<i>Stromateus</i> .	268	<i>loricata</i> , <i>Alysia</i> .	169
<i>longirostris</i> , <i>Alburnops</i> .	135	<i>loricatus</i> , <i>Gasterosteus</i> .	237
<i>Chætodon</i> .	359	<i>Macrognathus</i> .	36
<i>Esox</i> .	198	<i>Phalangistes</i> .	304
<i>Euleptorhamphus</i> .	198	<i>lorito</i> , <i>Sparsiona</i> .	432
<i>Forcipiger</i> .	359	<i>Loro</i> .	434, 435
<i>Hemirhamphus</i> .	198	<i>Lotia</i> .	212
<i>Hybopsis</i> .	135	<i>brosulaana</i> .	212
<i>Hypopion</i> .	17	<i>inornata</i> .	212
<i>Lepisosteus</i> .	36	<i>maculosa</i> .	212
<i>Leuciscus</i> .	120	<i>Lotella</i> .	211
<i>Lumpenella</i> .	471	<i>maxillaris</i> .	211
<i>Lumpenus</i> .	471	<i>lotharingus</i> , <i>Gasterosteus</i> .	239
<i>Saurus</i> .	163	<i>loubina</i> , <i>Perca</i> .	305
<i>longirostrum</i> , <i>Catostomus</i> .	106	<i>louisiana</i> , <i>Engraulis</i> .	48
<i>longispatha</i> , <i>Peristedion</i> .	405	<i>louisiane</i> , <i>Notropis</i> .	124
<i>longispatum</i> , <i>Peristedion</i> .	405	<i>Syngnathus</i> .	242
<i>longispinis</i> , <i>Aelurichthys</i> .	148	<i>Syrichtes</i> .	242
<i>Lepomis</i> .	301	<i>louisianensis</i> , <i>Lepidosteus</i> .	36
<i>Pontinus</i> .	372	<i>lowii</i> , <i>Polytmixia</i> .	234
<i>longulus</i> , <i>Pomotis</i> .	209	<i>lowii</i> , <i>Onosodus</i> .	171
<i>longum</i> , <i>Gobiosoma</i> .	446	<i>loxias</i> , <i>Gurnardus</i> .	406
<i>longurio</i> , <i>Carcharias</i> .	15	<i>Prionotus</i> .	406
<i>Scoliodon</i> .	15	<i>Lucania</i> .	175, 182
<i>longus</i> , <i>Balistes</i> .	493	<i>affinis</i> .	175
<i>Dasyatis</i> .	28	<i>browni</i> .	181
<i>Ophisurus</i> .	85	<i>goodelli</i> .	175
<i>lophar</i> , <i>Perca</i> .	280	<i>parva</i> .	175
<i>Lopharis</i> .	280	<i>lucasana</i> , <i>Sphyraena</i> .	255
<i>mediterraneus</i> .	280	<i>lucasanum</i> , <i>Thalassoma</i> .	429
<i>Lophiidae</i> .	504	<i>lucasanus</i> , <i>Julis</i> .	420
<i>Lophiomus</i> .	504	<i>lucasi</i> , <i>Ceratocottus</i> .	382
<i>setigerus</i> .	504	<i>lucayanum</i> , <i>Asymmetron</i> .	7
<i>Lophius</i> .	504	<i>lucens</i> , <i>Ceratichthys</i> .	137
<i>aculeatus</i> .	507	<i>Dacentrus</i> .	412
<i>americanus</i> .	504	<i>lucetia</i> , <i>Vinciguerra</i> .	73
<i>bufo</i> .	487	<i>lucetius</i> , <i>Maurolicus</i> .	73
<i>gibbus</i> .	505	<i>lucie</i> , <i>Hydrargyreus</i> .	178
<i>histrio ocellatus</i> .	505	<i>Zygonectes</i> .	178
<i>piscatorius</i> .	504	<i>lucida</i> , <i>Aethroprion</i> .	168
<i>radiatus</i> .	507	<i>Anchoviella</i> .	48
<i>setigerus</i> .	504	<i>lucidus</i> , <i>Leucichthys</i> .	62
<i>spectrum</i> .	505	<i>Luxilus</i> .	125
<i>tumidus</i> .	504	<i>Salmo</i> (<i>Coregonus</i>).	62
<i>vespertilio</i> .	507	<i>Stolephorus</i> .	48
<i>viviparus</i> .	504	<i>Lucifer</i> .	70
<i>lophius</i> , <i>Amiurus</i> .	163	<i>lucifer</i> , <i>Linophryne</i> .	510
<i>Haustor</i> .	153	<i>Lucifuga</i> .	479
<i>Lophobranchii</i> .	240	<i>dentatus</i> .	479
<i>Lophodolus</i> .	509	<i>subterranea</i> .	479
<i>acanthognathus</i> .	509	<i>subterraneus</i> .	479
<i>Lophogobius</i> .	438	<i>Lucioblemnius</i> .	465
<i>cyrinoides</i> .	438	<i>alepidotus</i> .	465
<i>Lopholatilis chamaleonticeps</i> .	357	<i>lucus</i> .	465
<i>Lopholatilis</i> .	357	<i>lucioceps</i> , <i>Saurus</i> .	163

List of scientific names—Continued

	Page		Page
<i>lucioceps</i> , <i>Synodus</i> .	163	<i>Lutianus analis</i> .	328
<i>Luciocharax</i> .	100	<i>apodus</i> .	325, 326
<i>insculptus</i> .	100	<i>argentiventris</i> .	326
<i>striatus</i> .	100	<i>aya</i> .	326
<i>Lucioperca americana</i> .	282	<i>blackfordii</i> .	326
<i>borea</i> .	282	<i>brachypterus</i> .	326
<i>canadensis</i> .	282	<i>buccanellia</i> .	326
<i>grisea</i> .	282	<i>canipechanus</i> .	326
<i>pepinis</i> .	282	<i>colorado</i> .	326
<i>Lucioperca undecimalis</i> , <i>Centropomus</i> .	305	<i>cyanopterus</i> .	325
<i>lucioperca</i> , <i>Mentipercra</i> .	319	<i>griseus</i> .	325, 326
<i>Serranus</i> .	319	<i>guttatus</i> .	326
<i>Lucius</i> .	172	<i>hastingsi</i> .	326
<i>lucius americanus</i> , <i>Esox</i> .	173	<i>jocú</i> .	325
<i>lucius</i> , <i>Atractosteus</i> .	38	<i>jordani</i> .	324
<i>Chagnoensis</i> .	465	<i>luteus</i> .	335
<i>Chirostoma</i> .	251	<i>lutianus</i> .	324
<i>Luciobrama</i> .	465	<i>lutjanoides</i> .	326
<i>Ptychocheilus</i> .	114	<i>mahogoni</i> .	327
<i>lucoides</i> , <i>Esox</i> .	173	<i>megalophthalmus</i> .	327
<i>lucretiae</i> , <i>Aboma</i> .	443	<i>novemfasciatus</i> .	325
<i>Gobius</i> .	443	<i>synagris</i> .	320, 327
<i>ludibunda</i> , <i>Cyprinella</i> .	130	<i>viridis</i> .	324
<i>luetkeni</i> , <i>Dolopichthys</i> .	510	<i>vivatus</i> .	326
<i>lugubris</i> , <i>Caranx</i> .	273	<i>lutianus</i> , <i>Lutianus</i> .	324
<i>Cyprinella</i> .	131	<i>lutipinnis</i> , <i>Hydrophiox</i> .	127
<i>Esox</i> .	173	<i>Pristigaster</i> .	45
<i>Malacoctenus</i> .	460	<i>lutjanoides</i> , <i>Lutianus</i> .	326
<i>Melampatius</i> .	233	<i>Ocyurus</i> .	328
<i>Myrodes</i> .	460	<i>Lutjanus</i> .	324
<i>Plectromus</i> .	233	<i>acutirostris</i> .	325
<i>Xurel</i> .	273	<i>auberti</i> .	327
<i>luitpoldi</i> , <i>Characodon</i> .	184	<i>blackfordi</i> .	326
<i>Goodea</i> .	184	<i>brachypterus</i> .	326
<i>lumbricus</i> , <i>Muraena</i> .	76	<i>cayanensis</i> .	354
<i>Myrophis</i> .	83	<i>colorado</i> .	326
<i>luminosum</i> , <i>Myctophum</i> .	170	<i>cubera</i> .	325
<i>Lumpenella</i> .	471	<i>dentatus</i> .	325
<i>longirostris</i> .	471	<i>lunulatus</i> .	311
<i>Lumpenidae</i> .	470	<i>novemfasciatus</i> .	325
<i>Lumpenus</i> .	470	<i>prieto</i> .	325
<i>fabricii</i> .	471	<i>purpureus</i> .	328
<i>gracilis</i> .	470	<i>stearnsi</i> .	325
<i>lampteraformis</i> .	471	<i>surinamensis</i> .	332
<i>longirostris</i> .	471	<i>torridus</i> .	326
<i>mackayi</i> .	471	<i>triangulum</i> .	348
<i>nobilis</i> .	471	<i>tridens</i> .	318
<i>Lumpus</i> .	397	<i>trilobatus</i> .	317
<i>anglerum</i> .	398	<i>verres</i> .	425
<i>vulgaris</i> .	398	<i>lutkeni</i> , <i>Cypselurus</i> .	201
<i>lumpus</i> , <i>Cyclopterus</i> .	397	<i>Exocetus</i> .	201
<i>luna</i> , <i>Pomotis</i> .	301	<i>Lutodaira</i> .	40
<i>Zeus</i> .	216	<i>lutrensis</i> , <i>Leuciscus</i> .	131
<i>lunarisi</i> , <i>Orthagoriscus</i> .	503	<i>Mormona</i> .	131
<i>unata</i> , <i>Centropyge</i> .	361	<i>lutzi</i> , <i>Heterandria</i> .	19C
<i>Echeneis</i> .	448	<i>Luvaridiae</i> .	264
<i>Solea</i> .	217	<i>Luvariformes</i> .	264
<i>lunaticus</i> , <i>Dactyloscopus</i> .	455	<i>Luvarus</i> .	264
<i>lunatus</i> , <i>Holocanthus</i> .	361	<i>imperialis</i> .	264
<i>Platyphrys</i> .	217	<i>luxatus</i> , <i>Chasmistes</i> .	108
<i>Pleuronectes</i> .	217	<i>Deltistes</i> .	108
<i>Rhinichthys</i> .	140	<i>Luxilinus</i> .	113
<i>lunulatus</i> , <i>Cynias</i> .	13	<i>luxiloides</i> , <i>Cyprinella</i> .	131
<i>Lutjanus</i> .	310	<i>Luxilus</i> .	125, 128
<i>Mustelus</i> .	13	<i>albofusca</i> .	129
<i>lupus</i> , <i>Anarhichas</i> .	472	<i>cerasinus</i> .	128
<i>Haustor</i> .	153	<i>chickasawensis</i> .	130
<i>Pimelodus</i> .	153	<i>cbrysocephalus</i> .	128
<i>Lusca</i> , <i>Cyclothone</i> .	72	<i>cornutus</i> .	128
<i>luscosus</i> , <i>Leucicorus</i> .	483	<i>cbrysocephalus</i> .	128
<i>lutea</i> , <i>Anguilla</i> .	77	<i>isolepis</i> .	128
<i>luteovinctum</i> , <i>Etheostoma</i> .	291	<i>dissimilis</i> .	138
<i>luteovinctus</i> , <i>Oligocephalus</i> .	291	<i>elongatus</i> .	120
<i>lutescens</i> , <i>Chætodon</i> .	360	<i>erythrogaster</i> .	113
<i>Kyphosus</i> .	340	<i>interruptus</i> .	128
<i>Pimeleopterus</i> .	340	<i>kentuckiensis</i> .	137
<i>luteum</i> , <i>Hæmulon</i> .	330	<i>lacertosus</i> .	129
<i>luteus</i> , <i>Centropomus</i> .	281	<i>leptosomus</i> .	116
<i>Gadus tomcodus</i> .	210	<i>lucidus</i> .	125
<i>Genyatremus</i> .	335	<i>roseus</i> .	127
<i>Lutianus</i> .	335	<i>seco</i> .	115
<i>Noturus</i> .	155	<i>selene</i> .	132
<i>Rhinichthys</i> .	139	<i>zonistius</i> .	128
<i>Lutianidae</i> .	324	<i>lycaodon</i> , <i>Salmo</i> .	55
<i>Lutianus</i> .	324	<i>Lycenchelys</i> .	475
<i>ambiguus</i> .	327	<i>paxillus</i> .	476

List of scientific names—Continued

	Page		Page
<i>Lycenchelys porifer</i>	476	<i>Lyosphæra</i>	503
<i>verrillii</i>	475	<i>globosa</i>	703
<i>Lycengraulis</i>	51	<i>lyricus</i> , <i>Gobionellus</i>	441
<i>grossidens</i>	51	<i>Gobius</i>	441
<i>poeyi</i>	51	<i>lysimus</i> , <i>Bryophorus</i>	465
<i>Lynchianopole</i>	72	<i>lythrochloris</i> , <i>Xenotis</i>	300
<i>argenteolus</i>	73	<i>Lythrulon</i>	331
<i>lynchus</i> , <i>Argyropelecus</i>	74	<i>flaviguttatum</i>	331
<i>Lycius</i>	475	<i>opalescens</i>	331
<i>nebulosus</i>	475	<i>Lythrurus</i>	125
<i>seminudus</i>	475	<i>ardens</i>	128
<i>Lycichthys</i>	472	<i>atripes</i>	126
<i>latifrons</i>	472	<i>bellus</i>	126
<i>minor</i>	472	<i>cyancephalus</i>	126
<i>Lycocara</i>	478	<i>fasciolaris</i>	126
<i>parrisi</i>	478	<i>lirus</i>	126
<i>Lycodalepis</i>	476	<i>matutinus</i>	126
<i>mucosus</i>	476	<i>notropis</i>	126
<i>polaris</i>	476	<i>punctulatus</i>	126
<i>Lycodapodidae</i>	478	<i>roseipinnis</i>	126
<i>Lycodapus</i>	478	<i>umbratilis</i>	125
<i>attenuatus</i>	479	<i>Lythrypnus</i>	442
<i>dermatinus</i>	478	<i>dalli</i>	442
<i>extensus</i>	478	<i>Macabia</i>	40
<i>ferasfer</i>	478	<i>macchellus</i> , <i>Caraux</i>	270
<i>grossidens</i>	479	<i>Decapterus</i>	270
<i>lycodon</i>	478	<i>macrusekeyi</i> , <i>Ictalurus</i>	153
<i>mandibularis</i>	479	<i>macclendoni</i> , <i>Bregmaceros</i>	208
<i>parviceps</i>	478	<i>macdonaldi</i> , <i>Acutomentum</i>	366
<i>Lycodes</i>	474	<i>Coccotis</i>	127
<i>anguis</i>	475	<i>Conocara</i>	53
<i>brevipes</i>	475	<i>Cynoscion</i>	356
<i>cicatriger</i>	475	<i>Eriscion</i>	356
<i>coccineus</i>	476	<i>Nannobrachium</i>	168
<i>concolor</i>	476	<i>Notropis</i>	127
<i>diapterus</i>	476	<i>Penopus</i>	483
<i>digitatus</i>	475	<i>Salmō</i>	57
<i>esmarkii</i>	474	<i>mykiss</i>	57
<i>frigidus</i>	474	<i>Zygonectes</i>	177
<i>gracilis</i>	475	<i>Macdonaldia</i>	76
<i>incisus</i>	475	<i>macellus</i> , <i>Prionistius</i>	380
<i>jordani</i>	474	<i>macer</i> , <i>Polyprosopus</i>	20
(<i>Lycodopsis</i>) <i>pacificus</i>	474	<i>mackayi</i> , <i>Lumpenus</i>	471
<i>mucoosus</i>	476	<i>Siphostoma</i>	241
<i>nebulosus</i>	475	<i>Syngnathus</i>	241
<i>palearis</i>	475	<i>mackenzii</i> , <i>Salmo</i>	61
<i>pixilloides</i>	476	<i>Stenodus</i>	61
<i>paxillus</i>	476	<i>macloei</i> , <i>Carcharhius</i>	17
<i>perspicillum</i>	474	<i>maculura</i> , <i>Raiā</i>	29
<i>porifer</i>	476	<i>macouni</i> , <i>Chauliodus</i>	71
<i>reticulatus</i>	474	<i>Pterygiocottus</i>	380
<i>serpens</i>	475	<i>macracanthum</i> , <i>Pristipoma</i>	334
<i>terra-novae</i>	474	<i>macracanthus</i> , <i>Alutarius</i>	496
<i>turneri</i>	476	<i>Macrhybopsis</i>	334
<i>vahlii</i>	474	<i>astivalis</i>	138
<i>verrillii</i>	475	<i>gelida</i>	139
<i>zonarchus</i>	475	<i>hyostoma</i>	139
<i>lycodon</i> , <i>Lycodapus</i>	478	<i>marconis</i>	138
<i>Lycodontis albimentis</i>	93	<i>meeki</i>	139
<i>Lycodonus</i>	476	<i>montana</i>	130
<i>mirabilis</i>	476	<i>sterletus</i>	139
<i>Lycodopsis</i>	474	<i>macrocephala</i> , <i>Clupea</i>	40
<i>crassilabris</i>	474	<i>Muraena</i>	77
<i>crotalinius</i>	473	<i>macrocephalum</i> , <i>Etheostoma</i>	284
<i>pacificus</i>	474	<i>macrocephalus</i> , <i>Alvordius</i>	284
<i>securus</i>	474	<i>Clinus</i>	455
<i>Lycogramma</i>	476	<i>Gadus</i>	210
<i>brunnea</i>	476	<i>Gymnophthalmus</i>	409
<i>Lycondontis</i>	91	<i>Mnieres</i>	455
<i>jordani</i>	92	<i>Sebastes</i>	372
<i>Lyconectes</i>	471	<i>Semotilus</i>	117
<i>aleutensis</i>	471	<i>macrocerus</i> , <i>Monacanthus</i>	404
<i>Lyconema</i>	476	<i>macrocheilus</i> , <i>Catostomus</i>	106
<i>barbatum</i>	476	<i>macrochir</i> , <i>Aldrovandia</i>	75
<i>lynx</i> , <i>Pimelodus</i>	153	<i>Halosaurus</i>	75
<i>Lycoetus</i>	510	<i>Sobastes</i>	364
<i>murrayi</i>	611	<i>macrochira</i> , <i>Holopercá</i>	301
<i>lyolepis</i> , <i>Anchoviella</i>	49	<i>macrochirus</i> , <i>Hemirhamphus</i>	199
<i>Stolephorus</i>	49	<i>Lepomis</i>	301
<i>Lyolliparis</i>	400	<i>Macrodon</i>	90, 364
<i>pulchellus</i>	400	<i>aimara</i>	99
<i>Lyomeri</i>	95	<i>ancylodon</i>	354
<i>Lyopomai</i>	74	<i>auritus</i>	90
<i>Lyopsetta</i>	222		
<i>jordani exilis</i>	222		

List of scientific names—Continued

	Page		Page
<i>Macrodon ferox</i>	99	<i>Macrostoma</i>	166
<i>microlepis</i>	99	<i>angustidens</i>	166
<i>mordax</i>	354	<i>castaneum</i>	167
<i>patana</i>	99	<i>margaritiferum</i>	166
<i>tereira</i>	99	<i>quercentum</i>	166
<i>tores</i>	99	<i>macrostoma</i> , <i>Cyprinella</i>	131
<i>trahira</i>	99	<i>Hæmulum</i>	329
<i>macrodon</i> , <i>Chelodipterus</i>	296	<i>macrostomum</i> , <i>Hæmulum</i>	329
<i>Erythrinus</i>	99	<i>Macrostomus</i>	166
<i>Linophryne</i>	510	<i>macrostomus</i> , <i>Accipenser</i>	35
<i>Macrodonopbis</i>	88	<i>Macrouridae</i>	202
<i>mordax ophiurus</i>	88	<i>Macrourus</i>	204, 205
<i>macroodus</i> , <i>Squalus</i>	19	<i>albatrossus</i>	206
<i>macrogenis</i> , <i>Cerna</i>	315	<i>anguliceps</i>	206
<i>macrognathus</i> , <i>Opistognathus</i>	452	<i>bairdii</i>	205
<i>Macrognathus</i>	245	<i>barbiger</i>	205
<i>brevirostris</i>	199	<i>berglax</i>	205
<i>loricatus</i>	36	<i>boöps</i>	205
<i>macrognathus</i> , <i>Leucichthys</i>	62	<i>bucephalus</i>	205
<i>macrolepidota</i> , <i>Anchovia</i>	50	<i>bullocki</i>	205
<i>Cobitis</i>	176	<i>canus</i>	206
<i>macrolepidotum</i> , <i>Moxostoma</i>	111	<i>cajito</i>	205
<i>macrolepidotus</i> , <i>Catostomus</i>	111	<i>carminalifer</i>	206
<i>Engraulis</i>	50	<i>cinereus</i>	235
<i>Fundulus heteroclitus</i>	176	<i>couvergens</i>	206
<i>Leuciscus</i>	116	<i>cuspis</i>	206
<i>Notropis</i>	125	<i>fragilis</i>	206
<i>Pogonichthys</i>	116	<i>gracilicauda</i>	206
<i>Macrolepis</i>	295	<i>ingolfi</i>	206
<i>macrolepis</i> , <i>Phenagoniates</i>	95	<i>latinasutus</i>	206
<i>Pontinus</i>	372	<i>latirostratus</i>	206
<i>Rroboides</i>	95	<i>lepturus</i>	207
<i>macronema</i> , <i>Pimelodus</i>	150	<i>leucophagus</i>	205
<i>macronemus</i> , <i>Polynemus</i>	256	(<i>Malacocephalus</i>) <i>pectoralis</i>	204
<i>macrophthalmus</i> , <i>Astyianax</i>	97	<i>orbitalis</i>	206
<i>Caranx</i>	271	<i>pectoralis</i>	204
<i>Centropristes</i>	328	<i>stegidolepis</i>	205
<i>Epigonus</i>	296	<i>tenuicauda</i>	206
<i>Gobiesox</i>	489	<i>trichurus</i>	206
<i>Hypentelium</i>	107	(<i>Macrourus</i>) <i>stegidolepis</i> , <i>Macrurus</i>	205
<i>macropoma</i> , <i>Bollmannia</i>	443	<i>macrourus</i> , <i>Alopias</i>	18
<i>Centropristes</i>	318	<i>Gymnotus</i>	101
<i>Diplectrum</i>	318	<i>Macrozoarcæ</i>	473
<i>Macrops</i>	328	<i>americanus</i>	473
<i>macrops</i> , <i>Balistes</i>	492	<i>macrurum</i> , <i>Ophisoma</i>	80
<i>Bathygadus</i>	203	<i>Macrurus</i>	205
<i>Calamus</i>	337	<i>acrolepis</i>	203
<i>Citharichthys</i>	219	<i>anguliceps</i>	205
<i>Conger</i>	79	<i>asper</i>	203
<i>Corvina</i>	345	<i>barbiger</i>	205
<i>Cottus</i>	384	<i>boöps</i>	205
<i>Gnathopogon</i>	453	<i>bucephalus</i>	205
<i>Hippoglossina</i>	222, 223	<i>bullocki</i>	205
<i>Opistognathus</i>	453	<i>canus</i>	206
<i>Opisthoteropus</i>	45	<i>capito</i>	205
<i>Pristigaster</i>	45	<i>caribicus</i>	207
<i>Vacuouqua</i>	345	<i>carminalatus</i>	207
<i>macroptera</i> , <i>Conocara</i>	53	<i>carminalifer</i>	206
<i>Gonlobatis</i>	30	<i>cinereus</i>	205
<i>macropterus</i> , <i>Hypentelium</i>	107	(<i>Cælorhynchus</i>) <i>scaphocephalus</i>	207
<i>macropterus</i> , <i>Alepocephalus</i>	53	<i>convergens</i>	206
<i>Balistes</i>	493	<i>cuspis</i>	206
<i>Centrarchus</i>	304	<i>dorsalis</i>	207
<i>Labrus</i>	304	<i>fimbriquamis</i>	204
<i>Thynnus</i>	260	<i>fragilis</i>	206
<i>macrophthalmus</i> , <i>Astyianax</i>	97	<i>goodii</i>	203
<i>Centropristes</i>	328	<i>gracilicauda</i>	206
<i>Dormitator</i>	436	<i>ingolfi</i>	206
<i>Gobiesox</i>	489	<i>latinasutus</i>	203
<i>Harengula</i>	43	<i>latirostratus</i>	206
<i>Priacanthus</i>	322	<i>leucophagus</i>	205
<i>Pristipomoides</i>	328	(<i>Lionurus</i>) <i>holepis</i>	207
<i>Scomber</i>	258	<i>longifilis</i>	203
<i>Tetragonopterus</i>	97	(<i>Macrourus</i>) <i>stegidolepis</i>	205
<i>macrothalamia</i> , <i>Cluia</i>	43	(<i>Nematonurus</i>) <i>magnus</i>	204
<i>macrops</i> , <i>Malacoctenus</i>	460	<i>occa</i>	207
<i>Myrodes</i>	460	<i>orbitalis</i>	206
<i>Macrorhamphosidae</i>	245	<i>suborbitalis</i>	203
<i>Macrorhamphosidae</i>	245	<i>tenuicauda</i>	206
<i>macrurus</i> , <i>Leptoferasfer</i>	486	<i>trichurus</i>	206
<i>Monacanthus</i>	493	<i>villosus</i>	207
<i>Macrorhamphosidae</i> , <i>hawaiiensis</i>	246	<i>macrurus</i> , <i>Anago</i>	80
<i>scolopax</i>	246	<i>Gymnotus</i>	101
<i>macrorhinus</i> , <i>Acipenser</i>	34	<i>Oxydontaichthys</i>	88
<i>macrosipha</i> , <i>Piramutana</i>	159	<i>Oxyodontichthys</i>	87

List of scientific names—Continued

	Page		Page
macularius baileyi, Cyprinodon.....	181	maculosus, Serpens marinus.....	87
macularius, Cyprinodon.....	181	Serranus.....	311
maculata, Belone.....	197	madeirensis, Ceratopeplus.....	167
<i>Etheostoma</i>	290	Pseudonuræna.....	91
<i>Hemitremia</i>	135	Scopelus.....	167
<i>Hybopsis</i>	135	maderaspatensis, <i>Prionanthichthys</i>	310
<i>Lophopsetta</i>	217	maderensis, <i>Helicolenus</i>	370
<i>Morone</i>	302	meandricus, <i>Caularchus</i>	488
<i>Nerophis</i>	243	Lepadogaster.....	488
<i>nigra</i> , <i>Muraena</i>	91	Mænidae.....	340
<i>nigra et viridis</i> , <i>Muraena</i>	92	magasema, <i>Geophagus</i>	424
<i>Percæ</i>	310	magdaleneæ, <i>Curimata</i>	100
<i>Sciæna</i>	436	<i>Curimatus</i>	100
maculaticeps, <i>Boleosoma</i>	287	<i>Otolithus</i>	356
maculatofasciatus, <i>Paralabrax</i>	317	<i>Paralichthys</i>	223
<i>Serranus</i>	317	magistralis, <i>Epinudia</i>	261
maculatostritus, <i>Diodon</i>	502	magniculus, <i>Ophichthus</i>	88
maculatum, <i>Aulostomus</i>	245	<i>Scytophalophis</i>	87, 88
<i>Boleosoma</i>	287	<i>magnus</i> , <i>Macrurus</i> (<i>Nematonurus</i>).....	204
maculatus, <i>Alvordius</i>	283	<i>mahogoni</i> , <i>Lutianus</i>	327
<i>Anarrhichas</i>	472	<i>Mesopriion</i>	327
<i>A pogon</i>	295	<i>mainensis</i> , <i>Gasterosteus</i>	230
<i>Aulostomus</i>	245	<i>Pungitius</i>	239
<i>Balistes</i>	493	<i>majalis</i> , <i>Cobitis</i>	176
<i>Canthidermis</i>	493	<i>Fundulus</i>	175, 176
<i>Cirrhites</i>	358	<i>major</i> , <i>Actinocirrhus</i>	401
<i>Clinus</i>	470	<i>Blepharis</i>	274
<i>Cogrus</i>	87	<i>Cyclopterus liparis</i>	401
<i>Cottus</i>	386	<i>Gallichthys</i>	274
<i>Cryptacanthodes</i>	471	<i>Ptychocheilus</i>	114
<i>Diodon</i>	501	<i>Makaira</i>	203
<i>Dormitator</i>	436	<i>amph</i>	263
<i>Galeus</i>	14	<i>holei</i>	263
<i>Gasteropelecus</i>	99	<i>marlina</i>	263
<i>Hadropterus</i>	283	<i>mazara</i>	263
<i>immaculatus</i> , <i>Platypterus</i>	193	<i>mitsukurii</i>	263
<i>Labrus scometerque</i>	355	<i>nigricans</i>	263
<i>Lembus</i>	435	<i>makua</i> , <i>Raniganza</i>	504
<i>Leptoclinus</i>	470	<i>malabarica</i> , <i>Elaeotris</i>	281
<i>Monopriion</i>	295	<i>malabaricus</i> , <i>Esox</i>	99
<i>Mullus</i>	343	<i>Hoplias</i>	99
<i>Nothonotus</i>	290	<i>Synodus</i>	99
<i>Notorynchus</i>	11	<i>Malacanthidae</i>	356
<i>Pimedodus</i>	151, 152, 158, 159	<i>Malacanthus</i>	356
<i>Platypterus</i>	103	<i>plumieri</i>	356
<i>Pleuronectes</i>	217	<i>Malacoccephalus occidentalis</i>	204
<i>Proceros</i>	30	<i>Malacocottus</i>	388
<i>Psenes</i>	268	<i>kincaldi</i>	388
<i>Rhypticus</i>	316, 317	<i>zonurus</i>	388
<i>Rypiticus</i>	310	<i>Malacocetus</i>	459
<i>Scomber</i>	257	<i>bifasciatus</i>	460
<i>Scomberomorus</i>	257, 258	<i>bimaculatus</i>	460
<i>Serranus</i>	311	<i>culebra</i>	400
<i>Sphaeroides</i>	498	<i>deletlandi</i>	459
<i>Tetronod hispidus var.</i>	498	<i>gilli</i>	400
<i>Upeneus</i>	343	<i>lugubris</i>	400
<i>Zoogoneticus</i>	182	<i>macropodus</i>	400
maculicauda, <i>Astatheros</i>	421	<i>moorei</i>	400
<i>Cichlosoma</i>	421	<i>ocellatus</i>	459
<i>Orthostechus</i>	331	<i>puertoricensis</i>	400
maculifer, <i>Diodon</i>	501	<i>varius</i>	459
<i>Platophrys</i>	217	<i>versicolor</i>	400
maculiferus, <i>Hippocampus</i>	321	<i>Malacostomidae</i>	71
<i>Pleuronectes</i>	217	<i>Malacosteus</i>	71
maculipinnis, <i>Iridio</i>	427	<i>niger</i>	71
<i>Julis</i>	427	<i>malarinat</i> , <i>Peristedion</i>	405
maculipinnis, <i>Exocetus</i>	201	<i>maliger</i> , <i>Pteropodus</i>	369
<i>Heros</i>	422	<i>Sebastichthys</i>	369
<i>Isogomphodon</i>	16	<i>mallispinosus</i> , <i>Monomeropus</i>	18
<i>Monochir</i>	230	<i>malleus</i> , <i>Zygema</i>	66
<i>Thyrosidea</i>	92	<i>Mallotus</i>	66
maculo-cinctus, <i>Sarotherodon</i>	360	<i>villus</i>	66
maculosa, <i>Harengula</i>	43	<i>(Mallotus) pacificus</i> , <i>Salmo</i>	66
<i>Lota</i>	212	<i>malma</i> , <i>Salmo</i>	61
<i>Muraena</i>	87	<i>Salvelinus</i>	61
<i>Thalassophryne</i>	488	<i>spectabilis</i> , <i>Salvelinus</i>	61
maculo-seriatus, <i>Chirus</i>	373	<i>Malthea</i>	506
maculoseum, <i>Hæmulon</i>	329	<i>truncata</i>	507
maculosus, <i>Acentrolophus</i>	267	<i>Malthe</i>	506
<i>Acipenser</i>	35	<i>cubifrons</i>	507
<i>Catostomus</i>	107	<i>elater</i>	507
<i>Centrarchus</i>	303	<i>Maltoea nasuta</i>	507
<i>Gadus</i>	212	<i>notata</i>	507
<i>Nomeus</i>	268	<i>Malthopsis</i>	507
<i>Oligocottus</i>	360	<i>spinosa</i>	507
<i>Paralichthys</i>	223	<i>spinulosa</i>	507
<i>Pimephales</i>	145	<i>managuense</i> , <i>Parapetenia</i>	418

List of scientific names—Continued

	Page		Page
managuensis, <i>Heros</i> .	418	marina, <i>Pastinaca</i> .	29
<i>Pimelodus</i> .	157	<i>Perca</i> .	363
<i>Rhamdia</i> .	157	<i>rubra, Perca</i> .	236
manatia, <i>Raiia</i> .	32	<i>sectatrix, Perca</i> .	340
manatinus, <i>Barathrodemus</i> .	482	<i>Strongylura</i> .	166
Mancalias.	508	<i>Vulpacula</i> .	18
<i>shufeldti</i> .	508	marinus, <i>Diabolus</i> .	32
<i>uranoscopus</i> .	508	<i>dorsatus, Petromyzon</i> .	9
mandibularis, <i>Lycodapus</i> .	479	<i>Esox</i> .	196
manglicola, <i>Gobius</i> .	439	<i>fer quadratus, Faber</i> .	358
<i>Rhinogobius</i> .	439	<i>Gasterosteus</i> .	239
mango, <i>Polynemus</i> .	256	<i>maculosus, Serpens</i> .	87
manitou, <i>Percina</i> .	283	<i>Mugilops</i> .	252
manitoulinus, <i>Leucichthys</i> .	64	<i>Petromyzon</i> .	8
manjuarli, <i>Lepidosteus</i> .	38	<i>Sebastes</i> .	363
manni, <i>Gambusia</i> .	186	<i>Sillurus</i> .	148
maomii, <i>Zygoctenes</i> .	174	marki, <i>Emblemaria</i> .	465
Manta.	32	<i>marlina, Makala</i> .	263
<i>americana</i> .	32	<i>marmorata, Amia</i> .	38
<i>birostris</i> .	32	<i>Pteroplata</i> .	29
<i>hamiltoni</i> .	32	<i>Ulca</i> .	393
manta, <i>Cephaloptera</i> .	32	<i>Unibranchapertura</i> .	76
<i>Cephalopterus</i> .	32	marmoratum, <i>Corallitoa</i> .	456
Mapo.	440	<i>Lepophidium</i> .	481
mapo, <i>Gobius</i> .	440	<i>Leptophidium</i> .	484
maraldi, <i>Gadella</i> .	211	<i>marinoratus, Ameiurus</i> .	154
<i>Gadus</i> .	211	<i>Cottus</i> .	387
marcgravii, <i>Aulostome</i> .	245	<i>Creminobates</i> .	456
<i>Coryphena</i> .	245	<i>Gobiesox</i> .	488
<i>Rhinobatus</i> .	23	<i>Hemitripterus</i> .	382, 393
<i>Salmo</i> .	100	<i>Plinelodus</i> .	154
marconis, <i>Hybopsis festivalis</i> .	138	<i>Rhinichthys</i> .	139
<i>Macrhybopsis</i> .	138	<i>Rivulus</i> .	179
Margariscus.	116	<i>Scorpaenichthys</i> .	382
<i>margarita</i> .	116	<i>Sphoeroides</i> .	498
<i>nachtriebli</i> .	116	<i>Synbranchus</i> .	76
margarita, <i>Clinostomus</i> .	116	<i>Tetrodon</i> .	498
<i>Echistoma</i> .	70	<i>Tirus</i> .	163
<i>Margariscus</i> .	116	marmorata, <i>Rabula</i> .	91
margaritaceus, <i>Entomacrodus</i> .	463	<i>marinoreus, Blennius</i> .	461
Margariscus.	463	<i>Muraenopsis</i> .	91
<i>Margarita</i> .	166	<i>Syngnathus</i> .	241
<i>nachtriebli</i> .	166	<i>marshi, Characidium</i> .	100
margarita, <i>Clinostomus</i> .	116	<i>marstoni, Salmo</i> .	61
<i>Echistoma</i> .	70	<i>Salvelinus</i> .	61
<i>Margariscus</i> .	116	<i>marta, Cyprinodon</i> .	181
margaritaceus, <i>Entomacrodus</i> .	463	<i>marti, Pristigaster</i> .	45
Salaris.	463	<i>martinica, Athrina</i> .	248
margaritifer, <i>Myctophum</i> .	169	<i>Spileara</i> .	340
margaritatus, <i>Batrachus</i> .	487	<i>martinicensis, Acus</i> .	243
<i>Porichthys</i> .	487	<i>Aylopon</i> .	320
<i>Tropidichthys</i> .	500	<i>Menticirrhus</i> .	251
margaritifer, <i>Astatheros</i> .	421	<i>Nerophis</i> .	243
<i>Heros</i> .	421	<i>Novaculichthys</i> .	430
<i>Serranus</i> .	311	<i>Ocyanthias</i> .	323
margaritifer, <i>Heimulon</i> .	331	<i>Umbrina</i> .	351
<i>Aplodon</i> .	339	<i>Vomer</i> .	275
<i>Macrostromia</i> .	166	<i>Xyrichthys</i> .	430
margaritiferus, <i>Notoscopelus</i> .	166	<i>martinius, Gobius</i> .	442
margaritum, <i>Echiostomum</i> .	70	<i>Membras</i> .	248
margarotis, <i>Enneacanthus</i> .	304	<i>Smaris</i> .	340
marginalis, <i>Epinephelus</i> .	310	<i>Upeneus</i> .	343
<i>Hippocampus</i> .	244	mas, <i>Salmo</i> .	56
marginata, <i>Aporistia</i> .	231	<i>Mascalongus</i> .	172
<i>Helioperca</i> .	301	<i>maschalepsilus, Scarus</i> .	433
<i>Rissoa</i> .	485	<i>Sparisoma</i> .	433
<i>Uranidea</i> .	384	<i>mascotte, Moxostoma</i> .	110
marginatum, <i>Ophidium</i> .	485	<i>masoni, Salmo</i> .	68
marginatus, <i>Abudusdul</i> .	415	<i>masquinougy, Esox</i> .	172
<i>Bodianus</i> .	313	<i>immaculatus, Esox</i> .	173
<i>Brosnius</i> .	479	<i>ohiensis, Esox</i> .	172
<i>Brosmophycis</i> .	479	massachusetts, <i>Monacanthus</i> .	494
<i>Carenx</i> .	272	<i>massiliensis, Polyprion</i> .	308
<i>Chætodon</i> .	415	<i>Scorpæna</i> .	308
<i>Cottus</i> .	384	<i>Masticura</i> .	28
<i>Heliases</i> .	413	<i>matamorensis, Limia</i> .	192
<i>Hemicarax</i> .	271	<i>matejuelo, Amphiprion</i> .	235
<i>Neobithites</i> .	481	<i>mathematicus, Tetrodon</i> .	497
<i>Noturus</i> .	156	<i>mattowacca, Clupea</i> .	42
<i>Phycis</i> .	212, 213	<i>matutinus, Alburnellus</i> .	126
<i>Pimelodus</i> .	152	<i>Lythrurus</i> .	126
<i>Pomotis</i> .	301	<i>mauriceii, Argyrotaenia</i> .	275
<i>Serranus</i> .	311	<i>mauritii, Chaetodon</i> .	415
<i>Syphurus</i> .	231	<i>eleotris</i> .	268
<i>Xurel</i> .	272	<i>Grammlistes</i> .	332
marianum, <i>Holocentrum</i> .	236	<i>Maurolicidae</i> .	73
marianus, <i>Flammeo</i> .	236		
marina, <i>Bagre</i> .	148		
<i>capite striato, Perca</i> .	330		
<i>cauda forcipata, Brania</i> .	265		
<i>cauda nigra, Perca</i> .	330		
<i>gibbosa cinerea, Perca</i> .	329		
<i>Mugilops</i> .	252		

List of scientific names—Continued

	Page		Page
Maurolicus.....	73	megacephalus, Cyclogaster.....	400
amrhystino punctatus.....	73	Liparis.....	400
attenuatus.....	73	Mogaderia.....	94
lucotinus.....	73	megalops, Acipenser (Huso).....	35
oculatus.....	73	Doratonotus.....	429
pennanti.....	73	Megalobrycon.....	98
tripunctulatus.....	73	Megalocottus.....	387
maxilaris, <i>Iotella</i>	211	laticeps.....	388
<i>Murenoides</i>	468	platycephalus.....	388
Maxillingua.....	146	megalodon, <i>Pristis</i>	23
maxillingua, <i>Cyprinus</i>	146	Megalomea punctatum.....	159
Exoglossum.....	146	robustum.....	159
maxillosa, <i>Gnathopogon</i>	453	Megapereca.....	308
maxillosus, <i>Opistognathus</i>	453	Megalops atlanticus.....	39
<i>Rhinichthys</i>	139	cepediana.....	48
maximus, <i>Cetorhinus</i>	29	elongatus.....	39
<i>Hippoglossus</i>	221	notata.....	44
<i>Labrus</i>	425	ogilvia.....	44
<i>Lachnolaimus</i>	425	megalops albeolus, <i>Notropis</i>	129
<i>Squalus</i>	20	megalops, <i>Alburnus</i>	125
maxinkuckensis, <i>Alvordius</i>	284	<i>Clupea</i>	42
<i>Hadropterus</i>	284	<i>Cylindrostreus</i>	37
mayeri, <i>Mayerina</i>	81	<i>Cyprinus</i>	128
<i>mayeri</i>	81	<i>Eunichthys</i>	410
Maynea.....	476	<i>Holconotus</i>	410
<i>brunnina</i>	476	<i>Microponer</i>	349
<i>bulbiiceps</i>	377	<i>Opsopocodus</i>	115
<i>californica</i>	477	<i>Paranotropis</i>	125
<i>patagonica</i>	476	<i>Pimeleducus</i>	152
<i>pusilla</i>	477	<i>Trycharchodon</i>	115
mazara, <i>Makaira</i>	163	megalophthalmus, <i>Lutianus</i>	327
<i>Tetrapturus</i>	263	<i>Neomegus</i>	327
mazatlanica, <i>Seriola</i>	279	megalotilis, <i>Ichotis</i>	300
<i>Solea</i>	230	<i>Xenotis</i>	300
mazatlanum, <i>Hæmulon</i>	331	Megapercra.....	308
mazatlanus, <i>Achirus</i>	230	<i>ishinagi</i>	308
<i>Zonichthys</i>	270	Megaphalus.....	488
mciell, <i>Gambusia</i>	186	megastoma, <i>Opistognathus</i>	452
meadiae, <i>Imostoma</i>	286	Megastomatobus.....	102
<i>Ulocoentra</i>	286	<i>cyprinella</i>	102
meanyi, <i>Ruscarius</i>	378	megastomus, <i>Catostomus</i>	107
meurnsi, <i>Hesperoleucus</i>	122	Meladerma.....	281
<i>Notropis</i>	122	<i>nigerrima</i>	281
Meda.....	147	Melamphax beanii.....	233
<i>tulgida</i>	147	<i>bispinosus</i>	233
media, <i>Palometa</i>	267	<i>cristiceps</i>	233
Medialuna.....	358	<i>lugubris</i>	233
<i>californiensis</i>	358	<i>nycterinus</i>	233
mediana, <i>Ancistrus</i>	161	Melamphaidæa.....	232
medians, <i>Ancistrus</i>	161	melampodus, <i>Apogonichthys</i>	296
Medidat.....	147	melampterus, <i>Salmo</i>	55
mediocris, <i>Clupea</i>	41, 42	melampygus, <i>Caranx</i>	272
<i>Pomolobus</i>	42	<i>Xurel</i>	272
mediorum, <i>Antennarius puncto</i>	505	Melanichthys.....	338
Medipollis.....	45	Melaniris.....	248
medirostris, <i>Acipenser</i>	34	<i>balsanus</i>	248, 249
mediterranea, <i>Chimera</i>	32	<i>sardina</i>	249
<i>Saurida</i>	163	Melanobranchus.....	203
mediterraneus, <i>Enchelyopus</i>	208	<i>Bathygadus</i>	203
<i>Hoplostethus</i>	234	<i>melanocephalus</i> , <i>Plurgyrus</i>	145
<i>Lopharis</i>	280	Melanocetidae.....	610
<i>Scomber</i>	259	<i>Melanocetus</i>	610
<i>Sternoptyx</i>	74	<i>johsoni</i>	510
medius, <i>Anisarchus</i>	470	<i>(Liometus) murrayi</i>	511
<i>Calamus</i>	337	<i>niger</i>	510
<i>Centropterus</i>	306	<i>polyactis</i>	510
<i>Clinus</i>	470	<i>melanochir</i> , <i>Julius</i>	429
<i>Grammatous</i>	37	<i>Stellifer</i>	347
<i>Stromateus</i>	267	<i>Zestis</i>	347
medinus, <i>Myoxocephalus</i>	387	melanochira, <i>Belone</i>	196
medusicola, <i>Caranx</i>	272	melanogaster, <i>Pleuronectes</i>	224
<i>Xurel</i>	272	<i>Precilla</i>	191
medusophagus, <i>Schedophilus</i>	268	Melanogrammus.....	210
meeki, <i>Clethosoma</i>	416	<i>seglestius</i>	211
<i>Cichlaurus</i>	416	melanopogon, <i>Heros</i>	421
<i>Fucinostomus</i>	341	<i>melanopoma</i> , <i>Polynemus</i>	250
<i>Fundulus</i>	178	melanops, <i>Callirurus</i>	302, 303
<i>Holocentrus</i>	236	<i>Catostomus</i>	109
<i>Hybopsis</i>	139	<i>Dionda</i>	144
<i>Macropygia</i>	130	<i>Haplochilus</i>	185
<i>Microgobius</i>	444	<i>Ictiellus</i>	208
<i>Thyrsina</i>	249	<i>Minytrema</i>	109
<i>Ulæma</i>	341	<i>Sebastes</i>	364
megacephalus, <i>Calamus</i>	336	<i>Sebastodes</i>	365
<i>Chitonotus</i>	375	<i>Sebastosomus</i>	365

List of scientific names—Continued

Page	Page
melanopsis, <i>Diodon</i>	501
melanoptera, <i>Alepes</i>	271
melanopterum, <i>Pristipoma</i>	332
melanopterus, <i>Balistes</i>	493
melanopus, <i>Arius</i>	150
melanorhina, <i>Plectropoma</i>	321
Melanorhinus.....	249
bucki	249
melanospilum, <i>Diagramma</i>	332
melanostichus, <i>Psettichthys</i>	222
Melanostigma.....	478
gelatinosus	478
pammelas	478
melanostomus, <i>Eosebastes</i>	366
Galeus	12
Sebastodes	366
melanotata, <i>Limia</i>	191
melanothos, <i>Holocanthus</i>	497
melanotis, <i>Lamnuræna</i>	90
Muraena	90
Pseudojulis	428
Scarus	432
melanopterum, <i>Pristipoma</i>	332
melanotus, <i>Catostomus</i>	137
melanozonus, <i>Acanthophacelus</i>	191
Melanura.....	174
annulata	174
melanura, <i>Perca</i>	330
Faecilia	178
melanurum, <i>Hæmulon</i>	330
Læmonema	213
melanurus, <i>Careproctus</i>	402, 403
Cichlaurus	416
Cyprinus	128
Exocetus	201
Heros	416
Rutilus	110
melanzonus, <i>Acanthophacelus</i>	191
melapleura, <i>Gambusia</i>	185
Pecilia	185
melapleurus, <i>Zygonectes</i>	178
melas, <i>Ameiurus</i>	154
Silurus	154
melastomas, <i>Pristurus</i>	13
melastomus, <i>Galeus</i>	13
meleagris, <i>Muraena</i>	93
Rhinichthys	140
Meletta cerulea.....	43
libertate	44
petenensis	47
sueurii	42
venosa	42
Melicthys.....	403
bispinosus	493
piecus	493
radula	493
Mellotes.....	380
papilio	380
Memarchus.....	101
Membras.....	248
beani	248
martinicus	248
membras, <i>Clupea</i>	41
Menephorus.....	309
dubius	309
punctiferus	309
menhaden, <i>Alosa</i>	44
<i>Clupea</i>	44
Menidia.....	48, 247
audens	247
beryllina	247
cerea	247
chagresi	249
clara	247
conchorum	247
dentex	247
gilberti	248
lisa	249
menidia	247
notata	247
peninsulae	247
atrimentis	247
gardina	247
starksii	247
vagrans lacifata	248
menidia, <i>Atherina</i>	247
Meudia	247
menisorrah, <i>Carcharias</i>	15
Carcharhinus	16
Menona.....	178
menona, <i>Fundulus</i>	178
Zygonectes diaphanus	178
mentalalis, <i>Platycephalus</i>	192
Menticirrhus.....	350
americanus	350
martinicensis	351
nasus	351
panamensis	351
saxatilis	350
sinus	351
undulatus	350
Mentipercara.....	319
lucioperca	319
mento, <i>Balistes</i>	493
Bramopsis	410
Chlorophthalmus	165
Exocoetus	200
Heros	419
Mustelus	13
Paraliparis	404
Parapetenia	419
mentzelli, <i>Chironectes</i>	505
mentzelli, <i>Chironectes</i>	506
Serranus	311
meridionalis, <i>Amiurus</i>	152
Aphia	246
Carpioidea	104
Cottus	385
Ictalurus	182
Sclerognathus	104
Merinthæ.....	372
nematophtalma	372
Merlangus.....	214
leptocephalus	209
polaris	209
productus	214
merlangus, <i>Anoplopona</i>	373
Merluccius.....	214
ambiguus	214
augustimanus	214
billinearis	214
merluccius	214
productus	214
smilidus	214
merluccius, <i>Gadus</i>	214
Merluccius	214
Merlucciidae.....	214
Merluccius esculentus.....	214
lanatus	214
Merlus.....	214
gayi	214
merlus, <i>Gadus</i>	214
merou, <i>Holoconthus</i>	310
Merriamella.....	237
dorysia	237
merriami, <i>Empetrichthys</i>	182
mertensi, <i>Cottus</i>	387
Merulinus.....	406
carolinus	407
roseus	407
salmonicolor	407
scitulus	407
Merus.....	310
merus, <i>Centropristi</i>	312
mesæum, <i>Boleosoma</i>	287
mesæum, <i>Poecilichthys</i>	287
mesogaster, <i>Exocoetus</i>	200
Parexocoetus	200
Mesogonistius.....	303
chaetodon	303
Mesopion.....	324
ambiguus	327
analis	326
aratus	327
argentiniventralis	326
arnillo	328
aurovittatus	327
bucanella	326
campechanus	326
caudanotatus	326

List of scientific names—Continued

	Page		Page
<i>Mesopriion cyanopterus</i>	325	<i>microdon</i> , <i>Cheonda</i>	118
<i>cynodon</i>	325	<i>Cyclothona</i>	72
<i>elegans</i>	323	<i>Gobionellus</i>	441
<i>flavescens</i>	325	<i>Gobius</i>	441
<i>griseus</i>	325	<i>Gonostoma</i>	72
<i>guttatus</i>	326	<i>Osmerus</i>	66
<i>inermis</i>	327	<i>Pseudotriakis</i>	12
<i>isodon</i>	328	<i>Richardsonius</i>	118
<i>linea</i>	325	<i>Microdonotphis</i>	87
<i>litura</i>	325	<i>altipinnis</i>	87
<i>mahogoni</i>	327	<i>Microleotris</i>	437
<i>olaneo</i>	327	<i>mindii</i>	437
<i>pacificus</i>	325	<i>panamensis</i>	437
<i>pargus</i>	325	<i>Microgadus</i>	210
<i>pomacanthus</i>	324	<i>proximus</i>	210
<i>profundus</i>	326	<i>tomcod</i>	210
<i>retrospinis</i>	324	<i>micrognathus</i> , <i>Lonchopisthus</i>	453
<i>ricardi</i>	327	<i>Opistognathus</i>	453
<i>rosaceus</i>	328	<i>Microgobius</i>	443
<i>sobra</i>	326	<i>cyclolepis</i>	443
<i>uninotatus</i>	327	<i>emblematicus</i>	443
<i>vivanus</i>	326	<i>eulepis</i>	444
<i>vorax</i>	328	<i>gulosus</i>	443
<i>Mesops</i>	424	<i>meeki</i>	444
<i>Mesopus</i>	67	<i>miraflorensis</i>	444
<i>mesotrema</i> , <i>Asternotremia</i>	215	<i>signatus</i>	443
<i>messiori</i> , <i>Sirembo</i>	480	<i>thalassinus</i>	444
<i>messiniensis</i> , <i>Gymnocephalus</i>	267	<i>microlepidotus</i> , <i>Gila</i>	112
<i>metallica</i> , <i>Agosia</i>	140	<i>Microlepidotum</i>	211
<i>Echoneis</i>	449	<i>grandiceps</i>	211
<i>metallicus</i> , <i>Girardinus</i>	189	<i>Microlepidotus</i>	335
<i>Notropis</i>	124	<i>brevipinnis</i>	335
<i>Metaponops cooperi</i>	218	<i>inornatus</i>	335
<i>Metanops</i>	218	<i>microlepidotus</i> , <i>Cynodon</i>	355
<i>cooperi</i>	218	<i>Labrisomus</i>	459
<i>metopias</i> , <i>Triglops</i>	380	<i>Orthodon</i>	112
<i>metopocampus</i> , <i>Scopelus</i>	168	<i>Otolithus</i>	355
<i>Metrogaster</i>	409	<i>microlepis</i> , <i>Antimora</i>	211
<i>lineolatus</i>	409	<i>Epyrrhus</i>	98
<i>Metzelaaria</i>	370	<i>Hoplias</i>	99
<i>tridecmispinosa</i>	370	<i>Lepophidium</i>	485
<i>metzelaari</i> , <i>Evermannichthys</i>	447	<i>Leptophidium</i>	455
<i>Kyphosus</i>	339	<i>Macrodon</i>	90
<i>metzi</i> , <i>Gibbonsia</i>	458	<i>Trisopteris</i>	314
<i>meulenii</i> , <i>Diodon</i>	502	<i>microlepsis</i> , <i>Bathytrectes</i>	52
<i>mexicana</i> , <i>Aztecula</i>	143	<i>microleptes</i> , <i>Ariphaton</i>	309
<i>campi</i> , <i>Signalosa</i>	47	<i>microlophus</i> , <i>Dolophilichthys</i>	509
<i>Pompheris</i>	323	<i>Pomotis</i>	302
<i>Pecilia</i>	192	<i>Micromesus</i>	31
<i>Signalosa</i>	47	<i>Micrometrus</i>	409
<i>mexicanum</i> , <i>Myctophum</i>	168	<i>minimus</i>	409
<i>Nannobrachium</i>	168	<i>micronemus</i> , <i>Peristedion</i>	406
<i>mexicanus</i> , <i>Amblyopsis</i>	448	<i>Microporcas</i>	282, 294
<i>Astyanax</i>	98	<i>praeocularis</i>	294
<i>Auchenipterus</i>	457	<i>praelaris</i>	294
<i>Centroponus</i>	306	<i>punctulata</i>	294
<i>Chatotessus</i>	47	<i>micropora</i> , <i>Etheostoma</i>	294
<i>Chonophorus</i>	442	<i>micropthalmus</i> , <i>Haemulon</i>	320
<i>Fuguerae</i>	343	<i>Micropogon</i>	349
<i>Gerres</i>	343	<i>altipinnis</i>	349
<i>Gobius</i>	442	<i>argenteus</i>	349
<i>Mugil</i>	253	<i>ectenes</i>	349
<i>Pempheris</i>	323	<i>furnieri</i>	349
<i>Sauris</i>	163	<i>lineatus</i>	349
<i>Tetragonopterus</i>	96	<i>megalops</i>	349
<i>meyeri</i> , <i>Iridio</i>	427	<i>trifilis</i>	349
<i>mezquital</i> , <i>Chirostoma</i>	250	<i>undulatus</i>	349
<i>miacanthus</i> , <i>Artedielius</i>	378	<i>micropogon</i> , <i>Ceratichthys</i>	137
<i>miarchus</i> , <i>Anchoviella</i>	47	<i>Noconis</i>	137
<i>Stolephorus</i>	47	<i>Micropogonias</i>	349
<i>micraspidophorus</i> , <i>Cyclogaster</i> (<i>Neoliopsis</i>)	309	<i>micropogonias</i> , <i>Agonostoma</i>	254
<i>Neoliopsis</i>	309	<i>Atherina</i>	246
(<i>Micristius</i>) <i>brachypterus</i> , <i>Zygonctes</i>	185	<i>Belone</i>	198
<i>Micristodus</i>	18	<i>Catilurus</i>	299
<i>punctatus</i>	18	<i>Callyodon</i>	435
<i>microcephalus</i> , <i>Erythrinus</i>	99	<i>Carcharias</i>	16
<i>Gasterosteus</i>	237	<i>Caulolatilus</i>	357
<i>williamsoni</i>	237	<i>Catostomus</i>	106
<i>Pleuronectes</i>	228	<i>Corvina</i>	346
<i>Somniosus</i>	22	<i>Cottunculus</i>	392
<i>Squalus</i>	21, 22	<i>Dajaus</i>	254
<i>Microconger</i>	81	<i>Nebris</i>	354
<i>caudalis</i>	81	<i>Otolithus</i>	356
(<i>Microconger</i>) <i>caudalis</i> , <i>Leptocephalus</i>	81	<i>Pagellus</i>	336, 337
<i>Microdesmus</i>	472	<i>Pseudoscarus</i>	435
<i>dipus</i>	472	<i>Stellifer</i>	346
<i>retropinnis</i>	472	<i>Strongylura</i>	196

List of scientific names—Continued

	Page		Page
microptera, <i>Evolantia</i>	199	miniatus, <i>Sclerotis</i>	300
<i>Rhamdia</i>	157	<i>Sebastichthys</i>	365
Micropterus.....	297, 298	<i>Zoogneticus</i>	182
<i>dolomieu</i>	298	Minellus.....	133
<i>pseudoplites</i>	298	minima, <i>Capriscus</i>	496
<i>pseudotoides</i>	298	Cichla.....	298
<i>salmoides</i>	297	Clupea.....	41
Micropterus, <i>Etheostoma</i>	290	<i>Perca</i>	287
<i>Exocoetus</i>	199	minimus, <i>Cymatogaster</i>	409
<i>Pimelodus</i>	157	<i>Micrometrus</i>	409
Micropteryx.....	270, 279	miniopterus, <i>Scarus</i>	432
<i>cosmopolita</i>	276	Minnius.....	123
micropteryx, <i>Notropis</i>	124	<i>bellus</i>	126
<i>Alburnellus</i>	124	<i>dinemus</i>	123
<i>Platysomus</i>	275	<i>microstomus</i>	133, 134
microphthalmum, <i>Hæmulon</i>	329	<i>notatus</i>	145
microphthalmus, <i>Cichlaurus</i>	416	<i>nigripinnis</i>	125
<i>Dormitator</i>	436	<i>punctulatus</i>	126
<i>Heros</i>	416	<i>scepticus</i>	123
<i>Tetrabranchus</i>	76	<i>xenurus</i>	130
<i>Tetragonopterus</i>	96	Miromus.....	105
micropus, <i>Clupea</i>	46	<i>bardus</i>	104
<i>Etrumeus</i>	46	<i>delphinus</i>	104
<i>Gasterosteus</i>	238	<i>Jarrovii</i>	104
microrhynchus, <i>Acipenser</i>	34	<i>platyryhynchus</i>	104
micorrhinos, <i>Pseudoscarus</i>	435	minor, <i>Anarrhichthys</i>	472
micros, <i>Citharichthys</i>	219	Aphanopus.....	202
<i>Etropus</i>	219	Couchia.....	208
Microspathodon.....	415	Cyclopteruslisi paris.....	400
<i>azurissimus</i>	416	Lycichthys.....	472
<i>cinerous</i>	418	<i>Umbra</i>	255
microstigmus, <i>Myrophis</i>	83	minutillis, <i>Chirolepis</i>	438
Microstoma.....	228	minutus, <i>Cyclopterus</i>	397
microstoma, <i>Crenilabrus</i>	424	<i>Hemigrammus</i>	97
<i>Atherina</i>	246	Minytrema.....	109
<i>Scartella</i>	461	<i>melanops</i>	109
<i>Uranidea</i>	385	minytrema, <i>Gyrinichthys</i>	403
Microstomidae.....	68	mirabilis, <i>Exoglossum</i>	142
Microstomus.....	228	<i>mirabilis</i> , <i>Clupea</i>	33
<i>grönlandica</i>	68	<i>Crystallichthys</i>	41
<i>kitt</i>	228	<i>Gilllichthys</i>	444
<i>latidens</i>	228	<i>Lycodonus</i>	476
<i>pacificus</i>	228	<i>Phenacobius</i>	142
microstomus, <i>Blennius</i>	461	<i>miraflorensis</i> , <i>Microgobius</i>	444
<i>Carpoides</i>	104	<i>mirum</i> , <i>Calloptilum</i>	208
<i>Citharichthys</i>	219	<i>mispillensis</i> , <i>Amoiorus</i>	154
<i>Conger</i>	80	<i>mississippiensis</i> , <i>Morone</i>	307
<i>Etropus</i>	219	<i>Pristis</i>	23
<i>Iridio</i>	427	<i>missouriensis</i> , <i>Ilybopsis</i>	133
<i>Minnius</i>	133, 134	<i>mitchilli</i> , <i>Acanthocottus</i>	386
<i>Pleuronectes</i>	228	<i>Acipenser</i>	34
microtrachys, <i>Raja</i>	25	<i>altervata</i> , <i>Perca</i>	307
micrura, <i>Pteroplatea</i>	29	<i>Anchoviella</i>	48
<i>Raia</i>	29	<i>Argyrius</i>	275
micrurum, <i>Syacium</i>	218	<i>Cottus</i>	388
milberti, <i>Acipenser</i>	34	<i>Engraulis</i>	48
<i>Arius</i>	149	<i>interrupta</i> , <i>Perca</i>	307
<i>Carcharias</i>	16	<i>Perca</i>	307
<i>(Prionodon)</i>	16	<i>Stolephorus</i>	48
<i>Lepidosteus</i>	37	<i>Mitchillina</i>	52
milbertianus, <i>Syngnathus</i>	242	<i>hairdii</i>	52
miles, <i>Porogadus</i>	482	<i>mitrarius</i> , <i>Hesperoleucus</i>	122
<i>pectoralis</i> , <i>Prionotus</i>	407	<i>mitsukurii</i> , <i>Makaira</i>	263
milesi, <i>Pimephales</i>	145	<i>Tetraprurus</i>	203
millaris, <i>Gymnothorax</i>	92	<i>mitura</i> , <i>Rabida</i>	156
<i>Thrysoidea</i>	92	<i>miurus</i> , <i>Noturus</i>	156
millitaris, <i>Bellator</i>	408	<i>Ophichthys</i>	89
<i>Prionotus</i>	408	<i>Scyhalichthys</i>	89
<i>Silurus</i>	159	<i>Mixonus</i>	483
milktschitch, <i>Salmo</i>	54	<i>caudalis</i>	483
milleri, <i>Bathylagus</i>	69	<i>laticeps</i>	483
<i>Cichlasoma</i>	417	<i>mixtus</i> , <i>Gadus tomcodus</i>	210
<i>Theraps</i>	417	<i>mirolepis</i> , <i>Plectromus</i>	233
millipunctatus, <i>Gasterosteus</i>	239	<i>Scopelus</i>	233
milneri, <i>Noconis</i>	136	<i>Mnioperas</i>	455
<i>Pogellus</i>	337	<i>macrocephalus</i>	455
milneriana, <i>Hemitremia</i>	121	<i>Mobula</i>	31
milnerianus, <i>Phoxinus</i>	121	<i>auriculata</i>	31
minacea, <i>Gila</i>	114	<i>hypostoma</i>	32
mindii, <i>Stellifer</i>	347	<i>olfersi</i>	32
<i>Syngnathus</i>	242	<i>thyopostoma</i>	32
mineopas, <i>Brytus</i>	299	<i>mobular</i> , <i>Raja</i>	31
miniatum, <i>Peristedion</i>	405	<i>Mobilidae</i>	31
miniatus, <i>Leptopomus</i>	300	<i>modesta</i> , <i>Cheonda</i>	120
<i>Lepomis</i>	300	<i>Gambusia</i>	185, 192
<i>Rosicola</i>	365		

List of scientific names—Continued

Page	Page
modestus, <i>Julis</i>	428
Xyrichthys.....	430
<i>Mecbia</i>	481
<i>promelas</i>	481
<i>moresus</i> , <i>Gonopterus</i>	362
<i>Moharra</i>	342
<i>evermanni</i>	342
<i>rhombea</i>	342
<i>mohavensis</i> , <i>Siphateles</i>	123
<i>mohri</i> , <i>Clinus</i>	471
<i>mojarra</i> , <i>Cichlasoma</i>	417
<i>Theraps</i>	417
<i>Mola</i>	503
<i>mola</i>	503
<i>rotunda</i>	503
<i>mola</i> , <i>Dicdon</i>	503
<i>Mola</i>	503
<i>Tetronodon</i>	503
<i>Molacanthus</i>	503
<i>palasi</i>	503
<i>Molarii</i>	94
<i>molestrum</i> , <i>Gobiesox</i>	446
<i>Molidae</i>	503
<i>Molinessia fasciata</i>	192
<i>Molinessia</i>	191
<i>formica</i>	192
<i>gracilis</i>	102
<i>jonesi</i>	188
<i>latipinna</i>	192
<i>petenensis</i>	192
<i>sphenops</i>	192
<i>cuneata</i>	193
<i>vanderpolli</i>	193
<i>spilura</i>	102
<i>velifera</i>	192
<i>Mollinesia gracilis</i>	192
<i>latipinna</i>	191
<i>petenensis</i>	192
<i>velifera</i>	192
<i>molls</i> , <i>Aphyonus</i>	484
<i>Bathysaurus</i>	164
<i>Bothrocara</i>	477
<i>Careproctus</i>	403
<i>Pleuronectes</i>	229
<i>moluccensis</i> , <i>Apogon</i>	295
<i>Molva</i>	212
<i>huntila</i>	212
<i>molva</i>	212
<i>vulgaris</i>	212
<i>molva</i> , <i>Gadus</i>	212
<i>Molva</i>	212
<i>Molvella</i>	208
<i>borealis</i>	208
<i>Monacanthidae</i>	493
<i>Monacanthum proboscideum</i>	495
<i>Monacanthus</i>	494
<i>amphioxys</i>	494
<i>auriga</i>	494
<i>ciliatus</i>	494
<i>davidsonii</i>	494
<i>filamentosus</i>	494
<i>gallinula</i>	494
<i>hispidus</i>	494
<i>irroratus</i>	494
<i>macrocerus</i>	494
<i>macrurus</i>	493
<i>massachusettsensis</i>	494
<i>occidentalis</i>	494
<i>oppositus</i>	495
<i>pardalis</i>	493
<i>parranulus</i>	494
<i>piraenæ</i>	494
<i>pullus</i>	494
<i>punctatus</i>	494
<i>rupelli</i>	494
<i>setifer</i>	494
<i>signifer</i>	494
<i>spilonotus</i>	495
<i>striatus</i>	494
<i>tuckeri</i>	495
<i>varius</i>	494
<i>monacanthus</i> , <i>Plectropomia</i>	313
<i>monacus</i> , <i>Ceratichthys</i>	138
<i>Erimonax</i>	138
<i>monte</i> , <i>Stephanoberyx</i>	232
<i>monensis</i> , <i>Squalus</i>	19
<i>Moniana</i>	131
<i>aurata</i>	132
<i>boucardi</i>	132
<i>callisema</i>	132
<i>chihuahuensis</i>	132
<i>complanata</i>	131
<i>couchii</i>	131
<i>delicosa</i>	134
<i>formosa</i>	129, 131
<i>frigida</i>	131
<i>gibbosa</i>	131
<i>gracilis</i>	131
<i>jugalis</i>	131
<i>lataabilis</i>	131
<i>leonina</i>	131
<i>lutrensis</i>	131
<i>nitida</i>	135
<i>prosperina</i>	132
<i>pulchella</i>	131
<i>robusta</i>	132
<i>rutila</i>	131
<i>santamarin</i>	131
<i>tristis</i>	132
<i>Monoceratias</i>	508
<i>acanthias</i>	508
<i>monoceros</i> , <i>Aluterus</i>	496
<i>Balistes</i>	495, 496
<i>unicolor</i> , <i>Balistes</i>	496
<i>Monochir maculipinnis</i>	230
<i>reticulatus</i>	229
<i>Monolene</i>	220
<i>atrimana</i>	220
<i>sessilicauda</i>	220
<i>Monoimeropus</i>	480
<i>mallspinosus</i>	480
<i>Monomitonpus</i>	480
<i>torvus</i>	480
<i>Monomitra</i>	404
<i>Monoprion</i>	205
<i>maculatus</i>	205
<i>pigmentarius</i>	205
<i>Monopterhinus</i>	11
<i>monopterygius</i> , <i>Aspidophoroides</i>	397
<i>Cottus</i>	397
<i>Labrax</i>	374
<i>Pleurogrammus</i>	374
<i>monophthalmus</i> , <i>Auchenipterus</i>	456, 457
<i>Monostria</i>	344
<i>stahli</i>	344
<i>monstrosa</i> , <i>Chimera</i>	32
<i>montagui</i> , <i>Liparis</i>	399
<i>montana</i> , <i>Clueoanda</i>	119
<i>Macrhybopsis</i>	139
<i>montanus</i> , <i>Clinostomus</i>	119
<i>Hybopsis</i>	139
<i>Thymallus</i>	66
<i>Monterey</i>	391
<i>recalva</i>	391
<i>montereyensis</i> , <i>Gibbonsia elegans</i>	458
<i>Raja</i>	26
<i>montezuma</i> , <i>Astatheros</i>	421
<i>Ileros</i>	421
<i>montezumæ</i> , <i>Xiphophorus</i>	194
<i>monticola</i> , <i>Dajaus</i>	254
<i>Mugil</i>	254
<i>montregis</i> , <i>Clioia</i>	131
<i>moorei</i> , <i>Acteis</i>	400
<i>Congrammus</i>	454
<i>Malacoctenus</i>	460
<i>mordax abbotti</i> , <i>Osmerus</i>	67
<i>mordax</i> , <i>Atherina</i>	68
<i>Conger</i>	88
<i>Engraulis</i>	47
<i>Gymnophorax</i>	91
<i>Macrodon</i>	354
<i>Muraena</i>	91
<i>ophiurus</i> , <i>Macrodonophis</i>	88
<i>Osmerus</i>	67
<i>Sagenichthys</i>	354
<i>spectrum</i> , <i>Osmerus</i>	67
<i>morbibundus</i> , <i>Balistes</i>	492
<i>moringa</i> , <i>Gymnothorax</i>	91
<i>Muraena</i>	91
<i>Moringua</i>	89
<i>boekel</i>	89
<i>lateralis</i>	89

List of scientific names—Continued

	Page		Page
<i>Moringua linearis</i>	89	<i>Mugil berlandieri</i>	253
<i>lumbricoides</i>	89	<i>brasiliensis</i>	253
<i>moringua, Murena</i>	91	<i>cephalus</i>	252, 253
<i>Moringuidae</i>	89	<i>chanos</i>	40
<i>morlo, Centrolophus</i>	267	<i>cinereus</i>	341
<i>Epinephelus</i>	312	<i>curare</i>	253
<i>Serranus</i>	312	<i>gaimardianus</i>	254
<i>mormyrus, Campostoma</i>	149	<i>guentheri</i>	253
<i>Morone</i>	307	<i>guntheri</i>	253
<i>americanus</i>	307	<i>grunniens</i>	352
<i>flavescens</i>	281	<i>hospes</i>	254
<i>interrupta</i>	307	<i>incilis</i>	253
<i>maeulata</i>	302	<i>irretitus</i>	254
<i>mississippiensis</i>	307	<i>lebranchus</i>	253
<i>pallida</i>	307	<i>lineatus</i>	253
<i>rufa</i>	307	<i>liza</i>	253
<i>Moropidae</i>	307	<i>mexicanus</i>	253
<i>Morrhua</i>	210	<i>monticola</i>	254
<i>americanus</i>	210	<i>obliquus</i>	349
<i>californicus</i>	210	<i>petrosus</i>	253
<i>punctatus</i>	211	<i>plumieri</i>	253
<i>morrhua, Gadus</i>	210	<i>proboscideus</i>	254
<i>Serranus</i>	310	<i>rammelsbergii</i>	253
<i>morisli, Leptocephalus</i>	79	<i>salinoneus</i>	40
<i>Moseleya</i>	203	<i>setosus</i>	254
<i>moseri, Verasper</i>	223	<i>tang</i>	253
<i>motaguensis, Heros</i>	418	<i>thoburni</i>	253
<i>Parapetenis</i>	418	<i>trichodon</i>	254
<i>Pimelodus</i>	157	<i>Mugillidae</i>	252
<i>Rhamdia</i>	157	<i>mugiloides, Electris</i>	436
<i>Motella</i>	208	<i>Mugilogomorus</i>	39
<i>argentata</i>	208	<i>anna-carolina</i>	39
<i>caudacuta</i>	209	<i>Mugilops</i>	252
<i>ensis</i>	208	<i>cyanella</i>	252
<i>reinhardti</i>	208	<i>cyanellus</i>	252
<i>septentrionalis</i>	208	<i>marina</i>	252
<i>motta, Elacate</i>	281	<i>merinus</i>	252
<i>mouchetta, Glyphisodon</i>	415	<i>Müller, Benthosema</i>	170
<i>mowbrayi, Hsemulon</i>	329	<i>Leporinus</i>	98
<i>Rhinogobius</i>	329	<i>Pompheris</i>	323
<i>Moxostoma</i>	109	<i>Rhamphichthys</i>	101
<i>album</i>	110	<i>Salmo</i>	170
<i>alleghaniensis</i>	111	<i>Müller, Carcharias (Prionodon)</i>	18
<i>anisurum</i>	109	<i>Mullhypeneus</i>	343
<i>aureolum</i>	110	<i>Mullidae</i>	343
<i>austrinum</i>	110	<i>Mulloides</i>	344
<i>breviceps</i>	111	<i>rathbuni</i>	344
<i>bucco</i>	110	<i>Mullus</i>	343
<i>campbelli</i>	109	<i>auratus</i>	343
<i>claviformis</i>	109	<i>barbatus</i>	343
<i>collapsum</i>	110	<i>auratus</i>	343
<i>congestum</i>	110	<i>bifasciatus</i>	343
<i>conus</i>	111	<i>flavolineatus</i>	344
<i>coregonus</i>	110	<i>imberbis</i>	295
<i>crassilabre</i>	111	<i>maculatus</i>	343
<i>kennerlyi</i>	109	<i>multifasciata, Adinia</i>	180
<i>lesueuri</i>	111	<i>Hydryarga</i>	178
<i>macrolepidotum</i>	111	<i>Parapetenia</i>	418
<i>mascotte</i>	110	<i>multifasciatum, Cichlasoma</i>	418
<i>papillosum</i>	109	<i>Gobiosoma</i>	446
<i>parvidens</i>	110	<i>multifasciatus, Pronotogrammus</i>	322
<i>pidiense</i>	110	<i>multifilis, Blennius</i>	461, 462
<i>pœciliurum</i>	111	<i>multiguttatum, Plectropoma</i>	313
<i>robustum</i>	111	<i>multiguttatus, Alphestes</i>	313
<i>rupiscartes</i>	111	<i>multilineata, Furcaria</i>	412
<i>tenue</i>	109	<i>Poecilia</i>	192
<i>thalassinum</i>	110	<i>multilineatum, Hæmulon</i>	330
<i>trisignatum</i>	105	<i>multilineatus, Heliases</i>	412
<i>valenciennei</i>	109	<i>Labrax</i>	307
<i>victorie</i>	109	<i>multimaculatus, Diodon</i>	601
<i>mucosum, Xiphidion</i>	469	<i>multilocellatus, Antennarius</i>	506
<i>mucusum, Acanthocottus</i>	387	<i>Chironectes</i>	506
<i>Liparis</i>	399	<i>multipunctata, Skiffia</i>	183
<i>Lycodesalepis</i>	476	<i>multipunctatum, Xenendum</i>	183
<i>Lycodes</i>	476	<i>multiradiatus, Arius</i>	151
<i>Neoliparis</i>	399	<i>Cbaracodon</i>	183
<i>Xiphister</i>	469	<i>Cubiceps</i>	288
<i>mucronata, Perca</i>	308	<i>Lermichthys</i>	183
<i>mucronatum, Ophidium</i>	467	<i>multispinosa, Herotilapia</i>	423
<i>mucronatus, Neoconger</i>	81	<i>multispinosus, Heros</i>	423
<i>Odontognathus</i>	45	<i>multicellata, Murena</i>	92
<i>mudfish, Fundulus</i>	175	<i>multofasciata, Sciaena</i>	349
<i>muelleri, Lepidomegas</i>	279	<i>mundeoila, Anchoviella</i>	50
<i>Scopelus</i>	170	<i>mundulus, Stolephorus</i>	50
<i>Mugil</i>	252, 253	<i>mundiceps, Xyrichtys</i>	430
<i>albus</i>	253	<i>mundicorpus, Inistius</i>	430

List of scientific names—Continued

	Page		Page
<i>mundus</i> , <i>Dactylagnus</i>	455	<i>Murænophis punctata</i>	92
<i>Oligoplites</i>	278	<i>vicina</i>	92
<i>Urotrygon</i>	30	<i>Murænopsis</i>	87
<i>Muræna</i>	77, 90	<i>triserialis</i>	88
<i>acuminata</i>	85	<i>murina</i> , <i>Callirurus</i>	299
<i>africana</i>	91	<i>murrayi</i> , <i>Ipnops</i>	170
<i>anguilla</i>	77	<i>Lycoetus</i>	511
<i>anguina</i>	90	<i>Melanocetus</i> (<i>Lioctetus</i>).....	511
<i>aqua-dulcis</i>	91, 93	<i>muscarum</i> , <i>Gobiesox</i>	490
<i>argentea</i>	77	<i>musculus</i> , <i>Cyclopterus</i>	400
<i>balaericana</i>	79	<i>muskoka</i> , <i>Notropis</i>	134
<i>bostonensis</i>	77	<i>mustacinus</i> , <i>Serranus</i>	310
<i>clepsydra</i>	90	<i>Mustela</i>	208
<i>cœca</i>	84	<i>mustela</i> , <i>Gadus</i>	208
<i>conger</i>	79	<i>Mustèles</i> , <i>Les</i>	208
<i>dovii</i>	92	<i>mustellurus</i> , <i>Gaidropsarurus</i>	208
<i>elaborata</i>	93	<i>Mustellus stellatus</i>	13
<i>erebus</i>	92, 93	<i>Mustelus</i>	13, 14, 17
<i>havenensis</i>	87	<i>asterias</i>	13
<i>heleena</i>	90	<i>californicus</i>	13
<i>infernalis</i>	92	<i>dorsalis</i>	13
<i>insularum</i>	90	<i>folis</i>	14
<i>lentiginosa</i>	90	<i>lævis</i>	13
<i>lineopinnis</i>	92	<i>lunulatus</i>	13
<i>lita</i>	93	<i>mento</i>	13
<i>longicauda</i>	91	<i>mustelus</i>	13
<i>lumbircus</i>	78	<i>plebejus</i>	13
<i>macrocephala</i>	77	<i>Mustelus</i> , <i>Galeus</i>	13
<i>maculata nigra</i>	91	<i>Squalus</i>	13
<i>et viridis</i>	92	<i>muticus</i> , <i>Lethotrema</i>	13, 14
<i>maculosa</i>	87	<i>Mycteroperca</i>	314
<i>melanotis</i>	90	<i>bonaci xanthosticta</i>	314
<i>meleagris</i>	93	<i>boulegeri</i>	313
<i>mordax</i>	91	<i>bowersi</i>	314
<i>moringa</i>	91	<i>calliura</i>	315
<i>multicellata</i>	92	<i>camelopardalis</i>	315
<i>myialeucosticta</i>	90	<i>falcata</i>	315
<i>nigra</i>	92	<i>phenax</i>	315
<i>nigriceps</i>	79	<i>hopkinsi</i>	314
<i>ophis</i>	90	<i>olfax</i>	314
<i>panamepsis</i>	87, 94	<i>pardalis</i>	315
<i>pantherina</i>	91	<i>phenax</i>	315
<i>pfeifferi</i>	93	<i>rosacea</i>	314
<i>plata</i>	93	<i>tigris</i>	315
<i>pinata</i>	78	<i>venadatum</i>	314
<i>pinta</i>	90	<i>xenarcha</i>	314
<i>pintita</i>	92	<i>Myrophidæ</i>	166
<i>polyuranodon</i>	91	<i>Myctophum</i>	169
<i>punctata</i>	91	<i>affine</i>	169
<i>ratiabouri</i>	89	<i>atratum</i>	160
<i>retifera</i>	90	<i>aurolateratum</i>	170
<i>rostrata</i>	77	<i>benolii</i>	169
<i>sancæ-helene</i>	92	<i>boops</i>	169
<i>savanna</i>	78	<i>californiense</i>	169
<i>serpentina</i>	77	<i>crenulare</i>	170
<i>seu conger brasiliensis</i>	93	<i>gracile</i>	169
<i>siderea</i>	93	<i>hians</i>	169
<i>sordida</i>	93	<i>humboldtii</i>	169
<i>supremo</i> etc.....	79	<i>hygomi</i>	169
<i>unicolor</i> etc.....	90	<i>laternatum</i>	170
<i>variegata</i>	93	<i>luminosum</i>	170
<i>Murexæna</i> , <i>Callechelys</i>	86	<i>margaritatum</i>	169
<i>Lycodes</i>	475	<i>mexicanum</i>	168
<i>Murexænidæ</i>	78	<i>nannochir</i>	167
<i>Murexænosidæ</i>	78	<i>nitidulum</i>	169
<i>Murexænosor</i>	78	<i>nocturnum</i>	168
<i>coniceps</i>	79	<i>oculeum</i>	169
<i>savanna</i>	78	<i>punctatum</i>	169
<i>tricuspidata</i>	78	<i>procularium</i>	170
<i>Murexænidæ</i>	8, 94	<i>protoctulus</i>	168
<i>Murexænoblennia</i>	8	<i>radmésquel</i>	168
<i>olivacea</i>	8	<i>regale</i>	168
<i>Murexænoidæ</i>	467	<i>remiger</i>	169
<i>maxillaris</i>	468	(<i>Stenobrachius</i>) <i>leucopsarum</i>	168
<i>sujei</i>	467	<i>tenaculum</i>	170
<i>Murexænoidæ</i> , <i>Bleennius</i>	467	<i>towsendi</i>	167
<i>Gymnelus</i>	468	<i>mydrus</i> , <i>Cyprinodon</i>	180
<i>Murexænophis</i>	90	<i>myersi</i> , <i>Gambusia</i>	185
<i>caramuru</i>	91	<i>Mykiss</i>	56
<i>catenula</i>	94	<i>mykiss</i> , <i>Salmo</i>	50
<i>curvilineata</i>	91	<i>agua-bonita</i> , <i>Salmo</i>	50
<i>marmoratus</i>	91	<i>Myliobatis</i>	31
<i>ocellatus</i>	87	<i>acuta</i>	31
<i>pantherina</i>	93	<i>asperrimus</i>	30
		<i>bispinosus</i>	31

List of scientific names—Continued

	Page		Page
<i>Myliobatis californicus</i>	31	<i>Myxostoma euryops</i>	110
<i>eeitenkee</i>	30	<i>peccilura</i>	111
<i>fremainvilli</i>	31	<i>Myxus harengus</i>	253
<i>goodei</i>	31	<i>Myzopsetta</i>	226
<i>marginitatus</i>	31		
<i>vespertilio</i>	31		
<i>Mylocheilus</i>	112	<i>nachtriebi</i> , <i>Leuciscus</i>	116
<i>caurinus</i>	118	<i>Margariscus</i>	116
<i>fraterculus</i>	112	<i>nacrea</i> , <i>Gila</i>	114
<i>lateralis</i>	112	<i>naimai</i> , <i>Sparactodon</i>	280
<i>Myloleucus</i>	119, 120	<i>namaycush</i> , <i>Cristivomer</i>	59
<i>parovanus</i>	121	<i>Saimo</i>	59
<i>pulverentulus</i>	119, 120	<i>Nannobrachium</i>	167
<i>thalassinus</i>	122	<i>leucopsarum</i>	168
<i>Mylopharodon</i>	112	<i>macdonaldi</i>	168
<i>conocephalus</i>	112, 113	<i>mexicanum</i>	168
<i>robustus</i>	112, 113	<i>mannochir</i>	167
<i>Myrophina</i>	31	<i>nigrum</i>	167
<i>myops</i> , <i>Salmo</i>	163	<i>regale</i>	168
<i>Saurus</i>	163	<i>Nannocampus</i>	244
<i>Trachinocephalus</i>	163	<i>nanus</i>	244
<i>Myrocephalus</i>	387	<i>subroseus</i>	244
<i>axillaris</i>	387	<i>nannochar</i> , <i>Myctophum</i>	167
<i>jaok</i>	387	<i>Nannobrachium</i>	167
<i>medinii</i>	387	<i>Nannostomus</i>	289
<i>niger</i>	387	<i>nanomyzon</i> , <i>Catostomus</i>	106
<i>parvulus</i>	387	<i>Nanostoma</i>	289
<i>polyacanthocephalus</i>	387	<i>arcansanum</i>	289
<i>stelleri</i>	387	<i>elegans</i>	289
<i>ve.rucosus</i>	387	<i>inscriptum</i>	289
<i>myrialeucosticta</i> , <i>Muraena</i>	90	<i>rupestre</i>	289
<i>myrialeucostictus</i> , <i>Muraena</i>	92	<i>swanana</i>	289
<i>myriaster</i> , <i>Rhomboichthys</i>	217	<i>thalassatum</i>	289
<i>Myrichthys</i>	85	<i>vinetipes</i>	289
<i>acuminatus</i>	85	<i>zonale</i>	289
<i>oculatus</i>	85	<i>Nansenia</i>	68
<i>pantostigmus</i>	85	<i>greenlandica</i>	68
<i>tigrinus</i>	85	<i>nanus</i> , <i>Engraulis</i>	47
<i>xysturus</i>	85	<i>Gadus</i>	210
<i>Myrolepis</i>	373	<i>Liparis</i>	401
<i>zonifer</i>	373	<i>Nannocampus</i>	244
<i>Myripristis</i>	235	<i>Narcacion</i>	27
<i>fulgens</i>	234	<i>Narcaciontes</i>	27
<i>occidentalis</i>	235	<i>Narcetes</i>	52
<i>trachypoma</i>	234	<i>erimelas</i>	52
<i>Myripristis</i>	235	<i>pluriserialis</i>	52
<i>humilis</i>	235	<i>stomias</i>	52
<i>jacobus</i>	235	<i>Narcine</i>	27
<i>japonicus</i>	234	<i>brasiliensis</i>	27
<i>lima</i>	235	<i>corallina</i>	27
<i>occidentalis</i>	235	<i>ntemedor</i>	27
<i>poecilopus</i>	235	<i>indica</i>	27
<i>Myrophis</i>	82	<i>nigra</i>	27
<i>egmontis</i>	82	<i>umbrosa</i>	27
<i>lumbriacus</i>	83	<i>naresi</i> , <i>Salmo</i>	61
<i>microstigmus</i>	83	<i>Salvelinus</i>	61
<i>punctatus</i>	82, 83	<i>narinari</i> , <i>Raia</i>	30, 31
<i>vafet</i>	83	<i>Stoasonodon</i>	30
<i>mystacalis</i> , <i>Lepomis</i>	300	<i>naso</i> , <i>Anchoviella</i>	49
<i>mystacina</i> , <i>Gnathoprops</i>	453	<i>Stolephorus</i>	49
<i>mystacinus</i> , <i>Epinephelus</i>	210	<i>nasus</i> , <i>Acanthonotus</i>	75
<i>Gnathoprops</i>	453	<i>Bassozetus</i>	481
<i>Serranus</i>	310	<i>Diplanchus</i>	503
<i>mystes</i> , <i>Scorpaena</i>	371	<i>Lamna</i>	19
<i>mysticetus</i> , <i>Cetengraulis</i>	51	<i>Menticirrhus</i>	351
<i>Engraulis</i>	51	<i>Notacanthus</i>	75
<i>mystinus</i> , <i>Sebastichthys</i>	364	<i>Squalus</i>	19
<i>Sebastosomus</i>	365	<i>Umbrina</i>	351
<i>Mystrophis</i>	88	<i>nasuta</i> , <i>Malthcea</i>	507
<i>interstitius</i>	88	<i>Netuma</i>	149
<i>Mytilophagus</i>	410	<i>nasutum</i> , <i>Agonostoma</i>	254
<i>fasciatus</i>	410	<i>Campistoma</i>	146
<i>Myxine</i>	8, 94	<i>nasutus</i> , <i>Astyanax</i>	97
<i>glutinosa</i>	8	<i>Cantherhines</i>	493, 494
<i>limosa</i>	8	<i>Leuciscus</i>	139
<i>Myxinidae</i>	8	<i>Ogcocephalus</i>	507
<i>Myxodagnus</i>	455	<i>Trachynotus</i>	276
<i>opercularis</i>	455	<i>natalis</i> , <i>Ameiurus</i>	153
<i>Myxodes elegans</i>	458	<i>analis</i> , <i>Ameiurus</i>	153
<i>lugubris</i>	460	<i>Pimelodus</i>	153
<i>macropus</i>	460	<i>natator</i> , <i>Carcharhinus</i>	17
<i>varius</i>	459	<i>Carcharias</i>	17
<i>versicolor</i>	460	<i>Nauchelus</i>	278
<i>Myxostoma</i>	109	<i>abbreviatus</i>	278
<i>austrina</i>	110	<i>annularis</i>	278
		<i>brachycentrus</i>	278

List of scientific names—Continued

	Page		Page
<i>Myxostoma compressus</i>	278	<i>Nemabrama</i>	234
<i>lecurus</i>	278	<i>Nematistiidae</i>	280
<i>triacanthus</i>	278	<i>Nematistius</i>	280
<i>Naucrates</i>	239, 278	<i>pectoralis</i>	280
<i>cyanophrys</i>	278	<i>Nematognathi</i>	147
<i>duotor</i>	278	<i>Nematonurus</i>	203
<i>fanfarus</i>	278	<i>abyssorum</i>	203
<i>noveboracensis</i>	278	<i>acrolepis</i>	203
<i>polysarcus</i>	278	<i>armatus</i>	203
<i>serratus</i>	278	<i>cyclolepis</i>	203
<i>naucrates</i> , <i>Echeneis</i>	448	<i>goodii</i>	203
<i>nauplagium</i> , <i>Balistes</i>	451	<i>suborbitalis</i>	203
<i>Nautichthys</i>	382	(<i>Nematonurus</i>) <i>magnus</i> , <i>Macrurus</i>	204
<i>oculofasciatus</i>	382	<i>Nematonurus</i>	482
<i>pribilovius</i>	382	<i>pectoralis</i>	482
<i>Nautiscus</i>	382	<i>nematophthalma</i> , <i>Merinthos</i>	372
<i>pribilovius</i>	382	<i>nematophthalmus</i> , <i>Sebastes</i>	372
<i>Nautopædium</i>	487	<i>nematopus</i> , <i>Neotropis</i>	423
<i>porosissimum</i>	487	<i>Physiculus</i>	212
<i>nantopædium</i> , <i>Porichthys</i>	487	<i>Nemicichthysidae</i>	83
<i>navarroensis</i> , <i>Hesperoleucus</i>	121	<i>Nemicichthys</i>	83
<i>newaga</i> , <i>Eleginops</i>	210	<i>avocetta</i>	83
<i>Gadus</i>	239, 210	<i>fronto</i>	83
<i>nazuz</i> , <i>Nototropis</i>	126	<i>infans</i>	83
<i>Oreella</i>	126	<i>scolopacea</i>	83
<i>Neatolus</i>	251	<i>scolopaceus</i>	83
<i>tripes</i>	261	<i>Nemocampsis</i>	297, 298
<i>neapolitanus</i> , <i>Lampugus</i>	265	<i>nemoptera</i> , <i>Dixonina</i>	40
<i>nebrasicensis</i> , <i>Nocomis</i>	137	<i>neo-boracensis</i> , <i>Fistularia</i>	245
<i>Nebris</i>	354	<i>Neobrythites</i>	481
<i>microps</i>	354	<i>gilli</i>	481
<i>occidentalis</i>	354	<i>marginatus</i>	481
<i>zestus</i>	354	<i>ocellatus</i>	481
<i>Nebrilius concolor</i>	94	<i>robustus</i>	481
<i>nebularis</i> , <i>Platophrrys</i>	217	<i>stelliferoides</i>	482
<i>nebulifer</i> , <i>Heros</i>	417	<i>Neoclinus</i>	458
<i>Labrax</i>	317	<i>blanchardi</i>	458
<i>Paralabrax</i>	317	<i>satricus</i>	458
<i>Theraps</i>	417	<i>Neoconger</i>	81
<i>nebuliferus</i> , <i>Catostomus</i>	104	<i>neuronotus</i>	81
<i>nebulopunctatus</i> , <i>Gobius</i>	440	<i>perlongus</i>	82
<i>nebulosa</i> , <i>Acedia</i>	231	<i>vermiformis</i>	81
<i>Aphoristia</i>	231	<i>neogaea</i> , <i>Pfrille</i>	121
<i>Cerna</i>	315	<i>neogaeus</i> , <i>Phoxinus</i>	121
<i>Perca (Percina)</i>	283	<i>neoguinaica</i> , <i>Albula</i>	40
<i>Saurida</i>	164	<i>neohantonus</i> , <i>Coregonus</i>	64
<i>nebulosus</i> , <i>Ameturus</i>	154	<i>Neolidaris</i>	399
<i>Apionichthys</i>	230	<i>atlanticus</i>	399
<i>catulus</i> , <i>Ameiurus</i>	154	<i>beringianus</i>	399
<i>Chiropsis</i>	373, 374	<i>bristolensis</i>	399
<i>Clinus</i>	471	<i>callyodon</i>	399
<i>Erisson</i>	355	<i>fissuratus</i>	399
<i>Gasterosteus</i>	239	<i>floræ</i>	399
<i>Lycias</i>	475	<i>grebniutzkii</i>	399
<i>Lycodes</i>	475	<i>greeni</i>	399
<i>Otolithus</i>	354, 355	<i>micraspidophorus</i>	399
<i>Pimelodus</i>	154, 155	<i>mucosus</i>	399
<i>Pteropodus</i>	369	<i>rutteri</i>	399
<i>Sebastes</i>	369	(<i>Neoliparis</i>) <i>beringianus</i> , <i>Cyclogaster</i>	399
<i>Silurus</i>	155	<i>micraspidophorus</i> , <i>Cyclogaster</i>	399
<i>Urobatis</i>	30	<i>Neomänis</i>	324
<i>Urolophus</i>	30	<i>blackfordi</i>	453
<i>Zeus</i>	232	<i>hastingsi</i>	326
<i>Nectoliparis</i>	402	<i>jordani</i>	324
<i>pelagicus</i>	402	<i>megalophthalmus</i>	327
<i>Nector</i>	346	<i>Neomugil</i>	254
<i>alutus</i>	346	<i>digueti</i>	254
<i>armatus</i>	346	<i>Neomursea</i>	91
<i>chrysoleucus</i>	346	<i>nigromarginata</i>	93
<i>verte-crucis</i>	346	<i>Neopeltilia</i>	190
<i>nectaria</i> , <i>Gymnomurrena</i>	95	<i>holacanthus</i>	190
<i>necturus</i> , <i>Uropterygius</i>	94, 95	<i>Neostoma</i>	72
<i>Neotropus</i>	423	<i>bathophilum</i>	72
<i>bocourti</i>	423	<i>Neothunnus</i>	260
<i>carpintis</i>	423	<i>albaces</i>	260
<i>nematopus</i>	423	<i>allisoni</i>	260
<i>sieboldii</i>	423	<i>catalinæ</i>	260
<i>nefastus</i> , <i>Pomotis</i>	300	<i>nepenthe</i> , <i>Eupomacentrus</i>	414
<i>neglecta</i> , <i>Clupea</i>	44	<i>nephela</i> , <i>Helicoperca</i>	301
<i>neglectus</i> , <i>Amblyodon</i>	352	<i>nephelus</i> , <i>Lepomis</i>	301
<i>Aploidonitus</i>	352	<i>Pleuronichthys</i>	225
<i>nelsoni</i> , <i>Awsous</i>	442	<i>Spherooides</i>	498
<i>Chonophorus</i>	442	<i>Tetradon</i>	498
<i>Conorhynchus</i>	166	<i>nerka</i> , <i>Oncorhynchus</i>	55
<i>Coregonus</i>	65	<i>Salmo</i>	55
<i>Salmo</i>	59	<i>Nerophis angulineus</i>	243

List of scientific names—Continued

	Page		Page
<i>Nerophis heckeli</i>	244	<i>nigricans</i> , <i>Cottus</i>	386
<i>maculata</i>	243	<i>Cyprinellus</i>	201
<i>murtimicensis</i>	243	<i>Enchelycore</i>	90
<i>nertinianus</i> , <i>Acipenser (Huuso)</i>	35	<i>Entomacrodus</i>	463
<i>Nettastoma procerum</i>	82	<i>etowanus</i> , <i>Catostomus</i>	107
<i>Nettastomidae</i>	82	<i>Exocetus</i>	201
<i>Netuma</i>	149, 150	<i>Girella</i>	339
<i>elattura</i>	149	<i>Heros</i>	419
<i>insculpta</i>	149	<i>Huro</i>	297
<i>kessleri</i>	149	<i>Hypentelium</i>	107
<i>nasuta</i>	149	<i>Hypoplectrus unicolor</i>	321
<i>oscula</i>	149	<i>Labrax</i>	308
<i>planiceps</i>	149	<i>Makaira</i>	263
<i>platypogon</i>	149	<i>Muraena</i>	90
<i>thalassina</i>	149	<i>Parapetenia</i>	419
<i>neurates</i> , <i>Echeneis</i>	448	<i>Petromyzon</i>	8
<i>neucratoides</i> , <i>Echeneis</i>	449	<i>Pimeledus</i>	152
<i>neustratus</i> , <i>Gasterosteus</i>	238	<i>Plectropoma</i>	321
<i>nevadensis</i> , <i>Apocope</i>	141	<i>nigriculus</i> , <i>Acrorurus</i>	363
<i>Cyprinodon</i>	181	<i>Serranus</i>	310
<i>Rhinichthys (Apocope)</i>	141	<i>nigrilabris</i> , <i>Oriomis</i>	154
<i>nevicense</i> , <i>Etheostoma</i>	283	<i>nigripinne</i> , <i>Corallicola</i>	456
<i>newberrii</i> , <i>Fario</i>	58	<i>nigripinnis</i> , <i>Argyrosomus</i>	62, 63
<i>Salmo</i>	58	<i>Bathyagonus</i>	305, 398
<i>newberryi</i> , <i>Eucyclogobius</i>	444	<i>Cephalopolis</i>	309
<i>Gobius</i>	444	<i>Clinus</i>	456
<i>newmani</i> , <i>Hyostoma</i>	286	<i>Coregonus</i>	63
<i>Nexilarius</i>	414	<i>Cotylis</i>	489
<i>concolor</i>	414	<i>Gobiesox</i>	489
<i>nianguae</i> , <i>Etheostoma (Hudropterus)</i>	285	<i>Leucichthys</i>	63
<i>Hypobomus</i>	285	<i>Minilus</i>	125
<i>spilotum</i> , <i>Etheostoma</i>	285	<i>Promicropodus</i>	317
<i>nicaraguensis</i> , <i>Carcharias</i>	16	<i>Rhypticus</i>	317
<i>Eulamia</i>	16	<i>Sirembo</i>	480
<i>Gambusia</i>	184, 186	<i>nigrirostris</i> , <i>Chetodon</i>	300
<i>Heros</i>	418	<i>Sarotherodon</i>	360
<i>Pimeledus</i>	157	<i>nigrita</i> , <i>Garrupa</i>	312
<i>Rhamdia</i>	157	<i>nigritum</i> , <i>Clechosoma</i>	418
<i>Theraps</i>	418	<i>nigritus</i> , <i>Serranus</i>	312
<i>nichoisi</i> , <i>Irigio</i>	428	<i>Theraps</i>	418
<i>Platyglossus</i>	428	<i>nigrocastaneus</i> , <i>Enchelycore</i>	90
<i>nicholsi</i> , <i>Gobius</i>	440	<i>Gymnothorax</i>	90
<i>Rhinogobius</i>	440	<i>nigrocinctus</i> , <i>Sebastes</i>	370
<i>Nicholsina</i>	431	<i>Sebastichthys</i>	370
<i>nieuhoffi</i> , <i>Echenoës</i>	449	<i>Sebastodes</i>	304
<i>niger</i> , <i>Astrostethes</i>	71	<i>nigrofasciata</i> , <i>Hydrarchyra</i>	176
<i>Bubalicthys</i>	102	<i>Limia</i>	191
<i>Carpioles</i>	102	<i>nigrofasciatus</i> , <i>Archocentrus</i>	420
<i>Centrolophus</i>	267	<i>Hadropterus</i>	283
<i>Chiasonodon</i>	452	<i>Heros</i>	420
<i>Cottus</i>	387	<i>nigrolineatus</i> , <i>Diodon</i>	602
<i>Esoclar</i>	261	<i>nigro-maculatus</i> , <i>Cantharus</i>	305
<i>Esox</i>	173	<i>nigromaculatus</i> , <i>Rhypticus</i>	316
<i>Gobius</i>	440	<i>nigromarginata</i> , <i>Neomururaena</i>	91, 93
<i>Lamprogrammus</i>	484	<i>nigromarginatus</i> , <i>Gymnothorax ocellatus</i>	93
<i>Malacosteus</i>	71	<i>nigrotæniata</i> , <i>Hybognathus</i>	144
<i>Melanocetus</i>	510	<i>nigrotæniatus</i> , <i>Graodus</i>	132, 133
<i>Myoxocephalus</i>	387	<i>nigrum</i> , <i>Boleosoma</i>	287
<i>Sparus</i>	265	<i>Petromyzon</i>	9
<i>Tautoga</i>	424	<i>nimbaria</i> , <i>Vinciguerra</i>	73
<i>Thyrsites</i>	281	<i>nimbarius</i> , <i>Zalargas</i>	73
<i>nigerrima</i> , <i>Meladerma</i>	281	<i>niphobles</i> , <i>Epinephelus</i>	312
<i>nigra</i> , <i>Astrostethes</i>	71	<i>Sparisoma</i>	432
<i>Balistes</i>	493	<i>niphonius</i> , <i>Priacanthus</i>	323
<i>Dicrolene</i>	483	<i>nipigon</i> , <i>Leucichthys</i>	64
<i>Etheostoma</i>	287	<i>nisorius</i> , <i>Fundulus</i>	175
<i>Gila</i>	119	<i>nitens</i> , <i>Anago</i>	80
<i>Muraena</i>	79	<i>Ophidion</i>	80
<i>maculata</i>	91, 92	<i>Trichiurus</i>	282
<i>Narcine</i>	27	<i>nitida</i> , <i>Hybopsis</i>	135
<i>Perca</i>	267	<i>Julis</i>	429
<i>marina cauda</i>	330	<i>Moniana</i>	135
<i>nigrellus</i> , <i>Hyborhynchus</i>	145	<i>Pomotis</i>	300
<i>nigrescens</i> , <i>Centropomus</i>	306	<i>nitidissima</i> , <i>Julis</i>	429
<i>Coryphaena</i>	317	<i>nitidulum</i> , <i>Myctophum</i>	169
<i>Cycleptus</i>	102	<i>nitidum</i> , <i>Pristipomus (Hemulopsis)</i>	333
<i>Dorsuarus</i>	339	<i>Thalassoma</i>	429
<i>Exoglossum</i>	146	<i>nitidus</i> , <i>Alburnus</i>	123
<i>Salmo</i>	60	<i>Balistes</i>	493
<i>Syphurus</i>	230	<i>Brachydeuterus</i>	333
<i>Tigoma</i>	120	<i>Leuciscus</i>	117
<i>nigricans</i> , <i>Bathylaco</i>	104	<i>Pomoxis</i>	304
<i>Camarina</i>	338, 339	<i>Salmo</i>	60
<i>Catostomus</i>	107	<i>Tetragonopterus</i>	96
<i>Centropristes</i>	317	<i>nivea</i> , <i>Eregala</i>	120

List of scientific names—Continued

	Page		Page
<i>niveatus</i> , <i>Epinephelus</i>	311	<i>notatus</i> , <i>Pomotis</i>	302
<i>Pomacentrus</i>	416	<i>Porichthys</i>	487
<i>Serranus</i>	310, 311	<i>Semotilus</i>	177
<i>Stegastes</i>	410	<i>Zygonectes</i>	177
<i>nivelventris</i> , <i>Amiurus</i>	153	<i>notenigrooides</i> , <i>Notropis</i>	125
<i>niveus</i> , <i>Hybopsis</i>	129	<i>Notemigonus</i>	114
<i>Nivicola</i>	201	<i>auratus</i>	114, 115
<i>artesiae</i>	202	<i>bosci</i>	115
<i>borealis</i>	201	<i>crysoleucas</i>	115
<i>cragini</i>	202	<i>gardonae</i>	115
<i>lepidogenys</i>	202	<i>ischanus</i>	115
<i>obeyense</i>	202	<i>nothochir</i> , <i>Ophichthys</i>	87
<i>pagei</i>	202	<i>Quassiremus</i>	87
<i>punctulata</i>	202	<i>Nothonotus</i>	282, 288, 290
<i>tippecanoe</i>	202	<i>camurus</i>	290
<i>virgata</i>	202	<i>cinerous</i>	290
<i>nivipes</i> , <i>Emblemaria</i>	404	<i>inscriptus</i>	289
<i>nobiliana</i> , <i>Torpedo</i>	27	<i>jordani</i>	290
<i>nobilior</i> , <i>Esox</i>	172	<i>maculatus</i>	290
<i>nobilis</i> , <i>Atractoscion</i>	356	<i>rufilineatus</i>	290
<i>Conodon</i>	333	<i>tesselatus</i>	289
<i>Esox</i>	172	<i>thalassinus</i>	289
<i>Gymnibus</i>	185	<i>vulneratus</i>	289
<i>Heterandria</i>	185	(<i>Nothonotus</i>) <i>jordanii</i> , <i>Etheostoma</i>	290
<i>Johnius</i>	356	<i>nothus</i> , <i>Cynoscion</i>	355
<i>Perca</i>	333	<i>Otolithus</i>	355
<i>Polyurixia</i>	234	<i>Notidani</i>	11
<i>Nocomis</i>	137	<i>Notidanus</i>	11
<i>bellicus</i>	137	<i>Notistium</i>	263
<i>hyostomus</i>	139	<i>gladius</i>	263
<i>kentuckiensis</i>	137	<i>Notogrammus</i>	469
<i>micropogon</i>	137	<i>rothrocki</i>	469
<i>milleri</i>	136	<i>Notolepidomyzon</i>	104
<i>nebrascanis</i>	137	<i>clarkii</i>	104
<i>rubrifrons</i>	138	<i>generosus</i>	104
<i>nocomis</i> , <i>Ifyopsis</i>	135	<i>plebeius</i>	104
<i>Notropis</i>	135	<i>santae-anne</i>	104
<i>nocturna</i> , <i>Colletta</i>	168	<i>Notorhynchus borealis</i>	11
<i>Echidna</i>	94	<i>Notorynchus</i>	11
<i>Rabida</i>	136	<i>maculatus</i>	11
<i>nocturnum</i> , <i>Myctophum</i>	168	<i>platycephalus</i>	11
<i>nocturnus</i> , <i>Noturus</i>	156	<i>Notoscopelus</i>	166
<i>Paeophis</i>	94	<i>castaneus</i>	167
<i>nodulosus</i> , <i>Xenodermichthys</i>	53	<i>margaritiferus</i>	166
<i>Nomeidae</i>	324	<i>querulus</i>	166
<i>Nomeus</i>	268	<i>Notosema</i>	223
<i>gronovii</i>	268	<i>dicta</i>	225
<i>maculosus</i>	268	<i>dictuum</i>	225
<i>oxyurus</i>	268	<i>notospilotus</i> , <i>Artedius</i>	376
<i>normalis</i> , <i>Bassozetus</i>	481	<i>Parastrolytes</i>	376
<i>norvegica</i> , <i>Perca</i>	363	<i>notospilus</i> , <i>Jullidio</i>	428
<i>norvegicus</i> , <i>Lepidolepus</i>	205	<i>Pseudolulis</i>	428
<i>norwegianus</i> , <i>Squalus</i>	22	<i>Notropis</i>	428
<i>notabilis</i> , <i>Argyreus</i>	141	<i>torfoniensis</i>	131
<i>Notacanthidæ</i>	75	<i>lirus</i>	126
<i>Notacanthus</i>	75	<i>nasus</i>	126
<i>analis</i>	76	<i>stilbius</i>	126
<i>challengeri</i>	76	<i>Notropis</i>	123
<i>chemnitzi</i>	75	<i>eneoculus</i>	143
<i>nasus</i>	75	<i>alabamae</i>	126
<i>phaganaeus</i>	76	<i>albicolor</i>	136
<i>rissorinus</i>	76	<i>ambassis</i>	124
<i>rostratus</i>	76	<i>amoenus</i>	123
<i>spinosus</i>	76	<i>amogenus</i>	136
<i>Notarius</i>	149	<i>arcansanus</i>	124
<i>notata</i> , <i>Atherina</i>	247	<i>argo</i>	124
<i>Belone</i>	195	<i>atherinoides</i>	123
<i>Cyprinella</i>	130	<i>aztecus</i>	143
<i>Dorosoma</i>	46	<i>boops</i>	134
<i>Erigone</i>	130	<i>braytoni</i>	135
<i>Lepomis</i>	298	<i>brimleyi</i>	127
<i>Megalops</i>	44	<i>buchananii</i>	134
<i>Menidia</i>	247	<i>caliente</i>	134
<i>Perca</i>	261	<i>cayuga</i>	133
<i>Maihrea</i>	507	<i>atrocaudalis</i>	133
<i>Strongylura</i>	105	<i>chalibucus abbotti</i>	127
<i>notatum</i> , <i>Hæmulon</i>	329	<i>chamberlaini</i>	132
<i>Pristipoma</i>	332	<i>chihuhua</i>	132
<i>notatus</i> , <i>Balistes</i>	403	<i>cornutus isolepis</i>	128
<i>Hyborhynchus</i>	145	<i>cummingsi</i>	133
<i>Labrax</i>	307	<i>dilectus</i>	125
<i>Minilius</i>	145	<i>fumeus</i>	125
<i>Passer lineatus transversis</i>	230	<i>garmani</i>	131
<i>Plimelodus</i>	152	<i>germanus</i>	136
		<i>gilberti</i>	134
		<i>heterodon richardsoni</i>	136

List of scientific names—Continued

	Page		Page
<i>Notropis heterolepis</i>	134	<i>nox</i> , <i>Auchenopterus</i>	457
<i>horatii</i>	134	<i>Crennobates</i>	457
<i>jordani</i>	136	<i>novesi</i> , <i>Scarus</i>	434
<i>kapawha</i>	134	<i>nubila</i> , <i>Apocope</i>	140
<i>keimi</i>	133	<i>Dlonda</i>	144
<i>kendalli</i>	135	<i>nubilus</i> , <i>Alburnops</i>	144
<i>lerina</i>	143	<i>Argyreus</i>	140
<i>leuciodus</i>	124	<i>Lumpenus</i>	471
<i>louisiane</i>	124	<i>nuchalis</i> , <i>Hybognathus</i>	143
<i>lythurus</i>	126	<i>Scarus</i>	434
<i>macdonaldi</i>	127	<i>nuchifilus</i> , <i>Bleennius</i>	461
<i>macrolepidotus</i>	125	<i>nichtipinnis</i> , <i>Clinus</i>	459
<i>mearnsi</i>	122	<i>Labrisomus</i>	450
<i>megalops albeolus</i>	129	<i>nudus</i> , <i>Cyclopterus</i>	489
<i>metallicus</i>	124	<i>Gymnachirus</i>	220
<i>micropteryx</i>	124	<i>nugator</i> , <i>Bryostemma</i>	466
<i>muskoka</i>	134	<i>nummifer</i> , <i>Carpioidea</i>	103
<i>noconis</i>	135	<i>Salmo</i>	61
<i>notatus</i>	124	<i>nuttalli</i> , <i>Exocetus</i>	201
<i>notemigonoides</i>	125	<i>nuttingi</i> , <i>Chunax</i>	506
<i>nux</i>	135	<i>Scorpea</i>	371
<i>orca</i>	120	<i>nuttingi</i> , <i>Antennarius</i>	506
<i>ortenburgeri</i>	135	<i>nux</i> , <i>Hybopsis</i>	165
<i>ozarkanus</i>	135	<i>Notropis</i>	135
<i>phenacobius</i>	134	<i>nycterinus</i> , <i>Melanophaës</i>	233
<i>photogenys</i>	123	<i>Plectromus</i>	233
<i>pilsbryi</i>	124	<i>Nyctimaster</i>	167
<i>reticulatus</i>	133	<i>alatus</i>	107
<i>robustus</i>	132	<i>townsendi</i>	167
<i>roseipinnis</i>	126	<i>Nyctophus</i>	169
<i>rubrifrons</i>	124	<i>rafinesquei</i>	188, 189
<i>sabinae</i>	135	<i>oaxacae</i> , <i>Fundulus</i>	174
<i>santamarie</i>	131	<i>Profoundulus</i>	174
<i>scepticus</i>	123	<i>Rhamdia</i>	157
<i>scopiferus</i>	132	<i>obesa</i> , <i>Algensea</i>	122
<i>socius</i>	124	<i>obesus</i> , <i>Amiurus</i>	154
<i>stilbius</i>	125	<i>Carcharitus</i>	14
<i>stonei</i>	127	<i>Echinorhinus</i>	22
<i>telescopus</i>	124	<i>Enneacanthus</i>	304
<i>arcansanus</i>	124	<i>Leuciscus</i>	128
<i>topeka</i>	143	<i>Pomotis</i>	304
<i>umbratilis fasciolaris</i>	126	<i>Siphateles</i>	122
<i>umbrifer</i>	131	<i>Triodonodon</i>	14
<i>universitatis</i>	124	<i>obeyense</i> , <i>Etheostoma</i>	292
<i>welaka</i>	124	<i>Nivicola</i>	292
<i>notropoides</i> , <i>Creagrutus</i>	98	<i>obliquatus</i> , <i>Cynoscion</i>	355
<i>nottii</i> , <i>Zygoneutes</i>	178	<i>Dinbasis</i>	330
<i>Noturus</i>	155	<i>Otolithus</i>	355
<i>elusochir</i>	156	<i>obliquus</i> , <i>Mugil</i>	349
<i>eleutherus</i>	156	<i>obliteratus</i> , <i>Alutarius</i>	498
<i>exilis</i>	156	<i>oblonga</i> , <i>Platessa</i>	224
<i>flavus</i>	155	<i>oblongor</i> , <i>Plimalepterus</i>	340
<i>funebris</i>	156	<i>oblongus</i> , <i>Astatheros</i>	421
<i>furious</i>	155, 156	<i>Cyprinus</i>	109
<i>gilberti</i>	156	<i>Erimyzon</i>	109
<i>leptacanthus</i>	155	<i>Gobius</i>	448
<i>luteus</i>	155	<i>Heros</i>	421
<i>marginatus</i>	156	<i>Paralichthys</i>	224
<i>mirurus</i>	156	<i>Pleuronectes</i>	224
<i>nocturnus</i>	150	<i>obularius</i> , <i>Gasterosteus</i>	237
<i>occidentalis</i>	155	<i>obscura</i> , <i>Liosoma</i>	228
<i>platycephalus</i>	155	<i>Sternoptix</i>	73
<i>sialis</i>	155	<i>Sternoptyx</i>	73
<i>nova-boracensis</i> , <i>Vomer</i>	275	<i>obscuratus</i> , <i>Gymnothorax</i>	93
<i>Novacula</i>	430	<i>Pomacentrus</i>	414
<i>kallosoma</i>	430	<i>obscurus</i> , <i>Carcharlas</i>	16
<i>novacula</i> , <i>Coryphena</i>	430	<i>Centrarchus</i>	208
<i>Xyrichthys</i>	430	<i>Pleuronectes</i>	228
<i>Novaculichthys</i>	430	<i>Pomotis</i>	301
<i>infrimis</i>	430	<i>Squalus</i>	16
<i>martinicensis</i>	430	<i>obtusa</i> , <i>Ophisooma</i>	79
<i>rosiques</i>	430	<i>Raiia</i>	26
<i>ventralis</i>	430	<i>obtusirostris</i> , <i>Exocetus</i>	200
<i>nova-angliae</i> , <i>Coregonus</i>	65	<i>obtusum</i> , <i>Hemulon</i>	332
<i>noveboracensis</i> , <i>Anguilla</i>	77	<i>obtusum</i> , <i>Rhinichthys</i>	140
<i>noveterre</i> , <i>Anguilla</i>	77	<i>Scarus</i>	434
<i>noveboracensis</i> , <i>Exocetus</i>	201	<i>Squamus</i>	16
<i>Gasterosteus</i>	237	<i>obvelatus</i> , <i>Prionodon</i>	16
<i>Naufrates</i>	278	<i>Occa</i>	394
<i>novenasciatus</i> , <i>Lutjanus</i>	325	<i>verrucosus</i>	394
<i>Lutjanus</i>	325	<i>occa</i> , <i>Celorophryncus</i>	207
<i>novenlineatus</i> , <i>Chasmodes</i>	402	<i>Macrurus</i>	207
<i>Pholis</i>	402	<i>Pristis</i>	23
<i>novenmaculatus</i> , <i>Diodon</i>	501	<i>Occella</i>	394
<i>novenradiata</i> , <i>Agonus</i>	141	<i>dodecadron</i>	394

List of scientific names—Continued

	Page		Page
<i>occidentalis</i> , <i>Amia</i> .	38	<i>oculocirris</i> , <i>Enblemaria</i> .	464
<i>Catostomus</i> .	106	<i>oculofasciatus</i> , <i>Blepsias</i> .	382
<i>Chultumra</i> .	204	<i>Nautichthys</i> .	382
<i>Epigonus</i> .	296	<i>oculo-radiato</i> , <i>Turdus</i> .	426
<i>Gasterosteus</i> .	238, 239	<i>Ocyanthias</i> .	320
<i>Heterandria</i> .	180	<i>martinicensis</i> .	320
<i>Hypopomus</i> .	101	<i>Ocyurus</i> .	327
<i>Lepomis</i> .	301	<i>chrysurus</i> .	327
<i>Letucosomus</i> .	413	<i>litjanoides</i> .	327
<i>Malacocephalus</i> .	204	<i>rijgersmelei</i> .	327
<i>Monacanthus</i> .	494	<i>ocurus</i> , <i>Centropristes</i> .	318
<i>Myripristis</i> .	235	<i>Pinnelepterus</i> .	340
<i>Myripristis</i> .	235	<i>Sectator</i> .	340
<i>Nebris</i> .	354	<i>Serranus</i> .	318
<i>Noturus</i> .	155	<i>Odontaspidae</i> .	19
<i>Poeciliopsis</i> .	189	<i>Odontaspis</i> .	19
<i>Roeboides</i> .	98	<i>litoralis</i> .	19
<i>Tetranarce</i> .	27	<i>Odontognathus</i> .	45
<i>Torpedo</i> .	27	<i>macronota</i> .	45
<i>Xenotis</i> .	301	<i>panamensis</i> .	45
<i>occipitalis</i> , <i>Scorpaena</i> .	371	(<i>Odontognathus</i>) <i>panamensis</i> , <i>Pristigaster</i> .	45
<i>oceania</i> , <i>Anguilla</i> .	79	<i>Odontopyxis</i> .	306
<i>oceanius</i> , <i>Blennius</i> .	461	<i>frenatus</i> .	395
<i>Gobionellus</i> .	441	<i>leptorhynchus</i> .	395
<i>Gobius</i> .	441	<i>trispinosus</i> .	396
<i>Rhinoscopelus</i> .	169	<i>Odontoscion</i> .	345
<i>Oceanomyzon</i> .	8	<i>archidium</i> .	345
<i>wilsoni</i> .	8	<i>dentex</i> .	334
<i>oceanoops</i> , <i>Elacatinus</i> .	442	<i>xanthops</i> .	334
<i>ocella</i> , <i>Fretmichthys</i> .	481	ørstedi, <i>Argyreiosus</i> .	275
<i>Rhinchthys</i> .	139	<i>Selene</i> .	275
<i>Venefica</i> .	82	<i>Tetragonopterus</i> .	98
<i>ocellaris</i> , <i>Fundulus</i> .	176	ogac, <i>Gadus</i> .	210
<i>Gobius</i> .	442	Ogcocephalidae.	506
<i>Platessa</i> .	224	Ogcocephalus.	506
<i>ocellata</i> , <i>Bolimannia</i> .	443	<i>nasutus</i> .	507
<i>Chenopsis</i> .	465	<i>radiatus</i> .	507
<i>Perca</i> .	348	<i>vespertilio</i> .	607
<i>Raiia</i> .	25	Ogilbiidae.	479
<i>Scianops</i> .	348	<i>cayorum</i> .	480
<i>Zenopsis</i> .	232	<i>ventrals</i> .	480
<i>ocellatus</i> , <i>Acanthocottus</i> .	387	<i>verrillii</i> .	480
<i>Anarrhichthys</i> .	472	ogilbya, <i>Megalops</i> .	44
<i>Antennarius</i> .	505	oglinum, <i>Ophisthonia</i> .	44
<i>Chenopsis</i> .	465	ohiensis, <i>Accipenser</i> .	35
<i>Chetodon</i> .	300	<i>Alosa</i> .	42
<i>Clinus</i> .	459	<i>Cithla</i> .	298
<i>Gymnothorax</i> .	91, 93	<i>Esox</i> .	172
<i>Hippoglossus</i> .	218	<i>masquinony</i> .	172
<i>Lophius histrio</i> .	505	ojanco, <i>Mesopriion</i> .	327
<i>Malacocetus</i> .	450	okeechobeensis, <i>Amieus lacustris</i> .	152
<i>Murennophis</i> .	87	Haustor.	152
<i>Neobrythites</i> .	481	<i>Ictalurus</i> .	152
<i>nigromarginatus</i> , <i>Gymnothorax</i> .	93	Okkelbergia.	10
<i>Opichtthus</i> .	87	<i>lamotteti</i> .	10
<i>Platophrys</i> .	217	olfax, <i>Mycteroperca</i> .	314
<i>Pleuronichthys</i> .	225	<i>Serranus</i> .	314
<i>Rhabodus</i> .	217	olfersi, <i>Argyropelecus</i> .	74
<i>saxicola</i> , <i>Gymnothorax</i> .	93	<i>Cephalopterus</i> .	32
<i>Zeus</i> .	232	<i>Nobula</i> .	32
<i>ocellata</i> , <i>Amia</i> .	38	<i>Sternoptyx</i> .	74
<i>ocellifer</i> , <i>Clinus</i> .	458	olidus, <i>Hypomesus</i> .	68
<i>ocotlane</i> , <i>Chirostoma</i> .	251	<i>Salmo</i> (<i>Osmurus</i>).	68
<i>octodecemspinosis</i> , <i>Acanthocottus</i> .	386	oligaspis, <i>Alburnus</i> .	125
<i>octodecem-spinosus</i> , <i>Cottus</i> .	386	Oligocephalus.	290
<i>octofasciatus</i> , <i>Astatheros</i> .	421	<i>auteenau bei</i> .	291
<i>Heros</i> .	421	<i>ceruleus</i> .	291
<i>octofills</i> , <i>Trichidion</i> .	256	<i>fricksi</i> .	291
<i>Octogrammus</i> .	373	<i>grahami</i> .	291
<i>pallasi</i> .	374	<i>hildebrandti</i> .	291
<i>octogrammus</i> , <i>Hexagrammos</i> .	374	<i>humeralis</i> .	293
<i>Labrax</i> .	374	<i>lowei</i> .	291
<i>octonemus</i> , <i>Polydenus</i> .	266	<i>jessiae</i> .	291
<i>oculata</i> , <i>Squatina</i> .	22	<i>leoniensis</i> .	291
<i>oculatus</i> , <i>Brytius</i> .	300	<i>lepidus</i> .	291
<i>Etelis</i> .	328	<i>luteovinctus</i> .	291
<i>Icelinus</i> .	376	<i>pulchellus</i> .	291
<i>Latebrus</i> .	297	Oligocottus.	390
<i>Lepisosteus</i> (<i>Cylindrostoeus</i>).	37	<i>acuticeps</i> .	391
<i>Maurolicus</i> .	73	<i>analis</i> .	390
<i>Myrichthys</i> .	85	<i>borealis</i> .	390
<i>Pisodonophis</i> .	85	<i>embryum</i> .	391
<i>Serranus</i> .	328	<i>globiceps</i> .	391
<i>oculeatus</i> , <i>Orthagoriscus</i> .	503	<i>masculosus</i> .	390
<i>oculeum</i> , <i>Myctophum</i> .	169	<i>synderi</i> .	391

List of scientific names—Continued

	Page		Page
oligodon, <i>Osmerus</i>	68	Ophichthus <i>chamensis</i>	88
<i>Polyniemus</i>	256	<i>evionthas</i>	87
oligolepis, <i>Brycon</i>	99	<i>gomesii</i>	88
oligopeplus, <i>Acipenser</i> (<i>Antaceus</i>).....	34	<i>guttifer</i>	87
Oligoplites.....	277	<i>magniculus</i>	88
<i>altus</i>	278	<i>ocellatus</i>	87
<i>inornatus</i>	277	<i>ophis</i>	87
<i>mundus</i>	278	<i>parilis</i>	88
<i>palometta</i>	278	<i>retropinnis</i>	87
<i>refulgens</i>	278	<i>rugifer</i>	88
<i>sallens</i>	277	<i>triserialis</i>	88
<i>saurus</i>	277	<i>zophochir</i>	88
Oligopodes.....	266	Ophichthyidae.....	84
Oligopodus.....	266	Ophichthys.....	87
Olipogon.....	266	biserials	88
oligosticta, <i>Gambusia</i>	186	(<i>Cryptopterus</i>) <i>frontalis</i>	85
olishostoma, <i>Gerres</i>	342	<i>guttifer</i>	87
olishostomus, <i>Dipterus</i>	342	<i>miurus</i>	89
olivacea, <i>Poecilia</i>	177	<i>nothochir</i>	87
olivateus, <i>Leucus</i>	122	<i>pauciporus</i>	88
<i>Siphateles</i>	122	<i>pisavarius</i>	85
olivarisi, <i>Opiadelus</i>	155	<i>retropinnis</i>	87
<i>Silurus</i>	155	<i>schneideri</i>	88
olmstedi, <i>Boleosoma</i>	287	(<i>Sphagebranchus</i>) <i>anguiformis</i>	84
<i>brevipinnis</i> , <i>Boleosoma</i>	287	<i>xysturus</i>	85
<i>Etheostoma</i>	287	<i>zopochir</i>	88
olomina, <i>Brachyrhaphis</i>	187	Ophichthus <i>evionthas</i>	87
<i>Prispichthys</i>	187	ophidianus, <i>Gempylus</i>	262
olrikii, <i>Aspidophoroides</i>	397	Ophididae.....	484
<i>Ulcina</i>	397	Ophidion.....	485
omiscomaycus, <i>Percopsis</i>	215	<i>beani</i>	485
<i>Salmo</i>	215	<i>grellsi</i>	485
ommata, <i>Discopyge</i>	27	<i>holbrookii</i>	485
<i>Heterandria</i>	175	<i>imberbe</i>	487
<i>Leptolucania</i>	175	Ophidium.....	485
<i>Opistognathus</i>	453	<i>atropurpureum</i>	485
ommastistius <i>terre-nova</i> , <i>Triglops</i>	380	<i>brevibarbe</i>	484
ommatum, <i>Opistognathus</i>	453	<i>holbrookii</i>	485
ommatus, <i>Paralichthys</i>	224	<i>josephi</i>	485
omochelys.....	86	<i>marginatum</i>	485
<i>cruentifer</i>	86	<i>mucronatum</i>	487
omocyaneus, <i>Eleotris</i>	438	<i>parrisi</i>	478
omostigma, <i>Genypterus</i>	485	<i>stigma</i>	477
<i>Otopodium</i>	485	<i>taylori</i>	485
Omosudidae.....	171	<i>unenak</i>	477
Omosuds.....	171	<i>viridis</i>	477
<i>lowii</i>	171	Ophioblennius.....	464
Oncococephalus.....	570	<i>steindachneri</i>	464
Oncocottus.....	388	<i>webbi</i>	464
<i>gilberti</i>	389	Ophiodon.....	374
<i>hexacornis</i>	388	<i>elongatus</i>	374
<i>gilberti</i>	389	Ophiodontidae.....	374
<i>quadricornis</i>	388	Ophiognathus.....	95
Oncorhynchus.....	54	<i>ampullaceus</i>	95
<i>gorbuscha</i>	54	ophioneus, <i>Sphagebranchus</i>	84
<i>kamloops</i>	58	Ophioscion.....	347
<i>kennerlyi</i>	55	<i>adustus</i>	347
<i>keta</i>	54	<i>imiceps</i>	348
<i>klisutch</i>	55	<i>punctatissimus</i>	348
<i>nerka</i>	55	<i>scirurus</i>	348
<i>tschawytscha</i>	55	<i>stimulus</i>	348
oneida, <i>Catostomus</i>	110	<i>strabo</i>	347
Oneirodes.....	609	<i>typicus</i>	347
<i>eschrichtii</i>	509	ophioscion, <i>Corvina</i>	347
onitis, <i>Labrus</i>	424	ophis, <i>Muraena</i>	87
Onos.....	208	<i>Ophichthus</i>	87
<i>rufus</i>	208	Ophisoma.....	79
ontariensis, <i>Leucichthys</i>	62	<i>acuta</i>	79
<i>Thymallus</i>	66	<i>balearicum</i>	80
<i>Triglopsis</i>	389	<i>macrurum</i>	80
Onus.....	214	<i>nitens</i>	80
<i>rialli</i>	214	<i>obtusa</i>	79
onychus, <i>Cottus</i>	386	<i>prorigerum</i>	80
Oonidus.....	499	Ophisomus.....	467
opan, <i>Zeus</i>	217	Ophisternon.....	76
opalescens, <i>Lythrum</i>	331	Ophisuraphis.....	84
opalina, <i>Julis</i>	426	Ophisurus.....	85
opercularis, <i>Acantholiparis</i>	401	<i>californiensis</i>	88
<i>Myxodagnus</i>	455	<i>chrysops</i>	88
<i>Polydactylus</i>	256	<i>crocodilinus</i>	89
<i>Stolephorus</i>	51	<i>gomesii</i>	88
<i>Trichidion</i>	256	<i>guttatus</i>	87
Ophichthitus <i>regius</i>	87	<i>interstitiatus</i>	88
Ophichthitus.....	87	<i>latimaculatus</i>	88
<i>biserials</i>	88		

List of scientific names—Continued

	Page		Page
<i>Ophisurus longus</i>	85	<i>oregonia</i> , <i>Atherinops</i>	252
<i>ophis</i>	87	<i>Oregonichthys</i>	138
<i>parilis</i>	88	<i>crameri</i>	138
<i>rostellatus</i>	88	<i>cumingii</i>	138
<i>sugillatus</i>	88	<i>oregonius</i> , <i>Coregonus</i>	66
<i>xysturus</i>	85	<i>Iridion</i>	66
<i>ophryas</i> , <i>Paralichthys</i>	224	<i>Oreosoma</i>	232
<i>Prionotus</i>	407	<i>atlanticum</i>	232
<i>ophthalmicus</i> , <i>Lepomis</i>	299	<i>conferrum</i>	232
<i>Opistharthri</i>	11	<i>orientalis</i> , <i>Anarhichas</i>	472
<i>Opisthostius</i>	339	<i>Pelamys</i>	259
<i>Opistognathidae</i>	452	<i>Salmo</i>	55
<i>Opistognathus</i>	452	<i>ornata</i> , <i>Amia</i>	88
<i>cuvieri</i>	453	<i>Aphoristia</i>	231
<i>lonchurum</i>	453	<i>Codoma</i>	143
<i>lonchurus</i>	453	<i>Embiotoca</i>	411
<i>macrognathum</i>	452	<i>Hydrargyra</i>	176
<i>macrops</i>	453	<i>Limia</i>	191
<i>maxillosus</i>	453	<i>Raja</i>	28
<i>megastoma</i>	452	<i>ornatum</i> , <i>Campsochima</i>	146
<i>micrognathus</i>	453	<i>ornatus</i> , <i>Achirus</i>	231
<i>ommatum</i>	453	<i>Cochlognathus</i>	146
<i>punctatum</i>	453	<i>Esox</i>	173
<i>punctatus</i>	453	<i>Gillellus</i>	455
<i>rhomaleus</i>	453	<i>Gunnelius</i>	468
<i>scaphiurus</i>	452	<i>Labrus</i>	429
<i>Opisthourema</i>	44	<i>Pholis</i>	408
<i>libertate</i>	44	<i>Tetradon</i>	600
<i>oglinum</i>	44	<i>Ornithichthys</i>	407
<i>Opisthoterpterus</i>	45	<i>orqueta</i> , <i>Chloroscomorus</i>	276
<i>dovii</i>	45	<i>orsini</i> , <i>Ozodus</i>	503
<i>macrops</i>	45	<i>ortenburgeri</i> , <i>Hybopsis</i>	135
<i>opisthopthalmus</i> , <i>Ariosoma</i>	80	<i>Notropis</i>	135
<i>opisthotremus</i> , <i>Careproctus</i>	403	<i>Orthocolus</i>	66
<i>opistophthalmus</i> , <i>Conger</i>	80	<i>Orthodon</i>	112
<i>Opiaedelus</i>	155	<i>microlepidotus</i>	112
<i>olivaris</i>	155	<i>Orthogorisicus</i>	503
<i>Opiopoma</i>	374	<i>orthogramma</i> , <i>Citula</i>	274
<i>pantherina</i>	374	<i>Orthoprias</i>	391
<i>oppositus</i> , <i>Monacanthus</i>	495	<i>triacis</i>	392
<i>Opsanus</i>	487	<i>orthonotus</i> , <i>Ditrema</i>	411
<i>cerapalus</i>	487	<i>Orthopristis</i>	335
<i>hildebrandi</i>	487	<i>chalcetus</i>	335
<i>pardus</i>	487	<i>chrysopterus</i>	335
<i>tud</i>	487	<i>duplex</i>	335
<i>Opsopoea</i>	115	<i>poeyi</i>	335
<i>bollmani</i>	115	<i>reddingi</i>	335
<i>Opsopoeodus</i>	115	<i>Orthopsetta</i>	218
<i>bollmani</i>	115	<i>Orthostethus</i>	331
<i>borealis</i>	115	<i>maculicauda</i>	331
<i>emiliae</i>	115	<i>Orthragoriscus alexandrinus</i>	503
<i>megalops</i>	115	<i>analis</i>	503
<i>oculus</i>	115	<i>battaræ</i>	503
<i>oquassa</i> , <i>Salmo</i>	61	<i>blochii</i>	503
<i>Salvelinus</i>	61	<i>elegans</i>	503
<i>crbe</i> , <i>L.</i> <i>Diodon</i>	502	<i>ghini</i>	503
<i>Orbis</i>	498	<i>lunaris</i>	503
<i>orbignianus</i> , <i>Exocetus</i>	200	<i>oculeatus</i>	503
<i>orbignyanus</i> , <i>Conger</i>	79	<i>redi</i>	503
<i>Orbis levis variegatus</i>	499	<i>ratzii</i>	503
<i>orbis</i> , <i>Cyclopterus</i>	398	<i>roulei</i>	503
<i>Eumicrotremus</i>	398	<i>solaris</i>	503
<i>orbitalis</i> , <i>Macrourus</i>	203	<i>Orthragus</i>	503
<i>Macrurus</i>	206	<i>Oryctenus</i>	260
<i>orbitarius</i> , <i>Pagellus</i>	336	<i>subulatus</i>	260
<i>orca</i> , <i>Notropis</i>	126	<i>Osbeck</i> , <i>Trachinus</i>	310
<i>Orcella</i>	126	<i>Osbeckia</i>	495
<i>Orcella</i>	126	<i>scripta</i>	495
<i>nasas</i>	126	<i>osborni</i> , <i>Liparis</i>	400
<i>orca</i>	126	<i>Typhlichthys</i>	195
<i>orcini</i> , <i>Brama</i>	265	<i>oscitans</i> , <i>Scienea</i>	347
<i>Orcula</i>	126	<i>oscitans</i> , <i>Zestis</i>	347
<i>creuti</i> , <i>Phoxinus</i>	120	<i>oscula</i> , <i>Apocope</i>	141
<i>Tigoma</i>	120	<i>Netuma</i>	149
<i>Orcynus</i>	260	<i>Scienea</i>	352
<i>subulatus</i>	280	<i>osculatilis</i> , <i>Labrax</i>	307
<i>ordinatus</i> , <i>Chirus</i>	374	<i>osculum</i> , <i>Holocentrum</i>	236
<i>Epinephelus</i>	311	<i>osculus</i> , <i>Argyreus</i>	141
<i>ores</i> , <i>Chromosomus</i>	113	<i>Arius</i>	149
<i>oregonensis</i> , <i>Cyprinus</i> (<i>Leuciscus</i>)	114	<i>Holocentrus</i>	236
<i>Ptychocheilus</i>	114	<i>Opsopoeodus</i>	115
<i>Rutilus</i>	122	<i>Osmeridae</i>	66
<i>Siphateles</i>	122	<i>osmeriformis</i> , <i>Coregonus</i>	62
		<i>Leucichthys</i>	62

List of scientific names—Continued

	Page		Page
<i>osmerinus</i> , <i>Hybognathus</i> .	143	Otrynter.	336
<i>Osmerus</i> .	67	<i>caprinus</i> .	336
<i>abbotti</i> .	67	Otrynter, <i>Caranx</i> .	274
<i>albatrossis</i> .	67	otsago, <i>Salmo</i> .	64
<i>albidus</i> .	163	ouachite, <i>Alvardius</i> .	284
<i>attenuatus</i> .	67	<i>Etheostoma</i> (<i>Hadropterus</i>).	284
<i>dentex</i> .	67	ouananiche, <i>Salmo</i> .	56
<i>elongatus</i> .	67	<i>Salmo salar</i> .	56
<i>lemniscatus</i> .	163	ounialibi, <i>Serranus</i> .	309
<i>microdon</i> .	66	ovale, <i>Acutomentum</i> .	366
<i>mordax</i> .	67	<i>Syacium</i> .	218
<i>abbotti</i> .	67	ovalis, <i>Hemibombus</i> .	218
<i>spectrum</i> .	67	<i>Sebastodes</i> .	366
<i>oligodon</i> .	68	ovicephalus, <i>Sparus</i> .	338
<i>sargentii</i> .	67	oviformis, <i>Chatodon</i> .	358
<i>spectrum</i> .	67	ovigerum, <i>Bathyphasma</i> .	401
<i>thaleichthys</i> .	67	ovinus, <i>Esox</i> .	181
<i>viridescens</i> .	67	ovis, <i>Sargus</i> .	338
<i>Osphylax</i> .	243	Ovoides.	490, 500
<i>pellucidus</i> .	244	Ovum.	499
<i>osseus</i> , <i>Esox</i> .	36	Oxybeles.	486
<i>Lepisosteus</i> .	36	oxybrachium, <i>Sparisoma</i> .	432
<i>Ostariophys</i> .	95	oxybrachius, <i>Scarus</i> .	432
<i>ostentum</i> , <i>Careproctus</i> .	402	Oxychætodon.	359
<i>osteochir</i> , <i>Echeneis</i> .	450	Oycottus.	391
<i>Rhombocheirus</i> .	450	<i>acuticeps</i> .	391
<i>osteosticta</i> , <i>Trygon</i> .	29	Oxygenium.	113
<i>Ostichthys</i> .	234	<i>pulverulentum</i> .	113
<i>trachypomus</i> .	234	Oxyjulis.	428
<i>Ostocrinhus</i> .	295	<i>californicus</i> .	428
<i>Ostracidae</i> .	496	Oxylabrax.	305
<i>Ostracion</i> .	496	Oxylebiidae.	374
<i>bicaudalis</i> .	497	Oxylebius.	374
<i>boops</i> .	503	<i>pictus</i> .	374
<i>clippertonense</i> .	496	Oxyloricaria.	162
<i>concatenatus</i> .	496	<i>citrenensis</i> .	162
<i>gronovii</i> .	497	<i>dariensis</i> .	162
<i>guineensis</i> .	497	<i>panamensis</i> .	162
<i>lentiginosum</i> .	496	Oxyodontichthys.	87
<i>lister</i> .	497	<i>brachyurus</i> .	88
<i>punctatum</i> .	496	<i>limbatus</i> .	88
<i>punctatus</i> .	496	<i>macrurus</i> .	88
<i>quadricornis</i> .	497	<i>oxyptera</i> , <i>Corvina</i> .	320
<i>sexcornutus</i> .	497	Oxyrhina.	19
<i>trigonus</i> .	497	<i>dækayi</i> .	19
<i>triqueter</i> .	496	<i>gomphodon</i> .	19
<i>undulatus</i> .	497	<i>oxyrhynchus</i> , <i>Acipenser</i> .	34
<i>yalei</i> .	497	<i>Carcharias</i> .	15, 16
<i>Ostracion</i> .	496	<i>Tetradon</i> .	500
<i>expansum</i> .	497	<i>oxyrinchus</i> , <i>Isurus</i> .	19
<i>tricornis</i> .	497	Oxyurus.	79
<i>Ostracoderini</i> .	496	<i>oxyurus</i> , <i>Lepisosteus</i> .	37
<i>otarius</i> , <i>Lepidosteus</i> .	37	<i>Nameus</i> .	268
<i>Othonops</i> .	447	Oxyzygocetes.	179
<i>eos</i> .	46	<i>baibae</i> .	179
<i>othonoporus</i> , <i>Perkinsia</i> .	46	<i>dovii</i> .	179
<i>othonopterus</i> , <i>Cynoscion</i> .	355	Ozadura.	503
<i>Otolithidæ</i> .	353	<i>ozarcanus</i> , <i>Hybopsis</i> .	135
<i>Otolithus albus</i> .	356	<i>Notropis</i> .	135
<i>californiensis</i> .	355	Ozdura orsinii.	503
<i>carolinensis</i> .	355		
<i>cayennensis</i> .	355	pachycephalus, <i>Fundulus</i> .	174
<i>drummondi</i> .	355	<i>Lagocephalus</i> .	497
<i>jamaicensis</i> .	355	<i>Tetradon</i> .	497
<i>leiorhynchus</i> .	355	pachygaster, <i>Cheilichthys</i> .	499
<i>magdalene</i> .	356	<i>Tetradon</i> (<i>Cheilichthys</i>).	499
<i>micolepidotus</i> .	355	Pachygnatha.	492
<i>microps</i> .	356	Pachygnathus.	492
<i>nebulosus</i> .	355	Pachylabrus.	411
<i>nothus</i> .	355	<i>variegatus</i> .	411
<i>obliquatus</i> .	355	pachylepis, <i>Atherinichthys</i> .	240
<i>reticulatus</i> .	355	<i>Thyrinops</i> .	249
<i>rhomboidealis</i> .	354	Pachynathus.	492
<i>squamipinnis</i> .	355	Pachypops.	349
<i>stolzmanni</i> .	358	<i>furcraea</i> .	349
<i>thalassinus</i> .	355	pachypus, <i>Cottus</i> .	386
<i>toe-roe</i> .	354	Pachyurus.	89
<i>viridescens</i> .	350	<i>pacifica</i> , <i>Belone</i> .	197
<i>Otophidian</i> .	485	<i>pacifici</i> , <i>Barbatoichthoides</i> .	487
<i>galeoides</i> .	485	<i>Batrachus</i> .	487
<i>omostigma</i> .	485	<i>Conodon</i> .	331
<i>scrippsi</i> .	485	<i>Paraconodon</i> .	331
<i>taylori</i> .	485	<i>pacificum</i> , <i>Diplectrum</i> .	318
<i>welshi</i> .	485	<i>pacificus</i> , <i>Argyrelosus</i> .	276
<i>otophorus</i> , <i>Eupomacentrus</i> .	414	<i>Artediellus</i> .	377
<i>Pomacentrus</i> .	414	<i>Bathylagus</i> .	69

List of scientific names—Continued

Page	Page
<i>pacificus</i> , <i>Gastrostomus</i>	95
<i>Glyptocephalus</i>	228
<i>Larimus</i>	344
<i>Letharchus</i>	85
<i>Lobotes</i>	323, 324
<i>Lycodes</i>	474
(<i>Lycodopsis</i>).....	474
<i>Lycodopsis</i>	474
<i>Mesoprion</i>	325
<i>Microstomus</i>	228
<i>Psenes</i>	269
<i>Salmo</i> (<i>Mallotus</i>).....	66
<i>Thaleichthys</i>	67
<i>Thynnis</i>	260
<i>Tylosurus</i>	197
<i>Verrugato</i>	324
<i>padangensis</i> , <i>Gobius</i>	440
<i>petetus</i> , <i>Citharichthys</i>	218
<i>pagei</i> , <i>Etheostoma</i>	292
<i>Nivicolus</i>	292
<i>Pagellus calamus</i>	336
<i>humilis</i>	337
<i>microps</i>	337
<i>milneri</i>	337
<i>orbitarius</i>	336
<i>pentu</i>	336, 337
<i>Pagrus</i>	337
<i>pagrus</i>	337
<i>vulgaris</i>	337
<i>pagrus</i> , <i>Pagrus</i>	337
<i>Sparus</i>	337
<i>Pajelus caninus</i>	336
<i>Palamita</i>	259
<i>palaeirus</i> , <i>Lycodes</i>	475
<i>Palinurichthys</i>	267
<i>perciformis</i>	268
<i>Palinurus</i>	267
<i>palladus</i> , <i>Labrus</i>	270, 302
<i>pallasi</i> , <i>Molaacanthus</i>	503
<i>Oogrammus</i>	373, 374
<i>Pallasia</i>	503
<i>Pallasia</i>	503
<i>pallasi</i>	503
<i>pallasi</i> , <i>Clupea</i>	41
<i>Pleuronectes</i>	226
<i>Pallasina</i>	394
<i>aix</i>	394
<i>barbata</i>	394
<i>pallida</i> , <i>Aldrovandia</i>	75
<i>Echenels</i>	449
<i>Lepomis</i>	298
<i>Morone</i>	307
<i>pallidus</i> , <i>Bodianus</i>	345
<i>Catostomus</i>	105
<i>Fundulus</i>	176
<i>Labrus</i>	301
<i>Lepomis</i>	299
<i>Leucosomus</i>	117
<i>Pimelodus</i>	152
<i>Platygobio</i>	136
<i>Pomotis</i>	301, 302
<i>Salmo</i>	59
<i>Xystropistes</i>	302
<i>palmaripes</i> , <i>Trigla</i>	407
<i>paloma</i> , <i>Trachinotus</i>	277
<i>Palometa</i>	266
<i>media</i>	267
<i>palometa</i>	267
<i>simillima</i>	266
<i>snyderi</i>	267
<i>palometa</i> , <i>Chorinomus</i>	278
<i>Oligoplites</i>	278
<i>Palometra</i>	267
<i>Stromateus</i>	266, 267
<i>Trachinotus</i>	276
<i>palustris</i> , <i>Pecilichthys</i>	294
<i>Pammelas</i>	267
<i>pammelas</i> , <i>Melanostigma</i>	478
<i>pampus</i> , <i>Trachinotus</i>	276
<i>panamense</i> , <i>Pristipoma</i>	333, 334
<i>panamensis</i> , <i>Ælurichthys</i>	148
<i>Anchoviella</i>	50
<i>Antherinella</i>	249
<i>Azovia</i>	221
<i>Bagre</i>	148
<i>Bascanichthys</i>	86
<i>panamensis</i> , <i>Caranx</i>	273
<i>Citharichthys</i>	221
<i>Engraulis</i>	50
<i>Evermannia</i>	445
<i>Hyphephosbrycon</i>	97
<i>Loricaria</i>	162
<i>Menticirrhus</i>	351
<i>Microleotris</i>	437
<i>Muraena</i>	45
<i>Odontognathus</i>	91
<i>Oxyloricaria</i>	162
<i>Panamicichthys</i>	157
<i>Parapssettus</i>	369
<i>Petrometopon</i>	308
<i>Piabucina</i>	99
<i>Priapichthys</i>	187
<i>Pristigaster</i> (<i>Odontognathus</i>).....	45
<i>Rabula</i>	91
<i>Ithencus</i>	334
<i>Serranus</i>	305
<i>Solea</i>	230
<i>Tetragonopterus</i>	96
<i>Umbrina</i>	351
<i>Panamicichthys</i>	187
<i>panamensis</i>	187
<i>pandionis</i> , <i>Apogon</i>	290
<i>Emblemaria</i>	464
<i>Xystrania</i>	206
<i>pandora</i> , <i>Clinostomus</i>	120
<i>panduratus</i> , <i>Leiobatus</i>	23
<i>panbosa</i> , <i>Scorpiena</i>	371
<i>pantherina</i> , <i>Muraenophis</i>	93
<i>Oplopoma</i>	374
<i>pantherinus</i> , <i>Cestracion</i>	10
<i>Ichthyoiphis</i>	94
<i>Pseudaroides</i>	159
<i>Pantosteus</i>	104
<i>arizone</i>	104
<i>columbianus</i>	105
<i>dolphinus</i>	104
<i>jordani</i>	104
<i>lahontan</i>	105
<i>platyrhynchus</i>	104
<i>santacruzae</i>	104
<i>virescens</i>	105
<i>pantostigmus</i> , <i>Myrichthys</i>	85
<i>pantoco</i> , <i>Gambusia</i>	186
<i>papalis</i> , <i>Diodona</i>	144
<i>papillifer</i> , <i>Caulistius</i>	488
<i>Gobiosox</i>	488
<i>papillifera</i> , <i>Forbesella</i>	114
<i>papilliferus</i> , <i>Chologaster</i>	154
<i>papilio</i> , <i>Melletta</i>	580
<i>papillosum</i> , <i>Moxostoma</i>	109
<i>Syactum</i>	218
<i>papillosus</i> , <i>Barbulifer</i>	446
<i>Pleurotoctes</i>	218
<i>Ptychostomus</i>	100
<i>Parachromis</i>	418
<i>gulosus</i>	418
<i>Paraclinus</i>	457
<i>chapleri</i>	457
<i>Paraconodon</i>	331
<i>crescius</i>	331
<i>dovii</i>	332
<i>pacifici</i>	331
<i>Paradiodon</i>	501
<i>paradoxa</i> , <i>Garmannia</i>	445
<i>paradoxus</i> , <i>Gobius</i>	445
<i>Psychrolutes</i>	392
<i>Stylophthalmus</i>	71
<i>Parafundulus</i>	182
<i>Paragambusia</i>	184
<i>Paragonus</i>	364
<i>Parahemiodon</i>	161
<i>typus</i>	161
<i>Parajulis</i>	426
<i>Paralabrax</i>	317
<i>clathratus</i>	317
<i>humeralis</i>	317
<i>maculatusfasciatus</i>	317
<i>nebulifer</i>	317
<i>(Paralabrax) dewegeri</i> , <i>Serranus</i>	319
<i>Paralepis</i>	165
<i>barracudina</i>	165
<i>borealis</i>	165

List of scientific names—Continued

	Page		Page
<i>Paralepis coregonoides</i>	165	<i>Pararterdius</i>	377
<i>coruscans</i>	165	<i>hankinsoni</i>	377
<i>intermedius</i>	165	<i>Parascaphirhynchus</i>	35
<i>paralepis</i> , <i>Coregonus</i>	165	<i>albus</i>	35
<i>Paralichthysidæ</i>	222	<i>Parascorprena</i>	370
<i>Paralichthys</i>	223	<i>Paraserranus</i>	318
<i>estuarinus</i>	224	<i>parasiticus</i> , <i>Simenchelys</i>	78
<i>albiguttus</i>	224	<i>atrimanus</i>	297
<i>albiguttus</i>	224	<i>Parastroylytes</i>	376
<i>brasiliensis</i>	224	<i>notospilotus</i>	376
<i>californicus</i>	223	<i>Paratractus</i>	272
<i>dentatus</i>	224	<i>caballus</i>	272
<i>lethostigma</i>	224	<i>crysos</i>	272
<i>maculosus</i>	223	<i>Parauchenipterus</i>	159
<i>magdalene</i>	224	<i>pardale</i> , <i>Lepophidium</i>	485
<i>oblongus</i>	224	<i>Leptophidium</i>	485
<i>ommatus</i>	224	<i>pardalis</i> , <i>Monacanthus</i>	493
<i>ophryas</i>	224	<i>Mycteroperca</i>	315
<i>sinaloae</i>	224	<i>Xystroperca</i>	315
<i>squamulentus</i>	224	<i>pardus</i> , <i>Batrachus tau</i>	487
<i>woolmani</i>	224	<i>Opsanus</i>	487
<i>Paraliparis</i>	404	<i>Parehippus</i>	358
<i>albescens</i>	404	<i>Parepinephelus</i>	315
<i>angustifrons</i>	405	<i>acutirostris</i>	315
<i>attenuatus</i>	404	<i>Pareques</i>	353
<i>bathybii</i>	404	<i>acuminatus</i>	353
<i>caudatus</i>	404	<i>pulcher</i>	353
<i>cephalus</i>	404	<i>punctatus</i>	353
<i>copel</i>	405	<i>umbrosus</i>	353
<i>dactylosus</i>	404	<i>viola</i>	353
<i>deani</i>	404	<i>Parexocetus</i>	200
<i>fimbriatus</i>	404	<i>mesogaster</i>	200
<i>garmani</i>	405	<i>pargus</i> , <i>Mesopriion</i>	325
<i>grandiceps</i>	404	<i>Paricelinus</i>	375
<i>holomesias</i>	404	<i>hopliticus</i>	375
<i>latifrons</i>	405	<i>thoburni</i>	375
<i>liparinus</i>	405	<i>paraleialis</i> , <i>Coliscus</i>	145
<i>membranaceus</i>	404	<i>parilis</i> , <i>Ophichthus</i>	88
<i>mento</i>	404	<i>Ophisurus</i>	88
<i>roseaceus</i>	404	<i>parismima</i> , <i>Brachyrhaphis</i>	187
<i>ulochir</i>	405	<i>Gambusia</i>	187
<i>parallelus</i> , <i>Centropomus</i>	306	<i>parkel</i> , <i>Salmo</i>	61
<i>Embryx</i>	474	<i>parma</i> , <i>Heros</i>	419
<i>Paralongichurus</i>	351	<i>Parapetenia</i>	419
<i>goodet</i>	351	<i>parmatus</i> , <i>Setarches</i>	372
<i>petersi</i>	351	<i>parmifera</i> , <i>Raja</i>	26
<i>Paramia</i>	296	<i>Parodon duriensis</i>	100
<i>paranasiomos</i> , <i>Acienser</i> (<i>Hiuso</i>).....	35	<i>piracicabae</i>	100
<i>Paraneetroplus</i>	423	<i>Parophrys</i>	225
<i>bulleri</i>	423	<i>ayresi</i>	225
<i>Paranotropis</i>	125	<i>hubbardi</i>	226
<i>ariommus</i>	125	<i>ischyrus</i>	226
<i>jejunus</i>	125	<i>vetulus</i>	225, 226
<i>megalops</i>	125	<i>parovanus</i> , <i>Myloleucus</i>	121
<i>scabriceps</i>	125	<i>parra</i> , <i>Diabasis</i>	329
<i>Paranthias</i>	320	<i>Hæmulon</i>	329
<i>furcifer</i>	320	<i>parreæ</i> , <i>Albula</i>	40
<i>Parapetenia</i>	418	<i>Brama</i>	426
<i>adspersa</i>	418	<i>Chætodon</i>	361
<i>alfari</i>	419	<i>Clepticus</i>	426
<i>bartoni</i>	419	<i>Exocottus</i>	202
<i>beani</i>	419	<i>parraianus</i> , <i>Monacanthus</i>	494
<i>centralis</i>	419	<i>parræ</i> , <i>Lycocara</i>	478
<i>dovii</i>	418	<i>Ophidium</i>	188
<i>friedrichstahli</i>	418	<i>parryi</i> , <i>Rhamdia</i>	414
<i>istlana</i>	418	<i>partitus</i> , <i>Eupomacentrus</i>	414
<i>labridens</i>	418	<i>Pomacentrus</i>	360
<i>managuense</i>	418	<i>paru</i> , <i>Chætodon</i>	360
<i>mento</i>	419	<i>Peprilus</i>	266
<i>motaguensis</i>	418	<i>Pomacanthus</i>	360
<i>multifasciata</i>	418	<i>Stromateus</i>	266
<i>nigricans</i>	410	<i>Parupeneus</i>	343
<i>parma</i>	419	<i>parvæ</i> , <i>Anchovia</i>	49
<i>salvinii</i>	418	<i>Anchoviella</i>	49
<i>steindachneri</i>	419	<i>Echenis</i>	449
<i>tenuis</i>	419	<i>Lucania</i>	175
<i>tetracantha</i>	419	<i>Pleuronectes satis</i>	217
<i>cincta</i>	419	<i>parviceps</i> , <i>Lycodapus</i>	478
<i>grisea</i>	419	<i>parvidens</i> , <i>Moxostoma</i>	110
<i>late</i>	419	<i>parvipinne</i> , <i>Etheostoma</i>	292
<i>torvalbasi</i>	419	<i>parvipinnis</i> , <i>Ancylodon</i>	353
<i>maculata</i>	420	<i>brevis</i> , <i>Fundulus</i>	176
<i>urophthalmna</i>	418	<i>Cynoscion</i>	356
<i>Parapsettus</i>	359	<i>Dicrotus</i>	261
<i>panamensis</i>	359		

List of scientific names—Continued

	Page	Page	
<i>parvipinnis</i> , <i>Eretscion</i>	356	<i>pelamides</i> , <i>Scomber</i>	258
<i>Fundulus</i>	176	<i>pelamidis</i> , <i>Rhynchichthys</i>	236
<i>Hesperoleucus</i>	122	<i>pelamis</i> , <i>Katsuwonus</i>	258
<i>Isopisthus</i>	353	<i>Scomber</i>	258
<i>Ptax</i>	281	<i>pelamitus</i> , <i>Scomber</i>	259
<i>parvula</i> , <i>Clupea</i>	41, 42	<i>Pelamys</i>	259
<i>parvulus</i> , <i>Myoxocephalus</i>	387	<i>chilensis</i>	259
<i>parvus</i> , <i>Cottopsis</i>	383	<i>lineolata</i>	259
<i>Cyprinodon</i>	175	<i>orientalis</i>	259
<i>Gobius</i>	430	<i>pelegrinus</i> , <i>Squalus</i>	20
<i>Rhinogobius</i>	430	<i>Pellona</i>	44
<i>Sebastes ruber</i>	368	<i>bleekeriana</i>	45
<i>Upeneus</i>	343	<i>orbignyana</i>	44
<i>passan</i> , <i>Apterontotus</i>	101	<i>pellucida</i> , <i>Salmo</i>	214, 215
<i>Passer</i> <i>lipels transversis notatus</i>	230	<i>pellucidum</i> , <i>Etheostoma</i>	283
<i>passer</i> , <i>Centropyge</i>	361	<i>Ophidium</i>	79
<i>Holacanthus</i>	361	<i>pellucidus</i> , <i>Actis</i>	244
<i>Pastinaca</i>	28	<i>Osphryxolax</i>	243, 244
<i>hastata</i>	28	<i>Pleurolepis</i>	288
<i>marina</i>	29	<i>Psenes</i>	269
<i>olivacea</i>	28	<i>Thyris</i>	220
<i>pastinaca</i> , <i>Raja</i>	28	<i>Vigil</i>	288
<i>Pastinachus</i>	28	<i>Pelodichthys</i>	155
<i>centrourus</i>	28	<i>Peloria</i>	217
<i>schmardae</i>	28	<i>heckeli</i>	217
<i>torrel</i>	28	<i>pelta</i> , <i>Chelmo</i>	359
<i>patagonica</i> , <i>Mayne</i>	476	<i>peiltastes</i> , <i>Lepomis</i>	300
<i>patana</i> , <i>Macrodon</i>	99	<i>peiltatum</i> , <i>Etheostoma</i>	283
<i>patao</i> , <i>Gerres</i>	342	<i>peiltatus</i> , <i>Alvordius</i>	283
<i>patatus</i> , <i>Julis</i>	426	<i>Turdus cinereus</i>	341
<i>patris</i> , <i>Acanthocottus</i>	389	<i>Pempheridae</i>	223
<i>patronus</i> , <i>Brevoortia</i>	44	<i>Pempheris</i>	323
<i>patruells</i> , <i>Gambusia</i>	185	<i>mexicana</i>	323
<i>Heterandria</i>	184	<i>mexicanus</i>	323
<i>pattersoni</i> , <i>Troglolagris</i>	155	<i>milleri</i>	323
<i>patzcuaro</i> , <i>Chirostoma</i>	250	<i>poeyi</i>	323
<i>paucidens</i> , <i>Leuryurus</i>	474	<i>schoembergi</i>	323
<i>Salmo</i>	65	<i>tourea</i>	323
<i>pauciflorus</i> , <i>Ophichthys</i>	88	<i>peninsulæ</i> , <i>Bascanichthys</i>	80
<i>pauciradiatus</i> , <i>Callionymus</i>	450	<i>Callicheilus</i>	86
<i>Pseudoxiphophorus</i>	188	<i>Chirostoma</i>	247
<i>paucispinis</i> , <i>Sebastodes</i>	364	<i>Menidia</i>	247
<i>Sebastodes</i>	364	<i>peninsulae atrimentis</i> , <i>Menidia</i>	247
<i>payo</i> , <i>Xyrichthys</i>	430	<i>penna</i> , <i>Calamus</i>	337
<i>pavonaceus</i> , <i>Herichthys</i>	422	<i>Pagellus</i>	337
<i>Heros</i>	422	<i>pennanti</i> , <i>Argentina</i>	73
<i>pavonina</i> , <i>Limia</i>	191	<i>Maurolicus</i>	73
<i>pavoninus</i> , <i>Adonis</i>	460	<i>Squalus</i>	19
<i>Cyclopterus</i>	397	<i>pennatula</i> , <i>Calamus</i>	336
<i>Inistioides</i>	430	<i>Penopus</i>	482
<i>Xyrichthys</i>	430	<i>macdonaldi</i>	482, 483
<i>pauroloides</i> , <i>Lycodes</i>	476	<i>pensacola</i> , <i>Harengula</i>	43
<i>pautilus</i> , <i>Lycenchelys</i>	476	<i>penshinensis</i> , <i>Salmo</i>	56
<i>Lycodes</i>	476	<i>pentacanthus</i> , <i>Asterotheca</i>	235
<i>peckianus</i> , <i>Syngnathus</i>	242	<i>pentacanthus</i> , <i>Bodianus</i>	303
<i>peckii</i> , <i>Syngnathus</i>	242	<i>Centrarchus</i>	303
<i>pectinatus</i> , <i>Centropomus</i>	306	<i>Labrus</i>	424
<i>Pristis</i>	23	<i>Xenochirius</i>	396
<i>pectinifer</i> , <i>Clinus</i>	459	<i>Pentanemus</i>	256
<i>Leucidius</i>	123	<i>quinquartius</i>	256
<i>pectoralis</i> , <i>Albatrossia</i>	204	<i>pepinus</i> , <i>Lucioperca</i>	282
<i>Bathyonus</i>	482	<i>pepo</i> , <i>Scarus</i>	433
<i>Bathypterois</i>	166	<i>Peprilus</i>	266
<i>Dactyloscopus</i>	454	<i>paru</i>	266
<i>Dallia</i>	172	<i>snyderi</i>	267
<i>Harpe</i>	425	<i>perarcutatus</i> , <i>Pleuronectes</i>	226
<i>Hemigaleus</i>	14	<i>Perca</i>	281
<i>Macrourus</i>	204	<i>aberrans</i>	316
(<i>Malacocephalus</i>).....	204	<i>acuta</i>	281
<i>Nematistius</i>	280	<i>alburnus</i>	350
<i>Nematonus</i>	482	<i>americana</i>	281, 307
<i>Prionotus</i>	407	<i>apoda</i>	325
<i>miles</i>	407	<i>ascensionis</i>	235
<i>Pedallion</i>	503	<i>atratia</i>	317
<i>pedallota</i> , <i>Bonapartia</i>	72	<i>chrysops</i>	307
<i>pedallotus</i> , <i>Zaphotias</i>	72	<i>chrysoptera</i>	336
<i>pedicellaris</i> , <i>Sciadonus</i>	483	<i>chrysopterus</i>	335
<i>Pediculatus</i>	504	<i>flavescens</i>	281
<i>pedimacula</i> , <i>Centropomus</i>	306	<i>flavopurpurea</i>	310
<i>Pediglyctis</i>	383, 385	<i>fluviatilis</i>	281
<i>ictalops</i>	383, 385	<i>gibbosa</i>	302
<i>Pelxe aquila</i>	195	<i>formosa</i>	348, 330
<i>pelagicus</i> , <i>Callionymus</i>	409	<i>furcata</i>	349
<i>Nectoliparis</i>	402	<i>furva</i>	317
<i>Scomber</i>	264	<i>gibbosa</i>	302, 329
<i>Syngnathus</i>	240-242	<i>gigas</i>	310

List of scientific names—Continued

	Page		Page
<i>Percia gracilis</i>	281	<i>Peristedion brevirostre</i>	405
<i>granulata</i>	281	<i>erustosum</i>	406
<i>guttata</i>	308, 311	<i>gracile</i>	405
<i>guttatus</i>	308	<i>longispathum</i>	405
<i>immaculata</i>	307	<i>malurmat</i>	405
<i>juba</i>	332	<i>microneurus</i>	406
<i>lanceolata</i>	352	<i>miniatum</i>	405
<i>lophar</i>	280	<i>platycephalum</i>	406
<i>loubina</i>	305	<i>Peristedium barbiger</i>	406
<i>maculata</i>	310	<i>erustosum</i>	406
<i>marina</i>	363	<i>longispatha</i>	405
<i>capito striato</i>	330	<i>platycephalum</i>	406
<i>cauda nigra</i>	330	<i>Peristethus</i>	405
<i>gibbosa cinerea</i>	329	<i>brevirostre</i>	405
<i>rubra</i>	236	<i>peristethus</i> , <i>Podothecus</i>	394
<i>secatrix</i>	340	<i>Perkinsia</i>	46
<i>melanura</i>	330	<i>othomops</i>	46
<i>minima</i>	287	<i>perlatum</i> , <i>Holocentrum</i>	236
<i>mitchilli</i>	307	<i>perlongus</i> , <i>Leptoconger</i>	82
<i>alternata</i>	307	<i>Neoconger</i>	82
<i>interrupta</i>	307	<i>perniger</i> , <i>Culius</i>	437
<i>muconata</i>	308	<i>Eleotris</i>	437
<i>nigra</i>	267	<i>Peropis</i>	332
<i>nobiils</i>	333	<i>perplexus</i> , <i>Cottus</i>	334
<i>norwegica</i>	363	<i>perrico</i> , <i>Pseudoscarus</i>	435
<i>notata</i>	281	<i>Scarus</i>	435
<i>ocellata</i>	348	<i>Perro colorado</i>	425
(<i>Percina</i>) <i>nebulosa</i>	283	<i>perrotteti</i> , <i>Pristis</i>	23
<i>philadelphica</i>	318	<i>personata</i> , <i>Eviota</i>	438
<i>punctata</i>	309, 345	<i>personatus</i> , <i>Ammodytes</i>	451
<i>pusilla</i>	295	<i>perspicabilis</i> , <i>Embletoeca</i>	411
<i>rhomboidalis</i>	338	<i>perspicillaris</i> , <i>Tetraodon</i>	500
<i>robusta</i>	311	<i>perspicillum</i> , <i>Lycodes</i>	474
<i>rufa</i>	236	<i>perspicuus</i> , <i>Hyborhynchus</i>	145
<i>salmonea</i>	282	<i>pertheata</i> , <i>Anchoviella</i>	47
<i>saltatrix</i>	280	<i>pertheatus</i> , <i>Stolephorus</i>	47
<i>saxatilis</i>	307	<i>perlanus</i> , <i>Anthus</i> (<i>Hemianthias</i>)	322
<i>secatrix</i>	340	<i>Hemianthias</i>	322
<i>septentrionalis</i>	307	<i>perugiae</i> , <i>Limia</i>	191
<i>serrato-granulata</i>	281	<i>Platypræciulus</i>	191
<i>stellio</i>	310	<i>peruvianus</i> , <i>Diapterus</i>	342
<i>striata</i>	331	<i>Galeichthys</i>	149
<i>tota</i>	310	<i>Geres</i>	342
<i>trifurca</i>	318	<i>Pescador</i>	505
<i>undulata</i>	349	<i>Petalosoma</i>	193
<i>unicolor</i>	321	<i>cultratum</i>	193
<i>unimaculata</i>	338	<i>Petalurichthys</i>	193
<i>varia</i>	317	<i>petenensis</i> , <i>Astyanax</i>	97
<i>variabilis</i>	365	<i>Moletta</i>	47
<i>venenosus</i>	313	<i>Mollinges</i>	102
<i>ventre luteo</i>	302	<i>Pimeledous</i>	158
<i>vitrea</i>	282	<i>Poecilia</i>	192
<i>perciliens</i> , <i>Raja</i>	23	<i>Rhamdia</i>	158
<i>Rhinobatos</i>	23	<i>Signlosa</i>	47
<i>Percoscoes</i>	281, 282	<i>Tetragonopterus</i>	97
<i>Percidae</i>	267, 268	<i>Petenia</i>	423
<i>percoformis</i> , <i>Coryphaena</i>	268	<i>splendida</i>	423
<i>Palinurichthys</i>	282	<i>petersi</i> , <i>Centropholis</i>	266
<i>Percina</i>		<i>Paralonchurus</i>	351
<i>bimaculata</i>	283	<i>petimbina</i> , <i>Fistularia</i>	245
<i>caprodes</i>	283	<i>Petrometopon</i>	308
<i>carbonaria</i>	283	<i>cruentatus</i>	308
<i>manitou</i>	283	<i>coronatus</i>	308
<i>nebulosa</i>	282	<i>panamensis</i>	308
<i>rex</i>	283	<i>Petromyzon</i>	8
<i>zebra</i>	283	<i>americanus</i>	8
(<i>Percina</i>) <i>nebulosa</i> , <i>Percia</i>	283	<i>appendix</i>	9
<i>percobromus</i> , <i>Alburnellus</i>	124	<i>argenteus</i>	9
<i>percoides</i> , <i>Agonostoma</i>	254	(<i>Artedi</i>)	8
<i>Dajus</i>	254	<i>astori</i>	10
<i>Percomorphi</i>	246	<i>ayresii</i>	10
<i>Percopsidae</i>	214	(<i>Bathymyzon</i>) <i>bairdii</i>	8
<i>Percopsis</i>	214	<i>bdephilum</i>	9
<i>guttatus</i>	214, 215	<i>borealis</i>	10
<i>hammondi</i>	215	<i>ciliatus</i>	10
<i>omiscomaycus</i>	215	<i>epiphexodon</i>	10
<i>peregrinus</i> , <i>Leptopus</i>	215	<i>fluvialis</i>	10
<i>perezi</i> , <i>Platypondon</i>	16	<i>lamotieni</i>	10
<i>perfasciata</i> , <i>Anchoviella</i>	47	<i>lividus</i>	10
<i>perfasciatus</i> , <i>Engraulis</i>	47	<i>marinus</i>	8
<i>periscopius</i> , <i>Gadus</i>	209	<i>dorsatus</i>	9
<i>Perissas</i>	220	<i>nigricans</i>	8
<i>tæniopterus</i>	220	<i>nlgrum</i>	9
<i>Peristediidae</i>	405	<i>plumbeus</i>	10
<i>Peristedion</i>	405	<i>tridentatus</i>	9, 10
<i>barbiger</i>	406	<i>Petromyzonidae</i>	8

List of scientific names—Continued

	Page		Page
Petronas...	433	Photogonls, <i>scabriceps</i> ...	125
petrosus, <i>Brycon</i> ...	98	spilopterus...	129
Mugil...	253	stigmaturus...	129
Serranus...	313	telescopus...	124
petus, <i>Acanthocybium</i> ...	250	photogenis, <i>Notropis</i> ...	123, 128
Cybium...	250	Squalius...	123
pfeifferi, <i>Muraena</i> ...	93	Photonectes...	70
Pfrille...	121	gracilis...	70
neogaea...	121	Phoxnagyrea...	247
phenina, <i>Hybopsis</i> ...	132	daii...	247
Phænodon...	71	Phoxinus clevelandi...	118
ringens...	71	flammeus...	121
Phæthonichthys...	264	milnerianus...	121
tuberculatus...	264	neogaeus...	121
phæton, <i>Pristigaster</i> ...	45	orcuttii...	120
Serranus...	309	phoxocephalus, <i>Cynoscion</i> ...	356
phalæna, <i>Umbrina</i> ...	350	Etheostoma...	284
Phalangistes...	395	phoxocephalus, <i>Alvordius</i> ...	284
loricatus...	394	Atractoscion...	356
phaleratus, <i>Esox</i> ...	173	phrynooides, <i>Eumicrotromus</i> ...	398
Phallichthys...	190	Pitheirichthys...	449
anates...	190	tropicus...	449
pittieri...	190	Phycis...	212, 213
Phanerodon...	411	chesteri...	213
atripes...	411	cirratus...	212
furcatus...	411	dekeyi...	213
Pharyngognathi...	424	earilli...	212
phasganorius, <i>Notacanthus</i> ...	76	floridanus...	213
phasma, <i>Careproctus</i> ...	402	marginatus...	212, 213
Phenacobius...	142	rostratus...	213
catostomus...	142	tinca...	212
mirabilis...	142	varrellii...	213
scopifer...	142	Phycis, <i>Lota</i> ...	211
teretulus...	142	Phycocetes...	478
<i>hosternus</i> ...	142	latitans...	478
uranops...	142	suspectus...	478
phenacobius, <i>Hybopsis</i> ...	134	Physiculus...	211
Notropis...	134	dalwigkii...	211
Phenacimates...	95	fulvus...	212
macrolepis...	95	kaupi...	211
wilsoni...	95	longipes...	212
phenax, <i>Aponotis</i> ...	299	nematopus...	212
Mycetoperca...	315	rastrelliger...	212
<i>falcata</i> ...	315	physignathus, <i>Ceratichthys</i> ...	136
philadelphicus, <i>Perca</i> ...	318	Platygobio...	136
philadelphicus, <i>Triloburus</i> ...	318	Physostomii...	39
philonips, <i>Cottus</i> ...	384	Phytichthys...	409
Philypnus...	435	<i>chirus</i> ...	409
<i>lateralis</i> ...	436	<i>versicolor</i> ...	409
phlebotomus, <i>Acanthurus</i> ...	362	Pistibrama...	99
phlegethonitis, <i>Clinostomus</i> ...	121	<i>erythrinoides</i> ...	99
<i>loti</i> ...	121	<i>festae</i> ...	99
phlegothontis, <i>Clinostomus</i> ...	121	<i>panamensis</i> ...	99
phlox, <i>Boleosoma</i> ...	287	Plectartli, <i>Hemiramphus</i> ...	198
<i>Ulocentra</i> ...	287	piceus, <i>Balistes</i> ...	493
Phobetor...	389	Melichthys...	493
phoibe, <i>Prionodes</i> ...	319	pichardi, <i>Joturus</i> ...	254
<i>Serranus</i> ...	319	Plicorellus...	172
Pholididae...	466	picta, <i>Muraena</i> ...	93
Pholidichthys...	465	Scorpaena...	370
<i>anguilliformis</i> ...	465	Siderea...	93
<i>leucotenia</i> ...	465	pleratura, <i>Aluteria</i> ...	495
Pholis...	461, 467	Gambusia...	185
<i>carolinus</i> ...	461	picturatus, <i>Gymnothorax</i> ...	91
<i>fasciatus</i> ...	468	Syngnathus...	241
<i>gilli</i> ...	468	pictus, <i>Chaura</i> ...	500
<i>gunnellus</i> ...	467	Chironectes...	504
<i>novemlineatus</i> ...	462	Chirurus...	374
<i>ornatus</i> ...	468	Eleotris...	437
<i>quadrifasciatus</i> ...	462	Gymnophorus...	477
<i>subblifurcatus</i> ...	470	Histrion...	504
pholis, <i>Blennius</i> ...	461	Iridio...	427
Photocorynus...	509	Julis...	427
<i>spiniceps</i> ...	509	Oxyblebius...	374
Photogenis...	123	Sciades...	148
<i>ariomimus</i> ...	125	Torpedo...	27
<i>cæruleus</i> ...	130	Urocentrus...	467
<i>callistius</i> ...	130	picuda, <i>Sphyraena sphyraena</i> var...	255
<i>eurystomus</i> ...	130	picudilla, <i>Sphyraena</i> ...	255
<i>grandipinnis</i> ...	129	pidiens, <i>Moxostoma</i> ...	110
<i>leucodus</i> ...	124	pidiensis, <i>Ptychostomus</i> ...	110
<i>leucops</i> ...	123	piger, <i>Syphurus</i> ...	230
<i>engraulinus</i> ...	130	pigmentarius, <i>Apogon</i> ...	295
<i>leucopus</i> ...	130	<i>Monopriion</i> ...	295
<i>piptolepis</i> ...	134	plgra, <i>Aporistia</i> ...	230
<i>pyrrhomelas</i> ...	129		

List of scientific names—Continued

	Page		Page
<i>pilchardus</i> , <i>Clupea</i>	43	<i>Pimelodus</i> , <i>marmoratus</i>	154
<i>pileatus</i> , <i>Prionotus</i>	282	<i>megalops</i>	152
<i>Plioema</i>	282	<i>micropterus</i>	157
<i>carbonaria</i>	283	<i>motaguensis</i>	157
<i>cymatogramma</i>	286	<i>natalis</i>	153
<i>semifasciata</i>	282	<i>nebulosus</i>	154, 155
<i>semifasciatum</i>	283	<i>nicaraguensis</i>	157
<i>zebra</i>	283	<i>nigricans</i>	152
<i>Pilicorvis</i> , <i>Bleennius</i>	461	<i>notatus</i>	152
<i>Pilodictis</i>	155	<i>pallidus</i>	152
<i>limosa</i>	155	<i>potenensis</i>	158
<i>pilosa</i> , <i>Solea</i>	230	<i>platycephalus</i>	154
<i>pilosus</i> , <i>Diodon</i>	501	<i>polycaulus</i>	158
<i>Trichododon</i>	501	<i>pulius</i>	154
<i>pilsbryi</i> , <i>Notropis</i>	124	<i>punctulatus</i>	155
<i>Pimeleopterus</i>	339	<i>quelon</i>	156
<i>analogus</i>	339	<i>rigidus</i>	159
<i>boscii</i>	339	<i>rogersi</i>	158
<i>sicula</i>	340	<i>salvini</i>	157
<i>bosquii</i>	340	<i>spixii</i>	150
<i>cornubiensis</i>	268	<i>viscosus</i>	155
<i>elegans</i>	339	<i>vulgaris</i>	153
<i>flavo-lineatus</i>	339	<i>vulpeculus</i>	154
<i>incisor</i>	339	<i>vulpes</i>	152
<i>lutescens</i>	340	<i>wagneri</i>	157
<i>oblongior</i>	340	<i>Pimelometopon</i>	420
<i>occurus</i>	340	<i>pulcher</i>	420
<i>sandvicensis</i>	339	<i>Pimephales</i>	145
<i>Pimelode insigne</i>	155	<i>agassizii</i>	145
<i>livree</i>	156	<i>anuli</i>	146
<i>Pimelodella</i>	158	<i>confertus</i>	145
<i>chagresi</i>	158	<i>fasciatus</i>	145
<i>Pimelodidae</i>	156	<i>maculosus</i>	145
<i>Pimelodon</i>	155	<i>milesi</i>	145
<i>insignarius</i>	155, 150	<i>promelas</i>	145
<i>Pimelodus</i>	158	<i>pingelli</i> , <i>Triglops</i>	379
<i>genaeus</i>	155	<i>pinguis</i> , <i>Pleuronectes</i>	221
<i>affinis</i>	152	<i>pinnima</i> , <i>Acara</i>	332
<i>allurus</i>	153	<i>pinnata</i> , <i>Muraena</i>	78
<i>albidus</i>	150, 153	<i>pinnatus</i> , <i>Eretmichthys</i>	481
<i>antoniensis</i>	153	<i>Synaphobranchus</i>	78
<i>argentinus</i>	152	<i>pinniger</i> , <i>Brysseteres</i>	488
<i>argyrus</i>	152	<i>Enneacanthus</i>	304
<i>ascita</i>	158	<i>Gobiesox</i>	488
<i>atrarius</i>	154	<i>Rosicola</i>	365
<i>baroni-mülleri</i>	157	<i>Sebastosomus</i>	365
<i>borealis</i>	152	<i>pinnimaculatus</i> , <i>Ailurichthys</i>	148
<i>brachycephalus</i>	157	<i>Bagre</i>	148
<i>brachypterus</i>	157	<i>pinnivaria</i> , <i>Hypoplectrus</i>	321
<i>catulus</i>	154	<i>pinnivarius</i> , <i>Hypoplectrus unicolor</i>	321
<i>caudafurcatus</i>	152	<i>pinnulata</i> , <i>Seriola</i>	279
<i>chagresi</i>	158	<i>pinta</i> , <i>Muraena</i>	90
<i>claritas</i>	159	<i>pintata</i> , <i>Muraena</i>	92
<i>conosus</i>	153	<i>iptolepis</i> , <i>Hybopsis</i>	134
<i>confinis</i>	153	<i>Photogenis</i>	134
<i>conirostris</i>	156	<i>Piquitinga</i>	48
<i>cristatus</i>	156	<i>piquotianus</i> , <i>Lepiosteus</i>	37
<i>cupresoides</i>	153	<i>piquotii</i> , <i>Amia</i>	38
<i>cupreus</i>	153	<i>piraca</i> , <i>Monacanthus</i>	494
<i>cyclopum</i>	153	<i>practicalbe</i> , <i>Parodon</i>	100
<i>deKayi</i>	152	<i>Piramutanga</i> , <i>blochii</i>	150
<i>erythrophtera</i>	153	<i>macrospila</i>	159
<i>felinus</i>	153	<i>pirapeda</i> , <i>Dactylopterus</i>	409
<i>febris</i>	154	<i>Pirati apia</i>	313
<i>furcatus</i>	152	<i>pirrense</i> , <i>Cyclopium</i>	160
<i>furcifer</i>	152	<i>pirrensis</i> , <i>Astroblepus</i>	160
<i>godmani</i>	157	<i>pisavarlus</i> , <i>Ophichthys</i>	85
<i>gracilis</i>	152	<i>piscatorius</i> , <i>Lophius</i>	504
<i>graciosus</i>	152	<i>piscatrix</i> , <i>Pseudorhamphus</i>	159
<i>guatemalensis</i>	157	<i>Pisces</i>	33
<i>hammondi</i>	152	<i>pisciculus</i> , <i>Esox</i>	176
<i>houghi</i>	152	<i>piscis viridis bahamensis</i>	432
<i>hoylei</i>	154	<i>piscis bahamensis</i> , <i>Unicornis</i>	495
<i>hypselurus</i>	157	<i>pisculentus</i> , <i>Esox</i>	176
<i>insigne</i>	155	<i>Pisodonophis</i>	86
<i>insignis</i>	156	<i>cancrivorus</i>	86
<i>lateralis</i>	152	<i>daspilatus</i>	86
<i>laticaudus</i>	157	<i>guttilatus</i>	85
<i>leucoptera</i>	152	<i>Pisodonophis</i> , <i>Electris</i>	86
<i>lupus</i>	153	<i>Pisodonophis</i> , <i>Gobius</i>	436
<i>lynx</i>	153	<i>Pisodonophis</i> , <i>eruentifer</i>	86
<i>macronema</i>	159	<i>oculatus</i>	85
<i>maculatus</i>	151, 152, 158, 159	<i>pisquetus</i> , <i>Caranx</i>	272
<i>managuensis</i>	157		
<i>marginatus</i>	152		

List of scientific names—Continued

	Page	Page	
pistilliger, <i>Cottus</i>	389	platorynchus, <i>Acipenser</i>	35
<i>Gymnoanthus</i>	389	<i>Scaphirhynchops</i>	35
<i>pitamba, Acara</i>	327	<i>platostomus, Cylindrostomus</i>	37
Pithecomyzon.....	103	<i>Lepisosteus</i>	37
pitieri, <i>Phallichthys</i>	190	Platotichthys.....	221
<i>Præcilia</i>	190	<i>chartes</i>	221
pitulatus, <i>Rhypticus</i>	316	<i>platyargyreus, Anchovia</i>	49
pixanga, <i>Serranus</i>	310	<i>Anchovela</i>	49
pixuma, <i>Amore</i>	438	<i>platycephalum, Peristedion</i>	406
placitus, <i>Hybognathus</i>	143	<i>Peristedium</i>	406
Placopharynx.....	111	<i>Platycephalus dormitor</i>	435
<i>carinatus</i>	111	<i>platycephalus, Amelurus</i>	154
Plagiogrammus.....	469	<i>Cottus</i>	387, 388
<i>hopkinsi</i>	469	<i>Megalocottus</i>	388
plagioplateo, <i>Electris capite</i>	436	<i>Noturus</i>	155
Plagiostoma.....	10	<i>Notorynchus</i>	11
Plagiusa.....	230	<i>Pimelodus</i>	154
<i>lactea</i>	230	Platygaster.....	44
plagiusa, <i>Pleuronectes</i>	231	Platyglossus <i>dipilus</i>	427
<i>Syphurnus</i>	231	<i>fluorealis</i>	428
Plagopterus.....	147	<i>nicholsi</i>	428
<i>argentissimus</i>	147	<i>poeyi</i>	427
Plagusia.....	230, 231	Platygobio.....	136
<i>brasiliensis</i>	231	<i>gracilis</i>	130
<i>fasciata</i>	231	<i>paulli</i>	136
<i>lactea</i>	230	<i>physignathus</i>	136
<i>tessellata</i>	231	(<i>Platygobio</i>) <i>gulonellus, Pogonichthys</i>	136
plagiusa, <i>Pleuronectes</i>	230, 231	Platyinius.....	328
<i>Syphurnus</i>	231	<i>platyodon, Carcharias</i>	15
Plagiodontem.....	171	<i>Squalus</i>	15
Plagyodus.....	171	Platypodon.....	15
<i>ascalapius</i>	171	<i>perezii</i>	16
plana, <i>Platea</i>	227	Platypocilius.....	193
plancl, <i>Tympanomium</i>	603	<i>couchianus</i>	193
Plancterus.....	179	<i>dominicensis</i>	191
<i>kansae</i>	179	<i>maculatus</i>	193
<i>zebra</i>	179	<i>immaculatus</i>	193
planiceps, <i>Ancistrus</i>	161	<i>mentalis</i>	192
<i>Arius</i>	149	<i>perugiae</i>	191
<i>Catostomus</i>	107	<i>quitzeoensis</i>	182
<i>Lasiancistrus</i>	161	<i>tropicus</i>	192
<i>Netuma</i>	149	<i>variatus</i>	193
planiceps, <i>Netuma</i>	149	playtypogon, <i>Arius</i>	149
planifrons, <i>Eupomacentrus</i>	414	<i>Netuma</i>	149
<i>Pomacentrus</i>	414	Platyhina <i>exasperata</i>	24
Planirostra.....	36	<i>triseriata</i>	24
<i>edentula</i>	36	Platyrhinidae.....	24
planus, <i>Pleuronectes</i>	227	Platyrhinoidis.....	24
Plaryurus <i>argentatus</i>	128	<i>triseriatus</i>	24
<i>bowmani</i>	128	platyrhinus, <i>Acipenser (Huso)</i>	35
<i>melanocephalus</i>	145	platyrhynchus, <i>Carcharias</i>	15
<i>typicus</i>	128	<i>Eulamia</i>	15
plaryurus, <i>Rutilus</i>	128	<i>Minomus</i>	104
plate, <i>Branchiostoma</i>	7	<i>Pantosteus</i>	104
Platessa.....	226	Platysomatichthys.....	221
<i>bilineata</i>	226	<i>stomias</i>	221
<i>dentata</i>	222	Platysomus.....	274
<i>dvinensis</i>	227	<i>micropteryx</i>	275
<i>ferruginea</i>	226, 227	<i>spixii</i>	275
<i>glabra</i>	227	Platysqualus.....	17
<i>oblonga</i>	223, 224	<i>tudes</i>	17
<i>ocellaris</i>	224	Platytroctes.....	53
<i>plana</i>	227	<i>apus</i>	53
<i>pusilla</i>	227	plebelus, <i>Catostomus</i>	104
<i>rostrata</i>	227	<i>Gasterosteus</i>	237
platessa, <i>Pleuronectes</i>	226	<i>Notolepidomyzon</i>	104
platessoides, <i>Hippoglossoides</i>	222	plebejus, <i>Mustelus</i>	13
<i>Pleuronectes</i>	222	Plecostomus.....	160
Platichthys.....	228	<i>boulengeri</i>	160
<i>rugosus</i>	226	<i>dorsopterygio</i>	160
<i>stellatus</i>	226	<i>seminudus</i>	160
<i>umbrosus</i>	226	plecostomus, <i>Hypostomus</i>	160
Platirostra.....	36	<i>Loricaria</i>	160
<i>edentula</i>	36	Plectobranchus.....	468, 469
Platophrys.....	217	<i>evides</i>	468, 469
<i>constellatus</i>	217	Plectognath.....	491
<i>ellipticus</i>	217	<i>plectrodon, Porichthys</i>	488
<i>leopardinus</i>	218	Plectromus.....	233
<i>lunatus</i>	217	<i>beani</i>	233
<i>maculifer</i>	217	<i>bispinosus</i>	233
<i>nebularis</i>	217	<i>crassiceps</i>	233
<i>ocellatus</i>	217	<i>cristiceps</i>	233
<i>spinous</i>	217	<i>lugubris</i>	233
<i>teichopterus</i>	220	<i>mizolepis</i>	233
platophrys, <i>Citharichthys</i>	219	<i>nycterinus</i>	233
Platopterus.....	24	<i>suborbitalis</i>	233

List of scientific names—Continued

	Page		Page
Plectropoma accensum	321	Pleuronichthys guttulatus	225
affine	321	nephelus	225
bovinum	321	ocellatus	225
chloropterum	312	ritteri	225
chlorurum	321	verticalis	225
crocota	321	pleurophthalmus, Antennarius	505
ephippium	321	pleurospilus, Girardinus	190
gummigutta	321	Poeciliastes	190
guttavarium	321	pleurostictus, Triglops	379
hispanum	308	Pleurothyris	74
indigo	321	Phloeostomna	45
melanorhina	321	lutipinnis	45
monecanthus	313	plumatula, Calamus	336
multiguttatum	313	plumbea, Chimæra	33
nigricans	321	Dionda	144
puella	320, 321	Gambusia	192
vitulinum	321	plumbeolus, Alburnops	128
Plectrypops	236	plumbeus, Bathylalopex	33
retrospinis	236	Cousesius	136
pleianus, Pseudoscarus	435	Gobio	136
Scarus	435	Petromyzon	10
pleii, Hemiramphus	199	Pseudoscarus	435
pleuriticus, Salmo	57	rubrilateralis, Cousesius	116
Pleuracromyion	13	plumiceanus, Caranxomorus	270
Pleurogrammus	374	plumieri, Le Diodon	501
monopterygius	374	plumieri, Chaetodon	358
Pleurolepis	288	Coryphaena	356
asprellus	288	Eugerres	343
pellucidus	288	Gerres	342, 343
Pleuronectes	226	Gobius	447
achirus	229	Himantulon	330
americanus	227	Lobrus	330
apoda	229	Mahicanthus	356
aquosus	217	Mugil	253
aramaca	218	Polydactylus	256
argus	217	Sciaena	333
asper	227	Scomber	271
cicatricosus	227	Scomberomorous	257
cornutus	225	Scorpaena	370
cynoglossus	221	Sicydiun	447
dentatus	224	plumierianus, Caranxomorus	270
diagrammus	226	plumieri, Polydactylus	256
ellipticus	217	plurisetialis, Narctes	52
franklini	227	plutonia, Raja	26
fuscus subtrotundus glaber	230	pluvialis, Tetronodon	409
gilli	228	pluvius, Salmo	61
glabra	217	Pneumatophorus	257
glacialis	227	australiasicus	258
hippoglossoides	221	diego	258
hippoglossus	221	grex	258
kitt	228	pneumatophorus, Scomber	257, 258
limanda	222, 226	pocatello, Catostomus	107
limandooides	222	Podager	341
lineatus	230	podus, Bothus	217
linguatula	222	Podoletus	215
lunatus	217	podostemone, Ethostoma	286
maculatus	217	Imostoma	289
maculiferus	217	Podothecus	304
melanogaster	224	acipectenirinus	394
microcephalus	228	gilberti	305
microstomus	228	peristethus	394
mollis	229	Pœcilia	100
oblongus	224	(Acropœcilia) tridens	100
obscurus	228	amates	100
ocellatus	217	boucardi	192
oculis etc.	229	butleri	192
pallasii	226	catenata	179
papillosus	218	chisoensis	102
perarcutus	226	cönica	175
pinguis	221	cuneata	193
plagiusa	231	dominicensis	191
plagiusa	230, 231	dovii	192
planus	227	elongata	130
platessa	226	limantouri	192
piatessoides	222	lineolata	192
quadridens	228	melanogaster	101
quadrifurculatus	226	molapleura	178, 185
quenseli	228	mexicana	102
satis parva	217	multilineata	192
saxicola	228	olivacea	177
stellatus	226	petenensis	192
surinamensis	217	pittieri	190
Pleruronectidae	225	presidionis	180
Pleuronichthys	225	reticulata	191
cœnosus	225	reticulatus	191
decurrans	225	salvatoris	192

List of scientific names—Continued

	Page		Page
Poecilia schneideri.....	100	Pogonias.....	352
sphenops.....	102	chromis.....	352
spilonota.....	102	courbina.....	352
spilurus.....	102	fuscatus.....	352
surinamensis.....	100	Pogonichthys.....	116
tenuis.....	102	argyrotaenia.....	118
thermales.....	102	communis.....	136
tridens.....	100	inæquilobus.....	116
vanderpoli.....	103	macrolepidotus.....	116
arubensis.....	103	(<i>Platygobio</i>) gulosellus.....	136
vittata.....	101	symmetricus.....	121
vivipara.....	100	Pogonius chromis.....	352
Poeciliichthys.....	282, 289	polaris, <i>Blennius</i>	476
artesiae.....	292	Cottus.....	388
aspregenis.....	291	Lycodalepis.....	476
beani.....	287	Merlangus.....	209
borealis.....	291	Porocottus.....	388
butlerianus.....	294	Polipturus.....	257
camurus.....	290	Polistiotrema.....	8
jessiae.....	291	curtissi-jamesii.....	8
mesaeus.....	287	deani.....	8
palustris.....	294	stoutii.....	8
punctulatus.....	292	politus, <i>Chelichthys</i>	499
quiесcens.....	294	Seriphis.....	353
rufilineatus.....	290	Tetraodon.....	409
sagitta.....	293	Pollachius.....	209
sanguifluus.....	290	chalcopterus fuscensis.....	209
saxatilis.....	287	virens.....	209
spectabilis.....	291	Pollachius, <i>Gadus</i>	209
swaini.....	291	<i>bullicaris, Uranidea</i>	388
variatus.....	289	Pollux, <i>Pontius</i>	372
versicolor.....	291	Polyacanthocephalus, <i>Cottus</i>	387
virgatus.....	292	<i>Myoxocephalus</i>	387
vitreus.....	288	Polyacanthonotus.....	76
vulnératus.....	290	challengeri.....	76
zonalis.....	289	rostratus.....	76
Poeciliidae.....	184	Polyactis, <i>Melanocetus</i>	510
Poeciliopsis.....	189	Polyactocephalus, <i>Bryostomina</i>	460
balsas.....	189	Polyactocephalus, <i>Blennius</i>	460
fasciata.....	189	Polycanthus.....	239
infans.....	189	polycaula, <i>Rhamdia</i>	158
isthmensis.....	190	polycaulus, <i>Pimelodus</i>	158
latidens.....	189	Polycirrhus.....	351
occidentalis.....	189	dumerili.....	351
presidionis.....	189	rathbuni.....	351
turubarensis.....	189	Polyclemus.....	351
Poeciliastes.....	189	Polydactylus.....	250
pleurospilus.....	190	opercularis.....	256
Poecilocephalus.....	87	plumieri.....	256
bonapartei.....	87	plumieri.....	256
Poecilodes.....	187	Polygonius, <i>Acanthostracion</i>	497
bimaculatus.....	187	Gymnothorax.....	92
poeciloides, <i>Lebiasina</i>	191	Polylepis, <i>Balistes</i>	492
Limia.....	192	Scorpiena.....	370
Poecilophis.....	94	Polyipnus.....	74
nocturnus.....	94	laternatus.....	74
Poeciliopsis isthmensis.....	150	spinifer.....	74
Poeciliopterus, <i>Julis</i>	426	Polyipnus, <i>Verrunculus</i>	492
Poeciliopus, <i>Myripristis</i>	235	Polymixia.....	234
<i>Rhamphoberyx</i>	235	lowei.....	234
Poecilosoma.....	289	nobilis.....	234
erythrogasterum.....	291	Polymixitidæ.....	234
transversum.....	291	polymorphus, <i>Gadus</i>	210
Poeciliura, <i>Myoxostoma</i>	111	Polyinemidae.....	250
Poecilurichthys.....	96	Polyinemus.....	250
brevirostris.....	96	americanus.....	250
pulcher.....	97	approximans.....	250
taenurus.....	97	artedi.....	250
unilineatus.....	98	californiensis.....	250
Poecilurum, <i>Moxostoma</i>	111	macronomus.....	250
Poecilus.....	100	mango.....	250
Poeyi, <i>Alepidosaurus</i> (<i>Caulopus</i>).....	171	melanopoma.....	250
Dactyloscopus.....	455	octonemus.....	250
Engraulis.....	51	oligodon.....	250
Euctenogobius.....	442	paradiseus.....	250
Gobius.....	442	quinquevittatus.....	250
Iridio.....	427	sexradiatus.....	409
Lycengraulis.....	51	virginicus.....	250
Orthopriscis.....	335	Polyodon.....	36
Pempheris.....	323	feuille.....	36
Platyglossus.....	427	folium.....	36
Siphostoma.....	241	spatula.....	36
Syngnathus.....	241	Polyodon, <i>Tetragonopterus</i> (<i>Hemibrycon</i>).....	97
Synodus.....	104	Polyodontidae.....	36
Pogonathus.....	352		
courbina.....	352		

List of scientific names—Continued

	Page		Page
<i>Polypera</i>	399	<i>Pomotis converifrons</i>	300
<i>greeni</i>	399	<i>elongatus</i>	300
<i>polygon</i> , <i>Linophryne</i>	510	<i>fallax</i>	300
<i>Polyprion</i>	308	<i>gibbosus</i>	301
<i>americanus</i>	308	<i>gulosus</i>	302, 303
<i>cernuum</i>	308	<i>guttatus</i>	304
<i>massilensis</i>	308	<i>heros</i>	302
<i>Polyprosopus</i>	20	<i>holbrooki</i>	302
<i>macer</i>	20	<i>incisor</i>	301
<i>Polypterichthys</i>	244	<i>inscriptus</i>	300
<i>valentini</i>	244	<i>longulus</i>	299
<i>polysarcus</i> , <i>Naucrates</i>	278	<i>luna</i>	301
<i>polyspilos</i> , <i>Rhombus</i>	223	<i>microlophus</i>	302
<i>Polyuranodon</i>	91	<i>nefastus</i>	300
<i>Polyuranodon</i> , <i>Muraena</i>	91	<i>nitida</i>	300
<i>Pomacanthodes</i>	360	<i>notatus</i>	302
<i>zonipictus</i>	360, 361	<i>obesus</i>	303
<i>Pomacanthus</i>	360	<i>obscurus</i>	301
<i>arcuatus</i>	360	<i>pallidus</i>	301
<i>balteatus</i>	360	<i>popeii</i>	300
<i>cingulatus</i>	360	<i>raveneli</i>	302
<i>crescentialis</i>	361	<i>rubri-cauda</i>	299
<i>paru</i>	360	<i>sanguinolentus</i>	300
<i>quinquecinctus</i>	360	<i>solis</i>	300
<i>pomacentrus</i> , <i>Mesopriion</i>	324	<i>speciosus</i>	301, 302
<i>Pomacentridæ</i>	412	<i>vulgaris</i>	302
<i>Pomacentrus</i>	413	<i>pomotis</i> , <i>Acantharchus</i>	303
<i>adustus</i>	414	<i>Centrarchus</i>	303
<i>analigutta</i>	414	<i>Pomoxis</i>	304
<i>analis</i>	413	<i>annularis</i>	304
<i>atro-cyanus</i>	413	<i>nitidus</i>	304
<i>bairdii</i>	415	<i>sparoides</i>	305
<i>breviceps</i>	413	<i>Pomoxys</i> <i>bericanda</i>	304
<i>caudalis</i>	413	<i>intermedius</i>	304
<i>denegatus</i>	415	<i>protacanthus</i>	304
<i>dorsopunicans</i>	413	<i>pompilius</i> , <i>Centrolophus</i>	267
<i>flavilatus</i>	414	<i>Pomphilus</i>	267
<i>flaviventer</i>	414	<i>pomphilus</i> , <i>Coryphaena</i>	267
<i>fuscus</i>	413	<i>Thynnus</i>	278
<i>gillii</i>	414	<i>ponderosus</i> , <i>Amiaurus</i>	152
<i>leucorus</i>	414	<i>pondicerriana</i> , <i>Elaeotis</i>	281
<i>leucostictus</i>	413	<i>ponticus</i> , <i>Gasterosteus</i>	237
<i>lividus</i>	413	<i>Hippoglossus</i>	221
<i>niveatus</i>	410	<i>Pontinus</i>	372
<i>obscuratus</i>	414	<i>beanorum</i>	372
<i>otophorus</i>	414	<i>bergii</i>	372
<i>partitus</i>	414	<i>castor</i>	372
<i>planifrons</i>	414	<i>furciferinus</i>	372
<i>quadrigutta</i>	416	<i>longispinus</i>	372
<i>rectifrenum</i>	414	<i>macrolepis</i>	372
<i>variabilis</i>	413	<i>pollux</i>	372
<i>xanthurus</i>	413	<i>rathbuni</i>	372
<i>Pomadasys</i> <i>bayanus</i>	334	<i>sierra</i>	372
<i>elongatus</i>	333	<i>popeii</i> , <i>Pomotis</i>	300
<i>grandis</i>	334	<i>popoche</i> , <i>Xystrosus</i>	142
<i>Pomadasys</i>	334	<i>porcus</i> , <i>Caprichus</i>	491
<i>approximans</i>	334	<i>Scorpea</i>	370
<i>bayanus</i>	334	<i>Porichthys</i>	487
<i>boschmai</i>	334	<i>greeni</i>	487
<i>cæsius</i>	331	<i>margaritatus</i>	487
<i>grandis</i>	334	<i>nautopedium</i>	487
<i>macracanthus</i>	334	<i>notatus</i>	487
<i>productus</i>	334	<i>plectrodon</i>	488
<i>sinuosus</i>	335	<i>porifer</i> , <i>Lycenchelys</i>	476
<i>starri</i>	334	<i>Lycodes</i>	476
<i>templei</i>	334	<i>Porobronchus</i>	486
<i>Pomatomichthys</i>	206	<i>linearis</i>	486
<i>constanciae</i>	206	<i>Poroclinus</i>	470
<i>Pomatomidae</i>	280	<i>rothrocki</i>	470
<i>Pomatomus</i>	280	<i>Porocottus</i>	388
<i>saltatrix</i>	280	<i>bradfordi</i>	388
<i>skib</i>	280	<i>polaris</i>	388
<i>telescopium</i>	296	<i>quadratus</i>	388
<i>Pomataprion</i>	415	<i>quadrifilis</i>	388
<i>Pomatopsetta</i>	222	<i>sellaris</i>	388
<i>Pomolobus</i>	41, 42	<i>Porogadus</i>	482
<i>aestivalis</i>	42	<i>atripectus</i>	482
<i>chrysichloris</i>	41	<i>breviceps</i>	482
<i>mediochlora</i>	42	<i>longiceps</i>	482
<i>pseudoharengus</i>	42	<i>miles</i>	482
<i>lacustris</i>	42	<i>pomelas</i>	481
<i>Pomotis</i>	299	<i>Poromitra</i>	233
<i>bombifrons</i>	301	<i>capito</i>	233
<i>breviceps</i>	300	<i>Poronotus</i>	267
<i>catesbeii</i>	302	<i>simillimus</i>	266
<i>chaetodon</i>	303	<i>triacanthus</i>	267

List of scientific names—Continued

Page	Page
porosissimum, <i>Nantopædium</i>	488
porosissimus, <i>Batrachus</i>	487, 488
porosus, <i>Carcharias</i>	16
<i>Cottus</i>	387
<i>Esox</i>	173
<i>Scoliodon</i>	15
<i>Squalus</i>	15
postica, <i>Echeneis</i>	449
<i>Potamocottus</i>	383
<i>bendirei</i>	384
<i>carolinae</i>	383, 385
<i>punctulatus</i>	383
<i>zopherus</i>	385
pottsii, <i>Aplexion</i>	290
<i>Ranhesquielius</i>	290
powelli, <i>Balistes</i>	492
præcinctus, <i>Clinus</i>	470
<i>Eumesogrammus</i>	470
prædatoria, <i>Buccone</i>	354
prædatorius, <i>Cestreus</i>	354
præstabilis, <i>Alosa</i>	42
præstiglator, <i>Serranus</i>	319
prayensis, <i>Pseudupeneus</i>	343
preblei, <i>Coregonus</i>	65
<i>Prosoptium</i>	65
prenadilla, <i>Brontes</i>	189
presidionis, <i>Pacilia</i>	189
<i>Puciliopsis</i>	189
pretiosa, <i>Argentina</i>	67
pretiosus, <i>Hypomesus</i>	67
<i>Ruvettus</i>	261
<i>Trachichthys</i>	234
Pricanthichthys.....	310
<i>maderaspatensis</i>	310
Pricanthidæ.....	322
Priacanthus.....	322
<i>altus</i>	323
<i>arenatus</i>	322
<i>carolinus</i>	322
<i>catalufa</i>	323
<i>cepedelanus</i>	323
<i>cruentatus</i>	322
<i>fulgens</i>	322
<i>macrophthalmus</i>	322
<i>niphonius</i>	322
<i>serrula</i>	323
Priapella.....	188
<i>bonita</i>	188
Priapichthys.....	187
<i>annectans</i>	187
<i>hesperis</i>	187
<i>fosteri</i>	189
<i>huberi</i>	187
<i>letonat</i>	190
<i>olomina</i>	187
<i>panamensis</i>	187
<i>pribilovius</i> , <i>Nautichthys</i>	382
<i>Nauticus</i>	382
<i>pricel</i> , <i>Campostoma</i>	147
<i>Haemulon</i>	153
<i>Villarius</i>	152, 153
Pricus.....	10
prieto, <i>Lutjanus</i>	325
Prionodon.....	500
(<i>Anchisomus</i>) <i>caudacinctus</i>	500
Primospina.....	364
princeps, <i>Caulophilus</i>	357
<i>Cottus</i>	384
<i>Latilis</i>	357
principis, <i>Autenarius</i>	505
<i>Chironectes</i>	505
<i>Julis</i>	420
Prionodon.....	180
Priodontophis.....	91
Prionace.....	17
<i>glauca</i>	17
Prionatus <i>egretta</i>	408
Prionistius.....	380
<i>macellus</i>	380
Prionodes.....	319
<i>equidens</i>	319
<i>baldwini</i>	310
<i>bulleri</i>	319
<i>dewaegei</i>	319
<i>fuscatus</i>	319
<i>flavescens</i>	319
Prionodes fusculus.....	319
<i>phœbe</i>	319
<i>tabacaria</i>	319
<i>tigrinus</i>	319
Prionodon.....	17
<i>cucurl</i>	16
<i>obvelatus</i>	16
(<i>Prionodon</i>) <i>limbatus</i> , <i>Carcharias</i>	16
<i>milberti</i> , <i>Carcharias</i>	16
Prionotus.....	497
<i>alatus</i>	408
<i>albirostris</i>	408
<i>beanii</i>	408
<i>birostratus</i>	408
<i>evolans</i>	407
<i>frontalis</i>	408
<i>gymnostethus</i>	408
<i>horrens</i>	408
<i>loxias</i>	408
<i>miles pectoralis</i>	407
<i>militaris</i>	408
<i>ophryas</i>	407
<i>pectoralis</i>	407
<i>pileatus</i>	407
<i>punctatus</i>	407
<i>quiescens</i>	408
<i>rosicus</i>	407
<i>rubio</i>	407
<i>rusarius</i>	408
<i>sarrtor</i>	407
<i>scitulus</i>	407
<i>stearnsi</i>	406
<i>stephanophrys</i>	407
<i>strigatus</i>	407
<i>tribulus</i>	408
<i>xenisma</i>	406
Prionurus punctatus.....	363
Pristidae.....	23
Pristidurus.....	12
Pristigaster.....	45
<i>cayanus</i>	45
<i>dovii</i>	45
<i>flavipinnis</i>	44
<i>lutipinnis</i>	45
<i>macrops</i>	45
<i>martii</i>	45
(<i>Odontognathus</i>) <i>panamensis</i>	45
<i>phatton</i>	45
<i>tarto</i>	45
Pristipoma.....	334
<i>acara-pinima</i>	333
<i>auratum</i>	333
<i>bicolor</i>	332
<i>bilineatum</i>	332
<i>boschmai</i>	334
<i>boucardi</i>	334
<i>branickii</i>	332
<i>brasiliense</i>	332
<i>brevipinne</i>	335
<i>camtharinum</i>	335
<i>catharinæ</i>	333
<i>chalecum</i>	335
<i>crocro</i>	334
<i>cultiferum</i>	334
<i>davidsoni</i>	332
<i>dovii</i>	332
<i>fasciatum</i>	335
<i>furthii</i>	332
(<i>Hæmulopsis</i>) <i>axillare</i>	333
<i>brevipinne</i>	335
<i>nitidum</i>	333
<i>humile</i>	334
<i>humilis</i>	335
<i>kneri</i>	333
<i>leuciscus</i>	333
<i>macracanthum</i>	334
<i>melanopterum</i>	332
<i>notatum</i>	332
<i>panamense</i>	333
<i>productum</i>	334
<i>ramosum</i>	334
<i>rodo</i>	333
<i>scapulare</i>	332
<i>spleniatum</i>	332
<i>serrula</i>	333
<i>trilineatum</i>	332

List of scientific names—Continued

	Page		Page
Pristipomes, les.	334	Promicrops.	312
Pristipomus.	334	italara.	312
Pristipomoides.	328	quinquefasciata.	312
macropthalmus.	328	Promicropodus.	316
typus.	328	bistrispinus.	316
Pristis.	23	decoratus.	317
acutirostris.	23	nigripinnis.	317
granulosa.	23	Pronotogrammus.	317
megalodon.	23	eos.	322
microdon.	23	multifasciatus.	322
mississippiensis.	23	proops, Bagrus.	322
occa.	23	Sciadeichthys.	322
pectinatus.	23	Propterygia.	324
zephyreus.	23	hyposticta.	324
Pristis, Squalus.	23	prorates, Lepophidium.	324
Pristurus.	12, 13	Leptophidium.	324
are.	12	proridens, Calanus.	326
melastomas.	12	proriger, Gnathophis.	326
xanthurus.	12	Sebastichthys.	326
Pristocantharus.	335	var. brevispinis, Sebastichthys.	326
Pristopomus.	334	Squalius.	326
Proarthri.	10	prorigerum, Acutomentum.	326
probatocephalus, Archosargus.	338	Ophisoma.	326
Sparus.	338	proserpina, Moniana.	326
proboscidea, Limanda.	227	Prosopium.	326
proboscideum, Monacanthum.	495	abyssicola.	326
proboscideus, Chænomugil.	254	couesi.	326
Mugil.	254	coulteri.	326
procellarium, Myctophum.	170	kennicotti.	326
procera, Venefica.	82	prehlei.	326
Proceros.	36	quadrilaterale.	326
inaculatus.	36	spilonotus.	326
vittatus.	36	stanleyi.	326
procerum, Nettastoma.	82	williamsoni.	326
Prochilus.	436	cismontanus.	326
procne, Exocetus (Cypsilurus).	201	Prospinus.	312
Hybognathus.	133	prosthemius, Ceratichthys.	316
Hybopsis.	133	prostheustis, Ameiurus.	316
Proctostegus.	204	protacanthus, Pomoxys.	304
proctostegus.	264	proteus, Salmo.	54
proctostegus, Proctostegus.	264	protoculus, Nyctophum.	168
producta, Rhinobatus.	24	Protoperus.	119
Ulæma.	341	dominicus.	119
productum, Pristopoma.	334	proxima, Seriola.	279
productus, Aleopcephalus.	51	proximus, Gadus.	210
Cylindrostoeus.	37	Microgadus.	210
Engraulis.	50	ruinosus, Gadus tomcodus.	210
Eucinostomus.	341	Pseudoblennius.	405
Leuciscus.	120	hypacanthus.	405
Merlangus.	214	Psempilius prometheus.	261
Merluccius.	214	Psenes.	268
Pomadasys.	334	auratus.	268
Rhinobatos.	24	chapmani.	268
preclaris, Microperca.	204	cyanophrys.	268
profundorum, Acanthocottus.	292	edwardsi.	268
Apiristurus.	12	fuscus.	268
Lepophidium.	484	javanicus.	268
Leptophidium.	484	maculatus.	268
Scyliorhinus.	12	pacificus.	269
Xenognathus.	52	rellucidus.	269
Zesticeles.	302	regulus.	269
Profundulus.	174	Psettidichthys.	222
candalarius.	174	molanostictus.	222
labialis.	174	sordidus.	218
oaxaceae.	174	umbrosus.	228
punctatus.	174	Pseudagenetosus.	159
protinus, Mesopriion.	326	pseudaphites, Micropterus.	248
Prognathodes.	359	Pseudariooides.	158
aculeatus.	359	pantherinus.	159
prognathus, Coregonus.	63	Pseudarius.	160
Leucichthys.	63	Pseudocheirodon.	96
Prognurus.	403	affinis.	96
cypselurus.	403	Pseudochromidae.	322
prolixus, Leuciscus.	146	pseudoguia, Eucinostomus.	341
promelas, Chirostoma.	251	pseudo-gula, Eucinostomus.	341
McCbia.	481	pseudoharengus, Clupea.	42
Pimephales.	446	lacustris, Pomolobus.	42
Porogadus.	481	Pomolobus.	42
Prometheus.	261	Pseudohemiodon.	161
atlanticus.	261	Pseudohispanica, Sardinia.	43
prometheus, Gampylus.	261	Pseudojulis.	428
Promethichthys.	261	adustus.	428
Psempylus.	261	girardi.	428
Promethichthys.	261	inornatus.	428
prometheus.	261	melanotis.	428

List of scientific names—Continued

	Page		Page
<i>Pseudojulis notospilus</i>	428	<i>Pteroplates micrura</i>	29
<i>venustus</i>	428	<i>rava</i>	29
<i>Pseudoloricaria</i>	161	<i>vallianti</i>	29
<i>Pseudomonacanthus</i>	403	<i>Pteropodus</i>	369
<i>Pseudomoringua</i>	89	<i>carnatus</i>	369
<i>Pseudomuricea</i>	91	<i>caurinus</i>	369
<i>maderensis</i>	91	<i>chrysomelas</i>	369
<i>Pseudopleuronectes</i>	227	<i>dalli</i>	369
<i>americanus</i>	227	<i>gilberti</i>	369
<i>dignabilis</i>	227	<i>maliger</i>	369
<i>pseudoplatis</i> , <i>Micropterus</i>	298	<i>nebulosus</i>	369
<i>Pseudopriacanthus</i>	323	<i>rastrelliger</i>	369
<i>altus</i>	323	<i>vxillaris</i>	369
<i>serrula</i>	323	<i>Pteropsideidae</i>	451
<i>pseudopterus</i> , <i>Micropterus</i>	298	<i>Pteropus</i>	369
<i>Pseudorhamdia</i>	158	<i>pterotum</i> , <i>Ateleobrachium</i>	207
<i>piscatrix</i>	159	<i>Pterura</i> , <i>Strongylura</i>	196
<i>Pseudorhombus</i>	223	<i>Pterurus</i> , <i>Tylosurus</i>	196
<i>vorax</i>	224	<i>Pterygiocottus</i>	380
<i>Pseudoscarus</i>	435	<i>macouni</i>	380
<i>aracanga</i>	434	<i>Pterostichon imberbe</i>	406
<i>coelestinus</i>	435	<i>Ptilichthysidae</i>	471
<i>gnathodes</i>	434	<i>Ptilichthys</i>	471
<i>guacamai</i>	434	<i>goodei</i>	471
<i>lineolatus</i>	434	<i>Ptychocheilus</i>	113
<i>micros</i>	435	<i>gracilis</i>	113, 114
<i>microrhinos</i>	435	<i>grandis</i>	114
<i>perrico</i>	435	<i>harfordi</i>	114
<i>pleianus</i>	436	<i>lucius</i>	114
<i>plumbeus</i>	435	<i>major</i>	114
<i>simplex</i>	435	<i>oregonensis</i>	114
<i>tetraopterus</i>	433	<i>umpquae</i>	114
<i>Pseudoscopelus</i>	462	<i>vorax</i>	114
<i>scriptus</i>	452	<i>Ptycholepis</i>	40
<i>Pseudosebastes</i>	370	<i>Ptychostomus</i>	109
<i>Pseudotriakidius</i>	12	<i>alibidus</i>	110
<i>Pseudotriakis</i>	12	<i>albus</i>	110
<i>microdon</i>	12	<i>breviceps</i>	111
<i>Pseudoxiphophorus</i>	187	<i>bucco</i>	110
<i>bimaculatus</i>	187	<i>cervinus</i>	111
<i>jonesi</i>	188	<i>collapsus</i>	110
<i>teniatulus</i>	188	<i>conus</i>	111
<i>paticiradiatus</i>	188	<i>coregonus</i>	110
<i>reticulatus</i>	187	<i>crassilabris</i>	111
<i>Pseudupeneus</i>	343	<i>haydeni</i>	109
<i>prayensis</i>	343	<i>lacrymalis</i>	111
<i>Psilonotus</i>	500	<i>papillosum</i>	109
<i>psittacinus</i> , <i>Serranus</i>	319	<i>pidiensis</i>	110
<i>psittacus</i> , <i>Labrus</i>	428	<i>robustus</i>	111
<i>psittacus</i> , <i>Colomesus</i>	500	<i>thalassinus</i>	110
<i>Coryphena</i>	430, 433	<i>velutus</i>	109
<i>Lachnolaimus</i>	425	<i>Pudiano verde</i>	428
<i>Tetrodon</i>	500	<i>vermelho</i>	425
<i>Xyrichtyhs</i>	430	<i>puella</i> , <i>Hypoplectrus unicolor</i>	321
<i>Psychichthys</i>	33	<i>Plectropoma</i>	320, 321
<i>affinis</i>	33	<i>puellaris</i> , <i>Cossyphus</i>	425
<i>Psychrolutes</i>	392	<i>Decodon</i>	425
<i>paradoxus</i>	392	<i>puertoricensis</i> , <i>Malacoctenus</i>	460
<i>zebra</i>	392	<i>pugentus</i> , <i>Ariedius</i>	375
<i>Psychromaster</i>	293	<i>Chitonotus</i>	375
<i>tuscumbia</i>	293	<i>pugetti</i> , <i>Gasterosteus</i>	238
<i>Ptax</i>	261	<i>pulchella</i> , <i>Gilia</i>	119, 120
<i>parvipinnis</i>	261	<i>Leuciscus</i>	120
<i>Pteraclidae</i>	266	<i>Montiana</i>	131
<i>Pteraclidus</i>	266	<i>Tigoma</i>	120
<i>Pteracilus</i>	266	<i>pulchelloides</i> , <i>Leuciscus</i>	117
<i>carolinus</i>	266	<i>pulchellus</i> , <i>Bodianus</i>	425
<i>trichipterus</i>	266	<i>Cossyphus</i>	425
<i>Pterimates canariensis</i>	29	<i>Cymatogaster</i>	410
<i>valencienni</i>	29	<i>Leuciscus</i>	117
<i>Pterengraulis</i>	51	<i>Liparis</i>	400
<i>atherinoides</i>	51	<i>Lyoliparis</i>	400
<i>Pteridium</i>	266	<i>Oligocephalus</i>	291
<i>Pterocephala</i>	31	<i>Zygonectes</i>	177
<i>Pterognathus</i>	458	<i>pulcher</i> , <i>Eques</i>	363
<i>satricus</i>	458	<i>Labrus</i>	426
<i>Pteromylaeus</i>	30	<i>Pareques</i>	553
<i>asperrimus</i>	30	<i>Pimelometopon</i>	426
<i>Pteronotus</i>	156	<i>Poecilurichthys</i>	97
<i>Pterophryne</i>	504	<i>pulchra</i> , <i>Tigoma</i>	120
<i>Pterophrynoidea</i>	504	<i>pullata</i> , <i>Dicrolene</i>	483
<i>Pteroplatea</i>	29	<i>nullum</i> , <i>Chondrostoma</i>	146
<i>altavala</i>	29	<i>pullus</i> , <i>Cantherines</i>	404
<i>crebripunctata</i>	29	<i>Monacanthus</i>	494
<i>marmorata</i>	29	<i>Pimelodus</i>	154

List of scientific names—Continued

	Page		Page
<i>pulverentulus</i> , <i>Myloleucus</i>	119-121	<i>punctulatus</i> , <i>Calliurus</i>	298
<i>pulveres</i> , <i>Zygonectes</i>	178	<i>Cottus</i>	383
<i>pulverulenta</i> , <i>Tigoma</i>	120	<i>Hippocampus</i>	244
<i>pulverulentum</i> , <i>Oxygeneum</i>	113	<i>Lampugus</i>	265
<i>puncta</i> , <i>Furcaria</i>	412	<i>Lythrurus</i>	126
<i>puncticeps</i> , <i>Cryptopterus</i>	85	<i>Minnilus</i>	126
<i>punctarenæ</i> , <i>Gymnothorax</i>	92	<i>Pimelodus</i>	155
<i>punctata</i> , <i>Alutera</i>	495	<i>Poecilichthys</i>	292
<i>Davidia</i>	495	<i>Potomocottus</i>	383
<i>Gambusia</i>	184, 185	<i>Scarus</i>	433
<i>Girella</i>	338	<i>Sicyases</i>	490
<i>Gorgasia</i>	77	<i>Squalus</i>	12
<i>Lamna</i>	19	<i>Pungitius</i>	238
<i>Muraena</i>	91	<i>brachypoda</i>	239
<i>Muraenophis</i>	92	<i>mainensis</i>	239
<i>Perca</i>	309, 345	<i>pungitius</i>	239
<i>punctulata</i> , <i>Gambusia</i>	185	<i>pungitius brachypoda</i> , <i>Gasterosteus</i>	239
<i>Solea</i>	217	<i>pungitius</i> , <i>Gasterosteus</i>	238, 239
<i>Trigla</i>	407	<i>Pungitius</i>	239
<i>punctatissima</i> , <i>Anguilla</i>	77	<i>puniceus</i> , <i>Hyborhynchus</i>	145
<i>punctatissimus</i> , <i>Canthigaster</i>	500	<i>purpurascens</i> , <i>Lepomis</i>	301
<i>Ophioscion</i>	347	<i>variegata</i> , <i>Salpa</i>	327
<i>Tetronodon</i>	500	<i>purpuratus bouvieri</i> , <i>Salmo</i>	56
<i>punctatum</i> , <i>Cichlasoma</i>	417	<i>Salmo</i>	56
<i>Hoplosternum</i>	160	<i>purpurea</i> , <i>Tigoma</i>	119
<i>Hyperprosopon argenteum</i>	410	<i>purpurens</i> , <i>Anoplarchus</i>	408
<i>Megaloneema</i>	159	<i>archolepis</i> , <i>Anoplarchus</i>	468
<i>Myctophum</i>	169	<i>purpureus</i> , <i>Gadus</i>	209
<i>Opistognathus</i>	453	<i>Labrus</i>	428
<i>Ostracion</i>	496	<i>Lutjanus</i>	226
<i>Sicydium</i>	447	<i>pusilla</i> , <i>Aporistila</i>	231
<i>Theraps</i>	417	<i>Bothrocara</i>	477
<i>punctatus</i> , <i>Aporomatis</i>	299	<i>Clupea</i>	42
<i>Aprionodon</i>	15	<i>Maynea</i>	477
<i>Ballistes</i>	492	<i>Perca</i>	295
<i>Blennius</i>	462, 469	<i>Platesa</i>	227
<i>Bryttus</i>	298, 299	<i>pusillus</i> , <i>Coregonus</i>	62
<i>Caranx</i>	270, 273	<i>Enneapterygius</i>	458
<i>Carcharius</i>	15	<i>Leucichtthys</i>	62
<i>Cephalopholis fulvus</i>	309	<i>Syphurus</i>	231
<i>Cotylopus</i>	448	<i>putaoi</i> , <i>Gymnotus</i>	101
<i>Decapterus</i>	270	<i>putnami</i> , <i>Acipenser</i>	34
<i>Dermatolepis</i>	313	<i>Cottogaster</i>	285
<i>Diodon</i>	501	<i>Euchalarodus</i>	227
<i>Eques</i>	353	<i>Liopsetta</i>	227
<i>Fundulus</i>	174	<i>Pycnomma</i>	438
<i>Gadus</i>	213	<i>semisquamatum</i>	438
<i>Gymnothorax</i>	92	<i>Pygidiliæ</i>	159
<i>Hoplosternum</i>	160	<i>Pygidium</i>	159
<i>Ictalurus</i>	152	<i>fuscum</i>	159
<i>Isurus</i>	19	<i>striatum</i>	159
<i>Labrus</i>	416	<i>pygmaea bilineata</i> , <i>Umbrina</i>	174
<i>Maurolicus amethystinus</i>	73	<i>pygmaea</i> , <i>Eucalia</i>	238
<i>Micristodus</i>	18	<i>Pygmaeus</i> , <i>Gadus</i>	210
<i>Monacanthus</i>	494	<i>Gasterosteus</i>	238
<i>Morrhua</i>	211	<i>Leuciscus</i>	174
<i>Myrophis</i>	82, 83	<i>Pygosteus</i>	238
<i>Opistognathus</i>	453	<i>Pylodictis limous</i>	155
<i>Ostracion</i>	496	<i>pyramidatus</i> , <i>Cyclopterus</i>	397
<i>Pareques</i>	353	<i>pyrrhogaster</i> , <i>Chrosomus</i>	113
<i>Prionotus</i>	407	<i>pyrrhomelas</i> , <i>Ergalaea</i>	129
<i>Prionurus</i>	363	<i>Photogenis</i>	129
<i>Profundulus</i>	174	<i>Pythonichthys</i>	90
<i>Silurus</i>	151, 152	<i>sanguineus</i>	90, 91
<i>Silurus</i>	151, 152		
<i>Squalus</i>	12, 14, 15		
<i>Stichæus</i>	469		
<i>Tetronodon</i>	499		
<i>Trachinus</i>	310		
<i>Upeneus</i>	343		
<i>Xesurus</i>	363		
<i>puncticeps</i> , <i>Cryptopterichthys</i>	85		
<i>Cryptopterus</i>	85		
<i>puncticulata</i> , <i>Gambusia</i>	185		
<i>puncticulatus</i> , <i>Apogonichthys</i>	298		
<i>Chiromycterus</i>	502		
<i>Holocentrum</i>	236		
<i>punctifer</i> , <i>Crotalopsis</i>	88		
<i>Hybognathus (Dionda)</i>	144		
<i>Trigonophalus</i>	187		
<i>punctiferus</i> , <i>Menophorus</i>	309		
<i>punctillatus</i> , <i>Gobius</i>	440		
<i>punctipinnis</i> , <i>Ayrasia</i>	413		
<i>punctostethus</i>	240		
<i>puncto-mediorum</i> , <i>Antennarius</i>	505		
<i>punctulata</i> , <i>Gambusia punctata</i>	185		
<i>Microperca</i>	294		
<i>Nivicola</i>	292		

List of scientific names—Continued

	Page		Page
<i>quadrilineatum</i> , <i>Hæmulon</i>	331	<i>radiatus</i> , <i>Lophius</i>	507
<i>quadriloba</i> , <i>Raja</i>	31	<i>Ogcocephalus</i>	507
<i>quadriporus</i> , <i>Gobius</i>	439	<i>Sparus</i>	428
<i>Rhinogobius</i>	439	<i>radostus</i> , <i>Antennarius</i>	506
<i>quadripunctatus</i> , <i>Scomber</i>	259	<i>radula</i> , <i>Balistes</i>	493
<i>quadremus</i> , <i>Exocetus</i>	201	<i>Melichthys</i>	493
<i>quadrigeratus</i> , <i>Artedius</i>	376	<i>Radulinus</i>	379
<i>Icelinus</i>	376	<i>asprellus</i>	379
<i>quadrispinosus</i> , <i>Scarus</i>	434	<i>boleoides</i>	379
<i>quadrituberculatus</i> , <i>Pleuronectes</i>	226	<i>rafinesquel</i> , <i>Acipenser</i> (<i>Huso</i>).....	35
<i>quadrocellatus</i> , <i>Ancylosetta</i>	224	<i>Colletta</i>	168
<i>quappella</i> , <i>Etheostoma</i>	291	<i>Cylindrostomus</i>	37
<i>quartus rondeleti</i> , <i>Anthius</i>	326	<i>Nyctophorus</i>	168
<i>Quassilabia</i>	112	<i>Rafinesquillius</i>	290
<i>Quassiremurus</i>	87	<i>pottsi</i>	290
<i>evionthas</i>	87	<i>rafinesquili</i> , <i>Acipenser</i> (<i>Huso</i>).....	35
<i>nothochir</i>	87	<i>Cylindrostomus</i>	37
<i>quatuordecimlaminatus</i> , <i>Echeneis</i>	449	<i>raleighianus</i> , <i>Squalus</i>	20
<i>quenseli</i> , <i>Pleuronectes</i>	228	<i>Raja</i>	24
<i>quericum</i> , <i>Macrostoma</i>	160	<i>americana</i>	25
<i>querinus</i> , <i>Notoscopelus</i>	166	<i>aquila</i>	31
<i>Querimana</i>	253	<i>birostris</i>	32
<i>curema</i>	253	<i>chanteray</i>	25
<i>gaimardianus</i>	254	<i>cooperi</i>	26
<i>gyrans</i>	253	<i>desmarestia</i>	25
<i>hospes</i>	254	<i>diaphanes</i>	25
<i>incilis</i>	253	<i>eglantaria</i>	25
<i>setosus</i>	254	<i>erinacea</i>	25
<i>thoburni</i>	253	<i>fimbriata</i>	32
<i>trichodon</i>	254	<i>flagellum</i>	30
<i>querina</i> , <i>Azevha</i>	221	<i>granulata</i>	25
<i>Cyclopsetta</i>	221	<i>guttata</i>	28
<i>quiebra</i> , <i>Lichia</i>	277	<i>inornata inermis</i>	26
<i>quiescens</i> , <i>Pedichthys</i>	294	<i>interrupta</i>	26
<i>Prionotus</i>	408	<i>jordani</i>	26
<i>Uranidea</i>	385	<i>maculata</i>	29
<i>Quietula</i>	444	<i>manatia</i>	32
<i>y-cauda</i>	444	<i>micrura</i>	29
<i>quinctifer</i> , <i>Crotolopsis</i>	88	<i>narinari</i>	30
<i>quinnat</i> , <i>Salmo</i>	55	<i>obtusa</i>	26
<i>quinquarius</i> , <i>Pentanemus</i>	256	<i>ocellata</i>	25
<i>Poly nemus</i>	256	<i>quadriloba</i>	31
<i>quinquecinctus</i> , <i>Pomacanthus</i>	360	<i>radiata</i>	25
<i>quinquefasciata</i> , <i>Promicrops</i>	312	<i>rosispinis</i>	26
<i>quinquefasciatus</i> , <i>Serranus</i>	312	<i>scabrata</i>	25
<i>quinquelineatum</i> , <i>Hæmulon</i>	331	<i>Rall</i> , <i>Brama</i>	265
<i>quitzeoensis</i> , <i>Platyceclus</i>	182	<i>Sparus</i>	265
<i>Zoogeneticus</i>	182	<i>Rai zero</i>	327
<i>quoyi</i> , <i>Cestracion</i>	10,	<i>aratus</i>	327
<i>Rabdophorus</i>	359	<i>Raja</i>	24
<i>Rabida</i>	155	<i>abyssicola</i>	26
<i>eleuthera</i>	158	<i>ackleyi</i>	25
<i>exilis</i>	158	<i>aleutica</i>	26
<i>funebris</i>	158	<i>altavela</i>	29
<i>furiosa</i>	156	<i>binoculata</i>	28
<i>gilberti</i>	156	<i>centroura</i>	28
<i>insignis</i>	156	<i>diaphanes</i>	25
<i>leptacantha</i>	155	<i>eglantaria</i>	25
<i>miura</i>	158	<i>equatorialis</i>	26
<i>nocturna</i>	158	<i>erinacea</i>	25
<i>Rabirubia</i>	327	<i>fullae</i>	25
<i>genizara</i>	426	<i>giorna</i>	31
<i>inermis</i>	327	<i>granulata</i>	26
<i>rabirubia</i> , <i>Anthias</i>	327	<i>inornata</i>	26
<i>Rabu</i>	91, 93	<i>interrupta</i>	26
<i>davisi</i>	91	<i>jamaicensis</i>	29
<i>longicauda</i>	91	<i>levius</i>	25
<i>marmorea</i>	91	<i>microtrachys</i>	25
<i>panamensis</i>	91	<i>mobular</i>	31
<i>rachovii</i> , <i>Heterophallus</i>	184	<i>montereyensis</i>	26
<i>Rachycentridæ</i>	281	<i>ornata</i>	26
<i>Rachycentron</i>	281	<i>parnifera</i>	26
<i>canadus</i>	281	<i>pastinaca</i>	28
<i>Radcliffeiella</i>	446	<i>parcellens</i>	23
<i>radiale</i> , <i>Diplectrum</i>	318	<i>plutonia</i>	26
<i>radials</i> , <i>Serranus</i>	318	<i>radiata</i>	25
<i>radians</i> , <i>Scarus</i>	432	<i>rhina</i>	26
<i>Serranus</i>	318	<i>rosispinus</i>	26
<i>Sparisoma</i>	432	<i>say</i>	26
<i>radiata</i> , <i>Iridio</i>	420	<i>scabrata</i>	25
<i>Raja</i>	25	<i>senta</i>	25
<i>Rajæ</i>	25	<i>stabulliforis</i>	25
<i>radiato</i> , <i>Turdus oculo</i>	426	<i>stellulata</i>	26
<i>radiatus</i> , <i>Halosaurus</i>	74	<i>trachura</i>	26
<i>Labrus</i>	426	<i>tuberulata</i>	28
		<i>Rajidae</i>	24

List of scientific names—Continued

	Page		Page
raja, <i>Squatina</i>	23	Reinhardtius.....	221
raleighana, <i>Harriotta</i>	33	hippoglossoides.....	221
rammelsbergi, <i>Mugil</i>	253	remifer, <i>Ispisithus</i>	353
ramosum, <i>Pristipoma</i>	334	remiger, <i>Myctophum</i>	163
ramosus, <i>Rhombiscus</i>	334	remigera, <i>Bothrocara</i>	477
Ramularia.....	224	Remigelia.....	449
dendrica.....	224	australis.....	449
ranchovii, <i>Heterophalalus</i>	184	Remora.....	449
helleri.....	193	remora.....	449
ranula, <i>Liparis</i>	402	remora, <i>Echeneis</i>	449
ranulus, <i>Careproctus</i>	402	Remora.....	449
Ranzania.....	504	albescens.....	449
makua.....	504	remoroidea, <i>Echeneis</i>	449
truncata.....	504	Remoropsis.....	449
raphidoma, <i>Belone</i>	198	brachyptera.....	449
<i>Tylosurus</i>	198	remotus, <i>Carcharias</i>	16
Raro.....	95	Serranus.....	312
arus, <i>Rhinoscopelus</i>	168	Reniceps.....	17
<i>Scopelus</i>	168	tiburo.....	17
rascacio, <i>Scorpaena</i>	370	repandus, <i>Serranus</i>	315
rashleighanus, <i>Squalus</i>	20	reticularis, <i>Antennarius</i>	505
rastralis, <i>Anchoviella</i>	50	reticulata, <i>Amia</i>	33
<i>Stolephorus</i>	50	Poecilia.....	191
rastrelliger, <i>Physiculus</i>	212	Spatularia.....	36
<i>Pteropodus</i>	369	Thalassophryne.....	488
<i>Sebastichthys</i>	369	reticulatus, <i>Brytius</i>	299
Rastrinus.....	378	Catostomus.....	105
scutiger.....	378	Diodon.....	502
Rataboura.....	89	Erision.....	355
Rathbunella.....	452	Esox.....	173
<i>alleni</i>	452	Haliichthys.....	507
<i>hypoplecta</i>	452	Lebistes.....	191
rathbuni, <i>Fundulus</i>	179	Lepidogaster.....	488
<i>Muloides</i>	344	Lycodes.....	474
<i>Polycirrus</i>	351	Monochir.....	229
<i>Pontinus</i>	372	Notropis.....	133
<i>Upeneus</i>	344	Otolithus.....	355
<i>Zonoscion</i>	351	Pseudoxiphophorus.....	187
<i>Zyngonectes</i>	179	Trisotrops.....	315
rauchi, <i>Acipenser</i> (<i>Huso</i>).....	35	retifer, <i>Scyllorhinus</i>	11
<i>Ichthyobus</i>	102	retifera, <i>Muraena</i>	90
raucus, <i>Sargus</i>	338	retiferum, <i>Scyllium</i>	11
rava, <i>Pteroplatea</i>	29	retractus, <i>Catodon</i>	431
raveneli, <i>Esox</i>	173	<i>Cryptotomus</i>	431
<i>Pomotis</i>	302	retrocurrens, <i>Hæmulon</i>	329
recalva, <i>Monterey</i>	391	retropinnis, <i>Catostomus</i>	107
recalvus, <i>Blenniottus</i>	391	Microdesmus.....	473
rectangulare, <i>Acara</i>	421	Ophichthus.....	87
rectangularis, <i>Astatheros</i>	421	Ophichthys.....	87
rectifrenum, <i>Eupomacentrus</i>	414	rotrosella, <i>Amia</i>	295
<i>Pomacentrus</i>	414	<i>Apogon</i>	295
recurvirostrata, <i>Sayris</i>	197	retrospinis, <i>Holocentrum</i>	236
reddingi, <i>Orthopristis</i>	335	<i>Plectrypops</i>	236
redi, <i>Orthagoriscus</i>	503	retzli, <i>Orthagoriscus</i>	503
refugiens, <i>Oligoplites</i>	278	Catostomus.....	103
regale, <i>Myctophum</i>	168	Etheostoma.....	233
<i>Nannobrachium</i>	168	Percina.....	283
regalis, <i>Cynoscion</i>	355	rex-mullorum, <i>Apogon</i>	235
<i>Enchelyopus</i>	213	rex-salmonorum, <i>Trachipterus</i>	216
<i>Johnius</i>	355	<i>Trachypterus</i>	216
<i>Salmo</i>	56	rhabdophora, <i>Brachyrhaphis</i>	188
<i>Scomber</i>	257	Gambusia.....	186
<i>Scomberomorus</i>	257	Rhacochilus.....	411
regani, <i>Chiostoma</i>	250	toxotes.....	411
<i>Gambusia</i>	186	Rhamdalla.....	156
<i>Rhamdia</i>	188	Rhamdia.....	156
regis, <i>Atherinops</i>	252	<i>barbata</i>	158
<i>Colpichthys</i>	252	<i>baroni-mulleri</i>	157
regius, <i>Bleennius</i>	213	<i>brachycephala</i>	157
<i>Hybognathus</i>	143	<i>bransfordi</i>	157
<i>Lampris</i>	216	<i>cabrerae</i>	157
<i>Ophichthius</i>	87	<i>depressa</i>	157
<i>Urophycis</i>	213	<i>godmani</i>	157
<i>Zeus</i>	216	<i>guatemalensis</i>	157
regulus, <i>Fsenes</i>	288	<i>heteracantha</i>	157
<i>Sebastes</i>	363	<i>laticauda</i>	157
reichardi, <i>Leucichthys</i>	63	<i>managuensis</i>	157
Reichardina.....	9	<i>microptera</i>	157
<i>unicolor</i>	9	<i>motaguensis</i>	157
reinhardti, <i>Careproctus</i>	402	<i>nicaraguensis</i>	157
<i>Liparis</i>	402	<i>oaxacea</i>	157
reinhardtii, <i>Motella</i>	208	<i>parryi</i>	158
reinhardtii, <i>Corynophorus</i>	509	<i>petenensis</i>	158
<i>Himantolophus</i>	509	<i>polycaula</i>	158

List of scientific names—Continued

	Page		Page
<i>Rhamdia regani</i>	158	<i>Rhinogobius</i>	439
<i>rogersi</i>	158	<i>boleosoma</i>	439
<i>sacrificii</i>	158	<i>claytoni</i>	439
<i>salvinii</i>	157	<i>curacao</i>	439
<i>underwoodi</i>	158	<i>eigenmanni</i>	439
<i>wagneri</i>	157	<i>encaecumus</i>	439
<i>Rhamphichthys brevirostris</i>	101	<i>garmani</i>	439
<i>Rhamphoberyx</i>	236	<i>glaucofrenum</i>	439
<i>leucopus</i>	235	<i>mangilcola</i>	439
<i>pselopodus</i>	235	<i>mowbrayi</i>	439
<i>Rhamphocottidae</i>	393	<i>parvus</i>	439
<i>Rhamphocottus</i>	393	<i>quadriporus</i>	439
<i>richardsoni</i>	393	<i>shufeldti</i>	439
<i>Rhegma</i>	322	<i>similis</i>	439
<i>thamnaustum</i>	322	<i>stigmaturus</i>	439
<i>Rhegmatis</i>	322	<i>tortuga</i>	439
<i>Rhegnopteri</i>	256	<i>Rhinoliparis</i>	405
<i>Rhenus</i>	333	<i>attenaueri</i>	405
<i>panamensis</i>	334	<i>barbulifer</i>	405
<i>Rheocrypta</i>	285	<i>Rhinonemus</i>	209
<i>copelandi</i>	285	<i>bonasus</i>	31
<i>Rhessodon</i> , <i>Arbaciosa</i>	490	<i>encephalæ</i>	31
<i>Gobiesox</i>	490	<i>lalandi</i>	31
<i>Rhina</i>	22	<i>steinbachi</i>	31
<i>rhina</i> , <i>Antimora</i>	211	<i>vespertilio</i>	31
<i>Raja</i>	26	<i>Rhinopteridae</i>	31
<i>Rhincodon</i>	18	<i>Rhinoscion</i>	348
<i>typicus</i>	18	<i>saturnus</i>	348
<i>Rhineodon</i>	18	<i>Rhinoscopelus</i>	168
<i>typicus</i>	18	<i>andreae</i>	168
<i>Rhineodontidae</i>	18	<i>coccoi</i>	169
<i>Rhinesomus</i>	496	<i>oceanicus</i>	169
<i>triqueter</i>	496	<i>rarus</i>	168
<i>Rhinichthys</i>	139	<i>Rhinotriaci</i>	13
(<i>Apococe</i>) <i>nevadensis</i>	141	<i>henlei</i>	14
<i>velifer</i>	141	<i>rhizophora</i> , <i>Fundulus</i>	175
<i>arenatus</i>	140	<i>rhodochloris</i> , <i>Sebastichthys</i>	367
<i>stronsassus</i>	140	<i>Sebastomus</i>	367
<i>badius</i>	140	<i>rhodopus</i> , <i>Trachinotus</i>	276
<i>bowersi</i>	139	<i>rhodus</i> , <i>Ascelichthys</i>	392
<i>cataractæ</i>	139	<i>rhodospilus</i> , <i>Gobiesox</i>	489
<i>crocus</i>	140	<i>rhodoterus</i> , <i>Holocotus</i>	410
<i>dulcis</i>	139	<i>Rhodymenichthys</i>	468
<i>henshawi</i>	141	<i>dolichogaster</i>	468
<i>lunatus</i>	140	<i>rhomalea</i> , <i>Gnathopops</i>	452
<i>luteus</i>	139	<i>rhomaleus</i> , <i>Olistognathus</i>	453
<i>marmoratus</i>	139	<i>Squalius</i>	119
<i>maxillosus</i>	139	<i>rhombaea</i> , <i>Moharra</i>	342
<i>meleagris</i>	140	<i>rhombaea</i> , <i>Gerres</i>	342
<i>obtusus</i>	140	<i>Rhombochirius</i>	460
<i>ocellus</i>	139	<i>osteochir</i>	460
<i>simus</i>	140	<i>rhomboidalis</i> , <i>Lebias</i>	181
<i>transmontanus</i>	139	<i>Otolithus</i>	354
<i>Rhinichthyoides</i> , <i>Tigoma</i>	141	<i>Perca</i>	338
<i>Rhinobatidae</i>	23	<i>Salema</i>	338
<i>Rhinobatos</i>	23	<i>rhomboides</i> , <i>Chetodon</i>	277
<i>glaucostigma</i>	24	<i>rhomboides</i> , <i>Hemicarax</i>	271
<i>forkelli</i>	23	<i>Lagodon</i>	337
<i>leptiginosus</i>	23	<i>Sparus</i>	337
<i>leucorhynchus</i>	24	<i>Trachinotus</i>	227
<i>percellens</i>	23	<i>Rhomboidichthys</i>	217
<i>productus</i>	24	<i>leopardinus</i>	218
<i>spinulosus</i>	24	<i>spiniferus</i>	217
<i>stellio</i>	23	<i>Rhomboplites</i>	327
<i>Rhinobatus</i>	23	<i>aurorubens</i>	328
<i>columna</i>	23	<i>Rhombotides</i>	362
<i>electricus</i>	24	<i>Rhombus</i>	266
<i>glaucoctonus</i>	23	<i>bahianus</i>	217
<i>glaucoctigna</i>	23	<i>levis cornubiensis</i>	228
<i>horkei</i>	23	<i>ocellatus</i>	217
<i>leptiginosus</i>	23	<i>polypilus</i>	223
<i>leucorhynchus</i>	24	<i>soleiformis</i>	218
<i>margravii</i>	23	<i>Rhondiscus</i>	334
<i>producta</i>	24	<i>braniicki</i>	334
<i>rhinobatus</i>	23	<i>croco</i>	334
<i>seu</i>	23	<i>ramosus</i>	334
<i>spinulosus</i>	24	<i>sinuosus</i>	336
<i>stellio</i>	23	<i>rhothea</i> , <i>Cottus</i>	363
<i>undulatus</i>	23	<i>Uranidea</i>	383
<i>Rhinoberyx</i>	235	<i>rhotheus</i> , <i>Leucosomus</i>	117
<i>chrysseus</i>	235	<i>Rhotheca</i>	289
<i>Rhinocheimeridae</i>	33	(<i>Rhotheca</i>) <i>biennius</i> , <i>Etheostoma</i>	257
<i>Rhinocymnus</i>	21	<i>rhotheca</i> , <i>Thoburnia</i>	108
<i>Rhinodon</i>	18	<i>rhothocetus</i> , <i>Castoromus</i>	108
<i>Rhinogobius</i>	440	<i>rhynchosus</i> , <i>Acipenser</i>	35
<i>nichosi</i>	440		

List of scientific names—Continued

	Page		Page
Rhynchias	451	Rivulus hartl.	180
septipinnis	451	heyei	180
Rhynchichthys	235	isthmensis	180
brachyrhynchus	235	marmoratus	179
Rhynchoeratias	511	tenuis	180
bravirostris	511	roanoka, <i>Alvordius</i>	284
leucorhinus	511	<i>Hadropterus</i>	284
rostratus	511	robalo, <i>Centropomus</i>	306
Rhrypticus	318	Robalo	306
maculatus	317	<i>prieto</i>	306
nigripinnis	317	roberti, <i>Exocetus</i>	201
nigromaculatus	316	<i>Hemirhamphus</i>	198
pituitosus	316	<i>Hyporhamphus</i>	198
subbifrenatus	316	robertsi, <i>Anchoviella</i>	49
xanti	316	<i>Ceratobatis</i>	32
Rhytidostomus	102	<i>Siphostoma</i>	241
rali, <i>Onus</i>	214	<i>Stolephorus</i>	49
ricardi, <i>Mesopion</i>	327	<i>Syngnathus</i>	241
rivel, <i>Cottopsis</i>	386	robertsoni, <i>Astatheros</i>	420
<i>Cottus</i>	386	<i>Cichlasoma</i>	420
richardi, <i>Caranx</i>	272	robini, <i>Hypostomus</i>	160
<i>Hemirhamphus</i>	198	robusta, <i>Agosia</i>	140
Salmo	55	<i>Apocope</i>	140
richardsoni, <i>Acipenser (Huso)</i>	35	Gila	114
<i>Astronesthes</i>	71	<i>Hermosilla</i>	340
<i>Catostomus</i>	103	<i>Moniana</i>	132
<i>Chauliodus</i>	71	<i>Perca</i>	311
<i>Corvina</i>	352	robustum, <i>Megalonema</i>	159
<i>Cottus</i>	385	<i>Moxostoma</i>	111
<i>Hybopsis heterodon</i>	136	robustus, <i>Benthocometes</i>	482
<i>Rhamphocottus</i>	393	<i>Clypeocottus</i>	381
<i>Trachidermis</i>	383	<i>Hippoglossoides</i>	222
richardsonii, <i>Coregonus</i>	64	<i>Ictalurus</i>	152
Richardsonius	117	<i>Mylopharodon</i>	113
<i>balteatus</i>	118	<i>Neobythites</i>	482
<i>lateralis</i>	118	<i>Notropis</i>	132
<i>microdon</i>	118	<i>Ptychostomus</i>	111
<i>susiawli</i>	118	<i>Zoogeneticus</i>	182
<i>thermophilus</i>	118	Roccus	307
Richia	293	<i>comes</i>	355
<i>brevispina</i>	293	<i>saxitilis</i>	307
Richiella	293	<i>striatus</i>	307
<i>rictolata, Bothrocara (sis-</i>	477	rochei, <i>Auxis</i>	259
<i>rictus, Xenomystax</i>	81	rodo, <i>Pristipoma</i>	333
riddlei, <i>Toxus</i>	188	Rœboïdes	98
rigidus, <i>Pimelodus</i>	159	<i>guatamalensis</i>	98
rijersmei, <i>Ocyurus</i>	327	<i>macrolepis</i>	95
rim, <i>Scomber</i>	273	<i>occidentalis</i>	98
rimator, <i>Bathystoma</i>	320, 331	<i>salvadoris</i>	98
<i>Hæmulon</i>	331	Rogenia	41
rimensis, <i>Ruscarius</i>	391	<i>alba</i>	41
Rimicola	490	rogersi, <i>Pimelodus</i>	158
<i>beryllinus</i>	490	<i>Rhamdia</i>	158
<i>egenmanni</i>	490	<i>Uropholus</i>	30
(Rimicola) <i>beryllinus</i> , <i>Go-fishox</i>	490	Roncador	348
rimosus, <i>Etropus</i>	220	<i>stearnsi</i>	348
Rineloricaria	162	roncador, <i>Umbrina</i>	350
<i>lima</i>	162	ronchus, <i>Bairdiella</i>	346
ringens, <i>Atopoclinus</i>	465	<i>Corvina</i>	346
<i>Balistes</i>	493	rondeleti, <i>Anthias quartus</i>	324
<i>Phenodon</i>	71	<i>Carcharodon</i>	20
<i>Sudis</i>	165	Rondeletia	171
<i>Xanthichthys</i>	493	<i>bicolor</i>	171
riparium, <i>Holocentrum</i>	236	Rondeletidae	171
risso, <i>Alpismaris</i>	163	<i>rondeletti, Exocetus</i>	200
rossianus, <i>Notacanthus</i>	76	Exonautae	200
Rissola	485	<i>Orthagorisicus</i>	503
<i>marginata</i>	485	Sargus	338
ritteri, <i>Lampauptyctus</i>	167	<i>Scombras</i>	197
<i>Pleuronichthys</i>	225	<i>Xiphias</i>	204
<i>Xeneretmus</i>	306	Ronquilus	452
<i>Xenopryx</i>	306	<i>jordanii</i>	452
rivalis, <i>Salmo</i>	60	roosevelti, <i>Salmo</i>	59
<i>riverendi, Cyprinodon</i>	181	rosacea, <i>Myctoperca</i>	314
<i>Trifasciatus</i>	181	rosaceus, <i>Cymatogaster</i>	410
rivoliana, <i>Seriola</i>	280	Epinephelus	314
rivolianus, <i>Zonichthys</i>	280	<i>Mesopion</i>	328
rivularis, <i>Salmo</i>	58	<i>Paraliparis</i>	404
rivulatus, <i>Cirrhites</i>	368	Sebastes	367
<i>Diodon</i>	502	<i>Sebastomus</i>	367
Rivulus	179	<i>Zalembius</i>	410
<i>brunneus</i>	180	rosea, <i>Clevelandia</i>	446
<i>chucunaquea succubta</i>	180	<i>Hemirhamphus</i>	198
<i>chucunaquea</i>	180	<i>Hyporhamphus</i>	198
<i>cylindraceus</i>	179	Troglichthys	195
<i>fibellicauda</i>	180	<i>rosarium, Acipenser (Huso)</i>	36
godmanni	180		

List of scientific names—Continued

	Page		Page
rosei, <i>Salmo</i>	59	rubrifrons, <i>Alburnus</i>	124
roseipinnis, <i>Lythurus</i>	126	Erinemus.....	138
<i>Notropis</i>	128	Noconis.....	138
roseus, <i>Cryptotomus</i>	431	Notropis.....	124, 128
<i>Hydrophox</i>	127	Zygonectes.....	177
Luxilus.....	127	rubrilateralis, <i>Coneusus plumbeus</i>	116
Merulinus.....	407	rubripinna, <i>Cyprinella</i>	131
Prionotus.....	407	rubripinna, <i>Sparisoma</i>	433
Rosicola alexandri.....	365	rubripinnis, <i>Argyreus</i>	128
<i>babcocki</i>	365	Scarus.....	433
<i>miniatu</i> s.....	365	rubrirostris, <i>Catetix</i>	480
<i>pinniger</i>	365	rubrivinctus, <i>Hispanicus</i>	368
rosipes, <i>Novaculichthys</i>	430	Sebastichthys.....	368
<i>Xyrichtys</i>	430	rubrofuvatilis, <i>Cyprinodon</i>	181
rosiplinis, <i>Rala</i>	26	<i>Cyprinodon bovinus</i>	181
<i>Raja</i>	28	rubropunctatus, <i>Salaris</i>	463
rossii, <i>Salmo</i>	60	rubrum, <i>Choristium</i>	315
Salvelinus.....	60	rudis, <i>Centrochromis</i>	415
rostratus, <i>Albula</i>	40	<i>Glyphidodon</i>	415
<i>Aldrovandia</i>	75	rufa, <i>Morone</i>	307
<i>Anguilla</i>	77	Perca.....	236
<i>Loricaria</i>	161, 162	rufilineatus, <i>Nothonotus</i>	290
<i>Muraena</i>	77	<i>Poecilichthys</i>	290
<i>Platessa</i>	227	rufipinnis, <i>Exocetus</i>	201
<i>Sturisoma</i>	161	<i>Exonutes</i>	201
rostratum, <i>Astatheros</i>	421	rufum, <i>Acutomentum</i>	366
<i>Holocentrum</i>	236	rufus, <i>Balistes</i>	493
rostratus, <i>Apterontotus</i>	101	<i>Bodianus</i>	425
<i>Canthigaster</i>	600	<i>Centropristes</i>	318
<i>Gymnothorax</i>	91	<i>Holocentrus</i>	236
<i>Halosaurus</i>	75	<i>Labrus</i>	425
<i>Heros</i>	421	<i>Onos</i>	208
<i>Heterostichus</i>	458	<i>Sebastodes</i>	366
<i>Holocentrus</i>	235	rugifer, <i>Ophichthus</i>	88
<i>Notacanthus</i>	76	rugosus, <i>Platichthys</i>	226
<i>Phycis</i>	213	rupertianus, <i>Acipenser</i>	35
<i>Polyacanthonotus</i>	76	rupestre, <i>Etheostoma</i> (<i>Niphodexa</i>).....	290
<i>Rhynchoseratus</i>	511	<i>Nanostoma</i>	290
<i>Scarus</i>	435	rupestris, <i>Ambloplites</i>	308
<i>Squalus</i>	19	<i>Arbacioides</i>	491
<i>Stenarchus</i>	101	<i>Bodianus</i>	303
<i>Tetronodon</i>	500	<i>Coryphena</i>	205
<i>Zeus</i>	275	<i>Coryphaenoides</i>	205
rotengulus, <i>Leuciscus</i>	117	<i>Gadus</i>	210
rothrocki, <i>Notogrammus</i>	469	<i>Gobiesox</i>	491
<i>Poroclinus</i>	470	<i>Sebastichthys</i>	368
rousseau, <i>Syngnathus</i>	241	<i>Sebastomus</i>	368
rubello, <i>Eximia</i>	391	<i>Serranus</i>	313
<i>Greeleyi</i>	391	<i>Xiphister</i>	469
rubellus, <i>Alburnus</i>	123	Rupisartes.....	463
rubens, <i>Tautoga</i>	424	<i>atlanticus</i>	463
rubber, <i>Apogon</i>	295	rupisartes, <i>Moxostoma</i>	111
<i>Cephalopholis fulvus</i>	309	<i>Scartomyzon</i>	111
<i>Coregonus</i>	143	ruppell, <i>Monacanthus</i>	494
<i>Elaphotroxon</i>	271	ruptus, <i>Julis</i>	427
<i>Epinephelus</i>	315	Rusciops.....	378
<i>Gadus</i>	214	<i>creaseri</i>	378
<i>Gymnocephalus</i>	309	Rusciarius.....	378
<i>parvus</i> , <i>Sebastes</i>	368	<i>meanyi</i>	378
<i>Rutilus</i>	126	Ruscarius, <i>Gurnardius</i>	406
<i>Scomber</i>	271	<i>Prionotus</i>	406
<i>Sebastes</i>	368	Ruscarius.....	391
ruberrimus, <i>Astyanax</i>	97	<i>rimensis</i>	391
<i>Gunnellus</i>	408	Rusulus.....	390
<i>Sebastodes</i>	368	<i>saburrae</i>	390
<i>Sebastopyr</i>	308	russula atlantica, <i>Scorpaena</i>	270
rubescens, <i>Algancea</i>	116	russula, <i>Scorpaena</i>	371
<i>Auchenopterus</i>	457	ruthenus, <i>Acipenser</i>	34
<i>Conger</i>	79	rutila, <i>Monlana</i>	131
<i>Exocetus</i>	201	Rutilus amblopolis.....	123
<i>Exonutes</i>	201	<i>anomalus</i>	146
<i>Steinegeria</i>	286	<i>columbianus</i>	122
rubicundus, <i>Acipenser</i>	35	<i>compressus</i>	123, 128
<i>Glyptodon</i>	415	<i>melanurus</i>	110
<i>Hypsypops</i>	415	<i>oregonensis</i>	122
rubiginosus, <i>Sicyas</i>	490	<i>plagyrurus</i>	128
Rublo volador.....	407	<i>ruber</i>	126
rubio, <i>Prionotus</i>	407	<i>storerasianus</i>	137
rubra, <i>Garninia</i>	445	Rutter, <i>Neoliparis</i>	399
<i>Lipopomona</i>	316	Ruvettus.....	261
<i>Sciensa</i>	235	<i>pretiosus</i>	261
<i>Liopomona</i>	315	<i>temminckii</i>	261
rubri-cauda, <i>Pomotis</i>	299	Rypticus.....	316
rubricleucus, <i>Hybopsis</i>	126, 127	<i>arenatus</i>	316
<i>Hydrophox</i>	127		

List of scientific names—Continued

	Page		Page
<i>Rypticus bicolor</i>	316	<i>Salmo canadensis</i>	60
<i>corlaceus</i>	316	<i>canis</i>	54
<i>maculatus</i>	316	<i>callaris</i>	61
<i>saponaceus</i>	316	<i>campbelli</i>	61
<i>xanti</i>	316	<i>canadensis</i>	60
<i>sabalo</i> , <i>Arges</i>	159	<i>canis</i>	54
<i>sabanensis</i> , <i>Hippoglossina</i>	223	<i>carlinus</i>	56
<i>sabine</i> , <i>Trygon</i>	29	<i>clarkii</i>	56
<i>sabine</i> , <i>Hybopsis</i>	135	<i>declivifrons</i>	57
<i>Notropis</i>	136	<i>jordani</i>	57
<i>sabinus</i> , <i>Amphotistius</i>	29	<i>tahoensis</i>	57
<i>saburra</i> , <i>Chasmodes</i>	462	<i>clupeaformis</i>	64
<i>Rusulus</i>	390	<i>confins</i>	59
<i>Saccopharyngidae</i>	95	<i>confuentus</i>	65
<i>Saccopharynx</i>	95	<i>consuetus</i>	54
<i>ampullaceus</i>	95	<i>cooperi</i>	55
<i>chordatus</i>	95	(<i>Coregonus</i>) <i>barengus</i>	62
<i>flagellum</i>	95	<i>lucidus</i>	62
<i>saccostoma</i> , <i>Lasiongathus</i>	510	<i>sapidissimus</i>	64
<i>sacrifici</i> , <i>Rhamdia</i>	158	<i>tullibee</i>	64
<i>sadina</i> , <i>Clupes</i>	46	<i>crescentis</i>	57
<i>Etrumeus</i>	46	<i>curlis</i>	61
<i>Safole</i>	305	<i>declivifrons</i>	57
<i>teniura</i>	305	<i>dermatinus</i>	64
<i>sagamius</i> , <i>Ebisus</i>	373	<i>edentulus</i>	100
<i>Sagenichthys</i>	354	<i>esperlanus</i>	67
<i>mordax</i>	354	<i>eremogenes</i>	68
<i>sagitta</i> , <i>Amblyopus</i>	448	<i>erythrorhynchos</i>	61
<i>Poecilichthys</i>	293	<i>evermanni</i>	59
<i>Torrentaria</i>	293	<i>fario</i>	55
<i>Tylosurus</i>	195	<i>fontens</i>	163
<i>Tyntlasses</i>	448	<i>fontinalis</i>	60
<i>sagittula</i> , <i>Euctenogobius</i>	441	<i>gairdneri</i> beardsleyi	57
<i>Gobionelius</i>	441	<i>crescentis</i>	57
<i>Bagrarcas</i>	235	<i>gilberti</i>	58
<i>salida</i> , <i>Boreogadus</i>	209	<i>shasta</i>	58
<i>Gadus</i>	209	<i>stonei</i>	58
<i>saira</i> , <i>Cololabis</i>	198	<i>gairdnerii</i>	58
<i>Salar</i>	55	<i>gibber</i>	64
<i>lewisi</i>	56	<i>gibbsii</i>	57
<i>virginalis</i>	57	<i>gilberti</i>	58
<i>salar</i> , <i>Salmo</i>	55, 58	<i>gloveri</i>	56
<i>Salaria</i>	460	<i>gorbuscha</i>	54
<i>Salarias alticus</i>	463	<i>grönlandicus</i>	66
<i>atlanticus</i>	463	<i>harengus</i>	61
<i>chllostictus</i>	463	<i>hearni</i>	60
<i>margaritaceus</i>	463	<i>henshawi</i>	57
<i>rubro-punctatus</i>	463	<i>hudsonicus</i>	60
<i>salarias rubropunctatus</i>	463	<i>immaculatus</i>	60
<i>textilis</i>	463	<i>irideus</i>	58
<i>vomerinus</i>	463	<i>iridia</i>	58
<i>salarias rubropunctatus</i> , <i>Salarias</i>	463	<i>japonicus</i>	64
<i>Salarichthys</i>	463	<i>jordanii</i>	57
<i>textilis</i>	463	<i>kennerlyi</i>	54, 55
<i>Salema surata</i>	339	<i>keta</i> vef <i>kayko</i>	54
<i>Salema</i>	337	<i>kisutch</i>	54, 55
<i>rhomboidalis</i>	338	<i>levigatus</i>	61
<i>tridens</i>	338	<i>lagocephalus</i>	54
<i>sallens</i> , <i>Blennius</i>	463	<i>latior</i>	64
<i>Oilgapilites</i>	277	<i>lavaretus</i>	64
<i>Scomber</i>	277	<i>lewisi</i>	56
<i>Thunnus</i>	269	<i>lordil</i>	61
<i>salm</i> , <i>Sparus</i>	338	<i>lycaodon</i>	55
<i>salla</i> , <i>Atherin'chthys</i>	248	<i>macleodii</i>	57
<i>salla</i> , <i>Algassea</i>	116	<i>mackenzii</i>	61
<i>Cerat'chthys</i>	116	(<i>Mallotus</i>) <i>pacificus</i>	66
<i>salla</i> , <i>Archomenidia</i>	248	<i>malma</i>	61
<i>Atherinichthys</i>	248	<i>marcgravii</i>	100
<i>Salmo</i>	55	<i>marstoni</i>	61
<i>adirondacus</i>	59	<i>mas</i>	56
<i>agassizii</i>	60	<i>masoni</i>	63
<i>aqua-bonita</i>	59	<i>melanopterus</i>	55
<i>alipes</i>	60	<i>miltkischitch</i>	54
<i>alleghanianus</i>	60	<i>miltschutsch</i>	54
<i>amethystinus</i>	59	<i>miltschitsch</i>	64
<i>ansonii</i>	55	<i>müllerii</i>	170
<i>aqullarum</i>	58	<i>mykiss</i>	56
<i>arabatch</i>	56	<i>aguia-bonita</i>	59
<i>arcticus</i>	66	<i>macleodii</i>	57
<i>arcturus</i>	61	<i>myops</i>	163
<i>bairdii</i>	61	<i>namaycush</i>	60
<i>bathceator</i>	57	<i>narasi</i>	61
<i>beardsleyi</i>	57	<i>narks</i>	55
<i>bouvieri</i>	56	<i>nelsoni</i>	69
<i>breviceauda</i>	56	<i>nerka</i>	55
	56	<i>newberryi</i>	58

List of scientific names—Continued

	Page		Page
<i>Salmo nigrescens</i>	60	<i>saludanus</i> , <i>Alburnops</i>	132
<i>nitidus</i>	60	<i>Hudsonius</i>	132
<i>nummifer</i>	61	<i>salvadoris</i> , <i>Rohoides</i>	98
<i>omiscomaycus</i>	214, 215	<i>salvatoris</i> , <i>Pocilia</i>	102
<i>omul</i>	61	<i>Salvelinus</i>	60
<i>oquassa</i>	61	<i>alpinus</i>	60
<i>orientalis</i>	55	<i>Salvelinus</i>	60
(<i>Osmurus</i>) <i>olidus</i>	68	<i>agassizii</i>	60
<i>otsego</i>	64	<i>arcturus</i>	61
<i>ouananiche</i>	58	<i>aureolus</i>	60
<i>pallidus</i>	59	<i>fontinalis</i>	60
<i>parki</i>	61	<i>malma</i>	61
<i>paucidens</i>	55	<i>spectabilis</i>	61
<i>pepsinensis</i>	56	<i>marstoni</i>	61
<i>pleuriticus</i>	57	<i>naresi</i>	61
<i>pluvius</i>	61	<i>oquassa</i>	61
<i>proteus</i>	54	<i>rossi</i>	60
<i>purpuratus</i>	58	<i>spectabilis</i>	61
<i>bouvieri</i>	58	<i>stagnalis</i>	60
<i>quinnat</i>	55	<i>timagamensis</i>	60
<i>regalis</i>	56	<i>salvelinus</i> , <i>Salmo</i>	60
<i>richardii</i>	55	<i>salvinii</i> , <i>Agonostomus</i>	264
<i>rivals</i>	60	<i>Cotylopus</i>	448
<i>rivularis</i>	58	<i>Heros</i>	418
<i>roosevelti</i>	59	<i>Parapetenia</i>	418
<i>rosel</i>	59	<i>Pimelodus</i>	157
<i>rossi</i>	60	<i>Rhamdia</i>	157
<i>salar</i>	55, 56	<i>Sicydium</i>	448
<i>ouananiche</i>	56	<i>sancta-helenae</i> , <i>Caranx</i>	270
<i>salvelinus</i>	60	<i>Decapterus</i>	270
<i>sanguinolentus</i>	55	<i>Gymnothorax</i>	92
<i>saurus</i>	163	<i>Muraena</i>	92
<i>scouleri</i>	54	<i>sancta-luciae</i> , <i>Corvula</i>	245
<i>sebago</i>	56	<i>Vacuqua</i>	345
<i>shasta</i>	58	<i>sancta-martae</i> , <i>Vomer</i>	275
<i>siscowet</i>	60	<i>sancta-petri</i> , <i>Vomer</i>	275
<i>siskowitz</i>	60	<i>sancta-roseae</i> , <i>Ulvicola</i>	467
<i>smaragdus</i>	58	<i>sancti-eustatii</i> , <i>Callionymus</i>	460
<i>socialis</i>	66	<i>sancti-laurentii</i> , <i>Engyophrys</i>	220
<i>spectabilis</i>	61	<i>sancti-martini</i> , <i>Gobiesox</i>	490
<i>spilurus</i>	57	<i>sancti-pauli</i> , <i>Holocentrum</i>	236
<i>stagnalis</i>	60	<i>Holocentrus</i>	236
<i>stomias</i>	57	<i>Sander</i>	282
<i>stonei</i>	58	<i>sandwichensis</i> , <i>Gobius</i>	440
<i>striatus</i>	54	<i>Pimelepterus</i>	339
<i>symmetricus</i>	59	<i>sandwichensis</i> , <i>Ballistes</i>	493, 494
<i>taboensis</i>	57	<i>sanguinifluus</i> , <i>Pocilloichthys</i>	290
<i>tadpisma</i>	55	<i>sanguineus</i> , <i>Antennarius</i>	505
<i>thymallus</i>	66	<i>Apodichthys</i>	467
<i>toma</i>	59	<i>Holocentrus</i>	363
<i>truncatus</i>	58	<i>Pythonichthys</i>	90, 91
<i>trutta</i>	55	<i>Sicyas</i>	490
<i>tschawytscha</i>	56	<i>sanguinolentus</i> , <i>Pomotis</i>	300
<i>tschawytschiformis</i>	54	<i>Salmo</i>	55
<i>tsuppitch</i>	55	<i>santa-anae</i> , <i>Gasterosteus</i>	238
<i>tudes</i>	61	<i>santa-anae</i> , <i>Notolepidomyzon</i>	104
<i>tumbil</i>	164	<i>Pantosteus</i>	104
<i>umbla</i>	60	<i>santamariae</i> , <i>Monopterus</i>	131
<i>ursinus</i>	60	<i>Notropis</i>	131
<i>utah</i>	56	<i>sapidissima</i>	42
<i>virginialis</i>	57	<i>Clupea</i>	42
<i>warreni</i>	55	<i>sapidissimus</i> , <i>Salmo</i> (<i>Coregonus</i>)	64
<i>whitel</i>	59	<i>saponaceus</i> , <i>Anthias</i>	316
<i>salmoides</i> , <i>Labrus</i>	207, 298	<i>Rypticus</i>	316
<i>Micropterus</i>	297	<i>sara</i> , <i>Cyhlum</i>	256
<i>salmonæ</i> , <i>Ericara</i>	53, 54	<i>saragus</i> , <i>Lepodus</i>	285
<i>Lepomis</i>	298	<i>Saranus apilarus</i>	308
<i>Perca</i>	282	<i>Sarchirus</i>	36
<i>salmoreus</i> , <i>Esox</i>	163, 172	<i>argenteus</i>	36
<i>Mugil</i>	40	<i>vittatus</i>	36
<i>Scombrocotus</i>	373	<i>Sarcidium</i>	142
<i>salmonicolor</i> , <i>Merulinus</i>	407	<i>scopiferum</i>	142
<i>Salmonidae</i>	54	<i>Sarcura</i>	23
<i>Salmonoidæ</i>	54	<i>Sarda</i>	259
<i>Salmoperca</i>	214	<i>chilensis</i>	259
<i>pellucida</i>	215	<i>lineolata</i>	259
<i>Salmoperca</i>	214	<i>orientalis</i>	259
<i>Salpa purpurascens variegata</i>	327	<i>sarda</i>	259
<i>saltans</i> , <i>Chorinemus</i>	277	<i>velox</i>	259
<i>saltator</i> , <i>Hemiramphus</i>	199	<i>sarda</i> , <i>Sarda</i>	259
<i>Scomberoides</i>	277	<i>Scomber</i>	259
<i>saltatorius</i> , <i>Alticus</i>	463	<i>Sardina</i>	43
<i>saltatrix</i> , <i>Gasterosteus</i>	280	<i>pseudohispanica</i>	43
<i>Perca</i>	280	<i>sardina</i> , <i>Atherina</i>	247
<i>Pomatomus</i>	280	<i>Harengula</i>	43

List of scientific names—Continued

	Page		Page
sardina, <i>Melaniris</i>	249	saxicola, <i>Pleuronectes</i>	228
Menidia.....	247	Sebastichthys.....	367
Thyrsis.....	249	say, <i>Amphotistius</i>	28
Sardinella.....	43	Raja.....	28
anchovia.....	43	sayanus, <i>Aphredoderus</i>	215
aurita.....	43	Scolopsis.....	215
Sardinia.....	43	Sayris.....	197
anchovia.....	43	bimaculatus.....	197
cærulea.....	43	biannus.....	197
pseudohispanica.....	43	recurvirostra.....	197
sargoides, <i>Chetodon</i>	415	serata.....	197
Sargosomus.....	412	scaber, <i>Antennarius</i>	505
finiavittata.....	412	Centropomus.....	505
Sargus.....	338	Chironectes.....	505
ambassis.....	338	Hexagrammus.....	374
argenteus.....	338	scabre, <i>Echidna</i>	94
aries.....	338	Lophius acute.....	505
caribeus.....	338	Trinectes.....	229
caudimacula.....	338	scabratu, <i>Raija</i>	25
flavolineatus.....	338	Raja.....	25
holbrookii.....	338	scabriceps, <i>Cylindrostetus</i>	37
humeralmaculatus.....	338	Paranotropis.....	125
ovis.....	338	Photogamis.....	125
raucus.....	338	scabripinnis, <i>Astyanax</i>	97
rondeletii.....	338	Tetragonopterus.....	97
sargus.....	338	Scataria.....	264
tridens.....	338	Scaphirhynchops.....	35
variegatus.....	338	Scaphirhynchus.....	35
vittata.....	338	platorynchus.....	35
sargus, <i>Diplodus</i>	338	rafinesquel.....	35
Sargus.....	338	scaphilurus, <i>Opistognathus</i>	462
Sparus.....	338	scaphopsis, <i>Colorhynchus</i>	207
Sarothrodus.....	359	Macrurus (<i>Ceslorhynchus</i>).....	207
amplexicollis.....	360	Scaphirhynchus.....	35
atænotatus.....	360	scapulare, <i>Pristipoma</i>	332
maculocinctus.....	360	scapulare, <i>Anisotremus</i>	332
nigrostris.....	360	Strongylura.....	196
Sarrtor.....	365	Tylosurus.....	195
frenatus.....	365	Scaridae.....	433
leptorhynchus.....	365	Scarcina.....	262
sarrtor, <i>Prionotus</i>	407	argyrea.....	262
Satanoperca.....	424	Scartella.....	401
(Satanoperca) crassilabris, <i>Geophagus</i>	424	brevipinnis.....	461
saturicus, <i>Neoclinus</i>	458	microstoma.....	461
Pterognathus.....	458	Scartes.....	403
satis parva et glabra, <i>Pleuronectes</i>	217	scartes, <i>Fundulus</i>	177
saturnus, <i>Amblodon</i>	348	Scartichthys.....	463
Rhinoscion.....	348	rubropunctatus.....	463
Saurenelches.....	82	Scartomyzon.....	111
cancrivora.....	82	cervinus.....	111
Saurida.....	164	rupisartes.....	111
mediterranea.....	163	Scarus.....	432, 433
nebulosa.....	164	abildgardi.....	431
Saurus.....	163	acutus.....	434
anolis.....	164	alternans.....	434
argenteus.....	277	amplus.....	432
brevostris.....	163	aracanga.....	433, 434
griseus.....	163	atomarius.....	431
intermedius.....	164	autostrenatus.....	432
longirostris.....	163	bollmani.....	433
lucioceps.....	163	brachialis.....	433
mexicanus.....	163	cæruleus.....	434
myops.....	163	(Catesby).....	432
spixianus.....	163	chloris.....	432
truncatus.....	163	chrysopterus.....	432
Saurus, <i>Elops</i>	39	circumnotatus.....	433
Esox.....	197	coccineus.....	432
Oligoplites.....	277	celestinus.....	435
Salmo.....	163	compressus.....	434
Scomber.....	277	croicensis.....	433, 434
Scomberesox.....	197	cuzamilæ.....	434
Synodus.....	163	dentiens.....	431
seusurii, <i>Brama</i>	265	diadema.....	433
Taractes.....	265	distinctus.....	432
savanna, <i>Muraena</i>	78	emarginatum.....	433
Muraenesox.....	78	emblematicus.....	435
saxatilis, <i>Catulus</i>	11	erythrinoides.....	432
Chætodon.....	414	evermanni.....	434
Glyphtodon.....	415	flavescens.....	433
Johnius.....	350	flavomarginatus.....	434
Menticirrhus.....	350	flavo marginatus.....	434
Perca.....	307	frondosus.....	433
Poecilichthys.....	287	gibbus.....	433
Roccus.....	307	gnathodus.....	484
• sardola, <i>Eosebastes</i>	367	guacamala.....	435
Gymnothorax ocellatus.....	93	holocyaneos.....	434

List of scientific names—Continued

	Page		Page
<i>Scarus hoplostomus</i>	432	<i>Scolops ensifera</i>	345
<i>humeralis</i>	433	<i>ericymba</i>	346
<i>insulæ-santo-crucis</i>	434	<i>fusca</i>	352
<i>lacrimosus</i>	432	<i>gigas</i>	352
<i>lateralis</i>	432	<i>grisea</i>	346
<i>maschalepsilos</i>	433	<i>icistia</i>	346
<i>melanotis</i>	432	<i>imberbis</i>	348
<i>miniforenatus</i>	432	<i>imleeps</i>	348
<i>noyesi</i>	434	<i>lineata</i>	307
<i>nuchalis</i>	434	<i>maculata</i>	436
<i>obtusus</i>	434	<i>multifasciata</i>	349
<i>oxybrachius</i>	432	<i>oscitans</i>	347
<i>pepo</i>	433	<i>oscula</i>	352
<i>perrico</i>	435	<i>plumieri</i>	333
<i>pleianus</i>	435	<i>rubra</i>	236
<i>punctulatus</i>	433	<i>sciera</i>	348
<i>quadrispinosus</i>	434	<i>tahmeli</i>	339
<i>radians</i>	432	<i>undecimalis</i>	305
<i>rostratus</i>	435	Sciaenidae	98, 344
<i>rubripinnis</i>	433	Sciaenops	348
<i>spinidens</i>	431, 432	<i>ocellata</i>	348
<i>squalidus</i>	433	<i>sclera, Sciaena</i>	348
<i>strigatus</i>	433	<i>Serraria</i>	284
<i>superbus</i>	434	<i>scelerus, Hadropterus</i>	284
<i>tænopterus</i>	433	<i>Ophidion</i>	348
<i>trilobatus</i>	434	<i>serula, Hadropterus</i>	284
<i>trispinosus</i>	434	<i>scintillans, Zostomias</i>	70
<i>truncatus</i>	433	Sciaeniscus	478
<i>turchesius</i>	435	<i>scitula, Anchovia</i>	49
<i>valginiensis</i>	431	<i>Anchoviella</i>	49
<i>vetula</i>	433, 434	<i>scutuliceps, Synodus</i>	164
<i>virginalis</i>	433	<i>scitulus, Merulius</i>	407
<i>virens</i>	433	<i>Prionotus</i>	407
<i>viridis</i>	432	<i>scururus, Diplectrum</i>	318
<i>scaurus, Lycodopsis</i>		<i>Hæmulon</i>	330
<i>scapus, Minnus</i>	123	<i>Sparus</i>	329, 330
<i>Notropis</i>	123	Sclerodermi	491
<i>Triglops</i>	380	Sclerognathus	102, 103
<i>Schedophilopsis</i>		<i>cypinus</i>	102
<i>spinosus</i>	269	<i>cyprinella</i>	102, 103
<i>Schedophilus</i>		<i>meridionalis</i>	104
<i>heathi</i>	268	Sclerotis	300
<i>medusophagus</i>	268	<i>miniatu</i>	300
<i>Schilbeodes</i>		scopelid, Anchoviella	49
<i>gyrinus</i>	155	<i>Stolephorus</i>	49
<i>Schipa</i>	34	Scelacosoma	9
<i>Schistorus</i>	310	Scoliodon	14
<i>Schizophalus</i>	184	<i>lalandi</i>	15
<i>schmardæ, Dasyatis</i>	28	<i>laticaudus</i>	14
<i>Pastinachus</i>	27	<i>longurio</i>	16
<i>schmidti, Edriolychnus</i>	508	<i>porosus</i>	16
<i>schmidti, Hoplunnis</i>	81	<i>terre-novæ</i>	14, 15
<i>schneideri, Ophichthys</i>	88	scopaces, Nemichthys	83
<i>Pæcellia</i>	190	scopaceus, Nemichthys	83
<i>schœpfi, Cyclichthys</i>	502	scopelates, Balistes	245, 246
<i>Diodon</i>	502	<i>Centriscus</i>	245
<i>schœpfi, Balistes</i>	495	<i>Macrorhamphosus</i>	246
<i>Ceratacanthus</i>	495	Scopelopsis sayanus	215
<i>scholaris, Thysites</i>	201	Scomber	257
<i>schomburgkii, Pempheris</i>	322	<i>alalunga</i>	260
<i>schoneveldii, Cataphractus</i>	396	<i>albicaores</i>	260
<i>schrankii, Hæmulon</i>	330	<i>alleteratus</i>	258
<i>schypa, Antaeus</i>	34	<i>auratus</i>	258
<i>Sciadichthys</i>		<i>australicus</i>	258
<i>emphysetus</i>	148	<i>balantioptthalmus</i>	271
<i>proops</i>	148	<i>bicus</i>	259
<i>trocheli</i>	148	<i>boops</i>	270
<i>Sciaedes pictus</i>	148	<i>carangus</i>	273
<i>trocheli</i>	148	<i>chrysurus</i>	276
(<i>Sciaedes</i>) <i>emphysetus</i> , <i>Bagrus</i>	148	<i>collas</i>	257, 258
<i>sciadicus, Fundulus</i>	177	<i>crumenophthalmus</i>	270, 271
<i>Zygonectes</i>	177	<i>cryos</i>	272
<i>Sciadonus</i>	483	<i>dehayi</i>	258
<i>pedicellaris</i>	483	<i>diego</i>	258
<i>Sciæna</i>	348	<i>fuscatus</i>	279
<i>aliuta</i>	346	<i>germo</i>	280
<i>aquila</i>	61	<i>gladius</i>	263
<i>argentea</i>	334	<i>gracilis</i>	258
<i>argyrea</i>	341	<i>rex</i>	258
<i>armata</i>	273	<i>gunneri</i>	217
<i>(Bairdiella) bedotis</i>	346	<i>hippos</i>	273
<i>caprodes</i>	282, 283	<i>ignobilis</i>	273
<i>cirrhosa</i>	380	<i>japonicus</i>	258
<i>coro</i>	333	<i>lacertus</i>	258
<i>crokeri</i>	349	<i>latus</i>	276
<i>edwardi</i>	353	<i>macropthalmus</i>	258

List of scientific names—Continued.

	Page		Page
<i>Scomber maculatus</i>	257, 258	<i>Scorpaena calcarata</i>	371
<i>mediterraneus</i>	259	<i>colesi</i>	370
<i>pelagicus</i>	264	<i>cristulata</i>	372
<i>pelamides</i>	258	<i>dactyloptera</i>	370
<i>pelamis</i>	258	<i>dactylopterus</i>	370
<i>pelamitus</i>	259	<i>echinata</i>	372
<i>pelamys</i>	259	<i>grandicornis</i>	371
<i>plumieri</i>	271	<i>guttata</i>	371
<i>pneumatophorus</i>	257, 258	<i>histrio</i>	371
<i>quadripunctatus</i>	258	<i>inermis</i>	371
<i>regalis</i>	257	<i>massilensis</i>	308
<i>rim</i>	273	<i>mystes</i>	371
<i>rochel</i>	258, 259	<i>nuttingi</i>	371
<i>ruber</i>	271	<i>occipitalis</i>	371
<i>saliens</i>	277	<i>pannosa</i>	371
<i>sarda</i>	259	<i>picta</i>	370
<i>saurus</i>	277	<i>plumieri</i>	370
<i>scombrus</i>	257	<i>polylepis</i>	370
<i>sloani</i>	260	<i>porcus</i>	370
<i>speciosus</i>	273	<i>rascacio</i>	370
<i>thazard</i>	259	<i>russula</i>	371
<i>thynnus</i>	250	<i>atlantica</i>	370
<i>trachurus</i>	270	<i>(Sebastoplus) sierra</i>	372
<i>undulatus</i>	258	<i>sonoræ</i>	371
<i>vernalis</i>	257	<i>stearnsi</i>	371
<i>zonatus</i>	277	<i>tridecemspinosa</i>	370
<i>Scomberesocidae</i>	197	<i>Scorpaenichthyidae</i>	382
<i>Scomberesox</i>	197	<i>Scorpaenichthys</i>	382
<i>brevirostris</i>	198	<i>lateralis</i>	377
<i>camperi</i>	197	<i>marmoratus</i>	382
<i>equirostrum</i>	197	<i>Scorpaenidae</i>	363
<i>forsteri</i>	197	<i>Scorpaenodes</i>	370
<i>rondeletii</i>	197	<i>xyris</i>	370
<i>saurus</i>	197	<i>Scorpidæ</i>	358
<i>scutellatum</i>	197	<i>scorpioides</i> , <i>Acanthocottus</i>	386
<i>storeri</i>	197	<i>Cottus</i>	386
<i>scomberius</i> , <i>Esox</i>	173	<i>Scorpius californiensis</i>	358
<i>Scomberoides</i>	260	<i>scorpius</i> , <i>Acanthocottus</i>	386
<i>saltator</i>	277	<i>Cottus</i>	386
<i>scomberoides</i> , <i>Coryphaena</i>	265	<i>scouleri</i> , <i>Salmo</i>	54
<i>Scomberomorus</i>	257	<i>scovelli</i> , <i>Siphonostoma</i>	242
<i>concolor</i>	257	<i>Syrichtes</i>	242
<i>maculatus</i>	257	<i>scovellii</i> , <i>Etheostoma</i>	293
<i>plumieri</i>	257	<i>scrippsi</i> , <i>Otopholidium</i>	483
<i>regalis</i>	257	<i>scripta</i> , <i>Balistes</i>	495
<i>sierra</i>	257	<i>Osbeckia</i>	495
<i>Scombridæ</i>	257	<i>scriptus</i> , <i>Balistes</i>	495
<i>Scombriformes</i>	256	<i>Gymnothorax</i>	92
<i>Scombrocttus</i>	373	<i>Pseudoscopelus</i>	452
<i>salmoneus</i>	373	<i>scrutator</i> , <i>Isesthes</i>	482
<i>Scombrocttidae</i>	297	<i>scudderii</i> , <i>Hæmulon</i>	330
<i>Scombrus alalonga</i>	260	<i>scutata</i> , <i>Echeneis</i>	449
<i>scombrus</i> , <i>Scomber</i>	257	<i>scutator</i> , <i>Belone</i>	196
<i>Scopelengys</i>	170	<i>scutatus</i> , <i>Ælurichthys</i>	148
<i>dispar</i>	170	<i>Bagre</i>	148
<i>tristis</i>	170	<i>scutellatum</i> , <i>Scomberesox</i>	197
<i>Scopelinus caninianus</i>	169	<i>Scuticaria</i>	95
<i>Scopelus</i>	169	<i>scuticaris</i> , <i>Bascanichthys</i>	86
<i>affine</i>	169	<i>Sphagebranchus</i>	86
<i>andreae</i>	168	<i>scutiger</i> , <i>Icelus</i>	378
<i>arcticus</i>	170	<i>Rastrinus</i>	378
<i>benoiti</i>	169	<i>scutum</i> , <i>Achirus</i>	229
<i>coccoi</i>	168,	<i>Soles</i>	229
<i>crassiceps</i>	233	<i>Scylliorhinus</i>	11
<i>engraulis</i>	168	<i>scylla</i> , <i>Exocetus</i>	201
<i>glacialis</i>	170	<i>Hybopsis</i>	135
<i>gracilis</i>	169	<i>Scylliorhinidae</i>	11
<i>hygomii</i>	169	<i>Scylliorhinus</i>	11
<i>madeirensis</i>	167	<i>catulus</i> (<i>canicula</i>)	13
<i>metapocampus</i>	168	<i>cephalus</i>	11
<i>mizolepis</i>	233	<i>indicus</i>	12
<i>muelleri</i>	170	<i>profundorum</i>	12
<i>rarus</i>	168	<i>retifer</i>	11
<i>scopifer</i> , <i>Phenacobius</i>	142	<i>Scyllium</i>	11
<i>scopifera</i> , <i>Sarcidium</i>	142	<i>capense</i>	11
<i>scopiferus</i> , <i>Notropis</i>	132	<i>laticeps</i>	11
<i>scops</i> , <i>Gnathopops</i>	453	<i>retiferum</i>	11
<i>Scorpaena</i>	370	<i>scyllium</i> , <i>Triakis</i>	14
<i>agassizii</i>	371	<i>Scymnus rostratus</i>	14
<i>albifimbria</i>	371	<i>Scyris</i>	21
<i>albofasciata</i>	393	<i>analis</i>	274
<i>americana</i>	393	<i>indica</i>	274
<i>americanus</i>	370	<i>Scyllalichthys</i>	89
<i>atlantica</i>	372	<i>miurus</i>	89
<i>bergii</i>	371	<i>Scytalina</i>	473
<i>brasiliensis</i>	371	<i>cerdale</i>	473
<i>bufo</i>	370	<i>Scytaliniidae</i>	473

List of scientific names—Continued

	Page		Page
<i>Scybalinus</i>	473	<i>Sebastodes semicinctus</i>	367
<i>Scytalophis</i>	87, 88	<i>serranoides</i>	368
<i>magniculus</i>	87	<i>swifl</i>	367
<i>sebago, Salmo</i>	56	<i>wilsoni</i>	368
<i>Sebastes</i>	363	<i>Sebastolobus</i>	364
<i>auriculatus</i>	368	<i>aliascanus</i>	364
<i>bougainvillae</i>	370	<i>altivellis</i>	364
<i>caurinus</i>	369	<i>Sebastomus</i>	367, 368
<i>elongatus</i>	368	<i>ayresii</i>	367
<i>fasciatus</i>	363, 369	<i>chlorostictus</i>	367
<i>glaucus</i>	364	<i>constellatus</i>	367
<i>helvomaculatus</i>	367	<i>eos</i>	368
<i>imperialis</i>	370	<i>gilli</i>	368
<i>macrochir</i>	364	<i>rhodochloris</i>	367
<i>macrocephalus</i>	372	<i>rosaceus</i>	367
<i>marinus</i>	363	<i>rupestris</i>	368
<i>melanops</i>	364	<i>umbrosus</i>	367
<i>minutus</i>	370	<i>Sebastopsis</i>	370
<i>nebulosus</i>	369	<i>xyris</i>	370
<i>nemastophthalmus</i>	322	<i>Sebastopyr</i>	368
<i>nigrocinctus</i>	370	<i>ruberrimus</i>	368
<i>ovalis</i>	365	<i>Sebastosomus</i>	364
<i>paucispinis</i>	364	<i>aleuticus</i>	365
<i>regulus</i>	363	<i>ciliatus</i>	365
<i>rosaceus</i>	367	<i>flavidus</i>	364
<i>ruber</i>	368	<i>melanops</i>	365
<i>parvus</i>	368	<i>mystinus</i>	365
<i>viviparus</i>	364	<i>pinniger</i>	365
<i>Sebastichthys</i>	369, 370	<i>serranoides</i>	365
<i>alutus</i>	366	<i>simulans</i>	365
<i>atrovirens</i>	365	<i>seco, Luxilus</i>	115
<i>aurora</i>	366	<i>Sectator</i>	340
<i>carnatus</i>	369	<i>ocyrurus</i>	340
<i>chlorostictus</i>	369	<i>sectatrix, Kyphosus</i>	340
<i>constellatus</i>	367	<i>Perca</i>	340
<i>diploproa</i>	367	<i>marina</i>	340
<i>entomelas</i>	366	<i>sector, Serrivomer</i>	84
<i>fasciolaris</i>	369	<i>secundodorsalis, Thunnus</i>	260
<i>introniger</i>	366	<i>secundus, Caranx</i>	360
<i>levis</i>	368	<i>sedentarius, Chetodon</i>	160
<i>maliger</i>	369	<i>seemani, Arius</i>	160
<i>miniatus</i>	365	<i>Galeichthys</i>	160
<i>mystinus</i>	364, 365	<i>Selache</i>	20
<i>nigrocinctus</i>	370	<i>Selachli</i>	10
<i>proriger</i>	366	<i>selachops, Apterichtus</i>	84
<i>var. brevispinis</i>	366	<i>Sphagebranchus</i>	84
<i>rastrelliger</i>	369	<i>Selachostom</i>	36
<i>rhodochloris</i>	367	<i>Selanonus</i>	19
<i>rubriventris</i>	368	<i>walkeri</i>	19
<i>rupestris</i>	368	<i>Selar</i>	270
<i>saxicola</i>	367	<i>Seleima</i>	337, 339
<i>sericeps</i>	369	<i>aurata</i>	339
<i>sinensis</i>	368	<i>Selenaspis</i>	148
<i>umbrosus</i>	367	<i>dowii</i>	148
<i>verricularis</i>	369	<i>Selene</i>	275
<i>zacentrus</i>	368	<i>argentea</i>	275
<i>Sebastocarthus</i>	369	<i>cerstedi</i>	275
<i>serriceps</i>	369	<i>quadrangularis</i>	368
<i>Sebastodes</i>	108, 304, 370	<i>selene, Carpiodes</i>	103
<i>aereus</i>	367	<i>Luxilus</i>	132
<i>aleutianus</i>	368	<i>Selenichthys</i>	216
<i>alexandri</i>	365	<i>seleniris, Trachypterus</i>	216
<i>atrorubens</i>	365	<i>selanops, Hiodon</i>	41
<i>ayresii</i>	367	<i>Hyodon</i>	41
<i>babcocki</i>	365	<i>sellaris, Acanthocottus</i>	388
<i>clavilatus</i>	366	<i>Alvordius</i>	284
<i>crameri</i>	367	<i>Hadropterus</i>	284
<i>deani</i>	367	<i>Porocottus</i>	388
<i>eigenmanni</i>	366	<i>sellicauda, Apogon</i>	295
<i>emphæus</i>	367	<i>Epinephelus</i>	311
<i>eos</i>	368	<i>sellifer, Halichoeres</i>	428
<i>flavidus</i>	364	<i>Iridio</i>	428
<i>gilberti</i>	369	<i>Sems</i>	409
<i>gilli</i>	368	<i>signifer</i>	409
<i>goodei</i>	364	<i>semicarmatus, Gasterosteus</i>	237
<i>hopkinsi</i>	366	<i>semicinctus, Eosebastes</i>	367
<i>Jordan</i>	364	<i>Gillellus</i>	456
<i>melanops</i>	365	<i>Iridio</i>	427
<i>melanostomus</i>	366	<i>Julis</i>	427
<i>nigrocinctus</i>	364	<i>semicorona, Sariola</i>	279
<i>ovalis</i>	366	<i>semifasciata, Pleoma</i>	282
<i>paucispinus</i>	364	<i>semifasciatus, Pleoma</i>	283
<i>rosaceus</i>	368	<i>Triakis</i>	14
<i>ruberrimus</i>	368	<i>semifasciatus, Gobius</i>	441
<i>rufus</i>	366	<i>Serranus</i>	317
		<i>semiloricatus, Gasterosteus</i>	237

List of scientific names—Continued

	Page		Page
semiluna, <i>Sparus</i>	327	serra, <i>Alepidosaurus</i>	171
semilunata, <i>Diana</i>	264	Gonionotus.....	280
seminolis, <i>Fundulus</i>	178	Serrana.....	363
<i>Zygonectes</i>	178	hispana	363
seminuda, <i>Albula</i>	40	Serranidae.....	317
<i>Gila</i>	114	serranoides, <i>Sebastosomus</i>	365
<i>Gymnoleotris</i>	438	Serranus <i>acanthophorus</i>	317
seminudus, <i>Eleotris</i>	438	aculeatus (<i>oculatus</i>).....	328
Enypniás.....	445	<i>acutirostris</i>	315
Gobius.....	445	aequidens	319
Lycias.....	475	<i>albomaculatus</i>	317
Lycodes.....	475	<i>altivella</i>	310
Plecostomus.....	160	<i>angustifrons</i>	311
semiradiatus, <i>Lepidotesteus</i>	38	<i>arara</i>	311
semiruber, <i>Labrus</i>	425	<i>armatus</i>	313
semiscaber, <i>Cottopsis</i>	384	<i>aspersus</i>	310
<i>Cottus</i>	384	bivittatus	318
semiscabra centropleura, <i>Uranidea</i>	383	<i>bonaci</i>	314
semispinosus, <i>Caranx</i>	270	<i>brunneus</i>	314
semisquamatum, <i>Pycnomma</i>	438	<i>bulleri</i>	319
Semitapicus.....	100	<i>calopteryx</i>	319
Semotilus.....	116, 117	<i>camelopardalis</i>	315
<i>atromaculatus</i>	117	<i>capuena</i>	331
<i>thoreauanus</i>	117	<i>carauna</i>	309
<i>biguttatus</i>	137	<i>catus</i>	311
<i>cephalus</i>	117	<i>colonus</i>	320
<i>diplemis</i>	125	<i>conspersus</i>	311
<i>diplemia</i>	117	<i>coronatus</i>	308
<i>dorsalis</i>	117	<i>couchii</i>	308
<i>hammondi</i>	117	<i>courtadei</i>	310
<i>macrocephalus</i>	117	<i>creolus</i>	320
<i>notatus</i>	177	<i>cyclopomatus</i>	314
<i>olivaceus</i>	177	<i>decimalis</i>	314
<i>speciosus</i>	117	<i>dimidiatus</i>	314
<i>thoreauanus</i>	117	<i>dubius</i>	309
senegalensis, <i>Vomer</i>	275	<i>emarginatus</i>	315
senilis, <i>Gambusia</i>	185	<i>erythrogaster</i>	312
senta, <i>Raja</i>	25	<i>falcatus</i>	315
sephen, <i>Raja</i>	28	<i>fascicularis</i>	318, 319
septemfasciatus, <i>Cichlosoma</i>	421	<i>felinus</i>	315
septemfasciatus, <i>Astatheros</i>	421	<i>fimbriatus</i>	311
septentrionalis, <i>Gaidropsarus</i>	208	<i>flavescens</i>	319
<i>Motella</i>	208	<i>furcifer</i>	320
<i>Perca</i>	307	<i>fuscus</i>	315
septiphinis, <i>Ammodytes</i>	451	<i>galeus</i>	312
<i>Rhynchias</i>	451	<i>guasa</i>	312
Serannus bonaci.....	314	<i>gigas</i>	310
serena, <i>Dionda</i>	144	<i>humeralis</i>	317
sergeant, <i>Osmerus</i>	67	<i>impetiginosus</i>	310
Seriola.....	269, 276, 279	<i>inermis</i>	313
<i>bipinnata</i>	279	<i>interstitialis</i>	314
<i>bonariensis</i>	280	<i>iradians</i>	319
<i>boscii</i>	280	<i>italara</i>	312
<i>carolinensis</i>	279	<i>jacome</i>	319
<i>coronata</i>	280	<i>labriformis</i>	311
<i>declivis</i>	280	<i>lamprurus</i>	321
<i>dorsalis</i>	279	<i>latepictus</i>	314
<i>dubia</i>	280	<i>lucopercaus</i>	319
<i>dumerili</i>	279	<i>meculetofasciatus</i>	317
<i>dussumieri</i>	278	<i>maculosus</i>	311
<i>falcata</i>	280	<i>margaritifer</i>	311
<i>gigas</i>	279	<i>marginatus</i>	311
<i>lalandi</i>	279	<i>mentzelli</i>	311
<i>ligulata</i>	280	<i>morio</i>	312
<i>mazatlanica</i>	279	<i>morrhua</i>	310
<i>pinnulata</i>	278	<i>mystacinus</i>	310
<i>proxima</i>	279	<i>nigriculus</i>	310
<i>rivoliana</i>	280	<i>nigritus</i>	312
<i>semicoronata</i>	279	<i>niveatus</i>	310, 311
<i>stearnsi</i>	279	<i>oculatus</i>	328
<i>succincta</i>	278	<i>occurus</i>	318
<i>zonata</i>	279	<i>olax</i>	314
Seriolichthys.....	279	<i>ouatalibi</i>	309
Seriolidae.....	278	<i>panamensis</i>	308
Seriolophus.....	280	<i>(Paralabrax) dewegeri</i>	319
<i>carangoides</i>	280	<i>petrosus</i>	313
Seriphus.....	353	<i>phaeton</i>	309
<i>politus</i>	353	<i>phœbe</i>	319
Serotinus, <i>Acipenser</i>	35	<i>pixanga</i>	310
<i>Acipenser</i>	33	<i>prestigiator</i>	319
Serpens marinus maculosus.....	87	<i>psittacinus</i>	319
Serpens, <i>Gempylus</i>	261, 262	<i>quinquefasciatus</i>	312
<i>Lycodes</i>	476	<i>radians</i>	318
<i>Ophisurus</i>	85	<i>radialis</i>	318
serpentina, <i>Muraena</i>	77	<i>remotus</i>	312
serpentinus, <i>Blennius</i>	470, 471	<i>repandus</i>	315
<i>Derichthys</i>	77	<i>rupestris</i>	313

List of scientific names—Continued

	Page		Page
<i>Serranus semifasciatus</i>	317	<i>Sicyases fasciatus</i>	490
stadhouderi	311	<i>haeres</i>	490
tænios	309	<i>punctulatus</i>	490
tigris	315	<i>rubiginosus</i>	490
tinca	315	<i>sanguineus</i>	490
undulosus	315	<i>Sicydium</i>	447
varius	310	<i>antillarum</i>	447
<i>Serraria</i>	284	<i>buscki</i>	447
<i>sciera</i>	284	<i>caguite</i>	447
<i>serula</i>	284	<i>gymnogaester</i>	447
<i>serrata</i> , <i>Fistularia</i>	245	<i>plumieri</i>	447
<i>Sayris</i>	197	<i>punctatum</i>	447
<i>serrato-granulata</i> , <i>Perca</i>	281	<i>salvini</i>	448
<i>serratum</i> , <i>Hæmulon</i>	329	<i>siragus</i>	447
<i>serratus</i> , <i>Gasterosteus</i>	237	<i>vincente</i>	447
<i>Nauirates</i>	278	<i>Sicyogaster</i>	488
<i>sericeps</i> , <i>Sebastichthys</i>	369	<i>Sidera</i>	93
<i>Sebastocarthus</i>	369	<i>castanea</i>	92
<i>serifer</i> , <i>Conodon</i>	333	<i>chlevastes</i>	93
<i>Serrivomer</i>	84	<i>verrilli</i>	93
<i>beani</i>	84	<i>Sidera</i>	93
<i>sector</i>	84	<i>picta</i>	93
<i>serula</i> , <i>Anisotremus</i>	333	<i>sideris</i> , <i>Muraena</i>	93
<i>Chalinura</i>	204	<i>sideritus</i> , <i>Hybognathus</i>	140
<i>Hadropterus sclerurus</i>	284	<i>Hyborhynchus</i> (<i>Hybognathus</i>)	140
<i>Priacanthus</i>	323	<i>sleboldii</i> , <i>Heros</i>	423
<i>Pristipoma</i>	333	<i>Sierra</i>	257
<i>Pseudopriacanthus</i>	323	<i>cavalla</i>	257
<i>Serraris</i>	284	<i>sierra</i> , <i>Doryrhamphus</i>	243
<i>sessilicauda</i> , <i>Monolepis</i>	220	<i>Pontinus</i>	372
<i>setapinnis</i> , <i>Vomer</i>	275	<i>Scomberomorus</i>	257
<i>Zeus</i>	275	<i>Scorpaena</i> (<i>Sebastoplus</i>)	372
<i>setapinnis cubensis</i> , <i>Vomer</i>	275	<i>slerrita</i> , <i>Tylosurus</i>	196
<i>Setarches</i>	372	<i>sigalutes</i> , <i>Gilbertidia</i>	393
<i>guntheri</i>	372	<i>Gilbertina</i>	392, 393
<i>parmatus</i>	372	<i>Sigmistes</i>	391
<i>setifer</i> , <i>Argyrosomus</i>	275	<i>caulias</i>	391
<i>Monacanthus</i>	494	<i>Sigmops</i>	72
<i>setiger</i> , <i>Dasy cottus</i>	392	<i>stigmaticus</i>	72
<i>setigerus</i> , <i>Lophi omus</i>	494	<i>Sigmurus</i>	348
<i>Lophius</i>	504	<i>vermicularis</i>	348
<i>setipinnis</i> , <i>Argyroreius</i>	604	<i>Signalosa</i>	46
<i>setosus</i> , <i>Mugil</i>	275	<i>atchafalayae</i>	46
<i>Querimana</i>	254	<i>vanhynningi</i>	47
<i>Tetraodon</i>	254	<i>mexicanus</i>	47
<i>seu conger</i> , etc., <i>Murena</i>	500	<i>campi</i>	47
<i>severus</i> , <i>Heros</i>	94	<i>petenonis</i>	47
<i>sexcornutus</i> , <i>Ostracion</i>	497	<i>signata</i> , <i>Cyclothona</i>	72
<i>sexdecim-lamellatus</i> , <i>Echeneis</i>	449	<i>signatus</i> , <i>Bathymaster</i>	451
<i>sexfasciatum</i> , <i>Clinchosoma</i>	421	<i>Hypopion</i>	17
<i>Hæmulon</i>	329	<i>Microgobius</i>	443
<i>sexfasciatus</i> , <i>Astatheros</i>	421	<i>signifer</i> , <i>Bryttus</i>	209
<i>Hæmulon</i>	329	<i>Centrithmus</i>	319, 320
<i>sex-maculatus</i> , <i>Diodon</i>	501	<i>Chatotessus</i>	44
<i>sexradiatus</i> , <i>Polynemus</i>	409	<i>Coregonus</i>	68
<i>shasta</i> , <i>Cottus</i>	383	<i>Monacanthus</i>	494
<i>Salmo</i>	58	<i>Neetroplus</i>	423
<i>gaudneri</i>	58	<i>Sema</i>	409
<i>shavlani</i> , <i>Cetorhinus</i>	20	<i>Stypondon</i>	113
<i>shufeldti</i> , <i>Gobius</i>	439	<i>Tynhmalius</i>	68
<i>Mancallas</i>	508	<i>silenus</i> , <i>Zaprora</i>	452
<i>Rhinogobius</i>	439	<i>Siluridae</i>	147
<i>Typhlosoras</i>	508	<i>Silurus bagre</i>	147
<i>shumardi</i> , <i>Alburnops</i>	134	<i>bells</i>	149
<i>Hadropterus</i>	285	<i>catus</i>	153
<i>Hybopsis</i>	134	<i>clarus</i>	158, 159
<i>Imostoma</i>	285	<i>cornutus</i>	245, 246
<i>sialis</i> , <i>Argentina</i>	68	<i>cupreus</i>	153
<i>Corvula</i>	345	<i>felis</i>	148
<i>Noturus</i>	155	<i>fuscus</i>	153
<i>Vacuoqua</i>	345	<i>galeatus</i>	159
<i>sibbaldi</i> , <i>Syngnathus</i>	343	<i>gyrinus</i>	155
<i>Siboma</i>	120	<i>hertzbergi</i>	148
<i>atrraria</i>	119	<i>limosus</i>	155
<i>conformis</i>	120	<i>lividus</i>	153
<i>crassicauda</i>	120	<i>marinus</i>	147, 148
<i>sicana</i> , <i>Cerna</i>	312	<i>melas</i>	154
<i>siccerum</i> , <i>Holocentrum</i>	236	<i>militaris</i>	159
<i>siculum</i> , <i>Chi rostoma</i>	251	<i>nebulosus</i>	156
<i>siculus</i> , <i>Labrisomus</i>	251	<i>olivaris</i>	155
<i>sicula</i> , <i>Pimelopterus basci</i>	340	<i>punctatus</i>	161, 162
<i>siculus</i> , <i>Lampugus</i>	265	<i>viscosus</i>	155
<i>Sicya</i>	447	<i>xanthocephalus</i>	154
<i>Sicyases</i>	490	<i>var. fucus</i>	153
<i>carneus</i>	490	<i>Silurus</i>	68
<i>erythrops</i>	490	<i>ascani</i>	68

List of scientific names—Continued

Page	Page
<i>silus</i> , <i>Argentina</i>	68
<i>sima</i> , <i>Careproctus</i>	401
<i>Careproctus</i>	401
<i>Electris</i>	436
<i>Simenichelys</i>	78
<i>Simenichelys</i>	78
<i>parasiticus</i>	78
<i>similis</i> , <i>Amphistichus</i>	411
<i>Fundulus</i>	177
<i>Hæmulon</i>	330
<i>Hydrargyre</i>	175, 177
<i>Iridio</i>	427
<i>Rhinogobius</i>	439
<i>simillima</i> , <i>Palometa</i>	266
<i>simillimus</i> , <i>Geres</i>	342
<i>Poronotus</i>	266
<i>simotera</i> , <i>Ulocentra</i>	286
<i>simoterum</i> , <i>Hyostoma</i>	286
<i>simplex</i> , <i>Aplurus</i>	261
<i>Cranidens</i>	339
<i>Pseudoscarus</i>	435
<i>Tetragonurus</i>	261
<i>simplicidens</i> , <i>Girella</i>	339
<i>Incisidens</i>	339
<i>simpsoni</i> , <i>Ictalurus</i>	152
<i>simula</i> , <i>Challinura</i>	204
<i>simulans</i> , <i>Hemiplites</i>	304
<i>Sebastosomus</i>	265
<i>Synodus</i>	164
<i>simulus</i> , <i>Ophioscion</i>	348
<i>simus</i> , <i>Alburnellus</i>	135
<i>Careproctus</i>	401
<i>Catetix</i>	480
<i>Creagrus</i>	98
<i>Hypobopsis</i>	135
<i>Monticirrus</i>	351
<i>Rhinichthys</i>	140
<i>sinuosus</i> , <i>Paralichthys</i>	224
<i>Siphonotoma</i>	241
<i>Syngnathus</i>	241
<i>Umbrina</i>	350
<i>sinensis</i> , <i>Hispanicus</i>	368
<i>Sebastichthys</i>	368
<i>Trygon</i>	28
<i>sinuosa</i> , <i>Pomadasys</i>	335
<i>Rhomboides</i>	335
<i>Siphagonus barbata</i>	394
<i>barbatus</i>	394
<i>Siphatales</i>	119, 120, 122
<i>bicolor</i>	122
<i>columbianus</i>	122
<i>formosus</i>	122
<i>mohavensis</i>	123
<i>obesus</i>	122
<i>olivaceus</i>	122
<i>oregonensis</i>	122
<i>vittatus</i>	122
<i>Siphonotoma</i>	240
<i>artum</i>	242
<i>auliscus</i>	240
<i>barbara</i>	240
<i>brevicaudum</i>	241
<i>carinatum</i>	240
<i>crinigerum</i>	241
<i>dendriticum</i>	242
<i>exile</i>	243
<i>floride</i>	241
<i>mackayi</i>	240
<i>poeyi</i>	241
<i>robertsi</i>	241
<i>scovelli</i>	242
<i>sinaloae</i>	241
<i>starksii</i>	242
<i>torrei</i>	242
<i>zatropis</i>	243
<i>siragus</i> , <i>Scydium</i>	447
<i>Sirembo guentheri</i>	488
<i>messieri</i>	480
<i>nigripinnis</i>	62
<i>sisco</i> , <i>Argyrosonus</i>	62
<i>huromius</i> , <i>Leucichthys</i>	62
<i>Leucichthys</i>	62
<i>siscowet</i> , <i>Cristivomer</i>	60
<i>Salmo</i>	60
<i>Salmo</i>	60
<i>skawitz</i> , <i>Salmo</i>	118
<i>siuslawi</i> , <i>Leuciscus</i>	118
<i>Richardsonius</i>	118
<i>skib</i> , <i>Pomatomus</i>	280
<i>Skiffia</i>	183
<i>bilineata</i>	183
<i>lermae</i>	183
<i>multipunctata</i>	183
<i>variegata</i>	183
<i>sloani</i> , <i>Chauliodus</i>	71
<i>Lophotus</i>	29
<i>Scomber</i>	260
<i>Urobatis</i>	29
<i>sloani vermiculatus</i> , <i>Urobatis</i>	29
<i>sluiteri</i> , <i>Bramnerella</i>	464
<i>Smaragdus</i>	440
<i>costales</i>	441
<i>stigmaticus</i>	441
<i>valenciennesi</i>	441
<i>smaragdus</i> , <i>Eleotris</i>	437
<i>Erotelis</i>	437
<i>Gobius</i>	440, 441
<i>Gobionellus</i>	441
<i>Salmo</i>	58
<i>Smaris</i>	340
<i>lineatus</i>	342
<i>martinicus</i>	340
<i>smaris</i> , <i>Sparus</i>	340
<i>Smecticus</i>	316
<i>bicolor</i>	316
<i>smiridus</i> , <i>Merluccius</i>	214
<i>smithi</i> , <i>Ceratichthys</i>	145
<i>Lepidotremus</i>	37
<i>smithi</i> , <i>Ciliola</i>	145
<i>Cyprinus</i> (<i>Abramis</i>).....	41
<i>Halieutichthys</i>	507
<i>snyderi</i> , <i>Catostomus</i>	106
<i>Dialarchus</i>	391
<i>Gnathoprops</i>	453
<i>Hyperoplus</i>	199
<i>Oligocottus</i>	391
<i>Palometa</i>	267
<i>Peprius</i>	267
<i>Sobaco</i>	492
<i>sobaco</i> , <i>Balistes</i>	492
<i>Canthidermis</i>	492
<i>sobra</i> , <i>Mesoprion</i>	326
<i>socialis</i> , <i>Salmo</i>	66
<i>socius</i> , <i>Auburnus</i>	124
<i>Notropis</i>	124
<i>socorroense</i> , <i>Thalassoma</i>	429
<i>socorroensis</i> , <i>Thalassoma</i>	429
<i>sogo</i> , <i>Holocentrus</i>	235
<i>sohar</i> , <i>Chætodon</i>	362
<i>solandri</i> , <i>Acanthocybium</i>	256
<i>Cybium</i>	256
<i>solaris</i> , <i>Orthragoriscus</i>	503
<i>Solea browni</i>	229
<i>fischeri</i>	230
<i>fonsecai</i>	230
<i>gronovii</i>	229
<i>kunzingeri</i>	229
<i>lunata</i>	217
<i>mazatlanus</i>	230
<i>panamensis</i>	230
<i>pilosa</i>	230
<i>punctata</i>	217
<i>scutum</i>	229
<i>soleaformis</i> , <i>Rhombus</i>	219
<i>Solenostomus</i>	245
<i>Soleotalpa</i>	230
<i>unicolor</i>	230
<i>solis</i> , <i>Lepomis</i>	300
<i>Pomotis</i>	300
<i>Somniodes</i>	21
<i>Somnioides</i>	21
<i>brevipinna</i>	21, 22
<i>microcephalus</i>	22
<i>somnolentus</i> , <i>Eleotris</i>	436
<i>Lobotes</i>	323
<i>sonnerati</i> , <i>Opistognathus</i>	462
<i>sonnagii</i> , <i>Antennarius</i>	508
<i>Chironectes</i>	508
<i>sonorensis</i> , <i>Atherinopsis</i>	252
<i>Scorpaena</i>	371
<i>sonoriensis</i> , <i>Catostomus</i>	106
<i>Girardinus</i>	189
<i>soporator</i> , <i>Bathygobius</i>	440
<i>Gobius</i>	440
<i>sordida</i> , <i>Murresna</i>	94

CHECK LIST OF FISHES

653

List of scientific names—Continued

	Page		Page
<i>sordidus</i> , <i>Chætodon</i>	414	<i>speciosus</i> , <i>Pomotis</i>	301
<i>Citharichthys</i>	218	<i>Scomber</i>	273
<i>Psetticthys</i>	218	<i>Semotilus</i>	117
<i>Verilius</i>	328	<i>spectabilis</i> , <i>Pocilichthys</i>	291
<i>spadicea</i> , <i>Dionda</i>	144	<i>Salmo</i>	61
<i>Lamptetra</i>	9	<i>Salvelinus</i>	61
<i>spadiceus</i> , <i>Centrurophis</i>	87	<i>malma</i>	61
<i>Tetrapleurodon</i>	9	<i>spectrum</i> , <i>Careproctus</i>	402
<i>spallanzani</i> , <i>Iurus</i>	19	<i>Lophius</i>	505
<i>Leptoccephalus</i>	79	<i>Osmerus</i>	67
<i>Sparactodon</i>	280	<i>mordax</i>	67
<i>nainai</i>	280	<i>specrunculus</i> , <i>Hybopsis</i>	134
<i>Sparidae</i>	336	<i>speculiger</i> , <i>Exocoetus</i>	201
<i>Sparisoma</i>	431, 433	<i>speculigera</i> , <i>Lampradena</i>	167
<i>abildgaardi</i>	432	<i>spaleatus</i> , <i>Amblyopsis</i>	195
<i>atomarium</i>	432	<i>spangleri</i> , <i>Sphoeroides</i>	498
<i>austrorhynchum</i>	432	<i>Tetronod</i>	498
<i>brachiale</i>	433	<i>spat</i> , <i>Esox</i>	255
<i>chrysopterum</i>	432	<i>Spherooides tuberculatus</i>	498
<i>cyanoleuca</i>	432	<i>Sphagebranchus</i>	84
<i>distinctum</i>	432	<i>anguiformis</i>	84, 91
<i>flavescens</i>	433	<i>kendalli</i>	84, 85
<i>frondosum</i>	433	<i>ophioneus</i>	84
<i>hoplomystax</i>	432	<i>rostratus</i>	84
<i>lorito</i>	432	<i>scuticaris</i>	84
<i>macrachelipios</i>	433	<i>selachops</i>	84
<i>niphobles</i>	432	<i>teres</i>	86
<i>oxybrachium</i>	432	(<i>Sphagebranchus</i>) <i>anguiformis</i> , <i>Ophichthys</i>	84
<i>radians</i>	432	<i>sphenops cuneata</i> , <i>Molliesenia</i>	193
<i>rubripinne</i>	433	<i>sphenops</i> , <i>Molliesenia</i>	192
<i>strigatum</i>	433	<i>Faecilla</i>	192
<i>viride</i>	432	<i>vandepolli</i> , <i>Molliesenia</i>	193
<i>xystodon</i>	431	<i>Spheroide tubercula</i>	498
<i>Sparisomidae</i>	431	<i>Spherooides</i>	498
<i>spario</i> , <i>Labrus</i>	304, 305	<i>asterias</i>	498
<i>Pomoxis</i>	305	<i>harperi</i>	498
<i>Sparis</i>	418	<i>les</i>	498
<i>abildgaardi</i>	432	(<i>Spherooides</i>) <i>eulepidotus</i> , <i>Tetronod</i>	498
<i>annularis</i>	338	<i>Spherooides</i>	498
<i>argenteus</i>	337	<i>angusticeps</i>	498
<i>argyrops</i>	336	<i>asterias</i>	498
<i>atlanticus</i>	310	<i>eulepidotus</i>	498
<i>aureo-ruber</i>	432	<i>harperi</i>	498
<i>aureus</i>	302	<i>lobatus</i>	498
<i>bajonado</i>	336	<i>maculatus</i>	498
<i>brachysomus</i>	337	<i>marmoratus</i>	498
<i>castaneola</i>	285	<i>nephelus</i>	498
<i>caris</i>	325	<i>spengleri</i>	498
<i>chromis</i>	413	<i>tuberculatus</i>	498
<i>chrysomelanurus</i>	311	<i>Sphyraena</i>	255
<i>clarysops</i>	336	<i>acus</i>	106, 197
<i>chrysurus</i>	327	<i>argentea</i>	255
<i>ercentatus</i>	308	<i>aureoviridis</i>	305
<i>falcatus</i>	425	<i>baracuda</i>	255
<i>niger</i>	265	<i>beccuna</i>	255
<i>ovicephalus</i>	338	<i>borealis</i>	255
<i>pagrus</i>	337	<i>ensis</i>	255
<i>protocephalus</i>	338	<i>guachanco</i>	255
<i>radiatus</i>	428	<i>günterii</i>	255
<i>rai</i>	265	<i>lucasana</i>	255
<i>rhomboides</i>	337	<i>picudilla</i>	255
<i>salin</i>	338	<i>sphyraena</i>	255
<i>sargus</i>	338	<i>picuda</i>	255
<i>sciturus</i>	329, 330	<i>viridensis</i>	255
<i>semiluna</i>	327	<i>vulgaris</i>	255
<i>smaris</i>	340	<i>sphyraena</i> , <i>Argentina</i>	68
<i>surinamensis</i>	424	<i>Chirostoma</i>	251
<i>synagrus</i>	327	<i>Esox</i>	255
<i>tetraacanthus</i>	325	<i>Sphyraena</i>	255
<i>vermicularis</i>	327	<i>sphyraenarum</i> , <i>Echenels</i>	449
<i>virginicus</i>	332	<i>Sphyraenidae</i>	225
<i>vitatus</i>	332	<i>Sphyraenops</i>	297
<i>xanthurus</i>	336	<i>bairdiana</i>	297
<i>Spathula</i>	36	<i>biardianus</i>	297
<i>Spathula</i> , <i>Polyodon</i>	36	<i>sphyraenops</i> , <i>Lestidiops</i>	166
<i>Squalus</i>	36	<i>Sphyriichthys</i>	18
<i>Spathularia</i>	36	<i>Sphyraena</i>	18
<i>reticulata</i>	36	<i>zygema</i>	18
<i>spatula</i> , <i>Atractosteus</i>	38	<i>Sphyriidae</i>	17
<i>Icelus</i>	379	<i>Spicara</i>	340
<i>Lepisosteus</i>	38	<i>flexuosa</i>	340
<i>Spatularia reticulata</i>	36	<i>martinica</i>	340
<i>speciosa</i> , <i>Gambusia</i>	183	<i>Spicifer</i> , <i>Hippichthys</i>	243
<i>speciosus</i> , <i>Gnathodon</i>	273	<i>Synagnathus</i>	243
<i>Gnathodon</i>	273	<i>spillmanii</i> , <i>Alvordius</i>	283

List of scientific names—Continued

	Page		Page
<i>spilonotopterus</i> , <i>Cypselurus</i>	202	<i>Squalidæ</i>	21
<i>Erocastus</i>	202	<i>qualidus</i> , <i>Scarus</i>	433
<i>spilonota</i> , <i>Poccilia</i>	192	<i>squalipeta</i> , <i>Echenelis</i>	449
<i>spilonotus</i> , <i>Coregonus</i>	65	<i>Squidius</i>	117
<i>Monacanthus</i>	495	<i>aliciae</i>	119
<i>Prosopium</i>	65	<i>copei</i>	119
<i>spilopterus</i> , <i>Citharichthys</i>	218, 219	<i>cruoratus</i>	119
<i>Photogenis</i>	129	<i>galtière</i>	118
<i>spilopus</i> , <i>Exocoetus</i>	201	<i>hyalope</i>	117
<i>spilota</i> , <i>Uranidea</i>	385	<i>lemmoni</i>	119
<i>spilotopterygius</i> , <i>Ballistes</i>	492	<i>photogenis</i>	123
<i>spilotum</i> , <i>Etheostoma nianküe</i>	285	<i>proriger</i>	120
<i>spilotus</i> , <i>Hypophomus</i>	285	<i>rhomaleus</i>	119
<i>spilura</i> , <i>Molliesenia</i>	192	<i>Squaloidei</i>	21
<i>spilurus</i> , <i>Archocentrus</i>	420	<i>Squalus</i>	13, 21
<i>Curimatus</i>	100	<i>acanthias</i>	21
<i>Heros</i>	420	<i>acronotus</i>	16
<i>Poccilia</i>	192	<i>americanus</i>	19
<i>Salmo</i>	57	<i>arcticus</i>	14
<i>Spinachia</i>	239	<i>argus</i>	12
<i>spinachia</i>	239	<i>borealis</i>	21, 22
<i>vulgaris</i>	230	<i>brucus</i>	22
<i>spinachia</i> , <i>Gasterosteus</i>	239	<i>caeruleus</i>	17
<i>Spinachilæ</i>	239	<i>canis</i>	11
<i>Spinax</i> (<i>Acanthias</i>) <i>suckleyi</i>	21	<i>carcharias</i>	15, 20
<i>Spinax</i> <i>fabricii</i>	21	(<i>Carcharias</i>) <i>terre-novæ</i>	14
<i>billianus</i>	21	<i>cetaceus</i>	20
<i>spinax</i> , <i>Squalus</i>	21	<i>cirratus</i>	12
<i>spinoscens</i> , <i>Auliscops</i>	239	<i>commersonii</i>	15
<i>spinicauda</i> , <i>Acanthenchelys</i>	89	<i>cornubicus</i>	19
<i>spinicephalus</i> , <i>Exoglossum</i>	146	<i>edentulus</i>	31
<i>Spinicephalus</i>	145	<i>elephas</i>	20
<i>fibulatus</i>	145	<i>ferox</i>	19
<i>spiniceps</i> , <i>Photocorynus</i>	509	<i>galeus</i>	13, 14
<i>spinidens</i> , <i>Scarus</i>	431, 432	<i>glacialis</i>	22
<i>spinifer</i> , <i>Acanthonus</i>	482	<i>glaucus</i>	17
<i>Engraulis</i>	50	<i>griseus</i>	11
<i>spinifera</i> , <i>Anchoviella</i>	50	<i>gunneri</i>	20
<i>spiniger</i> , <i>Icelus</i>	378	<i>hirundinaceus</i>	17
<i>Spinivomer</i>	84	<i>homianus</i>	20
<i>goodell</i>	84	<i>isodus</i>	20
<i>spinosa</i> , <i>Acanthemblemaria</i>	465	<i>litoralis</i>	19
<i>Malthopsis</i>	507	<i>longimanus</i>	15
<i>Thysanocara</i>	160	<i>macrodus</i>	19
<i>spinosissimum</i> , <i>Immaculatum</i> , <i>Cichlasoma</i>	420	<i>maximus</i>	20
<i>spinosissimus</i> , <i>Archocentrus</i>	420	<i>microcephalus</i>	21, 22
<i>Aspidophorus</i>	394	<i>monensis</i>	19
<i>Diodon</i>	501	<i>mustelus</i>	13, 14
<i>Heros</i> (<i>Cichlasoma</i>).....	420	<i>nasus</i>	19
<i>spinosus</i> , <i>Anicistrus</i>	160	<i>norwegianus</i>	22
<i>Calyclepidotus</i>	381	<i>obscurus</i>	10
<i>Centronotus</i>	281	<i>obtusus</i>	15
<i>Cyclichthys</i>	502	<i>peregrinus</i>	20
<i>Cyclopterus</i>	398	<i>pennanti</i>	19
<i>Diodon</i>	502	<i>platyodon</i>	15
<i>Eumicrotremus</i>	398	<i>porosus</i>	15
<i>Notacanthus</i>	76	<i>pristis</i>	23
<i>Platophrys</i>	217	<i>punctatus</i>	12, 14, 15
<i>Polyipnus</i>	74	<i>punctulatus</i>	12
<i>Rhinobatos</i>	24	<i>rashleighanus</i>	20
<i>Rhinobatus</i>	24	<i>rostratus</i>	19
<i>Rhomboldichthys</i>	217	<i>spathula</i>	36
<i>Schedophilopris</i>	269	<i>spinax</i>	21
<i>Squalus</i>	22	<i>spinosus</i>	22
<i>Trachinotus</i>	276	<i>squatina</i>	22
<i>spinulosa</i> , <i>Chalinura</i>	204	<i>stellaris</i>	11
<i>Malthopsis</i>	507	<i>suckleyi</i>	21
<i>spinulosus</i> , <i>Gasterosteus</i>	237	<i>tiburo</i>	10, 17
<i>Spirinchus</i>	67	<i>vulpes</i>	19
<i>thaleichthys</i>	67	<i>vulpinus</i>	18
<i>springilus</i> , <i>Leuciscus</i>	128	<i>zygæna</i>	18
<i>spixianus</i> , <i>Saurus</i>	163	<i>squamata</i> , <i>Swainia</i>	285
<i>spizii</i> , <i>Artus</i>	150	<i>Tigoma</i>	119
<i>Pimeledus</i>	150	<i>squamatus</i> , <i>Etheostoma</i> (<i>Iladropodus</i>).....	285
<i>Platysomus</i>	275	<i>squamiceps</i> , <i>Claricola</i>	292
<i>Vomer</i>	275	<i>Etheostoma</i>	292
<i>splendens</i> , <i>Beryx</i>	234	<i>squamileatus</i> , <i>Ceratichthys</i>	136
<i>Exocetus</i>	200	<i>Couesius</i>	136
<i>splendida</i> , <i>Petenia</i>	423	<i>Paralichthys</i>	224
<i>splenistum</i> , <i>Fristipoma</i>	332	<i>Squamipennes</i>	358
<i>splenistus</i> , <i>Anisotremus</i>	332	<i>squamipinnis</i> , <i>Cynoscion</i>	355
<i>spongicola</i> , <i>Evermannichthys</i>	440, 447	<i>Gerres</i>	342
<i>Garmannia</i>	446	<i>Otolithus</i>	355
<i>spongiosa</i> , <i>Helleutaa</i>	508	<i>squamosa viridis</i> , <i>Acus maxima</i>	36
<i>Spratelloides bryoporos</i>	41	<i>squamosus</i> , <i>Trachurus</i>	272

List of scientific names—Continued

	Page		Page
<i>squamulosus</i> , <i>Chætodon</i>	361	<i>stellatus</i> , <i>Fario</i>	56
<i>Squatina</i>	22	<i>Mustelius</i>	13
<i>angelus</i>	22	<i>Platichthys</i>	226
<i>californica</i>	22	<i>Pleuronectes</i>	226
<i>dumeril</i>	22	<i>Tetraodon</i>	497, 499
<i>fimbriata</i>	23	<i>Valencienneanus</i>	73
<i>levis</i>	22	<i>Xurel</i>	273
<i>lewis</i>	22	<i>stelleri</i> , <i>Cottus</i>	381
<i>oculata</i>	22	<i>Cyclopterus</i>	398
<i>raja</i>	23	<i>Hexagrammos</i>	373, 374
<i>squatina</i>	22	<i>Liparops</i>	398
<i>vulgaris</i>	22	<i>Myoxocephalus</i>	387
<i>Squatina</i> , <i>Squalus</i>	22	<i>Stellerina</i>	394
<i>Squatina</i>	22	<i>xyosterna</i>	394
<i>Squatinae</i>	22	<i>Stellicarens</i>	347
<i>Squatinoidei</i>	22	<i>zestocarous</i>	347
<i>squeteague</i> , <i>Labrus</i>	355	<i>Stellifer</i>	346
<i>maculatus</i> , <i>Labrus</i>	355	<i>colonensis</i>	347
<i>stabulifrons</i> , <i>Raja</i>	25	<i>ericymbia</i>	346
<i>stadhouderi</i> , <i>Serranus</i>	311	<i>illecebrosus</i>	347
<i>stagnalis</i> , <i>Salvelinus</i>	60	<i>lanceolatus</i>	346
<i>stahl</i> , <i>Auchenitius</i>	457	<i>melanochir</i>	347
<i>Monocira</i>	344	<i>microps</i>	346
<i>stanleyi</i> , <i>Coregonus</i>	65	<i>mindli</i>	347
<i>Prosoptium</i>	65	<i>oscitans</i>	347
<i>starksi</i> , <i>Menidia</i>	247	<i>stellifer</i>	346
<i>Siphostoma</i>	242	<i>zestocarous</i>	347
<i>Stolephorus</i>	48	<i>stellifer</i> , <i>Bodianus</i>	346
<i>Syngnathus</i>	242	<i>stellifer</i>	346
<i>Starksia</i>	456	<i>Xenisma</i>	179
<i>creminobates</i>	456	<i>stellifera</i> , <i>Corvina</i>	346
<i>holderi</i>	450	<i>Xenisma</i>	179
<i>starri</i> , <i>Pomadasys</i>	334	<i>stelliferoidea</i> , <i>Bassogigas</i>	482
<i>Stathmonotus</i>	465	<i>Neobythites</i>	482
<i>hemphillii</i>	465	<i>Stelliferus</i>	346
<i>teki</i>	466	<i>illecebrosus</i>	347
<i>stearnsi</i> , <i>Bleennius</i>	460	<i>zestocarous</i>	347
<i>Colotrigla</i>	406	<i>stellio</i> , <i>Perca</i>	310
<i>stearnsi</i> , <i>Corvina</i>	348	<i>Rhinobatos</i>	23
<i>Lutjanus</i>	325	<i>Rhinobatus</i>	23
<i>Prionotus</i>	406	<i>stellulata</i> , <i>Raja</i>	26
<i>Roncador</i>	348	<i>Stenesthes</i>	336
<i>Scorpaena</i>	371	(<i>Stenobrachius</i>) <i>leucopsarum</i> , <i>Myctophum</i>	168
<i>Seriola</i>	279	<i>Stenodus</i>	61
<i>Stegastes</i>	415	<i>mackenzii</i>	61
<i>bairdii</i>	415	<i>Stenometopus</i>	498
<i>chrysurus</i>	415	<i>Stenotoma</i>	336
<i>dorsalis</i>	418	<i>Stenotomus</i>	336
<i>imbricatus</i>	415	<i>caprinus</i>	336
<i>niveatus</i>	416	<i>chrysops</i>	336
<i>Steindachnerilia</i>	203	<i>versicolor</i>	336
<i>steindachneri</i> , <i>Argo</i>	265	<i>Stephanoberyidae</i>	222
<i>Arius</i>	151	<i>Stephanoberyx</i>	232
<i>Cithlasoma</i>	419	<i>gilli</i>	232
<i>Diasbasis</i>	330	<i>mona</i>	232
<i>Hæmulon</i>	330	<i>Stephanolepis</i>	494
<i>Ophibolennius</i>	464	<i>stephanophrys</i> , <i>Prionotus</i>	407
<i>Parapetenia</i>	419	<i>Stereolepis</i>	308
<i>Rhinoptera</i>	31	<i>californicus</i>	308
<i>Tachysurus</i>	151	<i>gigas</i>	308
<i>Thalassoma</i>	429	<i>Sterletta</i>	33
<i>Steindachneria</i>	203	<i>Sterletus</i>	33, 34
<i>argentea</i>	203	<i>sterletus</i> , <i>A verruncosus</i>	395
<i>Steindachneridion</i>	203	<i>Ceratichthys</i>	139
<i>Steinegeria</i>	268	<i>Maorhybopsis</i>	139
<i>rubescens</i>	268	<i>Sternarchus</i>	101
<i>Steinegerigeria</i>	268	<i>rostratus</i>	101
<i>stegidolepis</i> , <i>Macrourus</i>	205	<i>virescens</i>	101
<i>Macrourus</i> (<i>Macrourus</i>)	205	<i>Sternias</i>	380
<i>Stegidonotus</i>	379	<i>xenostethus</i>	380
<i>latifrons</i>	379	<i>Sternicla</i> , <i>Gasteropelecus</i>	100
<i>Stegis</i>	385	<i>Sternoptichidae</i>	73
<i>vulsum</i>	385	<i>Sternoptix</i>	73
<i>Stegistrops</i>	379	<i>diaphana</i>	73
<i>beringiana</i>	379	<i>obscura</i>	73
<i>Stegistrum</i> <i>beringianum</i>	379	<i>Sternoptix</i>	73
<i>Stelliferes</i> , <i>les</i>	346	<i>gardenii</i>	266
<i>stellaris</i> , <i>Squalus</i>	11	<i>mediterraneus</i>	74
<i>stellata</i> , <i>Halieutaea</i>	507	<i>obscura</i>	73
<i>stellatus</i> , <i>Acipenser</i>	34	<i>olfersi</i>	74
<i>Apogonichthys</i>	298	<i>Sternopygus</i>	101
<i>Astrapogon</i>	298	<i>dariensis</i>	101
<i>Bodianus</i>	309	<i>humboldtii</i>	101
<i>Caranx</i>	273	<i>lineatus</i>	101
<i>Cephalopholis</i>	309	<i>tumifrons</i>	101

List of scientific names—Continued

	Page		Page
<i>Sternotremia</i>	215	<i>stomias</i> , <i>Atheresthes</i>	221
<i>isolepis</i>	215	<i>Bathyroctes</i>	52
<i>stevensi</i> , <i>Thaileichthys</i>	68	<i>Chasmistes</i>	108
<i>Stichaeidae</i>	469	<i>Narcetes</i>	52
<i>Stichæus</i>	469	<i>Platysomichthys</i>	221
<i>punctatus</i>	469	<i>Salmo</i>	57
<i>stigma</i> , <i>Gymnelis</i>	477	<i>Trisopteris</i>	314
<i>Ophidium</i>	477	<i>Stomatidae</i>	69
<i>stigmæa</i> , <i>Ulcocentra</i>	287	<i>Stomiatoidei</i>	69
<i>stigmæum</i> , <i>Boleosoma</i>	287	<i>Stomocatus</i>	105
<i>stigmæus</i> , <i>Citharichthys</i>	219	<i>Stomosom bilinearis</i>	214
<i>stigmatodus</i> , <i>Acynnodontophis</i>	94	<i>stonei</i> , <i>Hydrophox</i>	127
<i>stigmaticus</i> , <i>Acyrachthys</i>	137	<i>Notropis</i>	127
<i>Gobionellus</i>	441	<i>Salmo</i>	58
<i>stigmatistum</i> , <i>Lepophidium</i>	485	<i>gaiderneri</i>	58
<i>Leptophidium</i>	485	<i>storeri</i> , <i>Acipenser</i>	34
<i>stigmatius</i> , <i>Sigmops</i>	72	<i>Aledon</i>	503
<i>Smaragdus</i>	441	<i>Cetomimus</i>	172
<i>stigmatura</i> , <i>Algansea</i>	116	<i>Leuciscus</i>	117
<i>Bollmannia</i>	443	<i>Scomberesox</i>	197
<i>Erogala</i>	129	<i>storeria</i> , <i>Cichla</i>	304
<i>stigmaturus</i> , <i>Gobius</i>	439	<i>storieranus</i> , <i>Erinemus</i>	137
<i>Photogenis</i>	129	<i>Rutilus</i>	137
<i>Rhinogobius</i>	439	<i>stormi</i> , <i>Cephalocassis</i>	149
<i>stilbe</i> , <i>Zalocys</i>	277	<i>stoutii</i> , <i>Bdellostoma</i>	8
<i>Stilbiscus</i>	89	<i>Polistotrema</i>	8
<i>edwardsi</i>	89	<i>strabo</i> , <i>Icelinus</i>	376
<i>stilbius</i> , <i>Leuroglossus</i>	68	<i>Ophioscion</i>	347
<i>Nototropis</i>	125	<i>straminea</i> , <i>Hybognathus delicosa</i>	134
<i>Notropis</i>	125	<i>stramineus</i> , <i>Hybognathus</i>	133
<i>stimpsoni</i> , <i>Triglopsis</i>	389	<i>Hypopsis</i>	134
<i>stipes</i> , <i>Atherina</i>	246	<i>Strephon</i>	159
<i>Heptseta</i>	246	<i>striata</i> , <i>Alausa</i>	43
<i>Joturus</i>	254, 255	<i>Argentina</i>	68
<i>stirurus</i> , <i>Chloroscombrus</i>	276	<i>Perca</i>	331
<i>Stizostedion</i>	282	<i>striato</i> , <i>Perca marina capite</i>	330
<i>glacum</i>	282	<i>striatulus</i> , <i>Brycon</i>	98
<i>vitreum</i>	282	<i>Chalcinopsis</i>	98
<i>Stizostethium</i>	282	<i>striatum</i> , <i>Bathystoma</i>	331
<i>Stoasodon</i>	30	<i>Pygidium</i>	159
<i>narinari</i>	30	<i>striatus</i> , <i>Acanthurus</i>	363
<i>Stolephorus</i>	47	<i>Anthias</i>	311, 318
<i>astilbe</i>	49	<i>Blennius</i>	462
<i>branchiomelas</i>	49	<i>Bodianus</i>	325
<i>brevirostrum</i>	50	<i>Centropristes</i>	317
<i>cultratus</i>	48	<i>Chætodon</i>	359
<i>curtus</i>	48	<i>Ctenochætus</i>	363
<i>epsetus</i>	47	<i>Diodon maculato</i>	502
<i>eurystole</i>	48	<i>Epinephelus</i>	311, 318
<i>exiguus</i>	47	<i>Holocentrus</i>	235
<i>garmani</i>	51	<i>Hypsoblennius</i>	462
<i>giberti</i>	51	<i>Labrus</i>	317
<i>gulanensis</i>	49	<i>Luciocharax</i>	100
<i>hiulcus</i>	48	<i>Monacanthus</i>	404
<i>ischanus</i>	48	<i>Roccus</i>	307
<i>lucidus</i>	48	<i>Salmo</i>	54
<i>lyolepis</i>	49	<i>strigata</i> , <i>Trigla</i>	407
<i>mlarchus</i>	47	<i>strigatum</i> , <i>Sparisoma</i>	433
<i>mundulus</i>	50	<i>strigatus</i> , <i>Attenuarius</i>	505
<i>naso</i>	49	<i>Euctenogobius</i>	441
<i>opercularis</i>	51	<i>Gobionellus</i>	441
<i>perfasciatus</i>	47	<i>Holacanthus</i>	361
<i>perthecatus</i>	47	<i>Prionotus</i>	407
<i>rastralis</i>	50	<i>Scarus</i>	433
<i>robertsi</i>	49	<i>Xiphophorus</i>	193
<i>rostralis</i>	47	<i>helleri</i>	193
<i>scofieldi</i>	49	<i>strigillata</i> , <i>Loricaria</i>	162
<i>spinifer</i>	47	<i>strigosus</i> , <i>Acanthurus</i>	363
<i>starksii</i>	48	<i>Anarhichas</i>	472
<i>surlamensis</i>	50	<i>Stromæi</i> , <i>Zeus</i>	217
<i>stolifera</i> , <i>Clupea</i>	44	<i>Stromatidae</i>	266
<i>Dussumieriæ</i>	45, 46	<i>Stromateiformes</i>	266
<i>Jenkinsia</i>	46	<i>Stromateus cryptosus</i>	267
<i>Lile</i>	44	<i>longipinnis</i>	266
<i>stolzmanni</i> , <i>Belone</i>	196	<i>medius</i>	267
<i>Ericson</i>	356	<i>palometæ</i>	267
<i>Otolithus</i>	356	<i>paru</i>	266
<i>Strongylura</i>	196	<i>triacanthus</i>	267
<i>stomata</i> , <i>Hippoglossina</i>	223	<i>strombi</i> , <i>Apogonichthys</i>	226
<i>Stomias</i>	69	<i>Strongylura</i>	195
<i>affinis</i>	69	<i>angusticeps</i>	196
<i>atriventer</i>	69	<i>argalus</i>	196
<i>colubrinus</i>	69	<i>caudimaculata</i>	196
<i>ferox</i>	69	<i>diplostena</i>	196
<i>hexagonatus</i>	69	<i>euryops</i>	196

List of scientific names—Continued

	Page		Page
<i>Strongylura exilis</i>	196	superbus, <i>Scarus</i>	434
<i>guianensis</i>	196	supercliosus, <i>Hexagrammos</i>	374
<i>merina</i>	196	<i>Hyborhynchus</i>	145
<i>microps</i>	196	<i>Labrax</i>	373
<i>notata</i>	195	supernas, <i>Leucichthys</i>	63
<i>pterura</i>	195	surinamensis, <i>Anisotremus</i>	332
<i>scapularis</i>	195	<i>Batrachoides</i>	486
<i>stolzmanni</i>	195	<i>Batrachus</i>	486
<i>timucu</i>	195	<i>Galeichthys</i>	150
<i>strongylura</i> , <i>Belone</i>	195	<i>Hexanematichthys</i>	150
<i>strumosus</i> , <i>Goblesox</i>	489	<i>Holocentrus</i>	323
<i>Sturio</i>	33, 34	<i>Lobotes</i>	323
<i>vulgaris</i>	33	<i>Lutjanus</i>	332
<i>sturlo</i> , <i>Acipenser</i>	33, 34	<i>Pleuronectes</i>	217
<i>Sturisoma</i>	161	<i>Poecilia</i>	190
<i>rostrata</i>	161	<i>Sparus</i>	424
<i>Stygicola</i>	479	<i>Stolephorus</i>	50
<i>dentatus</i>	479	<i>susane</i> , <i>Boleosoma</i>	288
<i>Stygogenes</i>	159	<i>spectus</i> , <i>Phycocetes</i>	478
<i>humboldti</i>	159	<i>sutor</i> , <i>Blepharis</i>	274
<i>Stylephoridae</i>	216	<i>swaini</i> , <i>Poecilichthys</i>	201
<i>Stylephorus</i>	216	<i>Swainia</i>	285
<i>chordatus</i>	216	<i>squamata</i>	285
<i>Styletor</i>	88	<i>swampina</i> , <i>Hydrargyre</i>	175
<i>stylifer</i> , <i>Hippocampus</i>	244	<i>swanii</i> , <i>Bothragonus</i>	396
<i>Stylocephalidae</i>	71	<i>Hypsagonus</i>	396
<i>Stylophthalmus</i>	71	<i>swannanoa</i> , <i>Etheostoma</i>	289
<i>paradoxus</i>	71	<i>Nanostoma</i>	289
<i>Stypondon</i>	113	<i>swifti</i> , <i>Eosebastes</i>	367
<i>signifer</i>	113	<i>Sebastodes</i>	367
<i>suavis</i> , <i>Cyprinella</i>	131	<i>Syacium</i>	218
<i>subaequalis</i> , <i>Corvina</i>	345	<i>latifrons</i>	218
<i>Vacuogn</i>	345	<i>micrurum</i>	218
<i>subarcuata</i> , <i>Zygæna</i>	18	<i>papillosum</i>	218
<i>subarcuatum</i> , <i>Hæmulon</i>	330	<i>Symbranchia</i>	76
<i>subarmatus</i> , <i>Acanthurus</i>	362	<i>Symbranchus</i>	76
<i>subbifrenatus</i> , <i>Rhypticus</i>	316	<i>symmetricus</i> , <i>Apomotis</i>	299
<i>subbifurcata</i> , <i>Ulvaria</i>	470	<i>Caranx</i>	270
<i>subbifurcatus</i> , <i>Pholis</i>	470	<i>Hesperoleucus</i>	121
<i>subcærulea</i> , <i>Amla</i>	38	<i>Lepomis</i>	209
<i>subditus</i> , <i>Hesperiroleucus</i>	122	<i>Pogonichthys</i>	121
<i>subfuscus</i> , <i>Labrus</i>	424	<i>Salmo</i>	50
<i>subfurcatus</i> , <i>Bermudichthys</i>	430	<i>Trachurus</i>	270
<i>subfilarius</i> , <i>Centropristes</i>	320	<i>Symmeturus</i>	116
<i>subniger</i> , <i>Chiassodon</i>	452	<i>Symbranchus vittatus</i>	76
<i>suborbitalis</i> , <i>Holocentrum</i>	236	<i>Symphurus</i>	230
<i>suborbitalis</i> , <i>Adioryx</i>	236	<i>atramentatus</i>	231
<i>Macrurus</i>	203	<i>atricaudus</i>	231
<i>Nematonurus</i>	203	<i>diomedianus</i>	231
<i>Plectromus</i>	233	<i>elongatus</i>	231
<i>subosseus</i> , <i>Nannocampus</i>	244	<i>fasciolaris</i>	231
<i>subterranea</i> , <i>Lucifuga</i>	479	<i>lei</i>	231
<i>subterranea</i> , <i>Lucifuga</i>	479	<i>marginalis</i>	231
<i>Typhlichthys</i>	104, 195	<i>nigrescens</i>	230
<i>subtruncata</i> , <i>Belone</i>	195	<i>piger</i>	230
<i>subtruncatus</i> , <i>Homopriion</i>	345	<i>plagiusa</i>	231
<i>subulatus</i> , <i>Orcynus</i>	280	<i>plagusia</i>	231
<i>Thunnus</i>	280	<i>pusillus</i>	231
<i>succincta</i> , <i>Seriola</i>	278	<i>williamsi</i>	231
<i>suecica</i> , <i>Cyprinus</i>	109	<i>synagris</i> , <i>Lutjanus</i>	3, 327
<i>Erimyzon</i>	109	<i>synagrus</i> , <i>Sparus</i>	327
<i>suckleyi</i> , <i>Spinax (Acanthias)</i>	21	<i>Synanceia cervus</i>	381
<i>Squalus</i>	21	<i>Synanceia curvifrons</i>	381
<i>sucklili</i> , <i>Catostomus</i>	105	<i>Synaphobranchidae</i>	78
<i>sucubti</i> , <i>Rivulus chucunaque</i>	180	<i>Synaphobranchus</i>	78
<i>Sudidæ</i>	165	<i>bathyplus</i>	78
<i>Suds</i>	165	<i>kaupii</i>	78
<i>hyalina</i>	165	<i>pinnatus</i>	78
<i>intermedius</i>	165	<i>Synapteretmus</i>	166
<i>ringeni</i>	165	<i>Synbranchidae</i>	76
<i>suensonii</i> , <i>Chlorhinus</i>	93	<i>Synbranchus</i>	76
<i>suerii</i> , <i>Coryphaena</i>	264	<i>fuliginosus</i>	76
<i>sueuri</i> , <i>Meletta</i>	42	<i>immaculatus</i>	76
<i>Sufflamen</i>	492	<i>marinoratus</i>	76
<i>verres</i>	492	<i>transversalis</i>	76
<i>Sufflamen</i> , <i>Balistes</i>	492	<i>Synchiridae</i>	393
<i>Canthidermis</i>	492	<i>Synchirus</i>	393
<i>sugillatus</i> , <i>Ophisurus</i>	88	<i>gilli</i>	393
<i>Sullius</i>	425	<i>Synechoglanis</i>	151
<i>sullius</i> , <i>Lachnolaimus</i>	425	<i>beedlei</i>	151, 162
<i>sujei</i> , <i>Muraenoides</i>	407	<i>Synentognathus</i>	195
<i>sulcatus</i> , <i>Coryphænoides</i>	207	<i>Syngnathidae</i>	240
<i>Herpetoichthys</i>	87	<i>Syngnathus</i>	240
<i>Trachonurus</i>	207	<i>abboti</i>	240
<i>sunichrasti</i> , <i>Citharichthys</i>	219		
<i>sundiacus</i> , <i>Bagrus</i>	149		

List of scientific names—Continued

	Page		Page
<i>Syngnathus acus</i>	240	<i>tabacarius</i> , <i>Centropristes</i>	310
<i>sequareus</i>	243	<i>tacheté</i> , <i>le Diodon</i>	501
<i>affinis</i>	242	<i>tania</i> , <i>Bassozetus</i>	481
<i>arctus</i>	242	<i>Bathyonus</i>	481
<i>arundinaceus</i>	240	<i>Blennius</i>	468
<i>ascendens</i>	241	<i>Clinostomus</i>	119
<i>auilscus</i>	240	<i>temata</i> , <i>Atherina</i>	246
<i>bairdianus</i>	242	<i>templatum</i> , <i>Hæmulon</i>	332
<i>brachycephalus</i>	241	<i>temniatus</i> , <i>Anisotremus</i>	332
<i>barbare</i>	240	<i>Chirocentrodon</i>	45
<i>brachycephalus</i>	241	<i>Evoxymetopon</i>	262
<i>bervicaudus</i>	241	<i>Pseudoxiphophorus bimaculatus</i>	188
<i>brevirostris</i>	240	<i>Tropidichthys</i>	500
<i>californiensis</i>	240	<i>Tænoconger</i>	82
<i>carinatus</i>	240	<i>digueti</i>	82
<i>crinigerum</i>	241	<i>Tæniosomps</i> , <i>Cephalopholis</i>	309
<i>dekayi</i>	242	<i>Serranus</i>	309
<i>dendriticus</i>	242	<i>Tæniopterus</i> , <i>Balistes</i>	491
<i>dimidiatus</i>	240	<i>Cottus</i>	388
<i>elcapitanense</i>	242	<i>Perissas</i>	220
<i>eicapitensis</i>	242	<i>Platophrys</i>	220
<i>eluens</i>	241	<i>Pseudoscarus</i>	433
<i>ethon</i>	241	<i>Scarus</i>	433
<i>fasciatus</i>	242	<i>Tæniomembras</i>	246
<i>flavulatus</i>	241	<i>Tæniophis</i>	91
<i>flavirostris</i>	241	<i>westphali</i>	91
<i>floridæ</i>	241	<i>Tæniomomi</i>	216
<i>fuscus</i>	242	<i>Tæniotoca</i>	411
<i>griseolineatus</i>	240	<i>lateralis</i>	411
<i>hippocampus</i>	244	<i>Tenuira</i> , <i>Safole</i>	305
<i>jonesi</i>	241	<i>Tenuirus</i> , <i>Dules</i>	305
<i>leptorhynchus</i>	240	<i>Labrus</i>	430
<i>linea</i>	241	<i>Poecilurichthys</i>	97
<i>louisianæ</i>	242	<i>Tachisurus</i> , <i>gulosus</i>	151
<i>mackayi</i>	241	<i>jordani</i>	150
<i>marmoratus</i>	241	<i>lentiginosus</i>	149
<i>milbertianus</i>	242	<i>longicephalus</i>	150
<i>mindii</i>	242	<i>Tachysurus</i>	150
<i>ophidion</i>	243	<i>chinensis</i>	150
<i>peckianus</i>	242	<i>emmeline</i>	151
<i>peckii</i>	242	<i>evermanni</i>	151
<i>pelagicus</i>	210	<i>Ilampus</i>	151
<i>picturatus</i>	242	<i>steindachneri</i>	151
<i>poeyi</i>	241	<i>tahoensis</i> , <i>Catostomus</i>	10 ⁺
<i>robertsi</i>	241	<i>Salmo</i>	5
<i>rousseau</i>	241	<i>clarkii</i>	57
<i>sibbaldi</i>	243	<i>talasica</i> , <i>Gobius</i>	442
<i>sinaloae</i>	241	<i>Talismania</i>	53
<i>spicifer</i>	243	<i>equatoris</i>	53
<i>starksii</i>	242	<i>antillarum</i>	53
<i>tenuis</i>	241	<i>tang</i> , <i>Mugil</i>	253
<i>torrel</i>	242	<i>tanneri</i> , <i>Hyperchoristus</i>	70
<i>tweddleii</i>	242	<i>tapidisma</i> , <i>Salmo</i>	55
<i>viridescens</i>	242	<i>Taractes</i>	265
<i>Synodontidae</i>		<i>asper</i>	265
<i>Synodus</i>	163	<i>longipinnis</i>	265
<i>acutus</i>	163	<i>saussurii</i>	265
<i>argenteus</i>	164	<i>Taranichthys</i>	373
<i>evermanni</i>	40	<i>cavifrons</i>	375
<i>foetens</i>	164	<i>filamentosus</i>	376
<i>intermedius</i>	164	<i>tenuis</i>	376
<i>Jenkinsi</i>	163	<i>tarascorum</i> , <i>Algansea</i>	116
<i>lacerta</i>	163	<i>Tarletonbeania</i>	170
<i>lacertinus</i>	164	<i>cranularis</i>	170
<i>lucioceps</i>	163	<i>tenua</i>	170
<i>malabaricus</i>	99	<i>Tarpon</i>	39
<i>poeyi</i>	164	<i>atlanticus</i>	39
<i>saurus</i>	163	<i>Tarpum</i>	39
<i>scituliceps</i>	164	<i>tarsodes</i> , <i>Bryostemma</i>	466
<i>simulans</i>	163	<i>tassellatum</i> , <i>Boleosoma</i>	287
<i>synodus</i>	163	<i>tassellatus</i> , <i>Labrus</i>	424
<i>Synodus</i> . <i>Esox</i>	163	<i>tau</i> , <i>Batrachoides</i>	486
<i>Synodus</i>	163	<i>Gadus</i>	487
<i>Sypterus</i>	210	<i>Opsanus</i>	487
<i>Syrtices</i>	242	<i>pardus</i> , <i>Batrachus</i>	487
<i>affinis</i>	242	<i>Tauridea</i>	383, 386
<i>bairdianus</i>	242	<i>taurinus</i> , <i>Aspicottus</i>	381
<i>exilis</i>	243	<i>Enophrrys</i>	381
<i>fuscus</i>	242	<i>taurocephalus</i> , <i>Alburnops</i>	145
<i>louisianæ</i>	242	<i>taurus</i> , <i>Abudedefduf</i>	415
<i>scovelli</i>	242	<i>Carcharias</i>	15
<i>Syraxis</i>	27	<i>Carpoides</i>	102
<i>Syrrhina</i>	23	<i>Glyphidodon</i>	415
<i>syrtentium</i> , <i>Argentina</i>	68	<i>Tautoga</i>	424
<i>tabacaria</i> , <i>Fistularia</i>	245	<i>ala</i>	424
<i>Prionodes</i>	319	<i>cerulea</i>	424
		<i>fusca</i>	424

List of scientific names—Continued

	Page		Page
Tautoga niger	424	testudineus, <i>Arothron</i>	499
nigra	424	<i>Chelichthys</i>	499
onitis	424	<i>Tetraodon</i>	499
rubens	424	Tetrabranchus	76
tautoga, <i>Labrus</i>	424	tetracantha cincta, <i>Parapetenia</i>	419
Tautogolabrus	424	<i>grisea</i> , <i>Parapetenia</i>	419
adspersus	424	<i>lata</i> , <i>Parapetenia</i>	419
taylori, <i>Arius</i>	151	<i>Parapetenia</i>	419
Asematichthys	300	<i>torralbasii</i> , <i>Parapetenia</i>	419
Ophidium	485	tetracanthus, <i>Centrarchus</i>	419
Otopinidium	485	<i>cinctus</i> , <i>Heros</i>	419
tepaæ, <i>Cieblasoma</i>	417	<i>griseus</i> , <i>Heros</i>	419
Theraps	417	<i>lotus</i> , <i>Heros</i>	419
Tectospondylæ	21	<i>Sparus</i>	325
Tekla	456	<i>torralbasii</i> , <i>Heros</i>	419
tokla, <i>Stathmonotus</i>	466	Tetragonopterus	369
Teleostol	39	<i>genes</i>	96
telescopium, <i>Pomatomus</i>	206	<i>brevimanus</i>	96
Telescops	206	<i>fischeri</i>	97
telescopus arcansanus, <i>Notropis</i>	124	<i>fulgens</i>	96
telescopus, <i>Netropis</i>	124	(<i>Hemibrycon</i>) <i>polyodon</i>	97
Photogenis	124	<i>humilis</i>	96
Telipomis	298	<i>macrophthalmus</i>	97
Temiculina	110, 120	<i>mexicanus</i>	96
temminckii, <i>Carcharias</i>	15	<i>microphthalmus</i>	96
temminckii, <i>Ruvettus</i>	261	<i>nitidus</i>	96
Teminista	380	<i>corstedi</i>	96
Tenmnodon	280	<i>petenensis</i>	97
tempel, <i>Pomadasys</i>	334	<i>scabripinnis</i>	97
tenebrosus, <i>Alepocephalus</i>	52	Tetragonuridae	269
<i>Antennarius</i>	505	Tetragonurus	269
<i>Caranx</i>	273	<i>atlanticus</i>	269
<i>Xurel</i>	273	<i>cuvieri</i>	269
tenellus, <i>Fundulus</i>	177	<i>simplex</i>	261
<i>Hyborhynchus</i>	145	Tetranarce	27
tentabunda, <i>Trigla</i>	409	<i>californica</i>	27
tentaculata, <i>Venefica</i>	82	<i>occidentalis</i>	27
tenua, <i>Tarletonbeania</i>	170	tetranemus, <i>Extrarius</i>	139
tenui, <i>Cichlasoma</i>	419	<i>Hybopsis</i>	139
<i>Moxostoma</i>	109	Tetraodon	499
tenuicauda, <i>Macrourus</i>	206	<i>ammocryptus</i>	499
<i>Macrurus</i>	206	<i>hispidus</i>	500
tenuiculum, <i>Mycophthum</i>	170	<i>levigatus</i>	497
tenuifilis, <i>Antennarous</i>	505	<i>lineatus</i>	499
tenuirostris, <i>Anguilla</i>	77	<i>mola</i>	503
tenuis, <i>Anthias</i>	322	<i>perspicillaris</i>	500
<i>Atherinopsis</i>	251	<i>politus</i>	499
<i>Cottus</i>	384	<i>pluviatilis</i>	499
<i>Cynodontichthys</i>	180	<i>reticularis</i>	499
<i>Gadus</i>	213	<i>rostratus</i>	500
<i>Icelinus</i>	376	<i>setosus</i>	500
<i>Leurosthes</i>	251	<i>stellatus</i>	407, 499
<i>Parapetenia</i>	419	<i>testudineus</i>	499
<i>Pœclia</i>	192	<i>trichocephalus</i>	499
<i>Rivulus</i>	180	Tetraodontidae	497
<i>Syngnathus</i>	241	Tetraleurodon	9
<i>Tarandichthys</i>	370	<i>spadiceus</i>	9
<i>Uranidea</i>	384	tetrapturorum, <i>Echenels</i>	450
<i>Urophycis</i>	213	Tetrapodus amplius	263
teporatus, <i>Heros</i>	422	<i>indicus</i>	263
terculatus, <i>Gasterosteus</i>	237	<i>mazara</i>	263
tereira, <i>Macrodon</i>	99	<i>mitsukurii</i>	263
teres, <i>Alosa</i>	46	tetraspilus, <i>Upeneus</i>	344
<i>Cyprinus</i>	105	Tetrodon angusticeps	498
<i>Macrodon</i>	99	<i>annulatus</i>	499
<i>Sphagebranchus</i>	86	<i>capistratus</i>	500
Teretulus	109	(<i>Chelichthys</i>) <i>pachygaster</i>	499
<i>cervinus</i>	111	<i>curvus</i>	497
teretulus liosternus, <i>Phenacobius</i>	142	<i>fluviatilis</i>	499
teretulus, <i>Phenacobius</i>	142	<i>formosus</i>	500
tergisus, <i>Hodon</i>	40	<i>fürthii</i>	499
terre-nove, <i>Lycodes</i>	474	<i>geometricus</i>	499
<i>Scolodon</i>	14	<i>heraldi</i>	499
<i>Squalus</i> (<i>Carcharitus</i>)	14	<i>hispidus maculatus</i>	498
<i>Triglops</i>	380	<i>imperatus</i>	500
<i>ommastistius</i>	550	<i>levigatus</i>	497
terræbæ, <i>Theraps</i>	417	<i>laterna</i>	500
terræbæns, <i>Brachyrhaphis</i>	184	<i>le</i>	498
<i>Gambusia</i>	184	<i>lineolatus</i>	497
tesseletata, <i>Etheostoma</i>	200	<i>marmoratus</i>	498
<i>Plagiusia</i>	231	<i>mathematicus</i>	497
tesellatus, <i>Hadropterus</i>	259	<i>mola</i>	503
<i>Labrus</i>	424	<i>nephelus</i>	498
<i>Nothonotus</i>	290	<i>ornatus</i>	500
te tar, <i>Lepadogaster</i>	450	<i>oxyrhynchus</i>	500

List of scientific names—Continued

	Page		Page
Tetronarce pachycephalus.....	497	Thoburnia.....	107
psittacus.....	500	rhothoeca.....	108
punctatissimus.....	500	Tholichthys.....	359
punctatus.....	499	osseus.....	359
rostratus.....	500	thompsoni, Amia.....	38
spengleri.....	498	Carpoides.....	103
(Spheroïdes) eulepidotus.....	498	Gnatholepis.....	443
truncatus.....	504	Lepidosteus.....	37
turgidus.....	498	Triglopsis.....	389
Tetronarce.....	27	thomsonii, Cottunculus.....	392
tetrophtalmia, Lioglossina.....	223	Cottus.....	392
tetrophtalmus, Anableps.....	194	thoracatus, Callichthys (Holosternum).....	160
Lioglossina.....	223	Thoracosteles.....	237
Teuthis.....	362	thoreauanus, Semotilus.....	117
australis.....	362	Semotilus atromaculatus.....	117
crestonis.....	363	Thorichthys.....	422
hepatus.....	362	acutus.....	422
terana, Anguilla.....	77	affinis.....	422
Cyprinella.....	130	aureus.....	422
Eregula.....	130	callolepis.....	422
teranus, Catostomus.....	108	elliotti.....	422
Xyrauchen.....	108	helleri.....	422
texensis, Dionda.....	144	thrissa, Clupea.....	44
textilis, Salaris.....	464	thrissina, Clupea.....	44
Salarichthys.....	464	Harengula.....	44
thalassinum, Moxostoma.....	110	Thrisomimus.....	62
Nanostoma.....	289	Thronodontis.....	91
thalassinus, Cynoscion.....	355	reticulata, tesselata.....	91
Doratonotus.....	420	Thunnidae.....	259
Gobius.....	444	thunnina, Thynnus.....	258
Microgobius.....	444	Thunnus.....	259
Myloleucus.....	122	allisoni.....	260
Nothonotus.....	289	coreta.....	260
Otolithus.....	355	sallens.....	259
Ptychostomus.....	110	secundodorsalis.....	260
Thalassoma.....	428	subulatus.....	260
bifasciatum.....	429	Thymallidae.....	66
grammaticum.....	429	thymalloides, Coregonus.....	66
lucasium.....	429	Thymallus.....	66
nitidum.....	429	montanus.....	66
socorroense.....	429	ontariensis.....	66
socorroensis.....	429	signifer.....	66
steindachneri.....	429	tricolor.....	66
virens.....	429	thymalus, Salmo.....	66
Thalassophryne.....	488	Thynnus.....	259
dowi.....	488	affinis.....	258
maculosa.....	488	atlanticus.....	260
reticulata.....	488	baiteatus.....	260
Thaileichthys.....	66	brasiliensis.....	258
albatrossis.....	67	brevipinnis.....	258
pacificus.....	66	coreta.....	260
stevensi.....	66	leachianus.....	258
thaleichthys, Osmerus.....	67	pacificus.....	260
Spirinchus.....	67	pompilus.....	278
thaumasiun, Rhegma.....	322	thunnina.....	258
Thecopterus.....	392	vagrana.....	258
aleuticus.....	392	thygnus, Scomber.....	259
Theragra.....	209	Thyrsoidea concolor.....	93
chalogramma.....	209	Thyrina.....	248
fucensis.....	209	balsana.....	249
Theraps.....	417	chagresi.....	249
balteatus.....	418	crystallina.....	248
eigenmanni.....	417	evermanni.....	248
frontalis.....	417	guatemalensis.....	249
gadovii.....	417	gujia.....	248
gibbiceps.....	417	meeki.....	249
irregularis.....	417	sardina.....	249
lentiginosus.....	417	Thyrinops.....	249
milleri.....	417	pachylepis.....	249
mojarra.....	417	Thyris.....	220
nebulifer.....	417	pellucidus.....	220
nicaraguensis.....	417	Thyrsicus.....	379
nigritus.....	417	anoplus.....	379
teape.....	417	Thyrosidea maculipinnis.....	92
terrabae.....	417	thyrsitoides, Lemnisoma.....	261
zonatus.....	417	militaris.....	92
thermals, Poecilia.....	192	Thyrsites acanthoderma.....	261
thermophilus, Richardsonius.....	118	niger.....	261
Therobromus.....	68	scholaris.....	261
callorrhini.....	68	Thyristops violaceus.....	260
theta, Diaphus.....	168	Thysanocara.....	160
thoburni, Alcides.....	376	spinosa.....	160
Mugil.....	263	Tiaroga.....	143
Paricelinus.....	376	cobitis.....	143
Querimana.....	253		

CHECK LIST OF FISHES

661

List of scientific names—Continued

	Page		Page
tiburo, <i>Reniceps</i>	17	torsk, <i>Gadus</i>	214
Squalus.....	17	tortuga, <i>Ctenogobius</i>	439
Zygæna.....	17	Holocentrus.....	236
Tiburón.....	17	Rhinogobius.....	439
Tigoma.....	119	torvus, <i>Cottunculus</i>	392
atraria.....	119	Monomitopus.....	480
bicolor.....	120	tota, <i>Perca</i>	310
crassa.....	120	toweri, <i>Goodea</i>	184
egregia.....	118	townsendi, <i>Angelichthys</i>	361
gibbosa.....	119	Gobius.....	444
gracilis.....	119	Mystophum.....	167
humboldtii.....	110	Nyctimaster.....	167
intermedia.....	119	toxotes, <i>Rhachochilus</i>	411
lineata.....	119	Toxus.....	188
nigrascens.....	120	cubensis.....	188
orcuttii.....	120	riddlei.....	188
pulchella.....	120	Trachycorystes.....	159
pulchra.....	120	amblops.....	159
pulverentula.....	120	typus.....	159
purpurea.....	119	Trachichthyidae.....	234
squamata.....	119	Trachichthys pretiosus.....	234
rhinichthyoïdes.....	141	Trachidermis richardsoni.....	383
tigrina, <i>Ichthypophis</i>	94	Trachinocephalus.....	163
tigrinus, <i>Galeocerdo</i>	14	myops.....	163
Holocentrus.....	319	Trachinotus.....	276
Myrichthys.....	85	argenteus.....	277
Prionodes.....	319	bailloni.....	276
tigris, <i>Antennarius</i>	508	carolinus.....	277
Carcharias.....	19	culveri.....	277
Chironectes.....	506	cupreus.....	277
Isurus.....	19	falcatus.....	276
Mycteroperca.....	315	fuscus.....	276
Serranus.....	315	goodell.....	277
tilapia, <i>Hemilepidotus</i>	381	kennedyi.....	277
timagamensis, <i>Salvelinus</i>	60	paloma.....	277
timiel, <i>Raja</i>	27	palometta.....	276
timpanognathus, <i>Hybopsis</i>	119	pampanus.....	276
Timucu.....	105, 199	rhodopus.....	276
timucu, <i>Belone</i>	195	spiniferous.....	276
Esox.....	195	Trachinus adscensionis.....	310
Strongylura.....	195	cirrhosus.....	382
tinca, <i>Serranus</i>	315	gasteropeteus.....	357
tincella, <i>Algansea</i>	115	osbecki.....	310
Leuciscus.....	115	punctatus.....	310
tippecanoe, <i>Etheostoma</i>	292	trichodon.....	357
Nivicola.....	292	Trachipteridae.....	216
Tirus.....	163	Trachipterus.....	216
marmoratus.....	163	arcticus.....	216
tlahuacencis, <i>Evarra</i>	142	tex-salmonorum.....	216
Todichthys.....	410	trachyurus.....	216
agassizii.....	410	Trachonurus.....	207
toe-roe, <i>Otolithus</i>	354	silicatus.....	207
toma, <i>Salmo</i>	59	trachura, <i>Gasterosteus</i>	237
tomcod, <i>Gadus</i>	210	Raja.....	26
Microgadus.....	210	Trachurops.....	270
tomcodus fuscus, <i>Gadus</i>	210	brachychira.....	271
luteus, <i>Gadus</i>	210	brachychir.....	271
mixtus, <i>Gadus</i>	210	crumenophthalmus.....	271
pruinosus, <i>Gadus</i>	210	Trachurus.....	270
Tomicodon.....	490	alicolus.....	279
Tomocichla.....	423	boops.....	272
geddesi.....	423	europeus.....	270
underwoodi.....	423	fasciatus.....	279
topeka, <i>Codoma</i>	143	japonicus.....	270
Notrops.....	143	lathami.....	270
Torpedinidae.....	27	linnei.....	270
torpedinus, <i>Trygonobatus</i>	29	squamosus.....	272
Torpedo.....	27	symmetricus.....	270
bancrofti.....	27	trachurus.....	270
brasiliensis.....	27	Trachurus, <i>Cottus</i>	381
californica.....	27	japonicus, <i>Caranx</i>	270
nobiliana.....	27	Scomber.....	270
occidentalis.....	27	Trachurus.....	276
pictus.....	27	Trachynotus fasciatus.....	276
torquatus, <i>Labrus bifasciatus</i>	429	glacoides.....	276
torralbasi, <i>Olaridichthys</i>	188	nasutus.....	276
Heros tetracanthus.....	188	rhodopus.....	276
Parapetenia tetracantha.....	419	trachypoma, <i>Myripristis</i>	234
torrei, <i>Dasyatis</i>	419	trachypomus, <i>Ostichthys</i>	234
Pastinachus.....	28	Trachypterus.....	216
Siphostoma.....	242	bogmarus.....	216
Syngnathus.....	242	rex-salmonorum.....	216
Torrentaria.....	203	seleniris.....	216
fasciata.....	203	vogmarus.....	216
sagitta.....	203	Trachyrincus.....	203
torridus, <i>Lutjanus</i>	326	helolepis.....	203

List of scientific names—Continued

	Page		Page
<i>trachyrincus</i> , <i>Lepidolepsus</i>	203	<i>tridentatus</i> , <i>Entosphenus</i>	10
<i>trachyurus</i> , <i>Trachipterus</i>	216	<i>Petromyzon</i>	10
<i>tractus</i> , <i>Acanthurus</i>	363	<i>tridentiger</i> , <i>Allogambusia</i>	188
<i>trahira</i> , <i>Erythrinus</i>	99	<i>Gambusia</i>	188
<i>Macrodon</i>	99	<i>tridigitatus</i> , <i>Dactyloscopus</i>	454
<i>tranquebar</i> , <i>Aspidophoroides</i>	397	<i>Trifasciatus</i>	180
<i>translucens</i> , <i>Gobius</i>	440	<i>felicianus</i>	181
<i>transmontanus</i> , <i>Columbia</i>	215	<i>riverendi</i>	181
<i>transmontanus</i> , <i>Asipenser</i>	34	<i>trifasciata</i> , <i>Hydrargyra</i>	176
<i>Rhinichthys</i>	139	<i>Lepomis</i>	298
<i>transversalis</i> , <i>Synbranchus</i>	76	<i>trifurca</i> , <i>Perca</i>	318
<i>transversum</i> , <i>Poecilosoma</i>	201	<i>Trigla carolina</i>	407, 408
<i>traski</i> , <i>Hystericarpus</i>	412	<i>evolans</i>	407
<i>Trematopsis</i>	503	<i>fasciata</i>	409
<i>willughbei</i>	503	<i>palmipes</i>	407
<i>willughbei</i>	503	<i>strigata</i>	407
<i>tremebundus</i> , <i>Elassodiscus</i>	402	<i>tentabunda</i>	409
<i>Triacanthidae</i>	401	<i>volitans</i>	408, 409
<i>triacanthus</i> , <i>Argyriosus</i>	275	<i>Triglidae</i>	406
<i>Nauclerus</i>	278	<i>Triglops</i>	379
<i>Poronotus</i>	267	<i>beani</i>	380
<i>Stromateus</i>	267	<i>mctopias</i>	380
<i>Xeneretmus</i>	396	<i>ommatistius</i>	379
<i>Xenochirius</i>	396	<i>terre-novae</i>	380
<i>Tricaris</i>	14	<i>pingeli</i>	379
<i>tricaris</i> , <i>Orthoplias</i>	301, 302	<i>pleurostictus</i>	379
<i>Trilepidodon</i>	14	<i>scepticus</i>	380
<i>obesus</i>	14	<i>terra-novae</i>	380
<i>triagramma</i> , <i>Heros</i>	418	<i>xenostethus</i>	380
<i>Triakis</i>	14	<i>Triglopsis</i>	389
<i>scyllium</i>	14	<i>ontaricensis</i>	389
<i>semifasciatum</i>	14	<i>simpsoni</i>	389
<i>triangularis</i> , <i>Balistes</i>	402	<i>Triglopsis thompsoni</i>	389
<i>triangulum</i> , <i>Chatodon</i>	359	<i>Trigonobatus</i>	28
<i>Lutjanus</i>	348	<i>vulgaris</i>	28
<i>Tribranchus</i>	77	<i>Trigonophallus</i>	187
<i>tribulus</i> , <i>Prionotus</i>	408	<i>punctifer</i>	187
<i>Trichidion</i>	256	<i>Trigonorhinus alveatus</i>	24
<i>octofilis</i>	256	<i>trigonus</i> , <i>Lactophrys</i>	497
<i>opercularis</i>	256	<i>Ostracion</i>	497
<i>trichopterus</i> , <i>Pteraclis</i>	206	<i>trigrammus</i> , <i>Chirus</i>	374
<i>Trichuridae</i>	262	<i>trilineatum</i> , <i>Pristipoma</i>	332
<i>Trichurus</i>	262	<i>trilobatus</i> , <i>Lutjanus</i>	317
<i>argenteus</i>	262	<i>Scarus</i>	434
<i>lepturus</i>	262	<i>Triloburus</i>	318
<i>nitens</i>	262	<i>philadelphicus</i>	318
<i>trichocephalus</i> , <i>Cheilichthys</i>	499	<i>trilobus</i> , <i>Blepsias</i>	382
<i>Tetraodon</i>	499	<i>trimaculata</i> , <i>Parapetenia</i>	420
<i>Trichocyclopus</i>	501	<i>trimaculatum</i> , <i>Heros</i>	420
<i>erinaceus</i>	501	<i>Trinectes</i>	220
<i>Trichoderma</i>	494	<i>scabra</i>	229
<i>Trichodiodon</i>	501	<i>trinitatis</i> , <i>Anchovia</i>	50
<i>pilosus</i>	501	<i>Anchoviella</i>	50
<i>Trichodon</i>	357	<i>tristegus</i> , <i>Acanthurus</i>	362
<i>japonicus</i>	357	<i>Chatodon</i>	362
<i>lineatus</i>	357	<i>trilourus</i> , <i>Bodianus</i>	323
<i>trichodon</i>	357	<i>tripes</i> , <i>Nealotus</i>	761
<i>trichodon</i> , <i>Mugil</i>	254	<i>Tripterodon</i>	64
<i>Querimana</i>	254	<i>hautin</i>	64
<i>Trachinus</i>	357	<i>Tripterygium</i> , <i>carminalis</i>	458
<i>Trichodon</i>	357	<i>triptyerus</i> , <i>Cottus</i>	393
<i>Trichodontidae</i>	39	<i>tripunctulatus</i> , <i>Maurolicus</i>	73
<i>Trichonotus</i>	220	<i>Valenciennellus</i>	73
<i>Trichopsetta</i>	220	<i>triquier</i> , <i>Ostracion</i>	496
<i>ventralis</i>	220	<i>Rhimesomus</i>	496
<i>trichoptista</i> , <i>Codoma</i>	130	<i>triserialis</i> , <i>Murenenophis</i>	88
<i>Ergala</i>	130	<i>Ophichthus</i>	88
<i>trichurus</i> , <i>Macrourus</i>	206	<i>triseriatus</i> , <i>Platyrhina</i>	24
<i>Macrurus</i>	206	<i>triseriatus</i> , <i>Platyrhinoidis</i>	24
<i>tricolor</i> , <i>Chatodon</i>	361	<i>trisignatum</i> , <i>Moxostoma</i>	105
<i>Engraulis</i>	48	<i>Trisotropis</i>	313
<i>Holacanthus</i>	361	<i>anguaii</i>	314
<i>Thynnalus</i>	60	<i>bonaui</i>	314
<i>Tricopterus</i>	273	<i>boulengeri</i>	313
<i>tricornis</i> , <i>Ostracion</i>	497	<i>chlorostomus</i>	314
<i>tricuspidatus</i> , <i>Hyperhamphus</i>	198	<i>dimidiatus</i>	314
<i>tricuspidis</i> , <i>Cottus</i>	389	<i>interstitialis</i>	314
<i>Gymnoanthus</i>	389	<i>jordani</i>	314
<i>tridecemlineatus</i> , <i>Esor</i>	173	<i>microlepis</i>	314
<i>tridecemspinosa</i> , <i>Metzclaria</i>	370	<i>reticulatus</i>	315
<i>tridecemspinosa</i> , <i>Scorpenea</i>	370	<i>stomias</i>	314
<i>tridens</i> , <i>Lutjanus</i>	318	<i>venenosus</i>	313
<i>Pœcilia</i>	190	<i>apua</i>	313
<i>(Acropœcilia)</i>	190	<i>xanthostictus</i>	314
<i>Salema</i>	328	<i>xenarchus</i>	314
<i>Sargus</i>	328	<i>trispinosa</i> , <i>Corvina</i>	346

List of scientific names—Continued

	Page		Page
<i>trispinosus</i> , <i>Odontopyxis</i>	390	<i>turgidus</i> , <i>Tetronotus</i>	498
<i>Scarus</i>	434	<i>turneri</i> , <i>Lycodes</i>	476
<i>tristis</i> , <i>Moniana</i>	132	<i>turrubarensis</i> , <i>Gambusia</i>	189
<i>tristechus</i> , <i>Atractosteus</i>	38	<i>Poeciliopsis</i>	189
<i>Esox</i>	38	<i>tuscumbia</i> , <i>Etheostoma</i>	293
<i>trivittatus</i> , <i>Graministes</i>	331	<i>Psychromaster</i>	293
<i>Troglichthys</i>	195	<i>tuvensis</i> , <i>Astatheros</i>	421
<i>rosea</i>	195	<i>Cichlasoma</i>	421
<i>Trogloglanis</i>	155	<i>Loricaria</i>	162
<i>pattersoni</i>	155	<i>tweediei</i> , <i>Syngnathus</i>	242
<i>Trompa</i>	434	<i>Tylosurus</i>	198
<i>troostii</i> , <i>Lepidosteus</i>	37	<i>acus</i>	197
<i>tropica</i> , <i>Echeneis</i>	449	<i>cantralini</i>	197
<i>tropicus</i> , <i>Atractosteus</i>	38	<i>caribbeus</i>	197
<i>Phtheirichthys</i>	449	<i>curvops</i>	196
<i>Platypterus</i>	192	<i>fodiator</i>	198
<i>Tropidichthys</i>	500	<i>galeatus</i>	197
<i>tenuatus</i>	500	<i>gladius</i>	196
<i>Tropidinius</i>	328	<i>gulaniensis</i>	196
<i>dentatus</i>	328	<i>pacificus</i>	197
<i>Tropidodus</i>	10	<i>pterurus</i>	196
<i>troscelli</i> , <i>Hericthys</i> : <i>Heros</i>	427	<i>rashidoma</i>	196
<i>Sciaes</i>	423	<i>sagitta</i>	195
<i>Sciaideichthys</i>	148	<i>scapularis</i>	195
<i>troscelli</i> , <i>Abudedefduf</i>	145	<i>sierrita</i>	196
<i>Glypheidodon</i>	415	<i>Tympanonotum</i>	503
<i>trowbridgei</i> , <i>Holoconotus</i>	415	<i>planci</i>	503
<i>trowbridgei</i> , <i>Homalopomus</i>	409	<i>Tyntastes</i>	448
<i>triculenta</i> , <i>Chlorophthalimus</i>	214	<i>brevis</i>	448
<i>truncata</i> , <i>Belone</i> : <i>Muthera</i>	165	<i>sagitta</i>	448
<i>Ranzania</i>	504	<i>Typhlichthys</i>	194
<i>truncatus</i> , <i>Blennius</i>	461	<i>osborni</i>	195
<i>Dinctus</i>	35	<i>subterraneus</i>	195
<i>Salmo</i>	58	<i>wyandotte</i>	195
<i>Saurus</i>	163	<i>Typhlogobius</i>	447
<i>Scarus</i>	433	<i>californiensis</i>	447
<i>Tetronotus</i>	504	<i>Typhlosarasa</i>	508
<i>Trutta</i>	55, 50	<i>shufeldti</i>	508
<i>trutta</i> , <i>Salmo</i>	55	<i>typicus</i> , <i>Ophioscion</i>	347
<i>Trutta</i>	55	<i>Platygirus</i>	128
<i>corpo variegato</i>	55	<i>Rhinocodon</i>	18
<i>Trychodon</i>	115	<i>typhus</i> , <i>Achirophichthys</i>	89
<i>megalops</i>	115	<i>Carangichthys</i>	273
<i>Trygon</i>	28	<i>tyrannus</i> , <i>Anguilla</i>	77
<i>gymnura</i>	28	<i>brevicaudata</i> , <i>Brevoortia</i>	44
<i>longa</i>	28	<i>Brevoortia</i>	44
<i>osteosticta</i>	29	<i>Clupea</i>	44
<i>sabina</i>	29	<i>uhleri</i> , <i>Citharichthys</i>	219
<i>sinensis</i>	28	<i>Uhæma</i>	341
<i>Trygonobatus torpedinus</i>	29	<i>lefroyi</i>	341
<i>tschawytscha</i> , <i>Oncorhynchus</i>	55	<i>meeki</i>	341
<i>Salmo</i>	55	<i>producta</i>	341
<i>tschawytschiformis</i> , <i>Salmo</i>	54	<i>Ulca</i>	393
<i>tschudii</i> , <i>Discopyge</i>	27	<i>marmorata</i>	393
<i>tsilcoensis</i> , <i>Catostomus</i>	106	<i>Ulcina</i>	397
<i>tsuppitchi</i> , <i>Fario</i>	57	<i>guntheri</i>	397
<i>Salmo</i>	55	<i>Ulocentra</i>	286
<i>tuberculata</i> , <i>Raja</i>	28	<i>blennius</i>	287
<i>tuberculatus</i> , <i>Catostomus</i>	109	<i>gilberti</i>	286
<i>Phæthonichthys</i>	264	<i>histrio</i>	286
<i>Sphæroides</i>	498	<i>meadiæ</i>	286
<i>tubercule</i> , <i>le Sphéroïde</i>	498	<i>phlox</i>	287
<i>tuberculée</i> , <i>le Sphéroïde</i>	498	<i>simotera</i>	286
<i>tubularis</i> , <i>Electris</i>	498	<i>stigmæa</i>	287
<i>tuckeri</i> , <i>Monacanthus</i>	437	<i>verecunda</i>	287
<i>tudes</i> , <i>Gobiesox</i>	495	<i>(Ulocentra) histrio</i> , <i>Etheostoma</i>	286
<i>Platysqualus</i>	489	<i>ulochir</i> , <i>Paraliparis</i>	405
<i>Salmo</i>	18	<i>ulvae</i> , <i>Xiphistes</i>	469
<i>Zygæna</i>	61	<i>Ulvaria</i>	470
<i>tuditanus</i> , <i>Hybopsis</i>	145	<i>subbifurcata</i>	470
<i>tullibee</i> , <i>Argyrosomus</i>	64	<i>Ulvicola</i>	467
<i>Coregonus</i>	61	<i>sanctæ rosæ</i>	467
<i>Leucichthys</i>	64	<i>umatilla</i> , <i>Agosia</i>	141
<i>Salmo</i> (<i>Coregonus</i>).....	64	<i>Umbria</i>	60
<i>tumidus</i> , <i>Carpoides</i>	103	<i>Umbra</i>	174
<i>Chironectes</i>	604	<i>delicatissima</i>	172
<i>Lophius</i>	604	<i>limi</i>	174
<i>tumifrons</i> , <i>Sternopygus</i>	101	<i>minor</i>	255
<i>tunicata</i> , <i>Liparis</i>	400	<i>pygmæa</i>	174
<i>tunicatus</i> , <i>Carelloparis</i>	400	<i>bilineata</i>	174
<i>turchesius</i> , <i>Scarus</i>	435	<i>umbratilis</i> , <i>Alburnus</i>	125
<i>Turdus cineraceus</i> <i>peltatus</i>	341	<i>Gambusia</i>	100
<i>flavus</i>	425	<i>Lythrurus</i>	125
<i>oculo-radiata</i>	426	<i>Xenophallus</i>	100
<i>pinnis branchiulibus carens</i>	325	<i>Umbridae</i>	174

List of scientific names—Continued

Page	Page		
umbrifer, <i>Cichlaurus</i>	417	unicolor <i>Reighardina</i>	9
<i>Notropis</i>	131	<i>Soleotalpa</i>	230
<i>Urobatis</i>	30	<i>vitulinus, Hypoplectrus</i>	321
<i>Urolophus</i>	30	<i>unicornis, Citharichthys</i>	219
umbriferum, <i>Cichlasoma</i>	417	<i>Unicornis pisces bahamensis</i>	495
Umbrina.....	350	<i>unicornu, Balistes</i>	496
<i>analis</i>	350	<i>unifasciatus, Hemirhamphus</i>	198
<i>arenata</i>	351	<i>Hyporhamphus</i>	198
<i>broussonnetii</i>	350	<i>unilineatus, Hemigrammus</i>	98
<i>coroides</i>	350	<i>Poecilurichthys</i>	98
<i>dorsalis</i>	350	<i>unimaculata, Perca</i>	338
<i>elongata</i>	351	<i>unimaculatus, Argyreiosus</i>	275
<i>furnieri</i>	349	<i>Clinus</i>	470
<i>gracilicirrhus</i>	350	<i>Gymnells virdis</i>	477
<i>gracilis</i>	351	<i>uninotatus, Ctenolabrus</i>	424
<i>Januaria</i>	351	<i>Girardinus</i>	188
<i>littoralis</i>	351	<i>Glaucichthys</i>	188
<i>martinicensis</i>	351	<i>Mesopriion</i>	327
<i>nanus</i>	351	<i>unionensis, Centropomus</i>	306
<i>panamensis</i>	351	<i>universitatis, Notropis</i>	124
<i>phalæna</i>	350	<i>univittatus, Apodichthys</i>	467
<i>roncador</i>	350	<i>Upeneus</i>	343
<i>simaloe</i>	350	<i>baiteatus</i>	343
<i>undulata</i>	350	<i>dentatus</i>	343
<i>xanti</i>	350	<i>flavovittatus</i>	343
umbrosa, <i>Cyprinella</i>	130	<i>grandisquamis</i>	344
<i>Narcine</i>	27	<i>maculatus</i>	343
umbrosus, <i>Eques acuminatus</i>	353	<i>martinicus</i>	343
<i>Esox</i>	173	<i>parvus</i>	343
<i>Gymnothorax</i>	90	<i>punctatus</i>	343
<i>Pareques</i>	353	<i>rathbuni</i>	344
<i>Platichthys</i>	226	<i>tetraspisus</i>	344
<i>Psettichthys</i>	226	<i>xanthogrammus</i>	343
<i>Sebastichthys</i>	367	<i>uracantha, Loricaria</i>	162
<i>Sebastomus</i>	367	<i>Uraleptus</i>	211
Umbrula.....	351	<i>Uranichthys</i>	87
<i>elongata</i>	351	<i>brachycephalus</i>	87
<i>littoralis</i>	351	<i>Uranidea</i>	383
umpque, <i>Ptychocheilus</i>	114	<i>greeni</i>	384
Unbarana.....	39	<i>hoylei</i>	385
uncinalis, <i>Icelus</i>	379	<i>kumlienii</i>	385
uncinatus, <i>Artedielius</i>	377	<i>marginata</i>	384
<i>Cottus</i>	377	<i>microstoma</i>	385
uncomphagre, <i>Xyrauchen</i>	108	<i>policaris</i>	386
undecimale, <i>Hæmulon</i>	330	<i>quiescens</i>	385
undecimals, <i>Centropomus</i>	305, 306	<i>rhothea</i>	383
<i>Centropomus lucioperca</i>	305	<i>semiscabra centropleura</i>	383
<i>Sciæna</i>	305	<i>spilota</i>	385
undecim-raditus, <i>Centropomus</i>	305	<i>tenuis</i>	384
underwoodi, <i>Herichthys</i>	423	<i>wheeleri</i>	384
<i>Rhamdia</i>	423	<i>uranidea, Cottogaster</i>	285
<i>Tomocichla</i>	423	<i>Etheostoma (Cottogaster)</i>	285
undulata, <i>Perca</i>	349	<i>uranops, Phenacobius</i>	142
<i>Umbrina</i>	350	<i>Urano scopidae</i>	454
undulatus, <i>Menticirrhus</i>	350	<i>Urano scopus anoplos</i>	454
<i>Micropogon</i>	349	<i>y-græcum</i>	454
<i>Ostracion</i>	497	<i>uranoscopus, Ceratius</i>	508
<i>Rhinobatoides</i>	23	<i>Mancalias</i>	508
undulatus, <i>Serranus</i>	258	<i>urichi, Cæcorhamdia</i>	158
undulosus, <i>Serranus</i>	315	<i>Uribatoides</i>	309
unerak, <i>Ophidium</i>	477	<i>Urobatis</i>	29
Unibranchiatura.....	76	<i>concentricus</i>	30
<i>grisea</i>	76	<i>halleri</i>	29
<i>immaculata</i>	76	<i>nebulosus</i>	30
<i>lineata</i>	76	<i>sloani</i>	29
<i>marmorata</i>	76	<i>vermiculatus</i>	29
unicolor aberrans, <i>Hypoplectrus</i>	321	<i>umbrifer</i>	30
<i>accensus</i> , <i>Hypoplectrus</i>	321	<i>vermiculatus</i>	30
<i>affinis</i> , <i>Hypoplectrus</i>	321	<i>Urocentrus</i>	467
<i>Ammocoetes</i>	0	<i>Uroconger</i>	81
<i>Balistes monoceros</i>	496	<i>varidens</i>	81
<i>bovinus</i> , <i>Hypoplectrus</i>	321	<i>vicus</i>	81
<i>Bryttus</i>	299	<i>Urolophus aspidurus</i>	30
<i>Chætodon</i>	360	<i>asterius</i>	30
<i>chlorurus</i> , <i>Hypoplectrus</i>	321	<i>goodei</i>	30
<i>crocatus</i> , <i>Hypoplectrus</i>	321	<i>halleri</i>	30
<i>gemma</i> , <i>Hypoplectrus</i>	321	<i>nebulosus</i>	30
<i>gummigutta</i> , <i>Hypoplectrus</i>	321	<i>rogersi</i>	30
<i>guttavarius</i> , <i>Hypoplectrus</i>	321	<i>umbrifer</i>	30
<i>Hypoplectrus</i>	321	<i>Uronectes</i>	478
<i>indigo</i> , <i>Hypoplectrus</i>	321	<i>Urophycis</i>	213
<i>Muræna</i>	90	<i>chesteri</i>	213
<i>nigricans</i> , <i>Hypoplectrus</i>	321	<i>chuss</i>	213
<i>Perca</i>	321	<i>floridae</i>	213
<i>pinnularius</i> , <i>Hypoplectrus</i>	321	<i>regius</i>	213
<i>puelia</i> , <i>Hypoplectrus</i>	321	<i>tenius</i>	213

List of scientific names—Continued

	Page		Page
urophthalmus, <i>Heros</i> .	418	veliger, <i>Histioclinus</i> .	456
Uropsetta.	223	velox, <i>Carcharhinus</i> .	16
Uropterygius.	94	Ciliola.	145
neurus.	94, 95	Euleptorhamphus.	198
urophalma, <i>Parapetenis</i> .	418	Sarda.	259
Urotrygon.	30	venadorum, <i>Myctoperca</i> .	314
aspidurus.	30	Venefica.	82
goodei.	30	ocells.	82
mundus.	30	procera.	82
Uroxis.	28	tentaculata.	82
ursinus, <i>Salmo</i> .	60	venenosus, <i>Perca</i> .	313
urus, <i>Bubalichthys</i> .	102	venosa, <i>Meletta</i> .	42
Carpoides.	102	venosus, <i>Aluterus</i> .	495
Ictiobus.	102	Trisopteris.	313
ustus, <i>Callyodon</i> .	431	ventralis, <i>Bathypterois</i> .	196
Cryptotomus.	431	Bromophycis.	480
utah, <i>Salmo</i> .	56	Citharichthys.	220
utawana, <i>Catostomus</i> .	105	Cottus.	389
uter, <i>Catulus</i> .	12	Eucrotus.	267
Cephaloscyllium.	12	Novaculichthys.	430
vacea, <i>Carpoides</i> .	103	Ogilbia.	480
Darnalichthys.		Trichopsetta.	220
Vacuqua.	345	Xyrichthys.	430
macrops.	345	ventricosus, <i>Aptocyclus</i> .	398
sancta-luciae.	345	Blepsias.	380, 381
sialis.	345	Ceratichthys.	141
subaequalis.	345	Cyclopterus.	398
wafer, <i>Myrophis</i> .	83	venusta, <i>Cyprinella</i> .	130
vagans, <i>Katsuwonus</i> .	298	Emmeekia.	428
Thynnus.	298	Erogala.	130
vagrans, <i>Chirostoma</i> .	248	Limia.	175
Kirtlandia.	248	venustus, <i>Dinemus</i> .	234
lacinata, <i>Menidia</i> .	248	Hesperoleucus.	122
vahlii, <i>Lycodes</i> .	474	Pseudojulis.	428
valginiensis, <i>Scarus</i> .	431	Xyrichthys.	430
vallanti, <i>Pteroplatea</i> .	29	vere-crucis, <i>Bairdiella</i> .	346
Vaillantia.	288	Nector.	346
camura.	288	verany, <i>Cybium</i> .	256
chlorosoma.	288	Verasper.	223
valenciennei, <i>Pteraplatea</i> .	29	moseri.	223
valenciennei, <i>Erotelis</i> .	437	verde, <i>Pudiano</i> .	426
Moxostoma.	441	verecunda, <i>Ulocentra</i> .	287
Smaragdus.	441	verecundum, <i>Etheostoma</i> .	287
Valencienneellus.	73	Lepidion.	211
stellatus.	73	Verilidae.	328
tripunctatus.	73	Verilius.	328
vampyrus, <i>Cephalopterus</i> .	32	sordidus.	328
Vandellius.	262	Verma.	328
vandepolli arubensis, <i>Procellia</i> .	193	kendalli.	85
Mollinesia sphenoops.	193	vermelho, <i>Pudiano</i> .	425
Poecilia.	193	vermicularis, <i>Corvina</i> .	328
vandoisulus, <i>Clinostomus</i> .	121	Signature.	348
Leuciscus.	121	Sparus.	327
vanhyningi, <i>Signalosa atchafalayae</i> .	47	vermiculatus, <i>Esox</i> .	173
varia, <i>Perca</i> .	317	Exocetus.	202
variabilis, <i>Diploites</i> .	298	Urobatis.	29
Perca.	298	sloani.	29
Pomacentrus.	413	Xyrichthys.	430
variata, <i>Ethostoma</i> .	289	vermiformis, <i>Neoconger</i> .	81
variatus, <i>Characodon</i> .	183	Oxyurus.	79
Platyoperculus.	193	vernalis, <i>Clupea</i> .	42
Pecilichthys.	239	Gobio.	137
varidens, <i>Uroconger</i> .	81	Hydrargyre.	176
variegata, <i>Batrachoides</i> .	487	Hyodon.	40
Loricaria.	181	Scomber.	257
Murena.	183	vernillas, <i>Batrachoides</i> .	487
Salpa purpurea.	327	verreatxi, <i>Conger</i> .	70
Skiffia.	183	verres, <i>Balistes</i> .	492
variegatus, <i>Cyprinodon</i> .	181	Lutjanus.	425
Limnurgus.	182	Sufflamen.	402
Orbis levius.	499	verrilli, <i>Gymnothorax</i> .	93
Pachylabrus.	411	Sidera.	93
Sargus.	338	verrillii, <i>Brosmophycis</i> .	490
varius, <i>Malacoctenus</i> .	459	Lycenchelys.	475
Monacanthus.	494	Lycodes.	475
Myrodes.	459	Ogilbia.	480
Serranus.	310	verrucosa, <i>Occa</i> .	304
velatus, <i>Ptychostomus</i> .	109	Brachyopsis.	306
velleana, <i>Atherina</i> .	246	Cottus.	394
velifer, <i>Carpoides</i> .	103	Diodon.	387
Catostomus.	103	Myoxocephalus.	302
Letharchus.	88	Verrugato.	323
Rhinchithys (Apococe).	141	pacificus.	324
velifera, <i>Apococe</i> .	141	Verrunculus.	492
Mollinessa.	141	polylepis.	492

List of scientific names—Continued

Page	Page		
<i>versicolor</i> , <i>Abramis</i>	115	<i>virginæ</i> , <i>Brachistoma</i>	7
<i>Girardinus</i>	191	<i>virginis</i> , <i>Salmo</i>	57
<i>Labrus</i>	336	<i>Scarus</i>	433
<i>Limia</i>	191	<i>virginicus</i> , <i>Anisotremus</i>	332
<i>Malacoctenus</i>	460	<i>Polynemus</i>	256
<i>Myxodes</i>	460	<i>Sparus</i>	322
<i>Phytichthys</i>	469	<i>viride</i> , <i>Sparisoma</i>	432
<i>Pœciliichthys</i>	291	<i>viridensis</i> , <i>Sphyræna</i>	255
<i>Stenotomus</i>	336	<i>viridescens</i> , <i>Atherina</i>	247
<i>Xiphistes</i>	469	<i>Fundulus</i>	176
<i>versipunctatus</i> , <i>Gymnothorax</i>	92	<i>Osmerus</i>	67
<i>vertagus</i> , <i>Gadus</i>	210	<i>Syngnathus</i>	242
<i>verticalis</i> , <i>Echeneis</i>	449	<i>virdipalidus</i> , <i>Gobius</i>	446
<i>Pleuronichthys</i>	225	<i>viridis</i> , <i>Amia</i>	38
<i>verus</i> , <i>Conger</i>	79	<i>bahamensis</i> , <i>Piscis</i>	433
<i>vespertilio</i> , <i>Lophius</i>	507	<i>Centropomus</i>	306
<i>Ogcocephalus</i>	507	<i>Diacope</i>	324
<i>Rhinoptera</i>	31	<i>Gymnells</i>	477
<i>verus</i> , <i>Carcharodon</i>	20	<i>Lutianus</i>	324
<i>vetula</i> , <i>Balistes</i>	491	<i>Ophidium</i>	477
<i>Scarus</i>	433, 434	<i>Scarus</i>	432
<i>vetulus</i> , <i>Parophrys</i>	226	<i>unimaculatus</i> , <i>Gymnæsis</i>	477
<i>vezillare</i> , <i>Boleosoma</i>	288	<i>viscosus</i> , <i>Cottus</i>	385
<i>verillaris</i> , <i>Pteropodus</i>	369	<i>Silurus</i>	156
<i>Sebastichthys</i>	369	<i>vitrea</i> , <i>Ioa</i>	288
<i>verillarium</i> , <i>Holocentrum</i>	236	<i>Perca</i>	282
<i>verillarius</i> , <i>Holocentrus</i>	236	<i>vitreum</i> , <i>Stizostedion</i>	282
<i>Verillifer</i>	480	<i>vitreus</i> , <i>Pœciliichthys</i>	288
<i>vicina</i> , <i>Murennopsis</i>	92	<i>vitta</i> , <i>Xyrichthys</i>	430
<i>vicinalis</i> , <i>Ictius</i>	378	<i>vittata</i> , <i>Aztecua</i>	143
<i>vicinus</i> , <i>Gymnothorax</i>	92	<i>Channomuraena</i>	95
<i>Uroconger</i>	81	<i>Clupea</i>	41
<i>victoriae</i> , <i>Moxostoma</i>	109	<i>Codoma</i>	143
<i>Vieja</i>	433, 434	<i>Echeneis</i>	449
<i>Vigil</i>	288	<i>Gambusia</i>	186
<i>clarus</i>	288	<i>Hemitremia</i>	121
<i>pellucidus</i>	288	<i>Inermia</i>	340
<i>vivax</i>	288	<i>Lepidomedes</i>	147
<i>vigil</i> , <i>Ioa</i>	288	<i>Limia</i>	191
<i>vigilax</i> , <i>Ceratichthys</i>	144, 145	<i>Poecilia</i>	191
<i>Villarius</i>	152	<i>vittatus</i> , <i>Ammodytes</i>	451
<i>pricei</i>	153	<i>Catostomus</i>	100
<i>willosa</i> , <i>Clupea</i>	68	<i>Cyprinus</i>	140
<i>Fistularia</i>	245	<i>Esox</i>	173
<i>willosus</i> , <i>Cottus</i>	393	<i>Gobiesox</i>	480
<i>Hemipteritus</i>	393	<i>Ichthyophis</i>	95
<i>Mallotus</i>	66	<i>Leuciscus</i>	128
<i>vincente</i> , <i>Sicydium</i>	447	<i>Proceros</i>	30
<i>vincigueræ</i> , <i>Exocetus</i>	200	<i>Sarchirus</i>	38
<i>Exonastes</i>	200	<i>Siphametes</i>	122
<i>Vinciguerra</i>	73	<i>Sparus</i>	332
<i>attenuata</i>	73	<i>Symbolanichthys</i>	76
<i>lucetia</i>	73	<i>vittula</i> , <i>Sargus</i>	338
<i>nimbaria</i>	73	<i>vittulinum</i> , <i>Plectropoma</i>	321
<i>vincta</i> , <i>Xuræ</i>	272	<i>vittulus</i> , <i>Hypoplectrus unicolor</i>	321
<i>vinctipes</i> , <i>Nonostoma</i>	289	<i>vittulus</i> , <i>Carpoides</i>	102
<i>vinctus</i> , <i>Blennius</i>	461	<i>vivanet</i> , <i>Bodianus</i>	325
<i>Caranx</i>	272	<i>vivanus</i> , <i>Anthias</i>	322
<i>Fundulus</i>	176	<i>Hemianthias</i>	322
<i>vincentus</i> , <i>Lethotrema</i>	398	<i>Lutianus</i>	326
<i>viola</i> , <i>Antimora</i>	211	<i>Mesopriion</i>	320
<i>Eques</i>	353	<i>vixax</i> , <i>Ammocrypta</i>	288
<i>Haloporphyrus</i>	211	<i>Cliola</i>	145
<i>Pareques</i>	353	<i>Vigil</i>	288
<i>violaceus</i> , <i>Apodichthys</i>	466	<i>Vivero</i>	312
<i>Cebidichthys</i>	466	<i>vivipara</i> , <i>Poecilia</i>	190
<i>Escolar</i>	466	<i>viviparus</i> , <i>Lophius</i>	504
<i>Thysritops</i>	260	<i>Sebastas</i>	364
<i>virens</i> , <i>Gadus</i>	209	<i>viamingii</i> , <i>Coryphaena</i>	265
<i>Pollachius</i>	209	<i>Vogmarus</i>	216
<i>Scarus</i>	433	<i>vogmarus</i> , <i>Trachypterus</i>	216
<i>Thalassoma</i>	429	<i>voador</i> , <i>Exocetus</i>	200
<i>virescens</i> , <i>Apodichthys</i>	407	<i>Istiophorus</i>	263
<i>Clupea</i>	42	<i>Rubio</i>	407
<i>Eigenmannia</i>	101	<i>voltans</i> , <i>Dactylopterus</i>	409
<i>Erlsdion</i>	356	<i>Exocetus</i>	200
<i>Gymnothorax</i>	92	<i>Trigla</i>	409
<i>Otolithus</i>	256	<i>volucella</i> , <i>Hybopsis</i>	133
<i>Pantosteus</i>	105	<i>volucellus</i> , <i>Hybognathus</i>	133
<i>Sternarchus</i>	101	<i>Vomer</i>	274
<i>virgata</i> , <i>Coryphaena</i>	265	<i>brasiliensis</i>	275
<i>Nivicola</i>	292	<i>brownii</i>	275
<i>virgatus</i> , <i>Daleolepis</i>	472	<i>cayennensis</i>	275
<i>Poecilichthys</i>	292	<i>columbianus</i>	275
<i>virgatulus</i> , <i>Gobiesox</i>	489	<i>cubæ</i>	275

List of scientific names—Continued

	Page		Page
Vomer curtus.	275	williamsoni clsmontanus, <i>Coregonus</i> .	65
declivifrons.	275	<i>Prosoplum</i> .	65
dominicensis.	275	williamsoni, <i>Coregonus</i> .	65
dorsalis.	275	<i>Ctenodax</i> .	269
gabonensis.	275	<i>Gasterosteis</i> .	237, 238
goreensis.	275	<i>Prosoplum</i> .	65
martinicensis.	275	willoughbyi, <i>Acrotus</i> .	240
nova-horacensis.	275	willughbyi, <i>Trematopsis</i> .	503
sancta-marthae.	275	willughbyi, <i>Balistes</i> .	493
sanctae-petri.	275	<i>Canthidermis</i> .	493
senegalensis.	275	<i>Trematopsis</i> .	503
setapinnis.	275	wilsoni, <i>Acutomentum</i> .	286
cubensis.	275	<i>Cottus</i> .	285
spixii.	275	<i>Oceanomyzon</i> .	8
vomer, <i>Argyreiosus</i> .	275	<i>Phenagoniates</i> .	96
* Zeus.	275	<i>Sebastodes</i> .	366
vomerina, <i>Atherina</i> .	250	winehilli, <i>Hybopsis</i> .	133
<i>Chiromista</i> .	250	woolmani, <i>Paralichthys</i> .	224
vomerinus, <i>Anarrhichthys</i> .	472	wrayi, <i>Gambusia</i> .	186
<i>Salaris</i> .	464	wrighti, <i>Etheostoma</i> .	285
vorax, <i>Mesoprlon</i> .	328	<i>Istiophorus</i> .	263
<i>Pseudorhombus</i> .	224	wurdemanni, <i>Gobius</i> .	441
<i>Ptychocheilus</i> .	114	wyandotte, <i>Typhlichthys</i> .	195
vulgaris, <i>Acanthias</i> .	21		
<i>Amelurus</i> .	153	xænocephalus, <i>Hybopsis</i> .	127
<i>Conger</i> .	79	<i>Hydrophorus</i> .	127
<i>Gunnellus</i> .	467	xænura, <i>Ergala</i> .	130
<i>Lumpus</i> .	398	xænurus, <i>Minilius</i> .	130
<i>Molva</i> .	212	xaliscone, <i>Xenendum</i> .	184
<i>Pagrus</i> .	337	xanthurus, <i>Catulus</i> .	12
<i>Pimelodus</i> .	153	<i>Pristurus</i> .	12
<i>Pomotis</i> .	302	xanthichthys.	493
<i>Sphyraena</i> .	255	<i>lineopunctatus</i> .	493
<i>Spinachia</i> .	22	<i>ringens</i> .	493
<i>Squatina</i> .	28	xanthocephalus, <i>Silurus</i> .	154
<i>Trigonobatus</i> .	28	xanthogrammus, <i>Upeneus</i> .	243
vulnerata, <i>Apocope</i> .	141	xanthomes, <i>Anguilla</i> .	77
vulneratus, <i>Nothonotus</i> .	200	xanthops, <i>Odontoscion</i> .	344
<i>Pecilichthys</i> .	200	xanthopteron, <i>Hæmulon</i> .	330
Vulpecula.	18	xanthopus, <i>Catostomus</i> .	107
<i>marina</i> .	18	xanthosticta, <i>Mycteroperca bonaci</i> .	314
vulpecula, <i>Galeus</i> .	18	xanthostictus, <i>Trisotropis bonaci</i> .	314
vulpeculus, <i>Pimelodus</i> .	154	xanthostigma, <i>Citharichthys</i> .	218
Vulpes bahamensis.	39	xanthostigma, <i>Citharichthys</i> .	218
vulpes, <i>Albula</i> .	39	xanthulum, <i>Cynoscion</i> .	356
<i>Esox</i> .	152	xanthulus, <i>Ericson</i> .	256
<i>Pimelodus</i> .	152	<i>Lelostomus</i> .	249
<i>Squalus</i> .	18	<i>Pomacentrus</i> .	413
vulpinus, <i>Alopas</i> .	18	<i>Sparus</i> .	336
<i>Squalus</i> .	18	xanti, <i>Labrisomus</i> .	459
Vulsculus.	406	<i>Rypticus</i> .	316
<i>imberbis</i> .	406	<i>Umbrina</i> .	350
vulsus, <i>Agonus</i> .	395	<i>Xenichthys</i> .	324
<i>Stelgis</i> .	395	Xantocles.	375
vultharius, <i>Acipenser</i> .	34	<i>frenatus</i> .	375
		xantops, <i>Odontoscion</i> .	334
wabashensis, <i>Anguilla</i> .	77	xantusi, <i>Lepidopus</i> .	262
wachna, <i>Gadus</i> .	210	xenarcha, <i>Mycteroperca</i> .	314
wagneri, <i>Pimelodus</i> .	157	Xenarchi.	215
<i>Rhamdia</i> .	157	xenarchus, <i>Trisotropis</i> .	314
walkeri, <i>Selanonius</i> .	19	Xenatherina.	249
<i>Salmo</i> .	294	<i>lisa</i> .	249
watauga, <i>Erimystax</i> .	138	xenauchen, <i>Galeichthys</i> .	150
<i>Hybopsis</i> .	138	Xenendum.	183
webbi, <i>Nemabrama</i> .	234	<i>caliente</i> .	184
<i>webbi</i> , <i>Blennophis</i> .	464	<i>multipunctatum</i> .	183
<i>Embletoeca</i> .	411	<i>xaliscone</i> .	184
<i>Ophiolebennius</i> .	464	Xeneretmus.	396
weberi, <i>Xenogobius</i> .	442	<i>infraspinatus</i> .	396
welaka, <i>Notropis</i> .	124	<i>lelops</i> .	396
welshi, <i>Otophodium</i> .	486	<i>ritteri</i> .	396
wheatlandi, <i>Gasterosteus</i> .	238	<i>trisacanthus</i> .	396
wheeleri, <i>Uranidea</i> .	384	Xenichthyidae.	324
whipplei alabamae, <i>Etheostoma</i> .	292	Xenichthys.	324
whipplei alabamae, <i>Etheostoma</i> .	292	<i>californiensis</i> .	324
<i>Boleichtys</i> .	292	<i>xanti</i> .	324
<i>Claricola</i> .	292	<i>xenops</i> .	324
<i>Cyprinella</i> .	129	Xenicus, <i>Fundulus</i> .	180
<i>Erogala</i> .	129	Xenisma.	179
whitei, <i>Balsichthys</i> .	183	<i>catenatum</i> .	179
<i>Goodea</i> .	183	<i>stellifer</i> .	179
<i>Salmo</i> .	59	Xenigma, <i>Gurnardus</i> .	406
wilderi, <i>Lampetra</i> .	9	<i>Prionotus</i> .	406
williamsi, <i>Sympnurus</i> .	231	Xenistius.	324
		<i>californiensis</i> .	324

List of scientific names—Continued

	Page
Xenoberyces.	232
Xenocara.	180
chagresi.	180
Xenochirus.	386
alsacanthus.	386
latirostris.	386
pentacanthus.	386
triacanthus.	386
Xenodermichthys.	83
copel.	53
nodulosus.	431
Xenodon, <i>Catostomus</i> .	431
Leptoscarus.	431
Xenognathus.	52
profundorum.	52
Xenogobius.	442
weberi.	442
Xenom.	172
Xenomyxus.	81
atrarius.	81
rietius.	81
Xenophallus.	190
umbratilis.	190
Xenops, <i>Xenichthys</i> .	324
Xenopsetta, <i>Cypselurus</i> .	202
Exocetus.	202
Xenopterygia.	488
Xenopthalmichthys.	69
daneæ.	69
Xenopyxis.	396
latifrons.	396
lelops.	396
ritteri.	396
Xenorhynchichthys.	254
xenostethus, <i>Sternias</i> .	380
Triglops.	380
Xenotilapia.	300
aureolus.	300
lythrocchloris.	300
megalotis.	300
occidentalis.	301
Xeropetes.	467
fuscum.	467
Xesurus.	363
clariensis.	363
punctatus.	363
Xiphias.	264
gladius.	264
imperator.	264
rondeletii.	264
Xiphidion.	309
mucosum.	469
Xiphidium cruentum.	469
Xiphidae.	264
Xiphiformes.	263
Xiphister.	469
chirurgus.	469
mucosus.	469
rupaestris.	469
Xiphisteridae.	469
Xiphistes.	193
uvula.	193
versicolor.	193
Xiphophorus.	187
bimaculatus.	187
brevis.	187
gracilis.	187
guntheri.	187
halleri.	187
brevis.	187
guntheri.	187
rachovii.	187
strigatus.	187
jalapae.	184
montezuma.	183
rachovii.	183
strigatus.	183
Xiphostomidae.	100
Xiphotheca.	262
Xurel.	272
fasciatus.	272
lata.	272
lugubris.	273
marginatus.	272
medusicola.	273
Xurel melampygus.	272
stellatus.	273
tenebrosus.	273
vincens.	272
Xylosterina, <i>Stellerina</i> .	304
Xylosternus, <i>Brachyopsis</i> .	394
Xyrauchen.	108
texanus.	108
uncomphagre.	108
Xyrichthys.	430
binghami.	430
infirmus.	430
jessiae.	429
martinicensis.	430
modestus.	430
mundiceps.	430
pavoninus.	430
psittacus.	430
rosipes.	430
ventralis.	430
venustus.	430
vermiculatus.	430
vitta.	430
Xyris, <i>Scorpioides</i> .	370
Sebastopsis.	370
Xyrule.	429
jessiae.	429
Xysterna.	341
havana.	342
Xyster.	339
Xysterus.	339
Xystes.	395
axinophrys.	395
Xystophorus.	278
Xystriama.	298
pandionis.	298
Xystreurus.	223
liolepis.	223
Xystrodon, <i>Sparisoma</i> .	431
Xystroperca.	315
pardalis.	315
Xystropitides.	301
gilli.	302
longimanus.	302
pallidus.	302
Xystrouros.	142
popoche.	142
Xysturus, <i>Myrichthys</i> .	85
Opifuris.	85
Yaito, <i>Euthynnus</i> .	259
Yale, <i>Ostracion</i> .	497
Yarrella.	72
blackfordi.	72
Yarrowi, <i>Agosia</i> .	141
Apococe.	141
y-cauda, <i>Gilllichthys</i> .	444
Quietula.	444
y-græcum, <i>Astroscopus</i> .	454
Yucautana, <i>Gambusia</i> .	186
Yuma, <i>Goblesox</i> .	490
Yurinia.	137
alta.	137
Zacentrus, <i>Hispanicus</i> .	388
Sebastichthys.	388
Zachirus, <i>Errex</i> .	228
Glyptocephalus.	228
Zaclemus.	351
goodsi.	361
Zadocki, <i>Cylindrosteus</i> .	37
Zalargas.	73
nimbarius.	73
Zalembius.	410
roseaceus.	410
Zalleutes.	507
elater.	507
Zalocys.	277
stilbe.	277
Zalopys.	865
aleutianus.	865
strorubens.	865
astrovirens.	865

List of scientific names—Continued

	Page		Page
Zalypnus.	443	zonale arcansanum, <i>Etheostoma</i> .	289
Zanclidæ.	362	<i>Nanostoma</i> .	289
Zancilurus.	263	zonalis, <i>Poecilichthys</i> .	289
Zancius.	362	zonata, <i>Elassoma</i> .	305
centrognathos.	362	<i>Heterandria</i> .	191
cornutus.	362	<i>Seriola</i> .	279
zanclius, <i>Dermatolepis</i> .	313	zonatum, <i>Cichlasoma</i> .	417
zanemus, <i>Ceratichthys</i> .	138	<i>Elassoma</i> .	305
Zaniolepidæ.	375	zonatus, <i>Alburnus</i> .	128
Zaniolepis.	375	<i>Chetodipterus</i> .	368
frenatus.	375	<i>Cycloctis</i> .	128
latipinnis.	375	<i>Ephippus</i> .	358
Zanotacanthus.	76	<i>Esox</i> .	178
Zaphotias.	72	<i>Fundulus</i> .	178
pedialotus.	72	<i>Scomber</i> .	279
Zaprota.	452	<i>Theraps</i> .	417
silemus.	452	<i>Zygonectes</i> .	177
Zaprotride.	452	Zonichthys.	279
Zapteryx.	24	<i>falcatus</i> .	280
exasperatus.	24	<i>fasciatus</i> .	279
xyster.	24	<i>mazatlanus</i> .	279
zapus, <i>Hemilepidotus</i> .	381	<i>rivolianus</i> .	280
Zastomias.	70	zonifer, <i>Clinus</i> .	459
scintillans.	70	<i>Erilepis</i> .	373
zatropis, <i>Siphostoma</i> .	243	<i>Myriolepis</i> .	373
zebra, <i>Acanthurus</i> .	362	<i>Zygonectes</i> .	178
Arbaciota.	491	zonipactus, <i>Pomacanthodes</i> .	361
Fundulus.	176	zonistius, <i>Coccotis</i> .	128
Gerris.	342	<i>Luxilus</i> .	128
Gobiesox.	401	Zonogobius.	441
Gobius.	441	<i>zebra</i> .	441
Hydrargyra.	179	zonope, <i>Jordania</i> .	375
Percina.	283	Zonoscion.	351
Pleoma.	283	<i>rathbuni</i> .	351
Plancterus.	179	zonus, <i>Malacocottus</i> .	388
Psychrotules.	392	Zoogoneticus.	182
Zonogobius.	441	<i>diazi</i> .	182
zebrinus, <i>Fundulus</i> .	179	<i>dugesii</i> .	182
Zeidie.	222	<i>maculatus</i> .	182
zelotes, <i>Dactyloscopus</i> .	455	<i>minifatus</i> .	182
Esioscopus.	455	<i>quitzeoensis</i> .	182
Hemicarax.	271	<i>robustus</i> .	182
Zenion.	232	Zophendum.	140
hololepis.	232	<i>australe</i> .	116
zenithicus, <i>Argyrosomus</i> .	64	zopherus, <i>Potamocottus</i> .	385
Leucichthys.	64	zophochir, <i>Ophichthus</i> .	88
Zenopsis.	232	<i>Ophichthys</i> .	88
ocellata.	232	zostera, <i>Hippocampus</i> .	244
Zenoidæ.	232	zosteria, <i>Evermannia</i> .	445
zephyreus, <i>Astroscopus</i> .	454	zosterurum, <i>Gobiosoma</i> .	445
Fristis.	23	zunensis, <i>Leuciscus</i> .	114
Zesticelus.	392	Zygema.	18
profundorum.	392	<i>lowini</i> .	18
Zestidium.	347	<i>mallous</i> .	18
colonense.	347	<i>subarcuata</i> .	18
flecebrosum.	347	<i>tudes</i> .	18
Zestis.	347	zygrena, <i>Sphyraena</i> .	18
furthii.	347	<i>Squalus</i> .	18
melanochir.	347	Zygotabitis.	31
oscitans.	347	Zygonectes.	177
zestocaros, <i>Stelliferens</i> .	347	<i>atrilatus</i> .	183
Stelliferus.	347	<i>auroguttatus</i> .	177
zestus, <i>Nebri</i> .	354	<i>chrysotus</i> .	171
Zeus capillarii.	275	<i>cingulatus</i> .	171
crinitus.	274	<i>craticula</i> .	171
gallus.	274, 275	<i>diaphanus</i> .	175
geometricus.	275	<i>menona</i> .	175
guttatus.	216	<i>dispar</i> .	179
imperialis.	217	<i>escambie</i> .	178
luna.	216	<i>extensus</i> .	179
ocellatus.	232	<i>funduloides</i> .	176
opah.	217	<i>guttatus</i> .	178
quadrotus.	358	<i>henshalli</i> .	177
regius.	216	<i>hieroglyphicus</i> .	178
rostratus.	275	<i>inurus</i> .	185
setapinnis.	275	<i>jenkinsi</i> .	178
stremii.	217	<i>internalis</i> .	177
vomer.	275	<i>limi</i> .	178
zirahuen, <i>Chiromystoma</i> .	260	<i>lineatus</i> .	177
Zoarces gronovii.	473	<i>lincolatus</i> .	178
labrosus.	473	<i>luciae</i> .	178
Zoarcidæ.	173	<i>macdonaldi</i> .	177
zoarchus, <i>Lycodes</i> .	475	<i>manni</i> .	175

List of scientific names—Continued

	Page		Page
<i>Zygonectes melapleurus</i>	178	<i>Zygonectes scartes</i>	177
(<i>Micristius</i>) <i>brachypterus</i>	185	<i>sciadicus</i>	177
<i>notatus</i>	177	<i>seminolus</i>	178
<i>nottii</i>	178	<i>zonatus</i>	177
<i>pulchellus</i>	177	<i>zonifer</i>	178
<i>pulvereus</i>	178	<i>zygopterus</i> , <i>Galeorhinus</i>	14
<i>rathbuni</i>	179	<i>Zyphothycus</i>	261
<i>rubrifrons</i>	177		

