

U. S. DEPARTMENT OF COMMERCE
BUREAU OF FISHERIES

Rare BK
514
11
A15
1935

U. S. Bureau of Commercial Fisheries
REPORT

OF THE



UNITED STATES
COMMISSIONER OF FISHERIES

FOR THE FISCAL YEAR 1935

WITH

APPENDIXES

FRANK T. BELL
Commissioner



UNITED STATES
GOVERNMENT PRINTING OFFICE
WASHINGTON : 1936

172 5522

National Oceanic and Atmospheric Administration

Report of the United States Commissioner of Fisheries

ERRATA NOTICE

One or more conditions of the original document may affect the quality of the image, such as:

Discolored pages

Faded or light ink

Binding intrudes into the text

This has been a co-operative project between the NOAA Central Library and the National Marine Fisheries Service (NOAA Fisheries). To view the original document, please contact the NOAA Central Library in Silver Spring, MD at (301) 713-2607 x124 or www.reference@nodc.noaa.gov.

LASON

Imaging Contractor

12200 Kiln Court

Beltsville, MD 20704-1387

November 19, 2004

NOTE

The first section of this volume, entitled "Bureau of Fisheries", constitutes what was known in years prior to 1933 as "Report of the Commissioner of Fisheries." Since then, in the interests of economy, it is a reprint from the "Annual Report of the Secretary of Commerce." The pagination, therefore, is the same as that of the Secretary's Report, rather than beginning with page I.

ERRATA

Page 114: The date in footnote 2 should be 1933 instead of 1833.
Page 287: The footnote at the bottom of the page should be omitted.
Page 331: Page numbers in Table of Contents, Mackerel should be page 344 and Pike-petchea should be 370.

CONTENTS

| | Page |
|---|---------|
| BUREAU OF FISHERIES. By Frank T. Bell. (Issued Jan. 29, 1936.) (See Note on p. ii)----- | 87-111 |
| ALASKA FISHERY AND FUR-SEAL INDUSTRIES IN 1934. By Ward T. Bower. Appendix I. (Issued Sept. 16, 1935)----- | 1-73 |
| FISHERY INDUSTRIES OF THE UNITED STATES, 1934. By R. H. Fiedler. Appendix II. (Issued Oct. 11, 1935)----- | 75-330 |
| PROGRESS IN BIOLOGICAL INQUIRIES, 1934. By Elmer Higgins. Appen- dix III, Administrative Report No. 21. (Issued Jan. 29, 1936)--- | 331-399 |
| PROPAGATION AND DISTRIBUTION OF FOOD FISHES, 1935. By Glen C. Leach and M. C. James. Appendix IV, Administrative Report No. 22. (Issued May 7, 1936)----- | 401-427 |

U. S. DEPARTMENT OF COMMERCE

BUREAU OF FISHERIES

HEADQUARTERS STAFF, 1934-35

Commissioner

FRANK T. BELL

Deputy Commissioner.—CHARLES E. JACKSON.

Chief Clerk.—FLOSSIE WHITE.

Chiefs of Divisions:

Fish Culture.—GLEN C. LEACH.

Inquiry Respecting Food Fishes.—ELMER HIGGINS.

Fishery Industries.—R. H. FIEDLER.

Alaska Service.—WARD T. BOWER.

Black Bass and Anglers.—TALBOTT DENMEAD.

Director of Aquarium.—FRED G. ORSINGER.

Editor.—CLIFFORD F. MAYNE.

Librarian.—LOUISE BEITZELL.

Publications.—BARBARA ALLER.



BUREAU OF FISHERIES

The fishery industry has not shown improvement in recent years to the extent apparent in several other food industries. In fact, during the fiscal year 1935 the fishery industry lost some of the gains which were made during the previous fiscal year. This has been especially true with the fishermen and distributors of so-called "market" fishes. Included in this category are such fishes as haddock, mackerel, halibut, mullet, red snappers, lake herring, and others. As a rule, these fishes are marketed in the fresh and frozen condition. Partial reports from the principal ports indicate a larger quantity of some of these fishes have entered the market during the past year, but the prices received by the fishermen have been less. By far the greater number of our fishermen are engaged in catching market fishes and in many instances these fishermen are unorganized.

During the past fiscal year some improvement has been noted among fishermen catching and selling "cannery" fishes, such as salmon, pilchards (sardines), Pacific coast mackerel, tunas, and others. This may be due to better organization among these fishermen as well as to the fact that canned fish is finding a better demand among consumers. Improvement among the sardine, Alaska herring, and menhaden fishermen, no doubt, has been due to a recent act of Congress, which has brought about a brisk demand in this country for domestically produced fish oils.

The fisheries of the United States and Alaska are prosecuted on the high seas and in the territorial waters of the Atlantic and Pacific Oceans and in the Gulf of Mexico and their adjacent waters, as well as in the Great Lakes and other interior waters. Available data for 1933, when the most recent detailed catch surveys were made, indicate an appreciable increase in both the volume and the value of the catch as compared with the previous year. Statistics of the catch in the New England, Middle Atlantic, Chesapeake, and Pacific States, and Alaska were collected for 1933, and when considering the combined catch of these sections alone an increase of 13 percent in the volume and also 13 percent in the value of the catch in 1933 is indicated as compared with the same sections in the previous year.

Based on the most recent surveys, our commercial fisheries gave employment to about 118,000 fishermen, and their catch aggregated 2,899,908,000 pounds, valued at \$60,218,000. There were increases in most of the groups of prepared products. The output of canned products amounted to 533,212,000 pounds, valued at \$59,800,000, representing an increase of 28 percent in volume and 37 percent in value as compared with 1932; the output of fishery byproducts was valued at \$17,466,000, representing an increase of 40 percent; and the production of frozen fishery products amounted to 95,874,000 pounds,

estimated to be valued at \$8,000,000, representing an increase of 4 percent in quantity and 14 percent in the estimated value. The output of fresh and frozen packaged fish and shellfish amounted to 129,608,000 pounds, valued at \$17,294,000, and cured fish, 104,310,000 pounds, valued at \$12,823,000. It is estimated that about 500,000,000 pounds of fresh fishery products (excluding packaged fish and shellfish), valued at about \$40,000,000, were marketed during 1933. The total marketed value to domestic primary handlers of all fishery products in 1933 is estimated at about \$155,000,000.

Imports of fishery products for consumption in the calendar year 1933 were valued at \$30,462,000, which is 3 percent more than in the previous year, while exports of fishery products were valued at \$8,339,000, or 7 percent more than in 1932.

FISHERY ADVISORY COMMITTEE

During the year the Secretary of Commerce appointed a Fishery Advisory Committee, representing various branches of the fishing industry, various geographical sections of the country, and related interests including science, medicine, and transportation, for the purpose of considering fishery problems from a national standpoint and offering recommendations for promotion and development.

The first meeting was held on May 6 under the chairmanship of Bernarr MacFadden and various subcommittees were appointed to give detailed study and to later report on specific fields. Committees were appointed on the protection of migratory fish, on game fish, on the relation of the fisheries to other water uses, and on scientific research. Committees were also appointed on food standards, on marketing and distribution, on international trade, and on education and publicity. All of these committees presented preliminary reports and drafted a program of future activities to be the subject of reports at the fall meeting in 1935.

EFFECTUATING A BETTER RELATIONSHIP BETWEEN COMMERCIAL AND SPORT FISHERMEN

From time to time considerable animosity has existed between commercial and sport fishermen. At times the sport fishermen have accused the commercial fishermen of depleting certain fisheries while, on the other hand, the commercial fishermen have censured the sport fishermen for trying to take away their means of livelihood. This, no doubt, has led to the enactment of some legislation parading under the guise of "conservation", which in some cases has resulted only in transferring the fishery problems from the hands of one group to another. For this reason there is a need for greater amity between these two groups, to the end that our great natural fishery resources will be prosecuted wisely. It is important that both groups give coordinated consideration to measures intended to sustain our fish supply. During the past year headway was made in this direction.

Members of the National Fishery Code Authority, representing the commercial fisheries, met with members of the National Planning Council of State Fish and Game Commissioners at the council's meeting at Montreal in September 1934. At this meeting various common problems facing the fisheries were discussed and the ground-

work laid for further cooperation in effectuating the conservation and sustained production of our fishery resources.

A local meeting similar to the above was held in Baltimore early in 1935 which was attended by commercial fishermen and anglers from the Chesapeake Bay section. Here various measures were discussed which were directly connected with the activities of these groups, and the discussion resulted in the meeting sponsoring fishery legislation of benefit to all concerned. Other such meetings have been held in various other sections of the country.

COOPERATION WITH STATES AND EDUCATIONAL INSTITUTIONS

For many years the Bureau has had valuable informal cooperation in fishery investigations with many institutions, both public and private, but the acceptance of formal cooperation was not authorized until the passage of the act of Congress approved May 21, 1930.

In connection with fish-cultural operations the closer cooperative relations with the States as initiated and developed by the National Planning Council of Commercial and Game Fish Commissioners has been unusually effective. During the past year developments of this nature have been largely a consolidation of cooperative procedure and a refinement in the working details of the various cooperative arrangements existing with a majority of the States. Arrangements have been made for several additional States, particularly Iowa, to review Federal applications for fish. The State of Vermont has afforded generous help in a financial way in meeting the costs of operation at the extensive brook-trout egg-producing plant maintained by the Bureau in the White Mountain National Forest, N. H. Tennessee and West Virginia have furnished food for fish held at the Bureau's hatcheries and have handled the distribution of these fish when they were ready for release. A number of the Western States have continued to supply helpful cooperation in the collection of trout eggs and are compensated therefor by receiving a proportion of the eggs taken. There has been a joint operation of trout and shad hatcheries in the State of South Carolina.

Many of the cooperative fish-cultural arrangements are largely a matter of administrative detail and are too voluminous for a full recital, but their effects have been plainly evident in a greater efficiency. In fact, it may be said that in several instances State and Federal fish-cultural facilities are practically interchangeable as far as stocking is concerned, and real distinction lies only in their being administered by different agencies.

California has continued its cooperative scientific investigations with the Bureau concerned with the trout supply, looking toward the more adequate supply of streams and the more rational regulation of fishing within its borders. New York State has continued its cooperation in the conduct of studies of the nutritional requirements of trout to improve hatchery practices in feeding and rearing. Oregon has cooperated with a Bureau investigator in a study of fish diseases and hatchery practices. North Carolina, Connecticut, and Washington have assisted in investigations for the restoration of oyster beds in their coastal waters; and Georgia, Louisiana, and Texas have taken an active part in the study of the important shell fisheries of the South Atlantic and Gulf coasts. Despite the fact

that State budgets have been reduced proportional to Federal budgets for these projects, this cooperation has been very effective and is greatly appreciated.

Another source of most valuable cooperation is provided by educational institutions, chiefly State universities. Laboratory quarters for the Bureau's investigative staffs are provided by Harvard University as headquarters for North and Middle Atlantic fishery investigations. University of Michigan has headquarters for the Great Lakes fishery investigations, University of Utah for fishery investigations in the intermountain section, University of Missouri for investigation of interior waters, and Stanford University for California trout investigations. The Wisconsin Natural History Geologic Survey has cooperated with the Bureau in many ways. Yale University, Cornell University, the University of Washington, and the Oregon State Agricultural College have likewise provided quarters or other facilities for investigative work.

In the technological work of the Bureau many State agencies have cooperated in extending their facilities for the prosecution of these studies. State universities, hospitals, agricultural experiment stations, and other State institutions of research have contributed personnel and laboratories in various projects. Among the institutions represented in this work during the past year were the South Carolina Food Research Commission and State Medical College, the Massachusetts State Agricultural College, the New York State College of Agriculture, Washington State College and Agricultural Experiment Station, the University of Washington, George Washington University, and the University of Maryland.

The value of this type of cooperation in the scientific fields cannot be overestimated. The Bureau's investigators receive, in addition to actual laboratory and office quarters, the use of university libraries, advice and assistance from the university faculties, and many other courtesies which stimulate a community of interest in technical problems of the fisheries. The universities thus contribute to research of practical value and application to their own communities, and their graduate students receive stimulation and advice in research problems similar to those of the Bureau and frequently part-time or temporary employment in Bureau projects, all of which contributes to the progress of aquatic biology and technology in the United States.

In the Bureau's statistical research of the fisheries of the Great Lakes and Pacific Coast States and of the States of Maryland and Virginia such exceptional cooperation has been obtained from State fishery agencies in recent years that the Bureau has conducted only limited surveys to supplement the data available from the States.

COOPERATION WITH OTHER FEDERAL AGENCIES

The Bureau's program has always been closely correlated with work of the various Bureaus of the Department of Commerce, from which it secures assistance of various kinds, chiefly relating to the promotive aspects of commercial fishery investigations.

At the close of the year plans were being developed for one project in North Carolina whereby a hatchery would be constructed and turned over to the Bureau for subsequent operation, with the object of providing fish for the surrounding territory.

The Bureau has been called on to make a survey in the Tennessee Valley to aid the Tennessee Valley Authority in developing a program of conserving fishery resources and establishing a hatchery system.

The United States Army Engineers have given sympathetic consideration to the Bureau's recommendations, with regard to developments in the upper Mississippi River area. This has been based upon a hope that the dams comprising part of the 9-foot channel development might be modified so as to provide extensive propagating ponds for the production of fish native to that area. In connection with the work on the Bonneville Dam (later discussed in more detail) the War Department allotted funds from its construction appropriations that the Bureau might carry on studies and design ways for passing the run of salmon over this new dam.

Through the cooperation of the Navy Department the annual supplies for the Pribilof Islands were forwarded from Seattle on the U. S. S. *Sirius*, and the season's take of sealskins was brought out by this vessel on its return voyage. The United States Coast Guard also rendered valuable assistance in maintaining a patrol for the protection of the fur seals and in performing other services.

The nature of the work performed by the National Park Service, Forest Service, the Office of Indian Affairs, the Bureau of Reclamation, and the Bureau of Biological Survey gives them of necessity an interest in the Bureau's fish-cultural work. The stocking of streams and lakes under the control of the foregoing agencies was an important feature of their administration. Consequently, it has been necessary to maintain closer contact with these establishments, and it is felt that the Bureau has been of real service in numerous instances.

While these cooperative relationships are of a more or less standing nature, there have been contacts with the newer emergency organizations which offer prospects of mutual value. The Agricultural Adjustment Administration, insofar as its work covers the utilization of marginal lands, has called on the Bureau in several instances to plan a program for fish propagation and the provision of angling.

Various members of the Division of Fishery Industries assisted other Federal agencies in the conduct of technological and economic studies relative to the fishery industry. In this connection special aid was rendered the National Recovery Administration in the development of fishery codes, a member of the Bureau being on detail to that Administration until October 1, 1934, to supervise and assist in this work; the Department of Agriculture, in various nutrition studies; the Federal Emergency Relief Administration, in fishery relief projects; the Federal Surplus Relief Corporation, in supplying data on the fishery food situation; and the Reconstruction Finance Corporation, on loans to the fishery industry. One member of the Bureau has been appointed a member of the food survey committee of the Department of Agriculture, which investigates the supply and price situation of surplus foods.

The Bureau has had the cooperation of the Bureau of Agricultural Economics in the collection of statistics on the volume of cold-storage holdings of fish; and the cooperation of the health authorities in Washington, D. C., in obtaining the volume of fish handled at the Municipal Fish Wharf and Market in this city. In another instance the Bureau obtained figures on the volume of the quarterly holdings of fish oils for the Bureau of the Census.

CONSTRUCTION ACTIVITIES

At the start of the year there were small unexpended balances in approximately 12 allotments for the repair and reconditioning of fish hatcheries. These allotments had been made by the Public Works Administration during the previous year, and the work was carried to a conclusion during the fiscal year 1935.

A considerable portion of a similar allotment for the continued development of the Leetown (W. Va.) experimental hatchery remained available and was expended in part for the construction of bass and trout ponds, remodeling of one of the buildings, and construction of a new dwelling.

Early in the year a new Public Works allotment of \$75,000 was received for the construction of a pond-fish hatchery at Harrison Lake, Va., 26 miles southeast of Richmond. Work was started during the winter and at the close of the year approximately 15 acres of ponds were virtually completed and other developments were well along to the point where the establishment could go into active operation during the fall of 1935. Arrangements were being made for the propagation of shad as well as pond fish at this point. Two dwellings as well as various service buildings comprised the structural improvements. The water supply, which is obtained from a lake, is fed through a canal.

In general, the construction and improvements effected during the fiscal years 1934 and 1935 can be credited in part for the increased hatchery output attained during the latter year.

CONSERVATION OF WHALES

The Multilateral Convention for the Regulation of Whaling agreed to by the economic committee of the Council of the League of Nations on September 24, 1931, became effective on January 16, 1935, in consequence of the deposit of the ratification of the convention by Great Britain and Northern Ireland on October 18, 1934. The convention was ratified by other nations as follows: United States, July 7, 1932; Norway, July 18, 1932; Union of South Africa, January 11, 1933; Switzerland, February 16, 1933; and Mexico, March 13, 1933. In addition to these ratifications, the following have signified adherence to the convention: Nicaragua on April 30, 1932; Sudan, April 13, 1932; Monaco, June 17, 1932; Brazil, November 21, 1932; and Egypt, January 25, 1933.

A bill has been introduced in Congress to give effect to the convention.

BIOLOGICAL FISHERY INVESTIGATIONS

The major portion of the research looking toward the conservation of the fishery resources of the United States is conducted by the Division of Scientific Inquiry. Some of the States maintain research stations and laboratories and some a single fishery biologist, but as a general rule the State governments do not conduct fishery research on any considerable scale. Many, however, contribute or cooperate effectively in scientific work conducted by the Bureau as has been mentioned previously. This cooperation should be developed and extended.

Research activities are about equally divided among three important fields. These are (1) aquicultural investigations including studies on the improvement of hatchery technique for both cold and warm water fishes and the planning of rational stocking policies in interior waters, (2) commercial fishery investigations concerned with the changes in abundance of the food fishes of the coastal areas and the Great Lakes, the detection of overfishing, and the correction of abuses in the commercial fisheries, and (3) shell-fishery investigations directed toward improving the quality of oysters, increasing production by cultural methods and combating oyster pests.

The technical staff of the Division of Scientific Inquiry numbers 45 trained experts, with perhaps an equal number of temporary assistants at some times of the year, but the problems of the fisheries are as numerous as the species that are exploited. Some 30 important food and game fishes are the subjects of continuing research, but to give adequate attention to the fisheries in all sections of the United States—coastal, interior, and in Alaska—would require double the number of present investigators.

AQUICULTURAL INVESTIGATIONS

The problems of the fresh-water fisheries throughout the country at present are concerned chiefly with increasing hatchery output and properly stocking interior waters. In carrying out the latter program, the discovery of suitable natural conditions by means of stream surveys, the conditioning of streams to improve their fish carrying capacity, and the control of stream pollution are vital to its success.

Pond-fish culture.—Investigations in pond-fish culture were conducted during the past year at the Natchitoches (La.) fish hatchery. These investigations were designed primarily to throw light on the problems which have arisen in connection with pond culture in the Southern States where conditions in many respects are quite different from those found in hatcheries in the North. Experiments during 1934, designed primarily to furnish a comparison of the value of different fertilizers show that cottonseed meal and other fertilizers are beneficial in producing a large crop of forage organisms on which the fish feed, but that food production alone is not directly correlated with fish production. In heavily fertilized ponds large numbers of small fish may be produced for early distribution, but additional forage food is required to carry the bass through the summer until October or November.

Trout culture.—Trout cultural investigations have been conducted at the experimental hatcheries located at Pittsford, Vt., and Leetown, W. Va. Here experiments were designed to determine the most economical ration for the feeding of both fingerling and older trout. Detailed observations were made on the food requirements of fish of the same species at different ages and of the different species of trout as a guide to more economical general hatchery practices. Selective breeding experiments have also been conducted at these two stations to secure a superior strain from the point of view of growth, egg production, and disease resistance. Future brood stock was selected from fish of known lineage for propagation of future supplies.

Fish nutrition.—At the United States Fishery Station, Cortland, N. Y., cooperative investigations of fundamental problems of trout

nutrition, which have been under way for several years with the State of New York, Cornell University, and the Bureau, have been continued. As a result of these investigations, improved trout diets have been devised and numerous problems regarding the digestibility of food materials have been solved that should aid in reducing cost of hatchery operations throughout the country. Similar study of hatchery and stocking problems have been undertaken on a cooperative basis in California, concerned primarily with problems of managing the supply of steelhead trout in coastal streams and of the various trout species in the high mountain regions.

Fish pathology.—Progress has been made during the year in the study of fish diseases that annually take heavy toll of the fish in hatcheries. Experiments on two of them, the so-called "ulcer disease" and "blue sac disease", have been made in an effort to find preventives or cures, and a clinic or disease service has been established to aid fish culturists in their practical problems of disease prevention. In this service prompt diagnosis of diseases is given by mail when fishes are sent to the laboratory for examination.

Stream surveys.—Utilizing funds provided by the Public Works, stream surveys were conducted by 16 parties operating in forests and parks in various parts of the country. The primary purpose of the surveys was to provide an inventory of conditions that affect the fish population in each lake and stream. With this information at hand, it has been possible to determine to what species of fish each body of water is best adapted and the number it can support most advantageously. One-sixth of the waters of our national forests have been surveyed during the past year, with the result that many miles of water formerly stocked have been shown to be worthless and in some localities suitable waters are neglected. Greater economy and efficiency can, therefore, be introduced into the stocking program.

In connection with the stream surveys, extensive work in stream improvement was carried out in the national forests for the purpose of increasing the carrying capacity and food production in streams. Since there are few reliable data on which to base an estimate of the true value of stream improvement, the work was primarily experimental, and its real result will be apparent only after several years.

Pollution of waters.—One of the most serious limitations on the maintenance of abundant stocks of food and game fishes in interior waters and the chief hindrance in stocking these waters with additional supplies is pollution from domestic and industrial sources. The destruction of fish life by polluting substances, already acute in many localities, is rapidly growing throughout the more densely settled and industrialized sections of the country.

Without authority, until the passage of the "Coordination Act" March 10, 1934, the Bureau has made during the past year, with an allotment of funds from the Public Works Administration, a very promising start on an investigation of pollution problems related to the conservation of aquatic life. The standards of water purities for the protection of aquatic life differ from those applying to industrial uses of water or for the protection of public health, and are to a certain degree more exacting.

Lacking suitable standards for the evaluation of polluted streams as fish habitats, research has been directed during the past year to a detailed study of the physiological effects of various polluting substances and the establishment of standards of purity for the maintenance of an abundant fish fauna. A manuscript has been completed for publication that establishes these minimum standards for fish as a guide to efforts at stream-pollution abatement.

In addition to the laboratory studies involved in this work detailed surveys of actual conditions in the stream inimical to fish life because of pollution have been made in widely scattered areas throughout the Mississippi drainage.

COMMERCIAL FISHERY INVESTIGATIONS

Although the total yield of the commercial fisheries in the United States has been maintained, and even increased during the past quarter of a century, many of the important commercial species are undergoing progressive depletion. The Bureau's investigations of commercial fisheries are organized, therefore, as continuing observations of the condition and trend of the important commercial fisheries, rather than as disconnected and temporary inquiries or surveys. Their purpose is to trace from year to year variations in the supply of food fishes resulting from natural causes, the early detection of the first signs of serious depletion, and the recommendation of legislation or control of fishing operations and the correction of abuses to maintain the fisheries on the basis of sustained yield. Such investigations are the sole guide to wise conservation policies.

Haddock.—In the North and Middle Atlantic sections biological and statistical studies have been continued on the fisheries for haddock, mackerel, squeteague, scup, and other shore fishes from the offshore Nova Scotian banks to North Carolina. Chief attention has been given to measuring accurately the fluctuations in the supply of haddock available for fishermen. This catch analysis shows that the haddock population on the principal northeast banks was at a low level of abundance in 1935. In fact, marketable haddock were not more than one-third as abundant as during the period 1926–28. As a result of this scarcity, the United States fleet is now forced to fish on banks off the Nova Scotian coast, a distance upward of 600 miles from their home ports. Numerous biological data on age composition, growth rates, and mass movements of the haddock population indicate that the scarcity is due to intensive commercial fishing which is removing haddock from the sea at a more rapid rate than natural replacement by reproduction and growth. One of the causes of this scarcity is the loss of large quantities of undersized and immature fish destroyed in the course of regular trawling operations.

Mackerel.—Mackerel investigations have been designed to discover the best possible use of the fluctuating supply that nature provides. One means is to protect the future abundance of mackerel so that the industry may adjust itself to the expected supply, and the other is to find out if possible whether a more stable and more valuable supply of large mackerel would result from reducing the intensity of fishing on the young. Predictions of the general abundance of mackerel in advance of each season have been made

annually for some years. During the fiscal year 1935 predictions for the previous season were checked by analyzing catch statistics to note the effect of the curtailment program of the industry under the code of fair competition. Results of the check indicate that the prediction of abundance was correct within 10 percent. A prediction for the 1935 season was prepared in May of the current year. The rate of catching mackerel in the early part of the season was entirely in accord with the prediction which anticipated an unusually heavy run during the early months. Extensive biological observations on the life history of the mackerel have been continued and a comprehensive report that will guide future conservation policies is in the process of preparation.

Shore fish of the Middle Atlantic States.—Further observations have been made regarding the state of the supply of the squeteague, scup, sea bass, flounders, and minor species taken in great quantities in the inshore regions from Cape Cod to North Carolina. Further details of the life history and migrations of the most important species, the squeteague or weakfish, perfect our understanding on the relationship between the centers of production in the southern portion of the range and the yield of the fisheries in New Jersey and New York. Fully one-half of the total supply taken north of Delaware Bay is derived from spawning in southern waters. Conservation, therefore, must be promoted by regulating the strain of exploitation from Chesapeake Bay southward, rather than by drastic restriction of the catch in northern waters.

Regulation of the New Jersey shore fishery is complicated by the rapid development of the offshore trawl fishery of Virginia. The catch of this important branch of the industry has been shown to depend not only on the abundance of fish but on the degree of availability which is affected by changes in water temperatures. Wastage of immature fish in southern waters constitutes a heavy drain upon the supply. Depletion has not yet been demonstrated, although its possibility as a result of severe exploitation in the shore areas in the summer and the offshore areas in the winter is clearly recognized.

Shrimp investigations in the South Atlantic and Gulf States.—Landings of shrimp, the most important fishery in the South Atlantic and Gulf States, have declined since 1930. No evidence has been secured of the depletion of the supply in the Gulf area, but on the South Atlantic coast great fears are felt for the future of the fisheries.

Previous investigations have outlined in some detail the life history and development of the most important species of shrimp. During the past year chief attention has been given to a study of mass movements of this species, since local variations in supply, particularly on the Atlantic coast, are very evident. Efforts are being made to trace the movements of the shrimp by the changing composition of populations and by direct tagging experiments. Tagging, however, is a difficult process and considerable attention has been given to the perfecting of a technique for this purpose. It is suspected that a considerable supply of shrimp may exist in offshore waters during the winter period, but the lack of a suitable research vessel prevents an immediate attempt to survey offshore fishing grounds or to follow shrimp migrations.

Great Lakes fisheries.—Owing to the continued curtailment of the budget, no field work was conducted on the Great Lakes during the past year. The staff, however, has been engaged in analyzing material gained in previous years' observations and in assisting in legislative matters with the States.

The most urgent need in this region today is concerned with uniform regulations of commercial fishing. Many attempts in the past have failed to secure coordinated legislation among the States, but progress in improving conservation laws has resulted from many conferences during the past year.

An intensive statistical study of the commercial fisheries in Great Lakes waters of the State of Michigan has been continued. These studies, based on detailed records secured by the State Conservation Department, demonstrate the rapid depletion of whitefish supply by deep trap nets, and the statistical evidence of their destructive action was directly instrumental in securing legislation regulating and restricting the use of this gear.

A study of the life history of the important commercial species of the pike perches is nearing completion, and similar studies of the yellow perch were pursued.

Pacific coast fishery investigations.—One of the most important undertakings of the Division of Scientific Inquiry is a comprehensive program of investigation of the salmon fisheries of the Columbia River, begun in July 1934. The aims of this investigation are:

1. To provide for rational regulation of a commercial fishery in the lower Columbia in order to permit an adequate escapement of fish for spawning purposes.
2. To assure free passage of spawning migrations over the various dams in the Columbia River.
3. To provide for increased spawning in the upper tributaries by removing all unnecessary obstructions and by rehabilitating spawning grounds formerly productive.
4. To improve artificial propagation of salmon and to increase production by reestablishing runs in suitable tributaries.
5. To improve survival by reducing the pollution hazards.
6. To prevent loss of downstream migrants in unscreened irrigation diversions.

A complete survey of the river system has been undertaken in order to assess these various problems, including a detailed analysis of the commercial catch from earliest times to the present. Detailed recommendations were offered by the Bureau to the War Department for the construction of fish protective works at Bonneville Dam including four gravity fish ladders of an improved type and three fish locks or hydraulic lifts for conveying spawning fish over the dam, and a series of fingerling bypasses to protect downstream migrants; \$3,200,000 has been allotted for this construction and although the problems of fish protection are by no means solved, the Bureau is confident that effective protection of the Columbia River salmon runs at this point will be afforded.

With an allotment from the Public Works Administration, the Bureau has constructed screens for three important diversions on the Yakima River for the protection of young salmon from loss in irrigation canals of the Reclamation Service. Two additional screens were constructed on other watersheds for a similar purpose.

The studies recently undertaken on the salmon fisheries of Puget Sound have also been continued with special attention given to the sockeye and coho salmon fisheries. Details of the life history of these fishes have been studied as a basis for future protection, and a statistical analysis of the sockeye salmon fishery of the Frazer River area was made which demonstrates the severe depletion of this important stock of fish and emphasizes the need for international regulation of the fishery.

Studies have been continued on the two most important species of salmon in Alaska, namely, the red salmon and the pink salmon. Studies of the red salmon of Karluk River system, which require continued observations for many years, were conducted during the past year for the purpose of determining production from known escapements of spawning fish. Records for eight generations are now available, showing wide variations in production, varying from a ratio of return to escapement from 0.6 to 1 to 5.6 to 1. There seems to be little possibility of regulating spawning escapement so as to produce consistently a large population. Hence, efforts are being directed to the determining of factors which determine survival in order to control natural forces and to regulate the fishery in the interest of conservation.

Similar studies of the changing abundance of pink salmon have also been continued in southeastern Alaska. Results of this study show that wide fluctuations may be expected in the yearly abundance of this species. Studies of the effect of rainfall and stream flow, as it affects spawning conditions, and population density in the sea, affecting rate of growth and time of migration, hold the key to better regulation of the fishery.

Further studies have been pursued of the depletion of the herring in limited areas in Alaska, with a view to determining the units of population that make up the herring supply. Supplementing bio-statistical analyses of the races of herring, large-scale tagging experiments have been undertaken to trace the migrations and to find the limits of movement of the various stocks. Recovery of tags has been facilitated by the perfection of an electric device for detecting tagged fish in commercial catches. Detailed records of actual migrations are now being secured to guide the drafting of more effective fishery regulations.

SHELL-FISHERIES INVESTIGATIONS

Investigation of the various problems relating to oyster culture were carried out in all the principal centers of the industry of the Atlantic and Pacific Coastal States and the Gulf of Mexico. The work was expanded by an experimental study of the effects of oil pollution in Louisiana waters, which demonstrates the dangers to aquatic life of the development of petroleum supplies in coastal waters.

In the New England area the larger part of the work was a continuation of the investigation of the growth and fattening of oysters started by the Bureau in the spring of 1932. The effects of water temperature and varying amounts of various chemical constituents of sea water were studied in relation to nutrition of oysters and the storing of glycogen or animal starch. These studies are funda-

mental to practical methods of improving the quality of market oysters through artificial feeding or the control or selection of the environment.

A number of investigations and surveys were conducted in the South Atlantic States with a view to rehabilitating public oyster beds in North Carolina and Florida, and studies on the cultivation of the native *Olympia* oyster in Puget Sound were continued. A practical result of this latter undertaking has been the prediction of the precise time of setting of oyster larva in Puget Sound, enabling the oystermen to regulate their planting activities so as to secure a maximum set of seed oysters.

In order to answer many requests regarding the cultivation of hard clams, a minor research project was undertaken on the life history of this valuable shellfish. Much information has been secured in Long Island Sound regarding its life history, habits, and physiology, as a basis for developing practical methods of propagation.

Oyster pest control.—Near the end of the year a special appropriation of \$100,000 became available for a study of means of controlling various pests of the oyster that have reached epidemic proportions, menacing the existence of cultivated and natural beds. A staff of competent investigators was recruited and headquarters were established at Milford, Conn., to study means of controlling the depredations of starfish; at Bivalve, N. J., and Beaufort, N. C., for the control of the drill, which is the most important oyster enemy in the Middle Atlantic section; and at Apalachicola, Fla., for combating the leech, a flat worm which has recently become extremely destructive of valuable oyster supplies in Florida. The steamship *Kittery* was transferred from the Shipping Board, without exchange of funds, to the Federal Emergency Relief Administration for use in the drill control work in the vicinity of Norfolk, Va. This investigation will continue throughout the next fiscal year with the hope of demonstrating practicable methods that can be applied on a large scale by the Federal or State Governments and by private oyster planters in controlling the most important enemies of the oyster industry.

ALASKA FISHERIES SERVICE

ADMINISTRATION OF FISHERY LAWS AND REGULATIONS

The control of commercial fishing in Alaska was administered as usual in accordance with general laws and regulations for the conservation of the fisheries. A consistent program has been followed, the main object of which is to assure an adequate escapement of brood fish to maintain a maximum supply. A breeding reserve of 50 percent of the salmon runs is regarded as the minimum requirement. Careful observations of the runs are made each season, and when signs of depletion are evident in any locality appropriate remedial measures are initiated.

An unusual abundance of salmon in practically all areas characterized the 1934 season, and few additional restrictions on commercial operations were imposed. In some instances existing restrictions were relaxed as the season advanced. Revised regulations for the protection of the fisheries in 1935 were issued on January 19, the most important changes of which were the prohibition of com-

mercial fishing in the Bristol Bay region during the red-salmon runs in order to build up the weak cycle appearing in the calendar years divisible by 5, and the closing of 31 trap sites to permit a greater escapement of salmon to the spawning grounds and to check the tendency of this form of gear to monopolize fishing in certain localities.

Twelve regular and 164 temporary employees were engaged in patrolling the fishing grounds in the 1934 season in addition to the crews of 12 vessels of the Bureau and 3 chartered vessels. More extensive use than in previous years was made of chartered airplanes as an auxiliary patrol and for general supervision of the Bureau's work and inspection of the spawning areas.

Weirs for counting the escapement of spawning salmon were operated in 12 typical salmon streams in the calendar year 1934. An allotment of Public Works Administration funds enabled the reestablishment of several of these structures, both in 1934 and 1935. The counts of salmon thus made furnish valuable information in connection with life-history studies, as well as providing a means of determining the ratio of escape to catch. Scientific investigations in regard to salmon, herring, and clams were continued.

Some work was accomplished in the removal of log jams and other barriers that prevented the passage of salmon upstream to the spawning beds. In certain localities, also, natural propagation conditions were improved by the destruction of predatory fishes that feed upon the salmon eggs and fingerlings. This work was done partly by weir operators and stream guards, but chiefly through the appropriation of funds for the purpose by the Territorial Legislature and by contributions of local packers in the Bristol Bay and Yakutat regions. At its 1935 session the Territorial Legislature again appropriated \$15,000 for clearing salmon spawning streams and for the destruction of predatory enemies of salmon, to be expended under the same provisions as in the case of similar previous appropriations.

ALASKA SALMON HATCHERIES

Only one salmon hatchery was operated in Alaska in the fiscal year 1935—that of the Pacific American Fisheries at Hugh Smith Lake, in the southeastern area. At this hatchery 10,221,000 red-salmon eggs were collected, from which 9,860,000 fry were produced and liberated in Alaska waters. Under the provisions of the Alaska fisheries act of June 26, 1906, the owners of private hatcheries receive a rebate on license fees and taxes on their catch and pack of salmon at the rate of 40 cents for each 1,000 red or king salmon fry liberated.

PRODUCTS OF THE FISHERIES

Notwithstanding the fact that some branches of the industry were handicapped by reason of the longshoremen's strike on the Pacific coast, which for a time caused a shortage of containers and other supplies, and by price disagreements between fishermen and packers in southeast Alaska and the Copper River area, the total output of fishery products was the largest in the history of the Territory. This exceptional harvest may be attributed especially to the abundance of salmon in virtually all producing centers and also to increased activity.

Salmon products comprised about 80 percent in quantity and 92 percent in value of the entire output of the Alaska fisheries. The total production of salmon amounted to 372,585,000 pounds valued at \$38,749,000, or an increase of about 42 percent in quantity and 32 percent in value over the output of the preceding year. About 96 percent of the salmon products in the 1934 season consisted of canned salmon, the pack amounting to 7,481,830 cases, valued at \$37,611,950. Red salmon represented 35 percent and pinks 51 percent of the total pack of canned salmon, as compared with 42 percent for each of these species in 1933.

The production of herring was also above the average, with a record output of meal and oil, inasmuch as large numbers of the fish taken were too small for satisfactory Scotch-curing. Halibut products declined in quantity but increased in value. A whaling station that had been closed since 1930 was reopened and the total output of whale products was more than double that for 1933. All minor species except clams showed an increased production in the 1934 season as compared with the previous year, the most marked gain being in the crab industry.

The total output of Alaska fishery products in the calendar year 1934 was 468,424,000 pounds, valued at \$41,963,000, as compared with an average of 360,097,000 pounds valued at \$35,845,000 for the 5-year period from 1929 to 1933, inclusive. The value of the catch to the fishermen was approximately \$11,707,000, or about \$2,618,000 more than in the preceding year. There were 26,190 persons employed in the various branches of the industry, as against 21,695 in 1933.

ALASKA FUR-SEAL SERVICE

GENERAL ACTIVITIES

The chief activity in the fur-seal service is the annual take of sealskins for commercial purposes during the months of June and July. Killings are confined as far as possible to 3-year-old male seals, of which approximately 50,000 are secured each season. Of the skins obtained on St. Paul Island, the majority are taken by the stripping process and blubbered before curing. The method of skinning by hand is still used altogether on St. George Island, where less than one-fourth as many skins are taken as on St. Paul Island.

Sealing operations were performed by Pribilof Islands natives and by approximately 60 natives from the Aleutian Islands under the direction of a staff of regular employees. Nineteen employees of the Fouke Fur Co. were at St. Paul Island during the season to assist with the washing and blubbering of the sealskins in accordance with the fur-seal contract.

An allotment of \$3,000, by the Public Works Administration, enabled the reconditioning and repair of several buildings used in the sealing industry. Some improvements were made also in the way of construction of new buildings and boatways and the extension of surfaced roads to facilitate transportation of sealskins from the killing grounds to the curing stations.

Delivery of 15 percent of the season's take of sealskins was again made to the Dominion of Canada under the provisions of the

international treaty for the protection of the fur seals. The skins were delivered to a representative of that Government at Seattle in August 1934. The Government of Japan, which is also entitled to 15 percent of the sealskins taken at the Pribilof Islands, continued to have its skins handled with those of the United States, receiving an appropriate share of the proceeds of the sale.

Two hundred sealskins were allotted to the United States as its share of the take by the Japanese Government on Robben Island in 1934. They were received by the Department's selling agents at St. Louis, Mo., in December.

SEAL HERD

As of August 10, 1934, the computed number of animals in the Pribilof Islands fur-seal herd was 1,430,418, an increase of 111,850, or 8.48 percent, over the corresponding figure for 1933.

TAKE OF SEALSKINS

In the calendar year 1934 there were taken on the Pribilof Islands 53,468 fur-seal skins, of which 42,972 were from St. Paul Island and 10,496 from St. George Island. This is a decrease of 1,082 from the number taken in 1933.

SALE OF SEALSKINS

Two public-auction sales of fur-seal skins taken on the Pribilof Islands were held at St. Louis, Mo., in the fiscal year 1935. On October 15, 1934, 10,000 black-dyed, 3,000 logwood brown-dyed, and 6,000 of the new safari brown-dyed skins were sold for \$336,846.25. At the same time three confiscated fur-seal skins, parchments, were sold for \$7.50.

The second sale, held on April 8, 1935, consisted of 12,600 black-dyed, 11,500 safari brown-dyed, and 2,303 logwood brown-dyed skins, which brought a gross sum of \$616,990.75.

Special sales of Pribilof Islands sealskins authorized by the Secretary of Commerce in the fiscal year 1935 consisted of 1,607 black-dyed, 1,843 safari brown-dyed, 1,027 logwood brown-dyed, and 300 raw salted skins, which brought a total gross sum of \$98,460.44. In addition, the Department advertised and sold to the highest bidders for \$621.22 two fur-seal coats, 1 black and 1 logwood brown, that had been used for display purposes. Each coat was made up of seven Pribilof Islands sealskins.

FOXES

The management of blue fox herds on St. Paul and St. George Islands is incidental to the fur-seal industry and requires attention at a time when sealing activities are light. Beginning about the first of December the animals are fed prepared rations and are trapped for their pelts.

During the 1934-35 season there were taken 184 blue and 16 white fox skins on St. Paul Island and 799 blue and 3 white skins on St. George Island, a total of 1,002. Eleven foxes on St. Paul Island and 266 on St. George Island were marked and released for breeding stock. The reserve includes also a considerable number of animals that were not caught in the traps.

In the fiscal year 1935 there were sold 914 blue fox skins, raw, for \$21,935.50; 25 white fox skins, raw, for \$333; 1 blue fox skin, dressed, for \$38; and 1 white fox skin, dressed, for \$24; a total of \$22,330.50. The raw skins were taken on the Pribilof Islands in the 1933-34 season, and the dressed skins were of the 1932-33 season's take which had been used for exhibition purposes.

FUR-SEAL SKINS TAKEN BY NATIVES

Indians dwelling on the American coast of the North Pacific Ocean are privileged to engage in limited pelagic sealing under the provisions of the North Pacific Sealing Convention of 1911. The sealskins thus taken must be authenticated by Government officials before they can enter into commerce. In the 1934 season there were taken and duly authenticated by officials of the respective Governments 290 fur-seal skins, of which 23 were taken by Indians of Washington, 11 by Indians of southeast Alaska, and 256 by Indians of British Columbia. Reports have been received, also, of the authentication of 6 additional sealskins taken in 1933 by Indians of the State of Washington, making a total of 35 taken by them that year.

FUR-SEAL PATROL

A patrol for the protection of the fur seals was maintained by the United States Coast Guard, which detailed seven cutters and two 125-foot patrol vessels to this work in the calendar year 1934. Two vessels of the Bureau also participated successively in the patrol in the vicinity of Neah Bay, Wash., during the spring migration of the herd.

PROTECTION OF SEA OTTERS, WALRUSES, AND SEA LIONS

Regulations issued on July 1, 1934, permit certain limited killing of walruses and sea lions in Alaska under specified conditions. The killing of sea otters is prohibited at all times.

PROPAGATION AND DISTRIBUTION OF FOOD AND GAME FISHES

It is gratifying to report that a very moderate increase in the appropriation available for the propagation of food and game fishes resulted in a noticeable increase in output for the fiscal year 1935. The total distribution of fish and eggs, including fry hatched on a cooperative basis in State hatcheries, approximated 5,000,000,000 in comparison with an output of 3,258,000,000 for the previous year. The 1935 figures are still materially below the maximum output of several years ago which exceeded 7,000,000,000. To a considerable extent, this increase over the previous year was due to the resumption of the activities at the Gloucester (Mass.) station which propagates marine forms. Coupled with its increase there was a slight step-up in the output of some of the varieties, both game and commercial, handled at interior stations.

Of the hatcheries which were closed the previous year owing to economy requirements, those in Alaska and the two Michigan commercial hatcheries remained closed. The output of game fish was closely comparable to the production of these species in 1934.

The output of small mouth black bass, however, a variety most eagerly sought by the sportsmen, was almost twice the production of the previous year. There is no evidence of slackening in the demand for hatchery-produced game fish, and in fact the increased requirements for stocking areas now coming under the control of the Federal Government constitute a serious problem.

There was also a moderate increase in the output of fingerlings, constituting the larger fish with a greater stocking value. The distribution of these larger fish was approximately 4,000,000 greater than last year. The list of the species propagated was practically the same as during the previous year with the exception that no white perch were handled during 1935, but a considerable number of pollock was included in the figures for this year.

The sportsmen's organizations and the general public have largely accepted the requirement that they must themselves meet the cost of transporting the fish from the hatcheries. The problem of distribution has been further simplified by the readiness with which several State fish and game departments have undertaken to handle the fish produced at Federal hatcheries.

PROPAGATION OF COMMERCIAL SPECIES

Marine species, Atlantic coast.—An increase of approximately 1,400,000,000 in the output of cod, haddock, pollock, and winter flounder is to a large extent the result of the reopening of the Gloucester (Mass.) station. However, a large percentage of this increase is represented by eggs fertilized on the fishing boats and planted on the spawning grounds, since funds were not available for full-scale operation of the hatchery. The output of these marine forms represents between 85 and 90 percent of the total output, a ratio which is similar to the average over a number of years. No cod were propagated at the Woods Hole (Mass.) station during the year.

Pacific salmon.—While increases were registered for the output of silver and chum salmon, there was a slight decline in the distribution of chinook variety, and a very large decline in the production of sockeye salmon. This was due to negligible egg collections at the Quinault (Wash.) station, ordinarily the biggest producer of sockeyes since closure of the Alaska stations. The run of fish did not materialize, presumably on account of overfishing in previous years. The number of all species of Pacific salmon distributed in the fingerling stage compares very favorably with the record of previous years.

Anadromous species, Atlantic coast.—The production of shad was virtually the same as in 1934. An output of 860,000 Atlantic salmon in Maine was significantly larger than in the previous year when Canadian eggs were not obtainable. Land-locked salmon were produced in lesser numbers. An increase in output of yellow perch to 450,000,000 was derived from resumption of this activity at the Potomac River station at Fort Belvoir, Va.

Commercial species, interior waters.—Restriction on the activities of the hatcheries handling commercial species of the Great Lakes has continued. Consequently, the output of whitefish and lake trout has remained small, even though the production of whitefish was approximately four times the 1934 figure. The Put-in-Bay (Ohio)

station again handled large numbers of pike perch, collecting over 500,000,000 eggs, the majority of which were incubated and distributed by the State. Two hatcheries in Michigan remained closed.

GAME SPECIES

As a whole, operations with the game species yielded satisfactory results. Inasmuch as some 15 species are included in this category, there were fluctuations among individual forms in comparison with last year's production. Special effort was made to augment the output of bass, owing to their high popularity in all parts of the country. It would be possible to increase the output of trout, particularly by distributing them at smaller sizes, but this would lower their stocking value. However, the problems surrounding the rearing of trout have been greatly intensified by the sharp increase in the price of food for fishes which became evident toward the close of the fiscal year.

RESCUE OPERATIONS

An increase in the amount appropriated for the administration of the Bureau's activities in the Upper Mississippi River Wild Life Refuge was reflected in a virtual doubling of the number of fish rescued. There was handled a total of 47,286,000 fish, comprising 10 species. This is still below the average number which can be rescued when funds are available to cover the entire territory where water conditions make the salvaging of fish trapped in land-locked areas an urgent need. Only 115,900 fish were distributed to other sections, the balance being returned directly to open waters.

AQUARIUM

The aquarium located under the main lobby in the Department of Commerce Building is maintaining its reputation as one of Washington's popular points of interest to visitors. The numerous organizations which gather for conventions in Washington, as a rule, include the aquarium in their itinerary of sight-seeing.

The general exhibits were maintained in much the same manner as during the previous year, and the display of trout has continued to be a point of particular interest. However, from time to time during the year it has been the practice to introduce more novel exhibits, one of which was the 500-pound green turtle. Another interesting novelty was a pair of Piranha—the small man-eating fish of the Amazon Basin.

The aquarium also was of great service in preparing and forwarding foreign shipments of live fish and eggs which were made to Cuba, Puerto Rico, Panama, and Venezuela.

As usual, the display of model hatching apparatus was kept in operation for the purpose of illustrating the methods followed in incubating various types of eggs.

MARKETING INVESTIGATIONS

Per capita consumption of fish.—A study made during the year indicates the domestic per capita consumption of fish amounted to 13.3 pounds in 1931. This is compared with a per capita consumption of meats amounting to 133.2 pounds in the same year.

STATISTICAL INVESTIGATIONS

FISHERIES OF THE UNITED STATES, CALENDAR YEAR 1933

New England States.—During 1933 the commercial fisheries of Maine, New Hampshire, Massachusetts, Rhode Island, and Connecticut employed 17,073 fishermen. Their catch amounted to 499,936,000 pounds, valued at \$13,486,000, an increase of 4 percent in volume, but a decrease of 4 percent in value as compared with the catch in 1932. Landings of fish by American fishing vessels at Boston and Gloucester, Mass., and Portland, Maine, amounted to 267,157,000 pounds as landed, valued at \$6,851,000—an increase of 6 percent in volume, and 13 percent in value, as compared with the preceding year.

Middle Atlantic States.—The commercial fisheries of New York, New Jersey, Pennsylvania, and Delaware in 1933 gave employment to 8,574 fishermen. Their catch amounted to 169,754,000 pounds, valued at \$4,811,000—an increase of 20 percent in volume, and 3 percent in value as compared with 1932. Landings of fish at New York City during 1933 amounted to 25,455,000 pounds. The shad fishery on the Hudson River was conducted by 317 fishermen, who caught 519,000 pounds of shad, valued at \$41,000—a decrease of 2 percent in volume and 20 percent in value as compared with 1931.

Chesapeake Bay States.—In 1933 the commercial fisheries of Maryland and Virginia employed 20,142 fishermen. Their catch amounted to 272,380,000 pounds, valued at \$5,061,000—a decrease of 24 percent in volume and 14 percent in value as compared with the previous year. The shad and alewife fisheries of the Potomac River were prosecuted by 651 fishermen, who caught 1,838,000 pounds of shad, valued at \$149,000 and 6,896,000 pounds of alewives, valued at \$24,000—a decrease of 19 percent in the catch of shad, but an increase of 1 percent in the catch of alewives.

South Atlantic and Gulf States.—No survey was made of the fisheries of this area for the year 1933.

Lake States.—No survey was made of the Lake fisheries (Lakes Ontario, Erie, Huron, Michigan, and Superior, and Namakan and Rainy Lakes, and Lake of the Woods) for 1933.

Pacific Coast States.—The commercial fisheries of Washington, Oregon, and California in 1933 employed 18,673 fishermen. Their catch amounted to 860,161,000 pounds, valued at \$13,988,000—an increase of 53 percent in volume and 47 percent in value, as compared with 1932. The total catch of halibut by United States and Canadian vessels amounted to 45,951,000 pounds, valued at \$2,582,000—an increase of 6 percent in volume and 48 percent in value, as compared with the preceding year.

Mississippi River and tributaries.—No survey was made of this area in 1933.

MANUFACTURED PRODUCTS IN UNITED STATES AND ALASKA, CALENDAR YEAR 1933

Fresh and frozen packaged fishery products.—Based on the most recent data available, the domestic production of fresh and frozen packaged fishery products amounted to 129,608,000 pounds, valued at

\$17,294,000. Important products in this group consist of fresh shucked oysters, 5,687,000 gallons, valued at \$6,864,000; packaged haddock, 35,149,000 pounds; valued at \$3,457,000; and fresh-cooked packaged crab meat, 6,864,000 pounds, valued at \$1,662,000.

Frozen products.—The production of frozen fishery products in 1933 amounted to 95,874,000 pounds, estimated to be valued at about \$8,000,300. The volume of the production was 4 percent greater than in 1932. The most important products frozen with respect to volume were ground fish, halibut, salmon, mackerel, whiting, and shellfish.

Cured products.—Based on the most recent data available, the domestic production of cured fishery products amounted to 104,310,000 pounds, valued at \$12,823,000. Important products in this group were boneless cod, 9,517,000 pounds, valued at \$1,646,000; and smoked salmon, 8,229,000 pounds, valued at \$2,256,000.

Canned products.—Canned fishery products produced in 1933 amounted to 533,212,000 pounds, valued at \$59,800,000—an increase of 37 percent in the value as compared with 1932. Canned salmon alone amounted to 305,398,000 pounds, valued at \$36,242,000. Other important products were tuna and tunalike fishes, sardines, shrimp, mackerel, clam products, and oysters.

Byproducts.—During 1933 the value of the production of fishery byproducts amounted to \$17,466,000—an increase of 40 percent as compared with the preceding year. Important products in this group consist of marine animal oils and meals, and aquatic shell products.

TECHNOLOGICAL INVESTIGATIONS

These investigations cover the general field of food technology as applied to fishery products. They involve the application of the sciences of chemistry, engineering, bacteriology, and general technology to the manufacture, preservation, and utilization of our fishery harvest. Chemical and bacteriological investigations dealing with the preservation of fishery products for food, the utilization of fishery byproducts, the nutritive value of aquatic products, and fish cookery investigations constituted the general scope of studies in fishery technology. Some of the outstanding accomplishments during the past year were: The development of methods for the home canning of some species of fish; increasing the practical value of and simplifying methods for determining the relative freshness of fish; the discovery of further facts concerning the high vitamin potency of fish oils and fish-liver oils; expansion of the possibilities for utilizing salmon cannery waste and other fish waste; discovery of further data on the role of the mineral constituents of fishery products in nutrition; and the issuance of a pamphlet on practical fish cookery.

Preservation of fishery products for food.—Studies in this field, in general, pertain to the chemistry and bacteriology of fish preservation and spoilage, the development of improved methods for handling fresh and frozen fish, studies on the smoking of fish, and the development of methods for canning fish in the home. During the past year, a better method for packing fresh mackerel for shipment was devised which is considered to be superior to the present com-

mercial method of "floating." One difficulty encountered in the storage of frozen mackerel is the tendency of the oil or fat in the fish to become rancid, thereby making the mackerel unfit for use. Obviously, the intensity of rancidity is dependent upon the percentage of oil in the fish, which percentage is subject to considerable seasonal fluctuation, the maximum percentage occurring in August. Studies are being made to prevent or minimize this difficulty. A process was developed for smoking mackerel which produced a high quality product. This process has promise of commercial application and should be of aid in expanding the market for mackerel during periods of unusual abundance. Other studies revealed that losses of food value through leaching would be prevented if fish are covered with waterproof wrappers when packed in ice for shipment. Progress was made in developing a practical method for determining the relative freshness of cod, pollock, and mackerel under commercial conditions. A pamphlet was issued containing practical and tested methods for canning some species of fish in the home.

Bacteriological investigations.—Bacteriology plays a role in nearly all technological investigations of fishery products in that the various experimental products or methods are under bacteriological control. For instance, bacteriological examinations were made of the experimental packs of fish canned by the Bureau to determine which processes produced sterility, on the keeping quality of fresh mackerel packed in ice, in determining the preservative value of smoke, and on the tests developed for determining the relative freshness of fish. In addition, special bacteriological investigations were conducted for the development of disinfectants for sponges in household use and for determining the effect of antiseptics in ice on the keeping quality of fresh haddock when packed in such ice for temporary preservation.

Preservation of fishery byproducts.—The value of the Bureau's investigations in the field of fishery waste utilization has been very apparent during the past year. Studies with respect to the utilization of salmon cannery waste have demonstrated that valuable vitamin-active oils and a high quality edible oil can be recovered from this type of material by proper methods of preparation. The vitamin oils have been proved as desirable sources of fat soluble vitamins for poultry feeding and also as a new and reasonable source of these vitamins for human nutrition. The edible salmon oil may be added to canned salmon to improve both the nutritive value and appearance of the pack. Definite commercial application has been made of the results of the above study with advantage.

Studies on the manufacture of fish meal have demonstrated the possibility of producing materials of higher quality, and the fundamental information obtained in the course of these investigations has contributed much to a better understanding of the value of fish meals in animal feeding.

By investigation of the oil contained in swordfish livers, the Bureau discovered the richest known natural source of both vitamins A and D, and as a result of this work the fishing industry is obtaining additional revenue from the sale of livers which were formerly discarded. Swordfish-liver oils can be prepared that contain as high as 300,000 U. S. P. vitamin A units and 9,500 U. S. P. vitamin D

units as compared to the minimum requirements for cod-liver oil of 600 vitamin A units and 95 vitamin D units. Besides giving added revenue to the fishing industry, these studies contributed to the domestic supply of vitamin-bearing materials.

Nutritive value of aquatic products.—Studies of the nutritive value of aquatic products during the past year included a determination of the vitamin content of swordfish-liver oils, salmon oils, and other fish and fish-liver oils extracted according to various experimental methods as previously discussed; a determination of vitamins A and G in crab meat, a study of the food value of sodium alginate, analyses of conch meat; and data on the role of the mineral constituents of fishery products in nutrition. In making these studies, our technologists made use of chemical methods and practical feeding tests with laboratory animals as well as biological tests with human subjects. Our investigations reveal that fish oils and fish-liver oils are of great potential sources of vitamins for both human and animal nutrition; that crab meat is a relatively good source of vitamins A and G; that sodium alginate is not only of value as a stabilizer in the preparation of dairy products, but that it has food value comparable to the kelp from which it is made; and that the mineral constituents found in fishery products and byproducts are of great importance in human and animal nutrition.

Fish cookery investigations.—During the past year, the Bureau established a fish cookery laboratory and developed by actual tests simple and practical recipes for the preparation and cooking of fish and shellfish. A very popular publication on practical fish cookery was issued and widely distributed by the Bureau, based on these recipes.

BLACK BASS AND ANGLERS DIVISION

The black-bass law was enforced by 2 regular field officers, 1 or 2 temporary employees, 1 permanent employee in the Washington office, and approximately 100 deputy black-bass-law inspectors who are regularly employed State fish and game protectors, and who serve the Federal Government without pay under the direction of the Chief of the Division. Several violations of the Federal black-bass law were found, but in only one instance was it necessary to report the case for prosecution in Federal court. Active field work has been conducted in well-known black-bass areas east of the Rocky Mountains, and fish markets of the principal cities have been inspected for illegal shipments. The Division receives the hearty cooperation of all the State authorities in those States where it functions.

During the year the legislatures of 44 States met in regular session, affording an opportunity to obtain much needed State legislation protecting black bass, without which the Federal law cannot be fully effective. Three States passed laws prohibiting the sale of black bass at all times, regardless of where taken, making a total of 37 States where these game fish are afforded this means of protection. One State adopted a closed season, leaving but 7 States that do not now protect the bass on the nest during at least a part of the spawning period. The closed season was also extended in several States.

Several States increased the size limit or decreased the daily limit on black bass; 6 States adopted a commission form of administering

the State fish and game laws, and several increased the powers of the commission to promulgate regulations. Part-time nonresident tourist licenses were adopted in 8 States.

The adoption of all these measures was advocated by the Bureau in line with its policies for the protection and increase of the black bass.

Because of its administration of the Federal black-bass law, the Division is daily called upon for information about seasons, angler's licenses, and every subject pertaining to angling. Especially is it requested by letter, telephone, and personal call to advise the applicants, how, when, and where to fish. It has been necessary for the Division to prepare a number of small pamphlets, as follows: List of Books on Angling, List of State Fish Agencies, Compilation of Anglers Licenses Issued by the States and Revenue Derived Therefrom, Tabulation of the Laws Covering the Sale of Black Bass in the Various States, List of States that have Adopted the Part-Time Angler's License; and a leaflet entitled "Angling" which supplies many items of interest to the angler. There has been a large demand for all of these publications, which may be obtained only by applying to the Bureau of Fisheries.

Owing to the many changes in the game-fish laws in 1935, Fishery Circular No. 16 containing the game-fish laws for 1933-34 has been revised, and will be published early in the next fiscal year.

Twenty-seven permits for taking certain species of small fish in the District of Columbia for use as bait, have been issued in the Division during the year, a decrease of 11 over the last fiscal year.

VESSELS

Thirteen vessels of the Alaska service cruised about 116,000 nautical miles in the fiscal year 1935, as compared with 127,000 nautical miles by 15 vessels in the preceding year. The *Penguin* logged the greatest distance, with approximately 27,700 miles; the *Brant* and *Teal* each covered about 13,300 miles; and the *Crane*, about 11,000 miles.

The *Penguin* made five round-trip voyages from Seattle to the Pribilof Islands, transporting personnel and emergency supplies. Supplies also were landed in the Aleutian Islands for the Navy Department and the Coast and Geodetic Survey.

The *Auklet*, *Kittiwake*, and *Teal* were engaged in fishery protective work in southeast Alaska during the 1934 season. The *Murre* operated in the Seward-Katalla district and the *Eider* in the Kodiak area. The *Scoter* patrolled the waters of Cook Inlet and in the fall, after the departure of the *Eider*, was assigned to the Kodiak area. The *Crane* and *Red Wing* were in the Alaska Peninsula region, and the *Blue Wing* was in Bristol Bay. After the close of the Bristol Bay season the *Blue Wing* assisted with the patrol in the Alaska Peninsula area and in southeast Alaska. As formerly, the *Ibis* was stationed at Chignik, and the *Coot* was on the Yukon River.

The *Widgeon* and *Petrel*, which were operated in southeast Alaska in the fiscal year 1934, were out of commission at Seattle.

The *Brant* was used in general supervisory work, chiefly in southeast Alaska, although two trips were made to Bristol Bay in west-

ern Alaska. Fur-seal patrol off Neah Bay, Wash., also was maintained by the *Brant*. A similar patrol in the vicinity of Sitka, Alaska, was carried on by the *Scoter*.

APPROPRIATIONS

Appropriations for the Bureau for the fiscal year aggregated \$1,325,327, as follows:

| | |
|---|--------------------------------|
| Salaries, Commissioner's office..... | \$136, 600 |
| Propagation of food fishes..... | 570, 600 |
| Maintenance of vessels..... | 135, 380 |
| Inquiry respecting food fishes..... | 151, 214 |
| Fishery industries..... | 57, 125 |
| Protecting seal and salmon fisheries of Alaska..... | 245, 693 |
| Upper Mississippi Wildlife and Fish Itefuge..... | 15, 000 |
| Enforcement of black-bass law..... | 13, 715 |
| Total..... | ¹1, 325, 327 |

¹Exclusive of indefinite deficiency appropriation of approximately \$25,000 for restoration of salaries. Also exclusive of an appropriation of \$100,000 for the fiscal year 1936 for oyster-pest control work, of which \$50,000 was made available immediately following the approval of the act on Mar. 22, 1935.

ALASKA FISHERY AND FUR-SEAL INDUSTRIES IN 1934¹

By WARD T. BOWER, Chief, Division of Alaska Fisheries

CONTENTS

| | Page | | Page |
|---|------|--|------|
| INTRODUCTION..... | 2 | FISHERY INDUSTRIES—Continued..... | |
| Visit of the Secretary of Commerce and other officials to Alaska..... | 3 | Herring..... | 44 |
| Protection of walruses and sea lions..... | 3 | Statistical summary..... | 46 |
| Public Works projects..... | 4 | Halibut..... | 47 |
| Civil Works project..... | 4 | Statistical summary..... | 47 |
| FISHERY INDUSTRIES..... | 4 | Cod..... | 48 |
| Alaska fisheries legislation..... | 5 | Whales..... | 49 |
| New fishery regulations..... | 5 | Clams..... | 49 |
| Annette Island Fishery Reserve..... | 11 | Shrimp..... | 50 |
| Stream improvement..... | 12 | Crabs..... | 50 |
| Stream marking..... | 12 | Japanese vessels in Bering Sea..... | 51 |
| Stream guards..... | 13 | Trout..... | 51 |
| Vessel patrol..... | 13 | Miscellaneous fishery products..... | 52 |
| Aerial patrol..... | 14 | FUR-SEAL INDUSTRY..... | 52 |
| Complaints and prosecutions..... | 14 | Pribilof Islands..... | 52 |
| Decision regarding license fee for nonresident fishermen..... | 16 | General administrative work..... | 52 |
| Territorial license tax..... | 16 | Transportation of supplies..... | 52 |
| Water-power projects in Alaska..... | 17 | Power vessel <i>Penguin</i> | 53 |
| Kuskokwim River..... | 17 | Roads..... | 53 |
| Yukon River..... | 18 | Buildings..... | 54 |
| Wells for counting salmon escapement..... | 18 | Natives..... | 54 |
| Klawak Creek..... | 19 | Census..... | 54 |
| Little Port Walter..... | 19 | Medical service..... | 54 |
| Situk River..... | 19 | Schools..... | 54 |
| Alitak Bay..... | 19 | Savings accounts..... | 55 |
| Chignik River..... | 20 | Payments for taking fur-seal skins..... | 55 |
| Chinik Creek..... | 20 | Payments for taking fox skins..... | 56 |
| English Bay..... | 20 | Fur seals..... | 56 |
| Kadla Bay..... | 21 | Killings..... | 56 |
| Karluk River..... | 21 | Age classes..... | 57 |
| Morzhovoi Bay..... | 22 | Reserving operations..... | 58 |
| Orzenoi River..... | 22 | Computation of fur-seal herd..... | 58 |
| Red River..... | 22 | Foxes..... | 59 |
| Salmon life-history studies..... | 22 | Trapping season of 1934-35..... | 59 |
| Observations on the escapement of salmon..... | 23 | Reindeer..... | 59 |
| Hatcheries..... | 24 | Fur-seal skins..... | 59 |
| General statistics of the fisheries..... | 25 | Shipments..... | 59 |
| Salmon..... | 26 | Sales..... | 60 |
| Catch and apparatus..... | 29 | Disposition of fur-seal skins taken at Pribilof Islands..... | 64 |
| Canning..... | 30 | Shipment and sale of fox skins..... | 65 |
| Changes in canneries..... | 30 | Sea-otter skin..... | 66 |
| New canneries..... | 31 | Fur-seal patrol..... | 66 |
| Canneries not operated..... | 31 | United States Coast Guard..... | 66 |
| Total canneries operated..... | 33 | Bureau of Fisheries..... | 66 |
| Losses and disasters..... | 35 | Sealing privileges accorded aborigines..... | 67 |
| Statistics..... | 35 | Japanese sealskins delivered to the United States..... | 67 |
| Pack in certain districts..... | 39 | COMPUTATION OF FUR SEALS, PRIBILOF ISLANDS, 1934..... | 67 |
| Mild curing..... | 40 | Bulls..... | 69 |
| Pickling..... | 42 | Average harem..... | 69 |
| Fresh salmon..... | 42 | Pups and cubs..... | 70 |
| Freezing..... | 43 | Mortality of seals at sea..... | 71 |
| Dry-salted, dried, and other miscellaneous salmon products..... | 43 | Complete computation..... | 72 |
| Byproducts..... | 44 | | |

¹ Appendix I to the Report of the U. S. Commissioner of Fisheries for 1935. Approved for publication June 14, 1935.

INTRODUCTION

The Bureau's work in connection with the conservation of the fisheries of Alaska and the protection and management of the Pribilof Islands fur-seal herd was continued along the usual lines. An inspection of the fishery and fur-seal activities was again made by the Commissioner of Fisheries. The Secretary of Commerce also observed these activities in his tour of the Territory to survey the Department's work.

As the runs of salmon in general were satisfactory, comparatively few additional restrictions on commercial fishing were necessary during the season. Among the modifications of the regulations were those permitting limited commercial fishing for king salmon in the Yukon and Kuskokwim Rivers by residents of the locality, in accordance with provisions of an act of Congress of April 16, 1934.

Although some branches of the industry were adversely affected by the water-front strike on the Pacific coast and by fishermen's strikes in southeast Alaska and the Copper River area, the output of fishery products for the season showed a considerable gain, both in quantity and value, over that for 1933. A new record was set in the production of canned salmon, with a total pack of 7,481,830 cases, or 828,948 cases more than the previous record pack of 1926.

A patrol was maintained during the most active part of the fishing season, 164 temporary employees being identified with this work for varying periods, in addition to the regular employees and crews of the Bureau's patrol vessels. Effective use was made of chartered airplanes as a supplementary patrol, chiefly during the weekly closed periods, and for supervisory work and the inspection of spawning areas.

Life-history studies of the salmon and herring were continued. Weirs for counting the escapement of salmon to the spawning grounds were operated in 12 typical salmon streams. At several of these weirs attention was given to the destruction of trout that feed upon young salmon, and in the Bristol Bay region bounties were paid on predatory trout through funds provided by the Territory and by local packers.

Allotments by the Public Works Administration for reestablishing weirs, overhauling and repairing Alaska vessels, and reconditioning and repairing buildings at the Pribilof Islands were of material assistance in furthering the Bureau's work. A stream-clearing project to improve salmon-spawning conditions in southeast Alaska was made possible by the Civil Works Administration and gave temporary employment to approximately 200 persons during the winter.

Sealing activities at the Pribilof Islands resulted in the take of 53,468 fur-seal skins. Killings were confined as far as possible to 3-year-old male seals, and sufficient animals of this age class were reserved for future breeding stock. A computation of the fur-seal herd as of August 10, 1934, showed 1,430,418 animals of all classes, or an increase of 8.48 percent over the previous year.

Attention was given to the feeding and care of blue-fox herds on St. Paul and St. George Islands during the winter, and 983 blue and 19 white-fox skins were taken on the two islands in the 1934-35 season.

Some permanent improvements were made at the islands in the way of new buildings and extensions of surfaced roads to expedite the fur-seal work.

Valuable assistance was rendered by the Navy Department in detailing the U. S. S. *Sirius* to transport the annual shipment of supplies to the Pribilof Islands and to bring out the season's take of seal-skins. The United States Coast Guard also rendered important service in maintaining a patrol of the fur seals along the Pacific coast during the northward migration and in Bering Sea throughout the breeding season.

Acknowledgment is made of the assistance rendered by members of the Bureau's staff in the preparation of this document.

VISIT OF THE SECRETARY OF COMMERCE AND OTHER OFFICIALS TO ALASKA

Secretary of Commerce Daniel C. Roper made a detailed investigation of Department of Commerce activities in Alaska in 1934. Departure from Seattle was made on board the Coast Guard cutter *Shoshone* on June 25. All the principal points in southeast, central, and western Alaska were visited, and a brief stop was made at the Pribilof Islands. The return voyage from Dutch Harbor to Seattle was made on the Coast Guard cutter *Tahoe*.

Commissioner Bell was in Alaska during much of the active salmon-fishing season in the summer of 1934, giving personal attention to the conservation needs of the industry. Visits were made to all the important fishing districts throughout the Territory, and conferences with fishermen and operators of canneries and other fishery establishments were held at various places. A trip was also made to the Pribilof Islands to observe the activities there.

Much of this inspection trip was made aboard the Bureau's vessel *Brant*, although some travel occurred on the *Penguin*, and on the Coast Guard cutter *Shoshone*. A number of extended airplane flights were made to reach some of the more inaccessible places, thus saving considerable time and making possible a more extensive inspection of the fisheries.

Commissioner Bell returned to Washington on August 18.

Second Assistant Postmaster General Harilee Branch was furnished transportation to southeast Alaska on the *Penguin*, sailing from Seattle on August 25. Julian W. Blount, secretary to Senator Peter Norbeck, likewise was a passenger on the vessel at that time, making the round trip from Seattle to the Pribilof Islands.

Through an oversight the visit of United States Senator Arthur R. Robinson, of Indiana, to the Pribilof Islands in 1933 was omitted from the corresponding report for that year. Senator Robinson was at St. Paul Island on July 8, 1933, observing sealing operations. Transportation to the Pribilofs was afforded by the U. S. Coast Guard cutter *Chelan*.

PROTECTION OF WALRUSES AND SEA LIONS

A new (seventh) edition of Department of Commerce Circular No. 286 was issued under date of July 1, 1934, containing the laws and regulations for the protection of walruses and sea lions in Alaska. The regulation governing the killing of walruses is as follows:

The killing of walruses in the Territory of Alaska or in any of the waters of Alaska over which the United States has jurisdiction is prohibited from July 1, 1934, to June 30, 1936, both dates inclusive. This prohibition shall not apply

to the killing of walruses by natives for food or clothing, by miners or explorers when in need of food, or to the collection of specimens for scientific purposes under permits issued by the Secretary of Commerce.

On June 16, 1934, an act was passed pertaining to the killing of sea lions in Alaska waters, the text of which is as follows:

Be it enacted by the Senate and House of Representatives of the United States of America in Congress assembled, That all acts and parts of acts making it unlawful to kill sea lions, as game animals or otherwise, in the waters of the Territory of Alaska are repealed: Provided, however, That sea lions shall not be killed in the waters of Alaska except under such rules and regulations as the Secretary of Commerce may prescribe, in order to prevent the extinction of sea lions as a species of interesting sea life in the waters of Alaska.

The following departmental regulations governing the killing of sea lions in the waters of Alaska were made effective on July 1, 1934:

The killing of sea lions in the Territory of Alaska, or in any of the waters of Alaska over which the United States has jurisdiction, is permitted as follows:

1. (a) By natives for food or clothing, and by miners or explorers when in need of food.

(b) By anyone in the necessary protection of property, or while such animals are destroying salmon or other food fish.

2. The killing of sea lions as specimens for scientific purposes will be under permits issued by the Secretary of Commerce.

PUBLIC WORKS PROJECTS

Allotments to the Bureau by the Public Works Administration in the fiscal year 1934 included \$29,000 for projects having to do with the Alaska service. Of this amount, \$20,000 was for reconditioning and repairing 12 Alaska vessels, \$3,000 for reconditioning and repairing buildings at the Pribilof Islands, and \$6,000 for reestablishing weirs to tally escapements of spawning salmon. These funds helped materially to further the Bureau's program for the conservation of the fisheries and the management of the fur-seal industry. A part of the allotment for weirs was carried over for reestablishing the structures in 1935.

CIVIL WORKS PROJECT

Under date of December 6, 1933, the Department was notified of the approval of the Civil Works project for the improvement of salmon spawning streams in southeast Alaska. The Territorial Civil Works Administrator was authorized to furnish 198 unskilled laborers for this work. Operations were under the supervision of three regular employees of the Bureau of Fisheries in the Alaska field service. A report of the work accomplished is included under the section regarding stream improvement.

FISHERY INDUSTRIES

As in corresponding reports for previous years, the Territory of Alaska is here considered in the three coastal geographic sections generally recognized, as follows: (1) Southeast Alaska—embracing all that narrow strip of mainland and the numerous adjacent islands from Portland Canal northwestward to and including Yakutat Bay; (2) central Alaska—the region on the Pacific from Yakutat Bay westward, including Prince William Sound, Cook Inlet, and the southern coast of Alaska Peninsula, to Unimak Pass; and (3) western Alaska—the north shore of the Alaska Peninsula, including the Aleutian Islands westward from Unimak Pass, Bristol Bay, and

the Kuskokwim and Yukon Rivers. These divisions are solely for statistical purposes and do not coincide with areas established in departmental regulations.

Detailed reports and statistical tables dealing with the various fishery industries are presented herewith, and there are also given the important features of certain subjects of special investigation or inquiry.

ALASKA FISHERIES LEGISLATION

On April 16, 1934, the President approved an act amending the Fisheries Act of June 26, 1906, as amended by the act of June 6, 1924. This modification permits commercial fishing for king salmon in the Yukon and Kuskokwim Rivers by native Indians and bona fide permanent white inhabitants under such restrictions as may be specified by the Secretary of Commerce. Prior to the enactment of this measure all commercial fishing was prohibited in the Yukon and Kuskokwim Rivers and within 500 yards of their mouths. The text of the amendment is as follows:

Be it enacted by the Senate and House of Representatives of the United States of America in Congress assembled, That section 3 of the act of Congress entitled "An act for the protection and regulation of the fisheries of Alaska", approved June 26, 1906, as amended by the act of Congress entitled "An act for the protection of the fisheries of Alaska, and for other purposes", approved June 6, 1924, be, and the same is hereby, amended to read as follows:

"**SEC. 3.** That it shall be unlawful to erect or maintain any dam, barricade, fence, trap, fishwheel, or other fixed or stationary obstruction except for purposes of fish culture, in any of the waters of Alaska at any point where the distance from shore to shore is less than one thousand feet, or within five hundred yards of the mouth of any creek, stream, or river into which salmon run, excepting the Karluk, Ugashik, Kuskokwim, and Yukon Rivers, with the purpose or result of capturing salmon or preventing or impeding their ascent to the spawning grounds, and the Secretary of Commerce is hereby authorized and directed to have any and all such unlawful obstructions removed or destroyed: *Provided, however,* That the exception hereinabove contained with reference to the Kuskokwim and Yukon Rivers shall be solely for the purpose of enabling native Indians and bona fide permanent white inhabitants along the said rivers to take from said rivers for commercial purposes and for export from the Territory of Alaska king salmon in such manner and such quantities, and at such times as the Secretary of Commerce may, by suitable regulations, from time to time permit: *Provided further,* That no person shall be deemed to be a bona fide permanent inhabitant of the said rivers who has not resided thereon, or within fifty miles thereof for a period of over one year, and that the term 'native Indians' as used herein shall be taken to mean members of the aboriginal races inhabiting Alaska when annexed to the United States, and their descendants of the whole or half blood. For the purposes of this section, the mouth of such creek, stream, or river shall be taken to be the point determined as such mouth by the Secretary of Commerce and marked in accordance with this determination. It shall be unlawful to lay or set any seine or net of any kind within one hundred yards of any other seine, net, or other fishing appliance which is being or which has been laid or set in any of the waters of Alaska, or to drive or to construct any trap or any other fixed fishing appliance within six hundred yards laterally or within one hundred yards endwise of any other trap or fixed fishing appliance."

SEC. 2. That section 4 of the act of Congress entitled "An Act for the protection and regulation of the fisheries of Alaska", approved June 26, 1906, as amended by the act of Congress entitled "An Act for the protection of the fisheries of Alaska, and for other purposes", approved June 6, 1924, be, and the same hereby is, amended to read as follows:

"**SEC. 4.** That it shall be unlawful to fish for, take, or kill any salmon of any species or by any means except by hand rod, spear, or gaff in any of the creeks, streams, or rivers of Alaska; or within five hundred yards of the mouth of any such creek, stream, or river over which the United States has jurisdiction, excepting the Karluk, Ugashik, Yukon, and Kuskokwim Rivers: *Provided,* That nothing herein contained shall prevent the taking of fish for local food requirements or for use as dog feed: *Provided further,* That the exception hereinabove contained

with reference to the Kuskokwim and Yukon Rivers shall be solely for the purpose of enabling native Indians and bona fide permanent white inhabitants along the said rivers to take from said rivers for commercial purposes and for export from the Territory of Alaska king salmon in such manner and such quantities, and at such times as the Secretary of Commerce may, by suitable regulations, from time to time permit: *Provided further*, That no person shall be deemed to be a bona fide permanent inhabitant of said rivers who has not resided thereon or within fifty miles thereof for a period of over one year, and that the term 'native Indians' as used herein shall be taken to mean members of the aboriginal races inhabiting Alaska when annexed to the United States, and their descendants of the whole or half blood."

Approved April 16, 1934.

NEW FISHERY REGULATIONS

The regulations for the protection of the fisheries of Alaska, issued December 21, 1933, were amended by the following regulations issued by the Acting Secretary of Commerce under the dates indicated:

[February 1, 1934]

ALASKA PENINSULA AREA

Salmon fishery.—1. Regulation No. 21 (*n*) is amended to read as follows: Unga Island: North and east coasts (1) within 2,500 feet of a point at 55 degrees 24 minutes 15 seconds north latitude, 160 degrees 41 minutes 24 seconds west longitude, (2) within 2,500 feet of a point at 55 degrees 11 minutes 42 seconds north latitude, 160 degrees 27 minutes 38 seconds west longitude, and (3) within 2,500 feet of a point at 55 degrees 13 minutes 29 seconds north latitude, 160 degrees 29 minutes 37 seconds west longitude.

2. Regulation No. 21 (*p*) is amended to read as follows: (1) Within 5,000 feet of a point on a reef near Guillemot Island at 55 degrees 33 minutes 10 seconds north latitude, 160 degrees 25 minutes 17 seconds west longitude, (2) within 1,000 feet of a point on Korovin Island at 55 degrees 24 minutes 24 seconds north latitude, 160 degrees 8 minutes 57 seconds west longitude, and (3) within 2,500 feet of a point on Nagai Island at 55 degrees 15 minutes 47 seconds north latitude, 159 degrees 54 minutes 46 seconds west longitude.

KODIAK AREA

Salmon fishery.—1. Regulation No. 14 (*h*) is amended to read as follows: Kodiak Island: (1) Within 2,500 feet of a point on the west side of Uganik Bay at 57 degrees 50 minutes 45 seconds north latitude, 153 degrees 38 minutes west longitude, and (2) within 5,000 feet easterly of a point at 57 degrees 52 minutes 5 seconds north latitude, 153 degrees 51 minutes 9 seconds west longitude.

2. Regulation No. 14 (*k*) is amended to read as follows: Kodiak Island: (1) Within 2,500 feet of a point on the east side of Uyak Bay at 57 degrees 35 minutes 40 seconds north latitude, 153 degrees 49 minutes 30 seconds west longitude, and (2) within 2,500 feet of a point northwest of Cape Uyak at 57 degrees 38 minutes 6 seconds north latitude, 154 degrees 20 minutes 3 seconds west longitude.

PRINCE WILLIAM SOUND AREA

Salmon fishery.—1. Regulation No. 12 (*a*) is amended to read as follows: Knight Island: From a point on the southeast coast at 60 degrees 9 minutes 50 seconds north latitude southerly to Point Helen, thence northerly to 60 degrees 11 minutes 15 seconds north latitude, 147 degrees 50 minutes 40 seconds west longitude, exclusive of all waters in Little Bay.

2. Regulation No. 12 (*e*) is amended to read as follows: Eastern coast of Culross Island: From the northern side of the entrance to Hidden Bay northerly to a point at 60 degrees 43 minutes 45 seconds north latitude, 148 degrees 6 minutes 35 seconds west longitude.

3. Regulation No. 12 (*g*) is amended to read as follows: Along the mainland coast within 1 statute mile easterly from the outermost extremity of Point Pellew.

4. Regulation No. 12 (*cc*) is amended to read as follows: Montague Island: Northern coast (1) from Graveyard Point to a point at 60 degrees 21 minutes 41 seconds north latitude, 147 degrees 9 minutes 47 seconds west longitude, and (2) within 2,500 feet of a point on the southeast side of Zaikof Bay at 60 degrees 17 minutes 40 seconds north latitude, 146 degrees 59 minutes west longitude.

SOUTHEASTERN ALASKA AREA

ICY STRAIT DISTRICT

Salmon fishery.—1. Regulation No. 14 (b) is amended to read as follows: Along the coast (1) within 2,500 feet of a point at 58 degrees 16 minutes 18 seconds north latitude, 136 degrees 20 minutes 27 seconds west longitude, (2) from a point on the east coast of the northwestern island at 58 degrees 15 minutes 50 seconds north latitude, northwesterly to the northernmost point of the island, thence southerly to a point at 58 degrees 15 minutes 26 seconds north latitude, and (3) within 1,000 feet of a point at 58 degrees 14 minutes 9 seconds north latitude, 136 degrees 20 minutes 6 seconds west longitude.

2. Regulation No. 14 (l) is amended to read as follows: Chichagof Island: Northeastern coast (1) within 2,500 feet of a point at 58 degrees 8 minutes 40 seconds north latitude, 135 degrees 23 minutes 59 seconds west longitude, (2) from 135 degrees 20 minutes west longitude to 135 degrees 11 minutes 11 seconds west longitude, and (3) within 2,500 feet of a point at 58 degrees 5 minutes 20 seconds north latitude, 135 degrees 9 minutes 29 seconds west longitude.

WESTERN DISTRICT

Salmon fishery.—1. Regulation No. 20 (e) is amended to read as follows: Chichagof Island: East coast (1) from 57 degrees 48 minutes 45 seconds north latitude to 57 degrees 48 minutes 18 seconds north latitude, (2) from 57 degrees 45 minutes north latitude to 57 degrees 42 minutes 40 seconds north latitude, (3) from 57 degrees 36 minutes 30 seconds north latitude to 57 degrees 34 minutes north latitude, and (4) from 57 degrees 29 minutes 30 seconds north latitude to Point Hayes.

2. Regulation No. 20 (k) is amended to read as follows: Mansfield Peninsula: West coast (1) from the south point of entrance to Funter Bay to 58 degrees 10 minutes 45 seconds north latitude, and (2) from 58 degrees 9 minutes 30 seconds north latitude to the southern extremity of the peninsula at the north side of the entrance to Hawk Inlet.

3. Regulation No. 20 (l) is amended to read as follows: Admiralty Island: West coast (1) from a point $\frac{3}{4}$ statute mile north of Parker Point to 57 degrees 45 minutes north latitude, (2) from 57 degrees 49 minutes 55 seconds north latitude to 57 degrees 51 minutes north latitude, (3) from 57 degrees 54 minutes 45 seconds north latitude to 57 degrees 56 minutes 20 seconds north latitude, and (4) from 57 degrees 57 minutes 50 seconds north latitude to 58 degrees 2 minutes north latitude.

NORTH PRINCE OF WALES ISLAND DISTRICT

Salmon fishery.—1. Regulation No. 16 (n) is amended to read as follows: (1) Kuiu Island: East coast of peninsula between Port Beauclerc and Reid Bay from 56 degrees 18 minutes 30 seconds north latitude northward to 56 degrees 19 minutes north latitude, and (2) unnamed island east of Kuiu Island: Within 2,500 feet measured along the east coast from a point at 56 degrees 28 minutes 30 seconds north latitude, 133 degrees 50 minutes 50 seconds west longitude.

2. Regulation No. 16 (s) is amended to read as follows: (1) Kupreanof Island: Southern coast within 2,500 feet of a point at 56 degrees 26 minutes 8 seconds north latitude, 133 degrees 29 minutes 35 seconds west longitude, and (2) Zarembo Island: Southwest coast within 2,500 feet of a point at 56 degrees 15 minutes north latitude, 132 degrees 53 minutes 30 seconds west longitude.

3. Regulation No. 16 (cc) is amended to read as follows: Prince of Wales Island: East coast from a point at 55 degrees 48 minutes 35 seconds north latitude southerly to a point at 55 degrees 46 minutes 45 seconds north latitude.

4. Regulation No. 16 (gg) is amended to read as follows: Cleveland Peninsula: (1) South side of Ernest Sound within 1,500 feet of a point on the northwestern extremity of Union Point at 55 degrees 48 minutes 10 seconds north latitude, 132 degrees 11 minutes west longitude, and (2) east side of Clarence Strait within 2,000 feet southerly of a point at 55 degrees 45 minutes 48 seconds north latitude, 132 degrees 17 minutes 5 seconds west longitude.

5. Regulation No. 16 (jj) is amended to read as follows: Gravina Island: West coast (1) from South Vallenar Point to 55 degrees 20 minutes 48 seconds north latitude, and (2) from 55 degrees 18 minutes 50 seconds north latitude to 55 degrees 8 minutes 15 seconds north latitude, including the Bronaugh Islands and rocky islets adjacent to this coast.

SOUTH PRINCE OF WALES ISLAND DISTRICT

Salmon fishery.—1. Regulation No. 13 (i) is amended to read as follows: Prince of Wales Island: West coast (1) within 2,500 feet of a point at 55 degrees 2 minutes 20 seconds north latitude, 132 degrees 35 minutes 18 seconds west longitude, and (2) from Point Webster southeasterly to 54 degrees 55 minutes 52 seconds north latitude, 132 degrees 32 minutes 30 seconds west longitude.

2. Regulation No. 13 (j) is amended to read as follows: Along the coast (1) within 900 feet west of the southern extremity of the eastern large island of the Ship Islands group at 54 degrees 53 minutes 42 seconds north latitude, 132 degrees 30 minutes 21 seconds west longitude, and (2) within 2,500 feet of a point on a small island near the entrance to Ruth Bay at 54 degrees 53 minutes 40 seconds north latitude, 132 degrees 26 minutes 27 seconds west longitude.

ICY STRAIT, WESTERN, EASTERN, NORTH PRINCE OF WALES ISLAND, SOUTH PRINCE OF WALES ISLAND, AND SOUTHERN DISTRICTS

Salmon fishery.—Regulation No. 4 is amended to read as follows: No floating trap shall exceed 900 feet in length when any part of such trap is in a greater depth of water than 100 feet at mean high tide. The length of any such trap shall be as measured along the lead from shore at mean high tide to the outer face of the pot.

[March 17, 1934]

SOUTHEASTERN ALASKA AREA

Shrimp fishery.—Commercial fishing for shrimps is prohibited in the period from March 20 to April 30, both dates inclusive, in each year.

[June 4, 1934]

YUKON-KUSKOKWIM AREA

Salmon fishery.—1. Regulation No. 1 (a) is amended to read as follows: Port Clarence district: Waters within a line extending from Cape Douglas to Cape York.

2. Regulation No. 1 (b) is amended to read as follows: Yukon district: All waters of the Yukon River and all waters along the coast between 62 degrees north latitude and 63 degrees 15 minutes north latitude.

3. Regulation No. 1 (c) is amended to read as follows: Kuskokwim district: All waters of Kuskokwim River, and waters of Kuskokwim Bay, exclusive of Goodnews Bay, within a line extending from the east shore of Kuskokwim Bay at the parallel of 59 degrees north latitude to Cape Avinof. The mouth of the Kuskokwim River shall be considered as at a straight line extending from a marker erected for the purpose at Beacon Point to another marker at Popokamute.

4. Regulation No. 2 is amended to read as follows: Commercial fishing for salmon shall be conducted solely by drift gill nets and set nets: *Provided*, That this shall not apply (a) to the use of purse seines in Kuskokwim Bay, exclusive of Goodnews Bay, between 59 degrees north latitude and 59 degrees 40 minutes north latitude westward to Cape Avinof, and (b) to the use of fish wheels in the Yukon and Kuskokwim Rivers by native Indians and bona fide permanent white residents along those rivers for the capture of king salmon.

5. Regulation No. 5 is amended to read as follows: Commercial fishing for salmon in the Yukon district shall be conducted solely by king-salmon gill nets having mesh of at least 8½ inches stretched measure between knots as measured when actually in use: *Provided*, That this shall not apply to the use of fish wheels in the Yukon River by native Indians and bona fide permanent white residents along the river for the capture of king salmon.

6. Regulation No. 6 is amended to read as follows: Commercial fishing for salmon is prohibited except in the period from 6 o'clock antemeridian June 1 to 6 o'clock postmeridian July 31, in each year.

7. Regulation No. 7 is amended to read as follows: The combined take of king and red salmon for commercial purposes shall not exceed 350,000 fish in any calendar year: *Provided*, That the take of king salmon in the Yukon district shall not exceed 100,000 fish, of which not more than 50,000 may be taken outside of the mouth of the Yukon River, in any calendar year: *And provided further*, That the combined take of red and king salmon in the Kuskokwim district shall not exceed 250,000 fish in any calendar year.

ALASKA PENINSULA AREA

Salmon fishery.—1. Regulation No. 2 is amended to read as follows: In the waters along the south side of the Alaska Peninsula, including the waters of the Shumagin and other adjacent islands, the 36-hour closed period for salmon fishing prescribed by section 5 of the act of June 6, 1924, is hereby extended to include the periods from 6 o'clock postmeridian Tuesday to 6 o'clock antemeridian Wednesday and from 6 o'clock postmeridian Wednesday to 6 o'clock antemeridian Thursday of each week making a weekly closed period of 60 hours: *Provided*, That this extension of 24 hours' closed period each week shall not be effective after 6 o'clock antemeridian of July 25 of each year.

2. Regulation No. 14 prohibiting commercial fishing for salmon prior to July 15 along the coast from the west side of the entrance to Sankin Bay to Morgan Point is revoked.

3. Regulation No. 15 prohibiting commercial fishing for salmon prior to July 1 along the coast from a point 1 statute mile northwest of Moss Cape to Castle Cape is revoked.

4. Regulation No. 16 prohibiting commercial fishing for salmon, except by beach seines, in the waters between Cape Tachini and Bold Cape and between Cape Tolstoi and Kupreanof Point is revoked.

ALEUTIAN ISLANDS AREA

Herring fishery.—1. Commercial fishing for herring, except for bait purposes, by means of any seine is prohibited except in the period from July 15 to October 31, both dates inclusive.

2. Regulation No. 5 is amended to read as follows: Commercial fishing for herring, except for bait purposes, by means of any seine is prohibited west of 166 degrees west longitude.

3. Regulation No. 6 is amended to read as follows: Commercial fishing for herring, including bait fishing, by means of any purse seine more than 1,400 meshes in depth, more than 180 fathoms in length, or of mesh less than 1½ inches stretched measure between knots is prohibited.

KODIAK AREA

Salmon fishery.—1. Regulation No. 11 is amended to read as follows: Commercial fishing for salmon between Cape Karluk and Cape Uyak except by beach seines and purse seines is prohibited.

2. Regulation No. 12 is amended to read as follows: Commercial fishing for salmon is prohibited from August 15 to August 31, both dates inclusive: *Provided*, That this prohibition shall not apply (1) to beach seines and purse seines on the north coast of Kodiak Island from Cape Karluk to Cape Uyak, (2) to set gill nets on the north coast of Kodiak Island from Cape Uyak to West Point, and (3) to traps on the west coast of Kodiak Island from Cape Uyak to Uyak Post Office, and from Chief Point to West Point.

3. Regulation No. 15 (p) is amended to read as follows: Kafia Bay, on north shore of Shelikof Strait: All waters within 1 statute mile outside the entrance of the outer lagoon.

COOK INLET AREA

Salmon fishery.—1. Regulation No. 1 is amended so as to permit commercial fishing for salmon south of the latitude of Anchor Point, except in Chinik Inlet and Kamishak Bay south of 59 degrees 9 minutes north latitude, until 6 o'clock postmeridian August 10.

2. Regulation No. 9 is amended to read as follows: No set or anchored gill net shall exceed 35 fathoms in length measured on the cork line.

3. Regulation No. 10 is amended to read as follows: Stake and set or anchored gill nets shall be operated in substantially a straight line: *Provided*, That not to exceed 20 yards of each net may be used as a hook. Only one such hook is permitted on a net.

4. Regulation No. 15 (q) is amended to read as follows: Along the mainland coast on the east side of Cook Inlet within 1,200 feet of a point on the west side of Nubble Point Spit at 59 degrees 28 minutes 45 seconds north latitude, 151 degrees 35 minutes 6 seconds west longitude.

RESURRECTION BAY AREA

Salmon fishery.—1. Regulation No. 6 is amended to read as follows: Prior to 6 o'clock antemeridian June 4 in each year commercial fishing for salmon with nets of mesh less than 8½ inches stretched measure between knots is prohibited.

2. Regulation No. 8 is amended to read as follows: In the waters of Resurrection Bay within a line from Cape Resurrection to the western side of Bear Glacier at its mouth, the 36-hour closed period for salmon fishing prescribed by section 5 of the Act of June 6, 1924, is hereby extended to include the period from 6 o'clock antemeridian of Saturday of each week until 6 o'clock antemeridian of the Monday following, making a weekly closed period of 48 hours: *Provided*, That this extension shall not be effective after August 23 in each year.

SOUTHEASTERN ALASKA AREA

ICY STRAIT DISTRICT

Salmon fishery.—1. Regulation No. 6 is amended so as to permit commercial fishing for salmon, except by means of traps, in the period from 6 o'clock antemeridian October 1 to 6 o'clock postmeridian October 15.

2. Regulation No. 16 is amended to read as follows: All commercial fishing for salmon is prohibited in Glacier Bay north of 58 degrees 27 minutes 54 seconds north latitude.

NORTH PRINCE OF WALES ISLAND DISTRICT

Salmon fishery.—1. Regulation No. 6 is amended to read as follows:

(a) Commercial fishing for salmon, other than trolling, is prohibited in all waters of the district off the west and north coasts of Prince of Wales Island, and in the waters north and east of a line extending from Union Point to the southern extremity of Ernest Point, thence northerly to a point on Etolin Island at 55 degrees 54 minutes 45 seconds north latitude, 132 degrees 21 minutes west longitude, thence northerly and westerly along the watershed of Etolin Island to a point northwest of the head of Mosman Inlet at 56 degrees 9 minutes 45 seconds north latitude, 132 degrees 37 minutes 15 seconds west longitude, thence southerly to a point at 56 degrees 6 minutes north latitude, 132 degrees 37 minutes 15 seconds west longitude, thence in a northerly direction along the watershed to the northern extremity of Point Harrington, thence in a westerly direction to the northern extremity of East Island, thence to the southern extremity of West Island, and thence to a point on the east coast of Prince of Wales Island at 56 degrees 9 minutes 15 seconds north latitude, 133 degrees 2 minutes 45 seconds west longitude, prior to 6 o'clock antemeridian July 10, from 6 o'clock postmeridian August 22 to 6 o'clock antemeridian October 1, and for the remainder of each calendar year after 6 o'clock postmeridian October 15.

(b) Commercial fishing for salmon, other than trolling, is prohibited in all waters between a line extending from the northern extremity of Point Harrington to the northern extremity of East Island, thence to the southern extremity of West Island, and thence to a point on the east coast of Prince of Wales Island at 56 degrees 9 minutes 15 seconds north latitude, 133 degrees 2 minutes 45 seconds west longitude, and a line extending from Narrow Point to Ernest Point prior to 6 o'clock antemeridian July 15, from 6 o'clock postmeridian August 28 to 6 o'clock antemeridian October 1, and for the remainder of each calendar year after 6 o'clock postmeridian October 15.

(c) Commercial fishing for salmon, other than trolling, is prohibited in all waters between a line extending from Narrow Point to Ernest Point, thence to Union Point and a line extending from Approach Point to Caamano Point prior to 6 o'clock antemeridian July 15, from 6 o'clock postmeridian August 25 to 6 o'clock antemeridian October 1, and for the remainder of each calendar year after October 15.

(d) Commercial fishing for salmon, other than trolling, is prohibited in all waters south of a line extending from Approach Point to Caamano Point, including all waters of Kasaan Bay, prior to 6 o'clock antemeridian July 10, from 6 o'clock postmeridian August 22 to 6 o'clock antemeridian October 1, and for the remainder of each calendar year after 6 o'clock postmeridian October 15.

2. Regulation No. 7 is amended to read as follows: Commercial fishing for salmon by means of any trap is prohibited in the period from 6 o'clock antemeridian October 1 to 6 o'clock postmeridian October 15.

3. Regulation No. 16 (*aa*) is amended to read as follows: Prince of Wales Island: East coast from a point $\frac{1}{2}$ statute mile northwest of Luck Point to 55 degrees 55 minutes 20 seconds north latitude, exclusive of 1 statute mile each side of the mouth of Eagle Creek.

4. Regulation No. 16 (*cc*) as modified by supplement No. 251-20-1 is amended to read as follows: Prince of Wales Island: East coast from a point at 55 degrees 47 minutes 35 seconds north latitude southerly to a point at 55 degrees 46 minutes 45 seconds north latitude.

5. Regulation No. 16 (*pp*) is amended so as to prohibit the operation of any trap along the east coast of Prince of Wales Island from 55 degrees 36 minutes 5 seconds north latitude to 55 degrees 37 minutes 5 seconds north latitude.

SOUTHEASTERN ALASKA AREA

Herring fishery.—Regulation No. 3 is amended to read as follows: All commercial fishing for herring, including bait fishing, is prohibited, as follows: All waters of Chatham Strait and contiguous waters along the western shore of Admiralty Island between Point Gardner and Marble Bluffs: *Provided*, That this prohibition shall not apply (a) to commercial fishing for herring, including bait fishing, by means of gill nets of mesh not less than $2\frac{1}{2}$ inches stretched measure between knots in the period from June 1 to December 31, both dates inclusive, and (b) to the taking of herring for bait by salmon trolling boats by means of any gill net of mesh not more than $2\frac{1}{2}$ inches stretched measure between knots, of not greater than No. 20 gill net thread, and not exceeding 10 fathoms in length and 100 meshes in depth.

GENERAL REGULATIONS

Regulation No. 6 is amended so as to permit commercial fishing for flounders by means of any trawl when such fishing does not result in the capture, injury, or destruction of other food fish.

[July 2, 1934]

BRISTOL BAY AREA

Salmon fishery.—1. In addition to existing prohibitions, commercial fishing for salmon in the Nushagak district, which embraces the waters of Nushagak Bay within a line from Point Protection to Etolin Point, is prohibited on Wednesday of each week from 4 o'clock antemeridian to 6 o'clock antemeridian, in the period prior to 6 o'clock antemeridian August 3.

2. In addition to existing prohibitions, commercial fishing for salmon in the Ugashik district, which includes the coastal waters from a point 3 statute miles north of Cape Greig to a point on the coast 3 statute miles south of Cape Meshikof, is prohibited from 10 o'clock antemeridian of Tuesday of each week to 6 o'clock antemeridian of Wednesday of each week, in the period prior to 6 o'clock antemeridian August 3.

[July 7, 1934]

COPPER RIVER AREA

Salmon fishery.—Regulation No. 1 is amended so as to permit commercial fishing for salmon from 6 o'clock antemeridian July 10 to 6 o'clock postmeridian July 28.

Revised regulations effective in 1935 for the protection of the fisheries of Alaska were issued by the Secretary of Commerce under date of January 19, 1935, copies of which may be secured, without cost, on application to the Bureau of Fisheries, Washington, D. C.

ANNETTE ISLAND FISHERY RESERVE

The Annette Island Canning Co. again operated in the Annette Island Fishery Reserve under its lease from the Department of the Interior.

In 1934 the company operated 8 traps within the reservation, the catch of which totaled 1,653,347 salmon, and 1,738 salmon taken in

seines and gill nets were purchased from the natives. In addition, 640,114 salmon taken in purse seines outside the reserve were purchased. Except for 47,308 salmon that were sold to other companies, the entire catch was packed at the cannery. In the operation of the plant and the fish traps employment was given to 20 whites, 182 natives, and 2 Chinese.

Profits to the Metlakatlan Indians of the reserve on the cannery operations for the year 1933 under the provisions of the lease amounted to approximately \$37,000. Preliminary estimates for the year 1934 place the figure at about \$50,000.

STREAM IMPROVEMENT

The reclaiming of former spawning areas that have become inaccessible to the salmon by reason of an accumulation of debris from slides and windfalls is of recognized importance in the conservation of the fisheries. Some work of this kind is accomplished from time to time by Bureau employees in connection with patrol duties, but it is necessarily limited by the amount of funds available.

In 1934 an extensive program of stream improvement in southeastern Alaska was undertaken as a project of the Civil Works Administration. The work was under the direction of three regular employees of the Bureau and was participated in by approximately 200 persons for varying periods. It was carried on throughout the first 4 months of the year, in which time 468 streams were cleared for a distance of 621 miles, and more than 100 miles of trail were cut to facilitate making surveys of the spawning beds. Expenditures for this work amounted to approximately \$35,000 for pay rolls of temporary employees, and \$2,660 for miscellaneous supplies, including gas and oil for the operation of power boats.

The fishway in Ketchikan Creek was reconstructed and improved, the work being started under the C. W. A. project and completed by the Bureau. By means of this structure the salmon are able to ascend to the spawning grounds above the falls without difficulty.

Other work to increase the returns from the spawning escapement included the destruction of trout that feed upon salmon eggs and fingerlings. This work was carried on throughout the winter of 1934-35 in the Bristol Bay region, where a bounty of 2½ cents each was paid on predatory trout taken by bona fide residents of Alaska. Payments were made from an appropriation for the purpose by the Territorial legislature at its 1933 session and from funds contributed by Bristol Bay packers interested in securing a maximum production from the salmon-breeding reserve. In the Yakutat district, also, considerable work was accomplished in clearing streams and destroying predatory trout through funds supplied by one of the canning companies. Captures of trout were made likewise by weir crews, particularly in the Kodiak area.

STREAM MARKING

New markers defining areas closed to commercial fishing were erected to replace those which had become illegible or damaged, and changes were made in the positions of others to conform with changes made in the regulations with respect to closed areas.

STREAM GUARDS

The Bureau employed 164 men in 1934 as stream guards, weir operators, and special workmen in connection with law-enforcement duties. Of these, 81 were stationed in southeastern Alaska, 56 in central, and 27 in western Alaska. Some of the workers were engaged for only a few days, but the average period of employment ranged from 2 to 5 months.

In southeastern Alaska 48 stream watchmen furnished their own launches and were assigned to patrol larger bodies of water or in the vicinity of several streams.

In central Alaska 14 guards were stationed in the Seward-Katalla district, 10 on Cook Inlet, 21 in the Kodiak-Afognak district, 4 at Chignik, and 7 in the Ikatan-Shumagin district. Eight stream guards, most of whom were in the Seward-Katalla district, provided their own launches.

In western Alaska 24 were on Bristol Bay and 3 in the Yukon-Kuskokwim district.

There were also 7 special employees engaged in scientific work—2 on herring and 5 on salmon investigations, this work being carried on in southeastern and central Alaska.

In addition, there were 12 statutory employees, 51 men on the Bureau's vessels, and 3 on the 3 chartered boats.

The foregoing makes a grand total of 237 persons identified with fishery-protective work in Alaska in 1934, as compared with 203 in 1933.

VESSEL PATROL

Twelve vessels of the Bureau were operated in the Alaska fisheries patrol in 1934, and two, the *Widgeon* and *Petrel*, were out of commission at Seattle. The vessels on duty were assigned as follows: the *Auklet*, *Kittiwake*, and *Teal* in southeast Alaska throughout the season; the *Eider* in the Kodiak area; the *Ibis* at Chignik; the *Murre* in the Seward-Katalla district; the *Red Wing* in the Alaska Peninsula area; and the *Coot* on the Yukon River. The *Scoter* was on Cook Inlet until August 25 and afterwards in the Kodiak area. The *Crane* was engaged in the patrol of the Alaska Peninsula area and carried employees to and from Bristol Bay at the beginning and end of the season. The *Blue Wing* operated in Bristol Bay waters during the fishing season there and later in the Alaska Peninsula area; in the fall it was used in the vicinity of Wrangell. The *Brant* made a cruise as far west as Unalaska in connection with general supervisory work, and from August to October it was engaged in the patrol of southeast Alaska.

The *Eider* and *Brant* also participated successively in the seal patrol in the vicinity of Neah Bay during the spring migration of the fur-seal herd. The *Auklet*, *Crane*, and *Scoter* were used in the C. W. A. project of clearing salmon spawning streams in southeast Alaska in the winter of 1933-34.

Seven small boats, including a speed boat, were operated by the Bureau in the patrol of the Bristol Bay area, and 3 other speed boats were used elsewhere—1 on Copper River and Prince William Sound, 1 in the vicinity of Wrangell, and 1 in the general patrol of southeast Alaska.

In addition to the vessels belonging to the Bureau, several boats were chartered for patrolling fishing areas, as follows: the *Sterling* in the Ketchikan district, the *Lady Luck* in the Wrangell district, and the *Wingham* in the Seward-Katalla district, chiefly on Copper and Bering Rivers. The launch *Marie S* was used on the Kuskokwim River.

Special attention was given to reconditioning and repairing vessels of the Alaska fleet at Seattle in the winter of 1933-34 under an allotment of \$20,000 for this purpose by the Public Works Administration. Nearly all the vessels were overhauled; those receiving the most extensive repairs were the *Brant*, *Teal*, *Eider*, *Kittiwake*, and *Murre*, of the fisheries patrol service, and the Pribilof Islands tender *Penguin*.

The *Merganser*, which for a number of years was used in the patrol of the Ikatan-Shumagin district and more recently was employed by the Division of Fish Culture as local tender for the Yes Bay hatchery, was formally transferred back to the Alaska Division in September 1934. It was reconditioned at Seattle during the winter preparatory to being assigned to patrol duty in the Wrangell district next season.

AERIAL PATROL

The use of aircraft to supplement the regular patrol of the fishing grounds by vessels was on a considerably larger scale in 1934 than in the previous year. Planes were chartered from a commercial company and used for general patrol during weekly closed periods in various parts of Alaska, particularly in the southeastern area and on Copper River Delta and Prince William Sound. Such a patrol has proved very effective, both in detecting and in deterring violations of the law and regulations. Several flights were made also in connection with general supervisory work and observations of the spawning grounds in some sections.

COMPLAINTS AND PROSECUTIONS

A floating trap operated by the Deep Sea Salmon Co. in southeast Alaska was seized on August 9 for illegal fishing. This trap was located on the shore of Harry Bay at a point approximately 3 nautical miles northeast of Cape Fox, where the use of traps was prohibited. The case was tried before the United States Commissioner at Ketchikan, and a fine of \$1,000 was imposed, together with costs of \$70, upon payment of which the trap was released.

Five purse-seine boats, manned by 19 Indians and 1 white, were seized for fishing in closed waters tributary to Portland Canal, the violations having been detected by airplane patrol. The cases were tried in the Commissioner's court and convictions were secured in each instance. Fines amounting to \$150, \$200, and \$250, respectively, were imposed on the crews of the *Prince T-1184*, *Bear T-623*, and *Norma*, which fished near the mouth of a salmon stream in Fillmore Inlet on August 7; while in the case of the *Ulloa* and *Herbert B*, which fished near a salmon stream in Tombstone Bay on August 9, the fines were \$375 and \$300, respectively. When the infringement of the *Herbert B* was observed, the seine had been set to encircle the mouth of the creek, and a member of the crew was upstream, apparently driving the salmon down to the net.

On the west coast of Prince of Wales Island, also, 5 purse-seine boats, each with a crew of 5 men, most of whom were Indians, were seized for fishing in closed waters during August and September. Of these, the *Thelma Jane* and the *Adelia* were found fishing near the mouth of Stoney Creek, the *Lois D* near the mouth of Calder Creek, the *Da-Kee-Noo* in Little Soda Bay, a small indentation of Trocadero Bay, and the *Lilly* near the mouth of a stream at the head of Keete Inlet. Brought to trial before the United States Commissioner at Craig, the defendants pleaded guilty and were assessed fines amounting to \$375 for the crew of each boat, upon payment of which the boats were released to the owners. Each of the boats had a catch of salmon on board at the time of seizure. These fish were sold to local buyers and the proceeds, amounting to a total of \$587.18, were turned over to the Department of Justice.

In the Seward-Katalla district 5 fishermen were arrested for fishing in waters closed to commercial operations. One of these men, charged with fishing within 500 yards of the grass banks on the Copper River delta between Copper River and Pete Dahl sloughs, was taken before the United States Commissioner at Cordova, where he pleaded not guilty and was bound over to the grand jury. The case was later dismissed. Another was fined \$100 for fishing near the mouth of Bear Creek; the sentence was suspended, and the seized gill net was forfeited to the United States. Three of the fishermen were charged with fishing in Cochrane Bay; the case against one was dismissed, as it was found that he had not been connected with the activity; one was fined \$40, and upon failure to pay he was sentenced to serve 20 days in jail; and one was fined \$50, although this was afterwards suspended.

A complaint was filed by local fishermen in Prince William Sound against the Copper River Packing Co. for illegal operation of three salmon traps after the close of the fishing season. The allegation was that the tunnels from pots to spillers had not been disconnected within 36 hours after the beginning of the closed period. At the end of the year the case was still pending.

Two gill-net fishermen were arrested for fishing from a small gas boat in closed waters at the mouth of Kenai River on July 17. They were tried before the local commissioner at Kenai and fined \$15 each. Twenty-five red salmon which were in the net at the time of the seizure were sold to a local buyer and the proceeds were turned over to the Department of Justice.

Four fishermen operating beach seines in the Kodiak area for the Washington Fish & Oyster Co. at Port Williams were charged with violation of the act prohibiting aliens from fishing in the waters of Alaska. Upon trial in the Commissioner's court at Kodiak they pleaded guilty and were fined \$100 each; they elected to serve out their time in jail. Four shackles of gill net, found illegally fishing during a weekly closed period, were seized on July 1 on the mainland shore in the Kodiak area. The owners were not apprehended, and the nets were turned over to the United States deputy marshal at Kodiak for disposition.

Charges were brought against the Alaska Packers Association at Chignik and the superintendent of the cannery there for wanton waste of salmon, as large numbers of short cuts of red salmon were

found dumped on the beach outside the cannery on June 23. Proceedings were started by the United States district attorney at Valdez, but the company requested that action be brought in court at Seattle, Wash., and the request was granted. The charges were accordingly brought in the United States district court at Seattle in December, and on January 19, 1935, the company's attorney appeared before the judge and pleaded guilty for the defendants. Each defendant was fined \$100 and costs, and the fines were paid.

By order of the district attorney at Valdez, the gill net and other fishing gear that had been seized from Harry W. Crosby in July 1933 were taken to Kodiak on the *Penguin* in November 1934 and turned over to the marshal for condemnation.

Four Bristol Bay fishermen operating two gill-net boats of the Red Salmon Canning Co. were arrested for fishing about 7 miles above the markers in Ugashik River on July 14. Upon trial before the local commissioner, they pleaded guilty and were fined, 2 of them at \$50 each, and 2 at \$30 each. The salmon which they had taken were confiscated for the Government and were sold for \$173.37.

DECISION REGARDING LICENSE FEE FOR NONRESIDENT FISHERMEN

On June 4, 1934, the United States Circuit Court of Appeals for the Ninth Circuit at San Francisco rendered a decision affirming that of the District Court of Alaska upholding the right of the Territorial Government of Alaska to impose certain license taxes on fishermen.

The act of the Territorial Legislature approved April 20, 1933, fixed license fees at \$1 for each resident fisherman of all classes and \$25 for each nonresident fisherman.

The suit was brought by a nonresident gill-net fisherman. In seeking to have the decision of the district court set aside, it was contended that the Alaska Legislature has no right to discriminate between citizens of the United States who are residents of Alaska and those who are not.

As bearing upon the matter, the decision of the court of appeals stated, in part, as follows:

The question involved here, then, is not the power of the legislature to discriminate between residents and nonresidents, but the question is whether or not the license fee imposed by the Territorial Legislature is an unreasonable interference with a right granted by Congress, and, therefore, impliedly prohibited by Congress. It is clear, then, that so long as the license tax imposed by the Territorial Legislature upon the citizens of the United States who are not residents of Alaska is not so exorbitant as to practically prohibit, or so unreasonable as to interfere with, the exercise of the right granted by Congress, it is within the power of the Territorial Legislature. We cannot say that the license fee imposed by Territorial Legislature in 1933 and now under attack is so unreasonable as to conflict with the act of Congress granting the right to fish.

Pending this decision nonresident fishermen of Alaska had been paying their licenses under protest, but as the law has been held valid they will not be able to recover such payments.

TERRITORIAL LICENSE TAX

Fisheries license taxes were collected by the Territory under the general revenue law of 1921, as amended in subsequent sessions of the Territorial Legislature. A statement from Oscar G. Olson, Territorial treasurer, under date of May 15, 1935, gives the collections made to

that date for the year 1934. It was stated that collections under the several schedules were fairly complete, although a number of the fisheries companies had not yet made full settlement. The outstanding salmon pack taxes amounted to approximately \$75,000 and about \$10,000 was still due on fish traps, while \$6,000 was still to be collected on fish oil and fertilizer.

Fishery license taxes collected by Territory for fiscal year ended Dec. 31, 1934

| Schedule | Division no. 1 | Division no. 2 | Division no. 3 | Total |
|---|----------------|----------------|----------------|--------------|
| Salmon canneries (pack)..... | \$166,624.86 | \$14.40 | \$574,389.22 | \$741,028.48 |
| Clam canneries..... | 1.00 | | 557.04 | 558.10 |
| Salteries..... | 1,439.69 | 181.82 | 1,542.95 | 3,164.46 |
| Cold-storage plants..... | 900.00 | | | 900.00 |
| Fish-oil works and fertilizer and fish-meal plants..... | 22,279.60 | | 7,672.71 | 29,952.31 |
| Whale oil and fertilizer stations..... | | | 9,549.00 | 9,549.00 |
| Fish traps..... | 81,691.88 | | 44,393.05 | 126,084.93 |
| Gill nets..... | 444.30 | 51.00 | 4,464.40 | 4,959.70 |
| Seines..... | 4,795.00 | | 2,520.00 | 7,315.00 |
| Total..... | 278,176.39 | 247.22 | 645,088.37 | 923,511.98 |
| Salmon canneries (net income), not possible of segregation as to judicial division..... | | | | 19,371.30 |
| Total collections..... | | | | 942,883.28 |

WATER-POWER PROJECTS IN ALASKA

An application for a permit to utilize the waters of Tunnel Creek tributary to Paul Lake, Prince of Wales Island, was referred to the Bureau by the Federal Power Commission for report as to the requirements which should be stipulated in the permit to protect migratory fish. It was found upon investigation that Tunnel Creek is not a salmon spawning stream, and the Commission was informed that no special conditions need be required of the licensee.

KUSKOKWIM RIVER

As in previous years, Stream Guard Charles McGonagall patrolled the Kuskokwim River district for about 3 months, using a chartered launch. Several of the local inhabitants had planned to mild-cure and hard-salt limited amounts of salmon for export, but the shipment of barrels and salt from Seattle was delayed several weeks by the longshoremen's strike, and by the time these supplies reached the Kuskokwim area the run of king salmon was over. There was no commercial fishing for export from Alaska at the mouth of the Kuskokwim in the 1934 season.

Fishing conditions at the mouth of the river were ideal, as few storms occurred during the runs, especially during the king-salmon run, which reached its peak from June 15 to 20. The peak of the runs of chum and red salmon was from June 20 to July 1. Both natives and whites had their catches in the fish houses by the middle of July. Little fishing was done after that date, as there was practically no market for dried fish, the use of airplanes instead of dog teams having been extended to additional routes. Three hundred and eighteen natives fished in the river for local requirements, using 705 gill nets of 10,580 fathoms, 47 wheels, and a number of small boats. They prepared 388 tons of dried chums.

YUKON RIVER

Four outfits engaged in commercial fishing at the mouth of the Yukon River in 1934 and prepared products for the outside market amounting to 447 tierces of mild-cured and 134 barrels of pickled king salmon. The output was considerably curtailed by reason of a shortage of tierces and salt, due to the water-front strike in Seattle which necessitated the discontinuance of operations before the end of June. By the time additional supplies were received, the run of king salmon had passed.

In the fall of 1933 some of the operators in this district had holes dug in the tundra and covered with poles and moss. High tides filled these holes with water, which froze solid during the winter, thus furnishing excellent cold-storage facilities in the curing season.

Operations were begun at the opening of the season on June 5, but catches were light until June 14; a good run of king salmon occurred thereafter, which reached its peak between June 29 and July 7. Chum salmon began to arrive at the same time as the kings, but very little fishing for this species was carried on until after the salteries had closed. A second run of chums entered the river on July 29; this run was exceptionally heavy at the mouth and as far upstream as Pilot Station. The catch from Pilot Station to Tanana, except at a few camps, was the lightest in years, which was attributed largely to the fact that the river was high and full of driftwood, interfering with operation of the wheels.

A patrol of the district was maintained by Inspector C. F. Townsend and a stream guard aboard the *Coot*. The vessel departed from the Government ways at Nenana on May 23 for the mouth of the river and returned to Nenana on September 11.

Products of the Yukon and Tanana fisheries, including the commercial output, were as follows: 144 cases of kings canned, 624 pounds of canned smoked kings, 447 tierces of mild-cured kings, 28,200 pounds of kings and 650 pounds of chums pickled, 200 pounds of kippered kings, and 308 tons of dried chums. Apparatus consisted of 215 wheels, 180 gill nets of 3,561 fathoms, 1 motor vessel of 40 tons, 3 launches, 4 scows, and miscellaneous small boats. There were 34 whites and 393 natives engaged in the fishery.

WEIRS FOR COUNTING SALMON ESCAPEMENT

Twelve weirs for counting the escapement of spawning salmon were operated in Alaska in 1934 to provide information needed in connection with the biological studies of the salmon and the regulation of commercial fishing. This represents an expansion of the weir work over that of the previous year, which was made possible through an allotment of \$6,000 from the Public Works Administration to be used in the fiscal years 1934 and 1935 for reconditioning and reestablishing Alaska weirs. Those reestablished under this appropriation were the Klawak Creek weir in southeast Alaska and the Red River, English Bay, Morzhovoi Bay, and Orzenoi Bay weirs in central Alaska.

Weirs were again maintained at Karluk River, Chinik Creek, Chignik River, and Alitak Bay, where the escapement records have been kept for a series of years. In addition, counting operations were resumed on Situk River and in the salmon stream tributary to

Kaffia Bay, after having been suspended in the 1933 season. Upon the recommendation of Dr. Frederick A. Davidson, who has conducted scientific studies of the pink salmon in southeast Alaska since 1929, operation of the Olive Cove weir was discontinued, and instead thereof a new weir was erected in 1934 at Little Port Walter.

Reports of operations of the weirs and of the counts of salmon in 1934 are as follows:

KLAWAK CREEK

The weir in Klawak Creek was erected in the same location as in the years 1930 to 1932, inclusive. Red salmon began to appear on June 4, but it was not until July 30 that pink salmon passed upstream. The escapement was small until August 18, after which it held up well for several weeks, reaching its peak on September 8, when 51,352 pink salmon were tallied. The total count for the season through October 8 consisted of 406,163 pink salmon, 16,402 chums, 16,374 reds, and 7,341 cohos. L. M. Johnson was the weir operator, under the direction of Warden Alexander P. Romine.

LITTLE PORT WALTER

From August 18 to September 17, inclusive, 6,952 pink salmon, 15 chums, and 1 coho were counted through the new weir in the stream at the head of Little Port Walter. This weir was established primarily to furnish information in connection with the biological studies of pink salmon in southeast Alaska. S. J. Hutchinson was in charge of operations, under the direction of Dr. F. A. Davidson.

SITUK RIVER

The weir in Situk River was completed on June 4, but salmon did not begin to pass through until 2 weeks later. More than 6,000 ascended to the spawning grounds on June 19, and the peak of the run occurred on July 10, with an escapement of 14,548 red salmon. In order to assure the escapement of at least 50 percent of the run of red salmon in this locality, the 60-hour weekly closed period specified by the regulations was extended 48 hours from 6 o'clock antemeridian July 1 and 72 hours beyond 6 o'clock antemeridian of the following Monday, in the Situk and Ahrnklin River locality. When counting was discontinued on August 5 the total escapement numbered 121,529 red salmon, 13,817 pinks, 1,486 kings, and 3 cohos. Assistant Agent Walter W. Kinsey directed the weir operations at this place.

ALITAK BAY

Construction of the cannery station weir on Olga Bay, in the Alitak Bay region, was begun on May 11 and completed on May 14. A trap for the capture of predatory trout was also installed during this period and was operated until the weir was removed. During the season 21,612 trout were taken by the trap and by hook and line. The seaward migration of young red salmon was the largest observed at the cannery stream since 1927.

The first red salmon were tallied on May 23, and counting was continued through August 20, when the total escapement consisted of 70,219 reds and 15,495 pinks. The largest count for any one day

was on August 18, when 7,366 red salmon and 4,722 pinks went through the weir. It was estimated that 75,000 red salmon were in the lagoon and bay at the time the weir was removed; they passed upstream to the spawning beds within the next few days. Many pink salmon also ascended the stream after counting was discontinued.

No count was made of the salmon in the upper station stream, which normally harbors the largest run in the Olga Bay region, but there was a good escapement, estimated to be fully 400,000 red salmon.

The work at this weir was in charge of Henry B. Loeff, under the supervision of Warden Charles P. Turner.

CHIGNIK RIVER

The Chignik weir was located about 150 feet below the site used in the previous year, where the river is approximately 450 feet wide and from 2 to 4½ feet deep. It was of the usual tripod type, with four 22-inch counting gates and a 76-inch gate to permit the passage of small boats. Construction work was started on April 30 and completed on May 21. The river was low at that time and for most of the season, and there were no freshets to damage the structure and interrupt counting operations.

Red salmon began to pass upstream on June 5, but no appreciable numbers appeared until June 10, when a good run started, which continued steadily to July 16 and then gradually diminished to the end of the season. The peak of the run occurred in the last week of June, when the total escapement for the week numbered 370,620 red salmon. The fish of this species were of exceptionally large size, while pinks were smaller than usual. No count was made of the pink salmon, as these fish spawn chiefly in streams emptying into Chignik Bay and Lagoon below the weir.

From June 5 to September 16, inclusive, 1,282,514 reds, 42,174 cohos, and 1,390 kings were counted through the weir. The total commercial catch of red salmon from the Chignik run was 1,032,059. Warden Charles Petry was in charge of the Bureau's work at this place.

CHINIK CREEK

A weir was again operated in Chinik Creek, tributary to Kamishak Bay, and from July 2 to July 30, inclusive, 35,778 red salmon and 162 pinks were counted. As the reported commercial catch exceeded the escapement at the weir on July 6, the locality was closed to commercial fishing until July 24 in order to permit an escapement of 50 percent of the run of red salmon. The work at this weir was in charge of Warden William B. Berry.

ENGLISH BAY

From June 25 to August 3 there were counted 1,655 red salmon and 170 pinks through the weir in the stream at the head of English Bay. A fair run of reds had ascended the stream earlier in June. A trap for the capture of Dolly Varden trout was operated near the weir, and 6,025 of these predatory enemies of the salmon were caught and destroyed. Warden William B. Berry supervised the work at this weir.

KAFLIA BAY

The Kafia Bay weir, located between the second lagoon and the first lake at the head of the bay, was installed on June 9, and the first count of salmon was made on that day. The run was very light until June 29, when a heavy rain raised the river 16 inches. At that time all the fish that had been collecting in the lagoon went up the river. The peak of the run occurred in the first week of July, after which it tapered off gradually. When the weir was removed on August 9 the total count numbered 13,295 red salmon and 375 pinks. This weir was in charge of Joe Zimmerman, who also acted as watchman in the bay, under the direction of Warden Charles P. Turner.

KARLUK RIVER

Construction of the Karluk weir was begun on May 18 and, being facilitated by good weather and a low stage of the river, was completed on May 22. While this work was under way a number of red salmon were already ascending to the spawning grounds, and natives had made some good catches in the lower lagoon the previous week. The escapement was very light, however, until a heavy rain on June 3 caused the fish to move upstream. On the following day 113,421 red salmon were counted, which was by far the largest escapement for any one day during the season. The largest weekly escapement of red salmon was 220,828 for the week ending June 23. Throughout July and August the run was light.

The run of pink salmon began about the middle of July and reached its peak in the week ending August 11. Spent pinks began to come downstream in large numbers on August 18, when the river was rising as a result of heavy rains. On August 20 it was necessary to install additional gates in the weir to allow the dead pinks to pass through, and on August 22 half of the structure was removed, as the crew was unable to handle the increasing number of spent fish. The river at this time was 20 inches above its normal summer level. It was not until September 17 that the weir was reestablished, and only a small escapement occurred thereafter until the weir was removed on October 5.

During the period in which the weir was in operation—from May 23 to August 21 and from September 18 to October 5, inclusive—846,299 red salmon, 978,171 pinks, 9,579 kings, and 6,141 cohos were counted through the gates. It was estimated that 300,000 red salmon ascended to the spawning grounds in the interval from August 22 to September 17, inclusive, which would bring the total escapement of that species to 1,146,299. The reported commercial catch of red salmon from Cape Karluk to West Point was 928,502, indicating that 45 percent of the Karluk run was caught and 55 percent escaped to the spawning grounds.

Young red salmon migrating seaward were seen in the river from May 21 to June 15. Fifty thousand red-salmon fingerlings were marked at Karluk during the season.

Morris Rafn was in charge of the Karluk weir, under the direction of Warden Charles P. Turner.

MORZHOVOI BAY

Red salmon ascending the stream that empties into Middle Lagoon, Morzhovoi Bay, were counted through a weir from June 24 to September 8, inclusive, during which time a total escapement of 81,748 was recorded. This is more than twice as many as the largest count previously made at this place in the years 1926 to 1932, inclusive. At the beginning of the season counting operations were carried on by David E. Hughes, and later by Mike Clipper, under the supervision of Warden Fred R. Lucas.

ORZENOI RIVER

The weir in Orzenoi River was ready for operation on June 23 and the first salmon passed upstream to the spawning grounds on July 4. Counting was continued through July 22, on which date the structure was washed out by a freshet, with the loss of so much material that it could not be replaced for the remainder of the season. The total count numbered 6,634 red salmon and 172 pinks. Thomas B. West was the weir operator, under the direction of Warden Fred R. Lucas.

RED RIVER

From May 23 to August 15, inclusive, 1,159,466 red salmon, 1,369,392 pinks, and 1,994 kings were counted through the weir in Red River, and many fish were still ascending the stream at the time the rack was removed. More than 100,000 predatory trout were exterminated during the season. Henry B. Loeff directed operations at this weir, under the supervision of Warden Charles P. Turner.

SALMON LIFE-HISTORY STUDIES

Studies of the biology of the Alaska salmon were continued in 1934 by the staff of investigators of the Fisheries Biological Station, at Seattle, Wash. Although these studies are carried on each year in more or less definite localities in Alaska, they have as their common goal the determination of the natural and artificial factors responsible for the fluctuations in the abundance of the salmon throughout the entire Territory. Two major investigations dealing with the red salmon, at Karluk and Chignik, and one pertaining to the pink salmon, at Little Port Walter, were in progress during the year. Data pertaining to the runs in all sections of the Territory were likewise collected for the purpose of keeping account of the general biological trend in the salmon fisheries as a whole.

The results from these investigations and the data collected at various points throughout the Territory indicate that the mortality of the young during their life in fresh water and their rate of growth in the ocean are factors which greatly influence the abundance of the salmon that return to spawn each season. Efforts are being made to evaluate these natural factors in order to provide information relative to the probable abundance of each year's salmon runs.

Studies dealing with the seasonal change in the quality of the salmon that enter the commercial fisheries of Alaska were also continued dur-

ing the year. These studies are being carried on in cooperation with the National Cannery Association of Seattle, and a report dealing with the seasonal change in the quality of the pink salmon in southeastern Alaska is being prepared for publication in 1935.

OBSERVATIONS ON THE ESCAPEMENT OF SALMON

Throughout the fishing season the runs of salmon were observed closely to assure the maintenance of an adequate breeding reserve in the streams of all districts, and the spawning grounds of representative streams were inspected after the close of commercial operations.

Southeast Alaska.—The pink salmon runs were late in entering the inside waters of southeastern Alaska, except in the southern district, where they were early. The runs generally were large, but because of low water in the streams the heavy early run in the vicinity of Ketchikan was prevented from reaching the spawning grounds. This run continued until after the close of the season, however, and good escapements were obtained when the fall rains began. The Behm Canal region was particularly well seeded. Streams along the east coast of Prince of Wales Island were well stocked with spawning pinks from the late run in that section. The tributaries of Ernest Sound and Bradford Canal were satisfactorily seeded in spite of heavy drains on the pink salmon runs in those waters by commercial operations. The escapement of pink salmon was good in all important streams on the west coast of Prince of Wales Island, and in the Icy Strait, western, and eastern districts the escapements of this species were reported to be above average. In the Yakutat district the pink run was light, and low water in the streams contributed to unsatisfactory spawning for this species.

The red salmon runs were below normal in all districts except Yakutat and in the vicinity of Wrangell. The run of reds in the Yakutat district was reported to have been better than in 1933, and good escapements were observed in all streams. Good escapements of kings also were obtained in the Yakutat district.

The early runs of chums were small in all districts, and the escapements were below normal in the Wrangell-Petersburg region and on the west coast of Prince of Wales Island. Fair escapements of chums in other districts were derived from late runs, the bulk of which reached the spawning grounds because of the relatively small amount of fall fishing.

The coho runs in all parts of southeastern Alaska were heavier than usual, and the escapements were considerably above average.

Prince William Sound and Copper River region.—The escapement of all species in Prince William Sound streams was very satisfactory. Of particular interest was the much improved pink salmon escapement in streams tributary to Port Wells, where the runs are so seriously depleted as to require prohibition of all commercial fishing. There was a good run of red salmon to Copper River and a good escapement was obtained. The escapement of kings in Copper River also was good. The run of reds to Bering River was much better than in 1933, and it is believed that a 50-percent escapement was obtained. There was no

fall fishing off the mouths of Bering River or Copper River and the entire coho runs escaped to the spawning grounds.

Cook Inlet.—There was a heavy run of red salmon to the Kenai and Kasilof Rivers, and exceptionally good escapements were obtained. The individual fish of this species were said to be of large size. These rivers also had an ample seeding of pinks and kings. Very good escapements of reds were observed in McNeil and Chinik Creeks in Kamishak Bay. Fair seeding of all species was reported in Turnagain Arm and Knik Arm in the northern part of the inlet.

Kodiak area.—Good runs of red salmon appeared in Shelikof Strait, and the spawning grounds of all important streams were well seeded. An exceptionally large escapement of this species was observed in Uganik Bay. Spawning escapements of red salmon in other streams throughout the district were satisfactory except in Kafia Bay, where a light run and small escapement were experienced.

The pink salmon runs to Karluk and Red Rivers were heavy, and on the south side of Kodiak Island the runs of this species were better than had been anticipated. As a result, good escapements were obtained. Light runs and escapements of pinks appeared in Danger Bay, Buskin River, Kalsin Bay, and Uyak Bay and its tributaries. The escapement of chums was generally fair.

Chignik.—The red salmon run at Chignik was heavy during June and July and resulted in a very satisfactory escapement. These fish were reported to be unusually large in size. The spawning grounds were fairly well seeded with pinks and chums, but the coho escapement was small.

Alaska Peninsula.—The runs of all species in this region were normal, but because of the increased intensity of commercial operations the spawning grounds of many streams were inadequately seeded. A late run of pink salmon entering the bays after the close of fishing was largely responsible for the escapement of that species in streams along the south side of the peninsula.

Bristol Bay.—The Kvichak, Naknek, and Egegik Rivers had exceptionally heavy runs of red salmon. The runs began in these rivers in advance of the fishing season and continued at a relatively high level until after the close of operations. Fish were so plentiful that many of the canneries completed their packs before the end of the season. Not only were the reds unusually abundant but they were also larger than average in size. The escapements to these rivers were very large. The run to Nushagak River was smaller in volume and later in appearing than that in the Egegik and Kvichak-Naknek waters. Although not large, the escapement was considered adequate. A run of satisfactory proportions appeared in the Ugashik River, and the escapement was considerably better than for many years.

HATCHERIES

Only one hatchery was engaged in the propagation of salmon in Alaska in 1934. This was the privately owned hatchery on Hugh Smith Lake, which was operated by the Pacific American Fisheries.

From the 20,650,000 red-salmon eggs that were collected at the Hugh Smith Lake hatchery in 1933 there were produced and liberated in Alaska waters 20,030,000 salmon fry. A collection of 10,221,000 red-salmon eggs was made at this hatchery in 1934.

Under provisions of the act of June 26, 1906, the owners of private hatcheries in Alaska who are also packers of canned salmon receive a rebate on license fees and taxes of every nature on their catch and pack of salmon at the rate of 40 cents per 1,000 king or red salmon fry liberated by them in Alaska waters. In the fiscal year ended June 30, 1934, therefore, the rebate due the Pacific American Fisheries on the 20,030,000 red-salmon fry liberated at Hugh Smith Lake during the year amounted to \$8,012.

GENERAL STATISTICS OF THE FISHERIES

The total number of persons engaged in the fisheries of Alaska in 1934 was 26,190, or 4,495 more than in 1933. Fishery products were valued at \$41,963,293, an increase of \$9,836,705, or about 31 percent over the preceding year. Of the total amount, 92.4 percent represented the value of salmon products; 3.8 percent, herring; 1.9 percent, halibut; and 1.9 percent, the value of all other fishery products.

Summary of persons engaged and products of the Alaska fisheries in 1934

| Items | Southeast Alaska | | Central Alaska | | Western Alaska | | Total | |
|---|------------------|--------------|----------------|--------------|----------------|--------------|---------------|--------------|
| | Number | Value | Number | Value | Number | Value | Number | Value |
| PERSONS ENGAGED | | | | | | | | |
| Whites..... | 6,026 | | 4,154 | | 3,936 | | 14,116 | |
| Natives..... | 3,394 | | 1,305 | | 1,355 | | 6,054 | |
| Chinese..... | 162 | | 238 | | 477 | | 877 | |
| Japanese..... | 591 | | 352 | | 370 | | 1,313 | |
| Filipinos..... | 1,317 | | 1,095 | | 599 | | 3,011 | |
| Mexicans..... | 17 | | 1 | | 713 | | 731 | |
| Negroes..... | 2 | | 1 | | 22 | | 25 | |
| Miscellaneous..... | 17 | | 3 | | 43 | | 63 | |
| Total..... | 11,526 | | 7,149 | | 7,515 | | 26,190 | |
| PRODUCTS | | | | | | | | |
| Salmon: | | | | | | | | |
| Canned.....cases..... | 3,295,093 | \$13,981,043 | 2,327,418 | \$11,452,568 | 1,859,319 | \$12,178,339 | 7,481,830 | \$37,611,950 |
| Mild cured.....pounds..... | 3,162,400 | 559,329 | 12,800 | 1,288 | 357,600 | 55,434 | 3,532,800 | 616,111 |
| Pickled.....do..... | | | 254,400 | 17,432 | 133,950 | 11,555 | 387,450 | 28,967 |
| Fresh, for food.....do..... | 1,385,789 | 69,837 | | | | | 1,385,789 | 69,837 |
| Frozen, for food.....do..... | 5,316,574 | 334,812 | | | | | 5,316,574 | 334,812 |
| Fresh, for bait and fox feed.....do..... | 24,600 | 150 | 2,000 | 10 | | | 26,600 | 160 |
| Frozen, for bait.....do..... | 99,700 | 450 | | | | | 99,700 | 450 |
| Dry-salted, dried, and kippered.....do..... | | | 35,450 | 1,794 | 1,404,200 | 56,300 | 1,439,650 | 58,094 |
| Smoked and canned.....cases..... | | | | | 18 | 140 | 18 | 140 |
| Fertilizer.....pounds..... | 702,000 | 10,530 | 313,560 | 4,703 | | | 1,015,560 | 15,233 |
| Oil.....gallons..... | 15,450 | 9,590 | 18,270 | 4,111 | | | 33,720 | 13,701 |
| Herring: | | | | | | | | |
| Fresh, for bait.....pounds..... | 2,117,160 | 21,201 | 1,241,625 | 10,663 | | | 3,358,785 | 31,864 |
| Frozen, for bait.....do..... | 3,009,765 | 20,229 | | | | | 3,009,765 | 20,229 |
| Pickled, for bait.....do..... | | | 23,400 | 166 | | | 23,400 | 166 |
| Pickled, for food: | | | | | | | | |
| Scotch cure.....do..... | 2,137,075 | 119,791 | 3,251,875 | 225,323 | 2,020,250 | 100,362 | 7,409,200 | 445,476 |
| Norwegian cure.....do..... | 21,790 | 965 | | | 208,750 | 7,333 | 230,540 | 8,298 |
| Roused, for food (bloaters stock).....do..... | | | | | 513,050 | 16,854 | 513,050 | 16,854 |
| Spiced.....do..... | 2,600 | 390 | 40,400 | 770 | | | 43,000 | 1,180 |
| Dry-salted.....do..... | | | | | 80,400 | 2,701 | 80,400 | 2,701 |
| Meal.....do..... | 19,956,468 | 316,604 | 7,954,000 | 125,613 | | | 27,910,468 | 442,217 |
| Oil.....gallons..... | 2,615,898 | 446,557 | 1,094,450 | 187,502 | | | 3,710,348 | 634,059 |
| Halibut: | | | | | | | | |
| Fresh.....pounds..... | 7,253,775 | 444,525 | | | | | 7,253,775 | 444,525 |
| Frozen.....do..... | 5,967,563 | 360,260 | | | | | 5,967,563 | 360,260 |

| | | | | | | | | | |
|----------------------|----------|---------|------------|-----------|------------|-----------|------------|-----------|-------------|
| Cod: | | | | | | | | | |
| Dry-salted..... | do. | | | 170,800 | 6,119 | 181,545 | 7,039 | 352,345 | 13,158 |
| Pickled..... | do. | | | 115,160 | 4,245 | 2,000 | 100 | 117,160 | 4,345 |
| Stock fish..... | do. | | | 23,400 | 2,403 | | | 23,400 | 2,403 |
| Whale: | | | | | | | | | |
| Oil..... | gallons. | | | 379,250 | 88,745 | 431,700 | 107,925 | 810,950 | 196,670 |
| Sperm oil..... | do. | | | 9,600 | 2,112 | 66,900 | 14,718 | 76,500 | 16,830 |
| Fertilizer..... | pounds. | | | 1,330,000 | 18,389 | 1,368,000 | 19,116 | 2,698,000 | 37,505 |
| Clam: Canned..... | cases | 71 | 144 | 31,725 | 188,522 | | | 31,796 | 188,666 |
| Crab: | | | | | | | | | |
| Canned..... | do. | 8,736 | 71,566 | 13,067 | 112,486 | | | 21,803 | 184,052 |
| Meat..... | pounds. | 68,082 | 21,254 | 3,880 | 1,164 | | | 71,962 | 22,418 |
| Whole in shell..... | dozen | 545 | 696 | 392 | 497 | | | 937 | 1,193 |
| Shrimp: | | | | | | | | | |
| Canned..... | cases | 93 | 279 | | | | | 93 | 279 |
| Meat..... | pounds. | 375,015 | 118,619 | 2,208 | 688 | | | 377,223 | 119,307 |
| Whole in shell..... | do. | 3,780 | 525 | | | | | 3,780 | 525 |
| Trout: | | | | | | | | | |
| Canned..... | cases | 475 | 1,103 | | | | | 475 | 1,103 |
| Fresh..... | pounds. | 23,372 | 1,441 | | | | | 23,372 | 1,441 |
| Frozen..... | do. | 31,341 | 2,443 | | | | | 31,341 | 2,443 |
| Dried..... | do. | | | 1,500 | 30 | | | 1,500 | 30 |
| Sablefish: | | | | | | | | | |
| Fresh..... | do. | 9,075 | 349 | | | | | 9,075 | 349 |
| Frozen..... | do. | 206,069 | 9,138 | | | | | 206,069 | 9,138 |
| Pickled..... | do. | 200 | 12 | | | | | 200 | 12 |
| Rockfish: | | | | | | | | | |
| Fresh..... | do. | 1,100 | 30 | | | | | 1,100 | 30 |
| Frozen..... | do. | 3,194 | 112 | | | | | 3,194 | 112 |
| Flounder: Fresh..... | do. | 270,000 | 4,000 | | | | | 270,000 | 4,000 |
| Total..... | | | 16,927,974 | | 12,457,353 | | 12,577,966 | | 141,963,293 |

¹ These figures represent the value of the manufactured product. It is estimated that the value of the catch, exclusive of whales, to the fishermen was approximately \$11,707,000. The round weight of the salmon catch landed by the fishermen was approximately 624,651,000 pounds, and the corresponding figure for herring was about 165,637,000 pounds. The cod figures given above do not include the offshore catch from waters adjacent to Alaska, which amounted to 4,930,701 pounds of dry-salted cod and 11,675 pounds of tongues, having a total value of \$213,366, landed at ports of the Pacific Coast States.

SALMON

Considering the salmon runs in Alaska as a whole, there was apparently a greater abundance and more even distribution in all districts than for any preceding year. The output of canned salmon was the largest in the history of the industry, exceeding the previous record pack for 1926 by 828,948 cases. Because of the large production during the summer, few canneries were operated in the fall season.

Heavy runs of red salmon again entered Bristol Bay, and although the catch in these waters was somewhat less than in the previous year, the decrease in numbers was more than offset by the larger size of the fish in 1934. Many of the Bristol Bay packers again had all their cans filled and discontinued operations before the close of the fishing season. Red salmon were plentiful also in the Chignik, Kodiak, Cook Inlet, and Copper River areas. The total catch of this species in Alaska was the largest since 1922, while the amount canned exceeded that for any previous year, which may be attributed in part to a decline in the pickled salmon industry, making almost the entire catch available for canning.

Enormous runs of pink salmon appeared both in southeastern and central Alaska, and the catch and pack of this species were the largest on record. Reports indicated that the fish were unusually small in the early runs in some sections but those that appeared later were of larger size.

Cohos, chums, and kings were fairly abundant, but no unusual catches were made of these species during the season. The total number of king salmon taken was the smallest for any year since 1914, which may be attributed in part to the suspension of operations in certain districts because of a shortage of containers and other supplies due to the longshoremen's strike on the Pacific coast and in part to strikes of fishermen for higher prices. The fishing season for king salmon in the Copper River area opened on May 1, but it was not until May 30 that an agreement regarding prices was reached between cannerymen and fishermen, largely through the mediation of Hugh J. Wade, representative of the National Industrial Recovery Administration in Alaska. When fishing was begun on May 31, the greater part of the king salmon run in that region was over. In southeastern Alaska a strike of the trollers continued through June and July and was finally settled through the organization of a cooperative marketing association. The voluntary curtailment of fall fishing likewise lessened the catch, particularly of cohos and chums.

The total catch of salmon increased 42 percent over that for 1933. By districts, southeastern Alaska and central Alaska showed gains of about 75 percent and 49 percent, respectively, while in western Alaska the catch decreased about 6 percent.

There was an increase of about 30 percent for the whole of Alaska in the number of fathoms of seines used, 5 percent in the number of fathoms of gill nets, and about 15 percent in the number of traps, as compared with those in operation in 1933.

CATCH AND APPARATUS

The total number of seines used in the salmon industry in 1934 was 633, of which 493 were purse seines and 140 beach seines. The purse seines aggregated 76,445 fathoms of webbing, and the beach seines 13,678 fathoms. The number of gill nets used was 3,664, having a total length of 234,456 fathoms. There were 162 driven and 297 floating traps—a total of 459.

Southeastern Alaska was accredited with 346 seines, or a total of 57,690 fathoms, an increase of 22 seines and 5,415 fathoms of webbing over the number used in 1933; also with 304 gill nets, aggregating 21,015 fathoms, an increase of 39 nets, but a decrease of 3,610 fathoms of webbing; and with 28 driven and 262 floating traps, an increase of 9 driven and 20 floating traps, as compared with the number operated in 1933.

Corresponding figures for central Alaska show 281 seines, or 30,753 fathoms, as compared with 158 seines, or 15,632 fathoms, in 1933; 984 gill nets, or 52,815 fathoms, as compared with 956 gill nets, or 44,410 fathoms, in 1933; and 134 driven and 35 floating traps, as compared with 119 driven and 19 floating traps in 1933.

In western Alaska, 6 seines, or 1,680 fathoms of webbing, were used, a decrease of 3 seines, but an increase of 140 fathoms of webbing, as compared with the figures for 1933. There were 2,376 gill nets used, or an aggregate of 160,626 fathoms, an increase of 315 nets and 6,001 fathoms of webbing. No traps were operated in this district.

Seines caught 25 percent of the salmon taken in 1934, gill nets 22 percent, and traps 52 percent, while lines and wheels took the remaining 1 percent.

Percentage of salmon caught in each Alaska district, by principal forms of apparatus

| Apparatus | Southeast Alaska | | Central Alaska | | Western Alaska | |
|----------------|------------------|------|----------------|------|----------------|------|
| | 1933 | 1934 | 1933 | 1934 | 1933 | 1934 |
| Seines..... | 31 | 29 | 28 | 33 | 1 | 5 |
| Gill nets..... | 2 | 1 | 8 | 7 | 97 | 94 |
| Traps..... | 65 | 68 | 64 | 60 | | |
| Lines..... | 2 | 2 | | | | |
| Wheels..... | | | | | 2 | 1 |

The total catch of salmon in 1934 was 116,649,072, an increase of 34,772,652, or 42 percent, over the number taken in 1933. The southeastern and central districts showed gains of 24,679,002 and 11,526,995, respectively, while there was a decrease of 1,433,345 in western Alaska. By species, the catch of cohos increased 750,565; chums, 234,056; pinks, 33,265,900; and reds, 571,217; while the catch of kings decreased 49,086.

Salmon taken in 1934, by apparatus and species, in each geographic section of Alaska

| Apparatus and species | Southeast Alaska | Central Alaska | Western Alaska | Total |
|------------------------|------------------|----------------|----------------|-------------|
| Seines: | | | | |
| Coho, or silver..... | 233,352 | 39,507 | | 272,859 |
| Chum, or keta..... | 2,271,837 | 1,289,960 | 10,860 | 3,572,657 |
| Pink, or humpback..... | 14,039,724 | 8,904,675 | | 22,944,399 |
| King, or spring..... | 1,357 | 1,489 | 1,619 | 4,465 |
| Red, or sockeye..... | 256,492 | 1,172,144 | 1,072,341 | 2,500,977 |
| Total..... | 16,802,762 | 11,407,775 | 1,084,820 | 29,295,357 |
| Gill nets: | | | | |
| Coho, or silver..... | 237,638 | 134,986 | 12,656 | 385,280 |
| Chum, or keta..... | 77,244 | 59,649 | 894,049 | 1,030,942 |
| Pink, or humpback..... | 91,024 | 409,231 | 41,430 | 541,685 |
| King, or spring..... | 32,294 | 72,980 | 93,541 | 198,815 |
| Red, or sockeye..... | 293,072 | 1,799,569 | 21,620,570 | 23,713,211 |
| Total..... | 731,272 | 2,476,415 | 22,662,246 | 25,869,933 |
| Traps: | | | | |
| Coho, or silver..... | 925,261 | 573,560 | | 1,498,821 |
| Chum, or keta..... | 1,456,429 | 1,778,248 | | 3,234,677 |
| Pink, or humpback..... | 36,197,142 | 13,355,087 | | 49,552,229 |
| King, or spring..... | 7,198 | 33,061 | | 40,259 |
| Red, or sockeye..... | 692,319 | 5,301,596 | | 5,993,915 |
| Total..... | 39,278,349 | 21,041,582 | | 60,319,931 |
| Lines: | | | | |
| Coho, or silver..... | 559,763 | | | 559,763 |
| King, or spring..... | 298,183 | | | 298,183 |
| Total..... | 857,946 | | | 857,946 |
| Wheels: | | | | |
| Chum, or keta..... | | | 288,000 | 288,000 |
| King, or spring..... | | | 17,905 | 17,905 |
| Total..... | | | 305,905 | 305,905 |
| Total: | | | | |
| Coho, or silver..... | 1,956,014 | 748,053 | 12,656 | 2,716,723 |
| Chum, or keta..... | 3,805,510 | 3,127,857 | 1,192,909 | 8,126,276 |
| Pink, or humpback..... | 50,327,890 | 22,668,993 | 41,430 | 73,038,313 |
| King, or spring..... | 339,032 | 107,560 | 113,065 | 559,657 |
| Red, or sockeye..... | 1,241,883 | 8,273,309 | 22,692,911 | 32,208,103 |
| Grand total..... | 57,670,329 | 34,925,772 | 24,052,971 | 116,649,072 |

CANNING

CHANGES IN CANNERIES

A new organization, the Scow Bay Packing Co., operated the plant at Scow Bay that had been leased by Hanseth Bros. in 1933. The cannery at Hoonah formerly belonging to the Hoonah Packing Co. was purchased and operated in 1934 by the recently incorporated Icy Straits Salmon Co. The Red Salmon Packers Association bought back the floating plant *Retriever*, which it had sold in 1930 to the New England Fish Co., and operated it in the Yakutat district. A new up-to-date 5-line cannery built by Libby, McNeill & Libby at Taku to replace the plant lost by fire in 1932 was in operation during the season. The New England Fish Co. and the Pacific American Fisheries reopened their canneries at Chatham and Excursion Inlet, respectively, which had been idle since 1931.

In central Alaska the operations of A. S. Day at Fort Liscum were carried on under the name of North Pacific Sea Foods. The Washington Fish & Oyster Co. leased and operated the cannery of Port Williams Packing Co. at Port Williams which had been closed since

1930. The Drier Bay plant of the Prince Packing Co. that had been operated under lease by the New England Fish Co. from 1928 to 1930, inclusive, was taken over and operated by the Alaska Pacific Salmon Co. Other plants that were reopened after having been idle for a year or more were the Kadiak Fisheries Co. at Shearwater Bay, the New England Fish Co. at Cordova, the Shelikof Packing Co. at Zachar Bay, the Shepard Point Packing Co. at Port Ashton, the Strand-Jensen Fisheries Co. at Cordova, and the Fidalgo Island Packing Co. at Port Graham. The Pacific American Fisheries operated its King Cove plant instead of the one at Ikatán.

The Kustatan Packing Co. removed from Kustatan to Anchorage, taking over the plant of the Toman Packing Co. at the latter place. The Farwest Fisheries, Inc., at Anchorage was succeeded by the General Fish Co., but the plant remained idle during the season, the pack being put up by Snug Harbor Packing Co. Only a limited pack of canned salmon was prepared by the Redoubt Bay Packing Co. at Redoubt Bay, and the plant is therefore not included in the list of canneries operated.

The Herendeen Bay Consolidated Canneries did not pack aboard the floating plant *Mazama*, but operated the Port Moller plant of the Pacific American Fisheries which had been closed in 1933.

Direct ownership has been acquired by the Pacific American Fisheries, Inc., of the plants formerly operated by subsidiary concerns, as follows: Petersburg Packing Co. at Petersburg, Alitak Fish Co. at Alitak and Zachar Bay, Shumagin Packing Co. at Squaw Harbor, and Alaska-Portland Packers Association at Naknek River and Nushagak Bay.

NEW CANNERIES

Eight new canneries were operated in 1934, of which 5 were shore plants, as follows: Lane Bros., a hand cannery at Moira Sound; Lindenberger Packing Co., at Craig; Cordova Fisheries Co., Inc., a hand cannery at Cordova; Herbert T. Domenici, at Uyak; and the Puget and Alaska Canning Co., which leased the plant at Seldovia that had formerly been used by the North Pacific Packing Co. for canning clams. New floating canneries were the *La Merced*, which was operated by the Alaska Southern Packing Co., Inc., in the Ikatán-Shumagin region and, in the fall season, at Tenakee in southeast Alaska; the *Memnon*, operated by the Columbia River Packers Association on the north and south sides of the Alaska Peninsula; and a large barge operated by the Western Pacific Packing Co. at Mist Harbor in the Shumagin Islands.

Lauritz Pedersen, of Unga, and Fred Phillips, of Valdez, put up small amounts of hand-packed salmon, but these have not been included in the list of canneries.

CANNERIES NOT OPERATED

Of the plants that had been operated in the previous year, only three were closed in 1934—the Cook Inlet Packing Co. at Seldovia, the Enterprise Seafood Co. at Ninilchik, and the Pacific American Fisheries, Inc., at Ikatán. It is not likely that the Ikatán plant will be operated again, as the machinery has been removed.

The Uzinki cannery of the International Packing Co. has been dismantled, and the cannery of Charles W. Pajoman at Iron Creek

was used as a herring saltery by the Apex Fish Co. There have also been dropped from the list of idle canneries the Tutka Bay cannery of the San Juan Fishing & Packing Co., which was purchased by the Fidalgo Island Packing Co. and dismantled, and the Drier Bay plant of the Alaska Pacific Salmon Co., from which the company moved equipment to the cannery it acquired from the Prince Packing Co. The plant of the Trinity Packing Co. at Three Saints Bay was destroyed by fire in 1931 and has not been rebuilt.

The following canneries were closed during the year but may be reopened:

Southeast Alaska:

| | |
|---|--|
| Alaska Pacific Fisheries..... | Burnett Inlet. Boca de Quadra. Chomly. |
| Alaska Pacific Salmon Co..... | Funter Bay. Pybus Bay. Tenakee. |
| Alaska Packers Association..... | Loring. Wrangell. |
| Alaska Sanitary Packing Co..... | Cape Fanshaw. |
| Columbia River Packers Association..... | Lake Bay. |
| Hoonah Packing Co..... | Gambier Bay. |
| Icy Straits Fisheries, Inc. (floating plant)..... | Idaho Inlet. |
| Libby, McNeill & Libby..... | Klawak. |
| The Nakat Packing Corporation..... | Ketchikan. Boca de Quadra. Dundas Bay. Hunter Bay. |
| Pacific American Fisheries, Inc..... | Kasaan. Ketchikan. Port Walter. Santa Ana. Shakan. |

Central Alaska:

| | |
|--------------------------------------|--|
| Alaska Packers Association..... | Alitak. Chignik. Kasilof. |
| Anderson Mercantile Co., Inc..... | Deep Creek. |
| Blue Island Packing Co..... | Blue Fox Bay. |
| Cook Inlet Packing Co..... | Seldovia. |
| W. G. Culver..... | Point McManus. |
| Enterprise Seafood Co..... | Ninilchik. |
| General Fish Co..... | Anchorage. |
| Gustan & Vogel..... | Point Possession. |
| Hemrich Packing Co..... | Kukak Bay. |
| North Coast Packing Co..... | Ninilchik. |
| Northern Light Packing Co..... | Mountain Slough. Bering River. Chignik. Kenai. |
| Pacific American Fisheries, Inc..... | Orca. Unakwik Inlet. Uyak. Valdez. Zachar Bay. |
| Point Possession Fish Co..... | Point Possession. |
| Redoubt Bay Packing Co..... | Redoubt Bay. |
| E. Sandvik..... | Swansons Creek. |
| Harvey J. Smith..... | West Foreland. |
| Spur Fish Corporation..... | Nikishka Bay. |
| Sunset Packing Co..... | Otter Creek. |
| John Wik..... | Kenai. |
| Jake Young..... | Port Chatham. |

Western Alaska:

| | |
|---|-----------------|
| Alaska Packers Association..... | { Naknek River. |
| Columbia River Packers Association..... | { Nushagak Bay. |
| Herendeen Bay Consolidated Canneries..... | Do. |
| Pacific American Fisheries, Inc..... | Herendeen Bay. |
| Red Salmon Canning Co..... | { Naknek River. |
| | { Nushagak. |
| | Naknek River. |

TOTAL CANNERIES OPERATED

There were 110 canneries operated in Alaska in 1934—44 in southeast, 44 in central, and 22 in western Alaska—which is 7 more in the southeast and 12 more in the central district than in 1933, a net gain of 19 plants. The floating plant *Memnon* of the Columbia River Packers Association was operated both in western and central Alaska, and the Alaska Southern Packing Co.'s cannery *La Merced* put up a small pack in southeastern Alaska in the fall on its return from operations in the Ikatan-Shumagin region. However, these plants are included but once in the total, both being credited to the central district.

Companies that canned salmon in Alaska, number and location of canneries operated, and number of traps owned by each, 1934

[New canneries indicated by (*)]

| Company | Canneries | | Traps | | |
|--|-----------|-----------------------------------|--------|----------|-------|
| | Number | Location | Driven | Floating | Total |
| Southeast Alaska: | | | | | |
| Alaska Pacific Salmon Co..... | 4 | { Kake..... | 1 | 10 | 11 |
| | | { Ketchikan..... | 5 | 6 | 11 |
| | | { Port Althorp..... | | 19 | 19 |
| | | { Rose Inlet..... | | 9 | 9 |
| Alaska Southern Packing Co., Inc..... | 1 | { Tanakee (floating)*..... | | | |
| Annette Island Canning Co..... | 1 | { Metlakatla..... | 1 | 7 | 8 |
| Astoria & Puget Sound Canning Co..... | 1 | { Excursion Inlet..... | | 7 | 7 |
| Beagle Packing Co..... | 1 | { Ketchikan..... | 1 | 4 | 5 |
| Berg Packing Co..... | 1 | { Tongass Narrows (floating)..... | | | |
| Deep Sea Salmon Co..... | 1 | { Skowl Arm..... | | 12 | 12 |
| Diamond K Packing Co..... | 1 | { Wrangell..... | 2 | 3 | 5 |
| Douglas Fisheries Co..... | 1 | { Douglas..... | | | |
| Fidalgo Island Packing Co..... | 2 | { Bay of Pillars..... | 5 | | 5 |
| | | { Ketchikan..... | 3 | 2 | 5 |
| Haines Packing Co..... | 1 | { Letnikof Cove..... | | | |
| P. E. Harris & Co..... | 1 | { Hawk Inlet..... | | 8 | 8 |
| Hood Bay Canning Co..... | 1 | { Hood Bay..... | | 5 | 5 |
| Icy Straits Salmon Co..... | 1 | { Hoonah..... | | | |
| Independent Salmon Canneries, Inc..... | 1 | { Ketchikan..... | | 1 | 1 |
| Kelly Packing Co..... | 1 | { do..... | | | |
| Ketchikan Packing Co..... | 1 | { do..... | | 2 | 2 |
| Klawock Packing Co..... | 1 | { Klawak..... | | | |
| Lane Bros..... | 1 | { Molra Sound *..... | | | |
| | | { Craig..... | 1 | 9 | 10 |
| Libby, McNeill & Libby..... | 4 | { George Inlet..... | | 6 | 6 |
| | | { Taku Harbor..... | 4 | 7 | 11 |
| | | { Yakutat..... | | | |
| Lindenberger Packing Co..... | 1 | { Craig *..... | | | |
| Nakat Packing Corporation, The..... | 3 | { Hidden Inlet..... | | 6 | 6 |
| | | { Union Bay..... | | 6 | 6 |
| | | { Waterfall..... | | 9 | 9 |
| | | { Chatham..... | | 5 | 5 |
| | | { Ketchikan..... | | 6 | 6 |
| | | { Noyes Island..... | | 0 | 0 |
| New England Fish Co..... | 3 | { Klawak..... | | | |
| Ocean Packing Co..... | 1 | { Excursion Inlet..... | 1 | 7 | 8 |
| Pacific American Fisheries, Inc..... | 2 | { Petersburg..... | 2 | 5 | 7 |
| Peril Straits Packing Co..... | 1 | { Todd..... | | 6 | 6 |
| Pyramid Packing Co., Inc..... | 1 | { Sitka..... | | 4 | 4 |
| Red Salmon Packers Association..... | 1 | { Yakutat (floating)..... | | 2 | 2 |
| Scow Bay Packing Co..... | 1 | { Scow Bay..... | | 6 | 6 |
| Sebastian Stuart Fish Co..... | 1 | { Tyee..... | | 6 | 6 |
| Superior Packing Co..... | 1 | { Tenakee..... | | 6 | 6 |
| Ward's Cove Packing Co..... | 1 | { Ward Cove..... | | 4 | 4 |
| Wrangell Packing Co..... | 1 | { Wrangell..... | | 3 | 3 |

Companies that canned salmon in Alaska, number and location of canneries operated, and number of traps owned by each, 1934—Continued

[New canneries indicated by (*)]

| Company | Canneries | | Traps | | |
|---|-----------|---|--------|----------|-------|
| | Number | Location | Driven | Floating | Total |
| Central Alaska: | | | | | |
| Alaska Pacific Salmon Co..... | 2 | Drier Bay..... | | 5 | 5 |
| | | Sand Point..... | 5 | | 5 |
| Alaska Packers Association..... | 2 | Chignik..... | 2 | | 2 |
| | | Karluk..... | 3 | | 3 |
| Alaska Southern Packing Co..... | 1 | Traders Cove and Kupreanof Harbor (floating)* | | | |
| Alaska Year-Round Canneries Co..... | 1 | Seldovia..... | 3 | | 3 |
| | | Chignik..... | 3 | | 3 |
| Columbia River Packers Association..... | 1 | Sankin Bay and Belkofski Bay (floating)* | | | |
| Copper River Packing Co..... | 1 | McClure Bay..... | | 6 | 6 |
| Cordova Fisheries Co., Inc..... | 1 | Cordova *..... | | | |
| Harry W. Crosby..... | 1 | Chignik..... | | | |
| Herbert T. Domenici..... | 1 | Uyak..... | | | |
| Emard Packing Co..... | 1 | Anchorage..... | 10 | | 10 |
| Fidalgo Island Packing Co..... | 1 | Port Graham..... | 9 | | 9 |
| General Fish Co..... | 1 | Anchorage..... | 2 | | 2 |
| Glacier Sea Foods Co..... | 1 | Cordova (floating)..... | 1 | | 1 |
| Grimes Packing Co..... | 1 | Uzinkl..... | | | |
| P. E. Harris & Co..... | 1 | False Pass..... | 8 | | 8 |
| International Packing Co..... | 1 | Sankin Bay and Ivanof Bay (floating)..... | | | |
| Kadiak Fisheries Co..... | 2 | Kodiak..... | 5 | | 5 |
| | | Shearwater Bay..... | 1 | | 1 |
| Kustatan Packing Co..... | 1 | Anchorage..... | | | |
| Libby, McNeill & Libby..... | 1 | Kenai..... | 15 | | 15 |
| New England Fish Co..... | 1 | Cordova..... | 4 | 1 | 5 |
| A. N. Nilson..... | 1 | Portlock..... | | | |
| Ninilchik Packing Co..... | 1 | Ninilchik..... | 2 | | 2 |
| North Pacific Sea Foods..... | 1 | Port Lisicum..... | | | |
| | | Alitak..... | 6 | | 6 |
| | | Chignik..... | 2 | | 2 |
| Pacific American Fisheries, Inc..... | 3 | King Cove..... | 16 | | 16 |
| | | Squaw Harbor..... | 3 | | 3 |
| | | Cordova..... | 2 | 1 | 3 |
| Pioneer Canneries, Inc..... | 1 | Eyak River..... | 3 | 3 | 6 |
| Pioneer Sea Foods Co..... | 1 | Stevens Creek..... | 2 | 1 | 3 |
| Premier Salmon Co..... | 1 | Seldovia *..... | | | |
| Puget and Alaska Canning Co..... | 1 | Port San Juan..... | 2 | 2 | 4 |
| San Juan Fishing & Packing Co..... | 2 | Uganik Bay..... | 4 | | 4 |
| | | Uganik Village..... | | | |
| Albert and Josie Sandvik..... | 1 | Seward..... | | | |
| Seward Fisheries, Inc..... | 1 | Zachar Bay..... | | | |
| Shellkof Packing Co., Inc..... | 1 | Port Ashton..... | | 4 | 4 |
| Shepard Point Packing Co..... | 2 | Shepard Point..... | | 5 | 5 |
| Snug Harbor Packing Co..... | 1 | Snug Harbor..... | 5 | | 5 |
| Strand-Jensen Fisheries Co..... | 1 | Cordova..... | | | |
| Uganik Fisheries, Inc..... | 1 | Uganik..... | 3 | | 3 |
| Washington Fish and Oyster Co., Inc..... | 1 | Port Williams..... | | | |
| Western Pacific Packing Co..... | 1 | Mist Harbor (floating)*..... | | | |
| Western Alaska: | | | | | |
| | | Egegik River..... | | | |
| | | Kvichak Bay (2)..... | | | |
| Alaska Packers Association..... | 7 | Naknek River (2)..... | | | |
| | | Nushagak Bay..... | | | |
| | | Ugashik River..... | | | |
| Alaska Salmon Co..... | 1 | Wood River..... | | | |
| Bristol Bay Packing Co..... | 1 | Kvichak Bay..... | | | |
| Columbia River Packers Association..... | 1 | Port Moller (floating)*..... | | | |
| Herendeen Bay Consolidated Canneries..... | 1 | Port Moller..... | | | |
| | | Egegik River..... | | | |
| | | Ekuk..... | | | |
| Libby, McNeill & Libby..... | 6 | Koggiung..... | | | |
| | | Libbyville..... | | | |
| | | Lockanok..... | | | |
| | | Nushagak..... | | | |
| Lowé Trading Co..... | 1 | Nushagak River (floating)..... | | | |
| Nakat Packing Corporation, The..... | 1 | Nakeen..... | | | |
| Pacific American Fisheries, Inc..... | 2 | Naknek River..... | | | |
| | | Nushagak Bay..... | | | |
| Red Salmon Canning Co..... | 2 | Naknek River..... | | | |
| | | Ugashik River..... | | | |

* Traps only were operated, the fish being packed at other canneries.

LOSSES AND DISASTERS

In southeastern Alaska the reduction plant of the Salmoil Corporation at Mud Bay was destroyed by fire in July, with a loss of \$25,400. Other losses reported for the district included the gas boat *White Bear* and miscellaneous fishing apparatus, the total value of which was \$14,540.

Reported losses in central Alaska were various fishing boats, apparatus, and equipment, as well as damage to buildings and docks, amounting to \$24,361; and in the western district, small boats, nets, and miscellaneous equipment, having a total value of \$26,462.

Twenty-five lives were lost—3 in southeast Alaska, 12 in central, and 10 in western Alaska. Three shoresmen in the southeastern district died of disease. In central Alaska, 3 fishermen, 1 shoresman, and 3 transporters were drowned; 1 fisherman and 3 shoresmen died of disease; and 1 fisherman met death by accident. Five fishermen and 1 transporter in western Alaska were drowned; 1 fisherman died of disease; and 1 fisherman, 1 shoresman, and 1 transporter were killed in accidents.

STATISTICS

There were 110 canneries operated in Alaska in 1934, or 19 more than in the previous year. Employment was given to 21,654 persons, as compared with 17,130 in 1933, an increase of 4,524. White employees increased 2,585; natives, 735; Chinese, 279; Japanese, 330; Filipinos, 774; and miscellaneous (Kanakas, Koreans, and Puerto Ricans), 28; while Mexicans decreased 175, and Negroes 32.

The total pack of canned salmon was 7,481,830 cases, valued at \$37,611,950. This was an increase of 2,256,226 cases, or 43 percent, over the pack of 1933, and an increase in value of \$9,235,936, or about 33 percent. All districts shared in the gain; in southeast Alaska the output increased from 2,087,951 to 3,295,093 cases, or about 58 percent; in central Alaska from 1,485,994 cases to 2,327,418 cases, or about 57 percent; and in western Alaska from 1,651,659 cases to 1,859,319 cases, or about 13 percent. In Alaska as a whole the pack of each species was larger in 1934 than in the preceding year; reds increased from 2,180,283 cases to 2,628,016 cases, or nearly 21 percent; pinks from 2,182,551 to 3,824,193 cases, or 75 percent; cohos from 162,568 to 236,117 cases, or 45 percent; chums from 658,789 to 740,641 cases, or 12 percent; and kings from 41,413 to 52,863 cases, or about 28 percent.

Data are included in the following tables to show comparison of the 1934 pack with the average for the 5 preceding years, 1929 to 1933, by cases of each species and by districts. All species except king salmon show a gain over the 5-year average. By districts, the pack in southeast Alaska increased 38 percent, in central Alaska 37 percent, and in western Alaska 58 percent, making a net increase of 42 percent over the 5-year average for the entire output.

Persons engaged, wages paid, and operating units of Alaska salmon canning industry,
1934

| Items | Southeast Alaska | Central Alaska | Western Alaska | Total |
|----------------------------------|------------------|----------------|----------------|---------------|
| PERSONS ENGAGED | | | | |
| Fishermen: | | | | |
| Whites..... | 1,117 | 1,228 | 1,803 | 4,148 |
| Natives..... | 1,125 | 496 | 438 | 2,061 |
| Chinese..... | 2 | | | 2 |
| Japanese..... | | | 3 | 3 |
| Filipinos..... | 6 | | 3 | 9 |
| Mexican..... | 1 | | | 1 |
| Miscellaneous ¹ | | 3 | | 3 |
| Total..... | 2,251 | 1,729 | 2,247 | 6,227 |
| Shoresmen: | | | | |
| Whites..... | 1,893 | 1,525 | 1,555 | 4,973 |
| Natives..... | 2,009 | 696 | 131 | 2,836 |
| Chinese..... | 157 | 238 | 477 | 872 |
| Japanese..... | 556 | 342 | 367 | 1,265 |
| Filipinos..... | 1,276 | 1,090 | 596 | 2,962 |
| Mexicans..... | 12 | 1 | 713 | 726 |
| Negroes..... | 2 | 1 | 22 | 25 |
| Miscellaneous ¹ | 14 | | 48 | 67 |
| Total..... | 5,919 | 3,893 | 3,904 | 13,716 |
| Transporters: | | | | |
| Whites..... | 658 | 597 | 375 | 1,630 |
| Natives..... | 6 | 62 | 3 | 71 |
| Japanese..... | 6 | | | 6 |
| Filipinos..... | 4 | | | 4 |
| Total..... | 674 | 659 | 378 | 1,711 |
| Total: | | | | |
| Whites..... | 3,668 | 3,350 | 3,733 | 10,751 |
| Natives..... | 3,140 | 1,256 | 572 | 4,968 |
| Chinese..... | 159 | 238 | 477 | 874 |
| Japanese..... | 562 | 342 | 370 | 1,274 |
| Filipinos..... | 1,286 | 1,090 | 599 | 2,975 |
| Mexicans..... | 13 | 1 | 713 | 727 |
| Negroes..... | 2 | 1 | 22 | 25 |
| Miscellaneous ¹ | 14 | 3 | 43 | 60 |
| Grand total..... | 8,844 | 6,281 | 6,529 | 21,654 |
| Wages paid shoresmen..... | \$1,406,624 | \$1,097,830 | \$1,118,220 | \$3,624,684 |
| Wages paid transporters..... | 272,401 | 284,246 | 150,583 | 707,280 |
| OPERATING UNITS | | | | |
| Plants: | | | | |
| Shore canneries..... | 42 | 39 | 21 | 102 |
| Floating canneries: | | | | |
| Power vessels..... | 1 | 3 | 1 | 5 |
| Net tonnage..... | 245 | 5,258 | 1,424 | 6,927 |
| Barges..... | 1 | 2 | | 3 |
| Net tonnage..... | 1,092 | 883 | | 1,975 |
| Total plants operated..... | 44 | 44 | 22 | 110 |
| Vessels: | | | | |
| Power, over 5 tons..... | 361 | 165 | 82 | 608 |
| Net tonnage..... | 7,298 | 5,507 | 24,556 | 37,361 |
| Launches..... | 105 | 161 | 26 | 292 |
| Power dories..... | 23 | 27 | 1 | 51 |
| Gill-net boats..... | 108 | 135 | 1,027 | 1,270 |
| Seine skiffs..... | 126 | 169 | 6 | 301 |
| Other rowboats and skiffs..... | 678 | 675 | 166 | 1,519 |
| Lighters and scows..... | 189 | 190 | 139 | 518 |
| Houseboats..... | 15 | 2 | 27 | 44 |
| Pile drivers..... | 22 | 29 | 14 | 65 |
| Pile pullers..... | 2 | 6 | | 8 |
| Rigging sbows..... | 35 | 10 | | 45 |
| Apparatus: | | | | |
| Purse seines..... | 342 | 144 | 6 | 492 |
| Fathoms..... | 57,290 | 17,400 | 1,680 | 76,370 |
| Beach seines..... | 4 | 121 | | 125 |
| Fathoms..... | 400 | 12,417 | | 12,817 |
| Gill nets..... | 265 | 958 | 1,489 | 2,742 |
| Fathoms..... | 20,790 | 52,110 | 146,360 | 219,260 |
| Traps, driven..... | 28 | 134 | | 162 |
| Traps, floating..... | 262 | 35 | | 297 |

¹ Kanakas, Koreans, Puerto Ricans, etc.

Output and value of canned salmon in Alaska in 1934¹

| Product | Southeast Alaska | | Central Alaska | | Western Alaska | | Total | |
|--------------------|------------------|------------|----------------|------------|----------------|------------|-----------|------------|
| | Cases | Value | Cases | Value | Cases | Value | Cases | Value |
| Coho, or silver: | | | | | | | | |
| ½-pound flat..... | 5,067 | \$40,540 | 718 | \$5,470 | | | 5,785 | \$46,010 |
| 1-pound flat..... | 6,826 | 28,958 | 1,457 | 7,701 | | | 8,283 | 46,659 |
| 1-pound tall..... | 146,634 | 755,434 | 74,196 | 879,374 | 1,219 | \$6,273 | 222,049 | 1,141,081 |
| Total..... | 158,527 | 834,932 | 76,371 | 392,545 | 1,219 | 6,273 | 236,117 | 1,233,750 |
| Chum, or keta: | | | | | | | | |
| ½-pound flat..... | 2,237 | 12,941 | 61 | 346 | | | 2,298 | 13,287 |
| 1-pound tall..... | 391,976 | 1,422,980 | 818,172 | 1,138,196 | 33,196 | 127,383 | 738,343 | 2,688,558 |
| Total..... | 394,212 | 1,435,921 | 818,233 | 1,138,541 | 33,196 | 127,383 | 740,641 | 2,701,845 |
| Pink, or humpback: | | | | | | | | |
| ½-pound flat..... | 27,675 | 182,478 | 1,118 | 6,981 | | | 28,793 | 189,459 |
| 1-pound flat..... | 1,621 | 7,134 | 47 | 188 | | | 1,668 | 7,322 |
| 1-pound tall..... | 2,593,066 | 10,633,959 | 1,198,707 | 4,826,930 | 1,959 | 8,132 | 3,793,732 | 15,469,021 |
| Total..... | 2,622,362 | 10,823,571 | 1,199,872 | 4,834,069 | 1,959 | 8,132 | 3,824,193 | 15,665,802 |
| King, or spring: | | | | | | | | |
| ½-pound flat..... | 4,257 | 42,719 | 4,240 | 43,717 | 1,466 | 13,081 | 9,983 | 99,517 |
| 1-pound flat..... | 4,633 | 34,426 | 3,609 | 33,695 | 1,972 | 12,226 | 10,214 | 80,347 |
| 1-pound tall..... | 6,704 | 37,288 | 20,623 | 113,795 | 5,339 | 30,952 | 32,696 | 182,035 |
| Total..... | 15,594 | 114,433 | 28,472 | 191,207 | 8,797 | 56,259 | 52,893 | 361,899 |
| Red, or sockeye: | | | | | | | | |
| ½-pound flat..... | 22,196 | 220,224 | 62,516 | 615,161 | 3,339 | 32,379 | 88,051 | 867,764 |
| 1-pound flat..... | 11,904 | 92,854 | 58,589 | 428,947 | 2,937 | 22,338 | 78,430 | 544,139 |
| 1-pound tall..... | 70,298 | 456,108 | 588,365 | 3,852,068 | 1,807,872 | 11,925,575 | 2,466,535 | 16,236,751 |
| Total..... | 104,398 | 772,186 | 709,470 | 4,896,176 | 1,814,148 | 11,980,292 | 2,628,016 | 17,648,654 |
| Grand total..... | 3,295,093 | 13,981,043 | 2,327,418 | 11,452,568 | 1,859,319 | 12,178,339 | 7,481,830 | 37,611,950 |

¹ Cases containing ½-pound cans have been reduced ½ in number, and thus, for the purpose of affording fair comparison, all are put upon the basis of forty-eight 1-pound cans to the case.

Output of canned salmon in Alaska, in cases, 1929 to 1934¹

BY SPECIES

| Product | 1929 | 1930 | 1931 | 1932 | 1933 | Average for 5-year period, 1929-1933 | 1934 | Percentage increase or decrease in 1934, as compared with 5-year average |
|--------------------|-----------|-----------|-----------|-----------|-----------|--------------------------------------|-----------|--|
| Coho, or silver: | | | | | | | | |
| ½-pound flat..... | | 271 | | | | 74 | | -100.00 |
| ½-pound flat..... | 7,880 | 18,808 | 9,962 | 3,442 | 3,367 | 8,692 | 5,785 | -33.43 |
| 1-pound flat..... | 6,730 | 5,926 | 2,902 | 1,763 | 4,657 | 4,396 | 8,283 | +88.42 |
| 1-pound tall..... | 157,846 | 307,317 | 167,014 | 142,970 | 154,544 | 183,838 | 222,049 | +20.79 |
| Total..... | 171,956 | 332,422 | 169,878 | 148,175 | 162,568 | 197,000 | 236,117 | +19.86 |
| Chum, or keta: | | | | | | | | |
| ½-pound flat..... | 4,961 | 8,384 | 4,242 | 624 | 658 | 3,774 | 2,298 | -39.11 |
| 1-pound tall..... | 859,551 | 591,550 | 529,579 | 819,932 | 658,131 | 691,748 | 738,343 | +6.74 |
| Total..... | 864,512 | 599,934 | 533,866 | 820,556 | 658,789 | 695,529 | 740,641 | +6.49 |
| Pink, or humpback: | | | | | | | | |
| ½-pound flat..... | | 1,113 | | | | 222 | | -100.00 |
| ½-pound flat..... | 44,762 | 81,064 | 46,524 | 7,166 | 14,857 | 38,875 | 28,793 | -25.93 |
| 1-pound flat..... | 3,910 | 4,887 | 4,410 | | | 2,637 | 1,668 | -36.74 |
| 1-pound tall..... | 2,522,965 | 3,101,490 | 2,902,926 | 2,105,979 | 2,167,694 | 2,560,215 | 3,793,732 | +48.18 |
| Total..... | 2,571,637 | 3,188,534 | 2,953,860 | 2,113,145 | 2,182,551 | 2,601,949 | 3,824,193 | +46.97 |

¹ The number of cases shown has been put upon the common basis of forty-eight 1-pound cans per case.

Output of canned salmon in Alaska, in cases, 1929 to 1934—Continued

BY SPECIES—Continued

| Product | 1929 | 1930 | 1931 | 1932 | 1933 | Average for 5-year period, 1929-1933 | 1934 | Percentage increase or decrease in 1934, as compared with 5-year average |
|-------------------|-----------|-----------|-----------|-----------|-----------|--------------------------------------|-----------|--|
| King, or spring: | | | | | | | | |
| ½-pound flat..... | 16,320 | 17,840 | 13,208 | 11,713 | 9,955 | 13,807 | 9,983 | -27.70 |
| 1-pound flat..... | 26,808 | 23,689 | 16,721 | 14,800 | 10,021 | 18,407 | 10,214 | -44.51 |
| 1-pound tall..... | 28,979 | 18,366 | 21,938 | 43,013 | 21,437 | 26,733 | 32,666 | +22.10 |
| Total..... | 72,107 | 59,922 | 51,867 | 69,526 | 41,413 | 58,967 | 52,863 | -10.35 |
| Red, or sockeye: | | | | | | | | |
| ½-pound flat..... | | 370 | | | | 74 | | -100.00 |
| ¾-pound flat..... | 100,136 | 110,605 | 58,178 | 47,707 | 53,638 | 74,053 | 88,051 | +18.90 |
| 1-pound flat..... | 75,326 | 62,972 | 41,002 | 75,524 | 60,052 | 62,975 | 73,430 | +16.60 |
| 1-pound tall..... | 1,514,465 | 677,567 | 1,595,098 | 1,979,850 | 2,066,593 | 1,566,715 | 2,466,535 | +57.43 |
| Total..... | 1,689,927 | 851,514 | 1,694,278 | 2,103,081 | 2,180,283 | 1,703,817 | 2,628,016 | +54.24 |
| Grand total..... | 5,370,159 | 5,032,326 | 5,403,739 | 5,254,483 | 5,225,604 | 5,257,262 | 7,481,830 | +42.31 |

BY DISTRICTS AND SPECIES

| | | | | | | | | |
|------------------------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|--------|
| Southeast Alaska: | | | | | | | | |
| Coho, or silver..... | 97,847 | 155,652 | 88,455 | 87,038 | 95,805 | 104,959 | 158,527 | +51.04 |
| Chum, or keta..... | 290,797 | 283,478 | 274,248 | 579,443 | 424,861 | 370,565 | 394,212 | +6.38 |
| Pink, or humpback..... | 1,542,615 | 2,309,976 | 2,013,442 | 1,379,066 | 1,478,013 | 1,744,611 | 2,622,362 | +50.31 |
| King, or spring..... | 7,000 | 6,939 | 14,896 | 23,624 | 8,146 | 12,121 | 15,594 | +28.65 |
| Red, or sockeye..... | 162,952 | 221,241 | 147,895 | 138,942 | 81,126 | 160,431 | 104,398 | -30.60 |
| Total..... | 2,101,211 | 2,977,286 | 2,538,936 | 2,208,053 | 2,087,051 | 2,382,687 | 3,265,093 | +38.29 |
| Central Alaska: | | | | | | | | |
| Coho, or silver..... | 71,330 | 173,352 | 81,331 | 60,674 | 65,307 | 60,399 | 76,371 | -15.52 |
| Chum, or keta..... | 497,774 | 284,751 | 193,053 | 147,410 | 207,879 | 266,173 | 313,233 | +17.68 |
| Pink, or humpback..... | 1,025,652 | 859,761 | 940,418 | 724,051 | 704,538 | 850,884 | 1,199,872 | +41.01 |
| King, or spring..... | 35,661 | 32,060 | 27,599 | 32,302 | 23,786 | 30,282 | 28,472 | -5.98 |
| Red, or sockeye..... | 454,086 | 268,621 | 430,153 | 660,161 | 484,484 | 481,301 | 709,470 | +53.80 |
| Total..... | 2,084,503 | 1,618,545 | 1,681,554 | 1,624,598 | 1,485,994 | 1,699,039 | 2,327,418 | +36.98 |
| Western Alaska: | | | | | | | | |
| Coho, or silver..... | 2,779 | 3,418 | 92 | 463 | 1,456 | 1,642 | 1,219 | -25.76 |
| Chum, or keta..... | 75,941 | 31,705 | 66,555 | 93,703 | 26,049 | 58,791 | 33,196 | -43.54 |
| Pink, or humpback..... | 3,390 | 18,797 | | 10,988 | | 6,455 | 1,959 | -69.65 |
| King, or spring..... | 29,446 | 20,923 | 9,372 | 13,600 | 9,481 | 16,564 | 8,797 | -46.89 |
| Red, or sockeye..... | 1,072,889 | 361,652 | 1,107,230 | 1,303,978 | 1,614,073 | 1,092,084 | 1,814,148 | +69.12 |
| Total..... | 1,184,445 | 436,495 | 1,183,249 | 1,421,832 | 1,651,658 | 1,175,536 | 1,859,319 | +58.17 |
| Grand total..... | 5,370,159 | 5,032,326 | 5,403,739 | 5,254,483 | 5,225,604 | 5,257,262 | 7,481,830 | +42.31 |

Relative importance of each species of canned salmon within each district in 1934

| District | Coho | Chum | Pink | King | Red | Total, all species |
|-----------------------|---------|---------|---------|---------|---------|--------------------|
| | Percent | Percent | Percent | Percent | Percent | Percent |
| Southeast Alaska..... | 4.8 | 11.9 | 79.6 | 0.5 | 3.2 | 100.0 |
| Central Alaska..... | 3.3 | 13.5 | 51.5 | 1.2 | 30.5 | 100.0 |
| Western Alaska..... | .0 | 1.8 | .1 | .5 | 97.6 | 100.0 |
| All Alaska..... | 3.2 | 9.9 | 51.1 | .7 | 35.1 | 100.0 |

Relative importance of each district in the production of each species of salmon canned in 1934

| District | Coho | Chum | Pink | King | Red | Total, all species |
|-----------------------|---------|---------|---------|---------|---------|--------------------|
| | Percent | Percent | Percent | Percent | Percent | Percent |
| Southeast Alaska..... | 67.1 | 53.2 | 68.6 | 29.5 | 4.0 | 44.0 |
| Central Alaska..... | 32.4 | 42.3 | 31.4 | 53.9 | 27.0 | 31.1 |
| Western Alaska..... | .5 | 4.5 | .0 | 16.6 | 69.0 | 24.9 |
| Total..... | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 |

Average annual price per case of forty-eight 1-pound cans of salmon, 1924 to 1934

| Product | 1924 | 1925 | 1926 | 1927 | 1928 | 1929 | 1930 | 1931 | 1932 | 1933 | 1934 |
|------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Coho, or silver..... | \$6.83 | \$9.72 | \$8.40 | \$8.51 | \$7.12 | \$7.59 | \$8.26 | \$6.51 | \$4.12 | \$5.20 | \$5.23 |
| Chum, or keta..... | 4.68 | 4.44 | 5.01 | 5.47 | 6.06 | 5.35 | 3.60 | 3.19 | 2.79 | 4.12 | 3.65 |
| Pink, or humpback..... | 4.93 | 5.28 | 5.39 | 5.87 | 6.56 | 6.06 | 4.17 | 3.46 | 3.14 | 4.52 | 4.10 |
| King, or spring..... | 8.89 | 11.91 | 10.37 | 11.25 | 11.13 | 11.92 | 13.32 | 9.40 | 5.46 | 7.51 | 6.85 |
| Red, or sockeye..... | 9.53 | 13.12 | 9.89 | 12.08 | 9.41 | 10.71 | 12.57 | 9.20 | 5.61 | 6.71 | 6.72 |

PACK IN CERTAIN DISTRICTS

Statistics of the salmon pack are again presented for subdivisions of the three main districts of Alaska, and comparison is made with similar statistics for 1933. Where the pack at a given cannery is made up of fish from more than one district, as in the case of that at certain Cordova canneries which pack fish caught both in Prince William Sound and in the Copper River area or at various plants in southeastern Alaska which draw for their supply on the catch of more than one district, due segregation has been made in order to credit each district with the pack from salmon caught therein. These districts are described as follows:

WESTERN ALASKA

Bristol Bay.—The Bering Sea shore, east and north of the Ugashik River.
Port Moller and Herendeen Bay.—Port Moller, Herendeen Bay, and Nelson Lagoon.

CENTRAL ALASKA

Ikatan-Shumagin Islands.—False Pass, Ikatan Bay, King Cove, and the Shumagin Islands.

Chignik.—Canneries located at Chignik.

Kodiak-Afognak Islands.—Kodiak, Spruce, and Raspberry Islands.

Cook Inlet.—The shores of Cook Inlet.

Prince William Sound.—Resurrection Bay to Point Whittsed.

Copper and Bering Rivers.—Point Whittsed to Bering River.

SOUTHEASTERN ALASKA

Yakutat and Dry Bay.—Yakutat Bay to and including Dry Bay.

Icy Strait-Lynn Canal.—West coast of Baranof and Chichagof Islands, the shores of Cross Sound, Icy Strait, Lynn Canal, and Stephens Passage, south to Taku Harbor.

Chatham Strait-Frederick Sound.—Both shores of Chatham Strait and its bays from Point Augusta to Cape Ommaney, and through Frederick Sound and its bays northward to Taku Harbor, including Kake.

Sumner Strait-Dixon Entrance.—Southward from Petersburg and eastward from Port Beauclerc to Cape Chacon and Dixon Entrance, and including all canneries on the mainland and intervening islands from the Stikine River to Portland Canal.

West coast, Prince of Wales Island.—Territory west and south of a line from Cape Chacon to Point Baker and Cape Ommaney.

Pack of canned salmon in Alaska in 1934, by districts ¹

| District | Coho | Chum | Pink | King | Red | Total | Percentage increase over 1933 |
|---|--------------|--------------|--------------|--------------|--------------|--------------|-------------------------------|
| | <i>Cases</i> | <i>Cases</i> | <i>Cases</i> | <i>Cases</i> | <i>Cases</i> | <i>Cases</i> | |
| Bristol Bay..... | 1,219 | 32,222 | 1,959 | 8,580 | 1,726,769 | 1,770,749 | 9.09 |
| Port Moller and Herendeen Bay..... | | 974 | | 217 | 87,379 | 88,570 | 211.44 |
| Ikatan-Shumagin Islands..... | 25,418 | 182,066 | 312,762 | 4,639 | 173,136 | 698,021 | 59.34 |
| Chignik..... | 2,476 | 13,417 | 28,406 | 157 | 122,326 | 166,782 | 90.83 |
| Kodiak-Afognak Islands..... | 9,151 | 73,588 | 420,684 | 344 | 151,924 | 656,691 | 31.55 |
| Cook Inlet..... | 25,665 | 9,141 | 48,186 | 19,148 | 154,697 | 256,837 | 82.33 |
| Prince William Sound..... | 13,661 | 34,421 | 389,676 | 715 | 14,549 | 453,022 | 82.56 |
| Copper and Bering Rivers..... | | | 158 | 3,469 | 92,838 | 96,465 | 32.67 |
| Yakutat and Dry Bay..... | 30,316 | 311 | 4,788 | 4,633 | 19,407 | 59,455 | 63.87 |
| Icy Strait-Lynn Canal..... | 20,291 | 97,572 | 340,032 | 3,216 | 32,035 | 493,146 | 14.21 |
| Chatham Strait-Frederick Sound..... | 27,627 | 185,613 | 442,709 | 912 | 9,237 | 666,096 | 32.37 |
| Sumner Strait-Dixon Entrance..... | 46,493 | 82,038 | 1,278,619 | 850 | 35,409 | 1,443,409 | 83.51 |
| West coast, Prince of Wales Island..... | 33,800 | 28,678 | 566,214 | 5,983 | 8,310 | 632,985 | 91.77 |
| Total..... | 236,117 | 740,641 | 3,824,193 | 52,863 | 2,628,016 | 7,481,830 | 43.18 |

¹ Pack reduced to the basis of forty-eight 1-pound cans per case.

MILD CURING

Operations in the salmon mild-curing business in Alaska were considerably affected by the longshoremen's strike on the Pacific coast and by disagreements over prices between the trollers and packers. Because of the delay in receiving shipments of tierces and salt, no mild curing was undertaken at the mouth of the Kuskokwim River during the season, and the salteries on the Yukon were forced to close down by June 29. The industry in southeast Alaska also was hindered by this disruption of shipping, and although a settlement with respect to Alaska shipments was reached early in June, operations were not resumed because the trollers were dissatisfied with prices offered.

The trollers' strike continued during June and July and was finally settled through an arrangement whereby many of the trollers formed a cooperative marketing association, which contracted for the sale of salmon to the Alaska Coast Fisheries, a subsidiary of the Atlantic & Pacific Packing Co. Under this agreement the fishermen received an advance payment at a specified rate upon delivery of the fish, and a further payment from the proceeds of sale after deduction of handling charges. The motor ship *Donna Lane*, which had been engaged in the herring fishery in the previous year, was used by the cooperative association for splitting and freezing purposes. When fishing was resumed on August 6 the fish were of excellent quality and brought good prices. Large catches were made for about 2 weeks, after which the run decreased.

Eighteen plants in southeast Alaska and four in the western district were engaged in the industry. The reports of these companies, together with the census of the trolling fleet by the Bureau, show a total of 1,281 persons employed.

The total output of mild-cured salmon was 3,532,800 pounds, valued at \$616,111, a decrease of 390,400 pounds in quantity and \$6,717 in value, as compared with the production in 1933.

Persons engaged, wages paid, and operating units, Alaska salmon mild-curing industry, 1934

| Item | Southeast Alaska | Central Alaska | Western Alaska | Total |
|------------------------------|------------------|----------------|----------------|----------|
| PERSONS ENGAGED | | | | |
| Fishermen: | | | | |
| Whites..... | 901 | 2 | 15 | 918 |
| Natives..... | 139 | | 23 | 162 |
| Chinese..... | 2 | | | 2 |
| Japanese..... | 1 | | | 1 |
| Filipinos..... | 12 | | | 12 |
| Mexicans..... | 2 | | | 2 |
| Korean..... | 1 | | | 1 |
| Total..... | 1,088 | 2 | 38 | 1,098 |
| Shoresmen: | | | | |
| Whites..... | 106 | | 6 | 112 |
| Natives..... | 4 | | 49 | 53 |
| Total..... | 110 | | 55 | 165 |
| Transporters: | | | | |
| Whites..... | 13 | | 1 | 14 |
| Natives..... | | | 4 | 4 |
| Total..... | 13 | | 5 | 18 |
| Grand total..... | 1,181 | 2 | 98 | 1,281 |
| Wages paid shoresmen..... | \$48,984 | | \$5,215 | \$54,199 |
| Wages paid transporters..... | \$6,727 | | \$1,178 | \$7,905 |
| OPERATING UNITS | | | | |
| Plants: | | | | |
| Shore..... | 14 | | 3 | 17 |
| Floating: | | | | |
| Power vessel..... | 1 | | | 1 |
| Net tonnage..... | 1,597 | | | 1,597 |
| Barges..... | 3 | | 1 | 4 |
| Net tonnage..... | 720 | | 633 | 1,353 |
| Total plants operated..... | 18 | | 4 | 22 |
| Vessels: | | | | |
| Power, over 5 tons..... | 212 | | 1 | 213 |
| Net tonnage..... | 1,579 | | 40 | 1,619 |
| Launches..... | 555 | 1 | 3 | 559 |
| Power dories..... | | | 2 | 2 |
| Gill-net boats..... | | | 14 | 14 |
| Rowboats and skiffs..... | 106 | 1 | 6 | 113 |
| Lights and scows..... | 2 | | 4 | 6 |
| House-boat..... | 1 | | | 1 |
| Apparatus: | | | | |
| Purse seines..... | | 1 | | 1 |
| Fathoms..... | | 75 | | 75 |
| Gill nets..... | 9 | 1 | 51 | 61 |
| Fathoms..... | 225 | 25 | 2,271 | 2,521 |
| Ltnes..... | 3,207 | | | 3,207 |

Products of Alaska salmon mild-curing industry in 1934

| Products | Southeast Alaska | | Central Alaska | | Western Alaska | | Total | |
|----------------------|------------------|----------|----------------|-------|----------------|----------|-------------|----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Coho, or silver..... | 1 230,400 | \$22,953 | 2 5,600 | \$400 | | | 3 236,000 | \$23,353 |
| Chum, or keta..... | 105,600 | 7,280 | | | | | 4 105,600 | 7,280 |
| King, or spring..... | 2 2,821,600 | 528,556 | 3 7,200 | 898 | 7 357,600 | \$58,484 | 5 3,186,400 | 584,938 |
| Red, or sockeye..... | 4,800 | 540 | | | | | 6 4,800 | 540 |
| Total..... | 3,162,400 | 559,329 | 12,800 | 1,298 | 357,600 | 58,484 | 3,532,800 | 616,111 |

1 239 tierces.
2 7 tierces.
3 295 tierces.

4 132 tierces.
5 3,527 tierces.
6 9 tierces.

7 447 tierces.
8 3,983 tierces.
9 6 tierces.

PICKLING

Several small plants in the central district, chiefly on islands along the south side of the Alaska Peninsula, were engaged in the pickling of salmon in 1934, but the industry as a whole was on a very limited scale. Although red salmon were plentiful in western Alaska, almost the entire catch went into the canned product, and the pickling of salmon was entirely incidental to canning and other branches of the fisheries. The production in the western district was but little more than half that for central Alaska, and no output whatever was reported for southeast Alaska.

Thirty-one persons were employed in the industry, a decrease of 75 from the previous year. The total output was 387,450 pounds, valued at \$28,987, as compared with 1,034,950 pounds valued at \$73,920 in 1933.

*Persons engaged, wages paid, and operating units, Alaska salmon-pickling industry, 1934*¹

| Items | Number | Items | Number | Items | Number |
|------------------------|--------|-----------------------------|---------|-----------------------------|--------|
| PERSONS ENGAGED | | PERSONS ENGAGED—CON. | | OPERATING UNITS—CON. | |
| Fishermen: | | Wages paid shoresmen... | \$1,251 | Apparatus: | |
| Whites..... | 22 | | | Beach seines..... | 15 |
| Natives..... | 4 | OPERATING UNITS | | Fathoms..... | 861 |
| Total..... | 26 | Plants, shore..... | 18 | Gill nets..... | 27 |
| | | Vessels: | | Fathoms..... | 805 |
| Shoresmen: | | Power, over 5 tons... | 1 | | |
| Whites..... | 4 | Net tonnage..... | 5 | | |
| Native..... | 1 | Launches..... | 4 | | |
| Total..... | 5 | Power dories..... | 7 | | |
| Grand total..... | 31 | Gill-net boats..... | 3 | | |
| | | Seine skiffs..... | 13 | | |
| | | Rowboats and skiffs.. | 12 | | |
| | | Lighters and scows.. | 2 | | |

¹ Except for 2 gill nets, aggregating 126 fathoms, from western Alaska, all items are credited to the central district.

Products of Alaska salmon-pickling industry in 1934

| Species | Central Alaska | | Western Alaska | | Total | |
|----------------------|----------------|---------|----------------|--------|---------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Coho, or silver..... | 62,800 | \$3,743 | | | 62,800 | \$3,743 |
| Chum, or keta..... | | | 1,450 | \$72 | 1,450 | 72 |
| King, or spring..... | 12,000 | 1,115 | 34,400 | 3,473 | 46,400 | 4,588 |
| Red, or sockeye..... | 179,600 | 12,574 | 97,200 | 8,010 | 276,800 | 20,584 |
| Total..... | 254,400 | 17,432 | 133,050 | 11,555 | 387,450 | 28,987 |

FRESH SALMON

Two operators in southeast Alaska were engaged primarily in the fresh-salmon business and gave employment to 16 shoresmen and 6 transporters. In addition, 7 operators in the southeastern district produced fresh salmon in connection with mild curing or other branches of the fisheries. The total products amounted to 1,385,789

pounds, valued at \$69,837, as compared with 559,287 pounds valued at \$30,601 in 1933—an increase of about 148 percent in quantity and 128 percent in value.

These figures do not include the salmon sold to halibut boats for bait nor those used for fox feed, which are shown under miscellaneous salmon products.

Products of the Alaska fresh-salmon industry in 1934

| Species | Pounds | Value |
|------------------------|-------------|-----------|
| Coho, or silver..... | 1, 010, 754 | \$50, 479 |
| Chum, or keta..... | 6, 196 | 116 |
| Pink, or humpback..... | 125, 148 | 1, 790 |
| King, or spring..... | 76, 340 | 7, 896 |
| Red, or sockeye..... | 167, 351 | 9, 556 |
| Total..... | 1, 385, 780 | 69, 837 |

FREEZING

The freezing of salmon in Alaska was largely incidental to mild curing and was carried on only in the southeastern district. One cold-storage plant whose chief product was frozen salmon gave employment to 20 white shoresmen. The total output was 5,316,574 pounds, valued at \$334,812, as compared with 4,236,252 pounds, valued at \$221,382, in 1933—an increase of approximately 26 percent in quantity and 51 percent in value.

Products of the Alaska frozen-salmon industry in 1934

| Species | Pounds | Value |
|------------------------|-------------|------------|
| Coho, or silver..... | 3, 930, 192 | \$229, 852 |
| Chum, or keta..... | 67, 730 | 2, 332 |
| Pink, or humpback..... | 95 | 1 |
| King, or spring..... | 1, 274, 257 | 100, 096 |
| Red, or sockeye..... | 44, 300 | 2, 531 |
| Total..... | 5, 316, 574 | 334, 812 |

DRY-SALTED, DRIED, AND OTHER MISCELLANEOUS SALMON PRODUCTS

Several operators in southeast Alaska reported the sale of salmon to halibut boats for bait. Small quantities of dry-salted, dried, and kippered salmon were prepared in central Alaska in conjunction with canning and pickling operations. In the central district, also, a limited amount of fresh salmon for fox feed was reported.

In the fishery of the Yukon, Tanana, and Kuskokwim Rivers, which is carried on principally by natives, 1,404,000 pounds of chum salmon were dried, valued at \$56,280; 624 pounds of kings were smoked and canned, valued at \$140; and 200 pounds of kings were kippered, valued at \$20. In this region 12 whites and 635 natives engaged in the fishery, and the apparatus used consisted of 262 wheels, 834 gill nets of 11,870 fathoms, 1 dory, and 50 rowboats and skiffs.

Production of dry-salted, dried, and other miscellaneous salmon products in Alaska in 1934

| Products | Southeast Alaska | | Central Alaska | | Western Alaska | | Total | |
|---|------------------|-------|----------------|---------|----------------|----------|-----------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Dry-salted: Red, or sockeye..... | | | 26,400 | \$1,584 | | | 26,400 | \$1,584 |
| Dried: | | | | | | | | |
| Coho, or silver..... | | | 1,500 | 30 | | | 1,500 | 30 |
| Chum, or keta..... | | | 3,500 | 70 | 1,404,000 | \$54,280 | 1,407,500 | 56,350 |
| Pink, or humpback..... | | | 3,500 | 70 | | | 3,500 | 70 |
| Total..... | | | 8,500 | 170 | 1,404,000 | 56,280 | 1,412,500 | 56,450 |
| Klippered: | | | | | | | | |
| Coho, or silver..... | | | 300 | 15 | | | 300 | 15 |
| King, or spring..... | | | | | 200 | 20 | 200 | 20 |
| Red, or sockeye..... | | | 250 | 25 | | | 250 | 25 |
| Total..... | | | 550 | 40 | 200 | 20 | 750 | 60 |
| Smoked and canned: King, or spring..... | | | | | 624 | 140 | 624 | 140 |
| Fresh, for bait: Pink, or humpback..... | 24,600 | \$150 | | | | | 24,600 | 150 |
| Frozen, for bait: Chum, or keta..... | 99,700 | 450 | | | | | 99,700 | 450 |
| Fresh, for fox feed: Pink, or humpback..... | | | 2,000 | 10 | | | 2,000 | 10 |
| Grand total..... | 124,300 | 600 | 37,450 | 1,804 | 1,404,824 | 56,440 | 1,566,574 | 58,844 |

BYPRODUCTS

Two reduction plants in southeast Alaska prepared salmon by-products, giving employment to 17 white shoresmen and 7 white transporters. One of these plants, that of the Salmoil Corporation at Mud Bay, was destroyed by fire in July. In central Alaska the manufacture of meal and oil was again carried on by a salmon cannery at Karluk in conjunction with its canning operations. The total production was 1,015,560 pounds of fertilizer, valued at \$15,233, and 33,720 gallons of oil, valued at \$13,701, as compared with 913,358 pounds of fertilizer, valued at \$14,679, and 35,700 gallons of oil, valued at \$5,748, in 1933—an increase of 11 percent in the amount of fertilizer and a decrease of about 6 percent in the output of oil.

HERRING

Although abundant runs of herring appeared on most of the important fishing grounds of Alaska in 1934, many of the fish were not of suitable size for curing. As a result, there was a sharp decline in the output of Scotch-cured herring, while the production of meal and oil was the largest in the history of the industry. The increase in the latter products was particularly marked in the Prince William Sound region.

The Kodiak Island area failed to maintain the position it had held for 2 years as the chief producing center for Scotch-cured herring, and instead dropped to the lowest place, not counting the Cook Inlet area, where the output was negligible. It was said that salteries in the Kodiak region found a much heavier waste in packing than in previous years, because a large percentage of the herring were too small for

curing. As yet, no herring reduction plants have been operated in this part of Alaska. The herring run in Unalaska Bay began on June 15 and continued to July 25, when it stopped abruptly.

The total production of Scotch-cured herring in the Kodiak area in 1934 was 1,455,875 pounds, as compared with 4,130,875 pounds in the previous year. In southeast Alaska the output declined from 3,874,703 pounds in 1933 to 2,137,075 pounds in 1934, and in the Prince William Sound area from 3,046,125 pounds to 1,790,750 pounds. Only in the Aleutian Islands district was there a noticeable increase, the pack for 1934 amounting to 2,015,375 pounds, as compared with 1,589,250 pounds in 1933. Small quantities of cured herring were prepared at Golovin Bay and Cook Inlet. No production was reported for Chignik.

Twenty-two concerns handled herring in southeastern Alaska in 1934, an increase of three over the number shown for the previous year. Five were cold-storage plants that froze herring for bait, and eight engaged only in the production of herring for bait. The following companies operated saltery and reduction plants in this district:

| | |
|---------------------------------|------------------|
| Arentsen & Co. | Big Port Walter. |
| Atlas Packing Corporation | Deep Cove. |
| Buchan & Heinen Packing Co. | Port Armstrong. |
| Chatham Strait Fish Co. | New Port Walter. |
| Northwestern Herring Co. | Port Conclusion. |
| Port Herbert Packing Co. | Port Herbert. |
| Storfold & Grondahl Packing Co. | Washington Bay. |

In central Alaska nine operators engaged in pickling herring, practically all the pack being prepared in the Prince William Sound and Kodiak areas. One firm in the latter district produced bait herring only. The more important operators in the district were as follows:

| | |
|-------------------------------|----------------|
| Salteries: | |
| Apex Fish Co. | Iron Creek. |
| Blue Island Packing Co. | Blue Fox Bay. |
| David Buvick | Shuyak Strait. |
| San Marco Fish Co. (Floating) | Kodiak Island. |
| Saltery and reduction plants: | |
| Chatham Strait Fish Co. | Crab Bay. |
| Evans Bay Packing Co., Inc. | Port Benny. |
| Johnson Fisheries Co. | Thumb Bay. |

The chief operators in the western district were the following, all of whom produced Scotch-cured herring, while two prepared Norwegian-cured herring also, and several put up bloater stock:

| | |
|----------------------------------|---------------|
| Austnes & Rod | Unalaska. |
| Campbell & Dougal | Dutch Harbor. |
| Ed Jacobsen & Co. | Do. |
| Jorden Colombos Lamadies | Do. |
| Northwestern Herring Co. | Do. |
| Peterson & Jorgensen Packing Co. | Do. |
| Polar Packing Co. | Unalaska. |
| Olaf Olsen | Do. |
| Odin Isaacson | Do. |

Biological studies of the Alaska herring were continued by Dr. George A. Rounsefell and Edwin H. Dahlgren. Their investigations were confined mainly to the herring fisheries of Chatham Strait, in the southeastern district.

STATISTICAL SUMMARY

Eight hundred and eighty-five persons engaged in the herring industry in 1934, as compared with 988 in 1933. The number of plants decreased from 31 to 26. Products of the fishery were valued at \$1,603,024, an increase of \$200,830, or 14 percent over 1933, when the total value was \$1,402,194. Scotch-cured herring decreased from 12,651,328 pounds, valued at \$586,331, to 7,409,200 pounds, valued at \$445,476, or 41 percent in quantity and 24 percent in value. Herring for bait increased from 4,471,890 pounds, valued at \$38,509, to 6,391,950 pounds, valued at \$52,259, or 43 percent in quantity and 36 percent in value. Meal increased about 27 percent both in quantity and in value, and oil increased about 20 percent in quantity and 61 percent in value.

Persons engaged, wages paid, and operating units, Alaska herring industry, 1934

| Item | Southeast Alaska | Central Alaska | Western Alaska | Total |
|--------------------------------|------------------|----------------|----------------|-----------|
| PERSONS ENGAGED | | | | |
| Fishermen: | | | | |
| Whites..... | 301 | 84 | 17 | 402 |
| Natives..... | 1 | 3 | 9 | 13 |
| Total..... | 302 | 87 | 26 | 415 |
| Shoresmen: | | | | |
| Whites..... | 234 | 147 | 42 | 423 |
| Natives..... | | 10 | 20 | 36 |
| Total..... | 234 | 157 | 68 | 459 |
| Transporters: Whites..... | | 9 | 2 | 11 |
| Grand total..... | 536 | 253 | 96 | 885 |
| Wages paid shoresmen..... | \$89,550 | \$51,166 | \$17,425 | \$158,141 |
| Wages paid transporters..... | | \$9,544 | \$500 | \$10,044 |
| OPERATING UNITS | | | | |
| Plants: | | | | |
| Shore..... | 7 | 6 | 10 | 23 |
| Floating: | | | | |
| Power vessel..... | | 1 | | 1 |
| Net tonnage..... | | 7 | | 7 |
| Sailing vessel..... | | | | 1 |
| Net tonnage..... | | 1,068 | | 1,068 |
| Scow..... | | 1 | | 1 |
| Total plants operated..... | 7 | 9 | 10 | 26 |
| Vessels: | | | | |
| Power, over 5 tons..... | 44 | 18 | 1 | 63 |
| Net tonnage..... | 1,410 | 461 | 5 | 1,876 |
| Launches..... | 2 | 1 | 1 | 4 |
| Power dories..... | | | 6 | 6 |
| Gill-net boats..... | | | 2 | 2 |
| Seine skiffs..... | 20 | 11 | | 31 |
| Other rowboats and skiffs..... | 20 | 8 | 4 | 32 |
| Lighters and scows..... | 1 | 1 | | 2 |
| Pile drivers..... | 1 | 1 | | 2 |
| Apparatus: | | | | |
| Purse seines..... | 46 | 14 | | 60 |
| Fathoms..... | 7,567 | 2,245 | | 9,812 |
| Gill nets..... | | 1 | 44 | 45 |
| Fathoms..... | | 50 | 1,770 | 1,820 |
| Pound seines..... | 9 | 2 | | 11 |
| Pounds..... | 9 | 11 | | 20 |

Products of Alaska herring industry in 1934

| Item | Southeast Alaska | | Central Alaska | | Western Alaska | | Total | |
|--|------------------|-----------|----------------|-----------|----------------|------------|--------------|-------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Fresh, for bait..... | 2, 117, 160 | \$21, 201 | 1, 241, 025 | \$10, 663 | ----- | ----- | 3, 358, 785 | \$31, 864 |
| Frozen, for bait..... | 3, 009, 765 | 20, 229 | ----- | ----- | ----- | ----- | 3, 009, 765 | 20, 229 |
| Pickled, for bait..... | ----- | ----- | 23, 400 | 166 | ----- | ----- | 23, 400 | 166 |
| Pickled, for food: | | | | | | | | |
| Scotch cure..... | 2, 137, 075 | 119, 791 | 3, 251, 875 | 225, 323 | 2, 020, 250 | \$100, 362 | 7, 409, 200 | 445, 476 |
| Norwegian cure..... | 21, 790 | 965 | ----- | ----- | ----- | 7, 333 | 230, 540 | 8, 298 |
| Roused for food (bloaters stock)..... | ----- | ----- | ----- | ----- | 513, 050 | 16, 854 | 513, 050 | 16, 854 |
| Spiced..... | 2, 600 | 390 | 40, 400 | 770 | ----- | ----- | 43, 000 | 1, 160 |
| Dry salted..... | ----- | ----- | ----- | ----- | 80, 400 | 2, 701 | 80, 400 | 2, 701 |
| Meal..... | 19, 956, 468 | 316, 604 | 7, 954, 000 | 125, 613 | ----- | ----- | 27, 910, 468 | 442, 217 |
| Oil..... | 19, 619, 235 | 446, 557 | 8, 208, 375 | 187, 502 | ----- | ----- | 27, 827, 610 | 634, 059 |
| Total..... | 46, 864, 093 | 925, 737 | 20, 719, 075 | 550, 037 | 2, 822, 450 | 127, 250 | 70, 406, 218 | 1, 603, 024 |

1 2,615,898 gallons.

2 1,094,450 gallons.

3 3,710,348 gallons.

HALIBUT

The American halibut fleet operated in 1934 under a program for controlling production which was a modification of the plan followed in 1933. This program involved restrictions on the number of trips made by each vessel and on the catch per man, but specific dates upon which the vessels should land their catches were not assigned as in 1933. In general this program was only moderately successful, as it was carried out fully in but one area. In spite of the lack of complete cooperation, the market for halibut in 1934 was much more stable than in 1933.

There was a very active demand for halibut livers during 1934, and virtually all of those landed were disposed of under contract at a price about 37 percent above that offered in the preceding year. Considerable profit has resulted to the fishermen in recent years through the pharmaceutical use of halibut liver extracts.

The halibut fishery was governed by regulations of the International Fisheries Commission. Fishing began on March 1 and ended in area no. 2 on August 19 and in areas 1 and 3 on October 27, at which dates the quotas prescribed by regulation were reached.

Biological investigations of the Pacific halibut were continued under the direction of Dr. William F. Thompson. These investigations are being directed primarily toward a determination of the extent to which the stock of halibut must yet be increased in order that there will be sufficient spawn to populate the banks to their proper maximum. The halibut vessel *Eagle* was chartered during the winter of 1933-34 in connection with studies in the Gulf of Alaska.

STATISTICAL SUMMARY

There were 602 persons employed in the Alaska halibut industry in 1934—an increase of 33 over the number reported for the preceding year, and products amounted to 13,221,338 pounds, valued at \$804,785. This production represents the total fares of the Alaska halibut

fleet, which comprises all American vessels landing more than one-half of their catch in Alaska or British Columbia ports rather than in the States. Landings of halibut in Alaska totaled 7,151,669 pounds having a value of \$398,337. In 1933 the landings of the Alaska fleet were 14,068,911 pounds valued at \$726,362, and landings in Alaska amounted to 6,779,768 pounds valued at \$316,310. Fares of the Alaska fleet in 1934 declined 847,573 pounds, or 6 percent in quantity, but increased 11 percent in value over 1933. The landings in Alaska ports in 1934 increased 371,901 pounds or about 5 percent in quantity and about 26 percent in value over the preceding year.

These statistics were compiled from data collected by the International Fisheries Commission and by agents of the Bureau. The amount of halibut livers landed by the Alaska fleet was not reported, but it was stated that there were altogether about 850,000 pounds of halibut, sablefish, "lingcod", and rockfish livers, valued at about \$175,000, landed at Alaska and Pacific coast ports during 1934 by American vessels.

Persons engaged, wages paid, and operating units, Alaska halibut industry, 1934

| Items | Total | Items | Total |
|----------------------------|----------|--------------------------|-------|
| PERSONS ENGAGED | | OPERATING UNITS | |
| Fishermen: Whites | 526 | Vessels: | |
| Shoresmen: | | Power, over 5 tons | 92 |
| Whites | 75 | Net tonnage | 1,630 |
| Native | 1 | Launches | 32 |
| Total | 76 | Skates of lines | 2,498 |
| Grand total | 602 | | |
| Wages paid shoresmen | \$32,805 | | |

NOTE.—No dories are shown, as none were actually engaged in fishing. Heretofore the figures have included life boats.

Products of the Alaska halibut fishery in 1934

| Products | Pounds | Value |
|-------------------------------|------------|-----------|
| Fresh (including local) | 7,253,775 | \$444,625 |
| Frozen | 5,967,563 | 360,260 |
| Total | 13,221,338 | 804,785 |

COD

Cod-fishing operations were again carried on from shore stations in the Shumagin and Aleutian Islands region and resulted in a larger output than for any other year since 1929. Fifty-four persons, consisting of 29 whites and 25 natives, were engaged in the industry, a gain of 19 over the number employed in 1933. Products of the shore fishery were as follows: 352,345 pounds of dry-salted cod, valued at \$13,158; 117,160 pounds of pickled cod, valued at \$4,345; and 23,400 pounds of stockfish, valued at \$2,403—a total of 492,905 pounds, valued at \$19,906, as compared with 338,475 pounds, valued at \$12,907, in 1933.

Seven sailing vessels comprised the Bering Sea offshore fleet, the products of which are not included with the Alaska fisheries output because the vessels operate from and land their fares in ports of the Pacific Coast States. Of these, the *C. A. Thayer* (390 tons), *John A* (235 tons), and *Sophie Christenson* (570 tons) were operated by the Pacific Coast Codfish Co.; the *Azalea* (365 tons) and *Wawona* (413 tons) by the Robinson Fisheries Co.; and the *Louise* (328 tons) and *William H. Smith* (496 tons) by the Union Fish Co.

Products of the offshore fishery were 4,930,701 pounds of dry-salted cod, valued at \$212,510, and 11,675 pounds of tongues, valued at \$856—a total of 4,942,376 pounds, valued at \$213,366, as compared with 4,890,469 pounds, valued at \$166,601, in 1933. The offshore fishery employed 199 persons, or 3 more than in the previous year.

WHALES

The American Pacific Whaling Co. reopened its whaling station at Akutan, which had been idle since 1930, and also operated again at Port Hobron. Seven steam whalers were used by the 2 plants, and employment was given to 210 persons, of whom 188 were whites and 22 natives. There were 465 whales taken during the season, consisting of 232 finback, 166 humpback, 44 sulphur bottom, 21 sperm, and 2 sei whales. This is an increase of 283 over the number taken in the previous year.

Products of the whale fishery were 810,950 gallons of whale oil, valued at \$196,670; 76,500 gallons of sperm oil, valued at \$16,830; 945 tons of fertilizer from meat, valued at \$27,405; and 404 tons of bone fertilizer, valued at \$10,100—a total value of \$251,005, as compared with \$68,989 in 1933.

CLAMS

In the Prince William Sound area 6 plants canned clams, producing about 97 percent of the total Alaska output. The pack limit, as specified by the regulations, of 552,000 pounds for the first half year was reached on June 21, and the summer closed season extended from that date through August 15. The limit of 168,000 pounds for the fall pack was reached in less than 1 month and the season closed on September 11. Operators in this district were the Pioneer Canneries, Inc. and Strand-Jensen Fisheries Co., who prepared the bulk of the pack in connection with their salmon-canning activities, the Gulf Packing Co., which engaged primarily in canning crabs, and Dale and Deville, E. A. Haltness, and S. E. Smith Packing Co.

Notwithstanding the good showing from the Prince William Sound district, the total output of clams in Alaska in 1934 dropped to the lowest level since 1929, due to almost negligible operations in the Kodiak and Cook Inlet areas. Reports indicate that clams were abundant on the mainland beaches in the Kodiak area, but only one plant, the Hand Packing Co., at Halibut Bay, was engaged in the industry there, and that for but a short time. On Cook Inlet the clam pack was produced by the Kustatan Packing Co. and the Ninilchik Packing Co., both of which were engaged primarily in canning salmon. A small quantity of hard-shell clams was packed

in southeast Alaska at the salmon cannery of Lane Brothers, on Moira Sound.

Employment was given to 342 persons, of whom 306 were whites, 21 natives, 10 Japanese, and 5 Filipinos. The output consisted of 31,796 cases, containing 782,388 pounds (780,684 pounds of razor clams, and 1,704 pounds of butter clams), valued at \$188,666. This is a decrease of about 25 percent in quantity and 23 percent in value as compared with figures for 1933, when clam products amounted to 1,045,920 pounds, valued at \$246,338.

Products of the Alaska clam industry in 1934

| Item | Cases | Pounds | Value |
|---------------------------------|---------------|----------------|----------------|
| Mincéd: | | | |
| ½-pound cans (48 to case)..... | 29,151 | 699,624 | \$171,508 |
| 10-ounce cans (48 to case)..... | 1,947 | 58,410 | 11,022 |
| 1-pound cans (48 to case)..... | 81 | 3,888 | 624 |
| 20-ounce cans (24 to case)..... | 97 | 2,910 | 582 |
| Whole: | | | |
| ½-pound cans (48 to case)..... | 76 | 1,824 | 912 |
| 10-ounce cans (48 to case)..... | 310 | 9,300 | 2,410 |
| 1-pound cans (48 to case)..... | 134 | 6,432 | 1,608 |
| Total | 31,796 | 782,388 | 188,666 |

SHRIMP

Operations in the shrimp industry were carried on in 1934 by the same companies as in the previous year. Three plants in southeast Alaska (the Alaskan Glacier Sea Food Co. at Petersburg, and the Reliance Shrimp Co. and Stikine Sea Food Co. at Wrangell) engaged primarily in this business and prepared the bulk of the product. A small pack of canned shrimp in glass jars was put up by John F. Willers, who operated at the plant of the Alaskan Glacier Sea Food Co. Some prospecting was carried on in the Yakutat district during July and August, and it was reported that shrimp were found in many places but not on suitable trawling grounds. In the central district the Northern Sea Food Co. at Cordova produced a limited quantity of shrimp meat in connection with the canning of crabs.

There were 138 persons engaged in the industry, of whom 22 were whites, 70 natives, 27 Japanese, 14 Filipinos, 2 Mexicans, 2 Koreans, and 1 Chinese. Products consisted of 377,223 pounds of shrimp meat, valued at \$119,307; 3,780 pounds of fresh shrimp in shell, valued at \$525; and 453 pounds of canned shrimp meat (93 cases of 6½-ounce jars, 12 jars to the case), valued at \$279—a total of 381,456 pounds, valued at \$120,111. Comparable figures for 1933 show a production of 319,052 pounds, valued at \$102,382.

CRABS

Influenced by favorable market conditions, the production of canned crabs in Alaska in 1934 was the largest in the history of the industry. Eleven companies handled crabs, as compared with 8 in the previous year. Of these, 6 operated in southeast Alaska, as follows—the Alaskan Glacier Sea Food Co., at Petersburg and Hooniah; V. L. Boardway & Co., and Polar Sea Foods, Inc., new outfits at Wrangell and Tenakee Inlet, respectively; Kaylor-Otness, Inc., which pur-

chased the cannery of the Northern Sea Food Co. at Petersburg; Polar Star Packing Co., Ltd., in Peril Strait, aboard a scow that had formerly been used as a salmon cannery by the Diamond K Packing Co.; and the Stikine Sea Food Co., primarily in the shrimp business, at Wrangell.

Five companies operating in the vicinity of Cordova were the Alaska Icepak Corporation, which took over the plant previously belonging to the Alaska Sea Products, Inc.; the Gulf Packing Co.; the Northern Sea Food Co.; the S. E. Smith Packing Co., whose operations were incidental to the canning of clams; and the Strand-Jensen Fisheries Co., which put up crabs for the first time in conjunction with its salmon-canning operations.

Two hundred and eighty persons were engaged in the industry, of whom 235 were whites, 39 natives, 5 Filipinos, and 1 Japanese. Products consisted of 544,311 pounds canned (19,727 cases of ½-pound cans, 1,066 cases of 1-pound cans, and 1,010 cases of 6½-ounce cans, 48 cans to the case), valued at \$184,052; 71,962 pounds of cold-packed meat, valued at \$22,418; and 937 dozen crabs in the shell, valued at \$1,193. The total value of products in 1934 was \$207,663, as compared with \$155,580 in 1933, an increase of 33 percent.

JAPANESE VESSELS IN BERING SEA

In 1934 there were 3 Japanese cannery vessels engaged in canning crab meat in offshore waters of Bering Sea, as compared with 4 such vessels in the previous year. The floating canneries reported in Bering Sea were the *Toten Maru* (formerly the *Nagato Maru*), the *Kasada Maru*, and the *Taihoku Maru*, each of which was accompanied by the usual number of tenders. Two beam trawlers, the *Kitami Maru* and *Adzuchi Maru*, of about 350 tons each, were operated in connection with the cannery vessel *Kasada Maru*.

Fishing operations were chiefly in the vicinity of Amak Island and off the entrance to Port Moller. Crab nets marked with buoys and flags were seen over areas as great as 320 square miles around one cannery vessel. The crab-net area southwest of Amak Island was estimated to be 23 miles long and 2 miles wide. Fishing and canning operations were continuous from about May 15 to August 15.

The training ship *Hakuyo Maru*, of the Imperial Fisheries Institute, again made a trip to Bering Sea for the purpose of instructing students in pelagic-fishery methods and to afford them practice in navigation and seamanship. The vessel was at Dutch Harbor from July 3 to July 7, and on July 8 and 9 the captain and a party visited the fur-seal rookeries at St. Paul Island.

TROUT

A small amount of trout was prepared in Alaska by operators engaged primarily in other branches of the fisheries. The products were as follows: Dolly Vardens, 6,795 pounds fresh, valued at \$612; 5,792 pounds frozen, valued at \$811; 11,400 pounds canned, valued at \$1,103; and 1,500 pounds dried, valued at \$30; steelheads, 16,577 pounds fresh, valued at \$829; and 25,549 pounds frozen, valued at \$1,632. The total output of both species was 67,613 pounds, valued at \$5,017, as compared with 40,355 pounds valued at \$1,965 in 1933.

MISCELLANEOUS FISHERY PRODUCTS

Several species of fish of minor commercial importance are taken in small quantities, chiefly in connection with the halibut fishery, and are landed at ports of Alaska and British Columbia and at Seattle. Such products landed in Alaska in 1934 were as follows: Sablefish, 9,075 pounds fresh, valued at \$349; 206,069 pounds frozen, valued at \$9,138; and 200 pounds pickled, valued at \$12; rockfish, 1,100 pounds fresh, valued at \$30; and 3,194 pounds frozen, valued at \$112; and flounders, 270,000 pounds fresh, for mink feed, valued at \$4,000.

FUR-SEAL INDUSTRY**PRIBILOF ISLANDS****GENERAL ADMINISTRATIVE WORK**

As usual, the major activity on the Pribilof Islands in 1934 was the taking of sealskins for commercial purposes. A total of 53,468 skins was taken, of which St. Paul Island produced 42,972, and St. George Island 10,496. Eighty-five percent of these skins were shipped to the Fouke Fur Co. at St. Louis, Mo., and 15 percent were delivered to a representative of the Canadian Government at Seattle as the share of Great Britain under the terms of the treaty of July 7, 1911. The Japanese share of 15 percent under the same agreement was included in the shipment to St. Louis to be processed and sold along with the share of the United States, settlement being made in money from the net proceeds of the sales.

A seal patrol was maintained by the United States Coast Guard during the northward migration of the seals from central latitudes to the Pribilof Islands and in Bering Sea during the summer sojourn of the seals in those waters.

The natives of the Pribilof Islands, as wards of the Government, were given the customary medical attention, housing, and sustenance, and schooling for the children.

Increased facilities for the handling of sealskins by enlargement of plants, construction of new roads and reconditioning of old ones, repair of equipment and buildings, and the erection of tanks for the storage of gasoline and oil are listed among the improvements during the year. Additional housing for the natives was also provided.

The Navy Department again rendered valuable service to the Bureau by detailing the U. S. S. *Sirius* for one voyage to transport the annual shipment of supplies to the islands and the season's take of sealskins to Seattle. Perishable foodstuffs and a considerable quantity of general merchandise were shipped from Seattle to the Pribilofs on the several voyages of the *Penguin* between those points.

TRANSPORTATION OF SUPPLIES

The U. S. S. *Sirius*, supply vessel of the Navy, sailed from Seattle on July 24 with 768 tons of general supplies for St. George Island and 1,493 tons for St. Paul Island. Other supplies were landed at St. Paul Island for the naval radio station, at Unalaska for the Coast and

Geodetic Survey, and on the Rat Islands for the Navy Department. The ship reached the Pribilof Islands on August 1. Seven days were required in discharging the cargo and loading the outgoing freight which consisted of 510 barrels of sealskins, 73 barrels of blubber, and 537 empty oil drums from St. Paul Island, and 212 barrels of sealskins and 135 empty drums from St. George Island. The *Sirius* sailed for Seattle on August 8 and reached its destination 11 days later.

Additional supplies for the islands, approximating 1,000 tons, were handled by the *Penguin* on five voyages from Seattle in the year.

POWER VESSEL "PENGUIN"

In 1934 the *Penguin* made five trips from Seattle to the Pribilof Islands and return, transporting passengers and freight to and from the islands and intermediate points, and also performing interisland service. The first voyage began January 4 and was completed on March 22, the next was from April 23 to May 28, and the third from June 10 to August 10. On August 25 the *Penguin* sailed from Seattle on its fourth voyage to the Pribilofs and returned on September 22. The fifth and last voyage began October 6 and ended November 15. During the remainder of the year the vessel was docked at Seattle for a general overhauling.

In May, trips were made to Indian settlements along the Alaska Peninsula and on the Aleutian Islands where native laborers were picked up and transported to St. Paul Island to work during the summer months. At the end of the season these natives were returned to their homes.

Commissioner Bell and his party were transported from Ikatan to the Pribilof Islands on July 11-13 for a short inspection and observation of sealing operations. They left the islands on July 14 and were landed at King Cove on July 16.

Service was rendered to the Navy Department, the Steamboat Inspection Service, and the Post Office Department in the transportation of officials and employees to various points in Alaska. Similar service was extended to a number of unofficial travelers. Passenger lists also included a number of Bureau employees and 20 sealing assistants from St. Louis who were engaged in the cleaning and curing of sealskins at the Pribilof Islands.

The *Penguin* cruised 28,502 nautical miles and consumed 69,409 gallons of fuel oil and 958 gallons of lubricating oil during the year.

ROADS

St. Paul Island.—Material progress in the construction of roads was made in 1934 by extending the Zapadni road 11,560 feet and by adding 2,580 feet to the Northeast Point road between the watch house and the killing field. Good roads to all the rookeries are of prime importance in facilitating the rapid transportation of skins from the various killing fields to the washing and blubbering plant in the village. Eventually all the rookeries will be connected with this plant by scoria-surfaced roads.

St. George Island.—The road to Zapadni was extended 1,800 feet.

BUILDINGS

St. Paul Island.—The three houses for natives for which foundations were laid in 1933 were completed in March. An extension was made to the salt house, a concrete floor was laid in the warehouse, and a platform for gasoline storage tanks was constructed. Boat ways at the west landing were installed and a seal driveway was built at Reef. A watch house was also built at Marunich.

Washhouses A and B were roofed, the coal house was enlarged, and the bunk house for white employees at Northeast Point was rebuilt. This work was accomplished under an allotment of funds by the Public Works Administration.

St. George Island.—Further work on the new schoolhouse which was begun in 1933 resulted in completion of the concrete work and outside walls. Satisfactory progress on the interior of the building was also reported.

NATIVES

CENSUS

On December 31, 1934, the total native population on St. Paul Island was 244, including 10 persons temporarily absent from the island, of whom 4 were on St. George Island, 4 at Seattle, 1 at Unalaska, and 1 at Belkofski. Births numbered 12, deaths 8, permanent departures 6, and permanent additions 4, making a net increase in population of 2.

There were 158 native residents on St. George Island as of December 31, 1934. Four natives of St. George Island became permanent residents of St. Paul Island, and there were 5 births and no deaths during the year.

The total population of both islands at the end of 1934 was 402, an increase of 3 over the total for 1933.

MEDICAL SERVICE

Two physicians were stationed at the Pribilof Islands during the year to render medical aid to the native population and to Government employees and their families.

The native girl who had been placed in the Orthopedic Hospital at Seattle in 1931 for treatment for a tubercular hip returned to St. Paul Island on September 6, 1934. The Bureau of Indian Affairs, which had charge of the child while she was convalescent, delivered her at Unalaska and she was transferred from there to St. Paul Island on the *Penguin*.

SCHOOLS

The Bureau maintains schools for the native children on St. Paul and St. George Islands, two teachers being employed on each island. All children between the ages of 6 and 16 are required to attend.

The 1933-34 school year on each island began September 11, 1933, and closed May 11, 1934. On St. Paul Island 16 boys and 18 girls were enrolled in the junior school and 16 boys and 13 girls in the senior school, a total of 63 pupils. On St. George Island 9 boys and 8 girls were enrolled in the junior and 12 boys and 11 girls in the senior school, a total of 40 pupils.

SAVINGS ACCOUNTS

Twelve Pribilof Islands natives have savings accounts in the bank of the Washington Loan & Trust Co., Washington, D. C., on which they receive interest, compounded semiannually. Effective July 1, 1934, the interest rate was reduced from 3 percent to 2½ percent. The Commissioner of Fisheries is the trustee of their funds. The following statement shows the condition of these accounts for the year:

| | |
|---|--------------|
| On hand Jan. 1, 1934..... | \$6, 121. 73 |
| Interest earned from Jan. 1 to Dec. 31, 1934..... | 158. 91 |
| | 6, 280. 64 |
| Withdrawn by natives..... | 520. 00 |
| | 5, 760. 64 |

The following statement shows the amount of money in the individual accounts:

Funds of the Pribilof Islands natives in the custody of the United States Commissioner of Fisheries, as trustee, Dec. 31, 1934

| | | | |
|---------------------------|------------|---------------------------|------------|
| Gromoff, Iuliania..... | \$380. 82 | Merculief, Elizabeth..... | \$68. 44 |
| Kochutin, Alexandra..... | 2, 546. 67 | Merculief, Erena..... | 632. 69 |
| Kozloff, Marina..... | 127. 86 | Merculief, George..... | 104. 30 |
| Kozloff, Raisa..... | 68. 23 | Merculief, Tatiana..... | 650. 88 |
| Lestenkof, Michael..... | 404. 71 | Pankoff, Agrippina..... | 196. 06 |
| Merculief, Alexandra..... | 88. 31 | | |
| Merculief, Daniel..... | 491. 67 | Total..... | 5, 760. 64 |

PAYMENTS FOR TAKING FUR-SEAL SKINS

The natives of the Pribilof Islands are divided into classes according to their ability to perform definite work in the killing and skinning of seals. Six classifications were made, 5 of men and 1 of boys, speed and skill in removing the skins being the determining factor in the personnel of each class. The most experienced and skillful workers were graded as first-class men, while those less experienced and skilled were placed in the lower classes. Boys were employed as apprentices. Advancement through the several grades is governed by the degree of proficiency attained in the specialized work each man is required to perform.

Payments were made at the rate of 50 cents per skin for the total number of skins taken in the season upon the allocation of a definite number of skins per man in each class. In 1934, St. Paul Island produced 42,972 skins and St. George Island 10,496 skins, resulting in a monetary return of \$26,734 to the six classes of workmen. Additional compensation amounting to \$280 was paid to 4 foremen and 4 mess attendants, making a gross income to the natives of \$27,014 on account of sealing operations. The details of these payments are shown in the following table:

Payments to Pribilof Islands natives for taking fur-seal skins, calendar year 1934

| Classification | St. Paul Island | | | St. George Island | | |
|--|-----------------|---------------|------------------|-------------------|---------------|-----------------|
| | Number of men | Share of each | Total | Number of men | Share of each | Total |
| First class..... | 30 | \$483.50 | \$14,505.00 | 27 | \$153.00 | \$4,131.00 |
| Second class..... | 11 | 386.50 | 4,251.50 | 3 | 115.00 | 345.00 |
| Third class..... | 5 | 267.00 | 1,335.00 | 4 | 90.00 | 360.00 |
| Fourth class..... | 4 | 193.50 | 774.00 | 4 | 67.00 | 268.00 |
| Fifth class..... | 9 | 53.00 | 477.00 | 2 | 32.00 | 64.00 |
| Boys' class..... | 7 | 20.50 | 143.50 | 4 | 20.00 | 80.00 |
| Foreman (additional compensation)..... | | | 60.00 | | | 55.00 |
| Do..... | | | 40.00 | | | 45.00 |
| Mess attendants (4)..... | | | 80.00 | | | |
| Total..... | 66 | | 21,666.00 | 44 | | 5,348.00 |

PAYMENTS FOR TAKING FOX SKINS

The natives were paid \$4 in cash for each fox skin taken on the Pribilof Islands in the 1933-34 season. The total payments amounted to \$948 for the 237 skins taken on St. Paul Island and \$2,808 for the 702 skins taken on St. George Island, a total of \$3,756.

FUR SEALS

KILLINGS

In 1934, 53,468 fur seals were killed, of which 42,972 were taken on St. Paul Island and 10,496 on St. George Island. Details in regard to the killings are shown in the following tabulations:

Seal killings on Pribilof Islands in 1934

ST. PAUL ISLAND

| Date | Serial no. of drive | Hauling ground | Skins secured |
|---------|---------------------|-----------------------------------|---------------|
| June 12 | 1 | Reef..... | 141 |
| 15 | 2 | Sea Lion Rock (Sivutch)..... | 66 |
| 18 | 3 | Zapadni and Little Zapadni..... | 79 |
| 19 | 4 | Vostochni and Morjovi..... | 266 |
| 20 | 5 | Polovina and Little Polovina..... | 28 |
| 21 | 6 | Tolstoi and Lukanin..... | 65 |
| 22 | 7 | Reef and Gorbatch..... | 757 |
| 23 | 8 | Zapadni and Little Zapadni..... | 388 |
| 24 | 9 | Polovina and Little Polovina..... | 82 |
| 25 | 10 | Vostochni and Morjovi..... | 1,150 |
| 26 | 11 | Tolstoi, Lukanin, and Kitovi..... | 232 |
| 27 | 12 | Reef and Gorbatch..... | 2,117 |
| 28 | 13 | Zapadni and Little Zapadni..... | 714 |
| 29 | 14 | Polovina and Little Polovina..... | 307 |
| 30 | 15 | Vostochni and Morjovi..... | 1,162 |
| July 1 | 16 | Reef and Gorbatch..... | 1,464 |
| 2 | 17 | Tolstoi, Lukanin, and Kitovi..... | 687 |
| 3 | 18 | Zapadni and Little Zapadni..... | 822 |
| 5 | 19 | Vostochni and Morjovi..... | 1,344 |
| 6 | 20 | Tolstoi, Lukanin, and Kitovi..... | 988 |
| 7 | 21 | Reef and Gorbatch..... | 2,773 |
| 8 | 22 | Zapadni and Little Zapadni..... | 1,745 |
| 9 | 23 | Polovina and Little Polovina..... | 902 |
| 10 | 24 | Vostochni and Morjovi..... | 2,455 |
| 11 | 25 | Tolstoi, Lukanin, and Kitovi..... | 353 |
| 12 | 26 | Reef and Gorbatch..... | 2,839 |
| 13 | 27 | Zapadni and Little Zapadni..... | 1,374 |
| 14 | 28 | Polovina and Little Polovina..... | 788 |
| 15 | 29 | Vostochni and Morjovi..... | 1,828 |
| 16 | 30 | Tolstoi, Lukanin, and Kitovi..... | 755 |
| 17 | 31 | Reef and Gorbatch..... | 1,820 |

Seal killings on Pribilof Islands in 1934—Continued

ST. PAUL ISLAND—Continued

| Date | Serial no. of drive | Hauling ground | Skins secured |
|------|---------------------|--|---------------|
| 18 | 32 | Zapadni and Little Zapadni..... | 599 |
| 19 | 33 | Polovina and Little Polovina..... | 658 |
| 20 | 34 | Vostochni and Morjovi..... | 1,519 |
| 21 | 35 | Tolstoi, Lukanin, and Kitovi..... | 680 |
| 22 | 36 | Reef and Gorbatch..... | 1,521 |
| 23 | 37 | Zapadni and Little Zapadni..... | 1,110 |
| 24 | 38 | Vostochni, Morjovi, and Polovina..... | 1,619 |
| 25 | 39 | Reef and Gorbatch..... | 1,191 |
| 26 | 40 | Zapadni, Little Zapadni, Tolstoi, Lukanin, and Kitovi..... | 1,148 |
| 27 | 41 | Vostochni, Morjovi, Polovina, and Little Polovina..... | 737 |
| 28 | 42 | Reef and Gorbatch..... | 901 |
| 29 | 43 | Zapadni, Little Zapadni, Tolstoi, Lukanin, and Kitovi..... | 809 |
| | | Total..... | 42,972 |

ST. GEORGE ISLAND

| | | | | |
|------|----|------------------------------|------------------------------|-------|
| June | 9 | 1 | North..... | 37 |
| | 16 | 2 | do..... | 30 |
| | 19 | 3 | East..... | 68 |
| | 20 | 4 | North and Staraya Artil..... | 155 |
| | 23 | 5 | East..... | 45 |
| | 24 | 6 | North and Staraya Artil..... | 375 |
| | 25 | 7 | Zapadni..... | 52 |
| | 27 | 8 | East..... | 69 |
| | 28 | 9 | North and Staraya Artil..... | 620 |
| July | 29 | 10 | Zapadni..... | 50 |
| | 1 | 11 | East..... | 336 |
| | 2 | 12 | North and Staraya Artil..... | 723 |
| | 3 | 13 | Zapadni..... | 135 |
| | 5 | 14 | East..... | 242 |
| | 6 | 15 | North and Staraya Artil..... | 1,308 |
| | 7 | 16 | Zapadni..... | 118 |
| | 9 | 17 | East..... | 122 |
| | 10 | 18 | North and Staraya Artil..... | 796 |
| | 11 | 19 | Zapadni..... | 86 |
| | 13 | 20 | East..... | 207 |
| | 14 | 21 | North and Staraya Artil..... | 1,015 |
| 15 | 22 | Zapadni..... | 147 | |
| 17 | 23 | East..... | 347 | |
| 18 | 24 | North and Staraya Artil..... | 498 | |
| 19 | 25 | Zapadni..... | 84 | |
| 21 | 26 | East..... | 247 | |
| 22 | 27 | North and Staraya Artil..... | 1,100 | |
| 24 | 28 | East..... | 186 | |
| 25 | 29 | North and Staraya Artil..... | 542 | |
| 27 | 30 | East and Zapadni..... | 175 | |
| 28 | 31 | North and Staraya Artil..... | 298 | |
| 29 | 32 | East..... | 193 | |
| | | Total..... | 10,496 | |

AGE CLASSES

The age class of a male seal of the Pribilof Islands herd is determined from the length of its body. The classification was derived from measurements of a large number of pups branded in 1912 and killed in subsequent years. The limits of the various age classes are shown in the table following:

Age classes of male seals, Pribilof Islands

| Age | Length (inches) | Age | Length (inches) |
|------------------|-----------------|------------------|-----------------|
| Yearlings..... | Up to 36.75. | 4-year-olds..... | 46 to 51.75. |
| 2-year-olds..... | 37 to 40.75. | 5-year-olds..... | 52 to 57.75. |
| 3-year-olds..... | 41 to 45.75. | 6-year-olds..... | 58 to 63.75. |

Ages of seals killed on Pribilof Islands, calendar year 1934

[On basis of classification shown in preceding table]

| Age | St. Paul Island | St. George Island | Total |
|-------------------------|-----------------|-------------------|--------|
| 2-year-old males..... | 777 | 164 | 941 |
| 3-year-old males..... | 41,818 | 10,117 | 51,935 |
| 4-year-old males..... | 330 | 202 | 532 |
| Cows ¹ | 47 | 13 | 60 |
| Total..... | 42,972 | 10,496 | 53,468 |

¹ Cows unavoidably and accidentally killed or found dead.

Some of the seals recorded in the above tabulation as 2-year-olds and 4-year-olds probably were 3-year-olds, as not all male seals of a given age fall within the length limits assigned for the males of that age. As far as possible, the killings in 1934 were confined to 3-year-old males.

RESERVING OPERATIONS

No 3-year-old male seals were marked for the breeding reserve in 1934. It was evident at the close of the season that the number of adolescent males of this age class was ample to maintain in subsequent years the supply of bulls in sufficient strength to meet all breeding requirements of the herd.

COMPUTATION OF FUR-SEAL HERD

The computation of the fur-seal herd in 1934 was made by Supt. H. J. Christoffers. As of August 10 the total of all classes was 1,430,418—a numerical increase of 111,850 over the figures for the preceding year. The detailed report will be found on pages 67 to 73 of this document. Following is a comparative statement of the numerical strength of the various elements of the fur-seal herd in the years 1923 to 1934, inclusive:

General comparison of computations of the seal herd on the Pribilof Islands, 1923 to 1934

| Classes | 1923 | 1924 | 1925 | 1926 | 1927 | 1928 |
|-----------------------|---------|---------|---------|---------|---------|---------|
| Harem bulls..... | 3,412 | 3,516 | 3,526 | 4,034 | 4,643 | 6,050 |
| Breeding cows..... | 197,659 | 208,396 | 226,090 | 244,114 | 263,566 | 284,725 |
| Surplus bulls..... | 1,891 | 2,043 | 3,558 | 2,002 | 4,827 | 5,285 |
| Idle bulls..... | 312 | 390 | 311 | 423 | 972 | 1,449 |
| 6-year-old males..... | 4,863 | 8,489 | 4,105 | 13,434 | 13,460 | 12,857 |
| 5-year-old males..... | 10,612 | 5,132 | 16,792 | 16,812 | 16,073 | 13,001 |
| 4-year-old males..... | 5,710 | 18,670 | 18,692 | 17,872 | 14,448 | 7,798 |
| 3-year-old males..... | 22,786 | 21,551 | 21,185 | 17,189 | 9,730 | 11,133 |
| 2-year-old males..... | 43,112 | 45,665 | 43,515 | 38,183 | 41,252 | 49,087 |
| Yearling males..... | 55,769 | 59,291 | 52,091 | 56,514 | 61,026 | 65,861 |
| 2-year-old cows..... | 48,801 | 51,359 | 49,786 | 44,415 | 48,186 | 57,061 |
| Yearling cows..... | 60,422 | 64,240 | 57,309 | 62,175 | 67,131 | 72,481 |
| Pups..... | 197,659 | 208,396 | 226,090 | 244,114 | 263,566 | 284,725 |
| Total..... | 653,008 | 697,158 | 723,050 | 761,281 | 808,870 | 871,513 |

General comparison of computations of the seal herd on the Pribilof Islands, 1929
to 1934—Continued

| Classes | 1929 | 1930 | 1931 | 1932 | 1933 | 1934 |
|-----------------------|-----------------|--------------------|--------------------|--------------------|--------------------|--------------------|
| Harem bulls..... | 7, 187 | 8, 312 | 9, 233 | 10, 088 | 10, 213 | 10, 770 |
| Breeding cows..... | 307, 491 | 332, 064 | 358, 642 | 387, 320 | 418, 299 | 451, 751 |
| Surplus bulls..... | 5, 207 | 3, 963 | 3, 201 | 2, 893 | 4, 700 | 6, 494 |
| Idle bulls..... | 1, 633 | 1, 899 | 1, 888 | 2, 349 | 2, 341 | 2, 282 |
| 6-year-old males..... | 10, 399 | 5, 612 | 6, 553 | 8, 164 | 9, 335 | 8, 173 |
| 5-year-old males..... | 7, 016 | 8, 191 | 10, 193 | 11, 689 | 10, 216 | 13, 897 |
| 4-year-old males..... | 9, 102 | 11, 327 | 12, 966 | 11, 351 | 16, 441 | 15, 862 |
| 3-year-old males..... | 13, 639 | 14, 871 | 13, 198 | 17, 849 | 18, 216 | 24, 770 |
| 2-year-old males..... | 64, 354 | 69, 674 | 74, 828 | 81, 101 | 87, 662 | 94, 920 |
| Yearling males..... | 86, 381 | 92, 232 | 99, 612 | 107, 592 | 116, 195 | 125, 490 |
| 2-year-old cows..... | 67, 210 | 72, 605 | 78, 410 | 84, 682 | 91, 454 | 98, 788 |
| Yearling cows..... | 85, 417 | 92, 247 | 99, 626 | 107, 593 | 116, 197 | 125, 490 |
| Pups..... | 307, 491 | 332, 064 | 358, 642 | 387, 320 | 418, 299 | 451, 751 |
| Total..... | 971, 527 | 1, 045, 101 | 1, 127, 082 | 1, 219, 961 | 1, 318, 568 | 1, 430, 418 |

FOXES

Sizable herds of blue foxes are found on both St. Paul and St. George Islands, from which several hundred pelts are obtained each year. The management of these herds is incidental to the sealing industry and requires attention only during the winter, when sealing activities are at a minimum. Beginning about December 1 the foxes are fed specially prepared rations to supplement the natural food supply, which has then become rather scarce. The trapping season also begins early in December, as soon as the skins become prime, and is continued for several weeks.

TRAPPING SEASON OF 1934-35

In the 1934-35 season there were taken 1,002 fox pelts, of which 983 were blue and 19 white. One hundred and eighty-four blue and 16 white pelts were taken on St. Paul Island and 799 blue and 3 white pelts on St. George Island. There were also trapped, marked, and released for breeding stock 11 foxes on St. Paul Island and 266 on St. George Island. The breeding reserve includes likewise a considerable number of animals that were not captured during the season.

REINDEER

St. Paul Island.—During the year ended September 30, 1934, 14 reindeer were killed and used for food. A count of the animals in the herd on October 10 showed a total of 820, of which 147 were the young of the season. The herd was apparently in good condition.

St. George Island.—No reindeer were killed for food during the year ended September 30, 1934. A count made on September 29 showed 72 animals in the herd, of which 8 were the young of the season.

FUR-SEAL SKINS

SHIPMENTS

Seven hundred and twenty-two barrels containing 53,462 fur-seal skins taken on the Pribilof Islands in 1934 were shipped on the U. S. S. *Sirius*, which sailed from the islands on August 8 and reached Seattle,

Wash., on August 19. Eight thousand and twenty-two of these skins, packed in 107 barrels, were delivered to a representative of the Canadian Government at Seattle on August 21, in accordance with provisions of the fur-seal treaty; the remainder were forwarded by freight to the Fouke Fur Co. at St. Louis, Mo., and arrived there on August 29.

Six skins taken in 1934, specially treated for experimental purposes, were turned over to an employee of the Fouke Fur Co. at the Pribilof Islands and were brought out by him on the *Penguin*, sailing from St. Paul Island on July 31.

SALES

Two public auctions of fur-seal skins taken on the Pribilof Islands were held at St. Louis in 1934—on April 30 and October 15, respectively—at which a total of 47,101 skins were sold. During the year, also, 3,652 sealskins taken on the Pribilof Islands were disposed of at special sales. In the following detailed statements the sales of other sealskins by the Department of Commerce for the account of the Government are included in order that the records may be complete.

April 30, 1934.—At this sale 28,101 Pribilof Islands fur-seal skins were sold for \$575,041.25. Of these, 27,656 dressed, dyed, and machined skins brought \$573,935.50, and 445 miscellaneous raw-salted and unhaired and dressed skins brought \$1,105.75. In addition, 170 raw-salted fur-seal skins taken by the Japanese Government on Robben Island in 1933 and allotted to the United States as its share of such skins under provisions of the fur-seal treaty were sold for \$467.50.

October 15, 1934.—Nineteen thousand Pribilof Islands fur-seal skins, dressed, dyed, and machined, were disposed of at this sale for \$336,846.25. These skins consisted of 10,000 dyed black, 3,000 dyed logwood brown, and 6,000 dyed Safari brown. This was the first auction of the new Safari brown, and the average price was higher than that obtained for skins dyed black or logwood brown. There were also sold 3 confiscated sealskins, parchments, for \$7.50, making a total of \$336,853.75 for fur-seal skins at this sale.

Special sales.—The Department authorized a number of special sales of sealskins in 1934 for advertising and promotional purposes, in accordance with which 3,652 skins were sold for \$74,812.05. Of these skins, 1,274 dyed black brought \$24,427.08; 1,009 dyed logwood brown, \$22,064.41; 1,069 dyed Safari brown, \$25,104.56; and 300 raw-salted, \$3,216.

Further details in regard to the sales of sealskins by the Department of Commerce for the account of the Government in 1934 are given in the following tables:

Comparative values, by sizes and grades, with percentages each size, of Pribilof sealskins sold at public auction in 1934

| Classes and sales | Grade | Number | High | Low | Average | Total | Total number | Average price | Total price | Percentage |
|--------------------|---------------------------|--------|---------|---------|---------|------------|--------------|---------------|-------------|------------|
| [DYED BLACK | | | | | | | | | | |
| Extra extra large: | | | | | | | | | | |
| Apr. 30..... | I and II..... | 15 | \$35.00 | \$35.00 | \$35.00 | \$525.00 | 44 | \$26.18 | \$1,152.00 | 0.25 |
| | Scarred, faulty, etc..... | 24 | 25.50 | 20.50 | 23.63 | 567.00 | | | | |
| | III..... | 5 | 12.00 | 12.00 | 12.00 | 60.00 | | | | |
| Extra large: | | | | | | | | | | |
| Apr. 30..... | I and II..... | 111 | 35.00 | 28.00 | 30.41 | 3,375.00 | 245 | 25.60 | 6,272.00 | 1.39 |
| | Scarred, faulty, etc..... | 120 | 25.50 | 20.50 | 22.83 | 2,739.00 | | | | |
| | III..... | 14 | 12.00 | 7.00 | 11.29 | 158.00 | | | | |
| Oct. 15..... | I and II..... | 210 | 24.00 | 21.00 | 22.14 | 4,650.00 | 450 | 19.83 | 8,925.00 | 4.50 |
| | Scarred, faulty, etc..... | 240 | 18.50 | 17.25 | 17.81 | 4,275.00 | | | | |
| Large: | | | | | | | | | | |
| Apr. 30..... | I and II..... | 2,004 | 36.50 | 25.50 | 30.12 | 60,356.00 | 4,341 | 24.18 | 104,977.00 | 24.64 |
| | Scarred, faulty, etc..... | 2,212 | 27.50 | 14.50 | 19.60 | 43,366.00 | | | | |
| | III..... | 125 | 12.00 | 7.00 | 10.04 | 1,255.00 | | | | |
| Oct. 15..... | I and II..... | 1,730 | 23.00 | 20.00 | 21.10 | 36,505.00 | 3,510 | 18.52 | 65,005.00 | 35.10 |
| | Scarred, faulty, etc..... | 1,690 | 18.00 | 15.50 | 16.35 | 27,475.00 | | | | |
| | III..... | 100 | 10.25 | 10.25 | 10.25 | 1,025.00 | | | | |
| Medium: | | | | | | | | | | |
| Apr. 30..... | I and II..... | 5,179 | 28.25 | 18.25 | 20.98 | 108,660.50 | 10,613 | 17.66 | 187,395.25 | 60.24 |
| | Scarred, faulty, etc..... | 5,118 | 19.00 | 12.50 | 14.97 | 76,610.00 | | | | |
| | III..... | 316 | 8.00 | 5.25 | 6.72 | 2,124.75 | | | | |
| Oct. 15..... | I and II..... | 2,620 | 18.50 | 15.50 | 16.35 | 42,830.00 | 5,383 | 14.68 | 79,029.50 | 53.83 |
| | Scarred, faulty, etc..... | 2,610 | 14.00 | 12.50 | 13.39 | 34,955.00 | | | | |
| | III..... | 153 | 9.00 | 6.50 | 8.13 | 1,244.50 | | | | |
| Small medium: | | | | | | | | | | |
| Apr. 30..... | I and II..... | 1,147 | 17.00 | 15.50 | 16.27 | 18,659.50 | 2,374 | 13.82 | 32,801.75 | 13.48 |
| | Scarred, faulty, etc..... | 1,136 | 13.50 | 9.50 | 11.94 | 13,563.00 | | | | |
| | III..... | 91 | 7.75 | 6.00 | 6.37 | 579.25 | | | | |
| Oct. 15..... | I and II..... | 340 | 13.00 | 12.00 | 12.37 | 4,205.00 | 657 | 11.56 | 7,593.00 | 6.57 |
| | Scarred, faulty, etc..... | 270 | 11.75 | 11.00 | 11.42 | 3,082.50 | | | | |
| | III..... | 47 | 6.50 | 6.50 | 6.50 | 305.50 | | | | |
| All classes: | | | | | | | | | | |
| Apr. 30..... | | | | | | | 17,617 | 18.88 | 332,598.00 | 100.00 |
| Oct. 15..... | | | | | | | 10,000 | 16.06 | 160,552.50 | 100.00 |

Comparative values, by sizes and grades, with percentages each size, of Pribilof sealskins sold at public auction in 1934—Continued

| Classes and sales | Grade | Number | High | Low | Average | Total | Total number | Average price | Total price | Percentage |
|---------------------------|-----------------------|--------|---------|---------|---------|-----------|--------------|---------------|-------------|------------|
| DYED LOGWOOD BROWN | | | | | | | | | | |
| Extra extra large: | | | | | | | | | | |
| Apr. 30 | I and II | 8 | \$34.50 | \$34.50 | \$34.50 | \$276.00 | 13 | \$30.85 | \$401.00 | 0.13 |
| | Scarred, faulty, etc. | 5 | 25.00 | 25.00 | 25.00 | 125.00 | | | | |
| Extra large: | | | | | | | | | | |
| Apr. 30 | I and II | 113 | 36.75 | 26.50 | 32.61 | 3,684.50 | 247 | 28.08 | 6,936.50 | 2.46 |
| | Scarred, faulty, etc. | 129 | 27.50 | 22.50 | 24.92 | 3,214.50 | | | | |
| | III | 5 | 7.50 | 7.50 | 7.50 | 37.50 | | | | |
| Oct. 15 | I and II | 26 | 22.50 | 22.50 | 22.50 | 586.00 | 146 | 18.39 | 2,685.00 | 4.87 |
| | Scarred, faulty, etc. | 120 | 19.00 | 16.00 | 17.60 | 2,100.00 | | | | |
| Large: | | | | | | | | | | |
| Apr. 30 | I and II | 1,293 | 36.75 | 26.50 | 31.25 | 40,402.75 | 2,692 | 27.32 | 73,536.25 | 26.82 |
| | Scarred, faulty, etc. | 1,348 | 26.50 | 22.50 | 24.30 | 32,751.00 | | | | |
| | III | 51 | 7.50 | 7.50 | 7.50 | 382.50 | | | | |
| Oct. 15 | I and II | 274 | 23.90 | 20.50 | 21.16 | 5,798.75 | 1,044 | 17.49 | 18,258.75 | 34.80 |
| | Scarred, faulty, etc. | 770 | 17.00 | 15.50 | 16.18 | 12,460.00 | | | | |
| Medium: | | | | | | | | | | |
| Apr. 30 | I and II | 3,076 | 33.75 | 21.50 | 26.20 | 80,589.50 | 5,911 | 23.82 | 140,775.50 | 58.88 |
| | Scarred, faulty, etc. | 2,740 | 23.75 | 20.00 | 21.70 | 59,444.50 | | | | |
| | III | 95 | 8.00 | 7.50 | 7.81 | 741.50 | | | | |
| Oct. 15 | I and II | 571 | 21.00 | 17.50 | 18.77 | 10,719.00 | 1,531 | 16.08 | 24,619.00 | 51.03 |
| | Scarred, faulty, etc. | 960 | 16.25 | 13.50 | 14.48 | 13,900.00 | | | | |
| Small medium: | | | | | | | | | | |
| Apr. 30 | I and II | 697 | 21.50 | 18.00 | 18.50 | 12,897.25 | 1,176 | 16.74 | 19,688.25 | 11.71 |
| | Scarred, faulty, etc. | 444 | 15.25 | 14.00 | 14.66 | 6,511.00 | | | | |
| | III | 35 | 8.00 | 8.00 | 8.00 | 280.00 | | | | |
| Oct. 15 | I and II | 99 | 19.00 | 13.75 | 15.29 | 1,513.50 | 279 | 12.20 | 3,403.50 | 9.30 |
| | Scarred, faulty, etc. | 180 | 11.00 | 10.00 | 10.50 | 1,890.00 | | | | |
| All classes: | | | | | | | | | | |
| Apr. 30 | | | | | | | 10,039 | 24.04 | 241,337.50 | 100.00 |
| Oct. 15 | | | | | | | 3,000 | 16.32 | 48,966.25 | 100.00 |
| DYED SAFARI BROWN | | | | | | | | | | |
| Extra large: | | | | | | | | | | |
| Oct. 15 | I and II | 99 | 34.00 | 25.50 | 31.29 | 3,098.00 | 277 | 25.60 | 7,092.00 | 4.62 |
| | Scarred, faulty, etc. | 178 | 23.00 | 21.00 | 22.44 | 3,994.00 | | | | |
| Large: | | | | | | | | | | |
| Oct. 15 | I and II | 846 | 34.00 | 24.50 | 28.53 | 24,139.50 | 2,138 | 24.39 | 52,151.75 | 35.63 |
| | Scarred, faulty, etc. | 1,292 | 23.50 | 20.00 | 21.68 | 28,012.25 | | | | |

| | | | | | | | | | | | |
|---------------|----------------------------|-------|-------|-------|-------|-----------|---|-------|-------|------------|--------|
| Medium: | | | | | | | | | | | |
| Oct. 15..... | {I and II..... | 1,175 | 25.00 | 16.00 | 23.40 | 27,498.25 | } | 3,115 | 19.77 | 61,568.25 | 51.92 |
| | {Scarred, faulty, etc..... | 1,940 | 19.00 | 15.00 | 17.57 | 34,090.00 | | | | | |
| Small medium: | | | | | | | | | | | |
| Oct. 15..... | {I and II..... | 180 | 21.25 | 16.00 | 17.48 | 3,145.50 | } | 470 | 13.82 | 6,495.50 | 7.83 |
| | {Scarred, faulty, etc..... | 290 | 12.75 | 10.50 | 11.55 | 3,350.00 | | | | | |
| All classes: | | | | | | | | | | | |
| Oct. 15..... | | | | | | | | 6,000 | 21.22 | 127,327.50 | 100.00 |
| | | | | | | | | | | | |
| | MISCELLANEOUS | | | | | | | | | | |
| Apr. 30..... | {Unhaired and dressed..... | 202 | 5.50 | 1.00 | 4.36 | 881.50 | } | 445 | 2.48 | 1,105.75 | 100.00 |
| | {Raw..... | 243 | 1.00 | .50 | .92 | 224.25 | | | | | |

Special sales of Pribilof Islands fur-seal skins in 1934

| Date | Number of skins | Description | Price per skin | Total |
|-----------|-----------------|---|----------------|-------------|
| Jan. 30.. | 3 | Dyed black, small medium..... | \$16. 23 | \$48. 69 |
| | 6 | Dyed logwood brown, medium..... | 22. 99 | 137. 94 |
| June 30.. | 49 | Dyed black, large..... | 30. 12 | 1, 475. 88 |
| | 205 | Dyed black, medium..... | 20. 98 | 4, 300. 90 |
| | 15 | Dyed black, small medium..... | 16. 27 | 244. 05 |
| | 2 | Dyed logwood brown, large..... | 31. 25 | 62. 50 |
| | 28 | Dyed Safari brown, large..... | 31. 25 | 812. 60 |
| Aug. 8.. | 94 | Dyed Safari brown, medium..... | 26. 20 | 2, 462. 80 |
| | 140 | Dyed black, large..... | 25. 53 | 3, 574. 20 |
| | 140 | Dyed black, large, scarred and faulty..... | 19. 20 | 2, 688. 00 |
| | 300 | Dyed black, medium..... | 19. 95 | 5, 985. 00 |
| | 300 | Dyed black, medium, scarred and faulty..... | 14. 42 | 4, 326. 00 |
| | 60 | Dyed black, small medium..... | 16. 23 | 973. 80 |
| | 60 | Dyed black, small medium, scarred and faulty..... | 12. 81 | 768. 60 |
| | 220 | Dyed logwood brown, large..... | 26. 10 | 5, 742. 00 |
| | 140 | Dyed logwood brown, large, scarred and faulty..... | 19. 08 | 2, 667. 00 |
| | 320 | Dyed logwood brown, medium..... | 22. 99 | 7, 358. 80 |
| | 200 | Dyed logwood brown, medium, scarred and faulty..... | 20. 05 | 4, 010. 00 |
| | 60 | Dyed logwood brown, small medium..... | 19. 08 | 1, 144. 80 |
| | 60 | Dyed logwood brown, small medium, scarred and faulty..... | 15. 41 | 924. 60 |
| | 2 | Dyed black, medium..... | 20. 88 | 41. 96 |
| | 25 | Dyed Safari brown, large..... | 31. 25 | 781. 25 |
| | 25 | Dyed Safari brown, medium..... | 26. 20 | 655. 00 |
| Aug. 31.. | 62 | Dyed Safari brown, large..... | 31. 25 | 1, 937. 50 |
| | 62 | Dyed Safari brown, medium..... | 26. 20 | 1, 624. 40 |
| Dec. 31.. | 153 | Dyed Safari brown, large..... | 28. 53 | 4, 365. 09 |
| | 220 | Dyed Safari brown, medium..... | 23. 40 | 5, 148. 00 |
| | 62 | Dyed Safari brown, small medium..... | 17. 47 | 1, 083. 14 |
| | 106 | Dyed Safari brown, large, scarred and faulty..... | 21. 68 | 2, 298. 08 |
| | 205 | Dyed Safari brown, medium, scarred and faulty..... | 17. 57 | 3, 601. 85 |
| | 29 | Dyed Safari brown, small medium, scarred and faulty..... | 11. 55 | 334. 95 |
| | 1 | Dyed logwood brown, medium..... | 18. 77 | 18. 77 |
| | 300 | Raw..... | 10. 72 | 3, 216. 00 |
| | 3, 652 | | | 74, 812. 05 |

Sale at St. Louis, Mo., April 30, 1934, of 170 fur-seal skins received from Japanese Government under treaty provisions

| Number of skins | Trade classification | Price per skin | Total |
|-----------------|----------------------|----------------|-----------|
| 170..... | Raw salted..... | \$2. 75 | \$467. 50 |

DISPOSITION OF FUR-SEAL SKINS TAKEN AT PRIBILOF ISLANDS

On January 1, 1934, there were on hand 73,420 fur-seal skins taken on the Pribilof Islands. Of these, 73,388 were at St. Louis, Mo., and 32 at Washington. The foregoing figures include two skins which were unaccounted for at the close of 1933 but which were subsequently found when the shipment for that year was unpacked at St. Louis.

In 1934 there were taken on the Pribilof Islands 53,468 fur-seal skins, of which 8,022 were allotted to the Government of the Dominion of Canada. The latter represent the Canadian share, or 15 percent, of the season's take, in accordance with treaty provisions, and 2 additional skins to offset a shortage in the previous year's shipment.

Sales of Pribilof Islands sealskins in 1934 totaled 50,753 skins, consisting of 50,553 that were on hand on January 1 and 200 that were taken during the year, leaving 68,113 on hand on December 31, 1934. The following tables show further details in regard to fur-seal skins taken on the Pribilof Islands, as well as details in regard to other Government-owned fur-seal skins under the control of the Department of Commerce:

Summary of Government-owned fur-seal skins in the custody of Fouke Fur Co., at St. Louis, Mo., calendar year 1934

| Source | On hand Jan. 1 | Receipts in 1934 | Sales in 1934 | On hand Dec. 31 |
|---|---------------------|------------------|---------------|-----------------|
| Taken on Pribilof Islands: | | | | |
| Calendar year 1931..... | 301 | ----- | 301 | ----- |
| Calendar year 1932..... | 26,718 | ----- | 26,718 | ----- |
| Calendar year 1933..... | ¹ 46,369 | ----- | 23,534 | 22,835 |
| Calendar year 1934..... | ----- | 45,446 | 200 | 45,246 |
| United States, share of Japanese fur-seal skins: | | | | |
| Season of 1933..... | 170 | ----- | 170 | ----- |
| Season of 1934..... | ----- | 200 | ----- | 200 |
| Confiscated fur-seal skins..... | | 3 | 3 | |
| Total..... | 73,558 | 45,649 | 50,926 | 68,281 |

¹ Includes 2 skins unaccounted for at close of 1933 which were subsequently found when shipment was unpacked at St. Louis.

Summary of all Government-owned fur-seal skins under control of Department of Commerce, calendar year 1934

| Source | On hand Jan. 1 | | | Re-ceipts in 1934 | Disposed of in 1934 | | On hand Dec. 31 | | |
|---|---------------------|--------------------|---------------|-------------------|---------------------|-----------------------|-----------------|--------------------|---------------|
| | Fouke Fur Co. | Wash-ington office | Total | | Sales | Deliv-ered to Can-ada | Fouke Fur Co. | Wash-ington office | Total |
| Taken on Pribilof Islands: | | | | | | | | | |
| Calendar year 1918, held for reference purposes..... | | 7 | 7 | ----- | ----- | ----- | 7 | 7 | |
| Calendar year 1923..... | | 3 | 3 | ----- | ----- | ----- | 3 | 3 | |
| Calendar year 1924..... | | 1 | 1 | ----- | ----- | ----- | 1 | 1 | |
| Calendar year 1929..... | | 5 | 5 | ----- | ----- | ----- | 5 | 5 | |
| Calendar year 1930..... | | 2 | 2 | ----- | ----- | ----- | 2 | 2 | |
| Calendar year 1931..... | 301 | ¹ 14 | 315 | ----- | 301 | ----- | 14 | ¹ 14 | |
| Calendar year 1932..... | 26,718 | ----- | 26,718 | ----- | 26,718 | ----- | ----- | ----- | |
| Calendar year 1933..... | ¹ 46,369 | ----- | 46,369 | ----- | 23,534 | ----- | 22,835 | 22,835 | |
| Calendar year 1934..... | ----- | ----- | ----- | 53,468 | 200 | 8,022 | 45,246 | 45,246 | |
| Miscellaneous skins held for reference purposes..... | | 4 | 4 | ----- | ----- | ----- | 4 | 4 | |
| United States share of Japanese sealskins: | | | | | | | | | |
| Season of 1933..... | 170 | ----- | 170 | ----- | 170 | ----- | ----- | ----- | |
| Season of 1934..... | ----- | ----- | ----- | 200 | ----- | ----- | 200 | 200 | |
| Confiscated skins..... | | | | 3 | 3 | ----- | ----- | ----- | |
| Total..... | 73,558 | 36 | 73,594 | 53,671 | 50,926 | 8,022 | 68,281 | 36 | 68,317 |

¹ Skins made up into coats for display purposes.

² Includes 2 skins unaccounted for at close of 1933 which were subsequently found when shipment was unpacked at St. Louis.

SHIPMENT AND SALE OF FOX SKINS

The 214 blue and 23 white fox skins taken on St. Paul Island and the 700 blue and 2 white fox skins taken on St. George Island in the season of 1933-34 were shipped from the islands on the *Penguin*, sailing on May 17. The vessel reached Seattle on May 28, and the fox skins were forwarded promptly by express to the Department's selling agents at St. Louis, Mo.

At the public auction sale in St. Louis on April 30, 1934, there were sold 559 blue fox skins that were taken on the Pribilof Islands in the 1932-33 season. These skins brought \$19,967.50, an average of \$35.72 per skin. The maximum price was \$75, obtained for a pelt sold singly.

On October 15, 1934, there were sold at public auction at St. Louis 457 blue and 25 white fox skins taken on the Pribilof Islands in the 1933-34 season. The blue pelts brought \$9,345, an average of \$20.45 each; and the white pelts brought \$333, an average of \$13.32 each. The maximum price was \$69, obtained for a single blue pelt.

SEA-OTTER SKIN

The skin of a sea-otter pup, said to have been found dead on the beach of Caton Island, Alaska, in June 1934, was surrendered to the Bureau. This skin was sold for Government account at public auction at St. Louis, Mo., on October 15, 1934, for \$3.

FUR-SEAL PATROL

UNITED STATES COAST GUARD

Seven Coast Guard cutters and two 125-foot patrol boats were assigned by the Secretary of the Treasury to the patrol for the protection of fur seals of the North Pacific which have their breeding grounds at the Pribilof Islands.

Beginning April 5 the *Redwing* patrolled from the southern boundary of Washington to Dixon Entrance until the fur-seal herd had passed. The *Tallapoosa* covered the waters from Dixon Entrance to Kodiak Island in the last half of April and from Kodiak Island to Unimak Pass during the first half of May. The *Haida* sailed from Seattle on April 16 and the patrol boats *Bonham* and *Ewing* on April 18 for Unalaska, the base for the patrol boats during the remainder of the season and for the *Haida* until July 17, when it was relieved by the *Chelan*. The cutters *Tahoe* and *Shoshone* likewise took turns in the patrol from Unalaska as base, the former having sailed from San Francisco on April 10 and the latter on June 20. The *Northland* left Oakland on May 5 for its usual cruise to the Arctic Ocean and patrolled in Bering Sea and adjacent waters en route. The patrol by Coast Guard vessels extended as far westward as Attu, the westernmost of the Aleutian chain, and was carried on in each locality as long as circumstances required.

BUREAU OF FISHERIES

On April 4 the *Eider* sailed from Seattle for Neah Bay, Wash., to patrol waters in that vicinity during the northward migration of the fur-seal herd. The vessel remained on this patrol duty for about 3 weeks, and after its return to Seattle the work was continued by the *Brant* from April 28 to May 18.

A representative of the Bureau was again stationed at La Push to secure compliance with the regulations prohibiting the use of firearms and motor boats in the taking of fur-seal skins by the Indians there. Very little sealing was done during the season, which was attributed not so much to poor weather conditions as to the low prices for skins. An additional factor was the employment of needy Indians on the C. W. A. road-building project, which proved a more attractive and profitable occupation than the uncertain sealing activities.

SEALING PRIVILEGES ACCORDED ABORIGINES

Under the provisions of the North Pacific Sealing Convention of July 7, 1911, Indians and other aborigines dwelling on the coasts of the waters designated by the convention may take fur-seal skins under limited conditions. In 1934 there were taken and duly authenticated by officials of the respective Governments 290 fur-seal skins, of which 34 were taken by Indians under the jurisdiction of the United States and 256 by Indians of Canada. Reports have also been received of the authentication of 6 additional skins, 3 from male and 3 from female seals, taken in 1933 by Indians of the State of Washington. These, together with the number previously reported, make a total of 98 skins taken in that year by Indians under the jurisdiction of the United States. The details for 1934 are as follows:

Washington.—Twenty-three sealskins taken by Indians of Washington were authenticated. Of these, 6 were from male and 17 from female seals. The skins were taken by Indians of La Push and Neah Bay in the months from February to May, inclusive, and were authenticated by Paul E. Thompson, special agent of the Bureau, and by N. O. Nicholson, superintendent of the Taholah Indian Agency, Hoquiam, Wash.

Alaska.—Eleven sealskins taken by natives of Sitka were authenticated by Warden Donald S. Haley. Of these skins, 3 were from male seals and 8 from females. The seals were taken in the waters off Biorka Island in the months of April and May.

British Columbia.—An official report received by the Bureau stated that 256 fur-seal skins were taken by Indians along the British Columbia coast in 1934.

JAPANESE SEALSKINS DELIVERED TO THE UNITED STATES

The treaty of July 7, 1911, for the protection of fur seals of the North Pacific Ocean provides that the United States shall receive 10 percent of the fur-seal skins taken annually from the Japanese herd. In accordance with that provision 200 sealskins were allotted to the United States as its share of the take on Robben Island in 1934. They were received by the Department's selling agents at St. Louis, Mo., on December 18, 1934.

COMPUTATION OF FUR SEALS, PRIBILOF ISLANDS, 1934

By HARRY J. CHRISTOFFERS

In order to determine whether sufficient male seals are being reserved for breeding purposes, a census of harem and idle bulls is taken annually at the Pribilof Islands. Using this census as a basis, and estimating the number of cows at the islands by applying the rate of increase ascertained by actual counts of pups through a series of years, the average harem is determined.

The size of the average harem should be suitable to provide for a maximum increase in the growth of the herd. When the average is maintained between 40 and 45 it is safe to assume that sufficient animals have been reserved for breeding purposes. If the average is much less than 40 there is a question whether the number of males

reserved may not be larger than necessary. With an average harem of about 40 it seems certain that all females arriving during the rookery breeding period are being properly served.

The virgin females arriving after the rookery breeding period is practically over are for the most part taken care of near the end of the season by the surplus and young bulls that have not held harems or were in the idle bull class. These bulls are strong and virile, while the older bulls which have been holding harems throughout the season are ready to retire before the virgin cows arrive. It is necessary, therefore, that personal observations be made in order to determine whether sufficient surplus bulls have been reserved to take care of the virgin females.

When there is an average harem of less than 40 females, and at the same time a large number of idle and surplus bulls, the breeding bulls do not retain their positions throughout the entire season. The stronger members of the idle and surplus bull class replace the harem bulls as soon as the latter begin to weaken. This does not necessarily mean that the replaced harem bull is not still able to procreate. If still strong and virile, he usually does not retire any farther than the hauling grounds or the rear of the rookery, where he is able to secure transient females.

In 1934, a total of 51,935 animals classed as 3-year-olds were killed, as compared with 52,747 in 1933. This decrease in the total number of 3-year-old males secured was undoubtedly due to the continual stormy weather, accompanied by heavy driving rains, which prevailed throughout the season.

The captain of the *Penguin* reported that large numbers of seals were in Ikatan Bay about July 20, and the captain of a Coast Guard cutter gave a similar report with respect to Akutan Bay. When conditions are favorable at the islands, seals do not usually remain for any considerable time around the Aleutians. The 3-year-old males have no particular reason for going to the Pribilof Islands, as they are not of breeding age. When they do arrive, they like to haul out and sleep and play. Warm foggy weather is favorable for the hauling out of bachelors, while driving rain storms cause them to return to the water.

If weather conditions had been favorable, a shortage of 3-year-old males would have indicated an abnormal death rate at sea. As stormy weather prevailed throughout the entire season, however, it was impossible to determine whether there was an actual shortage of seals. It may be assumed that there was a normal increase in the number of killable seals, but because of adverse weather they did not arrive at the islands.

An annual increase of approximately 8 percent in the past does not necessarily signify that this rate will continue as the herd increases in size. In some years there may be more than the average number of natural enemies of the seals at sea. If the natural enemies increase at the same rate as the herd, the death rate at sea would remain about constant. If the natural enemies increase at a lower rate, then it would soon be possible to have abnormally increased killings in a year or a series of years. It appears that a series of years in which there is a large increase may be followed by a series of years when the growth of the herd is at a standstill, gradually preparing for another period of forging ahead.

BULLS

A census of harem and idle bulls was taken at the usual time. Where large areas or unfavorable conditions made accurate counts impossible, a conservative estimate was made at the time the remainder of the rookery was counted.

Sivutch rookery, on Sea Lion Rock, was not counted. As the small area of Sea Lion Rock is now densely populated with seals during the breeding season, this rookery is hardly capable of continued expansion.

Considerably more bulls of the idle and surplus class were on the rookeries and hauling grounds during the last few days of July than there were about 2 weeks later when the census was taken. This may have been because unfavorable weather in the latter period caused the mature bulls without cows to remain in the water.

A number of bulls branded with a hot iron in 1923, as 3-year-olds, were observed holding harems on various rookeries. These bulls, though 14 years of age, were still active and robust.

Number of harem and idle bulls, approximate ratio of idle bulls to harem bulls, and average harem, 1934

| Rookery | Date | Harem bulls | Idle bulls | Total | Approximate ratio of idle bulls to harem bulls | Average harem |
|----------------------------|---------|-------------|------------|--------|--|---------------|
| St. Paul Island: | | | | | | |
| Kitovi..... | July 17 | 389 | 86 | 475 | 1:5 | 38.42 |
| Lukanin..... | do | 177 | 21 | 198 | 1:8 | 40.82 |
| Gorbach..... | July 18 | 793 | 149 | 942 | 1:5 | 47.85 |
| Ardiguen..... | do | 89 | 15 | 104 | 1:6 | 40.33 |
| Reef..... | do | 1,493 | 337 | 1,830 | 1:4 | 51.05 |
| Sivutch (estimated)..... | do | 400 | 85 | 485 | 1:5 | 58.30 |
| Lagoon (actual count)..... | July 17 | 3 | 1 | 4 | 1:3 | 37.33 |
| Tolstol..... | do | 997 | 214 | 1,211 | 1:5 | 45.00 |
| Zapadni..... | July 19 | 866 | 201 | 1,067 | 1:4 | 49.78 |
| Little Zapadni..... | do | 478 | 71 | 549 | 1:7 | 45.98 |
| Zapadni Reef..... | do | 62 | 12 | 74 | 1:5 | 12.29 |
| Polovina..... | July 20 | 372 | 98 | 470 | 1:4 | 41.65 |
| Polovina Cliffs..... | do | 317 | 55 | 372 | 1:6 | 26.76 |
| Little Polovina..... | do | 103 | 37 | 140 | 1:3 | 30.00 |
| Morjovi..... | July 21 | 309 | 89 | 378 | 1:4 | 17.97 |
| Vostochni..... | do | 1,993 | 409 | 2,402 | 1:5 | 31.02 |
| Total..... | | 8,841 | 1,860 | 10,701 | 1:5 | 41.68 |
| St. George Island: | | | | | | |
| North..... | July 23 | 691 | 164 | 855 | 1:4 | 43.51 |
| Staraya Artil..... | do | 478 | 92 | 570 | 1:5 | 47.40 |
| Zapadni..... | July 22 | 171 | 55 | 226 | 1:3 | 18.70 |
| South..... | do | 123 | 6 | 129 | 1:21 | 6.10 |
| East Reef..... | do | 157 | 33 | 190 | 1:5 | 43.88 |
| East Cliffs..... | do | 309 | 72 | 381 | 1:4 | 63.66 |
| Total..... | | 1,929 | 422 | 2,351 | 1:5 | 43.15 |
| Total (both islands)..... | | 10,770 | 2,282 | 13,052 | 1:5 | 41.95 |

AVERAGE HAREM

The maintaining of a proper average harem is necessary to provide for a maximum increase in growth of the herd. The average harem should be maintained at about 40. A slight variation from this figure would be of no consequence. The average harem for the two islands was 41.95 in 1934, as compared with 40.96 in 1933.

The number of cows, estimated by applying an increase of 8 percent to the number computed for the previous season, and the number of harem bulls, determined by actual count, form the basis for calculating the number of animals in the average harem.

The present average harem should give very satisfactory results. Sufficient harem and idle bulls were present to take care of all females arriving. In addition, there were at the end of the season large numbers of robust mature surplus bulls on the hauling grounds. Besides taking care of virgin females, these mature surplus bulls act as a breeding reserve, for they are at all times prepared to take up harem bull positions, if necessary, and are able to replace harem bulls which have weakened to such an extent that they are unable to hold their cows.

Computation of breeding cows, based on annual increase of 8 percent, and of average harem, in 1934

| Rookery | Breeding cows | | Harem bulls, 1934 | Average harem | | |
|---------------------------------|---------------|---------|-------------------|---------------|-------|--|
| | 1933 | 1934 | | 1934 | 1933 | Increase (+) or decrease (-) in 1934 from 1933 |
| St. Paul Island: | | | | | | |
| Kitovi..... | 13,837 | 14,944 | 389 | 38.42 | 36.80 | +1.62 |
| Lukanin..... | 6,690 | 7,225 | 177 | 40.82 | 45.20 | -4.38 |
| Gorbach..... | 35,132 | 37,943 | 793 | 47.85 | 47.22 | +0.63 |
| Ardiguen..... | 3,323 | 3,589 | 89 | 40.33 | 42.06 | -1.73 |
| Reef..... | 70,568 | 76,213 | 1,493 | 51.05 | 51.25 | -0.20 |
| Sivutch..... | 21,594 | 23,322 | 400 | 58.30 | 53.99 | +4.31 |
| Lagoon (actual count pups)..... | 114 | 112 | 3 | 37.33 | 22.80 | +14.53 |
| Tolstol..... | 41,542 | 44,865 | 997 | 45.00 | 43.68 | +1.32 |
| Zapadni..... | 39,923 | 43,117 | 886 | 49.78 | 50.34 | -0.56 |
| Little Zapadni..... | 20,360 | 21,978 | 478 | 45.98 | 44.92 | +1.06 |
| Zapadni Reef..... | 706 | 762 | 62 | 12.29 | 16.81 | -4.52 |
| Polovina..... | 14,847 | 16,495 | 372 | 41.65 | 43.61 | -1.96 |
| Polovina Cliffs..... | 7,856 | 8,484 | 317 | 26.76 | 28.16 | -1.40 |
| Little Polovina..... | 2,801 | 3,060 | 103 | 30.00 | 23.28 | +6.74 |
| Morjovi..... | 5,143 | 5,554 | 309 | 17.97 | 16.97 | +1.00 |
| Vostochni..... | 57,246 | 61,826 | 1,993 | 31.02 | 29.63 | +1.39 |
| Total..... | 341,232 | 368,519 | 8,841 | 41.68 | 40.94 | +0.74 |
| St. George Island: | | | | | | |
| North..... | 27,841 | 30,068 | 691 | 43.51 | 40.76 | +2.75 |
| Staraya Artil..... | 20,978 | 22,656 | 478 | 47.40 | 44.92 | +2.48 |
| Zapadni..... | 2,960 | 3,197 | 171 | 18.70 | 18.59 | +0.11 |
| South..... | 664 | 750 | 123 | 0.10 | 5.74 | +5.64 |
| East Reef..... | 6,380 | 6,800 | 157 | 43.88 | 41.16 | +2.72 |
| East Cliffs..... | 18,214 | 19,671 | 309 | 63.66 | 62.38 | +1.28 |
| Total..... | 77,067 | 83,232 | 1,929 | 43.15 | 41.01 | +2.14 |
| Total (both islands)..... | 418,299 | 451,751 | 10,770 | 41.95 | 40.96 | +0.99 |

PUPS AND COWS

The estimated number of pups and cows was determined, as heretofore, by applying an increase of 8 percent to the numbers computed for the previous season.

The percentage of dead pups determined by actual count in 1922 was applied to show the number dead on each rookery. For comparative purposes the dead pups are included in the total number of pups. It is possible that the percentage of dead pups may increase slightly

as the rookeries increase in size, especially on flat areas where there are no rocks to afford protection for the young until they are able to find their way to the edge of the breeding area. Inasmuch as the season of 1934 was exceptionally rainy and cold, it is believed that the death rate of pups was considerably larger than usual. For some time after birth the pups appear to suffer severely when it is wet and cold.

For the first time, two stillbirths were noted on Polovina rookery during the same day. Both pups were fully matured and appeared perfect in every way. The mothers exerted every effort to start life by shaking and biting the pups, but failed to do so.

Distribution of pups on the Pribilof Islands, Aug. 10, 1934, and comparison with distribution in 1933

| Rookery | 1934 | | | | 1933 | 1934 |
|----------------------------------|----------------|--------------|----------------|-------------------|----------------|---------------|
| | Living pups | Dead pups | Total pups | Percent dead pups | Total pups | Increase |
| St. Paul Island: | | | | | | |
| Kitovi..... | 14,724 | 220 | 14,944 | 1.47 | 13,837 | 1,107 |
| Lukanin..... | 7,068 | 157 | 7,225 | 2.17 | 6,690 | 535 |
| Gorbateh..... | 37,617 | 326 | 37,943 | .86 | 35,132 | 2,811 |
| Ardiguen..... | 3,503 | 86 | 3,589 | 2.39 | 3,323 | 266 |
| Reef..... | 75,100 | 1,113 | 76,213 | 1.46 | 70,668 | 5,545 |
| Sivuteh..... | 22,753 | 569 | 23,322 | 2.44 | 21,594 | 1,728 |
| Lagoon (actual count)..... | | 112 | 112 | | 114 | -2 |
| Tolstol..... | 44,241 | 624 | 44,865 | 1.39 | 41,542 | 3,323 |
| Zapadni..... | 42,375 | 742 | 43,117 | 1.72 | 39,923 | 3,194 |
| Little Zapadni..... | 21,429 | 549 | 21,978 | 2.50 | 20,350 | 1,628 |
| Zapadni Reef..... | 756 | 6 | 762 | .80 | 706 | 56 |
| Polovina..... | 15,258 | 237 | 15,495 | 1.53 | 14,347 | 1,148 |
| Polovina Cliffs..... | 8,327 | 157 | 8,484 | 1.85 | 7,856 | 628 |
| Little Polovina..... | 3,012 | 78 | 3,090 | 2.51 | 2,861 | 229 |
| Morjovi..... | 5,442 | 112 | 5,554 | 2.02 | 5,143 | 411 |
| Vostochni..... | 60,540 | 1,286 | 61,826 | 2.08 | 57,246 | 4,580 |
| Total..... | 362,257 | 6,262 | 368,519 | 1.70 | 341,232 | 27,287 |
| St. George Island: | | | | | | |
| North..... | 29,647 | 421 | 30,068 | 1.40 | 27,841 | 2,227 |
| Staraya Artil..... | 22,071 | 585 | 22,656 | 2.58 | 20,978 | 1,678 |
| Zapadni..... | 3,161 | 36 | 3,197 | 1.12 | 2,960 | 237 |
| South..... | 737 | 13 | 750 | 1.72 | 694 | 56 |
| East Reef..... | 6,786 | 104 | 6,890 | 1.51 | 6,380 | 510 |
| East Cliffs..... | 19,378 | 293 | 19,671 | 1.49 | 18,214 | 1,457 |
| Total..... | 81,780 | 1,452 | 83,232 | 1.74 | 77,067 | 6,165 |
| Total (both Islands)..... | 444,037 | 7,714 | 451,751 | 1.71 | 418,299 | 33,452 |

MORTALITY OF SEALS AT SEA

Inasmuch as there was no information at hand to indicate that a change was required, the same mortality rates used for computing the number of animals in the herd in 1933 were applied in this year's computation. The fact that there was no increase over the previous year in the number of 3-year-old males arriving during the killing season does not indicate that there has been a change in the average growth of the herd or of any particular class of animals. A decrease might occur for several years and then be followed by a sufficient increase to provide for the estimated average increase as computed from past experience.

COMPLETE COMPUTATION

The following summary shows the methods used for computing the number of animals in the Pribilof Islands fur-seal herd in 1934. The total number of seals of all classes is 1,430,418, or 111,850 more than were in the herd in 1933. This is an increase of 8.48 percent over the preceding year.

Complete computation of fur seals, Pribilof Islands, as of Aug. 10, 1934

| Class | St. Paul Island | St. George Island | Total |
|--|-----------------|-------------------|---------|
| Pups, estimated..... | 368,519 | 83,232 | 451,751 |
| Breeding cows, 3 years old and over, by inference..... | 368,519 | 83,232 | 451,751 |
| Harem bulls, counted..... | 8,841 | 1,929 | 10,770 |
| Idle bulls, counted..... | 1,860 | 422 | 2,282 |
| Yearlings, male and female, estimated: | | | |
| Females born in 1933..... | 170,616 | 38,534 | 209,150 |
| Natural mortality, 40 percent..... | 68,246 | 15,414 | 83,660 |
| Yearling females, Aug. 10, 1934..... | 102,370 | 23,120 | 125,490 |
| Males born in 1933..... | 170,616 | 38,533 | 209,149 |
| Natural mortality, 40 percent..... | 68,246 | 15,413 | 83,659 |
| Yearling males, Aug. 10, 1934..... | 102,370 | 23,120 | 125,490 |
| 2-year-olds, male and female, estimated: | | | |
| Yearling females, Aug. 10, 1933..... | 94,789 | 21,408 | 116,197 |
| Natural mortality, 15 percent..... | 14,218 | 3,211 | 17,429 |
| 2-year-old females, Aug. 10, 1934..... | 80,571 | 18,197 | 98,768 |
| Yearling males, Aug. 10, 1933..... | 94,788 | 21,407 | 116,195 |
| Natural mortality, 17.5 percent..... | 16,588 | 3,746 | 20,334 |
| 2-year-old males beginning 1934..... | 78,200 | 17,661 | 95,861 |
| 2-year-old males killed in 1934..... | 777 | 164 | 941 |
| 2-year-old males, Aug. 10, 1934..... | 77,423 | 17,497 | 94,920 |
| 3-year-old males, estimated: | | | |
| 2-year-old males, Aug. 10, 1933..... | 71,498 | 16,164 | 87,662 |
| Natural mortality, 12.5 percent..... | 8,937 | 2,020 | 10,957 |
| 3-year-old males beginning 1934..... | 62,561 | 14,144 | 76,705 |
| 3-year-old males killed in 1934..... | 41,818 | 10,117 | 51,935 |
| 3-year-old males, Aug. 10, 1934..... | 20,743 | 4,027 | 24,770 |
| 4-year-old males, estimated: | | | |
| 3-year-old males, Aug. 10, 1933..... | 14,695 | 3,521 | 18,216 |
| Natural mortality, 10 percent..... | 1,470 | 352 | 1,822 |
| 4-year-old males beginning 1934..... | 13,225 | 3,169 | 16,394 |
| 4-year-old males killed in 1934..... | 330 | 202 | 532 |
| 4-year-old males, Aug. 10, 1934..... | 12,895 | 2,967 | 15,862 |
| 5-year-old males, estimated: | | | |
| 4-year-old males, Aug. 10, 1933..... | 13,399 | 2,042 | 15,441 |
| Natural mortality, 10 percent..... | 1,340 | 204 | 1,544 |
| 5-year-old males, Aug. 10, 1934..... | 12,059 | 1,838 | 13,897 |
| 6-year-old males, estimated: | | | |
| 5-year-old males, Aug. 10, 1933..... | 8,616 | 1,600 | 10,216 |
| Natural mortality, 20 percent..... | 1,723 | 320 | 2,043 |
| 6-year-old males, Aug. 10, 1934..... | 6,893 | 1,280 | 8,173 |

Complete computation of fur seals, Pribilof Islands, as of Aug. 10, 1934—Continued

| Class | St. Paul Island | St. George Island | Total |
|---|-----------------|-------------------|--------|
| Surplus bulls, 7 years old and over, estimated: | | | |
| 6-year-old males, Aug. 10, 1933..... | 7,856 | 1,479 | 9,335 |
| Natural mortality, 20 percent..... | 1,571 | 296 | 1,867 |
| 7-year-old males, Aug. 10, 1934..... | 6,285 | 1,183 | 7,468 |
| Surplus bulls, Aug. 10, 1933..... | (1) | (1) | 4,700 |
| Natural mortality, 30 percent..... | | | 1,410 |
| Remaining surplus for 1934..... | | | 3,290 |
| Breeding bulls of 1933..... | 10,287 | 2,287 | 12,554 |
| Natural mortality, 30 percent..... | 3,080 | 686 | 3,766 |
| 1933 bulls remaining in 1934..... | 7,187 | 1,601 | 8,788 |
| Breeding bulls of 1934..... | 10,701 | 2,351 | 13,052 |
| 1933 bulls remaining, deducted..... | 7,187 | 1,601 | 8,788 |
| Increment of new bulls in 1934..... | 3,514 | 750 | 4,264 |
| 7-year-old males computed for 1934..... | 6,285 | 1,183 | 7,468 |
| Surplus bulls computed for 1934..... | | | 3,290 |
| Total theoretical bull stock for 1934..... | | | 10,758 |
| New increment of breeding bulls deducted..... | | | 4,264 |
| Surplus bulls, Aug. 10, 1934..... | | | 6,494 |

¹ Estimates have been worked out, insofar as possible, to show the approximate number of seals of each class which should be credited to each island. Seals do not, however, haul out in accordance with figures given. Seals born on either island frequent the other island. They travel promiscuously between and haul out on either of the 2 islands. The total for both islands, however, is approximately correct.

RECAPITULATION

| Class | Total | Class | Total |
|-------------------------|---------|-------------------------------|-----------|
| Pups..... | 451,751 | 5-year-old males..... | 13,897 |
| Cows..... | 451,751 | 6-year-old males..... | 8,173 |
| Harem bulls..... | 10,770 | Surplus bulls..... | 6,494 |
| Idle bulls..... | 2,282 | | |
| Yearling females..... | 125,490 | Total, 1934..... | 1,430,418 |
| Yearling males..... | 125,490 | | |
| 2-year-old females..... | 98,768 | Total, 1933..... | 1,318,568 |
| 2-year-old males..... | 94,920 | Numerical increase, 1934..... | 111,850 |
| 3-year-old males..... | 24,770 | Percent increase, 1934..... | 8.48 |
| 4-year-old males..... | 15,862 | | |



FISHERY INDUSTRIES OF THE UNITED STATES, 1934 ¹

By R. H. FIEDLER, *Chief, Division of Fishery Industries*

CONTENTS

| | Page |
|--|------|
| Foreword..... | 76 |
| Part 1.—Operations of the Division | |
| Cooperation with other Federal agencies..... | 76 |
| Cooperation with State agencies..... | 77 |
| Exhibit at "A Century of Progress"..... | 78 |
| Marketing investigations..... | 78 |
| Market survey in southeastern States..... | 79 |
| Per capita consumption of fish..... | 80 |
| Statistical investigations..... | 80 |
| Biological aspect..... | 80 |
| Economic aspect..... | 81 |
| Surveys conducted..... | 81 |
| Technological investigations..... | 82 |
| Laboratories..... | 82 |
| Preservation of fishery products for food..... | 83 |
| Handling fresh mackerel..... | 83 |
| Storage of frozen mackerel..... | 83 |
| Smoked mackerel and storage of this product..... | 84 |
| Shrinkage of fresh fish packed in ice..... | 84 |
| Determinations for the relative freshness of fish..... | 85 |
| Canning aquatic foods..... | 85 |
| Bacteriological investigations..... | 86 |
| Disinfectants for sponges..... | 86 |
| Preservation of fishery byproducts..... | 87 |
| Technological investigations—Con..... | 87 |
| Preservation of fishery byproducts—Con..... | 87 |
| Utilization of salmon cannery waste..... | 88 |
| Effect of manufacture on the quality of nonoily fish meals..... | 89 |
| Studies of the drying of nonoily fish waste..... | 89 |
| Utilization of swordfish livers..... | 89 |
| Haddock-liver oil..... | 89 |
| Nutritive value of aquatic products..... | 90 |
| Swordfish-liver oil..... | 90 |
| Salmon oil..... | 90 |
| Horse mackerel-liver oil..... | 91 |
| Crab meat..... | 91 |
| Sodium alginate..... | 91 |
| Conch meat..... | 91 |
| Mineral constituents of fishery products and byproducts..... | 92 |
| Fish cookery investigations..... | 93 |
| Research associates and student assistants..... | 93 |
| Educational and consulting services..... | 94 |
| Publications of the division..... | 94 |
| Documents, reports, and circulars..... | 95 |
| Special articles and addresses..... | 95 |
| Statistical bulletins..... | 95 |
| Part 2.—Fishery statistics, 1933 | |
| General review..... | 97 |
| Manufactured fishery products..... | 111 |
| Canned fishery products and byproducts trade..... | 115 |
| Frozen-fish trade..... | 123 |
| Fish frozen..... | 123 |
| Holdings..... | 126 |
| Cold-storage holdings of cured fish..... | 128 |
| Foreign fishery trade..... | 128 |
| Fisheries of the New England States..... | 132 |
| Maine..... | 141 |
| New Hampshire..... | 147 |
| Massachusetts..... | 147 |
| Rhode Island..... | 155 |
| Connecticut..... | 160 |
| Vessel fisheries at principal New England ports..... | 165 |
| Economic aspect..... | 165 |
| Biological aspect..... | 173 |
| Mackerel fishery of the Atlantic coast..... | 184 |
| Fisheries of the Middle Atlantic States..... | 186 |
| New York..... | 194 |
| New Jersey..... | 202 |
| Pennsylvania..... | 213 |
| Delaware..... | 214 |
| Vessel fisheries at New York City..... | 218 |
| Shad fishery of the Hudson River..... | 219 |
| Fisheries of the Chesapeake Bay States..... | 219 |
| Maryland..... | 224 |
| Virginia..... | 230 |
| Winter trawl fishery off New Jersey, Maryland, Virginia, and North Carolina..... | 241 |
| Fisheries of the Chesapeake Bay States—Continued..... | 242 |
| Shad and alewife fisheries of the Potomac River..... | 242 |
| Trade in fishery products in Washington, D. C..... | 243 |
| Fisheries of the South Atlantic and Gulf States..... | 245 |
| Sponges sold at the exchange, Tarpon Springs, Fla..... | 254 |
| Fisheries of the Pacific Coast States..... | 254 |
| Washington..... | 261 |
| Oregon..... | 265 |
| California..... | 269 |
| Halibut fishery of the Pacific coast..... | 284 |
| Vessel fisheries at Seattle, Wash..... | 285 |
| Lake fisheries..... | 289 |
| Fisheries of the Mississippi River and tributaries..... | 294 |
| Lake Pepin..... | 299 |
| Lake Keokuk..... | 301 |
| Mississippi River between Lake Pepin and Lake Keokuk..... | 302 |
| Fisheries of Alaska..... | 304 |
| Statistical survey procedure..... | 311 |
| Sectional surveys..... | 311 |
| Local and special surveys..... | 317 |
| Practices and terms..... | 320 |
| Conversion factors..... | 321 |
| Common and scientific names of fishery products..... | 323 |

¹ Appendix II to the Report of the U. S. Commissioner of Fisheries, 1935. Approved for publication May 14, 1935.

FOREWORD

This report constitutes a summary of the activities of the Division of Fishery Industries as well as an annual review on fishery statistics of the United States. As its name indicates, this division of the Bureau is concerned with the activities and welfare of the commercial fishery and fishery industries, the trade in fishery products, and the sh canning and preserving industries. Its functions include the ollection and publication of fishery statistics, the conducting of arket surveys, the prosecution of research designed to solve the echnical problems of the industry, and the dissemination of authoritative and practical information to the fishery industries and the ublic. Results of technological investigations and marketing studies are published in separate documents as each project is completed. The information obtained from statistical surveys is published in part 2 of this report, which includes all the detailed statistical information that has become available since the issuance of the previous report,² together with such summarized statements and interpretations of the statistics as are deemed significant and useful. In line with the general policy of economy of the Federal Government during the past year, the division's funds and personnel were greatly curtailed during this period. This made it necessary to curtail our economic and technological work and also made it impossible to conduct statistical surveys of the fisheries in several geographical sections. In the preparation of this report, members of the division's staff have taken part and their assistance is appreciatively acknowledged.

Part I. OPERATIONS OF THE DIVISION

COOPERATION WITH OTHER FEDERAL AGENCIES

As in 1933, various members of the division's staff assisted other Federal agencies in the conduct of technological, economic, and statistical work or studies having a bearing on the fishery industry. Such cooperation was rendered especially to the National Recovery Administration, the Department of Agriculture, the Federal Emergency Relief Administration, the Federal Surplus Relief Corporation, Reconstruction Finance Corporation, and others. The writer continued his detail with the National Recovery Administration, begun in August 1933, until September 30, 1934, as deputy administrator in charge of fishery codes, and on October 1, returned to the Bureau, resuming his regular duties. Since his return to the Bureau, the writer was appointed a member of the Food Survey Committee of the Department of Agriculture. This committee investigates the supply and price situation of surplus agricultural and fishery products to guide the Federal Surplus Relief Corporation in making purchases of surplus food commodities. Considerable assistance was given the Federal Emergency Relief Administration in connection with various relief activities among fishermen.

Also, the Bureau has cooperated with various Federal agencies in obtaining statistical data on our fisheries. The Bureau has the cooperation of the Bureau of Agricultural Economics, Department of Agriculture, in the collection of statistics on the volume of cold-

² Fishery Industries of the United States, 1933. By R. H. Fiedler, John Ruel Manning, and F. F. Johnson, Appendix I to the Report of the U. S. Commissioner of Fisheries for 1934, pp. 1-237.

storage holdings of fish, and the health authorities in Washington, D. C., in obtaining the volume of fish handled at the Municipal Fish Wharf and Market in this city. In another instance the Bureau obtains figures on the volume of the quarterly holdings of fish oils for the Bureau of the Census. During the past year, the Bureau aided the National Recovery Administration in obtaining data on methods for sharing the proceeds of a commercial fishing venture.

COOPERATION WITH STATE AGENCIES

In the conduct of its technological investigations, the Bureau has always encouraged and fostered cooperation with the States. By working in close conjunction with the members of the research staffs of various State laboratories and institutions, we have been able to increase the productivity of our relatively small technological staff and have been able to carry out such cooperative investigations at considerably less cost. During the past year, the following cooperative investigations were conducted in the State institutions listed:

A member of our technological staff was stationed in the laboratories of the State Medical College, Charleston, S. C., where members of the staff of the State Medical College have given valuable cooperation in a study of the mineral content of aquatic products. Dr. Roe E. Remington and Dr. Kenneth M. Lynch participated in these investigations.

At Massachusetts State College, Amherst, Mass., laboratory facilities were provided for our bacteriologist to study methods for handling fish. Members of the Massachusetts State College staff cooperating in these investigations were Doctors Fellers, Fuller, and Bradley.

At George Washington University, Washington, D. C., Dr. Leland W. Parr, associate professor of bacteriology in the school of medicine, assisted in the supervision of one of our cooperative investigations on sponge disinfectants.

At Cornell University, Ithaca, N. Y., Doctors H. S. Wilgus, Jr., L. C. Norris, and G. F. Heuser, cooperated in making feeding tests of fish meals experimentally prepared in our technological investigations.

Dr. J. S. Carver of Washington State College, Pullman, Wash., cooperated during the past year in carrying on tests with poultry in the feeding of salmon oils and meals experimentally prepared in our Seattle and Alaska laboratories.

The University of Washington, Seattle, Wash., placed space at the disposal of members of our Seattle technological laboratory for the conduct of certain byproducts investigations.

All of the above cooperative investigations are discussed in detail elsewhere in this report.

In the conduct of its statistical research work, the Bureau also obtains unusual cooperation from various States. The surveys of the fisheries in the various States bordering on the Great Lakes, in the Pacific Coast States, and in Maryland and Virginia, have been greatly facilitated by special cooperation obtained from the State fishery agencies in these States. With this aid, it is now only necessary for the Bureau to conduct partial surveys in these States to supplement the data available from the fishery agencies.

In addition, in nearly every other State where commercial fishing is prosecuted, some type of cooperation on its statistical work is rendered the Bureau by the State fishery or other agencies. This makes it possible for the Bureau to make statistical surveys of a greater portion of our fishery industries than otherwise would be possible.

EXHIBIT AT "A CENTURY OF PROGRESS"

The division continued during 1934 to have supervision of the Bureau's exhibit at "A Century of Progress", the world's fair at Chicago, Ill. The exhibit showed generally the activities of the Bureau in conserving our natural fishery resources and in fostering the fishery industry of the United States and Alaska.

Some of the activities of the Division of Fish Culture were depicted by models of a trout hatchery, fish ladder, and fish lock, and with several aquaria and pools containing some of the common species of food fishes taken in our fresh waters. In addition, a series of glass transparent pictures showed other activities of this division.

The Division of Scientific Inquiry exhibited apparatus used for deep-sea biological investigations, charts of a mackerel investigation being conducted by that Division, and methods of oyster culture. The biology of the Great Lakes fisheries was depicted by pictures and models of nets used in these fisheries.

The Alaska Division had an exhibit of natural and dyed fur-seal skins taken from the Pribiloff Islands.

The Black Bass and Angling Division exhibited fishing tackle appropriate for the taking of bass, trout, salt-water fishes, and surf fishes, thereby depicting the sport fishing angle of our fisheries.

The Division of Fishery Industries exhibited methods for the preservation of nets as recently developed by the Bureau. This was shown by charts, pieces of treated netting, and some of the chemicals used. A century of progress in the utilization of products and byproducts was exhibited by samples of such commodities marketed in 1833 as compared with those on the markets in 1933. Other exhibits of this Division included a display depicting the food value of fishery products, a display of canned and frozen aquatic foods, a model depicting the oyster fishery, and a display of the mussel-shell and pearl-button industry.

MARKETING INVESTIGATIONS

During 1934 an unusual interest was evidenced in the economic and marketing phases of the fishery industry. This industry has suffered to a very great degree during the depression, owing (1) to a curtailment of the movement of fishery products into consumption, and (2) to a somewhat lower price structure for the products that entered the market. This has resulted in a most demoralized situation, and various public and private agencies have attempted to develop plans which might be effective in removing the obstruction to the free flow of fishery products from the fishing port to the consumer. However, those attempting to develop a plan of action were invariably confronted with a lack of economic and marketing data on the fisheries upon which to base a foundation. This was most apparent in developing codes for the industry under the National Industrial Recovery Act, in effectuating relief to the industry through the Federal Emergency

Relief Administration, in extending credit to the industry through the Reconstruction Finance Corporation, in the development of cooperative associations under the act approved by the President on June 25, 1934, and in other ways.

During past years the Bureau conducted some studies on the marketing and distribution of fishery products. However, because of limited funds, these were of somewhat local nature and did not cover the situation in as detailed a manner as present necessity dictates. In view of this, and because fishery economic studies are needed, the writer has encouraged the various States to undertake marketing studies designed to aid the fishery industry within their States. This was urged upon the National Planning Council of Fish and Game Commissioners at the organization meeting of the council in St. Louis, Mo., in April 1934. At this time it was explained that fishermen are in dire need of assistance to market their catch profitably and it was suggested that the State fishery agencies might take a leading part in studying the marketing of fishery products. Some State fishery agencies have taken steps to conduct such studies, and in addition considerable fishery marketing work is being done by several State marketing departments. For instance, the Massachusetts Department of Agriculture has recently conducted schools in various cities of the State to instruct retail fish dealers in the proper care and handling of fresh fish and shellfish.

MARKET SURVEY IN SOUTHEASTERN STATES

With the limited funds at the disposal of the division for fishery marketing work, a study was made during the past year, by W. T. Conn and Herbert E. Munger, on the marketing of fish and shellfish in certain inland cities in the States of Maryland, Virginia, North Carolina, South Carolina, Georgia, Alabama, Mississippi, Tennessee, Kentucky, West Virginia, Ohio, and Indiana. In all, 39 of the larger cities in these States were visited by these men, who interviewed members of the wholesale and retail fish trade, restaurant owners, health officers, and others.

As a result of this study it was found that per capita consumption of aquatic foods west of the Appalachian Mountains in the area visited is about half of that east of the mountains. Great variations in trade practices and domestic demands were found in the several cities, almost every locality having special characteristics. In general, it was revealed that the usual domestic method for cooking fish is by frying, and that housekeepers and domestic help do not want to clean fish. There apparently are two distinct classes of domestic consumers in this area, one demanding quantity and the other demanding quality. A considerable proportion of fish is sold by retailers who have no interest in promoting the industry and who have taken business formerly handled through retail fish markets. Further, it was found that seafood can be retailed at a profit by progressive merchants; motor-truck peddlers have altered former trade practices in many localities; Negroes in the area apparently have double the per capita consumption of whites; there is a general lack of information regarding seafood and its domestic cookery; consumption through public eating places, especially cafeterias, is growing steadily; and that there is need of research to produce improved commercial handling of stocks.

PER CAPITA CONSUMPTION OF FISH

A study made during the year by Fred F. Johnson, in charge of statistical investigations, shows that the domestic per capita consumption of fish and shellfish during 1931 amounted to about 13.3 pounds in terms of the weight as prepared for market. This total figure was comprised of fresh and frozen products, 8.2 pounds; canned, 3.6 pounds; salted, 1 pound; smoked, 0.3 pound; and other products, 0.2 pound. It was interesting to note that only four species or groups of species contributed more than 0.5 pound to the per capita consumption. Of these salmon led with 2.7 pounds and following in order were the group consisting of cod, haddock, hake, pollock, and cusk, 1.5 pounds; sardines, 0.7 pound; and oysters, 0.6 pound. Eight additional species or groups contributed from 0.25 to 0.5 pound. These were in order of their importance—sea herring (excluding sardines), mackerel, flounders, halibut, clams, crabs, tuna and tunalike fishes (including Pacific yellowtail), and shrimp.

In 1931 the per capita consumption of meats, according to figures of the Department of Agriculture amounted to 133.2 pounds or about 10 times that of fish and shellfish. The total consumption figure of meats consisted of 69.6 pounds of pork, 49.6 pounds of beef, 7.1 pounds of lamb and mutton, and 6.9 pounds of veal.

STATISTICAL INVESTIGATIONS

Fishery statistics are collected by the Bureau to serve two principal purposes—first, biological; and second, economic. For this reason the Bureau must plan its statistical surveys to obtain comprehensive data for furnishing a complete and reliable picture of the condition and trend of the fisheries. The collection and compilation of the great mass of data necessary, involves many problems. The fisheries are broad in scope including over 160 varieties of aquatic products which enter into commercial production. These, many of which are migratory, are taken by a great variety of types of gear in water areas along our seacoast and in our interior lakes and streams. If the biological aspect is to be served, complete annual statistics are needed on each of these phases in every section. If the economic aspect is to be served statistics are needed on not only the phases listed above relative to the biological aspect but also on the price structure, the processing function, and the marketing and distributing functions.

BIOLOGICAL ASPECT

The biological aspect must consider two problems—the conservation and sustained supply of the resource and the prediction of future trends or yield. Since our fisheries are usually prosecuted in areas not under private ownership, the problem of the conservation of these fisheries is of national concern. It, therefore, is important that close watch be kept of the condition of the various fisheries to detect depletion so that remedial measures can be promulgated timely and wisely. For this reason it is imperative that current statistical data be obtained on the yield of our fisheries.

These statistics then furnish the biologist with the background upon which to base his prediction of future trends and yields. This he does

by coupling the statistical data with studies of the life history of the species. Difficulty is experienced in making these predictions because the supply (or population) of the species cannot be seen, as is the case with farm animals or crops. The more complete and more reliable the statistics on yield are, the better foundation the biologist has for conducting his studies. The Bureau, therefore, aims to obtain a complete picture of each individual fishery to further these biological studies.

ECONOMIC ASPECT

With the fishery conserved and trends and yields of the fishery predicted there still remains the problem of supplying the fishery trade with that information so essential to the conduct of its business activities. In these days of increasing competition the very existence of the fishery industry must depend upon reliable economic statistical information. Such material has been especially valuable during the recovery period when it has been required in national planning. The Bureau, therefore, aims to make its statistical survey so complete that the industry and the various governmental organizations may turn to it for reliable fishery statistics.

SURVEYS CONDUCTED

The statistical surveys during 1934 were conducted under the direction of Fred F. Johnson, in charge of statistical investigations. These surveys included the collection and dissemination of statistics of the commercial catch and its value, operating units, and employment in the fisheries. In addition, data were collected on employment and compensation of those engaged in, and products of fishery wholesale and manufacturing establishments.

Unfortunately, continued curtailment of funds and personnel made it necessary to eliminate surveys of the South Atlantic and Gulf States, the Lake States, and the States of the Mississippi River and tributaries; however, summaries of the production in these sections for the most recent years available are included in part 2 of this report.

In addition to the above, statistics were collected on the following special phases: The landings of fish by American fishing vessels at the ports of Boston and Gloucester, Mass., Portland, Maine, and Seattle, Wash. (published monthly); landings of halibut at North Pacific coast ports (published monthly); catch of mackerel in the North Atlantic fishery; cold-storage holdings of frozen and cured fish and amount of fish frozen, which are furnished by the Bureau of Agricultural Economics (published monthly); production, consumption, and holdings of marine-animal oils of the United States and Alaska (published quarterly by the Bureau of the Census); production of canned fishery products and byproducts of the United States and Alaska; transactions on the sponge exchange at Tarpon Springs, Fla.; volume of fishery products handled at the Municipal Fish Wharf and Market, Washington, D. C.; and the volume of the United States foreign trade in fishery products, furnished by the Bureau of Foreign and Domestic Commerce.

The following statistical and marketing agents assisted in the collection and compilation of the statistical data: H. F. Brown, W. H. Brown, F. F. Dimick, W. H. Dumont, R. L. Greer, V. E. Heffelfinger,

B. E. Lindgren, E. A. Power, W. H. Rich, V. J. Samson, C. B. Tendick, and A. S. Young.

The reader is especially referred to the section in the latter part of this report entitled "Statistical survey procedure" which gives in detail the methods employed in the collection of fishery statistics and other pertinent information.

TECHNOLOGICAL INVESTIGATIONS

Some of the food industries of this country during the past 15 years have made rapid strides in developing new products and new methods of manufacture and in developing a more orderly plan of marketing and merchandising. These developments are no doubt directly traceable to such factors as changes in consumer demand, education of the consumer, shifts in population, occupational changes among our people, and others. Many of these developments have leaned toward the production of prepared and packaged foods of new types, with consequent concentration of trimmings or waste which is available for conversion into valuable byproducts. In turn, methods have been found for converting much of this waste into useful commodities. These developments have clearly demonstrated the need for technical advancement in the aquatic food industry if it is to keep pace with the other food industries. In order to aid in this connection the Bureau is continuously conducting a series of technological investigations covering problems in the manufacture, preservation, and handling of fishery products. A summary of the accomplishments in these fields during the past year appears in the next few pages. For details regarding this work, the reader is referred to published reports and special articles of the various members of the staff, or in case the investigation is not completed, by direct communication with the Bureau of Fisheries in Washington.

LABORATORIES

During 1934 the Division carried on its technological studies under the direction of J. R. Manning, in charge of technological investigations, at its laboratories located in Washington, D. C., Gloucester, Mass., Seattle, Wash., College Park, Md., and at a field laboratory on Kodiak Island, Alaska. In addition, certain cooperative investigations were conducted by our technologists in the chemical laboratory of the State Medical College at Charleston, S. C., the food products laboratory of the Massachusetts State College at Amherst, Mass., the bacteriological and chemical laboratories at George Washington University, Washington, D. C., and the laboratory of the Agricultural Experiment Station, College Park, Md. A résumé of the facilities and equipment at the laboratories in Washington, D. C., Gloucester, Mass., Seattle, Wash., and Charleston, S. C., is contained in the report of this Division for 1933. Late in 1934, under a cooperative arrangement with the University of Maryland, our nutrition laboratory was moved from Washington, D. C., to quarters provided free of charge by the university at College Park, Md. This laboratory is now equipped with all the appliances which were formerly in the nutrition laboratory in Washington, D. C.

PRESERVATION OF FISHERY PRODUCTS FOR FOOD

Experimental work during the past year in the preservation of fishery products for food was conducted at the Gloucester laboratory by James M. Lemon, technologist in charge, assisted by Francis P. Griffiths, bacteriologist, and Maurice E. Stansby, chemist, and at the Washington laboratory by Norman D. Jarvis, technologist in charge of canning work. During the latter quarter of the year, Mr. Griffiths was detailed to conduct a series of cooperative bacteriological experiments at the food products laboratory of the Massachusetts State College, Amherst, Mass.

HANDLING FRESH MACKEREL

Mackerel is an extremely delicate fish and is easily bruised and crushed during packing and in shipment. For this reason special care is needed in the handling of this fish so that it will arrive at the market in good condition. One of the methods employed by the industry to overcome this difficulty is "floating," which consists of packing the mackerel in a watertight barrel with sea water and ice. In some cases, the sea water is drawn from the harbor under the packing plant, where it is more or less contaminated. While this method of packing retards bruising and crushing, it leads, in many cases, to decomposition of the mackerel en route to market, with consequent loss to the industry. In order to aid in minimizing these losses, a series of tests were conducted to develop a method for packing and shipping this fish which would have promise of commercial application. As a result of this work, it was found that when fresh mackerel were eviscerated, washed in clean water, and packed in boxes with finely crushed ice, a better product could be placed in the hands of consumers than with the "floating" method.

While this method is superior to the commercial method of "floating", it still is not ideal, as water from the melting ice has a tendency to leach the minerals from the fish.

STORAGE OF FROZEN MACKEREL

When frozen mackerel have been held in cold storage for a certain period, there is a tendency for the oil or fat in the fish to become rancid, thereby making the mackerel unfit for sale. In order to aid in correcting this condition, a series of experiments were conducted on the relationship between temperature of storage and the development of rancidity in mackerel oil, the action of enzymes on mackerel oil, and the seasonal variation in the oil content of mackerel. While these experiments have not progressed to the point where final results can be reported, they reveal that the percentage of oil in mackerel flesh varies from 2 to 5 percent from early spring catches to nearly 30 percent in the August catch, then declines to about 10 percent in the December catch. This study further showed that the development of rancidity is more or less directly correlated with the oil content. This would indicate that mackerel of low oil content, that is, those caught during the early spring, would be less apt to develop rancidity in cold storage. However, economic factors may tend to neutralize some of these benefits, for mackerel commands a relatively high price

in the spring, and rental charges for holding the frozen fish in cold storage over a longer period of months, as would be necessary, would be increased.

SMOKED MACKEREL AND STORAGE OF THIS PRODUCT

At periods during the peak of the mackerel season, it often becomes difficult for the trade to dispose of this fish in the fresh condition. This has led to depressing prices with consequent lowering of the income of mackerel fishermen. It has become increasingly evident that some different method of preservation is needed for mackerel which will present a new "appetite" appeal to consumers for this fish. In view of this, a study was begun, during the past year, at the Gloucester laboratory to develop methods for smoking mackerel and storing smoked mackerel under optimum conditions. While smoked mackerel is not what might be called a new product, as it can be obtained in the market at intervals, no extensive trade has been developed in product owing to the fact that it spoils easily during storage. At this writing, a technique has not yet been developed which has promise of commercial application for overcoming this spoilage. However, tests to date indicate that a practical method can be developed. In this connection, it is interesting to note that smoked mackerel can be used to make delicious and appetizing salads. It, therefore, appears that if a practical method can be developed, a new use can be found for this fish which may be of material aid to the mackerel fishery.

SHRINKAGE OF FRESH FISH PACKED IN ICE

During the past several years, considerable technical study has been given to the effect of ice when used as a preservative on fresh fish. The importance of this work may be seen from the fact that normally about 600,000,000 pounds of iced fresh fish are shipped to market annually. As a rule, the fish in these shipments are packed in direct contact with the ice. Previous study has shown that this method of icing results in dissipating a large proportion of the mineral salts in the fish, with consequent loss in flavor and food value. This loss is caused by general hydrolysis of the complex protein molecules of the flesh induced by the action of enzymes and bacteria. Thus, the soluble protein and mineral salts are drawn from the fish by the water formed from the melting ice. The actual loss in weight is of little consequence, although it may be as high as 10 pounds per ton over a period of 7 days, as further experiments show.

The present investigation was directed toward learning the effect on the weight of fresh fish when packed in ice. As a result, it was found that other than the loss of the mineral salts, and water soluble proteins, as referred to above, there is no appreciable loss in weight, provided the fish are entirely covered with ice. Apparently, there is an exchange of the mineral salts from the fish with water from the melting ice thereby holding the weight of the fish nearly constant. Thus, while this method of preservation results in little or no actual loss in weight, the flavor may be somewhat impaired. The latter may be corrected by packing the round or dressed fish or cuts therefrom in waterproof wrappers before being packed in the ice.

DETERMINATIONS FOR THE RELATIVE FRESHNESS OF FISH

For many years, there has been a need for a standard method for determining the relative freshness of fish or conversely for determining the stages of decomposition in fresh fish. A method has been desired by such governmental authorities as public health inspectors, pure food inspectors, grading inspectors, and the like, as well as many members of the fishery industry. The usual practice has been to determine freshness by an organoleptic test of the product, sometimes accompanied by bacteriological analysis. The former tests oftentimes have resulted in as many verdicts as to the quality as there are judges and the latter requires considerable time as it is necessary to make a culture from the bacteria obtained from the fish. It, therefore, appeared that a need existed for a test that could be used with uniform results by the layman and one which could be performed with a minimum of delay. In order to aid in this direction, a series of technical studies have been conducted by members of the technological staff which resulted in the development of an electrometric method for the detection of the relative freshness of haddock. This method, which was described in brief in last year's report of this division, consists in measuring electrometrically the quantity of a standard acid absorbed by the protein in a given quantity of the flesh of the fish and checking results with a previously developed chart of standardized values.

As the property for absorbing acid by the protein varies slightly in different species, it would be necessary to develop a technique for the various species. In view of this, studies were conducted, during the past year, for determining the relative freshness of cod, pollock, and mackerel. While these have not progressed to the point where a chart can be prepared against which to check the tests, they do indicate that this type of test is applicable to all species of nonoily fish, such as cod and pollock, but that possibly a different technique will need to be developed for such oily fish as mackerel. It appears that the technique for the oily fish will need to include a test for determining the relative rancidity of the oil in such fish.

CANNING AQUATIC FOODS

At the present time, only about 20 of the 160 varieties or species of fish and shellfish taken in our waters are canned (hermetically sealed in tin cans or glass jars) on a commercial scale. This has been due largely to the economic factors of supply and price. However, during late years, there has been a demand on the part of housewives, home economic and relief workers, and others for information on methods for canning aquatic foods. With this in mind, a series of technological and bacteriological investigations have been undertaken during the past several years, to develop methods for canning these products which could be safely and easily used in the home as well as by home economic and relief workers in community kitchens or in commercial establishments. This work has been conducted toward three principal goals. First, toward the selection of species taken in the fishery harvest which would be suitable for canning from the standpoint of availability and adaptability. Second, toward the determination of proper cooking temperatures to create sterility.

And, third, toward packing the product so it will present a pleasing appearance when removed from the can. During the conduct of this work, the following aquatic foods have been experimentally canned: Fish flakes; fish cakes; fish chowder; fish balls; fish pastes; mullet, plain, in tomato sauce, and spiced; mackerel, plain, in tomato sauce, spiced, and smoked; amberfish; salmon, plain and spiced; grouper; squeteagues; croaker; eels, plain and pickled; catfish; carp, plain and spiced; lake trout; whitefish; shrimp; crab meat; clams, whole, minced, and chowder; oysters; alewives; and shad.

In canning the crab meat, particular emphasis has been placed on developing a method for commercial application in canning meat taken from the blue crab caught along the Atlantic coast. Attention also has been given to packing fresh crab meat in various ways in cans for shipment to market under ice.

To date, successful methods have been developed for the home canning of salmon, shad, mackerel, lake trout, whitefish, mullet, and for canning certain types of spiced fish and fish chowder. These methods may be adapted with modifications to the commercial or semicommercial canning of these species. A report on this work has been recently issued by the Bureau and copies are available on request.

As regards the canning methods developed, it has been definitely shown that to obtain sterility in the final product, when canned in the home, it should be packed in no container larger than a no. 2 tin can or pint glass jar, and the product while in the container should be cooked in a steam-pressure retort equipped with a thermometer as well as a pressure gage.

BACTERIOLOGICAL INVESTIGATIONS

In the development of improved processes and methods for the preservation, storage, and other handling of aquatic products, the investigator is constantly confronted with the problems of bacterial spoilage. Therefore, in various of the investigations discussed previously, bacteriological studies are correlated with chemical and engineering studies. For instance, bacteriological examinations were made of the experimental packs of canned aquatic foods to determine which processes produced sterility, on the keeping qualities of fresh mackerel packed in ice, in determining the preservative value of smoke, on the tests developed for determining the relative freshness of fish, and others.

During the latter part of the past year, a special cooperative bacteriological study was begun to determine the effect of antiseptics in ice on the keeping quality of fresh haddock when packed in such ice for temporary preservation. This study is still in progress. In addition, a bacteriological investigation is in progress on disinfecting sponges used in the household. This will be discussed in the following section.

DISINFECTANTS FOR SPONGES

Sponges used in the household for cleansing purposes may be a possible medium for the spread of infectious diseases among the members of the family. In view of this situation, the sponge industry has been meeting with consumer resistance in the purchase of sponges. In order to learn whether this resistance was justified, and if so, what

steps might be taken to remedy the situation, the Sponge and Chamois Institute, New York City, requested the Bureau to undertake a study of this problem. This was undertaken as a cooperative investigation with George Washington University, Washington, D. C., where two students from the medical school receiving aid from the Federal Emergency Relief Administration, were detailed to our technological staff. This investigation, which has been conducted in the laboratory of the medical school, has not progressed to the point where definite conclusions can be drawn. However, thus far, it was found that thorough washing of the sponges in hot soapy water sterilizes sponges which had been experimentally infected with certain kinds of bacteria, but such washing has not proven effective in sterilizing the sponges infected with such pathogenic organisms as that causing ringworm. Further studies are being made of the value and possible application of other disinfectants such as phenol.

PRESERVATION OF FISHERY BYPRODUCTS

Because of the increasing demand for information with respect to the utilization of fishery waste materials, technologists of the division have continued an active program of research on the preservation of fishery byproducts. In 1934, investigations were in progress at the Gloucester technological laboratory, the Seattle technological laboratory, a field laboratory in Alaska, and at several State agricultural experiment stations. These studies were carried on under the direction of R. W. Harrison, in charge of our Seattle technological laboratory, with the assistance of A. W. Anderson, technologist of the Seattle laboratory, and S. R. Pottinger, chemist of the Gloucester technological laboratory.

UTILIZATION OF SALMON CANNERY WASTE

Our catch of salmon in the Pacific Coast States and Alaska averages between 500,000,000 and 600,000,000 pounds, annually. Of this amount, at least one-third, or approximately 200,000,000 pounds is waste material, the greater portion of which is discarded and represents a complete loss as well as a sanitary nuisance. Because of the great potential value of this amount of material, the Bureau has continued the studies on the utilization of salmon cannery waste as its major byproducts problem during 1934. The data obtained to date indicate that salmon waste is a potential source of vitamin active oils and highly nutritious fish meals.

Studies concerned with the distribution of the oil in the fish and their vitamin content show that the greater portion of the oil is in the head and flesh portions, that vitamin D is quite uniformly distributed in all the waste fat, although the visceral fats are richest in this vitamin, and that vitamin A is confined almost entirely to the visceral oils.

Oils from the total waste of the various species of salmon are quite uniform in vitamin D, but vary considerably in vitamin A. Chinook oil has the highest vitamin A potency, but contains less vitamin D than the other salmon oils. Sockeye and silver salmon are good sources of both vitamins A and D; and pink and dog salmon are good sources of vitamin D, but relatively poor sources of vitamin A. All salmon oils, therefore, are highly desirable for poultry nutrition, while oil from

chinook, sockeye, and silver salmon, if prepared properly, will compare favorably with U. S. P. requirements for medicinal cod-liver oil.

Studies with regard to methods for manufacturing the fish oils indicate that the common practice of cooking is not necessarily destructive to the vitamins extracted by the fats. Rather the vitamin potency of the oils seems to depend upon the freshness of the raw material, the extent to which the visceral parts are utilized in making the oil, and the efficient mechanical extraction of the press liquors.

As a corollary to these studies, our technologists gave assistance to the industry in the development of edible salmon oils for use in canning and conducted tests on the effect of adding high-grade salmon oil to salmon during the canning process. Preliminary results indicate that the addition of this oil is effective in improving quality.

Studies with respect to effect of manufacture on salmon meals were started, but sufficient data are not available for making a preliminary report.

EFFECT OF MANUFACTURE ON THE QUALITY OF NONOILY FISH MEALS

Studies concerned with the effect of manufacture on the composition and nutritive properties of nonoily fish meals, which have been in progress several years, were completed in 1933 and during the past year the data have been assembled for use in reports. These studies have shown:

(1) Fish meal proteins, as shown by tests for feeding efficiency, biological value, digestibility, and destruction of essential amino acids, are not affected as much by conditions of drying as is vitamin G.

(2) Vitamin G is water soluble and is removed with the water soluble protein by the wet process.

(3) The water soluble proteins removed by the wet process are not of as high quality as the residual flesh proteins.

(4) The head portion of the waste is the richest source of vitamin G, while the backbone and flesh portion has the highest quality protein.

(5) Flame drying is definitely destructive to vitamin G and decreases the value of the protein.

From these data it is possible to draw certain practical conclusions, some of which are:

(1) Dry process meals have greater general nutritional value than wet process meals, owing to the advantage given by vitamin G, in the former, overbalancing the slightly higher quality proteins of the latter.

(2) Steam dried meals are superior to flame dried meals.

(3) Steam drying under vacuum improves the value of a meal slightly over steam drying at higher temperature. This is due more to effect on vitamin G than on the protein.

(4) Wet process oily fish meals cannot be expected to be important sources of vitamin G.

(5) From the standpoint of quality, there would be no particular advantage in drying wet process meals under vacuum if normal steam drying at atmospheric pressure can be accomplished.

STUDIES OF THE DRYING OF NONOILY FISH WASTE

The studies on effect of manufacture on the quality of nonoily fish meals demonstrated the need for more information on a satisfactory dry rendering process. The experimental data obtained as a result of studies in this connection have shown that dry reduction can be accomplished without detrimental "sticking" and "caking" of the glue-like material in the dryer, if steam pressure and vacuum are maintained at proper levels as drying progresses. From a strictly production standpoint, the greater yield of meal by the dry process counterbalances the less difficult and apparently lower operating costs of the wet process. When nutritive quality of final products are considered, the dry process has an additional advantage over the wet process.

UTILIZATION OF SWORDFISH LIVERS

The work on swordfish-liver oil which was begun in 1933, to develop a method for the commercial manufacture of this product and thereby make use of a product not fully utilized, was continued during the past year. On the basis of the data obtained the following conclusions may be drawn:

Swordfish livers vary in fat content between 15 and 20 percent during the course of the fishing season. The livers do not give up oil readily either by steaming or boiling in water. To obtain efficient oil removal either solvent or improved mechanical extraction methods must be used.

Depending on the method of extraction, swordfish-liver oil will vary in color from amber to dark brown and from a liquid at room temperature to a solid fat.

The manner of fat and vitamin storage in swordfish livers is such that the mere extraction of oil does not give efficient vitamin extraction unless conditions are such that the oil comes in intimate contact with the liver tissue. This brings out the role of fat as a vitamin solvent during liver extraction and suggests the possibility of using fish and other oils commercially in extracting vitamins from low fat content livers.

Solvent-extracted swordfish-liver oils were prepared during the course of the investigation which were 100 times more potent than the U. S. P. standard of reference cod-liver oil, which contains 3,000 U. S. P. vitamin A units per gram and 95 U. S. P. vitamin D units per gram. These samples indicate that swordfish-liver oil is the richest known natural concentrate of both vitamins A and D.

A preliminary report on this investigation has been prepared and as a result of these studies, swordfish livers are now being used in liver-oil manufacture and fishermen are obtaining a new source of revenue from materials formerly discarded.

HADDOCK-LIVER OIL

The general results of the Bureau's haddock-liver oil investigation were given in the 1933 report. During the past year these data have been assembled in a form suitable for publication. In last year's report, it was stated that haddock-liver oil corresponded favorably

with U. S. P. specifications for cod-liver oil in every respect except iodine number. The variation with respect to iodine number is not important when unchilled oils are considered. However, during the past year, a new U. S. P. specification has appeared, in which U. S. P. cod-liver oil is designated as a partially destearinized oil. Partially destearinized haddock-liver oils will exceed U. S. P. iodine number specifications in many cases. Therefore, since it is quite generally agreed that haddock-liver oil was meant for inclusion with cod-liver oil and similar liver oils in the class of U. S. P. cod-liver oil, publication of these data should serve as a guide in changing specifications to cover adequately the materials that are supposed to be included under the classification.

NUTRITIVE VALUE OF AQUATIC PRODUCTS

Because we are concerned primarily with food products, our nutrition studies are a very necessary and important phase of our technological work. Not only is it essential to determine the food value of fish and shellfish products and byproducts now on the market, but the relative nutritive value of any experimentally prepared food product is the only true yardstick or standard of measurement for evaluating improvements in methods of manufacture, preservation, handling, and storage of such products.

During the past year, various phases of the nutrition investigations were carried on, respectively, in the nutrition laboratory in Washington, D. C., which in the early fall was moved to College Park, Md., and in the chemical laboratory of the State Medical College at Charleston, S. C., by Charles F. Lee and E. J. Coulson, of the technological staff.

SWORDFISH-LIVER OIL

In order to evaluate the adequacy of experimental methods for producing oils from swordfish livers, as described previously in this report, such oils as manufactured in the laboratory by various methods were tested for their vitamin potency.

Livers from last season's (1934) catch of swordfish were extracted with a number of different organic solvents, such as petroleum ether, ethyl ether, ethylene dichloride, carbon tetrachloride, ethyl acetate, acetone, and toluene. Extractions were made of both cooked and uncooked livers. Results thus far indicate that ethylene dichloride, petroleum ether, and ethyl ether, seem to yield the oil with the highest vitamin potency from both.

Including samples produced both in 1933 and 1934, the variation in vitamin A potency ranges from 36,000 to 300,000 international vitamin A units per gram. These samples vary in vitamin D potency from 2,850 to 9,500 international vitamin D units per gram. Expressing it another way, swordfish-liver oil runs as high as 100 times the U. S. P. standard reference cod-liver oil in vitamins A and D.

SALMON OIL

Vitamin tests of oils prepared from the trimmings remaining from salmon canning operations, from salmon eggs, and from salmon livers, were continued during 1934. The oil samples used in these tests were

prepared according to various experimental methods of manufacture by technologists in our Seattle and Alaska laboratories. In general, oils prepared from salmon livers are approximately 5 to 20 times as potent in vitamin A and approximately 2 to 3 times as potent in vitamin D as an average medicinal cod-liver oil. The oils prepared from salmon eggs and from trimmings compare favorably in vitamin potency with an average medicinal cod-liver oil. The results obtained in 1934 averaged about the same vitamin potency as the samples prepared experimentally in 1933 insofar as their vitamin D content was concerned, although some of them were somewhat low in vitamin A potency.

A more detailed discussion of the variation in potencies of salmon oils appears in a previous section of this report under the heading of "Utilization of salmon cannery waste."

HORSE MACKEREL-LIVER OIL

During the past year, a sample of oil extracted by ethylene dichloride from the liver of the horse mackerel or Atlantic tuna was tested for vitamins A and D. As a result, it was found that this oil contained 7,100 international vitamin D units and 60,000 international vitamin A units.

CRAB MEAT

Samples of fresh crab meat as prepared commercially from the blue crab of Chesapeake Bay were tested for vitamins A and G. In the vitamin A tests, the crab meat was found to contain 40 to 55 U. S. P. vitamin A units per ounce of crab meat. In the vitamin G tests, the crab meat was found to contain about one-half a unit of vitamin G per gram of crab meat. The so-called "curative technique" was used for the vitamin A assay, while the "Sherman" method was used for the vitamin G assay.

SODIUM ALGINATE

Because, in recent years, sodium alginate, a product made from the Pacific coast kelp, has found widespread use in dairy products as a stabilizer, the Bureau was requested by the industry to determine its food value. Accordingly, tests were made of this product with laboratory animals. Fed at the level of 3 percent of the ration, sodium alginate showed definite food value. Our tests indicated that this product possesses the same general properties in stimulating growth and appetite in the experimental animals (white rats) as was found in the kelp-meal tests conducted by our technologists, a few years ago (Technological Report No. 5).

CONCH MEAT

Considerable quantities of conchs are found along the coast of Florida and in certain of our possessions in the West Indies. The meat of this animal is highly prized for food in different localities and in recent years an effort has been made to widen distribution. This has been done by developing a canning industry in Florida which produces such products from conch meat as cocktails, juices, concentrates, etc. Since no analyses of conch meat were available, a study

was made of the chemical composition of this product. This revealed that conch meat, on a fresh basis, contained the following: moisture, 74.6 percent; protein, 18.6 percent; fat, 0.3 percent; and ash, 1.7 percent. Also it was found that conch meat contains 290 parts of iodine per billion.

MINERAL CONSTITUENTS OF FISHERY PRODUCTS AND BYPRODUCTS

Much has been written in the scientific literature in recent years concerning the increasing importance of minerals in nutrition. Probably no other class of foods offers so attractive a field of study, in this respect as fishery products, since it is commonly known that these products contain minerals in quantity and variety, many of which have been shown by scientific investigators to be of great importance in both human and animal nutrition. During recent years the Bureau has carried on an extensive study of the nutritive value of minerals in fishery products. Toward this end, chemical analyses of the quantity of these minerals in various fishery products of commercial importance are being made. Following this, these fishery products are fed to laboratory animals to determine the biological value of such minerals.

The biochemical investigations during the past year have been of two types; first, a study of those mineral elements contained in sea foods which are considered to be of great importance in both human and animal nutrition; and second, investigations of the characteristics of those elements, also contained in sea foods, which are usually considered to be deleterious to health. Investigations of the iodine content of various fishery products of commercial importance by a more recent method has revealed that the iodine content of some of them is actually many times higher than has previously been reported in this country. For instance, the following average results are reported of the iodine content, in parts per billion, of some of the important species of fish, on a fresh basis: haddock, 5,130; cod, 1,030; and mullet, 4,850.

The problem of the removal of arsenical spray residues from fruits and vegetables in order to make those foods safe for human consumption has recently focused the attention of investigators on the naturally high arsenic content of sea foods. Because of increasing concern which is being shown by scientific workers at the present time, an investigation was undertaken to study the characteristics of organic arsenic as it is contained in marine products when fed to laboratory animals.

The results of these investigations have revealed that there is a difference in the metabolism of organic arsenic as it occurs in shrimp as compared to inorganic arsenic and that while the inorganic arsenic is absorbed and stored in the body, the organic arsenic from shrimp is eliminated. Careful histological studies of the organs of the experimental animals have failed to reveal any deleterious effects of this element when fed in the form of shrimp. Like studies on the effect of feeding organic copper in the form of "coppery green" oysters have yielded similar results.

Considering the results of the above-described work on arsenic and copper, as well as the findings of other investigators that only a part of the iron and copper of some foods is available for hemoglobin production, it is becoming apparent that not only must the presence of an element in a food be proven, but its availability to the body must be tested by animal experimentation methods. Consequently, it can

no longer be assumed that the chemical analysis of a food, with regard to its mineral content, is an accurate measure of its value as a source of those minerals in the diet.

FISH COOKERY INVESTIGATIONS

As is related elsewhere in this report, the per capita consumption of aquatic foods in this country is very low compared with many other countries, being only about 13.3 pounds annually. This, of course, has a direct effect on the industry in that capacity production cannot be attained by fishermen and processors and also an effect on our people in that they are depriving themselves of a wholesome nutritious food of valuable healthful properties. Believing that consumption is retarded because of the lack of informative methods for cooking fish and shellfish for the table, the Bureau installed a fish cookery kitchen in its laboratory in the Department of Commerce Building in Washington. Here a series of simple recipes for cooking aquatic foods in the home were tested by Miss Agnes I. Webster and W. T. Conn.

During the course of this work, many kinds of fish and shellfish were cooked by frying, broiling, baking, boiling or steaming, and planking. Also recipes were tested for making fish chowders, and flaked fish, for cooking fish with vegetables, for cooking salted and smoked fish, and for cooking oysters, shrimp, crabs, clams, lobsters, and scallops. In addition, recipes were tested for making sauces to be used with fish and shellfish.

As a result of these tests, a cookbook has been prepared giving simple recipes for cooking aquatic foods which can be followed with ease in the home by the housewife.

RESEARCH ASSOCIATES AND STUDENT ASSISTANTS

In the above lines of industrial research the Bureau has attacked those fundamental problems which promise to be of greatest value to the largest number and which are possible with the funds and personnel available for the purpose. For this reason, the division has not been able to study special problems affecting certain products, processes, or methods. In order to serve the industry in this connection, the Bureau by congressional authorization has provided research associate facilities whereby firms or groups having special industrial problems to solve will furnish the investigator and pay his salary and expenses. The investigation is carried out in cooperation with the Bureau's staff in its laboratories and under its control. Thus the industry can be provided with laboratory, consultation, and library facilities which in many instances it is unable to obtain elsewhere.

In addition to the above, the Bureau has opened its technological laboratories to research students pursuing courses in several universities. In this manner the student is able to apply in a practical manner the theoretical knowledge obtained at the university. These men are usually assigned to aid our investigators in the conduct of investigations already initiated by the Bureau. For instance, during the past year, Donald A. Bean and Richard M. Locke, students at Northeastern University in Boston, spent alternately 5 weeks in the university pursuing the regular-school curriculum and 5 weeks in our technological laboratory at Gloucester, Mass., aiding in the conduct of research problems pertaining to fish preservation and spoilage.

James W. McCurley, W. B. Matthews, John W. Webster, Roscoe Dwiggin, Arthur Kahn, and F. L. Clavelaus, jr. (deceased), students at the University of Maryland receiving aid from the Federal Emergency Relief Administration, worked in our nutrition laboratory at College Park, assisting in the conduct of studies on the vitamin content of fish oils. These men, during the school year, spent several hours per day in the laboratory.

At George Washington University, Washington, D. C., William H. Conway and William J. P. Howard, medical school students receiving aid from the Federal Emergency Relief Administration, were detailed to a study of the development of disinfectants for sponges in household use. This project was under the joint supervision of the Bureau and the university, and the details of the investigation are described elsewhere in this report.

EDUCATIONAL AND CONSULTING SERVICES

In addition to the activities described in this report, our economic and technological staff conducts an educational and consulting service for those interested in the fisheries. Some of these functions and services have been discussed or referred to in previous paragraphs of the report. In addition, various members of the staff have conducted lectures and practical demonstrations relative to the fisheries and the fishery industries at various State educational institutions and at public gatherings of various sorts. Further, members of the staff have delivered radio addresses on various fishery subjects. The Bureau or the division also answered many thousands of letters on fishery subjects and has supplied information to persons who have called at the Bureau, personally. Many of the latter have come from foreign lands to seek information on the conduct of the fishery industries in the United States, which might be useful in the more orderly conduct of the fisheries of their native lands.

PUBLICATIONS OF THE DIVISION

During the calendar year 1934 the following publications were prepared and addresses delivered by members of the division's staff. These do not include the monthly statistical bulletins of the landings of fishery products at Boston and Gloucester, Mass., Portland, Maine, and Seattle, Wash., nor the monthly reports on cold-storage holdings of frozen fish and quantities of fish frozen. The fishery reports and circulars may be purchased at the prices shown from the Superintendent of Documents, Government Printing Office, Washington, D. C. The statistical bulletins and special or S-memoranda are distributed free of charge upon request to the Bureau. The special articles may be obtained from the sources of publication.

Those wishing to receive current copies of this report and statistical bulletins issued by the Bureau should request that their names be placed on the Bureau's mailing lists nos. 128 for the annual statistical report, 128a for general statistical bulletins, and 128b for monthly cold-storage reports. Those desiring historical statistical data on the domestic fisheries for the period 1880 to 1929 should consult the report entitled "Fishery industries of the United States, 1930" by R. H. Fiedler, Appendix II to the Report of the United States Commissioner of Fisheries for the fiscal year 1931.

DOCUMENTS, REPORTS, AND CIRCULARS

- COULSON, E. J.
The iodine content of oysters. 8°, 10 pp., 1 fig. Investigational Report No. 18. 5 cents.
- COULSON, E. J., ROE E. REMINGTON, and KENNETH M. LYNCH.
Studies on the metabolism of copper. 8°, 12 pp., 2 figs. Investigational Report No. 23. 5 cents.
- FIEDLER, R. H., JOHN RUEL MANNING, and F. F. JOHNSON.
Fishery industries of the United States, 1933. 8°, 237 pp. Appendix I Report of Commissioner, 1934. 20 cents.
- GRIFFITHS, FRANCIS P., and J. M. LEMON.
Studies on the smoking of haddock. 8°, 12 pp., 1 fig. Investigational Report No. 20. 5 cents.
- JOHNSON, F. F.
Aquatic shell industries. 8°, 17 pp., 5 figs. Fishery Circular No. 15. 5 cents.
- JOHNSON, FRED F., and MILTON J. LINDNER.
Shrimp industry of the South Atlantic and Gulf States, with notes on other domestic and foreign areas. 8°, 83 pp., 31 figs. Investigational Report No. 21. 10 cents.

SPECIAL ARTICLES AND ADDRESSES

- CONN, W. T.
Net preservative research, 1933, with recommendations. Bureau of Fisheries' Memorandum S-335, February 9, 1934. Published in Fishing Magazine, February 1934, New York City.
Retailing fish. Bureau of Fisheries' Special Memorandum 2450-G, December 19, 1934. Published in The Fisherman, February-March 1935, Grand Haven, Mich.
A tip to coast fishermen on nets. Alabama Game and Fish News, August 1934, Montgomery, Ala.
Save nets—save dollars. The Florida Conservator, December 1934, Tallahassee, Fla.
More life from fish nets. (Bureau of Fisheries' Special Memorandum 1651-H, December 1933.) Reproduced (without trade lists) in Fish and Oyster Reporter, February 1934, Tampa, Fla.; Atlantic Fisherman, January 1934, Goffstown; Fishing Magazine, February 1934, New York City; abstracted in Pacific Fisherman, February 1934, Seattle, Wash.; and recommended in the Fishing Gazette, January 1934, New York City.
Atmospheric exposure of linen and cotton with special reference to fish nets. (Bureau of Fisheries' Special Memorandum 1651-G, published November 1933.) Reproduced in substance in The Fisherman, November 1934, Grand Haven, Mich.
- COULSON, E. J.
The oyster as a source of minerals. Fishing Magazine, May 1934, New York City.
Oysters in anemia. Address before the joint convention of the National Shellfisheries Association and the Oyster Growers and Dealers Association of North America, Inc., Baltimore, Md., August 7, 1934.
Chemical composition of conch meat. Bureau of Fisheries' Special Memorandum 2065-B.
- COULSON, E. J., ROE E. REMINGTON, and KENNETH M. LYNCH.
Studies on the toxicity of copper. Address before the Biological Chemistry Section, American Chemical Society, St. Petersburg, Fla., March 27, 1934. Published as Fisheries' Investigational Report No. 23.
Toxicity of naturally occurring arsenic in foods. Science. September 7, 1934, New York City. Bureau of Fisheries' Special Memorandum 2524.
- FIEDLER, R. H.
Purpose of the oyster code. Address before the joint convention of the National Shellfisheries Association and the Oyster Growers and Dealers Association of North America, Inc., Baltimore, Md., August 6, 1934.
Relationship between the sportsmen anglers and the commercial fishermen. Radio address delivered over Station WMAL, November 30, 1934. Washington.
Codifying our fisheries under the Recovery Act. Address before the National Planning Council of Fish and Game Commissioners, St. Louis, Mo., April 23, 1934.

Griffiths, Francis P., and Maurice E. Stansby.

The significance of bacterial count and chemical tests in determining the relative freshness of haddock. Address before the Montreal meeting of the American Fisheries Society, September 13, 1934.

Harrison, Roger W.

Recent developments in fisheries technology. Address before Fisheries Section, Pacific Northwest Regional Planning Council, Seattle, Wash., December 13, 1934.

Jarvis, Norman D.

Cold smoked Florida fish. Florida Conservator, vol. 1, no. 5, pp. 11-12, November 1, 1934. Tallahassee, Fla.

Lee, Charles F., and Chester D. Tolle.

Salmon liver and salmon egg oils, vitamin content, and chemical and physical properties. Industrial and Engineering Chemistry, vol. 26, p. 446, April 1934. Bureau of Fisheries' Special Memorandum 1738-23.

Manning, J. R.

Relation of the fishery industries as a source of raw material in the manufacture of dog foods and related products for animal feeding and general information on the need for standards and marketing grades in this industry. (Special report prepared for the National Recovery Administration and delivered at the public hearing on the Code of Fair Competition for the Dog Food Industry on Feb. 16, 1934.) Bureau of Fisheries' Special Memorandum 2450-1.

Interchangeability of uses of oils and fats. (Copied from p. 5925 of Cong. Rec., 73d Cong., 2d sess., vol. 78, no. 70, Mar. 29, 1934, Washington.) Bureau of Fisheries' Special Memorandum 1417-B.

Technological investigation in the Great Lakes fisheries. Published in The Fisherman, vol. 3, no. 3, May-June 1934. Grand Haven, Mich.

Sumario del valor de los productos de la pesca en la nutrición humana y animal. Published in Boletín de la Oficina Sanitaria Panamericana, Año 13: no. 6, Junio de 1934.

Fish and shellfish for food. Bureau of Fisheries' Special Memorandum 2256-B.

Domestic fish oils now available as sources of vitamins for human nutrition. Prepared for and issued by the Division of Cooperative Extension, United States Department of Agriculture, December 6, 1934.

Remington, Roe E., E. J. Coulson, and H. von Kolnitz.

Application of enclosed torch to estimation of arsenic in foods. Industrial and Engineering Chemistry, Analytical Edition, July 15, 1934. Easton, Pa.

STATISTICAL BULLETINS

- Fisheries of the New England States, 1932. Statistical Bulletin No. 1074.
 Fisheries of the Middle Atlantic States, 1932. Statistical Bulletin No. 1061.
 Fisheries of the Chesapeake Bay States, 1932. Statistical Bulletin No. 1062.
 Fisheries of the South Atlantic and Gulf States, 1932. Statistical Bulletin No. 1073.
 Fisheries of the Pacific Coast States, 1932. Statistical Bulletin No. 1068.
 Lake fisheries, 1932. Statistical Bulletin No. 1069.
 Fisheries of the United States and Alaska, 1932. Statistical Bulletin No. 1075.
 Fishery products frozen and cold storage holdings of frozen and cured fishery products in the United States and Alaska, 1933. Statistical Bulletin No. 1067.
 Production of fresh and frozen packaged fish in the United States, 1933. Statistical Bulletin No. 1080.
 Canned fishery products and byproducts of the United States and Alaska, 1933. Statistical Bulletin No. 1086.
 Fisheries of Alaska, 1933. Statistical Bulletin No. 1085.
 Landings by fishing vessels at principal New England ports, 1933—by months. Statistical Bulletin No. 1065.
 Landings by fishing vessels at principal New England ports, 1933—by gear and fishing grounds. Statistical Bulletin No. 1079.
 Fishery products landed by United States fishing vessels at Seattle, Wash., 1933. Statistical Bulletin No. 1081.

Part 2. FISHERY STATISTICS, 1933**GENERAL REVIEW**

Available data for 1933 indicate an appreciable increase in both the volume and the value of the catch of fishery products in the United States and Alaska as compared with the preceding year. Statistics of the catch in the New England, Middle Atlantic, Chesapeake, and Pacific States, and Alaska were collected for 1933, and when considering the combined catch of these sections alone an increase of 13 percent in the volume and also 13 percent in the value of the catch in 1933 is indicated as compared with the same sections the previous year. This increase was accounted for principally in the Pacific Coast States and was reflected especially in increased catches of salmon, pilchard, and mackerel. The value of the production of canned fishery products in all sections increased 37 percent as compared with 1932; byproducts increased 40 percent; and frozen fish, 14 percent. The value of the production of packaged fish also increased.

The total catch of fishery products in the United States and Alaska as based on the most recent surveys, amounted to 2,899,048,000 pounds, valued at \$60,218,000. About 117,000 fishermen were employed in making this catch.

In 1933 in the United States and Alaska, the production of canned fishery products amounted to 533,212,154 pounds, valued at \$59,799,963; the output of byproducts was valued at \$17,465,986; and the production of frozen fishery products amounted to 95,873,507 pounds, estimated to be valued at \$8,000,000. Based on the most recent surveys the production of cured fishery products amounted to 104,310,213 pounds, valued at \$12,823,491, and fresh and frozen packaged fish and shellfish, 129,608,348 pounds, valued at \$17,294,092. It is estimated that about 500,000,000 pounds of fresh fishery products (excluding packaged fish and shellfish), valued at about \$40,000,000, were marketed during 1933. The total marketed value to domestic primary handlers of all fishery products in 1933 is estimated at about \$155,000,000.

Fishery products imported for consumption were valued at \$30,462,341 and domestic exports were valued at \$8,338,723.

New England States.—The catch in these States for 1933 showed an increase in volume as compared with the previous year but a decrease in value. The value was less than in any year for which there are records since 1902. There were increases in both the volume and value of the combined landings of fishery products by vessels at Boston and Gloucester, Mass., and Portland, Maine, and there was a large increase in production of Maine sardines. The quantity of fish frozen was practically the same as in 1932.

Middle Atlantic States.—The catch statistics for the Middle Atlantic States in 1933 showed an increase in both volume and value as compared with the previous year; however, with the exception of 1932 the value was less than in any year for which there are records. There was an increase in the production of frozen fish but a small decrease in the production of packaged fish. There was a decrease in the production of shad on the Hudson River.

Chesapeake Bay States.—In 1933 the catch of fishery products in the Chesapeake Bay States decreased sharply, the volume being less than in any year since 1888 and the value to the fishermen was less than in any year for which there are records. Menhaden products,

which are produced in Virginia, decreased in both volume and value and there was a decrease in the catch of shad in the Potomac River.

South Atlantic and Gulf States.—No survey for statistics of the catch of fishery products in these States was made for 1933. In 1932 the volume of the catch showed a small increase over the preceding year, but the value was less than in any year for which records were available since 1902. There was a considerable increase in the output of canned shrimp in 1933 but a decrease in the production of canned oysters.

Pacific Coast States.—Statistics of the Pacific Coast States for 1933 showed the largest volume of catch of fishery products of any year since 1929 and the largest value since 1930. There were increases in the packs of canned salmon, sardines, tuna and tunalike fishes, and mackerel and frozen fish.

Lake States.—No survey has yet been made for catch statistics of the fisheries of the Great Lakes and the international lakes of northern Minnesota for 1933. In 1932 the catch decreased somewhat under that of 1931.

Mississippi River and tributaries.—The most recent complete catch statistics of the fisheries of the Mississippi River and tributaries are those collected for the year 1931. As compared with 1922 when the most recent preceding survey was made, there was a decrease in the catch which was reflected principally in a smaller catch of fresh-water mussels. A survey made for Lakes Pepin and Keokuk and the Mississippi River between these two lakes showed a decrease in 1933 as compared with the preceding year. The production of mussel-shell buttons increased appreciably in 1933.

Alaska.—The catch of fishery products in Alaska in 1933 increased in both volume and value as compared with 1932. The pack of canned salmon was somewhat less in volume in 1933, but the value was considerably greater; the production of frozen fish increased in both volume and value; and the output of cured products decreased in volume and increased in value.

Fisheries of the United States and Alaska

SUMMARY OF CATCH: BY SECTIONS

[Expressed in thousands of pounds and thousands of dollars; that is, 000 omitted]

| Products | New England, 1933, Area XXII | | Middle Atlantic, 1933, Area XXIII | | Chesapeake, 1933, Area XXIII | | South Atlantic and Gulf, 1932, Areas XXIV and XXV | | Pacific, 1933 | |
|---------------------|------------------------------|--------|-----------------------------------|-------|------------------------------|-------|---|-------|---------------|--------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| Fish..... | 461,621 | 9,593 | 142,771 | 2,266 | 189,708 | 2,055 | 184,158 | 2,715 | 840,523 | 12,793 |
| Shellfish, etc..... | 38,315 | 3,893 | 28,982 | 2,545 | 82,672 | 3,006 | 115,759 | 3,713 | 15,307 | 1,111 |
| Whale products..... | | | | | | | | | 4,331 | 84 |
| Total..... | 499,936 | 13,486 | 169,753 | 4,811 | 272,380 | 5,061 | 299,917 | 6,428 | 860,161 | 13,988 |

| Products | Lakes, 1932 | | Mississippi River and tributaries, 1931 | | Alaska, 1933 | | Total | |
|---------------------|-------------|-------|---|-------|--------------|-------|-----------|--------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| Fish..... | 81,829 | 4,361 | 44,062 | 2,257 | 624,687 | 8,941 | 2,569,359 | 44,961 |
| Shellfish, etc..... | 1,915 | 28 | 38,321 | 640 | 2,709 | 148 | 321,980 | 15,084 |
| Whale products..... | | | | | 3,378 | 69 | 7,709 | 153 |
| Total..... | 83,744 | 4,389 | 82,383 | 2,897 | 630,774 | 9,158 | 2,899,048 | 60,218 |

NOTE.—The roman numerals appearing under the names of the sections are the numbers given these areas by the North American Council on Fishery Investigations. It should be explained that there are included under these areas craft whose principal fishing ports are in the respective areas but at times they may fish elsewhere.

Fisheries of the United States and Alaska—Continued

OPERATING UNITS: BY SECTIONS

| Item | New England, 1933 | Middle Atlantic, 1933 | Chesapeake, 1933 | South Atlantic and Gulf, ¹ 1932 | Pacific, 1933 |
|---|-------------------|-----------------------|------------------|--|---------------|
| Fishermen: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| On vessels..... | 5,049 | 2,442 | 2,125 | 2,409 | 6,512 |
| On boats and shore..... | 12,024 | 6,132 | 18,017 | 19,151 | 12,161 |
| Total | 17,073 | 8,574 | 20,142 | 21,560 | 18,673 |
| Vessels: | | | | | |
| Steam..... | 24 | 19 | 25 | ----- | 5 |
| Net tonnage..... | 2,879 | 3,010 | 2,831 | ----- | 116 |
| Motor..... | 570 | 384 | 105 | 441 | 919 |
| Net tonnage..... | 16,602 | 6,121 | 1,791 | 6,646 | 24,866 |
| Sail..... | 1 | 4 | 157 | 71 | 5 |
| Net tonnage..... | 47 | 32 | 1,749 | 841 | 2,173 |
| Total vessels | 595 | 407 | 287 | 512 | 929 |
| Total net tonnage | 19,528 | 9,163 | 6,341 | 7,457 | 27,155 |
| Boats: | | | | | |
| Motor..... | 4,581 | 1,747 | 7,492 | 5,052 | 5,283 |
| Other..... | 3,819 | 2,123 | 5,937 | 7,797 | 1,238 |
| Accessory boats..... | 1,227 | 157 | 102 | 85 | ----- |
| Apparatus: | | | | | |
| Haul seines..... | 79 | 240 | 524 | 891 | 131 |
| Purse seines..... | 178 | 27 | 34 | 42 | 430 |
| Lampara nets..... | ----- | ----- | ----- | ----- | 201 |
| Otter trawls (including all types and sizes)..... | 566 | 197 | 27 | 1,680 | 18 |
| Beam trawls..... | ----- | ----- | ----- | ----- | 32 |
| Paranzella nets..... | ----- | ----- | ----- | ----- | 13 |
| Gill nets..... | 7,348 | 1,520 | 9,376 | 10,860 | 4,171 |
| Trammel nets..... | ----- | ----- | ----- | 358 | 52 |
| Pound nets, trap nets, and weirs..... | 457 | 574 | 2,570 | 1,737 | 436 |
| Stop nets..... | ----- | 96 | 6 | 7 | ----- |
| Fyke nets..... | 121 | 2,667 | 2,830 | 1,085 | 2,591 |
| Bag nets and pocket nets..... | 144 | ----- | ----- | ----- | 44 |
| Other nets ² | 423 | 433 | 3,133 | 2,062 | 370 |
| Hooks, baits, or snoods..... | 3,617,680 | 622,304 | 1,437,984 | 324,694 | 1,245,282 |
| Fish wheels..... | ----- | ----- | ----- | 21 | 20 |
| Eel pots and traps..... | 4,774 | 7,324 | 8,870 | 1,325 | ----- |
| Lobster pots..... | 319,460 | 46,576 | ----- | ----- | 6,099 |
| Crab, shrimp, crawfish pots, traps, drags, etc..... | 3,407 | 10 | ----- | 4,635 | 20,334 |
| Clam dredges..... | 75 | 90 | ----- | 1 | ----- |
| Crab dredges..... | ----- | 86 | 130 | ----- | ----- |
| Mussel dredges..... | 1 | 2 | ----- | ----- | ----- |
| Oyster dredges..... | 107 | 353 | 716 | 577 | 4 |
| Scallop dredges and drags..... | 2,569 | 260 | 2 | 64 | ----- |
| Crab scrapes..... | ----- | ----- | 683 | ----- | ----- |
| Tongs, rakes, hoes, forks, picks, grabs, etc..... | 4,509 | 3,229 | 9,423 | 2,920 | ----- |
| Abulone diving outfits..... | ----- | ----- | ----- | ----- | 17 |
| Sponge diving outfits..... | ----- | ----- | ----- | 54 | ----- |
| Other apparatus ³ | 3,007 | 200 | 2 | 2,593 | 54 |

| Item | Lakes, 1932 | Mississippi River and tributaries, 1931 | Alaska, 1933 | Total |
|--------------------------------|---------------|---|---------------|----------------|
| Fishermen: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| On vessels..... | 1,705 | ----- | 8,656 | 28,898 |
| On boats and shore..... | 5,227 | 15,884 | ----- | 88,598 |
| Total | 6,932 | 15,884 | 8,656 | 117,494 |
| Vessels: | | | | |
| Steam..... | 106 | ----- | 3 | 182 |
| Net tonnage..... | 2,364 | ----- | 217 | 11,417 |
| Motor..... | 392 | ----- | 504 | 3,315 |
| Net tonnage..... | 4,055 | ----- | 7,370 | 67,421 |
| Sail..... | ----- | ----- | ----- | 238 |
| Net tonnage..... | ----- | ----- | ----- | 4,842 |
| Total vessels | 498 | ----- | 507 | 3,735 |
| Total net tonnage | 6,419 | ----- | 7,587 | 83,680 |

¹ Includes the fisheries of Lake Okeechobee, Fla.

² Includes persons in boat and shore fisheries.

³ Includes dip nets, push nets, reef nets, cast nets, scap nets, and drag nets.

⁴ Includes periwinkle, cockle, and fish pots; harpoons; spears; baskets; and box traps.

Fisheries of the United States and Alaska—Continued

OPERATING UNITS: BY SECTIONS—Continued

| Item | Lakes, 1932 | Mississippi River and tributaries, 1931 | Alaska, 1933 | Total |
|---|-------------|---|------------------|--------------|
| | Number | Number | Number | Number |
| Boats: | | | | |
| Motor..... | 1, 624 | 4, 426 | 1, 032 | 31, 237 |
| Other..... | 1, 535 | 10, 120 | 3, 186 | 36, 755 |
| Accessory boats..... | | | | 1, 571 |
| Apparatus: | | | | |
| Haul seines..... | 332 | 1, 013 | 102 | 3, 321 |
| Purse seines..... | | | 445 | 1, 156 |
| Lampara nets..... | | | | 201 |
| Otter trawls (including all types and sizes)..... | | | | 2, 488 |
| Beam trawls..... | | | 11 | 43 |
| Paranzella nets..... | | | | 13 |
| Gill nets..... | 103, 518 | 101 | 3, 346 | 140, 249 |
| Trammel nets..... | 226 | 518 | | 1, 154 |
| Pound nets, trap nets, and weirs..... | 9, 259 | 374 | 416 | 15, 823 |
| Stop nets..... | | | | 109 |
| Fyke nets..... | 2, 574 | 32, 541 | | 44, 409 |
| Bag nets and pocket nets..... | | | | 188 |
| Other nets ¹ | | 191 | | 6, 612 |
| Hooks, baits, or snoods..... | 900, 513 | 2, 459, 179 | (⁴) | 10, 667, 636 |
| Fish wheels..... | | | 280 | 330 |
| Eel pots and traps..... | | | | 22, 293 |
| Loyster pots..... | | | | 372, 135 |
| Crab, shrimp, crawfish pots, traps, drags, etc..... | 2, 910 | 456 | 992 | 32, 744 |
| Clam dredges..... | | | | 166 |
| Crab dredges..... | | | | 216 |
| Mussel dredges..... | | 440 | | 443 |
| Oyster dredges..... | | | | 1, 757 |
| Scallop dredges and drags..... | | | | 2, 895 |
| Crab scrapes..... | | | | 683 |
| Toms, rakes, hoes, forks, picks, grabs, etc..... | 126 | 3, 994 | | 24, 201 |
| A balone diving outfits..... | | | | 17 |
| Sponge diving outfits..... | | | | 54 |
| Crowfoot bars..... | 360 | 4, 480 | | 4, 810 |
| Other apparatus ² | | 3, 781 | | 9, 637 |

CATCH: BY SECTIONS

[Expressed in thousands of pounds and thousands of dollars; that is, 000 omitted]

| Species | New England, 1933 | | Middle Atlantic, 1933 | | Chesapeake, 1933 | | South Atlantic and Gulf, 1932 ¹ | | Pacific, 1933 | |
|------------------------------|-------------------|------------------|-----------------------|------------------|------------------|------------------|--|------------------|------------------|------------------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| FISH | | | | | | | | | | |
| Alewives..... | 2, 817 | 17 | | 10 | | | | | | |
| Amberjack..... | 2 | (⁵) | | | | | | | | |
| Anchovies..... | | | | | | | | | 317 | 4 |
| Barracuda..... | | | | | | | | | 3, 073 | 123 |
| Black bass..... | | | | | 77 | 6 | 4 | (⁵) | | |
| Bluefish..... | 921 | 76 | 3, 252 | 132 | 803 | 35 | 310 | 22 | | |
| Blue runner or hardtail..... | | | | | | | 2, 131 | 78 | | |
| Bonito..... | 52 | 2 | 247 | 14 | 19 | 1 | 163 | 2 | | |
| Bowfin..... | | | | | | | 2 | (⁵) | | |
| Buffalofish..... | | | | | | | 12 | 1 | | |
| Butterfish..... | 1, 554 | 68 | 4, 410 | 180 | 2, 875 | 68 | 55 | 1 | | |
| Cabio or crab eater..... | | | | | 20 | 1 | 6 | (⁵) | | |
| Cabrilla..... | | | | | | | | | 55 | 3 |
| Carp..... | | | 390 | 35 | 598 | 28 | 128 | 6 | 411 | 3 |
| Catfish and bullheads..... | | | 76 | 6 | 982 | 32 | 4, 364 | 139 | 172 | 20 |
| Cero..... | | | 3 | (⁵) | | | 13 | 1 | | |
| Cigarfish..... | | | | | | | 9 | (⁵) | | |
| Cod..... | 99, 632 | 1, 856 | 7, 493 | 213 | 7 | (⁵) | 2 | (⁵) | 16, 036 | 156 |
| Corbina..... | | | | | | | | | (⁵) | (⁵) |
| Crappie..... | | | | | 1 | (⁵) | 405 | 12 | | |
| Crevalle..... | 2 | (⁵) | 7 | (⁵) | | | 25 | 1 | | |
| Croaker..... | 2, 497 | 36 | 2, 048 | 48 | 16, 042 | 211 | 4, 675 | 50 | | |
| Cunner..... | 42 | 1 | | | | | | | | |
| Cusk..... | 6, 109 | 73 | | | | | | | | |
| Dolphin..... | | | (⁵) | (⁵) | 1 | (⁵) | 12 | (⁵) | | |
| Drum: | | | | | | | | | | |
| Black..... | (⁵) | (⁵) | (⁵) | (⁵) | 123 | 2 | 1, 077 | 21 | | |
| Red or redfish..... | 2 | (⁵) | 9 | (⁵) | 59 | 2 | 2, 063 | 78 | | |

¹ Includes the fisheries of Lake Oksechobee, Fla.² Includes dip nets, push nets, reef nets, cast nets, scrap nets, and drag nets.³ Number not determined.⁴ Includes periwinkle, cockle, and fish pots; harpoons; spears; baskets and box traps.⁵ Less than 500 pounds or dollars.

Fisheries of the United States and Alaska—Continued

CATCH: BY SECTIONS—Continued

[Expressed in thousands of pounds and thousands of dollars; that is, 000 omitted]

| Species | New England, 1933 | | Middle Atlantic, 1933 | | Chesapeake, 1933 | | South Atlantic and Gulf, 1932 | | Pacific, 1933 | |
|------------------------------------|-------------------|-------|-----------------------|-------|------------------|-------|-------------------------------|-------|---------------|-------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| FISH | | | | | | | | | | |
| Eels: | | | | | | | | | | |
| Common..... | 618 | 35 | 680 | 60 | 312 | 16 | 65 | 2 | | |
| Conger..... | 41 | 1 | 16 | 1 | | | | | | |
| Flounders..... | 37,795 | 1,173 | 9,233 | 863 | 1,153 | 55 | 1,396 | 54 | 11,036 | 457 |
| Flyingfish..... | | | | | | | | | 16 | 1 |
| Frigate mackerel..... | 125 | 4 | 102 | 2 | | | 2 | (*) | | |
| Garfish..... | | | | | | | (*) | (*) | | |
| Gizzard shad..... | | | | | 144 | 3 | 19 | (*) | | |
| Goosefish..... | | | 10 | (*) | | | | | | |
| Graysfish..... | 13 | (*) | 7 | (*) | | | | | 471 | 9 |
| Groupers..... | | | | | | | 3,301 | 67 | | |
| Grunts..... | | | | | | | 51 | 2 | | |
| Haddock..... | 160,106 | 3,646 | 8,507 | 248 | (*) | (*) | | | | |
| Hake..... | 15,320 | 203 | 182 | 3 | 23 | 1 | 10 | (*) | 38 | (*) |
| Halibut..... | 2,457 | 236 | 53 | 7 | | | | | 24,497 | 1,568 |
| Hardhead..... | | | | | | | | | 157 | 8 |
| Harvestfish or "starfish"..... | | | | | 180 | 4 | 1,077 | 12 | | |
| Herring: | | | | | | | | | | |
| Round..... | | | 1 | (*) | | | | | | |
| Sea..... | 48,086 | 211 | 666 | 6 | | | | | 1,214 | 9 |
| Herring smelt..... | 12 | (*) | | | | | | | | |
| Hickory shad..... | 2 | (*) | 1 | (*) | 68 | 2 | 167 | 6 | | |
| Hogfish..... | | | | | | | 29 | 1 | | |
| Horse mackerel..... | | | | | | | | | 1,011 | 12 |
| Jewfish..... | | | | | | | 38 | 1 | | |
| Kingfish (California)..... | | | | | | | | | 664 | 12 |
| Kingfish or "king mackerel"..... | | | (*) | (*) | | | 3,301 | 120 | | |
| King whiting or "kingfish"..... | 40 | 1 | 158 | 15 | 74 | 2 | 652 | 13 | | |
| Ladyfish..... | | | | | | | 3 | (*) | | |
| Launce..... | 21 | (*) | | | | | | | | |
| "Lingcod"..... | | | | | | | | | 1,972 | 58 |
| Mackerel..... | 40,832 | 878 | 602 | 22 | 43 | 1 | | | 60,615 | 420 |
| Marlin..... | | | | | | | | | 7 | (*) |
| Menhaden..... | 1,029 | 3 | 79,576 | 236 | 115,990 | 386 | 89,346 | 132 | | |
| Mojarro..... | | | | | | | 36 | 1 | | |
| Mullet..... | 9 | (*) | 318 | 9 | 64 | 3 | 25,088 | 417 | 24 | 1 |
| Mummichog..... | | | 51 | 4 | | | | | | |
| Muttonfish..... | | | | | | | 203 | 9 | | |
| Paddlefish or "spoonbill cat"..... | | | | | | | 1 | (*) | | |
| Permit..... | | | | | | | 3 | (*) | | |
| Pigfish..... | 4 | (*) | (*) | (*) | 61 | (*) | 129 | 2 | | |
| Pike or pickerel (jacks)..... | | | (*) | (*) | 29 | 4 | 5 | (*) | | |
| Pilchard..... | | | | | | | | | 500,805 | 1,505 |
| Pinfish..... | | | | | | | 295 | 2 | | |
| Pollock..... | 15,027 | 164 | 776 | 12 | (*) | (*) | 590 | 81 | 5 | 2 |
| Pompano..... | | | (*) | (*) | 4 | (*) | | | | |
| Porkfish..... | | | | | | | (*) | (*) | | |
| Rock bass..... | | | | | | | | | 249 | 15 |
| Rockfishes..... | | | | | | | | | 5,138 | 164 |
| Rosefish..... | 264 | 3 | | | | | | | | |
| Rudderfish..... | | | | | | | | | 13 | 1 |
| Sablefish..... | | | | | | | | | 2,716 | 74 |
| Salmon: | | | | | | | | | | |
| Atlantic..... | 25 | 6 | | | | | | | | |
| Blueback, red or sockeye..... | | | | | | | | | 9,866 | 875 |
| Chinook or king..... | | | | | | | | | 35,114 | 2,105 |
| Chum or keta..... | | | | | | | | | 8,071 | 181 |
| Humpback or pink..... | | | | | | | | | 38,599 | 907 |
| Silver or coho..... | | | | | | | | | 15,042 | 600 |
| Sculpin..... | | | | | | | | | 68 | 4 |
| Scup or porgies..... | 4,105 | 89 | 6,359 | 92 | 1,589 | 38 | 279 | 7 | | |
| Sea bass..... | 3,999 | 121 | 2,475 | 87 | 342 | 10 | 704 | 24 | 449 | 18 |
| Sea bass, white (California)..... | | | | | | | | | 1,163 | 69 |
| Sea robin..... | 77 | 1 | 30 | (*) | (*) | (*) | | | | |
| Shad..... | 356 | 16 | 834 | 76 | 6,191 | 573 | 1,882 | 240 | 1,606 | 36 |
| Sharks..... | 66 | 1 | 12 | (*) | 10 | (*) | 5,051 | 12 | | |
| Sheepshead, salt-water..... | (*) | (*) | (*) | (*) | | | 673 | 14 | 59 | 1 |
| Silver perch..... | | | | | 1 | (*) | | | | |
| Silversides..... | | | 23 | 3 | | | | | | |
| Slates..... | 240 | 2 | 84 | 1 | | | | | 194 | 2 |
| Skipper or "billfish"..... | 5 | (*) | | | | | | | | |

* Less than 500 pounds or dollars.

Fisheries of the United States and Alaska—Continued

CATCH: BY SECTIONS—Continued

[Expressed in thousands of pounds and thousands of dollars; that is, 000 omitted]

| Species | New England, 1933 | | Middle Atlantic, 1933 | | Chesapeake, 1933 | | South Atlantic and Gulf, 1932 | | Pacific, 1933 | |
|-----------------------------------|----------------------|--------------|--------------------------|--------------|---------------------|--------------|----------------------------------|--------------|------------------------|---------------|
| | Quan- tity | Value | Quan- tity (*) | Value (*) | Quan- tity | Value | Quan- tity | Value | Quan- tity 2,580 | Value 64 |
| FISH—continued | | | | | | | | | | |
| Smelt..... | 551 | 60 | | | | | | | | |
| Snapper: | | | | | | | | | | |
| Mangrove..... | | | | | | | 96 | 2 | | |
| Red..... | | | | | | | 6,359 | 315 | | |
| Snook..... | | | | | | | 323 | 8 | | |
| Spanish mackerel..... | | | 7 | (*) | 68 | 4 | 6,465 | 217 | 4 | (*) |
| Spilltail..... | | | | | | | | | 17 | (*) |
| Spot..... | 33 | (*) | 528 | 8 | 747 | 19 | 1,680 | 20 | | (*) |
| Squawfish..... | | | | | | | | | 1 | (*) |
| Squeteague or "sea trout": | | | | | | | | | | |
| Gray..... | 369 | 14 | 7,874 | 227 | 13,464 | 263 | 3,992 | 74 | | |
| Spotted..... | 2 | (*) | (*) | (*) | 173 | 14 | 6,239 | 298 | | |
| Squirrel hake..... | | | 53 | (*) | | | | | | |
| Steelhead trout..... | | | | | | | | | 2,702 | 142 |
| Striped bass..... | 61 | 7 | 40 | 6 | 833 | 101 | 507 | 54 | 510 | 28 |
| Sturgeon..... | 8 | 1 | 29 | 3 | 9 | 2 | 45 | 5 | 91 | 2 |
| Suckers..... | 52 | 2 | 172 | 13 | 8 | (*) | (*) | (*) | 14 | (*) |
| Sunfish..... | | | 1 | (*) | 2 | (*) | 718 | 18 | | |
| Surfshines (perch)..... | | | | | | | | | 250 | 9 |
| Swallowfish..... | | | 2 | (*) | 1 | (*) | | | | |
| Swordfish..... | 3,381 | 404 | 103 | 13 | | | | | 851 | 71 |
| Tai..... | | | | | | | | | (*) | (*) |
| Tautog..... | 484 | 16 | 132 | 12 | 6 | (*) | | | | |
| Tenpounder..... | | | | | | | 80 | 1 | | |
| Thimble-eyed mackerel..... | 78 | 2 | 25 | (*) | 11 | (*) | | | | |
| Tilfish..... | 207 | 10 | 1,360 | 68 | | | | | | |
| Tomcod..... | 2 | (*) | 8 | (*) | | | | | 1 | (*) |
| Tripletail..... | | | | | | | 2 | (*) | | |
| Tuna and tunalike fishes: | | | | | | | | | | |
| Albacore..... | | | | | | | | | 2 | (*) |
| Bluefin or horse mackerel..... | 402 | 18 | 43 | 2 | | | 3 | (*) | 561 | 29 |
| Bonito..... | | | | | | | | | 2,252 | 40 |
| Skipjack or striped..... | | | | | | | | | 16,687 | 613 |
| Yellowfin..... | | | | | | | | | 51,076 | 2,275 |
| Turbot..... | | | | | | | 4 | (*) | | |
| Whitebait..... | | | 3 | 1 | | | | | 96 | 3 |
| Whitefish..... | | | | | | | | | 96 | 4 |
| White perch..... | 50 | 7 | 75 | 5 | 533 | 26 | 832 | 21 | | |
| Whiting..... | 9,419 | 97 | 2,147 | 21 | 1 | (*) | | | | |
| Wolfish..... | 2,197 | 32 | 26 | 1 | | | | | | |
| Yellow perch..... | 1 | (*) | 8 | 1 | 240 | 12 | 180 | 5 | | |
| Yellowtail..... | | | | | | | 92 | 4 | 3,899 | 88 |
| Miscellaneous fish..... | | | | | | | | | 151 | 2 |
| Total..... | 461,621 | 9,593 | 142,771 | 2,266 | 189,706 | 2,055 | 184,158 | 2,715 | 840,523 | 12,793 |
| SHELLFISH, ETC. | | | | | | | | | | |
| Abalone..... | | | | | | | | | 551 | 80 |
| Clams: | | | | | | | | | | |
| Coquina..... | | | | | | | 5 | (*) | | |
| Hard..... | 3,039 | 375 | 3,621 | 532 | 1,211 | 268 | 1,387 | 61 | 578 | 31 |
| Pismo..... | | | | | | | | | 28 | 6 |
| Razor..... | 358 | 16 | | | | | | | 587 | 84 |
| Soft..... | 9,431 | 476 | 1,135 | 64 | | | | | 63 | 14 |
| Surf..... | 59 | 4 | 526 | 24 | | | | | | |
| Mixed..... | | | | | | | | | 16 | 1 |
| Conchs..... | | | 35 | 2 | | | 2 | (*) | | |
| Crabs: | | | | | | | | | | |
| Hard..... | 5,629 | 40 | 955 | 25 | 50,559 | 616 | 8,484 | 89 | 6,176 | 355 |
| King..... | 7 | (*) | 2,750 | 7 | | | | | | |
| Soft..... | | | 94 | 20 | 5,517 | 382 | 413 | 60 | | |
| Stone..... | | | | | | | 154 | 8 | | |
| Crawfish..... | | | | | | | | | 99 | 10 |
| Lobsters: | | | | | | | | | | |
| Common..... | 9,088 | 1,608 | 724 | 138 | (*) | (*) | 445 | 32 | 1,050 | 145 |
| Spiny..... | | | | | | | | | (*) | (*) |
| Mussels, sea..... | 141 | 6 | 53 | 3 | | | | | | |
| Octopus..... | | | | | | | 1 | (*) | 56 | 3 |

* Less than 500 pounds or dollars.

Fisheries of the United States and Alaska—Continued

CATCH: BY SECTIONS—Continued

[Expressed in thousands of pounds and thousands of dollars; that is, 000 omitted]

| Species | New England, 1933 | | Middle Atlantic, 1933 | | Chesapeake, 1933 | | South Atlantic and Gulf, 1932 | | Pacific, 1933 | |
|--|-------------------|------------------|-----------------------|------------------|------------------|------------------|-------------------------------|------------------|---------------|------------------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| SHELLFISH ETC.—CON. | | | | | | | | | | |
| Oysters: | | | | | | | | | | |
| Eastern, market, public..... | 87 | 12 | 178 | 20 | 14,900 | 914 | 11,382 | 447 | | |
| Eastern, market, private..... | 5,070 | 735 | 13,755 | 1,428 | 10,253 | 812 | 4,368 | 264 | 60 | 23 |
| Western, market, Japanese, market..... | | | | | | | | | 226 | 120 |
| Periwinkles and "cockles"..... | 191 | 12 | (⁶) | (⁶) | | | | | 2,843 | 187 |
| Scallops: | | | | | | | | | | |
| Bay..... | 621 | 204 | 49 | 17 | | | 153 | 14 | 10 | 2 |
| Sea..... | 2,158 | 267 | 2,197 | 216 | 73 | 8 | | | | |
| Shrimp..... | 41 | 2 | 86 | 15 | (⁶) | (⁶) | 88,262 | 2,036 | 2,141 | 36 |
| Squid..... | 1,076 | 19 | 792 | 16 | 145 | 3 | 9 | (⁶) | 824 | 14 |
| Terrapin..... | | | 1 | (⁶) | 12 | 3 | 23 | 4 | | |
| Turtles..... | | | 11 | 1 | 2 | (⁶) | 1 | (⁶) | 3 | (⁶) |
| Frogs..... | | | | | | | | | | |
| Irish moss..... | 12 | 1 | | | | | | | | |
| Sponges..... | | | | | | | 612 | 697 | | |
| Bloodworms..... | 661 | 71 | 11 | 10 | | | | | | |
| Sandworms..... | 640 | 45 | 9 | 7 | | | | | | |
| Sea urchins..... | 6 | (⁶) | | | | | | | | |
| Total..... | 38,315 | 3,893 | 26,982 | 2,545 | 82,672 | 3,006 | 115,759 | 3,713 | 15,307 | 1,111 |
| WHALE PRODUCTS ⁷ | | | | | | | | | | |
| Meat..... | | | | | | | | | 2,214 | 43 |
| Oil, whale..... | | | | | | | | | 2,117 | 41 |
| Total..... | | | | | | | | | 4,331 | 84 |
| Grand total..... | 499,936 | 13,486 | 169,753 | 4,811 | 272,380 | 5,061 | 299,917 | 6,428 | 860,161 | 13,988 |

| Species | Lakes, 1932 | | Mississippi River and tributaries, 1931 | | Alaska, 1933 | | Total | |
|------------------------------|-------------|------------------|---|-------|--------------|-------|------------------|------------------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| FISH | | | | | | | | |
| Alewives..... | | | | | | | 36,698 | 199 |
| Amberjack..... | | | | | | | 7 | (⁶) |
| Anchovies..... | | | | | | | 317 | 4 |
| Barracuda..... | | | | | | | 3,077 | 123 |
| Black bass..... | | | | 14 | 1 | | 401 | 29 |
| Bluefish..... | | | | | | | 7,107 | 321 |
| Blue pike..... | 9,947 | 411 | | | | | 9,947 | 411 |
| Blue runner or hardtail..... | | | | | | | 163 | 2 |
| Bonito..... | | | | | | | 318 | 17 |
| Bowfin..... | 3 | (⁶) | 428 | 9 | | | 433 | 9 |
| Buffalo fish..... | 2 | (⁶) | 15,772 | 687 | | | 15,786 | 688 |
| Butterfish..... | | | | | | | 8,894 | 317 |
| Burbot..... | 331 | 4 | | | | | 331 | 4 |
| Cable or crab eater..... | | | | | | | 26 | 1 |
| Cabrilla..... | | | | | | | 85 | 3 |
| Carp..... | 4,294 | 119 | 11,892 | 456 | | | 17,433 | 647 |
| Catfish and bullheads..... | 833 | 46 | 10,267 | 878 | | | 16,894 | 1,121 |
| Cero..... | | | | | | | 16 | 1 |
| Chubs..... | 4,056 | 248 | | | | | 4,056 | 248 |
| Cigarfish..... | | | | | | | 9 | (⁶) |
| Cisco..... | 160 | 17 | | | | | 160 | 17 |
| Cod..... | | | | | 828 | 6 | 123,908 | 2,231 |
| Corbina..... | | | | | | | (⁶) | (⁶) |
| Crappie..... | 1 | (⁶) | 41 | 3 | | | 448 | 15 |
| Crevalle..... | | | | | | | 34 | 1 |
| Croaker..... | | | | | | | 25,262 | 345 |
| Cunner..... | | | | | | | 42 | 1 |
| Cusk..... | | | | | | | 6,113 | 73 |
| Dolly Varden trout..... | | | | | 44 | 2 | 44 | 2 |
| Dolphin..... | | | | | | | 13 | (⁶) |

⁶ Less than 500 pounds or dollars.

⁷ The weight of whales caught was not determined; therefore, the weight of the manufactured products is shown.

Fisheries of the United States and Alaska—Continued

CATCH: BY SECTIONS—Continued

[Expressed in thousands of pounds and thousands of dollars; that is, 000 omitted]

| Species | Lakes, 1932 | | Mississippi River and tributaries, 1931 | | Alaska, 1933 | | Total | |
|------------------------------------|-------------|-------|---|-------|--------------|-------|----------|-------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| FISH—continued | | | | | | | | |
| Drum: | | | | | | | | |
| Black..... | | | | | | | 1,200 | 23 |
| Red or redbfish..... | | | | | | | 2,153 | 80 |
| Eels: | | | | | | | | |
| Common..... | 43 | 1 | 7 | 1 | | | 1,625 | 115 |
| Conger..... | | | | | | | 57 | 2 |
| Flounders..... | | | | | 83 | 1 | 60,716 | 2,103 |
| Flyingfish..... | | | | | | | 16 | 1 |
| Frigate mackerel..... | | | | | | | 229 | 6 |
| Garfish..... | | | 73 | 1 | | | 73 | 1 |
| Gizzard shad..... | | | | | | | 163 | 3 |
| Goldfish..... | 49 | 1 | | | | | 49 | 1 |
| Goosefish..... | | | | | | | 10 | (*) |
| Grayfish..... | | | | | | | 491 | 9 |
| Groupers..... | | | | | | | 3,301 | 67 |
| Grunts..... | | | | | | | 51 | 2 |
| Haddock..... | | | | | | | 168,613 | 3,894 |
| Hake..... | | | | | | | 15,563 | 207 |
| Halibut..... | | | | | 15,632 | 726 | 42,639 | 2,537 |
| Hardhead..... | | | | | | | 157 | 8 |
| Harvestfish or "starfish"..... | | | | | | | 1,257 | 16 |
| Herring: | | | | | | | | |
| Round..... | | | | | | | 1 | (*) |
| Lake..... | 11,686 | 181 | | | | | 11,686 | 181 |
| Sea..... | | | | | 140,581 | 703 | 190,547 | 929 |
| Herring smelt..... | | | | | | | 12 | (*) |
| Hickory shad..... | | | | | | | 238 | 8 |
| Hogfish..... | | | | | | | 29 | 1 |
| Horse mackerel..... | | | | | | | 1,011 | 12 |
| Jewfish..... | | | | | | | 38 | 1 |
| Kingfish (California)..... | | | | | | | 564 | 12 |
| Kingfish or "king mackerel"..... | | | | | | | 3,301 | 120 |
| King whiting or "kingfish"..... | | | | | | | 924 | 31 |
| Ladyfish..... | | | | | | | 3 | (*) |
| Lake trout..... | 10,662 | 920 | | | | | 10,662 | 920 |
| Launce..... | | | | | | | 21 | (*) |
| "Lingcod"..... | | | | | | | 1,972 | 58 |
| Mackerel..... | | | | | | | 111,152 | 1,321 |
| Marlin..... | | | | | | | 7 | (*) |
| Menhaden..... | | | | | | | 285,941 | 757 |
| Minnows..... | | | 1 | (*) | | | 1 | (*) |
| Mojarro..... | | | | | | | 36 | 1 |
| Mooneye..... | 16 | (*) | 3 | (*) | | | 19 | (*) |
| Mullet..... | | | | | | | 25,503 | 430 |
| Mummichog..... | | | | | | | 51 | 4 |
| Muttonfish..... | | | | | | | 203 | 9 |
| Noddlefish or "spoonbill cat"..... | | | 952 | 43 | | | 953 | 43 |
| Permit..... | | | | | | | 3 | (*) |
| Pigfish..... | | | | | | | 194 | 2 |
| Pike or pickerel (jacks)..... | 373 | 16 | 5 | 1 | | | 412 | 21 |
| Pilchard..... | | | | | | | 509,805 | 1,505 |
| Pinfish..... | | | | | | | 295 | 2 |
| Pollock..... | | | | | | | 15,803 | 176 |
| Pompano..... | | | | | | | 599 | 83 |
| Porkfish..... | | | | | | | (*) | (*) |
| Quillback..... | | | 268 | 11 | | | 268 | 11 |
| Rock bass..... | 17 | 1 | | | | | 366 | 16 |
| Rockfishes..... | | | | | | | 5 | (*) |
| Rosefish..... | | | | | | | 5,143 | 164 |
| Rudderfish..... | | | | | | | 264 | 3 |
| Sablefish..... | | | | | 152 | 5 | 13 | 1 |
| Salmon: | | | | | | | 2,868 | 79 |
| Atlantic..... | | | | | | | 25 | 6 |
| Blueback, red or sockeye..... | | | | | 209,325 | 3,749 | 219,191 | 4,624 |
| Chinook or king..... | | | | | 12,175 | 264 | 47,289 | 2,849 |
| Chum or keta..... | | | | | 71,030 | 707 | 79,101 | 838 |
| Humpback or pink..... | | | | | 159,090 | 2,522 | 197,689 | 3,429 |
| Silver or coho..... | | | | | 15,729 | 256 | 30,771 | 856 |
| Sauger..... | 3,448 | 136 | 2 | (*) | | | 3,450 | 136 |
| Sculpin..... | | | | | | | 68 | 4 |
| Scup or porgies..... | | | | | | | 12,422 | 226 |
| Sea bass..... | | | | | | | 7,969 | 260 |
| Sea bass, white (California)..... | | | | | | | 1,163 | 69 |
| Sea robin..... | | | | | | | 107 | 1 |
| Shad..... | | | | | | | 10,899 | 941 |

* Less than 500 pounds or dollars.

Fisheries of the United States and Alaska—Continued

CATCH: BY SECTIONS—Continued

[Expressed in thousands of pounds and thousands of dollars; that is, 000 omitted]

| Species | Lakes, 1932 | | Mississippi River and tributaries, 1931 | | Alaska, 1933 | | Total | |
|--------------------------------|---------------|--------------|---|--------------|----------------|--------------|-------------------|---------------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity 6,139 | Value 13 |
| FISH—continued | | | | | | | | |
| Sharks..... | | | | | | | | |
| Sheepshead: | | | | | | | | |
| Fresh-water..... | 2,158 | 44 | 3,905 | 143 | | | 6,063 | 187 |
| Salt-water..... | | | | | | | 732 | 15 |
| Silver perch..... | | | | | | | 1 | (*) |
| Silversides..... | | | | | | | 23 | 3 |
| Skates..... | | | | | | | 518 | 5 |
| Skipper or "billfish"..... | | | | | | | 5 | (*) |
| Smelt..... | 98 | 3 | | | 1 | (*) | 2,210 | 127 |
| Snapper: | | | | | | | | |
| Mangrove..... | | | | | | | 96 | 2 |
| Red..... | | | | | | | 6,359 | 815 |
| Snook..... | | | | | | | 323 | 8 |
| Spanish mackerel..... | | | | | | | 6,544 | 221 |
| Spittail..... | | | | | | | 17 | (*) |
| Spot..... | | | | | | | 2,988 | 47 |
| Squawfish..... | | | | | | | 1 | (*) |
| Squeteague or "sea trout": | | | | | | | | |
| Gray..... | | | | | | | 25,699 | 578 |
| Spotted..... | | | | | | | 6,414 | 312 |
| Squirrel hake..... | | | | | | | 53 | (*) |
| Steelhead trout..... | 5 | 1 | | | 12 | (*) | 2,719 | 143 |
| Striped bass..... | | | | | | | 1,951 | 196 |
| Sturgeon..... | 30 | 6 | | | | | 212 | 19 |
| Sturgeon, shovelnose..... | | | 87 | 8 | | | 87 | 8 |
| Suckers..... | 6,192 | 137 | 315 | 13 | | | 6,768 | 165 |
| Sunfish..... | 8 | (*) | 22 | 1 | | | 761 | 19 |
| Surffishes (perch)..... | | | | | | | 260 | 9 |
| Swellfish..... | | | | | | | 3 | (*) |
| Swordfish..... | | | | | | | 4,336 | 488 |
| Tal..... | | | | | | | (*) | (*) |
| Tautog..... | | | | | | | 622 | 28 |
| Tenpounder..... | | | | | | | 80 | 1 |
| Thimble-eyed mackerel..... | | | | | | | 114 | 2 |
| Tilfish..... | | | | | | | 1,557 | 78 |
| Tomcod..... | | | | | | | 11 | (*) |
| Tripletail..... | | | | | | | 2 | (*) |
| Tullibee..... | 1,297 | 16 | | | | | 1,297 | 16 |
| Tuna and tunalike fishes: | | | | | | | | |
| Albacore..... | | | | | | | 2 | (*) |
| Bluefin or horse mackerel..... | | | | | | | 1,009 | 49 |
| Bonito..... | | | | | | | 2,252 | 40 |
| Skipjack or striped..... | | | | | | | 16,687 | 613 |
| Yellowfin..... | | | | | | | 51,076 | 2,275 |
| Turbot..... | | | | | | | 4 | (*) |
| White bass..... | 253 | 10 | 3 | (*) | | | 258 | 10 |
| Whitebait..... | | | | | | | 99 | 4 |
| Whitefish..... | 9,731 | 1,102 | | | | | 9,826 | 1,106 |
| Whitefish, Menominee..... | 233 | 17 | | | | | 233 | 17 |
| White perch..... | | | | | | | 1,490 | 59 |
| Whiting..... | | | | | | | 11,567 | 118 |
| Wolfish..... | | | | | | | 2,223 | 33 |
| Yellow perch..... | 11,472 | 467 | | | | | 11,901 | 485 |
| Yellow pike..... | 4,441 | 457 | 5 | 1 | | | 4,446 | 458 |
| Yellowtail..... | | | | | | | 3,991 | 92 |
| Miscellaneous fish..... | | | | | | | 151 | 2 |
| Total..... | 81,829 | 4,361 | 44,062 | 2,257 | 624,687 | 8,941 | 2,569,359 | 44,981 |
| SHELLFISH, ETC. | | | | | | | | |
| Abalone..... | | | | | | | 551 | 80 |
| Clams: | | | | | | | | |
| Coquina..... | | | | | | | 5 | (*) |
| Hard..... | | | | | 4 | (*) | 9,838 | 1,267 |
| Pismo..... | | | | | | | 26 | 6 |
| Razor..... | | | | | 1,042 | 49 | 1,987 | 149 |
| Soft..... | | | | | | | 10,629 | 554 |
| Surf..... | | | | | | | 585 | 28 |
| Mixed..... | | | | | | | 16 | 1 |
| Conchs..... | | | | | | | 37 | 2 |
| Crabs: | | | | | | | | |
| Hard..... | | | | | 1,063 | 78 | 72,886 | 1,203 |
| King..... | | | | | | | 2,757 | 7 |
| Soft..... | | | | | | | 6,024 | 462 |
| Stone..... | | | | | | | 154 | 8 |

* Less than 500 pounds or dollars.

Fisheries of the United States and Alaska—Continued

CATCH: BY SECTIONS—Continued

[Expressed in thousands of pounds and thousands of dollars; that is, 000 omitted]

| Species | Lakes, 1932 | | Mississippi River and tributaries, 1931 | | Alaska, 1933 | | Total | |
|----------------------------------|---------------|--------------|---|--------------|----------------|--------------|------------------|---------------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| SHELLFISH, ETC.—continued | | | | | | | | |
| Crawfish..... | 20 | 1 | 29 | (*) | | | 148 | 11 |
| Lobsters: | | | | | | | | |
| Common..... | | | | | | | 9,812 | 1,746 |
| Spiny..... | | | | | | | 1,495 | 177 |
| Mussels, sea..... | | | | | | | 194 | 9 |
| Mussel shells..... | 1,895 | 26 | 37,255 | 421 | | | 39,150 | 447 |
| Octopus..... | | | | | | | 57 | 3 |
| Oysters: | | | | | | | | |
| Eastern, market, public..... | | | | | | | 26,547 | 1,393 |
| Eastern, market, private..... | | | | | | | 33,506 | 3,282 |
| Western, market..... | | | | | | | 226 | 120 |
| Japanese, market..... | | | | | | | 2,843 | 187 |
| Periwinkles and "cockles"..... | | | | | | | 191 | 12 |
| Scallops: | | | | | | | | |
| Bay..... | | | | | | | 833 | 237 |
| Sea..... | | | | | | | 4,428 | 491 |
| Shrimp..... | | | 49 | 4 | 580 | 21 | 91,159 | 2,114 |
| Squid..... | | | | | | | 2,846 | 52 |
| Terrapin..... | | | 19 | 1 | | | 55 | 8 |
| Turtles..... | | | 94 | 3 | | | 168 | 5 |
| Frogs..... | | | 875 | 131 | | | 876 | 131 |
| Irish moss..... | | | | | | | 12 | 1 |
| Sponges..... | | | | | | | 612 | 697 |
| Bloodworms..... | | | | | | | 672 | 81 |
| Sandworms..... | | | | | | | 649 | 52 |
| Pearls and slugs..... | | 1 | | 80 | | | | 81 |
| Sea urchins..... | | | | | | | 6 | (*) |
| Total..... | 1,915 | 28 | 38,321 | 640 | 2,709 | 148 | 321,980 | 15,084 |
| WHALE PRODUCTS | | | | | | | | |
| Meat..... | | | | | | | 2,214 | 43 |
| Fertilizer..... | | | | | 1,034 | 14 | 1,034 | 14 |
| Oil, sperm..... | | | | | 84 | 2 | 84 | 2 |
| Oil, whale..... | | | | | 2,290 | 53 | 4,377 | 94 |
| Total..... | | | | | 3,378 | 69 | 7,709 | 153 |
| Grand total..... | 83,744 | 4,389 | 82,383 | 2,897 | 630,774 | 9,158 | 2,899,048 | 60,218 |

CATCH: BY STATES *

[Expressed in thousands of pounds and thousands of dollars; that is, 000 omitted]

| States | Marine and coastal rivers | | Mississippi River and tributaries | | Lakes * | | Total | |
|------------------|---------------------------|-------|-----------------------------------|-------|----------|-------|----------|-------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| Alabama..... | 6,107 | 169 | 1,822 | 33 | | | 7,929 | 202 |
| Arkansas..... | | | 15,733 | 412 | | | 15,733 | 412 |
| California..... | 706,899 | 7,094 | | | | | 706,899 | 7,094 |
| Connecticut..... | 9,878 | 613 | | | | | 9,878 | 613 |
| Delaware..... | 36,527 | 209 | | | | | 36,527 | 209 |
| Florida..... | 101,920 | 2,917 | | | | | 103,290 | 2,973 |
| Georgia..... | 16,523 | 186 | | | 1,370 | 56 | 16,523 | 186 |
| Illinois..... | | | 14,263 | 367 | 885 | 58 | 15,148 | 425 |
| Indiana..... | | | 7,718 | 157 | 630 | 37 | 8,348 | 194 |
| Iowa..... | | | 7,778 | 303 | | | 7,778 | 303 |
| Kansas..... | | | 455 | 17 | | | 455 | 17 |
| Kentucky..... | | | 1,622 | 61 | | | 1,622 | 61 |
| Louisiana..... | 48,340 | 1,181 | 19,213 | 994 | | | 67,553 | 2,175 |
| Maine..... | 98,498 | 2,307 | | | | | 98,498 | 2,307 |

* Less than 500 pounds or dollars.

* The catch under "Marine and coastal rivers" is for 1933 except in the South Atlantic and Gulf States which is for 1932; the catch of the "Mississippi River and tributaries" is for 1931; and the catch of the "Lakes" is for 1932.

* Includes Lake Ontario, Lake Erie, Lake Huron, Lake Michigan, Lake Superior, Rainy Lake, Namakan Lake, Lake of the Woods, Lake Okeechobee, and several mussel-bearing streams tributary to Lakes Huron, Erie, and Michigan.

Fisheries of the United States and Alaska—Continued

CATCH: BY STATES

[Expressed in thousands of pounds and thousands of dollars; that is, 000 omitted]

| States | Marine and coastal rivers | | Mississippi River and tributaries | | Lakes | | Total | |
|---------------------|---------------------------|--------|-----------------------------------|-------|----------|-------|-----------|--------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| Maryland..... | 55,362 | 1,734 | | | | | 55,362 | 1,734 |
| Massachusetts..... | 373,670 | 9,507 | | | | | 373,670 | 9,507 |
| Michigan..... | | | | | 30,130 | 2,162 | 30,130 | 2,162 |
| Minnesota..... | | | 3,498 | 138 | 8,507 | 199 | 12,005 | 337 |
| Mississippi..... | 20,603 | 497 | 2,650 | 123 | | | 23,253 | 620 |
| Missouri..... | | | 928 | 77 | | | 928 | 77 |
| Nebraska..... | | | 145 | 16 | | | 145 | 16 |
| New Hampshire..... | 523 | 57 | | | | | 523 | 57 |
| New Jersey..... | 93,263 | 2,140 | | | | | 93,263 | 2,140 |
| New York..... | 30,911 | 2,453 | | | 1,435 | 112 | 41,346 | 2,565 |
| North Carolina..... | 86,214 | 827 | | | | | 86,214 | 827 |
| Ohio..... | | | 185 | 7 | 28,516 | 1,161 | 28,701 | 1,168 |
| Oklahoma..... | | | 40 | 4 | | | 40 | 4 |
| Oregon..... | 24,533 | 1,210 | | | | | 24,533 | 1,210 |
| Pennsylvania..... | 52 | 3 | | | 2,535 | 167 | 2,587 | 170 |
| Rhode Island..... | 17,366 | 1,001 | | | | | 17,366 | 1,001 |
| South Carolina..... | 4,536 | 123 | | | | | 4,536 | 123 |
| South Dakota..... | | | 114 | 11 | | | 114 | 11 |
| Tennessee..... | | | 3,435 | 104 | | | 3,435 | 104 |
| Texas..... | 14,304 | 472 | 139 | 6 | | | 14,443 | 478 |
| Virginia..... | 217,018 | 3,327 | | | | | 217,018 | 3,327 |
| Washington..... | 128,730 | 5,684 | | | | | 128,730 | 5,684 |
| Wisconsin..... | | | 2,645 | 68 | 11,107 | 493 | 13,752 | 561 |
| Alaska..... | 630,773 | 9,158 | | | | | 630,773 | 9,158 |
| Total..... | 2,731,550 | 52,875 | 82,383 | 2,398 | 85,115 | 4,446 | 2,899,048 | 60,218 |

SEED OYSTER FISHERY

| Item | New England, 1933 | | Middle Atlantic, 1933 | |
|----------------------------|-------------------|--------------|-----------------------|--------------|
| | Number | | Number | |
| OPERATING UNITS | | | | |
| Fishermen: | | | | |
| On vessels..... | 93 | | 1,586 | |
| On boats and shore: | | | | |
| Regular..... | 5 | | 298 | |
| Casual..... | 160 | | 150 | |
| Total..... | 258 | | 2,034 | |
| Vessels: | | | | |
| Steam..... | 3 | | | |
| Net tonnage..... | 245 | | | |
| Motor..... | 18 | | 7 | |
| Net tonnage..... | 271 | | 69 | |
| Sail..... | 2 | | 146 | |
| Net tonnage..... | 17 | | 2,977 | |
| Total vessels..... | 23 | | 163 | |
| Total net tonnage..... | 533 | | 3,046 | |
| Boats: | | | | |
| Motor..... | 7 | | 206 | |
| Other..... | 90 | | 152 | |
| Apparatus: | | | | |
| Dredges, oyster..... | 101 | | 300 | |
| Yards at mouth..... | 97 | | 363 | |
| Tongs..... | 96 | | 374 | |
| Rakes..... | 46 | | 79 | |
| CATCH | | | | |
| Oysters: | <i>Bushels</i> | <i>Value</i> | <i>Bushels</i> | <i>Value</i> |
| Seed, public, spring..... | 31,458 | \$12,884 | 1,003,540 | \$261,862 |
| Seed, public, fall..... | 43,183 | 17,285 | 23,500 | 8,250 |
| Seed, private, spring..... | 207,185 | 69,617 | 40,290 | 39,540 |
| Seed, private, fall..... | | | 38,290 | 38,290 |
| Total..... | 281,806 | 99,566 | 1,105,620 | 347,942 |

Fisheries of the United States and Alaska—Continued

SEED OYSTER FISHERY—Continued

| Item | Chesapeake, 1933 | | South Atlantic and Gulf, 1932 | | Total | |
|----------------------------|------------------|--------------|-------------------------------|--------------|----------------|--------------|
| | Number | | Number | | Number | |
| OPERATING UNITS | | | | | | |
| Fishermen: | | | | | | |
| On vessels..... | | | | | 1,679 | |
| On boats and shore: | | | | | | |
| Regular..... | 1,650 | | 12 | | 1,665 | |
| Casual..... | 67 | | | | 377 | |
| Total..... | 1,717 | | 12 | | 4,021 | |
| Vessels: | | | | | | |
| Steam: | | | | | 3 | |
| Net tonnage..... | | | | | 245 | |
| Motor..... | | | | | 26 | |
| Net tonnage..... | | | | | 340 | |
| Sail..... | | | | | 148 | |
| Net tonnage..... | | | | | 2,994 | |
| Total vessels..... | | | | | 176 | |
| Total net tonnage..... | | | | | 3,879 | |
| Boats: | | | | | | |
| Motor..... | 804 | | 6 | | 1,023 | |
| Other..... | 286 | | | | 628 | |
| Apparatus: | | | | | | |
| Dredges, oyster..... | | | 12 | | 413 | |
| Yards at mouth..... | | | 12 | | 472 | |
| Tongs..... | 1,474 | | | | 1,944 | |
| Rakes..... | 130 | | | | 265 | |
| CATCH | | | | | | |
| Oysters: | <i>Bushels</i> | <i>Value</i> | <i>Bushels</i> | <i>Value</i> | <i>Bushels</i> | <i>Value</i> |
| Seed, public, spring..... | 574,620 | \$60,880 | | \$8,280 | 1,649,359 | \$343,706 |
| Seed, public, fall..... | 815,240 | 85,934 | | | 881,903 | 111,449 |
| Seed, private, spring..... | 35,600 | 7,720 | | | 283,075 | 116,877 |
| Seed, private, fall..... | 56,800 | 11,360 | | | 95,090 | 49,650 |
| Total..... | 1,482,260 | 165,894 | 39,741 | 8,280 | 2,909,427 | 621,682 |

NOTE.—Of the number of persons fishing for seed oysters, a total of 3,652 are duplicated among those fishing for market oysters or other species. Similarly, the following craft and gear are duplicated: 87 vessels, 823 motor boats, 437 other boats, 172 dredges, 1,570 tongs, and 61 rakes.

Yield of the fisheries of the United States: By gear

| Gear | New England, 1933 | | Middle Atlantic, 1933 | | Chesapeake, 1933 | |
|--------------------------|-------------------|------------|-----------------------|-----------|------------------|-----------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Purse seines..... | 56,586,422 | \$796,362 | 77,945,874 | \$205,603 | 115,343,100 | \$384,227 |
| Haul seines..... | 910,017 | 32,041 | 1,845,567 | 70,668 | 3,669,524 | 154,091 |
| Gill nets..... | 18,593,520 | 359,016 | 2,517,827 | 117,750 | 1,846,636 | 147,762 |
| Lines..... | 99,382,952 | 2,167,473 | 7,633,174 | 292,745 | 42,619,655 | 489,041 |
| Pound nets..... | 14,419,422 | 227,622 | 28,325,529 | 692,557 | 62,619,356 | 1,215,833 |
| Floating traps..... | 8,458,029 | 159,754 | | | | |
| Other traps..... | 28,910 | 2,512 | | | | |
| Weirs..... | 21,801,005 | 92,599 | 898,336 | 2,485 | | |
| Stop nets..... | | | 208,874 | 20,472 | 63,504 | 2,308 |
| Fyke nets..... | 81,216 | 3,192 | 694,092 | 21,736 | 923,307 | 40,709 |
| Dip nets..... | 3,640,232 | 45,571 | 86,022 | 16,852 | 4,662,036 | 305,638 |
| Cast nets..... | | | 400 | 48 | | |
| Scap nets..... | | | 149,664 | 8,574 | | |
| Bag and pocket nets..... | 116,068 | 13,039 | 51,900 | 10,855 | | |
| Drag nets..... | 18,000 | 7,000 | | | | |
| Push nets..... | 235,003,844 | 5,290,742 | 24,410,956 | 780,076 | 5,455,375 | 129,812 |
| Other trawls..... | 15,081,503 | 1,673,370 | 12,369,400 | 1,203,269 | 167,225 | 9,452 |
| Pots..... | 3,481,594 | 406,727 | 102,388 | 12,549 | | |
| Harpoons..... | 106,663 | 7,537 | 128,568 | 12,662 | 6,000 | 300 |
| Spears..... | | | | | 2,001,916 | 65,929 |
| Scrapes, crab..... | | | | | 13,573,634 | 638,012 |
| Dredges..... | 8,311,071 | 1,224,218 | 17,238,166 | 1,698,835 | 18,357,340 | 1,309,318 |
| Tongs..... | 1,443,961 | 202,993 | 2,618,049 | 372,867 | 690,140 | 75,133 |
| Rakes..... | 1,129,890 | 145,761 | 1,281,227 | 179,516 | | |
| Forks..... | 2,811,825 | 283,046 | 171,093 | 29,770 | | |
| Hoes..... | 8,378,980 | 333,750 | 913,580 | 45,752 | | |
| Picks..... | | | 750 | 75 | 279,864 | 69,494 |
| Gaffs..... | | | | | 101,440 | 23,800 |
| By hand..... | 151,016 | 5,225 | 169,331 | 15,519 | | |
| Total..... | 499,936,139 | 13,485,550 | 169,753,735 | 4,811,055 | 272,380,052 | 5,080,829 |

Yield of the fisheries of the United States: By gear—Continued

| Gear | South Atlantic and Gulf, 1932 | | Pacific, 1933 | | Lakes, 1932 | |
|---|-------------------------------|-----------|------------------|------------------|-------------|-----------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Purse seines..... | 89,971,248 | \$163,404 | 423,264,286 | \$2,672,000 | | |
| Haul seines..... | 24,175,742 | 602,678 | 3,824,178 | 199,434 | 4,538,616 | \$129,882 |
| Gill nets..... | 38,137,303 | 948,195 | 30,262,256 | 1,369,692 | 32,046,421 | 1,766,456 |
| Tammel nets..... | 3,159,789 | 118,055 | 661,335 | 41,492 | 205,546 | 4,140 |
| Lines..... | 25,218,356 | 764,227 | 136,182,745 | 5,680,665 | 2,307,481 | 206,357 |
| Pound nets..... | 10,763,098 | 190,839 | 26,648,553 | 1,297,707 | 8,207,975 | 466,941 |
| Traps (other than floating)..... | | | | | 30,068,291 | 1,628,039 |
| Weirs..... | 2,000 | 20 | 467,200 | 4,672 | | |
| Wheels..... | 283,000 | 1,505 | 763,235 | 56,339 | | |
| Stop nets..... | 891,793 | 15,055 | | | | |
| Fyke nets..... | 345,520 | 10,661 | 197,393 | 20,701 | 3,425,469 | 161,008 |
| Dip nets..... | 491,388 | 27,215 | 1,848,148 | 51,968 | | |
| Drag bag nets..... | | | 136,557 | 5,838 | | |
| Cast nets..... | 108,932 | 3,994 | | | | |
| Bag and pocket nets..... | | | 1,494,768 | 22,422 | | |
| Reef nets..... | | | 177,315 | 5,612 | | |
| Lampara nets..... | | | 203,895,501 | 863,526 | | |
| Paranzella nets..... | | | 11,549,104 | 451,971 | | |
| Otter trawls..... | 87,089,274 | 2,016,418 | 789,873 | 21,061 | | |
| Beam trawls..... | | | 656,187 | 15,548 | | |
| Pots..... | 1,117,738 | 50,446 | 7,264,551 | 506,478 | 19,677 | 984 |
| Harpoons..... | | | 5,190,295 | 155,663 | | |
| Spears..... | 155,803 | 9,036 | | | | |
| Drudges..... | 7,373,518 | 297,347 | (¹) | (²) | | |
| Tongs..... | 6,049,736 | 368,583 | \$ 4,396,438 | \$ 464,770 | | |
| Crowfoot bars..... | | | (¹) | (¹) | 1,468,430 | 21,071 |
| Rakes..... | 312,476 | 20,013 | | | | |
| Forks..... | 281,340 | 11,356 | | | | |
| Grabs..... | 2,692,154 | 56,533 | | | | |
| Picks..... | | | | | 246,966 | 8,647 |
| Hooks..... | 335,203 | 234,967 | | | | |
| Dividing apparatus, abalone and sponge..... | 278,824 | 462,077 | 351,268 | 80,433 | | |
| By hand..... | 701,463 | 50,861 | | | 179,518 | 2,536 |
| Total..... | 299,916,728 | 6,428,385 | 860,161,216 | 13,987,992 | 83,744,389 | 4,389,061 |

| Gear | Mississippi River and tributaries, 1931 | | Total | |
|---|---|-----------|---------------|-------------|
| | Pounds | Value | Pounds | Value |
| Purse seines..... | | | 763,050,930 | \$4,221,496 |
| Haul seines..... | 13,739,657 | \$574,641 | 52,703,291 | 1,763,235 |
| Gill nets..... | 166,598 | 6,547 | 123,570,561 | 4,710,418 |
| Tammel nets..... | 1,134,206 | 75,615 | 6,160,875 | 239,302 |
| Lines..... | 10,140,037 | 772,245 | 323,484,400 | 10,372,753 |
| Pound nets..... | 224,275 | 9,541 | 162,208,208 | 4,101,040 |
| Floating traps..... | | | 8,458,029 | 159,754 |
| Other traps..... | | | 30,127,301 | 1,628,551 |
| Weirs..... | | | 23,163,541 | 69,776 |
| Wheels..... | | | 1,026,235 | 57,544 |
| Stop nets..... | | | 1,164,171 | 37,835 |
| Fyke nets..... | 18,507,204 | 797,130 | 24,174,201 | 1,056,137 |
| Dip nets..... | 30,045 | 3,307 | 10,757,871 | 450,581 |
| Drag bag nets..... | | | 136,687 | 6,838 |
| Cast nets..... | | | 109,332 | 4,042 |
| Scap nets..... | | | 149,664 | 8,574 |
| Bag and pocket nets..... | | | 1,610,838 | 36,461 |
| Drag nets..... | | | 51,900 | 10,855 |
| Push nets..... | | | 18,000 | 7,000 |
| Reef nets..... | | | 177,315 | 5,612 |
| Lampara nets..... | | | 203,895,501 | 863,526 |
| Paranzella nets..... | | | 11,549,104 | 451,971 |
| Otter trawls..... | | | 352,749,322 | 8,244,109 |
| Beam trawls..... | | | 656,187 | 15,548 |
| Pots..... | 1,310,455 | 1,26,277 | 26,330,549 | 2,470,276 |
| Harpoons..... | | | 8,774,275 | 574,939 |
| Spears..... | 2,250 | 270 | 397,284 | 29,825 |
| Scrapes, crab..... | | | 2,001,916 | 65,929 |
| Drudges..... | 3,699,100 | 40,958 | 50,195,489 | 3,699,370 |
| Tongs..... | 1,601,876 | 21,091 | 34,467,400 | 2,739,622 |
| Crowfoot bars..... | 20,893,560 | 265,443 | 22,361,980 | 286,514 |
| Rakes..... | 370,130 | 4,029 | 3,783,863 | 425,352 |
| Forks..... | 4,812,737 | 76,214 | 8,076,995 | 400,386 |
| Hoops..... | | | 9,292,540 | 279,502 |
| Grabs..... | 873,099 | 130,621 | 3,566,253 | 187,154 |
| Picks..... | | | 526,830 | 78,111 |
| Hooks..... | | | 335,203 | 284,967 |
| Dividing apparatus, abalone and sponge..... | | | 830,092 | 642,510 |
| Gaffs..... | | | 750 | 75 |
| By hand..... | 5,877,304 | 93,528 | 7,180,101 | 200,469 |
| Total..... | 82,382,523 | 2,897,357 | 2,268,274,782 | 61,060,229 |

¹ Includes the catch by baskets.

² The catch by shovels, rakes, and dredges is included with tongs.

Industries related to the fisheries of the United States and Alaska

| Item | New England, 1933 | Middle Atlantic, 1933 | Chesapeake, 1933 | South Atlantic and Gulf, 1931 |
|---|-------------------|-----------------------|------------------|-------------------------------|
| Transporting: | | | | |
| Persons engaged: | | | | |
| On vessels..... | 256 | 110 | 869 | 354 |
| On boats..... | 54 | 92 | 25 | 389 |
| Total..... | 310 | 202 | 894 | 743 |
| Vessels: | | | | |
| Steam..... | 1 | | | |
| Net tonnage..... | 67 | | | |
| Motor..... | 104 | 30 | 421 | 152 |
| Net tonnage..... | 1,615 | 629 | 4,929 | 1,440 |
| Sail..... | | | 11 | 30 |
| Net tonnage..... | | | 307 | 279 |
| Total vessels..... | 105 | 30 | 432 | 182 |
| Total net tonnage..... | 1,682 | 629 | 5,236 | 1,719 |
| Boats..... | 31 | 87 | 102 | 251 |
| Wholesale and manufacturing: | | | | |
| Establishments..... | 362 | 398 | 502 | 585 |
| Persons engaged: | | | | |
| Proprietors..... | 266 | 358 | 664 | 700 |
| Salaried employees..... | 799 | 1,067 | 315 | 352 |
| Wage earners: | | | | |
| Average for season..... | 8,112 | 4,206 | 10,617 | 12,583 |
| Average for year..... | 4,170 | 2,948 | 4,704 | 4,581 |
| Salaries and wages paid..... | \$5,410,072 | \$6,085,981 | \$2,366,762 | \$2,821,543 |
| Manufactured products *..... | \$14,322,274 | \$10,603,564 | \$7,245,169 | \$7,680,627 |
| Fishermen's manufactured products: | | | | |
| Persons engaged..... | 2,223 | 512 | 14 | 1,379 |
| Products *..... | \$553,235 | \$260,146 | \$3,548 | \$258,805 |

| Item | Pacific, 1933 | Lakes, 1931 | Mississippi River and tributaries, 1931 | Alaska, 1933 | Total |
|---|---------------|-------------|---|--------------|---------------|
| Transporting: | | | | | |
| Persons engaged: | | | | | |
| On vessels..... | 245 | 19 | 29 | 1,283 | 3,165 |
| On boats..... | | | | (1) | 560 |
| Total..... | 245 | 19 | 29 | 1,283 | 3,725 |
| Vessels: | | | | | |
| Steam..... | | | | 10 | 11 |
| Net tonnage..... | | | | 18,367 | 18,434 |
| Motor..... | 104 | 9 | 8 | 282 | 1,110 |
| Net tonnage..... | 2,328 | 120 | 104 | 10,982 | 22,147 |
| Sail..... | | | | | 41 |
| Net tonnage..... | | | | | 586 |
| Total vessels..... | 104 | 9 | 8 | 292 | 1,162 |
| Total net tonnage..... | 2,328 | 120 | 104 | 29,349 | 41,167 |
| Boats..... | | | | 618 | 1,089 |
| Wholesale and manufacturing: | | | | | |
| Establishments..... | 313 | 230 | 217 | 224 | 2,831 |
| Persons engaged: | | | | | |
| Proprietors..... | 347 | 225 | 204 | | |
| Salaried employees..... | 747 | 471 | 355 | | |
| Wage earners: | | | | 11,756 | 70,824 |
| Average for season..... | 10,899 | 1,506 | 4,275 | | |
| Average for year..... | 3,667 | 1,034 | 3,483 | (*) | (*) |
| Salaries and wages paid..... | \$6,095,492 | \$2,610,439 | \$3,080,430 | (*) | (*) |
| Manufactured products *..... | \$28,772,185 | \$1,891,558 | \$3,723,414 | \$31,099,801 | \$105,338,592 |
| Fishermen's manufactured products: | | | | | |
| Persons engaged..... | 168 | 106 | 216 | (*) | (*) |
| Products *..... | \$174,669 | \$63,844 | \$8,761 | (*) | (*) |

* Included in vessels.

† Includes scows, houseboats, pile drivers, etc.

‡ Statistics not available.

§ Includes packaged, cured, and canned fishery products and also byproducts.

¶ Includes data for 1933 on packaged and canned products and byproducts.

NOTE.—Of the total number of persons engaged in the preparation of fishermen's manufactured products, 4,029 have also been included as fishermen and 866 of the persons shown on transporting craft have also been included as fishermen.

MANUFACTURED FISHERY PRODUCTS

Statistics of the output of the various manufactured fishery products have been collected in considerable more detail for 1930, 1931, and 1933 than in previous years. In addition to statistics of the production of cured fishery products, canned fishery products, and fishery byproducts, data also have been collected on the output of packaged fishery products, including shucked oysters, fresh-cooked crab meat, and similar commodities, in the more recent surveys.

Since only a part of the United States was surveyed for general fishery statistics for 1933, the following compilation of manufactured fishery products consists of composite data based on the most recent statistics. The years covered by the data are indicated by footnotes.

*Manufactured fishery products of the United States and Alaska*¹

| Item | Quantity | Value |
|---|------------|-----------|
| Alewives: | | |
| Salted: | | |
| Canned ¹pounds.. | 3,821,100 | \$59,522 |
| Tight-pack-cut.....do.. | 5,353,868 | 130,237 |
| Tight-pack roe ²do.. | 395,180 | 13,924 |
| Pickled ³do.. | 2,118,700 | 69,537 |
| Smoked ³do.. | 272,435 | 10,425 |
| Canned.....standard cases.. | 20,280 | 46,363 |
| Roe, canned.....do.. | 28,513 | 111,644 |
| Dry scrap.....do.. | 612 | 15,777 |
| Oil.....gallons.. | 30,900 | 3,518 |
| Bluefish, smoked.....pounds.. | 259 | 46 |
| Buffalo fish, smoked ³do.. | 469,700 | 135,855 |
| Butterfish, smoked ³do.. | 631,427 | 165,070 |
| Cabrilla, fresh fillets.....do.. | 18,000 | 2,700 |
| Carp, smoked ³do.. | 173,822 | 53,556 |
| Cisco, chubs, tullibee, and lake herring, smoked ³do.. | 6,601,825 | 1,403,953 |
| Cod: | | |
| Fresh fillets.....do.. | 7,316,708 | 847,413 |
| Frozen fillets.....do.. | 3,971,227 | 344,676 |
| Fresh steaks and sticks.....do.. | 288,778 | 49,434 |
| Salted: | | |
| Green.....do.. | 2,927,287 | 231,678 |
| Dry.....do.. | 1,446,514 | 84,064 |
| Boneless, including absolutely boneless.....do.. | 9,517,111 | 1,645,689 |
| Stockfish.....do.. | 31,220 | 3,085 |
| Smoked fillets.....do.. | 1,128,228 | 167,650 |
| Other smoked.....do.. | 10,440 | 1,116 |
| Oil: | | |
| Cod.....gallons.. | 45,331 | 22,152 |
| Cod liver.....do.. | 19,552 | 14,708 |
| Cronker, fresh fillets.....pounds.. | 30,500 | 3,720 |
| Cusk: | | |
| Fresh fillets.....do.. | 321,566 | 40,144 |
| Fresh steaks and sticks.....do.. | 140,177 | 25,387 |
| Salted: | | |
| Green.....do.. | 2,611,650 | 1,288 |
| Dry.....do.. | 33,972 | 1,593 |
| Boneless, including absolutely boneless.....do.. | 28,835 | 3,469 |
| Smoked fillets.....do.. | 195,803 | 25,334 |
| Eels: | | |
| Smoked ³do.. | 115,504 | 31,078 |
| Pickled.....do.. | 48 | 14 |
| Flounders: | | |
| Fresh fillets.....do.. | 3,488,222 | 532,812 |
| Frozen fillets.....do.. | 267,524 | 35,539 |
| Smoked fillets ²do.. | 5,331 | 1,072 |
| Groupers, fresh steaks and fillets.....do.. | 261,800 | 27,287 |
| Haddock: | | |
| Fresh fillets.....do.. | 16,693,543 | 1,986,554 |
| Frozen fillets.....do.. | 18,380,191 | 1,457,883 |
| Fresh sticks.....do.. | 69,168 | 12,559 |

¹ Data are for 1933 unless otherwise indicated.

² This item represents a combination of 1933 data in sections where available, and 1931 data in other sections.

³ This is usually an intermediate product and although included in the total, may also be shown in its final stage of processing elsewhere in the table.

Manufactured fishery products of the United States and Alaska—Continued

| Item | Quantity | Value | |
|--|---------------------|--------------|-------------|
| Haddock—Continued. | | | |
| Salted: | | | |
| Green..... | pounds..... | \$ 45, 765 | \$1, 091 |
| Dry..... | do..... | 52, 032 | 2, 550 |
| Boneless, including absolutely boneless..... | do..... | 98, 295 | 19, 220 |
| Smoked filets..... | do..... | 149, 214 | 26, 928 |
| Finn in haddie..... | do..... | 1, 171, 364 | 134, 818 |
| Canned finnan haddie..... | standard cases..... | 329 | 3, 505 |
| Bake: | | | |
| Fresh filets..... | pounds..... | 1, 386, 900 | 125, 891 |
| Frozen filets..... | do..... | 28, 821 | 2, 361 |
| Fresh sticks..... | do..... | 200, 630 | 29, 454 |
| Salted: | | | |
| Green..... | do..... | \$ 408, 653 | 11, 195 |
| Dry..... | do..... | 277, 670 | 8, 896 |
| Boneless, including absolutely boneless..... | do..... | 681, 201 | 47, 851 |
| Smoked filets..... | do..... | 33, 282 | 4, 586 |
| Halibut: | | | |
| Fresh filets..... | do..... | 109, 484 | 23, 408 |
| Frozen filets..... | do..... | 17, 480 | 3, 012 |
| Herring, lake, salted ¹ | do..... | 4, 262, 250 | 102, 925 |
| Herring, sea: | | | |
| Salted: | | | |
| Roused..... | do..... | 68, 400 | 1, 790 |
| Spiced..... | do..... | 509, 790 | 17, 474 |
| Pickled ² | do..... | 189, 758 | 10, 120 |
| Scotch-cure..... | do..... | 3, 256, 200 | 59, 244 |
| Norwegian-cure..... | do..... | 12, 651, 328 | 686, 331 |
| Dry-salted..... | do..... | 290, 650 | 14, 019 |
| Smoked: | do..... | 54, 200 | 2, 020 |
| Bloaters, hard..... | do..... | 948, 642 | 45, 853 |
| Bloaters, soft..... | do..... | 886, 904 | 64, 656 |
| Bloaters, unclassified..... | do..... | 121, 620 | 12, 971 |
| Boneless..... | do..... | 1, 953, 365 | 187, 020 |
| Lengthwise..... | do..... | 114, 915 | 9, 456 |
| Medium-scale..... | do..... | 171, 870 | 14, 597 |
| Kippered..... | do..... | 239, 855 | 29, 139 |
| Miscellaneous..... | do..... | 3, 850 | 231 |
| Canned "sardines"..... | standard cases..... | 980, 906 | 2, 397, 348 |
| Pearl essence..... | pounds..... | 7, 521 | 74, 210 |
| Me.l..... | tons..... | 11, 913 | 374, 123 |
| Oil..... | gallons..... | 3, 174, 212 | 402, 157 |
| Lake trout: | | | |
| Fresh filets..... | pounds..... | 12, 005 | 2, 676 |
| Smoked ² | do..... | 429, 630 | 102, 196 |
| "Jingcod" fresh filets..... | do..... | 157, 475 | 20, 220 |
| Mackerel: | | | |
| Fresh filets..... | do..... | 46, 066 | 7, 598 |
| Salted: | | | |
| Filets..... | do..... | 1, 787, 946 | 133, 266 |
| Split..... | do..... | 1, 390, 180 | 95, 396 |
| Miscellaneous..... | do..... | 1, 225 | 86 |
| Smoked ² | do..... | 175, 631 | 28, 368 |
| Canned..... | standard cases..... | 748, 567 | 1, 867, 915 |
| Meal..... | tons..... | 1, 278 | 35, 554 |
| Oil..... | gallons..... | 83, 778 | 10, 302 |
| Menhaden: | | | |
| Acid scrap..... | tons..... | 9, 481 | 173, 280 |
| Dry scrap..... | do..... | 25, 068 | 767, 878 |
| Meal..... | do..... | 6, 385 | 194, 734 |
| Oil..... | gallons..... | 3, 344, 343 | 460, 970 |
| Mullet: | | | |
| Salted ¹ | pounds..... | 1, 191, 362 | 48, 102 |
| Roe, salted ⁴ | do..... | 74, 704 | 17, 350 |
| Paddlefish: | | | |
| Smoked ¹ | do..... | 373, 800 | 166, 760 |
| Roe, salted ⁴ | do..... | 1, 595 | 812 |
| Pike: | | | |
| Fresh filets..... | do..... | 2, 744, 585 | 547, 646 |
| Frozen filets..... | do..... | 330, 888 | 63, 207 |
| Pilchard: | | | |
| Salted: | | | |
| Canned "sardines"..... | standard cases..... | 73, 400 | 4, 454 |
| Meal..... | tons..... | 1, 539, 440 | 8, 805, 168 |
| Meal..... | do..... | 60, 581 | 1, 530, 218 |
| Flour..... | do..... | 384 | 11, 834 |
| Oil..... | gallons..... | 10, 263, 776 | 1, 593, 088 |

¹ This item represents a combination of 1923 data in sections where available, and 1931 data in other sections.

² This is usually an intermediate product and although included in the total, may also be shown in its final stage of processing elsewhere in the table.

⁴ Data are for 1931.

Manufactured fishery products of the United States and Alaska—Continued

| Item | Quantity | Value |
|--|----------------|--------------|
| Pollock: | | |
| Fresh fillets and steaks.....pounds.. | 443, 631 | \$37, 479 |
| Frozen fillets.....do..... | 1, 091, 401 | 65, 147 |
| Salted: | | |
| Green.....do..... | 114, 145 | 2, 275 |
| Dry.....do..... | 983, 829 | 39, 119 |
| Rockfishes, fresh fillets.....do..... | 881, 800 | 132, 002 |
| Sablefish: | | |
| Fresh fillets.....do..... | 113, 000 | 12, 760 |
| Salted.....do..... | 144, 700 | 9, 055 |
| Kippered.....do..... | 290, 022 | 40, 854 |
| Smoked ¹do..... | 116, 030 | 29, 426 |
| Salmon: | | |
| Salted: | | |
| Dry.....do..... | 1, 399, 532 | 55, 281 |
| Pickled in brine.....do..... | 1, 034, 950 | 73, 920 |
| Mild-cured.....do..... | * 10, 494, 446 | 1, 891, 676 |
| Miscellaneous.....do..... | 15, 158 | 1, 584 |
| Smoked ¹do..... | 8, 228, 954 | 2, 256, 091 |
| Kippered.....do..... | 1, 662, 561 | 333, 912 |
| Canned: | | |
| Chinook or king.....standard cases.. | 322, 448 | 3, 071, 445 |
| Blueback, red or sockeye.....do..... | 2, 320, 886 | 10, 379, 350 |
| Silver or coho.....do..... | 243, 310 | 1, 352, 676 |
| Humpback or pink.....do..... | 2, 722, 723 | 12, 235, 661 |
| Chum or keta.....do..... | 735, 568 | 3, 012, 159 |
| Steelhead trout.....do..... | 17, 550 | 161, 228 |
| Meal: | | |
| Oil.....tons..... | 1, 657 | 50, 477 |
| Eggs, for bait, canned.....gallons.. | 187, 801 | 29, 761 |
| standard cases..... | 3, 273 | 93, 116 |
| Sea bass (Pacific coast): | | |
| Black, fresh fillets.....pounds.. | 260, 000 | 28, 600 |
| White, fresh fillets.....do..... | 147, 000 | 29, 900 |
| Shad: | | |
| Smoked and kippered ¹do..... | 242, 109 | 47, 745 |
| Canned.....standard cases.. | 1, 148 | 3, 837 |
| Roe, canned.....do..... | 1, 108 | 24, 823 |
| Sheepshead: | | |
| Fresh fillets.....pounds.. | 37, 502 | 4, 399 |
| Smoked ⁴do..... | 617 | 77 |
| Snapper, red: | | |
| Fresh fillets.....do..... | 25, 400 | 4, 500 |
| Fresh steaks.....do..... | 18, 600 | 3, 520 |
| Spanish mackerel, fresh fillets.....do..... | 2, 700 | 535 |
| Spot, salted ⁴do..... | 63, 500 | 2, 601 |
| Squeteagues: | | |
| Fresh fillets.....do..... | 80, 500 | 10, 638 |
| Sun-dried ⁴do..... | 33, 148 | 6, 166 |
| Squirrel hake, smoked.....do..... | 50 | 5 |
| Sturgeon: | | |
| Smoked and kippered.....do..... | 1, 529, 342 | 785, 970 |
| Salted roe.....do..... | 564 | 381 |
| Caviar, canned.....standard cases.. | 2, 596 | 333, 530 |
| Trout (Dolly Varden), dried.....pounds.. | 1, 500 | 30 |
| Tuna: | | |
| Canned: | | |
| Albacore.....standard cases.. | 54, 097 | 266, 301 |
| Bluefin.....do..... | 4, 179 | 21, 619 |
| Bonito.....do..... | 42, 439 | 170, 178 |
| Striped.....do..... | 225, 461 | 1, 022, 819 |
| "Tonno".....do..... | 136, 740 | 852, 236 |
| Yellowfin.....do..... | 936, 299 | 4, 417, 676 |
| Yellowtail.....do..... | 43, 918 | 163, 756 |
| Meal: | | |
| White bass, fresh fillets.....tons..... | 6, 004 | 153, 300 |
| pounds..... | 50, 873 | 10, 050 |
| Whitefish: | | |
| Fresh fillets.....do..... | 13, 988 | 3, 020 |
| Smoked ¹do..... | 2, 461, 295 | 647, 338 |
| Caviar, canned.....standard cases.. | 854 | 29, 754 |
| Whiting, smoked.....pounds.. | 50 | 5 |
| Wolfish, fresh fillets.....do..... | 85, 439 | 6, 262 |
| Yellow perch: | | |
| Fresh fillets.....do..... | 760, 113 | 155, 432 |
| Frozen fillets.....do..... | 87, 721 | 16, 854 |
| Crabs: | | |
| Meat, packaged, fresh-cooked ¹do..... | 6, 863, 844 | 1, 662, 497 |
| Canned.....standard cases.. | 9, 272 | 143, 517 |
| Meal.....tons..... | 800 | 11, 000 |
| Dry scrap.....do..... | 550 | 9, 412 |

¹ This item represents a combination of 1933 data in sections where available, and 1931 data in other sections.

² This is usually an intermediate product and although included in the total, may also be shown in its final stage of processing elsewhere in the table.

⁴ Data are for 1931.

Manufactured fishery products of the United States and Alaska—Continued

| Item | Quantity | Value | |
|--|---------------------|------------------------|-----------|
| Crabs, king, meal and dry scrap..... | tons.. | 572 | \$17,005 |
| Lobsters, meat, packaged, fresh-cooked..... | pounds.. | 79,020 | 86,420 |
| Shrimp: | | | |
| Fresh and frozen, packaged ¹ | do..... | 2,403,962 | 397,004 |
| Sun-dried ² | do..... | 1,469,152 | 321,857 |
| Cooked and peeled ³ | do..... | 779,050 | 153,388 |
| Canned..... | standard cases..... | 880,462 | 3,479,477 |
| Meal or "bran"..... | tons..... | 1,179 | 24,540 |
| Abalone, steaks..... | pounds..... | 531,698 | 143,282 |
| Clams, hard: | | | |
| Fresh-shucked ⁴ | gallons..... | 13,932 | 22,197 |
| Canned: | | | |
| Whole..... | standard cases..... | 32,704 | 82,044 |
| Chowder..... | do..... | 169,613 | 621,421 |
| Minced..... | do..... | 13,058 | 68,324 |
| Broth, bouillon, and cocktail..... | do..... | 4,503 | 25,485 |
| Juice..... | do..... | 7,327 | 28,938 |
| Clams, razor: | | | |
| Fresh-shucked..... | gallons..... | 36,078 | 17,729 |
| Canned: | | | |
| Whole and juice..... | standard cases..... | 2,214 | 17,061 |
| Minced..... | do..... | 62,010 | 498,472 |
| Clams, soft: | | | |
| Fresh-shucked..... | gallons..... | 129,286 | 120,024 |
| Steamed..... | pounds..... | ⁵ 161,887 | 11,279 |
| Canned: | | | |
| Whole..... | standard cases..... | 76,698 | 248,190 |
| Chowder..... | do..... | 46,732 | 131,314 |
| Juice and bouillon..... | do..... | 19,441 | 44,157 |
| Clams, surf, fresh-shucked..... | gallons..... | 5,922 | 4,050 |
| Clam shells, ground, poultry feed..... | tons..... | 1,630 | 15,340 |
| Marine-shell products: | | | |
| Buttons..... | gross..... | 5,486,180 | 3,299,907 |
| Novelties..... | do..... | | 463,093 |
| Mussels, fresh-water, shell products: | | | |
| Buttons..... | gross..... | 18,714,008 | 4,232,036 |
| Novelties..... | do..... | | 153,760 |
| Lime..... | tons..... | 3,083 | 8,920 |
| Poultry feed..... | do..... | 8,312 | 63,645 |
| Oysters: | | | |
| Eastern: | | | |
| Fresh-shucked ⁶ | gallons..... | 5,535,219 | 6,697,956 |
| Canned..... | standard cases..... | 317,630 | 927,128 |
| Japanese: | | | |
| Fresh-shucked..... | gallons..... | 151,384 | 166,361 |
| Canned..... | standard cases..... | 30,500 | 149,190 |
| Native, Pacific, fresh-shucked..... | gallons..... | 26,575 | 139,884 |
| Shell products: | | | |
| Poultry feed..... | tons..... | 276,342 | 1,430,398 |
| Lime..... | do..... | 45,660 | 104,524 |
| Lime, "burned"..... | do..... | 3,314 | 22,948 |
| Scallops, bay, fresh-shucked ⁷ | gallons..... | 126,162 | 306,921 |
| Scallops, sea, fresh-shucked..... | do..... | 381,650 | 397,733 |
| Turtles and terrapin, meat and soup, canned..... | standard cases..... | 2,607 | 42,008 |
| Whales: | | | |
| Meal, bone..... | tons..... | 207 | 4,453 |
| Meal, meat..... | do..... | 310 | 9,320 |
| Oil, whale..... | gallons..... | 569,850 | 93,341 |
| Oil, sperm..... | do..... | 16,200 | 2,900 |
| Unclassified products: | | | |
| Fillets, fresh..... | pounds..... | ⁸ 526,400 | 84,909 |
| Fillets, frozen..... | do..... | ⁹ 183,299 | 21,271 |
| Miscellaneous, packaged, fresh and frozen ¹ | do..... | ⁷ 1,637,453 | 164,533 |
| Salted ² | do..... | ⁴ 424,026 | 35,947 |
| Smoked ³ | do..... | ⁵ 43,407 | 10,561 |
| Canned: | | | |
| Fish for cat and dog food..... | standard cases..... | 213,811 | 588,090 |
| Fish cakes, balls, etc..... | do..... | 65,100 | 444,236 |

¹ This item represents a combination of 1833 data in sections where available, and 1931 data in other sections.

² This is usually an intermediate product and although included in the total, may also be shown in its final stage of processing elsewhere in the table.

³ Includes fresh fillets of barracuda, bluefish, grayfish, kingfish, lake herring, mullet, red drum, snook, and totuava.

⁴ Includes frozen fillets of cusk, mackerel, red snapper, rosefish, salmon, sheepshead, Spanish mackerel, squeteagues, white bass, and wolffish.

⁵ Includes fresh alewife roe, fresh-shucked conchs, pan-dressed flounders, fresh and frozen swordfish steaks, and frozen whiting sticks.

⁶ Includes salted anchovies, barracuda, bluefish, bonito, halibut, lake trout, black and white sea bass, Spanish mackerel, yellowtail, and tongues; spiced alewives; and mild-cured shad.

⁷ Includes smoked goosefish, halibut, salmon and shad weiners, suckers, and spiced salmon.

Manufactured fishery products of the United States and Alaska—Continued

| Item | Quantity | Value |
|---|-----------------------|------------------------|
| Unclassified products—Continued. | | |
| Fish flakes.....standard cases..... | 20, 076 | \$155, 424 |
| Other.....do..... | ¹⁰ 30, 592 | 212, 053 |
| Dry and green scrap.....tons..... | ¹¹ 3, 187 | 58, 926 |
| Meal.....do..... | ¹² 9, 843 | 435, 965 |
| Oil.....gallons..... | ¹³ 6, 479 | 1, 622 |
| Glue.....do..... | 355, 949 | 704, 101 |
| Other byproducts..... | | ¹⁴ 341, 289 |
| Total, fresh and frozen packaged products ¹pounds..... | 129, 608, 348 | 17, 294, 092 |
| Total, cured products ¹do..... | 104, 310, 213 | 12, 823, 491 |
| Total, canned.....do..... | 533, 212, 154 | 59, 799, 963 |
| Total, byproducts.....do..... | | 17, 465, 986 |
| Grand total..... | | 107, 383, 532 |

¹ This item represents a combination of 1933 data in sections where available, and 1931 data in other sections.

¹⁰ Includes canned anchovies, pickled eels, groundfish roe, haddock chowder, smoked lingcod, mackerel stew, salmon caviar, kippered salmon cheeks and steelhead trout, smoked sturgeon, conch cocktail, coquina broth, oyster puree, oyster soup, pickled sea mussels, rat poison, sea cucumber, squid, and miscellaneous canned fish.

¹¹ Includes scrap from sea herring, ground fish, and mixed fish.

¹² Includes meal from ground fish, salmon eggs, clams, and mixed fish.

¹³ Includes shark and miscellaneous fish oil.

¹⁴ Includes agar agar, alligator hides, concentrated powdered fish meat, isinglass, kelp products, mussel-shell stucco and chips, shark fins and hides, and miscellaneous novelties.

NOTE.—Some of the above products may have been manufactured from products imported from another country; therefore, they cannot be correlated directly with the catch within the United States and Alaska.

CANNED FISHERY PRODUCTS AND BYPRODUCTS TRADE

The output of canned fishery products and byproducts in the United States and Alaska in 1933 was valued at \$77,258,318. Of this total, canned products comprised \$59,799,963, and byproducts, \$17,458,355, an increase of 37 percent in the value of canned products and 40 percent in the value of byproducts when compared with the respective values of the same groups of commodities for the previous year.

Fishery products were canned at 384 establishments in the United States and Alaska during 1933. The combined output of these canneries amounted to 13,116,968 standard cases. The net weight of the products canned amounted to 533,212,154 pounds.

Canned fishery products or byproducts were prepared in 26 States and in Alaska during 1933. Alaska ranked first in the value of the products, accounting for 38 percent of the total; and California ranked second with 22 percent.

Canned fishery products and byproducts of the United States and Alaska, 1933

SUMMARY OF PRODUCTION: BY COMMODITIES

| Product | Number of plants | Standard cases | Quantity | Value |
|-------------------------------|------------------|----------------|---------------|---------------|
| Canned products: | | | | |
| Salmon: | | | | |
| United States..... | 43 | 1, 138, 861 | 54, 569, 328 | \$7, 865, 903 |
| Alaska..... | 91 | 5, 226, 604 | 260, 828, 992 | 28, 376, 014 |
| Sardines: | | | | |
| Maine..... | 23 | 980, 906 | 24, 522, 650 | 2, 397, 348 |
| California..... | 19 | 1, 539, 446 | 73, 693, 408 | 3, 805, 168 |
| Tuna and tunalike fishes..... | 13 | 1, 443, 133 | 34, 635, 192 | 6, 934, 485 |
| Alewife..... | 11 | 20, 280 | 973, 440 | 46, 363 |
| Alewife roe..... | 26 | 28, 513 | 1, 368, 624 | 111, 844 |
| Shad..... | 4 | 1, 148 | 55, 104 | 3, 837 |
| Shad roe..... | 9 | 1, 108 | 53, 184 | 24, 823 |
| Mackerel..... | 14 | 748, 567 | 35, 631, 216 | 1, 867, 915 |
| Fish flakes..... | 4 | 20, 076 | 963, 648 | 155, 424 |
| Fish cakes, balls, etc..... | 7 | 65, 100 | 3, 124, 800 | 444, 236 |

Canned fishery products and byproducts of the United States and Alaska, 1933—Con.

SUMMARY OF PRODUCTION: BY COMMODITIES—Continued

| Product | Number of plants | Standard cases | Quantity | Value |
|--|------------------|-------------------|------------------------------|-------------------|
| Canned products—Continued. | | | | |
| Cat and dog food..... | 6 | 213,811 | 10,262,928 | \$588,090 |
| Sturgeon caviar..... | 4 | 2,596 | 124,608 | 333,530 |
| Whitefish roe and caviar..... | 4 | 854 | 40,992 | 29,754 |
| Salmon eggs (for bait)..... | 8 | 3,273 | 157,104 | 93,145 |
| Miscellaneous fish, roe, and caviar..... | 17 | 12,110 | 581,280 | 85,729 |
| Oysters..... | 40 | 348,130 | 5,221,950 | 1,076,318 |
| Shrimp..... | 58 | 860,462 | 24,201,126 | 3,470,477 |
| Clam products..... | 65 | 434,500 | 10,239,060 | 1,766,406 |
| Crabs..... | 10 | 9,272 | 445,056 | 143,517 |
| Turtle and terrapin products..... | 5 | 2,607 | 125,136 | 42,008 |
| Miscellaneous shellfish..... | 6 | 18,611 | 893,328 | 128,529 |
| Total..... | 384 | 13,116,968 | 533,212,154 | 59,799,963 |
| Byproducts: | | | | |
| Oyster-shell products..... | | | <i>Tons</i> 326,946 | 1,623,210 |
| Fresh-water mussel-shell products..... | | | | 4,498,111 |
| Marine pearl-shell products..... | | | | 3,763,000 |
| Scrap, meal, etc..... | | | 129,991 | 3,877,296 |
| Marine animal oils..... | | | | |
| Miscellaneous byproducts..... | | | <i>Gallons</i> 17,742,222 | 2,624,519 |
| Total..... | | | | 17,458,355 |
| Grand total..... | | | | 77,258,318 |

¹ "Cutout" or "drained" weights of can contents are included for whole and minced clams, and gross can contents for other clam products.

² Exclusive of duplication.

VALUE OF PRODUCTION: BY STATES

| State | Canned products | Byproducts ¹ | Total |
|--|-------------------|-------------------------|-------------------|
| Maine..... | \$2,941,678 | \$246,713 | \$3,188,391 |
| Massachusetts..... | 791,864 | 1,261,165 | 2,062,752 |
| Rhode Island..... | | 9,723 | |
| Connecticut..... | 843,803 | 719,213 | 4,466,269 |
| New York..... | | 1,532,536 | |
| New Jersey..... | | 1,416,750 | |
| Pennsylvania..... | | 673,171 | |
| Delaware..... | | 202,531 | 202,531 |
| Maryland..... | 121,568 | 654,776 | 776,344 |
| Virginia..... | 91,765 | 867,039 | 958,804 |
| North Carolina and South Carolina..... | 363,176 | 239,542 | 602,718 |
| Georgia..... | 515,903 | 675,053 | 1,658,854 |
| Florida..... | 467,898 | | |
| Alabama..... | 320,911 | 62,450 | 1,366,654 |
| Mississippi..... | 974,293 | | |
| Louisiana..... | 1,572,942 | | |
| Texas and Wisconsin..... | 317,486 | 4,659 | 322,145 |
| Missouri, Illinois, and Kentucky..... | | 112,808 | 112,808 |
| Iowa..... | | 3,304,028 | 3,304,028 |
| Washington..... | 5,801,293 | 75,678 | 5,876,971 |
| Oregon..... | 2,674,923 | 16,240 | 2,691,163 |
| California..... | 13,231,561 | 3,868,736 | 17,100,297 |
| Alaska..... | 28,759,899 | 833,132 | 29,593,031 |
| Total..... | 59,799,963 | 17,458,355 | 77,258,318 |

¹ Includes menhaden, fresh-water mussel-shell, and marine-pearl shell products.

Canned fishery products and byproducts of the United States and Alaska, 1933—Con.

PACK OF CANNED SALMON: STANDARD CASES

| Product | Alaska | | | | | | | |
|---------------------------|-------------|-------------|-------------|-------------|-------------|--------------|-------------|--------------|
| | Southeast | | Central | | Western | | Total | |
| | Cases | Value | Cases | Value | Cases | Value | Cases | Value |
| Chinook or king: | | | | | | | | |
| 1-pound tall..... | 3, 873 | \$21, 111 | 12, 578 | \$69, 964 | 4, 986 | \$28, 929 | 21, 437 | \$120, 004 |
| 1-pound oval..... | 3, 189 | 27, 608 | 3, 811 | 31, 032 | 3, 021 | 28, 480 | 10, 021 | 87, 120 |
| 1-pound flat..... | 1, 084 | 9, 676 | 7, 397 | 79, 637 | 1, 474 | 14, 745 | 9, 955 | 104, 058 |
| 1/2-pound flat..... | | | | | | | | |
| Total..... | 8, 146 | 58, 395 | 23, 786 | 180, 633 | 9, 481 | 72, 154 | 41, 413 | 311, 182 |
| Blueback, red or sockeye: | | | | | | | | |
| 1-pound tall..... | 64, 467 | 421, 349 | 389, 169 | 2, 532, 373 | 1, 612, 957 | 10, 711, 792 | 2, 066, 593 | 13, 665, 514 |
| 1-pound flat..... | 3, 832 | 30, 656 | 54, 651 | 411, 717 | 1, 569 | 10, 277 | 60, 052 | 462, 650 |
| 1/2-pound flat..... | 12, 827 | 121, 842 | 40, 664 | 392, 209 | 147 | 1, 073 | 53, 638 | 515, 124 |
| Total..... | 81, 126 | 573, 847 | 484, 484 | 3, 336, 299 | 1, 614, 673 | 10, 723, 142 | 2, 180, 283 | 14, 633, 288 |
| Silver or coho: | | | | | | | | |
| 1-pound tall..... | 90, 737 | 472, 569 | 62, 351 | 313, 521 | 1, 466 | 7, 517 | 154, 544 | 793, 607 |
| 1-pound flat..... | 2, 096 | 12, 576 | 2, 561 | 14, 133 | | | 4, 657 | 26, 709 |
| 1/2-pound flat..... | 2, 972 | 22, 335 | 395 | 2, 368 | | | 3, 367 | 24, 703 |
| Total..... | 95, 805 | 507, 480 | 65, 307 | 330, 022 | 1, 466 | 7, 517 | 162, 568 | 845, 019 |
| Humpback or pink: | | | | | | | | |
| 1-pound tall..... | 1, 467, 473 | 6, 664, 922 | 700, 221 | 3, 116, 508 | | | 2, 167, 694 | 9, 781, 430 |
| 1/2-pound flat..... | 10, 540 | 64, 660 | 4, 317 | 27, 629 | | | 14, 857 | 92, 289 |
| Total..... | 1, 478, 013 | 6, 729, 582 | 704, 538 | 3, 144, 137 | | | 2, 182, 551 | 9, 873, 719 |
| Chum or keta: | | | | | | | | |
| 1-pound tall..... | 424, 611 | 1, 728, 120 | 207, 471 | 865, 780 | 26, 049 | 115, 254 | 658, 131 | 2, 709, 154 |
| 1/2-pound flat..... | 250 | 1, 355 | 408 | 2, 287 | | | 658 | 3, 652 |
| Total..... | 424, 861 | 1, 730, 485 | 207, 879 | 868, 067 | 26, 049 | 115, 254 | 658, 789 | 2, 712, 806 |
| Grand total..... | 2, 087, 951 | 9, 598, 789 | 1, 485, 994 | 7, 859, 158 | 1, 651, 659 | 10, 918, 067 | 5, 225, 604 | 28, 376, 014 |

| Product | United States | | | | | | Grand total, Alaska and United States | |
|---------------------------|---------------|-------------|-----------------------|-------------|----------|-------------|---------------------------------------|--------------|
| | Washington | | Oregon and California | | Total | | | |
| | Cases | Value | Cases | Value | Cases | Value | | |
| Chinook or king: | | | | | | | | |
| 1-pound tall..... | 15, 597 | \$69, 727 | 34, 210 | \$142, 644 | 49, 807 | \$212, 371 | 71, 244 | \$332, 375 |
| 1-pound oval..... | 210 | 3, 728 | 697 | 73, 372 | 907 | 16, 100 | 907 | 16, 100 |
| 1-pound flat..... | 8, 502 | 67, 451 | 39, 181 | 332, 854 | 47, 683 | 400, 305 | 57, 704 | 487, 425 |
| 1/2-pound oval..... | | | 106 | 2, 672 | 106 | 2, 672 | 106 | 2, 672 |
| 1/2-pound flat..... | 50, 258 | 536, 300 | 128, 259 | 1, 529, 797 | 178, 517 | 2, 066, 097 | 188, 472 | 2, 170, 155 |
| 1/4-pound flat..... | | | 4, 015 | 62, 718 | 4, 015 | 62, 718 | 4, 015 | 62, 718 |
| Total..... | 74, 567 | 677, 206 | 206, 468 | 2, 063, 057 | 281, 035 | 2, 760, 263 | 322, 448 | 3, 071, 445 |
| Blueback, red or sockeye: | | | | | | | | |
| 1-pound tall..... | 2, 473 | 21, 954 | | | 2, 473 | 21, 954 | 2, 099, 066 | 13, 687, 468 |
| 1-pound flat..... | 24, 800 | 285, 200 | | | 24, 800 | 285, 200 | 84, 852 | 737, 850 |
| 1/2-pound flat..... | 106, 807 | 1, 350, 899 | 6, 238 | 83, 901 | 113, 045 | 1, 434, 800 | 166, 683 | 1, 949, 924 |
| 1/4-pound flat..... | | | 265 | 4, 106 | 265 | 4, 106 | 265 | 4, 106 |
| Total..... | 134, 080 | 1, 658, 053 | 6, 503 | 88, 009 | 140, 583 | 1, 745, 062 | 2, 320, 866 | 16, 379, 350 |
| Silver or coho: | | | | | | | | |
| 1-pound tall..... | 17, 817 | 89, 477 | 4, 227 | 21, 135 | 22, 044 | 110, 612 | 176, 588 | 904, 219 |
| 1-pound flat..... | 10, 704 | 59, 942 | 9, 146 | 51, 218 | 19, 850 | 111, 160 | 24, 507 | 137, 866 |
| 1/2-pound flat..... | 17, 809 | 139, 801 | 15, 077 | 118, 354 | 32, 886 | 258, 155 | 36, 253 | 282, 858 |
| 1/4-pound flat..... | 495 | 4, 811 | 5, 467 | 52, 319 | 5, 962 | 57, 130 | 5, 962 | 67, 180 |
| Total..... | 46, 825 | 294, 031 | 33, 917 | 243, 026 | 80, 742 | 537, 057 | 243, 310 | 1, 382, 076 |
| Humpback or pink: | | | | | | | | |
| 1-pound tall..... | 505, 880 | 2, 173, 134 | | | 505, 880 | 2, 173, 134 | 2, 673, 074 | 11, 954, 564 |
| 1-pound flat..... | 15, 109 | 70, 710 | | | 15, 109 | 70, 710 | 15, 109 | 70, 710 |
| 1/2-pound flat..... | 19, 683 | 118, 098 | | | 19, 683 | 118, 098 | 34, 540 | 210, 387 |
| Total..... | 540, 172 | 2, 361, 942 | | | 540, 172 | 2, 361, 942 | 2, 722, 723 | 12, 235, 661 |

Canned fishery products and byproducts of the United States and Alaska, 1933—Con.

| Product | United States | | | | | | Grand total, Alaska and United States | |
|-------------------------|----------------|------------------|--------------------------|------------------|------------------|------------------|---|-------------------|
| | Washington | | Oregon and California | | Total | | | |
| Chum or keta: | <i>Cases</i> | <i>Value</i> | <i>Cases</i> | <i>Value</i> | <i>Cases</i> | <i>Value</i> | <i>Cases</i> | <i>Value</i> |
| 1-pound tall..... | 46,508 | \$181,593 | 26,191 | \$98,272 | 72,699 | \$279,865 | 730,830 | \$2,989,019 |
| 1-pound flat..... | 1,773 | 7,535 | 444 | 1,820 | 2,217 | 9,355 | 2,217 | 9,355 |
| ½-pound flat..... | 1,498 | 8,089 | 365 | 2,044 | 1,863 | 10,133 | 2,521 | 13,785 |
| Total..... | 49,779 | 197,217 | 27,000 | 102,136 | 76,779 | 299,353 | 735,568 | 3,012,159 |
| Steelhead: | | | | | | | | |
| 1-pound tall..... | 375 | 1,561 | 2,008 | 8,233 | 2,383 | 9,794 | 2,383 | 9,794 |
| 1-pound flat..... | 996 | 6,523 | 1,143 | 7,487 | 2,139 | 14,010 | 2,139 | 14,010 |
| ½-pound oval..... | 389 | 5,018 | 3,924 | 50,620 | 4,313 | 55,638 | 4,313 | 55,638 |
| ½-pound flat..... | 1,721 | 15,803 | 5,926 | 55,408 | 7,647 | 71,211 | 7,647 | 71,211 |
| ¼-pound flat..... | | | 1,068 | 10,573 | 1,068 | 10,573 | 1,068 | 10,573 |
| Total..... | 3,481 | 28,905 | 14,009 | 132,321 | 17,550 | 161,226 | 17,550 | 161,226 |
| Grand total..... | 848,904 | 5,217,354 | 287,957 | 2,648,549 | 1,136,861 | 7,865,903 | 6,362,465 | 36,241,917 |

NOTE.—“Standard cases” represent the various-sized cases converted to the equivalent of forty-eight 1-pound cans to the case. Salmon were canned at 31 plants in Washington, 11 in Oregon, 1 in California, and 91 in Alaska.

PACK OF CANNED SARDINES

| Sardines (herring) | Maine | | Sardines (pilchard) | California | |
|------------------------------------|----------------|------------------|--|------------------|------------------|
| Quarters, ¼-pound (100 cans): | <i>Cases</i> | <i>Value</i> | 1-pound oval (48 cans): | <i>Cases</i> | <i>Value</i> |
| In cottonseed oil..... | 821,388 | \$2,000,218 | In cottonseed oil..... | 13,871 | \$32,810 |
| In mustard..... | 89,061 | 252,237 | In mustard..... | 128,128 | 311,960 |
| In other sauces or oils..... | 7,786 | 29,673 | In tomato sauce..... | 1,160,029 | 2,819,755 |
| Three-quarters, ⅜-pound (48 cans): | | | In natural oil..... | 21,991 | 50,652 |
| In mustard..... | 43,522 | 115,220 | In other sauces or oils..... | 8,579 | 21,237 |
| | | | ½-pound oval (48 cans): | | |
| | | | In various sauces or oils.. | 4,604 | 13,571 |
| | | | 1-pound tall (48 cans): | | |
| | | | In natural oil..... | 105,087 | 227,456 |
| | | | In other sauces or oils..... | 20,403 | 49,255 |
| | | | 6-ounce (100 cans): | | |
| | | | In tomato sauce..... | 4,247 | 10,251 |
| | | | In natural oil..... | 83,183 | 201,640 |
| | | | In other sauces or oils..... | 4,576 | 11,843 |
| | | | Other sizes: | | |
| | | | In various sauces or oils (standard cases)..... | 7,176 | 64,738 |
| Total..... | 961,757 | 2,397,348 | Total..... | 1,561,874 | 3,805,168 |
| Total (standard cases)..... | 980,906 | | Total (standard cases)..... | 1,539,446 | |

† Includes a small amount packed in 6-ounce cans, 100 to the case, which have been converted to the basis of 4-ounce cans, 100 to the case.

NOTE.—“Standard cases” represent the various sized cases converted to the uniform basis of one hundred ¼-pound cans to the case of sardines (herring), and forty-eight 1-pound cans to the case of sardines (pilchard). Sardines were canned at 23 plants in Maine and 19 in California.

PACK OF CANNED TUNA AND TUNALIKE FISHES IN CALIFORNIA †

| Size | Albacore | | Yellowfin † | | Bluefin | | Striped | |
|------------------------------------|---------------|----------------|----------------|------------------|--------------|---------------|----------------|------------------|
| | <i>Cases</i> | <i>Value</i> | <i>Cases</i> | <i>Value</i> | <i>Cases</i> | <i>Value</i> | <i>Cases</i> | <i>Value</i> |
| ¼-pound (48 cans)..... | 1,242 | \$4,667 | 112,982 | \$380,829 | 1,345 | \$5,320 | 18,687 | \$60,107 |
| ½-pound (48 cans)..... | 41,404 | 227,023 | 643,373 | 3,071,470 | 2,539 | 12,063 | 187,963 | 852,653 |
| 1-pound (48 cans)..... | 4,287 | 7,39,713 | 7,69,383 | 7,592,069 | 244 | 2,122 | 10,040 | 81,023 |
| Flakes (standard cases)..... | 3,558 | 14,898 | 97,669 | 373,308 | 480 | 2,014 | 8,074 | 29,036 |
| Total..... | 50,461 | 286,301 | 923,407 | 4,417,676 | 4,008 | 21,519 | 224,764 | 1,022,819 |
| Total (standard cases)..... | 54,097 | | 936,299 | | 4,179 | | 225,461 | |

† Includes a very small amount of tuna packed in Oregon which has been included with albacore in ¼-pound cans, 48 to the case.

† Includes a small amount of creamed yellowfin tuna.

† Includes the pack in 4-pound cans, 12 to the case, which have been converted to the equivalent of 1-pound cans, 48 to the case.

Canned fishery products and byproducts of the United States and Alaska, 1933—Con.

PACK OF CANNED TUNA AND TUNALIKE FISHES IN CALIFORNIA—Continued

| Size | "Tonno" | | Bonito | | Yellowtail | | Total | |
|-------------------------------------|-----------------|-----------------|----------------|-----------------|----------------|-----------------|--------------------|--------------------|
| | Cases | Value | Cases | Value | Cases | Value | Cases | Value |
| ¼-pound (48 cans)..... | | | 21 | \$63 | | | 134, 277 | \$450, 986 |
| ¼-pound (100 cans)..... | 119, 620 | \$775, 423 | 4, 247 | 23, 867 | | | 123, 867 | 799, 290 |
| ½-pound (48 cans)..... | 12, 143 | 76, 813 | 28, 764 | 114, 761 | 23, 088 | \$91, 425 | 939, 274 | 4, 446, 198 |
| 1-pound (48 cans)..... | | | 4, 620 | 31, 497 | 10, 415 | 72, 331 | 98, 959 | 818, 753 |
| Flakes (standard cases)..... | | | (1) | (5) | | | 109, 781 | 419, 266 |
| Total | 131, 763 | 852, 236 | 37, 652 | 170, 178 | 33, 503 | 163, 756 | 1, 406, 158 | 6, 934, 485 |
| Total (standard cases) | 136, 740 | | 42, 439 | | 43, 918 | | 1, 443, 133 | |

* A small amount of bonito flakes is included with canned bonito in ¼-pound cans, 48 to the case.

NOTE.—"Standard cases" represent the various sized cases converted to the equivalent of forty-eight ¼-pound cans to the case. Tuna and tunalike fishes were canned at 12 plants in California and 1 in Oregon.

PACK OF CANNED ALEWIVES AND ALEWIFE ROE: STANDARD CASES

| Product | Maryland | | Virginia and North Carolina | | Total | |
|--------------------|----------------|----------------|-----------------------------|----------------|----------------|-----------------|
| | Cases | Value | Cases | Value | Cases | Value |
| Alewives..... | 16, 575 | \$36, 600 | 3, 705 | \$9, 853 | 20, 280 | \$46, 363 |
| Alewife roe..... | 7, 336 | 29, 390 | 21, 177 | 82, 254 | 28, 513 | 111, 644 |
| Total | 23, 911 | 65, 990 | 24, 882 | 92, 117 | 48, 793 | 158, 007 |

PACK OF CANNED ALEWIVES AND ALEWIFE ROE: ACTUAL CASES

| Product and size | Cases | Value | Product and size | | Cases | Value |
|---------------------------------------|---------|----------------|---------------------------------------|--|---------|-----------------|
| | | | | | | |
| Alewives: | | | Alewife roe—Continued. | | | |
| 16-ounce (48 cans)..... | 14, 678 | \$34, 831 | 10-, 16-, and 19-ounce (24 cans)..... | | 4, 988 | \$9, 061 |
| 16-, 17-, and 19-ounce (24 cans)..... | 10, 111 | 11, 532 | 17-ounce (24 cans)..... | | 43, 486 | 85, 974 |
| Total | | 46, 363 | Total | | | 111, 644 |
| Alewife roe: | | | Grand total | | | 158, 007 |
| 8-ounce (48 cans)..... | 4, 851 | 13, 583 | | | | |
| 7½, 8½, and 10-ounce (48 cans)..... | 875 | 3, 026 | | | | |

NOTE.—"Standard cases" represent the various sized cases converted to the equivalent of forty-eight 1-pound cans to the case. Alewives or alewife roe were canned at 7 plants in Maryland, 17 in Virginia, and 2 in North Carolina.

PACK OF CANNED SHRIMP: STANDARD CASES

| State | Dry pack (in tins) | | Wet pack (in tins) | | Wet pack (in glass) | | Total | |
|---------------------------------|--------------------|-----------------|--------------------|--------------------|---------------------|-----------------|-----------------|--------------------|
| | Cases | Value | Cases | Value | Cases | Value | Cases | Value |
| South Carolina and Alabama..... | 21, 813 | \$91, 358 | 71, 893 | \$291, 073 | | | 93, 696 | \$382, 431 |
| Georgia..... | 10, 675 | 33, 105 | 86, 813 | 344, 285 | 24, 114 | \$159, 875 | 121, 802 | 536, 265 |
| Florida..... | 3, 579 | 12, 387 | 65, 179 | 205, 698 | 17, 255 | 125, 194 | 86, 013 | 343, 209 |
| Mississippi..... | 38, 690 | 144, 759 | 81, 182 | 302, 313 | | | 119, 872 | 447, 072 |
| Louisiana..... | 108, 932 | 456, 833 | 263, 823 | 1, 044, 243 | (9) | (9) | 372, 755 | 1, 501, 076 |
| Texas..... | 22, 037 | 89, 676 | 44, 487 | 179, 748 | | | 66, 524 | 269, 424 |
| Total | 205, 726 | 828, 088 | 613, 367 | 2, 367, 320 | 41, 369 | 284, 069 | 860, 462 | 3, 479, 477 |

* The pack of shrimp in glass for Louisiana and Texas has been included with that of Georgia to avoid disclosure of private enterprise.

PACK OF CANNED SHRIMP: ACTUAL CASES

| Size | Cases | Value | Size | Cases | Value |
|-----------------------------------|----------|-------------|-----------------------------------|---------|--------------------|
| | | | | | |
| In tins, dry: | | | In glass, wet: | | |
| 4-ounce (48 cans)..... | 1, 111 | \$3, 393 | 4-ounce (24 jars)..... | 2, 140 | \$13, 607 |
| 5-ounce (48 cans)..... | 180, 930 | 719, 132 | 6-ounce (24 jars)..... | 46, 900 | 158, 834 |
| 8½-ounce (24 cans)..... | 26, 310 | 102, 425 | Other sizes (standard cases)..... | 16, 167 | 111, 623 |
| Other sizes (standard cases)..... | 2, 202 | 3, 138 | Total | | 3, 479, 477 |
| In tins, wet: | | | | | |
| 5¼-ounce (48 cans)..... | 610, 592 | 2, 353, 476 | | | |
| 9¼-ounce (24 cans)..... | 2, 478 | 9, 506 | | | |
| Other sizes (standard cases)..... | 674 | 4, 338 | | | |

NOTE.—"Standard cases" represent the various sized cases converted to the equivalent of forty-eight 5-ounce cans to the case in the dry pack and forty-eight 5¼-ounce cans to the case in the wet pack. Shrimp were canned at 1 plant in South Carolina, 6 in Georgia, 7 in Florida, 3 in Alabama, 15 in Mississippi, 21 in Louisiana, and 5 in Texas.

Canned fishery products and byproducts of the United States and Alaska, 1933—Con.

PACK OF CANNED OYSTERS: STANDARD CASES

| State | Cases | Value | State | Cases | Value |
|---|----------|------------|-----------------|----------|-------------|
| South Carolina..... | 93, 082 | \$274, 674 | Louisiana..... | 19, 439 | \$60, 164 |
| Georgia, Florida, Alabama, and New Jersey..... | 21, 982 | 65, 069 | Washington..... | 30, 500 | 149, 190 |
| Mississippi..... | 183, 127 | 527, 221 | Total..... | 348, 130 | 1, 076, 318 |

PACK OF CANNED OYSTERS: ACTUAL CASES

| Size | Cases | Value | Size | Cases | Value |
|-------------------------|----------|-----------|-----------------------------------|---------|-------------|
| 4-ounce (48 cans)..... | 24, 150 | \$69, 971 | 10-ounce (24 cans)..... | 38, 768 | \$103, 731 |
| 5-ounce (48 cans)..... | 252, 588 | 744, 211 | Other sizes (standard cases)..... | 2, 993 | 12, 880 |
| 7½-ounce (48 cans)..... | 14, 086 | 103, 389 | Total..... | | 1, 076, 318 |
| 8-ounce (24 cans)..... | 16, 875 | 42, 136 | | | |

NOTE.—“Standard cases” represent the various sized cases converted to the equivalent of forty-eight 5-ounce cans to the case. Oysters were canned at 1 plant in New Jersey, 4 in South Carolina, 2 in Georgia, 2 in Florida, 2 in Alabama, 15 in Mississippi, 5 in Louisiana, and 9 in Washington.

PACK OF CANNED CLAMS AND CLAM PRODUCTS

| Item and State | Cases | Value | Item and State | Cases | Value |
|---|----------|----------|---|----------|-------------|
| Razor clams (Washington, Oregon, and Alaska): | | | Hard clams—Continued. | | |
| Whole: | | | Juice: | | |
| ½-pound, 4-ounce (48 cans)..... | 116 | \$540 | No. 1, 10-ounce (48 cans).... | 971 | \$3, 581 |
| No. 1, 5-ounce (48 cans)..... | 1, 309 | 12, 731 | No. 10, 102-ounce (6 cans).... | 1, 990 | 5, 450 |
| 1-pound, 8-ounce (48 cans).... | 299 | 2, 410 | Other sizes (standard cases)... | 3, 818 | 19, 937 |
| Other sizes (standard cases).... | 46 | 382 | Cocktail: All sizes (standard cases)..... | 1, 167 | 6, 498 |
| Minced: | | | Broth and bouillon: | | |
| ½-pound, 4-ounce (48 cans).... | 52, 903 | 347, 267 | All sizes (standard cases).... | 3, 536 | 19, 987 |
| No. 1, 5-ounce (48 cans)..... | 19, 278 | 148, 292 | Total..... | 229, 017 | 827, 212 |
| No. 2, 10-ounce (24 cans)..... | 370 | 2, 713 | Total (standard cases)..... | 227, 405 | ----- |
| Other sizes (standard cases).... | 40 | 200 | Soft clams (Maine and Massachusetts): | | |
| Juice: | | | Whole: | | |
| All sizes (standard cases)..... | 288 | 998 | No. 1, 5-ounce (48 cans)..... | 55, 986 | 183, 907 |
| Total..... | 74, 649 | 515, 533 | 1-pound, 8-ounce (48 cans).... | 5, 324 | 22, 562 |
| Total (standard cases)..... | 64, 224 | ----- | No. 2, 10-ounce (24 cans).... | 8, 998 | 28, 517 |
| Hard clams (Rhode Island, New York, New Jersey, Pennsylvania, Maryland, Florida, Washington, and Alaska): ¹⁶ | | | Other sizes (standard cases)... | 3, 196 | 13, 204 |
| Whole: | | | Chowder: | | |
| No. 1, 5-ounce (48 cans)..... | 1, 598 | 11, 061 | No. 1, 10-ounce (48 cans).... | 16, 563 | 51, 542 |
| 1-pound, 8-ounce (48 cans).... | 1, 512 | 9, 292 | Other sizes (standard cases)... | 30, 169 | 79, 772 |
| No. 2, 10-ounce (24 cans)..... | 3, 293 | 18, 501 | Bouillon and juice: | | |
| No. 10, 52-ounce (6 cans).... | 3, 864 | 16, 422 | No. 1, 10-ounce (48 cans).... | 4, 867 | 14, 519 |
| Other sizes (standard cases).... | 20, 371 | 26, 745 | No. 2, 20-ounce (24 cans).... | 10, 077 | 21, 568 |
| Minced: | | | No. 10, 102-ounce (6 cans).... | 1, 851 | 2, 977 |
| ½-pound, 4-ounce (48 cans).... | 6, 167 | 20, 043 | Other sizes (standard cases)... | 2, 137 | 5, 093 |
| No. 1, 5-ounce (48 cans)..... | 1, 969 | 9, 386 | Total..... | 139, 168 | 423, 661 |
| Other sizes (standard cases).... | 6, 155 | 38, 895 | Total (standard cases)..... | 142, 871 | ----- |
| Chowder: | | | Grand total (standard cases)..... | 434, 500 | 1, 766, 406 |
| No. 1, 10-ounce (48 cans)..... | 114, 468 | 378, 931 | | | |
| 8½-ounce (12 cans)..... | 20, 330 | 74, 077 | | | |
| No. 10, 102-ounce (6 cans).... | 2, 052 | 12, 521 | | | |
| Other sizes (standard cases).... | 35, 756 | 155, 892 | | | |

¹⁶ Includes a small amount of coquina broth packed in Florida.

NOTE.—“Standard cases” represent the various sized cases converted to the equivalent of 48 no. 1, 5-ounce cans to the case for whole and minced clams; and 48 no. 1, 10-ounce cans to the case for other clam products. Razor-clam products were canned at 10 plants in Washington, 4 in Oregon, and 14 in Alaska; hard-clam products, at 1 plant in Rhode Island, 2 in New York, 2 in New Jersey, 1 in Pennsylvania, 2 in Maryland, 1 in Florida, 6 in Washington, and 1 in Alaska; soft clam products, at 18 plants in Maine and 2 in Massachusetts; and coquina clam products, at 1 plant in Florida.

Canned fishery products and byproducts of the United States and Alaska, 1933—Con.

PACK OF MISCELLANEOUS CANNED FISHERY PRODUCTS: STANDARD CASES

| Item | Cases | Value | Item | Cases | Value |
|-------------------------------------|---------|-------------|--|-----------|-----------|
| Mackerel..... | 748,567 | \$1,867,915 | Salmon eggs (for bait)..... | 3,273 | \$93,145 |
| Shad..... | 1,148 | 3,837 | Miscellaneous fish, roe and caviar ¹³ | 12,110 | 85,729 |
| Shad roe..... | 1,108 | 24,823 | Crabs..... | 9,272 | 143,517 |
| Fish flakes ¹¹ | 20,076 | 155,424 | Turtle and terrapin products..... | 2,607 | 42,008 |
| Cat and dog food..... | 213,811 | 598,090 | Miscellaneous shellfish ¹⁴ | 18,611 | 128,829 |
| Fish cakes, balls, etc..... | 65,100 | 444,236 | | | |
| Sturgeon caviar ¹² | 2,596 | 333,530 | Total..... | 1,099,133 | 3,940,837 |
| Whitefish roe and caviar..... | 854 | 20,754 | | | |

¹¹ Tuna flakes are not included in this table, but are included in the table for canned tuna and tunalike fishes.

¹² Produced principally from imported sturgeon.

¹³ Includes kippered and smoked salmon, finnan haddie, haddock, chowder, pickled eels, mackerel stew, smoked lingcod and sturgeon, anchovies, salmon roe and caviar, and miscellaneous roe.

¹⁴ Includes oyster puree and soup, pickled mussels, sea cucumbers, squid, and conch products.

PRODUCTION OF OYSTER-SHELL PRODUCTS¹⁵

| State | Crushed oyster-shell for poultry feed | | Oyster-shell lime | | Total | |
|--|---------------------------------------|-----------|----------------------|----------------------|---------|-----------|
| | Tons | Value | Tons | Value | Tons | Value |
| Rhode Island and Delaware..... | 945 | \$7,517 | 315 | \$1,316 | 1,260 | \$8,833 |
| New Jersey..... | 4,553 | 44,864 | 1,610 | 6,629 | 6,163 | 51,493 |
| Pennsylvania..... | 3,646 | 33,119 | 1,114 | 4,570 | 4,760 | 37,889 |
| Maryland..... | 34,298 | 175,909 | 20,998 | 30,017 | 55,296 | 205,986 |
| Virginia..... | 15,537 | 91,940 | ¹⁶ 14,898 | ¹⁶ 60,543 | 30,427 | 152,483 |
| North Carolina and South Carolina..... | 7,100 | 45,300 | ¹⁷ 1,850 | ¹⁷ 6,287 | 8,950 | 51,587 |
| Florida..... | 65,383 | 318,715 | (¹⁷) | (¹⁷) | 65,383 | 318,715 |
| Alabama, Louisiana, and Texas..... | 113,154 | 619,173 | 1,638 | 4,014 | 114,792 | 623,187 |
| Mississippi..... | 14,567 | 46,992 | 2,744 | 1,408 | 17,311 | 48,400 |
| Washington..... | 2,046 | 19,467 | | | 2,046 | 19,467 |
| California..... | 16,743 | 92,682 | 3,815 | 12,688 | 20,558 | 105,370 |
| Total..... | 277,972 | 1,495,738 | 48,974 | 127,472 | 326,946 | 1,623,210 |

¹⁵ Most of the production in Washington and a small part of the production in California was made from crushed clam shells.

¹⁶ Of this amount 3,314 tons, valued at \$22,948 were reported as "burned" lime.

¹⁷ The production of oyster-shell lime in Florida has been included with that of North Carolina and South Carolina.

NOTE.—The above crushed shell products were prepared at 2 plants in Rhode Island, 9 in New Jersey, 5 in Pennsylvania, 1 in Delaware, 5 in Maryland, 10 in Virginia, 2 in North Carolina, 2 in South Carolina, 3 in Florida, 2 in Alabama, 5 in Mississippi, 1 in Louisiana, 1 in Texas, 5 in California, and 5 in Washington.

PRODUCTION OF FRESH-WATER MUSSEL-SHELL PRODUCTS

| Item | Iowa | | New York | | Other States | | Total | |
|---|------------|-------------|-----------|-----------|--------------|-----------|------------|-------------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| Pearl buttons, gross..... | 13,698,445 | \$3,139,905 | 4,346,839 | \$959,337 | 668,724 | \$132,794 | 18,714,006 | \$4,232,036 |
| Crushed shell for poultry feed, tons..... | 7,675 | 59,639 | ----- | ----- | 692 | 4,006 | 8,312 | 63,645 |
| Lime.....do..... | 2,624 | 8,274 | 294 | 466 | 245 | 180 | 3,063 | 8,920 |
| Other products ¹⁴ | ----- | 156,210 | ----- | ----- | ----- | 32,300 | ----- | 188,510 |
| Total..... | ----- | 3,364,028 | ----- | 959,803 | ----- | 169,280 | ----- | 4,493,111 |

¹⁴ Includes stucco, colored shells, and "pearl novelties."

NOTE.—Mussel shells utilized in the above production amounted to 60,292,000 pounds, valued at \$873,138. Shells were taken in 16 States in the Mississippi Valley and Great Lakes region. The producing States in order of their importance were: Arkansas, which contributed 23 percent of the total quantity; Indiana, 16 percent; Tennessee, 14 percent; Illinois, 11 percent; Michigan, 9 percent; Kentucky, 6 percent; Iowa, 5 percent; Wisconsin, 5 percent; Minnesota, 3 percent; Alabama and Texas each 2 percent; Missouri, Ohio, and South Dakota, each 1 percent; and Mississippi and Oklahoma each less than one-half of 1 percent.

Canned fishery products and byproducts of the United States and Alaska, 1933—Con.

PRODUCTION OF MARINE PEARL-SHELL PRODUCTS ¹⁹

| Item | Maine, Massachusetts, Rhode Island, and Connecticut | | New York | | New Jersey | |
|-------------------------|---|------------|----------|------------|-------------|---------------|
| | Gross | Value | Gross | Value | Gross | Value |
| Pearl buttons | 1, 199, 684 | \$741, 969 | 825, 708 | \$509, 920 | 1, 617, 128 | \$1, 073, 275 |
| Novelties ²⁰ | | 215, 400 | | 46, 813 | | 78, 955 |
| Total | | 957, 369 | | 556, 733 | | 1, 152, 230 |

| Item | Pennsylvania, Maryland, and Florida | | Oregon and California | | Total | |
|-------------------------|-------------------------------------|-------------|-----------------------|-----------|-------------|---------------|
| | Gross | Value | Gross | Value | Gross | Value |
| Pearl buttons | 1, 843, 660 | \$974, 743 | | | 5, 486, 180 | \$3, 299, 907 |
| Novelties ²⁰ | | 101, 339 | | \$20, 586 | | 463, 093 |
| Total | | 1, 076, 082 | | 20, 586 | | 3, 763, 000 |

¹⁹ Produced principally from imported shells.²⁰ Includes buckles, inlays for jewelry, knife handles, lamps, handles for manicure sets, ornaments, etc.

NOTE.—Marine pearl-shell products were manufactured at 1 plant in Maine, 2 in Massachusetts, 1 in Rhode Island, 6 in Connecticut, 10 in New York, 21 in New Jersey, 3 in Pennsylvania, 1 in Maryland, 3 in Florida, 2 in Oregon, and 2 in California.

FISH UTILIZED AND PRODUCTS OF THE MENHADEN INDUSTRY

| State | Menhad-en utilized | Products | | | | | | |
|---|-----------------------------|-----------------------|------------------------|------------------|------------|-------------|------------|-------------|
| | | Dry scrap and meal | | Acidulated scrap | | Oil | | Total |
| | | Number | Tons | Value | Tons | Value | Gallons | |
| New Jersey, Delaware, Georgia and Florida | 270, 860, 000 ²¹ | 14, 304 | \$419, 675 | 5, 917 | \$116, 547 | 1, 554, 482 | \$196, 846 | \$733, 068 |
| Virginia | 192, 220, 000 | 13, 846 | 434, 647 | | | 1, 555, 995 | 231, 117 | 666, 764 |
| North Carolina | 70, 839, 000 | 3, 303 | 108, 290 | 3, 564 | 56, 733 | 233, 866 | 23, 007 | 188, 030 |
| Total | ²¹ 533, 919, 000 | ²² 31, 453 | ²³ 962, 612 | 9, 481 | 173, 280 | 3, 344, 343 | 450, 970 | 1, 586, 862 |

²¹ 320,352,000 pounds.²² Of this production, 25,068 tons, valued at \$767,878 were reported as dry scrap, and 6,385 tons, valued at \$194,734, as fish meal.

NOTE.—The menhaden factories were located as follows: 2 in New Jersey, 2 in Delaware, 11 in Virginia, 9 in North Carolina, 1 in Georgia, and 5 in Florida.

PRODUCTION OF MISCELLANEOUS BYPRODUCTS

| Product | Atlantic and Gulf coasts | | Pacific coast (including Alaska) | | Total | |
|-----------------------------|--------------------------|-----------|----------------------------------|-------------------|----------|-------------|
| | Quantity | Value | Quantity | Value | Quantity | Value |
| Dried scrap: | | | | | | |
| Alewife | 612 | \$15, 777 | | | 612 | \$15, 777 |
| Crab, blue and king | 1, 106 | 25, 777 | | | 1, 106 | 25, 777 |
| Miscellaneous ²⁴ | 2, 474 | 50, 919 | (²⁴) | (²⁴) | 2, 474 | 50, 919 |
| Acidulated and green scrap | 693 | 8, 007 | | | 693 | 8, 007 |
| Meal: | | | | | | |
| Herring (Alaska) | | | 11, 015 | \$349, 522 | 11, 015 | 349, 522 |
| Herring (Maine) | 898 | 24, 601 | | | 898 | 24, 601 |
| Pilchard | | | 50, 581 | 1, 530, 218 | 50, 581 | 1, 530, 218 |
| Salmon | | | 1, 657 | 50, 477 | 1, 657 | 50, 477 |
| Tuna | | | 6, 004 | 153, 300 | 6, 004 | 153, 300 |
| Mackerel | | | 1, 278 | 35, 554 | 1, 278 | 35, 554 |
| Ground fish "white fish" | 9, 083 | 402, 097 | | | 9, 083 | 402, 097 |
| Blue crab | 800 | 11, 000 | | | 800 | 11, 000 |
| Shrimp | 950 | 17, 655 | 229 | 6, 885 | 1, 179 | 24, 540 |
| Miscellaneous ²⁵ | 419 | 16, 940 | 1, 258 | 42, 075 | 1, 677 | 59, 615 |

²⁴ Includes ground fish, herring, and miscellaneous dried scrap.²⁵ A quantity of miscellaneous dried scrap produced by two firms in California is included with the production of the Atlantic and Gulf coasts.²⁶ Includes clam, king crab, salmon egg, whale meat and bone, and miscellaneous meal and fish flour.

Canned fishery products and byproducts of the United States and Alaska, 1933—Con.

PRODUCTION OF MISCELLANEOUS BYPRODUCTS—Con.

| Product | Atlantic and Gulf coasts | | Pacific coast (including Alaska) | | Total | |
|---|--------------------------|-----------------------|----------------------------------|-----------|------------|-----------|
| | Quantity | Value | Quantity | Value | Quantity | Value |
| Oil: | | | | | | |
| Alewife.....gallons.. | 30,900 | \$3,518 | | | 30,900 | \$3,518 |
| Cod and cod-liver: | | | | | | |
| Medicinal.....do..... | 19,552 | 14,708 | | | 19,552 | 14,708 |
| Industrial.....do..... | 45,331 | 22,152 | | | 45,331 | 22,152 |
| Herring (Alaska).....do..... | | | 3,104,821 | \$394,194 | 3,104,821 | 394,194 |
| Herring (Maine).....do..... | 69,391 | 7,963 | | | 69,391 | 7,963 |
| Pilchard.....do..... | | | 10,263,776 | 1,593,088 | 10,263,776 | 1,593,088 |
| Salmon.....do..... | | | 187,801 | 29,761 | 187,801 | 29,761 |
| Mackerel.....do..... | | | 83,778 | 10,302 | 83,778 | 10,302 |
| Whale: | | | | | | |
| Sperm.....do..... | | | 14,200 | 2,900 | 14,200 | 2,900 |
| Other.....do..... | | | 569,850 | 93,341 | 569,850 | 93,341 |
| Miscellaneous ¹⁸do..... | 17 6,479 | 17 1,622 | (17) | (17) | 6,479 | 1,622 |
| Liquid glue.....do..... | ¹⁸ 355,949 | ¹⁸ 704,101 | (18) | (18) | 355,949 | 704,101 |
| Miscellaneous byproducts ¹⁹do..... | | 93,405 | | 279,713 | | 373,118 |
| Total..... | | 1,420,242 | | 4,571,930 | | 5,992,172 |

¹⁸ Includes shark, shark-liver, and miscellaneous oil.

¹⁷ A quantity of shark-liver oil produced by one firm in California is included with the production of miscellaneous oil of the Atlantic and Gulf States.

¹⁸ A quantity of liquid glue produced by one firm in California is included with the production of liquid glue of the Atlantic and Gulf States.

¹⁹ Includes pearl essence, fish scale novelties, shark skins and fins, isinglass, agar, and kelp products.

NOTE.—The oils produced on the Pacific coast are reported in trade gallons (7¼ pounds) and those produced on the Atlantic and Gulf coasts are reported in United States gallons (about 7.74 pounds).

FROZEN FISH TRADE ³

FISH FROZEN

During 1933 the output of freezing plants which reported their activities to the Government, amounted to 95,873,507 pounds of frozen fishery products. These products at the time they were held in cold storage plants were estimated to be valued at about \$8,000,000. Compared with the pack in 1932 this was an increase of 4 percent. Six species or groups of fishery products comprised 68 percent of the pack. First in importance was the cod, haddock, haddock fillet, hake, and pollock group with 19 percent of the total, haddock fillets accounting for the bulk of the volume of this group. Next in importance was halibut, accounting for 14 percent of the total. Following was salmon with 12 percent of the total; mackerel, 11 percent; whiting, 8 percent; and shellfish, 4 percent. Other products frozen in considerable quantities during the year included sea herring, butterfish, cisco or lake herring, sablefish, smelts, weakfish or sea trout, and whitefish.

³ The statistics in this section have been furnished by the Bureau of Agricultural Economics, Department of Agriculture.

Production of frozen fishery products, 1933

BY SPECIES AND MONTHS

| Species | Month ended the 15th of— | | | | | | |
|--|--------------------------|------------------|------------------|------------------|------------------|-------------------|-------------------|
| | January | February | March | April | May | June | July |
| | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds |
| Bluefish (all trade sizes)..... | 6,987 | 8,969 | 7,106 | 3,403 | 13,881 | 25,391 | 18,997 |
| Butterfish (all trade sizes)..... | 3,012 | 4,648 | 3,995 | 1,012 | 89,821 | 156,826 | 106,244 |
| Catfish..... | 40,989 | 31,344 | 14,089 | 36,676 | 75,384 | 83,059 | 60,567 |
| Cisco (Lake Erie)..... | 9,500 | | | 1,329 | 336 | 8,487 | 6,218 |
| Cisco (lake herring), including bluefin, blackfin, and chub..... | 57,258 | 28,758 | 1,710 | 11,332 | 20,605 | 118,097 | 153,667 |
| Cisco (tullibees, Canadian lakes)..... | 38,451 | 70,695 | 2,298 | 17,300 | 9,925 | 20,001 | 8,453 |
| Cod, haddock, hake, and pollock..... | 159,543 | 178,839 | 143,675 | 334,516 | 375,788 | 195,921 | 497,582 |
| Croaker..... | 5,208 | 7,355 | 3,360 | 15,425 | 158,939 | 165,396 | 186,522 |
| Flounders..... | 20,308 | 13,475 | 2,454 | 34,385 | 131,818 | 97,721 | 46,197 |
| Haddock filets..... | 315,534 | 299,068 | 697,544 | 1,448,434 | 1,616,094 | 1,371,660 | 1,020,309 |
| Halibut (all trade sizes)..... | 172,842 | 214,224 | 955,911 | 377,502 | 813,457 | 1,819,825 | 2,609,855 |
| Herring, sea (including alewives and bluebacks)..... | 51,826 | 17,990 | 104,656 | 268,707 | 245,166 | 238,778 | 72,961 |
| Lake trout..... | 9,339 | 24,070 | 5,329 | 10,573 | 30,365 | 116,365 | 24,560 |
| Mackerel (except Spanish)..... | 60,751 | 101,913 | 69,234 | 85,765 | 130,637 | 1,860,424 | 1,096,802 |
| Pike, blue and sauger..... | 20,932 | 22,095 | 345 | | 143,906 | 101,927 | 40,965 |
| Pike, yellow or wall-eyed..... | 84,858 | 15,687 | 47,960 | 10,306 | 27,224 | 19,856 | 10,901 |
| Pike (including pickerel, jacks, and yellow jack)..... | 14,606 | 5,453 | 7,106 | 1,029 | 7,661 | 27,680 | 8,572 |
| Sablefish (black cod)..... | 17,769 | 13,719 | 18,638 | 11,245 | 69,716 | 17,762 | 36,314 |
| Salmon, chinook or king..... | 2,374 | 12,995 | 15 | 2,325 | 81,659 | 22,006 | 198,695 |
| Salmon, silver or coho..... | 34,547 | 34,889 | 60,008 | 36,390 | 7,165 | 906 | 45,171 |
| Salmon, fall and pink..... | 43,571 | 29,748 | 9,470 | 33,835 | 23,293 | 620 | 9,258 |
| Salmon, steelhead trout..... | 18,764 | 460 | 2,387 | | 7,775 | 4,963 | 72,148 |
| Salmon, red or sockeye..... | 32,009 | 63,053 | 81,342 | 14,918 | 15,621 | 53,211 | 47,988 |
| Scup (porgies)..... | 300 | 1,198 | 287 | | 11,570 | 157,918 | 20,439 |
| Shad and shad roe..... | 6,085 | 10,491 | 16,691 | 13,221 | 39,823 | 56,578 | 145,826 |
| Shellfish..... | 348,688 | 304,169 | 168,035 | 67,970 | 172,108 | 419,557 | 274,152 |
| Smelts, eulachon, etc..... | 257,142 | 287,673 | 157,959 | 32,837 | 18,790 | 4,176 | 4,902 |
| Squid..... | 8,461 | 7,366 | 6,028 | | 68,658 | 416,676 | 210,460 |
| Sturgeon and spoonbill cat..... | 2,390 | 3,422 | 11,996 | 14,875 | 3,059 | 26,865 | 22,020 |
| Stickers..... | 400 | 100 | 1,935 | 100 | 10,241 | 15,378 | 10,714 |
| Weakfish (including southern "sea trout")..... | 14,947 | 59,591 | 47,712 | 9,426 | 161,204 | 95,661 | 33,003 |
| Whitefish..... | 34,470 | 50,120 | 43,270 | 26,477 | 16,427 | 44,696 | 152,837 |
| Whiting..... | 85,302 | 83,560 | 76,237 | 46,730 | 38,120 | 1,870,586 | 3,215,618 |
| Miscellaneous fish..... | 732,671 | 744,212 | 348,090 | 647,111 | 1,029,650 | 872,624 | 773,723 |
| Total..... | 2,684,804 | 2,751,249 | 3,106,862 | 3,605,184 | 5,863,579 | 10,509,549 | 11,282,640 |

| Species | Month ended the 15th of— | | | | | |
|--|--------------------------|-----------|-----------|-----------|----------|------------|
| | August | September | October | November | December | Total |
| | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds |
| Bluefish (all trade sizes)..... | 185,488 | 92,209 | 206,571 | 107,768 | 13,951 | 690,821 |
| Butterfish (all trade sizes)..... | 25,263 | 224,446 | 790,496 | 198,362 | 7,364 | 1,611,509 |
| Catfish..... | 24,224 | 13,806 | 14,075 | 35,004 | 64,931 | 484,130 |
| Cisco (Lake Erie)..... | 17,470 | 28,146 | 4,755 | 4,532 | 36,046 | 116,819 |
| Cisco (lake herring), including bluefin, blackfin, and chub..... | 171,911 | 300,448 | 71,790 | 210,922 | 789,010 | 1,935,508 |
| Cisco (tullibees, Canadian lakes)..... | 14,503 | 10,394 | 5,080 | 10,000 | 33,922 | 242,932 |
| Cod, haddock, hake, and pollock..... | 818,695 | 356,680 | 348,738 | 558,289 | 662,744 | 4,628,890 |
| Croaker..... | 169,974 | 54,141 | 20,083 | 126,872 | 52,895 | 964,170 |
| Flounders..... | 77,963 | 26,745 | 38,866 | 59,669 | 91,769 | 639,993 |
| Haddock filets..... | 1,857,486 | 1,637,281 | 1,831,203 | 1,124,859 | 714,688 | 13,934,160 |
| Halibut (all trade sizes)..... | 2,809,652 | 1,641,819 | 962,976 | 1,165,043 | 438,231 | 13,881,337 |
| Herring, sea (including alewives and bluebacks)..... | 255,598 | 664,433 | 432,588 | 600,895 | 414,520 | 3,368,118 |
| Lake trout..... | 30,358 | 23,098 | 80,038 | 315,361 | 146,650 | 825,035 |
| Mackerel (except Spanish)..... | 3,164,049 | 2,314,254 | 1,146,051 | 391,168 | 113,286 | 10,534,336 |
| Pike, blue and sauger..... | 2,681 | 7,514 | 29,197 | 110,695 | 283,861 | 763,821 |
| Pike, yellow or wall-eyed..... | 15,496 | 30,142 | 78,399 | 28,522 | 8,280 | 377,661 |
| Pike (including pickerel, jacks, and yellow jack)..... | 32,394 | 35,658 | 16,872 | 25,438 | 7,114 | 189,488 |
| Sablefish (black cod)..... | 104,537 | 178,351 | 325,536 | 238,488 | 40,978 | 1,073,073 |
| Salmon, chinook or king..... | 912,615 | 632,703 | 394,187 | 75,682 | 70,657 | 2,405,913 |
| Salmon, silver or coho..... | 1,399,692 | 1,688,842 | 1,696,295 | 1,354,783 | 42,371 | 6,387,989 |
| Salmon, fall and pink..... | 59,399 | 98,166 | 153,200 | 991,473 | 67,482 | 1,524,445 |
| Salmon, steelhead trout..... | 318,598 | 40,855 | 30,195 | 16,820 | 50,276 | 562,251 |
| Salmon, red or sockeye..... | 77,441 | 141,015 | 180,149 | 43,567 | 96,039 | 646,353 |
| Scup (porgies)..... | 2,561 | 53,184 | 3,870 | 2,350 | 200 | 283,867 |
| Shad and shad roe..... | 91,142 | 83,171 | 2,474 | 12,592 | 3,899 | 491,993 |
| Shellfish..... | 215,224 | 435,134 | 497,062 | 606,967 | 716,180 | 4,217,246 |

Production of frozen fishery products, 1933—Continued

BY SPECIES AND MONTHS—Continued

| Species | Month ended the 15th of— | | | | | |
|---|--------------------------|---------------|---------------|---------------|---------------|---------------|
| | August | September | October | November | December | Total |
| | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> |
| Smelts, eulachon, etc. | 28,412 | 25,794 | 26,573 | 55,988 | 174,451 | 1,075,507 |
| Squid | 57,026 | 43,084 | 17,143 | 37,259 | 1,122 | 872,283 |
| Sturgeon and spoonbill cat | 41,869 | 23,213 | 39,159 | 13,102 | 5,122 | 207,092 |
| Suckers | 740 | 691 | 1,966 | 6,564 | 2,195 | 51,024 |
| Weakfish (including southern "sea trout") | 22,068 | 80,443 | 225,630 | 388,402 | 64,190 | 1,202,277 |
| Whitefish | 345,604 | 251,224 | 220,730 | 50,695 | 72,314 | 1,308,866 |
| Whiting | 1,171,751 | 258,186 | 170,947 | 136,046 | 159,495 | 7,285,578 |
| Miscellaneous fish | 1,018,282 | 1,234,096 | 927,206 | 1,321,189 | 1,280,111 | 10,928,967 |
| Total | 15,550,306 | 12,629,261 | 10,988,872 | 10,424,266 | 6,726,956 | 95,873,507 |

BY GEOGRAPHICAL SECTIONS AND SPECIES¹

[Expressed in thousands of pounds; that is, 000 omitted]

| Species | New England | Middle Atlantic | South Atlantic | North Central, East | North Central, West | South Central | Pacific | Total |
|---|-------------|-----------------|----------------|---------------------|---------------------|---------------|---------|--------|
| Bluefish (all trade sizes) | 40 | 584 | 3 | 28 | 3 | 23 | | 691 |
| Butterfish (all trade sizes) | 135 | 1,435 | 17 | 35 | | | | 1,612 |
| Catfish | 94 | | 67 | 72 | 207 | 44 | | 484 |
| Cisco (Lake Erie) | | 114 | | 3 | | | | 117 |
| Cisco (lake herring), including bluefin, blackfin, and chub | | 695 | | 966 | 275 | | | 1,936 |
| Cisco (tullibees, Canadian lakes) | 40 | 159 | | 30 | 14 | | | 423 |
| Cod, haddock, hake, and pollock | 4,178 | 116 | 1 | 55 | 180 | | 99 | 4,629 |
| Croaker | | 245 | 572 | 143 | | 4 | | 964 |
| Flounders | 273 | 335 | | 14 | | 3 | 15 | 640 |
| Haddock filets | 13,099 | 138 | 14 | 565 | 81 | | 37 | 13,934 |
| Hallbut (all trade sizes) | 114 | 557 | 53 | 904 | 30 | 25 | 12,198 | 13,881 |
| Herring, sea (including alewives and bluebacks) | 2,128 | 207 | 6 | 406 | 1 | 5 | 615 | 3,368 |
| Lake trout | 2 | 144 | | 565 | 106 | 8 | | 825 |
| Mackerel (except Spanish) | 9,034 | 1,121 | 3 | 263 | 20 | 28 | 66 | 10,535 |
| Pike, blue and sauger | | 490 | | 272 | 1 | 1 | | 764 |
| Pike, yellow or wall-eyed | | 157 | | 128 | 93 | | | 378 |
| Pike (including pickerel, jacks, and yellow jack) | | 17 | | 67 | 105 | | | 189 |
| Sablefish (black cod) | | | | 123 | 13 | | 937 | 1,073 |
| Salmon, king or chinook | 56 | 57 | | 40 | 9 | | 2,244 | 2,406 |
| Salmon, silver or coho | 20 | 219 | 2 | 19 | 4 | | 6,124 | 6,388 |
| Salmon, fall and pink | 6 | 27 | | 19 | 3 | | 1,409 | 1,524 |
| Salmon, steelhead trout | | 36 | 9 | | | | 517 | 562 |
| Salmon, sockeye or red | | 206 | | 122 | 23 | 3 | 492 | 846 |
| Soup (porgies) | 85 | 166 | 3 | | | | | 254 |
| Shad and shad roe | 321 | 77 | | 37 | 2 | 2 | 43 | 482 |
| Shellfish | 505 | 1,785 | 809 | 559 | 217 | 7 | 335 | 4,217 |
| Smelts, eulachon, etc. | 47 | 874 | 1 | 85 | 3 | | 116 | 1,076 |
| Squid | 510 | 350 | | 3 | | | 9 | 872 |
| Sturgeon and spoonbill cat | | 137 | 2 | | 21 | 27 | 12 | 207 |
| Suckers | | | | 50 | 1 | | | 51 |
| Weakfish (including southern "sea trout") | 1 | 916 | 285 | | | | | 1,202 |
| Whitefish | 5 | 930 | | 363 | 8 | 2 | 1 | 1,309 |
| Whiting | 6,472 | 591 | 2 | 11 | 210 | | | 7,286 |
| Miscellaneous fish | 2,439 | 1,373 | 1,207 | 2,202 | 518 | 1,619 | 1,571 | 10,929 |
| Total | 39,604 | 14,258 | 3,056 | 8,107 | 2,148 | 1,801 | 26,900 | 95,874 |

¹ New England includes the six States of that section; Middle Atlantic—New York, New Jersey, and Pennsylvania; South Atlantic—Delaware, Maryland, District of Columbia, Virginia, West Virginia, North Carolina, South Carolina, Georgia, and Florida; North Central, East—Ohio, Indiana, Illinois, Michigan, and Wisconsin; North Central, West—Minnesota, Iowa, Missouri, North Dakota, South Dakota, Nebraska, and Kansas; South Central—Kentucky, Tennessee, Alabama, Mississippi, Louisiana, Texas, Oklahoma, and Arkansas; Pacific—Washington, Oregon, California, and Alaska.

Production of frozen fishery products, 1933—Continued

BY GEOGRAPHICAL SECTIONS AND MONTHS

[Expressed in thousands of pounds; that is, 000 omitted]

| Month ended the 15th of— | New Eng-land | Middle Atlan-tic | South Atlan-tic | North Cen-tral, East | North Cen-tral, West | South Cen-tral | Pacific | Total |
|--------------------------|--------------|------------------|-----------------|----------------------|----------------------|----------------|---------|--------|
| January..... | 475 | 878 | 294 | 519 | 95 | 139 | 285 | 2,685 |
| February..... | 411 | 1,007 | 373 | 428 | 70 | 165 | 299 | 2,751 |
| March..... | 768 | 674 | 90 | 252 | 126 | 132 | 1,065 | 3,107 |
| April..... | 1,809 | 246 | 211 | 214 | 149 | 96 | 880 | 3,605 |
| May..... | 2,367 | 734 | 345 | 685 | 130 | 223 | 1,180 | 5,664 |
| June..... | 5,606 | 1,499 | 178 | 1,080 | 135 | 112 | 1,900 | 10,510 |
| July..... | 6,294 | 842 | 266 | 592 | 173 | 122 | 2,944 | 11,233 |
| August..... | 7,640 | 1,205 | 261 | 522 | 187 | 167 | 5,568 | 15,580 |
| September..... | 5,920 | 1,491 | 55 | 528 | 166 | 203 | 4,266 | 12,629 |
| October..... | 3,733 | 2,455 | 81 | 530 | 262 | 148 | 3,780 | 10,989 |
| November..... | 2,736 | 1,631 | 193 | 1,309 | 336 | 138 | 4,081 | 10,424 |
| December..... | 1,845 | 1,596 | 709 | 1,450 | 319 | 156 | 652 | 6,727 |
| Total..... | 39,604 | 14,258 | 3,056 | 8,107 | 2,148 | 1,801 | 26,900 | 95,874 |

HOLDINGS

During 1933 monthly holdings of frozen fish and shellfish averaged 39,386,000 pounds, which is a decrease of 17 percent as compared with the average monthly holdings in 1932. The holdings during November were largest, amounting to 58,338,000 pounds; however, the holdings during each of the months from September to December inclusive exceeded 50,000,000 pounds. The holdings during April were smallest when only 19,335,000 pounds of frozen fishery products were in storage.

Holdings of frozen fishery products, 1933

BY SPECIES AND MONTHS

| Species | Month ended the 15th of— | | | | | |
|--|--------------------------|------------|------------|------------|------------|------------|
| | January | February | March | April | May | June |
| Bluefish (all trade sizes)..... | 559,958 | 435,422 | 347,239 | 269,480 | 248,963 | 242,194 |
| Butterfish (all trade sizes)..... | 338,783 | 212,110 | 104,304 | 52,327 | 114,287 | 238,344 |
| Catfish..... | 298,454 | 310,491 | 184,515 | 115,312 | 116,312 | 205,495 |
| Cisco (Lake Erie)..... | 185,365 | 226,019 | 169,994 | 109,912 | 70,406 | 35,578 |
| Cisco (Lake herring), including bluefin, blackfin, and chub..... | 642,936 | 383,448 | 120,022 | 50,850 | 97,241 | 405,164 |
| Cisco (tullibees, Canadian lakes)..... | 519,842 | 574,534 | 399,555 | 369,894 | 365,597 | 293,063 |
| Cod, haddock, bake, and pollock..... | 711,656 | 572,020 | 307,200 | 378,341 | 640,836 | 701,109 |
| Croaker..... | 677,052 | 465,963 | 200,309 | 99,023 | 193,385 | 295,893 |
| Flounders..... | 213,749 | 167,560 | 91,124 | 92,739 | 184,057 | 191,793 |
| Haddock fillets..... | 2,119,821 | 1,440,810 | 868,494 | 1,492,497 | 2,549,574 | 2,892,345 |
| Hallbut (all trade sizes)..... | 2,817,401 | 1,785,179 | 1,531,530 | 1,458,041 | 1,877,057 | 3,534,446 |
| Herring, sea (including alewives and bluebacks)..... | 1,573,849 | 1,216,964 | 1,009,421 | 963,268 | 1,049,428 | 955,260 |
| Lake trout..... | 591,939 | 440,842 | 194,651 | 67,641 | 89,183 | 194,227 |
| Mackerel (except Spanish)..... | 10,328,815 | 7,914,664 | 5,617,720 | 4,131,263 | 3,326,512 | 4,522,169 |
| Pike, blue and sauger..... | 326,299 | 186,901 | 90,121 | 24,889 | 149,893 | 246,037 |
| Pike, yellow or wall-eyed..... | 283,738 | 510,054 | 412,613 | 173,974 | 169,628 | 95,776 |
| Pike (including pickerel, jacks, and yellow jack)..... | 241,928 | 223,587 | 233,229 | 149,677 | 142,447 | 157,669 |
| Sablefish (black cod)..... | 696,155 | 564,961 | 454,893 | 337,914 | 298,243 | 245,779 |
| Salmon, chinook or king..... | 1,634,799 | 1,392,399 | 961,432 | 517,196 | 263,207 | 190,445 |
| Salmon, silver or coho..... | 3,639,900 | 2,350,943 | 1,151,225 | 564,037 | 261,091 | 217,694 |
| Salmon, fall and pink..... | 913,521 | 601,835 | 467,685 | 405,231 | 398,988 | 286,894 |
| Salmon, steelhead trout..... | 402,223 | 352,581 | 176,530 | 149,293 | 119,810 | 112,911 |
| Salmon, red or sockeye..... | 650,882 | 586,203 | 307,500 | 250,383 | 174,003 | 123,236 |
| Scup (porgy)..... | 176,846 | 128,328 | 63,661 | 14,318 | 15,195 | 168,943 |
| Shad and shad roe..... | 224,448 | 187,738 | 139,641 | 89,706 | 107,700 | 126,789 |
| Shellfish..... | 1,911,491 | 1,347,992 | 1,387,739 | 939,533 | 832,412 | 1,043,363 |
| Smelts, eulachon, etc..... | 344,100 | 797,367 | 1,001,143 | 637,430 | 570,138 | 514,727 |
| Squid..... | 1,568,225 | 1,358,679 | 949,980 | 555,421 | 481,787 | 732,305 |
| Sturgeon and spoonbill cat..... | 1,015,871 | 944,398 | 844,254 | 687,991 | 604,613 | 580,370 |
| Suckers..... | 13,000 | 12,305 | 8,902 | 3,748 | 12,404 | 27,410 |
| Weakfish (including southern "sea trout")..... | 378,693 | 351,864 | 208,129 | 82,780 | 213,998 | 285,282 |
| Whitefish..... | 1,312,443 | 1,119,443 | 1,151,730 | 623,737 | 457,872 | 312,506 |
| Whiting..... | 2,919,007 | 1,987,544 | 1,285,482 | 583,907 | 325,040 | 2,123,489 |
| Miscellaneous fish..... | 5,241,234 | 4,327,068 | 3,179,268 | 2,893,112 | 3,123,304 | 3,473,048 |
| Total..... | 45,476,122 | 35,468,516 | 25,854,938 | 19,334,869 | 19,644,637 | 25,711,323 |

Holdings of frozen fishery products, 1933—Continued

BY SPECIES AND MONTHS—Continued

| Species | Month ended the 15th of— | | | | | |
|---|--------------------------|------------|------------|------------|------------|------------|
| | July | August | September | October | November | December |
| | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds |
| Bluefish (all trade sizes)..... | 200,182 | 403,013 | 397,554 | 530,399 | 569,529 | 463,416 |
| Butterfish (all trade sizes)..... | 304,718 | 305,680 | 493,657 | 1,166,770 | 1,166,206 | 996,213 |
| Catfish..... | 247,078 | 220,992 | 188,331 | 169,853 | 158,686 | 218,372 |
| Cisco (Lake Erie)..... | 43,676 | 54,903 | 80,299 | 80,902 | 70,921 | 96,969 |
| Cisco (lake herring), including bluefin, blackfin, and chub..... | 608,124 | 895,361 | 1,119,058 | 1,202,198 | 1,282,964 | 1,931,787 |
| Cisco (tullbees, Canadian lakes)..... | 280,144 | 282,035 | 223,554 | 225,609 | 169,036 | 303,956 |
| Cod, haddock, hake, and pollock..... | 1,117,222 | 1,688,080 | 1,531,219 | 1,212,754 | 1,429,789 | 1,843,168 |
| Croaker..... | 457,792 | 568,898 | 548,210 | 520,290 | 511,073 | 522,778 |
| Flounders..... | 188,961 | 244,312 | 210,006 | 196,141 | 223,986 | 265,347 |
| Haddock fillets..... | 3,039,853 | 4,255,986 | 4,575,522 | 5,590,470 | 6,349,530 | 4,980,536 |
| Halibut (all trade sizes)..... | 5,875,009 | 8,372,420 | 9,507,466 | 9,767,880 | 9,243,309 | 7,792,242 |
| Herring, sea (including alewives and blue- backs)..... | 796,715 | 845,985 | 1,263,856 | 1,435,579 | 2,023,058 | 2,016,529 |
| Lake trout..... | 180,340 | 175,023 | 174,598 | 269,536 | 566,689 | 702,037 |
| Mackerel (except Spanish)..... | 5,044,686 | 7,635,683 | 8,839,982 | 9,015,455 | 7,864,902 | 6,928,127 |
| Pike, blue and sauger..... | 171,917 | 29,832 | 27,785 | 131,874 | 360,420 | 754,228 |
| Pike, yellow or wall-eyed..... | 94,201 | 74,079 | 111,727 | 168,761 | 131,178 | 189,019 |
| Pike (including pickerel, jacks, and yel- low jack)..... | 121,832 | 130,037 | 156,886 | 153,858 | 152,016 | 174,791 |
| Sablefish, (black cod)..... | 264,425 | 315,896 | 392,060 | 614,749 | 789,817 | 744,083 |
| Salmon, chinook or king..... | 305,232 | 1,088,979 | 1,630,992 | 1,890,125 | 1,937,629 | 1,686,661 |
| Salmon, silver or coho..... | 251,069 | 1,352,722 | 2,807,859 | 4,175,741 | 5,302,877 | 3,508,861 |
| Salmon, fall and pink..... | 221,546 | 247,481 | 320,926 | 335,047 | 1,192,622 | 2,326,835 |
| Salmon, steelhead trout..... | 157,676 | 315,006 | 357,643 | 377,532 | 234,291 | 282,063 |
| Salmon, red or sockeye..... | 105,838 | 121,427 | 219,554 | 365,603 | 346,922 | 354,713 |
| Scup (porgy)..... | 204,253 | 337,660 | 415,015 | 379,399 | 386,933 | 389,991 |
| Shad and shad roe..... | 981,724 | 895,879 | 1,055,641 | 1,264,341 | 1,699,591 | 2,024,535 |
| Shellfish..... | 500,734 | 517,146 | 526,927 | 518,504 | 437,173 | 430,666 |
| Smelts, eulachon, etc..... | 828,188 | 731,668 | 633,377 | 537,180 | 500,149 | 363,533 |
| Squid..... | 460,495 | 357,055 | 276,883 | 261,110 | 272,584 | 405,263 |
| Sturgeon and spoonbill cat..... | 36,245 | 35,390 | 32,430 | 32,094 | 21,414 | 29,978 |
| Suckers..... | 272,347 | 279,232 | 328,761 | 531,953 | 782,728 | 668,234 |
| Weak fish (including southern "sea trout")..... | 1,229,976 | 2,180,867 | 2,525,231 | 2,502,796 | 2,511,392 | 2,729,242 |
| Whitefish..... | 5,020,126 | 5,652,475 | 5,634,893 | 5,283,745 | 4,977,054 | 4,799,664 |
| Whiting..... | 3,477,144 | 4,091,137 | 4,703,111 | 4,956,142 | 5,488,579 | 6,105,432 |
| Miscellaneous fish..... | | | | | | |
| Total..... | 33,330,914 | 44,882,133 | 51,474,979 | 55,927,781 | 58,338,360 | 57,188,489 |

BY GEOGRAPHICAL SECTIONS AND MONTHS ¹

[Expressed in thousands of pounds; that is, 000 omitted]

| Month ended the 15th of— | New England | Middle Atlantic | South Atlantic | North Central, East | North Central, West | South Central | Pacific ² | Total |
|--------------------------|-------------|-----------------|----------------|---------------------|---------------------|---------------|----------------------|--------|
| January..... | 16,120 | 9,937 | 2,781 | 4,760 | 2,402 | 524 | 8,952 | 45,476 |
| February..... | 12,271 | 8,843 | 2,045 | 3,825 | 2,246 | 456 | 5,783 | 35,469 |
| March..... | 8,348 | 6,877 | 1,298 | 2,370 | 1,857 | 340 | 4,765 | 25,855 |
| April..... | 6,173 | 4,847 | 1,048 | 1,610 | 1,441 | 284 | 3,932 | 19,335 |
| May..... | 6,130 | 4,462 | 1,066 | 2,277 | 1,493 | 396 | 3,821 | 19,645 |
| June..... | 9,916 | 4,988 | 1,060 | 2,883 | 1,592 | 330 | 4,962 | 25,711 |
| July..... | 14,474 | 5,811 | 1,231 | 2,604 | 1,435 | 268 | 7,508 | 33,331 |
| August..... | 20,071 | 7,064 | 1,427 | 2,608 | 1,408 | 329 | 11,956 | 44,882 |
| September..... | 22,598 | 7,572 | 1,257 | 2,717 | 1,587 | 404 | 15,340 | 51,475 |
| October..... | 23,015 | 9,378 | 1,306 | 2,972 | 1,708 | 397 | 17,162 | 55,928 |
| November..... | 21,389 | 10,436 | 1,361 | 4,119 | 2,083 | 427 | 18,523 | 68,338 |
| December..... | 19,342 | 11,988 | 1,980 | 5,622 | 2,383 | 485 | 16,388 | 67,188 |
| Average..... | 14,987 | 7,686 | 1,488 | 3,196 | 1,803 | 387 | 9,840 | 39,386 |

¹ New England includes the 6 States of that section; Middle Atlantic—New York, New Jersey, and Pennsylvania; South Atlantic—Delaware, Maryland, District of Columbia, Virginia, West Virginia, North Carolina, South Carolina, Georgia, and Florida; North Central, East—Ohio, Indiana, Illinois, Michigan, and Wisconsin; North Central, West—Minnesota, Iowa, Missouri, North Dakota, South Dakota, Nebraska, and Kansas; South Central—Kentucky, Tennessee, Alabama, Mississippi, Louisiana, Texas, Oklahoma, and Arkansas; Pacific—Washington, Oregon, California, and Alaska.

² Includes a very small amount of fish held in Colorado in the Mountain section.

COLD-STORAGE HOLDINGS OF CURED FISH

During 1933 monthly cold-storage holdings of cured herring and mild-cured salmon averaged 14,907,000 pounds, which is a decrease of 18 percent as compared with the average monthly holdings in 1932. The holdings during October were largest, amounting to 21,924,000 pounds, and the smallest were in April, amounting to 9,191,000 pounds.

Holdings of cured fish, 1933: By species and months

| Month ended the 15th of— | Cured herring | Mild-cured salmon | Total |
|--------------------------|---------------|-------------------|---------------|
| | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> |
| January..... | 12, 136, 374 | 3, 933, 005 | 16, 069, 379 |
| February..... | 11, 137, 364 | 3, 544, 376 | 14, 681, 740 |
| March..... | 9, 623, 604 | 2, 400, 345 | 12, 023, 949 |
| April..... | 7, 621, 553 | 1, 569, 788 | 9, 191, 341 |
| May..... | 8, 632, 589 | 2, 266, 722 | 10, 919, 311 |
| June..... | 9, 211, 715 | 1, 402, 142 | 10, 613, 857 |
| July..... | 9, 082, 692 | 1, 696, 780 | 10, 779, 472 |
| August..... | 9, 236, 158 | 5, 169, 901 | 14, 406, 059 |
| September..... | 12, 324, 553 | 6, 631, 918 | 18, 956, 471 |
| October..... | 15, 710, 119 | 6, 214, 266 | 21, 924, 385 |
| November..... | 15, 214, 185 | 5, 928, 630 | 21, 142, 815 |
| December..... | 13, 620, 771 | 4, 576, 118 | 18, 196, 889 |

FOREIGN FISHERY TRADE

The foreign trade in fishery products of the United States in 1933 amounted to \$38,801,064 of which \$30,462,341 represents the value of these products imported for consumption, and \$8,338,723, the value of exports of domestic fishery products. Compared with the previous year, there was an increase of 4 percent in the total trade, 3 percent in the value of the imports, and 7 percent in the value of exports.

Imports consisted of 284,306,782 pounds of edible products, valued at \$21,782,874, and nonedible products, valued at \$8,679,467. Fishery exports consisted of 80,007,878 pounds of edible products, valued at \$7,376,401 and nonedible products, valued at \$962,322.

Exports of domestic fishery products, 1933

| Item | Quantity | Value |
|--|--------------------|------------|
| EDIBLE FISHERY PRODUCTS | | |
| Fish, fresh, frozen, or packed in ice: | | |
| Salmon..... | pounds 5, 754, 305 | \$545, 946 |
| Other..... | do 2, 744, 270 | 207, 106 |
| Total..... | do 8, 498, 575 | 753, 052 |
| Fish, salted, smoked, or dry-cured: | | |
| Cod..... | do 647, 419 | 58, 472 |
| Salmon..... | do 779, 232 | 108, 456 |
| Other..... | do 989, 921 | 40, 513 |
| Total..... | do 2, 416, 572 | 213, 441 |
| Fish, pickled: | | |
| Salmon..... | do 1, 941, 800 | 268, 296 |
| Other..... | do 1, 088, 000 | 43, 424 |
| Total..... | do 3, 029, 800 | 311, 720 |

FISHERY INDUSTRIES OF THE UNITED STATES, 1934 129

Exports of domestic fishery products, 1933—Continued

| Item | Quantity | Value |
|--|-------------------|------------------|
| EDIBLE FISHERY PRODUCTS—continued | | |
| Fish, canned: | | |
| Mackerel..... pounds.. | 1,968,982 | \$93,071 |
| Salmon..... do..... | 26,539,379 | 3,289,924 |
| Sardines..... do..... | 25,641,265 | 1,380,680 |
| Other..... do..... | 320,132 | 47,316 |
| Total..... do..... | 54,469,758 | 4,816,991 |
| Shellfish, not canned: | | |
| Oysters, fresh, in the shell..... do..... | 2,901,860 | 97,134 |
| Oysters, fresh, shucked, frozen, or in ice..... do..... | 1,265,060 | 161,785 |
| Shrimp, fresh, frozen, or in ice..... do..... | 1,852,664 | 176,859 |
| Shrimp, dried..... do..... | 1,646,888 | 265,561 |
| Other shellfish, fresh, frozen, in ice or dried..... do..... | 129,108 | 13,833 |
| Total..... do..... | 7,795,580 | 705,172 |
| Shellfish, canned..... do..... | 3,481,702 | 512,632 |
| Other fish products..... do..... | 315,801 | 63,383 |
| Total edible products..... do..... | 80,007,878 | 7,376,401 |
| NONEDIBLE FISHERY PRODUCTS | | |
| Marine-animal oils..... do..... | 5,849,288 | 163,138 |
| Sponges..... do..... | 72,101 | 67,727 |
| Fish meal for feed..... tons..... | 7,918 | 346,133 |
| Oyster shells..... do..... | 57,000 | 386,324 |
| Total nonedible products..... do..... | | 962,322 |
| Grand total..... do..... | | 8,338,723 |

Imports of fishery products entered for consumption, 1933

| Item | Pounds | Value |
|--|------------|-----------|
| EDIBLE FISHERY PRODUCTS | | |
| Fish, fresh or frozen: | | |
| Whole, beheaded, or eviscerated, or both: | | |
| Salmon..... | 5,083,345 | \$364,786 |
| Fresh-water fish, not elsewhere specified: | | |
| Yellow pike..... | 8,498,767 | 589,015 |
| Whitefish..... | 10,029,807 | 871,847 |
| Tullibees..... | 1,535,130 | 46,425 |
| Jacks or grass pike..... | 1,825,550 | 86,837 |
| Lake trout..... | 1,766,268 | 167,389 |
| Yellow perch..... | 1,077,969 | 71,347 |
| Lake herring, ciscoes, and chubs..... | 992,572 | 101,667 |
| Fresh-water fish, not elsewhere specified..... | 14,434,347 | 839,508 |
| Eels..... | 237,735 | 19,626 |
| Cod, haddock, hake, pollock, and cusk..... | 593,186 | 26,769 |
| Halibut: | | |
| Fresh..... | 3,015,619 | 230,834 |
| Frozen..... | 1,070,753 | 89,367 |
| Mackerel..... | 507,816 | 22,719 |
| Swordfish..... | 2,633,935 | 232,349 |
| Sturgeon..... | 1,980,794 | 222,685 |
| Fish, not specially provided for..... | 2,154,022 | 108,205 |
| Whether or not whole: | | |
| Smelts..... | 6,726,196 | 672,068 |
| Tuna fish..... | 5,935,957 | 232,987 |
| See herring: | | |
| Fresh..... | 23,899,000 | 95,197 |
| Frozen..... | 1,963,474 | 52,535 |
| Fillets, skinned, boned, sliced, or divided, not specially provided for..... | 2,504,541 | 261,523 |
| Total..... | 98,446,783 | 5,440,675 |
| Fish, salted, dried, smoked, pickled or preserved: | | |
| Dried and unsalted: | | |
| Cod, haddock, hake, pollock, and cusk..... | 27,721 | 2,812 |
| Other..... | 3,150,758 | 295,311 |
| In oil or in oil and other substances: | | |
| Sardines..... | 32,555,513 | 2,811,197 |
| Anchovies..... | 1,724,766 | 455,718 |
| Antipasto..... | 158,866 | 55,587 |
| Tuna..... | 14,382,256 | 1,969,144 |
| Other..... | 136,416 | 20,027 |

Imports of fishery products entered for consumption, 1933—Continued

| Item | Pounds | Value |
|---|-------------|------------|
| EDIBLE FISHERY PRODUCTS—continued | | |
| Fish, salted, dried, smoked, pickled or preserved—Continued. | | |
| Not in oil or in oil and other substances: | | |
| In air-tight containers weighing with contents, not over 15 pounds each: | | |
| Anchovies..... | 2,025,739 | \$170,845 |
| Salmon..... | 4,815,768 | 227,312 |
| Herring and sardines..... | 7,699,857 | 446,378 |
| Fish cakes, balls, and pudding..... | 1,746,804 | 91,103 |
| Other..... | 1,538,446 | 133,499 |
| Pickled or salted: | | |
| Not in oil, etc., and not in air-tight containers weighing, with contents, 15 pounds or less each: | | |
| Salmon..... | 317,900 | 31,777 |
| Cod, haddock, hake, pollock, and cusk, neither skinned nor boned (except that vertebral column may be removed): | | |
| Containing not more than 43 percent moisture by weight..... | 12,164,979 | 564,847 |
| Containing more than 43 percent moisture by weight..... | 26,476,539 | 833,671 |
| Cod, haddock, hake, pollock and cusk, skinned or boned..... | 2,027,855 | 139,522 |
| Herring: | | |
| In bulk or in containers weighing, with contents, more than 15 pounds each (net weight)..... | 29,326,957 | 1,335,125 |
| In containers (not air-tight), weighing, with contents, not more than 15 pounds each..... | 133,638 | 4,704 |
| Mackerel: | | |
| In bulk or in containers weighing, with contents, more than 15 pounds each (net weight)..... | 4,403,451 | 176,341 |
| Pickled or salted, not specially provided for: | | |
| In bulk or in containers weighing, with contents, more than 15 pounds each (net weight)..... | 1,294,831 | 74,184 |
| In containers (not air-tight) weighing, with contents, not more than 15 pounds each..... | 8,750 | 881 |
| Smoked or kippered: | | |
| Not in oil, etc., and not in air-tight containers weighing, with contents, 15 pounds or less each: | | |
| Salmon..... | 2,270 | 1,109 |
| Herring: | | |
| Whole or beheaded..... | 638,607 | 27,044 |
| Eviscerated, split, skinned, boned, or divided..... | 871,021 | 73,631 |
| Cod, haddock, hake, pollock, and cusk: | | |
| Whole, or beheaded, or eviscerated, or both..... | 686,551 | 54,108 |
| Filleted, skinned, boned, sliced, or divided..... | 885,815 | 80,003 |
| Smoked or kippered, not specially provided for..... | 16,025 | 1,169 |
| Fish paste and fish sauce..... | 74,281 | 24,142 |
| Prepared or preserved, not specially provided for: | | |
| In containers weighing, with contents, not more than 15 pounds each..... | 57,504 | 6,529 |
| In bulk or in containers weighing, with contents, more than 15 pounds each (net weight)..... | 148,464 | 14,231 |
| Total..... | 150,498,348 | 10,122,941 |
| Caviar and other fish roe: | | |
| Not boiled, etc.: | | |
| Sturgeon..... | 263,147 | 248,657 |
| Fish roe, not specially provided for..... | 99,195 | 11,320 |
| Boiled, packed in air-tight containers..... | 49,227 | 3,774 |
| Total..... | 411,569 | 263,751 |
| Shellfish: | | |
| Crab meat, crab sauce, and crab paste..... | 9,526,113 | 2,893,435 |
| Clams, clam juice, or either in combination with other substances, in air-tight containers..... | 1,417,315 | 131,856 |
| Oysters, oyster juice, or either in combination with other substances in airtight containers..... | 97,381 | 19,659 |
| Lobsters (including spiny lobsters and crawfish): | | |
| Not canned..... | 10,622,543 | 1,733,988 |
| Canned..... | 1,468,796 | 485,038 |
| Clams not in airtight containers..... | 1,368,207 | 21,313 |
| Shrimp and prawn..... | 716,235 | 84,989 |
| Scallops..... | 834,556 | 114,695 |
| Oysters, not in airtight containers..... | 4,905,457 | 118,337 |
| Shellfish, not specially provided for..... | 3,622,816 | 326,087 |
| Pastes and sauces of shellfish, not specially provided for..... | 124,524 | 12,077 |
| Crabs..... | 11,380 | 920 |
| Turtles..... | 234,759 | 13,113 |
| Total..... | 34,950,082 | 5,955,507 |
| Total edible fishery products..... | 284,306,782 | 21,782,874 |

Imports of fishery products entered for consumption, 1933—Continued

| Item | Pounds | Value |
|--|--------------------|-----------------|
| NONEDIBLE FISHERY PRODUCTS | | |
| Marine-animal oils: | <i>Quantity</i> | |
| Cod oil.....gallons | 2, 115, 144 | \$449, 695 |
| Cod-liver oil.....do | 3, 432, 569 | 1, 711, 072 |
| Eulachon oil.....do | 760 | 326 |
| Herring oil.....do | 699, 296 | 42, 431 |
| Menhaden oil.....do | 1, 493 | 119 |
| Seal oil.....do | 106, 048 | 22, 939 |
| Sod oil.....do | 78, 652 | 11, 131 |
| Whale oil: | | |
| Sperm, crude.....do | 520, 212 | 119, 527 |
| Sperm, refined or otherwise processed.....do | 104, 308 | 38, 238 |
| Whale oil, not specially provided for.....do | 5, 735, 613 | 2, 398, 728 |
| Total.....do | 12, 796, 095 | 4, 794, 206 |
| Pearls and imitation pearls: | | |
| Pearls and parts, not strung or set.....do | | 681, 083 |
| Imitation pearls: | | |
| Half pearls and hollow or filled.....do | | 4, 509 |
| Solid pearls, not elsewhere specified: | | |
| Valued at not more than one-fourth cent an inch.....inches | 14, 880 | 10 |
| Valued at more than one-fourth cent and not more than 1 cent an inch.....inches | 54, 707 | 355 |
| Iridescent solid pearls: | | |
| Valued at not more than 10 cents per inch.....do | 3, 065 | 79 |
| Total.....do | | 686, 036 |
| Shells and buttons of pearl or shell: | | |
| Shells, unmanufactured: | | |
| Green snail shell.....pounds | 205, 538 | 16, 110 |
| Mother-of-pearl.....do | 4, 905, 025 | 1, 131, 197 |
| Shells, not specially provided for.....do | 4, 392, 206 | 39, 375 |
| Shells and mother-of-pearl, engraved, cut, ornamented, or manufactured.....do | | 50, 078 |
| Shell pearl buttons: | | |
| Ocean or trochus.....gross | 236, 829 | 66, 566 |
| Fresh-water.....do | 100 | 40 |
| Buttons (from Philippine Islands).....do | 749, 469 | 249, 170 |
| Total.....do | | 1, 552, 536 |
| Sponges: | | |
| Sheepswool.....pounds | 133, 477 | 202, 083 |
| Yellow, grass, or velvet.....do | 231, 621 | 79, 389 |
| Other.....do | 82, 786 | 69, 636 |
| Manufactures of.....do | 96 | 28 |
| Total.....do | 447, 980 | 351, 136 |
| Agar agar.....do | 629, 784 | 165, 627 |
| Ambergris.....do | 8 | 763 |
| Cod-liver oil cake and cod-liver oil cake meal.....do | 1, 614, 588 | 24, 744 |
| Cuttlefish bone.....do | 295, 182 | 35, 091 |
| Goldfish, live.....number | 1, 125, 012 | 7, 103 |
| Fish for other than human consumption, not elsewhere specified.....pounds | 98, 046 | 14, 223 |
| Fish sounds.....tons | 26, 526 | 689, 192 |
| Fish scrap and fish meal.....pounds | 38, 550 | 561 |
| Kelp.....square feet | 26 | 11 |
| Skins, fish, raw or salted.....pounds | 884, 412 | 31, 804 |
| Skins, seal, raw (not fur skins).....do | 1, 416, 355 | 125, 008 |
| Spermaceti wax.....do | 100, 843 | 10, 334 |
| Whalebone, unmanufactured.....do | 503 | 92 |
| Whalebone, manufactures of.....do | | 664 |
| Total.....do | | 1, 295, 553 |
| Total, nonedible fishery products.....do | | 8, 679, 467 |
| Grand total.....do | | 30, 462, 341 |

FISHERIES OF THE NEW ENGLAND STATES

(Area XXII) ⁴

The yield of the commercial fisheries of the New England States (Maine, New Hampshire, Massachusetts, Rhode Island, and Connecticut) during 1933, amounted to 499,936,139 pounds, valued at \$13,485,550 to the fishermen, representing an increase of 4 percent in volume, but a decrease of 4 percent in value as compared with the catch in the previous year. These fisheries gave employment to 17,073 fishermen as compared with 16,472, in 1932.

There were 362 fishery wholesale and manufacturing establishments in the 5 States in 1933 as compared with 436 in 1931 when the most recent previous survey of such concerns was made. In 1933 these establishments employed 9,177 persons, paid \$5,410,072 in salaries and wages, and produced manufactured products (canned, cured, packaged, and byproducts) valued at \$14,322,274. In 1931, the wholesale and manufacturing firms employed 10,273 persons, paid \$7,113,463 in salaries and wages, and produced manufactured products, valued at \$18,616,951.

Fisheries of the New England States, 1933

SUMMARY OF CATCH

| Product | Maine | | New Hampshire | | Massachusetts | |
|---------------------|------------|-----------|---------------|---------|---------------|-------------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Fish..... | 83,071,766 | \$848,662 | 306,475 | \$8,841 | 359,144,370 | \$8,196,646 |
| Shellfish, etc..... | 15,426,316 | 1,458,445 | 216,929 | 48,230 | 14,525,908 | 1,310,332 |
| Total..... | 98,498,082 | 2,307,107 | 523,404 | 57,071 | 373,670,278 | 9,506,978 |

| Product | Rhode Island | | Connecticut | | Total | |
|---------------------|--------------|-----------|-------------|-----------|-------------|-------------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Fish..... | 11,648,054 | \$282,642 | 7,450,301 | \$255,682 | 461,620,966 | \$9,692,453 |
| Shellfish, etc..... | 5,718,293 | 718,622 | 2,427,727 | 357,468 | 38,315,173 | 3,893,097 |
| Total..... | 17,366,347 | 1,001,264 | 9,878,028 | 613,130 | 499,936,139 | 13,485,550 |

⁴ This is the number given this area by the North American Council on Fishery Investigations. It should be explained that there are included under this area craft whose principal fishing ports are in the area but at times fish elsewhere. Notable examples are the ground fish fishery in area XXI and the mackerel and southern trawl fisheries in areas XIII and XIV. It should be observed that the persons engaged, gear and craft employed, and catch of the seed oyster fishery are not included among the statistics of the fishery for market oysters and other species but are shown in separate tables in this section. For a clearer understanding of the statistics published in this section, the reader is referred to the section in the latter part of this document entitled "Statistical survey procedure."

Fisheries of the New England States, 1933—Continued

OPERATING UNITS: BY STATES

| Item | Maine | New Hampshire | Massachusetts | Rhode Island | Connecticut | Total |
|-------------------------------|--------------|---------------|---------------|--------------|--------------|---------------|
| Fishermen: | | | | | | |
| On vessels..... | 443 | | 3,971 | 328 | 307 | 5,049 |
| On boats and shore: | | | | | | |
| Regular..... | 3,748 | 48 | 2,781 | 371 | 360 | 7,306 |
| Casual..... | 1,841 | 52 | 1,652 | 677 | 494 | 4,716 |
| Total..... | 6,032 | 100 | 8,404 | 1,376 | 1,161 | 17,073 |
| Vessels: | | | | | | |
| Steam..... | | | 13 | 8 | 3 | 24 |
| Net tonnage..... | | | 2,097 | 201 | 581 | 2,879 |
| Motor..... | 73 | | 365 | 65 | 67 | 570 |
| Net tonnage..... | 856 | | 13,995 | 695 | 1,056 | 16,602 |
| Sail..... | 1 | | | | | 1 |
| Net tonnage..... | 47 | | | | | 47 |
| Total vessels..... | 74 | | 378 | 73 | 70 | 595 |
| Total net tonnage..... | 903 | | 16,092 | 896 | 1,637 | 19,528 |
| Boats: | | | | | | |
| Motor..... | 2,342 | 34 | 1,478 | 433 | 294 | 4,581 |
| Other..... | 1,396 | 17 | 1,624 | 506 | 276 | 3,819 |
| A accessory boats..... | 172 | | 942 | 74 | 39 | 1,227 |
| Apparatus: | | | | | | |
| Purse seines: | | | | | | |
| Mackerel..... | 25 | | 97 | 2 | | 124 |
| Length, yards..... | 6,648 | | 45,720 | 420 | | 52,788 |
| Other..... | 51 | | 2 | 1 | | 54 |
| Length, yards..... | 7,040 | | 260 | 400 | | 7,700 |
| Haul seines..... | 31 | 1 | 15 | 20 | 12 | 79 |
| Length, yards..... | 3,100 | 85 | 1,655 | 2,512 | 1,668 | 8,970 |
| Gill nets: | | | | | | |
| Anchor..... | 1,736 | 4 | 704 | | 2 | 2,446 |
| Square yards..... | 556,491 | 2,160 | 280,655 | | 1,680 | 840,896 |
| Drift..... | 709 | | 4,105 | 44 | 43 | 4,901 |
| Square yards..... | 230,396 | | 1,530,960 | 17,800 | 12,360 | 1,791,516 |
| Runaround..... | | | 1 | | | 1 |
| Square yards..... | | | 1,800 | | | 1,800 |
| Lines: | | | | | | |
| Hand..... | 5,060 | 116 | 505 | 248 | 405 | 6,333 |
| Hooks..... | 5,259 | 116 | 671 | 319 | 443 | 6,808 |
| Trawl..... | 24,134 | 260 | 47,836 | 51 | 800 | 73,081 |
| Hooks..... | 1,155,574 | 13,000 | 2,380,062 | 24,709 | 35,680 | 3,009,016 |
| Troll..... | | | 8 | 8 | | 16 |
| Hooks..... | | | 8 | 48 | | 56 |
| Trot with hooks..... | 16 | | | 1 | | 17 |
| Hooks..... | 1,600 | | | 200 | | 1,800 |
| Pound nets..... | 3 | | 86 | 64 | 9 | 162 |
| Floating traps..... | 21 | | 17 | 40 | | 78 |
| Weirs..... | 213 | | 4 | | | 217 |
| Fyke nets..... | 56 | | 24 | 37 | | 121 |
| Dip nets..... | 124 | | 130 | | 119 | 373 |
| Bag nets..... | 112 | 30 | | | | 142 |
| Push nets..... | | | 50 | | | 50 |
| Pocket nets..... | 2 | | | | | 2 |
| Otter trawls..... | 50 | | 366 | 61 | 89 | 566 |
| Yards at mouth..... | 1,235 | | 21,084 | 1,635 | 2,373 | 26,327 |
| Box traps..... | 13 | | 3 | | | 16 |
| Pots: | | | | | | |
| Crab..... | 2,296 | | 1,095 | | | 3,391 |
| Eel..... | 222 | | 1,565 | 1,698 | 1,289 | 4,774 |
| Lobster..... | 180,439 | 2,990 | 65,147 | 53,672 | 17,212 | 319,460 |
| Periwinkle and cockle..... | | | 935 | 1,585 | | 2,520 |
| Harpoons..... | 43 | | 126 | 36 | 16 | 221 |
| Spears..... | 26 | | 185 | 13 | 42 | 266 |
| Dredges: | | | | | | |
| Oyster..... | | | 34 | 30 | 43 | 107 |
| Yards at mouth..... | | | 44 | 45 | 65 | 154 |
| Scallop..... | 155 | | 2,038 | 374 | 2 | 2,569 |
| Yards at mouth..... | 231 | | 1,895 | 324 | 7 | 2,457 |
| Clam..... | | | 63 | 12 | | 75 |
| Yards at mouth..... | | | 31 | 9 | | 40 |
| Mussel..... | | | 1 | 1 | | 2 |
| Yards at mouth..... | | | 1 | 1 | | 2 |
| Tongs..... | | | 166 | 543 | 118 | 827 |
| Rakes..... | | | 654 | 75 | 71 | 800 |
| Forks..... | | | 945 | 8 | | 953 |
| Hoes..... | 1,605 | | 260 | 26 | 38 | 1,929 |

Fisheries of the New England States, 1933—Continued

CATCH: BY STATES

| Species | Maine | | New Hampshire | | Massachusetts | | Rhode Island | | Connecticut | | Total | |
|---------------------------------|------------|---------|---------------|-------|---------------|-----------|--------------|---------|-------------|---------|-------------|-----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | | | | | |
| Alewives..... | 1,703,078 | \$6,371 | | | 923,056 | \$8,399 | 176,000 | \$1,915 | 15,162 | \$152 | 2,817,296 | \$16,837 |
| Amberjack..... | | | | | 2,275 | 68 | | | | | 2,275 | 68 |
| Bluefish..... | 85 | 11 | | | 430,040 | 42,882 | 194,386 | 10,806 | 296,454 | 22,058 | 920,965 | 75,757 |
| Bonito..... | | | | | 5,222 | 233 | 46,322 | 1,923 | | | 51,544 | 2,156 |
| Butterfish..... | 99,475 | 5,059 | | | 951,585 | 45,905 | 486,522 | 16,204 | 15,941 | 764 | 1,553,523 | 67,932 |
| Cod..... | 9,336,894 | 145,484 | 28,602 | \$858 | 89,210,361 | 1,688,460 | 908,241 | 16,735 | 147,925 | 4,457 | 99,632,023 | 1,855,994 |
| Crevaille..... | | | | | | | | 81 | | | 2,200 | 81 |
| Croaker..... | | | | | 2,491,145 | 35,716 | | | 5,738 | 136 | 2,496,883 | 35,852 |
| Cunners..... | 175 | 5 | | | 152 | 2 | 41,440 | 451 | | | 41,767 | 458 |
| Cusk..... | 1,583,366 | 18,342 | 17,161 | 343 | 4,509,004 | 53,766 | | | | | 6,109,531 | 72,451 |
| Drum: | | | | | | | | | | | | |
| Black..... | | | | | 32 | 1 | | | 100 | 1 | 132 | 2 |
| Red..... | | | | | 1,755 | 32 | | | | | 1,755 | 32 |
| Eels: | | | | | | | | | | | | |
| Common..... | 102,025 | 8,028 | | | 222,279 | 10,727 | 145,611 | 12,113 | 47,394 | 3,964 | 517,309 | 34,832 |
| Conger..... | | | | | 38,617 | 733 | 958 | 29 | 1,400 | 120 | 40,975 | 882 |
| Flounders..... | 1,178,669 | 27,517 | | | 27,187,427 | 878,601 | 3,253,077 | 90,930 | 6,176,038 | 175,968 | 37,795,211 | 1,173,016 |
| Frigate mackerel..... | | | | | 68,799 | 2,752 | 56,654 | 1,609 | | | 125,453 | 4,361 |
| Grayfish..... | 316 | 7 | | | 12,512 | 132 | 600 | 12 | | | 13,428 | 151 |
| Haddock..... | 9,306,555 | 215,744 | 60,063 | 2,703 | 150,680,476 | 3,425,852 | 23,968 | 703 | 35,450 | 1,018 | 160,106,512 | 3,646,020 |
| Hake..... | 7,084,013 | 66,270 | 188,749 | 2,531 | 8,056,678 | 133,691 | 9,171 | 183 | 1,081 | 20 | 15,319,692 | 202,695 |
| Halibut..... | 50,011 | 4,776 | | | 2,401,292 | 230,469 | | | 6,063 | 310 | 2,457,366 | 235,555 |
| Herring, sea..... | 43,774,988 | 160,816 | 2,000 | 20 | 3,572,960 | 38,784 | 737,006 | 11,467 | | | 48,086,954 | 211,087 |
| Herring smelt..... | | | | | 11,648 | 313 | | | | | 11,648 | 313 |
| Hickory shad..... | | | | | | | 2,308 | 23 | | | 2,308 | 23 |
| King whiting or "kingfish"..... | | | | | 38,296 | 1,035 | 1,440 | 53 | 451 | 12 | 40,187 | 1,100 |
| Launce..... | | | | | 21,000 | 420 | | | | | 21,000 | 420 |
| Mackerel..... | 4,490,107 | 57,740 | 1,050 | 53 | 35,611,363 | 798,108 | 719,289 | 22,137 | 870 | 27 | 40,831,679 | 878,065 |
| Menhaden..... | | | | | 334 | 4 | 1,003,510 | 2,535 | 25,250 | 252 | 1,029,094 | 2,791 |
| Mullet..... | | | | | 10 | 73 | 8,811 | 321 | | | 8,821 | 322 |
| Pigfish..... | | | | | 3,832 | 7 | | | | | 3,832 | 73 |
| Pollock..... | 3,006,065 | 23,414 | 11,441 | 114 | 11,991,798 | 140,076 | 17,242 | 451 | | | 15,026,546 | 164,035 |
| Rosefish..... | 2,063 | 14 | | | 262,255 | 2,827 | | | | | 264,318 | 2,841 |
| Salmon..... | 24,644 | 6,269 | | | 45 | 9 | | | | | 24,689 | 6,278 |
| Scup or porgy..... | | | | | 2,133,410 | 50,068 | 2,019,431 | 37,433 | 41,987 | 1,659 | 4,194,828 | 89,160 |
| Sea bass..... | | | | | 3,923,089 | 117,321 | 48,265 | 2,543 | 27,615 | 1,843 | 3,998,969 | 121,707 |
| Sea robin..... | | | | | 400 | 8 | 73,785 | 697 | 3,250 | 33 | 77,435 | 738 |
| Shad..... | 178,901 | 1,834 | | | 62,751 | 1,966 | 10,760 | 1,138 | 133,280 | 10,969 | 385,692 | 15,927 |

| | | | | | | | | | | | | |
|----------------------------------|-------------------|----------------|----------------|--------------|--------------------|------------------|-------------------|----------------|------------------|----------------|--------------------|------------------|
| Sharks..... | 29,923 | 230 | | | 34,148 | 287 | 1,947 | 28 | | | 66,018 | 545 |
| Sheepshead..... | | | | | 9 | 1 | | | | | 9 | 1 |
| Skates..... | 7,981 | 115 | | | 37,256 | 479 | 176,410 | 1,328 | 18,550 | 185 | 240,197 | 2,107 |
| Skipper or "billfish"..... | | | | | 4,110 | 41 | 730 | 14 | | | 4,840 | 55 |
| Smelt..... | 529,990 | 56,827 | 14,267 | 2,140 | | | 4,250 | 425 | 2,240 | 224 | 550,747 | 59,616 |
| Spot..... | | | | | 32,743 | 376 | 45 | 1 | 50 | 1 | 32,838 | 378 |
| Squeteagues or "sea trout": | | | | | | | | | | | | |
| Gray..... | | | | | 286,547 | 8,543 | 63,310 | 4,012 | 19,510 | 1,736 | 369,367 | 14,291 |
| Spotted..... | | | | | 2,048 | 204 | | | | | 2,048 | 204 |
| Stripped bass..... | | | | | 19,957 | 3,932 | 39,232 | 2,418 | 2,250 | 225 | 61,439 | 6,575 |
| Sturgeon..... | 2,308 | 224 | | | | | | | 44 | 5 | 7,931 | 905 |
| Suckers..... | 50,518 | 1,502 | | | | | | | 1,000 | 60 | 51,518 | 1,562 |
| Swordfish..... | 323,939 | 39,381 | | | 2,643,512 | 324,817 | 259,710 | 23,799 | 154,235 | 16,650 | 3,381,396 | 404,647 |
| Tantog..... | | | | | 173,543 | 6,766 | 267,123 | 7,402 | 43,098 | 1,895 | 483,764 | 16,063 |
| Thimble-eyed mackerel..... | | | | | | | 77,707 | 1,533 | | | 77,707 | 1,533 |
| Tilefish..... | | | | | | | | | 207,000 | 10,330 | 207,000 | 10,330 |
| Tomcod..... | 1,964 | 51 | | | | | 200 | 1 | | | 2,164 | 52 |
| Tuna or "horse mackerel"..... | 100,532 | 2,069 | | | 256,372 | 14,016 | 44,577 | 1,484 | | | 401,481 | 17,569 |
| White perch..... | | | | | 48,786 | 7,109 | 861 | 42 | | | 49,647 | 7,151 |
| Whiting..... | | | | | 8,677,742 | 88,648 | 724,506 | 7,626 | 16,775 | 547 | 9,419,023 | 96,821 |
| Wolfish..... | 93,647 | 482 | 3,142 | 79 | 2,096,569 | 31,334 | | | 4,100 | 41 | 2,197,458 | 31,936 |
| Yellow perch..... | 534 | 80 | | | | | | | | | 534 | 80 |
| Total..... | 83,071,766 | 848,662 | 306,475 | 8,841 | 359,144,370 | 8,196,646 | 11,648,054 | 282,642 | 7,450,301 | 255,662 | 461,620,966 | 9,592,453 |
| SHELLFISH, ETC. | | | | | | | | | | | | |
| Crabs: | | | | | | | | | | | | |
| Hard..... | 501,666 | 15,128 | | | 5,106,515 | 24,795 | 20,478 | 369 | 400 | 25 | 5,629,059 | 40,315 |
| King..... | | | | | | | 6,574 | 18 | | | 6,574 | 18 |
| Lobsters..... | 5,897,685 | 1,000,094 | 216,929 | 48,230 | 1,928,616 | 381,649 | 708,095 | 113,933 | 336,800 | 64,135 | 9,088,125 | 1,608,041 |
| Shrimp..... | | | | | 41,200 | 2,170 | | | | | 41,200 | 2,170 |
| Periwinkles and cockles..... | 15,516 | 825 | | | 65,700 | 6,350 | 109,442 | 5,201 | | | 190,658 | 12,376 |
| Clams: | | | | | | | | | | | | |
| Hard, public ¹ | 11,594 | 1,054 | | | 1,766,525 | 215,048 | 1,068,891 | 125,047 | 101,605 | 20,735 | 2,948,615 | 361,884 |
| Hard, private ¹ | | | | | 2,200 | 400 | 89,045 | 12,249 | | | 91,245 | 12,649 |
| Razor..... | | | | | 358,400 | 15,874 | | | | | 358,400 | 15,874 |
| Soft, public ² | 6,548,620 | 224,005 | | | 2,836,090 | 245,469 | 32,384 | 4,214 | 13,683 | 1,953 | 9,430,777 | 475,641 |
| Surf or skimmer..... | | | | | 59,240 | 3,575 | | | | | 59,240 | 3,575 |
| Mussels, sea..... | 110,596 | 2,534 | | | 30,000 | 3,000 | 650 | 38 | | | 141,246 | 5,572 |
| Oysters: ³ | | | | | | | | | | | | |
| Market, public, spring..... | | | | | | | 26,880 | 3,840 | 10,150 | 1,250 | 37,030 | 5,090 |
| Market, public, fall..... | | | | | | | 38,444 | 5,507 | 11,560 | 1,450 | 49,944 | 6,957 |
| Market, private, spring..... | | | | | 75,161 | 25,680 | 1,525,384 | 211,748 | 613,547 | 81,363 | 2,214,092 | 318,811 |
| Market, private, fall..... | | | | | 105,448 | 34,357 | 1,527,055 | 211,897 | 1,223,661 | 170,261 | 2,856,164 | 416,515 |

¹ Statistics on hard clams used in this table are based on yields of 11 pounds of meats per bushel in Maine, Massachusetts, and Rhode Island and 10 pounds in Connecticut.

² Statistics on soft clams used in this table are based on yield of 15 pounds of meats per bushel in Maine and Massachusetts; 16 pounds in Rhode Island, and 14 pounds in Connecticut.

³ Statistics on oysters used in this table are based on yields of 6.57 pounds of meats per bushel in Massachusetts, 6.96 pounds in Rhode Island and 6.81 pounds in Connecticut.

Fisheries of the New England States, 1933—Continued

CATCH: BY STATES—Continued

| Species | Maine | | New Hampshire | | Massachusetts | | Rhode Island | | Connecticut | | Total | |
|----------------------------------|--------------|-------------|---------------|-----------|---------------|-------------|--------------|-------------|-------------|-----------|---------------|--------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| SHELLFISH, ETC.—continued | | | | | | | | | | | | |
| Scallops: | | | | | | | | | | | | |
| Bay..... | | | | | 500, 115 | \$180, 975 | | | | | 620, 735 | \$203, 918 |
| Sea..... | 1, 073, 172 | \$145, 884 | | | 1, 029, 097 | 115, 344 | 63, 120 | \$12, 943 | 57, 500 | \$10, 000 | 2, 158, 195 | 267, 454 |
| Squid..... | 203 | 2 | | | 570, 536 | 7, 564 | 501, 851 | 11, 618 | 55, 926 | 6, 228 | 1, 075, 495 | 19, 234 |
| Irish moss..... | | | | | 11, 650 | 582 | | | 2, 905 | 50 | 11, 650 | 582 |
| Bloodworms..... | 642, 852 | 42, 724 | | | 17, 759 | 28, 000 | | | | | 660, 611 | 70, 724 |
| Sandworms..... | 618, 567 | 25, 980 | | | 21, 656 | 19, 500 | | | | | 640, 223 | 45, 480 |
| Sea urchins..... | 5, 845 | 217 | | | | | | | | | 5, 845 | 217 |
| Total..... | 15, 426, 316 | 1, 458, 445 | 216, 929 | \$48, 230 | 14, 525, 908 | 1, 310, 332 | 5, 718, 293 | 718, 622 | 2, 427, 727 | 357, 468 | 38, 315, 173 | 3, 893, 097 |
| Grand total..... | 98, 498, 082 | 2, 307, 107 | 523, 404 | 57, 071 | 373, 670, 278 | 9, 506, 978 | 17, 366, 347 | 1, 001, 264 | 9, 878, 028 | 613, 130 | 499, 936, 139 | 13, 485, 550 |

NOTE.—Of the total catch in Massachusetts, 10,619,503 pounds of fishery products, valued at \$281,810, were taken in the southern winter trawl fishery off southern New Jersey, Maryland, Virginia, and North Carolina. Of the total catch in Connecticut, 111,837 pounds of fishery products, valued at \$4,641, were taken in the same fishery. These products consisted principally of croaker, flounders, scup, and sea bass.

Fisheries of the New England States, 1933—Continued

PRODUCTION OF CERTAIN SHELLFISH IN NUMBER AND BUSHELS

| Product | Maine | | Massachusetts | | Rhode Island | | Connecticut | | Total | |
|-----------------------------|-----------|----------|---------------|----------|--------------|---------|-------------|---------|------------|----------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| Crabs: | | | | | | | | | | |
| Hard, number | 1,504,999 | \$15,126 | 15,319,546 | \$24,795 | 61,434 | \$369 | 1,200 | \$25 | 16,887,177 | \$40,315 |
| King, do | | | | | 1,644 | 18 | | | 1,644 | 18 |
| Clams: | | | | | | | | | | |
| Hard, public bushels | 1,054 | 1,054 | 160,593 | 215,048 | 97,172 | 125,047 | 10,160 | 20,735 | 268,979 | 361,884 |
| Hard, private do | | | 200 | 400 | 8,095 | 12,249 | | | 8,295 | 12,649 |
| Razor, do | | | 11,200 | 15,874 | | | | | 11,200 | 15,874 |
| Soft, public do | | | | | | | | | | |
| Surf or skimmer do | 436,575 | 224,005 | 189,073 | 245,469 | 2,024 | 4,214 | 977 | 1,953 | 628,649 | 475,641 |
| Mussels, sea, do | | | 3,485 | 3,575 | | | | | 3,485 | 3,575 |
| Oysters: | 9,216 | 2,534 | 3,000 | 3,000 | 50 | 38 | | | 12,266 | 5,572 |
| Market, public, spring, do | | | | | 3,862 | 3,840 | 1,490 | 1,250 | 5,352 | 5,090 |
| Market, public, fall, do | | | | | 5,524 | 5,507 | 1,696 | 1,450 | 7,220 | 6,957 |
| Market, private, spring, do | | | 11,440 | 25,680 | 219,164 | 211,748 | 90,095 | 81,383 | 320,699 | 318,811 |
| Market, private, fall, do | | | 16,050 | 34,357 | 219,404 | 211,897 | 179,686 | 170,261 | 415,140 | 416,515 |
| Periwinkles and cockles, do | 862 | 825 | 3,650 | 6,350 | 6,080 | 5,201 | | | 10,592 | 12,376 |
| Scallops: | | | | | | | | | | |
| Bay, do | | | 83,352 | 180,975 | 10,520 | 12,943 | 10,000 | 10,000 | 103,872 | 203,918 |
| Sea, do | 178,862 | 145,884 | 171,816 | 115,344 | | | 9,321 | 6,226 | 359,690 | 267,454 |

SEED OYSTER FISHERY

| Item | Rhode Island | | Connecticut | | Total | |
|------------------------|--------------|-------|-------------|----------|---------|----------|
| | Number | Value | Number | Value | Number | Value |
| OPERATING UNITS | | | | | | |
| Fishermen: | | | | | | |
| On vessels | | | 93 | | 93 | |
| On boats and shore: | | | | | | |
| Regular | 5 | | | | 5 | |
| Casual | 6 | | 154 | | 160 | |
| Total | 11 | | 247 | | 258 | |
| Vessels: | | | | | | |
| Steam | | | 3 | | 3 | |
| Net tonnage | | | 245 | | 245 | |
| Motor | | | 18 | | 18 | |
| Net tonnage | | | 271 | | 271 | |
| Sail | | | 2 | | 2 | |
| Net tonnage | | | 17 | | 17 | |
| Total vessels | | | 23 | | 23 | |
| Total net tonnage | | | 533 | | 533 | |
| Boats: | | | | | | |
| Motor | | | 7 | | 7 | |
| Other | 2 | | 88 | | 90 | |
| Apparatus: | | | | | | |
| Dredges | | | 101 | | 101 | |
| Yards at mouth | | | 97 | | 97 | |
| Tongs | 2 | | 94 | | 96 | |
| Rakes | | | 46 | | 46 | |
| CATCH | | | | | | |
| Oysters: | | | | | | |
| Seed, public, spring | 995 | \$499 | 30,463 | \$12,185 | 31,458 | \$12,684 |
| Seed, public, fall | | | 43,163 | 17,265 | 43,163 | 17,265 |
| Seed, private, spring | | | 207,185 | 69,617 | 207,185 | 69,617 |
| Total | 995 | 499 | 280,811 | 99,067 | 281,806 | 99,566 |

NOTE.—Of the total number of persons fishing for seed oysters, 135 are duplicated among those fishing for market oysters or other species. Similarly the following craft and gear are duplicated; 73 boats other than motor, 82 tongs, and 46 rakes.

Industries related to the fisheries of the New England States, 1933

OPERATING UNITS, SALARIES, AND WAGES

| Item | Maine and New Hampshire | Massachusetts | Rhode Island | Connecticut | Total |
|--------------------------------------|-------------------------|--------------------|------------------|------------------|--------------------|
| Transporting: | | | | | |
| Persons engaged: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| On vessels..... | 149 | 43 | 8 | 56 | 256 |
| On boats..... | 16 | | 38 | | 54 |
| Total..... | 165 | 43 | 46 | 56 | 310 |
| Vessels: | | | | | |
| Steam: | | | | 1 | 1 |
| Net tonnage..... | | | | 67 | 67 |
| Motor: | 74 | 12 | 4 | 14 | 104 |
| Net tonnage..... | 813 | 338 | 39 | 425 | 1,615 |
| Total vessels..... | 74 | 12 | 4 | 15 | 105 |
| Total net tonnage..... | 813 | 338 | 39 | 492 | 1,682 |
| Boats..... | 11 | | 20 | | 31 |
| Wholesale and manufacturing: | | | | | |
| Establishments..... | 131 | 165 | 35 | 81 | 392 |
| Persons engaged: | | | | | |
| Proprietors..... | 112 | 89 | 29 | 36 | 266 |
| Salaried employees..... | 142 | 587 | 35 | 35 | 799 |
| Wage earners: | | | | | |
| Average for season..... | 4,383 | 3,012 | 301 | 416 | 8,112 |
| Average for year..... | 1,298 | 2,338 | 207 | 329 | 4,170 |
| Paid to salaried employees..... | \$253,670 | \$1,291,380 | \$83,949 | \$85,198 | \$1,714,197 |
| Paid to wage earners..... | \$765,974 | \$2,392,013 | \$159,894 | \$387,994 | \$3,695,875 |
| Total salaries and wages..... | \$1,019,644 | \$3,673,393 | \$243,843 | \$473,192 | \$5,410,072 |
| Fishermen manufacturing..... | 612 | 1,480 | 109 | 22 | 2,223 |

PRODUCTS MANUFACTURED

| Item | Maine | | Massachusetts | | Rhode Island | | Connecticut | |
|---|----------------|----------------|---------------|-----------|--------------|-------|-------------|-------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| BY MANUFACTURING ESTABLISHMENTS | | | | | | | | |
| Alewives: Salted, tight pack pounds..... | 587,238 | \$9,669 | (1) | (1) | | | | |
| Cod: | | | | | | | | |
| Fresh sticks.....do..... | 251,778 | 46,844 | | | | | | |
| Fresh fillets.....do..... | 109,857 | 12,903 | 4,827,997 | \$325,929 | | | | |
| Frozen fillets.....do..... | (1) | (1) | 3,969,607 | 344,363 | | | | |
| Salted: | | | | | | | | |
| Green ¹do..... | 734,166 | 28,885 | 593,764 | 29,648 | | | | |
| Dry.....do..... | 59,196 | 2,054 | (1) | (1) | | | | |
| Boneless, including absolutely boneless pounds..... | 131,175 | 16,333 | 6,312,775 | 1,189,945 | | | | |
| Smoked fillets.....do..... | 44,889 | 6,718 | 1,017,034 | 147,241 | | | | |
| Oil, cod.....gallons..... | 3,350 | 670 | 41,981 | 21,482 | | | | |
| Oil, cod-liver.....do..... | | | 19,552 | 14,708 | | | | |
| Cusk: | | | | | | | | |
| Fresh sticks.....pounds..... | 136,177 | 25,067 | | | | | | |
| Fresh fillets.....do..... | 9,612 | 958 | 311,954 | 39,186 | | | | |
| Salted: | | | | | | | | |
| Green ¹do..... | 31,150 | 673 | (1) | (1) | | | | |
| Boneless, including absolutely boneless pounds..... | (1) | (1) | 23,785 | 2,762 | | | | |
| Flounders: | | | | | | | | |
| Fresh fillets.....do..... | | | 867,450 | 148,917 | | | | |
| Frozen fillets.....do..... | (1) | (1) | 264,924 | 34,984 | | | | |
| Haddock: | | | | | | | | |
| Fresh sticks.....do..... | 69,168 | 12,559 | | | | | | |
| Fresh fillets.....do..... | 163,908 | 26,982 | 15,016,474 | 1,750,062 | | | | |
| Frozen fillets.....do..... | (1) | (1) | 18,383,269 | 1,467,461 | | | | |
| Salted: | | | | | | | | |
| Green ¹do..... | 31,400 | 803 | | | | | | |
| Boneless, including absolutely boneless pounds..... | | | 98,295 | 19,220 | | | | |
| Finnan haddie.....do..... | 131,400 | 18,960 | 931,754 | 103,793 | (1) | (1) | | |

¹ This item has been included under "Unclassified products."

² This item is usually an intermediate product and although included in the total, may also be shown in its final stage of processing in this or another State.

Industries related to the fisheries of the New England States, 1933—Continued

PRODUCTS MANUFACTURED—Continued

| Item | Maine | | Massachusetts | | Rhode Island | | Connecticut | |
|---|-----------|-----------|---------------|----------|--------------|---------|-------------|-----------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| BY MANUFACTURING ESTABLISHMENTS—continued | | | | | | | | |
| Hake: | | | | | | | | |
| Fresh sticks.....pounds.. | 200,630 | \$29,454 | | | | | | |
| Fresh fillets.....do..... | 64,285 | 7,305 | 1,013,618 | \$85,709 | | | | |
| Frozen fillets.....do..... | (1) | (1) | 27,850 | 2,161 | | | | |
| Salted: | | | | | | | | |
| Green ¹do..... | 334,123 | 10,069 | (1) | (1) | | | | |
| Dry.....do..... | (1) | (1) | 271,172 | 8,723 | | | | |
| Halibut, fresh fillets.....do..... | | | 6,659 | 1,502 | | | | |
| Herring, sea: | | | | | | | | |
| Salted, round.....do..... | 560,700 | 11,439 | (1) | (1) | | | | |
| Smoked: | | | | | | | | |
| Bloaters: | | | | | | | | |
| Soft.....do..... | 286,662 | 10,954 | 585,332 | 52,782 | | | | |
| Hard.....do..... | 259,830 | 6,952 | (1) | (1) | | | | |
| Boneless.....do..... | 1,953,365 | 187,020 | | | | | | |
| Lengthwise.....do..... | 114,915 | 9,456 | | | | | | |
| Medium-scaled.....do..... | | | | | | | | |
| Kipperd.....do..... | 168,370 | 14,247 | (1) | (1) | | | | |
| Canned "sardines".....do..... | (1) | (1) | 169,505 | 17,131 | | | | |
| standard cases.....do..... | 980,906 | 2,397,348 | | | | | | |
| Scrap, green.....tons..... | 193 | 507 | | | | | | |
| Meal.....do..... | 898 | 24,601 | | | | | | |
| Oil.....gallons..... | 69,391 | 7,963 | | | | | | |
| Mackerel: | | | | | | | | |
| Fresh fillets.....pounds..... | | | 21,065 | 3,343 | | | | |
| Salted: | | | | | | | | |
| Fillets.....do..... | (1) | (1) | 1,787,746 | 133,250 | | | | |
| Split.....do..... | 84,420 | 2,711 | 1,285,360 | 91,823 | | | | |
| Pollock: | | | | | | | | |
| Fresh fillets.....do..... | | | 403,389 | 32,107 | | | | |
| Frozen fillets.....do..... | | | 1,091,401 | 65,147 | | | | |
| Salted: | | | | | | | | |
| Green ¹do..... | 89,056 | 1,824 | (1) | (1) | | | | |
| Dry.....do..... | 42,387 | 2,004 | 941,442 | 37,115 | | | | |
| Wolfish, fresh fillets.....do..... | | | 85,439 | 6,282 | | | | |
| Crab meat, packaged, fresh-cooked.....pounds..... | 67,441 | 33,571 | 151,087 | 62,979 | (1) | (1) | | |
| Lobster meat, packaged, fresh-cooked.....pounds..... | | | 79,020 | 86,420 | | | | |
| Clams, hard, fresh-shucked.....gallons..... | | | | | 2,660 | \$4,815 | (1) | (1) |
| Clams, soft: | | | | | | | | |
| Fresh-shucked.....do..... | | | | | 12,148 | 12,063 | | |
| Canned: | | | | | | | | |
| Whole.....do..... | | | | | | | | |
| standard cases.....do..... | 76,666 | 248,190 | | | | | | |
| Chowder.....do..... | 36,567 | 94,452 | (1) | (1) | | | | |
| Juice and bouillon.....do..... | | | | | | | | |
| standard cases.....do..... | 19,441 | 44,157 | | | | | | |
| Marine-shell products, buttons.....gross..... | (1) | (1) | (1) | (1) | | | 1,017,225 | \$635,213 |
| Oysters, fresh-shucked.....gallons..... | | | | | 301,942 | 514,470 | 175,313 | 313,445 |
| Unclassified products: | | | | | | | | |
| Fillets, fresh and frozen.....pounds..... | 7,673 | 1,310 | 1,637,522 | 136,046 | (1) | (1) | | |
| Salted.....do..... | 31,363 | 2,639 | 1,601,189 | 103,216 | | | | |
| Smoked.....do..... | 251,757 | 25,587 | 764,708 | 151,769 | (1) | (1) | (1) | (1) |

¹ This item has been included under "Unclassified products."

² This item is usually an intermediate product and although included in the total, may also be shown its final stage of processing in this or another State.

³ Includes frozen fillets of cod, cusk, flounders, haddock, and hake.

⁴ Includes pan-dressed flounders, fresh steaks of cod, cusk, and pollock; frozen swordfish steaks; frozen whiting sticks; and frozen fillets of halibut, mackerel, rosefish, salmon, and wolfish.

⁵ This item is included with miscellaneous.

⁶ Includes dry-salted cusk, haddock, and hake; boneless cusk and hake; and salted sea herring and mackerel fillets.

⁷ Includes green-salted cusk, hake, and pollock; dry-salted cod, cusk, haddock, and tongues; boneless hake; salted alewives, tight-pack; and salted sea herring, round and split.

⁸ Includes finnan cod; smoked cusk, haddock, and hake fillets; kippered herring; and spiced sea herring.

⁹ Includes smoked alewives, butterfish, carp, flounders, halibut, lake trout, mackerel, salmon, shad, and whitefish; smoked cusk, haddock, and hake fillets; hard smoked sea herring, bloaters, and medium scaled; spiced sea herring; and smoked and spiced salmon.

Industries related to the fisheries of the New England States, 1933—Continued

PRODUCTS MANUFACTURED—Continued

| Item | Maine | | Massachusetts | | Rhode Island | | Connecticut | |
|---|-----------------------|------------|---------------|------------|--------------|------------|-------------|------------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| BY MANUFACTURING ESTABLISHMENTS—Continued | | | | | | | | |
| Unclassified products—Con. | | | | | | | | |
| Canned: | | | | | | | | |
| Fish cakes, balls, etc. standard cases..... | 5,469 ⁽¹⁰⁾ | \$19,736 | 59,605 | \$424,318 | ----- | ----- | ----- | ----- |
| Fish flakes.....do..... | ----- | ----- | 2,183 | 18,489 | ----- | ----- | ----- | ----- |
| Other.....do..... | 11 17,089 | 11 137,795 | 13 34,186 | 13 131,958 | ----- | ----- | ----- | ----- |
| Meal, ground fish.....tons..... | 430 | 16,080 | 8,653 | 386,017 | ----- | ----- | ----- | ----- |
| Miscellaneous..... | ----- | 13 196,892 | ----- | 14 838,958 | ----- | 14 233,771 | ----- | 14 147,213 |
| Total..... | ----- | 3,754,341 | ----- | 8,706,923 | ----- | 765,139 | ----- | 1,095,871 |
| BY FISHERMEN | | | | | | | | |
| Alewives: | | | | | | | | |
| Corned.....pounds..... | ----- | ----- | 80,000 | 4,800 | ----- | ----- | ----- | ----- |
| Salted, tight-pack.....do..... | 263,100 | 439 | 420,000 | 12,600 | ----- | ----- | ----- | ----- |
| Smoked.....do..... | 165,085 | 4,848 | 12,000 | 600 | ----- | ----- | ----- | ----- |
| Cod, salted: | | | | | | | | |
| Green ¹do..... | ----- | ----- | 403,778 | 11,087 | ----- | ----- | ----- | ----- |
| Dry.....do..... | 5,860 | 226 | ----- | ----- | ----- | ----- | ----- | ----- |
| Cusk, dry-salted.....do..... | 620 | 11 | 2,714 | 52 | ----- | ----- | ----- | ----- |
| Haddock, green-salted² | | | | | | | | |
| do.....do..... | ----- | ----- | 14,365 | 288 | ----- | ----- | ----- | ----- |
| Hake, green-salted ³do..... | ----- | ----- | 3,030 | 31 | ----- | ----- | ----- | ----- |
| Hake, dry-salted.....do..... | 4,080 | 77 | ----- | ----- | ----- | ----- | ----- | ----- |
| Halibut, green-salted.....do..... | ----- | ----- | 2,115 | 97 | ----- | ----- | ----- | ----- |
| Mackerel, salted, split.....do..... | 600 | 45 | 25,800 | 817 | ----- | ----- | ----- | ----- |
| Pollock, green-salted ⁴do..... | ----- | ----- | 90 | 1 | ----- | ----- | ----- | ----- |
| Crab meat, packaged, fresh-cooked.....pounds..... | | | | | | | | |
| Clams, razor, fresh-shucked.....gallons..... | 3,459 | 1,558 | 5,000 | 2,500 | 1,200 | 480 | ----- | ----- |
| Clams, soft: | | | | | | | | |
| Fresh-shucked.....do..... | 70,281 | 48,225 | 19,000 | 24,000 | 50 | 75 | 500 | 1,000 |
| Steamed.....pounds..... | 161,887 | 11,279 | ----- | ----- | ----- | ----- | ----- | ----- |
| Clams, surf or skimmer, fresh-shucked.....gallons..... | | | | | | | | |
| Oysters, fresh-shucked.....do..... | ----- | ----- | 5,922 | 4,050 | ----- | ----- | 2,840 | 4,680 |
| Scallops, bay, fresh-shucked.....gallons..... | | | | | | | | |
| Scallops, sea, fresh-shucked.....gallons..... | 27,151 | 56,242 | 114,341 | 115,031 | 7,181 | 29,247 | 900 | 3,600 |
| Total..... | ----- | 122,950 | ----- | 391,203 | ----- | 29,802 | ----- | 9,280 |
| Grand total..... | ----- | 3,877,291 | ----- | 9,098,126 | ----- | 794,941 | ----- | 1,105,151 |

¹ This item is usually an intermediate product and although included in the total, may also be shown its final stage of processing in this or another State.

² This item is included with other canned fishery products.

³ Includes canned finnan haddie, fish flakes, and mackerel.

⁴ Includes canned finnan haddie; haddock chowder; mackerel; hard and soft clam chowder; dog food; ground fish roe; and rat bait.

⁵ Includes dry-scrap from herring; clam meal; marine shell buttons; and pearl essence.

⁶ Includes dry-scrap from groundfish; glue; isinglass; and marine and fresh-water shell products.

⁷ Includes packaged fresh-cooked crab meat; fresh-shucked bay scallops; finnan haddie; canned hard clam chowder and oyster puree; oyster-shell products; and marine-shell novelties.

⁸ Includes fresh-shucked hard clams; smoked butterfish, carp, mackerel, piddiefish or spoonbill cat, salmon, and whitefish; and marine-shell novelties.

NOTE.—The total value of manufactured products for the New England States was as follows: By manufacturing establishments, \$14,322,274; and by fishermen, \$563,235. Some of the above products may have been manufactured from products imported from another State or country; therefore they cannot be correlated directly with the catch within the State. Of the total number of persons engaged in the preparation of fishermen's manufactured products, 2,119 have also been included as fishermen and 45 of the persons shown on transporting craft have also been included as fishermen. This should be considered when computing the total number of persons in the fishery industries exclusive of duplication.

MAINE

Fisheries of Maine, 1933

OPERATING UNITS: BY GEAR

| Item | Purse seines | | Haul seines | Gill nets | | Lines | | |
|-------------------------------|--------------|------------|-------------|------------|-----------|--------------|------------|-----------------|
| | Mack-erel | Other | | Anchor | Drift | Hand | Trawl | Trot with hooks |
| | Number | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | | |
| On vessels..... | 86 | 65 | | 52 | 8 | 12 | 187 | |
| On boats and shore: | | | | | | | | |
| Regular..... | 36 | 118 | 54 | 104 | 39 | 210 | 519 | |
| Casual..... | | | 8 | 70 | 4 | 829 | 32 | 3 |
| Total..... | 122 | 183 | 62 | 226 | 51 | 1,051 | 738 | 3 |
| Vessels: | | | | | | | | |
| Motor..... | 16 | 14 | | 11 | 2 | 6 | 18 | |
| Net tonnage..... | 145 | 155 | | 95 | 23 | 45 | 321 | |
| Sail..... | | 1 | | | | | | |
| Net tonnage..... | | 47 | | | | | | |
| Total vessels..... | 16 | 15 | | 11 | 2 | 6 | 18 | |
| Total net tonnage..... | 145 | 202 | | 95 | 23 | 45 | 321 | |
| Boats: | | | | | | | | |
| Motor..... | 9 | 36 | 30 | 77 | 34 | 176 | 411 | |
| Other..... | 9 | 36 | 28 | 45 | 2 | 7 | 12 | 3 |
| Accessory boats..... | 12 | 11 | | 2 | 3 | | 80 | |
| Apparatus: | | | | | | | | |
| Number..... | 25 | 51 | 31 | 1,736 | 709 | 5,089 | 34,134 | 16 |
| Length yards..... | 6,648 | 7,040 | 3,100 | 556,491 | 230,396 | | | |
| Square yards..... | | | | | | | | |
| Hooks, baits or snoods..... | | | | | | 5,259 | 1,155,574 | 1,600 |

| Item | Pound nets | Floating traps | Weirs | Fyke nets | Dip nets | Bag nets | Pocket nets | Otter trawls | Box traps |
|---------------------|------------|----------------|------------|-----------|------------|-----------|-------------|--------------|-----------|
| | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | | | |
| On vessels..... | | | | | | | | 50 | |
| On boats and shore: | | | | | | | | | |
| Regular..... | 3 | 19 | 172 | 2 | | 7 | | 73 | |
| Casual..... | | 2 | 96 | 12 | 124 | 73 | 1 | 2 | 10 |
| Total..... | 3 | 21 | 268 | 14 | 124 | 80 | 1 | 125 | 10 |
| Vessels: | | | | | | | | | |
| Motor..... | | | | | | | | 13 | |
| Net tonnage..... | | | | | | | | 130 | |
| Boats: | | | | | | | | | |
| Motor..... | 3 | 16 | 1 | | | | | 37 | |
| Other..... | | 2 | 185 | 12 | | 9 | | | 4 |
| Apparatus: | | | | | | | | | |
| Number..... | 3 | 21 | 213 | 66 | 124 | 112 | 2 | 50 | 13 |
| Yards at mouth..... | | | | | | | | 1,235 | |

U. S. BUREAU OF FISHERIES

Fisheries of Maine, 1933—Continued

OPERATING UNITS: BY GEAR—Continued

| Item | Pots | | | Har- poons | Spears | Dredges, scallop | Hoos | By hand | Total, exclu- sive of dupli- cation |
|------------------------|--------|--------|---------|---------------|--------|---------------------|--------|------------|---|
| | Crab | Eel | Lobster | | | | | | |
| Fishermen: | | | | | | | | | |
| On vessels..... | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| On boats and shore: | | | 6 | 78 | | 43 | | | 443 |
| Regular..... | 43 | 3 | 2,426 | 46 | | 230 | 999 | 28 | 3,748 |
| Casual..... | 14 | 13 | 36 | 6 | 26 | 2 | 606 | 22 | 1,841 |
| Total..... | 57 | 16 | 2,468 | 130 | 26 | 275 | 1,605 | 50 | 6,032 |
| Vessels: | | | | | | | | | |
| Motor..... | | | 4 | 16 | | 6 | | | 73 |
| Net tonnage..... | | | 23 | 314 | | 69 | | | 856 |
| Sail..... | | | | | | | | | 1 |
| Net tonnage..... | | | | | | | | | 47 |
| Total vessels..... | | | 4 | 16 | | 6 | | | 74 |
| Total net tonnage..... | | | 23 | 314 | | 69 | | | 903 |
| Boats: | | | | | | | | | |
| Motor..... | 22 | | 1,794 | 27 | | 140 | 29 | | 2,342 |
| Other..... | 34 | 15 | 537 | | 21 | | 715 | 9 | 1,366 |
| Accessory boats..... | | | | 64 | | | | | 172 |
| Apparatus: | | | | | | | | | |
| Number..... | 2,296 | 222 | 180,439 | 43 | 26 | 155 | 1,605 | | |
| Yards at mouth..... | | | | | | 231 | | | |

CATCH: BY GEAR

| Species | Purse seines | | | | Haul seines | | Gill nets | | | |
|----------------------------------|--------------|---------|------------|---------|-------------|---------|-----------|---------|---------|-------|
| | Mackerel | | Other | | | | Anchor | | Drift | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | 199,453 | \$1,910 | 103,100 | \$507 | | | 47,000 | \$400 | | |
| Bluefish..... | | | | | | | | | 85 | \$11 |
| Butterfish..... | 17,936 | 764 | 6,630 | 248 | | | 20 | 1 | 109 | 7 |
| Cod..... | | | | | | | 4,106,347 | 73,662 | | |
| Cusk..... | | | | | | | 54,442 | 597 | | |
| Flounders..... | | | | | | | 31,497 | 778 | | |
| Grayfish..... | | | | | | | | | 316 | 7 |
| Haddock..... | | | | | | | 2,176,218 | 33,990 | | |
| Hake..... | | | | | | | 878,384 | 8,469 | | |
| Halibut..... | | | | | | | 4,660 | 50 | | |
| Herring, sea..... | 176,900 | 1,251 | 24,061,530 | 94,869 | | | 17,375 | 45 | | |
| Mackerel..... | 2,105,293 | 27,965 | 945,567 | 8,472 | | | 32,882 | 1,372 | 345,655 | 5,835 |
| Pollock..... | 910 | 3 | 75,302 | 377 | | | 2,089,751 | 17,979 | | |
| Salmon..... | | | | | | | 5,215 | 1,150 | 49 | 10 |
| Shad..... | 110,272 | 867 | 28,790 | 212 | | | 2,537 | 200 | | |
| Sharks..... | | | | | | | 19,945 | 135 | 9,725 | 92 |
| Smelt..... | 19 | 1 | | | 93,789 | \$7,431 | 66,496 | 8,910 | | |
| Sturgeon..... | | | | | | | 1,559 | 112 | 749 | 112 |
| Tomcod..... | | | | | 611 | 10 | | | | |
| Tuna or "horse mackerel"..... | | | | | | | 92 | 2 | | |
| Wolffish..... | | | | | | | 3,525 | 19 | | |
| Lobsters..... | | | | | | | 128 | 25 | | |
| Total..... | 2,610,783 | 32,761 | 25,220,919 | 104,685 | 94,400 | 7,441 | 9,538,073 | 147,896 | 356,688 | 6,074 |

Fisheries of Maine, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Lines | | | | | |
|-------------------|-------------|-----------|--------------|-----------|-----------------|-------|
| | Hand | | Trawl | | Trot with hooks | |
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Cod..... | 1, 108, 156 | \$11, 186 | 3, 356, 442 | \$50, 546 | ----- | ----- |
| Cunners..... | 175 | 5 | ----- | ----- | ----- | ----- |
| Cusk..... | 8, 800 | 69 | 1, 431, 293 | 16, 713 | 11, 346 | \$809 |
| Eels, common..... | ----- | ----- | ----- | ----- | ----- | ----- |
| Flounders..... | ----- | ----- | 16, 630 | 229 | ----- | ----- |
| Haddock..... | 624, 077 | 14, 715 | 4, 777, 398 | 126, 526 | ----- | ----- |
| Hake..... | 655, 601 | 4, 218 | 4, 729, 610 | 47, 045 | ----- | ----- |
| Halibut..... | 7, 143 | 880 | 30, 406 | 3, 198 | ----- | ----- |
| Mackerel..... | 5, 198 | 156 | 45 | 6 | ----- | ----- |
| Pollock..... | 390, 844 | 2, 345 | 359, 756 | 2, 218 | ----- | ----- |
| Rosefish..... | ----- | ----- | 233 | 2 | ----- | ----- |
| Sharks..... | ----- | ----- | 253 | 3 | ----- | ----- |
| Skates..... | ----- | ----- | 5, 013 | 85 | ----- | ----- |
| Smelt..... | 222, 014 | 24, 787 | 4, 394 | 27 | ----- | ----- |
| Wolfish..... | ----- | ----- | 80, 682 | 423 | ----- | ----- |
| Total..... | 3, 022, 008 | 58, 361 | 14, 792, 155 | 247, 021 | 11, 346 | 809 |

| Species | Pound nets | | Floating traps | | Weirs | |
|-------------------|------------|--------|----------------|----------|--------------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | ----- | ----- | ----- | ----- | 211, 450 | \$468 |
| Butterfish..... | 8, 603 | \$405 | 65, 872 | \$3, 615 | ----- | ----- |
| Herring, sea..... | 20, 435 | 186 | 184, 913 | 1, 490 | 19, 313, 835 | 62, 975 |
| Mackerel..... | 76, 876 | 1, 294 | 461, 025 | 7, 293 | 526, 566 | 5, 347 |
| Pollock..... | 1, 314 | 9 | 20, 873 | 140 | 1, 021 | 10 |
| Salmon..... | 112 | 22 | 4, 245 | 935 | 13, 138 | 3, 756 |
| Shad..... | 491 | 6 | 2, 339 | 17 | 34, 460 | 531 |
| Smelt..... | 1, 767 | 133 | 8, 218 | 610 | 7, 223 | 686 |
| Squid..... | 203 | 2 | ----- | ----- | ----- | ----- |
| Total..... | 109, 801 | 2, 057 | 747, 485 | 14, 100 | 20, 107, 693 | 73, 773 |

| Species | Fyke nets | | Dip nets | | Bag nets | | Pocket nets | | Otter trawls | |
|-------------------|-----------|--------|-------------|----------|----------|-----------|-------------|-------|--------------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | ----- | ----- | 1, 142, 075 | \$3, 086 | ----- | ----- | ----- | ----- | ----- | ----- |
| Butterfish..... | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | 305 | \$19 |
| Cod..... | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | 765, 949 | 10, 090 |
| Cusk..... | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | 88, 831 | 963 |
| Eels, common..... | 2, 086 | \$125 | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- |
| Flounders..... | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | 1, 128, 284 | 26, 487 |
| Haddock..... | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | 1, 728, 863 | 40, 513 |
| Hake..... | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | 820, 418 | 6, 538 |
| Halibut..... | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | 7, 802 | 648 |
| Pollock..... | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | 66, 294 | 333 |
| Rosefish..... | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | 1, 830 | 12 |
| Salmon..... | ----- | ----- | 1, 885 | 396 | ----- | ----- | ----- | ----- | ----- | ----- |
| Shad..... | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | 12 | 1 |
| Smelt..... | ----- | ----- | 15, 002 | 1, 905 | 109, 532 | \$12, 199 | 1, 536 | \$138 | ----- | ----- |
| Suckers..... | 50, 518 | 1, 502 | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- |
| Tomcod..... | 953 | 29 | ----- | ----- | 400 | 12 | ----- | ----- | ----- | ----- |
| Wolfish..... | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | 9, 440 | 40 |
| Yellow perch..... | 534 | 80 | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- |
| Total..... | 54, 091 | 1, 736 | 1, 158, 962 | 5, 387 | 109, 932 | 12, 211 | 1, 536 | 138 | 4, 618, 027 | 85, 644 |

U. S. BUREAU OF FISHERIES

Fisheries of Maine, 1933—Continued

CATCH: By GEAR—Continued

| Species | Box traps | | Pots | | | | | | Harpoons | |
|--------------------------|-----------|---------|---------|----------|--------|---------|-----------|-----------|----------|----------|
| | | | Crab | | Eel | | Lobster | | | |
| | | | Pounds | Value | Pounds | Value | Pounds | Value | | |
| Eels, common | 28,910 | \$2,512 | | | 36,284 | \$2,661 | | | | |
| Swordfish | | | | | | | | | 323,939 | \$39,381 |
| Tuna or "horse mackerel" | | | | | | | | | 100,440 | 2,067 |
| Crabs, hard | | | 449,134 | \$13,700 | | | | 52,632 | \$1,426 | |
| Lobsters | | | 256 | 64 | | | 5,897,301 | 1,000,005 | | |
| Total | 28,910 | 2,512 | 449,300 | 13,764 | 36,284 | 2,661 | 5,949,833 | 1,001,431 | 424,379 | 41,448 |

| Species | Spears | | Dredges, scallops | | Hoes | | By hand | |
|---------------|--------|---------|-------------------|-----------|-----------|---------|---------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Eels, common | 23,399 | \$1,921 | | | | | | |
| Flounders | 2,258 | 23 | | | | | | |
| Skates | 2,968 | 30 | | | | | | |
| Clams: | | | | | | | | |
| Hard, public | | | | | 11,594 | \$1,054 | | |
| Soft, public | | | | | 6,548,620 | 224,005 | | |
| Mussels, sea | | | | | | | 110,596 | \$2,534 |
| Periwinkles | | | | | | | 15,516 | 825 |
| Scallops, sea | | | 1,073,172 | \$145,884 | | | | |
| Bloodworms | | | | | 642,852 | 42,724 | | |
| Sandworms | | | | | 618,567 | 25,980 | | |
| Sea urchins | | | | | | | 5,845 | 217 |
| Total | 28,625 | 1,974 | 1,073,172 | 145,884 | 7,821,633 | 293,763 | 131,957 | 3,576 |

OPERATING UNITS: BY COUNTIES

| Item | Cumber-land | Hancock | Kennebec | Knox | Lincoln | Penobscot | Sagadahoc | Waldo | Washington | York |
|---------------------|-------------|---------|----------|--------|---------|-----------|-----------|--------|------------|---------|
| | Number | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | | | | |
| On vessels | 259 | 18 | | 78 | 56 | | | | 25 | 7 |
| On boats and shore: | | | | | | | | | | |
| Regular | 887 | 654 | | 620 | 433 | | 187 | 22 | 752 | 193 |
| Casual | 33 | 610 | 168 | 176 | 197 | 15 | 68 | 59 | 453 | 62 |
| Total | 1,179 | 1,282 | 168 | 874 | 686 | 15 | 255 | 81 | 1,230 | 262 |
| Vessels: | | | | | | | | | | |
| Motor | 34 | 6 | | 16 | 10 | | | | 5 | 2 |
| Net tonnage | 499 | 49 | | 144 | 86 | | | | 63 | 15 |
| Sail | | | | | | | | | 1 | |
| Net tonnage | | | | | | | | | 47 | |
| Total vessels | 34 | 6 | | 16 | 10 | | | | 6 | 2 |
| Total net tonnage | 499 | 49 | | 144 | 86 | | | | 110 | 15 |
| Boats: | | | | | | | | | | |
| Motor | 511 | 485 | | 439 | 226 | | 84 | 8 | 424 | 165 |
| Other | 217 | 294 | 7 | 129 | 163 | 12 | 67 | 37 | 420 | 50 |
| Accessory boats | 139 | 1 | | 12 | 14 | | | | 5 | 1 |
| Apparatus: | | | | | | | | | | |
| Purse seines: | | | | | | | | | | |
| Mackerel | 12 | | | 3 | 8 | | | | | 2 |
| Length, yards | 3,260 | | | 528 | 2,600 | | | | | 260 |
| Other | 10 | 2 | | 6 | 23 | | 1 | | 9 | |
| Length, yards | 1,760 | 360 | | 970 | 2,920 | | 200 | | 830 | |
| Haul seines | 30 | | | 1 | | | | | | |
| Length, yards | 3,000 | | | 100 | | | | | | |
| Gill nets: | | | | | | | | | | |
| Anchor | 1,069 | 64 | | 8 | 140 | 42 | 3 | 20 | 318 | 72 |
| Square yards | 421,879 | 13,320 | | 3,122 | 52,254 | 3,160 | 270 | 2,242 | 27,948 | 27,296 |
| Drift | 202 | | | | 5 | | 60 | | | 442 |
| Square yards | 96,123 | | | | 233 | | 27,960 | | | 106,080 |

Fisheries of Maine, 1933—Continued

OPERATING UNITS: BY COUNTIES—Continued

| Item | Cum-ber-land | Han-cock | Ken-nebec | Knor | Lin-coln | Pen-ob-scot | Saga-daboo | Waldo | Wash-ington | York |
|----------------------|--------------|----------|-----------|---------|----------|-------------|------------|--------|-------------|--------|
| Apparatus—Continued. | | | | | | | | | | |
| Lines: | Number | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| Hand..... | 68 | 1,724 | 820 | 993 | 1,089 | ----- | 275 | ----- | 114 | 6 |
| Hooks..... | 68 | 1,777 | 820 | 1,075 | 1,093 | ----- | 275 | ----- | 144 | 7 |
| Trawl..... | 12,380 | 2,352 | ----- | 2,310 | 1,000 | ----- | 1,780 | 72 | 2,980 | 1,280 |
| Hooks..... | 619,000 | 71,424 | ----- | 115,500 | 50,000 | ----- | 89,000 | 3,800 | 144,050 | 63,000 |
| Trot with hooks..... | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | 16 |
| Hooks..... | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | 1,600 |
| Pound nets..... | 3 | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- |
| Floating traps..... | 5 | ----- | ----- | ----- | 3 | ----- | 9 | 4 | ----- | ----- |
| Weirs..... | ----- | 84 | ----- | 9 | 3 | 13 | 1 | 27 | 76 | ----- |
| Fyke nets..... | ----- | 1 | 38 | ----- | 17 | ----- | ----- | ----- | ----- | ----- |
| Dip nets..... | ----- | 4 | ----- | 32 | 6 | ----- | ----- | 10 | 60 | 12 |
| Bag nets..... | ----- | 9 | ----- | ----- | ----- | 2 | ----- | 41 | 60 | ----- |
| Pocket nets..... | ----- | ----- | 2 | ----- | ----- | ----- | ----- | ----- | ----- | ----- |
| Otter trawls..... | 26 | 17 | ----- | 3 | 1 | ----- | 2 | ----- | 1 | ----- |
| Yards at mouth..... | 630 | 430 | ----- | 75 | 30 | ----- | 45 | ----- | 25 | ----- |
| Box traps..... | ----- | 2 | ----- | 6 | ----- | ----- | ----- | ----- | 3 | 2 |
| Pots: | | | | | | | | | | |
| Crab..... | 2,271 | ----- | ----- | ----- | 25 | ----- | ----- | ----- | ----- | ----- |
| Eel..... | 21 | ----- | ----- | ----- | 65 | 16 | 18 | 83 | 4 | 15 |
| Lobster..... | 22,089 | 40,116 | ----- | 37,913 | 18,813 | ----- | 4,879 | 125 | 46,042 | 10,462 |
| Harpoons..... | 30 | ----- | ----- | ----- | 1 | ----- | 5 | ----- | ----- | 7 |
| Spears..... | ----- | 10 | ----- | 2 | 2 | ----- | 6 | ----- | 3 | 3 |
| Dredges: | | | | | | | | | | |
| Scallop..... | 54 | 53 | ----- | 27 | 7 | ----- | ----- | ----- | 14 | ----- |
| Yards at mouth..... | 72 | 78 | ----- | 55 | 9 | ----- | ----- | ----- | 17 | ----- |
| Hoes..... | 339 | 338 | ----- | 159 | 202 | ----- | 102 | 17 | 410 | 38 |

CATCH: BY COUNTIES

| Species | Cumberland | | Hancock | | Kennebec | |
|-------------------------------|------------|---------|-----------|---------|----------|-------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | 52,586 | \$661 | 192,630 | \$572 | ----- | ----- |
| Butterfish..... | 40,488 | 2,032 | ----- | ----- | ----- | ----- |
| Cod..... | 5,218,503 | 96,283 | 1,120,583 | 10,571 | ----- | ----- |
| Cunners..... | 175 | 5 | ----- | ----- | ----- | ----- |
| Cusk..... | 1,214,160 | 14,547 | 83,411 | 519 | ----- | ----- |
| Eels, common..... | 924 | 34 | 8,915 | 857 | 2,065 | \$125 |
| Flounders..... | 560,008 | 11,115 | 397,414 | 11,136 | ----- | ----- |
| Grayfish..... | 316 | 7 | ----- | ----- | ----- | ----- |
| Haddock..... | 4,770,028 | 101,984 | 1,326,816 | 27,835 | ----- | ----- |
| Hake..... | 3,960,122 | 43,870 | 862,784 | 4,562 | ----- | ----- |
| Halibut..... | 15,200 | 1,255 | 10,097 | 992 | ----- | ----- |
| Herring, sea..... | 8,170,795 | 37,583 | 2,934,984 | 11,572 | ----- | ----- |
| Mackerel..... | 772,889 | 13,446 | 105,924 | 1,271 | ----- | ----- |
| Pollock..... | 2,327,787 | 19,486 | 68,114 | 331 | ----- | ----- |
| Rosefish..... | 2,063 | 14 | ----- | ----- | ----- | ----- |
| Salmon..... | 623 | 124 | 7,069 | 2,081 | ----- | ----- |
| Shad..... | 5,122 | 137 | ----- | ----- | ----- | ----- |
| Sharks..... | 22,988 | 164 | ----- | ----- | ----- | ----- |
| Smelt..... | 107,854 | 7,957 | 63,786 | 8,558 | 79,584 | 7,165 |
| Sturgeon..... | 708 | 68 | ----- | ----- | ----- | ----- |
| Suckers..... | ----- | ----- | ----- | ----- | 31,851 | 955 |
| Swordfish..... | 323,939 | 30,381 | ----- | ----- | ----- | ----- |
| Tomcod..... | 611 | 10 | ----- | ----- | 953 | 29 |
| Tuna or "horse mackerel"..... | 65,642 | 1,282 | ----- | ----- | ----- | ----- |
| Wolfish..... | 70,842 | 361 | ----- | ----- | ----- | ----- |
| Yellow perch..... | ----- | ----- | ----- | ----- | 142 | 21 |
| Crabs, hard..... | 473,016 | 14,268 | ----- | ----- | ----- | ----- |
| Lobsters..... | 648,284 | 116,974 | 1,393,672 | 227,273 | ----- | ----- |
| Clams: | | | | | | |
| Hard, public..... | 9,394 | 854 | ----- | ----- | ----- | ----- |
| Soft, public..... | 1,487,060 | 73,240 | 939,070 | 23,488 | ----- | ----- |
| Mussels, sea..... | 110,596 | 2,534 | ----- | ----- | ----- | ----- |
| Scallops, sea..... | 50,166 | 9,062 | 232,752 | 53,627 | ----- | ----- |
| Squid..... | 203 | 2 | ----- | ----- | ----- | ----- |
| Bloodworms..... | 136,762 | 21,468 | ----- | ----- | ----- | ----- |
| Total..... | 30,625,924 | 630,208 | 9,747,951 | 385,245 | 114,616 | 8,298 |

U. S. BUREAU OF FISHERIES

Fisheries of Maine, 1933—Continued

CATCH: BY COUNTIES—Continued

| Species | Knox | | Lincoln | | Penobscot | |
|-------------------------------|-----------|---------|------------|---------|-----------|-------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | 436,240 | \$1,099 | 581,440 | \$2,099 | | |
| Butterfish..... | | | 32,846 | 1,420 | | |
| Cod..... | 695,858 | 7,607 | 989,325 | 9,504 | | |
| Cusk..... | 46,181 | 473 | 58,533 | 362 | | |
| Eels, common..... | 7,398 | 550 | 11,127 | 801 | 5,025 | \$126 |
| Flounders..... | 119,094 | 3,206 | 41,944 | 735 | | |
| Haddock..... | 1,358,851 | 36,037 | 416,807 | 7,703 | | |
| Hake..... | 791,237 | 5,134 | 484,408 | 2,746 | | |
| Halibut..... | 8,681 | 910 | 808 | 81 | | |
| Herring, sea..... | 1,982,305 | 8,694 | 9,220,728 | 35,435 | | |
| Mackerel..... | 278,311 | 3,616 | 2,369,765 | 23,961 | | |
| Pollock..... | 305,209 | 1,490 | 86,707 | 455 | | |
| Salmon..... | | | 2,162 | 438 | 1,257 | 368 |
| Shad..... | 90 | 2 | 137,780 | 1,155 | | |
| Sharks..... | | | 347 | 3 | | |
| Smelt..... | 28,787 | 3,596 | 21,350 | 2,449 | 7,641 | 923 |
| Sturgeon..... | | | 18,667 | 647 | | |
| Suckers..... | | | 11 | 10 | | |
| Tuna or "horse mackerel"..... | | | 524 | 4 | | |
| Wolfish..... | | | 632 | 59 | | |
| Yellow perch..... | | | 392 | 858 | | |
| Crabs, hard..... | | | 28,650 | 104,247 | | |
| Lobsters..... | 1,595,485 | 267,980 | 619,488 | 20,908 | | |
| Clams, soft, public..... | 721,465 | 20,356 | 631,580 | 4,500 | | |
| Scallops, sea..... | 767,934 | 77,927 | 15,900 | 21,256 | | |
| Bloodworms..... | | | 508,100 | 25,980 | | |
| Sandworms..... | | | 618,567 | 60 | | |
| Sea urchins..... | | | 3,600 | | | |
| Total..... | 9,143,186 | 438,677 | 16,953,344 | 267,787 | 13,923 | 1,417 |

| Species | Sagadahoc | | Waldo | | Washington | | York | |
|-------------------------------|-----------|--------|---------|--------|------------|---------|-----------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | 15,000 | \$300 | | | 326,615 | \$640 | 98,567 | \$1,000 |
| Bluefish..... | | | | | | | 85 | 11 |
| Butterfish..... | 25,641 | 1,572 | | | | | 500 | 35 |
| Cod..... | 409,808 | 6,377 | 38,246 | \$780 | 620,740 | 9,782 | 243,831 | 4,580 |
| Cusk..... | 87,258 | 1,181 | | | 22,892 | 199 | 70,931 | 1,061 |
| Eels, common..... | 6,991 | 629 | 11,600 | 1,020 | 12,053 | 1,104 | 36,006 | 2,782 |
| Flounders..... | 28,953 | 501 | | | 28,282 | 779 | 2,974 | 45 |
| Haddock..... | 437,704 | 10,670 | 4,210 | 211 | 618,140 | 20,326 | 373,999 | 10,978 |
| Hake..... | 414,698 | 3,998 | 539 | 11 | 216,777 | 2,132 | 347,458 | 3,817 |
| Halibut..... | 1,453 | 129 | | | 7,402 | 1,230 | 6,290 | 179 |
| Herring, sea..... | 468,020 | 2,879 | 17,850 | 128 | 20,980,246 | 64,525 | | |
| Mackerel..... | 289,448 | 5,192 | 72,400 | 1,034 | 288,772 | 2,193 | 321,598 | 7,027 |
| Pollock..... | 69,798 | 457 | | | 111,138 | 924 | 37,312 | 271 |
| Salmon..... | 1,073 | 214 | 4,560 | 1,305 | 7,870 | 1,739 | | |
| Shad..... | 1,449 | 9 | | | 34,460 | 531 | | |
| Sharks..... | 1,565 | 16 | | | | | 5,023 | 48 |
| Skates..... | | | | | 7,981 | 115 | | |
| Smelt..... | 53,476 | 6,180 | 49,704 | 7,455 | 116,372 | 12,400 | 1,436 | 144 |
| Sturgeon..... | 749 | 112 | | | | | 684 | 33 |
| Tomcod..... | | | | | 400 | 12 | | |
| Tuna or "horse mackerel"..... | 19,535 | 375 | | | | | 14,831 | 402 |
| Wolfish..... | 11,655 | 61 | | | 1,173 | 5 | 9,345 | 51 |
| Lobsters..... | 133,570 | 23,809 | 2,509 | 602 | 1,057,423 | 173,009 | 447,254 | 86,200 |
| Clams: | | | | | | | | |
| Hard, public..... | 2,200 | 200 | | | | | | |
| Soft, public..... | 754,425 | 29,625 | 192,240 | 12,440 | 1,687,265 | 37,551 | 85,615 | 6,397 |
| Periwinkles..... | | | | | 15,516 | 825 | | |
| Scallops, sea..... | | | | | 3,420 | 768 | | |
| Sea urchins..... | | | | | 2,245 | 157 | | |
| Total..... | 3,234,459 | 94,485 | 393,858 | 24,986 | 26,167,182 | 330,946 | 2,103,639 | 125,061 |

NEW HAMPSHIRE

Fisheries of New Hampshire, 1933¹

OPERATING UNITS: BY GEAR

| Item | Haul seines | Gill nets, anchor | Lines | | Pots, lobster | Bag nets | Total, exclusive of duplication |
|--------------------------------|-------------|-------------------|--------|--------|---------------|----------|---------------------------------|
| | | | Hand | Trawl | | | |
| Fishermen, on boats and shore: | Number | Number | Number | Number | Number | Number | Number |
| Regular..... | 2 | 2 | 2 | 8 | 44 | 15 | 48 |
| Casual..... | | | 29 | | 8 | | 52 |
| Total..... | 2 | 2 | 29 | 8 | 52 | 15 | 100 |
| Boats: | | | | | | | |
| Motor..... | | 2 | | 5 | 32 | | 34 |
| Other..... | 1 | | | | 16 | | 17 |
| Apparatus: | | | | | | | |
| Number..... | 1 | 4 | 116 | 260 | 2,990 | 30 | |
| Length, yards..... | 35 | | | | | | |
| Square yards..... | | 2,160 | | | | | |
| Hooks..... | | | 116 | 13,000 | | | |

CATCH: BY GEAR

| Species | Haul seines | | Gill nets, anchor | | Lines | | | | Pots, lobster | | Bag nets | |
|-------------------|-------------|-------|-------------------|-------|-------|---------|---------|-------|---------------|----------|----------|-------|
| | Lbs. | Value | Lbs. | Value | Hand | | Trawl | | Lbs. | Value | Lbs. | Value |
| | | | | | Lbs. | Value | Lbs. | Value | | | | |
| Cod..... | | | | | | | 28,602 | \$858 | | | | |
| Cusk..... | | | | | | | 17,161 | 343 | | | | |
| Haddock..... | | | | | | | 60,063 | 2,703 | | | | |
| Hake..... | | | | | | | 168,749 | 2,531 | | | | |
| Herring, sea..... | 2,000 | \$20 | | | | | | | | | | |
| Mackerel..... | | | 1,050 | \$63 | | | | | | | | |
| Pollock..... | | | | | | | 11,441 | 114 | | | | |
| Smelt..... | | | | | 9,667 | \$1,460 | | | | | 4,600 | \$690 |
| Wolfish..... | | | | | | | 3,142 | 79 | | | | |
| Lobsters..... | | | | | | | | | 216,929 | \$48,230 | | |
| Total..... | 2,000 | 20 | 1,050 | 63 | 9,667 | 1,460 | 289,158 | 6,628 | 216,929 | 48,230 | 4,600 | 690 |

¹ The commercial fisheries of New Hampshire are confined to Rockingham County.

MASSACHUSETTS

Fisheries of Massachusetts, 1933

OPERATING UNITS: BY GEAR

| Item | Purse seines | | Haul seines | Gill nets | | | Lines | |
|------------------------------|--------------|--------|-------------|-----------|-----------|------------|--------|-----------|
| | Mack-erel | Other | | Anchor | Drift | Run-around | Hand | Trawl |
| Fishermen: | Number | Number | Number | Number | Number | Number | Number | Number |
| On vessels..... | 925 | | | 130 | 633 | 4 | 111 | 1,122 |
| On boats and shore: | | | | | | | | |
| Regular..... | 62 | | 37 | 14 | 168 | | 228 | 690 |
| Casual..... | | 6 | 8 | | 8 | | 72 | 12 |
| Total..... | 987 | 6 | 45 | 144 | 709 | 4 | 411 | 1,724 |
| Vessels: | | | | | | | | |
| Motor..... | 83 | | | 15 | 68 | 1 | 11 | 69 |
| Net tonnage..... | 2,729 | | | 285 | 1,653 | 5 | 301 | 3,326 |
| Boats: | | | | | | | | |
| Motor..... | 14 | 2 | 4 | 7 | 76 | | 176 | 227 |
| Other..... | 14 | 4 | 20 | 7 | 58 | | 55 | 31 |
| Accessory boats..... | 66 | | | 7 | 26 | | | 612 |
| Apparatus: | | | | | | | | |
| Number..... | 97 | 2 | 15 | 704 | 4,105 | 1 | 505 | 47,836 |
| Length, yards..... | 45,720 | 260 | 1,655 | | | | | |
| Square yards..... | | | | 280,655 | 1,530,960 | 1,800 | | |
| Hooks, baits, or snoods..... | | | | | | | 671 | 2,380,062 |

Fisheries of Massachusetts, 1935—Continued

OPERATING UNITS: BY GEAR—Continued

| Item | Lines— Con. | Pound nets | Float- ing traps | Weirs | Fyke nets | Dip nets | Push nets | Otter trawls |
|------------------------------|----------------|---------------|------------------------|--------|--------------|-------------|--------------|-----------------|
| | Troll | | | | | | | |
| | Number | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | | |
| On vessels..... | | | | | | | | 2,067 |
| On boats and shore: | | | | | | | | |
| Regular..... | 5 | 129 | 40 | 6 | 11 | 120 | | 199 |
| Casual..... | | 2 | | | 7 | 42 | 100 | |
| Total..... | 5 | 131 | 40 | 6 | 18 | 162 | 100 | 2,266 |
| Vessels: | | | | | | | | |
| Steam..... | | | | | | | | 13 |
| Net tonnage..... | | | | | | | | 2,097 |
| Motor..... | | | | | | | | 245 |
| Net tonnage..... | | | | | | | | 9,680 |
| Total vessels..... | | | | | | | | 258 |
| Total net tonnage..... | | | | | | | | 11,777 |
| Boats: | | | | | | | | |
| Motor..... | 5 | 35 | 16 | 2 | | 24 | | 79 |
| Other..... | | 57 | 15 | 2 | 6 | 35 | 50 | |
| Accessory boats..... | | | 6 | | | | | |
| Apparatus: | | | | | | | | |
| Number..... | 8 | 86 | 17 | 4 | 24 | 120 | 50 | 366 |
| Yards at mouth..... | | | | | | | | 21,084 |
| Hooks, baits, or snoods..... | 8 | | | | | | | |

| Item | Box traps | Pots | | | | Har- poons | Spears |
|------------------------|--------------|--------|--------|--------------|---------------------------------|---------------|--------|
| | | Crab | Eel | Lob- ster | Peri- winkle or cockle | | |
| | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | |
| On vessels..... | | | | | | 632 | |
| On boats and shore: | | | | | | | |
| Regular..... | 3 | 25 | 36 | 717 | 22 | 88 | 68 |
| Casual..... | | 4 | 15 | 294 | | | 117 |
| Total..... | 3 | 29 | 51 | 1,011 | 22 | 720 | 185 |
| Vessels: | | | | | | | |
| Steam..... | | | | | | 1 | |
| Net tonnage..... | | | | | | 180 | |
| Motor..... | | | | | | 88 | |
| Net tonnage..... | | | | | | 3,128 | |
| Total vessels..... | | | | | | 89 | |
| Total net tonnage..... | | | | | | 3,308 | |
| Boats: | | | | | | | |
| Motor..... | | 20 | 27 | 650 | 12 | 37 | |
| Other..... | | 2 | 27 | 383 | 8 | 3 | 125 |
| Accessory boats..... | | | | | | 225 | |
| Apparatus: | | | | | | | |
| Number..... | 3 | 1,095 | 1,565 | 65,147 | 935 | 126 | 185 |

Fisheries of Massachusetts, 1933—Continued

OPERATING UNITS: BY GEAR—Continued

| Item | Dredges | | | Tongs | Rakes | Forks | Hoes | By hand | Total, exclusive of duplication |
|------------------------|---------|--------|---------|--------|--------|--------|--------|---------|---------------------------------|
| | Clam | Oyster | Scallop | | | | | | |
| Fishermen: | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| On vessels..... | 5 | 8 | 116 | | | | | | 3,971 |
| On boats and shore: | | | | | | | | | |
| Regular..... | 110 | 25 | 391 | 132 | 403 | 452 | 183 | 18 | 2,781 |
| Casual..... | | | 383 | 34 | 251 | 493 | 77 | 8 | 1,652 |
| Total..... | 115 | 33 | 890 | 166 | 654 | 945 | 260 | 26 | 8,404 |
| Vessels: | | | | | | | | | |
| Steam..... | | | | | | | | | 13 |
| Net tonnage..... | | | | | | | | | 2,097 |
| Motor..... | 2 | 3 | 19 | | | | | | 365 |
| Net tonnage..... | 17 | 35 | 340 | | | | | | 13,995 |
| Total vessels..... | 2 | 3 | 19 | | | | | | 378 |
| Total net tonnage..... | 17 | 35 | 340 | | | | | | 16,092 |
| Boats: | | | | | | | | | |
| Motor..... | 59 | 12 | 378 | 28 | 45 | 66 | | | 1,478 |
| Other..... | | | 25 | 143 | 604 | 310 | 49 | | 1,624 |
| Accessory boats..... | | | | | | | | | 942 |
| Apparatus: | | | | | | | | | |
| Number..... | 63 | 34 | 2,038 | 166 | 654 | 945 | 260 | | |
| Yards at mouth..... | 31 | 44 | 1,895 | | | | | | |

CATCH: BY GEAR

| Species | Purse seines | | | | Haul seines | |
|---------------------------------|--------------|---------|--------|-------|-------------|---------|
| | Mackerel | | Other | | Pounds | Value |
| | Pounds | Value | Pounds | Value | | |
| Alewives..... | 442,300 | \$3,711 | | | 199,000 | \$1,990 |
| Bluefish..... | | | | | 1,000 | 190 |
| Butterfish..... | 6,155 | 390 | | | | |
| Eels, common..... | | | | | 90,000 | 2,700 |
| Herring, sea..... | 11,500 | 91 | | | 5,000 | 50 |
| King whiting or "kingfish"..... | 13,000 | 390 | | | | |
| Launces..... | | | 21,000 | \$420 | | |
| Mackerel..... | 27,133,285 | 648,833 | | | 20,000 | 600 |
| Rosefish..... | 2,175 | 17 | | | | |
| Shad..... | 12,420 | 188 | | | 22,500 | 1,125 |
| Sharks..... | 84 | 1 | | | | |
| Striped bass..... | | | | | 4,500 | 675 |
| "Tuna or "horse mackerel"..... | 301 | 30 | | | | |
| White perch..... | | | | | 48,500 | 7,100 |
| Total..... | 27,621,220 | 653,671 | 21,000 | 420 | 390,500 | 14,340 |

| Species | Gill nets | | | | | |
|-------------------|-----------|---------|-----------|---------|-----------|---------|
| | Anchor | | Drift | | Runaround | |
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | 25 | \$1 | | | | |
| Bluefish..... | | | 21,400 | \$2,860 | 22,000 | \$1,760 |
| Butterfish..... | 300 | 15 | 500 | 25 | | |
| Cod..... | 2,993,283 | 69,964 | 92,000 | 1,600 | | |
| Flounders..... | 6,655 | 275 | | | | |
| Grayfish..... | | | 4,000 | 40 | | |
| Haddock..... | 756,380 | 17,157 | 11,500 | 300 | | |
| Hake..... | 279,990 | 4,796 | | | | |
| Herring, sea..... | | | 5,000 | 50 | | |
| Mackerel..... | | | 2,253,796 | 68,268 | | |
| Pollock..... | 2,060,935 | 27,367 | | | | |
| Rosefish..... | 55 | 1 | | | | |
| Sea bass..... | | | | | 250 | 15 |
| Shad..... | 1,242 | 57 | 21,200 | 344 | | |
| Sharks..... | 1,490 | 45 | | | | |
| Wolfish..... | 12,820 | 139 | | | | |
| Total..... | 6,123,135 | 119,817 | 2,409,305 | 73,505 | 22,250 | 1,775 |

Fisheries of Massachusetts, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Lines | | | | | | Pound nets | | Floating traps | |
|-------------------------------------|-----------|----------|------------|-----------|-------|-------|------------|---------|----------------|--------|
| | Hand | | Trawl | | Troll | | Lb. | Value | Lb. | Value |
| | Lb. | Value | Lb. | Value | Lb. | Value | | | | |
| Alewives | | | | | | | 27,906 | \$284 | 8,825 | \$88 |
| Amberjack | | | | | | | 2,275 | 68 | | |
| Bluefish | 359,000 | \$35,248 | | | 1,500 | \$250 | 19,691 | 2,253 | 475 | 51 |
| Bonito | | | | | | | 5,204 | 232 | 18 | 1 |
| Butterfish | 10,010 | 501 | 100 | \$4 | | | 715,700 | 34,081 | 33,048 | 1,697 |
| Cod | 2,025,138 | 39,389 | 32,812,744 | 601,079 | | | 3,225 | 84 | 51,420 | 1,286 |
| Cusk | 46,753 | 584 | 4,004,062 | 47,282 | | | | | | |
| Eels: | | | | | | | | | | |
| Common | 1,150 | 72 | | | | | 7,179 | 431 | | |
| Conger | | | | | | | 597 | 30 | | |
| Flounders | 18,000 | 808 | 381,328 | 12,437 | | | 18,486 | 740 | 210 | 8 |
| Frigate mackerel | | | | | | | 52,239 | 2,090 | | |
| Grayfish | | | | | | | 5,580 | 56 | | |
| Haddock | 279,405 | 6,667 | 30,481,801 | 758,794 | | | | | | |
| Hake | 28,280 | 323 | 4,401,359 | 70,510 | | | | | | |
| Halibut | 2,613 | 306 | 1,863,422 | 176,928 | | | | | | |
| Herring, sea | | | | | | | 1,141,890 | 12,136 | 149,045 | 1,640 |
| Mackerel | 70,000 | 2,100 | | | | | 3,542,314 | 46,604 | 1,720,884 | 21,342 |
| Menhaden | | | | | | | 334 | 4 | | |
| Mullet | | | | | | | 10 | 1 | | |
| Pollock | 300,457 | 2,118 | 1,597,967 | 17,019 | | | 108,610 | 987 | 9,416 | 86 |
| Rosefish | | | 6,800 | 136 | | | 40 | 1 | | |
| Salmon | | | | | | | | | 45 | 9 |
| Scup or porgy | 228,300 | 4,814 | | | | | 110,783 | 3,398 | 3,049 | 92 |
| Sea bass | 90,500 | 3,825 | | | | | 2,172 | 110 | | |
| Shad | | | | | | | 5,099 | 257 | 107 | 5 |
| Sharks | 60 | 1 | 342 | 3 | | | 27,284 | 117 | 30 | 1 |
| Skates | | | 20,000 | 200 | | | 2,840 | 29 | | |
| Skipper or billfish | | | | | | | 4,110 | 41 | | |
| Spot | | | | | | | 985 | 30 | | |
| Squeteagues or "sea trout", spotted | | | | | | | 2,048 | 204 | | |
| Striped bass | 3,200 | 320 | | | | | 240 | 36 | | |
| Sturgeon | | | | | | | 825 | 148 | 64 | 10 |
| Tautog | 123,700 | 4,820 | | | | | 46,085 | 1,803 | 155 | 6 |
| Tuna or "horse mackerel" | | | | | | | 251,976 | 13,671 | 1,795 | 180 |
| White perch | | | | | | | 296 | 9 | | |
| Whiting | | | | | | | 6,413,067 | 64,131 | 1,252,695 | 12,527 |
| Wolfish | 7,397 | 146 | 408,150 | 5,813 | | | | | | |
| Lobsters | | | | | | | 11 | 2 | | |
| Squid | | | | | | | 482,145 | 6,081 | 33,201 | 664 |
| Total | 3,588,963 | 102,022 | 75,978,071 | 1,688,183 | 1,500 | 250 | 13,001,216 | 190,149 | 3,274,382 | 39,693 |

| Species | Weirs | | Fyke nets | | Dip nets | | Push nets | | Otter trawls | |
|----------------------------|---------|---------|-----------|-------|-----------|---------|-----------|-------|--------------|-----------|
| | Lb. | Value | Lb. | Value | Lb. | Value | Lb. | Value | Lb. | Value |
| Alewives | | | | | 235,000 | \$2,125 | | | 4,974 | \$390 |
| Bluefish | | | | | | | | | 158,707 | 7,884 |
| Butterfish | 26,165 | \$1,308 | | | | | | | 51,228,435 | 974,951 |
| Cod | 4,117 | 107 | | | | | | | 2,491,145 | 35,716 |
| Croaker | | | | | | | | | 152 | 2 |
| Cunners | | | | | | | | | 458,189 | 5,940 |
| Cusk | | | | | | | | | | |
| Drum: | | | | | | | | | | |
| Black | | | | | | | | | 32 | 1 |
| Red | | | | | | | | | 1,755 | 32 |
| Eels: | | | | | | | | | | |
| Common | | | 15,500 | \$930 | | | | | | |
| Conger | | | | | | | | | 38,020 | 703 |
| Flounders | 225 | 9 | | | | | | | 28,782,516 | 864,324 |
| Frigate mackerel | 16,560 | 662 | | | | | | | | |
| Grayfish | | | | | | | | | 2,932 | 36 |
| Haddock | | | | | | | | | 119,141,440 | 2,644,934 |
| Hake | | | | | | | | | 3,350,049 | 58,062 |
| Halibut | | | | | | | | | 535,257 | 53,237 |
| Herring, sea | 133,350 | 1,334 | | | 2,125,000 | 23,375 | | | 2,175 | 108 |
| Herring smelt | | | | | | | | | 11,648 | 313 |
| King whiting or "kingfish" | | | | | | | | | 25,296 | 645 |
| Mackerel | 800,205 | 8,002 | | | 50,000 | 1,500 | | | 11,970 | 821 |
| Pigfish | | | | | | | | | 3,832 | 73 |

Fisheries of Massachusetts, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Weirs | | Fyke nets | | Dip nets | | Push nets | | Otter trawls | |
|---------------------------------------|------------------|---------------|---------------|--------------|------------------|---------------|---------------|--------------|--------------------|------------------|
| | Lb. | Value | Lb. | Value | Lb. | Value | Lb. | Value | Lb. | Value |
| Pollock..... | 330 | \$3 | | | | | | | 7,914,083 | \$92,496 |
| Rosefish..... | | | | | | | | | 253,185 | 2,672 |
| Scup or porgy..... | 5,100 | 153 | | | | | | | 1,788,173 | 41,611 |
| Sea bass..... | | | | | | | | | 3,830,167 | 113,371 |
| Sea robin..... | | | | | | | | | 400 | 8 |
| Shad..... | 75 | 4 | | | | | | | 108 | 6 |
| Sharks..... | | | | | | | | | 4,878 | 119 |
| Sheepshead..... | | | | | | | | | 9 | 1 |
| Skates..... | | | | | | | | | 14,416 | 250 |
| Spot..... | | | | | | | | | 31,758 | 346 |
| Squeteagues or "sea trout", gray..... | | | | | | | | | 286,547 | 8,543 |
| Striped bass..... | | | | | 12,000 | \$2,900 | | | 17 | 1 |
| Sturgeon..... | | | | | | | | | 4,241 | 481 |
| Swordfish..... | | | | | | | | | 90 | 3 |
| Tautog..... | 2,915 | 117 | | | | | | | 688 | 20 |
| Tuna or "horse mackerel"..... | 2,100 | 105 | | | | | | | | |
| Whiting..... | 669,150 | 6,692 | | | | | | | 342,830 | 5,298 |
| Wolfish..... | | | | | | | | | 1,668,202 | 25,236 |
| Lobsters..... | | | | | | | | | 1,719 | 163 |
| Shrimp..... | | | | | 300 | 125 | | | 40,900 | 2,045 |
| Scallops: | | | | | | | | | | |
| Bay..... | | | | | | | 18,000 | \$7,000 | | |
| Sea..... | | | | | | | | | 810 | 149 |
| Squid..... | 33,020 | 330 | | | | | | | 22,170 | 459 |
| Total..... | 1,693,312 | 18,826 | 15,500 | \$930 | 2,422,300 | 30,025 | 18,000 | 7,000 | 220,433,920 | 4,941,450 |

| Species | Box traps | | Pots | | | | | | | | | |
|------------------------------|---------------|------------|------------------|---------------|---------------|--------------|------------------|----------------|-----------------------|--------------|--------|-------|
| | | | Crab | | Eel | | Lobster | | Periwinkle and Cockle | | | |
| | | | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | 10,000 | \$200 | | | | | | | | | | |
| Eels, common..... | | | | | 57,950 | \$3,477 | | | | | | |
| Crabs, hard..... | | | 3,273,887 | \$15,843 | | | 1,832,628 | \$8,952 | | | | |
| Lobsters..... | | | | | | | 1,926,886 | 381,484 | | | | |
| Periwinkles and cockles..... | | | | | | | | | 62,100 | \$5,950 | | |
| Total..... | 10,000 | 200 | 3,273,887 | 15,843 | 57,950 | 3,477 | 3,759,514 | 390,436 | 62,100 | 5,950 | | |

| Species | Harpoons | | Spears | | Dredges | | | | | | | |
|-------------------------------|------------------|----------------|---------------|--------------|----------------|---------------|---------------|---------------|------------------|----------------|--------|-------|
| | | | | | Clam | | Oyster | | Scallop | | | |
| | | | | | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Eels, common..... | | | 50,500 | \$3,117 | | | | | | | | |
| Swordfish..... | 2,643,422 | \$324,814 | | | | | | | | | | |
| Tuna or "horse mackerel"..... | 200 | 30 | | | | | | | | | | |
| Clams: | | | | | | | | | | | | |
| Hard, public..... | | | | | 471,900 | \$51,674 | | | | | | |
| Surf or skimmer..... | | | | | 3,240 | 325 | | | | | | |
| Mussels, sea..... | | | | | 30,000 | 3,000 | | | | | | |
| Oysters: | | | | | | | | | | | | |
| Market, private, spring..... | | | | | | | 29,762 | \$9,640 | | | | |
| Market, private, fall..... | | | | | | | 43,198 | 13,292 | | | | |
| Scallops: | | | | | | | | | | | | |
| Bay..... | | | | | | | | | 481,575 | \$173,795 | | |
| Sea..... | | | | | | | | | 1,028,287 | 115,195 | | |
| Total..... | 2,643,622 | 324,844 | 50,500 | 3,117 | 505,140 | 54,999 | 72,960 | 22,932 | 1,509,862 | 288,990 | | |

Fisheries of Massachusetts, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Tongs | | Rakes | | Forks | | Hoes | | By hand | |
|------------------------------|---------|----------|---------|-----------|-----------|---------|---------|---------|---------|-------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Periwinkles and cockles. | | | | | | | | | 3,600 | \$400 |
| Clams: | | | | | | | | | | |
| Hard, public..... | 370,750 | \$46,487 | 890,175 | \$113,687 | 5,100 | \$600 | 28,600 | \$2,600 | | |
| Hard, private..... | 2,200 | 400 | | | | | | | | |
| Razor..... | | | | | 54,400 | 2,124 | 304,000 | 13,750 | | |
| Soft, public..... | | | | | 2,680,090 | 228,744 | 176,000 | 16,725 | | |
| Surf or skimmer..... | | | 2,700 | 150 | 42,500 | 2,500 | | | 10,800 | 600 |
| Oysters: | | | | | | | | | | |
| Market, private, spring..... | 45,399 | 16,040 | | | | | | | | |
| Market, private, fall..... | 62,250 | 21,065 | | | | | | | | |
| Scallops, bay..... | | | 540 | 180 | | | | | | |
| Irish moss..... | | | 11,650 | 582 | | | | | | |
| Bloodworms..... | | | | | 17,759 | 28,000 | | | | |
| Sandworms..... | | | | | 21,656 | 19,500 | | | | |
| Total..... | 480,599 | 83,992 | 905,065 | 114,599 | 2,801,505 | 281,468 | 508,600 | 33,075 | 14,400 | 1,000 |

OPERATING UNITS: BY COUNTIES

| Item | Barnstable | Bristol | Dukes | Essex | Nantucket | Norfolk | Plymouth | Suffolk |
|----------------------------|------------|---------|--------|-----------|-----------|---------|----------|-----------|
| Fishermen: | Number | Number | Number | Number | Number | Number | Number | Number |
| On vessels..... | 158 | 225 | 59 | 1,736 | 95 | | | 1,698 |
| On boats and shore: | | | | | | | | |
| Regular..... | 761 | 248 | 241 | 633 | 62 | 71 | 251 | 514 |
| Casual..... | 266 | 91 | 330 | 372 | 150 | 159 | 182 | 102 |
| Total..... | 1,185 | 564 | 630 | 2,741 | 307 | 230 | 433 | 2,314 |
| Vessels: | | | | | | | | |
| Steam: | | | | | | | | 13 |
| Net tonnage..... | | | | | | | | 2,097 |
| Motor: | 30 | 31 | 10 | 132 | 20 | | | 142 |
| Net tonnage..... | 363 | 786 | 137 | 5,519 | 256 | | | 6,934 |
| Total vessels..... | 30 | 31 | 10 | 132 | 20 | | | 155 |
| Total net tonnage..... | 363 | 786 | 137 | 5,519 | 256 | | | 9,031 |
| Boats: | | | | | | | | |
| Motor..... | 341 | 148 | 197 | 322 | 105 | 46 | 183 | 136 |
| Other..... | 479 | 152 | 222 | 406 | 50 | 26 | 242 | 47 |
| Accessory boats..... | 48 | 32 | 22 | 530 | 2 | | | 308 |
| Apparatus: | | | | | | | | |
| Furse seines: | | | | | | | | |
| Mackerel..... | 4 | 2 | | 62 | | | | 29 |
| Length, yards..... | 1,720 | 240 | | 30,140 | | | | 13,620 |
| Other..... | 2 | | | | | | | |
| Length, yards..... | 260 | | | | | | | |
| Haul seines..... | 2 | | 1 | 5 | 6 | | 1 | |
| Length, yards..... | 55 | | 400 | 475 | 700 | | 25 | |
| Gill nets: | | | | | | | | |
| Anchor..... | | | | 704 | | | | |
| Square yards..... | | | | 280,655 | | | | |
| Drift..... | 1,070 | 45 | 45 | 2,070 | 5 | 6 | 14 | 850 |
| Square yards..... | 304,360 | 18,900 | 24,500 | 824,740 | 2,250 | 1,520 | 6,240 | 348,450 |
| Runaround..... | 1 | | | | | | | |
| Square yards..... | 1,800 | | | | | | | |
| Lines: | | | | | | | | |
| Hand..... | 105 | 104 | 39 | 170 | 30 | 10 | 10 | 37 |
| Hooks..... | 135 | 208 | 39 | 194 | 30 | 10 | 10 | 45 |
| Trawl..... | 3,640 | 200 | 280 | 22,067 | 60 | | 480 | 21,109 |
| Hooks..... | 186,600 | 7,200 | 11,800 | 1,095,124 | 2,100 | | 17,280 | 1,060,058 |
| Troll..... | 8 | | | | | | | |
| Hooks..... | 8 | | | | | | | |
| Pound nets..... | 62 | 14 | 7 | | 3 | | | |
| Floating traps..... | 4 | | | 13 | | | | |
| Weirs..... | 4 | | | | | | | |
| Fyke nets..... | 20 | 4 | | | | | | |
| Dip nets..... | 15 | | | 66 | | 2 | 17 | 30 |
| Push nets..... | | | 50 | | | | | |
| Otter trawls..... | 52 | 32 | 22 | 72 | 22 | | | 166 |
| Yards at mouth..... | 1,338 | 1,063 | 573 | 2,247 | 640 | | | 15,223 |
| Box traps..... | | | | | | | 3 | |
| Pots: | | | | | | | | |
| Crab..... | | | | 45 | | 204 | 41 | 805 |
| Eel..... | 495 | 118 | 320 | 115 | 450 | | 67 | |
| Lobster..... | 6,557 | 4,882 | 12,743 | 17,479 | 1,116 | 2,903 | 13,966 | 5,501 |
| Periwinkle and cockle..... | | 580 | | 200 | | | 156 | |

Fisheries of Massachusetts, 1933—Continued

OPERATING UNITS: BY COUNTIES—Continued

| Item | Barnstable | Bristol | Dukes | Essex | Nantucket | Norfolk | Plymouth | Suffolk |
|----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| Apparatus—Continued. | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| Harpoons..... | 17 | 25 | 18 | 49 | 3 | | | 14 |
| Spears..... | 76 | 19 | 6 | 20 | 12 | 14 | 38 | |
| Dredges: | | | | | | | | |
| Clam..... | 25 | 16 | 6 | | 9 | | 3 | 4 |
| Yards at mouth..... | 10 | 7 | 3 | | 5 | | 2 | 4 |
| Oyster..... | 14 | | | | | | 20 | |
| Yards at mouth..... | 24 | | | | | | 20 | |
| Scallop..... | 276 | 197 | 830 | 2 | 638 | | 95 | |
| Yards at mouth..... | 258 | 212 | 738 | 7 | 595 | | 85 | |
| Tongs..... | 46 | 76 | | | 15 | | 29 | |
| Rakes..... | 358 | 65 | 109 | | 35 | | 87 | |
| Forks..... | 26 | 6 | | 579 | | 160 | 11 | 163 |
| Hoes..... | 129 | | | | | 30 | 101 | |

CATCH: BY COUNTIES

| Species | Barnstable | | Bristol | | Dukes | | Essex | |
|---------------------------------|---------------|--------------|---------------|--------------|---------------|--------------|---------------|--------------|
| | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> |
| Alewives..... | 314,000 | \$2,540 | 4,506 | \$50 | 4,400 | \$44 | 396,350 | \$3,568 |
| Amberjack..... | 2,275 | 68 | | | | | | |
| Bluefish..... | 329,795 | 33,507 | 6,306 | 884 | 23,065 | 2,911 | 3,926 | 287 |
| Bonito..... | 1,183 | 40 | 526 | 21 | 3,178 | 159 | 18 | 1 |
| Butterfish..... | 597,693 | 28,256 | 41,395 | 2,093 | 100,482 | 5,025 | 112,644 | 5,105 |
| Cod..... | 2,167,492 | 47,329 | 2,075,929 | 37,125 | 120,002 | 2,968 | 22,437,126 | 421,879 |
| Croaker..... | 17,512 | 323 | 48,215 | 1,123 | 468 | 8 | 2,105,090 | 29,324 |
| Cunners..... | | | | | | | | 152 |
| Cusk..... | 142,772 | 1,275 | 1,208 | 19 | 345 | 3 | 1,767,972 | 21,176 |
| Drum: | | | | | | | | |
| Black..... | | | | | | | | 32 |
| Red..... | | | 175 | 2 | | | 1,580 | 30 |
| Eels: | | | | | | | | |
| Common..... | 67,230 | 4,034 | 17,299 | 1,063 | 16,200 | 972 | 101,500 | 3,345 |
| Conger..... | 70 | 1 | 3,216 | 83 | | | 26,772 | 514 |
| Flounders..... | 3,022,419 | 67,661 | 5,626,867 | 186,812 | 954,170 | 24,753 | 3,432,129 | 122,896 |
| Frigate mackerel..... | 68,799 | 2,752 | | | | | | |
| Grayfish..... | 9,580 | 96 | | | | | 2,932 | 36 |
| Haddock..... | 2,046,138 | 48,745 | 6,862,729 | 169,075 | 31,740 | 808 | 22,444,944 | 527,084 |
| Hake..... | 276,526 | 3,088 | 60,624 | 907 | | | 2,699,382 | 42,669 |
| Halibut..... | 8,407 | 1,004 | 24,408 | 2,332 | | | 1,618,750 | 154,589 |
| Herring, sea..... | 1,353,786 | 14,255 | 27,054 | 271 | 200 | 2 | 485,745 | 7,113 |
| King whiting or "kingfish"..... | 732 | 32 | 300 | 8 | 115 | 2 | 34,981 | 937 |
| Launce..... | 21,000 | 420 | | | | | | |
| Mackerel..... | 6,068,862 | 93,535 | 115,267 | 2,395 | 90,700 | 1,949 | 21,160,924 | 499,090 |
| Menhaden..... | 189 | 2 | 175 | 2 | | | | |
| Mullet..... | | | 10 | 1 | | | | |
| Pigfish..... | | | | | | | 3,654 | 68 |
| Pollock..... | 259,680 | 2,448 | 22,927 | 284 | 220 | 1 | 4,065,136 | 48,730 |
| Rosefish..... | 6,500 | 130 | | | 40 | 1 | 33,830 | 299 |
| Salmon..... | | | | | | | 45 | 9 |
| Scup or porgy..... | 43,143 | 1,209 | 305,821 | 6,302 | 70,635 | 2,119 | 1,401,941 | 33,881 |
| Sea bass..... | 37,116 | 1,232 | 187,741 | 6,092 | 6,215 | 311 | 3,108,326 | 88,458 |
| Sea robin..... | | | 400 | 8 | | | | |
| Shad..... | 4,877 | 246 | 106 | 5 | 19 | 1 | 57,265 | 1,716 |
| Sharks..... | 516 | 5 | 26,748 | 112 | | | 4,070 | 92 |
| Sheepshead..... | | | | | | | | 9 |
| Skates..... | 1,160 | 12 | 2,340 | 24 | 500 | 5 | 108 | 2 |
| Skipper or "billfish"..... | 4,110 | 41 | | | | | | |
| Smelt..... | | | | | | | 100 | 5 |
| Spot..... | | | 985 | 30 | | | 16,265 | 202 |
| Squeteagues or "sea trout": | | | | | | | | |
| Gray..... | 1,021 | 49 | 4,119 | 192 | | | 216,709 | 6,507 |
| Spotted..... | 97 | 9 | 1,644 | 164 | 307 | 31 | | |
| Striped bass..... | 4,500 | 675 | 240 | 36 | | | 12,017 | 2,901 |
| Sturgeon..... | 541 | 92 | 218 | 15 | 64 | 8 | 2,602 | 337 |
| Swordfish..... | 276,476 | 25,864 | 269,473 | 25,194 | 266,837 | 24,700 | 1,533,102 | 210,700 |
| Tautog..... | 34,757 | 1,242 | 119,732 | 4,770 | 8,211 | 328 | 803 | 25 |
| Tuna or "horse mackerel"..... | 254,076 | 13,776 | | | | | 2,296 | 240 |
| White perch..... | | | 286 | 9 | 3,500 | 350 | | |
| Whiting..... | 7,552,908 | 75,533 | 309 | 4 | | | 795,020 | 8,240 |
| Wolfish..... | 96,640 | 909 | 16,687 | 216 | | | 262,675 | 4,117 |
| Crabs, hard..... | | | | | | | 477,187 | 3,013 |
| Lobsters..... | 124,432 | 31,954 | 85,159 | 14,648 | 225,987 | 43,047 | 680,241 | 130,866 |
| Shrimp..... | 300 | 125 | | | 40,900 | 2,045 | | |
| Periwinkles and cockles..... | 32,400 | 3,200 | | | | | 6,300 | 700 |

Fisheries of Massachusetts, 1933—Continued

CATCH: BY COUNTIES—Continued

| Species | Barnstable | | Bristol | | Dukes | | Essex | |
|------------------------------|------------|----------|------------|----------|-----------|---------|------------|-----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Clams: | | | | | | | | |
| Hard, public..... | 704,385 | \$38,602 | 510,530 | \$64,655 | 154,000 | \$15,05 | | |
| Hard, private..... | 2,200 | 40 | | | | | | |
| Razor..... | 240,000 | 11 25 | | | | | 64,400 | \$2,124 |
| Soft, public..... | 77,570 | 9,340 | 3,740 | 440 | | | 1,892,200 | 173,157 |
| Surf or skimmer..... | 3,240 | 325 | | | 2,700 | 150 | 42,500 | 2,500 |
| Oysters: | | | | | | | | |
| Market, private, spring..... | 65,963 | 22,88 | | | | | | |
| Market, private, fall..... | 85,738 | 28,35 | | | | | | |
| Scallops: | | | | | | | | |
| Bay..... | 66,945 | 21,74 | 14,310 | 5,585 | 248,850 | 96,85 | | |
| Sea..... | 170,564 | 20,64 | 528,157 | 57,93 | 175,050 | 10,45 | 52,854 | 5,891 |
| Squid..... | 457,746 | 5,718 | 44,370 | 563 | 13,039 | 130 | 60,222 | 1,000 |
| Bloodworms..... | | | | | | | 2,544 | 4,000 |
| Sandworms..... | | | | | | | 4,778 | 4,300 |
| Total..... | 27,125,981 | 717,057 | 17,152,311 | 591,561 | 2,572,139 | 245,181 | 93,612,148 | 2,573,727 |

| Species | Nantucket | | Norfolk | | Plymouth | | Suffolk | |
|---------------------------------------|-----------|---------|---------|--------|-----------|---------|-------------|-----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | | | | | | | | |
| Bluefish..... | 31,379 | \$3,155 | 20,000 | \$200 | 175,000 | \$1,935 | 8,800 | \$62 |
| Bonito..... | 317 | 12 | | | | | 35,569 | 2,138 |
| Butterfish..... | 21,561 | 983 | | | | | 77,810 | 4,443 |
| Cod..... | 916,378 | 16,458 | 13,800 | 360 | 115,000 | 3,000 | 61,364,634 | 1,159,341 |
| Croaker..... | | | | | | | 319,860 | 4,938 |
| Cusk..... | | | | | | | 2,596,707 | 31,293 |
| Eels: | | | | | | | | |
| Common..... | 3,000 | 270 | 2,150 | 132 | 14,900 | 911 | | |
| Conger..... | 1,351 | 15 | | | | | 7,208 | 120 |
| Flounders..... | 3,217,166 | 95,856 | 3,000 | 150 | 4,600 | 230 | 10,887,046 | 380,243 |
| Haddock..... | 1,608,930 | 40,965 | | | 17,250 | 600 | 117,668,745 | 2,638,575 |
| Hake..... | 16,519 | 257 | | | | | 5,003,627 | 86,770 |
| Halibut..... | | | | | | | 749,727 | 72,544 |
| Herring, sea..... | | | | | 200,000 | 2,000 | 1,506,175 | 15,143 |
| Herring smelt..... | | | | | | | 2,630 | 74 |
| King whiting or "kingfish"..... | | | | | | | 2,168 | 56 |
| Mackerel..... | 17,135 | 514 | 6,000 | 180 | 21,000 | 480 | 8,131,475 | 199,935 |
| Pigfish..... | | | | | | | 178 | 5 |
| Pollock..... | 11,154 | 201 | | | 1,650 | 15 | 7,631,031 | 88,394 |
| Rosefish..... | 509 | 5 | | | | | 221,385 | 2,392 |
| Scup or porgy..... | 51,223 | 991 | | | 1,500 | 45 | 269,147 | 5,521 |
| Sea bass..... | 96,651 | 2,797 | | | | | 487,040 | 18,401 |
| Shad..... | 172 | 9 | | | | | 312 | 9 |
| Sharks..... | | | | | | | 2,814 | 436 |
| Skates..... | | | | | | | 33,158 | 436 |
| Smelt..... | | | | | | | 8,918 | 234 |
| Spot..... | | | | | | | 15,493 | 144 |
| Squeteagues or "sea trout", gray..... | 142 | 11 | | | | | 64,556 | 1,784 |
| Striped bass..... | 3,200 | 320 | | | | | 1,421 | 131 |
| Sturgeon..... | 284 | 56 | | | | | 213,002 | 31,383 |
| Swordfish..... | 83,722 | 6,976 | | | | | 40 | 1 |
| Tautog..... | | | | | 10,000 | 400 | | |
| White perch..... | 45,000 | 6,750 | | | | | | |
| Whiting..... | | | | | | | 329,505 | 4,871 |
| Wolfish..... | | | | | | | 1,720,567 | 26,092 |
| Crabs, hard..... | | | 117,050 | 1,075 | 145,037 | 1,072 | 4,367,241 | 19,635 |
| Lobsters..... | 8,358 | 2,356 | 114,137 | 23,993 | 531,702 | 109,266 | 138,600 | 25,489 |
| Periwinkles and cockles..... | | | | | 27,000 | 2,450 | | |
| Clams: | | | | | | | | |
| Hard, public..... | 200,750 | 24,662 | 28,600 | 2,600 | 138,200 | 18,449 | | |
| Razor..... | | | | | 64,000 | 2,500 | | |
| Soft, public..... | | | 365,500 | 21,500 | 113,900 | 9,325 | 383,250 | 31,687 |
| Surf or skimmer..... | | | | | 10,800 | 600 | | |
| Mussels, sea..... | | | | | | | 30,000 | 3,000 |
| Oysters: | | | | | | | | |
| Market, private, spring..... | | | | | 9,198 | 2,800 | | |
| Market, private, fall..... | | | | | 19,710 | 6,000 | | |
| Scallops: | | | | | | | | |
| Bay..... | 165,600 | 55,200 | | | 4,410 | 1,620 | | |
| Sea..... | 102,070 | 11,340 | | | | | 382 | 77 |
| Squid..... | 100 | 2 | | | | | 5,059 | 151 |
| Irish moss..... | | | | | 11,650 | 582 | | |
| Bloodworms..... | | | 11,400 | 18,000 | | | 3,815 | 6,000 |
| Sandworms..... | | | 11,100 | 10,000 | | | 5,778 | 5,200 |
| Total..... | 6,602,682 | 270,164 | 692,737 | 78,190 | 1,656,507 | 164,310 | 224,265,773 | 4,806,790 |

RHODE ISLAND

Fisheries of Rhode Island, 1933

OPERATING UNITS: BY GEAR

| Item | Purse seines | | Haul seines | Gill nets, drift | Lines | | | | Pounds nets |
|-------------------------------|---------------|---------------|---------------|------------------|---------------|---------------|---------------|-----------------|---------------|
| | Mack-erel | Men-haden | | | Hand | Trawl | Troll | Trot with hooks | |
| Fishermen: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| On vessels..... | 11 | 20 | | 3 | 28 | 4 | | | 13 |
| On boats and shore: | | | | | | | | | |
| Regular..... | | | 28 | 3 | 77 | 24 | 22 | | 35 |
| Casual..... | | | 37 | 2 | 35 | | 2 | 1 | 16 |
| Total..... | 11 | 20 | 65 | 8 | 140 | 28 | 24 | 1 | 64 |
| Vessels: | | | | | | | | | |
| Steam..... | | 1 | | | | | | | |
| Net tonnage..... | | 45 | | | | | | | |
| Motor..... | 2 | | | 1 | 13 | 2 | | | 3 |
| Net tonnage..... | 12 | | | 13 | 98 | 17 | | | 29 |
| Total vessels..... | 2 | 1 | | 1 | 13 | 2 | | | 3 |
| Total net tonnage..... | 12 | 45 | | 13 | 98 | 17 | | | 29 |
| Boats: | | | | | | | | | |
| Motor..... | | | | 2 | 83 | 12 | 22 | | 1 |
| Other..... | | | 26 | 1 | 13 | | | 1 | 38 |
| Accessory boats..... | 1 | 2 | | | | | | | 7 |
| Apparatus: | | | | | | | | | |
| Number..... | 2 | 1 | 20 | 44 | 248 | 51 | 48 | 1 | 64 |
| Length, yards..... | 420 | 400 | 2,512 | | | | | | |
| Square yards..... | | | | 17,800 | | | | | |
| Hooks, baits or snoods..... | | | | | 319 | 24,700 | 48 | 200 | |

| Item | Float- ing traps | Fyke nets | Otter trawls | Pots | | | Har- poons | Spears |
|-------------------------------|------------------------|---------------|-----------------|---------------|---------------|----------------------------------|---------------|---------------|
| | | | | Eel | Lob- ster | Peri- winkle and cockle | | |
| Fishermen: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| On vessels..... | 111 | | 103 | | 21 | | 72 | |
| On boats and shore: | | | | | | | | |
| Regular..... | 34 | 4 | 34 | 21 | 162 | 23 | 26 | 3 |
| Casual..... | 1 | 3 | 6 | 19 | 128 | 12 | 2 | 10 |
| Total..... | 146 | 7 | 143 | 40 | 311 | 35 | 100 | 13 |
| Vessels: | | | | | | | | |
| Steam..... | 4 | | 5 | | | | | 3 |
| Net tonnage..... | 45 | | 55 | | | | | |
| Motor..... | 7 | | 33 | | 10 | | | 22 |
| Net tonnage..... | 77 | | 292 | | 69 | | | 185 |
| Total vessels..... | 11 | | 38 | | 10 | | | 22 |
| Total net tonnage..... | 122 | | 317 | | 69 | | | 185 |
| Boats: | | | | | | | | |
| Motor..... | 1 | 4 | 23 | 21 | 203 | 27 | 14 | |
| Other..... | 21 | 1 | | 13 | 26 | | | 12 |
| Accessory boats..... | 37 | | | | | | 27 | |
| Apparatus: | | | | | | | | |
| Number..... | 40 | 37 | 61 | 1,698 | 53,672 | 1,585 | 36 | 13 |
| Yards at mouth..... | | | 1,635 | | | | | |

Fisheries of Rhode Island, 1933—Continued

OPERATING UNITS: BY GEAR—Continued

| Item | Dredges | | | | Tongs | Rakes | Forks | Hoes | By hand | Total, exclusive of duplication |
|------------------------|---------|--------|--------|---------|--------|--------|--------|--------|---------|---------------------------------|
| | Clam | Mussel | Oyster | Scallop | | | | | | |
| | Number | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | | | | |
| On vessels..... | 7 | | 63 | | | | | | | 328 |
| On boats and shore: | | | | | | | | | | |
| Regular..... | 17 | 1 | | 78 | 107 | 25 | 4 | 3 | 12 | 371 |
| Casual..... | 5 | | | 34 | 436 | 50 | 4 | 23 | 16 | 677 |
| Total..... | 29 | 1 | 63 | 112 | 543 | 75 | 8 | 26 | 28 | 1,376 |
| Vessels: | | | | | | | | | | |
| Steam..... | | | 2 | | | | | | | 8 |
| Net tonnage..... | | | 101 | | | | | | | 201 |
| Motor..... | 2 | | 13 | | | | | | | 65 |
| Net tonnage..... | 18 | | 249 | | | | | | | 695 |
| Total vessels..... | 2 | | 15 | | | | | | | 73 |
| Total net tonnage..... | 18 | | 350 | | | | | | | 896 |
| Boats: | | | | | | | | | | |
| Motor..... | 10 | 1 | | 70 | 162 | 24 | 1 | 1 | 1 | 433 |
| Other..... | | | | | 365 | 47 | 6 | 11 | | 506 |
| Accessory boats..... | | | | | | | | | | 74 |
| Apparatus: | | | | | | | | | | |
| Number..... | 12 | 1 | 30 | 374 | 543 | 75 | 8 | 26 | | |
| Yards at mouth..... | 9 | 1 | 45 | 324 | | | | | | |

CATCH: BY GEAR

| Species | Purse seines | | | | Haul seines | | Gill nets, drift | |
|---------------------------------------|--------------|-------|-----------|---------|-------------|---------|------------------|---------|
| | Mackerel | | Menhaden | | | | | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | | | | | 160,000 | \$1,600 | | |
| Bluefish..... | | | | | 16,250 | 1,040 | 19,375 | \$1,240 |
| Eels, common..... | | | | | 16,599 | 1,280 | | |
| Herring, sea..... | 30,000 | \$300 | | | 102,000 | 1,620 | | |
| Mackerel..... | 37,500 | 1,350 | | | | | 30,000 | 600 |
| Menhaden..... | | | 1,000,000 | \$2,500 | | | | |
| Mullet..... | | | | | 6,000 | 180 | | |
| Smelt..... | | | | | 2,000 | 200 | | |
| Squeteagues or "sea trout", gray..... | | | | | 8,750 | 560 | 2,500 | 160 |
| Striped bass..... | | | | | 2,800 | 280 | | |
| Tautog..... | | | | | 3,000 | 90 | | |
| Thimble-eyed mackerel..... | 45,000 | 675 | | | | | | |
| Total..... | 112,500 | 2,325 | 1,000,000 | 2,500 | 377,399 | 6,850 | 51,875 | 2,000 |

| Species | Lines | | | | | | | |
|---------------------------------------|---------|--------|---------|---------|--------|---------|-----------------|-------|
| | Hand | | Trawl | | Troll | | Trot with hooks | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Bluefish..... | 2,875 | \$184 | | | | | | |
| Cod..... | 610,425 | 10,682 | 164,500 | \$2,693 | 34,500 | \$2,048 | | |
| Eels, common..... | 3,638 | 295 | | | | | 2,400 | \$360 |
| Flounders..... | 4,000 | 320 | | | | | | |
| Haddock..... | | | 6,250 | 250 | | | | |
| Mackerel..... | | | | | 4,300 | 129 | | |
| Sea bass..... | 13,350 | 1,003 | | | | | | |
| Squeteagues or "sea trout", gray..... | 1,750 | 112 | | | | | | |
| Striped bass..... | 2,900 | 290 | | | | | | |
| Tautog..... | 125,550 | 3,895 | | | | | | |
| Thimble-eyed mackerel..... | 460 | 9 | | | | | | |
| Tuna or "horse mackerel"..... | | | | | 21,467 | 483 | | |
| Total..... | 764,948 | 16,790 | 170,750 | 2,943 | 60,267 | 2,680 | 2,400 | 360 |

Fisheries of Rhode Island, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Pound nets | | Floating traps | | Fyke nets | | Otter trawls | |
|--|------------------|---------------|------------------|----------------|---------------|------------|------------------|---------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | 16,000 | \$315 | | | | | | |
| Bluefish..... | 13,969 | 660 | 107,417 | \$5,634 | | | | |
| Bonito..... | 5,938 | 208 | 40,334 | 1,715 | | | | |
| Butterfish..... | 178,366 | 5,495 | 308,156 | 10,709 | | | | |
| Cod..... | 133 | 3 | 128,183 | 3,277 | | | 5,000 | \$80 |
| Creville..... | | | 2,200 | 81 | | | | |
| Cunners..... | | | 7,440 | 111 | | | 34,000 | 340 |
| Fels: | | | | | | | | |
| Common..... | 49,994 | 3,761 | | | 2,700 | \$200 | | |
| Conger..... | 643 | 23 | 315 | 6 | | | | |
| Flounders..... | 69,110 | 2,324 | 301,242 | 10,139 | 1,225 | 37 | 2,877,500 | 78,110 |
| Frigate mackerel..... | 2,640 | 26 | 54,014 | 1,583 | | | | |
| Grayfish..... | | | 600 | 12 | | | | |
| Haddock..... | | | 718 | 28 | | | 17,000 | 425 |
| Hake..... | 171 | 3 | 9,000 | 180 | | | | |
| Herring, sea..... | 212,746 | 3,327 | 109,800 | 1,135 | | | 222,400 | 5,065 |
| Hickory shad..... | 2,006 | 20 | 300 | 3 | | | | |
| King whiting or "kingfish"..... | 1,115 | 39 | 325 | 14 | | | | |
| Mackerel..... | 171,363 | 5,668 | 478,126 | 14,490 | | | | |
| Menhaden..... | 120 | 1 | 8,396 | 34 | | | | |
| Mullet..... | 11 | 1 | 2,800 | 140 | | | | |
| Pollock..... | 785 | 21 | 16,457 | 430 | | | | |
| Scup or porgy..... | 123,578 | 2,337 | 1,878,853 | 34,783 | | | 17,000 | 313 |
| Sea bass..... | 2,938 | 108 | 31,977 | 1,432 | | | | |
| Sea robin..... | 4,175 | 41 | 69,110 | 646 | | | 500 | 10 |
| Shad..... | 3,174 | 398 | 7,586 | 740 | | | | |
| Sharks..... | | | 1,947 | 28 | | | | |
| Skates..... | | | 8,410 | 83 | | | 168,000 | 1,245 |
| Skipper or "billfish"..... | | | 730 | 11 | | | | |
| Smolt..... | 2,250 | 225 | | | | | | |
| Spot..... | | | 45 | 1 | | | | |
| Squeteagues or "sea trout", gray..... | 18,636 | 1,136 | 31,624 | 2,044 | | | | |
| Striped bass..... | 2,110 | 99 | 31,423 | 1,749 | | | | |
| Sturgeon..... | | | 449 | 37 | | | | |
| Swordfish..... | | | 852 | 14 | | | | |
| Tautog..... | 98,371 | 2,314 | 33,902 | 914 | 6,800 | 189 | | |
| Thimble-eyed mackerel..... | 18,192 | 637 | 14,055 | 212 | | | | |
| Tomcod..... | 290 | 1 | | | | | | |
| Tuna or "horse mackerel"..... | 2,018 | 63 | 21,092 | 938 | | | | |
| White perch..... | 711 | 39 | 150 | 3 | | | | |
| Whiting..... | 162,567 | 1,026 | 338,664 | 3,905 | | | 223,275 | 2,095 |
| Crabs, king..... | 6,574 | 18 | | | | | | |
| Periwinkles and cockles..... | | | | | | | 272 | 4 |
| Squid..... | 52,259 | 1,189 | 396,867 | 8,697 | | | 52,725 | 1,732 |
| Total..... | 1,222,915 | 32,026 | 4,436,162 | 105,961 | 10,225 | 426 | 3,617,672 | 89,439 |

| Species | Pots | | | | | | Harpoons | | Spears | |
|---------------------------------|---------------|--------------|----------------|----------------|--------------------------|--------------|----------------|---------------|--------------|------------|
| | Eel | | Lobster | | Periwinkle and cockle | | | | | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Eels, common..... | 63,627 | \$5,466 | | | | | | | 6,653 | \$751 |
| Swordfish..... | | | | | | | 259,358 | \$23,785 | | |
| Crabs, hard..... | | | 20,478 | \$369 | | | | | | |
| Lobsters..... | | | 708,095 | 113,933 | | | | | | |
| Periwinkles and cockles..... | | | | | 109,170 | \$5,197 | | | | |
| Total..... | 63,627 | 5,466 | 728,573 | 114,302 | 109,170 | 5,197 | 259,358 | 23,785 | 6,653 | 751 |

| Species | Dredges | | | | | | | |
|------------------------------|---------------|---------------|------------|-----------|------------------|----------------|---------------|---------------|
| | Clam | | Mussel | | Oyster | | Scallop | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Clams: | | | | | | | | |
| Hard, public..... | 75,625 | \$8,594 | | | | | | |
| Hard, private..... | 24,200 | 3,300 | | | 64,845 | \$8,949 | | |
| Mussels, sea..... | | | 650 | \$38 | | | | |
| Oysters: | | | | | | | | |
| Market, private, spring..... | | | | | 1,525,104 | 211,688 | | |
| Market, private, fall..... | | | | | 1,528,635 | 211,807 | | |
| Scallops, bay..... | | | | | | | 63,120 | \$12,943 |
| Total..... | 99,825 | 11,894 | 650 | 38 | 3,116,584 | 432,444 | 63,120 | 12,943 |

Fisheries of Rhode Island, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Tongs | | Rakes | | Forks | | Hoes | | By hand | |
|------------------------------|---------|-----------|---------|----------|--------|---------|--------|-------|---------|-------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Clams: | | | | | | | | | | |
| Hard, public..... | 868,560 | \$100,554 | 128,601 | \$15,036 | | | 4,050 | \$713 | 1,155 | \$150 |
| Soft, public..... | | | | | 10,320 | \$1,578 | | | | |
| Oysters: | | | | | | | | | | |
| Market, public, spring..... | | | 26,880 | 3,840 | | | | | | |
| Market, public, fall..... | 7,242 | 1,051 | 27,699 | 3,957 | | | | | 3,503 | 499 |
| Market, private, spring..... | 280 | 60 | | | | | | | | |
| Market, private, fall..... | 420 | 90 | | | | | | | | |
| Total..... | 868,502 | 101,755 | 183,180 | 22,833 | 10,320 | 1,578 | 27,014 | 3,349 | 4,658 | 649 |

OPERATING UNITS: BY COUNTIES

| Item | Bristol | Kent | Newport | Providence | Washington |
|----------------------------|---------|--------|---------|------------|------------|
| | Number | Number | Number | Number | Number |
| Fishermen: | | | | | |
| On vessels..... | 26 | 3 | 267 | 24 | 18 |
| On boats and shore: | | | | | |
| Regular..... | 17 | 88 | 144 | 10 | 112 |
| Casual..... | 93 | 249 | 157 | 108 | 70 |
| Total..... | 136 | 340 | 558 | 142 | 200 |
| Vessels: | | | | | |
| Steam..... | | | 6 | 2 | |
| Net tonnage..... | | | 100 | 101 | |
| Motor..... | 7 | 1 | 48 | 2 | 7 |
| Net tonnage..... | 107 | 11 | 418 | 82 | 77 |
| Total vessels..... | 7 | 1 | 54 | 4 | 7 |
| Total net tonnage..... | 107 | 11 | 518 | 183 | 77 |
| Boats: | | | | | |
| Motor..... | 36 | 122 | 175 | 12 | 88 |
| Other..... | 67 | 197 | 67 | 105 | 70 |
| Accessory boats..... | | | 74 | | |
| Apparatus: | | | | | |
| Purse seines: | | | | | |
| Mackerel..... | | | 2 | | |
| Length, yards..... | | | 420 | | |
| Menhaden..... | | | 1 | | |
| Length, yards..... | | | 400 | | |
| Haul seines..... | 2 | 3 | | 9 | 6 |
| Length, yards..... | 100 | 225 | | 667 | 1,520 |
| Gill nets, drift..... | | | 42 | | 2 |
| Square yards..... | | | 15,300 | | 2,500 |
| Lines: | | | | | |
| Hand..... | 4 | | 172 | 11 | 61 |
| Hooks..... | 8 | | 213 | | 98 |
| Trawl..... | | | 43 | | 8 |
| Hooks..... | | | 20,700 | | 4,000 |
| Troll..... | | | 16 | | 32 |
| Hooks..... | | | 16 | | 32 |
| Trot with hooks..... | | | | 1 | |
| Hooks..... | | | | 200 | |
| Pound nets..... | 1 | | 45 | 1 | 17 |
| Floating traps..... | | | 30 | | 10 |
| Fyke nets..... | 11 | 26 | | | |
| Otter trawls..... | | | 45 | | 16 |
| Yards at mouth..... | | | 1,211 | | 424 |
| Pots: | | | | | |
| Eel..... | 90 | 343 | 430 | 145 | 690 |
| Lobster..... | 2,805 | 4,565 | 36,662 | 585 | 9,055 |
| Periwinkle and cockle..... | 500 | | 490 | | 595 |
| Harpoons..... | 1 | | 32 | | 3 |
| Spears..... | 1 | | | 10 | 2 |
| Dredges: | | | | | |
| Clam..... | | 3 | 9 | | |
| Yards at mouth..... | | 1 | 8 | | |
| Mussel..... | 1 | | | | |
| Yards at mouth..... | | | | | |
| Oyster..... | 14 | 2 | | 8 | 6 |
| Yards at mouth..... | 21 | 2 | | 12 | 9 |
| Scallop..... | | 259 | 61 | | 48 |
| Yards at mouth..... | | 230 | 48 | | 41 |
| Tongs..... | 84 | 203 | 26 | 97 | 43 |
| Rakes..... | | 40 | 13 | 1 | 21 |
| Forks..... | 2 | | | | 6 |
| Hoes..... | 7 | 4 | 2 | 7 | 6 |

Fisheries of Rhode Island, 1933—Continued

CATCH: BY COUNTIES

| Species | Bristol | | Kent | | Newport | | Providence | | Washington | |
|----------------------------------|------------------|----------------|----------------|----------------|-------------------|----------------|------------------|----------------|------------------|----------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives | 10,000 | \$75 | | | | | | | 166,000 | \$1,840 |
| Bluefish | | | | | 129,641 | \$6,662 | | | 64,745 | 4,144 |
| Bonito | | | | | 46,212 | 1,914 | | | 110 | 9 |
| Butterfish | | | | | 424,513 | 14,306 | | | 62,009 | 1,899 |
| Cod | 2,312 | 37 | | | 776,160 | 14,227 | | | 129,769 | 2,471 |
| Crevalle | | | | | 2,200 | 81 | | | | |
| Cunners | | | | | 41,440 | 451 | | | | |
| Eels: | | | | | | | | | | |
| Common | 2,983 | 175 | 11,815 | \$890 | 58,964 | 3,902 | 37,382 | \$3,985 | 34,467 | 3,161 |
| Conger | | | | | 958 | 29 | | | | |
| Flounders | 400 | 10 | 925 | 28 | 2,604,467 | 73,901 | | | 647,285 | 16,991 |
| Frigate mackerel | | | | | 56,294 | 1,598 | | | 360 | 11 |
| Grayfish | | | | | 600 | 12 | | | | |
| Haddock | | | | | 23,968 | 703 | | | | |
| Hake | | | | | 9,171 | 183 | | | | |
| Herring, sea | | | | | 527,486 | 8,922 | | | 209,520 | 2,545 |
| Hickory shad | | | | | 2,308 | 23 | | | | |
| King whiting or "kingfish" | | | | | 1,440 | 53 | | | | |
| Mackerel | | | | | 696,810 | 21,621 | | | 22,479 | 516 |
| Menhaden | | | | | 1,000,200 | 2,502 | | | 3,310 | 33 |
| Mullet | | | | | 2,811 | 141 | | | 6,000 | 180 |
| Pollock | | | | | 17,092 | 449 | | | 150 | 2 |
| Scup or porgy | | | | | 1,948,671 | 35,737 | | | 70,760 | 1,696 |
| Sea bass | | | | | 45,505 | 2,345 | | | 2,760 | 198 |
| Sea robin | | | | | 64,955 | 653 | | | 8,830 | 44 |
| Shad | 2,476 | 299 | | | 7,489 | 729 | 650 | 95 | 146 | 15 |
| Sharks | | | | | 600 | 18 | | | 1,347 | 10 |
| Skates | | | | | 139,620 | 1,110 | | | 36,790 | 218 |
| Skipper or "billfish" | | | | | 730 | 14 | | | | |
| Smelt | | | | | | | | | 4,250 | 425 |
| Spot | | | | | | | | | 45 | 1 |
| Squeteagues or "sea trout", gray | | | | | 14,041 | 920 | | | 49,260 | 3,063 |
| Striped bass | | | | | 30,087 | 1,501 | | | 9,165 | 917 |
| Sturgeon | | | | | 449 | 37 | | | | |
| Swordfish | 775 | 118 | | | 234,268 | 21,461 | | | 24,667 | 2,220 |
| Tautog | 5,850 | 161 | 6,300 | 189 | 140,163 | 3,608 | | | 114,810 | 3,444 |
| Thimble-eyed mackerel | | | | | 76,957 | 1,516 | | | 760 | 15 |
| Tomcod | | | | | 200 | 1 | | | | |
| Tuna or "horse mackerel" | | | | | 44,577 | 1,484 | | | | |
| White perch | | | | | 411 | 19 | 450 | 23 | | |
| Whiting | | | | | 696,416 | 7,354 | | | 28,090 | 272 |
| Crabs: | | | | | | | | | | |
| Hard | 10,478 | 189 | | | 10,000 | 180 | | | | |
| King | | | | | 6,574 | 18 | | | | |
| Lobsters | 31,127 | 4,981 | 33,011 | 5,942 | 459,246 | 70,238 | 2,484 | 397 | 182,227 | 32,375 |
| Clams: | | | | | | | | | | |
| Hard, public | 102,399 | 12,513 | 636,844 | 72,213 | 115,126 | 13,083 | 112,661 | 12,823 | 102,861 | 14,415 |
| Hard, private | 60,170 | 8,206 | 28,875 | 4,044 | | | | | | |
| Soft, public | 14,720 | 1,653 | 1,984 | 186 | 1,600 | 200 | 7,440 | 930 | 6,640 | 1,245 |
| Mussels, sea | 650 | 38 | | | | | | | | |
| Oysters: | | | | | | | | | | |
| Market, public, spring | | | | | | | | | 26,880 | 3,840 |
| Market, public, fall | | | 6,903 | 1,004 | | | | | 31,541 | 4,803 |
| Market, private, spring | 864,738 | 129,259 | 60,000 | 7,500 | | | 429,268 | 50,485 | 171,388 | 24,504 |
| Market, private, fall | 695,177 | 104,936 | 60,000 | 7,500 | | | 420,257 | 50,485 | 342,621 | 48,976 |
| Periwinkles and cockles | 28,620 | 1,840 | | | 40,772 | 1,692 | | | 40,050 | 1,669 |
| Scallops, bay | | | 47,250 | 9,844 | 10,920 | 2,274 | 630 | 106 | 4,320 | 720 |
| Squid | | | | | 338,161 | 8,044 | | | 163,690 | 8,574 |
| Total | 1,832,874 | 264,489 | 892,907 | 109,340 | 10,850,253 | 325,926 | 1,020,212 | 119,328 | 2,770,101 | 182,181 |

U. S. BUREAU OF FISHERIES

Fisheries of Rhode Island, 1933—Continued

SEED OYSTER FISHERY: BY GEAR

| Item | Tongs | | By hand | | Total, exclusive of duplication | |
|---------------------------------------|-----------------------|-----------------------|-----------------------|-----------------------|---------------------------------|-----------------------|
| OPERATING UNITS | | | | | | |
| Fishermen, on boats and shore: | <i>Number</i> | | <i>Number</i> | | <i>Number</i> | |
| Regular..... | 1 | | 4 | | 5 | |
| Casual..... | 1 | | 6 | | 6 | |
| Total..... | 2 | | 10 | | 11 | |
| Boats, other than motor..... | 2 | | | | 2 | |
| Apparatus, number..... | 2 | | | | | |
| CATCH | | | | | | |
| Oysters, seed, public, spring..... | <i>Bushels</i> 169 | <i>Value</i> \$135 | <i>Bushels</i> 826 | <i>Value</i> \$364 | <i>Bushels</i> 995 | <i>Value</i> \$499 |

SEED OYSTER FISHERY: BY COUNTIES

| Item | Bristol | | Kent | |
|---------------------------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| OPERATING UNITS | | | | |
| Fishermen, on boats and shore: | <i>Number</i> | | <i>Number</i> | |
| Regular..... | 3 | | 2 | |
| Casual..... | 1 | | 5 | |
| Total..... | 4 | | 7 | |
| Boats, other than motor..... | | | 2 | |
| Apparatus, tongs..... | | | 2 | |
| CATCH | | | | |
| Oysters, seed, public, spring..... | <i>Bushels</i> 629 | <i>Value</i> \$206 | <i>Bushels</i> 366 | <i>Value</i> \$263 |

NOTE.—Of the total number of persons fishing for seed oysters, 9 are duplicated among those fishing for market oysters or other species. The 2 tongs also are duplicated.

CONNECTICUT

Fisheries of Connecticut, 1933

OPERATING UNITS: BY GEAR

| Item | Haul seines | Gill nets | | Lines | | Pound nets | Fyke nets | Dip nets | Otter trawls |
|-----------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | | Anchor | Drift | Hand | Trawl | | | | |
| Fishermen: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| On vessels..... | | | | 49 | 15 | | | | 112 |
| On boats and shore: | | | | | | | | | |
| Regular..... | | 2 | | 140 | 8 | 11 | | 10 | 93 |
| Casual..... | 66 | | 70 | 30 | | 2 | | 109 | |
| Total..... | 66 | 2 | 70 | 219 | 23 | 11 | 2 | 119 | 205 |
| Vessels: | | | | | | | | | |
| Motor..... | | | | 7 | 1 | | | | 41 |
| Net tonnage..... | | | | 161 | 47 | | | | 515 |
| Boats: | | | | | | | | | |
| Motor..... | 1 | 2 | 31 | 98 | 4 | 5 | | | 48 |
| Other..... | 15 | | 16 | | | 5 | 1 | 86 | |
| Accessory boats: | | | | | 24 | | | | |
| Apparatus: | | | | | | | | | |
| Number..... | 12 | 2 | 43 | 405 | 800 | 9 | 4 | 119 | 89 |
| Length, yards..... | 1,668 | | | | | | | | |
| Square yards..... | | 1,680 | 12,360 | | | | | | |
| Yards at mouth..... | | | | | | | | | 2,373 |
| Hooks, baits or snoods..... | | | | 443 | 35,680 | | | | |

Fisheries of Connecticut, 1933—Continued

OPERATING UNITS: BY GEAR—Continued

| Item | Pots | | Har- poons | Spears | Dredges | | Tongs | Rakes | Hoes | Total, exclu- sive of dupli- cation |
|-------------------------------|-----------|--------------|---------------|-----------|------------|--------------|------------|-----------|-----------|--|
| | Eel | Lob- ster | | | Oyster | Scal- lop | | | | |
| | Number | Number | Number | Number | Number | Number | Number | Number | Number | |
| Fishermen: | | | | | | | | | | |
| On vessels..... | | 9 | 73 | | 115 | 8 | | | | 307 |
| On boats and shore: | | | | | | | | | | |
| Regular..... | 21 | 205 | 3 | 9 | 2 | | 16 | | 4 | 300 |
| Casual..... | 12 | 51 | 6 | 33 | | | 102 | 71 | 34 | 494 |
| Total..... | 33 | 265 | 82 | 42 | 117 | 8 | 118 | 71 | 38 | 1,161 |
| Vessels: | | | | | | | | | | |
| Steam: | | | | | 3 | | | | | 3 |
| Net tonnage..... | | | | | 581 | | | | | 581 |
| Motor: | | 4 | 13 | | 16 | 1 | | | | 67 |
| Net tonnage..... | | 30 | 189 | | 401 | 55 | | | | 1,056 |
| Total vessels..... | | 4 | 13 | | 19 | 1 | | | | 70 |
| Total net tonnage..... | | 30 | 189 | | 932 | 55 | | | | 1,637 |
| Boats: | | | | | | | | | | |
| Motor..... | 11 | 201 | 3 | | 1 | | 2 | | | 294 |
| Other..... | 22 | 35 | | 28 | | | 75 | 33 | 25 | 276 |
| Accessory boats..... | | | 16 | | | | | | | 39 |
| Apparatus: | | | | | | | | | | |
| Number..... | 1,289 | 17,212 | 16 | 42 | 43 | 2 | 118 | 71 | 38 | |
| Yards at mouth..... | | | | | 65 | 7 | | | | |

CATCH: BY GEAR

| Species | Haul seines | | Gill nets | | | | Lines | | | |
|--|---------------|--------------|--------------|------------|---------------|--------------|----------------|---------------|----------------|---------------|
| | | | Anchor | | Drift | | Hand | | Trawl | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | | | | | 162 | \$2 | | | | |
| Bluefish..... | | | 1,500 | \$150 | | | 292,351 | \$21,695 | | |
| Cod..... | | | | | | | 23,200 | 686 | 80,200 | \$2,559 |
| Croaker..... | | | | | | | 3,000 | 90 | | |
| Eels, conger..... | | | | | | | | | 1,000 | 100 |
| Haddock..... | | | | | | | | | 3,150 | 142 |
| Hake..... | | | | | | | | | 500 | 10 |
| Halibut..... | | | | | | | | | 6,000 | 300 |
| Scup or porky..... | | | | | | | 16,000 | 800 | | |
| Sea bass..... | | | | | | | 23,695 | 1,620 | | |
| Shad..... | 43,478 | \$3,166 | | | 88,732 | 7,669 | | | | |
| Smelt..... | 2,240 | 224 | | | | | | | | |
| Squeteagues or "sea trout", gray..... | | | 750 | 75 | | | 940 | 94 | | |
| Striped bass..... | | | | | | | 750 | 75 | | |
| Tautog..... | | | | | | | 33,933 | 1,465 | | |
| Tilefish..... | | | | | | | | | 207,000 | 10,330 |
| Total..... | 45,718 | 3,390 | 2,250 | 225 | 88,894 | 7,671 | 393,869 | 26,555 | 289,850 | 13,441 |

| Species | Pound nets | | Fyke nets | | Dip nets | | Otter trawls | |
|---------------------------------|------------|-------|-----------|-------|----------|-------|--------------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | 15,000 | \$150 | | | | | | |
| Bluefish..... | 2,490 | 204 | | | | | 113 | \$9 |
| Butterfish..... | 12,425 | 586 | | | | | 3,516 | 178 |
| Cod..... | | | | | | | 44,525 | 1,202 |
| Croaker..... | | | | | | | 2,738 | 46 |
| Drum, black..... | | | | | | | 100 | 1 |
| Eels: | | | | | | | | |
| Common..... | 450 | 34 | 400 | \$40 | | | 190 | 4 |
| Conger..... | | | | | | | 400 | 20 |
| Flounders..... | 3,850 | 205 | | | | | 6,172,188 | 175,753 |
| Haddock..... | | | | | | | 32,300 | 876 |
| Hake..... | | | | | | | 581 | 10 |
| Halibut..... | | | | | | | 63 | 10 |
| King whiting or "kingfish"..... | | | | | | | 451 | 12 |

Fisheries of Connecticut, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Pound nets | | Fyke nets | | Dip nets | | Otter trawls | |
|---------------------------------------|------------|-------|-----------|-------|----------|--------|--------------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Mackerel..... | | | | | | | 870 | \$27 |
| Menhaden..... | 25,250 | \$252 | | | | | | |
| Scup or porgy..... | | | | | | | 25,987 | 859 |
| Sea bass..... | | | | | | | 3,920 | 223 |
| Sea robin..... | | | | | | | 3,250 | 33 |
| Shad..... | | | | | 1,070 | \$134 | | |
| Skates..... | | | | | | | 18,550 | 185 |
| Spot..... | | | | | | | 50 | 1 |
| Squeteagues or "sea trout", gray..... | 17,500 | 1,550 | | | | | 320 | 17 |
| Striped bass..... | 1,500 | 150 | | | | | | |
| Sturgeon..... | | | | | | | 44 | 5 |
| Suckers..... | | | 1,000 | \$60 | | | | |
| Tautog..... | 4,600 | 222 | | | | | 2,425 | 81 |
| Whiting..... | | | | | | | 18,775 | 547 |
| Wolfish..... | | | | | | | 4,100 | 41 |
| Crabs, hard..... | | | | | 400 | 25 | | |
| Lobsters..... | | | | | | | 163 | 20 |
| Scallops: | | | | | | | | |
| Bay..... | | | | | 57,500 | 10,000 | | |
| Sea..... | | | | | | | 128 | 26 |
| Squid..... | 2,425 | 37 | | | | | 480 | 13 |
| Total..... | 85,490 | 3,380 | 1,400 | 100 | 58,970 | 10,159 | 6,334,225 | 180,209 |

| Species | Pots | | | | Harpoons | | Spears | |
|-------------------|--------|---------|---------|--------|----------|----------|--------|---------|
| | Eel | | Lobster | | | | | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Eels, common..... | 25,469 | \$2,191 | | | | | 20,885 | \$1,695 |
| Swordfish..... | | | | | 154,235 | \$18,650 | | |
| Tautog..... | | | 2,140 | \$107 | | | | |
| Lobsters..... | | | 336,637 | 64,115 | | | | |
| Total..... | 25,469 | 2,191 | 338,777 | 64,222 | 154,235 | 18,650 | 20,885 | 1,695 |

| Species | Dredges | | | | Tongs | | Rakes | | Hoes | |
|------------------------------|-----------|----------|---------|---------|--------|----------|--------|---------|--------|---------|
| | Oyster | | Scallop | | | | | | | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Clams: | | | | | | | | | | |
| Hard, public..... | | | | | 51,910 | \$10,796 | 41,645 | \$8,329 | | |
| Soft, public..... | | | | | | | | | 13,683 | \$1,953 |
| Oysters: | | | | | | | | | | |
| Market, public, spring..... | | | | | 10,150 | 1,250 | | | | |
| Market, public, fall..... | | | | | 11,550 | 4,450 | | | | |
| Market, private, spring..... | 605,447 | \$80,083 | | | 8,100 | 1,300 | | | | |
| Market, private, fall..... | 1,208,511 | 167,811 | | | 15,150 | 2,450 | | | | |
| Scallops, sea..... | | | 55,800 | \$6,200 | | | | | | |
| Total..... | 1,813,958 | 247,894 | 55,800 | 6,200 | 96,860 | 17,246 | 41,645 | 8,329 | 21,733 | 3,563 |

Fisheries of Connecticut, 1933—Continued

OPERATING UNITS: BY COUNTIES

| Item | Fairfield | Hartford | Middlesex | New Haven | New London |
|--------------------------------|---------------|---------------|---------------|---------------|---------------|
| Fishermen: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| On vessels..... | 99 | | | 60 | 148 |
| On boats and shore: | | | | | |
| Regular..... | 56 | | 76 | 59 | 160 |
| Casual..... | 163 | 61 | 113 | 20 | 137 |
| Total | 318 | 61 | 189 | 139 | 454 |
| Vessels: | | | | | |
| Steam..... | 2 | | | 1 | |
| Net tonnage..... | 187 | | | 394 | |
| Motor..... | 17 | | | 13 | 37 |
| Net tonnage..... | 421 | | | 172 | 463 |
| Total vessels | 19 | | | 14 | 37 |
| Total net tonnage | 608 | | | 566 | 463 |
| Boats: | | | | | |
| Motor..... | 42 | 2 | 80 | 42 | 128 |
| Other..... | 95 | 16 | 48 | 19 | 98 |
| Accessory boats | | | | | 39 |
| Apparatus: | | | | | |
| Haul seines..... | 3 | 8 | 1 | | |
| Length, yards..... | 300 | 900 | 468 | | |
| Gill nets: | | | | | |
| Anchor..... | | | | | 2 |
| Square yards..... | | | | | 1,680 |
| Drift..... | | 2 | 30 | | 11 |
| Square yards..... | | 560 | 8,680 | | 3,120 |
| Lines: | | | | | |
| Hand..... | 72 | | 116 | 65 | 152 |
| Hooks..... | 100 | | 116 | 65 | 162 |
| Trawl..... | | | | | 800 |
| Hooks..... | | | | | 35,650 |
| Total | | | | | 9 |
| Pound nets..... | | | | | |
| Fyke nets..... | | 4 | | | 92 |
| Dip nets..... | | 7 | | | 62 |
| Otter trawls..... | 13 | | 20 | 10 | 62 |
| Yards at mouth..... | 330 | | 100 | 237 | 1,706 |
| Pots: | | | | | |
| Eel..... | 115 | | 264 | 338 | 572 |
| Lobster..... | 5,007 | | 2,843 | 2,814 | 6,548 |
| Harpoons..... | 1 | | | 1 | 14 |
| Spears..... | 20 | | | 5 | 17 |
| Dredges: | | | | | |
| Oyster..... | 27 | | | 16 | |
| Yards at mouth..... | 40 | | | 25 | |
| Scallop..... | 2 | | | | |
| Yards at mouth..... | 7 | | | | |
| Tongs..... | 82 | | 32 | | 4 |
| Rakes..... | 71 | | | | |
| Hoes..... | | | 10 | 10 | 18 |

CATCH: BY COUNTIES

| Species | Fairfield | | Hartford | | Middlesex | | New Haven | | New London | |
|---------------------------------|-----------|-------|----------|-------|-----------|-------|-----------|---------|------------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | | | | | 162 | \$2 | | | 15,000 | \$150 |
| Bluefish..... | 12,759 | \$993 | | | 36,067 | 3,138 | 18,949 | \$1,597 | 228,689 | 16,330 |
| Butterfish..... | | | | | | | | | 15,941 | 764 |
| Cod..... | 2,000 | 60 | | | 540 | 16 | | | 145,385 | 4,381 |
| Croaker..... | 3,000 | 90 | | | | | 222 | 2 | 2,616 | 44 |
| Drum, black..... | | | | | | | | | 100 | 1 |
| Eels: | | | | | | | | | | |
| Common..... | 18,333 | 1,498 | 400 | \$40 | 3,240 | 324 | 3,810 | 381 | 21,611 | 1,721 |
| Conger..... | | | | | | | | | 1,400 | 120 |
| Flounders..... | 196,578 | 5,441 | | | 28,540 | 704 | 112,945 | 3,600 | 5,437,975 | 166,163 |
| Haddock..... | | | | | | | | | 35,450 | 1,018 |
| Hake..... | 275 | 5 | | | | | | | 808 | 15 |
| Halibut..... | | | | | | | | | 6,063 | 310 |
| King whiting or "kingfish"..... | | | | | | | | | 451 | 12 |
| Mackerel..... | | | | | | | | | 870 | 27 |
| Menhaden..... | | | | | | | | | 25,250 | 252 |
| Scup or porgy..... | 32,220 | 1,325 | | | | | | | 9,767 | 330 |
| Sea bass..... | 11,150 | 762 | | | 1,215 | 74 | 220 | 13 | 15,030 | 994 |
| Sea robin..... | | | | | | | | | 3,250 | 33 |

Fisheries of Connecticut, 1933—Continued

CATCH: BY COUNTIES—Continued

| Species | Fairfield | | Hartford | | Middlesex | | New Haven | | New London | |
|---------------------------------------|-------------|----------|----------|----------|-----------|----------|-------------|----------|-------------|----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Shad..... | | | 44, 148 | \$3, 275 | 67, 502 | \$5, 531 | | | 21, 630 | \$2, 183 |
| Skates..... | | | | | | | | | 18, 550 | 185 |
| Smelt..... | 2, 240 | \$224 | | | | | | | | |
| Spot..... | 50 | 1 | | | | | | | | |
| Squeteagues or "sea trout", gray..... | 58 | 4 | | | 940 | 94 | 10 | \$1 | 18, 502 | 1, 637 |
| Striped bass..... | | | | | | | | | 2, 250 | 225 |
| Sturgeon..... | | | | | | | 27 | 3 | 17 | 2 |
| Suckers..... | | | 1, 000 | 60 | | | | | | |
| Swordfish..... | 20, 500 | 1, 968 | | | | | 2, 280 | 130 | 131, 475 | 14, 552 |
| Tautog..... | 8, 231 | 453 | | | 13, 274 | 656 | 1, 191 | 72 | 20, 402 | 714 |
| Thiefish..... | | | | | | | | | 207, 000 | 10, 330 |
| Whiting..... | | | | | | | | | 16, 775 | 547 |
| Wolfish..... | | | | | | | | | 4, 100 | 41 |
| Crabs, hard..... | | | | | | | | | 400 | 25 |
| Lobsters..... | 53, 211 | 10, 635 | | | 46, 011 | 9, 202 | 62, 565 | 12, 513 | 175, 013 | 31, 785 |
| Clams: | | | | | | | | | | |
| Hard, public..... | 88, 455 | 17, 691 | | | 5, 100 | 1, 434 | 8, 050 | 1, 610 | | |
| Soft, public..... | | | | | 8, 400 | 1, 200 | 1, 500 | 150 | 3, 783 | 603 |
| Oysters: | | | | | | | | | | |
| Market, public, spring..... | | | | | 10, 150 | 1, 250 | | | | |
| Market, public, fall..... | | | | | 11, 550 | 1, 450 | | | | |
| Market, private, spring..... | 350, 972 | 47, 633 | | | 2, 100 | 300 | 248, 475 | 32, 450 | 6, 000 | 1, 000 |
| Market, private, fall..... | 569, 317 | 79, 394 | | | 3, 150 | 450 | 639, 194 | 88, 417 | 12, 000 | 2, 000 |
| Scallops: | | | | | | | | | | |
| Bay..... | | | | | 11, 500 | 2, 000 | | | 46, 000 | 8, 000 |
| Sea..... | 55, 800 | 6, 200 | | | | | | | 126 | 26 |
| Squid..... | 415 | 12 | | | | | | | 2, 490 | 38 |
| Total..... | 1, 431, 564 | 174, 393 | 45, 548 | 3, 375 | 249, 431 | 27, 825 | 1, 099, 418 | 140, 999 | 7, 052, 087 | 266, 538 |

SEED OYSTER FISHERY: BY GEAR

| Item | Dredges, oyster | | Tongs | | Rakes | | Total, exclusive of duplication | |
|---------------------------------|-----------------|--------------|----------------|--------------|----------------|--------------|---------------------------------|--------------|
| OPERATING UNITS | | | | | | | | |
| Fishermen: | <i>Number</i> | | <i>Number</i> | | <i>Number</i> | | <i>Number</i> | |
| On vessels..... | 93 | | 94 | | 46 | | 93 | |
| On boats and shore, casual..... | 14 | | | | | | 154 | |
| Total..... | 107 | | 94 | | 46 | | 247 | |
| Vessels: | | | | | | | | |
| Steam..... | 3 | | | | | | 3 | |
| Net tonnage..... | 245 | | | | | | 245 | |
| Motor..... | 18 | | | | | | 18 | |
| Net tonnage..... | 271 | | | | | | 271 | |
| Sail..... | 2 | | | | | | 2 | |
| Net tonnage..... | 17 | | | | | | 17 | |
| Total vessels..... | 23 | | | | | | 23 | |
| Total net tonnage..... | 533 | | | | | | 533 | |
| Boats: | | | | | | | | |
| Motor..... | 7 | | | | | | 7 | |
| Other..... | | | 60 | | 28 | | 88 | |
| Apparatus: | | | | | | | | |
| Number..... | 101 | | 94 | | 46 | | | |
| Yards at mouth..... | 97 | | | | | | | |
| CATCH | | | | | | | | |
| Oysters: | <i>Bushels</i> | <i>Value</i> | <i>Bushels</i> | <i>Value</i> | <i>Bushels</i> | <i>Value</i> | <i>Bushels</i> | <i>Value</i> |
| Seed, public, spring..... | 23, 463 | \$9, 385 | 5, 200 | \$2, 080 | 1, 800 | \$720 | 30, 463 | \$12, 185 |
| Seed, public, fall..... | 23, 663 | 9, 465 | 16, 900 | 6, 760 | 2, 600 | 1, 040 | 43, 163 | 17, 265 |
| Seed, private, spring..... | 207, 185 | 69, 617 | | | | | 207, 185 | 69, 617 |
| Total..... | 254, 311 | 88, 467 | 22, 100 | 8, 840 | 4, 400 | 1, 760 | 280, 811 | 99, 067 |

Fisheries of Connecticut, 1933—Continued

SEED OYSTER FISHERY: BY COUNTIES

| Item | Fairfield | | New Haven | |
|---------------------------------|----------------|--------------|----------------|--------------|
| OPERATING UNITS | | | | |
| Fishermen: | <i>Number</i> | | <i>Number</i> | |
| On vessels..... | 51 | | 42 | |
| On boats and shore, casual..... | 150 | | 4 | |
| Total..... | 201 | | 46 | |
| Vessels: | | | | |
| Steam..... | 2 | | 1 | |
| Net tonnage..... | 140 | | 105 | |
| Motor..... | 10 | | 8 | |
| Net tonnage..... | 113 | | 158 | |
| Sail..... | 2 | | | |
| Net tonnage..... | 17 | | | |
| Total vessels..... | 14 | | 9 | |
| Total net tonnage..... | 270 | | 263 | |
| Boats: | | | | |
| Motor..... | 5 | | 2 | |
| Other..... | 88 | | | |
| Apparatus: | | | | |
| Dredges..... | 75 | | 26 | |
| Yards at mouth..... | 64 | | 33 | |
| Tongs..... | 94 | | | |
| Rakes..... | 46 | | | |
| CATCH | | | | |
| Oysters: | <i>Bushels</i> | <i>Value</i> | <i>Bushels</i> | <i>Value</i> |
| Seed, public, spring..... | 26,863 | \$10,745 | 3,600 | \$1,440 |
| Seed, public, fall..... | 39,563 | 15,825 | 3,600 | 1,440 |
| Seed, private, spring..... | 74,817 | 41,099 | 132,368 | 26,518 |
| Total..... | 141,243 | 67,669 | 139,568 | 31,398 |

NOTE.—Of the total number of persons fishing for seed oysters, 126 are duplicated among those fishing for market oysters or other species. Similarly the following craft and gear are duplicated: 78 boats other than motor, 80 tongs, and 46 rakes.

VESSEL FISHERIES AT PRINCIPAL NEW ENGLAND PORTS

Due to the importance of the ports of Boston and Gloucester, Mass., and Portland, Maine, as landing points for fishery products, detailed monthly statistics are collected for these landings which are published in the following section. These landings are included in the catch by States appearing elsewhere in this document, but also are presented here for their value in detailed form.

ECONOMIC ASPECT

The landings of fishery products at the three principal New England ports (Boston and Gloucester, Mass., and Portland, Maine) by vessels of 5 net tons capacity or more, during 1933 amounted to 267,157,218 pounds as landed, valued at \$6,850,901. This is an increase of 6 percent in the quantity of the catch as compared with 1932, and an increase of 13 percent in the value of the catch. Of the total landings 99 percent consisted of fresh fish and 1 percent salted fish. The landings at Boston accounted for 232,583,049 pounds, valued at \$6,093,604, or 87 percent of the total landings. The landings at Gloucester amounted to 21,736,997 pounds, valued at \$441,937, or 8 percent of the total. Landings at Portland amounted to 12,837,172 pounds, valued at \$315,360, or 5 percent of the total landings.

Among the landings of fresh fish, haddock far outranked other species in volume landed, the amount of all sizes in 1933 being 129,847,257 pounds, or 49 percent of the total fresh fish.

Landings by fishing vessels at the 3 principal New England ports, 1933

BOSTON: BY MONTHS

| Species | January | | February | | March | | April | | May | | June | | July | |
|-----------------------|------------|----------|------------|----------|------------|-----------|------------|----------|------------|----------|------------|----------|------------|----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Cod, fresh: | | | | | | | | | | | | | | |
| Large..... | 2,585,139 | \$55,765 | 4,079,853 | \$79,391 | 4,638,620 | \$123,961 | 2,907,927 | \$48,245 | 1,873,315 | \$31,875 | 1,449,898 | \$39,203 | 1,759,968 | \$41,125 |
| Market..... | 1,700,490 | 38,482 | 1,782,396 | 43,506 | 2,247,765 | 57,831 | 3,015,715 | 37,841 | 2,435,900 | 31,572 | 1,952,585 | 37,552 | 4,369,380 | 75,054 |
| Scrod..... | 27,270 | 483 | 12,250 | 192 | 4,115 | 94 | 900 | 9 | 2,100 | 22 | 11,050 | 172 | 23,250 | 383 |
| Cod, salted: | | | | | | | | | | | | | | |
| Large..... | | | | | | | | | | | | | | |
| Market..... | | | | | | | | | | | | | | |
| Haddock, fresh: | | | | | | | | | | | | | | |
| Large..... | 6,623,185 | 234,991 | 8,521,960 | 288,389 | 9,645,090 | 333,564 | 11,661,785 | 175,953 | 9,600,536 | 161,449 | 7,310,335 | 177,536 | 7,818,740 | 183,996 |
| Scrod..... | 1,561,565 | 38,608 | 1,966,970 | 49,726 | 1,751,360 | 48,940 | 1,999,220 | 20,050 | 1,710,785 | 19,910 | 1,364,580 | 24,194 | 1,642,450 | 27,792 |
| Hake, fresh: | | | | | | | | | | | | | | |
| Large..... | 757.65 | 15,680 | 298,540 | 8,633 | 122,430 | 4,780 | 181,640 | 3,985 | 173,925 | 2,359 | 237,575 | 4,132 | 123,110 | 1,688 |
| Small..... | 16,000 | 344 | 10,760 | 338 | 9,800 | 330 | 3,775 | 106 | 37,950 | 569 | 19,630 | 401 | 83,190 | 1,359 |
| Pollock, fresh..... | 1,216,725 | 16,835 | 484,325 | 10,233 | 305,715 | 7,718 | 895,735 | 5,649 | 674,695 | 5,049 | 256,270 | 3,339 | 376,990 | 4,902 |
| Cusk, fresh..... | 416,510 | 6,083 | 330,736 | 5,159 | 108,325 | 2,044 | 205,330 | 1,809 | 119,905 | 1,021 | 143,575 | 1,562 | 113,540 | 1,439 |
| Halibut, fresh..... | 38,476 | 7,481 | 119,622 | 15,143 | 209,685 | 21,403 | 180,140 | 24,278 | 270,191 | 27,357 | 207,918 | 17,551 | 269,122 | 25,869 |
| Mackerel, fresh..... | 1,155 | 93 | 120 | 1 | | | | | 3,632,163 | 56,165 | 2,169,695 | 67,426 | 2,353,943 | 52,084 |
| Flounders, fresh..... | 761,390 | 36,663 | 729,510 | 31,617 | 631,250 | 28,067 | 587,000 | 23,582 | 1,059,400 | 20,769 | 789,940 | 18,714 | 710,304 | 22,565 |
| Swordfish, fresh..... | | | | | | | | | | | | | | |
| Herring, fresh..... | 775 | 39 | 75 | 2 | | | | | | | | | | |
| Other, fresh..... | 135,895 | 5,674 | 119,487 | 3,368 | 223,703 | 5,826 | 336,323 | 4,818 | 550,349 | 6,354 | 299,632 | 3,527 | 165,385 | 2,751 |
| Total, fresh..... | 15,841,740 | 457,221 | 18,436,500 | 515,698 | 19,798,458 | 634,558 | 21,975,490 | 346,325 | 22,140,614 | 364,471 | 16,375,176 | 436,288 | 20,272,648 | 530,592 |
| Total, salted..... | | | | | | | | | 11,000 | 253 | 12,500 | 253 | 11,000 | 305 |
| Grand total..... | 15,841,740 | 457,221 | 18,436,500 | 515,698 | 19,798,458 | 634,558 | 21,975,490 | 346,325 | 22,151,614 | 364,724 | 16,387,676 | 436,541 | 20,283,648 | 530,897 |
| Landed in 1932: | | | | | | | | | | | | | | |
| Fresh..... | 14,621,426 | 472,864 | 18,685,804 | 660,364 | 19,220,853 | 592,393 | 19,552,627 | 320,842 | 19,773,187 | 389,166 | 17,046,277 | 418,749 | 18,353,142 | 388,315 |
| Salted..... | | | | | | | 11,000 | 165 | 10,000 | 300 | 24,000 | 741 | 1,150 | 25 |
| Total..... | 14,621,426 | 472,864 | 18,685,804 | 660,364 | 19,220,853 | 592,393 | 19,563,627 | 321,007 | 19,783,187 | 389,466 | 17,070,277 | 419,490 | 18,354,292 | 388,340 |

NOTE.—The weights of fresh and salted fish given in these statistics represent the fish as landed from the vessels, and the values are those received by the fishermen. Large cod are classified as those weighing over 10 pounds; market cod, 2½ to 10 pounds; and scrod cod, 1 to 2½ pounds. Large haddock are those weighing over 2½ pounds and scrod haddock 1 to 2½ pounds. Large hake are those weighing over 6 pounds and small hake, under 6 pounds. Only landings by vessels having a capacity of 5 net tons or greater are used in this tabulation.

| Species | August | | September | | October | | November | | December | | Total, 1933 | | 1932 | |
|-----------------------------|------------|----------|------------|----------|------------|----------|------------|----------|------------|----------|-------------|-----------|-------------|-----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Cod, fresh: | | | | | | | | | | | | | | |
| Large..... | 1,596,001 | \$43,864 | 1,267,174 | \$41,283 | 1,933,240 | \$56,712 | 1,961,566 | \$82,806 | 1,910,170 | \$57,404 | 27,961,973 | \$681,634 | 21,445,174 | \$586,706 |
| Market..... | 4,103,189 | 69,330 | 2,292,725 | 53,983 | 3,028,470 | 70,792 | 2,909,520 | 81,077 | 3,207,700 | 77,077 | 33,025,829 | 674,097 | 26,917,646 | 454,035 |
| Scrod..... | 4,150 | 51 | 29,680 | 546 | 52,410 | 947 | 11,870 | 174 | 16,900 | 320 | 195,945 | 3,393 | 101,015 | 1,737 |
| Cod, salted: | | | | | | | | | | | | | | |
| Large..... | | | | | 30,500 | 1,028 | | | | | 50,500 | 1,540 | 47,815 | 1,518 |
| Market..... | | | | | 1,000 | 30 | | | | | 15,500 | 329 | 32,360 | 587 |
| Haddock, fresh: | | | | | | | | | | | | | | |
| Large..... | 7,487,140 | 201,163 | 6,516,290 | 205,917 | 7,114,815 | 234,421 | 4,520,533 | 217,142 | 3,532,885 | 170,781 | 90,353,894 | 2,565,302 | 87,083,975 | 2,368,147 |
| Scrod..... | 2,506,795 | 44,422 | 6,141,240 | 123,543 | 5,582,620 | 95,954 | 2,987,710 | 71,558 | 3,316,565 | 75,016 | 32,531,860 | 639,713 | 27,649,754 | 399,352 |
| Haddock, salted, large..... | 400 | 4 | | | | | | | | | 400 | 4 | | |
| Hake, fresh: | | | | | | | | | | | | | | |
| Large..... | 264,590 | 4,025 | 478,948 | 8,984 | 517,255 | 10,009 | 846,435 | 20,723 | 682,000 | 15,436 | 4,683,613 | 100,434 | 5,647,501 | 111,176 |
| Small..... | 54,200 | 948 | 85,800 | 1,958 | 102,800 | 2,212 | 75,640 | 1,833 | 38,100 | 933 | 537,645 | 11,331 | 19,000 | 318 |
| Hake, salted, small..... | | | | | | | | | | | | | 4,000 | 80 |
| Pollock, fresh..... | 427,420 | 5,334 | 320,050 | 4,552 | 752,000 | 9,920 | 1,297,645 | 17,549 | 1,254,710 | 14,781 | 8,261,580 | 105,861 | 5,775,443 | 70,602 |
| Pollock, salted..... | 50 | | | | | | | | | | 50 | | 7,300 | 73 |
| Cusk, fresh..... | 55,720 | 674 | 245,180 | 4,177 | 272,400 | 4,580 | 207,850 | 4,292 | 279,505 | 5,450 | 2,498,576 | 38,290 | 2,492,595 | 38,187 |
| Halibut, fresh..... | 128,409 | 13,256 | 115,446 | 12,674 | 108,803 | 12,335 | 58,663 | 7,274 | 49,189 | 6,091 | 1,755,664 | 191,012 | 2,084,176 | 239,176 |
| Mackerel, fresh..... | 4,197,145 | 80,304 | 2,003,641 | 50,638 | 2,520,690 | 72,943 | 276,110 | 14,770 | 193,455 | 7,886 | 17,348,117 | 402,310 | 25,274,474 | 450,214 |
| Mackerel, salted..... | | | 9,800 | 147 | | | | | | | 9,800 | 147 | | |
| Flounders, fresh..... | 800,445 | 27,413 | 888,336 | 36,831 | 807,865 | 33,069 | 896,472 | 42,487 | 716,630 | 27,336 | 9,278,542 | 349,113 | 6,796,804 | 274,679 |
| Swordfish, fresh..... | 529,335 | 73,748 | 512,176 | 74,158 | 12,749 | 3,500 | 375 | 24 | | | 1,681,304 | 281,694 | 2,257,522 | 315,092 |
| Herring, fresh..... | 1,100 | 11 | | | 675 | 27 | 830 | 17 | 650 | 40 | 4,105 | 136 | 7,500 | 90 |
| Other, fresh..... | 100,496 | 3,648 | 62,509 | 2,686 | 191,930 | 3,751 | 116,495 | 3,047 | 85,958 | 1,814 | 2,388,152 | 47,264 | 1,974,925 | 45,156 |
| Total, fresh..... | 22,256,135 | 568,191 | 20,959,195 | 621,930 | 22,998,722 | 611,172 | 16,167,704 | 544,773 | 15,284,417 | 460,365 | 232,506,799 | 6,091,584 | 215,527,504 | 5,364,667 |
| Total, salted..... | 450 | 4 | 9,800 | 147 | 31,500 | 1,058 | | | | | 76,250 | 2,020 | 91,475 | 2,258 |
| Grand total..... | 22,256,585 | 568,195 | 20,968,995 | 622,077 | 23,030,222 | 612,230 | 16,167,704 | 544,773 | 15,284,417 | 460,365 | 232,583,049 | 6,093,604 | 215,618,979 | 5,366,925 |
| Landed in 1932: | | | | | | | | | | | | | | |
| Fresh..... | 20,830,781 | 430,596 | 20,573,161 | 420,414 | 20,461,179 | 396,038 | 14,176,025 | 452,927 | 12,233,042 | 421,999 | | | 215,527,504 | 5,364,667 |
| Salted..... | 17,180 | 185 | | | 15,000 | 510 | 9,145 | 252 | 4,000 | 80 | | | 91,475 | 2,258 |
| Total..... | 20,847,961 | 430,781 | 20,573,161 | 420,414 | 20,476,179 | 396,548 | 14,185,170 | 453,179 | 12,237,042 | 422,079 | | | 215,618,979 | 5,366,925 |

Landings by fishing vessels at the 3 principal New England ports, 1933—Continued

GLOUCESTER: BY MONTHS

| Species | January | | February | | March | | April | | May | | June | | July | |
|---------------------|---------|---------|----------|---------|---------|---------|-----------|----------|-----------|----------|-----------|----------|-----------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Cod, fresh: | | | | | | | | | | | | | | |
| Large | 156,450 | \$3,897 | 396,240 | \$7,546 | 408,540 | \$9,770 | 1,089,505 | \$19,940 | 1,723,450 | \$30,380 | 379,490 | \$10,053 | 306,700 | \$6,174 |
| Market | 15,695 | 250 | 21,555 | 391 | 21,930 | 445 | 240,680 | 2,457 | 365,030 | 3,903 | 100,475 | 1,006 | 128,960 | 1,274 |
| Scrod | 1,455 | 14 | 460 | 7 | 5,290 | 81 | 660 | 4 | 3,665 | 19 | | | 650 | 5 |
| Cod, salted: | | | | | | | | | | | | | | |
| Large | | | | | | | 2,750 | 70 | 193,904 | 4,764 | 40,330 | 1,009 | 688,712 | 20,661 |
| Market | | | | | | | 2,290 | 46 | 94,892 | 1,423 | 12,292 | 185 | 266,458 | 5,329 |
| Scrod | | | | | | | | | 15,585 | 156 | | | 80,835 | 607 |
| Haddock, fresh: | | | | | | | | | | | | | | |
| Large | 58,575 | 2,402 | 37,965 | 1,211 | 80,635 | 3,109 | 1,096,165 | 19,639 | 767,420 | 13,385 | 15,215 | 228 | 76,765 | 1,180 |
| Scrod | 2,350 | 34 | 1,430 | 24 | 1,360 | 26 | 43,650 | 332 | 6,490 | 41 | 55 | 1 | 11,720 | 89 |
| Hake, fresh, large | 55,145 | 1,108 | 19,285 | 474 | 11,190 | 374 | 10,690 | 180 | 12,115 | 81 | 1,740 | 13 | 7,609 | 50 |
| Hake, salted, large | | | | | | | | | | | 30 | 1 | 1,325 | 13 |
| Pollock, fresh | 122,790 | 1,792 | 58,895 | 1,479 | 25,100 | 815 | 48,530 | 272 | 23,675 | 119 | 11,110 | 95 | 5,075 | 27 |
| Pollock, salted | | | | | | | 90 | 1 | | | | | | |
| Cusk, fresh | 1,940 | 24 | 1,930 | 27 | 3,565 | 71 | 104,740 | 362 | 29,885 | 189 | 25,195 | 189 | 21,220 | 145 |
| Halibut, fresh | | | 39 | 8 | 24 | 2 | | | 1,315 | 169 | 2,807 | 217 | | |
| Halibut, salted | | | | | | | | | 130 | 7 | | | | |
| Mackerel, fresh | | | | | | | | | 462,220 | 3,439 | 103,515 | 2,327 | 261,100 | 4,637 |
| Mackerel, salted | | | | | | | | | | | | | 4,400 | 264 |
| Flounders, fresh | 59,265 | 2,046 | 75,680 | 2,420 | 50,180 | 2,026 | 47,350 | 1,107 | 15,900 | 240 | 21,860 | 491 | 19,020 | 438 |
| Herring, fresh | | | | | | | | | | | 11,500 | 91 | | |
| Other, fresh | 5,510 | 53 | 7,150 | 91 | 8,065 | 188 | 4,170 | 102 | 490 | 4 | 715 | 6 | 12,170 | 143 |
| Total, fresh | 479,175 | 11,620 | 620,629 | 13,678 | 615,879 | 16,907 | 2,686,140 | 44,395 | 3,412,625 | 51,969 | 673,677 | 14,717 | 848,989 | 14,162 |
| Total, salted | | | | | | | 5,130 | 117 | 304,511 | 6,350 | 52,652 | 1,195 | 1,041,730 | 26,874 |
| Grand total | 479,175 | 11,620 | 620,629 | 13,678 | 615,879 | 16,907 | 2,691,270 | 44,512 | 3,717,136 | 58,319 | 726,329 | 15,912 | 1,890,719 | 41,036 |
| Landed in 1932: | | | | | | | | | | | | | | |
| Fresh | 356,379 | 13,520 | 457,680 | 21,320 | 773,065 | 22,126 | 2,126,060 | 37,088 | 2,015,490 | 31,081 | 1,546,305 | 25,391 | 2,550,828 | 29,019 |
| Salted | | | | | | | | | 255,480 | 8,312 | 118,070 | 3,429 | 303,304 | 8,454 |
| Total | 356,379 | 13,520 | 457,680 | 21,320 | 773,065 | 22,126 | 2,126,060 | 37,088 | 2,270,970 | 39,393 | 1,664,375 | 28,820 | 2,854,132 | 37,473 |

| Species | August | | September | | October | | November | | December | | Total, 1933 | | 1932 | |
|------------------------|-----------|---------|-----------|----------|-----------|----------|-----------|---------|-----------|---------|-------------|-----------|------------|-----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Cod, fresh: | | | | | | | | | | | | | | |
| Large | 143,905 | \$2,390 | 291,835 | \$11,338 | 521,575 | \$17,739 | 65,100 | \$2,161 | 51,770 | \$1,906 | 5,534,610 | \$123,294 | 4,883,161 | \$126,270 |
| Market | 113,008 | 1,131 | 17,585 | 186 | 181,225 | 4,313 | 55,185 | 1,177 | 44,165 | 751 | 1,304,473 | 17,284 | 1,813,670 | 18,113 |
| Scrod | | | 200 | 2 | 3,150 | 60 | 205 | 3 | 475 | 6 | 16,210 | 201 | 3,120 | 24 |
| Cod, salted: | | | | | | | | | | | | | | |
| Large | 349,857 | 10,374 | 110,439 | 3,720 | 147,436 | 5,777 | | | | | 1,533,428 | 46,375 | 690,265 | 20,242 |
| Market | 263,296 | 5,266 | 91,848 | 2,266 | 162,623 | 4,866 | | | | | 893,699 | 19,381 | 415,559 | 9,805 |
| Scrod | 98,041 | 1,225 | 18,859 | 283 | 22,357 | 447 | | | | | 235,677 | 2,718 | 121,965 | 1,347 |
| Haddock, fresh: | | | | | | | | | | | | | | |
| Large | 103,325 | 1,668 | 22,615 | 894 | 115,909 | 4,630 | 77,790 | 2,911 | 13,295 | 695 | 2,465,674 | 51,952 | 2,212,557 | 46,726 |
| Scrod | 1,715 | 16 | 5,000 | 68 | 11,920 | 266 | 750 | 5 | 3,395 | 35 | 89,805 | 937 | 218,085 | 1,752 |
| Haddock, salted, scrod | | | | | | | | | | | | | | |
| Large | 14,050 | 127 | 45,815 | 971 | 199,708 | 3,528 | 56,185 | 1,344 | 42,510 | 898 | 476,042 | 9,148 | 1,029,151 | 9,311 |
| Small | | | | | 9,170 | 127 | 2,635 | 77 | | | 11,805 | 204 | 1,045 | 14 |
| Hake, salted, large | | | | | | | | | | | | | | |
| Pollock, fresh | 27,945 | 267 | 356,205 | 6,027 | 739,857 | 9,990 | 1,320,055 | 18,289 | 658,875 | 8,331 | 3,398,112 | 47,553 | 1,174,950 | 9,644 |
| Pollock, salted | | | | | | | | | | | 90 | 1 | 1,295 | 13 |
| Cusk, fresh | 68,085 | 450 | 11,995 | 80 | 4,580 | 62 | 2,390 | 42 | 3,340 | 57 | 278,865 | 1,698 | 236,857 | 1,702 |
| Cusk, salted | 944 | 12 | 440 | 7 | | | | | | | 1,384 | 19 | 4,000 | 57 |
| Halibut, fresh | | | | | | | 95 | 14 | | | 4,280 | 410 | 162,320 | 11,787 |
| Halibut, salted | | | | | | | | | | | 190 | 7 | 840 | 59 |
| Mackerel, fresh | 745,500 | 11,535 | 1,071,156 | 16,498 | 868,627 | 20,636 | 340,001 | 17,405 | 128,335 | 4,812 | 3,980,544 | 81,289 | 11,031,008 | 138,343 |
| Mackerel, salted | | | 40,000 | 800 | 11,600 | 406 | | | | | 56,000 | 1,470 | 23,225 | 696 |
| Flounders, fresh | 8,360 | 200 | 12,855 | 583 | 12,540 | 531 | 15,415 | 719 | 38,485 | 1,793 | 376,910 | 12,594 | 415,995 | 14,980 |
| Swordfish, fresh | | | 771 | 110 | | | | | | | 771 | 110 | 7,032 | 7,899 |
| Herring, fresh | | | | | | | | | | | 11,500 | 91 | 21,000 | 210 |
| Herring, salted | | | | | | | | | | | 703,494 | 21,290 | 703,494 | 21,290 |
| Other, fresh | 207,200 | 2,051 | 94,700 | 750 | 7,412 | 220 | 6,112 | 223 | | | 5,995 | 54 | 359,689 | 3,885 |
| Total, fresh | 1,433,183 | 19,835 | 1,930,762 | 37,507 | 2,675,673 | 62,102 | 1,941,918 | 44,370 | 990,640 | 19,388 | 18,309,290 | 350,650 | 23,444,319 | 381,713 |
| Total, salted | 712,138 | 16,877 | 261,586 | 7,076 | 344,018 | 11,406 | | | 705,944 | 21,302 | 3,427,707 | 91,287 | 1,883,894 | 52,363 |
| Grand total | 2,145,321 | 36,712 | 2,192,348 | 44,583 | 3,019,689 | 73,598 | 1,941,918 | 44,370 | 1,696,584 | 40,690 | 21,736,997 | 441,937 | 25,328,213 | 434,076 |
| Landed in 1932: | | | | | | | | | | | | | | |
| Fresh | 3,467,071 | 39,047 | 3,984,238 | 56,620 | 4,161,740 | 44,647 | 1,190,792 | 34,613 | 814,671 | 27,241 | | | 23,444,319 | 381,713 |
| Salted | 97,925 | 2,958 | 332,560 | 7,321 | 255,655 | 6,814 | | | 520,900 | 15,075 | | | 1,883,894 | 52,363 |
| Total | 3,564,996 | 42,005 | 4,316,798 | 63,941 | 4,417,395 | 51,461 | 1,190,792 | 34,613 | 1,335,571 | 42,316 | | | 25,328,213 | 434,076 |

137070-36-7

Landings by fishing vessels at the 3 principal New England ports, 1933—Continued

PORTLAND: BY MONTHS

| Species | January | | February | | March | | April | | May | | June | | July | |
|--------------------------|---------|---------|----------|---------|---------|---------|-----------|---------|-----------|---------|-----------|---------|-----------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Cod, fresh: | | | | | | | | | | | | | | |
| Large..... | 72,965 | \$1,855 | 70,442 | \$1,746 | 156,496 | \$3,976 | 233,387 | \$2,954 | 237,627 | \$3,143 | 339,113 | \$8,439 | 386,421 | \$9,271 |
| Market..... | 37,304 | 570 | 38,633 | 740 | 71,730 | 1,369 | 113,710 | 1,139 | 90,475 | 914 | 80,498 | 1,321 | 124,741 | 2,421 |
| Scrod..... | 3,423 | 26 | 1,580 | 15 | 1,040 | 10 | 2,165 | 11 | 320 | 1 | 270 | 1 | 425 | 2 |
| Cod, salted: | | | | | | | | | | | | | | |
| Large..... | | | | | | | | | 21,645 | 555 | 13,460 | 336 | | |
| Market..... | | | | | | | | | 3,800 | 75 | 6,230 | 125 | | |
| Haddock, fresh: | | | | | | | | | | | | | | |
| Large..... | 196,729 | 9,170 | 224,685 | 6,683 | 306,669 | 10,684 | 557,723 | 7,340 | 697,212 | 9,671 | 233,433 | 6,521 | 680,467 | 15,525 |
| Scrod..... | 6,482 | 81 | 5,767 | 71 | 2,476 | 24 | 1,320 | 8 | 257 | 1 | 78,675 | 1,494 | 188,784 | 3,668 |
| Haddock, salted: | | | | | | | | | | | | | | |
| Large..... | | | | | | | | | | | 9,900 | 248 | | |
| Scrod..... | | | | | | | | | | | 1,615 | 24 | | |
| Hake, fresh: | | | | | | | | | | | | | | |
| Large..... | 3,030 | 58 | 500 | 10 | 145 | 4 | | | 20 | | 10 | | 3,965 | 29 |
| Small..... | 184,240 | 2,841 | 96,693 | 1,929 | 72,515 | 1,543 | 99,773 | 1,769 | 102,044 | 878 | 79,200 | 970 | 123,165 | 1,386 |
| Hake, salted, large..... | | | | | | | | | 1,675 | 17 | | | | |
| Pollock, fresh..... | 50,832 | 341 | 32,585 | 396 | 57,001 | 721 | 24,285 | 98 | 36,602 | 148 | 80,685 | 393 | 52,802 | 347 |
| Pollock, salted..... | | | | | | | | | | | | | | |
| Cusk, fresh..... | 79,184 | 1,225 | 65,146 | 1,193 | 89,885 | 1,589 | 113,155 | 1,032 | 96,380 | 729 | 1,876 | 26 | 873 | 14 |
| Cusk, salted..... | | | | | | | | | 330 | 3 | | | | |
| Halibut, fresh..... | 394 | 59 | 216 | 27 | 67 | 10 | 32,252 | 3,551 | 93,313 | 6,982 | 86,402 | 8,451 | 24,769 | 2,798 |
| Halibut, salted..... | | | | | | | | | 1,985 | 90 | | | | |
| Mackerel, fresh..... | | | | | | | | | 10,350 | 214 | 25,995 | 412 | 391,558 | 3,639 |
| Flounders, fresh..... | 29,219 | 906 | 35,773 | 998 | 93,508 | 3,841 | 26,525 | 567 | 40,017 | 485 | 98,691 | 1,104 | 93,483 | 1,897 |
| Swordfish, fresh..... | | | | | | | | | | | 15,861 | 3,423 | 53,899 | 9,099 |
| Herring, fresh..... | | | | | | | | | | | 4,200 | 32 | 28,400 | 250 |
| Other, fresh..... | 26,464 | 592 | 14,978 | 446 | 15,476 | 464 | 2,451 | 33 | 2,566 | 30 | 17,986 | 188 | 46,237 | 382 |
| Total, fresh..... | 690,266 | 17,726 | 586,398 | 16,254 | 867,002 | 24,235 | 1,206,746 | 18,502 | 1,407,183 | 23,196 | 1,142,895 | 32,765 | 2,200,039 | 50,728 |
| Total, salted..... | | | | | | | | | 29,435 | 740 | 31,205 | 733 | | |
| Grand total..... | 690,266 | 17,726 | 586,398 | 16,254 | 867,002 | 24,235 | 1,206,746 | 18,502 | 1,436,618 | 23,936 | 1,174,100 | 33,498 | 2,200,039 | 50,726 |
| Landed in 1932: | | | | | | | | | | | | | | |
| Fresh..... | 584,943 | 17,884 | 515,287 | 17,952 | 844,076 | 23,264 | 1,678,850 | 25,482 | 788,605 | 11,812 | 730,969 | 23,183 | 1,072,120 | 32,923 |
| Salted..... | | | | | | | 1,850 | 41 | 5,455 | 148 | 3,395 | 95 | 556 | 16 |
| Total..... | 584,943 | 17,884 | 515,287 | 17,952 | 844,076 | 23,264 | 1,678,700 | 25,523 | 794,060 | 11,960 | 734,364 | 23,278 | 1,072,676 | 32,939 |

| Species | August | | September | | October | | November | | December | | Total, 1933 | | 1932 | |
|--------------------|-----------|---------|-----------|---------|---------|---------|----------|---------|----------|---------|-------------|----------|------------|----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Cod, fresh: | | | | | | | | | | | | | | |
| Large..... | 300,398 | \$9,783 | 219,249 | \$9,146 | 93,106 | \$3,087 | 48,200 | \$1,643 | 49,330 | \$1,488 | 2,206,728 | \$56,531 | 2,507,439 | \$72,657 |
| Market..... | 98,562 | 1,916 | 26,859 | 501 | 34,635 | 642 | 28,035 | 613 | 29,330 | 716 | 773,912 | 12,862 | 567,519 | 8,002 |
| Scrod..... | 290 | 1 | 2,885 | 18 | 1,815 | 11 | 2,665 | 22 | 1,985 | 21 | 18,863 | 139 | 23,932 | 176 |
| Cod, salted: | | | | | | | | | | | | | | |
| Large..... | 25,571 | 792 | | | 430 | 15 | | | | | 61,106 | 1,698 | 6,306 | 205 |
| Market..... | 3,135 | 66 | | | 160 | 3 | | | | | 13,325 | 269 | 3,805 | 87 |
| Scrod..... | | | | | | | | | | | | | 90 | 2 |
| Haddock, fresh: | | | | | | | | | | | | | | |
| Large..... | 567,743 | 14,161 | 156,206 | 7,760 | 152,209 | 7,568 | 102,578 | 5,570 | 86,890 | 4,906 | 3,963,564 | 107,559 | 2,882,137 | 90,213 |
| Scrod..... | 131,655 | 2,572 | 10,471 | 98 | 6,319 | 45 | 6,668 | 84 | 3,587 | 53 | 442,460 | 8,204 | 70,366 | 618 |
| Haddock, salted: | | | | | | | | | | | | | | |
| Large..... | | | | | | | | | | | 9,900 | 248 | | |
| Scrod..... | | | | | | | | | | | 1,615 | 24 | | |
| Hake, fresh: | | | | | | | | | | | | | | |
| Large..... | 68,205 | 817 | 143,056 | 2,184 | 217,722 | 3,704 | 126,640 | 2,633 | 73,738 | 1,791 | 637,031 | 11,230 | 195,436 | 3,551 |
| Small..... | 76,505 | 609 | 3,575 | 24 | 6,940 | 46 | 1,450 | 12 | 1,293 | 14 | 847,393 | 11,921 | 848,531 | 12,138 |
| Hake, salted: | | | | | | | | | | | | | | |
| Large..... | | | | | | | | | | | 1,675 | 17 | | |
| Small..... | | | | | | | | | | | | | 455 | 4 |
| Pollock, fresh | | | | | | | | | | | | | | |
| Large..... | 133,556 | 815 | 133,008 | 910 | 45,335 | 877 | 48,492 | 520 | 40,332 | 334 | 735,515 | 5,400 | 890,639 | 5,354 |
| Small..... | | | | | | | | | | | | | 615 | 3 |
| Pollock, salted | | | | | | | | | | | | | | |
| Large..... | 30,045 | 235 | 28,043 | 450 | 70,822 | 1,145 | 42,069 | 837 | 33,693 | 773 | 651,171 | 9,248 | 333,388 | 5,817 |
| Small..... | | | | | | | | | | | 330 | 3 | 160 | 2 |
| Cusk, fresh | | | | | | | | | | | | | | |
| Large..... | 110,623 | 9,967 | 24,784 | 2,293 | 1,171 | 116 | 1,670 | 210 | 967 | 140 | 376,628 | 34,604 | 108,533 | 10,191 |
| Small..... | | | | | | | | | | | 1,985 | 90 | 195 | 6 |
| Cusk, salted | | | | | | | | | | | | | | |
| Large..... | 353,476 | 3,352 | 163,824 | 2,180 | 56,398 | 2,224 | 10,297 | 460 | | | 1,011,928 | 12,481 | 1,677,321 | 18,563 |
| Small..... | 30,100 | 396 | 8,000 | 80 | | | | | | | 38,100 | 476 | 90,500 | 591 |
| Flounders, fresh | | | | | | | | | | | | | | |
| Large..... | 79,559 | 2,106 | 39,913 | 1,507 | 10,017 | 386 | 5,443 | 203 | 30,883 | 710 | 583,031 | 14,802 | 230,034 | 5,588 |
| Small..... | 79,780 | 10,631 | 6,299 | 1,032 | | | | | | | 155,839 | 24,185 | 339,361 | 41,052 |
| Swordfish, fresh | | | | | | | | | | | | | | |
| Large..... | 20,000 | 100 | 90,400 | 380 | | | | | | | 143,000 | 762 | 71,000 | 355 |
| Small..... | 4,928 | 43 | 24,778 | 211 | 1,702 | 46 | 4,015 | 80 | 492 | 92 | 162,073 | 2,607 | 537,371 | 7,405 |
| Other, fresh | | | | | | | | | | | | | | |
| Total, fresh..... | 2,055,325 | 57,108 | 1,073,350 | 23,694 | 699,190 | 19,397 | 428,222 | 12,887 | 352,520 | 11,043 | 12,709,136 | 312,535 | 11,285,007 | 281,950 |
| Total, salted..... | 58,806 | 1,254 | 8,000 | 80 | 590 | 18 | | | | | 128,036 | 2,825 | 102,126 | 900 |
| Grand total..... | 2,114,131 | 58,362 | 1,081,350 | 28,774 | 699,780 | 19,415 | 428,222 | 12,887 | 352,520 | 11,043 | 12,837,172 | 315,360 | 11,387,133 | 282,850 |
| Landed in 1932: | | | | | | | | | | | | | | |
| Fresh..... | 1,576,395 | 42,849 | 1,694,588 | 36,828 | 679,803 | 17,840 | 538,505 | 16,607 | 582,866 | 15,326 | | | 11,285,007 | 281,950 |
| Salted..... | 55,000 | 413 | 35,500 | 178 | | | 370 | 9 | | | | | 102,126 | 900 |
| Total..... | 1,631,395 | 43,262 | 1,730,088 | 37,006 | 679,803 | 17,840 | 538,875 | 16,616 | 582,866 | 15,326 | | | 11,387,133 | 282,850 |

Landings by fishing vessels at the 3 principal New England ports, 1933—Continued

SUMMARY: BY PORTS

| Species | Boston | | Gloucester | | Portland | | Total, 1933 | | 1932 | |
|-----------------------|-------------|-----------|------------|-----------|------------|----------|-------------|-----------|-------------|-----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Cod, fresh: | | | | | | | | | | |
| Large..... | 27,961,973 | \$681,634 | 5,534,610 | \$123,294 | 2,206,728 | \$56,531 | 35,703,311 | \$861,459 | 28,835,774 | \$785,633 |
| Market..... | 33,025,829 | 674,097 | 1,304,473 | 17,284 | 773,912 | 12,862 | 35,104,214 | 704,243 | 29,298,835 | 480,150 |
| Scrod..... | 195,945 | 3,393 | 16,210 | 201 | 18,863 | 139 | 231,018 | 3,733 | 128,067 | 1,937 |
| Cod, salted: | | | | | | | | | | |
| Large..... | 50,500 | 1,540 | 1,533,428 | 46,375 | 61,106 | 1,698 | 1,645,034 | 49,613 | 714,386 | 21,965 |
| Market..... | 15,500 | 329 | 893,699 | 19,381 | 13,325 | 269 | 922,524 | 19,979 | 451,724 | 10,479 |
| Scrod..... | | | 235,677 | 2,718 | | | 235,677 | 2,718 | 122,055 | 1,349 |
| Haddock, fresh: | | | | | | | | | | |
| Large..... | 90,353,894 | 2,565,302 | 2,465,674 | 51,952 | 3,963,564 | 107,559 | 96,783,132 | 2,724,813 | 92,178,669 | 2,505,086 |
| Scrod..... | 32,531,860 | 639,713 | 89,805 | 937 | 442,460 | 8,204 | 33,064,125 | 648,854 | 27,938,205 | 401,722 |
| Haddock, salted: | | | | | | | | | | |
| Large..... | 400 | 4 | | | 9,900 | 248 | 10,300 | 252 | | |
| Scrod..... | | | 2,450 | 12 | 1,615 | 24 | 4,065 | 36 | | |
| Hake, fresh: | | | | | | | | | | |
| Large..... | 4,683,613 | 100,434 | 476,042 | 9,148 | 637,031 | 11,230 | 5,795,686 | 120,812 | 6,872,088 | 124,038 |
| Small..... | 537,645 | 11,331 | 11,805 | 204 | 847,393 | 11,921 | 1,396,843 | 23,456 | 867,531 | 12,456 |
| Hake, salted: | | | | | | | | | | |
| Large..... | | | 1,355 | 14 | 1,675 | 17 | 3,030 | 31 | 1,045 | 14 |
| Small..... | | | | | | | | | 4,455 | 84 |
| Pollock, fresh..... | 8,261,580 | 105,861 | 3,398,112 | 47,553 | 735,515 | 5,400 | 12,395,207 | 168,814 | 7,841,032 | 85,600 |
| Pollock, salted..... | 50 | | 90 | 1 | | | 140 | 1 | 9,210 | 89 |
| Cusk, fresh..... | 2,498,576 | 38,290 | 278,865 | 1,698 | 651,171 | 9,248 | 3,428,612 | 49,236 | 3,064,840 | 45,706 |
| Cusk, salted..... | | | 1,394 | 19 | 330 | 3 | 1,714 | 22 | 4,160 | 69 |
| Halibut, fresh..... | 1,755,664 | 191,012 | 4,280 | 410 | 376,628 | 34,604 | 2,136,572 | 226,026 | 2,355,029 | 261,154 |
| Halibut, salted..... | | | 130 | 7 | 1,985 | 90 | 2,115 | 97 | 1,035 | 65 |
| Mackerel, fresh..... | 17,348,117 | 402,310 | 3,980,544 | 81,289 | 1,011,928 | 12,481 | 22,340,589 | 496,050 | 37,982,803 | 617,120 |
| Mackerel, salted..... | 9,800 | 147 | 56,000 | 1,470 | 38,100 | 476 | 103,900 | 2,093 | 113,725 | 1,267 |
| Flounders, fresh..... | 9,278,542 | 349,113 | 376,910 | 12,594 | 583,031 | 14,802 | 10,238,483 | 376,509 | 7,442,833 | 295,517 |
| Swordfish, fresh..... | 1,681,304 | 281,694 | 771 | 110 | 155,839 | 24,185 | 1,837,914 | 305,989 | 2,603,915 | 356,933 |
| Herring, fresh..... | 4,105 | 136 | 11,500 | 91 | 143,000 | 762 | 158,605 | 989 | 99,500 | 655 |
| Herring, salted..... | | | 703,494 | 21,290 | | | 703,494 | 21,290 | 655,700 | 20,130 |
| Other, fresh..... | 2,388,152 | 47,264 | 359,689 | 3,885 | 162,073 | 2,607 | 2,909,914 | 53,756 | 2,747,709 | 54,623 |
| Total, fresh..... | 232,506,799 | 6,091,584 | 18,309,290 | 350,650 | 12,709,136 | 312,535 | 263,525,225 | 6,754,769 | 250,256,830 | 6,028,330 |
| Total, salted..... | 76,250 | 2,020 | 3,427,707 | 91,287 | 128,036 | 2,825 | 3,631,993 | 96,132 | 2,077,495 | 55,521 |
| Grand total..... | 232,583,049 | 6,093,604 | 21,736,997 | 441,937 | 12,837,172 | 315,360 | 267,157,218 | 6,850,901 | 252,334,325 | 6,083,851 |
| Landed in 1932: | | | | | | | | | | |
| Fresh..... | 215,527,504 | 5,364,667 | 23,444,319 | 381,713 | 11,285,007 | 281,950 | | | 250,256,830 | 6,028,330 |
| Salted..... | 91,475 | 2,258 | 1,883,894 | 52,363 | 102,126 | 900 | | | 2,077,495 | 55,521 |
| Total..... | 215,618,979 | 5,366,925 | 25,328,213 | 434,076 | 11,387,133 | 282,850 | | | 252,334,325 | 6,083,851 |

¹ The items under "Other, fresh" include alewives, 341,725 pounds, value \$2,910; butterfish, 97,320 pounds, value \$5,654; rosefish, 250,075 pounds, value \$2,639; scup, 6,700 pounds, value \$100; sea robins, 400 pounds, value \$4; shad, 70,871 pounds, value \$917; sharks, 21,363 pounds, value \$313; skates, 10,308 pounds, value \$208; smelt, 11,767 pounds, value \$327; sturgeon, 1,322 pounds, value \$178; tuna or "horse mackerel", 4,826 pounds, value \$200; whiting, 137,340 pounds, value \$3,191; wolfish, 1,814,264 pounds, value \$30,470; lobsters, 259 pounds, value \$58; scallops, 5,018 pounds, value \$649; livers, 36,100 pounds, value \$722; sounds, 350 pounds, value \$14; spawn, 95,355 pounds, value \$5,055; and mixed fish, 3,550 pounds, value \$143.

BIOLOGICAL ASPECT

In 1933 the fishing fleet landing fares at Boston and Gloucester, Mass., and Portland, Maine, and operating on the fishing banks of the North Atlantic numbered 390 steam, motor, and sail vessels of 5 net tons capacity or greater as measured by the United States Customs Service. These made 12,214 trips to the fishing grounds, and were absent from port 52,651 days, or an average of 4.3 days per trip. This is 0.1 of a day less than the average length of a trip during 1932. The catches of edible fish landed at the 3 ports amounted to 270,145,894 pounds when the salted fish had been converted to the basis of fresh gutted or round fish as landed. This however does not represent the entire catch of edible fish of these vessels, for small quantities estimated at not more than 5 percent of their total catch were landed at ports in New England other than these three, at New York City, and at ports in New Jersey.

Otter trawls on all sizes of vessels accounted for 169,683,377 pounds, or 63 percent of the total landings. Line trawls were next in importance, accounting for 62,260,813 pounds, or 23 percent of the total landings.

The catch taken on Georges Bank and landed at the 3 ports amounted to 90,850,098 pounds, or 34 percent of the total; that on shore grounds, 45,602,905 pounds, or 17 percent; Browns Bank, 37,249,263 pounds, or 14 percent; Sable Island Bank, 30,415,492 pounds, or 11 percent; and South Channel, 27,617,824 pounds, or 10 percent. No other bank accounted for as much as 10,000,000 pounds in the landings at the 3 ports.

Landings by fishing vessels at the 3 principal New England ports, 1933

BY GEAR AND FISHING GROUNDS

| Gear and fishing grounds | Vessels fishing | Trips | Days absent | Cod | | | Haddock | | Hake | |
|---------------------------------------|-----------------|--------------|---------------|-------------------|-------------------|----------------|-------------------|------------------|------------------|----------------|
| | | | | Large | Market | Scrod | Large | Scrod | Large | Small |
| | Number | Number | Number | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds |
| Line trawls: | | | | | | | | | | |
| Off Funks..... | 1 | 1 | 14 | | | | | | | |
| Grand Bank..... | 6 | 13 | 283 | 122,871 | 24,501 | | 20,394 | 3,392 | | |
| St. Peters Bank..... | 2 | 2 | 31 | 3,700 | 7,740 | | 8,400 | 425 | | |
| Off Newfoundland (Treaty Coast)..... | 1 | 1 | 7 | 52,250 | 35,413 | | | | | |
| Gulf of St. Lawrence..... | 6 | 13 | 366 | 2,605,238 | 1,651,667 | 466,641 | | | 3,182 | 2,700 |
| Quereau Bank..... | 6 | 15 | 312 | 152,349 | 23,764 | | | | 6,717 | |
| The Gully..... | 1 | 2 | 35 | 12,946 | 1,017 | | | | | |
| Sable Island Bank (Western Bank)..... | 21 | 35 | 528 | 706,595 | 595,330 | | 913,375 | 29,900 | 34,590 | 5,175 |
| Cape Shore..... | 36 | 108 | 1,363 | 1,074,260 | 1,949,015 | 1,500 | 1,655,775 | 517,990 | 329,530 | |
| Emerald Bank..... | 8 | 9 | 85 | 101,040 | 115,575 | | 252,900 | 20,825 | 13,545 | |
| La Have Bank..... | 23 | 51 | 618 | 942,705 | 929,094 | | 1,225,650 | 90,650 | 134,875 | |
| Roseway Bank..... | 5 | 6 | 54 | 80,570 | 118,900 | | 126,100 | 19,700 | 14,300 | |
| Browns Bank..... | 44 | 270 | 3,008 | 3,409,825 | 2,984,177 | 11,595 | 8,353,630 | 729,495 | 442,909 | 4,100 |
| Georges Bank..... | 46 | 160 | 1,562 | 2,916,335 | 1,184,478 | | 3,447,125 | 197,810 | 158,293 | 300 |
| Georges Bank (occasional)..... | | | | 54,790 | 6,600 | 740 | 10,700 | | | |
| South Channel..... | 33 | 193 | 1,488 | 1,441,952 | 1,703,218 | 2,350 | 4,084,004 | 172,200 | 568,425 | |
| Off Highland Light..... | 6 | 13 | 81 | 15,735 | 8,015 | | 112,830 | 7,040 | 38,940 | |
| Off Chatham..... | 9 | 13 | 80 | 37,915 | 35,165 | 175 | 190,950 | 2,060 | 23,040 | |
| Nantucket Shoals..... | 3 | 5 | 52 | 19,470 | 58,400 | | 14,200 | 1,300 | | |
| Cashes Bank..... | 18 | 39 | 160 | 94,800 | 77,355 | 6,490 | 116,245 | 9,886 | 133,285 | 115,465 |
| Fippenies Bank..... | 13 | 23 | 101 | 48,180 | 27,025 | 620 | 109,395 | 17,172 | 99,180 | 750 |
| Platts Bank..... | 11 | 65 | 151 | 82,865 | 43,367 | 5,025 | 137,287 | 13,917 | 87,520 | 94,505 |
| Jeffreys Ledge..... | 17 | 149 | 312 | 83,605 | 47,693 | 4,060 | 327,984 | 16,666 | 135,478 | 138,915 |
| Tillies Bank..... | 1 | 1 | 12 | 190 | 70 | | 8,600 | 700 | 1,400 | |
| Middle Bank (Stellwagen Bank)..... | 16 | 71 | 317 | 46,315 | 41,320 | 225 | 400,875 | 19,500 | 338,393 | 2,575 |
| South..... | 1 | 1 | 3 | 200 | 300 | | 600 | | | |
| Shore, general..... | 89 | 863 | 2,174 | 647,501 | 471,443 | 28,140 | 1,268,550 | 40,618 | 499,500 | 353,564 |
| Shore, general (occasional)..... | | | | 100 | 108 | | 217 | | | 50 |
| Total..... | 1 135 | 2,122 | 13,197 | 14,754,282 | 12,140,840 | 527,581 | 22,885,586 | 1,911,356 | 3,063,102 | 718,099 |
| Hand lines: | | | | | | | | | | |
| Cape Shore..... | 3 | 12 | 115 | 156,700 | 263,500 | | 11,215 | | 1,900 | |
| Browns Bank..... | 1 | 1 | 11 | 16,415 | 21,800 | | 200 | | | |
| Georges Bank..... | 3 | 15 | 139 | 241,175 | 209,595 | | 1,610 | | 30 | |
| South Channel..... | 2 | 2 | 11 | 9,200 | 10,550 | | 19,900 | 150 | 10,850 | |
| Nantucket Shoals..... | 3 | 8 | 74 | 31,355 | 45,625 | | 100 | | | |

| | 8 | 3 | 13 | 11,130 | 14,950 | 365 | 145 | | | |
|----------------------------------|------------|--------------|---------------|-------------------|-------------------|----------------|-------------------|-------------------|------------------|----------------|
| Shore, general | | | | 11,130 | 14,950 | | | | | |
| Shore, general (occasional) | | | | 6,047 | 2,290 | 5 | | | 15 | |
| Total | 19 | 41 | 363 | 472,022 | 568,310 | 5 | 33,390 | 150 | 12,925 | 15 |
| Harpoons: | | | | | | | | | | |
| Cape Shore | 34 | 37 | 910 | | | | | | | |
| Georges Bank | 71 | 199 | 3,834 | | | | | | | |
| South Channel | 1 | 1 | 30 | | | | | | | |
| Nantucket Shoals | 5 | 5 | 96 | | | | | | | |
| Shore, general | 9 | 11 | 152 | | | | | | | |
| Total | 174 | 253 | 5,022 | | | | | | | |
| Otter trawls, large: | | | | | | | | | | |
| Gulf of St. Lawrence | 1 | 3 | 35 | 12,675 | 33,900 | | 85,750 | 68,600 | 55,100 | |
| Sable Island Bank (Western Bank) | 50 | 238 | 2,854 | 3,500,208 | 3,974,745 | 26,680 | 6,117,125 | 10,114,270 | 374,435 | 1,000 |
| Cape Shore | 8 | 9 | 109 | 76,240 | 99,950 | | 133,370 | 198,370 | 13,870 | |
| Emerald Bank | 3 | 3 | 30 | 18,750 | 49,500 | 2,800 | 139,810 | 47,700 | 8,020 | |
| La Have Bank | 18 | 31 | 318 | 599,990 | 922,290 | 17,920 | 2,128,620 | 642,200 | 26,365 | |
| Browns Bank | 42 | 137 | 1,362 | 1,907,000 | 2,571,225 | 1,350 | 7,929,270 | 2,370,395 | 88,335 | 200 |
| Georges Bank | 46 | 672 | 6,583 | 8,238,780 | 9,508,000 | 78,100 | 29,820,990 | 11,428,905 | 690,225 | 1,400 |
| South Channel | 37 | 131 | 1,146 | 769,795 | 1,069,145 | 4,700 | 6,417,273 | 1,357,050 | 353,550 | 2,010 |
| Off Chatham | 1 | 1 | 9 | 3,800 | 20,200 | | 84,200 | 43,800 | 300 | |
| Nantucket Shoals | 2 | 2 | 23 | 4,800 | 3,400 | | 81,200 | 10,600 | 2,400 | |
| Cashes Bank | 1 | 1 | 3 | 13,480 | 30,800 | | 79,100 | 17,900 | 700 | |
| Shore, general | 6 | 8 | 68 | 50,800 | 139,500 | | 302,200 | 182,100 | 4,280 | |
| Total | 152 | 1,236 | 12,540 | 15,196,318 | 18,423,655 | 131,650 | 53,320,908 | 26,381,890 | 1,617,580 | 4,610 |
| Otter trawls, medium: | | | | | | | | | | |
| Sable Island Bank (Western Bank) | 13 | 29 | 342 | 318,680 | 344,950 | | 460,750 | 386,410 | 38,015 | |
| Cape Shore | 7 | 7 | 78 | 26,030 | 96,265 | | 76,360 | 96,750 | 7,280 | |
| Emerald Bank | 1 | 1 | 13 | 2,380 | 6,000 | | 3,600 | 6,800 | 3,000 | |
| La Have Bank | 6 | 11 | 124 | 154,150 | 171,425 | | 198,700 | 80,100 | 5,690 | |
| Browns Bank | 20 | 47 | 458 | 353,000 | 435,165 | | 1,472,870 | 360,825 | 22,700 | |
| Georges Bank | 41 | 375 | 3,401 | 1,872,835 | 1,919,510 | 3,295 | 7,183,296 | 2,373,385 | 124,775 | 80 |
| Georges Bank (occasional) | | | | | | | 21,700 | | | |
| South Channel | 34 | 199 | 1,613 | 372,684 | 814,285 | 2,000 | 4,346,610 | 1,054,655 | 127,685 | |
| Off Highland Light | 11 | 19 | 146 | 34,395 | 20,180 | | 197,850 | 29,910 | 29,225 | |
| Off Chatham | 15 | 67 | 511 | 175,926 | 587,370 | 1,200 | 1,090,250 | 171,740 | 75,245 | 800 |
| Nantucket Shoals | 13 | 16 | 111 | 6,740 | 67,935 | 700 | 194,820 | 47,125 | 9,910 | |
| Middle Bank (Stellwagen Bank) | 2 | 11 | 46 | 13,455 | 14,290 | | 105,865 | 4,010 | 45,306 | |
| South | 1 | 1 | 4 | | | | | | | |
| Shore, general | 46 | 628 | 1,515 | 161,080 | 288,619 | 12,655 | 1,158,665 | 79,655 | 69,595 | 111,058 |
| Total | 177 | 1,311 | 8,362 | 3,491,355 | 4,755,994 | 19,850 | 16,511,426 | 4,687,365 | 558,435 | 111,938 |

Landings by fishing vessels at the 3 principal New England ports, 1933—Continued

BY GEAR AND FISHING GROUNDS—Continued

| Gear and fishing grounds | Vessels fishing | Trips | Days absent | Cod | | | Haddock | | Hake | |
|-----------------------------------|-----------------|--------|-------------|-----------|---------|--------|-----------|--------|---------|---------|
| | | | | Large | Market | Scrod | Large | Scrod | Large | Small |
| | Number | Number | Number | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds |
| Other trawls, small: | | | | | | | | | | |
| Georges Bank | 3 | 5 | 31 | 850 | 5,500 | | 26,565 | 6,000 | 1,550 | |
| South Channel | 2 | 14 | 100 | 8,595 | 66,300 | | 260,780 | 42,950 | 2,550 | |
| Off Chatham | 2 | 2 | 11 | 735 | 29,300 | 300 | 13,990 | 1,700 | 400 | |
| Nantucket Shoals | 2 | 2 | 14 | | 5,500 | | 3,900 | 6,850 | | |
| Cashes Bank | 1 | 1 | 9 | 1,200 | 1,500 | | 23,500 | 9,000 | 300 | |
| Fippenies Bank | 1 | 1 | 6 | 1,530 | 440 | | 2,225 | | 4,500 | |
| Shore, general | 88 | 1,581 | 4,422 | 417,913 | 486,488 | 15,313 | 2,089,192 | 23,441 | 127,065 | 373,355 |
| Total | 190 | 1,606 | 4,593 | 430,823 | 595,028 | 15,613 | 2,420,152 | 89,941 | 136,365 | 373,355 |
| Sink gill nets: | | | | | | | | | | |
| Jeffreys Ledge | 1 | 3 | 3 | 3,697 | 1,230 | | 1,460 | | | 10 |
| Shore, general | 32 | 3,765 | 3,783 | 4,480,355 | 408,848 | 3,060 | 1,631,078 | 1,920 | 413,886 | 188,816 |
| Total | 132 | 3,768 | 3,786 | 4,484,052 | 410,078 | 3,060 | 1,632,538 | 1,920 | 413,886 | 188,826 |
| Drift gill nets: | | | | | | | | | | |
| Bay of Islands | 2 | 2 | 122 | | | | | | | |
| Sable Island Bank (Western Bank) | 1 | 1 | 4 | | | | | | | |
| Cape Shore | 1 | 1 | 8 | | | | | | | |
| Off Chatham | 1 | 1 | 2 | | | | | | | |
| Shore, general | 60 | 356 | 733 | | | | | | | |
| Total | 162 | 361 | 869 | | | | | | | |
| Purse seines: | | | | | | | | | | |
| Gulf of St. Lawrence (occasional) | | | | | | | | | | |
| Georges Bank | 3 | 3 | 8 | | | | | | | |
| Off Chatham | 32 | 43 | 146 | | | | | | | |
| Nantucket Shoals | 11 | 12 | 52 | | | | | | | |
| Middle Bank (Stellwagen Bank) | 20 | 21 | 59 | | | | | | | |
| South | 48 | 70 | 240 | | | | | | | |
| Shore, general | 101 | 1,363 | 3,405 | 20 | 5 | | 350 | 40 | 150 | |
| Total | 1102 | 1,512 | 3,910 | 20 | 5 | | 350 | 40 | 150 | |

| | | | | | | | | | | |
|----------------|-------|---------|---------|--------------|--------------|----------|--------------|--------------|-------------|-------------|
| Scallop drags: | | | | | | | | | | |
| Georges Bank | 1 | 1 | 6 | | | | | | | |
| Shore, general | 1 | 3 | 3 | | | | | | | |
| Total | 2 | 4 | 9 | | | | | | | |
| Grand total | 1 390 | 12, 214 | 52, 651 | 38, 828, 872 | 36, 893, 910 | 697, 659 | 96, 904, 350 | 33, 072, 662 | 5, 802, 443 | 1, 396, 843 |

1 Exclusive of duplication.

2 Incidental catch.

NOTE.—The three principal New England ports are Boston and Gloucester, Mass., and Portland, Maine. Otter trawls (including V-D trawls) are classified according to the size of the vessel. The weight of salted fish landed has been converted to the equivalent of fresh fish as landed. Only landings by vessels having a capacity of 5 net tons or greater are used in this tabulation. "Occasional" after the name of a bank or ground indicates that the vessel or vessels contributing to the catch as shown fished chiefly with another type of gear. In such cases the number of vessels fishing, number of trips, and number of days absent are shown under the principal types of gear used.

| Gear and fishing grounds | Pollock | Cusk | Halibut | Flounders | Swordfish | Mackerel | Herring | Other | Total |
|----------------------------------|-------------|-------------|-------------|-----------|-----------|----------|---------|----------|--------------|
| | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds |
| Line trawls: | | | | | | | | | |
| Off Funks | | | 26, 171 | | | | | | 26, 171 |
| Grand Bank | | | 426, 606 | | | | | | 597, 764 |
| St. Peters Bank | | | 52, 032 | | | | | | 72, 297 |
| Off Newfoundland (Treaty Coast) | | | | | | | | | 87, 663 |
| Gulf of St. Lawrence | | 1, 397 | 141, 198 | 220 | | | | | 4, 872, 213 |
| Quereau Bank | | 15, 565 | 265, 547 | | 441 | | | | 464, 383 |
| The Gully | | | 51, 196 | | | | | | 65, 156 |
| Sable Island Bank (Western Bank) | 20, 035 | 84, 230 | 142, 706 | | | 662 | | 350 | 2, 532, 908 |
| Cape Shore | 75, 065 | 432, 210 | 11, 350 | 2, 200 | | | | 28, 675 | 6, 077, 570 |
| Emerald Bank | 6, 200 | 8, 400 | 6, 718 | | | | | 2, 050 | 527, 153 |
| La Have Bank | 34, 120 | 199, 610 | 46, 935 | | | 1, 573 | | 14, 440 | 3, 619, 552 |
| Roseway Bank | 6, 050 | 21, 290 | 2, 972 | | | | | 1, 375 | 391, 347 |
| Browns Bank | 361, 421 | 1, 086, 750 | 167, 732 | 2, 825 | 1, 975 | | | 102, 580 | 17, 659, 014 |
| Georges Bank | 157, 005 | 202, 330 | 225, 873 | 2, 075 | 2, 493 | | | 8, 135 | 8, 302, 252 |
| Georges Bank (occasional) | 3, 620 | 430 | | | | | | | 76, 890 |
| South Channel | 214, 008 | 86, 825 | 23, 579 | 11, 610 | 148 | 85 | | 15, 150 | 8, 323, 644 |
| Off Highland Light | 4, 875 | 24, 950 | 323 | 4, 357 | | | | 470 | 217, 535 |
| Off Chatham | 12, 965 | 3, 150 | 1, 118 | | | | | 3, 755 | 310, 313 |
| Nantucket Shoals | | | 12, 230 | | 176 | | | | 105, 776 |
| Cashes Bank | 23, 355 | 173, 190 | 2, 191 | | | | | 2, 783 | 755, 045 |
| Fippenies Bank | 18, 785 | 33, 695 | 3, 139 | | | | | 275 | 358, 196 |
| Platts Bank | 33, 675 | 164, 600 | 2, 188 | | | | | 11, 414 | 676, 363 |
| Jeffreys Ledge | 50, 520 | 147, 670 | 2, 107 | | | | | 18, 826 | 973, 544 |
| Tillies Bank | 100 | 350 | 15 | | | | | | 11, 425 |
| Middle Bank (Stellwagen Bank) | 30, 885 | 70, 621 | 882 | | | | | 3, 985 | 965, 576 |
| South | 50 | 300 | | | | | | | 1, 450 |
| Shore, general | 143, 590 | 354, 320 | 5, 968 | 46, 977 | | 136 | | 38, 233 | 3, 998, 540 |
| Shore, general (occasional) | | 10 | 5 | 500 | | | | | 990 |
| Total | 1, 196, 374 | 3, 111, 863 | 1, 620, 781 | 70, 764 | 7, 468 | 221 | | 252, 496 | 62, 260, 813 |

Landings by fishing vessels at the 5 principal New England ports, 1933—Continued

BY GEAR AND FISHING GROUNDS—Continued

| Gear and fishing grounds | Pollock | Cusk | Halibut | Flounders | Swordfish | Mackerel | Herring | Other | Total |
|--|---------------|---------------|---------------|---------------|------------------|---------------|---------------|---------------|------------------|
| | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> |
| Hand lines: | | | | | | | | | |
| Cape Shore..... | 30,300 | 11,500 | 279 | | | | | 20,705 | 496,099 |
| Browns Bank..... | 1,700 | 700 | | | | | | 200 | 41,015 |
| Browns Bank (occasional)..... | | | 232 | | | | | | 232 |
| Georges Bank..... | 32,200 | 9,500 | 185 | | | | | 5,030 | 499,325 |
| Georges Bank (occasional)..... | | | 301 | | | | | | 301 |
| South Channel..... | 1,450 | 175 | 103 | 100 | | | | | 52,478 |
| Nantucket Shoals..... | 3,770 | | 351 | | | | | 175 | 81,376 |
| Shore, general..... | 630 | | | | | | | | 27,220 |
| Shore, general (occasional)..... | 30 | | | | | | | | 8,387 |
| Total..... | 70,080 | 21,875 | 1,451 | 100 | | | | 28,110 | 1,206,433 |
| Harpoons: | | | | | | | | | |
| Quereau Bank (occasional)..... | | | | | 3,930 | | | | 3,930 |
| The Gully (occasional)..... | | | | | 1,216 | | | | 1,216 |
| Sable Island Bank (Western Bank) (occasional)..... | | | | | 1,874 | | | | 1,874 |
| Cape Shore..... | | | | | 449,247 | | | | 449,247 |
| Cape Shore (occasional)..... | | | | | 294 | | | | 294 |
| La Have Bank (occasional)..... | | | | | 902 | | | | 902 |
| Browns Bank (occasional)..... | | | | | 7,306 | | | | 7,306 |
| Georges Bank..... | | | | | 1,257,907 | | | 3,367 | 1,261,274 |
| Georges Bank (occasional)..... | | | | | 5,160 | | | | 5,160 |
| South Channel..... | | | | | 12,602 | | | | 12,602 |
| Nantucket Shoals..... | | | | | 42,554 | | | | 42,554 |
| Shore, general..... | | | | | 46,934 | | | 987 | 47,921 |
| Total..... | | | | | 1,829,926 | | | 4,354 | 1,834,280 |
| Otter trawls, large: | | | | | | | | | |
| Gulf of St. Lawrence..... | 470 | | 3,174 | 31,950 | | | | 285 | 292,004 |
| Sable Island Bank (Western Bank)..... | 1,396,780 | 4,405 | 128,271 | 362,045 | | 1,400 | | 164,435 | 26,165,799 |
| Cape Shore..... | 34,730 | | 1,678 | 20,775 | | | | 7,170 | 588,153 |
| Emerald Bank..... | 12,800 | 500 | 703 | 5,300 | | | | 5,500 | 291,383 |
| La Have Bank..... | 334,300 | 11,355 | 25,472 | 45,075 | | | | 76,525 | 4,730,112 |
| Browns Bank..... | 1,105,040 | 30,130 | 79,045 | 173,400 | | | | 344,660 | 16,600,050 |
| Georges Bank..... | 2,637,875 | 161,700 | 163,853 | 1,394,045 | | 3,675 | 2,605 | 595,924 | 64,727,077 |
| South Channel..... | 540,375 | 9,955 | 33,228 | 318,634 | 300 | | | 180,175 | 11,054,515 |
| Off Chatham..... | 900 | | 200 | 9,800 | | | | | 163,208 |
| Nantucket Shoals..... | 1,900 | | | 5,700 | | | | 1,350 | 111,350 |

| | | | | | | | | | |
|----------------------------------|-----------|---------|---------|-----------|--------|-----------|-----------|-----------|-------------|
| Cashes Bank | 16,656 | | 230 | 1,075 | | | | 240 | 159,575 |
| Shore, general | 10,100 | 100 | 2,453 | 8,775 | | | | 9,802 | 710,110 |
| Total | 6,091,320 | 218,145 | 438,313 | 2,376,574 | \$ 300 | \$ 5,400 | \$ 2,005 | 1,386,166 | 125,595,334 |
| Otter trawls, medium: | | | | | | | | | |
| Sable Island Bank (Western Bank) | 76,730 | | 7,196 | 68,740 | | | | 4,250 | 1,705,721 |
| Cape Shore | 5,275 | 7,200 | 2,487 | 3,100 | | | | 1,800 | 327,547 |
| Emerald Bank | 1,200 | | 116 | 2,400 | | | | 250 | 25,746 |
| La Have Bank | 29,500 | 10,250 | 4,784 | 28,900 | | | | 6,390 | 688,879 |
| Browns Bank | 139,120 | 13,500 | 11,254 | 68,085 | | | | 75,125 | 2,941,646 |
| Georges Bank | 246,560 | 14,625 | 32,527 | 1,749,875 | 220 | 7,900 | 850 | 107,486 | 15,637,219 |
| Georges Bank (occasional) | | | 30 | | | | | | 21,738 |
| South Channel | 148,530 | 3,135 | 12,931 | 753,611 | | 1,150 | 650 | 127,723 | 7,763,649 |
| Off Highland Light | 236,600 | | 133 | 22,350 | | 300 | | 5,410 | 2,722,453 |
| Off Chatham | 51,250 | | 685 | 133,220 | | 625 | | 106,310 | 2,394,421 |
| Nantucket Shoals | 6,420 | | | 84,535 | | 145 | | 3,130 | 411,480 |
| Middle Bank (Stellwagen Bank) | 31,450 | 14,050 | 739 | 500 | | | | 1,195 | 230,359 |
| Shore, general | 149,515 | 1,875 | 869 | 1,226,335 | | | | 6,700 | 6,700 |
| Total | 1,120,150 | 64,635 | 73,661 | 4,136,651 | \$ 220 | \$ 10,020 | \$ 1,500 | 555,847 | 36,099,047 |
| Otter trawls, small: | | | | | | | | | |
| Georges Bank | 150 | | 785 | 37,280 | | | | 785 | 79,445 |
| South Channel | 100 | | 721 | 21,220 | | | | 5,720 | 408,936 |
| Off Chatham | 370 | | | 4,010 | | | | 430 | 51,235 |
| Nantucket Shoals | | | 13 | 23,330 | | | | | 39,593 |
| Cashes Bank | 400 | | 157 | 2,700 | | | | 500 | 39,257 |
| Fippenies Bank | 150 | 1,450 | | | | | | | 10,325 |
| Shore, general | 55,949 | 5,903 | 4,710 | 3,544,100 | | | | 216,776 | 7,360,205 |
| Total | 57,119 | 7,383 | 6,366 | 3,632,640 | | | | 224,211 | 7,988,996 |
| Sink gill nets: | | | | | | | | | |
| Jeffreys Ledge | 2,690 | 10 | | | | | | 87 | 9,184 |
| Shore, general | 3,847,380 | 7,858 | 230 | 21,504 | | 751 | | 31,227 | 11,036,913 |
| Total | 3,850,070 | 7,868 | 230 | 21,504 | | \$ 751 | | 31,314 | 11,046,097 |
| Drift gill nets: | | | | | | | | | |
| Bay of Islands | | | | | | | 1,055,241 | | 1,055,241 |
| Sable Island Bank (Western Bank) | | | | | | 9,100 | | | 9,100 |
| Cape Shore | | | | | | 13,980 | | | 13,980 |
| Off Chatham | | | | | | 600 | | 900 | 1,500 |
| Shore, general | | | | | | 980,030 | | 1,723 | 981,753 |
| Total | | | | | | 1,003,710 | 1,055,241 | 2,623 | 2,061,574 |

*Incidental catch.

Landings by fishing vessels at the 3 principal New England ports, 1933—Continued

BY GEAR AND FISHING GROUNDS—Continued

| Gear and fishing grounds | Pollock | Cusk | Halibut | Flounders | Swordfish | Mackerel | Herring | Other | Total |
|--|-------------------|------------------|------------------|-------------------|------------------|-------------------|------------------|------------------|--------------------|
| | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> |
| Purse seines: | | | | | | | | | |
| Gulf of St. Lawrence (occasional)..... | | | | | | 54,000 | | | 54,000 |
| Georges Bank..... | | | | | | 35,405 | | | 35,405 |
| Off Chatham..... | | | | | | 716,238 | | | 716,238 |
| Nantucket Shoals..... | | | | | | 214,885 | | | 214,885 |
| Middle Bank (Stellwagen Bank)..... | | | | | | 323,775 | | | 323,775 |
| South..... | | | | | | 2,645,138 | | | 2,645,138 |
| Shore, general..... | 10,360 | 100 | | 250 | | 17,471,311 | 154,500 | 422,375 | 18,059,401 |
| Total..... | 10,360 | 100 | | 250 | | 21,460,752 | 154,500 | 422,375 | 22,048,902 |
| Scallop drags: | | | | | | | | | |
| Georges Bank..... | | | | | | | | 4,022 | 4,022 |
| Shore, general..... | | | | | | | | 396 | 396 |
| Total..... | | | | | | | | 4,418 | 4,418 |
| Grand total..... | 12,395,473 | 3,431,869 | 2,140,902 | 10,238,483 | 1,837,914 | 22,480,854 | 1,213,846 | 2,909,914 | 270,145,894 |

SUMMARY: BY FISHING GROUNDS

| Fishing grounds | Vessels fishing | Trips | Days absent | Cod | | | Haddock | | Hake | |
|--------------------------------------|-----------------|---------------|---------------|----------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | | | | Large | Market | Scrod | Large | Scrod | Large | Small |
| | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> |
| Off Newfoundland: | | | | | | | | | | |
| Area XVIII: Off Funks..... | 1 | 1 | 14 | | | | | | | |
| Area XIX: | | | | | | | | | | |
| Bay of Islands..... | 2 | 2 | 122 | | | | | | | |
| Off Newfoundland (Treaty Coast)..... | 1 | 1 | 7 | 52,250 | 35,413 | | | | | |
| Area XX: | | | | | | | | | | |
| Grand Bank..... | 6 | 13 | 283 | 122,871 | 24,501 | | 20,394 | 3,392 | | |
| St. Peters Bank..... | 2 | 2 | 31 | 3,700 | 7,740 | | 8,400 | 425 | | |
| Total..... | 19 | 19 | 457 | 178,821 | 67,654 | | 28,794 | 3,817 | | |

| | | | | | | | | | | |
|---------------------------------------|-----|--------|--------|------------|------------|---------|------------|------------|-----------|-----------|
| Off Canada: | | | | | | | | | | |
| Area XIX: Gulf of St. Lawrence..... | 7 | 16 | 401 | 2,617,913 | 1,685,567 | 466,641 | 85,750 | 68,600 | 58,282 | 2,700 |
| Area XXI: | | | | | | | | | | |
| Quereu Bank..... | 6 | 15 | 312 | 152,349 | 23,764 | | | | 6,717 | |
| The Gully..... | 1 | 2 | 35 | 12,946 | 12,017 | | | | | |
| Sable Island Bank (Western Bank)..... | 84 | 303 | 3,726 | 4,525,483 | 4,915,025 | 26,690 | 7,491,250 | 10,530,580 | 447,040 | 6,175 |
| Cape Shore..... | 81 | 174 | 2,583 | 1,333,230 | 2,408,730 | 1,500 | 1,878,710 | 813,110 | 352,590 | |
| Emerald Bank..... | 12 | 13 | 126 | 122,170 | 171,075 | 2,800 | 396,210 | 75,325 | 24,565 | |
| La Have Bank..... | 45 | 93 | 1,090 | 1,696,845 | 2,022,809 | 17,920 | 3,552,870 | 712,950 | 166,930 | |
| Roseway Bank..... | 5 | 6 | 54 | 80,570 | 118,990 | | 126,100 | 19,700 | 14,300 | |
| Browns Bank..... | 104 | 455 | 4,839 | 5,686,240 | 6,012,267 | 12,945 | 17,755,970 | 3,460,715 | 553,944 | 4,300 |
| Total..... | 149 | 1,077 | 13,140 | 16,227,746 | 17,359,344 | 528,486 | 31,286,860 | 15,680,980 | 1,624,368 | 13,175 |
| Off United States: | | | | | | | | | | |
| Area XXII: | | | | | | | | | | |
| Georges Bank..... | 187 | 1,430 | 15,564 | 13,324,765 | 12,834,683 | 82,135 | 40,511,986 | 14,006,100 | 974,873 | 1,780 |
| South Channel..... | 103 | 540 | 4,358 | 2,602,226 | 3,663,498 | 9,050 | 15,128,587 | 2,627,095 | 1,063,090 | 2,010 |
| Off Highland Light..... | 17 | 32 | 227 | 80,130 | 28,198 | | 310,780 | 32,950 | 66,163 | |
| Off Chatham..... | 59 | 127 | 759 | 218,376 | 672,035 | 1,675 | 1,379,390 | 219,320 | 98,965 | 800 |
| Nantucket Shoals..... | 38 | 50 | 422 | 62,365 | 170,860 | 700 | 264,220 | 65,876 | 12,310 | |
| Cashes Bank..... | 20 | 41 | 172 | 109,480 | 109,655 | 6,490 | 218,845 | 36,786 | 134,265 | 115,465 |
| Flippenies Bank..... | 13 | 24 | 107 | 49,690 | 27,466 | 620 | 111,620 | 17,172 | 103,680 | 750 |
| Platts Bank..... | 11 | 65 | 151 | 82,865 | 43,367 | 6,025 | 137,287 | 13,917 | 87,520 | 94,505 |
| Jeffreys Ledge..... | 18 | 152 | 815 | 87,302 | 48,923 | 4,080 | 329,444 | 16,666 | 135,478 | 138,925 |
| Tillies Bank..... | 1 | 1 | 12 | 190 | 70 | | 8,600 | 700 | 1,400 | |
| Middle Bank (Stellwagen Bank)..... | 36 | 103 | 422 | 59,770 | 55,610 | 225 | 506,740 | 23,510 | 383,698 | 2,575 |
| Shore, general..... | 279 | 8,481 | 16,266 | 5,774,946 | 1,812,261 | 59,173 | 6,550,617 | 327,774 | 1,114,621 | 1,026,858 |
| Area XXIII: South..... | 49 | 72 | 247 | 200 | 300 | | 600 | | | |
| Total..... | 382 | 11,118 | 39,064 | 22,422,306 | 19,466,912 | 169,173 | 65,488,696 | 17,387,865 | 4,178,075 | 1,383,668 |
| Grand total..... | 390 | 12,214 | 52,651 | 38,826,872 | 35,893,910 | 697,659 | 96,804,550 | 33,072,662 | 5,802,443 | 1,396,843 |

¹ Exclusive of duplication.

² Incidental catch.

NOTE.—The weight of salted fish landed has been converted to the equivalent of fresh fish as landed. The roman numerals appearing in the stub of the above table refer to the numbers given these regions by the North American Council on Fishery Investigations.

Landings by fishing vessels at the 5 principal New England ports, 1938—Continued

SUMMARY: BY FISHING GROUNDS—Continued

| Fishing grounds | Pollock | Cusk | Halibut | Flounders | Swordfish | Mackerel | Herring | Other | Total |
|---------------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> | <i>Pounds</i> |
| Off Newfoundland: | | | | | | | | | |
| Area XVIII: Off Fuuks..... | | | 26,171 | | | | | | 26,171 |
| Area XIX: | | | | | | | | | |
| Bay of Islands..... | | | | | | | 1,055,241 | | 1,055,241 |
| Off Newfoundland (Treaty Coast) | | | | | | | | | 87,663 |
| Area XX: | | | | | | | | | |
| Grand Bank..... | | | 426,606 | | | | | | 597,764 |
| St. Peters Bank..... | | | 52,032 | | | | | | 72,297 |
| Total..... | | | 504,809 | | | | 1,055,241 | | 1,839,136 |
| Off Canada: | | | | | | | | | |
| Area XIX: Gulf of St. Lawrence..... | 470 | 1,367 | 144,372 | 32,170 | | 54,000 | | 385 | 5,218,217 |
| Area XXI: | | | | | | | | | |
| Quereau Bank..... | | 15,565 | 265,547 | | 4,371 | | | | 468,313 |
| The Gully..... | | | 51,196 | | 1,216 | | | | 66,375 |
| Sable Island Bank (Western Bank)..... | 1,493,595 | 88,635 | 278,173 | 430,785 | 2,536 | 10,500 | | 169,035 | 30,415,492 |
| Cape Shore..... | 145,370 | 450,910 | 15,794 | 31,075 | 449,541 | 13,980 | | 58,350 | 7,952,890 |
| Emerald Bank..... | 20,200 | 8,900 | 7,537 | 7,700 | | | | 7,800 | 844,282 |
| La Have Bank..... | 397,920 | 221,215 | 77,191 | 73,975 | 2,475 | | | 96,345 | 9,039,445 |
| Roseway Bank..... | 6,050 | 21,200 | 2,972 | | | | | 1,375 | 391,347 |
| Browns Bank..... | 1,607,281 | 1,131,060 | 258,265 | 234,310 | 9,281 | | | 522,565 | 37,249,263 |
| Total..... | 3,670,899 | 1,938,962 | 1,101,047 | 810,015 | 469,420 | 78,480 | | 855,855 | 91,645,624 |
| Off United States: | | | | | | | | | |
| Area XXII: | | | | | | | | | |
| Georges Bank..... | 3,077,410 | 388,585 | 423,542 | 3,183,275 | 1,265,780 | 46,980 | 3,455 | 724,749 | 90,850,098 |
| South Channel..... | 902,463 | 100,090 | 70,563 | 1,105,175 | 13,050 | 1,590 | 650 | 328,768 | 27,617,624 |
| Off Highland Light..... | 241,475 | 24,960 | 456 | 26,707 | | 300 | | 5,880 | 789,968 |
| Off Chatham..... | 65,485 | 3,150 | 1,909 | 147,030 | | 717,363 | | 111,395 | 3,636,913 |
| Nantucket Shoals..... | 12,090 | | 12,594 | 113,565 | 42,730 | 215,030 | | 4,655 | 1,006,994 |
| Cashes Bank..... | 39,905 | 173,190 | 2,578 | 3,775 | | | | 3,523 | 953,577 |
| Fippenies Bank..... | 18,935 | 35,175 | 3,139 | | | | | 275 | 368,521 |
| Platts Bank..... | 33,675 | 164,600 | 2,188 | | | | | 11,414 | 676,363 |
| Jeffreys Ledge..... | 53,210 | 147,680 | 2,107 | | | | | 18,913 | 982,728 |
| Tillies Bank..... | 100 | 350 | 15 | | | | | | 11,425 |
| Middle Bank (Stellwagen Bank)..... | 62,335 | 84,671 | 1,621 | 500 | | 329,775 | | 5,180 | 1,510,210 |
| Shore, general..... | 4,217,554 | 370,166 | 14,235 | 4,848,441 | 46,934 | 18,452,228 | 154,600 | 832,607 | 45,602,905 |
| Area XXIII: South..... | 50 | 300 | | | | 2,645,138 | | 6,700 | 2,653,288 |
| Total..... | 8,724,587 | 1,492,907 | 534,946 | 9,428,468 | 1,368,494 | 22,402,374 | 158,605 | 2,054,059 | 178,661,134 |
| Grand total..... | 12,395,473 | 3,431,869 | 2,140,802 | 10,238,483 | 1,837,914 | 22,480,854 | 1,213,846 | 2,909,914 | 270,145,894 |

Days' absence from port of fishing vessels landing fish at Boston and Gloucester, Mass., and Portland, Maine, 1933

| Fishing grounds | January | February | March | April | May | June | July | August | September | October | November | December | Total |
|----------------------------------|---------|----------|-------|-------|-------|-------|-------|--------|-----------|---------|----------|----------|--------|
| Off Newfoundland: | | | | | | | | | | | | | |
| Area XVIII: Off Funks | | | | | | | | 14 | | | | | 14 |
| Area XIX: | | | | | | | | | | | | 122 | 122 |
| Bay of Islands | | | | | | | | | | | | | |
| Off Newfoundland (Treaty Coast) | | | | | | | 7 | | | | | | 7 |
| Area XX: | | | | | | | | | | | | | |
| Grand Bank | | | 29 | 61 | 18 | 131 | 37 | 7 | | | | | 283 |
| St. Peters Bank | | 21 | | | | | | | | | 10 | | 31 |
| Total | | 21 | 29 | 61 | 18 | 131 | 44 | 21 | | | 10 | 122 | 457 |
| Off Canada: | | | | | | | | | | | | | |
| Area XIX: Gulf of St. Lawrence | | | | | 109 | 32 | 117 | 54 | 44 | 31 | 14 | | 401 |
| Area XXI: | | | | | | | | | | | | | |
| Quereau Bank | | 20 | 43 | | 69 | 24 | 21 | 62 | 73 | | | | 312 |
| The Gully | | | 21 | | | | | 14 | | | | | 35 |
| Sable Island Bank (Western Bank) | 106 | 29 | 96 | 126 | 279 | 56 | 56 | 96 | 507 | 493 | 851 | 1,031 | 3,728 |
| Cape Shore | 90 | 41 | 6 | 9 | 54 | 65 | 86 | 128 | 940 | 192 | 466 | 506 | 2,583 |
| Emerald Bank | 8 | | 8 | 12 | 13 | 8 | 22 | | 26 | | | | 31 |
| La Have Bank | 170 | 7 | 86 | 122 | 112 | 48 | 93 | 80 | 61 | 78 | 35 | 168 | 1,060 |
| Roseway Bank | 11 | 16 | | | | | | | 10 | | | | 17 |
| Browns Bank | 538 | 671 | 540 | 1,115 | 749 | 245 | 188 | 49 | 42 | 236 | 185 | 281 | 4,839 |
| Total | 923 | 784 | 800 | 1,384 | 1,385 | 478 | 583 | 485 | 1,667 | 1,066 | 1,551 | 2,034 | 13,140 |
| Off United States: | | | | | | | | | | | | | |
| Area XXII: | | | | | | | | | | | | | |
| Georges Bank | 1,344 | 1,400 | 1,429 | 557 | 565 | 1,435 | 2,906 | 2,497 | 1,053 | 1,281 | 739 | 358 | 15,564 |
| South Channel | 246 | 228 | 347 | 216 | 480 | 658 | 290 | 671 | 587 | 216 | 361 | 88 | 4,388 |
| Off Highland Light | 53 | 27 | 8 | 5 | | 8 | 12 | | 6 | 4 | 41 | 63 | 227 |
| Off Chatham | 87 | 33 | 44 | 40 | 77 | 46 | 281 | 24 | 26 | 59 | 15 | 27 | 759 |
| Nantucket Shoals | 10 | | 6 | 24 | 35 | 56 | 13 | 46 | 132 | 66 | 34 | | 423 |
| Cashes Bank | 13 | 3 | 12 | 30 | 33 | | | | 9 | 15 | 29 | 28 | 172 |
| Fippenies Bank | 22 | 14 | | 2 | | | | | 3 | 9 | 13 | 40 | 107 |
| Platts Bank | 17 | 26 | 13 | 9 | | | | | 16 | 25 | 29 | 16 | 151 |
| Jeffries Ledge | 63 | 61 | 41 | 8 | | | | | 12 | 38 | 54 | 48 | 315 |
| Tillies Bank | | | | | | | | | | | | | 12 |
| Middle Bank (Stellwagen Bank) | 45 | 57 | 21 | | | 57 | 2 | | 19 | 60 | 70 | | 82 |
| Shore, general | 976 | 838 | 1,037 | 1,328 | 1,551 | 1,519 | 1,459 | 1,679 | 1,521 | 1,927 | 1,524 | 909 | 16,268 |
| Area XXIII: South | | | | | 247 | | | | | | | | 247 |
| Total | 2,876 | 2,689 | 2,953 | 2,219 | 2,968 | 3,779 | 4,963 | 4,917 | 3,384 | 3,713 | 2,909 | 1,659 | 39,054 |
| Grand total | 3,799 | 3,494 | 3,787 | 3,664 | 4,391 | 4,388 | 5,590 | 5,423 | 5,051 | 4,779 | 4,470 | 3,815 | 62,661 |

NOTE.—The Roman numerals appearing in the stub of the above table refer to the numbers given these areas by the North American Council on Fishery Investigations.

MACKEREL FISHERY OF THE ATLANTIC COAST ⁵

That part of the 1933 mackerel catch taken by the purse seines and drift gill nets amounted to 29,532,600 pounds, a decrease of 37 per cent as compared with corresponding statistics for the previous year. The decrease resulted mainly from voluntary curtailment of the fishery on the part of the purse-seine fleet. It is estimated that normal exploitation of the fishery would have produced an increase rather than a decrease in total catch in 1933 as compared with 1932.

The statistics in the following table were obtained by combining the figures of mackerel landed at Boston and Gloucester, Mass., and Portland, Maine, with those obtained by agents stationed at other ports where mackerel are landed. The figures include approximately 1,000,000 pounds of mackerel that were brought to port but not landed due to lack of market. The statistics include the catches made by all purse seine and drift gill net vessels over 5 net tons (Customs measure) and also the catch of the larger of the boats of less than 5 net tons.

Data in the mackerel fishery are also included in the catch of States.

Mackerel fishery of the Atlantic coast, 1933

CATCH: BY AREAS IN 7-DAY PERIODS

| Date | Southern (Area XXIII) | | Block Island (Area XXII, west of Nantucket Shoals) | | Gulf of Maine (Area XXII north of Nantucket Shoals) | | Cape Shore (Area XXI) | Total |
|----------------------|-----------------------|---------|--|---------|---|-----------|-----------------------|------------|
| | Selmers | Netters | Selmers | Netters | Selmers | Netters | Selmers | |
| | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds |
| Apr. 9-15..... | 40,300 | 8,300 | | | | | | 48,600 |
| Apr. 16-22..... | 1,160,600 | 10,900 | | | | | | 1,171,500 |
| Apr. 23-29..... | 390,500 | 3,900 | | | | | | 394,400 |
| Apr. 30-May 6..... | 989,600 | 228,300 | | | | | | 1,217,900 |
| May 7-13..... | 238,900 | 92,600 | | | | | | 331,400 |
| May 14-20..... | 731,200 | 295,500 | 2,552,200 | | | | | 3,488,900 |
| May 21-27..... | | 25,600 | 1,348,400 | | 10,400 | | | 1,384,600 |
| May 28-June 3..... | | 40,900 | 229,100 | 12,500 | | 22,700 | | 305,200 |
| June 4-10..... | | | 1,064,300 | 12,600 | 16,100 | 97,400 | 1,200 | 1,191,600 |
| June 11-17..... | | | 484,800 | | 55,000 | 48,900 | | 588,700 |
| June 18-24..... | | | 519,300 | 3,000 | 97,600 | 41,100 | | 661,000 |
| June 25-July 1..... | | | 70,300 | | 479,500 | 20,900 | | 570,700 |
| July 2-8..... | | | 39,800 | | 401,600 | 28,500 | | 467,900 |
| July 9-15..... | | | | | 901,500 | 14,100 | | 915,600 |
| July 16-22..... | | | | | 799,700 | 24,800 | | 824,500 |
| July 23-29..... | | | | | 1,066,200 | 9,700 | | 1,095,900 |
| July 30-Aug. 5..... | | | | | 1,476,500 | 11,600 | | 1,488,100 |
| Aug. 6-12..... | | | | | 680,300 | 1,600 | | 681,900 |
| Aug. 13-19..... | | | | | 1,377,400 | 3,300 | | 1,380,700 |
| Aug. 20-26..... | | | | | 1,668,600 | 1,000 | | 1,669,600 |
| Aug. 27-Sept. 2..... | | | | | 1,472,400 | | | 1,472,400 |
| Sept. 3-9..... | | | | | 1,082,800 | | | 1,082,800 |
| Sept. 10-16..... | | | | | 299,500 | | | 299,500 |
| Sept. 17-23..... | | | | | 519,900 | 2,000 | | 521,900 |
| Sept. 24-30..... | | | | | 1,591,700 | | | 1,591,700 |
| Oct. 1-7..... | | | | | 329,900 | | | 329,900 |
| Oct. 8-14..... | | | | | 1,755,900 | 300 | | 1,756,200 |
| Oct. 15-21..... | | | | | 1,139,700 | 600 | | 1,140,300 |
| Oct. 22-28..... | | | | | 361,300 | 1,500 | | 362,800 |
| Oct. 29-Nov. 4..... | | | | | 131,900 | 22,000 | | 153,900 |
| Nov. 5-11..... | | | | | 8,100 | 32,300 | | 40,400 |
| Nov. 12-18..... | | | | | | 138,700 | | 138,700 |
| Nov. 19-25..... | | | | | | 159,400 | | 159,400 |
| Nov. 26-Dec. 2..... | | | | | | 420,600 | | 420,600 |
| Dec. 3-9..... | | | | | | 182,200 | | 182,200 |
| Dec. 10-16..... | | | | | | 1,000 | | 1,000 |
| Dec. 17-23..... | | | | | | 300 | | 300 |
| Total..... | 3,551,000 | 615,900 | 6,308,200 | 28,100 | 17,743,500 | 1,284,700 | 1,200 | 29,532,600 |

⁵ This section was prepared by O. E. Sette of the Division of Scientific Inquiry of the Bureau.

Mackerel fishery of the Atlantic coast, 1933—Continued

OPERATING UNITS AND CATCH: BY FLEET CLASSIFICATION AND GROUNDS

| Designation | Vessels and boats | Tonnage | Crew | Trips | Total catch |
|--------------------------------------|-------------------------|-----------------|---------------|---------------|-------------------|
| SOUTHERN—AREA XXIII | | | | | |
| Seiners: | <i>Number</i> | <i>Net tons</i> | <i>Number</i> | <i>Number</i> | <i>Pounds</i> |
| Regular vessels..... | 35 | 1,561 | 445 | 160 | 3,310,790 |
| Miscellaneous vessels..... | 9 | 289 | 110 | 13 | 240,300 |
| Netters: | | | | | |
| Regular vessels..... | 18 | 354 | 125 | 120 | 587,100 |
| Miscellaneous vessels..... | 1 | 8 | 5 | 3 | 17,000 |
| Miscellaneous boats..... | 2 | | | 4 | 11,800 |
| Total..... | 1 63 | 2,212 | 685 | 300 | 4,166,990 |
| BLOCK ISLAND—AREA XXII | | | | | |
| (West of Nantucket Shoals only) | | | | | |
| Seiners: | | | | | |
| Regular vessels..... | 57 | 2,303 | 715 | 256 | 6,174,400 |
| Miscellaneous vessels..... | 5 | 102 | 50 | 8 | 133,000 |
| Netters: Regular vessels..... | 3 | 32 | 17 | 6 | 28,100 |
| Total..... | 1 65 | 2,437 | 782 | 270 | 6,336,200 |
| GULF OF MAINE—AREA XXII | | | | | |
| (North of Nantucket Shoals only) | | | | | |
| Seiners: | | | | | |
| Regular vessels..... | 61 | 2,057 | 715 | 1,160 | 15,649,500 |
| Miscellaneous vessels..... | 26 | 482 | 216 | 165 | 1,318,500 |
| Miscellaneous boats..... | 21 | | | 114 | 775,500 |
| Netters: | | | | | |
| Spring and summer: | | | | | |
| Miscellaneous vessels..... | 13 | 179 | 78 | 44 | 86,200 |
| Miscellaneous boats..... | 48 | | | 239 | 237,600 |
| Fall: | | | | | |
| Regular vessels..... | 34 | 856 | 262 | 287 | 877,400 |
| Miscellaneous vessels..... | 20 | 372 | 134 | 38 | 53,700 |
| Miscellaneous boats..... | 13 | | | 32 | 30,100 |
| Total..... | 1 154 | 3,946 | 1,465 | 2,079 | 19,028,200 |
| CAPE SHORE—AREA XXI | | | | | |
| Seiners..... | 1 | 67 | 14 | 1 | 1,200 |
| Total seiners..... | 1 63 | | | 1,877 | 27,603,900 |
| Total netters..... | 1 72 | | | 773 | 1,928,700 |
| Grand total..... | 1 129 | | | 2,650 | 29,532,600 |

¹ Exclusive of duplication and of boats.

² Of this total approximately 21,430,000 pounds were tinkers (under 1½ pounds each) and 8,102,000 pounds were of larger sizes. There were no bullseye mackerel landed by the fleet.

NOTE.—The roman numerals appearing in the stub of the above table refer to the numbers given these areas by the North American Council on Fishery Investigations.

FISHERIES OF THE MIDDLE ATLANTIC STATES

(Area XXIII) ⁶

The yield of the commercial fisheries of the Middle Atlantic States (New York, New Jersey, Pennsylvania, and Delaware) during 1933 amounted to 169,753,735 pounds, valued at \$4,811,055 to the fishermen, representing an increase of 20 percent in volume and 3 percent in value as compared with the catch in the previous year. These fisheries gave employment to 8,574 fishermen, as compared with 8,370 in 1932.

There were 398 fishery wholesale and manufacturing establishments in the 4 States in 1933 as compared with 418 in 1931 when the most recent previous survey of such concerns was made. In 1933 these establishments employed 5,631 persons, paid \$6,085,981 in salaries and wages, and produced manufactured products (canned, cured, packaged, and byproducts), valued at \$11,219,966. In 1931 the wholesale and manufacturing firms employed 4,989 persons, paid \$7,042,586 in salaries and wages, and produced manufactured products valued at \$12,451,810.

Fisheries of the Middle Atlantic States, 1933

SUMMARY OF CATCH

| Product | New York | | New Jersey | | Pennsylvania | |
|---------------------|------------|-----------|------------|-------------|--------------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Fish..... | 28,305,152 | \$988,984 | 79,038,724 | \$1,138,015 | 52,232 | \$2,090 |
| Shellfish, etc..... | 11,606,280 | 1,463,581 | 14,224,713 | 1,008,309 | ----- | ----- |
| Total..... | 39,911,432 | 2,452,565 | 93,263,437 | 2,146,324 | 52,232 | 2,090 |

| Product | Delaware | | Total | |
|---------------------|------------|-----------|-------------|-------------|
| | Pounds | Value | Pounds | Value |
| Fish..... | 35,375,390 | \$138,191 | 142,771,488 | \$2,266,170 |
| Shellfish, etc..... | 1,151,254 | 72,995 | 26,982,247 | 2,544,685 |
| Total..... | 36,526,634 | 209,186 | 169,753,735 | 4,811,055 |

OPERATING UNITS: BY STATES

| Item | New York | New Jersey | Pennsylvania | Delaware | Total |
|---------------------|----------|------------|--------------|----------|--------|
| | Number | Number | Number | Number | Number |
| Fishermen: | | | | | |
| On vessels..... | 825 | 1,267 | ----- | 350 | 2,442 |
| On boats and shore: | | | | | |
| Regular..... | 1,509 | 954 | ----- | 34 | 2,497 |
| Casual..... | 1,482 | 1,622 | 53 | 478 | 3,635 |
| Total..... | 3,816 | 3,843 | 53 | 862 | 8,574 |
| Vessels: | | | | | |
| Steam..... | 9 | ----- | ----- | 10 | 19 |
| Net tonnage..... | 1,773 | ----- | ----- | 1,237 | 3,010 |
| Motor..... | 160 | 208 | ----- | 16 | 384 |
| Net tonnage..... | 2,422 | 3,470 | ----- | 229 | 6,121 |

⁶ This is the number given to this area by the North American Council on Fishery Investigations. It should be explained that there are included in this area craft whose principal fishing ports are in the area but at times fish elsewhere. A notable example is the southern trawl fishery which extends into Area XXIV. It should be observed that the persons engaged, gear and craft employed, and catch of the seed oyster fishery are not included among the statistics of the fishery for market oysters and other species but are shown in separate tables in this section. For a clearer understanding of the statistics published in this section, the reader is referred to the section in the latter part of this document entitled "Statistical survey procedure."

Fisheries of the Middle Atlantic States, 1933—Continued

OPERATING UNITS: BY STATES—Continued

| Item | New York | New Jersey | Pennsylvania | Delaware | Total |
|--------------------------------|----------|------------|--------------|----------|-----------|
| | Number | Number | Number | Number | Number |
| Vessels—Continued. | | | | | |
| Sail..... | | 3 | | 1 | 4 |
| Net tonnage..... | | 24 | | 8 | 32 |
| Total vessels..... | 169 | 211 | | 27 | 407 |
| Total net tonnage..... | 4,195 | 3,494 | | 1,474 | 9,163 |
| Boats: | | | | | |
| Motor..... | 636 | 1,029 | 5 | 77 | 1,747 |
| Other..... | 1,188 | 740 | 14 | 181 | 2,123 |
| Accessory boats..... | 39 | 88 | | 30 | 157 |
| Apparatus: | | | | | |
| Purse seines: | | | | | |
| Menhaden..... | | 8 | | 10 | 18 |
| Length, yards..... | | 2,806 | | 3,140 | 5,946 |
| Other..... | 2 | 7 | | | 9 |
| Length, yards..... | 800 | 2,510 | | | 3,310 |
| Haul seines..... | 71 | 105 | 12 | 61 | 249 |
| Length, yards..... | 5,926 | 9,040 | 1,635 | 18,990 | 35,591 |
| Gill nets: | | | | | |
| Anchor..... | 46 | 10 | | | 56 |
| Square yards..... | 40,832 | 5,830 | | | 46,662 |
| Drift..... | 254 | 605 | 7 | 63 | 929 |
| Square yards..... | 423,369 | 448,333 | 5,200 | 145,720 | 1,022,622 |
| Runaround..... | 43 | 69 | | 33 | 145 |
| Square yards..... | 124,822 | 210,695 | | 41,450 | 376,967 |
| Stake..... | 49 | 251 | | 99 | 399 |
| Square yards..... | 23,161 | 46,882 | | 7,770 | 77,813 |
| Lines: | | | | | |
| Hand..... | 225 | 463 | | 12 | 700 |
| Hooks..... | 441 | 720 | | 24 | 1,185 |
| Trawl..... | 693 | 712 | | 2 | 1,407 |
| Hooks..... | 187,600 | 420,400 | | 1,300 | 609,300 |
| Troll..... | 20 | 404 | | | 424 |
| Hooks..... | 20 | 404 | | | 424 |
| Trot with baits or snoods..... | | 9 | | | 9 |
| Baits or snoods..... | | 9,900 | | | 9,900 |
| Trot with hooks..... | 14 | | | | 14 |
| Hooks..... | 1,495 | | | | 1,495 |
| Pound nets..... | 301 | 153 | | 32 | 486 |
| Weirs..... | | 88 | | | 88 |
| Stop nets..... | 24 | 65 | | 7 | 96 |
| Square yards..... | 11,258 | 50,615 | | 1,820 | 63,693 |
| Fyke nets..... | 849 | 1,369 | | 449 | 2,667 |
| Dip nets..... | 5 | 34 | | 51 | 90 |
| Cast nets..... | | 3 | | | 3 |
| Scap nets..... | 279 | | | | 279 |
| Drag nets..... | 47 | 14 | | | 61 |
| Yards at mouth..... | 94 | 28 | | | 122 |
| Otter trawls..... | 130 | 67 | | | 197 |
| Yards at mouth..... | 2,791 | 1,514 | | | 4,305 |
| Wire baskets..... | | 12 | | | 12 |
| Pots: | | | | | |
| Crab..... | | 10 | | | 10 |
| Eel..... | 3,877 | 2,571 | | 876 | 7,324 |
| Lobster..... | 18,340 | 28,071 | | 165 | 46,576 |
| Harpoons..... | 18 | 1 | | | 19 |
| Spears..... | 77 | 92 | | | 169 |
| Dredges: | | | | | |
| Clam..... | 14 | 43 | | 33 | 90 |
| Yards at mouth..... | 12 | 48 | | 47 | 107 |
| Crab..... | 6 | 69 | | 11 | 86 |
| Yards at mouth..... | 14 | 90 | | 20 | 124 |
| Mussel..... | 2 | | | | 2 |
| Yards at mouth..... | 2 | | | | 2 |
| Oyster..... | 78 | 259 | | 16 | 353 |
| Yards at mouth..... | 115 | 309 | | 23 | 447 |
| Scallop..... | 250 | 10 | | | 260 |
| Yards at mouth..... | 365 | 34 | | | 399 |
| Tongs..... | 837 | 783 | | 32 | 1,652 |
| Rakes..... | 568 | 509 | | | 1,077 |
| Forks..... | 271 | 51 | | | 322 |
| Hoes..... | | 176 | | | 176 |
| Gaffs..... | | 2 | | | 2 |

Fisheries of the Middle Atlantic States, 1933—Continued

CATCH: BY STATES

| Species | New York | | New Jersey | | Pennsylvania | | Delaware | | Total | |
|----------------------------------|-------------|----------|--------------|----------|--------------|-------|--------------|----------|--------------|-----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | | | |
| Alewives..... | 218, 257 | \$5, 023 | 13, 990 | \$244 | 8, 775 | \$88 | 1, 148, 980 | \$4, 786 | 1, 390, 002 | \$10, 141 |
| Bluefish..... | 1, 131, 721 | 56, 097 | 2, 115, 304 | 76, 146 | | | 4, 720 | 236 | 3, 251, 745 | 132, 479 |
| Bonito..... | 159, 007 | 9, 541 | 88, 407 | 4, 103 | | | | | 247, 414 | 13, 644 |
| Butterfish..... | 1, 498, 212 | 61, 339 | 2, 912, 214 | 119, 319 | | | | | 4, 410, 426 | 180, 658 |
| Carp..... | 219, 497 | 17, 574 | 143, 471 | 16, 047 | 640 | 64 | 26, 420 | 1, 589 | 390, 428 | 35, 274 |
| Catfish and bullheads..... | 19, 008 | 2, 561 | 55, 380 | 3, 321 | | | 1, 845 | 116 | 76, 238 | 5, 998 |
| Cero..... | | | 2, 500 | 75 | | | | | 2, 500 | 75 |
| Cod..... | 4, 260, 669 | 112, 583 | 3, 229, 886 | 99, 688 | | | 2, 240 | 69 | 7, 492, 795 | 212, 340 |
| Crevalle..... | | | 7, 254 | 141 | | | | | 7, 254 | 141 |
| Croaker..... | 80, 147 | 1, 531 | 1, 901, 073 | 44, 945 | | | | | 67, 130 | 1, 068 |
| Cusk..... | 3, 500 | 99 | | | | | | | 2, 048, 350 | 47, 544 |
| Dolphin..... | | | 111 | 9 | | | | | 3, 500 | 99 |
| Drum: | | | | | | | | | 111 | 9 |
| Black..... | | | | | | | | | | |
| Red or redfish..... | 447 | 9 | | | | | | | 447 | 9 |
| Eels: | | | 8, 770 | 126 | | | 520 | 16 | 9, 290 | 142 |
| Common..... | 290, 470 | 27, 446 | 336, 466 | 28, 565 | | | | | 680, 076 | 59, 903 |
| Conger..... | 5, 260 | 257 | 11, 163 | 432 | | | 53, 140 | 3, 892 | 16, 423 | 689 |
| Flounders..... | 5, 191, 128 | 192, 032 | 4, 051, 797 | 170, 601 | | | | | 9, 252, 375 | 363, 125 |
| Frigate mackerel..... | 81, 886 | 1, 234 | 19, 754 | 243 | | | 9, 450 | 492 | 101, 640 | 1, 477 |
| Goosefish..... | | | 10, 000 | 38 | | | | | 10, 000 | 38 |
| Grayfish..... | | | 6, 739 | 135 | | | | | 6, 739 | 135 |
| Haddock..... | 8, 506, 679 | 248, 282 | | | | | | | 8, 506, 679 | 248, 282 |
| Hake..... | 139, 954 | 2, 811 | 22, 171 | 360 | | | | | 162, 125 | 3, 171 |
| Halibut..... | 52, 508 | 7, 280 | | | | | | | 52, 508 | 7, 280 |
| Herring: | | | | | | | | | | |
| Round..... | | | 750 | 8 | | | | | 750 | 8 |
| Sea..... | 74, 987 | 610 | 590, 465 | 5, 171 | | | | | 665, 452 | 5, 781 |
| Hickory shad..... | 1, 300 | 25 | | | | | | | 1, 300 | 25 |
| Kingfish or "king mackerel"..... | | | 140 | 5 | | | | | 140 | 5 |
| King whiting or "kingfish"..... | 73, 223 | 4, 057 | 84, 039 | 10, 932 | | | | | 157, 262 | 14, 989 |
| Mackerel..... | 343, 717 | 12, 353 | 318, 047 | 9, 692 | | | | | 661, 764 | 22, 045 |
| Menhaden..... | 201, 176 | 938 | 45, 774, 117 | 122, 929 | | | 33, 600, 000 | 111, 700 | 79, 575, 293 | 235, 567 |
| Mullet..... | 2, 500 | 96 | 82, 747 | 6, 234 | | | 233, 000 | 2, 390 | 318, 247 | 8, 720 |
| Mummichog..... | 25, 050 | 2, 505 | 25, 925 | 1, 939 | | | | | 50, 975 | 4, 444 |
| Pigfish..... | | | 208 | 5 | | | | | 203 | 5 |
| Pike or pickerel..... | 420 | 53 | | | | | | | 420 | 53 |
| Pollock..... | 770, 231 | 11, 551 | 6, 280 | 94 | | | | | 776, 511 | 11, 645 |
| Pompano..... | | | 317 | 77 | | | | | 317 | 77 |
| Scup or porgy..... | 1, 478, 153 | 26, 049 | 4, 881, 364 | 65, 293 | | | | | 6, 359, 517 | 91, 942 |
| Sea bass..... | 505, 688 | 23, 068 | 1, 969, 143 | 63, 601 | | | | | 2, 474, 831 | 87, 269 |

| | | | | | | | | | | |
|----------------------------------|-------------------|----------------|-------------------|------------------|---------------|--------------|-------------------|----------------|--------------------|------------------|
| Sea robin..... | 18,758 | 276 | 10,805 | 109 | | | | | 29,563 | 385 |
| Sbad..... | 351,786 | 28,761 | 457,458 | 45,181 | 2,217 | 392 | 22,325 | 1,993 | 833,786 | 76,327 |
| Sharks..... | 2,500 | 35 | 2,500 | 203 | | | | | 12,110 | 238 |
| Sheepshead..... | | | 9 | 1 | | | | | 9 | 1 |
| Silversides..... | 18,500 | 1,838 | 4,830 | 676 | | | | | 23,330 | 2,514 |
| Skates..... | 725 | 7 | 82,976 | 712 | | | | | 83,701 | 719 |
| Smelts..... | 245 | 47 | | | | | | | 245 | 47 |
| Spanish mackerel..... | | | 6,419 | 287 | | | | | 6,419 | 287 |
| Spot..... | 22,765 | 463 | 473,565 | 6,958 | | | 31,920 | 903 | 528,250 | 8,324 |
| Squeteagues or "sea trout": | | | | | | | | | | |
| Gray..... | 823,634 | 38,147 | 6,927,381 | 185,597 | | | 123,180 | 3,341 | 7,874,195 | 227,085 |
| Spotted..... | | | 30 | 2 | | | | | 30 | 2 |
| Squirrel hake..... | | | 52,765 | 402 | | | | | 52,765 | 402 |
| Striped bass..... | 18,954 | 2,724 | 9,122 | 1,770 | | | 11,700 | 1,382 | 39,776 | 5,876 |
| Sturgeon..... | 1,851 | 311 | 26,713 | 2,130 | | | 1,200 | 300 | 29,764 | 2,741 |
| Suckers..... | 28,624 | 2,365 | 102,575 | 8,462 | 40,600 | 2,436 | | | 171,799 | 13,263 |
| Sunfish..... | 1,690 | 132 | | | | | | | 1,690 | 132 |
| Swellfish..... | 2,000 | 200 | | | | | | | 2,000 | 200 |
| Swordfish..... | 100,966 | 12,441 | 1,600 | 128 | | | | | 102,566 | 12,589 |
| Tautog..... | 61,158 | 1,584 | 70,923 | 10,048 | | | | | 132,081 | 11,632 |
| Thimble-eyed mackerel..... | | | 24,475 | 374 | | | | | 25,475 | 374 |
| Tilfish..... | 1,350,000 | 67,500 | 25 | 1 | | | | | 1,350,025 | 67,501 |
| Tomcod..... | 7,690 | 338 | | | | | | | 7,690 | 338 |
| Tuna or "horse mackerel"..... | 6,786 | 270 | 36,492 | 1,656 | | | | | 43,228 | 1,926 |
| Whitebait..... | 3,000 | 450 | | | | | | | 3,000 | 450 |
| White perch..... | 15,182 | 1,119 | 29,420 | 2,666 | | | 30,650 | 1,490 | 75,252 | 5,275 |
| Whiting..... | 106,111 | 1,023 | 2,040,544 | 20,134 | | | 480 | 30 | 2,147,135 | 21,192 |
| Wolfish..... | 25,625 | 576 | | | | | | | 25,625 | 576 |
| Yellow perch..... | 1,857 | 168 | | | | | 6,490 | 398 | 8,337 | 566 |
| Total..... | 28,306,152 | 988,984 | 79,038,724 | 1,138,015 | 52,232 | 2,980 | 35,375,380 | 136,191 | 142,771,488 | 2,266,170 |
| SHELLFISH, ETC. | | | | | | | | | | |
| Crabs: | | | | | | | | | | |
| Hard..... | 149,383 | 5,437 | 709,410 | 18,356 | | | 96,600 | 1,295 | 955,393 | 25,088 |
| King..... | | | 2,183,772 | 6,631 | | | 556,600 | 595 | 2,750,372 | 6,526 |
| Soft..... | 825 | 225 | 57,363 | 12,239 | | | 36,000 | 7,200 | 94,188 | 19,664 |
| Lobsters..... | 346,496 | 67,256 | 364,220 | 67,629 | | | 12,840 | 2,568 | 723,556 | 137,453 |
| Shrimp..... | 69,912 | 11,981 | 25,741 | 3,328 | | | | | 85,653 | 15,309 |
| Squid..... | 444,938 | 6,696 | 347,312 | 8,760 | | | | | 792,250 | 15,456 |
| Clams: | | | | | | | | | | |
| Hard, public ¹ | 1,923,610 | 271,295 | 1,405,204 | 215,579 | | | 191,300 | 30,310 | 3,520,114 | 517,194 |
| Hard, private ¹ | 21,040 | 4,601 | 70,461 | 8,721 | | | 9,640 | 1,898 | 101,141 | 15,220 |
| Soft, public ¹ | 222,150 | 18,760 | 913,560 | 46,752 | | | | | 1,135,710 | 64,512 |
| Surf or skimmer..... | 379,500 | 18,380 | 146,313 | 5,732 | | | | | 525,813 | 24,112 |
| Conchs..... | 35,000 | 2,065 | 54 | 2 | | | | | 35,054 | 2,097 |
| Mussels, sea..... | 47,400 | 3,150 | 5,200 | 200 | | | | | 62,600 | 3,350 |

¹Statistics on hard clams are based on yields of 8 pounds of meats to the bushel in New York, 8.96 in New Jersey, and 10 pounds in Delaware.

²Statistics on soft clams used in this table are based on yields of 16 pounds of meats to the bushel in New York and 20 pounds in New Jersey.

Fisheries of the Middle Atlantic States, 1933—Continued

CATCH: BY STATES—Continued

| Species | New York | | New Jersey | | Pennsylvania | | Delaware | | Total | |
|------------------------------|------------|-----------|------------|-----------|--------------|---------|------------|---------|-------------|-----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| SHELLFISH, ETC.—continued | | | | | | | | | | |
| Oysters: ¹ | | | | | | | | | | |
| Market, public, spring..... | 19,250 | \$2,662 | 10,966 | \$1,138 | ----- | ----- | 39,600 | \$5,140 | 69,816 | \$8,940 |
| Market, public, fall..... | 31,994 | 9,512 | 26,606 | 1,792 | ----- | ----- | ----- | ----- | 108,590 | 11,304 |
| Market, private, spring..... | 2,761,525 | 381,138 | 3,718,539 | 282,883 | ----- | ----- | 24,520 | 2,985 | 6,504,584 | 667,006 |
| Market, private, fall..... | 3,229,025 | 439,378 | 3,855,564 | 301,288 | ----- | ----- | 165,240 | 20,065 | 7,249,829 | 760,731 |
| Periwinkles and cockles..... | ----- | ----- | 100 | 2 | ----- | ----- | ----- | ----- | 100 | 2 |
| Scallops: | | | | | | | | | | |
| Bay..... | 48,000 | 17,100 | 594 | 30 | ----- | ----- | ----- | ----- | 48,594 | 17,130 |
| Sea..... | 1,823,492 | 193,445 | 373,589 | 22,402 | ----- | ----- | ----- | ----- | 2,197,081 | 215,847 |
| Terrapin, diamond-back..... | ----- | ----- | 502 | 145 | ----- | ----- | 410 | 290 | 912 | 405 |
| Turtles, snapper..... | ----- | ----- | 2,500 | 250 | ----- | ----- | 8,504 | 679 | 11,004 | 929 |
| Bloodworms..... | 9,250 | 8,390 | 1,740 | 1,505 | ----- | ----- | ----- | ----- | 10,990 | 9,895 |
| Sandworms..... | 3,500 | 2,060 | 5,403 | 4,645 | ----- | ----- | ----- | ----- | 8,903 | 6,725 |
| Total..... | 11,606,280 | 1,463,581 | 14,224,713 | 1,008,309 | ----- | ----- | 1,151,254 | 72,995 | 26,982,247 | 2,544,885 |
| Grand total..... | 39,911,432 | 2,452,565 | 93,263,437 | 2,146,324 | 52,232 | \$2,980 | 36,526,634 | 209,186 | 169,753,735 | 4,811,055 |

¹ Statistics on oysters used in this table are based on yields of 7 pounds of meats to the bushel in New York, 8.70 in New Jersey, and 7.88 in Delaware.

NOTE.—Of the total catch in New Jersey, 7,500 pounds of shad, valued at \$600, were caught in the St. Johns River in Florida, and 1,447,252 pounds of fishery products, valued at \$47,109, were taken in the southern trawl fishery off southern New Jersey, Maryland, Virginia, and North Carolina. Of the total catch in New York, 1,024,291 pounds of fishery products, valued at \$35,928, were taken in the southern trawl fishery. The products of the southern trawl fishery consisted principally of croaker, flounders, scup, and sea bass.

Fisheries of the Middle Atlantic States, 1933—Continued

PRODUCTION OF CERTAIN SHELLFISH IN NUMBER AND BUSHEL

| Product | New York | | New Jersey | | Delaware | | Total | |
|-----------------------------------|----------|---------|------------|----------|----------|---------|-----------|----------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| Crabs: | | | | | | | | |
| Hard.....number.. | 448,149 | \$5,437 | 2,128,230 | \$18,356 | 289,800 | \$1,295 | 2,866,179 | \$25,088 |
| King.....do..... | | | 546,943 | 5,931 | 141,650 | 595 | 687,563 | 6,526 |
| Soft.....do..... | 3,300 | 225 | 172,089 | 12,239 | 144,000 | 7,200 | 319,389 | 19,664 |
| Clams: | | | | | | | | |
| Hard, public...bushels.. | 240,451 | 271,295 | 156,831 | 215,579 | 19,130 | 30,310 | 416,412 | 517,184 |
| Hard, private...do..... | 2,630 | 4,601 | 7,864 | 8,721 | 964 | 1,898 | 11,453 | 15,220 |
| Soft, public...do..... | 13,884 | 18,760 | 45,678 | 45,752 | | | 59,562 | 64,512 |
| Surf or skimmer...do... | 31,625 | 18,380 | 11,705 | 5,732 | | | 43,330 | 24,112 |
| Conchs.....do..... | 1,944 | 2,095 | 3 | 2 | | | 1,947 | 2,097 |
| Mussels, sea...do..... | 4,740 | 3,150 | 400 | 200 | | | 5,140 | 3,350 |
| Oysters: | | | | | | | | |
| Market, public, spring bushels.. | 2,750 | 2,662 | 1,260 | 1,138 | 5,025 | 5,140 | 9,035 | 8,940 |
| Market, public, fall bushels.. | 11,712 | 9,512 | 8,058 | 1,792 | | | 14,770 | 11,304 |
| Market, private, spring bushels.. | 394,504 | 381,138 | 427,418 | 282,883 | 3,112 | 2,985 | 825,034 | 667,006 |
| Market, private, fall bushels.. | 461,289 | 439,378 | 443,168 | 301,288 | 20,969 | 20,065 | 925,426 | 760,731 |
| Periwinkles and cockles bushels.. | | | 8 | 2 | | | 8 | 2 |
| Scallops: | | | | | | | | |
| Bay.....do..... | 9,600 | 17,100 | 99 | 30 | | | 9,699 | 17,130 |
| Sea.....do..... | 303,915 | 193,445 | 92,265 | 22,402 | | | 396,180 | 215,847 |

SEED OYSTER FISHERY

| Item | New York | | New Jersey | | Delaware | | Total | |
|----------------------------|----------------|--------------|----------------|--------------|----------------|--------------|----------------|--------------|
| OPERATING UNITS | | | | | | | | |
| Fishermen: | <i>Number</i> | | <i>Number</i> | | <i>Number</i> | | <i>Number</i> | |
| On vessels..... | 24 | | 1,514 | | 48 | | 1,586 | |
| On boats and shore: | | | | | | | | |
| Regular..... | 230 | | 68 | | | | 298 | |
| Casual..... | | | 107 | | 43 | | 150 | |
| Total..... | 254 | | 1,689 | | 91 | | 2,034 | |
| Vessels: | | | | | | | | |
| Motor..... | 7 | | | | | | 7 | |
| Net tonnage..... | 69 | | | | | | 69 | |
| Sail..... | | | 139 | | 7 | | 146 | |
| Net tonnage..... | | | 2,881 | | 96 | | 2,977 | |
| Total vessels..... | 7 | | 139 | | 7 | | 153 | |
| Total net tonnage..... | 69 | | 2,881 | | 96 | | 3,046 | |
| Boats: | | | | | | | | |
| Motor..... | 120 | | 86 | | | | 206 | |
| Other..... | 30 | | 81 | | 41 | | 152 | |
| Apparatus: | | | | | | | | |
| Dredges..... | 6 | | 278 | | 16 | | 300 | |
| Yards at mouth..... | 9 | | 333 | | 21 | | 363 | |
| Tongs..... | 194 | | 140 | | 40 | | 374 | |
| Rakes..... | 45 | | 34 | | | | 79 | |
| CATCH | | | | | | | | |
| Oysters: | <i>Bushels</i> | <i>Value</i> | <i>Bushels</i> | <i>Value</i> | <i>Bushels</i> | <i>Value</i> | <i>Bushels</i> | <i>Value</i> |
| Seed, public, spring..... | 42,400 | \$20,550 | 926,000 | \$282,582 | 34,940 | \$8,730 | 1,003,510 | \$201,862 |
| Seed, public, fall..... | 12,500 | 5,500 | 11,000 | 2,750 | | | 23,500 | 8,250 |
| Seed, private, spring..... | 40,290 | 39,540 | | | | | 40,290 | 39,540 |
| Seed, private, fall..... | 38,290 | 38,290 | | | | | 38,290 | 38,290 |
| Total..... | 133,680 | 103,880 | 937,000 | 235,332 | 34,940 | 8,730 | 1,106,620 | 347,942 |

NOTE.—Of the total number of persons fishing for seed oysters, 1,881 are duplicated among those fishing for market oysters or other species. Similarly the following craft and gear are duplicated: 87 vessels, 79 motor boats, 92 other boats, 172 dredges, 327 tongs, and 15 rakes.

Industries related to the fisheries of the Middle Atlantic States, 1933

OPERATING UNITS, SALARIES, AND WAGES

| Item | New York | New Jersey | Pennsylvania | Delaware | Total |
|--------------------------------------|--------------------|--------------------|------------------|------------------|--------------------|
| Transporting: | | | | | |
| Persons engaged: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| On vessels..... | 100 | 10 | | | 110 |
| On boats..... | | 92 | | | 92 |
| Total..... | 100 | 102 | | | 202 |
| Vessels: | | | | | |
| Motor..... | 26 | 4 | | | 30 |
| Net tonnage..... | 580 | 49 | | | 629 |
| Boats..... | | 87 | | | 87 |
| Wholesale and manufacturing: | | | | | |
| Establishments..... | 210 | 124 | 49 | 15 | 398 |
| Persons engaged: | | | | | |
| Proprietors..... | 184 | 116 | 44 | 14 | 358 |
| Salaried employees..... | 775 | 159 | 113 | 20 | 1,067 |
| Wage earners: | | | | | |
| A verage for season..... | 1,806 | 1,381 | 477 | 452 | 4,206 |
| A verage for year..... | 1,457 | 906 | 396 | 189 | 2,948 |
| Paid to salaried employees..... | \$2,164,542 | \$311,040 | \$200,846 | \$12,460 | \$2,688,888 |
| Paid to wage earners..... | \$2,088,832 | \$785,657 | \$403,433 | \$119,171 | \$3,397,093 |
| Total salaries and wages..... | \$4,253,374 | \$1,096,697 | \$604,279 | \$131,631 | \$6,085,981 |
| Fishermen manufacturing..... | 348 | 157 | | 7 | 512 |

PRODUCTS MANUFACTURED

| Item | New York | | New Jersey | | Pennsylvania | | Delaware | |
|--|-----------|-----------|------------|----------|--------------|----------|----------|-------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| By manufacturing establishments: | | | | | | | | |
| Buffalo fish, smoked | 440,000 | \$131,750 | | | | | | |
| Butterfish, smoked, do..... | 408,600 | 100,050 | 80,806 | \$20,269 | (1) | (1) | | |
| Carp, smoked.....do..... | 91,000 | 27,200 | (1) | (1) | | | | |
| Chubs, elcso, lake herring, and tullibee, smoked | | | | | | | | |
| pounds..... | 2,495,880 | 534,914 | (1) | (1) | 333,000 | \$93,000 | | |
| Cod: | | | | | | | | |
| Fresh fillets.....do..... | 2,365,354 | 306,896 | | | | | | |
| Smoked fillets.....do..... | 66,305 | 13,700 | (1) | (1) | | | | |
| Flounders, fresh fillets | | | | | | | | |
| pounds..... | 669,780 | 109,678 | | | | | | |
| Haddock, fresh fillets | | | | | | | | |
| pounds..... | 1,500,661 | 207,200 | | | | | | |
| Hake, fresh fillets.....do..... | 297,000 | 31,440 | | | | | | |
| Herring, sea: | | | | | | | | |
| Kipperd.....do..... | 55,100 | 10,210 | (1) | (1) | (2) | (2) | | |
| Bloaters.....do..... | (1) | (1) | (1) | (1) | 540,000 | 30,250 | | |
| Lake trout, smoked, do..... | 90,400 | 28,650 | (1) | (1) | (1) | (1) | | |
| Mackerel, smoked.....do..... | 77,300 | 13,602 | (1) | (1) | (1) | (1) | | |
| Paddlefish or spoonbill cat, smoked.....pounds..... | 338,600 | 155,920 | | | | | | |
| Salmon: | | | | | | | | |
| Kipperd.....do..... | 269,000 | 82,700 | (2) | (2) | (1) | (1) | | |
| Smoked.....do..... | 5,427,102 | 1,494,924 | 478,966 | 130,804 | (1) | (1) | | |
| Shad, smoked.....do..... | 149,000 | 30,400 | (1) | (1) | (1) | (1) | | |
| Sturgeon, smoked.....do..... | 1,684,000 | 548,600 | 905,987 | 149,781 | (1) | (1) | | |
| Sturgeon, caviar, canned standard cases..... | 2,596 | 333,530 | | | | | | |
| Whitefish, smoked | | | | | | | | |
| pounds..... | 1,104,000 | 297,760 | (1) | (1) | 340,000 | 85,200 | | |
| Whitefish roe and caviar, canned standard cases..... | 122 | 4,569 | | | | | | |
| Crabs, king, scrap and meal.....tons..... | | | 278 | 8,150 | | | | |
| Clams: | | | | | | | | |
| Hard, fresh-shucked | | | | | | | | |
| gallons..... | | | 1,875 | 2,723 | 4,110 | 14,495 | | |
| Soft, fresh-shucked | | | | | | | | |
| gallons..... | | | 16,360 | 14,902 | | | | |

¹ The production of this item is included under "Unclassified products."

² Kipperd sea herring is included with bloaters.

³ Kipperd salmon is included with smoked salmon.

Industries related to the fisheries of the Middle Atlantic States, 1933—Continued

PRODUCTS MANUFACTURED—Continued

| Item | New York | | New Jersey | | Pennsylvania | | Delaware | |
|---|----------|-------------|-------------|---------------|---------------|-------------|----------|-----------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| By manufacturing establishments—Continued. | | | | | | | | |
| Marine-shell products: | | | | | | | | |
| Buttons—gross | 826, 705 | \$509, 920 | 1, 617, 128 | \$1, 073, 275 | 1, 143, 630 | \$589, 743 | | |
| Novelties | | 46, 813 | | 81, 255 | | 45, 739 | | |
| Oysters, fresh-shucked | | | | | | | | |
| gallons | 265, 967 | 469, 016 | 372, 848 | 577, 217 | 60, 958 | 139, 056 | 39, 671 | \$48, 434 |
| Oyster-shell products: | | | | | | | | |
| Poultry feed—tons | | | 4, 553 | 44, 804 | 3, 646 | 33, 119 | (1) | (1) |
| Lime—do | | | 1, 610 | 6, 629 | 1, 114 | 4, 570 | (1) | (1) |
| Unclassified products: | | | | | | | | |
| Fillets, fresh and frozen | | | | | | | | |
| pounds | 145, 492 | 33, 972 | | | | | | |
| Smoked—do | (6) | (6) | 7 581, 849 | 7 162, 300 | 1 1, 014, 000 | 8 291, 270 | | |
| Canned | | | | | | | | |
| standard cases | 6, 234 | 75, 343 | (6) | (6) | | | | |
| Miscellaneous | | 373, 501 | | 781, 226 | | | | 203, 331 |
| Total | | 5, 971, 402 | | 3, 053, 955 | | 1, 329, 442 | | 248, 765 |
| By fishermen: | | | | | | | | |
| Bluefish, smoked | | | | | | | | |
| pounds | | | 50 | 2 | | | | |
| Eels: | | | | | | | | |
| Smoked—do | 2, 150 | 537 | 5, 450 | 1, 675 | | | | |
| Pickled—do | | | 48 | 14 | | | | |
| Herring, sea, smoked | | | | | | | | |
| pounds | | | 4, 900 | 196 | | | | |
| Mackerel, smoked—do | | | 100 | 20 | | | | |
| Squirrel hake, smoked | | | | | | | | |
| pounds | | | 50 | 5 | | | | |
| Whiting, smoked—do | | | 50 | 5 | | | | |
| Clams, soft, fresh-shucked | | | | | | | | |
| gallons | | | 6, 840 | 5, 244 | | | | |
| Scallops: | | | | | | | | |
| Bay, fresh-shucked | | | | | | | | |
| gallons | 5, 400 | 17, 100 | | | | | | |
| Sea, fresh-shucked | | | | | | | | |
| gallons | 204, 565 | 192, 647 | 35, 603 | 33, 813 | | | | |
| Sturgeon roe, salted | | | | | | | | |
| pounds | | | 50 | 33 | | | | |
| King crab scrap—tons | | | 214 | 6, 655 | | | 80 | 2, 200 |
| Total | | 210, 284 | | 47, 662 | | | | 2, 200 |
| Grand total | | 6, 181, 686 | | 3, 101, 617 | | 1, 329, 442 | | 250, 965 |

¹ The production of this item is included under "Unclassified products."

² Includes the value of a small quantity of fresh-water mussel shell novelties.

³ Includes fresh fillets of bluefish, halibut, mackerel, pollock, red snapper, and frozen fillets of salmon.

⁴ This has been included under "Miscellaneous."

⁵ Includes smoked bluefish, carp, chubs, cisco, cod, eels, flounders, goosefish, haddock, lake trout, mackerel, sea herring (blouters), shad, tullibee, and whitefish and kippered sea herring.

⁶ Includes smoked alewives, butterfish, haddock fillets, haddock (finnan haddie), lake trout, mackerel, salmon, shad, and sturgeon, and kippered salmon and shad.

⁷ Includes canned clam products, pickled eels, terrapin and turtle products, sea mussels, and salmon caviar.

⁸ Includes smoked eels, haddock fillets, and sea herring (blouters); fish meal; and fresh-water mussel shell products.

⁹ Includes canned hard clams, canned hard clam chowder, canned oysters, shredded salt cod, and fresh-cooked crab meat.

¹⁰ Includes menhaden dry and acid scrap and oil, and oyster-shell products.

NOTE.—The total value of products for the Middle Atlantic States was as follows: By manufacturing establishments, \$10,603,564; and by fishermen, \$260,146. Some of the above products may have been manufactured from fishery products imported from another State or a foreign country; therefore, they cannot be correlated directly with the catch within the State. Of the total number of persons engaged on transporting craft, 135 have been included as fishermen and among the total number of persons engaged in the preparation of fishermen's prepared products, 511 have been included as fishermen.

U. S. BUREAU OF FISHERIES

NEW YORK

Fisheries of New York, 1933

OPERATING UNITS: BY GEAR

| Item | Purse seines, other than menhaden | Haul seines | Gill nets | | | | | Lines | | | |
|-------------------------------|-----------------------------------|-------------|-----------|------------|------------|-----------|--------------|------------|------------|-----------------|--|
| | | | Anchor | Drift | Run-round | Stake | Hand | Trawl | Troll | Trot with hooks | |
| | Number | Number | Number | Number | Number | Number | Number | Number | Number | Number | |
| Fishermen: | | | | | | | | | | | |
| On vessels..... | 17 | | 6 | 9 | 18 | | 26 | 77 | | | |
| On boats and shore: | | | | | | | | | | | |
| Regular..... | | 57 | 27 | 51 | 46 | 9 | 104 | 90 | 5 | | |
| Casual..... | | 98 | 6 | 174 | 1 | 62 | 32 | 18 | 5 | 14 | |
| Total..... | 17 | 155 | 39 | 234 | 65 | 71 | 172 | 185 | 10 | 14 | |
| Vessels: | | | | | | | | | | | |
| Motor..... | 2 | | 3 | 1 | 6 | | 6 | 8 | | | |
| Net tonnage..... | 41 | | 22 | 21 | 61 | | 135 | 219 | | | |
| Boats: | | | | | | | | | | | |
| Motor..... | | | 12 | | 23 | | 44 | 53 | 5 | | |
| Other..... | | 68 | 13 | 114 | 2 | 45 | 16 | 6 | | 14 | |
| Accessory boats..... | 2 | | | | | | | 37 | | | |
| Apparatus: | | | | | | | | | | | |
| Number..... | 2 | 71 | 46 | 254 | 43 | 49 | 225 | 663 | 20 | 14 | |
| Length, yards..... | 800 | 5,926 | | | | | | | | | |
| Square yards..... | | | 40,832 | 423,369 | 124,822 | 23,161 | | | | | |
| Hooks, baits, or snoods..... | | | | | | | 441 | 187,600 | 20 | 1,495 | |
| Item | Pound nets | Stop nets | Fyke nets | Dip nets | Scap nets | Drag nets | Otter trawls | Pots | | Harpoons | |
| | | | | | | | | Eel | Lobster | | |
| | Number | Number | Number | Number | Number | Number | Number | Number | Number | Number | |
| Fishermen: | | | | | | | | | | | |
| On vessels..... | 3 | | | | | | 357 | 2 | 7 | 40 | |
| On boats and shore: | | | | | | | | | | | |
| Regular..... | 174 | 2 | 40 | 5 | | 27 | 75 | 66 | 175 | 20 | |
| Casual..... | 17 | 29 | 59 | | 279 | 20 | 12 | 37 | 31 | | |
| Total..... | 194 | 31 | 99 | 5 | 279 | 47 | 444 | 105 | 213 | 60 | |
| Vessels: | | | | | | | | | | | |
| Steam..... | | | | | | | 9 | | | | |
| Net tonnage..... | | | | | | | 1,773 | | | | |
| Motor..... | 1 | | | | | | 56 | 1 | 5 | 8 | |
| Net tonnage..... | 10 | | | | | | 786 | 10 | 33 | 117 | |
| Total vessels..... | 1 | | | | | | 65 | 1 | 5 | 8 | |
| Total net tonnage..... | 10 | | | | | | 2,559 | 10 | 33 | 117 | |
| Boats: | | | | | | | | | | | |
| Motor..... | 43 | | 17 | | | 12 | 55 | 35 | 120 | 10 | |
| Other..... | 115 | 22 | 75 | 5 | 215 | | | 54 | 2 | | |
| Apparatus: | | | | | | | | | | | |
| Number..... | 301 | 24 | 849 | 5 | 279 | 47 | 130 | 3,877 | 18,340 | 18 | |
| Square yards..... | | 11,258 | | | | | | | | | |
| Yards at mouth..... | | | | | | 94 | 2,791 | | | | |

Fisheries of New York, 1933—Continued

OPERATING UNITS: BY GEAR—Continued

| Item | Spears | Dredges | | | | | Tongs | Rakes | Forks | Total, exclusive of duplication |
|-------------------------------|-----------|-----------|-----------|-----------|------------|------------|------------|------------|------------|---------------------------------|
| | | Clam | Crab | Mus-sel | Oyster | Scal-lop | | | | |
| | Num-ber | Num-ber | Num-ber | Num-ber | Num-ber | Num-ber | Num-ber | Num-ber | Num-ber | Num-ber |
| Fishermen: | | | | | | | | | | |
| On vessels..... | | 10 | 5 | 3 | 165 | 134 | 50 | | | 825 |
| On boats and shore: | | | | | | | | | | |
| Regular..... | 43 | 12 | | | | 41 | 584 | 302 | 133 | 1,509 |
| Casual..... | 34 | 2 | | | | 161 | 209 | 280 | 138 | 1,482 |
| Total..... | 77 | 24 | 5 | 3 | 165 | 336 | 843 | 582 | 271 | 3,816 |
| Vessels: | | | | | | | | | | |
| Steam..... | | | | | | | | | | 9 |
| Net tonnage..... | | | | | | | | | | 1,773 |
| Motor..... | | 5 | 1 | 1 | 39 | 25 | 23 | | | 160 |
| Net tonnage..... | | 43 | 23 | 13 | 791 | 382 | 148 | | | 2,422 |
| Total vessels..... | | 5 | 1 | 1 | 39 | 25 | 23 | | | 169 |
| Total net tonnage..... | | 43 | 23 | 13 | 791 | 382 | 148 | | | 4,195 |
| Boats: | | | | | | | | | | |
| Motor..... | | 9 | | | | 8 | 272 | 154 | | 636 |
| Other..... | 71 | | | | | 125 | 256 | 364 | 16 | 1,188 |
| Accessory boats..... | | | | | | | | | | 39 |
| Apparatus: | | | | | | | | | | |
| Number..... | 77 | 14 | 6 | 2 | 78 | 250 | 837 | 568 | 271 | |
| Yards at mouth..... | | 12 | 14 | 2 | 115 | 365 | | | | |

CATCH: BY GEAR

| Species | Purse seines, other than menhaden | | Haul seines | | Gill nets | | | | | | | | | |
|---------------------------------------|-----------------------------------|--------------|----------------|---------------|----------------|--------------|----------------|---------------|----------------|--------------|---------------|--------------|--------|-------|
| | | | | | Anchor | | Drift | | Runaround | | Stake | | | |
| | | | | | Lb. | Value | Lb. | Value | Lb. | Value | Lb. | Value | Lb. | Value |
| Alewives..... | | | 37,119 | \$540 | | | 12,850 | \$302 | | | | | 767 | \$31 |
| Bluefish..... | 27,700 | \$1,422 | 1,100 | 85 | 55,500 | \$2,720 | 26,000 | 1,300 | 102,200 | \$2,658 | | | 2,500 | 250 |
| Carp..... | | | 64,283 | 4,701 | 1,089 | 104 | 1,001 | 103 | | | | | 625 | 44 |
| Catfish and bull-heads..... | | | 2,695 | 401 | | | 1,280 | 159 | | | | | | |
| Eels, common..... | | | 5,500 | 725 | | | 287 | 49 | | | | | | |
| King whiting or "kingfish"..... | | | 12,500 | 1,000 | | | | | | | | | | |
| Mackerel..... | | | | | | | 36,400 | 1,092 | 127,000 | 5,820 | | | | |
| Mummichog..... | | | 18,000 | 1,800 | | | | | | | | | | |
| Scup or porgy..... | | | 600 | 12 | | | | | | | | | | |
| Shad..... | | | 4,040 | 448 | | | 297,753 | 23,844 | | | | | 34,143 | 3,660 |
| Silversides..... | | | 18,500 | 1,838 | | | | | | | | | | |
| Squeteagues or "sea trout," gray..... | 5,000 | 250 | 32,000 | 1,600 | 52,900 | 2,390 | | | 31,700 | 1,416 | | | 6,591 | 1,004 |
| Striped bass..... | | | 4,214 | 495 | | | 2,283 | 298 | | | | | 60 | 9 |
| Sturgeon..... | | | | | | | 800 | 200 | | | | | | |
| Suckers..... | | | 5,350 | 451 | | | 187 | 28 | | | | | | |
| Sunfish..... | | | 200 | 20 | | | 35 | 5 | | | | | | |
| White perch..... | | | 2,850 | 240 | | | 989 | 83 | | | | | 4,124 | 297 |
| Yellow perch..... | | | 350 | 27 | | | | | | | | | 500 | 28 |
| Shrimp..... | | | 17,500 | 3,500 | | | | | | | | | | |
| Total..... | 32,700 | 1,672 | 226,801 | 17,883 | 109,489 | 5,214 | 379,865 | 27,463 | 260,900 | 9,893 | 49,310 | 5,332 | | |

Fisheries of New York, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Lines | | | | | | | | Pound nets | |
|---------------------------------------|-----------|----------|-----------|----------|--------|---------|-----------------|-------|------------|---------|
| | Hand | | Trawl | | Troll | | Trot with hooks | | | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | | | | | | | | | 1,000 | \$10 |
| Bluefish..... | 734,850 | \$37,069 | | | 15,000 | \$1,200 | | | 166,579 | 8,474 |
| Bonito..... | | | | | | | | | 159,007 | 9,541 |
| Butterfish..... | | | | | | | | | 1,479,690 | 60,660 |
| Carp..... | | | | | | | 508 | \$61 | | |
| Catfish and bullheads..... | | | | | | | 400 | 63 | | |
| Cod..... | 64,802 | 1,840 | 396,500 | \$12,335 | | | | | 42,433 | 854 |
| Croaker..... | | | | | | | | | 3,842 | 174 |
| Eels: | | | | | | | | | | |
| Common..... | | | | | | | 2,371 | 303 | 85,065 | 9,728 |
| Conger..... | | | 2,000 | 200 | | | | | | |
| Flounders..... | 29,250 | 877 | | | | | | | 194,157 | 10,409 |
| Frigate mackerel..... | | | | | | | | | 81,886 | 1,234 |
| Haddock..... | 1,000 | 20 | 12,000 | 340 | | | | | | |
| Herring, sea..... | | | | | | | | | 74,987 | 610 |
| Hickory shad..... | | | | | | | | | 1,300 | 25 |
| King whiting or "kingfish"..... | | | | | | | | | 58,112 | 2,967 |
| Mackerel..... | | | | | | | | | 160,167 | 5,435 |
| Menhaden..... | | | | | | | | | 201,176 | 938 |
| Mullet..... | | | | | | | | | 2,500 | 96 |
| Pollock..... | 125,000 | 2,500 | | | | | | | 9,341 | 139 |
| Scup or porgy..... | 24,600 | 402 | | | | | | | 1,020,187 | 16,482 |
| Sea bass..... | 102,000 | 5,880 | | | | | | | 56,891 | 3,512 |
| Sea robin..... | | | | | | | | | 18,758 | 276 |
| Shad..... | | | | | | | | | 15,754 | 788 |
| Sharks..... | | | | | | | | | 2,500 | 35 |
| Spot..... | | | | | | | | | 21,029 | 423 |
| Squeteagues or "sea trout", gray..... | 10,000 | 470 | 23,500 | 1,045 | | | | | 654,793 | 30,377 |
| Striped bass..... | | | | | | | | | 4,579 | 726 |
| Sturgeon..... | | | | | | | | | 160 | 26 |
| Swellfish..... | | | | | | | | | 2,000 | 200 |
| Swordfish..... | | | | | | | | | 180 | 20 |
| Tautog..... | 25,000 | 750 | | | | | | | 34,598 | 803 |
| Tilefish..... | | | 1,350,000 | 67,500 | | | | | | |
| Tuna or "horse mackerel"..... | | | | | | | | | 6,736 | 270 |
| Whitebait..... | | | | | | | | | 3,000 | 450 |
| White perch..... | | | | | | | | | 450 | 85 |
| Whiting..... | | | | | | | | | 105,211 | 1,019 |
| Crabs, hard..... | | | | | | | | | 21,050 | 362 |
| Lobsters..... | | | | | | | | | 800 | 200 |
| Squid..... | | | | | | | | | 442,492 | 6,619 |
| Total..... | 1,116,502 | 50,798 | 1,784,000 | 81,420 | 15,000 | 1,200 | 3,279 | 432 | 5,152,384 | 173,947 |

| Species | Stop nets | | Fyke nets | | Dip nets | | Scap nets | | Drag nets | |
|----------------------------|-----------|-------|-----------|---------|----------|-------|-----------|---------|-----------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | 300 | \$3 | 81,655 | \$1,000 | | | 84,566 | \$3,137 | | |
| Carp..... | 92,258 | 7,717 | 7,654 | 675 | | | 52,079 | 4,169 | | |
| Catfish and bullheads..... | 545 | 95 | 12,246 | 1,569 | | | 1,862 | 270 | | |
| Eels, common..... | 820 | 70 | 1,896 | 231 | | | 76 | 6 | | |
| Flounders..... | | | 317,000 | 4,310 | | | | | | |
| Pike or pickerel..... | 30 | 3 | 390 | 50 | | | | | | |
| Shad..... | 40 | 6 | | | | | 52 | 6 | | |
| Smelt..... | | | | | | | 245 | 47 | | |
| Striped bass..... | 855 | 122 | 378 | 70 | | | 43 | 8 | | |
| Sturgeon..... | 400 | 32 | 16 | 1 | | | 50 | 10 | | |
| Suckers..... | 926 | 84 | 12,889 | 1,034 | | | 9,272 | 768 | | |
| Sunfish..... | 75 | 4 | 1,065 | 73 | | | 315 | 30 | | |
| Tautog..... | | | 1,500 | 30 | | | | | | |
| Tomcod..... | | | 7,560 | 333 | | | 130 | 5 | | |
| White perch..... | 50 | 5 | 5,830 | 328 | | | 889 | 101 | | |
| Yellow perch..... | | | 862 | 96 | | | 145 | 17 | | |
| Crabs: | | | | | | | | | | |
| Hard..... | | | 20,000 | 200 | | | | | | |
| Soft..... | | | | | 825 | \$225 | | | | |
| Shrimp..... | | | | | | | | | 42,490 | \$8,480 |
| Total..... | 96,299 | 8,141 | 470,941 | 10,000 | 825 | 225 | 140,664 | 8,574 | 42,490 | 8,480 |

Fisheries of New York, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Otter trawls | | Pots | | | | Harpoons | | Spears | |
|--|-------------------|----------------|----------------|---------------|----------------|---------------|----------------|---------------|---------------|--------------|
| | | | Eel | | Lobster | | | | | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Bluefish..... | 292 | \$19 | | | | | | | | |
| Butterfish..... | 18,552 | 679 | | | | | | | | |
| Catfish and bull- heads..... | | | 40 | \$4 | | | | | | |
| Cod..... | 3,756,934 | 97,554 | | | | | | | | |
| Croaker..... | 76,305 | 1,357 | | | | | | | | |
| Cusk..... | 3,500 | 99 | | | | | | | | |
| Drum, black..... | 447 | 9 | | | | | | | | |
| Eels: | | | | | | | | | | |
| Common..... | | | 173,205 | 14,214 | | | | | 21,250 | \$2,115 |
| Conger..... | 3,260 | 57 | | | | | | | | |
| Flounders..... | 4,650,721 | 176,436 | | | | | | | | |
| Haddock..... | 8,493,679 | 247,922 | | | | | | | | |
| Hake..... | 139,954 | 2,811 | | | | | | | | |
| Halibut..... | 52,508 | 7,280 | | | | | | | | |
| King whiting or "kingfish"..... | 2,611 | 90 | | | | | | | | |
| Mackerel..... | 150 | 6 | | | | | | | | |
| Mummichog..... | | | 7,050 | 705 | | | | | | |
| Pollock..... | 635,990 | 8,912 | | | | | | | | |
| Scup or porgy..... | 432,709 | 9,663 | | | | | | | | |
| Sea bass..... | 265,997 | 10,231 | | | 80,900 | \$4,045 | | | | |
| Skates..... | | | | | | | | | | |
| Spot..... | 1,736 | 40 | | | | | | | | |
| Squeteagues or "sea trout," gray..... | 13,741 | 600 | | | | | | | | |
| Striped bass..... | 11 | 1 | | | | | | | | |
| Sturgeon..... | 365 | 33 | | | | | | | | |
| Swordfish..... | | | | | | | 100,786 | \$12,421 | | |
| Tautog..... | 60 | 1 | | | | | | | | |
| Whiting..... | 900 | 9 | | | | | | | | |
| Wolfish..... | 25,625 | 576 | | | | | | | | |
| Lobsters..... | 4,621 | 607 | | | 341,075 | 66,449 | | | | |
| Shrimp..... | 12 | 1 | | | | | | | | |
| Conchs..... | | | | | 18,000 | 500 | | | | |
| Scallops, sea..... | 297 | 47 | | | | | | | | |
| Squid..... | 2,446 | 77 | | | | | | | | |
| Total..... | 18,584,005 | 565,124 | 180,295 | 14,923 | 439,975 | 70,994 | 100,786 | 12,421 | 21,250 | 2,115 |

| Species | Dredges | | | | | | | | | |
|-----------------------------------|----------------|---------------|----------------|--------------|---------------|--------------|------------------|----------------|------------------|----------------|
| | Clam | | Crab | | Mussel | | Oyster | | Scallop | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Crabs, hard..... | | | 108,333 | \$4,875 | | | | | | |
| Clams, surf or skim- mers..... | 359,000 | \$16,180 | | | | | | | 17,000 | \$1,596 |
| Conchs..... | | | | | 45,000 | \$3,000 | 2,400 | \$150 | | |
| Mussels, sea..... | | | | | | | | | | |
| Oysters: | | | | | | | | | | |
| Market, pri- vate, spring..... | | | | | | | 2,717,550 | 375,901 | | |
| Market, pri- vate, fall..... | | | | | | | 3,179,560 | 433,591 | | |
| Scallops: | | | | | | | | | | |
| Bay..... | | | | | | | | | 48,000 | 17,100 |
| Sea..... | | | | | | | | | 1,823,195 | 193,398 |
| Total..... | 359,000 | 16,180 | 108,333 | 4,875 | 45,000 | 3,000 | 5,899,500 | 809,642 | 1,888,195 | 212,093 |

| Species | Tongs | | | | Rakes | | Forks | |
|------------------------------|------------------|----------------|----------------|---------------|----------------|---------------|----------------|---------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Clams: | | | | | | | | |
| Hard, public..... | 1,345,710 | \$191,976 | | | 577,900 | \$79,319 | | |
| Hard, private..... | 21,040 | 4,601 | | | | | | |
| Soft, public..... | 6,400 | 725 | | | 64,550 | 4,885 | 151,200 | \$13,150 |
| Surf or skimmers..... | 20,600 | 2,200 | | | | | | |
| Oysters: | | | | | | | | |
| Market, public, spring..... | 19,250 | 2,662 | | | | | | |
| Market, public, fall..... | 71,484 | 8,462 | | | 10,500 | 1,050 | | |
| Market, private, spring..... | 43,975 | 5,237 | | | | | | |
| Market, private, fall..... | 49,475 | 5,787 | | | | | | |
| Bloodworms..... | | | | | | | 9,250 | 8,390 |
| Sandworms..... | | | | | | | 3,500 | 2,080 |
| Total..... | 1,877,834 | 221,650 | 662,950 | 85,254 | 163,950 | 85,254 | 163,950 | 23,620 |

Fisheries of New York, 1933—Continued

OPERATING UNITS: BY COUNTIES

| Item | Albany | Bronx | Colum- bia | Dutch- ess | Greene | Kings | Nas- sau | New York |
|--------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| Fishermen: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| On vessels..... | | | | | | 188 | 44 | 300 |
| On boats and shore: | | | | | | | | |
| Regular..... | | 3 | 2 | 14 | 2 | 185 | 259 | |
| Casual..... | 8 | | 72 | 154 | 56 | 130 | 97 | |
| Total | 8 | 3 | 74 | 168 | 58 | 503 | 400 | 300 |
| Vessels: | | | | | | | | |
| Steam..... | | | | | | | | 9 |
| Net tonnage..... | | | | | | | | 1,773 |
| Motor..... | | | | | | 39 | 10 | 16 |
| Net tonnage..... | | | | | | 594 | 152 | 424 |
| Total vessels | | | | | | 39 | 10 | 25 |
| Total net tonnage | | | | | | 594 | 152 | 2,197 |
| Boats: | | | | | | | | |
| Motor..... | | | | | | 80 | 48 | |
| Other..... | 8 | 3 | 54 | 97 | 32 | 10 | 202 | |
| Accessory boats | | | | | | 15 | 2 | 22 |
| Apparatus: | | | | | | | | |
| Purse seines: | | | | | | | | |
| Other than menhaden..... | | | | | | 1 | 1 | |
| Length, yards..... | | | | | | 400 | 400 | |
| Haul seines..... | | | 6 | 4 | 5 | 5 | 15 | |
| Length, yards..... | | | 650 | 560 | 585 | 60 | 860 | |
| Gill nets: | | | | | | | | |
| Drift..... | | | 3 | 31 | 1 | | | 140 |
| Square yards..... | | | 8,666 | 122,461 | 800 | | | 25,200 |
| Runaround..... | | | | | | 2 | 31 | |
| Square yards..... | | | | | | 16,000 | 93,200 | |
| Lines: | | | | | | | | |
| Hand..... | | | | | | 118 | | 4 |
| Hooks..... | | | | | | 231 | | 5 |
| Trawl..... | | | | | | 70 | 93 | 436 |
| Hooks..... | | | | | | 70,800 | 39,200 | 30,600 |
| Troll..... | | | | | | 20 | | |
| Hooks..... | | | | | | 20 | | |
| Trot with hooks..... | | | | 1 | | | | |
| Hooks..... | | | | 150 | | | | |
| Pound nets..... | | | | | | | 1 | |
| Stop nets..... | | | 2 | 1 | 12 | | | |
| Square yards..... | | | 140 | 100 | 7,775 | | | |
| Fyke nets..... | 2 | | 20 | 16 | 6 | | | |
| Dip nets..... | | | | | | | 5 | |
| Scap nets..... | 8 | | 44 | 93 | 31 | | | |
| Drag nets..... | | | | | | | 5 | |
| Yards at mouth..... | | | | | | | 10 | |
| Otter trawls..... | | | | | | 30 | 4 | 22 |
| Yards at mouth..... | | | | | | 693 | 68 | 393 |
| Pots: | | | | | | | | |
| Eel..... | | | | 4 | | | 296 | |
| Lobster..... | | | | | | 6,820 | 2,035 | |
| Harpoons..... | | | | | | | | 2 |
| Spears..... | | | | | | 12 | 30 | |
| Dredges: | | | | | | | | |
| Clam..... | | | | | | 12 | 1 | |
| Yards at mouth..... | | | | | | 10 | 1 | |
| Mussel..... | | | | | | | 2 | |
| Yards at mouth..... | | | | | | | 2 | |
| Oyster..... | | | | | | | 10 | 12 |
| Yards at mouth..... | | | | | | | 13 | 18 |
| Scallop..... | | | | | | 46 | 6 | 8 |
| Yards at mouth..... | | | | | | 122 | 17 | 24 |
| Tongs..... | | | | | | | 162 | |
| Rakes..... | | 3 | | | | | 84 | |
| Forks..... | | | | | | 100 | 95 | |

Fisheries of New York, 1933—Continued

OPERATING UNITS: BY COUNTIES—Continued

| Item | Orange | Putnam | Queens | Rensselaer | Richmond | Rockland | Suffolk | Ulster | Westchester |
|---------------------|-----------|----------|----------|------------|-----------|-----------|--------------|------------|-------------|
| | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | | | |
| On vessels | | | 2 | | 7 | | 284 | | |
| On boats and shore: | | | | | 8 | 5 | 991 | 24 | 9 |
| Regular | 7 | | | | 2 | 46 | 622 | 165 | 64 |
| Casual | 46 | 4 | | | | | | | |
| Total | 53 | 4 | 2 | 10 | 17 | 51 | 1,897 | 189 | 73 |
| Vessels: | | | | | | | | | |
| Motor | | | 1 | | 2 | | 92 | | |
| Net tonnage | | | 14 | | 30 | | 1,208 | | |
| Boats: | | | | | | | | | |
| Motor | | | | | 8 | | 499 | | 1 |
| Other | 32 | 4 | | | | 31 | 536 | 128 | 39 |
| Apparatus: | | | | | | | | | |
| Haul seines | 4 | 1 | | 3 | | | 20 | 5 | 3 |
| Length, yards | 533 | 125 | | 375 | | | 1,195 | 700 | 283 |
| Gill nets: | | | | | | | | | |
| Anchor | | | | | | | 44 | 2 | |
| Square yards | | | | | | | 40,497 | 335 | |
| Drift | 9 | | | 1 | | 13 | | 44 | 12 |
| Square yards | 37,928 | | | 3,200 | | 42,766 | | 142,731 | 39,617 |
| Run-around | | | | | | | 10 | | |
| Square yards | | | | | | | 15,622 | | |
| Stake | | 1 | | | | 10 | 23 | 1 | 14 |
| Square yards | | 165 | | | | 3,205 | 6,286 | 135 | 13,370 |
| Lines: | | | | | | | | | |
| Hand | | | | | | | 103 | | |
| Hooks | | | | | | | 202 | | |
| Trawl | | | | | | | 94 | | |
| Hooks | | | | | | | 47,000 | | |
| Trot with hooks | 1 | | | 1 | | 3 | | 6 | 2 |
| Hooks | 120 | | | 100 | | 325 | | 600 | 200 |
| Pound nets | | | | | | | 300 | | |
| Stop nets | | | | | | | | 7 | 1 |
| Square yards | 400 | | | | | | | 2,708 | 135 |
| Fyke nets | 6 | 1 | | 3 | | | 765 | 23 | 8 |
| Scap nets | 15 | 2 | | 6 | | | | 78 | 2 |
| Drag nets | | | | | | | | 42 | |
| Yards at mouth | | | | | | | | 84 | |
| Otter trawls | | | 1 | | 1 | | | 72 | |
| Yards at mouth | | | 30 | | 17 | | 1,590 | | |
| Pots: | | | | | | | | | |
| Eel | | | | | | 57 | 3,212 | 22 | 36 |
| Lobster | | | | | 1,150 | | 8,255 | | 80 |
| Harpoons | | | | | | | | 16 | |
| Spears | | | | | | | | 26 | |
| Dredges: | | | | | | | | | |
| Clam | | | | | 1 | | | | |
| Yards at mouth | | | | | 1 | | | | |
| Crab | | | | | 6 | | | | |
| Yards at mouth | | | | | 14 | | | | |
| Oyster | | | | | | | | 56 | |
| Yards at mouth | | | | | | | | 84 | |
| Scallop | | | | | 2 | | | 188 | |
| Yards at mouth | | | | | 5 | | | 197 | |
| Tongs | | | | | | | | 674 | 1 |
| Rakes | | | | | | | | 480 | 1 |
| Forks | | | | | | | | 76 | |

CATCH: BY COUNTIES

| Species | Albany | | Bronx | | Columbia | | Dutchess | | Greene | |
|-----------------------|--------------|------------|--------------|------------|---------------|--------------|----------------|---------------|----------------|--------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives | 1,250 | \$43 | | | 47,733 | \$773 | 36,363 | \$1,461 | 12,317 | \$400 |
| Carp | 3,850 | 274 | | | 11,930 | 821 | 11,510 | 1,018 | 91,215 | 7,454 |
| Catfish and bullheads | 90 | 10 | | | 4,398 | 574 | 3,092 | 307 | 1,892 | 295 |
| Eels, common | | | | | 480 | 45 | 870 | 111 | 726 | 59 |
| Pike or pickerel | | | | | | | | | 38 | 5 |
| Shad | | | | | 6,086 | 540 | 101,857 | 7,280 | 1,500 | 133 |
| Striped bass | 16 | 2 | | | 40 | 8 | 40 | 8 | 700 | 85 |
| Sturgeon | | | | | | | 850 | 210 | 16 | 1 |
| Suckers | 2,850 | 210 | | | 4,351 | 338 | 3,103 | 241 | 2,273 | 186 |
| Sunfish | | | | | 60 | 6 | 600 | 33 | 200 | 20 |
| Tomcod | | | | | | | 130 | 5 | | |
| White perch | | | | | 2,036 | 68 | 965 | 75 | 2,406 | 211 |
| Yellow perch | | | | | 50 | 10 | 117 | 12 | | |
| Clams, hard, public | | | 3,600 | \$675 | | | | | | |
| Total | 8,056 | 539 | 3,600 | 675 | 77,124 | 3,275 | 159,497 | 10,761 | 113,283 | 8,849 |

Fisheries of New York, 1933—Continued

CATCH: BY COUNTIES—Continued

| Species | Kings | | Nassau | | New York | | Orange | |
|----------------------------------|-------------|-----------|-------------|----------|--------------|----------|---------|--------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives | | | | | | | 6, 105 | \$175 |
| Bluefish | 726, 850 | \$38, 176 | 96, 774 | \$2, 317 | 63, 215 | \$3, 161 | | |
| Bonito | | | 79 | 5 | | | | |
| Butterfish | 3, 217 | 160 | 1, 535 | 61 | | | | |
| Carp | | | | | | | 8, 516 | 781 |
| Catfish and bullheads | | | | | | | 1, 768 | 339 |
| Cod | 332, 900 | 6, 837 | 118, 161 | 4, 843 | 3, 648, 444 | 95, 436 | | |
| Croaker | 36 | 1 | | | 7, 070 | 174 | | |
| Cusk | 200 | 4 | | | 3, 300 | 95 | | |
| Eels: | | | | | | | | |
| Common | 15, 000 | 1, 500 | 21, 000 | 2, 375 | | | 1, 132 | 136 |
| Conger | 178 | 4 | | | 2, 510 | 210 | | |
| Flounders | 1, 032, 766 | 30, 740 | 100, 059 | 3, 005 | 455, 422 | 16, 057 | | |
| Frigate mackerel | | | 11 | 1 | | | | |
| Haddock | 288, 300 | 6, 275 | | | 8, 179, 879 | 240, 827 | | |
| Hake | 2, 753 | 35 | | | 126, 608 | 2, 628 | | |
| Halibut | 200 | 8 | | | 52, 102 | 7, 243 | | |
| King whiting or "kingfish" | 25 | 1 | 26 | 1 | 22 | 1 | | |
| Mackerel | 67, 000 | 4, 020 | 60, 212 | 1, 806 | 36, 400 | 1, 062 | | |
| Mummichog | 12, 450 | 1, 245 | 6, 600 | 660 | | | | |
| Pollock | 700 | 22 | | | 635, 140 | 8, 889 | | |
| Scup or porgy | 12, 570 | 225 | 4, 828 | 86 | 65, 531 | 1, 376 | | |
| Sea bass | 54, 236 | 2, 264 | 85, 950 | 4, 298 | 62, 964 | 1, 802 | | |
| Shad | | | | | | | 17, 262 | 1, 621 |
| Silversides | 300 | 18 | 10, 200 | 1, 020 | | | | |
| Skates | 725 | 7 | | | | | | |
| Spot | | | 20 | 1 | 920 | 17 | | |
| Squeteagues or "sea trout", gray | 6, 000 | 270 | 26, 032 | 1, 093 | 294 | 20 | | |
| Striped bass | | | 3, 000 | 300 | | | 858 | 139 |
| Sturgeon | 8 | 1 | | | | | | |
| Suckers | | | | | | | 4, 175 | 438 |
| Sunfish | | | | | | | 260 | 26 |
| Swordfish | | | | | 35, 796 | 4, 856 | | |
| Tautog | 25, 000 | 750 | | | | | | |
| Tilfish | 322, 000 | 16, 100 | | | 1, 028, 000 | 51, 400 | | |
| Tuna or "horse mackerel" | | | 19 | 1 | | | | |
| Whiting | 900 | 9 | 70 | 1 | | | | |
| Wolfish | | | | | 25, 625 | 576 | | |
| Yellow perch | | | | | | | 720 | 74 |
| Crabs, soft | | | 825 | 225 | | | | |
| Lobsters | 174, 150 | 31, 347 | 19, 400 | 4, 850 | | | | |
| Shrimp | | | 16, 400 | 3, 280 | | | | |
| Clams: | | | | | | | | |
| Hard, public | | | 257, 750 | 60, 185 | | | | |
| Soft, public | | | 116, 000 | 11, 250 | | | | |
| Surf or skimmers | 335, 000 | 14, 680 | 23, 500 | 1, 960 | | | | |
| Conchs | 15, 200 | 1, 520 | | | | | | |
| Mussels, sea | | | 47, 400 | 3, 150 | | | | |
| Oysters: | | | | | | | | |
| Market, private, spring | | | 200, 600 | 27, 800 | 524, 498 | 74, 625 | | |
| Market, private, fall | | | 296, 400 | 42, 150 | 524, 498 | 74, 925 | | |
| Scallops, sea | 968, 420 | 107, 600 | 171, 990 | 21, 341 | 426, 105 | 47, 345 | | |
| Squid | | | 860 | 5 | 95 | 2 | | |
| Bloodworms | 9, 000 | 8, 140 | 250 | 250 | | | | |
| Total | 4, 406, 084 | 280, 959 | 1, 685, 951 | 188, 312 | 15, 904, 452 | 633, 056 | 40, 796 | 3, 729 |

| Species | Putnam | | Queens | | Rensselaer | | Richmond | |
|----------------------------|--------|-------|--------|-------|------------|--------|----------|----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives | | | | | 2, 269 | \$85 | | |
| Carp | 3, 855 | \$286 | | | 9, 228 | 870 | | |
| Catfish and bullheads | 240 | 12 | | | 2, 794 | 422 | | |
| Croaker | | | 140 | \$2 | | | | |
| Eels, common | | | | | 350 | 87 | | |
| Flounders | | | 5, 162 | 300 | | | 21, 000 | \$1, 050 |
| King whiting or "kingfish" | | | 20 | 1 | | | | |
| Pike or pickerel | | | | | 17 | 3 | | |
| Shad | | | | | 1, 215 | 110 | | |
| Suckers | 385 | 26 | | | 200 | 14 | | |
| Tomcod | 360 | 58 | | | | | | |
| White perch | 80 | 8 | | | | | | |
| Yellow perch | | | | | 80 | 5 | | |
| Crabs, hard | | | | | | | 108, 333 | 4, 875 |
| Lobsters | | | | | | | 14, 500 | 2, 610 |
| Clams, surf or skimmers | | | | | | | 12, 000 | 600 |
| Conchs | | | | | | | 1, 800 | 75 |
| Total | 4, 920 | 390 | 5, 322 | 303 | 16, 151 | 1, 566 | 157, 633 | 9, 110 |

Fisheries of New York, 1935—Continued

CATCH: BY COUNTIES—Continued

| Species | Rockland | | Suffolk | | Ulster | | Westchester | |
|----------------------------------|----------|------------|------------|-----------|---------|---------------|-------------|------------|
| | Pounds | Value \$39 | Pounds | Value | Pounds | Value \$1,051 | Pounds | Value \$26 |
| Alewives | 1,681 | | 81,000 | \$970 | | | | |
| Bluefish | | | 244,882 | 12,443 | | | | |
| Bonito | | | 158,928 | 9,536 | | | | |
| Butterfish | | | 1,493,460 | 61,118 | | | | |
| Carp | 250 | 25 | | | 74,885 | 4,636 | 4,200 | 409 |
| Catfish and bullheads | 140 | 14 | | | 3,504 | 472 | 1,090 | 116 |
| Cod | | | 161,164 | 5,467 | | | | |
| Croaker | | | 72,901 | 1,354 | | | | |
| Drum, black | | | 447 | 9 | | | | |
| Eels: | | | | | | | | |
| Common | 6,280 | 450 | 240,035 | 22,302 | 2,417 | 179 | 2,200 | 202 |
| Conger | | | 2,572 | 43 | | | | |
| Flounders | | | 3,576,719 | 131,880 | | | | |
| Frigate mackerel | | | 81,875 | 1,233 | | | | |
| Haddock | | | 38,500 | 1,180 | | | | |
| Hake | | | 10,593 | 148 | | | | |
| Halibut | | | 206 | 29 | | | | |
| Herring, sea | | | 74,987 | 610 | | | | |
| Hickory shad | | | 1,300 | 25 | | | | |
| King whiting or "kingfish" | | | 73,130 | 4,053 | | | | |
| Mackerel | | | 180,105 | 5,435 | | | | |
| Menhaden | | | 201,176 | 938 | | | | |
| Mullet | | | 2,500 | 96 | | | | |
| Mummichog | | | 6,000 | 600 | | | | |
| Pike or pickerel | | | | | 215 | 24 | 150 | 21 |
| Pollock | | | 134,391 | 2,640 | | | | |
| Porgy | | | 1,395,224 | 24,961 | | | | |
| Sea bass | | | 302,518 | 15,304 | | | | |
| Sea robin | | | 18,758 | 276 | | | | |
| Shad | 17,562 | 1,812 | 17,488 | 1,397 | 135,578 | 11,177 | 63,238 | 4,691 |
| Sharks | | | 2,500 | 35 | | | | |
| Silversides | | | 8,000 | 800 | | | | |
| Smelt | | | | | 245 | 47 | | |
| Spot | | | 21,825 | 445 | | | | |
| Squeteagues or "sea trout", gray | | | 791,308 | 36,764 | | | | |
| Striped bass | 7,878 | 1,135 | 4,590 | 727 | 27 | 6 | 1,845 | 322 |
| Sturgeon | | | 517 | 58 | | | 460 | 41 |
| Suckers | | | | | 7,697 | 562 | 3,590 | 350 |
| Sunfish | | | | | 570 | 47 | | |
| Swellfish | | | 2,000 | 200 | | | | |
| Swordfish | | | 65,170 | 7,585 | | | | |
| Tautog | | | 36,158 | 834 | | | | |
| Tomcod | | | 5,000 | 200 | 200 | 25 | 2,000 | 50 |
| Tuna or "horse mackerel" | | | 6,717 | 269 | | | | |
| White bait | | | 3,000 | 450 | | | | |
| White perch | 2,444 | 176 | 450 | 65 | 2,651 | 235 | 4,150 | 181 |
| Whiting | | | 106,141 | 1,018 | | | | |
| Yellow perch | 500 | 28 | | | 390 | 39 | | |
| Crabs, hard | | | 41,050 | 562 | | | | |
| Lobsters | | | 136,946 | 28,149 | | | 1,500 | 300 |
| Shrimp | | | 43,512 | 8,701 | | | | |
| Clams: | | | | | | | | |
| Hard, public | | | 1,659,460 | 219,785 | | | 2,800 | 650 |
| Hard, private | | | 21,040 | 4,601 | | | | |
| Soft, public | | | 106,150 | 7,510 | | | | |
| Surf or skimmer | | | 9,000 | 1,250 | | | | |
| Conchs | | | 18,000 | 500 | | | | |
| Oysters: | | | | | | | | |
| Market, public, spring | | | 19,250 | 2,662 | | | | |
| Market, public, fall | | | 81,984 | 9,512 | | | | |
| Market, private, spring | | | 2,036,430 | 278,413 | | | | |
| Market, private, fall | | | 2,408,130 | 322,303 | | | | |
| Scallops: | | | | | | | | |
| Bay | | | 48,000 | 17,100 | | | | |
| Sea | | | 256,977 | 17,159 | | | | |
| Squid | | | 443,983 | 6,689 | | | | |
| Sandworms | | | 3,500 | 2,080 | | | | |
| Total | 36,715 | 3,679 | 16,956,647 | 1,280,473 | 257,132 | 19,500 | 78,069 | 7,359 |

U. S. BUREAU OF FISHERIES

Fisheries of New York, 1933—Continued

SEED OYSTER FISHERY: BY GEAR

| Item | Dredges | | Tongs | | Rakes | | Total, exclusive of duplication | |
|----------------------------|---------|----------|---------|----------|---------|---------|---------------------------------|----------|
| OPERATING UNITS | | | | | | | | |
| Fishermen: | Number | | Number | | Number | | Number | |
| On vessels..... | 15 | | 9 | | | | 24 | |
| On boats and shore: | | | | | | | | |
| Regular..... | | | 185 | | 45 | | 230 | |
| Total..... | 15 | | 194 | | 45 | | 254 | |
| Vessels, motor..... | 3 | | 4 | | | | 7 | |
| Net tonnage..... | 45 | | 24 | | | | 69 | |
| Boats: | | | | | | | | |
| Motor..... | | | 100 | | 20 | | 120 | |
| Other..... | | | 5 | | 25 | | 30 | |
| Apparatus: | | | | | | | | |
| Number..... | 6 | | 194 | | 45 | | | |
| Yards at mouth..... | 9 | | | | | | | |
| CATCH | | | | | | | | |
| Oysters: | Bushels | Value | Bushels | Value | Bushels | Value | Bushels | Value |
| Seed, public, spring..... | | | 35,100 | \$17,550 | 7,500 | \$3,000 | 42,600 | \$20,550 |
| Seed, public, fall..... | | | 5,000 | 2,500 | 7,500 | 3,000 | 12,500 | 5,500 |
| Seed, private, spring..... | 40,290 | \$39,540 | | | | | 40,290 | \$39,540 |
| Seed, private, fall..... | 38,290 | 38,290 | | | | | 38,290 | 38,290 |
| Total..... | 78,580 | 77,830 | 40,100 | 20,050 | 15,000 | 6,000 | 133,680 | 103,880 |

NOTE.—The seed-oyster fishery of New York is confined to Suffolk County. Of the total number of persons fishing for seed oysters, 242 are duplicated among those fishing for market oysters or other species. Similarly the following craft and gear are duplicated; 1 motor vessel, 188 tongs, and 15 rakes.

NEW JERSEY

Fisheries of New Jersey, 1933

OPERATING UNITS: BY GEAR

| Item | Purse seines | | Haul seines | Gill nets | | | | Lines | |
|-----------------------------|--------------|--------|-------------|-----------|---------|------------|--------|--------|---------|
| | Men-haden | Other | | Anchor | Drift | Run-around | Stake | Hand | Trawl |
| Fishermen: | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| On vessels..... | 150 | 73 | | | 8 | | | 80 | 38 |
| On boats and shore: | | | | | | | | | |
| Regular..... | | | 65 | 2 | 68 | 57 | 37 | 156 | 228 |
| Casual..... | | | 188 | 5 | 202 | 53 | 62 | 134 | 56 |
| Total..... | 150 | 73 | 253 | 7 | 278 | 110 | 99 | 370 | 322 |
| Vessels: | | | | | | | | | |
| Motor..... | 8 | 7 | | | 1 | | | 14 | 10 |
| Net tonnage..... | 285 | 132 | | | 11 | | | 175 | 107 |
| Sail..... | | | | | 1 | | | 1 | 1 |
| Net tonnage..... | | | | | 7 | | | 7 | 7 |
| Total vessels..... | 8 | 7 | | | 2 | | | 15 | 11 |
| Total net tonnage..... | 285 | 132 | | | 18 | | | 182 | 114 |
| Boats: | | | | | | | | | |
| Motor..... | | | 6 | 3 | 93 | 51 | 27 | 150 | 129 |
| Other..... | | | 95 | | 54 | | 22 | 1 | |
| Accessory boats..... | 22 | 12 | | | | | | 30 | 10 |
| Apparatus: | | | | | | | | | |
| Number..... | 8 | 7 | 105 | 10 | 605 | 69 | 251 | 463 | 712 |
| Length, yards..... | 2,806 | 2,510 | 9,040 | | | | | | |
| Square yards..... | | | | 5,830 | 448,333 | 210,695 | 46,882 | | |
| Hooks, baits or snoods..... | | | | | | | | 720 | 420,400 |

Fisheries of New Jersey, 1933—Continued

OPERATING UNITS: BY GEAR—Continued

| Item | Lines | | Pound nets | Weirs | Stop nets | Fyke nets | Dip nets | Cast nets | Drag nets |
|-----------------------------|--------|---------------------------|------------|--------|-----------|-----------|----------|-----------|-----------|
| | Troll | Trot with baits or snoods | | | | | | | |
| Fishermen: | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| On vessels..... | 9 | | 200 | | | | | | |
| On boats and shore: | | | | | | | | | |
| Regular..... | 50 | 6 | 117 | | 20 | 35 | 9 | | 2 |
| Casual..... | 60 | 8 | 13 | 15 | 72 | 62 | 25 | 3 | 11 |
| Total..... | 119 | 14 | 330 | 15 | 92 | 97 | 34 | 3 | 13 |
| Vessels: | | | | | | | | | |
| Motor..... | 2 | | 30 | | | | | | |
| Net tonnage..... | 20 | | 183 | | | | | | |
| Boats: | | | | | | | | | |
| Motor..... | 101 | 7 | 20 | 4 | 19 | 31 | 9 | | 8 |
| Other..... | | | 11 | 4 | 34 | 37 | 21 | | |
| Accessory boats..... | 4 | | 10 | | | | | | |
| Apparatus: | | | | | | | | | |
| Number..... | 404 | 9 | 153 | 88 | 65 | 1,369 | 34 | 3 | 14 |
| Square yards..... | | | | | 50,615 | | | | |
| Yards at mouth..... | | | | | | | | | 28 |
| Hooks, baits or snoods..... | 404 | 9,900 | | | | | | | |

| Item | Otter trawls | Wire baskets | Pots | | | Harpoons | Spears | Dredges | |
|------------------------|--------------|--------------|--------|--------|---------|----------|--------|---------|--------|
| | | | Crab | El | Lobster | | | Clam | Crab |
| Fishermen: | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| On vessels..... | 123 | | | | | 5 | | 51 | 65 |
| On boats and shore: | | | | | | | | | |
| Regular..... | 61 | | 2 | 35 | 177 | | 48 | 8 | 2 |
| Casual..... | | 1 | | 51 | 55 | | 44 | | |
| Total..... | 184 | 1 | 2 | 86 | 232 | 5 | 92 | 59 | 67 |
| Vessels: | | | | | | | | | |
| Motor..... | 37 | | | | | 1 | | 15 | 15 |
| Net tonnage..... | 706 | | | | | 26 | | 229 | 312 |
| Sail..... | | | | | | | | 2 | 2 |
| Net tonnage..... | | | | | | | | 17 | 17 |
| Total vessels..... | 37 | | | | | 1 | | 17 | 17 |
| Total net tonnage..... | 706 | | | | | 26 | | 246 | 329 |
| Boats: | | | | | | | | | |
| Motor..... | 30 | | 1 | 39 | 126 | | 5 | 4 | 1 |
| Other..... | | | | 27 | | | 69 | | |
| Accessory boats..... | | | | | | | | | |
| Apparatus: | | | | | | | | | |
| Number..... | 67 | 12 | 10 | 2,571 | 28,071 | 1 | 92 | 43 | 69 |
| Yards at mouth..... | 1,514 | | | | | | | 48 | 90 |

| Item | Dredges | | Tongs | Rakes | Forks | Hoes | Gaffs | By hand | Total, exclusive of duplication |
|------------------------|---------|---------|--------|--------|--------|--------|--------|---------|---------------------------------|
| | Oyster | Scallop | | | | | | | |
| Fishermen: | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| On vessels..... | 635 | 27 | | | | | | | 1,267 |
| On boats and shore: | | | | | | | | | |
| Regular..... | 19 | | 275 | 178 | 47 | 56 | 2 | 19 | 954 |
| Casual..... | 15 | | 508 | 331 | 4 | 121 | | 121 | 1,622 |
| Total..... | 669 | 27 | 783 | 509 | 51 | 177 | 2 | 140 | 3,843 |
| Vessels: | | | | | | | | | |
| Motor..... | 109 | 5 | | | | | | | 208 |
| Net tonnage..... | 2,025 | 83 | | | | | | | 3,470 |
| Sail..... | | | | | | | | | 3 |
| Net tonnage..... | | | | | | | | | 24 |
| Total vessels..... | 109 | 5 | | | | | | | 211 |
| Total net tonnage..... | 2,025 | 83 | | | | | | | 3,494 |
| Boats: | | | | | | | | | |
| Motor..... | 22 | | 309 | 264 | | 28 | | | 1,029 |
| Other..... | | | 336 | 222 | | 59 | 2 | | 740 |
| Accessory boats..... | | | | | | | | | 88 |
| Apparatus: | | | | | | | | | |
| Number..... | 259 | 10 | 783 | 509 | 51 | 176 | 2 | | |
| Yards at mouth..... | 309 | 34 | | | | | | | |

U. S. BUREAU OF FISHERIES

Fisheries of New Jersey, 1933—Continued

CATCH: BY GEAR

| Species | Purse seines | | | | Haul seines | |
|--------------------------------------|--------------|----------|-----------|---------|-------------|--------|
| | Menhaden | | Other | | Pounds | Value |
| | Pounds | Value | Pounds | Value | | |
| Alewives..... | | | | | 4,450 | \$66 |
| Bluefish..... | | | 76,085 | \$3,053 | 1,100 | 66 |
| Butterfish..... | | | 55 | 2 | | |
| Carp..... | | | | | 35,871 | 4,003 |
| Catfish and bullheads..... | | | | | 19,300 | 915 |
| Croaker..... | | | 3,707 | 69 | 15,100 | 409 |
| Eels, common..... | | | | | 53,001 | 3,338 |
| Flounders..... | | | 7,828 | 152 | 41,050 | 2,562 |
| Herring, round..... | | | | | 750 | 8 |
| Kingfish or "king mackerel"..... | | | 60 | 1 | | |
| King whiting or "kingfish"..... | | | 53 | 2 | 1,300 | 86 |
| Mackerel..... | | | 1,350 | 34 | | |
| Menhaden..... | 40,557,900 | \$73,561 | 2,890,800 | 7,207 | | |
| Mullet..... | | | | | 82,480 | 6,228 |
| Mummichog..... | | | | | 10,100 | 445 |
| Scup or porgy..... | | | 407,241 | 2,576 | | |
| Sea bass..... | | | 437 | 17 | | |
| Shad..... | | | | | 66,709 | 7,803 |
| Silversides..... | | | | | 4,330 | 626 |
| Spanish mackerel..... | | | 5 | 1 | | |
| Spot..... | | | | | 2,000 | 80 |
| Squeteague or "sea trout", gray..... | | | 367,309 | 5,448 | 49,612 | 2,763 |
| Striped bass..... | | | | | 1,200 | 280 |
| Suckers..... | | | | | 101,600 | 8,395 |
| Thimble-eyed mackerel..... | | | 300 | 6 | | |
| Tuna or "horse mackerel"..... | | | 44 | 2 | | |
| White perch..... | | | | | 5,500 | 457 |
| Crabs: | | | | | | |
| Hard..... | | | | | 2,355 | 144 |
| Soft..... | | | | | 16,881 | 3,101 |
| Shrimp..... | | | | | 400 | 80 |
| Total..... | 40,557,900 | 73,561 | 3,755,574 | 18,570 | 516,089 | 41,883 |

| Species | Gill nets | | | | | | | |
|--------------------------------------|-----------|-------|---------|---------|-----------|---------|---------|--------|
| | Anchor | | Drift | | Runaround | | Stake | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | | | | | | | 5,800 | \$143 |
| Bluefish..... | 1,000 | \$50 | 15,719 | \$1,020 | 159,842 | \$6,877 | 6,425 | 361 |
| Bonito..... | | | | | 1,114 | 78 | | |
| Butterfish..... | | | 825 | 34 | 1,543 | 62 | | |
| Croaker..... | | | 113,806 | 2,068 | 33,952 | 939 | 1,500 | 80 |
| Mackerel..... | | | 175,968 | 4,890 | 13,131 | 421 | | |
| Scup or porgy..... | | | | | 2,292 | 46 | | |
| Shad..... | 638 | 85 | 75,772 | 10,989 | 19 | 2 | 251,211 | 20,703 |
| Spot..... | | | 55,000 | 1,430 | 11,520 | 330 | 7,500 | 375 |
| Squeteague or "sea trout", gray..... | 750 | 28 | 91,865 | 2,149 | 74,143 | 3,614 | 6,314 | 289 |
| Striped bass..... | | | 1,780 | 300 | | | 3,795 | 692 |
| Sturgeon..... | | | 498 | 98 | | | | |
| White perch..... | | | | | | | 10,300 | 1,200 |
| Whiting..... | | | | | 45 | 1 | | |
| Crabs, soft..... | | | 801 | 100 | | | | |
| Total..... | 2,388 | 173 | 532,034 | 23,078 | 297,601 | 12,370 | 292,845 | 23,843 |

Fisheries of New Jersey, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Lines | | | | | | | |
|--------------------------------------|------------------|---------------|------------------|---------------|----------------|---------------|---------------------------|--------------|
| | Hand | | Trawl | | Troll | | Trot with baits or snoods | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Bluefish..... | 1,062,448 | \$31,076 | | | 446,992 | \$20,176 | | |
| Bonito..... | 8,180 | 413 | | | 9,832 | 734 | | |
| Cero..... | 2,500 | 75 | | | | | | |
| Cod..... | 28,355 | 834 | 2,543,028 | \$80,294 | | | | |
| Croaker..... | 20,500 | 930 | | | | | | |
| Dolphin..... | | | | | 111 | 9 | | |
| Eels, common..... | 2,200 | 141 | | | | | | |
| Flounders..... | 12,769 | 464 | 6,533 | 357 | | | | |
| Grayfish..... | | | 1,500 | 30 | | | | |
| Hake..... | | | 409 | 14 | | | | |
| King whiting or "kingfish"..... | 31 | 3 | | | | | | |
| Scup or porgy..... | 31,458 | 393 | | | | | | |
| Sea bass..... | 342,430 | 16,017 | | | | | | |
| Sharks..... | | | 539 | 8 | | | | |
| Skates..... | | | 4,838 | 25 | | | | |
| Spanish mackerel..... | 5,600 | 168 | | | | | | |
| Squeteague or "sea trout", gray..... | 60,518 | 2,659 | | | | | | |
| Striped bass..... | 400 | 40 | | | | | | |
| Tautog..... | 5,635 | 412 | | | | | | |
| Tuna or "horse mackerel"..... | 1,063 | 81 | | | 3,324 | 157 | | |
| Crabs: | | | | | | | | |
| Hard..... | | | | | | | 102,102 | \$2,913 |
| Soft..... | | | | | | | 1,200 | 225 |
| Total..... | 1,584,385 | 63,706 | 2,556,847 | 80,728 | 460,259 | 21,076 | 103,302 | 3,138 |

| Species | Pound nets | | Weirs | | Stop nets | | Fyke nets | |
|---------------------------------------|-------------------|----------------|----------------|--------------|----------------|---------------|----------------|--------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | | | | | | | 3,740 | \$35 |
| Bluefish..... | 335,143 | \$12,752 | | | | | | |
| Bonito..... | 69,281 | 2,878 | | | | | | |
| Butterfish..... | 2,811,867 | 114,175 | | | | | | |
| Carp..... | | | | | 107,200 | \$11,996 | | |
| Catfish and bullheads..... | | | | | 2,000 | 120 | 34,080 | 2,286 |
| Cod..... | 642,251 | 18,207 | | | | | | |
| Crevalle..... | 7,254 | 141 | | | | | | |
| Croaker..... | 786,492 | 21,271 | | | | | | |
| Drum, red or redfish..... | 6,498 | 101 | | | | | | |
| Eels: | | | | | | | | |
| Common..... | 20,702 | 1,914 | | | | | 30,562 | 2,706 |
| Conger..... | 2,029 | 112 | | | | | | |
| Flounders..... | 386,831 | 20,237 | | | | | 68,283 | 1,771 |
| Frigate mackerel..... | 19,754 | 243 | | | | | | |
| Goosefish..... | 10,000 | 38 | | | | | | |
| Grayfish..... | 5,239 | 105 | | | | | | |
| Herring, sea..... | 590,465 | 5,171 | | | | | | |
| King whiting or "kingfish"..... | 72,140 | 10,437 | | | | | | |
| Mackerel..... | 119,861 | 3,942 | | | | | | |
| Menhaden..... | 2,325,061 | 42,150 | | | | | 500 | 100 |
| Mummichog..... | | | | | | | | |
| Pollock..... | 6,280 | 94 | | | | | | |
| Pompano..... | 317 | 77 | | | | | | |
| Scup or porgy..... | 3,708,999 | 39,041 | | | | | 55 | 1 |
| Sea bass..... | 198,523 | 6,622 | | | | | 4,237 | 150 |
| Sea robin..... | 10,805 | 109 | | | | | | |
| Shad..... | 63,109 | 5,599 | | | | | | |
| Sharks..... | 9,071 | 195 | | | | | | |
| Skates..... | 74,976 | 632 | | | | | | |
| Spanish mackerel..... | 814 | 118 | | | | | | |
| Spot..... | 287,662 | 4,565 | | | | | | |
| Squeteagues or "sea trout", gray..... | 6,140,500 | 162,415 | | | | | | |
| Squid..... | 52,765 | 402 | | | | | | |
| Striped bass..... | 420 | 31 | | | | | 1,500 | 423 |
| Sturgeon..... | 372 | 70 | | | | | | |
| Suckers..... | | | | | 975 | 67 | | |
| Tautog..... | 66,969 | 9,533 | | | | | | |
| Thimble-eyed mackerel..... | 25,175 | 368 | | | | | | |
| Tuna or "horse mackerel"..... | 32,051 | 1,416 | | | | | | |
| White perch..... | 500 | 25 | | | | | 13,120 | 954 |
| Whiting..... | 2,040,499 | 20,133 | | | | | | |
| Crabs: | | | | | | | | |
| Hard..... | 9,600 | 64 | | | | | | |
| King..... | 1,269,600 | 3,326 | 893,336 | \$2,485 | | | 7,500 | 80 |
| Lobsters..... | 2,610 | 470 | | | | | | |
| Squid..... | 337,000 | 8,480 | | | | | | |
| Turtles, snapper..... | | | | | | | 1,750 | 175 |
| Total..... | 22,630,465 | 517,698 | 893,336 | 2,485 | 110,175 | 12,183 | 165,327 | 8,681 |

Fisheries of New Jersey, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Dip nets | | Cast nets | | Drag nets | | Otter trawls | | Wire baskets | |
|-----------------------------|----------|---------|-----------|-------|-----------|---------|--------------|---------|--------------|-------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Bluefish | | | | | | | 10,550 | \$715 | | |
| Butterfish | | | | | | | 97,924 | 5,046 | | |
| Carp | | | 400 | \$48 | | | | | | |
| Cod | | | | | | | 16,262 | 353 | | |
| Croaker | | | | | | | 926,016 | 19,179 | | |
| Drum, red or redfish | | | | | | | 2,272 | 25 | | |
| Eels, conger | | | | | | | 9,134 | 320 | | |
| Flounders | | | | | | | 3,528,407 | 145,053 | | |
| Hake | | | | | | | 21,762 | 346 | | |
| Kingfish or "king mackerel" | | | | | | | 80 | 4 | | |
| King whiting or "kingfish" | | | | | | | 10,515 | 404 | | |
| Mackerel | | | | | | | 7,737 | 405 | | |
| Menhaden | | | | | | | 356 | 2 | | |
| Mullet | | | | | | | 267 | 8 | | |
| Pigfish | | | | | | | 203 | 5 | | |
| Scup or porgy | | | | | | | 731,349 | 23,236 | | |
| Sea bass | | | | | | | 267,214 | 9,889 | | |
| Sheephead | | | | | | | 9 | 1 | | |
| Skates | | | | | | | 3,162 | 55 | | |
| Spot | | | | | | | 9,883 | 178 | | |
| Squeteague or "sea trout": | | | | | | | | | | |
| Gray | | | | | | | 136,072 | 6,222 | | |
| Spotted | | | | | | | 30 | 2 | | |
| Striped bass | | | | | | | 27 | 4 | | |
| Sturgeon | | | | | | | 25,843 | 1,962 | | |
| Tautog | | | | | | | 1,617 | 34 | | |
| Tilfish | | | | | | | 25 | 1 | | |
| Crabs: | | | | | | | | | | |
| Hard | 17,658 | \$1,778 | | | | | 60 | 1 | | |
| Soft | 31,839 | 7,649 | | | | | | | 1,341 | \$168 |
| Lobsters | | | | | | | 1,749 | 215 | | |
| Shrimp | | | | | 9,500 | \$2,375 | 15,841 | 873 | | |
| Periwinkles and cockles | | | | | | | 100 | 2 | | |
| Scallops: | | | | | | | | | | |
| Bay | | | | | | | 594 | 30 | | |
| Sea | | | | | | | 1,589 | 102 | | |
| Squid | | | | | | | 10,312 | 280 | | |
| Total | 49,197 | 9,427 | 400 | 48 | 9,500 | 2,375 | 5,826,951 | 214,952 | 1,341 | 168 |

| Species | Pots | | | | | | Harpoons | | Spears | |
|--------------|--------|-------|---------|---------|-----------|--------|----------|-------|---------|--------|
| | Crab | | Kel | | Lobster | | Pounds | Value | Pounds | Value |
| | Pounds | Value | Pounds | Value | Pounds | Value | | | | |
| Eels, common | | | 124,683 | \$9,899 | | | | | | |
| Flounders | | | | | 96 | \$5 | | | | |
| Mummichog | | | 15,325 | 1,394 | | | | | | |
| Sea bass | | | | | 1,166,302 | 30,906 | | | | |
| Silversides | | | 500 | 50 | | | | | | |
| Swordfish | | | | | | | 1,600 | \$128 | | |
| Tautog | | | | | 6,702 | 69 | | | | |
| Crabs: | | | | | | | | | | |
| Hard | 12,000 | \$750 | | | 2,280 | 438 | | | | |
| Soft | 4,800 | 900 | | | | | | | | |
| Lobsters | | | | | 359,881 | 66,944 | | | | |
| Total | 16,800 | 1,650 | 140,508 | 11,343 | 1,535,241 | 98,362 | 1,600 | 128 | 105,318 | 10,567 |

Fisheries of New Jersey, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Dredges | | | | | | | | Tongs | |
|-------------------------|---------|---------|---------|----------|-----------|---------|---------|----------|---------|-----------|
| | Clam | | Crab | | Oyster | | Scallop | | | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Crabs, hard | | | 561,753 | \$12,220 | | | | | | |
| Clams: | | | | | | | | | | |
| Hard, public | 23,443 | \$3,333 | | | 2,625 | \$275 | | | 760,636 | \$115,843 |
| Hard, private | 63,143 | 7,027 | | | 1,336 | 550 | | | 5,107 | 1,021 |
| Surf or skimmers | 140,063 | 5,482 | | | | | | | | |
| Total | 54 | 2 | | | | | | | | |
| Conchs: | | | | | | | | | | |
| Oysters: | | | | | | | | | | |
| Market, public, spring | | | | | | | | | 10,330 | 992 |
| Market, public, fall | | | | | | | | | 25,589 | 1,601 |
| Market, private, spring | | | | | 3,622,470 | 269,885 | | | 56,919 | 7,676 |
| Market, private, fall | | | | | 3,668,151 | 275,843 | | | 137,834 | 18,419 |
| Scallops, sea | | | | | | | 372,000 | \$22,300 | | |
| Total | 226,703 | 15,844 | 561,753 | 12,220 | 7,294,582 | 546,653 | 372,000 | 22,300 | 996,415 | 145,552 |

| Species | Rakes | | Forks | | Hoos | | Gaffs | | By hand | |
|-------------------------|---------|--------|--------|---------|---------|----------|--------|-------|---------|--------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Crabs: | | | | | | | | | | |
| Hard | 1,602 | \$48 | | | | | | | 13,336 | \$40 |
| King | | | | | | | | | | |
| Soft | 801 | 96 | | | | | | | | |
| Clams: | | | | | | | | | | |
| Hard, public | 531,283 | 81,480 | | | | | | | 87,217 | 14,548 |
| Hard, private | 875 | 123 | | | | | | | | |
| Soft, public | | | | | 913,560 | \$45,752 | | | | |
| Surf or skimmers | 6,250 | 250 | | | | | | | | |
| Mussels, sea | | | | | | | | | 5,200 | 200 |
| Oysters: | | | | | | | | | | |
| Market, public, spring | 198 | 96 | | | | | | | 438 | 50 |
| Market, public, fall | 580 | 141 | | | | | | | 437 | 50 |
| Market, private, spring | 38,725 | 5,272 | | | | | | | 425 | 50 |
| Market, private, fall | 47,963 | 6,756 | | | | | | | 1,616 | 270 |
| Terrapin, diamond-back | | | | | | | 750 | \$75 | 502 | 145 |
| Turtles, snapper | | | | | | | | | | |
| Bloodworms | | | 1,740 | \$1,505 | | | | | | |
| Sandworms | | | 5,403 | 4,645 | | | | | | |
| Total | 628,277 | 94,262 | 7,143 | 6,150 | 913,560 | 45,752 | 750 | 75 | 109,171 | 15,353 |

OPERATING UNITS: BY COUNTIES

| Item | Atlantic | Bergen | Burlington | Camden | Cape May | Cumberland | Gloucester |
|---------------------|----------|--------|------------|--------|----------|------------|------------|
| | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | |
| On vessels | 64 | 3 | 6 | | 288 | 593 | |
| On boats and shore: | | | | | | | |
| Regular | 217 | 8 | 34 | | 154 | 39 | 2 |
| Casual | 211 | 47 | 102 | 17 | 342 | 143 | 19 |
| Total | 492 | 58 | 142 | 17 | 784 | 775 | 21 |
| Vessels: | | | | | | | |
| Motor | 15 | 1 | 2 | | 51 | 97 | |
| Net tonnage | 225 | 15 | 22 | | 839 | 1,877 | |
| Boats: | | | | | | | |
| Motor | 184 | 13 | 62 | 1 | 192 | 46 | 5 |
| Other | 178 | 13 | 35 | 8 | 149 | 95 | 7 |
| Accessory boats | 7 | | | | 56 | | |

Fisheries of New Jersey, 1933—Continued

OPERATING UNITS: BY COUNTIES—Continued

| Item | Atlantic | Bergen | Burlington | Camden | Cape May | Cumberland | Gloucester |
|--------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| Apparatus: | | | | | | | |
| Purse seines: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| Menhaden..... | | | | | 1 | | |
| Length, yards..... | | | | | 456 | | |
| Other..... | 1 | | | | 6 | | |
| Length, yards..... | 360 | | | | 2,150 | | |
| Haul seines..... | 25 | | 20 | 4 | 10 | 8 | 3 |
| Length, yards..... | 2,045 | | 1,811 | 400 | 495 | 800 | 240 |
| Gill nets: | | | | | | | |
| Drift..... | 71 | 1 | 18 | 7 | 124 | 78 | |
| Square yards..... | 22,680 | 1,998 | 20,450 | 840 | 41,368 | 61,997 | |
| Runaround..... | 1 | | | | 4 | 4 | |
| Square yards..... | 4,800 | | | | 7,200 | 1,200 | |
| Stake..... | | 13 | 4 | | 9 | 69 | |
| Square yards..... | | 15,390 | 100 | | 1,300 | 8,120 | |
| Lines: | | | | | | | |
| Hand..... | 39 | | | | 209 | 16 | |
| Hooks..... | 65 | | | | 374 | 22 | |
| Trawl..... | 137 | | | | 303 | 9 | |
| Hooks..... | 82,100 | | | | 198,850 | 5,400 | |
| Troll..... | 4 | | | | 268 | | |
| Hooks..... | 4 | | | | 268 | | |
| Trot with baits or snoods..... | | | | | | 8 | |
| Baits or snoods..... | | | | | | 9,700 | |
| Pound nets..... | 2 | | | | 63 | | |
| Weirs..... | | | | | 62 | 26 | |
| Stop nets..... | | | 8 | 5 | | 8 | 10 |
| Square yards..... | | | 2,920 | 495 | | 7,500 | 18,600 |
| Fyke nets..... | 21 | | 278 | 15 | 121 | 174 | 40 |
| Dip nets..... | 4 | | | | 6 | 1 | |
| Cast nets..... | | | | | | | 1 |
| Otter trawls..... | 23 | | | | 40 | | |
| Yards at mouth..... | 540 | | | | 872 | | |
| Pots: | | | | | | | |
| Eel..... | 6 | 105 | | | 207 | 32 | |
| Lobster..... | | | | | 470 | | |
| Spears..... | 2 | | | | 3 | | |
| Dredges: | | | | | | | |
| Clam..... | 3 | | | | 6 | 14 | |
| Yards at mouth..... | 3 | | | | 6 | 18 | |
| Crab..... | | 12 | | | | 20 | |
| Yards at mouth..... | | 12 | | | | 30 | |
| Oyster..... | 10 | | 16 | | 2 | 192 | |
| Yards at mouth..... | 11 | | 16 | | 3 | 236 | |
| Scallop..... | 4 | | | | | | |
| Yards at mouth..... | 14 | | | | | | |
| Tongs..... | 304 | | 53 | | 165 | 58 | |
| Rakes..... | 201 | | 25 | | 136 | | |
| Hoes..... | 2 | | | | | | |
| Gaffs..... | | | | | | 2 | |

| Item | Hudson | Hunterdon | Mercer | Mid-dlesex | Mon-mouth | Ocean | Salem |
|------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| Fishermen: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| On vessels..... | 2 | | | 2 | 162 | 147 | |
| On boats and shore: | | | | | | | |
| Regular..... | | | | 6 | 298 | 193 | 3 |
| Casual..... | | 24 | 26 | 11 | 323 | 265 | 92 |
| Total..... | 2 | 24 | 26 | 19 | 783 | 605 | 95 |
| Vessels: | | | | | | | |
| Motor..... | 1 | | | 1 | 14 | 26 | |
| Net tonnage..... | 9 | | | 8 | 276 | 199 | |
| Sail..... | | | | | 3 | | |
| Net tonnage..... | | | | | 24 | | |
| Total vessels..... | 1 | | | 1 | 17 | 26 | |
| Total net tonnage..... | 9 | | | 8 | 300 | 199 | |
| Boats: | | | | | | | |
| Motor..... | | | | 7 | 200 | 281 | 38 |
| Other..... | | 5 | 7 | 3 | 157 | 68 | 16 |

Fisheries of New Jersey, 1933—Continued

OPERATING UNITS: BY COUNTIES—Continued

| Item | Hud-son | Hun-terdon | Mercer | Mid-diesex | Mon-mouth | Ocean | Salem |
|--------------------------------|---------|------------|--------|------------|-----------|--------|---------|
| | Number | Number | Number | Number | Number | Number | Number |
| Accessory boats..... | | | | | 25 | | |
| Apparatus: | | | | | | | |
| Purse seines: | | | | | | | |
| Menhaden..... | | | | | 7 | | |
| Length, yards..... | | | | | 2,350 | | |
| Haul seines..... | | 5 | 4 | | 13 | 5 | 8 |
| Length, yards..... | | 1,005 | 740 | | 253 | 281 | 970 |
| Gill nets: | | | | | | | |
| Anchor..... | | | | 3 | 7 | | |
| Square yards..... | | | | 2,500 | 3,330 | | |
| Drift..... | | | | | 275 | | 31 |
| Square yards..... | | | | | 106,320 | | 192,690 |
| Runaround..... | | | | 6 | 36 | 18 | |
| Square yards..... | | | | 15,895 | 98,800 | 82,800 | |
| Stake..... | | | | | 4 | 152 | |
| Square yards..... | | | | | 5,360 | 16,606 | |
| Lines: | | | | | | | |
| Hand..... | | | | | 151 | 46 | 2 |
| Hooks..... | | | | | 202 | 53 | 4 |
| Trawl..... | | | | | 97 | 166 | |
| Hooks..... | | | | | 50,100 | 83,950 | |
| Troll..... | | | | | 96 | 36 | |
| Hooks..... | | | | | 96 | 36 | |
| Trot with baits or snoods..... | | | | | | | 1 |
| Baits or snoods..... | | | | | | | 200 |
| Pound nets..... | | | | | 47 | 41 | |
| Stop nets..... | | | | | | | 33 |
| Square yards..... | | | | | | | 21,100 |
| Fyke nets..... | | | 15 | | 92 | 517 | 96 |
| Dip nets..... | | | | | 19 | 4 | |
| Cast nets..... | | | | | | | 2 |
| Drag nets..... | | | | | | 14 | |
| Yards at mouth..... | | | | | | 28 | |
| Otter trawls..... | 1 | | | | 3 | | |
| Yards at mouth..... | 27 | | | | 75 | | |
| Wire baskets..... | | | | | | 12 | |
| Pots: | | | | | | | 10 |
| Crab..... | | | | | 195 | 1,158 | 868 |
| Eel..... | | | | | 245 | 16,887 | 10,369 |
| Lobster..... | | | | | | | 1 |
| Harpoons..... | | | | | | | 13 |
| Spears..... | | | | | 74 | | |
| Dredges: | | | | | | | |
| Clam..... | | | | 4 | 16 | | |
| Yards at mouth..... | | | | 5 | 16 | | |
| Crab..... | | | | | 37 | | |
| Yards at mouth..... | | | | | 48 | | |
| Oyster..... | | | | | | 39 | |
| Yards at mouth..... | | | | | | 43 | |
| Scallop..... | | | | | 6 | | |
| Yards at mouth..... | | | | | 20 | | |
| Tongs..... | | | | | 4 | 199 | |
| Rakes..... | | | | | 36 | 111 | |
| Forks..... | | | | 3 | 48 | | |
| Hoos..... | | | | | 174 | | |

CATCH: BY COUNTIES

| Species | Atlantic | | Bergen | | Burlington | | Camden | | Cape May | |
|----------------------------|----------|---------|--------|-------|------------|-------|--------|-------|-----------|----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | | | | | 150 | \$4 | | | | |
| Bluefish..... | 175,050 | \$7,072 | | | | | | | 599,782 | \$25,844 |
| Bonito..... | 550 | 22 | | | | | | | 15,626 | 849 |
| Butterfish..... | 9,270 | 309 | | | | | | | 764,404 | 24,888 |
| Carp..... | | | | | 15,471 | 1,661 | 8,500 | \$850 | | |
| Catfish and bullheads..... | 4,050 | 122 | | | 14,470 | 1,300 | 1,000 | 80 | | |
| Cero..... | | | | | | | | | 2,500 | 75 |
| Cod..... | 553,995 | 21,149 | | | | | | | 1,256,165 | 39,697 |
| Croaker..... | 67,548 | 1,198 | | | 1,050 | 21 | | | 1,121,482 | 21,814 |
| Dolphin..... | | | | | | | | | 111 | 9 |
| Drum, red or redfish..... | | | | | | | | | 5,272 | 55 |

Fisheries of New Jersey, 1933—Continued

CATCH: BY COUNTIES—Continued

| Species | Atlantic | | Bergen | | Burlington | | Camden | | Cape May | |
|-----------------------------|-----------|---------|---------|--------|------------|--------|--------|-------|------------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Eels: | | | | | | | | | | |
| Common | 68,100 | \$4,236 | 6,500 | \$303 | 8,112 | \$610 | 1,000 | \$160 | 23,108 | \$2,543 |
| Conger | 3,771 | 72 | | | | | | | 5,363 | 248 |
| Flounders | 1,182,122 | 47,772 | | | | | | | 2,323,790 | 98,570 |
| Frigate mackerel | | | | | | | | | 405 | 4 |
| Hake | 6,066 | 93 | | | | | | | 15,705 | 254 |
| Herring: | | | | | | | | | | |
| Round | 750 | 8 | | | | | | | | |
| Sea | 7,500 | 75 | | | | | | | 9,580 | 94 |
| Kingfish or "king mackerel" | 60 | 1 | | | | | | | 80 | 4 |
| King whiting or "kingfish" | 2,980 | 136 | | | | | | | 31,431 | 1,205 |
| Mackerel | 6,453 | 224 | | | | | | | 12,290 | 673 |
| Menhaden | | | | | | | | | 8,603,761 | 21,545 |
| Mullet | 27,600 | 1,656 | | | | | | | 51,267 | 4,158 |
| Mummichog | 8,000 | 200 | | | | | 500 | 100 | 2,225 | 84 |
| Pigfish | | | | | | | | | 263 | 5 |
| Pollock | | | | | | | | | 380 | 10 |
| Pompano | 200 | 24 | | | | | | | | |
| Scup or porgy | 76,367 | 1,299 | | | | | | | 2,095,045 | 30,147 |
| Sea bass | 155,448 | 4,939 | | | | | | | 467,152 | 22,085 |
| Shad | 1,960 | 124 | 171,024 | 12,573 | 11,249 | 2,020 | 2,330 | 405 | 10,818 | 931 |
| Sharks | | | | | | | | | 2,665 | 26 |
| Sheepshead | | | | | | | | | 9 | 1 |
| Skates | | | | | | | | | | |
| Spanish mackerel | 5 | 1 | | | | | | | 12,904 | 122 |
| Spot | 2,434 | 89 | | | | | | | 5,675 | 180 |
| Squeteagues or "sea trout": | | | | | | | | | 31,767 | 811 |
| Gray | 295,903 | 5,673 | | | 2,562 | 143 | | | 2,353,679 | 54,400 |
| Spotted | | | | | | | | | 30 | 2 |
| Squirrel hake | | | | | | | | | 3,000 | 10 |
| Striped bass | 2,037 | 492 | | | 650 | 195 | | | 610 | 61 |
| Sturgeon | 131 | 9 | | | | | | | 26,074 | 2,021 |
| Suckers | 850 | 26 | | | 59,100 | 4,613 | 150 | 11 | | |
| Tautog | 224 | 3 | | | | | | | 2,130 | 46 |
| Thimble-eyed mackerel | | | | | | | | | 9,004 | 92 |
| Tilefish | | | | | | | | | 25 | 1 |
| Tuna or "horse mackerel" | 192 | 4 | | | | | | | 3,376 | 160 |
| White perch | 6,850 | 568 | | | 2,100 | 210 | | | | |
| Whiting | 3,000 | 15 | | | | | | | 10,964 | 64 |
| Crabs: | | | | | | | | | | |
| Hard | 6,525 | 337 | 23,001 | 690 | | | | | 3,471 | 177 |
| King | | | | | | | | | 1,920,272 | 5,243 |
| Soft | 942 | 113 | | | | | | | 933 | 112 |
| Lobsters | 554 | 79 | | | | | | | 1,195 | 136 |
| Shrimp | 841 | 123 | | | | | | | 15,000 | 750 |
| Clams: | | | | | | | | | | |
| Hard, public | 446,134 | 71,434 | | | 94,106 | 13,229 | | | 285,557 | 53,333 |
| Hard, private | 375 | 165 | | | 1,094 | 154 | | | | |
| Soft, public | 2,000 | 250 | | | | | | | | |
| Surf or skimmer | 77,563 | 2,982 | | | | | | | | |
| Conchs | | | | | | | | | 68,750 | 2,750 |
| Mussels, sea | | | | | | | | | 54 | 2 |
| Oysters: | | | | | | | | | 5,200 | 200 |
| Market, public, spring | | | | | | | | | 2,796 | 546 |
| Market, public, fall | 425 | 58 | | | | | | | 197 | 96 |
| Market, private, spring | 113,900 | 14,992 | | | 108,801 | 13,500 | | | 21,468 | 1,907 |
| Market, private, fall | 199,291 | 26,474 | | | 108,801 | 13,500 | | | 18,508 | 1,626 |
| Periwinkles and cockles | | | | | | | | | 100 | 2 |
| Scallops: | | | | | | | | | | |
| Bay | | | | | | | | | 594 | 30 |
| Sea | 150,170 | 7,542 | | | | | | | 1,419 | 60 |
| Squid | 564 | 18 | | | | | | | 198,648 | 5,179 |
| Total | 3,669,259 | 222,278 | 200,525 | 13,566 | 427,716 | 51,060 | 13,480 | 1,596 | 22,424,319 | 425,944 |

Fisheries of New Jersey, 1933—Continued

CATCH: BY COUNTIES—Continued

| Species | Cumberland | | Gloucester | | Hudson | | Hunterdon | | Mercer | |
|---------------------------------------|------------|---------|------------|-------|--------|---------|-----------|-------|--------|-------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | 7,800 | \$170 | 2,000 | \$30 | | | | | | |
| Bluefish..... | 5,000 | 440 | | | | | | | | |
| Carp..... | 13,750 | 1,316 | 28,950 | 3,764 | | | 350 | \$35 | 1,100 | 132 |
| Catfish and bullheads..... | 14,780 | 823 | 7,000 | 210 | | | | | | |
| Cod..... | 13,334 | 600 | | | | | | | | |
| Croaker..... | 20,500 | 760 | | | | | | | | |
| Eels, common..... | 11,200 | 696 | 700 | 70 | | | | | 1,750 | 300 |
| Flounders..... | | | | | 50,000 | \$1,500 | | | | |
| Shad..... | 45,814 | 6,310 | 410 | 43 | | | 25,359 | 2,500 | 31,833 | 3,969 |
| Spot..... | 65,000 | 1,730 | | | | | | | | |
| Squeteagues or "sea trout", gray..... | 103,064 | 2,478 | | | | | | | | |
| Striped bass..... | 5,055 | 816 | | | | | | | | |
| Sturgeon..... | 498 | 98 | | | | | | | | |
| Suckers..... | | | 400 | 28 | | | 8,000 | 480 | 34,000 | 3,400 |
| White perch..... | 10,500 | 1,186 | | | | | | | | |
| Crabs: | | | | | | | | | | |
| Hard..... | 471,567 | 7,073 | | | | | | | | |
| King..... | 256,000 | 608 | | | | | | | | |
| Soft..... | 876 | 131 | | | | | | | | |
| Clams: | | | | | | | | | | |
| Hard, public..... | 4,375 | 930 | | | | | | | | |
| Hard, private..... | 63,143 | 7,027 | | | | | | | | |
| Oysters: | | | | | | | | | | |
| Market, public, spring..... | 7,200 | 480 | | | | | | | | |
| Market, public, fall..... | 23,400 | 1,235 | | | | | | | | |
| Market, private, spring..... | 3,408,582 | 241,928 | | | | | | | | |
| Market, private, fall..... | 3,412,176 | 242,167 | | | | | | | | |
| Terrapin, diamond-back..... | 502 | 145 | | | | | | | | |
| Turtles, snapper..... | 2,500 | 250 | | | | | | | | |
| Total..... | 7,966,596 | 519,397 | 39,460 | 4,145 | 50,000 | 1,500 | 33,709 | 3,018 | 68,983 | 7,806 |

| Species | Middlesex | | Monmouth | | Ocean | | Salem | |
|---------------------------------------|-----------|---------|------------|--------|-----------|--------|--------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | | | 1,500 | \$10 | 2,240 | \$25 | | |
| Bluefish..... | 81,931 | \$3,647 | 678,703 | 26,328 | 574,838 | 12,815 | | |
| Bonito..... | | | 24,701 | 1,325 | 47,530 | 1,907 | | |
| Butterfish..... | | | 556,743 | 23,459 | 1,681,788 | 70,573 | | |
| Carp..... | | | | | | | 75,350 | \$8,289 |
| Catfish and bullheads..... | | | | | | | 14,100 | 786 |
| Cod..... | | | 572,666 | 17,035 | 833,726 | 21,207 | | |
| Crevalle..... | | | 2,200 | 39 | 5,054 | 102 | | |
| Croaker..... | 15,031 | 489 | 194,369 | 10,820 | 472,063 | 9,443 | 9,000 | 400 |
| Drum, red or redfish..... | | | 538 | 11 | 2,960 | 60 | | |
| Eels: | | | | | | | | |
| Common..... | 4,375 | 384 | 158,753 | 15,196 | 51,868 | 3,837 | 3,000 | 290 |
| Conger..... | | | 2,029 | 112 | | | | |
| Flounders..... | | | 367,039 | 18,149 | 128,846 | 4,610 | | |
| Frigate mackerel..... | | | 13,911 | 139 | 5,438 | 100 | | |
| Goosefish..... | | | 10,000 | 38 | | | | |
| Grayfish..... | | | 6,739 | 135 | | | | |
| Hake..... | | | 400 | 13 | | | | |
| Herring, sea..... | | | 307,225 | 3,072 | 266,160 | 1,930 | | |
| King whiting or "kingfish"..... | | | 20,867 | 8,151 | 28,761 | 1,440 | | |
| Mackerel..... | | | 254,300 | 6,993 | 45,004 | 1,802 | | |
| Menhaden..... | | | 36,967,308 | 99,897 | 203,048 | 1,487 | | |
| Mullet..... | | | 3,880 | 420 | | | | |
| Mummichog..... | 3,650 | 365 | 11,350 | 1,170 | 200 | 26 | | |
| Pollock..... | | | 3,195 | 42 | 2,705 | 42 | | |
| Pompano..... | | | 117 | 53 | | | | |
| Scup or porgy..... | | | 1,308,771 | 19,836 | 1,401,181 | 14,011 | | |
| Sea bass..... | | | 224,006 | 8,200 | 1,122,457 | 28,376 | | |
| Sea robin..... | | | 4,705 | 47 | 6,100 | 62 | | |
| Shad..... | 276 | 52 | 119,267 | 11,519 | 21,470 | 1,900 | 16,648 | 2,835 |
| Sharks..... | | | 6,645 | 177 | | | | |
| Silversides..... | | | 3,930 | 426 | 900 | 250 | | |
| Skates..... | | | 38,900 | 453 | 31,172 | 137 | | |
| Spanish mackerel..... | | | 232 | 26 | 507 | 77 | | |
| Spot..... | | | 219,014 | 2,777 | 1,555,850 | 1,561 | | |
| Squeteagues or "sea trout", gray..... | 27,594 | 1,538 | 1,463,449 | 51,055 | 2,687,505 | 70,090 | 3,625 | 220 |

Fisheries of New Jersey, 1935—Continued

CATCH: BY COUNTIES—Continued

| Species | Middlesex | | Monmouth | | Ocean | | Salem | |
|-------------------------------|----------------|--------------|-------------------|----------------|-------------------|----------------|----------------|---------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Squirrel hake..... | | | 36,235 | \$241 | 14,632 | \$145 | | |
| Striped bass..... | | | | | 480 | 120 | 320 | \$96 |
| Sturgeon..... | | | 10 | 2 | | | | |
| Suckers..... | | | | | | | 75 | 4 |
| Swordfish..... | | | 1,600 | 128 | | | | |
| Tautog..... | | | 40,290 | 9,283 | 28,279 | 716 | | |
| Thimble-eyed mackerel..... | | | 8,908 | 129 | 7,563 | 153 | | |
| Tuna or "horse mackerel"..... | | | 17,991 | 932 | 14,933 | 560 | | |
| White perch..... | | | 20 | 2 | 9,950 | 700 | | |
| Whiting..... | | | 1,284,816 | 12,769 | 741,764 | 7,286 | | |
| Crabs: | | | | | | | | |
| Hard..... | | | 188,646 | 8,616 | 1,200 | 525 | 15,000 | 938 |
| King..... | | | 7,600 | 60 | | | | |
| Soft..... | | | 43,283 | 9,780 | 5,349 | 978 | 6,000 | 1,125 |
| Lobsters..... | 848 | \$213 | 342,123 | 62,526 | 19,500 | 4,675 | | |
| Shrimp..... | | | | | 9,900 | 2,455 | | |
| Clams: | | | | | | | | |
| Hard, public..... | 9,513 | 1,440 | 101,346 | 12,409 | 464,173 | 62,804 | | |
| Hard, private..... | | | 2,700 | 620 | 2,649 | 735 | | |
| Soft, public..... | | | 911,560 | 45,502 | | | | |
| Oysters: | | | | | | | | |
| Market, public, spring..... | | | | | 970 | 112 | | |
| Market, public, fall..... | | | | | 2,584 | 403 | | |
| Market, private, spring..... | | | 750 | 150 | 65,038 | 10,406 | | |
| Market, private, fall..... | | | 750 | 150 | 116,038 | 17,371 | | |
| Scallops, sea..... | | | 222,000 | 14,800 | | | | |
| Squid..... | | | 36,145 | 1,322 | 111,955 | 2,241 | | |
| Bloodworms..... | 6 | 8 | 1,734 | 1,497 | | | | |
| Sandworms..... | 3 | 3 | 5,400 | 4,642 | | | | |
| Total..... | 143,227 | 8,069 | 46,788,317 | 512,706 | 11,295,728 | 360,239 | 142,118 | 14,983 |

SEED OYSTER FISHERY: BY GEAR

| Item | Dredges | | Tongs | | Rakes | | By hand | | Total, exclusive of duplication |
|---------------------------|----------------|----------------|----------------|---------------|----------------|--------------|----------------|--------------|---------------------------------|
| OPERATING UNITS | | | | | | | | | |
| Fishermen: | <i>Number</i> | | <i>Number</i> | | <i>Number</i> | | <i>Number</i> | | <i>Number</i> |
| On vessels..... | 1,514 | | | | | | | | 1,514 |
| On boats and shore: | | | | | | | | | |
| Regular..... | | | 48 | | 20 | | | | 68 |
| Casual..... | | | 92 | | 14 | | 1 | | 107 |
| Total..... | 1,514 | | 140 | | 34 | | 1 | | 1,689 |
| Vessels, all..... | 139 | | | | | | | | 139 |
| Net tonnage..... | 2,881 | | | | | | | | 2,881 |
| Boats: | | | | | | | | | |
| V Motor..... | | | 60 | | 26 | | | | 86 |
| Other..... | | | 73 | | 8 | | | | 81 |
| Apparatus: | | | | | | | | | |
| Number..... | 278 | | 140 | | 34 | | | | |
| Yards at mouth..... | 333 | | | | | | | | |
| CATCH | | | | | | | | | |
| Oysters: | <i>Bushels</i> | <i>Value</i> | <i>Bushels</i> | <i>Value</i> | <i>Bushels</i> | <i>Value</i> | <i>Bushels</i> | <i>Value</i> | <i>Bushels</i> <i>Value</i> |
| Seed, public, spring..... | 860,500 | \$216,125 | 66,855 | \$14,282 | 8,545 | \$2,125 | 100 | \$50 | 926,000 \$232,582 |
| Seed, public, fall..... | | | 11,000 | 2,750 | | | | | 11,000 2,750 |
| Total..... | 860,500 | 216,125 | 67,865 | 17,032 | 8,545 | 2,125 | 100 | 50 | 937,000 235,332 |

Fisheries of New Jersey, 1933—Continued

SEED OYSTER FISHERY: BY COUNTIES

| Item | Atlantic | | Burlington | | Cumberland | | Ocean | |
|---------------------------|----------------|--------------|----------------|--------------|----------------|----------------|----------------|--------------|
| OPERATING UNITS | | | | | | | | |
| Fishermen: | <i>Number</i> | | <i>Number</i> | | <i>Number</i> | | <i>Number</i> | |
| On vessels..... | | | | | 1,514 | | | |
| On boats and shore: | | | | | | | | |
| Regular..... | 26 | | 20 | | 20 | | 2 | |
| Casual..... | 3 | | 30 | | 70 | | 4 | |
| Total..... | 29 | | 50 | | 1,604 | | 6 | |
| Vessels, sail..... | | | | | 139 | | | |
| Net tonnage..... | | | | | 2,881 | | | |
| Boats: | | | | | | | | |
| Motor..... | 23 | | 45 | | 14 | | 4 | |
| Other..... | 5 | | 3 | | 72 | | 1 | |
| Apparatus: | | | | | | | | |
| Dredges..... | | | | | 278 | | | |
| Yards at mouth..... | | | | | 333 | | | |
| Tongs..... | 19 | | 26 | | 90 | | 5 | |
| Rakes..... | 10 | | 24 | | | | | |
| CATCH | | | | | | | | |
| Oysters: | <i>Bushels</i> | <i>Value</i> | <i>Bushels</i> | <i>Value</i> | <i>Bushels</i> | <i>Value</i> | <i>Bushels</i> | <i>Value</i> |
| Seed, public, spring..... | 5,775 | \$1,427 | 13,150 | \$3,287 | 905,650 | \$227,455 | 1,425 | \$413 |
| Seed, public, fall..... | | | | | 11,000 | 2,750 | | |
| Total..... | 5,775 | 1,427 | 13,150 | 3,287 | 916,650 | 230,205 | 1,425 | 413 |

NOTE.—Of the total number of persons fishing for seed oysters, 1,583 are duplicated among those fishing for market oysters of other species. Similarly the following craft and gear are duplicated; 80 sail vessels, 79 motor boats, 61 other boats, 160 dredges, and 108 tongs.

PENNSYLVANIA

Fisheries of Pennsylvania, 1933

OPERATING UNITS: BY GEAR

| Item | Haul seines, drift | Gill nets | Total, exclusive of duplication |
|--|--------------------|---------------|---------------------------------|
| | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| Fishermen, on boats and shore, casual..... | 39 | 14 | 53 |
| Boats: | | | |
| Motor..... | | 5 | 5 |
| Other..... | 12 | 2 | 14 |
| Apparatus: | | | |
| Number..... | 12 | 7 | |
| Length, yards..... | 1,635 | | |
| Square yards..... | | 5,200 | |

CATCH: BY GEAR

| Species | Haul seines | | Gill nets, drift | |
|-------------------|---------------|--------------|------------------|--------------|
| | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> |
| Alewives..... | 8,775 | \$88 | | |
| Carp..... | 640 | 64 | | |
| Shad..... | 1,352 | 251 | 865 | \$141 |
| Suckers..... | 40,600 | 2,436 | | |
| Total..... | 51,367 | 2,839 | 865 | 141 |

Fisheries of Pennsylvania, 1933—Continued

OPERATING UNITS: BY COUNTIES

| Item | Bucks | Philadel- phia |
|---|--------|-------------------|
| | Number | Number |
| Fishermen, on boats and shores, casual..... | 43 | 10 |
| Boats: | | |
| Motor..... | | 5 |
| Other..... | 14 | |
| Apparatus: | | |
| Haul seines..... | 12 | |
| Length, yards..... | 1,635 | |
| Gill nets, drift..... | 2 | 5 |
| Square yards..... | 900 | 4,300 |

CATCH: BY COUNTIES

| Species | Bucks | | Philadelphia | |
|---------------|--------|-------|--------------|-------|
| | Pounds | Value | Pounds | Value |
| Alewives..... | 8,775 | \$88 | | |
| Carp..... | 640 | 64 | | |
| Shad..... | 1,515 | 278 | 702 | \$114 |
| Buckers..... | 40,600 | 2,436 | | |
| Total..... | 51,530 | 2,866 | 702 | 114 |

DELAWARE

Fisheries of Delaware, 1933

OPERATING UNITS: BY GEAR

| Item | Purse seiner, men- haden | Haul seines | Gill nets | | | Lines | | Pound nets | Stop nets |
|-----------------------------|-----------------------------------|----------------|-----------|----------------|--------|--------|--------|---------------|--------------|
| | | | Drift | Run- around | Stake | Hand | Trawl | | |
| Fishermen: | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| On vessels..... | 282 | | | | | | | | |
| On boats and shore: | | | | | | | | | |
| Regular..... | | 13 | 11 | 12 | 6 | 7 | 2 | | |
| Casual..... | | 217 | 82 | 32 | 18 | 5 | | 24 | 12 |
| Total..... | 282 | 230 | 93 | 44 | 24 | 12 | 2 | 24 | 12 |
| Vessels: | | | | | | | | | |
| Steam..... | 10 | | | | | | | | |
| Net tonnage..... | 1,237 | | | | | | | | |
| Boats: | | | | | | | | | |
| Motor..... | | | 25 | 16 | 4 | 5 | | 1 | |
| Other..... | | 61 | 26 | 15 | 12 | 7 | 2 | 11 | 7 |
| Accessory boats..... | 30 | | | | | | | | |
| Apparatus: | | | | | | | | | |
| Number..... | 10 | 61 | 63 | 33 | 99 | 12 | 2 | 32 | 7 |
| Length, yards..... | 3,140 | 18,990 | | | | | | | |
| Square yards..... | | | 145,720 | 41,450 | 7,770 | | | | 1,820 |
| Hooks, baits or snoods..... | | | | | | 24 | 1,300 | | |

Fisheries of Delaware, 1933—Continued

OPERATING UNITS: BY GEAR—Continued

| Item | Fyke nets | Dip nets | Pots | | Dredges | | | Tongs | By hand | Total, exclusive of duplication |
|---------------------|-----------|----------|--------|---------|---------|--------|--------|--------|---------|---------------------------------|
| | | | Eel | Lobster | Clam | Crab | Oyster | | | |
| | Number | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | | | | 350 |
| On vessels | | | | | 41 | 19 | 41 | | | |
| On boats and shore: | | | | | | | | | | |
| Regular | 19 | 10 | 11 | | | | | | | 34 |
| Casual | 43 | 41 | 17 | 8 | 29 | | | 32 | 24 | 478 |
| Total | 62 | 51 | 28 | 8 | 70 | 19 | 41 | 32 | 24 | 862 |
| Vessels: | | | | | | | | | | |
| Steam | | | | | | | | | | 10 |
| Net tonnage | | | | | | | | | | 1,237 |
| Motor | | | | | 12 | 6 | 7 | | | 16 |
| Net tonnage | | | | | 153 | 99 | 110 | | | 229 |
| Sail | | | | | | | 1 | | | 1 |
| Net tonnage | | | | | | | 8 | | | 8 |
| Total vessels | | | | | 12 | 6 | 8 | | | 27 |
| Total net tonnage | | | | | 163 | 99 | 118 | | | 1,474 |
| Boats: | | | | | | | | | | |
| Motor | 8 | | 3 | 4 | 13 | | | 8 | | 77 |
| Other | 40 | 23 | 16 | | | | | 24 | | 181 |
| Apparatus: | | | | | | | | | | |
| Number | 449 | 51 | 876 | 165 | 33 | 11 | 16 | 32 | | |
| Yards at mouth | | | | | 47 | 20 | 23 | | | |

CATCH: BY GEAR

| Species | Purse seines, menhaden | | Haul seines | | Gill nets | | | | | |
|----------------------------------|------------------------|-----------|-------------|---------|-----------|-------|------------|-------|---------|---------|
| | | | | | Drift | | Run-around | | Stake | |
| | | | | | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives | | | 879,720 | \$2,179 | 17,260 | \$183 | | | 215,460 | \$2,056 |
| Bluefish | | | | | 3,400 | 170 | | | 420 | 22 |
| Carp | | | 11,240 | 685 | | | | | | |
| Catfish and bullheads | | | 420 | 20 | | | | | | |
| Croaker | | | 31,160 | 487 | 19,260 | 307 | 11,540 | \$180 | 680 | 9 |
| Flounders | | | 1,130 | 62 | | | | | 6,240 | 326 |
| Menhaden | 33,600,000 | \$111,700 | | | | | | | | |
| Mullet | | | | | | | 233,000 | 2,390 | | |
| Shad | | | 2,310 | 242 | 19,720 | 1,715 | | | 250 | 34 |
| Spot | | | 13,920 | 373 | 15,600 | 478 | 2,400 | 52 | | |
| Squeteagues or "sea trout", gray | | | 90,580 | 2,565 | 12,400 | 184 | 9,290 | 144 | 9,400 | 415 |
| Striped bass | | | 2,220 | 301 | 860 | 105 | | | 7,540 | 861 |
| Sturgeon | | | | | 1,200 | 300 | | | | |
| White perch | | | 14,380 | 725 | 1,620 | 90 | | | 4,900 | 222 |
| Whiting | | | 480 | 30 | | | | | | |
| Yellow perch | | | 4,740 | 294 | | | | | | |
| Total | 33,600,000 | 111,700 | 1,052,300 | 7,963 | 91,320 | 3,532 | 256,230 | 2,766 | 244,980 | 3,945 |

| Species | Lines | | | | Pound nets | | Stop nets | |
|----------------------------------|--------|-------|--------|-------|------------|-------|-----------|-------|
| | Hand | | Trawl | | Pounds | Value | Pounds | Value |
| | Pounds | Value | Pounds | Value | | | | |
| Alewives | | | | | 22,400 | \$150 | | |
| Bluefish | 900 | \$44 | | | | | | |
| Carp | | | | | | | 2,220 | \$136 |
| Catfish and bullheads | | | | | | | 180 | 12 |
| Cod | | | | | | | | |
| Croaker | 4,490 | 85 | | | | | | |
| Drum, red or redfish | 520 | 16 | | | | | | |
| Eels, common | | | | | 840 | 59 | | |
| Flounders | | | | | 240 | 12 | | |
| Squeteagues or "sea trout", gray | 1,450 | 33 | | | | | | |
| Striped bass | | | | | 360 | 38 | | |
| White perch | | | | | 3,240 | 154 | | |
| Crabs, king | | | | | 506,600 | 529 | | |
| Total | 7,360 | 178 | 2,240 | 69 | 533,650 | 942 | 2,400 | 148 |

Fisheries of Delaware, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Fyke nets | | Dip nets | | Pots | | | |
|-----------------------------|-----------|--------|----------|----------|---------|----------|---------|----------|
| | Pounds | Value | Pounds | Value | Eel | | Lobster | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alwives..... | 14, 140 | \$218 | | | | | | |
| Carp..... | 12, 960 | 768 | | | | | | |
| Catfish and bullheads..... | 1, 245 | 84 | | | | | | |
| Eels, common..... | 9, 000 | 572 | | | 42, 400 | \$3, 261 | | |
| Flounders..... | 1, 840 | 92 | | | | | | |
| Shad..... | 15 | 2 | | | | | | |
| Striped bass..... | 720 | 77 | | | | | | |
| White perch..... | 6, 510 | 260 | | | | | | |
| Yellow perch..... | 1, 740 | 104 | | | | | | |
| Crabs, soft..... | | | 36, 000 | \$7, 200 | | | | |
| Lobsters..... | | | | | | | 12, 840 | \$2, 568 |
| Terrapin, diamond-back..... | 250 | 160 | | | | | | |
| Turtles, snapper..... | 8, 504 | 679 | | | | | | |
| Total..... | 57, 824 | 3, 055 | 36, 000 | 7, 200 | 42, 400 | 3, 261 | 12, 840 | 2, 568 |

| Species | Dredges | | | | | | Tongs | | By hand | |
|------------------------------|----------|-----------|----------|----------|----------|----------|---------|----------|---------|-------|
| | Clam | | Crab | | Oyster | | Pounds | Value | Pounds | Value |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Crabs: | | | | | | | | | | |
| Hard..... | | | 96, 600 | \$1, 205 | | | | | | |
| King..... | | | | | | | | | 60, 000 | \$66 |
| Clams: | | | | | | | | | | |
| Hard, public..... | 150, 500 | \$29, 895 | 40, 800 | 415 | | | | | | |
| Hard, private..... | 2, 640 | 538 | | | 7, 000 | \$1, 360 | | | | |
| Oysters: | | | | | | | | | | |
| Market, public, spring..... | | | | | | | 39, 600 | \$5, 140 | | |
| Market, private, spring..... | | | | | 20, 320 | 2, 460 | 4, 200 | 525 | | |
| Market, private, fall..... | | | | | 165, 240 | 20, 065 | | | | |
| Terrapin, diamond-back..... | | | | | | | | | 160 | 100 |
| Total..... | 153, 140 | 30, 433 | 137, 400 | 1, 710 | 192, 560 | 23, 885 | 43, 800 | 5, 665 | 60, 160 | 166 |

OPERATING UNITS: BY COUNTIES

| Item | Kent | New Castle | Sussex |
|-----------------------------|--------|------------|---------|
| | Number | Number | Number |
| Fishermen: | | | |
| On vessels..... | 58 | | 307 |
| On boats and shore: | | | |
| Regular..... | 2 | 3 | 29 |
| Casual..... | 179 | 48 | 251 |
| Total..... | 239 | 51 | 687 |
| Vessels: | | | |
| Steam: | | | |
| Net tonnage..... | | | 10 |
| Motor..... | | | 1, 237 |
| Net tonnage..... | 14 | | 3 |
| Sail..... | 195 | | 204 |
| Net tonnage..... | 1 | | |
| Net tonnage..... | 8 | | |
| Total vessels..... | 15 | | 13 |
| Total net tonnage..... | 203 | | 1, 441 |
| Boats: | | | |
| Motor..... | 31 | 10 | 36 |
| Other..... | 56 | 20 | 105 |
| Accessory boats..... | | | 30 |
| Apparatus: | | | |
| Purse seines, menhaden..... | | | 10 |
| Length, yards..... | | | 3, 140 |
| Haul seines..... | 18 | 10 | 33 |
| Length, yards..... | 6, 940 | 770 | 11, 280 |

Fisheries of Delaware, 1933—Continued

OPERATING UNITS: BY COUNTIES—Continued

| Item | Kent | New Castle | Sussex |
|-----------------------------|--------|------------|--------|
| | Number | Number | Number |
| Apparatus—Continued. | | | |
| Gill nets: | | | |
| Drift: | | | |
| Square yards | 7 | 18 | 38 |
| Runaround | 11,200 | 102,800 | 31,720 |
| Square yards | 3 | | 30 |
| Stake | 8,600 | | 32,850 |
| Square yards | 46 | | 53 |
| Lines: | 5,120 | | 2,650 |
| Hand | | | 12 |
| Hooks | | | 24 |
| Trawl | | | 2 |
| Hooks | | | 1,300 |
| Pound nets | 15 | | 17 |
| Stop nets | 2 | 5 | |
| Square yards | 800 | 1,020 | |
| Fyke nets | 80 | 122 | 247 |
| Dip nets | | | 51 |
| Pots: | | | |
| Eel | 30 | | 846 |
| Lobster | | | 165 |
| Dredges: | | | |
| Clam | 29 | | 4 |
| Yards at mouth | 41 | | 6 |
| Crab | 7 | | 4 |
| Yards at mouth | 13 | | 7 |
| Oyster | 14 | | 2 |
| Yards at mouth | 20 | | 3 |
| Tongs | 26 | | 6 |

CATCH: BY COUNTIES

| Species | Kent | | New Castle | | Sussex | |
|---------------------------------|------------------|---------------|---------------|--------------|-------------------|----------------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives | 1,580 | \$18 | 1,800 | \$23 | 1,145,600 | \$4,745 |
| Bluefish | | | | | 4,720 | 236 |
| Carp | 4,540 | 240 | 18,400 | 1,185 | 3,480 | 164 |
| Catfish and bullheads | 125 | 6 | 1,080 | 71 | 640 | 39 |
| Cod | | | | | 2,240 | 69 |
| Croaker | 23,500 | 307 | | | 43,630 | 761 |
| Drum, red or redfish | | | | | 620 | 16 |
| Eels, common | 2,160 | 86 | 7,640 | 382 | 43,340 | 3,424 |
| Flounders | | | | | 9,450 | 492 |
| Menhaden | | | | | 33,600,000 | 111,700 |
| Mullet | | | | | 233,000 | 2,390 |
| Shad | 265 | 30 | 7,260 | 718 | 14,770 | 1,239 |
| Spot | 9,080 | 265 | | | 22,840 | 638 |
| Squeteague or "sea trout", gray | 77,270 | 2,398 | | | 45,910 | 843 |
| Striped bass | 2,740 | 363 | 40 | 6 | 8,920 | 1,013 |
| Sturgeon | | | 1,200 | 300 | | |
| White perch | 6,060 | 295 | 600 | 44 | 23,990 | 1,151 |
| Whiting | | | 480 | 30 | | |
| Yellow perch | | | 340 | 20 | 6,140 | 378 |
| Crabs: | | | | | | |
| Hard | 44,600 | 545 | | | 52,000 | 750 |
| King | 566,600 | 595 | | | | |
| Soft | | | | | 36,000 | 7,200 |
| Lobsters | | | | | 12,840 | 2,568 |
| Clams: | | | | | | |
| Hard, public | 189,700 | 29,950 | | | 1,600 | 300 |
| Hard, private | 9,240 | 1,808 | | | 400 | 90 |
| Oysters: | | | | | | |
| Market, public, spring | 27,600 | 3,540 | | | 12,000 | 1,600 |
| Market, private, spring | 24,520 | 2,985 | | | | |
| Market, private, fall | 139,760 | 16,425 | | | 25,480 | 3,640 |
| Terrapin, diamond-back | 250 | 160 | 160 | 100 | | |
| Turtles, snapper | 4,504 | 439 | 4,000 | 240 | | |
| Total | 1,134,124 | 60,461 | 43,000 | 3,119 | 35,349,610 | 145,606 |

U. S. BUREAU OF FISHERIES

Fisheries of Delaware, 1933—Continued

SEED OYSTER FISHERY: BY GEAR

| Item | Dredges | | Tongs | | Total, exclusive of duplication | |
|-------------------------------------|--------------------------|-------------------------|-------------------------|-------------------------|---------------------------------|-------------------------|
| | Number | Number | Number | Number | Number | Number |
| OPERATING UNITS | | | | | | |
| Fishermen: | | | | | | |
| On vessels..... | 48 | | | | 48 | |
| On boats and shore, casual..... | 3 | | 40 | | 43 | |
| Total..... | 51 | | 40 | | 91 | |
| Vessels, sail..... | 7 | | | | 7 | |
| Net tonnage..... | 96 | | | | 96 | |
| Boats, other than motor..... | 1 | | 40 | | 41 | |
| Apparatus: | | | | | | |
| Number..... | 16 | | 40 | | | |
| Yards at mouth..... | 21 | | | | | |
| CATCH | | | | | | |
| Oysters, seed, public, spring..... | <i>Bushels</i> 25,340 | <i>Value</i> \$6,330 | <i>Bushels</i> 9,600 | <i>Value</i> \$2,400 | <i>Bushels</i> 34,940 | <i>Value</i> \$8,730 |

SEED OYSTER FISHERY: BY COUNTIES

| Item | Kent | | Sussex | |
|-------------------------------------|--------------------------|-------------------------|-------------------------|-------------------------|
| | Number | Number | Number | Number |
| OPERATING UNITS | | | | |
| Fishermen: | | | | |
| On vessels..... | 33 | | 15 | |
| On boats and shore, casual..... | 37 | | 6 | |
| Total..... | 70 | | 21 | |
| Vessels, sail..... | 5 | | 2 | |
| Net tonnage..... | 60 | | 36 | |
| Boats, other than motor..... | 35 | | 6 | |
| Apparatus: | | | | |
| Dredges..... | 12 | | 4 | |
| Yards at mouth..... | 16 | | 5 | |
| Tongs..... | 34 | | 6 | |
| CATCH | | | | |
| Oysters, seed, public, spring..... | <i>Bushels</i> 27,140 | <i>Value</i> \$6,780 | <i>Bushels</i> 7,800 | <i>Value</i> \$1,960 |

NOTE.—Of the total number of persons fishing for seed oysters, 56 are duplicated among those fishing for market oysters or other species. Similarly the following craft and gear are duplicated: 6 sail vessels, 31 small boats other than motor, 12 dredges, and 31 tongs.

VESSEL FISHERIES AT NEW YORK CITY⁷

During 1933, fishing vessels of 5 net tons capacity or greater landed 24,455,000 pounds of fishery products at New York City. The landings consisted of bluefish, 2,467,000 pounds; bonito, 9,000 pounds; butterfish, 45,000 pounds; cod, 2,548,000 pounds; croaker, 4,000 pounds; eels, 3,000 pounds; flounders, 6,712,000 pounds; haddock, 8,209,000 pounds; hake, 5,000 pounds; halibut, 9,000 pounds; sea herring, 3,000 pounds; king whiting, 6,000 pounds; mackerel, 1,288,000 pounds; pollock, 5,000 pounds; scup, 212,000 pounds; sea bass, 159,000 pounds; swordfish, 3,000 pounds; tilefish, 1,621,000 pounds; and sea scallops, 2,147,000 pounds. Since the landings at Groton, Conn., were inconsiderable they have not been included with those at New York City as has been the custom in the past several years. Data on the landings at New York City are included also in the catch by States.

⁷ Statistics on the landings at New York City are collected by J. H. Matthews, executive secretary, Middle Atlantic Fisheries Association.

SHAD FISHERY OF THE HUDSON RIVER

The shad fishery of the Hudson River in 1933 was prosecuted by 317 fishermen who used 11 motor boats, 145 other boats, 2 haul seines having a combined length of 280 yards, 119 drift gill nets having a total area of 406,871 square yards, and 27 stake gill nets having a total area of 28,760 square yards. The total commercial catch amounted to 154,437 shad having a weight of 518,680 pounds and a value to the fishermen of \$40,729. This is a decrease of 3 percent in the number of shad and 20 percent in their value as compared with 1932. The average price per pound received by the fishermen in 1933 was about 8 cents as compared with 10 cents in 1932.

Drift gill nets accounted for 61 percent of the weight of the shad taken while stake gill nets accounted for 39 percent. Small quantities, amounting to less than one-half of 1 percent, were taken by haul seines.

Statistics of the catch of shad in the Hudson River are included also in the catch data for New York and New Jersey which are published elsewhere in this report.

Shad fishery of the Hudson River, 1933

| Item | New York | | | New Jersey | | | Total | | |
|----------------------------|----------------|----------------|---------------|---------------|----------------|---------------|----------------|----------------|---------------|
| | Number | Pounds | Value | Number | Pounds | Value | Number | Pounds | Value |
| Fishermen: | | | | | | | | | |
| On boats and shore: | | | | | | | | | |
| Regular..... | 55 | | | 6 | | | 61 | | |
| Casual..... | 212 | | | 44 | | | 256 | | |
| Total..... | 267 | | | 50 | | | 317 | | |
| Boats: | | | | | | | | | |
| Motor..... | | | | 11 | | | 11 | | |
| Other..... | 133 | | | 12 | | | 145 | | |
| Apparatus: | | | | | | | | | |
| Haul seines..... | 2 | | | | | | 2 | | |
| Length, yards..... | 280 | | | | | | 280 | | |
| Gill nets: | | | | | | | | | |
| Drift..... | 118 | | | 1 | | | 119 | | |
| Square yards..... | 404,873 | | | 1,998 | | | 406,871 | | |
| Stake..... | 14 | | | 13 | | | 27 | | |
| Square yards..... | 13,370 | | | 15,390 | | | 28,760 | | |
| Shad caught: | | | | | | | | | |
| With haul seines..... | 576 | 1,670 | \$155 | | | | 576 | 1,670 | \$155 |
| With drift gill nets..... | 96,411 | 314,693 | 25,182 | 150 | 524 | \$40 | 96,561 | 315,217 | 25,222 |
| With stake gill nets..... | 8,300 | 31,293 | 2,319 | 49,000 | 170,500 | 12,533 | 57,300 | 201,793 | 15,352 |
| Total..... | 105,287 | 347,656 | 28,156 | 49,150 | 171,024 | 12,573 | 154,437 | 518,680 | 40,729 |

FISHERIES OF THE CHESAPEAKE BAY STATES

(Area XXIII) *

The yield of the commercial fisheries of the Chesapeake Bay States (Maryland and Virginia) during 1933 amounted to 272,380,052 pounds, valued at \$5,060,829 to the fishermen, representing a decrease

* This is the number given to this area by the North American Council on Fishery Investigations. It should be explained that there are included under this area, craft whose principal fishing ports are in the area but at times fish elsewhere. A notable example is the southern trawl fishery, which extends into Area XXIV. Data on the operating units and catch of the fisheries of the Chesapeake Bay States have been taken largely from statistics collected by the State fishery agencies of Maryland and Virginia. Supplementary surveys, compilations, and analyses have been made by agents of this Bureau in order that the figures may be presented in a manner comparable with those of other sections. It should be observed that the persons engaged, gear and craft employed, and catch of the seed oyster fishery are not included among the statistics of the fishery for market oysters and other species but are shown in separate tables in this section. For a clearer understanding of the statistics published in this section, the reader is referred to the section in the latter part of this document entitled "Statistical survey procedure."

of 24 percent in volume and 14 percent in value as compared with the catch in the previous year. These fisheries gave employment to 20,142 fishermen as compared with 20,946 in 1932.

There were 502 fishery wholesale and manufacturing establishments in the two States in 1933 as compared with 564 in 1931 when the most recent previous survey of such concerns was made. In 1933 these establishments employed 11,596 persons, paid \$2,366,762 in salaries and wages, and produced manufactured products (canned, cured, packaged, and byproducts), valued at \$7,245,169. In 1931 the wholesale and manufacturing firms employed 12,333 persons, paid \$2,802,420 in salaries and wages, and produced manufactured products, valued at \$7,905,626.

Fisheries of the Chesapeake Bay States, 1933

SUMMARY OF CATCH

| Product | Maryland | | Virginia | | Total | |
|---------------------|------------|-----------|-------------|-------------|-------------|-------------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Fish..... | 13,498,490 | \$370,360 | 176,209,213 | \$1,685,109 | 189,707,703 | \$2,055,469 |
| Shellfish, etc..... | 41,863,299 | 1,363,495 | 40,809,050 | 1,641,865 | 82,672,349 | 3,005,360 |
| Total..... | 55,361,789 | 1,733,855 | 217,018,263 | 3,326,974 | 272,380,052 | 5,060,829 |

OPERATING UNITS: BY STATES

| Item | Maryland | Virginia | Total |
|--------------------------------|---------------|---------------|---------------|
| Fishermen: | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| On vessels..... | 654 | 1,471 | 2,125 |
| On boats and shore: | | | |
| Regular..... | 5,569 | 7,073 | 12,632 |
| Casual..... | 2,314 | 3,071 | 5,385 |
| Total..... | 8,527 | 11,615 | 20,142 |
| Vessels: | | | |
| Steam..... | | 25 | 25 |
| Net tonnage..... | | 2,831 | 2,831 |
| Motor..... | | 105 | 105 |
| Net tonnage..... | | 1,761 | 1,761 |
| Sail..... | 154 | 3 | 157 |
| Net tonnage..... | 1,727 | 22 | 1,749 |
| Total vessels..... | 154 | 133 | 287 |
| Total net tonnage..... | 1,727 | 4,614 | 6,341 |
| Boats: | | | |
| Motor..... | 3,280 | 4,212 | 7,492 |
| Other..... | 2,316 | 3,621 | 5,937 |
| Accessory boats..... | | 102 | 102 |
| Apparatus: | | | |
| Furse seines: | | | |
| Menhaden..... | | 34 | 34 |
| Length, yards..... | | 10,160 | 10,160 |
| Haul seines..... | 333 | 191 | 524 |
| Length, yards..... | 36,849 | 52,063 | 88,912 |
| Gill nets: | | | |
| Anchor..... | 74 | | 74 |
| Square yards..... | 26,654 | | 26,654 |
| Drift..... | 296 | 410 | 706 |
| Square yards..... | 367,311 | 337,045 | 704,356 |
| Stake..... | 3,883 | 4,913 | 8,796 |
| Square yards..... | 208,963 | 286,576 | 565,538 |
| Lines: | | | |
| Hand..... | 12 | | 12 |
| Hooks..... | 24 | | 24 |
| Trot with baits or snoods..... | 1,547 | 1,075 | 2,620 |
| Baits or snoods..... | 938,350 | 499,610 | 1,437,966 |
| Pound nets..... | 690 | 1,880 | 2,572 |
| Stop nets..... | | 6 | 6 |
| Square yards..... | | 10,300 | 10,300 |

Fisheries of the Chesapeake Bay States, 1933—Continued

OPERATING UNITS: BY STATES—Continued

| Item | Maryland | Virginia | Total |
|---------------------|----------|----------|--------|
| | Number | Number | Number |
| Apparatus—Continued | | | |
| Fyke nets..... | 2,181 | 649 | 2,830 |
| Dip nets..... | 1,458 | 1,675 | 3,133 |
| Otter trawls..... | | 27 | 27 |
| Yards at mouth..... | | 803 | 803 |
| Pots, eel..... | 8,835 | 35 | 8,870 |
| Spears..... | 2 | | 2 |
| Scrapes..... | 642 | 41 | 683 |
| Yards at mouth..... | 642 | 41 | 683 |
| Dredges: | | | |
| Crab..... | | 130 | 130 |
| Yards at mouth..... | | 235 | 235 |
| Oyster..... | 418 | 298 | 716 |
| Yards at mouth..... | 454 | 339 | 793 |
| Scallop..... | | 2 | 2 |
| Yards at mouth..... | | 6 | 6 |
| Tongs..... | 4,753 | 3,748 | 8,501 |
| Rakes..... | 20 | 362 | 382 |
| Picks..... | | 540 | 540 |

CATCH: BY STATES

| Species | Maryland | | Virginia | | Total | |
|---------------------------------|-------------------|----------------|--------------------|------------------|--------------------|------------------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | |
| Alewives..... | 6,549,673 | \$43,164 | 19,177,448 | \$86,766 | 25,727,121 | \$129,930 |
| Black bass..... | 76,491 | 5,725 | | | 76,491 | 5,725 |
| Bluefish..... | 118,700 | 8,106 | 684,359 | 27,029 | 803,059 | 85,196 |
| Bonito..... | 9,100 | 194 | 10,092 | 455 | 19,192 | 649 |
| Butterfish..... | 589,263 | 12,709 | 2,285,353 | 55,203 | 2,874,616 | 67,912 |
| Cabio or crab eater..... | 1,100 | 44 | 18,942 | 791 | 19,942 | 835 |
| Carp..... | 169,833 | 11,406 | 428,464 | 16,877 | 598,297 | 28,283 |
| Catfish and bullheads..... | 263,088 | 10,261 | 718,408 | 22,351 | 981,476 | 32,612 |
| Cod..... | 100 | 2 | 6,866 | 137 | 6,966 | 139 |
| Crappie..... | 1,379 | 52 | | | 1,379 | 52 |
| Croaker..... | 1,806,886 | 24,349 | 14,235,182 | 186,720 | 16,042,048 | 211,069 |
| Dolphin..... | 500 | 75 | | | 500 | 75 |
| Drum: | | | | | | |
| Black..... | 42,000 | 350 | 80,480 | 1,594 | 122,480 | 1,944 |
| Red or redfish..... | 7,200 | 140 | 51,539 | 1,456 | 58,739 | 1,596 |
| Eels..... | 203,413 | 11,727 | 108,634 | 4,347 | 312,047 | 16,074 |
| Flounders..... | 69,564 | 2,194 | 1,083,413 | 52,031 | 1,152,977 | 55,225 |
| Gizzard shad..... | 27,504 | 584 | 116,820 | 2,399 | 144,324 | 2,983 |
| Haddock..... | | | 47 | 1 | 47 | 1 |
| Hake..... | | | 23,455 | 387 | 23,455 | 387 |
| Harvestfish..... | 19,535 | 831 | 160,050 | 3,585 | 179,585 | 4,416 |
| Hickory shad..... | 11,426 | 249 | 56,228 | 1,196 | 67,654 | 1,445 |
| King whiting or "kingfish"..... | 9,900 | 396 | 64,303 | 2,050 | 74,203 | 2,446 |
| Mackerel..... | 27,200 | 272 | 16,216 | 901 | 43,416 | 1,173 |
| Menhaden..... | | | 115,990,500 | 385,825 | 115,990,500 | 385,825 |
| Mullet..... | 20,100 | 1,206 | 44,541 | 1,648 | 64,641 | 2,854 |
| Pigfish..... | | | 60,765 | 520 | 60,765 | 520 |
| Pike or pickerel..... | 28,088 | 3,913 | 200 | 11 | 28,898 | 3,924 |
| Pollock..... | | | 25 | 1 | 26 | 1 |
| Pompano..... | | | 4,530 | 243 | 4,530 | 249 |
| Scup..... | 82,500 | 1,650 | 1,506,131 | 36,324 | 1,588,631 | 37,974 |
| Sea bass..... | 10,100 | 805 | 332,075 | 9,392 | 342,175 | 9,807 |
| Sea robin..... | 500 | 25 | | | 500 | 25 |
| Shad..... | 1,374,315 | 133,142 | 4,816,714 | 440,027 | 6,191,029 | 673,169 |
| Sharks..... | | | 10,225 | 159 | 10,225 | 159 |
| Silver perch..... | 1,062 | 22 | | | 1,062 | 22 |
| Spanish mackerel..... | 1,400 | 70 | 66,501 | 3,950 | 67,901 | 4,020 |
| Spot..... | 30,827 | 704 | 718,665 | 18,010 | 747,292 | 18,714 |
| Squeteague or "sea trout": | | | | | | |
| Gray..... | 1,163,654 | 32,043 | 12,310,138 | 230,981 | 13,463,792 | 263,024 |
| Spotted..... | 8,263 | 497 | 164,415 | 12,987 | 172,678 | 13,494 |
| Stripe bass..... | 313,795 | 37,739 | 518,900 | 63,474 | 832,695 | 101,218 |
| Sturgeon..... | | 115 | 8,141 | 1,598 | 8,226 | 1,711 |
| Suckers..... | 7,559 | 277 | | | 7,559 | 277 |
| Sunfish..... | 2,112 | 42 | | | 2,112 | 42 |
| Swellfish..... | | | 1,065 | 53 | 1,065 | 53 |
| Tautog..... | 5,000 | 150 | 1,048 | 21 | 6,048 | 171 |
| Thimble-eyed mackerel..... | | | 11,250 | 169 | 11,250 | 169 |
| White perch..... | 285,870 | 15,802 | 247,420 | 9,828 | 533,290 | 25,630 |
| Whiting..... | 500 | 25 | | | 500 | 25 |
| Yellow perch..... | 167,945 | 8,603 | 71,765 | 2,599 | 239,710 | 12,202 |
| Total..... | 13,498,490 | 370,360 | 176,209,213 | 1,685,109 | 189,707,703 | 2,055,469 |

Fisheries of the Chesapeake Bay States, 1933—Continued

CATCH: BY STATES—Continued

| Species | Maryland | | Virginia | | Total | |
|------------------------------|------------|-----------|-------------|-----------|-------------|-----------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| SHELLFISH, ETC. | | | | | | |
| Crabs: | | | | | | |
| Hard..... | 26,648,100 | \$331,747 | 23,911,375 | \$283,898 | 50,559,475 | \$615,645 |
| Soft..... | 3,449,029 | 238,401 | 2,067,957 | 143,351 | 5,516,986 | 381,752 |
| Lobsters..... | | | 131 | 2 | 131 | 2 |
| Shrimp..... | | | 371 | 20 | 371 | 20 |
| Clams, hard, public..... | 41,536 | 3,894 | 1,169,266 | 264,258 | 1,210,802 | 268,152 |
| Oysters: ¹ | | | | | | |
| Market, public, spring..... | 3,328,176 | 192,478 | 1,660,049 | 95,142 | 4,988,225 | 287,620 |
| Market, public, fall..... | 7,279,200 | 477,725 | 2,632,065 | 148,189 | 9,911,205 | 625,914 |
| Market, private, spring..... | 390,125 | 43,983 | 3,766,701 | 285,440 | 4,146,826 | 330,423 |
| Market, private, fall..... | 687,283 | 74,011 | 5,419,017 | 407,284 | 6,106,300 | 481,295 |
| Scallops, sea..... | | | 72,645 | 8,094 | 72,645 | 8,094 |
| Squid..... | 37,500 | 750 | 107,793 | 2,627 | 145,293 | 3,377 |
| Terrapin, diamond-back..... | 1,800 | 500 | 10,240 | 2,500 | 12,040 | 3,000 |
| Turtles, snapper..... | 550 | 6 | 1,500 | 60 | 2,050 | 66 |
| Total..... | 41,863,299 | 1,363,495 | 40,809,050 | 1,641,865 | 82,672,349 | 3,005,360 |
| Grand total..... | 55,361,789 | 1,733,855 | 217,018,263 | 3,326,974 | 272,380,052 | 5,060,829 |

¹ Statistics on oysters used in this table are based on yields of 6.57 pounds of meats to the bushel for market oysters in Maryland and 6.6 pounds in Virginia.

PRODUCTION OF CERTAIN SHELLFISH IN NUMBER AND BUSHELS

| Product | Maryland | | Virginia | | Total | |
|-------------------------------------|------------|-----------|------------|-----------|-------------|-----------|
| | Quantity | Value | Quantity | Value | Quantity | Value |
| Crabs: | | | | | | |
| Hard.....number..... | 79,944,300 | \$331,747 | 71,734,125 | \$283,898 | 151,678,425 | \$615,645 |
| Soft.....do..... | 13,796,116 | 238,401 | 8,271,828 | 143,351 | 22,067,944 | 381,752 |
| Clams, hard.....bushels..... | 5,192 | 3,894 | 146,158 | 264,258 | 151,350 | 268,152 |
| Oysters: | | | | | | |
| Market, public, spring.....do..... | 506,572 | 192,478 | 251,523 | 95,142 | 758,095 | 287,620 |
| Market, public, fall.....do..... | 1,107,945 | 477,725 | 398,789 | 148,189 | 1,506,734 | 625,914 |
| Market, private, spring.....do..... | 59,380 | 43,983 | 569,197 | 285,440 | 628,577 | 330,423 |
| Market, private, fall.....do..... | 104,609 | 74,011 | 821,063 | 407,284 | 925,672 | 481,295 |
| Scallops, sea.....do..... | | | 12,107 | 8,094 | 12,107 | 8,094 |

SEED OYSTER FISHERY

| Item | Virginia | Item | Virginia | | |
|------------------------|----------|----------------------------|-----------|----------|--|
| | Number | | Bushels | Value | |
| OPERATING UNITS | | CATCH | | | |
| Fishermen: | | Oysters: | | | |
| On boats and shore: | | Seed, public, spring..... | 574,620 | \$80,880 | |
| Regular..... | 1,650 | Seed, public, fall..... | 816,240 | 85,934 | |
| Casual..... | 67 | Seed, private, spring..... | 35,600 | 7,720 | |
| Total..... | 1,717 | Seed, private, fall..... | 56,800 | 11,360 | |
| | | Total..... | 1,482,260 | 165,894 | |
| Boats: | | | | | |
| Motor..... | 804 | | | | |
| Other..... | 286 | | | | |
| Apparatus: | | | | | |
| Tongs..... | 1,474 | | | | |
| Rakes..... | 130 | | | | |

NOTE.—Of the number of persons fishing for seed oysters, 1,636 are duplicated among those fishing for market oysters or other species. Similarly the following craft and gear are duplicated: 744 motor boats, 267 other boats, and 1,161 tongs.

Industries related to the fisheries of the Chesapeake Bay States, 1933

OPERATING UNITS, SALARIES, AND WAGES

| Item | Maryland | Virginia | Total |
|---------------------------------------|---------------|---------------|---------------|
| Transporting: | | | |
| Persons engaged: | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| On vessels..... | 291 | 578 | 869 |
| On boats..... | | 25 | 25 |
| Total..... | 291 | 603 | 894 |
| Vessels: | | | |
| Motor..... | 152 | 269 | 421 |
| Net tonnage..... | 2,177 | 2,752 | 4,929 |
| Sail..... | 11 | | 11 |
| Net tonnage..... | 307 | | 307 |
| Total vessels..... | 163 | 269 | 432 |
| Total net tonnage..... | 2,484 | 2,752 | 5,236 |
| Boats..... | | 102 | 102 |
| Wholesale and manufacturing: | | | |
| Establishments..... | 306 | 194 | 502 |
| Persons engaged: | | | |
| Proprietors..... | 438 | 226 | 664 |
| Salaried employees..... | 175 | 140 | 315 |
| Wage earners: | | | |
| Average for season..... | 5,822 | 4,795 | 10,617 |
| Average for year..... | 2,707 | 1,997 | 4,704 |
| Paid to salaried employees..... | \$322,981 | \$157,830 | \$480,811 |
| Paid to wage earners..... | \$1,034,205 | \$951,746 | \$1,985,951 |
| Total salaries and wages..... | \$1,357,186 | \$1,009,576 | \$2,366,762 |
| Fishermen's manufacturing..... | 14 | | 14 |

PRODUCTS MANUFACTURED

| Item | Maryland | | Virginia | |
|---|-----------|-----------|-----------|-----------|
| | Quantity | Value | Quantity | Value |
| By manufacturing establishments: | | | | |
| Alewives: | | | | |
| Salted: | | | | |
| Corned..... pounds..... | 108,500 | \$2,450 | 1,070,000 | \$17,297 |
| Pickled..... do..... | 908,200 | 30,429 | (?) | (?) |
| Tight-pack cut..... do..... | 474,000 | 10,700 | 3,583,020 | 91,036 |
| Tight-pack roe..... do..... | (?) | (?) | 171,780 | 4,422 |
| Canned..... standard cases..... | 16,575 | 36,500 | 3,705 | 9,863 |
| Roe, canned..... do..... | 7,336 | 29,390 | 20,375 | 78,002 |
| Dry scrap..... tons..... | (?) | (?) | 524 | 13,137 |
| Oil..... gallons..... | | | 29,900 | 3,368 |
| Butterfish, smoked..... pounds..... | 92,500 | 23,400 | | |
| Chubs, cisco, and tullibee, smoked..... do..... | 148,000 | 43,100 | | |
| Croaker, fresh fillets..... do..... | | | 30,500 | 3,720 |
| Flounders, fresh fillets..... do..... | | | 30,000 | 5,040 |
| Haddock, fresh fillets..... do..... | | | 9,500 | 1,470 |
| Menhaden products: | | | | |
| Dry scrap and meal..... tons..... | | | 13,846 | 434,647 |
| Oil..... gallons..... | | | 1,555,995 | 231,117 |
| Salmon, smoked..... pounds..... | 158,000 | 55,800 | | |
| Sea bass, fresh fillets..... do..... | | | 62,000 | 7,900 |
| Squeteagues, fresh fillets..... do..... | | | 77,200 | 10,038 |
| Sturgeon, smoked..... do..... | 10,250 | 9,138 | | |
| Whitefish, smoked..... do..... | 73,000 | 24,050 | | |
| Crabs, blue: | | | | |
| Meat, packaged, fresh-cooked..... do..... | 3,662,539 | 785,312 | 2,086,900 | 534,714 |
| Dry scrap and meal..... tons..... | (?) | (?) | 850 | 15,412 |
| Oysters, fresh-shucked..... gallons..... | 1,907,211 | 1,985,599 | 1,676,581 | 1,803,878 |
| Oyster-shell products: | | | | |
| Poultry feed..... tons..... | 34,298 | 175,969 | 15,537 | 91,940 |
| Lime..... do..... | 20,998 | 30,017 | 11,578 | 37,595 |
| Lime, "burned"..... do..... | | | 3,314 | 22,948 |

¹ This item is usually an intermediate product and, although included in the total, may be shown in its final stage of processing in this or another State.

² The production of this item is included under "Unclassified products".

Industries related to the fisheries of the Chesapeake Bay States, 1933—Continued

PRODUCTS MANUFACTURED—Continued

| Item | Maryland | | Virginia | |
|---|-------------------------|----------------------|-------------------------|----------------------|
| | Quantity ⁽¹⁾ | Value ⁽²⁾ | Quantity ⁽¹⁾ | Value ⁽²⁾ |
| By manufacturing establishments—Continued. | | | | |
| Unclassified products: | | | 64,438 | \$10,315 |
| Fresh-packaged.....pounds.. | | | | |
| Smoked.....do..... | 132,300 | \$16,565 | | |
| Scrap and meal.....tons.. | 588 | 7,640 | 792 | 7,635 |
| Miscellaneous..... | | 501,368 | | 33,248 |
| Total..... | | 3,767,427 | | 3,477,742 |
| By fishermen: | | | | |
| Alewives, salted, tight-pack cut.....pounds.. | 175,000 | 3,500 | | |
| Sturgeon roe, salted.....do..... | 80 | 48 | | |
| Total..... | 175,080 | 3,548 | | |
| Grand total..... | | 3,770,975 | | 3,477,742 |

¹ This has been included under "Miscellaneous".

² Includes fresh fillets of cod, red drum, hake and Spanish mackerel, and fresh-shucked hard clams.

³ Includes smoked alewives, carp, eels, shad, lake trout, kippered salmon, and sea herring bloomers.

⁴ Includes alewife dry scrap and blue crab meal.

⁵ Includes miscellaneous dry and acidulated scrap.

⁶ Includes canned hard clam chowder and terrapin products; fresh-packaged alewife roe; salted tight-pack alewife roe; alewife oil; pearl essence; and marine shell products.

⁷ Includes canned crab meat, pickled alewives, and miscellaneous oil.

NOTE.—The total value of manufactured products in the Chesapeake Bay States was as follows: By manufacturing establishments, \$7,245,169; and by fishermen, \$3,548. Some of the above products may have been imported from another State or a foreign country; therefore, they cannot be correlated directly with the catch within the State. Of the total number of persons engaged in transporting vessels and boats 295 have been included as fishermen and among the total number of persons engaged in the preparation of fishermen's prepared products, all have been included as fishermen.

MARYLAND

Fisheries of Maryland, 1933

OPERATING UNITS: BY GEAR

| Item | Haul seines | Gill nets | | | Lines | | Pound nets | Fyke nets |
|------------------------------|---------------|---------------|---------------|---------------|---------------|---------------------------|---------------|---------------|
| | | Anchor | Drift | Stake | Hand | Trot with baits or snoods | | |
| Fishermen: | | | | | | | | |
| On boats and shore: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| Regular..... | 426 | 6 | 38 | 76 | 6 | 1,180 | 453 | 63 |
| Casual..... | 360 | 10 | 282 | 148 | | 127 | 119 | 71 |
| Total..... | 816 | 16 | 298 | 223 | 6 | 1,307 | 572 | 134 |
| Boats: | | | | | | | | |
| Motor..... | 139 | 7 | 68 | 77 | 3 | 1,257 | 233 | 53 |
| Other..... | 300 | 4 | 117 | 116 | | 44 | 168 | 83 |
| Apparatus: | | | | | | | | |
| Number..... | 333 | 74 | 206 | 3,683 | 12 | 1,547 | 690 | 2,181 |
| Length, yards..... | 36,849 | | | | | | | |
| Square yards..... | | 26,684 | 367,311 | 298,963 | | | | |
| Hooks, baits, or snoods..... | | | | | 24 | 938,350 | | |

| Item | Dip nets | Pots, eel | Spears | Scrapes | Dredges, oyster | Tongs | Rakes | By hand | Total, exclusive of duplication |
|----------------------------|--------------|------------|----------|------------|-----------------|--------------|-----------|-----------|---------------------------------|
| | | | | | | | | | Number |
| Fishermen: | | | | | | | | | |
| On vessels..... | | | | | 654 | | | | 654 |
| On boats and shore: | | | | | | | | | |
| Regular..... | 985 | 80 | 2 | 397 | 83 | 3,821 | 20 | | 5,559 |
| Casual..... | 473 | 61 | | | | 932 | | 15 | 2,314 |
| Total..... | 1,458 | 141 | 2 | 397 | 737 | 4,753 | 20 | 15 | 8,627 |
| Vessels, sail..... | | | | | 164 | | | | 164 |
| Net tonnage..... | | | | | 1,727 | | | | 1,727 |
| Boats: | | | | | | | | | |
| Motor..... | 100 | 79 | 2 | | 23 | 2,301 | 2 | | 3,280 |
| Other..... | 1,343 | 86 | | 321 | 32 | 168 | 20 | 15 | 2,316 |
| Apparatus: | | | | | | | | | |
| Number..... | 1,458 | 8,835 | 2 | 642 | 418 | 4,753 | 20 | | |
| Yards at mouth..... | | | | 642 | 454 | | | | |

Fisheries of Maryland, 1933—Continued

CATCH: BY GEAR

| Species | Haul seines | | Gill nets | | | | | | | |
|-----------------------------|------------------|---------------|---------------|--------------|----------------|---------------|----------------|---------------|--------|-------|
| | | | Anchor | | Drift | | Stake | | | |
| | | | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives | 65,550 | \$834 | | | 20,950 | \$368 | 19,700 | \$496 | | |
| Black bass | 47,967 | 3,667 | | | | | 1,345 | 81 | | |
| Bluefish | 17,130 | 1,086 | | | 49,000 | 4,200 | 4,080 | 243 | | |
| Butterfish | 8,000 | 520 | | | | | | | | |
| Carp | 143,618 | 9,988 | | | 510 | 17 | 1,100 | 47 | | |
| Catfish and bullheads | 60,020 | 2,134 | | | 19 | 1 | 1,295 | 57 | | |
| Crapple | | 14 | | | | | | | | |
| Croaker | 317,373 | 5,431 | | | 8,200 | 250 | 12,708 | 390 | | |
| Drum, red or redfish | 2,700 | 40 | | | | | | | | |
| Eels, common | 1,314 | 68 | | | | | 40 | 2 | | |
| Flounders | 3,030 | 142 | | | | | 200 | 10 | | |
| Gizzard shad | 3,283 | 53 | | | | | 264 | 4 | | |
| Hickory shad | | | | | 160 | 3 | 108 | 11 | | |
| Mullet | 100 | 6 | | | 20,000 | 1,200 | | | | |
| Pike or pickerel | 15,241 | 1,994 | | | | | 1,031 | 154 | | |
| Shad | 5,918 | 616 | 19,600 | \$2,223 | 275,280 | 26,670 | 148,688 | 15,460 | | |
| Silver perch | | 17 | | | | | | | | |
| Spot | 5,710 | 162 | | | | | | | | |
| Squeteagues or "sea trout": | | | | | | | | | | |
| Gray | 148,603 | 7,653 | | | 3,500 | 158 | 2,240 | 108 | | |
| Spotted | 7,200 | 410 | | | | | | | | |
| Striped bass | 84,253 | 11,529 | 4,500 | 570 | 32,650 | 3,766 | 46,012 | 5,298 | | |
| Sturgeon | | | | | 285 | 35 | | | | |
| Suckers | 3,698 | 117 | | | 28 | 1 | | | | |
| Sunfish | 712 | 14 | | | | | | | | |
| White perch | 61,529 | 3,593 | | | 3,550 | 228 | 12,120 | 754 | | |
| Whiting | 500 | 25 | | | | | | | | |
| Yellow perch | 38,430 | 1,836 | | | 3,000 | 360 | 2,982 | 192 | | |
| Crabs, soft | 266,884 | 27,013 | | | | | | | | |
| Turtles, snapper | 50 | 3 | | | | | | | | |
| Total | 1,309,839 | 78,965 | 24,100 | 2,793 | 417,332 | 37,257 | 253,913 | 23,307 | | |

| Species | Lines | | | | Pound nets | | Fyke nets | |
|-----------------------------|--------|-------|---------------------------|-------|------------|----------|-----------|-------|
| | Hand | | Trot with baits or snoods | | Pounds | Value | Pounds | Value |
| | Pounds | Value | Pounds | Value | | | | |
| Alewives | | | | | 6,438,813 | \$41,357 | 4,660 | \$109 |
| Black bass | | | | | 325 | 26 | 26,854 | 1,951 |
| Bluefish | 10,000 | \$600 | | | 38,490 | 1,977 | | |
| Bonito | 400 | 20 | | | 8,700 | 174 | | |
| Butterfish | | | | | 581,263 | 12,189 | | |
| Cabio or crab eater | | | | | 1,100 | 44 | | |
| Carp | | | | | 12,532 | 615 | 12,073 | 739 |
| Catfish and bullheads | | | | | 94,997 | 3,592 | 106,737 | 4,477 |
| Cod | | | | | 100 | 2 | | |
| Croaker | | | | | | | 1,204 | 38 |
| Crapple | | | | | 1,464,512 | 18,193 | 4,073 | 85 |
| Dolphin | 500 | 75 | | | | | | |
| Drum: | | | | | | | | |
| Black | | | | | 42,000 | 350 | | |
| Red or redfish | | | | | 4,500 | 100 | | |
| Eels: | | | | | | | | |
| Common | | | | | 24,712 | 1,603 | 6,848 | 418 |
| Other | 200 | 2 | | | 50 | 1 | | |
| Flounders | 5,000 | 250 | | | 60,534 | 2,748 | 800 | 44 |
| Gizzard shad | | | | | 23,397 | 498 | 560 | 29 |
| Harvestfish | | | | | 19,535 | 831 | | |
| Hickory shad | | | | | 11,158 | 235 | | |
| King whiting or "kingfish" | | | | | 9,900 | 396 | | |
| Mackerel | | | | | 27,200 | 272 | | |
| Pike or pickerel | | | | | 1,134 | 171 | 11,282 | 1,594 |
| Scup | 2,500 | 50 | | | 80,000 | 1,600 | | |
| Sea bass | 10,000 | 500 | | | 100 | 5 | | |
| Sea robin | | | | | 500 | 25 | | |
| Shad | | | | | 924,829 | 88,173 | | |
| Silver perch | | | | | 111 | 3 | 100 | 2 |
| Spanish mackerel | | | | | 1,400 | 70 | | |
| Spot | | | | | 24,917 | 542 | | |
| Squeteagues or "sea trout": | | | | | | | | |
| Gray | | | | | 996,111 | 23,930 | 3,200 | 194 |
| Spotted | | | | | 1,063 | 87 | | |
| Striped bass | | | | | 141,708 | 16,011 | 4,471 | 565 |
| Sturgeon | | | | | 400 | 80 | | |

U. S. BUREAU OF FISHERIES

Fisheries of Maryland, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Lines | | | | Pound nets | | Fyke nets | |
|-----------------------|---------------|----------------|---------------------------|----------------|-------------------|----------------|----------------|---------------|
| | Hand | | Trot with baits or snoods | | Pounds | Value \$ | Pounds | Value |
| | Pounds | Value | Pounds | Value | | | | |
| Suckers..... | | | | | 360 | | 3,473 | \$150 |
| Sunfish..... | | | | | | | 1,400 | 28 |
| Tautog..... | | | | | 5,000 | 150 | | |
| White perch..... | | | | | 123,190 | 6,587 | 85,481 | 4,660 |
| Yellow perch..... | | | | | 11,259 | 686 | 112,274 | 5,529 |
| Crabs, hard..... | | | 25,543,900 | \$317,945 | | | | |
| Squid..... | | | | | 37,500 | 750 | | |
| Turtles, snapper..... | | | | | | | 500 | 3 |
| Total..... | 28,600 | \$1,497 | 25,543,900 | 317,945 | 11,213,401 | 224,062 | 385,690 | 20,610 |

| Species | Dip nets | | Pots, eel | | Spears | | Scrapes | |
|-------------------|------------------|----------------|----------------|--------------|--------------|------------|------------------|---------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Eels, common..... | | | 164,549 | \$9,338 | 6,000 | \$300 | | |
| Crabs: | | | | | | | | |
| Hard..... | 88,300 | \$1,104 | | | | | 1,015,900 | \$12,698 |
| Soft..... | 2,441,379 | 168,937 | | | | | 740,766 | 44,451 |
| Total..... | 2,529,679 | 168,041 | 164,549 | 9,338 | 6,000 | 300 | 1,756,666 | 57,149 |

| Species | Dredges, oyster | | Tongs | | Rakes | | By hand | |
|------------------------------|------------------|----------------|------------------|----------------|--------------|------------|--------------|--------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Clams, hard, public..... | | | 24,640 | \$2,310 | 9,856 | \$924 | 7,040 | \$660 |
| Oysters: | | | | | | | | |
| Market, public, spring..... | 627,196 | \$37,563 | 2,700,980 | 154,915 | | | | |
| Market, public, fall..... | 1,104,010 | 76,510 | 6,175,190 | 401,215 | | | | |
| Market, private, spring..... | 76,600 | 9,655 | 313,525 | 34,328 | | | | |
| Market, private, fall..... | 95,384 | 11,655 | 591,899 | 62,356 | | | | |
| Terrapin, diamond-back..... | | | | | | | 1,800 | 500 |
| Total..... | 1,903,190 | 135,383 | 9,806,234 | 655,124 | 9,856 | 924 | 8,840 | 1,160 |

OPERATING UNITS: BY COUNTIES

| Item | Anne Arundel | Baltimore | Calvert | Caroline | Cecil | Charles | Dorchester | Harford |
|--------------------------------|--------------|------------|------------|-----------|------------|------------|--------------|-----------|
| | Number | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | | |
| On vessels..... | | 46 | 12 | | | | 170 | |
| On boats and shore: | | | | | | | | |
| Regular..... | 534 | 20 | 212 | 2 | 55 | 112 | 773 | 13 |
| Casual..... | 366 | 60 | 94 | 46 | 88 | 244 | 198 | 55 |
| Total..... | 900 | 126 | 318 | 48 | 143 | 356 | 1,141 | 68 |
| Vessels, sail..... | | 8 | 3 | | | | 40 | |
| Net tonnage..... | | 196 | 83 | | | | 356 | |
| Boats: | | | | | | | | |
| Motor..... | 338 | 30 | 132 | 2 | 52 | 154 | 514 | 25 |
| Other..... | 200 | 42 | 139 | 33 | 51 | 90 | 219 | 29 |
| Apparatus: | | | | | | | | |
| Haul seines..... | 46 | 13 | 6 | 5 | 27 | 25 | 3 | 10 |
| Length, yards..... | 2,425 | 2,577 | 1,960 | 1,535 | 5,695 | 4,620 | 1,050 | 2,310 |
| Gill nets: | | | | | | | | |
| Anchor..... | | | | | 74 | | | |
| Square yards..... | | | | | 26,654 | | | |
| Drift..... | 5 | | | 37 | 21 | 40 | 15 | 20 |
| Square yards..... | 400 | | | 1,090 | 45,912 | 125,744 | 18,733 | 43,159 |
| Stake..... | 15 | 81 | | 1 | 113 | 1,202 | 261 | 310 |
| Square yards..... | 1,390 | 1,097 | | 24 | 7,527 | 94,428 | 6,411 | 40,200 |
| Lines: | | | | | | | | |
| Trot with baits or snoods..... | 77 | 41 | 43 | | | 19 | 548 | |
| Baits or snoods..... | 44,200 | 12,300 | 19,500 | | | 19,000 | 237,200 | |
| Pound nets..... | 35 | 10 | 19 | | 41 | 40 | 170 | 6 |
| Fyke nets..... | 5 | 248 | | 14 | 1,155 | 29 | | 119 |
| Dip nets..... | 150 | 10 | 126 | | | 27 | 168 | |
| Pots, eel..... | 305 | 381 | | 25 | 590 | 100 | 2,230 | 161 |
| Dredges, oyster..... | | 16 | 6 | | | | 88 | |
| Yards at mouth..... | | 24 | 9 | | | | 93 | |
| Tongs..... | 745 | | 226 | | | 218 | 609 | |

Fisheries of Maryland, 1933—Continued

OPERATING UNITS: BY COUNTIES—Continued

| Item | Kent | Prince Georges | Queen Annes | St. Marys | Somer-set | Talbot | Wiccom- ioo | Worces- ter |
|---------------------------|---------|----------------|-------------|-----------|-----------|---------|----------------|----------------|
| | Number | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | | |
| On vessels | | | | | 364 | 62 | | |
| On boats and shore: | | | | | | | | |
| Regular | 544 | 11 | 693 | 436 | 1,028 | 634 | 337 | 157 |
| Casual | 234 | 24 | 53 | 829 | 140 | 231 | 118 | 34 |
| Total | 778 | 35 | 746 | 765 | 1,530 | 927 | 455 | 191 |
| Vessels, sail | | | | | 87 | 16 | | |
| Net tonnage | | | | | 965 | 127 | | |
| Boats: | | | | | | | | |
| Motor | 297 | 10 | 340 | 325 | 395 | 424 | 176 | 66 |
| Other | 175 | 16 | 117 | 235 | 598 | 211 | 43 | 118 |
| Apparatus: | | | | | | | | |
| Haul seines | 105 | 9 | 75 | 4 | 3 | 1 | 1 | |
| Length, yards | 6,722 | 1,360 | 3,390 | 1,240 | 1,075 | 400 | 500 | |
| Gill nets: | | | | | | | | |
| Drift | 87 | 8 | | 1 | 13 | 3 | 43 | 3 |
| Square yards | 69,600 | 12,385 | | 4,000 | 3,460 | 4,998 | 36,900 | 330 |
| Stake | 934 | | 9 | | 189 | 24 | 544 | |
| Square yards | 112,000 | | 1,060 | | 3,148 | 4,890 | 26,768 | |
| Lines: | | | | | | | | |
| Hand | | | | | | | | 12 |
| Hooks | | | | | | | | 24 |
| Trot with baits or snoods | 189 | | 98 | 130 | 63 | | 7 | 56 |
| Baits or snoods | 109,800 | | 72,250 | 117,000 | 50,400 | 206,300 | 5,600 | 44,800 |
| Pound nets | 28 | 6 | 13 | 75 | 50 | 122 | 23 | 45 |
| Fyke nets | 420 | 39 | 38 | | 23 | 45 | 42 | 4 |
| Dip nets | 100 | | 103 | 198 | 418 | 165 | | 3 |
| Pots, eel | 513 | 7 | 470 | 18 | | 3,895 | | 150 |
| Spears | | | | | | | | 2 |
| Scrapes | | | | | 642 | | | |
| Yards at month | | | | | 642 | | | |
| Dredges, oyster | | | | | 230 | 32 | | 46 |
| Yards at month | | | | | 252 | 43 | | 33 |
| Tongs | 501 | | 627 | 607 | 310 | 437 | 355 | 118 |
| Rakes | | | | | | | | 20 |

CATCH: BY COUNTIES

| Species | Anne Arundel | | Baltimore | | Calvert | | Caroline | |
|----------------------------------|------------------|----------------|----------------|---------------|------------------|---------------|---------------|--------------|
| | Pounds | Value \$ | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives | 858,609 | \$6,541 | 114,180 | \$735 | 193,150 | \$1,010 | 8,005 | \$116 |
| Black bass | | | | | | | 200 | 20 |
| Bluefish | 2,000 | 141 | 6,800 | 500 | 1,575 | 143 | | |
| Butterfish | 7,500 | 225 | | | | | 3,000 | 240 |
| Carp | 2,140 | 107 | 15,964 | 1,063 | 660 | 18 | 1,325 | 79 |
| Catfish and bullheads | 2,645 | 136 | 21,013 | 620 | 5,185 | 240 | 10,806 | 381 |
| Croaker | 192,411 | 2,282 | 1,073 | 35 | 102,000 | 1,253 | 3,100 | 124 |
| Eels, common | 16,813 | 761 | 10,300 | 557 | 260 | 15 | 1,153 | 19 |
| Flounders | 1,981 | 111 | 500 | 20 | 575 | 38 | | |
| Gizzard shad | 5,322 | 62 | 5,160 | 87 | 954 | 19 | 4,500 | 180 |
| Harvestfish | 35 | 1 | | | | | | |
| Hickory shad | 1,632 | 34 | 200 | 4 | 200 | 4 | | |
| Pike or pickerel | | | 375 | 73 | 207 | 31 | 173 | 31 |
| Shad | 90,532 | 8,301 | 2,270 | 228 | 68,082 | 7,234 | 11,384 | 1,365 |
| Spot | 8,367 | 201 | | | | | 225 | 9 |
| Squeteagues or "sea trout", gray | 308,216 | 11,648 | 3,000 | 180 | 5,000 | 220 | 4,000 | 320 |
| Striped bass | 4,666 | 480 | 8,188 | 1,081 | 7,281 | 1,063 | 9,850 | 1,442 |
| Suckers | | | | | 133 | 5 | 548 | 41 |
| White perch | 10,089 | 672 | 28,186 | 1,371 | 3,875 | 264 | 8,914 | 464 |
| Yellow perch | 375 | 25 | 12,954 | 702 | 740 | 52 | 670 | 44 |
| Crabs: | | | | | | | | |
| Hard | 721,800 | 10,105 | 190,000 | 2,850 | 712,100 | 10,669 | | |
| Soft | 102,049 | 9,721 | 2,700 | 450 | 195,187 | 12,365 | | |
| Oysters: | | | | | | | | |
| Market, public, spring | 470,470 | 26,884 | 57,750 | 2,475 | 125,513 | 11,972 | | |
| Market, public, fall | 717,507 | 46,220 | 26,250 | 1,500 | 301,119 | 29,387 | | |
| Market, private, spring | | | | | 70,000 | 6,900 | | |
| Market, private, fall | | | | | 87,500 | 8,750 | | |
| Turtles, snapper | 500 | 3 | | | | | | |
| Total | 3,525,659 | 124,561 | 506,863 | 14,531 | 1,881,406 | 91,684 | 67,852 | 4,875 |

Fisheries of Maryland, 1933—Continued

CATCH: BY COUNTIES—Continued

| Species | Cecil | | Charles | | Dorchester | | Harford | |
|-----------------------------|-----------|---------|---------|--------|------------|---------|---------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alwives..... | 757,600 | \$4,719 | 148,000 | \$821 | 462,212 | \$4,493 | 144,500 | \$1,325 |
| Black bass..... | 48,356 | 3,170 | 9,012 | 881 | | | 3,658 | 279 |
| Bluefish..... | | | 400 | 24 | 28,552 | 2,000 | | |
| Butterfish..... | | | | | 200 | 20 | | |
| Carp..... | 64,910 | 4,305 | 29,799 | 2,484 | 3,702 | 152 | 17,044 | 1,242 |
| Catfish and bullheads..... | 79,444 | 3,093 | 23,968 | 852 | 6,700 | 298 | 13,597 | 522 |
| Crappie..... | 1,000 | 24 | | | | | | |
| Croaker..... | | | 5,000 | 100 | 268,125 | 3,893 | | |
| Eels, common..... | 16,590 | 895 | 7,176 | 407 | 46,775 | 2,792 | 5,719 | 152 |
| Flounders..... | | | 87 | 5 | 12,969 | 506 | | |
| Gizzard shad..... | | | 3,968 | 61 | | | | |
| Hickory shad..... | 1,660 | 33 | 108 | 11 | 800 | 12 | 200 | 5 |
| Pike or pickerel..... | 12,965 | 1,631 | 814 | 125 | 50 | 8 | 5,685 | 866 |
| Shad..... | 116,778 | 11,789 | 124,962 | 10,101 | 103,182 | 10,882 | 56,810 | 5,864 |
| Silver perch..... | | | 562 | 12 | | | | |
| Spot..... | | | 60 | 3 | 1,400 | 118 | | |
| Squeteagues or "sea trout": | | | | | | | | |
| Gray..... | 68 | 8 | 3,200 | 214 | 29,781 | 1,442 | | |
| Spotted..... | | | | | 1,000 | 55 | | |
| Striped bass..... | 36,139 | 4,284 | 39,909 | 4,727 | 29,356 | 3,394 | 16,408 | 2,100 |
| Sturgeon..... | | | 125 | 20 | | | | |
| Suckers..... | 2,153 | 49 | 40 | 1 | | | 1,570 | 46 |
| Sunfish..... | 1,400 | 28 | | | | | 712 | 14 |
| White perch..... | 60,532 | 3,246 | 20,169 | 1,118 | 23,777 | 1,224 | 11,353 | 471 |
| Yellow perch..... | 89,600 | 3,888 | 3,864 | 256 | 20 | 1 | 10,024 | 614 |
| Crabs: | | | | | | | | |
| Hard..... | | | 158,700 | 1,905 | 8,700,500 | 104,405 | | |
| Soft..... | | | 2,775 | 239 | 545,251 | 32,732 | | |
| Oysters: | | | | | | | | |
| Market, public, spring..... | | | 65,044 | 5,575 | 398,639 | 24,221 | | |
| Market, public, fall..... | | | 72,289 | 5,164 | 824,513 | 50,533 | | |
| Turtles, snapper..... | 50 | 3 | | | | | | |
| Total..... | 1,288,245 | 41,165 | 720,029 | 35,088 | 11,487,504 | 243,181 | 287,278 | 13,500 |

| Species | Kent | | Prince Georges | | Queen Annes | | St. Marys | |
|------------------------------|-----------|---------|----------------|-------|-------------|---------|-----------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alwives..... | 187,249 | \$1,420 | 995 | \$31 | 21,195 | \$133 | 618,572 | \$3,117 |
| Black bass..... | 22 | 1 | 15,220 | 1,372 | | | | |
| Bluefish..... | 4,013 | 185 | | | | | 4,340 | 270 |
| Butterfish..... | 100 | 10 | | | | | 2,900 | 152 |
| Carp..... | 9,282 | 390 | 17,243 | 1,119 | 1,919 | 139 | 2,076 | 88 |
| Catfish and bullheads..... | 50,760 | 1,845 | 7,100 | 280 | 2,096 | 80 | 4,000 | 160 |
| Crappie..... | | | 379 | 28 | | | | |
| Croaker..... | 54,620 | 933 | 4,500 | 180 | 65,600 | 682 | 70,670 | 1,286 |
| Drum, red or redfish..... | | | | | | | 200 | 10 |
| Eels, common..... | 9,604 | 671 | 315 | 20 | 4,518 | 371 | 600 | 30 |
| Flounders..... | 1,751 | 89 | | | | | 5,355 | 237 |
| Gizzard shad..... | 1,000 | 20 | | | | | 1,000 | 15 |
| Hickory shad..... | 150 | 6 | | | | | 1,020 | 21 |
| Pike or pickerel..... | 6,800 | 953 | 1,289 | 154 | 30 | 6 | | |
| Shad..... | 148,560 | 16,047 | 9,490 | 920 | 1,330 | 141 | 138,764 | 12,327 |
| Silver perch..... | | | 500 | 10 | | | | |
| Spot..... | 1,450 | 34 | | | 3,000 | 60 | 925 | 40 |
| Squeteagues or "sea trout": | | | | | | | | |
| Gray..... | 52,544 | 2,212 | 10 | 1 | 146 | 12 | 50,120 | 2,058 |
| Spotted..... | | | | | 450 | 25 | | |
| Striped bass..... | 71,695 | 9,053 | 1,895 | 203 | 3,253 | 363 | 31,150 | 3,479 |
| Suckers..... | 900 | 36 | 2,215 | 94 | | | | |
| White perch..... | 67,402 | 4,087 | 1,500 | 112 | 7,775 | 334 | 3,565 | 176 |
| Whiting..... | | | | | | | 500 | 25 |
| Yellow perch..... | 27,339 | 1,612 | 4,560 | 307 | 13,649 | 889 | | |
| Crabs: | | | | | | | | |
| Hard..... | 1,377,500 | 17,218 | | | 2,855,900 | 35,699 | 498,700 | 6,842 |
| Soft..... | 180,920 | 21,360 | | | 253,436 | 22,696 | 96,519 | 8,752 |
| Oysters: | | | | | | | | |
| Market, public, spring..... | 366,626 | 20,670 | | | 592,085 | 27,327 | 199,252 | 9,381 |
| Market, public, fall..... | 770,034 | 44,097 | | | 1,336,376 | 68,778 | 441,916 | 23,948 |
| Market, private, spring..... | | | | | | | 47,915 | 4,791 |
| Market, private, fall..... | | | | | | | 46,276 | 4,360 |
| Total..... | 3,390,302 | 142,949 | 67,221 | 4,831 | 5,162,747 | 157,735 | 2,266,415 | 81,565 |

Fisheries of Maryland, 1933—Continued

CATCH: BY COUNTIES—Continued

| Species | Somerset | | Talbot | | Wicomico | | Worcester | |
|---------------------------------|-----------|---------|------------|----------|----------|--------|-----------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | 423,900 | \$3,010 | 2,500,925 | \$14,307 | 69,100 | \$889 | 32,500 | \$500 |
| Black bass..... | | | | | 25 | | 19,500 | 885 |
| Bluefish..... | 2,750 | 212 | 41,670 | 3,301 | 7,100 | 445 | 9,100 | 194 |
| Bonito..... | | | 500 | 30 | 2,200 | 130 | 662,600 | 11,250 |
| Butterfish..... | 10,273 | 652 | | | | | 1,100 | 44 |
| Cable or crab eater..... | | | | | | | | |
| Carp..... | 1,764 | 53 | 505 | 37 | 1,500 | 150 | 1,000 | 40 |
| Catfish and bullheads..... | 5,090 | 252 | 11,897 | 462 | 17,800 | 1,000 | 100 | 2 |
| Cod..... | | | | | | | 760,200 | 7,710 |
| Croaker..... | 146,107 | 3,125 | 101,150 | 1,901 | 32,300 | 845 | 500 | 75 |
| Dolphin..... | | | | | | | | |
| Drum: | | | | | | | | |
| Black..... | | | | | 12,000 | 200 | 30,000 | 150 |
| Red or redfish..... | 3,500 | 60 | | | | | 3,500 | 70 |
| Eels: | | | | | | | | |
| Common..... | 380 | 22 | 59,360 | 3,502 | 2,600 | 130 | 21,000 | 1,380 |
| Others..... | | | | | | | 250 | 3 |
| Flounders..... | 18,001 | 574 | 2,845 | 155 | 7,000 | 534 | 18,500 | 925 |
| Gizzard shad..... | 2,100 | 63 | 500 | 22 | 3,000 | 55 | | |
| Harvestfish..... | 3,000 | 180 | | | 8,000 | 480 | 8,500 | 170 |
| Hickory shad..... | 1,406 | 36 | 3,050 | 63 | | | 1,000 | 20 |
| King whiting or "kingfish"..... | | | | | | | 9,600 | 396 |
| Mackerel..... | | | | | | | 27,200 | 272 |
| Mullet..... | 100 | 6 | | | | | 20,000 | 1,200 |
| Pike or pickerel..... | | | | | 200 | 35 | | |
| Scup..... | | | | | | | 52,500 | 1,650 |
| Sea bass..... | | | | | | | 10,100 | 505 |
| Sea robin..... | | | | | | | 500 | 25 |
| Shad..... | 155,048 | 16,266 | 270,131 | 24,310 | 60,103 | 5,738 | 17,899 | 1,629 |
| Spanish mackerel..... | | | | | | | 1,400 | 70 |
| Spot..... | 500 | 10 | 1,600 | 42 | 1,500 | 70 | 11,500 | 115 |
| Squeteagues or "sea trout": | | | | | | | | |
| Gray..... | 62,874 | 2,879 | 5,795 | 261 | 44,000 | 1,765 | 581,900 | 8,823 |
| Spotted..... | 6,313 | 367 | 500 | 50 | | | | |
| Striped bass..... | 7,012 | 985 | 22,986 | 2,165 | 23,497 | 2,812 | 600 | 78 |
| Sturgeon..... | | | | | 160 | 15 | 400 | 80 |
| Suckers..... | | | 100 | 5 | | | | |
| Tautog..... | | | | | | | 5,090 | 150 |
| White perch..... | 9,600 | 522 | 13,583 | 587 | 13,350 | 1,034 | 2,200 | 220 |
| Yellow perch..... | | | 4,000 | 200 | 150 | 13 | | |
| Crabs: | | | | | | | | |
| Hard..... | 4,417,100 | 55,213 | 6,534,000 | 81,674 | 31,800 | 573 | 450,000 | 4,594 |
| Soft..... | 1,852,915 | 111,180 | 209,152 | 18,700 | | | 2,125 | 206 |
| Clams, hard, public..... | | | | | | | 41,536 | 3,894 |
| Oysters: | | | | | | | | |
| Market, public, spring..... | 558,208 | 34,552 | 447,129 | 26,653 | 47,460 | 2,768 | | |
| Market, public, fall..... | 1,752,769 | 139,732 | 867,192 | 58,489 | 169,236 | 9,877 | | |
| Market, private, spring..... | 19,020 | 1,464 | | | 71,190 | 4,153 | 182,000 | 26,675 |
| Market, private, fall..... | 105,917 | 18,417 | | | 253,854 | 14,809 | 193,736 | 27,675 |
| Squid..... | | | | | | | 37,500 | 750 |
| Terrapin, diamond-back..... | 1,800 | 500 | | | | | | |
| Total..... | 9,567,417 | 390,332 | 11,107,480 | 236,916 | 879,125 | 48,519 | 3,150,246 | 102,425 |

VIRGINIA

Fisheries of Virginia, 1933

OPERATING UNITS: BY GEAR

| Item | Purse seines, menhaden | Haul seines | Gill nets | | Lines, trot with baits or snoods | Pound nets | Stop nets | Fyke nets | Dip nets | Otter trawls |
|------------------------------|------------------------|-------------|-----------|---------|----------------------------------|------------|-----------|-----------|----------|--------------|
| | | | Drift | Stake | | | | | | |
| Fishermen: | Number | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| On vessels..... | 1,103 | | | | | | | | | 106 |
| On boats and shore: | | | | | | | | | | |
| Regular..... | | 251 | 76 | 95 | 946 | 1,792 | 9 | 79 | 496 | |
| Casual..... | | 260 | 533 | 189 | 135 | 258 | 3 | 73 | 1,187 | |
| Total..... | 1,103 | 511 | 609 | 284 | 1,081 | 2,050 | 12 | 152 | 1,683 | 106 |
| Vessels: | | | | | | | | | | |
| Steam..... | 25 | | | | | | | | | |
| Net tonnage..... | 2,831 | | | | | | | | | |
| Motor..... | 9 | | | | | | | | | 27 |
| Net tonnage..... | 698 | | | | | | | | | 349 |
| Total vessels..... | 34 | | | | | | | | | 27 |
| Total tonnage..... | 3,529 | | | | | | | | | 399 |
| Boats: | | | | | | | | | | |
| Motor..... | | 163 | 69 | 115 | 946 | 557 | 6 | 63 | 174 | |
| Other..... | | 97 | 336 | 76 | 131 | 630 | 5 | 58 | 1,273 | |
| Accessory boats..... | 102 | | | | | | | | | |
| Apparatus: | | | | | | | | | | |
| Number..... | 34 | 191 | 410 | 4,913 | 1,075 | 1,880 | 6 | 649 | 1,675 | 27 |
| Length, yards..... | 10,160 | 52,063 | | | | | | | | |
| Square yards..... | | | 337,045 | 266,575 | | | | 10,300 | | |
| Yards at mouth..... | | | | | | | | | | 803 |
| Hooks, baits, or snoods..... | | | | | 499,610 | | | | | |

| Item | Pots, eel | Scrapes | Dredges | | | Tongs | Rakes | Picks | By hand | Total, exclusive of duplication |
|----------------------|-----------|---------|---------|--------|---------|--------|--------|--------|---------|---------------------------------|
| | | | Crab | Oyster | Scallop | | | | | |
| Fishermen: | Number | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| On vessels..... | | | 163 | 109 | 4 | 12 | | | | 1,471 |
| On boats and shore: | | | | | | | | | | |
| Regular..... | 6 | 41 | 36 | 208 | | 3,826 | 342 | 540 | 329 | 7,073 |
| Casual..... | | | 3 | | | 592 | 20 | 25 | | 3,071 |
| Total..... | 6 | 41 | 202 | 317 | 4 | 4,430 | 362 | 540 | 354 | 11,615 |
| Vessels: | | | | | | | | | | |
| Steam..... | | | | | | | | | | 25 |
| Net tonnage..... | | | | | | | | | | 2,831 |
| Motor..... | | | 52 | 20 | 1 | 4 | | | | 105 |
| Net tonnage..... | | | 420 | 302 | 18 | 22 | | | | 1,761 |
| Sail..... | | | | 3 | | | | | | 3 |
| Net tonnage..... | | | | 22 | | | | | | 22 |
| Total vessels..... | | | 52 | 23 | 1 | 4 | | | | 133 |
| Total tonnage..... | | | 420 | 324 | 18 | 22 | | | | 4,614 |
| Boats: | | | | | | | | | | |
| Motor..... | 2 | 20 | 13 | 125 | | 2,338 | 92 | 42 | 18 | 4,212 |
| Other..... | 2 | 21 | | | | 536 | 293 | 440 | 156 | 3,621 |
| Accessory boats..... | | | | | | | | | | 102 |
| Apparatus: | | | | | | | | | | |
| Number..... | 35 | 41 | 130 | 298 | 2 | 3,748 | 362 | 540 | | |
| Yards at mouth..... | | 41 | 235 | 339 | 6 | | | | | |

Fisheries of Virginia, 1933—Continued

CATCH: BY GEAR

| Species | Purse seines | | Haul seines | | Gill nets | | | |
|-----------------------------|--------------|-----------|-------------|--------|-----------|--------|---------|--------|
| | | | | | Drift | | Stake | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | | | 59,400 | \$573 | 15,205 | \$256 | 12,800 | \$182 |
| Bluefish..... | | | 59,040 | 2,358 | 2,100 | 105 | | |
| Butterfish..... | | | 1,000 | 25 | | | | |
| Carp..... | | | 310,554 | 12,772 | 440 | 9 | | |
| Catfish and bullheads..... | | | 240,510 | 7,531 | | | | |
| Croaker..... | | | 712,130 | 9,663 | 13,089 | 124 | 2,000 | 40 |
| Drum, red or redfish..... | | | 13,052 | 366 | | | | |
| Eels..... | | | 52,400 | 1,572 | | | | |
| Flounders..... | | | 23,730 | 953 | | | | |
| Gizzard shad..... | | | 26,240 | 517 | 1,740 | 30 | 7,340 | 147 |
| Hickory shad..... | | | 1,389 | 37 | 4,443 | 135 | | |
| Menhaden..... | 115,343,100 | \$384,227 | | | | | | |
| Mullet..... | | | 1,901 | 79 | 3,660 | 98 | 31,180 | 1,145 |
| Pigfish..... | | | 50,740 | 252 | | | | |
| Pike or pickerel..... | | | 25 | 2 | | | | |
| Shad..... | | | 4,564 | 451 | 530,493 | 39,699 | 372,755 | 28,367 |
| Spot..... | | | 296,120 | 6,696 | 33,970 | 1,017 | 1,160 | 35 |
| Squeteagues or "sea trout": | | | | | | | | |
| Gray..... | | | 173,815 | 5,462 | 3,400 | 170 | 4,090 | 178 |
| Spotted..... | | | 150,720 | 12,251 | 3,360 | 190 | | |
| Striped bass..... | | | 80,880 | 9,297 | 41,731 | 4,341 | 63,010 | 7,977 |
| White perch..... | | | 81,440 | 3,239 | 2,750 | 132 | | |
| Yellow perch..... | | | 20,035 | 1,000 | 675 | 28 | | |
| Total..... | 115,343,100 | 384,227 | 2,359,685 | 75,126 | 656,956 | 46,334 | 494,335 | 38,071 |

| Species | Lines, trot with baits or snoods | | Pound nets | | Stop nets | | Fyke nets | |
|---------------------------------|----------------------------------|-----------|------------|----------|-----------|---------|-----------|--------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | | | 19,071,306 | \$85,561 | | | 18,520 | \$192 |
| Bluefish..... | | | 595,160 | 22,361 | | | | |
| Bonito..... | | | 10,092 | 455 | | | | |
| Butterfish..... | | | 2,220,455 | 53,148 | | | 4,650 | 134 |
| Cabio or crab eater..... | | | 18,842 | 791 | | | | |
| Carp..... | | | 46,010 | 1,606 | 52,263 | \$1,856 | 19,197 | 634 |
| Catfish and bullheads..... | | | 202,855 | 6,156 | 11,241 | 452 | 263,802 | 8,212 |
| Cod..... | | | 6,525 | 130 | | | | |
| Croaker..... | | | 10,881,977 | 131,247 | | | 18,880 | 326 |
| Drum: | | | | | | | | |
| Black..... | | | 80,480 | 1,594 | | | | |
| Red or redfish..... | | | 34,069 | 1,018 | | | | |
| Eels..... | | | 41,180 | 2,161 | | | 8,328 | 434 |
| Flounders..... | | | 281,890 | 19,746 | | | 4,000 | 60 |
| Gizzard shad..... | | | 60,180 | 1,322 | | | 21,320 | 363 |
| Harvestfish..... | | | 160,050 | 3,585 | | | | |
| Hickory shad..... | | | 49,255 | 1,007 | | | 1,141 | 17 |
| King whiting or "kingfish"..... | | | 29,211 | 841 | | | | |
| Mackerel..... | | | 16,056 | 893 | | | | |
| Menhaden..... | | | 647,400 | 1,598 | | | | |
| Mullet..... | | | 5,695 | 231 | | | 2,205 | 95 |
| Pigfish..... | | | 6,430 | 129 | | | | |
| Pike or pickerel..... | | | | | | | 175 | 9 |
| Pompano..... | | | 4,530 | 248 | | | | |
| Scup..... | | | 337,540 | 13,005 | | | | |
| Sea bass..... | | | 4,980 | 199 | | | | |
| Shad..... | | | 3,902,955 | 370,916 | | | 5,947 | 594 |
| Spanish mackerel..... | | | 66,501 | 3,950 | | | | |
| Spot..... | | | 358,595 | 9,764 | | | | |
| Squeteagues or "sea trout": | | | | | | | | |
| Gray..... | | | 11,754,530 | 211,616 | | | 10,760 | 399 |
| Spotted..... | | | 9,740 | 525 | | | | |
| Striped bass..... | | | 303,151 | 38,758 | | | 30,128 | 3,101 |
| Sturgeon..... | | | 4,946 | 1,112 | | | | |
| Tautog..... | | | 560 | 12 | | | | |
| Thimble-eyed mackerel..... | | | 11,250 | 169 | | | | |
| White perch..... | | | 82,089 | 3,320 | | | 81,069 | 3,134 |
| Yellow perch..... | | | 5,150 | 258 | | | 46,005 | 2,313 |
| Crabs, hard..... | 17,047,155 | \$169,599 | | | | | | |
| Squid..... | | | 94,440 | 2,339 | | | 1,500 | 60 |
| Turtles, snapper..... | | | | | | | | |
| Total..... | 17,047,155 | 169,599 | 51,405,955 | 991,771 | 63,504 | 2,308 | 537,617 | 20,069 |

U. S. BUREAU OF FISHERIES

Fisheries of Virginia, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Dip nets | | Otter trawls | | Pots, eel | | Scrapes | |
|---------------------------------|-----------|---------|--------------|---------|-----------|-------|---------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | | | 217 | \$2 | | | | |
| Bluefish..... | | | 28,059 | 2,205 | | | | |
| Butterfish..... | | | 59,248 | 1,806 | | | | |
| Cod..... | | | 341 | 7 | | | | |
| Croaker..... | | | 2,607,106 | 45,318 | | | | |
| Drum, red or redfish..... | | | 4,418 | 72 | | | | |
| Eels..... | | | 4,050 | 66 | 2,676 | \$114 | | |
| Flounders..... | | | 773,793 | 31,272 | | | | |
| Haddock..... | | | 47 | 1 | | | | |
| Hake..... | | | 23,455 | 387 | | | | |
| King whiting or "kingfish"..... | | | 35,092 | 1,209 | | | | |
| Mackereel..... | | | 160 | 8 | | | | |
| Pigfish..... | | | 3,595 | 109 | | | | |
| Pollock..... | | | 25 | 1 | | | | |
| Scup..... | | | 1,168,591 | 23,319 | | | | |
| Sea bass..... | | | 327,095 | 9,193 | | | | |
| Sharks..... | | | 10,225 | 159 | | | | |
| Spot..... | | | 26,820 | 498 | | | | |
| Squeteagues or "sea trout": | | | | | | | | |
| Gray..... | | | 363,543 | 13,156 | | | | |
| Spotted..... | | | 595 | 31 | | | | |
| Sturgeon..... | | | 3,196 | 484 | | | | |
| Swellfish..... | | | 1,065 | 53 | | | | |
| Tautog..... | | | 488 | 9 | | | | |
| White perch..... | | | 102 | 3 | | | | |
| Crabs: | | | | | | | | |
| Hard..... | 193,100 | \$1,968 | | | | | 116,550 | \$1,058 |
| Soft..... | 1,939,257 | 135,629 | | | | | 128,700 | 7,722 |
| Lobsters..... | | | 131 | 20 | | | | |
| Shrimp..... | | | 371 | 2 | | | | |
| Scallops, sea..... | | | 195 | 44 | | | | |
| Squid..... | | | 13,353 | 288 | | | | |
| Total..... | 2,132,357 | 137,697 | 5,455,375 | 129,812 | 2,676 | 114 | 245,250 | 8,780 |

| Species | Dredges | | | | | | Tongs | |
|------------------------------|-----------|-----------|-----------|---------|---------|---------|-----------|-----------|
| | Crab | | Oyster | | Scallop | | Pounds | Value |
| | Pounds | Value | Pounds | Value | Pounds | Value | | |
| Crabs, hard..... | 6,554,570 | \$111,273 | | | | | | |
| Crabs, hard, public..... | | | | | | | 718,522 | \$152,074 |
| Oysters: | | | | | | | | |
| Market, public, spring..... | | | 12,102 | \$672 | | | 1,645,547 | 94,280 |
| Market, public, fall..... | | | 600 | 35 | | | 2,624,965 | 147,599 |
| Market, private, spring..... | | | 1,846,178 | 142,986 | | | 1,608,059 | 117,311 |
| Market, private, fall..... | | | 3,184,544 | 239,613 | | | 1,954,013 | 142,930 |
| Scallops, sea..... | | | | | 72,450 | \$8,050 | | |
| Total..... | 6,554,570 | 111,273 | 5,043,424 | 383,036 | 72,450 | 8,050 | 8,551,106 | 654,194 |

| Species | Rakes | | Picks | | By hand | |
|------------------------------|---------|----------|---------|----------|---------|----------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Clams, hard, public..... | 91,160 | \$22,790 | 279,864 | \$69,464 | 79,720 | \$19,930 |
| Oysters: | | | | | | |
| Market, public, spring..... | 1,200 | 100 | | | 1,200 | 90 |
| Market, public, fall..... | 5,000 | 435 | | | 1,440 | 120 |
| Market, private, spring..... | 302,464 | 26,143 | | | | |
| Market, private, fall..... | 280,460 | 24,741 | | | | |
| Terrapin, diamond-back..... | | | | | 10,240 | 2,500 |
| Total..... | 680,284 | 74,209 | 279,864 | 69,464 | 92,600 | 22,640 |

Fisheries of Virginia, 1933—Continued

OPERATING UNITS: BY COUNTIES

| Item | Acco- mac | Arling- ton | Caro- line | Charles City | Ches- ter- field | Din- wid- die | Eliza- beth City | Essex | Fair- fax |
|--|--------------|----------------|---------------|-----------------|------------------------|---------------------|------------------------|--------|--------------|
| | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | | | |
| On vessels..... | 19 | | | | | | 124 | | |
| On boats and shore: | | | | | | | | | |
| Regular..... | 1,432 | 4 | | 10 | | | 153 | 108 | 34 |
| Casual..... | 89 | 18 | 6 | 98 | 23 | 9 | | 24 | 41 |
| Total..... | 1,540 | 22 | 6 | 108 | 23 | 9 | 277 | 132 | 75 |
| Vessels: | | | | | | | | | |
| Motor..... | 2 | | | | | | 31 | | |
| Net tonnage..... | 11 | | | | | | 418 | | |
| Sail..... | 3 | | | | | | | | |
| Net tonnage..... | 22 | | | | | | | | |
| Total vessels..... | 5 | | | | | | 31 | | |
| Total net tonnage..... | 33 | | | | | | 418 | | |
| Boats: | | | | | | | | | |
| Motor..... | 727 | 10 | | 5 | | | 81 | 42 | 34 |
| Other..... | 727 | 2 | 3 | 68 | 14 | 5 | 7 | 49 | 15 |
| Apparatus: | | | | | | | | | |
| Haul seines..... | 13 | | 1 | 2 | 2 | 1 | | 3 | 4 |
| Length, yards..... | 8,640 | | 300 | 325 | 400 | 200 | | 1,000 | 840 |
| Gill nets: | | | | | | | | | |
| Dr ft..... | | 11 | 2 | 64 | 12 | 4 | | 7 | 30 |
| Square yards..... | | 22,415 | 1,800 | 39,040 | 3,960 | 2,080 | | 5,600 | 45,800 |
| Stake..... | 4 | | | 110 | | | | | |
| Square yards..... | 1,440 | | | 3,340 | | | | | |
| Lines, trot with baits or snoods..... | 168 | | | | | | 9 | 3 | |
| Baits or snoods..... | 86,600 | | | | | | 3,500 | 1,200 | |
| Pound nets..... | 214 | | | | | | 137 | 2 | |
| Stop nets..... | | | | 2 | | | | | |
| Square yards..... | | | | 1,600 | | | | | |
| Fyke nets..... | 3 | 2 | | 19 | 3 | | | | 196 |
| Dip nets..... | 364 | | | | | | | | |
| Otter trawls..... | | | | | | | 11 | | |
| Yards at mouth..... | | | | | | | 330 | | |
| Pots, eel..... | | | | 14 | | | | | |
| Scrapes..... | 41 | | | | | | | | |
| Yards at mouth..... | 41 | | | | | | | | |
| Dredges: | | | | | | | | | |
| Crab..... | 2 | | | | | | 32 | | |
| Yards at mouth..... | 3 | | | | | | 64 | | |
| Oyster..... | 256 | | | | | | 14 | | |
| Yards at mouth..... | 269 | | | | | | 25 | | |
| Scallop..... | | | | | | | 2 | | |
| Yards at mouth..... | | | | | | | 6 | | |
| Tongs..... | 684 | | | | | | 43 | 99 | |
| Rakes..... | 174 | | | | | | | | |
| Picks..... | 315 | | | | | | | | |

| Item | Gloucester | Henrico | Isle of Wight | James City | King and Queen | King George | King William | Lan- caster | Math- ews |
|------------------------|------------|---------|---------------------|---------------|----------------------|----------------|-----------------|----------------|--------------|
| | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | | | |
| On vessels..... | 6 | | 3 | | | | | 443 | 16 |
| On boats and shore: | | | | | | | | | |
| Regular..... | 338 | 3 | 519 | 37 | | 72 | | 536 | 642 |
| Casual..... | 39 | 65 | 59 | 31 | 51 | 74 | 83 | 417 | 211 |
| Total..... | 383 | 68 | 581 | 68 | 51 | 146 | 83 | 1,396 | 869 |
| Vessels: | | | | | | | | | |
| Steam..... | | | | | | | | 8 | |
| Net tonnage..... | | | | | | | | 820 | |
| Motor..... | 2 | | 1 | | | | | 9 | 4 |
| Net tonnage..... | 15 | | 5 | | | | | 385 | 36 |
| Total vessels..... | 2 | | 1 | | | | | 17 | 4 |
| Total net tonnage..... | 15 | | 5 | | | | | 1,205 | 36 |

Fisheries of Virginia, 1933—Continued

OPERATING UNITS: BY COUNTIES—Continued

| Item | Gloucester | Henrico | Isle of Wright | James City | King and Queen | King George | King William | Lancaster | Mathews |
|--|---------------|---------------|----------------|---------------|----------------|---------------|---------------|---------------|---------------|
| Boats: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| Motor..... | 218 | | 226 | 29 | | 61 | 5 | 374 | 356 |
| Other..... | 72 | 40 | 73 | 27 | 33 | 39 | 49 | 293 | 306 |
| Accessory boats..... | | | | | | | | 42 | |
| Apparatus: | | | | | | | | 14 | |
| Purse seines, menhaden | | | | | | | | 4,240 | |
| Length, yards..... | | | | | | | | 1 | 5 |
| Haul seines..... | | 3 | 1 | 2 | 1 | 5 | 2 | | |
| Length, yards..... | | 375 | 200 | 540 | 200 | 1,380 | 173 | 800 | 1,980 |
| Gill nets: | | | | | | | | | |
| Drift..... | 4 | 37 | | 13 | 31 | 17 | 49 | | 6 |
| Square yards..... | 3,200 | 19,240 | | 7,300 | 19,600 | 21,870 | 20,890 | | 3,100 |
| Stake..... | | | 1,720 | 1,024 | 600 | | | | |
| Square yards..... | | | 58,480 | 30,820 | 70,000 | | | | |
| Lines, trot, with baits or snoods..... | 52 | | 61 | | | 16 | | 47 | 88 |
| Baits or snoods..... | 26,400 | | 33,400 | | | 8,500 | | 25,300 | 46,400 |
| Pound nets..... | 107 | | 7 | 7 | | 36 | | 182 | 459 |
| Fyke nets..... | 13 | | 96 | 43 | 2 | 30 | 6 | | |
| Dip nets..... | | | | | | | | 290 | 86 |
| Dredges: | | | | | | | | | |
| Crab..... | 4 | | | | | | | | 6 |
| Yards at mouth..... | 8 | | | | | | | | 12 |
| Oyster..... | | | | | | | | 6 | 8 |
| Yards at mouth..... | | | | | | | | 9 | 12 |
| Tongs..... | 186 | | 468 | 7 | | 12 | | 426 | 172 |

| Item | Middlesex | Nansemond | New Kent | Norfolk | Northampton | Northumberland | Prince George | Princess Anne |
|--|---------------|---------------|---------------|---------------|---------------|----------------|---------------|---------------|
| Fishermen: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| On vessels..... | 4 | 3 | | 33 | | 670 | | |
| On boats and shore: | | | | | | | | |
| Regular..... | 619 | 141 | 6 | 73 | 503 | 822 | 7 | 72 |
| Casual..... | 286 | 7 | 67 | 58 | 63 | 744 | 24 | 147 |
| Total..... | 909 | 151 | 73 | 164 | 566 | 2,236 | 31 | 219 |
| Vessels: | | | | | | | | |
| Steam..... | | | | | | 17 | | |
| Net tonnage..... | | | | | | 2,011 | | |
| Motor..... | 1 | 1 | | 6 | | 3 | | |
| Net tonnage..... | 8 | 6 | | 112 | | 352 | | |
| Total vessels..... | 1 | 1 | | 6 | | 20 | | |
| Total net tonnage..... | 8 | 6 | | 112 | | 2,363 | | |
| Boats: | | | | | | | | |
| Motor..... | 475 | 56 | 2 | 47 | 209 | 469 | 4 | 81 |
| Other..... | 164 | | 57 | 71 | 404 | 823 | 17 | 69 |
| Accessory boats..... | | | | | | 60 | | |
| Apparatus: | | | | | | 20 | | |
| Purse seines, menhaden | | | | | | 5,920 | | |
| Length, yards..... | | | | | | | 4 | 81 |
| Haul seines..... | 7 | | 3 | 5 | 7 | | | |
| Length, yards..... | 2,180 | | 800 | 2,460 | 2,080 | | 520 | 16,820 |
| Gill nets: | | | | | | | | |
| Drift..... | | | 56 | 16 | | 3 | 13 | |
| Square yards..... | | | 44,800 | 3,200 | | 2,940 | 9,860 | |
| Stake..... | | 189 | | 210 | 4 | | | |
| Square yards..... | | 5,670 | | 6,300 | 2,080 | | | |
| Lines, trot, with baits or snoods..... | 49 | | | 50 | 113 | 204 | | 32 |
| Baits or snoods..... | 32,600 | | | 25,560 | 5,600 | 105,760 | | 9,600 |
| Pound nets..... | 14 | | | 28 | 120 | 411 | | 6 |
| Stop nets..... | | | | | | | 4 | |
| Square yards..... | | | | | | | 8,700 | |
| Fyke nets..... | | 2 | 15 | | 3 | 4 | 79 | |
| Dip nets..... | 148 | | | | 46 | 741 | | |
| Otter trawls..... | | | | 2 | | | | |
| Yards at mouth..... | | | | 60 | | | | |
| Pots, eel..... | | | 21 | | | | | |
| Dredges: | | | | | | | | |
| Oyster..... | 2 | | | 10 | | | | |
| Yards at mouth..... | 3 | | | 17 | | | | |
| Tongs..... | 539 | 104 | | 56 | 190 | 135 | | 34 |
| Rakes..... | | | | | 188 | | | |
| Picks..... | | | | | 225 | | | |

Fisheries of Virginia, 1933—Continued

OPERATING UNITS: BY COUNTIES—Continued

| Item | Prince Wil- liam | Rich- mond | Spot- syl- vania | Staf- ford | Surry | War- wick | West- more- land | York |
|--|------------------------|---------------|------------------------|---------------|--------|--------------|------------------------|--------|
| | Number | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | | 134 |
| On vessels..... | | | | | | 16 | | |
| On boats and shore: | | | | | | | | 238 |
| Regular..... | 24 | 121 | | 62 | 14 | 226 | 257 | |
| Casual..... | 40 | 67 | 2 | 34 | 29 | 43 | 118 | 4 |
| Total..... | 64 | 188 | 2 | 96 | 43 | 285 | 375 | 378 |
| Vessels: | | | | | | | | 41 |
| Motor..... | | | | | | 4 | | 360 |
| Net tonnage..... | | | | | | 53 | | |
| Boats: | | | | | | | | 163 |
| Motor..... | 35 | 48 | | 43 | 14 | 149 | 249 | 18 |
| Other..... | 5 | 74 | 1 | 16 | 14 | 6 | 60 | |
| Apparatus: | | | | | | | | 11 |
| Haul seines..... | 5 | | 1 | 12 | 3 | | 3 | 0,800 |
| Length, yards..... | 380 | | 300 | 1,180 | 780 | | 410 | |
| Gill nets: | | | | | | | | 2 |
| Drift..... | 13 | 16 | 1 | 2 | | | 1 | 1,860 |
| Square yards..... | 23,490 | 17,400 | 500 | 7,000 | | | 1,100 | 12 |
| Stake..... | 44 | | | 112 | 672 | 112 | 10 | 360 |
| Square yards..... | 20,145 | | | 42,900 | 20,160 | 4,480 | 400 | 46 |
| Lines, trot, with baits or snoods..... | 2 | 9 | | | | | 58,400 | 22,500 |
| Baits or snoods..... | 1,000 | 3,850 | | 3,500 | | | 10 | 29 |
| Pound nets..... | | 24 | | 14 | | | 73 | 6 |
| Fyke nets..... | 64 | 1 | | 35 | 21 | 3 | 3 | 10 |
| Otter trawls..... | | | | | | | | 296 |
| Yards at mouth..... | | | | | | 117 | | |
| Dredges: | | | | | | | | 86 |
| Crab..... | | | | | | | | 148 |
| Yards at mouth..... | | | | | | | | 2 |
| Oyster..... | | | | | | | | 4 |
| Yards at mouth..... | | | | | | | | 121 |
| Tongs..... | | 95 | | | | 219 | 148 | |

CATCH: BY COUNTIES

| Species | Accomac | | Arlington | | Caroline | | Charles City | |
|---------------------------------|------------|---------|-----------|-------|----------|-------|--------------|-------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alowives..... | 770,060 | \$3,850 | | | | | | |
| Bluefish..... | 64,350 | 3,088 | | | | | | |
| Bonito..... | 6,252 | 301 | | | | | | |
| Butterfish..... | 581,660 | 14,559 | | | | | | |
| Carp..... | | | | | 1,500 | 42 | 23,677 | 726 |
| Catfish and bullheads..... | 780 | 31 | 830 | \$33 | 900 | 27 | 32,197 | 1,452 |
| Cod..... | 1,800 | 36 | | | | | | |
| Croaker..... | 1,323,147 | 20,252 | | | | | | |
| Drum: | | | | | | | | |
| Black..... | 80,480 | 1,594 | | | | | | |
| Red or redfish..... | 25,559 | 574 | | | | | | |
| Eels..... | 32,040 | 1,615 | | | | | 2,196 | 86 |
| Flounders..... | 54,400 | 2,176 | | | | | | |
| Gizzard shad..... | | | | | 800 | 20 | 3,500 | 36 |
| Harvestfish..... | 8,060 | 98 | | | | | | |
| Hickory shad..... | | | | | 200 | 8 | | |
| King whiting or "kingfish"..... | 1,875 | 75 | | | | | | |
| Mackerel..... | 5,828 | 381 | | | | | | |
| Mullet..... | 4,640 | 232 | | | 500 | 15 | | |
| Scup..... | 105,620 | 3,774 | | | | | | |
| Shad..... | 185,827 | 17,115 | 69,188 | 3,731 | 900 | 135 | 66,657 | 6,820 |
| Spanish mackerel..... | 4,545 | 266 | | | | | | |
| Spot..... | 32,295 | 896 | | | | | | |
| Squeteagues or "sea trout": | | | | | | | | |
| Gray..... | 1,517,950 | 37,292 | | | | | | |
| Spotted..... | 2,640 | 137 | | | | | | |
| Striped bass..... | 8,540 | 864 | | | 200 | 20 | 2,688 | 338 |
| Sturgeon..... | 2,600 | 521 | | | | | | |
| Thimble-eyed mackerel..... | 11,250 | 109 | | | | | | |
| White perch..... | 560 | 16 | 225 | 10 | 1,000 | 50 | 1,454 | 64 |
| Yellow perch..... | | | 245 | 12 | 700 | 35 | 75 | 7 |
| Crabs: | | | | | | | | |
| Hard..... | 2,648,590 | 27,232 | | | | | | |
| Soft..... | 932,429 | 49,254 | | | | | | |
| Clams, hard, public..... | 606,518 | 126,852 | | | | | | |
| Oysters: | | | | | | | | |
| Market, public, spring..... | 122,174 | 7,749 | | | | | | |
| Market, public, fall..... | 197,444 | 13,721 | | | | | | |
| Market, private, spring..... | 682,140 | 92,016 | | | | | | |
| Market, private, fall..... | 1,095,240 | 103,817 | | | | | | |
| Squid..... | 55,500 | 1,560 | | | | | | |
| Terrapin, diamond-back..... | 10,210 | 2,500 | | | | | | |
| Total..... | 11,452,931 | 534,413 | 70,488 | 3,786 | 7,700 | 370 | 154,144 | 9,743 |

Fisheries of Virginia, 1933—Continued

CATCH: BY COUNTIES—Continued

| Species | Chesterfield | | Dinwiddie | | Elizabeth City | | Essex | |
|---------------------------------|---------------|------------|--------------|------------|------------------|----------------|----------------|---------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | 5,370 | \$85 | 4,650 | \$46 | 118,047 | \$621 | 4,800 | \$76 |
| Bluefish..... | | | | | 65,369 | 3,268 | | |
| Butterfish..... | | | | | 126,160 | 3,476 | | |
| Carp..... | 3,600 | 99 | 2,800 | 112 | | | 7,000 | 340 |
| Catfish and bullheads..... | 680 | 22 | | | | | 11,500 | 450 |
| Cod..... | | | | | 138 | 3 | | |
| Croaker..... | | | | | 2,966,978 | 45,474 | 7,500 | 155 |
| Drum, red or redfish..... | | | | | 4,188 | 67 | | |
| Eels..... | | | | | 3,768 | 62 | | |
| Flounders..... | | | | | 457,554 | 18,603 | | |
| Gizzard shad..... | | | | | 1,200 | 18 | 14,500 | 310 |
| Haddock..... | | | | | 47 | 1 | | |
| Hake..... | | | | | 20,524 | 326 | | |
| Harvestfish..... | | | | | 8,460 | 212 | | |
| Hickory shad..... | 615 | 15 | 250 | 5 | 14,100 | 282 | 1,050 | 29 |
| King whiting or "kingfish"..... | | | | | 21,602 | 736 | | |
| Mackerel..... | | | | | 150 | 7 | | |
| Mullet..... | 35 | 2 | | | 3,000 | 116 | 800 | 36 |
| Pigfish..... | | | | | 2,813 | 95 | | |
| Pollock..... | | | | | 25 | 1 | | |
| Scup..... | | | | | 605,291 | 12,112 | | |
| Sea bass..... | | | | | 239,992 | 7,017 | | |
| Shad..... | 3,115 | 321 | 480 | 48 | 191,900 | 19,190 | 5,800 | 545 |
| Sharks..... | | | | | 8,866 | 119 | | |
| Spanish mackerel..... | | | | | 1,350 | 54 | | |
| Spot..... | | | | | 80,999 | 2,257 | | |
| Squeteagues or "sea trout": | | | | | | | | |
| Gray..... | | | | | 643,911 | 13,282 | | |
| Spotted..... | | | | | 595 | 31 | | |
| Striped bass..... | 310 | 20 | 1,240 | 136 | 41,500 | 4,150 | 4,700 | 530 |
| Sturgeon..... | | | | | 2,879 | 450 | | |
| Swellfish..... | | | | | 491 | 25 | | |
| Tautog..... | | | | | 891 | 19 | | |
| White perch..... | | | | | 102 | 3 | 1,600 | 80 |
| Yellow perch..... | | | | | | | 800 | 40 |
| Crabs, hard..... | | | | | 2,533,750 | 44,196 | 32,760 | 312 |
| Lobsters..... | | | | | 131 | 2 | | |
| Shrimp..... | | | | | 172 | 9 | | |
| Clams, hard, public..... | | | | | 63,120 | 15,780 | | |
| Oysters: | | | | | | | | |
| Market, public, spring..... | | | | | 19,200 | 1,280 | 115,577 | 5,779 |
| Market, public, fall..... | | | | | 43,200 | 2,850 | 144,227 | 7,211 |
| Market, private, spring..... | | | | | 356,619 | 24,037 | 74,718 | 4,280 |
| Market, private, fall..... | | | | | 1,050,090 | 74,974 | 63,207 | 3,040 |
| Scallops, sea..... | | | | | 72,645 | 8,094 | | |
| Squid..... | | | | | 12,534 | 267 | | |
| Total..... | 13,725 | 544 | 9,420 | 347 | 9,764,261 | 303,596 | 480,539 | 23,193 |

| Species | Fairfax | | Gloucester | | Henrico | | Isle of Wight | |
|---------------------------------|----------------|---------------|------------------|----------------|---------------|--------------|----------------|---------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | | | 18,200 | \$114 | 1,355 | \$11 | 10,300 | \$206 |
| Bluefish..... | | | 105,735 | 4,229 | | | | |
| Butterfish..... | | | 295,980 | 5,919 | | | 5,150 | 154 |
| Cabio or crab eater..... | | | 820 | 49 | | | | |
| Carp..... | 7,620 | \$229 | | | 34,730 | 1,106 | 12,580 | 377 |
| Catfish and bullheads..... | 106,290 | 3,178 | 1,040 | 21 | 5,235 | 178 | 19,600 | 688 |
| Croaker..... | | | 1,633,960 | 24,514 | | | 60,500 | 1,210 |
| Eels..... | | | 370 | 18 | | | 2,000 | 50 |
| Flounders..... | | | 16,540 | 661 | | | | |
| Gizzard shad..... | 480 | 10 | | | | | 26,460 | 530 |
| Harvestfish..... | | | 62,400 | 1,248 | | | | |
| Hickory shad..... | | | 968 | 19 | | | | |
| King whiting or "kingfish"..... | | | 19,561 | 391 | | | | |
| Menhaden..... | | | 70,000 | 226 | | | | |
| Mullet..... | | | | | | | 9,000 | 450 |
| Shad..... | 205,366 | 12,516 | 172,940 | 17,294 | 9,508 | 1,152 | 62,720 | 6,527 |
| Spanish mackerel..... | | | 1,496 | 150 | | | | |
| Spot..... | | | 36,680 | 1,100 | | | | |
| Squeteagues or "sea trout": | | | | | | | | |
| Gray..... | | | 1,472,240 | 18,404 | | | 8,500 | 425 |
| Spotted..... | | | 940 | 51 | | | | |
| Striped bass..... | 26,830 | 1,945 | 427 | 47 | 615 | 73 | 9,270 | 943 |
| White perch..... | 32,830 | 1,249 | 260 | 10 | 100 | 5 | 12,300 | 461 |
| Yellow perch..... | 23,190 | 1,159 | 540 | 27 | 75 | 3 | | |
| Crabs, hard..... | | | 1,855,500 | 19,110 | | | 160,800 | 1,940 |
| Clams, hard, public..... | | | 166,400 | 39,200 | | | 9,600 | 2,400 |
| Oysters: | | | | | | | | |
| Market, public, spring..... | | | 104,340 | 7,350 | | | 235,440 | 15,706 |
| Market, public, fall..... | | | 177,400 | 12,560 | | | 255,120 | 17,008 |
| Market, private, spring..... | | | | | | | 34,800 | 2,906 |
| Market, private, fall..... | | | | | | | 49,200 | 4,100 |
| Total..... | 402,606 | 20,286 | 6,214,727 | 152,712 | 51,618 | 2,528 | 973,100 | 55,375 |

Fisheries of Virginia, 1933—Continued

CATCH: BY COUNTIES—Continued

| Species | James City | | King and Queen | | King George | | King William | |
|---------------------------------------|----------------|--------------|----------------|--------------|----------------|---------------|---------------|--------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | 7,400 | \$74 | 1,440 | \$28 | 142,000 | \$770 | 17,540 | \$132 |
| Bluefish..... | 150 | 8 | | | | | | |
| Butterfish..... | | | | | 2,500 | 85 | | |
| Carp..... | 20,740 | 709 | | | 12,440 | 429 | 20 | 1 |
| Catfish and bullheads..... | 20,330 | 519 | 840 | 25 | 185,550 | 4,768 | 458 | 23 |
| Croaker..... | 11,880 | 185 | | | 1,700 | 37 | | |
| Gizzard shad..... | 23,880 | 478 | | | 4,400 | 93 | | |
| Hickory shad..... | | | 480 | 15 | 2,500 | 83 | 620 | 6 |
| Mullet..... | | | 1,560 | 42 | 1,550 | 72 | 238 | 6 |
| Pike or pickerel..... | | | | | | | 25 | 2 |
| Shad..... | 32,860 | 3,286 | 10,148 | 1,066 | 125,454 | 9,872 | 15,936 | 1,342 |
| Squeteagues or "sea trout", gray..... | 1,000 | 30 | | | 9,520 | 286 | | |
| Striped bass..... | 21,570 | 2,167 | 530 | 56 | 106,180 | 15,811 | | 89 |
| White perch..... | 14,640 | 595 | 340 | 16 | 56,475 | 2,306 | 130 | 9 |
| Yellow perch..... | 950 | 34 | | | 3,890 | 213 | 100 | 4 |
| Crabs, hard..... | | | | | 192,000 | 3,200 | | |
| Clams, hard, public..... | 1,000 | 175 | | | | | | |
| Oysters: | | | | | | | | |
| Market, public, spring..... | | | | | 8,680 | 620 | | |
| Market, private, spring..... | | | | | 10,640 | 760 | | |
| Market, private, fall..... | | | | | 14,400 | 1,050 | | |
| Total..... | 156,400 | 8,264 | 15,338 | 1,248 | 879,879 | 40,423 | 35,152 | 1,584 |

| Species | Lancaster | | Mathews | | Middlesex | | Nansemond | |
|------------------------------|-------------------|----------------|-------------------|----------------|------------------|---------------|----------------|---------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | 1,711,400 | \$8,564 | 488,500 | \$2,441 | 30,000 | \$170 | | |
| Bluefish..... | 89,020 | 4,126 | 205,710 | 6,306 | 7,860 | 460 | | |
| Butterfish..... | | | 472,319 | 11,812 | | | | |
| Cabio or crab eater..... | | | 15,910 | 636 | | | | |
| Carp..... | | | | | 37,840 | 915 | | |
| Catfish and bullheads..... | | | | | 2,000 | 60 | | |
| Croaker..... | 502,720 | 5,327 | 3,492,100 | 34,921 | 136,635 | 1,472 | 2,640 | \$48 |
| Flounders..... | 27,580 | 1,134 | 53,900 | 10,528 | 500 | 20 | | |
| Gizzard shad..... | 480 | 24 | | | | | 780 | 9 |
| Harvestfish..... | | | 10,830 | 270 | | | | |
| Menhaden..... | 44,840,200 | 149,071 | 92,000 | 220 | | | | |
| Pigfish..... | | | 6,430 | 129 | | | | |
| Shad..... | 409,430 | 40,943 | 1,035,293 | 93,176 | 8,970 | 897 | 3,420 | 342 |
| Spanish mackerel..... | | | 13,460 | 673 | | | | |
| Spot..... | 31,920 | 791 | 172,610 | 4,284 | 41,640 | 1,009 | | |
| Squeteagues or "sea trout": | | | | | | | | |
| Gray..... | 569,970 | 11,399 | 3,177,600 | 39,788 | 42,100 | 1,047 | | |
| Spotted..... | 10,400 | 5,024 | 30,800 | 1,563 | | | | |
| Striped bass..... | 12,940 | 5,675 | 26,230 | 2,555 | 13,300 | 1,339 | | |
| Sturgeon..... | | | 918 | 367 | | | | |
| White perch..... | | | 4,590 | 138 | 1,160 | 58 | 980 | 89 |
| Crabs: | | | | | | | | |
| Hard..... | 633,000 | 3,165 | 1,223,520 | 11,802 | 764,520 | 6,370 | | |
| Soft..... | 174,600 | 23,200 | 35,520 | 1,954 | 356,200 | 28,496 | 70,400 | 17,600 |
| Clams, hard, public..... | | | 23,120 | 5,460 | | | | |
| Oysters: | | | | | | | | |
| Market, public, spring..... | 413,241 | 20,662 | 22,303 | 1,312 | 188,657 | 9,433 | 120,000 | 7,960 |
| Market, public, fall..... | 716,285 | 33,814 | 48,036 | 3,049 | 448,910 | 22,445 | 177,300 | 11,745 |
| Market, private, spring..... | 305,226 | 17,003 | 182,641 | 13,602 | 121,664 | 7,580 | 144,000 | 12,000 |
| Market, private, fall..... | 489,597 | 28,942 | 193,686 | 14,615 | 188,963 | 15,448 | 162,000 | 13,500 |
| Total..... | 50,938,709 | 358,864 | 11,027,986 | 261,699 | 2,390,909 | 95,219 | 681,520 | 63,243 |

| Species | New Kent | | Norfolk | | Northampton | | Northumberland | |
|----------------------------|----------|-------|---------|-------|-------------|--------|----------------|----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives..... | 5,000 | \$50 | 133,000 | \$665 | 43,200 | \$218 | 12,675,246 | \$57,682 |
| Bluefish..... | | | 7,992 | 406 | 27,250 | 1,200 | 63,025 | 2,206 |
| Bonito..... | | | | | 3,840 | 154 | | |
| Butterfish..... | | | 206,871 | 4,185 | 410,200 | 10,284 | 22,796 | 671 |
| Cabio or crab eater..... | | | | | | | 2,112 | 106 |
| Carp..... | 4,350 | 174 | | | | | | |
| Catfish and bullheads..... | 13,750 | 354 | | | | | | |
| Cod..... | | | 115 | 2 | 4,725 | 94 | | |
| Croaker..... | | | 349,022 | 3,689 | 747,520 | 12,837 | 1,604,955 | 16,041 |
| Drum, red or redfish..... | | | 1,340 | 53 | 7,302 | 278 | 2,640 | 106 |
| Eels..... | 580 | 35 | 282 | 4 | 19,130 | 655 | 1,800 | 128 |
| Flounders..... | | | 91,370 | 3,472 | 51,470 | 2,059 | 29,290 | 1,071 |

Fisheries of Virginia, 1933—Continued

CATCH: BY COUNTIES—Continued

| Species | New Kent | | Norfolk | | Northampton | | Northumberland | |
|-----------------------------|---------------|--------------|------------------|----------------|------------------|----------------|-------------------|----------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Glizzard shad | 7,040 | \$145 | | | | | | |
| Hake | | | 1,484 | \$28 | | | | |
| Harvestfish | | | 62,500 | 1,662 | | | | |
| Hickory shad | | | | | | | 19,425 | \$399 |
| King whiting or "kingfish" | | | 4,814 | 260 | 3,845 | \$154 | 350 | 11 |
| Mackerel | | | | | 6,810 | 341 | | |
| Menhaden | | | | | | | 70,988,300 | 236,308 |
| Mullet | 250 | 5 | 9,750 | 196 | 6,680 | 267 | | |
| Pigfish | | | 1,140 | 34 | | | | |
| Pike or pickerel | 175 | 9 | | | | | | |
| Pompano | | | 2,240 | 134 | | | 2,290 | 114 |
| Scup | | | 65,470 | 1,331 | 224,660 | 9,086 | | |
| Sea bass | | | 28,741 | 540 | 4,980 | 199 | | |
| Sbad | 28,542 | 3,042 | 71,160 | 7,203 | 27,760 | 2,776 | 1,510,240 | 146,999 |
| Sharks | | | 763 | 23 | | | | |
| Spanish mackerel | | | 24,890 | 1,990 | 13,040 | 573 | | |
| Spot | | | 87,080 | 2,319 | 29,450 | 884 | | |
| Squeteagues or "sea trout": | | | | | | | | |
| Gray | | | 113,321 | 2,426 | 3,621,010 | 80,440 | 371,460 | 8,961 |
| Spotted | | | 7,900 | 395 | 16,000 | 1,060 | 1,200 | 60 |
| Striped bass | 3,000 | 300 | 3,670 | 367 | 900 | 108 | 66,730 | 6,673 |
| Sturgeon | | | 1,453 | 228 | | | | |
| Swellfish | | | 259 | 13 | | | | |
| White perch | 2,740 | 118 | | | 4,914 | 147 | | |
| Yellow perch | 1,400 | 70 | | | | | | |
| Crabs: | | | | | | | | |
| Hard | | | 1,452,000 | 11,650 | 816,000 | 10,375 | 3,681,355 | 29,247 |
| Soft | | | | | 150,000 | 12,500 | 419,208 | 27,947 |
| Shrimp | | | 125 | 7 | | | | |
| Clams, hard, public | | | 16,000 | 4,000 | 139,636 | 34,422 | | |
| Oysters: | | | | | | | | |
| Market, public, spring | | | | | 1,200 | 100 | 3,900 | 200 |
| Market, public, fall | | | | | 10,476 | 873 | 39,700 | 2,430 |
| Market, private, spring | | | 639,164 | 49,228 | 262,892 | 20,241 | 139,790 | 9,668 |
| Market, private, fall | | | 1,103,907 | 78,850 | 142,240 | 11,854 | 240,800 | 16,280 |
| Squid | | | 257 | 6 | 38,940 | 779 | | |
| Turtles, snapper | 1,500 | 60 | | | | | | |
| Total | 68,367 | 4,362 | 4,538,080 | 175,144 | 6,827,160 | 214,928 | 91,912,972 | 562,899 |

| Species | Prince George | | Princess Anne | | Prince William | | Richmond | | Spotsylvania | |
|-----------------------------|----------------|--------------|------------------|---------------|----------------|---------------|----------------|---------------|--------------|------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives | 6,050 | \$80 | 21,000 | \$105 | 15,470 | \$232 | 46,750 | \$907 | | |
| Bluefish | | | 7,100 | 355 | | | | | | |
| Butterfish | | | 135,380 | 3,385 | | | 1,400 | 43 | | |
| Carp | 38,417 | 1,569 | 125,600 | 6,280 | 20,800 | 1,042 | 15,550 | 498 | 600 | \$15 |
| Catfish and bullheads | 50,795 | 1,759 | 2,800 | 84 | 43,300 | 1,299 | 68,100 | 2,889 | 1,000 | 40 |
| Croaker | | | 169,000 | 1,590 | | | 20,500 | 485 | | |
| Drum, red or redfish | | | 9,930 | 365 | | | | | | |
| Eels | 1,548 | 72 | 52,400 | 1,572 | | | 700 | 35 | | |
| Flounders | | | 3,400 | 160 | | | | | | |
| Glizzard shad | | | | | | | 23,500 | 585 | 1,000 | 30 |
| Harvestfish | | | 7,800 | 195 | | | | | | |
| Hickory shad | 300 | 8 | | | | | 300 | 15 | 100 | 5 |
| Mackerel | | | 3,420 | 171 | | | | | | |
| Mullet | 275 | 15 | | | | | 75 | 3 | | |
| Sbad | 8,530 | 880 | 17,800 | 1,798 | 126,420 | 8,393 | 23,050 | 3,035 | 300 | 15 |
| Spanish mackerel | | | 7,300 | 219 | | | | | | |
| Spot | | | 34,900 | 906 | | | | | | |
| Squeteagues or "sea trout": | | | | | | | | | | |
| Gray | | | 258,600 | 3,233 | | | 4,600 | 276 | | |
| Spotted | | | 24,600 | 1,230 | | | | | | |
| Striped bass | 860 | 87 | | | 23,400 | 2,340 | 8,500 | 1,250 | 100 | 15 |
| White perch | 900 | 35 | 8,300 | 415 | 17,340 | 566 | 15,850 | 793 | 500 | 25 |
| Yellow perch | 400 | 16 | | | 13,800 | 690 | 2,050 | 103 | 500 | 25 |
| Crabs, hard | | | 813,100 | 7,400 | 13,200 | 124 | 51,480 | 936 | | |
| Clams, hard, public | | | 2,632 | 659 | | | | | | |
| Oysters: | | | | | | | | | | |
| Market, public, spring | | | | | | | 115,577 | 5,779 | | |
| Market, public, fall | | | | | | | 144,227 | 7,211 | | |
| Market, private, spring | | | 10,360 | 2,220 | | | 74,725 | 4,261 | | |
| Market, private, fall | | | 14,140 | 3,030 | | | 53,207 | 3,040 | | |
| Total | 108,075 | 4,521 | 1,710,562 | 35,362 | 273,736 | 14,686 | 670,141 | 32,144 | 4,000 | 170 |

Fisheries of Virginia, 1933—Continued

CATCH: BY COUNTIES—Continued

| Species | Stafford | | Surry | | Warwick | | Westmoreland | | York | |
|----------------------------|----------------|---------------|---------------|--------------|------------------|---------------|------------------|---------------|------------------|----------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives | | | 7,000 | \$58 | 8,070 | \$73 | 2,846,200 | \$9,204 | 16,700 | \$84 |
| Bluefish | | | | | 731 | 58 | | | 40,967 | 1,310 |
| Butterfish | | | | | 13,340 | 360 | | | 11,697 | 330 |
| Carp | 28,160 | \$1,126 | 21,900 | 657 | | | 8,640 | 432 | | |
| Catfish and bullheads | 94,900 | 2,850 | 19,520 | 585 | | | 29,925 | 998 | 6,000 | 120 |
| Cod | | | | | 36 | 1 | | | 52 | 1 |
| Croaker | | | 4,010 | 60 | 455,624 | 7,734 | 102,870 | 1,286 | 652,021 | 9,523 |
| Drum, red or redfish | | | | | 200 | 4 | | | 290 | 9 |
| Eels | | | 820 | 16 | | | | | | |
| Flounders | | | | | 81,237 | 3,184 | | | 216,172 | 8,963 |
| Gizzard shad | | | 8,800 | 112 | | | | | | |
| Hake | | | | | 245 | 5 | | | 1,202 | 28 |
| Hickory shad | | | | | | | 15,330 | 307 | | |
| King whiting or "kingfish" | | | | | 5,127 | 141 | | | 7,129 | 282 |
| Mackerel | | | | | 10 | 1 | | | | |
| Mullet | | | | | 900 | 32 | | | 5,290 | 159 |
| Pickfish | | | | | 987 | 12 | | | 49,695 | 250 |
| Scup | | | | | 265,433 | 5,271 | | | 239,657 | 4,750 |
| Sea bass | | | | | 17,015 | 452 | | | 41,347 | 1,184 |
| Shad | 163,644 | 10,705 | 7,710 | 771 | 5,170 | 517 | 178,080 | 14,334 | 42,550 | 4,242 |
| Sharks | | | | | 396 | 11 | | | 200 | 6 |
| Spanish mackerel | | | | | | | | | 420 | 25 |
| Spot | | | | | 9,229 | 202 | | | 133,602 | 2,781 |
| Squeteagues or "sea trout" | | | | | | | | | | |
| Gray | | | | | 108,226 | 3,141 | 211,920 | 5,298 | 178,150 | 5,253 |
| Spotted | | | | | 8,420 | 842 | | | 68,940 | 3,447 |
| Striped bass | 20,060 | 2,342 | 7,810 | 781 | 263 | 27 | 93,563 | 11,227 | 4,638 | 464 |
| Sturgeon | | | | | | | | | 28 | 3 |
| Swellfish | | | | | 157 | 2 | | | 315 | 15 |
| Tautog | | | | | | | | | | |
| White perch | 35,855 | 1,434 | 6,980 | 216 | 3,800 | 115 | 21,495 | 860 | 600 | 30 |
| Yellow perch | 22,450 | 1,127 | | | | | | | 5,900,000 | 93,135 |
| Crabs, hard | 92,400 | 840 | | | | | 1,038,400 | 13,464 | 74 | 4 |
| Shrimp | | | | | | | | | 70,840 | 17,710 |
| Clams, hard, public | | | | | | | | | | |
| Oysters: | | | | | | | | | | |
| Market, public, spring | | | | | 69,780 | 4,652 | 24,700 | 1,270 | 95,280 | 5,290 |
| Market, public, fall | | | | | 110,520 | 7,368 | 7,800 | 400 | 111,360 | 5,474 |
| Market, private, spring | | | | | 16,752 | 838 | 149,940 | 8,568 | 250,800 | 17,260 |
| Market, private, fall | | | | | 7,200 | 360 | 236,470 | 14,084 | 324,400 | 22,500 |
| Squid | | | | | 370 | 10 | | | 192 | 5 |
| Total | 457,559 | 20,424 | 84,550 | 3,256 | 1,188,838 | 35,413 | 4,965,333 | 81,732 | 8,478,508 | 204,646 |

SEED OYSTER FISHERY: BY GEAR

| Item | Tongs | | Rakes | | By hand | | Total, exclusive of duplication | |
|--------------------------------|------------------|----------------|---------------|---------------|---------------|---------------|---------------------------------|----------------|
| | Number | Value | Number | Value | Number | Value | Number | Value |
| OPERATING UNITS | | | | | | | | |
| Fishermen, on boats and shore: | Number | | Number | | Number | | Number | |
| Regular | 1,562 | | 130 | | 178 | | 1,660 | |
| Casual | 67 | | | | | | 67 | |
| Total | 1,629 | | 130 | | 178 | | 1,717 | |
| Boats: | Number | | Number | | Number | | Number | |
| Motor | 802 | | | | 2 | | 804 | |
| Other | 226 | | 130 | | 80 | | 286 | |
| Apparatus: Number | 1,474 | | 130 | | | | | |
| CATCH | | | | | | | | |
| Oysters: | Bushels | Value | Bushels | Value | Bushels | Value | Bushels | Value |
| Seed, public, spring | 559,920 | \$57,940 | | | 14,700 | \$2,940 | 574,620 | \$60,880 |
| Seed, public, fall | 781,140 | 79,114 | | | 34,100 | 6,820 | 815,240 | 85,934 |
| Seed, private, spring | 7,300 | 1,760 | 21,000 | \$4,200 | 7,300 | 1,760 | 36,600 | 7,720 |
| Seed, private, fall | 12,400 | 2,480 | 41,400 | 8,280 | 3,000 | 600 | 56,800 | 11,360 |
| Total | 1,360,760 | 141,204 | 62,400 | 12,480 | 59,100 | 12,120 | 1,482,260 | 165,894 |

U. S. BUREAU OF FISHERIES

Fisheries of Virginia, 1933—Continued

SEED OYSTER FISHERY: BY COUNTIES

| Item | Accomac | | Elizabeth City | | Gloucester | | Isle of Wight | |
|--|---------|---------|----------------|---------|------------|---------|---------------|----------|
| OPERATING UNITS | | | | | | | | |
| Fishermen, on boats and shore: Regular | Number | | Number | | Number | | Number | |
| Boats: | 140 | | 43 | | 182 | | 512 | |
| Motor | 50 | | 24 | | 128 | | 222 | |
| Other | 20 | | 19 | | 54 | | 26 | |
| Apparatus: Tongs | 92 | | 33 | | 144 | | 466 | |
| CATCH | | | | | | | | |
| Oysters: | Bushels | Value | Bushels | Value | Bushels | Value | Bushels | Value |
| Seed, public, spring | 25,400 | \$5,080 | 12,100 | \$1,210 | 10,500 | \$1,050 | 256,600 | \$25,660 |
| Seed, public, fall | 40,600 | 8,120 | 21,300 | 2,130 | 31,700 | 3,170 | 326,400 | 32,640 |
| Total | 66,000 | 13,200 | 33,400 | 3,340 | 42,200 | 4,220 | 583,000 | 58,300 |

| Item | James City | | Mathew | | Middlesex | | Nansemond | | New Kent | |
|--------------------------------|------------|-------|---------|-------|-----------|---------|-----------|---------|----------|-------|
| OPERATING UNITS | | | | | | | | | | |
| Fishermen, on boats and shore: | Number | | Number | | Number | | Number | | Number | |
| Regular | 9 | | 43 | | 54 | | 108 | | 4 | |
| Casual | 5 | | 32 | | 54 | | 51 | | 2 | |
| Boats: Motor | 7 | | 48 | | 54 | | 87 | | 4 | |
| Apparatus: Tongs | 7 | | 48 | | 54 | | 87 | | 4 | |
| CATCH | | | | | | | | | | |
| Oysters: | Bushels | Value | Bushels | Value | Bushels | Value | Bushels | Value | Bushels | Value |
| Seed, public, spring | 3,700 | \$370 | 6,400 | \$640 | 5,480 | \$1,096 | 34,400 | \$3,440 | 900 | \$90 |
| Seed, public, fall | 7,800 | 780 | 14,600 | 1,460 | 5,480 | 1,096 | 31,800 | 3,180 | 2,900 | 290 |
| Total | 11,500 | 1,150 | 21,000 | 2,100 | 5,480 | 1,096 | 66,200 | 6,620 | 3,800 | 380 |

| Item | Norfolk | | Northampton | | Warwick | | York | |
|--------------------------------|---------|---------|-------------|--------|---------|----------|---------|---------|
| OPERATING UNITS | | | | | | | | |
| Fishermen, on boats and shore: | Number | | Number | | Number | | Number | |
| Regular | 51 | | 180 | | 215 | | 108 | |
| Casual | 28 | | 180 | | 35 | | 108 | |
| Total | 79 | | 180 | | 250 | | 108 | |
| Boats: | 19 | | 180 | | 143 | | 74 | |
| Motor | 60 | | 180 | | 143 | | 74 | |
| Other | 19 | | 180 | | 143 | | 74 | |
| Apparatus: | 79 | | 140 | | 219 | | 101 | |
| Tongs | 79 | | 140 | | 219 | | 101 | |
| Rakes | 19 | | 180 | | 143 | | 74 | |
| CATCH | | | | | | | | |
| Oysters: | Bushels | Value | Bushels | Value | Bushels | Value | Bushels | Value |
| Seed, public, spring | 21,740 | \$2,174 | 3,300 | \$660 | 166,800 | \$16,680 | 27,300 | \$2,730 |
| Seed, public, fall | 65,340 | 6,534 | 3,500 | 700 | 211,900 | 21,190 | 57,400 | 5,740 |
| Seed, private, spring | ----- | ----- | 35,000 | 7,720 | ----- | ----- | ----- | ----- |
| Seed, private, fall | ----- | ----- | 56,800 | 11,360 | ----- | ----- | ----- | ----- |
| Total | 87,080 | 8,708 | 99,200 | 20,440 | 378,700 | 37,870 | 84,700 | 8,470 |

NOTE.—Of the total number of persons fishing for seed oysters, 1,636 are duplicated among those fishing for market oysters or other species. Similarly, the following craft and gear are duplicated: 744 motorboats, 267 other boats, and 1,161 tongs.

WINTER TRAWL FISHERY OFF NEW JERSEY, MARYLAND, VIRGINIA, AND NORTH CAROLINA

The catch in the winter trawl fishery off the coasts of southern New Jersey, Maryland, Virginia, and North Carolina (excluding the shrimp fishery) in 1933 amounted to 18,789,969 pounds of fishery products, valued at \$506,513 to the fishermen. Craft whose home port was in Massachusetts accounted for 57 percent of the total volume and 56 percent of the value, while Virginia craft accounted for 29 percent of the volume and 26 percent of the value. The principal species contributing to the catch were croaker, sea bass, flounders, and scup.

Statistics of the winter trawl fishery are included also in the catch data for the various States published elsewhere in this report.

Winter trawl fishery off New Jersey, Maryland, Virginia, and North Carolina, 1933

 CATCH: BY STATES¹

| Species | Massachusetts | | Connecticut | | New York | | New Jersey | |
|---------------------------------|---------------|---------|-------------|-------|-----------|--------|------------|--------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Bluefish..... | 4,974 | \$360 | 113 | \$9 | 292 | \$19 | 3,434 | \$199 |
| Butterfish..... | 70,871 | 2,637 | 87 | 3 | 17,802 | 634 | 22,800 | 695 |
| Cod..... | 8,398 | 291 | | | 160 | 2 | 12,090 | 233 |
| Croaker..... | 2,490,145 | 35,696 | 2,738 | 46 | 76,305 | 1,357 | 367,623 | 7,125 |
| Cunners..... | 152 | 2 | | | | | | |
| Drum: | | | | | | | | |
| Black..... | 32 | 1 | 100 | 1 | 447 | 9 | | |
| Red..... | 1,755 | 32 | | | | | 555 | 9 |
| Eels, conger..... | 38,020 | 703 | 190 | 4 | 3,260 | 57 | 4,704 | 126 |
| Flounders..... | 2,147,253 | 82,289 | 87,969 | 3,827 | 523,966 | 22,355 | 628,634 | 26,622 |
| Grayfish..... | 2,932 | 36 | | | | | | |
| Haddock..... | 299 | 4 | | | | | | |
| Hake..... | 89,935 | 1,372 | 275 | 5 | 14,526 | 203 | 9,204 | 149 |
| King whiting or "kingfish"..... | 25,296 | 645 | 451 | 12 | 2,611 | 90 | 6,942 | 274 |
| Mackerel..... | 395 | 6 | | | 150 | 6 | 37 | 1 |
| Mullet..... | | | | | | | 267 | 5 |
| Pigfish..... | 3,832 | 73 | | | | | 151 | 4 |
| Pollock..... | 660 | 10 | | | | | | |
| Scup or porgy..... | 1,645,178 | 37,422 | 16,587 | 536 | 249,466 | 5,719 | 248,166 | 6,209 |
| Sea bass..... | 3,701,867 | 108,239 | 2,270 | 142 | 114,522 | 4,551 | 102,439 | 3,943 |
| Shad..... | 28 | 2 | | | | | | |
| Sheepshead..... | 9 | 1 | | | | | 9 | 1 |
| Skates..... | 108 | 2 | | | 725 | 7 | | |
| Spot..... | 31,758 | 346 | 50 | 1 | 816 | 23 | 1,126 | 22 |
| Squeteagues or "sea trout": | | | | | | | | |
| Gray..... | 286,547 | 8,543 | 320 | 17 | 13,741 | 600 | 32,715 | 1,065 |
| Spotted..... | | | | | | | 30 | 2 |
| Striped bass..... | 17 | 1 | | | 11 | 1 | | |
| Sturgeon..... | 2,960 | 371 | 44 | 5 | 365 | 33 | 641 | 47 |
| Tautog..... | 688 | 20 | | | 60 | 1 | 149 | 3 |
| Whiting..... | 590 | 7 | | | 900 | 9 | | |
| Crabs, hard..... | | | | | | | 60 | 1 |
| Lobsters..... | 1,588 | 130 | 163 | 20 | 1,421 | 127 | 1,105 | 121 |
| Shrimp..... | 40,900 | 2,045 | | | 12 | 1 | 841 | 128 |
| Scallops, sea..... | 158 | 35 | | | 297 | 47 | 170 | 42 |
| Squid..... | 22,170 | 489 | 480 | 13 | 2,446 | 77 | 3,370 | 85 |
| Total..... | 10,619,603 | 281,810 | 111,837 | 4,641 | 1,024,291 | 36,928 | 1,447,252 | 47,109 |

¹ The data as shown for the various States represent the catch of craft whose principal fishing ports are in such States.

Winter trawl fishery off New Jersey, Maryland, Virginia, and North Carolina,
1933—Continued

CATCH: BY STATES—Continued

| Species | Virginia | | Georgia | | Florida | | Total | |
|-----------------------------|-----------|---------|---------|-------|---------|-------|------------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Alewives | 217 | \$2 | | | | | 217 | \$2 |
| Bluefish | 28,069 | 2,205 | | | 153 | \$10 | 37,025 | 2,802 |
| Butterfish | 52,601 | 1,536 | | | 393 | 12 | 164,554 | 5,517 |
| Cod | 315 | 6 | | | | | 20,953 | |
| Croaker | 2,607,106 | 45,318 | | | 73,267 | 1,029 | 5,617,154 | 90,571 |
| Cunners | | | | | | | 152 | 2 |
| Drum: | | | | | | | | |
| Black | | | | | | | 579 | 11 |
| Red | 4,418 | 72 | 20 | \$1 | 180 | 6 | 6,028 | 120 |
| Eels, conger | 1,175 | 16 | | | | | 47,349 | 906 |
| Flounders | 676,579 | 27,014 | | | 79,137 | 2,852 | 4,143,538 | 164,959 |
| Grayfish | | | | | | | 2,932 | 36 |
| Haddock | 47 | 1 | | | | | 346 | 5 |
| Hake | 19,060 | 311 | | | 15 | 1 | 133,015 | 2,041 |
| King whiting or "kingfish" | 34,747 | 1,188 | | | 137 | 4 | 70,184 | 2,213 |
| Mackerel | 120 | 6 | | | | | 702 | 19 |
| Mullet | | | | | | | 267 | 8 |
| Pigfish | 3,300 | 65 | | | 45 | 1 | 7,328 | 143 |
| Pollock | | | | | | | 660 | 10 |
| Scup or porgy | 1,148,135 | 22,756 | | | 33,768 | 322 | 3,341,290 | 72,964 |
| Sea bass | 303,539 | 8,028 | | | 11,430 | 505 | 4,236,067 | 126,008 |
| Shad | | | | | | | 28 | 2 |
| Sharks | 10,225 | 159 | 955 | 28 | | | 11,180 | 187 |
| Sheepshead | | | | | | | 18 | 2 |
| Skates | | | | | | | 833 | 9 |
| Spot | 26,552 | 492 | | | 536 | 10 | 60,838 | 894 |
| Squeteagues or "sea trout": | | | | | | | | |
| Gray | 361,235 | 13,094 | | | 17,076 | 404 | 711,434 | 23,723 |
| Spotted | 595 | 31 | | | | | 625 | 33 |
| Striped bass | | | | | | | 28 | 2 |
| Sturgeon | 3,195 | 494 | | | 187 | 16 | 7,382 | 958 |
| Shellfish | 1,065 | 53 | 39 | 2 | | | 1,104 | 55 |
| Tautog | 488 | 9 | | | 83 | 1 | 1,468 | 34 |
| White perch | 102 | 3 | | | | | 102 | 3 |
| Whiting | | | | | | | 1,490 | 16 |
| Crabs, hard | | | | | | | 60 | 1 |
| Lobsters | 131 | 2 | | | | | 4,408 | 400 |
| Shrimp | 371 | 20 | | | | | 42,124 | 2,189 |
| Scallops, sea | 72,450 | 8,050 | | | | | 73,073 | 8,174 |
| Squid | 13,353 | 288 | 65 | 1 | 420 | 11 | 42,304 | 964 |
| Total | 5,369,180 | 131,809 | 1,079 | 32 | 216,827 | 5,184 | 18,789,969 | 506,513 |

SHAD AND ALEWIFE FISHERIES OF THE POTOMAC RIVER

The catch of shad in the Potomac River in 1933 amounted to 611,425 in number, 1,837,623 pounds in weight, and their total value to the fishermen was \$149,114. The catch of alewives for the same season amounted to 17,238,850 in number, with a total weight of 6,895,540 pounds, and a value to the fishermen of \$23,845. These figures show a decrease of 19 percent in the weight and 14 percent in the value of shad as compared with 1932, and an increase of 1 percent in the weight but a decrease of 1 percent in the value of alewives.

About 57 percent of the shad, in weight, were taken with pound nets, and 43 percent, with gill nets. Less than one-half of 1 percent were taken with haul seines. More than 99½ percent of alewives were taken with pound nets, only small quantities being taken with gill nets, haul seines, and fyke nets.

Statistics of the catch of shad and alewives in the Potomac River are included also in the catch data for Maryland and Virginia which are published elsewhere in this report.

Shad and alewife fisheries of the Potomac River, 1933

| Item | Maryland | | | Virginia | | | Total | | |
|--------------------------------------|------------------|----------------|---------------|-------------------|------------------|----------------|-------------------|------------------|----------------|
| | Number | Pounds | Value | Number | Pounds | Value | Number | Pounds | Value |
| Fishermen on boats and shore: | | | | | | | | | |
| Regular..... | 49 | | | 322 | | | 371 | | |
| Casual..... | 119 | | | 161 | | | 280 | | |
| Total..... | 168 | | | 483 | | | 651 | | |
| Boats: | | | | | | | | | |
| Motor..... | 58 | | | 183 | | | 241 | | |
| Other..... | 50 | | | 97 | | | 147 | | |
| Apparatus: | | | | | | | | | |
| Pound nets..... | 67 | | | 300 | | | 367 | | |
| Gill nets..... | 1,251 | | | 885 | | | 2,136 | | |
| Square yards..... | 236,557 | | | 230,790 | | | 467,347 | | |
| Haul seines..... | 1 | | | | | | 1 | | |
| Length, yards..... | 200 | | | | | | 200 | | |
| Fyke nets..... | | | | 3 | | | 3 | | |
| Shad caught: | | | | | | | | | |
| With pound nets.. | 17,714 | 60,833 | \$6,237 | 342,971 | 990,532 | \$90,705 | 360,685 | 1,051,365 | \$96,942 |
| With gill nets..... | 33,260 | 114,932 | 9,222 | 217,417 | 671,126 | 42,927 | 250,677 | 786,058 | 52,149 |
| With haul seines.. | 63 | 200 | 23 | | | | 63 | 200 | 23 |
| Total..... | 51,037 | 175,965 | 15,482 | 560,388 | 1,661,658 | 133,632 | 611,425 | 1,837,623 | 149,114 |
| Alewives caught: | | | | | | | | | |
| With pound nets.. | 1,200,060 | 483,624 | 2,277 | 15,961,865 | 6,384,746 | 21,207 | 17,170,925 | 6,868,370 | 23,484 |
| With gill nets..... | 10,500 | 4,200 | 60 | 48,675 | 19,470 | 282 | 59,175 | 23,670 | 342 |
| With haul seines.. | 1,250 | 500 | 10 | | | | 1,250 | 500 | 10 |
| With fyke nets..... | | | | 7,500 | 3,000 | 9 | 7,500 | 3,000 | 9 |
| Total..... | 1,220,810 | 488,324 | 2,347 | 16,018,040 | 6,407,216 | 21,498 | 17,238,850 | 6,895,540 | 23,846 |

TRADE IN FISHERY PRODUCTS IN WASHINGTON, D. C.¹

The municipal fish wharf and market in Washington, D. C., is located in the southwestern part of the city on an arm of the Potomac River. At the present time 17 firms have stalls in the market, 2 firms are in private buildings across the street, and 4 firms have stalls in the new Center Market. Altogether the 23 above firms employed 111 persons who received \$79,186 in salaries and wages during 1933. Of the total employees 95 were regularly employed. These firms conduct a wholesale and retail business, chiefly wholesale, however.

During the year 1933 the receipts of fresh and frozen fishery products as received at the municipal fish wharf amounted to 9,572,135 pounds. This is a decrease of 16 percent as compared with the year 1932 but an increase of 2 percent as compared with the 5-year average.

During 1933 three firms in Washington, D. C., smoked fishery products which amounted to 257,825 pounds, valued at \$27,806. Of this amount 230,000 pounds, valued at \$20,880, consisted of herring smoked as bloaters; 24,400 pounds, valued at \$6,292, were whitefish; while the remainder or 3,425 pounds, valued at \$634, consisted of alewives or "river herring", eels, and haddock.

There were three firms which shucked oysters mostly for hotel and restaurant trade. Their production amounted to 6,100 gallons, valued at \$9,600. In addition to this quantity there were 42,245 gallons of oysters, valued at \$78,160, shucked mostly by fishermen for retail trade. Virtually all of the fishery products were consumed in the city.

¹ Statistics of fishery products handled at the municipal wharf, Washington, D. C., are reported to the Bureau by agents of the Health Department, District of Columbia.

Fishery products received at municipal fish wharf and market, Washington, D. C., 1953

| Species | January | February | March | April | May | June | July | August | September | October | November | December | Total |
|---------------------------------|---------|----------|---------|---------|---------|---------|---------|---------|-----------|---------|----------|----------|-----------|
| | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds | Pounds |
| Alewives (river herring)..... | 6,700 | 18,000 | 73,400 | 318,800 | 142,700 | | | | | | | | 559,800 |
| Bluefish..... | 1,500 | 4,200 | 6,200 | 6,300 | 43,800 | 30,600 | 15,900 | 21,600 | 26,200 | 34,100 | 12,800 | 5,600 | 208,800 |
| Butterfish..... | 2,700 | 2,600 | 2,300 | 6,400 | 31,900 | 75,500 | 35,200 | 55,300 | 26,200 | 36,900 | 15,400 | 2,900 | 293,300 |
| Carp..... | 11,500 | 6,300 | 10,700 | 21,600 | 15,600 | 5,800 | 4,100 | 1,300 | 6,600 | 5,100 | 5,500 | 6,400 | 100,500 |
| Catfish..... | 2,300 | 8,400 | 21,700 | 26,600 | 8,500 | 5,100 | 2,300 | 300 | 4,800 | 12,700 | 11,000 | 2,600 | 106,300 |
| Cod..... | 1,000 | 1,200 | 600 | 1,100 | 700 | 200 | | | 500 | 800 | 600 | 800 | 7,900 |
| Croaker..... | 103,600 | 101,400 | 68,500 | 280,000 | 269,100 | 233,000 | 284,000 | 306,800 | 116,600 | 149,000 | 181,400 | 172,200 | 2,255,600 |
| Drum, red or redfish..... | 9,000 | | 1,400 | 6,000 | 680 | 110 | 500 | | 2,800 | 4,400 | 4,900 | 2,900 | 32,690 |
| Eels..... | 600 | | 600 | 3,800 | 400 | 100 | | | 1,100 | 1,400 | 1,800 | 200 | 10,000 |
| Flounders..... | 53,500 | 44,400 | 42,100 | 26,800 | 31,700 | 36,000 | 19,400 | 20,400 | 8,500 | 20,100 | 33,500 | 24,300 | 360,700 |
| Gizzard shad..... | 24,500 | 7,500 | 22,200 | 5,100 | | | | | 200 | 2,300 | 1,500 | 300 | 63,600 |
| Haddock..... | 28,525 | 33,310 | 27,490 | 38,105 | 32,970 | 25,120 | 16,230 | 30,560 | 21,520 | 21,330 | 31,720 | 20,180 | 327,060 |
| Hake..... | 1,700 | 1,200 | | | | | | | | | 400 | 200 | 3,500 |
| Halibut..... | 6,900 | 5,900 | 9,600 | 6,800 | 5,400 | 3,300 | 4,500 | 4,400 | 5,200 | 7,200 | 5,600 | 4,000 | 68,800 |
| Hickory shad or "jacks"..... | 5,300 | 2,300 | 2,300 | 1,300 | 200 | | | | | | | | 11,400 |
| Hogfish..... | | | | | | | | 200 | 200 | 3,300 | 300 | | 4,000 |
| King sh or "king mackerel"..... | 1,400 | 1,400 | 600 | 17,800 | | 200 | | 600 | | | 1,900 | 1,700 | 25,600 |
| Mackerel..... | 33,100 | 28,000 | 18,400 | 16,600 | 16,700 | 25,200 | 12,600 | 22,200 | 18,400 | 20,600 | 16,700 | 21,400 | 249,900 |
| Menhaden..... | | | | | | | | | | | 200 | | 200 |
| Mullet..... | 5,400 | 2,100 | 700 | 3,000 | | | | 800 | 5,300 | 7,600 | 8,700 | 4,000 | 37,600 |
| Perch..... | 4,700 | 6,800 | 24,700 | 30,900 | 4,400 | 800 | 900 | 900 | 2,100 | 6,300 | 5,900 | 5,800 | 94,200 |
| Pike or pickeral..... | 800 | | | 1,500 | 600 | | | | | | 700 | 600 | 4,200 |
| Pollock..... | 200 | | | | | 100 | 400 | 200 | | | 400 | 9,200 | 10,500 |
| Pompano..... | 200 | 100 | 800 | 200 | 600 | 200 | 700 | 1,300 | 300 | 200 | 300 | 100 | 5,000 |
| Salmon..... | 4,800 | 2,500 | 3,000 | 1,200 | 1,200 | 1,200 | 3,000 | 3,200 | 5,000 | 8,300 | 7,900 | 3,100 | 44,400 |
| Scup or porgy..... | 8,200 | 23,000 | 20,400 | 8,800 | 11,500 | 10,700 | | 1,600 | 2,800 | 2,300 | 1,700 | 200 | 91,200 |
| Sea bass..... | 70,400 | 82,500 | 119,600 | 30,800 | 7,500 | 19,000 | | | 2,500 | 4,300 | 4,700 | 7,500 | 363,900 |
| Shad..... | 22,200 | 26,165 | 70,700 | 194,200 | 244,100 | 15,600 | | 7,700 | | | 900 | | 673,865 |
| Sheepshead..... | | | | | | | | | | | 100 | | 100 |
| Smelt..... | 2,580 | 3,540 | 2,220 | 1,275 | | | | | 15 | | 180 | 1,020 | 10,830 |
| Snapper, red..... | 1,000 | 300 | 400 | 650 | | 450 | 300 | 300 | 300 | 200 | 300 | 600 | 5,000 |
| Spot..... | 1,100 | 200 | | | 6,200 | 16,300 | 35,200 | 26,200 | 27,900 | 42,000 | 14,400 | 500 | 170,000 |
| Squeteagus or "sea trout"..... | 95,900 | 66,000 | 100,100 | 83,800 | 287,300 | 232,700 | 108,300 | 178,600 | 228,500 | 279,800 | 128,800 | 106,100 | 1,893,900 |
| Striped bass..... | 8,500 | 5,900 | 27,100 | 33,220 | 11,200 | 2,600 | 900 | 1,100 | 4,900 | 23,700 | 23,300 | 7,100 | 149,520 |
| Sturgeon..... | 200 | | | 100 | 500 | | | | | 50 | 200 | 295 | 1,345 |
| Swordfish..... | | | | | | 100 | 350 | 600 | 507 | 585 | 452 | 125 | 2,719 |
| Tilefish..... | 200 | | | 100 | | | | | | | 200 | 100 | 600 |
| Whitefish..... | | 300 | | 200 | | | | | | | | | 500 |
| Whiting..... | 600 | | | | | | | | | | 800 | 500 | 1,900 |
| Yellowtail..... | 2,000 | | | 200 | | | | | | 800 | | | 2,800 |

| | | | | | | | | | | | | | | |
|----------------------------------|---------|---------|---------|-----------|-----------|---------|---------|---------|---------|---------|---------|---------|-----------|--|
| Crabs: | | | | | | | | | | | | | | |
| Hard..... | | | 160 | 3,375 | 24,750 | 58,500 | 62,025 | 33,925 | 32,175 | 4,500 | 450 | | 234,850 | |
| Soft..... | | | 2,250 | 4,455 | 19,800 | 31,500 | 10,485 | 13,135 | 22,065 | 12,015 | 990 | | 122,805 | |
| Oyster..... | | 25 | | | | | | | | | | | 25 | |
| Meat..... | 4,240 | 3,180 | 5,485 | 13,355 | 29,640 | 29,205 | 22,950 | 23,770 | 18,925 | 16,605 | 12,330 | 3,805 | 190,490 | |
| Sea crawfish or "spiny lobster": | | | | | | | | | | | | | | |
| Alive..... | 100 | 50 | | 50 | | | | | | 75 | 200 | 50 | 525 | |
| Meat..... | | 60 | 25 | | | | | | | | | | 85 | |
| Lobsters: | | | | | | | | | | | | | | |
| Alive..... | 1,060 | 800 | 720 | 1,240 | 2,920 | 1,410 | 440 | 750 | 750 | 950 | 910 | 600 | 12,570 | |
| Meat..... | 105 | 145 | 120 | 50 | 145 | 80 | 50 | 155 | 80 | 190 | 190 | 220 | 1,530 | |
| Shrimp..... | 5,650 | 5,400 | 5,100 | 7,250 | 9,600 | 11,475 | 9,385 | 21,515 | 12,268 | 13,362 | 8,085 | 6,050 | 115,140 | |
| 8 mid..... | 1,400 | | 700 | | | | | | 100 | 100 | 300 | 100 | 2,700 | |
| Clams: | | | | | | | | | | | | | | |
| Hard..... | 6,112 | 6,080 | 8,960 | 5,664 | 8,032 | 7,776 | 5,376 | 5,888 | 6,016 | 8,256 | 5,088 | 4,512 | 177,760 | |
| Soft..... | | 16 | | | | | | | | | | | 16 | |
| Oysters: | | | | | | | | | | | | | | |
| In the shell (meat) | 50,645 | 31,794 | 27,622 | 9,807 | 574 | | | 70 | 8,498 | 30,611 | 35,140 | 23,408 | * 218,169 | |
| Opened (meat) | 66,132 | 64,286 | 55,099 | 18,682 | 656 | | | | 17,859 | 58,739 | 75,504 | 66,062 | * 423,019 | |
| Scallops..... | 1,072 | 1,552 | 1,160 | 1,952 | 2,128 | 3,672 | 920 | 2,394 | 2,904 | 5,160 | 2,416 | 1,016 | 26,336 | |
| Frcgs..... | | | | 183 | 18 | 100 | 13 | | 63 | 60 | 48 | 15 | 500 | |
| Terrapin..... | | | | | | | | | | | 82 | | 82 | |
| Turtles: | | | | | | | | | | | | | | |
| Sea..... | | | | | 180 | | | | | | | | 180 | |
| Snapper..... | 1,528 | 140 | | 128 | 28 | | | | | | | | 1,824 | |
| Total..... | 660,885 | 599,027 | 775,201 | 1,241,441 | 1,274,371 | 883,638 | 663,724 | 803,152 | 639,665 | 845,788 | 666,885 | 518,358 | 9,572,135 | |

¹ 9,720 bushels.

² 31,167 bushels.

³ 48,345 gallons.

NOTE.—Hard clams have been converted to pounds on the basis of 8 pounds of meats to the bushel, and oysters on the basis of 7 pounds of meats to the bushel, and 8¾ pound to the gallon.

FISHERIES OF THE SOUTH ATLANTIC AND GULF STATES

(South Atlantic, Area XXIV; Gulf, Area XXV) ¹

The most recent complete catch statistics for the fisheries of the South Atlantic and Gulf States are those collected for the year 1932 and the most recent complete statistics on fisheries wholesale and manufacturing industries in the same region are for 1931. The yield of the commercial fisheries in the marine areas of these States, comprised of North Carolina, South Carolina, Georgia, Florida, Alabama, Mississippi, Louisiana, and Texas, during 1932, amounted to 299,916,728 pounds, valued at \$6,428,385, to the fishermen, representing an increase of 4 percent in volume but a decrease of 20 percent in value as compared with the catch in 1931. Detailed statistics of the operating units and catch of these fisheries for 1932 appear in "Fishery Industries of the United States, 1933", Appendix I to the Report of the Commissioner of Fisheries for the fiscal year 1934, while data on wholesale and manufacturing industries for 1931 are published in "Fishery Industries of the United States, 1932", Appendix III to the Report of the Commissioner of Fisheries for the fiscal year 1933. A summary of these fisheries appears in the following tables.

Fisheries of the South Atlantic and Gulf States, 1932

OPERATING UNITS: BY STATES

| Item | North Carolina | South Carolina | Georgia | Florida | Alabama | Mississippi | Louisiana | Texas | Total |
|-------------------------------|----------------|----------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| Fishermen: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| On vessels..... | 758 | 17 | 86 | 638 | 139 | 474 | 154 | 143 | 2,409 |
| On boats and shore: | | | | | | | | | |
| Regular..... | 2,754 | 615 | 427 | 5,988 | 360 | 829 | 1,864 | 1,223 | 14,060 |
| Casual..... | 1,411 | 843 | 539 | 1,331 | 90 | 205 | 289 | 383 | 5,091 |
| Total..... | 4,923 | 1,475 | 1,052 | 7,957 | 589 | 1,508 | 2,307 | 1,749 | 21,560 |
| Vessels: | | | | | | | | | |
| Motor..... | 79 | 4 | 20 | 98 | 31 | 114 | 62 | 33 | 441 |
| Net tonnage..... | 1,234 | 59 | 245 | 2,467 | 299 | 1,507 | 447 | 388 | 6,646 |
| Sail..... | 53 | | | 1 | | 15 | 2 | | 71 |
| Net tonnage..... | 502 | | | 64 | | 237 | 38 | | 841 |
| Total vessels..... | 132 | 4 | 20 | 99 | 31 | 129 | 64 | 33 | 512 |
| Total net tonnage..... | 1,736 | 59 | 245 | 2,531 | 299 | 1,744 | 485 | 388 | 7,487 |
| Boats: | | | | | | | | | |
| Motor..... | 1,154 | 84 | 119 | 2,318 | 153 | 268 | 574 | 382 | 5,052 |
| Other..... | 1,584 | 773 | 523 | 2,945 | 151 | 407 | 996 | 418 | 7,797 |
| Accessory boats..... | 70 | | 4 | 11 | | | | | 85 |
| Apparatus: | | | | | | | | | |
| Purse seines: | | | | | | | | | |
| Menhaden..... | 33 | | 2 | 5 | | | | | 40 |
| Length, yards..... | 8,025 | | 600 | 1,450 | | | | | 10,075 |
| Other..... | 1 | | | 1 | | | | | 2 |
| Length, yards..... | 175 | | | 400 | | | | | 575 |
| Haul seines: | | | | | | | | | |
| Common..... | 454 | 20 | 11 | 110 | 5 | 3 | 102 | 54 | 759 |
| Length, yards..... | 66,326 | 2,780 | 1,105 | 34,300 | 2,900 | 800 | 13,434 | 9,315 | 130,960 |
| Long..... | 56 | | | 76 | | | | | 132 |
| Length, yards..... | 58,275 | | | 59,200 | | | | | 117,475 |
| Gill nets: | | | | | | | | | |
| Anchor..... | 1,661 | 324 | 45 | 12 | | | | | 2,042 |
| Square yards..... | 908,610 | 154,872 | 10,625 | 9,600 | | | | | 1,083,707 |

¹ These are the numbers given to these areas by the North American Council on Fishery Investigations. For a clearer understanding of the statistics published in this section, the reader is referred to the section in the latter part of this document entitled "Statistical survey procedure."

Fisheries of the South Atlantic and Gulf States, 1932—Continued

OPERATING UNITS: BY STATES—Continued

| Item | North Carolina | South Carolina | Georgia | Florida | Alabama | Mississippi | Louisiana | Texas | Total |
|-----------------------------------|----------------|----------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| Apparatus—Con. | | | | | | | | | |
| Gill nets.—Con. | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| Drift..... | 227 | 323 | 158 | 146 | ----- | ----- | ----- | ----- | 854 |
| Square yards..... | 399,265 | 284,554 | 111,863 | 139,606 | ----- | ----- | ----- | ----- | 935,288 |
| Runaround..... | 188 | 7 | 10 | 2,111 | ----- | ----- | ----- | ----- | 47 |
| Square yards..... | 83,500 | 1,450 | 3,170 | 2,110,492 | ----- | ----- | ----- | 13,165 | 2,211,777 |
| Stake..... | 5,271 | ----- | 40 | 5 | 18 | ----- | ----- | ----- | 267 |
| Square yards..... | 561,965 | ----- | 10,050 | 1,250 | 2,880 | ----- | ----- | 74,845 | 650,990 |
| Trammel nets..... | ----- | ----- | ----- | 182 | 53 | 39 | 23 | 61 | 358 |
| Square yards..... | ----- | ----- | ----- | 122,469 | 17,365 | 15,775 | 6,985 | 22,071 | 184,665 |
| Lines: | | | | | | | | | |
| Hand..... | 86 | 180 | 43 | 1,620 | 149 | 132 | 167 | 467 | 2,844 |
| Hooks..... | 166 | 550 | 46 | 2,573 | 260 | 142 | 172 | 594 | 4,503 |
| Troll..... | 45 | ----- | ----- | 1,190 | 2 | ----- | ----- | 8 | 1,245 |
| Hooks..... | 45 | ----- | ----- | 1,485 | 2 | ----- | ----- | 8 | 1,540 |
| Trot with baits or snoods..... | 156 | 6 | 31 | 13 | 15 | 36 | 318 | 25 | 600 |
| Trot with hooks..... | 119,000 | 4,500 | 9,390 | 2,700 | 2,336 | 8,895 | 60,025 | 3,175 | 210,021 |
| Hooks..... | 26 | ----- | 40 | 198 | 101 | ----- | 3 | 48 | 416 |
| Pound nets..... | 3,200 | ----- | 2,840 | 85,005 | 10,370 | ----- | 300 | 6,915 | 108,630 |
| Weirs..... | 1,722 | ----- | ----- | 13 | ----- | ----- | ----- | ----- | 1,735 |
| Wheels..... | 2 | ----- | ----- | ----- | ----- | ----- | ----- | ----- | 2 |
| Stop nets..... | ----- | ----- | ----- | 7 | ----- | ----- | ----- | ----- | 21 |
| Square yards..... | ----- | ----- | ----- | 11,475 | ----- | ----- | ----- | ----- | 11,475 |
| Fyke nets..... | 801 | ----- | ----- | 278 | 6 | ----- | ----- | ----- | 1,085 |
| Dip nets: | | | | | | | | | |
| Common..... | 204 | ----- | ----- | 50 | ----- | ----- | ----- | ----- | 254 |
| Drop..... | ----- | ----- | ----- | 54 | ----- | 130 | 1,520 | ----- | 1,704 |
| Cast nets..... | ----- | ----- | 10 | 16 | ----- | 60 | 18 | ----- | 104 |
| Otter trawls: | | | | | | | | | |
| Fish..... | 2 | ----- | ----- | 4 | ----- | ----- | ----- | ----- | 6 |
| Yards at mouth..... | 41 | ----- | ----- | 109 | ----- | ----- | ----- | ----- | 150 |
| Shrimp..... | 51 | 28 | 125 | 376 | 112 | 255 | 476 | 251 | 1,674 |
| Yards at mouth..... | 1,007 | 560 | 2,510 | 7,135 | 1,465 | 3,154 | 5,942 | 3,634 | 25,407 |
| Pots: | | | | | | | | | |
| Crab..... | ----- | ----- | 12 | 1,433 | ----- | ----- | ----- | ----- | 1,445 |
| Eel..... | 1,285 | ----- | ----- | 40 | ----- | ----- | ----- | ----- | 1,325 |
| Fish..... | 465 | ----- | 81 | 1,515 | ----- | ----- | ----- | ----- | 2,061 |
| Sea crawfish..... | ----- | ----- | ----- | 3,190 | ----- | ----- | ----- | ----- | 3,190 |
| Spears..... | 50 | 6 | ----- | 27 | 30 | 63 | ----- | 152 | 328 |
| Dredges: | | | | | | | | | |
| Clam..... | ----- | ----- | ----- | 1 | ----- | ----- | ----- | ----- | 1 |
| Oyster..... | 182 | ----- | ----- | 2 | ----- | 328 | 26 | 39 | 577 |
| Yards at mouth..... | 182 | ----- | ----- | 2 | ----- | 329 | 26 | 38 | 577 |
| Scallop..... | 64 | ----- | ----- | ----- | ----- | ----- | ----- | ----- | 64 |
| Yards at mouth..... | 64 | ----- | ----- | ----- | ----- | ----- | ----- | ----- | 64 |
| Tongs..... | 387 | 6 | 120 | 413 | 142 | 245 | 465 | 222 | 2,000 |
| Rakes..... | 433 | ----- | 4 | ----- | ----- | ----- | ----- | ----- | 437 |
| Forks..... | ----- | ----- | ----- | 40 | ----- | ----- | ----- | ----- | 40 |
| Grabs..... | ----- | 333 | 60 | ----- | ----- | ----- | ----- | ----- | 393 |
| Coquina scoops..... | ----- | ----- | ----- | 3 | ----- | ----- | ----- | ----- | 3 |
| Hooks, sponge..... | ----- | ----- | ----- | 201 | ----- | ----- | ----- | ----- | 201 |
| Diving apparatus..... | ----- | ----- | ----- | 54 | ----- | ----- | ----- | ----- | 54 |

CATCH: BY STATES¹

| Species | North Carolina | | South Carolina | | Georgia | |
|----------------------------|----------------|--------------|----------------|--------------|---------------|--------------|
| | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> |
| FISH | | | | | | |
| Alwives..... | 6,584,000 | \$41,899 | ----- | ----- | ----- | ----- |
| Black bass..... | 31,800 | 3,180 | ----- | ----- | ----- | ----- |
| Bluefish..... | 686,597 | 16,409 | 4,062 | \$325 | ----- | ----- |
| Bowfin..... | 1,700 | 17 | ----- | ----- | ----- | ----- |
| Butterfish..... | 54,514 | 786 | ----- | ----- | ----- | ----- |
| Carp..... | 128,400 | 6,640 | ----- | ----- | ----- | ----- |
| Catfish and bullheads..... | 524,904 | 9,600 | ----- | ----- | 98,389 | \$5,841 |
| Cero..... | 13,000 | 520 | ----- | ----- | ----- | ----- |
| Croaker..... | 4,540,356 | 46,642 | ----- | ----- | 8,226 | 329 |
| Drum, red or redfish..... | 87,200 | 1,744 | 3,170 | 108 | 2,141 | 107 |
| Eels..... | 56,715 | 1,877 | ----- | ----- | 550 | 22 |
| Flounders..... | 789,767 | 32,797 | 5,175 | 284 | 2,904 | 88 |
| Gizzard shad..... | 19,200 | 161 | ----- | ----- | ----- | ----- |

See footnotes at end of table.

Fisheries of the South Atlantic and Gulf States, 1932—Continued

CATCH: BY STATES—Continued

| Species | North Carolina | | South Carolina | | Georgia | |
|---------------------------------|-------------------|----------------|------------------|----------------|-------------------|----------------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH—continued | | | | | | |
| Grunts..... | | | 6,300 | \$220 | | |
| Hake..... | 1,624 | \$22 | | | | |
| Harvestfish or "sta. fish"..... | 1,077,381 | 11,858 | | | | |
| Hickory shad..... | 117,325 | 4,055 | 11,068 | 886 | 9,841 | \$707 |
| Hogfish..... | 992 | 12 | | | | |
| King whiting or "kingfish"..... | 300,048 | 4,033 | 16,210 | 610 | 19,746 | 617 |
| Mechiniden..... | 54,476,000 | 75,135 | | | 11,520,000 | 16,000 |
| Mullet..... | 2,472,050 | 51,635 | 148,050 | 6,042 | 52,627 | 1,904 |
| Pigfish..... | 62,300 | 827 | | | | |
| Pike or pickerel..... | 5,300 | 393 | | | | |
| Pinfish or sailors choice..... | 270,000 | 1,012 | | | | |
| Pompano..... | 150 | 22 | | | | |
| Scup..... | 5,615 | 172 | | | | |
| Sea bass..... | 202,485 | 6,251 | 218,750 | 8,187 | 32,000 | 960 |
| Shad..... | 924,994 | 123,926 | 123,036 | 15,459 | 288,145 | 46,111 |
| Sharks..... | | | 8,000 | 80 | | |
| Sheepshead..... | 2,650 | 53 | | | | |
| Spanish mackerel..... | 77,900 | 3,660 | | | | |
| Spot..... | 1,587,555 | 17,821 | 10,000 | 400 | 9,542 | 351 |
| Squeezes or "sea trout": | | | | | | |
| Gray..... | 3,636,323 | 64,067 | 2,460 | 148 | 2,000 | 120 |
| Spotted..... | 1,895,700 | 78,363 | 14,355 | 1,048 | 46,210 | 3,357 |
| Striped bass..... | 506,760 | 54,516 | | | | |
| Sturgeon..... | 1,661 | 179 | 23,340 | 3,734 | 4,965 | 397 |
| Suckers..... | 450 | 9 | | | | |
| Sunfish..... | 55,250 | 1,105 | | | | |
| White perch..... | 831,600 | 21,302 | | | | |
| Yellow perch..... | 179,900 | 4,871 | | | | |
| Total..... | 82,209,976 | 689,421 | 593,974 | 37,531 | 12,097,286 | 75,911 |
| SHELLFISH, ETC. | | | | | | |
| Crabs: | | | | | | |
| Hard ³ | 1,847,600 | 18,448 | 16,000 | 320 | 226,492 | 3,383 |
| Soft..... | 308,555 | 33,921 | | | | |
| Shrimp..... | 292,104 | 9,393 | 1,500,687 | 32,529 | 3,601,564 | 89,547 |
| Clams: | | | | | | |
| Hard, public ³ | 260,624 | 17,278 | 4,800 | 600 | 600 | 75 |
| Oysters:⁴ | | | | | | |
| Market, public, spring..... | 626,462 | 26,067 | 1,205,886 | 21,569 | | |
| Market, public, fall..... | 563,478 | 26,613 | 475,704 | 10,175 | | |
| Market, private, spring..... | 10,216 | 659 | 429,460 | 9,646 | 413,121 | 8,789 |
| Market, private, fall..... | 1,200 | 100 | 306,791 | 10,466 | 175,287 | 6,881 |
| Scallops, bay..... | 91,458 | 6,560 | | | | |
| Octopus..... | | | 1,200 | 72 | | |
| Squid..... | 763 | 13 | | | | |
| Terrapin, diamond-back..... | 1,557 | 370 | 1,786 | 483 | 9,645 | 1,356 |
| Total..... | 4,004,017 | 187,322 | 3,942,314 | 85,860 | 4,425,709 | 110,031 |
| Grand total..... | 86,213,993 | 826,743 | 4,536,288 | 123,391 | 16,522,995 | 185,942 |

| Species | Florida | | Alabama | | Mississippi | |
|------------------------------|-----------|---------|---------|-------|-------------|-------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | |
| Alewives..... | 79,947 | \$437 | | | | |
| Amberjack..... | 4,577 | 122 | | | | |
| Barracuda..... | 4,245 | 180 | | | | |
| Black bass..... | 278,477 | 18,618 | | | | |
| Bluefish..... | 1,421,233 | 60,614 | 12,401 | \$563 | 4,750 | \$66 |
| Blue runner or hardtail..... | 162,807 | 2,311 | 924 | 17 | | |
| Buttalo fish..... | | | 11,829 | 323 | | |
| Butterfish..... | 997 | 47 | | | | |
| Caho or crab eater..... | 5,145 | 103 | 550 | 15 | 110 | 2 |
| Catfish and bullheads..... | 3,531,636 | 116,214 | 60,211 | 2,736 | 27,115 | 493 |
| Cero..... | 275 | 4 | | | | |
| Cigarfish..... | 9,350 | 170 | | | | |
| Cod..... | 2,039 | 43 | | | | |
| Crapple..... | 404,926 | 11,866 | | | | |
| Crevalle..... | 22,761 | 518 | 259 | 5 | 990 | 9 |
| Croaker..... | 25,775 | 431 | 18,111 | 330 | 10,835 | 191 |
| Dolphin..... | 12,060 | 361 | | | | |
| Drum: | | | | | | |
| Black..... | 48,010 | 817 | 742 | 14 | 8,937 | 115 |
| Red or redfish..... | 704,784 | 11,894 | 44,292 | 2,645 | 75,100 | 2,062 |
| Eels..... | 7,580 | 153 | | | | |
| Flounders..... | 465,131 | 12,365 | 21,490 | 1,668 | 46,540 | 2,129 |
| Frigate mackerel..... | 2,260 | 90 | | | | |

Fisheries of the South Atlantic and Gulf States, 1932—Continued

CATCH: BY STATES—Continued

| Species | Florida | | Alabama | | Mississippi | |
|------------------------------------|---------------|-------------|-------------|----------|--------------|----------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH—continued | | | | | | |
| Groupers..... | 3, 163, 878 | \$64, 600 | | | | |
| Grunts..... | 44, 391 | 1, 297 | 99, 746 | \$1, 998 | 16, 117 | |
| Hake..... | 8, 218 | 165 | | | | |
| Hickory shad..... | 28, 147 | 507 | | | | |
| Hogfish..... | 28, 430 | 853 | | | | |
| Jewfish..... | 30, 290 | 1, 034 | | | | |
| Kingfish or "king mackerel"..... | 3, 294, 501 | 119, 544 | 880 | 40 | | |
| King whiting or "kingfish"..... | 285, 069 | 6, 880 | 3, 718 | 68 | 2, 728 | 45 |
| Ladyfish..... | 2, 622 | 79 | | | | |
| Menhaden..... | 23, 349, 860 | 41, 220 | | | | |
| Mojarro..... | 35, 689 | 514 | | | | |
| Mullet..... | 21, 141, 449 | 338, 254 | 696, 968 | 10, 673 | 564, 970 | 8, 235 |
| Muttonfish..... | 203, 135 | 8, 811 | | | | |
| Paddlefish or spoonbill cat..... | | | 1, 320 | 60 | | |
| Permit..... | 2, 850 | 107 | | | | |
| Pigfish..... | 66, 648 | 948 | | | | |
| Pinfish or sailors choice..... | 24, 977 | 484 | | | | |
| Pompano..... | 631, 263 | 80, 067 | 8, 144 | 436 | 132 | 12 |
| Porgies..... | 25, 786 | 612 | | | | |
| Porkfish..... | 363 | 7 | | | | |
| Scup..... | 247, 792 | 5, 936 | | | | |
| Sea bass..... | 250, 995 | 8, 102 | | | | |
| Shad..... | 646, 066 | 52, 940 | | | | |
| Sharks..... | 6, 043, 000 | 12, 005 | | | | |
| Sheepshead..... | 635, 330 | 8, 466 | 4, 441 | 120 | 23, 815 | 639 |
| Snapper: | | | | | | |
| Mangrove..... | 95, 580 | 2, 407 | | | | |
| Red..... | 4, 688, 265 | 228, 536 | 691, 673 | 30, 263 | 36, 812 | 1, 841 |
| Snook or sergeantfish..... | 301, 780 | 6, 936 | | | | |
| Spanish mackerel..... | 6, 837, 588 | 209, 836 | 8, 028 | 292 | | |
| Spot..... | 68, 360 | 925 | 401 | 7 | | |
| Squeteagues or "sea trout": | | | | | | |
| Gray..... | 21, 418 | 674 | 6, 050 | 110 | 103, 015 | 1, 873 |
| Spotted..... | 2, 666, 525 | 106, 425 | 108, 224 | 9, 392 | 124, 394 | 4, 524 |
| Sturgeon..... | 4, 370 | 199 | 10, 742 | 977 | | |
| Sunfish..... | 662, 494 | 16, 831 | | | | |
| Tenpounder..... | 77, 845 | 1, 849 | 1, 400 | 14 | 350 | 4 |
| Trillettail..... | 890 | 18 | | | 176 | 4 |
| Tuna or "horse mackerel"..... | 3, 850 | 134 | | | | |
| Turbot..... | 4, 125 | 124 | | | | |
| Yellowtail..... | 91, 870 | 4, 441 | | | | |
| Total..... | 81, 108, 701 | 1, 599, 398 | 1, 792, 434 | 62, 766 | 1, 046, 966 | 22, 486 |
| SHELLFISH, ETC. | | | | | | |
| Crabs: | | | | | | |
| Hard..... | 82, 182 | 3, 519 | 70, 070 | 982 | 320, 107 | 4, 665 |
| Soft..... | | | 1, 280 | 236 | 3, 572 | 893 |
| Stone..... | 153, 825 | 8, 335 | | | | |
| Sea crawfish or spiny lobster..... | 445, 547 | 32, 078 | | | | |
| Shrimp..... | 18, 136, 334 | 535, 198 | 3, 381, 700 | 71, 910 | 14, 009, 720 | 267, 428 |
| Clams: | | | | | | |
| Coquina..... | 5, 400 | 335 | | | | |
| Hard, public..... | 1, 120, 812 | 42, 742 | | | | |
| Conchs..... | 1, 600 | 120 | | | | |
| Oysters: | | | | | | |
| Market, public, spring..... | 542, 438 | 27, 493 | 748, 962 | 27, 216 | 4, 472, 358 | 169, 783 |
| Market, public, fall..... | 669, 715 | 35, 668 | 88, 485 | 3, 802 | 749, 962 | 32, 162 |
| Market, private, spring..... | 186, 558 | 7, 886 | 3, 960 | 220 | | |
| Market, private, fall..... | 113, 495 | 6, 320 | 17, 820 | 990 | | |
| Scallops, bay..... | 61, 965 | 6, 885 | | | | |
| Squid..... | 7, 653 | 147 | | | | |
| Frogs..... | | | 697 | 104 | | |
| Terrapin, diamond-back..... | | | 1, 089 | 275 | | |
| Turtles, soft-shell..... | 51, 669 | 336 | | | | |
| Sponges: | | | | | | |
| Grass..... | 181, 867 | 37, 319 | | | | |
| Sheepswool..... | 277, 087 | 593, 674 | | | | |
| Velvet..... | 71 | 20 | | | | |
| Wire..... | 29, 466 | 13, 387 | | | | |
| Yellow..... | 124, 536 | 52, 524 | | | | |
| Total..... | 22, 181, 520 | 1, 403, 986 | 4, 314, 053 | 105, 825 | 19, 555, 719 | 474, 931 |
| Grand total..... | 103, 290, 221 | 2, 973, 384 | 6, 106, 487 | 168, 591 | 20, 602, 585 | 497, 417 |

See footnotes at end of table.

Fisheries of the South Atlantic and Gulf States, 1932—Continued

CATCH: BY STATES—Continued

| Species | Louisiana | | Texas | | Total | |
|----------------------------------|------------------|---------------|------------------|----------------|--------------------|------------------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | |
| Alewives..... | | | | | 6,063,947 | \$42,336 |
| Amberjack..... | | | | | 4,577 | 122 |
| Barracuda..... | | | | | 4,245 | 180 |
| Black bass..... | | | | | 310,277 | 21,698 |
| Bluefish..... | | | 1,760 | \$80 | 2,130,783 | 78,077 |
| Blue runner or hardtail..... | | | | | 163,431 | 2,328 |
| Howfin..... | | | | | 1,700 | 17 |
| Buffalofish..... | | | | | 11,829 | 323 |
| Butterfish..... | | | | | 55,511 | 833 |
| Cabio or crab eater..... | | | | | 5,805 | 120 |
| Carp..... | | | | | 128,400 | 6,640 |
| Catfish and bullheads..... | 44,850 | \$1,583 | 76,825 | 2,752 | 4,363,930 | 139,219 |
| Cero..... | | | | | 13,275 | 524 |
| Cigarfish..... | | | | | 9,350 | 170 |
| Cod..... | | | | | 2,039 | 43 |
| Crapple..... | | | | | 404,926 | 11,866 |
| Crevalle..... | 300 | 9 | | | 24,300 | 541 |
| Croaker..... | 44,470 | 1,924 | 27,025 | 676 | 4,674,798 | 50,423 |
| Dolphin..... | | | | | 12,050 | 361 |
| Drum: | | | | | | |
| Black..... | 87,412 | 2,704 | 932,091 | 17,153 | 1,077,192 | 20,803 |
| Red or redfish..... | 281,739 | 14,493 | 824,819 | 45,322 | 2,083,245 | 78,375 |
| Eels..... | | | | | 64,825 | 2,062 |
| Flounders..... | 4,405 | 314 | 70,515 | 4,614 | 1,395,927 | 54,259 |
| Frigate mackerel..... | | | | | 2,250 | 90 |
| Garfish..... | 300 | 15 | | | 300 | 15 |
| Gizzard shad..... | | | | | 19,200 | 161 |
| Groupers..... | 3,400 | 68 | 18,301 | 380 | 3,301,442 | 67,368 |
| Grunts..... | | | | | 50,691 | 1,517 |
| Hake..... | | | | | 9,842 | 187 |
| Harvestfish or "starfish"..... | | | | | 1,077,381 | 11,858 |
| Hickory shad..... | | | | | 160,379 | 6,156 |
| Hogfish..... | | | | | 29,422 | 866 |
| Jewfish..... | 2,400 | 48 | 5,750 | 165 | 38,440 | 1,247 |
| Kingfish or "king mackerel"..... | | | 5,280 | 162 | 3,300,661 | 119,746 |
| King whiting or "kingfish"..... | 16,000 | 374 | 8,535 | 155 | 652,044 | 12,782 |
| Ladyfish..... | | | | | 2,622 | 39 |
| Menhaden..... | | | | | 89,345,860 | 132,355 |
| Mojarro..... | | | | | 35,889 | 514 |
| Mullet..... | 6,300 | 155 | 4,950 | 90 | 25,087,354 | 417,008 |
| Muttonfish..... | | | | | 203,135 | 8,811 |
| Paddlefish or spoonbill cat..... | | | | | 1,320 | 60 |
| Permit..... | | | | | 2,850 | 107 |
| Pigfish..... | | | | | 128,748 | 1,575 |
| Pike or pickerel..... | | | | | 6,200 | 393 |
| Pinfish or sailors choice..... | | | | | 204,975 | 1,497 |
| Pompano..... | 90 | 11 | 5,159 | 469 | 589,938 | 81,037 |
| Porgies..... | | | | | 25,786 | 512 |
| Porkfish..... | | | | | 363 | 7 |
| Scup..... | | | | | 253,407 | 6,108 |
| Sea bass..... | | | | | 704,240 | 23,500 |
| Shad..... | | | | | 1,882,261 | 239,436 |
| Sharks..... | | | | | 5,051,000 | 12,085 |
| sheepshead..... | 77,673 | 4,019 | 29,154 | 599 | 673,063 | 13,786 |
| Snapper: | | | | | | |
| Mangrove..... | | | | | 95,580 | 2,407 |
| Red..... | 66,884 | 4,013 | 985,291 | 50,076 | 6,358,825 | 314,729 |
| Snook or sergeantfish..... | | | 20,893 | 569 | 322,673 | 7,505 |
| Spanish mackerel..... | 400 | 16 | 41,140 | 2,616 | 6,465,066 | 216,420 |
| Spot..... | 3,450 | 87 | | | 1,679,308 | 19,591 |
| Squeteagues or "sea trout": | | | | | | |
| Gray..... | 220,471 | 6,603 | | | 3,991,737 | 73,627 |
| Spotted..... | 412,427 | 31,607 | 976,344 | 63,660 | 6,239,179 | 298,376 |
| Striped bass..... | | | 495 | 18 | 507,255 | 54,534 |
| Sturgeon..... | | | | | 45,087 | 5,486 |
| Suckers..... | | | | | 450 | 9 |
| Sunfish..... | | | | | 717,744 | 17,936 |
| Tenpounder..... | | | | | 79,595 | 1,367 |
| Tripletail..... | 990 | 49 | | | 2,056 | 71 |
| Tuna or "horse mackerel"..... | | | | | 3,350 | 134 |
| Turbot..... | | | | | 4,125 | 124 |
| White perch..... | | | | | 831,600 | 21,302 |
| Yellow perch..... | | | | | 179,900 | 4,871 |
| Yellowtail..... | | | | | 91,870 | 4,441 |
| Total..... | 1,273,961 | 68,092 | 4,034,327 | 189,466 | 184,157,525 | 2,715,061 |

Fisheries of the South Atlantic and Gulf States, 1932—Continued

CATCH: BY STATES—Continued

| Species | Louisiana | | Texas | | Total | |
|------------------------------------|--------------|-------------|--------------|----------|---------------|-------------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| SHELLFISH, ETC. | | | | | | |
| Crabs: | | | | | | |
| Hard ¹ | 5, 877, 737 | \$56, 776 | 44, 660 | \$669 | 8, 483, 848 | \$58, 762 |
| Soft..... | 99, 340 | 25, 258 | | | 412, 747 | 60, 308 |
| Stone..... | | | | | 153, 825 | 8, 335 |
| Sea crawfish or spiny lobster..... | | | | | 446, 547 | 32, 078 |
| Shrimp..... | 38, 095, 780 | 800, 452 | 9, 244, 246 | 229, 529 | 88, 262, 135 | 2, 036, 986 |
| Clams: | | | | | | |
| Coquina..... | | | | | 5, 400 | 335 |
| Hard, public ² | | | | | 1, 366, 836 | 60, 695 |
| Conchs..... | | | | | 1, 600 | 120 |
| Oysters: ³ | | | | | | |
| Market, public, spring..... | 267, 672 | 16, 054 | 442, 932 | 25, 501 | 8, 306, 700 | 312, 773 |
| Market, public, fall..... | | | 537, 669 | 27, 019 | 3, 075, 013 | 134, 529 |
| Market, private, spring..... | 1, 164, 853 | 92, 616 | | | 2, 208, 168 | 119, 716 |
| Market, private, fall..... | 1, 545, 636 | 119, 657 | | | 2, 160, 129 | 144, 414 |
| Scallops, bay..... | | | | | 153, 423 | 13, 445 |
| Octopus..... | | | | | 1, 200 | 72 |
| Squid..... | | | | | 8, 316 | 160 |
| Frogs..... | | | | | 697 | 104 |
| Terrapin, diamond-back..... | 8, 996 | 1, 619 | | | 23, 073 | 4, 103 |
| Turtles: | | | | | | |
| Soft-shell..... | | | | | 51, 669 | 336 |
| Loggerhead..... | 6, 450 | 129 | | | 6, 450 | 129 |
| Sponges: | | | | | | |
| Grass..... | | | | | 181, 367 | 37, 319 |
| Sheepswool..... | | | | | 277, 087 | 593, 674 |
| Velvet..... | | | | | 71 | 20 |
| Wire..... | | | | | 29, 466 | 13, 387 |
| Yellow..... | | | | | 124, 636 | 52, 524 |
| Total..... | 47, 066, 364 | 1, 112, 561 | 10, 269, 507 | 282, 808 | 115, 759, 203 | 3, 713, 324 |
| Grand total..... | 48, 340, 325 | 1, 180, 653 | 14, 303, 834 | 472, 264 | 299, 916, 728 | 6, 428, 385 |

¹ Excluding seed oyster fishery. The seed oyster fishery was prosecuted in this section only in North Carolina where 12 regular fishermen using 6 motor boats and 12 dredges with an aggregate of 12 yards at the mouth took 39,741 bushels of seed oysters, valued at \$8,280, from public beds. None of these fishermen, craft, or gear was duplicated among those in the fisheries for market oysters or other species.

² Statistics on hard crabs used in this table are based on yields of 3 pounds per dozen in North Carolina; 6 pounds, in South Carolina and Georgia; 7.32 pounds, in Florida; 6.25 pounds, in Mississippi; 6.98 pounds, in Alabama and Texas; and 6.45 pounds, in Louisiana.

³ Statistics on hard clams used in this table are based on yields of 8 pounds of meats per bushel in all States.

⁴ Statistics on market oysters used in this table are based on yields of 5.71 pounds of meats per bushel in North Carolina; 4.76, in South Carolina; 5.69, in Georgia; 3.29, in Florida; 2.40, in Alabama; 2.19, in Mississippi; 4.14, in Louisiana; and 5.05, in Texas.

NOTE.—Of the total catch in North Carolina, 268,136 pounds of fishery products, valued at \$5,925, were taken in the winter trawl fishery of Maryland, Virginia, and North Carolina. Of the total catch in Florida, 942,791 pounds of fishery products, valued at \$20,607, were taken in the same fishery. These products consisted principally of scup, sea bass, flounders, croakers, and gray squeteague.

Industries related to the fisheries of the South Atlantic and Gulf States

OPERATING UNITS, SALARIES, AND WAGES, 1931

| Item | North Carolina | South Carolina | Georgia | Florida | Alabama | Mississippi | Louisiana | Texas | Total |
|------------------------|----------------|----------------|---------|---------|---------|-------------|-----------|-------|--------|
| Transporting: | | | | | | | | | |
| Persons engaged: | | | | | | | | | |
| On vessels..... | 60 | 85 | 8 | 43 | 9 | 16 | 147 | 2 | 354 |
| On boats..... | 4 | 39 | | 73 | 2 | | 255 | | 349 |
| Total..... | 64 | 124 | 8 | 116 | 11 | 16 | 402 | 2 | 743 |
| Vessels: | | | | | | | | | |
| Motor..... | 37 | 9 | 4 | 22 | 8 | | 71 | 1 | 152 |
| Net tonnage..... | 342 | 101 | 26 | 318 | 68 | | 579 | 6 | 1, 440 |
| Sail..... | | 29 | | 1 | | | | | 30 |
| Net tonnage..... | | 263 | | 16 | | | | | 279 |
| Total vessels..... | 37 | 38 | 4 | 23 | 8 | | 71 | 1 | 182 |
| Total net tonnage..... | 342 | 364 | 26 | 334 | 68 | | 579 | 6 | 1, 719 |
| Boats..... | 2 | 41 | | 72 | 1 | 8 | 127 | | 251 |

Industries related to the fisheries of the South Atlantic and Gulf States—Continued

OPERATING UNITS, SALARIES, AND WAGES, 1931—Continued

| Item | North Carolina | South Carolina | Georgia | Florida | Alabama | Mississippi | Louisiana | Texas | Total |
|---------------------------------|----------------|----------------|-----------|-----------|-----------|-------------|-----------|-----------|-------------|
| Wholesale and manufacturing: | | | | | | | | | |
| Establishments..... | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| Persons engaged: | 64 | 26 | 33 | 250 | 20 | 46 | 90 | 52 | 585 |
| Proprietors..... | 70 | 23 | 34 | 311 | 27 | 62 | 112 | 61 | 700 |
| Salariat employees..... | 25 | 24 | 22 | 105 | 16 | 31 | 96 | 33 | 352 |
| Wage earners: | | | | | | | | | |
| Average for season..... | 669 | 763 | 1,198 | 2,447 | 352 | 2,287 | 3,287 | 1,581 | 12,583 |
| Average for year..... | 265 | 269 | 273 | 927 | 183 | 968 | 1,280 | 416 | 4,581 |
| Paid to salaried employees..... | \$33,112 | \$58,986 | \$94,047 | \$257,707 | \$41,238 | \$82,525 | \$202,958 | \$45,436 | \$786,003 |
| Paid to wage earners..... | \$146,855 | \$109,845 | \$137,498 | \$561,198 | \$59,919 | \$301,101 | \$541,074 | \$178,050 | \$2,035,540 |
| Total salaries and wages..... | \$179,967 | \$168,831 | \$201,545 | \$818,905 | \$101,157 | \$383,626 | \$744,032 | \$223,486 | \$2,821,543 |
| Fishermen manufacturing..... | 729 | 10 | 17 | 254 | 46 | 86 | 10 | 227 | 1,379 |

PRODUCTS MANUFACTURED

| Item | North Carolina | | South Carolina | | Georgia | | Florida | |
|---|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| By manufacturing establishments: | | | | | | | | |
| Groupers, fresh steaks ¹ pounds..... | | | | | | | 254,000 | \$26,200 |
| Menhaden products: | | | | | | | | |
| Acid scrap ¹ tons..... | 3,564 | \$56,733 | | | (²) | (²) | 150 | 9,000 |
| Dry scrap ¹ do..... | 2,560 | 84,282 | | | | | 5,069 | 138,475 |
| Fish meal ¹ do..... | 737 | 24,008 | | | | | 2,948 | 81,776 |
| Oil ¹ gallons..... | 233,866 | 23,007 | | | (²) | (²) | 666,111 | 68,522 |
| Mullet: | | | | | | | | |
| Salted..... pounds..... | 278,000 | 11,335 | | | | | 320,167 | 12,754 |
| Roe, salted..... do..... | | | | | | | 30,590 | 8,485 |
| Crab meat, packaged, fresh-cooked..... pounds..... | 188,032 | 33,748 | | | 18,250 | \$5,844 | (²) | (²) |
| Shrimp: | | | | | | | | |
| Cooked and peeled do..... | | | | | (²) | (²) | | |
| Canned ¹ standard cases..... | | | (²) | (²) | 116,945 | 601,686 | 86,013 | 343,209 |
| Oysters: | | | | | | | | |
| Fresh-shucked..... gallons..... | 119,238 | 111,350 | 20,514 | \$21,668 | 23,467 | 25,897 | 113,629 | 142,827 |
| Canned ¹ standard cases..... | | | 93,082 | 274,674 | (²) | (²) | (²) | (²) |
| Crushed shells for poultry feed ¹ tons..... | (²) | (²) | (²) | (²) | | | 65,363 | 318,715 |
| Marine-shell novelties ¹ | | | | | | | | 60.0 |
| Unclassified products: | | | | | | | | |
| Steaks and fillets, fresh and frozen ¹ pounds..... | (²) | (²) | | | | | 4 29,000 | 4 5,194 |
| Canned ¹ standard cases..... | | (²) | | (²) | | (²) | 4 21,606 | 4 124,689 |
| Miscellaneous ² | | 10,365 | | 131,750 | | 19 69,495 | | 11 17,925 |
| Total..... | | 354,828 | | 428,092 | | 602,922 | | 1,296,371 |

¹ Data are for 1933.² This item has been included under "Unclassified products."³ This item has been included under "Miscellaneous."⁴ Includes fresh fillets of red drum, grouper, kingfish, mullet, red snapper, Spanish mackerel, squeteagues, and snook; and fresh steaks of red snapper.⁵ Includes canned coquina broth, hard clam products, conch cocktail, oysters, and turtle products.⁶ Both 1931 and 1933 data are included in these items.⁷ Includes canned alewife roe; salted spot; fresh fillets of squeteagues; fresh-shucked hard clams; and oyster shell products.⁸ Includes canned shrimp and oyster-shell products.⁹ Includes cooked and peeled shrimp; fresh-shucked hard clams; canned oysters; and menhaden products.¹⁰ Includes dry-salted bluefish and Spanish mackerel; fresh-cooked crab meat; fresh-shucked bay scallops, hard clams and conchs; shark hides, fins and oil; oyster-shell lime; and fish scale novelties.

Industries related to the fisheries of the South Atlantic and Gulf States—Continued

PRODUCTS MANUFACTURED—Continued

| Item | North Carolina | | South Carolina | | Georgia | | Florida | |
|--|----------------|---------|----------------|-------------------|----------|---------|---------------------|-----------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| By fishermen: | | | | | | | | |
| Alewives: | | | | | | | | |
| Corned.....pounds.. | 2,562,600 | \$34.97 | | | | | | |
| Tight-pack cut.....do.. | 23,100 | 84 | | | | | | |
| Tight-pack roe.....do.. | 182,000 | 7,86 | | | | | | |
| Smoked.....do..... | 6,000 | 45 | | | | | | |
| Mullet: | | | | | | | | |
| Salted.....do..... | 110,000 | 4,90 | | | | | 475,19 ¹ | \$18,793 |
| Roe, salted.....do..... | | | | | | | 44,114 | 10,855 |
| Spot, salted.....do..... | 25,000 | 1,125 | | | | | | |
| Sturgeon caviar.....do..... | | | 312 | \$15 ² | | | 12 | 10 |
| Crab meat, packaged, fresh-cooked.....pounds | | | | | | | 1,000 | 500 |
| Clams, hard, fresh-shucked.....gallons | 300 | 300 | | | | | | |
| Oysters, fresh-shucked.....gallons | 15,150 | 12,120 | 1,139 | 1,159 | 5,648 | \$5,648 | 27,915 | 35,870 |
| Scallops, bay, fresh-shucked.....gallons | 55,000 | 56,250 | | | | | 1,265 | 1,305 |
| Total..... | | 118,829 | | 1,315 | | 5,648 | | 67,343 |
| Grand total..... | | 473,657 | | 429,407 | | 608,570 | | 1,363,714 |

| Item | Alabama | | Mississippi | | Louisiana | | Texas | |
|--|------------------|------------------|-------------|---------------------|------------------------|------------------|---------------------|-----------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| By manufacturing establishments: | | | | | | | | |
| Groupers, fresh steaks¹ | | | | | | | | |
| Crab meat, packaged, fresh-cooked.....pounds.. | | | 22,500 | \$5,82 ² | 174,550 | \$47,184 | | |
| Shrimp: | | | | | | | | |
| Fresh and frozen packaged.....pounds.. | | | | | 1,333,568 ³ | 302,041 | 2,066,950 | \$294,903 |
| Sun-dried.....do..... | | | | | | | | |
| Cooked and peeled.....do..... | | | 326,550 | 81,63 ⁴ | | | | |
| Canned.....standard cases | 73,705 | \$298,151 | 119,872 | 447,072 | 374,481 | 1,512,778 | 70,455 ⁵ | 202,301 |
| Meal or "bran" ⁶tons.. | | | | | 943 | 17,550 | | |
| Oysters: | | | | | | | | |
| Fresh-shucked.....gallons.. | 20,370 | 23,424 | 41,436 | 42,198 ⁷ | 202,015 | 256,882 | 65,294 | 74,448 |
| Canned.....standard cases | | | 183,127 | 527,221 | 19,439 | 60,104 | | |
| Crushed shells for poultry feed ⁸tons.. | (⁹) | (⁹) | 14,567 | 46,992 | (⁹) | (⁹) | | |
| Unclassified products: | | | | | | | | |
| Steaks and fillets, fresh and frozen ¹⁰pounds.. | 16,900 | 3,757 | | | | | | |
| Canned.....standard cases | | | | | | | | |
| Miscellaneous ¹¹ | | 45,780 | | 1,728 | | 610,914 | | 5,433 |
| Total..... | | 371,142 | | 1,152,674 | | 2,807,513 | | 667,086 |
| By fishermen: | | | | | | | | |
| Crab meat, packaged, fresh-cooked.....pounds.. | | | | | | | 460 | 115 |
| Shrimp: | | | | | | | | |
| Sun-dried.....do..... | | | | | 13,900 | 2,780 | | |
| Meal or "bran" ⁶tons.. | | | | | 7 | 105 | | |
| Oysters, fresh-shucked.....gallons | 6,289 | 6,747 | 17,133 | 16,205 | | | 46,151 | 39,718 |
| Total..... | | 6,747 | | 16,205 | | 2,885 | | 39,833 |
| Grand total..... | | 377,889 | | 1,168,879 | | 2,810,398 | | 706,918 |

¹ This item has been included under "Unclassified products."

² This item has been included under "Miscellaneous."

³ Includes fresh steaks of grouper and red snapper; and fresh and frozen fillets of flounders, red snapper, Spanish mackerel, and squeteagues.

⁴ Includes canned oysters and oyster-shell products.

⁵ Includes salted mullet and oyster-shell lime.

⁶ Includes sun-dried squeteagues and oyster-shell products.

⁷ Includes fresh-cooked crab meat and oyster-shell poultry feed.

NOTE.—Unless otherwise indicated the data are for 1931. The total value of manufactured products for the South Atlantic and Gulf States was as follows: By manufacturing establishments, \$7,680,627; and by fishermen, \$258,805. Some of the above products may have been manufactured from products imported from another State or country, therefore, they cannot be correlated directly with the catch within the State.

SPONGES SOLD AT THE EXCHANGE, TARPON SPRINGS, FLA.

During 1933 sponges handled on the exchange at Tarpon Springs, Fla., amounted to 373,178 pounds, valued at \$420,481. This is a decrease of 13 percent in quantity and 19 percent in value as compared with the transactions on the exchange during 1932. Of the total sponges sold on the exchange during 1933, 107,642 pounds, valued at \$184,967, were large wool; 21,689 pounds, valued at \$23,983, were medium and small wool; 130,165 pounds, valued at \$143,548, were wool rags; 80,164 pounds, valued at \$51,487, were yellow; 20,105 pounds, valued at \$9,494, were grass; and 13,413 pounds, valued at \$7,002, were wire. It is estimated that sponges valued at \$95,000 were sold outside the exchange.

FISHERIES OF THE PACIFIC COAST STATES¹⁰

The yield of the commercial fisheries of the Pacific Coast States (Washington, Oregon, and California) during 1933 amounted to 860,161,216 pounds, valued at \$13,987,992 to the fishermen, representing an increase of 53 percent in volume and 47 percent in value as compared with the catch in the previous year. These fisheries gave employment to 18,673 fishermen as compared with 17,882 in 1932.

There were 313 fishery wholesale and manufacturing establishments in the three States in 1933 as compared with 316 in 1931, when the most recent previous survey of such concerns was made. In 1933 these establishments employed 11,993 persons, paid \$6,095,492 in salaries and wages, and produced manufactured products (canned, cured, packaged, and byproducts) valued at \$28,946,754. In 1931 the wholesale and manufacturing firms employed 11,651 persons, paid \$6,750,607 in salaries and wages, and produced manufactured products, valued at \$28,652,513.

Fisheries of the Pacific Coast States, 1933

SUMMARY OF CATCH

| Product | Washington | | Oregon | |
|---------------------|-------------|-------------|------------|-------------|
| | Pounds | Value | Pounds | Value |
| Fish..... | 123,420,393 | \$5,230,293 | 22,503,816 | \$1,131,977 |
| Shellfish, etc..... | 5,309,111 | 453,595 | 2,029,102 | 77,867 |
| Total..... | 128,729,504 | 5,683,891 | 24,532,918 | 1,209,844 |

| Product | California | | Total | |
|---------------------|-------------|-------------|-------------|--------------|
| | Pounds | Value | Pounds | Value |
| Fish..... | 694,594,850 | \$6,430,698 | 840,523,059 | \$12,792,908 |
| Shellfish, etc..... | 7,969,054 | 579,292 | 15,307,267 | 1,110,757 |
| Whale products..... | 4,330,890 | 84,267 | 4,330,890 | 84,267 |
| Total..... | 706,894,794 | 7,094,257 | 860,161,216 | 13,987,992 |

¹⁰ Data on the operating units and catch of the fisheries of the Pacific Coast States have been taken largely from statistics collected by the various State agencies. Supplementary surveys, compilations, and analyses have been made by agents of this Bureau in order that the figures may be presented in a manner comparable with those of other sections. While statistics of the fisheries of California are for the calendar year, those for Oregon and Washington are for the fiscal year ending March 31, except that statistics of the halibut fishery in these latter States are for the calendar year. For a clearer understanding of the statistics published in this section the reader is referred to the section in the latter part of this document entitled "Statistical survey procedure."

Fisheries of the Pacific Coast States, 1933—Continued

OPERATING UNITS: BY STATES

| Item | Washington | | | | Oregon | | |
|--------------------------------|----------------------|------------------|-------------------------|-------------|-------------------------|------------------|-------------|
| | Puget Sound district | Coastal district | Columbia River district | Total | Columbia River district | Coastal district | Total |
| | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | |
| On vessels..... | 3, 112 | 41 | 4 | 3, 157 | 42 | 33 | 75 |
| On boats and shore..... | 1, 787 | 2, 818 | 1, 251 | 5, 856 | 2, 011 | 1, 416 | 3, 427 |
| Total..... | 4, 899 | 2, 859 | 1, 255 | 9, 013 | 2, 063 | 1, 449 | 3, 502 |
| Vessels: | | | | | | | |
| Steam..... | 3 | | | 3 | | | |
| Net tonnage..... | 75 | | | 75 | | | |
| Motor..... | 484 | 20 | 1 | 505 | 21 | 12 | 33 |
| Net tonnage..... | 10, 421 | 166 | 9 | 10, 596 | 213 | 120 | 333 |
| Sail..... | 3 | | | 3 | | | |
| Net tonnage..... | 1, 349 | | | 1, 349 | | | |
| Total vessels..... | 490 | 20 | 1 | 511 | 21 | 12 | 33 |
| Total net tonnage..... | 11, 845 | 186 | 9 | 12, 020 | 213 | 120 | 333 |
| Boats: | | | | | | | |
| Motor..... | 793 | 442 | 640 | 1, 875 | 1, 079 | 943 | 2, 022 |
| Other..... | 444 | 164 | 89 | 697 | 51 | 156 | 207 |
| Apparatus: | | | | | | | |
| Purse seines: | | | | | | | |
| Salmon..... | 241 | | | 241 | | | |
| Length, yards..... | 144, 118 | | | 144, 118 | | | |
| Haul seines..... | 59 | | 31 | 90 | 32 | 2 | 34 |
| Length, yards..... | 5, 216 | | 9, 550 | 14, 766 | 16, 135 | 317 | 15, 452 |
| Gill nets: | | | | | | | |
| Drift: | | | | | | | |
| Salmon..... | 304 | 114 | 434 | 852 | 823 | 485 | 1, 308 |
| Square yards..... | 398, 488 | 207, 480 | 1, 206, 520 | 1, 812, 488 | 2, 538, 335 | 638, 280 | 3, 226, 595 |
| Sot: | | | | | | | |
| Salmon..... | 4 | 190 | 134 | 328 | 122 | 750 | 870 |
| Square yards..... | 1, 080 | 52, 500 | 34, 036 | 87, 616 | 32, 330 | 114, 000 | 146, 332 |
| Lines: | | | | | | | |
| Trawl, set, and hand..... | 25, 215 | | 166 | 25, 381 | 342 | 915 | 1, 257 |
| Hooks..... | 521, 129 | | 4, 725 | 525, 854 | 10, 450 | 19, 775 | 30, 225 |
| Troll..... | 1, 580 | 605 | 18 | 2, 203 | 755 | 515 | 1, 270 |
| Hooks..... | 7, 118 | 2, 722 | 85 | 9, 925 | 3, 397 | 2, 317 | 5, 714 |
| Pound nets..... | 88 | 105 | 204 | 397 | 35 | | 35 |
| Brush weirs..... | 4 | | | 4 | | | |
| Fish wheels..... | | | 29 | 29 | | | |
| Dip nets..... | 7 | 46 | 150 | 203 | 140 | | 140 |
| Drag bag nets..... | 29 | 4 | | 33 | | | |
| Length, yards..... | 2, 382 | 260 | | 2, 642 | | | |
| Reef nets..... | 8 | | | 8 | | | |
| Beam trawls..... | 14 | | | 14 | | | |
| Yards at mouth..... | 66 | | | 66 | | | |
| Otter trawls..... | 16 | | | 16 | | | |
| Yards at mouth..... | 187 | | | 187 | | 40 | 40 |
| Traps: | | | | | | | |
| Crab..... | 2, 800 | 2, 464 | | 5, 264 | | 8, 340 | 8, 340 |
| Crawfish..... | | | | | 1, 500 | | 1, 500 |
| Tongs, rakes, and shove's..... | 429 | 2, 273 | | 2, 702 | | 179 | 179 |
| Dredges, oyster..... | | 4 | | 4 | | | |
| Yards at mouth..... | | | | | | | |

| Item | California | | | | | Total | Grand total |
|-------------------------|--------------------|-------------------------|--------------------|--------------------|--------------------|---------|-------------|
| | North-ern district | San Fran-cisco district | Mon-terey district | San Pedro district | San Diego district | | |
| | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | |
| On vessels..... | 34 | 388 | 538 | 1, 610 | 710 | 3, 280 | 6, 512 |
| On boats and shore..... | 455 | 1, 004 | 539 | 660 | 214 | 2, 878 | 12, 181 |
| Total..... | 489 | 1, 392 | 1, 077 | 2, 276 | 924 | 6, 158 | 18, 673 |
| Vessels: | | | | | | | |
| Steam..... | | 2 | | | | 2 | 5 |
| Net tonnage..... | | 41 | | | | 41 | 119 |
| Motor..... | 15 | 41 | 60 | 184 | 81 | 381 | 919 |
| Net tonnage..... | 132 | 821 | 1, 560 | 6, 862 | 4, 562 | 13, 937 | 24, 866 |
| Sail..... | | 2 | | | | 2 | 5 |
| Net tonnage..... | | 824 | | | | 824 | 2, 173 |
| Total vessels..... | 15 | 46 | 60 | 184 | 81 | 385 | 929 |
| Total net tonnage..... | 132 | 1, 686 | 1, 560 | 6, 862 | 4, 562 | 14, 802 | 27, 155 |

Fisheries of the Pacific Coast States, 1933—Continued

OPERATING UNITS: BY STATES—Continued

| Item | California | | | | | | Grand total |
|--------------------------------|-------------------|------------------------|-------------------|--------------------|--------------------|---------|-------------|
| | Northern district | San Francisco district | Monterey district | San Pedro district | San Diego district | Total | |
| | Number | Number | Number | Number | Number | Number | Number |
| Boats: | | | | | | | |
| Motor..... | 20 | 564 | 208 | 315 | 97 | 1,386 | 5,283 |
| Other..... | 128 | 92 | 63 | 49 | 2 | 334 | 1,238 |
| Apparatus: | | | | | | | |
| Purse seines: | | | | | | | |
| Barracuda..... | | | | 19 | | 19 | 19 |
| Length, yards..... | | | | 7,942 | | 7,942 | 7,942 |
| Salmon..... | | | | | | | 241 |
| Length, yards..... | | | | | | | 144,118 |
| Sardine..... | | 10 | 33 | 80 | | 123 | 123 |
| Length, yards..... | | 3,287 | 10,960 | 30,240 | | 44,487 | 44,487 |
| Tuna..... | | | | 47 | | 47 | 47 |
| Length, yards..... | | | | 27,129 | | 27,129 | 27,129 |
| Lampara nets: | | | | | | | |
| Mackerel..... | | | | 58 | 17 | 75 | 75 |
| Length, yards..... | | | | 23,086 | 5,610 | 28,696 | 28,696 |
| Sardine..... | | 18 | 33 | 28 | 6 | 85 | 85 |
| Length, yards..... | | 6,165 | 10,101 | 12,437 | 2,560 | 31,263 | 31,263 |
| Squid..... | | | | 34 | | 34 | 34 |
| Length, yards..... | | | 6,607 | | | 6,607 | 6,607 |
| Other..... | | | | 7 | | 7 | 7 |
| Length, yards..... | | | | 2,220 | | 2,220 | 2,220 |
| Haul seines..... | 5 | 1 | | 1 | | 7 | 131 |
| Length, yards..... | 1,040 | 165 | | 214 | | 1,419 | 31,637 |
| Gill nets: | | | | | | | |
| Drift: | | | | | | | |
| Barracuda..... | | | | 26 | 17 | 43 | 43 |
| Square yards..... | | | | 240,237 | 137,003 | 377,240 | 377,240 |
| Salmon..... | 128 | 152 | | | | 280 | 2,440 |
| Square yards..... | 122,131 | 464,138 | | | | 586,269 | 5,625,352 |
| Sea bass..... | | 2 | 4 | | | 6 | 6 |
| Square yards..... | | 1,060 | 11,408 | | | 12,468 | 12,468 |
| Shad..... | | 188 | | | | 188 | 188 |
| Square yards..... | | 627,655 | | | | 627,655 | 627,655 |
| Set: | | | | | | | |
| "California halibut"..... | | | 11 | | | 11 | 11 |
| Square yards..... | | | 54,733 | | | 54,733 | 54,733 |
| Salmon..... | | | | | | | 1,200 |
| Square yards..... | | | | | | | 233,046 |
| Sea bass..... | | | | 31 | 21 | 52 | 52 |
| Square yards..... | | | | 167,487 | 110,061 | 277,548 | 277,548 |
| Miscellaneous..... | 15 | 91 | 82 | 30 | 13 | 231 | 231 |
| Square yards..... | 11,210 | 127,602 | 111,999 | 26,441 | 17,180 | 294,432 | 294,432 |
| Trammel nets..... | | | | 39 | 13 | 52 | 52 |
| Square yards..... | | | | 259,608 | 128,713 | 388,221 | 388,221 |
| Lines: | | | | | | | |
| Trawl, set, and hand..... | 265 | 1,257 | 980 | 1,520 | 1,006 | 5,028 | 31,666 |
| Hooks..... | 45,560 | 140,349 | 137,972 | 281,547 | 55,418 | 660,846 | 1,216,925 |
| Troll..... | 1,056 | 992 | 442 | 274 | 181 | 2,945 | 6,418 |
| Hooks..... | 4,601 | 5,066 | 2,596 | 274 | 181 | 12,718 | 26,357 |
| Pound nets..... | | | | | | | 432 |
| Brush weirs..... | | | | | | | 4 |
| Fish wheels..... | | | | | | | 29 |
| Fyke nets..... | | 2,591 | | | | 2,591 | 2,591 |
| Dip nets..... | 11 | 8 | | | | 19 | 362 |
| Bag nets, shrimp..... | | 11 | | | | 11 | 11 |
| Length, yards..... | | 7,312 | | | | 7,312 | 7,312 |
| Drag bag nets..... | | | | | | | 33 |
| Length, yards..... | | | | | | | 2,642 |
| Reef nets..... | | | | | | | 8 |
| Paranzella nets..... | | 8 | 1 | 4 | | 13 | 13 |
| Yards at mouth..... | | 133 | 17 | 67 | | 217 | 217 |
| Beam trawls..... | | 18 | | | | 18 | 32 |
| Yards at mouth..... | | 120 | | | | 120 | 186 |
| Otter trawls..... | | | | | | | 18 |
| Yards at mouth..... | | | | | | | 227 |
| Traps: | | | | | | | |
| Crab..... | 324 | 4,808 | 47 | | | 5,179 | 18,783 |
| Crawfish..... | | | | | | | 1,500 |
| Lobster..... | | | | 4,791 | 1,308 | 6,099 | 6,099 |
| Octopus..... | | | 51 | | | 51 | 51 |
| Harpoons: | | | | | | | |
| Swordfish and turtles..... | | | | 34 | 18 | 52 | 52 |
| Whales..... | | 2 | | | | 2 | 2 |
| Tongs, rakes, and shovels..... | 16 | 108 | 38 | 42 | | 204 | 3,085 |
| A balone outfits..... | | | 16 | 1 | | 17 | 17 |
| Dredges, oyster..... | | | | | | | 4 |
| Yards at mouth..... | | | | | | | 4 |

Fisheries of the Pacific Coast States, 1933—Continued

CATCH: BY STATES

| Species | Washington | | Oregon | |
|---|----------------------|--------------------|---------------------|--------------------|
| | Pounds | Value | Pounds | Value |
| FISH | | | | |
| Carp..... | 74, 240 | \$2, 227 | 8, 500 | \$170 |
| Cod 1..... | 10, 501, 381 | 106, 423 | | |
| Flounders: | | | | |
| " Sole "..... | 536, 004 | 15, 003 | 34, 412 | 676 |
| Other..... | 62, 027 | 1, 230 | 32, 406 | 433 |
| Hallbut..... | 23, 765, 161 | 1, 527, 349 | 410, 442 | 23, 766 |
| Herring..... | 582, 329 | 5, 823 | 30, 038 | 302 |
| " Lingcod "..... | 669, 644 | 18, 406 | 213, 502 | 3, 911 |
| Perch..... | 20, 178 | 732 | 15, 322 | 193 |
| Pilchard or sardine..... | | | 7, 090 | 71 |
| Rockfishes..... | 301, 621 | 9, 667 | 48, 709 | 859 |
| Sablefish..... | 1, 359, 724 | 41, 898 | 23, 816 | 607 |
| Salmon: | | | | |
| Blueback, red or sockeye..... | 9, 701, 564 | 865, 338 | 104, 849 | 8, 913 |
| Chinook or king..... | 17, 420, 915 | 1, 055, 289 | 13, 123, 423 | 794, 300 |
| Chum or keta..... | 6, 896, 595 | 166, 263 | 1, 174, 333 | 14, 628 |
| Humpback or pink..... | 38, 594, 513 | 900, 887 | 4, 667 | 184 |
| Silver or coho..... | 10, 109, 215 | 411, 364 | 4, 932, 093 | 189, 140 |
| Shad..... | 87, 529 | 1, 751 | 360, 744 | 6, 048 |
| Smelt..... | 1, 285, 648 | 24, 999 | 545, 319 | 11, 985 |
| Steelhead trout..... | 1, 345, 814 | 68, 632 | 1, 356, 421 | 73, 380 |
| Striped bass..... | | | 24, 021 | 1, 038 |
| Sturgeon..... | 39, 234 | 871 | 51, 729 | 1, 284 |
| Tuna and tunalike fishes, albacore..... | | | 1, 980 | 119 |
| Other fish..... | 7, 093 | 141 | | |
| Total..... | 123, 420, 393 | 5, 230, 293 | 22, 503, 816 | 1, 131, 977 |
| SHELLFISH, ETC. | | | | |
| Crabs..... | 1, 114, 424 | 46, 710 | 1, 838, 040 | 55, 996 |
| Crawfish..... | | | 99, 000 | 9, 900 |
| Shrimp..... | 52, 867 | 4, 229 | | |
| Clams: | | | | |
| Hard..... | 555, 067 | 25, 178 | | |
| Razor..... | 540, 271 | 77, 182 | 46, 347 | 6, 621 |
| Mixed..... | | | 15, 965 | 950 |
| Octopus..... | 24, 113 | 1, 108 | | |
| Oysters: | | | | |
| Eastern, market..... | 1, 466 | 844 | | |
| Japanese, market..... | 2, 780, 750 | 178, 858 | 25, 200 | 1, 800 |
| Native, market..... | 219, 968 | 117, 064 | 4, 500 | 2, 600 |
| Scallops..... | 10, 185 | 2, 425 | | |
| Total..... | 5, 309, 111 | 453, 598 | 2, 029, 102 | 77, 867 |
| Grand total..... | 128, 729, 504 | 5, 683, 891 | 24, 532, 918 | 1, 209, 844 |

| Species | California 2 | | Total | |
|-----------------------------|---------------|-------------|---------------|-------------|
| | Pounds | Value | Pounds | Value |
| FISH | | | | |
| Anchovies..... | 317, 292 | \$3, 855 | 317, 292 | \$3, 855 |
| Barracuda..... | 3, 072, 962 | 122, 601 | 3, 072, 962 | 122, 601 |
| Cabrilla..... | 84, 612 | 2, 806 | 84, 612 | 2, 806 |
| Carp..... | 87, 863 | 610 | 140, 516 | 3, 007 |
| Catfish..... | 172, 445 | 19, 849 | 172, 443 | 19, 849 |
| Cod 1..... | 5, 534, 600 | 49, 670 | 16, 035, 981 | 156, 093 |
| Corbina..... | 290 | 23 | 290 | 23 |
| Flounders: | | | | |
| " California halibut "..... | 989, 225 | 63, 328 | 989, 225 | 63, 328 |
| " Sole "..... | 8, 306, 970 | 332, 726 | 8, 877, 386 | 348, 406 |
| Other..... | 1, 074, 732 | 48, 179 | 1, 163, 165 | 44, 842 |
| Flyingfish..... | 16, 396 | 401 | 16, 306 | 501 |
| Grayfish..... | 471, 030 | 8, 645 | 471, 030 | 8, 685 |
| Hake..... | 37, 539 | 361 | 37, 339 | 361 |
| Halibut..... | 321, 664 | 16, 459 | 24, 497, 267 | 1, 567, 574 |
| Hardhead..... | 156, 687 | 7, 684 | 156, 687 | 7, 684 |
| Herring..... | 601, 445 | 3, 031 | 1, 213, 812 | 9, 156 |
| Horse mackerel..... | 1, 010, 800 | 11, 921 | 1, 010, 800 | 11, 921 |
| Kingfish..... | 564, 266 | 12, 153 | 564, 266 | 12, 153 |
| " Lingcod "..... | 1, 088, 955 | 35, 860 | 1, 972, 101 | 57, 677 |
| Mackerel..... | 69, 614, 899 | 420, 148 | 69, 614, 899 | 420, 148 |
| Marlin..... | 6, 859 | 281 | 6, 859 | 281 |
| Mullet..... | 24, 014 | 1, 001 | 24, 014 | 1, 001 |
| Perch..... | 215, 014 | 8, 522 | 250, 514 | 9, 447 |
| Pilchard or sardine..... | 509, 797, 481 | 1, 505, 117 | 509, 804, 571 | 1, 505, 188 |

1 The cod were taken off Alaska.

2 Taken off the Pacific coast including Latin America.

Fisheries of the Pacific Coast States, 1933—Continued

CATCH: BY STATES—Continued

| Species | California | | Total | |
|------------------------------------|---------------|-------------|---------------|--------------|
| | Pounds | Value | Pounds | Value |
| FISH—continued | | | | |
| Pompano..... | 4, 689 | \$1, 326 | 4, 589 | \$1, 326 |
| Rockbass..... | 348, 392 | 15, 432 | 348, 392 | 15, 432 |
| Rockfishes..... | 4, 787, 810 | 153, 219 | 5, 137, 940 | 163, 745 |
| Rudderfish..... | 13, 152 | 547 | 13, 152 | 547 |
| Sablefish..... | 1, 332, 573 | 31, 779 | 2, 716, 113 | 74, 284 |
| Salmon: | | | | |
| Blueback, red or sockeye..... | | | 9, 866, 413 | 874, 251 |
| Chinook or king..... | 4, 569, 808 | 255, 701 | 35, 114, 146 | 2, 105, 290 |
| Chum or keta..... | | | 8, 070, 892 | 180, 891 |
| Humpback or pink..... | | | 38, 599, 180 | 907, 071 |
| Silver or coho..... | | | 15, 041, 308 | 600, 504 |
| Sculpin..... | 68, 425 | 3, 806 | 68, 425 | 3, 806 |
| Sea bass: | | | | |
| Black..... | 449, 294 | 18, 054 | 449, 294 | 18, 054 |
| White..... | 1, 163, 092 | 68, 500 | 1, 163, 092 | 68, 500 |
| Shad..... | 1, 157, 489 | 27, 853 | 1, 605, 762 | 35, 652 |
| Sheepshead..... | 58, 609 | 1, 616 | 58, 609 | 1, 616 |
| Skates..... | 193, 711 | 2, 196 | 193, 711 | 2, 196 |
| Smelt..... | 729, 702 | 27, 443 | 2, 560, 669 | 64, 397 |
| Spanish mackerel..... | 4, 197 | 191 | 4, 197 | 191 |
| Splittail..... | 17, 509 | 199 | 17, 509 | 199 |
| Squawfish..... | 727 | 25 | 727 | 25 |
| Steelhead trout..... | | | 2, 702, 235 | 142, 012 |
| Striped bass..... | 485, 926 | 26, 764 | 509, 947 | 27, 802 |
| Sturgeon..... | | | 90, 963 | 2, 155 |
| Suckers..... | 14, 187 | 147 | 14, 187 | 147 |
| Swordfish..... | 850, 699 | 71, 078 | 850, 699 | 71, 078 |
| Tal..... | 138 | 4 | 138 | 4 |
| Tomcod..... | 729 | 15 | 729 | 15 |
| Tuna and tunalike fishes: | | | | |
| Albacore..... | 487 | 53 | 2, 467 | 172 |
| Bluefin..... | 560, 492 | 29, 352 | 560, 492 | 29, 352 |
| Bonito..... | 2, 252, 199 | 40, 232 | 2, 252, 199 | 40, 232 |
| Skipjack or striped tuna..... | 16, 687, 308 | 613, 091 | 16, 687, 308 | 613, 091 |
| Yellowfin..... | 51, 075, 630 | 2, 274, 831 | 51, 075, 630 | 2, 274, 831 |
| Whitebait..... | 95, 751 | 3, 197 | 95, 751 | 3, 197 |
| Whitefish..... | 95, 053 | 4, 430 | 95, 053 | 4, 430 |
| Yellowtail..... | 3, 898, 888 | 87, 636 | 3, 898, 888 | 87, 636 |
| Other fish..... | 144, 383 | 2, 110 | 151, 476 | 2, 251 |
| Total..... | 694, 598, 850 | 6, 430, 698 | 840, 523, 059 | 12, 792, 968 |
| SHELLFISH, ETC. | | | | |
| Crabs..... | 3, 223, 312 | 252, 934 | 6, 175, 776 | 355, 640 |
| Crawfish..... | | | 99, 000 | 9, 900 |
| Sea crawfish or spiny lobster..... | 1, 049, 905 | 144, 545 | 1, 049, 905 | 144, 545 |
| Shrimp..... | 2, 088, 750 | 31, 498 | 2, 141, 617 | 35, 727 |
| Abalone..... | 551, 268 | 80, 433 | 551, 268 | 80, 433 |
| Clams: | | | | |
| Hard..... | 21, 040 | 5, 321 | 576, 107 | 30, 499 |
| Pismo..... | 26, 472 | 5, 768 | 26, 472 | 5, 768 |
| Razor..... | | | 586, 618 | 83, 803 |
| Soft..... | 62, 834 | 13, 390 | 62, 834 | 13, 390 |
| Mixed..... | | | 15, 965 | 950 |
| Mussels, sea..... | 47 | 23 | 47 | 23 |
| Octopus..... | 31, 521 | 2, 145 | 55, 634 | 3, 253 |
| Oysters: | | | | |
| Eastern, market..... | 58, 419 | 21, 907 | 59, 885 | 22, 751 |
| Japanese, market..... | 26, 824 | 6, 706 | 2, 842, 774 | 187, 364 |
| Native, market..... | 1, 218 | 558 | 225, 736 | 120, 222 |
| Scallops..... | | | 10, 185 | 2, 425 |
| Squid..... | 824, 543 | 13, 939 | 824, 543 | 13, 939 |
| Turtles..... | 2, 901 | 125 | 2, 901 | 125 |
| Total..... | 7, 969, 054 | 579, 292 | 15, 307, 267 | 1, 110, 757 |
| WHALE PRODUCTS | | | | |
| Whale meat..... | 2, 214, 000 | 43, 242 | 2, 214, 000 | 43, 242 |
| Whale oil..... | 2, 116, 890 | 41, 025 | 2, 116, 890 | 41, 025 |
| Total..... | 4, 330, 890 | 84, 267 | 4, 330, 890 | 84, 267 |
| Grand total..... | 706, 898, 794 | 7, 094, 257 | 860, 161, 216 | 13, 987, 992 |

Industries related to the fisheries of the Pacific Coast States, 1933

OPERATING UNITS, SALARIES, AND WAGES

| Item | Washing- ton | Oregon | California | Total |
|-------------------------------------|-----------------|-----------|-------------|-------------|
| | Number | Number | Number | Number |
| Transporting: | | | | |
| Persons engaged..... | 179 | 48 | 18 | 245 |
| Vessels, motor..... | 74 | 26 | 4 | 104 |
| Net tonnage..... | 1,763 | 320 | 245 | 2,328 |
| Wholesale and manufacturing: | | | | |
| Establishments..... | 111 | 57 | 145 | 313 |
| Persons engaged: | | | | |
| Proprietors..... | 105 | 53 | 189 | 347 |
| Salaried employees..... | 205 | 72 | 470 | 747 |
| Wage earners: | | | | |
| Average for season..... | 3,010 | 806 | 7,017 | 10,899 |
| Average for year..... | 865 | 391 | 2,411 | 3,667 |
| Paid to salaried employees..... | \$540,577 | \$192,345 | \$2,277,735 | \$3,010,657 |
| Paid to wage earners..... | \$348,157 | \$456,628 | \$2,280,050 | \$3,084,835 |
| Total salaries and wages..... | \$888,734 | \$648,973 | \$4,557,785 | \$6,095,492 |
| Fishermen manufacturing..... | 38 | | 130 | 168 |

PRODUCTS MANUFACTURED

| Item | Washington | | Oregon | | California | |
|--|------------|-----------|-----------|-----------|------------|-----------|
| | Quantity | Value | Quantity | Value | Quantity | Value |
| By manufacturing firms: | | | | | | |
| Barracuda, fresh fillets..... pounds. | | | | | 50,000 | \$6,000 |
| Cabrilla, fresh fillets..... do. | | | | | 18,000 | 2,700 |
| Cod, salted: | | | | | | |
| Dry..... do. | 315,980 | \$17,846 | | | (1) | (1) |
| Boneless, including absolutely boneless..... pounds. | 1,651,013 | 119,347 | | | (1) | (1) |
| Flounders, fresh fillets..... do. | 126,392 | 15,552 | 15,300 | \$1,800 | 1,779,000 | 252,750 |
| Herring, smoked: | | | | | | |
| Bloaters..... do. | 121,620 | 12,971 | | | | |
| Other smoked and kippered..... pounds. | 6,850 | 484 | | | | |
| Lingcod, fresh fillets..... do. | (1) | (1) | 8,240 | 824 | 130,000 | 18,200 |
| Mackerel: | | | | | | |
| Canned..... standard cases | | | | | 747,542 | 1,858,630 |
| Meal..... tons | | | | | 1,278 | 35,554 |
| Oil..... gallons | | | | | 83,778 | 10,302 |
| Pilchard: | | | | | | |
| Salted..... pounds | | | | | 73,400 | 4,454 |
| Canned, "sardines"..... standard cases | | | | | 1,539,446 | 3,805,168 |
| Meal..... tons | | | | | 50,581 | 1,530,218 |
| Flour..... pounds | | | | | | 11,334 |
| Oil..... gallons | | | | | 10,263,776 | 1,593,088 |
| Rockfishes, fresh fillets..... pounds | | | | | 880,000 | 131,760 |
| Sablefish: | | | | | | |
| Fresh fillets..... do. | (1) | (1) | | | 103,000 | 11,760 |
| Salted..... do. | 143,300 | 8,955 | | | | |
| Kippered..... do. | 183,072 | 22,138 | | | | |
| Salmon: | | | | | | |
| Mild-cured ¹ do. | 3,062,271 | 530,038 | 2,486,800 | 524,154 | 1,022,175 | 214,668 |
| Kippered..... do. | 1,018,687 | 145,732 | (1) | (1) | | |
| Smoked..... do. | 45,904 | 8,591 | (1) | (1) | 220,233 | 78,966 |
| Canned: | | | | | | |
| Chinook or king..... standard cases | 74,567 | 677,206 | 204,320 | 2,052,119 | (1) | (1) |
| Blueback, red or sockeye..... standard cases | 134,080 | 1,658,052 | 6,503 | 88,009 | | |
| Silver or coho..... do. | 46,825 | 294,031 | 33,917 | 243,026 | | |
| Humpback or pink..... do. | 540,172 | 2,361,942 | | | | |
| Chum or keta..... do. | 49,779 | 197,217 | 27,000 | 102,136 | | |
| Steelhead trout..... do. | 3,481 | 28,905 | 14,069 | 132,321 | | |
| Eggs for bait..... do. | 8,637 | 75,281 | 649 | 17,804 | | |
| Meal..... tons | 990 | 29,098 | (1) | (1) | | |
| Oil..... gallons | 116,401 | 14,513 | 35,700 | 9,500 | | |
| Sea bass: | | | | | | |
| Black, fresh fillets..... pounds | | | | | 260,000 | 28,600 |
| White, fresh fillets..... do. | | | | | 85,000 | 16,000 |
| Shad: | | | | | | |
| Canned..... standard cases | | | 1,148 | 3,837 | | |
| Roe, canned..... do. | (1) | (1) | 321 | 7,259 | 71 | 15,787 |
| Sheepshead, fresh fillets..... pounds | | | | | 15,500 | 1,920 |

¹ The production of this item is included under "Unclassified products."

² This item is usually an intermediate product and, although included in the total, may be shown in its final stage of processing in this or another State.

Industries related to the fisheries of the Pacific Coast States, 1933—Continued

PRODUCTS MANUFACTURED—Continued

| Item | Washington | | Oregon | | California | |
|--|------------|-----------|-------------------------|----------------------|-------------|------------|
| | Quantity | Value | Quantity ⁽¹⁾ | Value ⁽¹⁾ | Quantity | Value |
| By manufacturing firms—Continued: | | | | | | |
| Tuna and tunalike fishes: | | | | | | |
| Canned: | | | | | | |
| Albacore..... standard cases..... | | | | | 54,087 | \$286,236 |
| Bluefin..... do..... | | | | | 4,179 | 21,519 |
| Bonito..... do..... | | | | | 42,439 | 170,178 |
| Striped..... do..... | | | | | 225,481 | 1,022,819 |
| "Tonno"..... do..... | | | | | 136,740 | 852,236 |
| Yellowfin..... do..... | | | | | 936,299 | \$,417,676 |
| Yellowtail..... do..... | | | | | 43,918 | 163,766 |
| Meal..... tons..... | | | | | 6,004 | 153,300 |
| Crabs, meat, packaged, fresh-cooked | | | | | | |
| pounds..... | | | 197,083 | \$55,575 | | |
| Shrimp, meal..... tons..... | | | | | 115 | 4,605 |
| Abalone steaks..... pounds..... | | | | | 531,698 | 143,282 |
| Clams, hard: | | | | | | |
| Fresh-shucked..... gallons..... | 2,513 | \$2,681 | | | | |
| Canned: | | | | | | |
| Whole..... standard cases..... | 28,986 | 66,173 | | | | |
| Minced..... do..... | 7,050 | 28,619 | | | | |
| Chowder, juice and cocktail | | | | | | |
| standard cases..... | 4,071 | 9,890 | | | | |
| Clams, razor, canned: | | | | | | |
| Minced..... do..... | 26,767 | 246,042 | 1,165 | 10,095 | | |
| Whole and juice..... do..... | 1,454 | 12,561 | 111 | 883 | | |
| Clam shells, crushed for poultry | | | | | | |
| feed..... tons..... | 1,630 | 15,340 | | | | |
| Oysters: | | | | | | |
| Japanese: | | | | | | |
| Fresh-shucked..... gallons..... | 100,949 | 116,776 | 47,336 | 43,385 | (1) | (1) |
| Canned..... standard cases..... | 30,500 | 149,190 | | | | |
| Eastern and native, fresh- | | | | | | |
| shucked..... gallons..... | 19,607 | 102,622 | 6,843 | 36,762 | 6,625 | 26,600 |
| Oyster-shell products: | | | | | | |
| Poultry feed..... tons..... | (1) | (1) | | | 16,743 | 92,682 |
| Lime..... do..... | | | | | 3,816 | 12,688 |
| Unclassified: | | | | | | |
| Packaged..... pounds..... | \$168,842 | \$35,956 | | (4) | \$1,067,125 | \$175,900 |
| Salted..... do..... | | (4) | | | \$1,226,361 | \$141,350 |
| Smoked..... do..... | | | 723,440 | 74,769 | \$194,350 | \$39,526 |
| Canned, cat and dog food | | | | | | |
| standard cases..... | | | | | 208,780 | 576,596 |
| Canned, other..... do..... | \$477 | \$6,183 | 10,736 | 10,17,374 | 115,063 | 1140,960 |
| Meal and scrap..... tons..... | | | | (4) | 19,419 | 24,410 |
| Oil..... gallons..... | | | | | 273,656 | 41,325 |
| Miscellaneous..... | | 19,559 | | 6,992 | | 358,058 |
| Total..... | | 7,019,992 | | 3,358,674 | | 18,393,519 |
| By fishermen: | | | | | | |
| Cod, green-salted ² pounds..... | 3,255,428 | 106,423 | | | 1,715,726 | 48,270 |
| Cod, tongues..... do..... | | | | | 18,000 | 1,400 |
| Shrimp: | | | | | | |
| Dried..... do..... | | | | | 115,684 | 16,196 |
| Meal..... tons..... | | | | | 114 | 2,280 |
| Total..... | | | | | 1,849,524 | 68,146 |
| Total..... | 3,255,428 | 106,423 | | | 1,849,524 | 68,146 |
| Grand total..... | | 7,126,415 | | 3,358,674 | | 18,461,666 |

¹ The production of this item is included under "Unclassified products."

² This item is usually an intermediate product and, although included in the total, may be shown in its final stage of processing in this or another State.

³ Includes fresh fillets of halibut, lingcod, and sablefish; packaged fresh-cooked crab meat; and fresh-shucked scallops.

⁴ This has been included under "Miscellaneous."

⁵ Includes fresh fillets of grayfish and totuava; swordfish steaks; fresh-cooked shrimp meats; and fresh-shucked Japanese oysters.

⁶ Includes salted anchovies, barracuda, bonito, mackerel, sea bass, black and white, and yellowtail; mild-cured shad; and dry and boneless salt cod.

⁷ Includes smoked salmon, shad, and salmon and shad wieners; and kippered sablefish, salmon, and sturgeon.

⁸ Includes smoked chubs, mackerel, and sablefish, and kippered sablefish.

⁹ Includes canned shad roe, sea cucumbers, and fish balls and loaf.

¹⁰ Includes canned crabs, smoked lingcod, kippered steelhead trout, kippered Chinook salmon cheeks, smoked sturgeon, and tuna.

¹¹ Includes canned anchovies, mackerel stew, salmon, and squid.

¹² Includes miscellaneous meal and scrap.

¹³ Includes whale, sperm, and miscellaneous oil.

¹⁴ Includes salted salmon, spiced herring, salmon-egg meal, and crushed oyster shell for poultry feed.

¹⁵ Includes fresh fillets of rockfishes, salmon meal, and marine-shell novelties.

¹⁶ Includes dried shrimp, concentrated powdered fish meal, marine-shell novelties, liquid glue, agar agar, and kelp products.

NOTE.—The total value of manufactured products in the Pacific Coast States was as follows: By manufacturing establishments, \$28,772,185; and by fishermen, \$174,569. Some of the above products may have been imported from another State or a foreign country; therefore, they cannot be correlated directly with the catch within the State. All of the persons engaged in the preparation of fishermen's manufactured products have also been included as fishermen.

WASHINGTON

Fisheries of Washington, 1933

CATCH: BY DISTRICTS

| Species | Puget Sound district | | Coastal district | | Columbia River district | |
|-------------------------------|----------------------|------------------|------------------|----------------|-------------------------|----------------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | 74,240 | \$2,227 |
| Carp..... | | | | | | |
| Cod ¹ | 10,501,381 | \$106,423 | | | | |
| Flounders: | | | | | 966 | 22 |
| " Sole"..... | 535,038 | 14,981 | | | | |
| Other..... | 62,027 | 1,230 | | | | |
| Hallbut..... | 23,704,547 | 1,623,874 | 1,950 | \$98 | 58,064 | 3,377 |
| Herring..... | 582,320 | 5,823 | | | | |
| " Lingcod"..... | 655,131 | 18,191 | 4,728 | 71 | 9,787 | 144 |
| Perch..... | 20,178 | 732 | | | | |
| Rockfishes..... | 293,811 | 9,540 | 1,340 | 20 | 6,470 | 107 |
| Sablefish..... | 1,358,345 | 41,863 | | | 1,379 | 35 |
| Salmon: | | | | | | |
| Blueback, red or sockeye..... | 8,989,856 | 809,073 | 444,272 | 28,433 | 327,436 | 27,832 |
| Chinook or king..... | 7,262,126 | 461,355 | 1,943,545 | 99,781 | 8,216,244 | 494,153 |
| Chum or keta..... | 5,139,810 | 143,914 | 1,100,919 | 13,761 | 655,830 | 8,588 |
| Humpback or pink..... | 38,539,383 | 906,784 | 5,130 | 103 | | |
| Silver or coho..... | 6,964,821 | 294,171 | 2,358,028 | 86,482 | 786,368 | 30,711 |
| Shad..... | | | | | 87,529 | 1,751 |
| Smelt..... | 170,713 | 10,243 | 60,700 | 2,125 | 1,054,235 | 12,631 |
| Steelhead trout..... | 42,408 | 2,969 | 81,640 | 3,266 | 1,221,766 | 62,397 |
| Sturgeon..... | 319 | 16 | | | 38,915 | 855 |
| Other fish..... | 7,093 | 141 | | | | |
| Total..... | 104,879,316 | 4,351,323 | 6,002,248 | 234,140 | 12,538,829 | 644,830 |
| SHELLFISH | | | | | | |
| Crabs..... | 410,564 | 16,796 | 703,860 | 29,914 | | |
| Shrimp..... | 52,867 | 4,229 | | | | |
| Clams: | | | | | | |
| Hard: | | | | | | |
| Butter..... | 239,031 | 9,903 | | | | |
| Little neck..... | 316,036 | 15,275 | | | | |
| Razor..... | | | 540,271 | 77,182 | | |
| Octopus..... | 24,113 | 1,108 | | | | |
| Oysters: | | | | | | |
| Eastern, market..... | | | 1,466 | 844 | | |
| Japanese, market..... | 1,165,101 | 63,732 | 1,625,649 | 115,126 | | |
| Native, market..... | 212,458 | 114,231 | 7,510 | 2,833 | | |
| Scallops..... | 10,185 | 2,425 | | | | |
| Total..... | 2,430,355 | 227,699 | 2,878,756 | 225,899 | | |
| Grand total..... | 107,309,671 | 4,579,022 | 8,881,004 | 460,039 | 12,538,829 | 644,830 |

¹ The catch of cod were taken off Alaska.

Fisheries of the Puget Sound district of Washington, 1933

OPERATING UNITS: BY GEAR

| Item | Purse seines | Haul seines | Gill nets | | Lines | | Pound nets | Brush weirs | Dip nets |
|-------------------------|--------------|-------------|---------------|-------------|----------------------|------------|------------|-------------|----------|
| | | | Drift, salmon | Set, salmon | Trawl, set, and hand | Troll | | | |
| | | | Number | Number | Number | Number | | | |
| Fishermen: | | | | | | | | | |
| On vessels..... | 1,869 | 13 | | | 1,106 | 176 | 15 | | |
| On boats and shore..... | 25 | 169 | 315 | 4 | 48 | 296 | 198 | 8 | 7 |
| Total..... | 1,894 | 182 | 315 | 4 | 1,151 | 472 | 213 | 8 | 7 |
| Vessels: | | | | | | | | | |
| Steam: | | | | | | | 1 | | |
| Net tonnage..... | | | | | | | 42 | | |
| Motor: | | | | | | | 4 | | |
| Net tonnage..... | 236 | 4 | | | 149 | 100 | 4 | | |
| Sail: | 5,565 | 35 | | | 4,144 | 815 | 119 | | |
| Net tonnage..... | | | | | 3 | | | | |
| Total vessels..... | 236 | 4 | | | 152 | 100 | 5 | | |
| Total net tonnage..... | 5,565 | 35 | | | 5,493 | 815 | 161 | | |
| Boats: | | | | | | | | | |
| Motor..... | 5 | 36 | 296 | 4 | 86 | 216 | 15 | | 3 |
| Other..... | | 23 | 8 | | 163 | | 102 | 4 | 4 |
| Apparatus: | | | | | | | | | |
| Number..... | 241 | 59 | 304 | 4 | 25,215 | 1,580 | 88 | 4 | 7 |
| Length, yards..... | 144,118 | 5,216 | | | | | | | |
| Square yards..... | | | 398,468 | 1,080 | | | | | |
| Hooks..... | | | | | 621,129 | 7,118 | | | |

Fisheries of the Puget Sound district of Washington, 1933—Continued

OPERATING UNITS: BY GEAR—Continued

| Item | Drag bag nets | Reef nets | Beam trawls | Otter trawls | Traps, crab | Tongs and rakes | Shovels | Total, exclusive of duplication |
|-------------------------------|---------------|-----------|-------------|--------------|-------------|-----------------|------------|---------------------------------|
| | Number | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | | |
| On vessels..... | | | 24 | 35 | | 2 | | 3, 112 |
| On boats and shore..... | 83 | 33 | 8 | 2 | 100 | 256 | 346 | 1, 787 |
| Total..... | 83 | 33 | 32 | 37 | 100 | 258 | 346 | 4, 899 |
| Vessels: | | | | | | | | |
| Steam..... | | | 2 | | | | | 3 |
| Net tonnage..... | | | 33 | | | | | 75 |
| Motor..... | | | 8 | 15 | | 1 | | 484 |
| Net tonnage..... | | | 59 | 232 | | 8 | | 10, 421 |
| Sail..... | | | | | | | | 3 |
| Net tonnage..... | | | | | | | | 1, 349 |
| Total vessels..... | | | 8 | 15 | | 1 | | 490 |
| Total net tonnage..... | | | 92 | 232 | | 8 | | 11, 845 |
| Boats: | | | | | | | | |
| Motor..... | 25 | 8 | 6 | 1 | 96 | 32 | | 793 |
| Other..... | 8 | 16 | | | 4 | 114 | | 444 |
| Apparatus: | | | | | | | | |
| Number..... | 29 | 8 | 14 | 16 | 2, 800 | 83 | 346 | |
| Length, yards..... | 2, 382 | | | | | | | |
| Yards at mouth..... | | | 66 | 187 | | | | |

CATCH: BY GEAR

| Species | Purse seines | | Haul seines | | Gill nets | | | |
|-------------------------------|---------------------|--------------------|-----------------|----------------|--------------------|----------------|---------------|------------|
| | Pounds | Value | Pounds | Value | Drift | | Set | |
| | | | | | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | |
| Flounders: | | | | | | | | |
| " Sole "..... | | | 708 | \$20 | | | | |
| Other..... | | | 467 | 7 | | | | |
| Herring..... | | | 92, 900 | 929 | | | | |
| " Lingcod "..... | | | 910 | 23 | | | 5, 345 | \$134 |
| Perch..... | | | 12, 728 | 509 | | | | |
| Rockfishes..... | | | 2, 374 | 68 | | | 305 | 9 |
| Salmon: | | | | | | | | |
| Blueback, red or sockeye..... | 3, 511, 600 | \$316, 044 | 27, 428 | 2, 469 | 63, 383 | \$5, 704 | 2, 973 | 268 |
| Chinook or king..... | 613, 940 | 27, 626 | 10, 626 | 616 | 550, 456 | 31, 926 | | |
| Chum or keta..... | 4, 267, 690 | 119, 495 | 2, 440 | 68 | 192, 990 | 5, 404 | | |
| Humpback or pink..... | 26, 803, 522 | 676, 883 | 35, 442 | 832 | 160, 832 | 3, 780 | 448 | 11 |
| Silver or coho..... | 3, 199, 328 | 111, 976 | 5, 776 | 277 | 338, 266 | 16, 238 | 24 | 1 |
| Smelt..... | | | 112, 692 | 6, 762 | | | | |
| Steelhead trout..... | 1, 080 | 76 | | | | | | |
| Total..... | 40, 397, 130 | 1, 252, 100 | 304, 489 | 12, 581 | 1, 305, 957 | 63, 052 | 9, 095 | 423 |
| SHELLFISH | | | | | | | | |
| Shrimp..... | | | 49 | 4 | | | | |
| Octopus..... | | | 95 | 4 | | | | |
| Total..... | | | 144 | 8 | | | | |
| Grand total..... | 40, 397, 130 | 1, 252, 100 | 304, 633 | 12, 589 | 1, 305, 957 | 63, 052 | 9, 095 | 423 |

Fisheries of the Puget Sound district of Washington, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Lines | | | | Pound nets | | Brush weirs | |
|-------------------------------|-----------------------------------|-----------|-----------|---------|------------|-----------|-------------|---------|
| | Trawl, set, and hand ¹ | | Troll | | Pounds | Value | Pounds | Value |
| | Pounds | Value | Pounds | Value | | | | |
| FISH | | | | | | | | |
| Cod ² | 10,501,381 | \$106,423 | | | | | | |
| Flounders: | | | | | | | | |
| "Solé"..... | 1,727 | 48 | | | | | | |
| Other..... | 1,500 | 22 | | | 126 | \$2 | | |
| Halibut..... | 23,682,511 | 1,522,759 | 20,706 | \$1,035 | 509 | 30 | | |
| Herring..... | | | | | | | 467,200 | \$4,672 |
| "Kingcod" ³ | 506,314 | 16,080 | 11,017 | 165 | 311 | 8 | | |
| Perch..... | 631 | 19 | | | | | | |
| Rockfishes..... | 240,343 | 8,040 | 3,427 | 51 | | | | |
| Sablefish..... | 1,358,295 | 41,861 | | | | | | |
| Salmon: | | | | | | | | |
| Blueback, red or sockeye..... | | | 2,498,758 | 177,412 | 5,366,054 | 482,045 | | |
| Chinook or king..... | | | | | 3,584,328 | 223,551 | | |
| Chum or keta..... | | | | | 674,080 | 18,874 | | |
| Humpback or pink..... | | | 19,400 | 388 | 9,424,212 | 221,469 | | |
| Silver or coho..... | | | 1,451,605 | 71,129 | 1,958,040 | 93,986 | | |
| Steelhead trout..... | | | | | 41,328 | 2,893 | | |
| Starreon..... | | | | | 319 | 16 | | |
| Other fish..... | | | | | 71 | 1 | | |
| Total..... | 36,352,702 | 1,695,252 | 4,004,913 | 250,180 | 21,049,369 | 1,043,775 | 467,200 | 4,672 |
| SHELLFISH | | | | | | | | |
| Octopus..... | 22,179 | 1,020 | | | | | | |
| Grand total..... | 36,374,881 | 1,696,272 | 4,004,913 | 250,180 | 21,049,369 | 1,043,775 | 467,200 | 4,672 |

| Species | Dip nets | | Drag bag nets | | Reef nets | | Beam trawls ⁴ | |
|-------------------------------|----------|-------|---------------|-------|-----------|---------|--------------------------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | |
| Herring..... | 1,829 | \$18 | 20,400 | \$204 | | | | |
| Perch..... | | | 6,414 | 192 | | | | |
| Rockfishes..... | | | 132 | 2 | | | | |
| Salmon: | | | | | | | | |
| Blueback, red or sockeye..... | | | 1,410 | 112 | 17,008 | \$1,531 | | |
| Chinook or king..... | | | 3,630 | 200 | 418 | 24 | | |
| Chum or keta..... | | | | | 2,610 | 73 | | |
| Humpback or pink..... | | | | | 145,527 | 3,420 | | |
| Silver or coho..... | | | | | 11,752 | 564 | | |
| Smelt..... | | | 58,021 | 3,481 | | | | |
| Total..... | 1,829 | 18 | 90,007 | 4,191 | 177,315 | 5,612 | | |
| SHELLFISH | | | | | | | | |
| Shrimp..... | | | | | | | 52,818 | \$4,225 |
| Octopus..... | | | 1,580 | 72 | | | | |
| Scallops..... | | | | | | | 10,185 | 2,425 |
| Total..... | | | 1,580 | 72 | | | 63,003 | 6,650 |
| Grand total..... | 1,829 | 18 | 91,587 | 4,263 | 177,315 | 5,612 | 63,003 | 6,650 |

¹ In addition, the vessels in the Pacific coast halibut fleet landed about 485,000 pounds of halibut, sablefish, and "lingcod" livers at Seattle, which were valued at about \$73,000.

² The cod were taken off Alaska.

³ Does not include a small amount of sea cucumbers taken by beam trawls. The poundage of scallops is based on a yield of 15 percent meat.

Fisheries of the coastal district of Washington, 1933—Continued

CATCH: BY GEAR

| Species | Gill nets | | | | Lines, troll | | Pound nets | | |
|--------------------------|-----------|----------|-----------|----------|--------------|---------|------------|---------|--------|
| | Drift | | Set | | Pounds | Value | Pounds | Value | |
| FISH | Pounds | Value | Pounds | Value | | | | | Pounds |
| Hallbut | | | | | 1,950 | \$98 | | | |
| "Lingcod" | | | | | 4,726 | 71 | | | |
| Rockfishes | | | | | 1,240 | 20 | | | |
| Salmon: | | | 444,272 | \$28,433 | | | | | |
| Blueback, red or sockeye | | | | | | | | | |
| Chinook or king | 473,136 | \$15,140 | 239,131 | 7,652 | 963,794 | 68,429 | 267,485 | \$8,560 | |
| Chum or keta | 137,398 | 1,717 | 325,762 | 4,072 | | | 637,779 | 7,972 | |
| Humpback or pink | | | | | 5,130 | 103 | | | |
| Silver or coho | 388,954 | 12,446 | 522,580 | 16,723 | 787,553 | 36,227 | 668,989 | 21,086 | |
| Steelhead trout | 1,460 | 58 | 71,890 | 2,876 | | | 8,290 | 332 | |
| Total | 1,000,937 | 29,361 | 1,603,625 | 59,756 | 1,764,463 | 104,948 | 1,572,493 | 37,950 | |

| Species | Dip nets | | Drag bag nets | | Traps | | Dredges, tongs, and rakes | | Shovels | |
|---------------------------|----------|-------|---------------|---------|---------|----------|---------------------------|---------|---------|----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | | | |
| Smelt | 15,700 | \$550 | 45,000 | \$1,575 | | | | | | |
| SHELLFISH | | | | | | | | | | |
| Crabs ¹ | | | | | 703,860 | \$29,914 | | | | |
| Clams, razor ² | | | | | | | | | 540,271 | \$77,182 |
| Oysters: ³ | | | | | | | | | | |
| Eastern, market | | | | | | | 1,466 | \$844 | | |
| Japanese, market | | | | | | | 1,625,049 | 115,126 | | |
| Native, market | | | | | | | 7,510 | 2,833 | | |
| Total | | | | | 703,860 | 29,914 | 1,634,625 | 118,803 | 540,271 | 77,182 |
| Grand total | 15,700 | 850 | 45,000 | 1,575 | 703,860 | 29,914 | 1,634,625 | 118,803 | 540,271 | 77,182 |

¹ The poundage of crabs is based on a weight of 20 pounds per dozen.
² The poundage of razor clams is based on a yield of 42 percent of cleaned clam meat.
³ The poundage of oysters is based on a weight of 13 percent oyster meat for eastern oysters, 14 percent for Japanese oysters, and 14 percent for native oysters.

Fisheries of the Columbia River district of Washington, 1933

OPERATING UNITS: BY GEAR

| Item | Haul seines | Gill nets | | Lines | | Pound nets | Fish wheels | Dip nets | Total, exclusive of duplication |
|--------------------|-------------|---------------|-------------|---------------|--------|------------|-------------|----------|---------------------------------|
| | | Drift, salmon | Set, salmon | Trawl and set | Troll | | | | |
| Fishermen: | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| On vessels | | | | 4 | | | | | 4 |
| On boats and shore | 326 | 596 | 65 | 7 | 6 | 182 | 20 | 150 | 1,251 |
| Total | 326 | 596 | 65 | 11 | 6 | 182 | 20 | 150 | 1,255 |
| Vessels, motor | | | | 1 | | | | | 1 |
| Net tonnage | | | | 9 | | | | | 9 |
| Boats: | | | | | | | | | |
| Motor | 20 | 434 | 50 | 7 | 4 | 88 | | 78 | 640 |
| Other | 30 | | 15 | | | 40 | | 6 | 89 |
| Apparatus: | | | | | | | | | |
| Number | 31 | 434 | 134 | 166 | 18 | 204 | 20 | 150 | |
| Length, yards | 9,560 | | | | | | | | |
| Square yards | | 1,206,520 | 34,086 | | | | | | |
| Hooks | | | | 4,725 | 95 | | | | |

Fisheries of the Columbia River district of Washington, 1933—Continued

CATCH: BY GEAR

| Species | Haul seines | | Gill nets | | | | Lines | |
|--------------------------|-------------|---------|-----------|---------|---------|---------|---------------|-------|
| | | | Drift | | Set | | Trawl and set | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | 74,240 | \$2,227 | | | | | | |
| Carp | | | | | | | 906 | \$22 |
| Flounders, "sole" | | | | | | | 58,684 | 3,377 |
| Halibut | | | | | | | 9,787 | 144 |
| "Lingscod" | | | | | | | 6,470 | 107 |
| Rockfishes | | | | | | | 1,379 | 35 |
| Sablefish | | | | | | | | |
| Salmon: | | | | | | | | |
| Blueback, red or sockeye | 8,911 | 757 | 24,334 | \$2,068 | 12,279 | \$1,044 | | |
| Chinook or king | 1,339,380 | 80,095 | 3,665,013 | 219,168 | 81,094 | 4,622 | | |
| Chum or keta | 5,690 | 71 | 483,122 | 6,329 | 18,244 | 239 | | |
| Silver or coho | 37,711 | 1,395 | 249,561 | 10,107 | 11,865 | 415 | | |
| Shad | 54,529 | 1,091 | 26,724 | 534 | | 128 | 3 | |
| Smelt | | | 167,100 | 3,760 | | | | |
| Steelhead trout | 236,594 | 11,830 | 404,115 | 21,256 | 17,742 | 915 | | |
| Sturgeon | 2,272 | 32 | 17,590 | 369 | 1,073 | 32 | 3,182 | 95 |
| Total | 1,759,327 | 97,498 | 5,037,559 | 263,591 | 142,425 | 7,270 | 80,448 | 3,780 |

| Species | Lines | | Pound nets | | Fish wheels | | Dip nets | |
|--------------------------|--------|---------|------------|---------|-------------|----------|-----------|---------|
| | Troll | | | | | | Pounds | Value |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | |
| Salmon: | | | | | | | | |
| Blueback, red or sockeye | | | 31,013 | \$2,638 | 204,406 | \$17,375 | 46,493 | \$3,052 |
| Chinook or king | 27,273 | \$2,057 | 2,345,635 | 140,269 | 478,444 | 35,692 | 278,405 | 12,250 |
| Chum or keta | | | 148,774 | 1,949 | | | | |
| Silver or coho | 51,376 | 2,248 | 430,374 | 16,354 | | | 5,355 | 187 |
| Shad | | | 1,838 | 37 | 4,310 | 86 | | |
| Smelt | | | | | | | 887,135 | 8,871 |
| Steelhead trout | | | 463,120 | 23,712 | 65,200 | 2,934 | 34,995 | 1,750 |
| Sturgeon | | | 3,913 | 76 | 10,749 | 247 | 136 | 4 |
| Total | 78,649 | 4,305 | 3,424,667 | 185,033 | 763,235 | 56,339 | 1,252,519 | 27,014 |

OREGON

Fisheries of Oregon, 1933

CATCH: BY DISTRICTS

| Species | Columbia River district | | Coastal district | |
|--------------------------|-------------------------|---------|------------------|---------|
| | Pounds | Value | Pounds | Value |
| FISH | | | | |
| Carp | 8,500 | \$170 | | |
| Flounders: | | | | |
| "sole" | 9,455 | 236 | 24,957 | \$440 |
| Other | | | 32,406 | 433 |
| Halibut | 181,030 | 10,540 | 229,412 | 13,226 |
| Herring | | | 30,038 | 302 |
| "Lingscod" | 107,794 | 1,684 | 105,708 | 2,227 |
| Perch | | | 15,322 | 193 |
| Pilchard | | | 7,090 | 71 |
| Rockfishes | 11,961 | 259 | 36,748 | 600 |
| Sablefish | 16,181 | 410 | 7,635 | 197 |
| Salmon: | | | | |
| Blueback, red or sockeye | 101,849 | 8,913 | | |
| Chinook or king | 11,500,763 | 699,388 | 1,622,660 | 94,012 |
| Chum or keta | 517,864 | 6,751 | 656,449 | 7,877 |
| Humpback or pink | 1,626 | 32 | 3,041 | 152 |
| Silver or coho | 1,719,855 | 75,420 | 3,212,238 | 113,720 |
| Shad | 127,322 | 2,547 | 233,422 | 3,501 |
| Smelt | 540,968 | 11,798 | 4,351 | 157 |
| Steelhead trout | 1,072,819 | 55,059 | 283,602 | 18,321 |
| Striped bass | | | 24,021 | 1,038 |

Fisheries of Oregon, 1933—Continued

CATCH: BY DISTRICTS—Continued

| Species | Columbia River district | | Coastal district | |
|-----------------------|-------------------------|---------|------------------|---------|
| | Pounds | Value | Pounds | Value |
| FISH—continued | | | | |
| Sturgeon..... | 45,563 | \$1,105 | 6,176 | \$179 |
| Tuna, albacore..... | | | 1,980 | 119 |
| Total..... | 15,966,560 | 874,312 | 6,537,256 | 257,665 |
| SHELLFISH | | | | |
| Crabs..... | | | 1,838,040 | 55,996 |
| Crawfish..... | 99,000 | 9,900 | | |
| Clams: | | | | |
| Razor..... | | | 46,347 | 6,621 |
| Mixed..... | | | 15,965 | 950 |
| Oysters: | | | | |
| Japanese, market..... | | | 25,200 | 1,800 |
| Native, market..... | | | 4,550 | 2,600 |
| Total..... | 99,000 | 9,900 | 1,930,102 | 67,967 |
| Grand total..... | 16,065,560 | 884,212 | 8,467,358 | 325,632 |

Fisheries of the Columbia River district of Oregon, 1933

OPERATING UNITS: BY GEAR

| Item | Haul seines | Gill nets | | Lines | | Pound nets | Dip nets | Traps, crawfish | Total, exclusive of duplication |
|-------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|-----------------|---------------------------------|
| | | Drift, salmon | Set, salmon | Trawl and set | Troll | | | | |
| Fisherman: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| On vessels..... | | | | 8 | 34 | | | | 42 |
| On boats and shore..... | 430 | 1,176 | 52 | 37 | 160 | 38 | 140 | 25 | 2,011 |
| Total..... | 430 | 1,176 | 52 | 45 | 194 | 38 | 140 | 25 | 2,053 |
| Vessels, motor..... | | | | 2 | 19 | | | | 21 |
| Net tonnage..... | | | | 34 | 179 | | | | 213 |
| Boats: | | | | | | | | | |
| Motor..... | 18 | 823 | 46 | 35 | 132 | 20 | 5 | 21 | 1,079 |
| Other..... | 32 | | 6 | 2 | | 9 | | 4 | 51 |
| Apparatus: | | | | | | | | | |
| Number..... | 32 | 823 | 122 | 342 | 755 | 35 | 140 | 1,500 | |
| Length, yards..... | 15,135 | | | | | | | | |
| Square yards..... | | 2,588,335 | 32,330 | | | | | | |
| Hooks..... | | | | 10,450 | 3,397 | | | | |

CATCH: BY GEAR

| Species | Haul seines | | Gill nets | | | | Lines | |
|-------------------------------|-------------|--------|------------|-----------|---------|---------|---------------|--------|
| | | | Drift | | Set | | Trawl and set | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | |
| Carp..... | 8,500 | \$170 | | | | | | |
| Flounders, "sole"..... | | | | | | | 9,456 | \$236 |
| Halibut..... | | | | | | | 171,771 | 10,108 |
| "Lingcod"..... | | | | | | | 47,157 | 961 |
| Rockfishes..... | | | | | | | 11,961 | 259 |
| Sablefish..... | | | | | | | 16,181 | 410 |
| Salmon: | | | | | | | | |
| Blueback, red or sockeye..... | 10,012 | 851 | 46,185 | \$3,926 | 27,224 | \$2,314 | | |
| Chinook or king..... | 974,608 | 58,282 | 8,351,281 | 1,499,407 | 87,452 | 4,985 | | |
| Chum or keta..... | 55,206 | 691 | 422,102 | 5,530 | 966 | 13 | | |
| Silver or coho..... | 69,284 | 2,563 | 272,597 | 11,040 | 4,180 | 146 | | |
| Shad..... | 11,477 | 230 | 114,309 | 2,286 | 460 | 9 | | |
| Smelt..... | | | 511,054 | 11,499 | | | | |
| Steelhead trout..... | 349,820 | 17,491 | 565,846 | 29,537 | 15,695 | 800 | | |
| Sturgeon..... | 441 | 12 | 32,612 | 725 | 2,410 | 72 | 8,008 | 280 |
| Total..... | 1,479,438 | 80,290 | 10,315,986 | 563,950 | 138,387 | 8,339 | 264,533 | 12,234 |

Fisheries of the Columbia River district of Oregon, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Lines | | Pound nets | | Dip nets | | Traps | | | |
|-------------------------------|-----------|---------|------------|--------|----------|--------|--------|--------|---------|-------|
| | Troll | | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | | | |
| Halibut..... | 9,259 | \$432 | | | | | | | | |
| "Lingcod"..... | 60,637 | 703 | | | | | | | | |
| Salmon: | | | | | | | | | | |
| Blueback, red or sockeye..... | | | 10,277 | \$874 | 11,151 | \$948 | | | | |
| Chinook or king..... | 1,329,163 | 98,491 | 307,559 | 18,392 | 450,700 | 19,831 | | | | |
| Chum or keta..... | | | 38,272 | 501 | 1,248 | 16 | | | | |
| Humpback or pink..... | 1,626 | 32 | | | | | | | | |
| Silver or coho..... | 1,247,312 | 56,628 | 123,209 | 4,928 | 3,273 | 115 | | | | |
| Shad..... | | | 1,076 | 22 | | | | | | |
| Smelt..... | | | | | 29,914 | 299 | | | | |
| Steelhead trout..... | 160 | 8 | 121,298 | 6,223 | 20,000 | 1,000 | | | | |
| Sturgeon..... | | | 333 | 9 | 1,749 | 47 | | | | |
| Total..... | 2,648,157 | 156,294 | 602,024 | 30,949 | 618,035 | 22,256 | | | | |
| SHELLFISH | | | | | | | | | | |
| Crawfish..... | | | | | | | | 99,000 | \$9,900 | |
| Grand total..... | 2,648,157 | 156,294 | 602,024 | 30,949 | 618,035 | 22,256 | | 99,000 | 9,900 | |

Fisheries of the coastal district of Oregon, 1933

OPERATING UNITS: BY GEAR

| Item | Haul seines | Gill nets | | Lines | | Otter trawls | Traps, crab | Tongs and rakes | Shovels | Total, exclusive of duplication |
|-------------------------|-------------|---------------|-------------|---------------|-------|--------------|-------------|-----------------|---------|---------------------------------|
| | | Drift, salmon | Set, salmon | Trawl and set | Troll | | | | | |
| Fishermen: | | | | | | | | | | |
| On vessels..... | | | | 20 | 10 | 7 | | | | 33 |
| On boats and shore..... | 5 | 631 | 374 | 18 | 130 | | 278 | 8 | 171 | 1,416 |
| Total..... | 5 | 631 | 374 | 38 | 140 | 7 | 278 | 8 | 171 | 1,449 |
| Vessels, motor: | | | | 5 | 6 | 2 | | | | 12 |
| Net tonnage..... | | | | 66 | 46 | 24 | | | | 120 |
| Boats: | | | | | | | | | | |
| Motor..... | 2 | 485 | 215 | 12 | 97 | | 250 | 2 | | 943 |
| Other..... | 2 | | 132 | | | | 28 | 2 | | 156 |
| Apparatus: | | | | | | | | | | |
| Number..... | 2 | 485 | 750 | 915 | 515 | 2 | 8,340 | 8 | 171 | |
| Length, yards..... | 317 | | | | | | | | | |
| Square yards..... | | 638,260 | 114,000 | | | | | | | |
| Yards at mouth..... | | | | 19,776 | 2,317 | 40 | | | | |
| Hooks..... | | | | | | | | | | |

CATCH: BY GEAR

| Species | Haul seines | | Gill nets | | Lines | | | |
|-----------------------|-------------|-------|-----------|---------|---------------|--------|-----------|--------|
| | Pounds | Value | Pounds | Value | Trawl and set | | Troll | |
| FISH | | | | | | | | |
| Flounders: | | | | | | | | |
| "Sole"..... | | | | | 12,386 | \$248 | 188 | \$3 |
| Other..... | 3,417 | \$61 | 5,063 | \$75 | 5,969 | 96 | | |
| Halibut..... | | | | | 213,733 | 12,407 | 5,122 | 291 |
| Herring..... | 16,732 | 167 | 13,306 | 136 | | | | |
| "Lingcod"..... | | | | | 52,780 | 1,148 | 27,459 | 538 |
| Perch..... | 9,142 | 138 | 6,180 | 55 | | | | |
| Pileard..... | 7,090 | 71 | | | | | | |
| Rockfishes..... | | | 830 | 12 | 31,637 | 514 | 1,914 | 31 |
| Sablefish..... | | | | | 7,210 | 184 | | |
| Salmon: | | | | | | | | |
| Chinook or king..... | | | 1,243,962 | 65,184 | | | 378,698 | 29,728 |
| Chum or keta..... | | | 656,449 | 7,877 | | | | |
| Humpback or pink..... | | | | | | | 3,041 | 152 |
| Silver or coho..... | | | 2,699,736 | 66,772 | | | 1,112,502 | 46,948 |
| Shad..... | | | 283,422 | 3,501 | | | | |
| Smelt..... | 81 | 3 | 4,270 | 154 | | | | |
| Steelhead trout..... | | | 283,602 | 18,321 | | | | |
| Striped bass..... | | | 24,021 | 1,038 | | | | |
| Sturgeon..... | | | 5,910 | 168 | | | | |
| Tuna, albacore..... | | | | | | | 1,980 | 119 |
| Total..... | 36,462 | 430 | 4,576,751 | 163,292 | 323,715 | 14,597 | 1,530,904 | 77,810 |

Fisheries of the coastal district of Oregon, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Otter trawls | | Traps | | Tongs and rakes | | Shovels | |
|--------------------------|--------------|-------|-----------|----------|-----------------|---------|---------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | |
| Flounders: | | | | | | | | |
| " Sole "..... | 12,383 | \$189 | | | | | | |
| Other..... | 17,957 | 211 | | | | | | |
| Halibut..... | 10,557 | 528 | | | | | | |
| "Lingcod"..... | 25,469 | 541 | | | | | | |
| Rockfishes..... | 2,367 | 43 | | | | | | |
| Sablefish..... | 425 | 13 | | | | | | |
| Sturgeon..... | 266 | 11 | | | | | | |
| Total..... | 69,424 | 1,536 | | | | | | |
| SHELLFISH | | | | | | | | |
| Crabs..... | 880 | 46 | 1,837,160 | \$55,950 | | | | |
| Clams: | | | | | | | | |
| Razor ¹ | | | | | | | 46,347 | \$6,621 |
| Mixed ² | | | | | | | 15,965 | 950 |
| Oysters: | | | | | | | | |
| Japanese, market..... | | | | | 25,200 | \$1,800 | | |
| Native, market..... | | | | | 4,550 | 2,600 | | |
| Total..... | 880 | 46 | 1,837,100 | 55,950 | 29,750 | 4,400 | 62,312 | 7,571 |
| Grand total..... | 70,304 | 1,582 | 1,837,100 | 55,950 | 29,750 | 4,400 | 62,312 | 7,571 |

¹ Razor clam poundage is the weight of the steam shucked cleaned meat, which is 42 percent of the round weight.

² Mixed clams consist principally of eastern soft-shelled clams. The weight shown is the fresh-shucked weight which is 21 percent of the round weight.

CALIFORNIA

Fisheries of California, 1933

CATCH: BY DISTRICTS

| Species | Northern district | | San Francisco district | | Monterey district | |
|---------------------------|-------------------|---------|------------------------|---------|-------------------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | |
| Anchovies..... | | | 185,095 | \$2,134 | 90,758 | \$1,161 |
| Barracuda..... | | | | | 29 | 8 |
| Carp..... | | | 57,856 | 610 | | |
| Catfish..... | | | 172,463 | 19,849 | | |
| Cod ¹ | | | 5,534,600 | 49,670 | | |
| Flounders: | | | | | | |
| "California halibut"..... | | | 23,812 | 2,005 | 22,065 | 1,574 |
| " Sole "..... | 3,091,061 | 128,462 | 3,973,339 | 163,847 | 569,440 | 21,572 |
| Other..... | 446,594 | 16,723 | 632,307 | 21,947 | 81,997 | 2,689 |
| Grayfish..... | 3,360 | 17 | 86,456 | 432 | 6,470 | 34 |
| Hake..... | 3,440 | 34 | 25,991 | 260 | 2,642 | 27 |
| Halibut..... | 321,664 | 16,459 | | | | |
| Hardhead..... | | | 156,687 | 7,684 | | |
| Herring..... | | | 544,995 | 2,725 | 51,695 | 236 |
| Horse mackerel..... | | | | | 33,479 | 1,682 |
| Kingfish..... | | | 13,442 | 538 | 189,285 | 5,308 |
| "Lingcod"..... | 491,493 | 14,432 | 396,368 | 13,868 | 188,827 | 6,798 |
| Mackerel..... | | | 218 | 6 | 1,382,754 | 19,789 |
| Perch..... | 4,261 | 106 | 110,675 | 4,376 | 55,399 | 1,589 |
| Pilchard or sardine..... | | | 62,214,480 | 195,674 | 254,311,594 | 784,486 |
| Pompano..... | | | 59 | 27 | 95 | 43 |
| Rock bass..... | | | 5 | 1 | 10 | 1 |
| Rockfishes..... | 486,384 | 18,590 | 654,513 | 22,417 | 1,644,126 | 45,674 |
| Sablefish..... | 483,089 | 12,095 | 121,857 | 3,047 | 238,515 | 3,516 |
| Salmon..... | 3,356,047 | 185,769 | 643,854 | 34,023 | 569,859 | 85,902 |
| Sculpin..... | | | 1,961 | 40 | 2,070 | 21 |
| Sea bass: | | | | | | |
| Black..... | | | | | 30 | 1 |
| White..... | | | 1,306 | 146 | 5,514 | 409 |
| Shad..... | | | 1,157,484 | 27,852 | 5 | 1 |
| Skates..... | 5,623 | 56 | 138,222 | 1,382 | 25,642 | 387 |
| Smelt..... | 32,630 | 1,007 | 316,089 | 12,676 | 117,054 | 4,907 |
| Spittail..... | | | 17,509 | 199 | | |

¹ The catch of cod was taken off Alaska.

NOTE.—The catch of pilchard by floating reduction plants off the coast of California is not included.

Fisheries of California, 1933—Continued

CATCH: BY DISTRICTS—Continued

| Species | Northern district | | San Francisco district | | Monterey district | |
|----------------------------------|-------------------|---------|------------------------|-----------|-------------------|-----------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH—continued | | | | | | |
| Squawfish..... | | | 727 | \$25 | | |
| Striped bass..... | | | 485,926 | 26,764 | | |
| Suckers..... | | | 14,187 | 147 | | |
| Tamcod..... | 438 | \$9 | 291 | 6 | | |
| Tuna and tunalike fishes: | | | | | | |
| Albacore..... | | | | | 497 | \$43 |
| Bonito..... | | | | | 10 | 1 |
| Whitebait..... | 49,460 | 1,736 | 44,682 | 1,372 | 1,609 | 89 |
| Other fish..... | 91,663 | 1,208 | 34,682 | 374 | 11,017 | 244 |
| Total..... | 8,867,277 | 396,705 | 77,662,138 | 616,113 | 259,572,297 | 938,087 |
| SHELLFISH | | | | | | |
| Crabs..... | 126,904 | 7,728 | 2,893,546 | 236,457 | 188,044 | 8,067 |
| Shrimp..... | | | 2,087,952 | 31,320 | 798 | 178 |
| A balone..... | | | | | 444,300 | 69,732 |
| Clams: | | | | | | |
| Hard..... | 113 | 14 | 16,522 | 4,245 | | |
| Pismo..... | | | | | 6,524 | 1,398 |
| Soft..... | 6,557 | 646 | 56,277 | 12,744 | | |
| Octopus..... | 337 | 25 | 8,500 | 602 | 22,535 | 1,512 |
| Oysters, market: | | | | | | |
| Eastern..... | | | 58,419 | 21,907 | | |
| Japanese..... | | | | | 26,824 | 6,706 |
| Native..... | | | 1,218 | 558 | | |
| Squid..... | | | | | 769,695 | 13,007 |
| Total..... | 133,911 | 8,413 | 5,122,524 | 307,833 | 1,458,720 | 100,600 |
| WHALE PRODUCTS | | | | | | |
| Whale meat..... | | | 2,214,000 | 43,242 | | |
| Whale oil..... | | | 2,116,890 | 41,025 | | |
| Total..... | | | 4,330,890 | 84,267 | | |
| Grand total..... | 9,001,188 | 405,118 | 87,115,552 | 1,008,213 | 261,031,017 | 1,038,687 |

| Species | San Pedro district | | | | | |
|---------------------------|--------------------|---------|-------------------|----------|-------------|---------|
| | Off California | | Off Latin America | | Total | |
| FISH | Pounds | Value | Pounds | Value | Pounds | Value |
| Anchovies..... | 41,369 | \$558 | | | 41,369 | \$558 |
| Barracuda..... | 2,433,934 | 93,661 | 126,263 | \$10,925 | 2,560,197 | 104,586 |
| Cabrilla..... | | | 8,275 | 233 | | 233 |
| Flounders: | | | | | | |
| "California halibut"..... | 780,521 | 48,722 | 91 | 8 | 780,612 | 48,730 |
| "Sole"..... | 669,214 | 18,514 | | | 669,214 | 18,514 |
| Other..... | 13,812 | 1,819 | | | 13,812 | 1,819 |
| Flyingfish..... | 16,396 | 501 | | | 16,396 | 501 |
| Grayfish..... | 318,394 | 7,880 | 61 | 2 | 318,455 | 7,882 |
| Hake..... | 5,466 | 40 | | | 5,466 | 40 |
| Herring..... | 190 | 8 | | | 190 | 8 |
| Horse mackerel..... | 976,771 | 10,233 | | | 976,771 | 10,233 |
| Kingfish..... | 359,896 | 6,273 | | | 359,896 | 6,273 |
| "Lingcod"..... | 12,267 | 272 | | | 12,267 | 272 |
| Mackerel..... | 58,922,507 | 344,828 | | | 58,922,597 | 344,828 |
| Marlin..... | 6,253 | 257 | | | 6,253 | 257 |
| Mullet..... | 1,374 | 88 | 423 | 17 | 1,797 | 86 |
| Perch..... | 44,218 | 2,476 | 112 | 11 | 44,330 | 2,487 |
| Pilchard or sardine..... | 192,738,816 | 521,607 | | | 192,738,816 | 521,607 |
| Pompano..... | 3,693 | 1,190 | | | 3,693 | 1,190 |
| Rock bass..... | 200,868 | 9,479 | 12,992 | 829 | 213,860 | 10,308 |
| Rockfishes..... | 1,401,711 | 46,729 | 7,099 | 218 | 1,408,810 | 46,947 |
| Rudderfish..... | 12,813 | 537 | | | 12,813 | 537 |
| Sablefish..... | 487,508 | 13,093 | | | 487,508 | 13,093 |
| Salmon..... | 48 | 7 | | | 48 | 7 |
| Sculpin..... | 50,356 | 3,393 | | | 50,356 | 3,393 |
| Sea bass: | | | | | | |
| Black..... | 46,047 | 2,128 | 172,116 | 7,798 | 218,163 | 9,926 |
| White..... | 632,539 | 37,353 | 39,390 | 2,528 | 671,929 | 39,881 |
| Sheepshead..... | 50,924 | 1,398 | | | 50,924 | 1,398 |
| Skates..... | 23,524 | 413 | | | 23,524 | 413 |
| Smelt..... | 240,943 | 8,104 | | | 240,943 | 8,104 |
| Spanish mackerel..... | | | 3,012 | 145 | 3,012 | 145 |
| Swordfish..... | 575,849 | 50,549 | | | 575,849 | 50,549 |
| Tal..... | | | 138 | 4 | 138 | 4 |

NOTE.—The catch of pilchard by floating reduction plants off the coast of California is not included.

Fisheries of California, 1933—Continued

CATCH: BY DISTRICTS—Continued

| Species | San Pedro district | | | | | |
|------------------------------------|--------------------|------------------|-------------------|----------------|--------------------|------------------|
| | Off California | | Off Latin America | | Total | |
| | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH—continued | | | | | | |
| Tuna and tunalike fishes: | | | | | | |
| Albacore..... | 80 | \$10 | | | 80 | \$10 |
| Bluefin..... | 316,914 | 18,429 | 236,276 | \$10,504 | 553,190 | 28,935 |
| Bonito..... | 919,999 | 12,981 | 186,819 | 4,440 | 1,106,818 | 17,421 |
| Skipjack or striped tuna..... | 10 | 1 | 5,375,294 | 193,248 | 5,375,304 | 193,249 |
| Yellowfin..... | | | 16,029,485 | 708,092 | 16,029,485 | 708,092 |
| Whitefish..... | 39,120 | 2,147 | | | 44,348 | 2,410 |
| Yellowtail..... | 245,890 | 8,766 | 430,875 | 14,018 | 676,765 | 22,784 |
| Other fish..... | 5,992 | 235 | | | 5,992 | 235 |
| Total..... | 262,602,316 | 1,274,654 | 22,633,949 | 953,283 | 285,236,265 | 2,227,937 |
| SHELLFISH | | | | | | |
| Crabs..... | 14,818 | 682 | | | 14,818 | 682 |
| Sea crawfish or spiny lobster..... | 306,687 | 42,775 | 21,298 | 3,731 | 327,985 | 46,506 |
| Abalone..... | 106,968 | 10,701 | | | 106,968 | 10,701 |
| Clams: | | | | | | |
| Hard..... | 4,405 | 1,062 | | | 4,405 | 1,062 |
| Pismo..... | 19,948 | 4,370 | | | 19,948 | 4,370 |
| Mussels..... | 47 | 23 | | | 47 | 23 |
| Octopus..... | 59 | 6 | | | 59 | 6 |
| Squid..... | 54,848 | 932 | | | 54,848 | 932 |
| Total..... | 507,780 | 60,551 | 21,298 | 3,731 | 529,078 | 64,282 |
| Grand total..... | 263,110,096 | 1,335,205 | 22,655,247 | 957,014 | 285,765,343 | 2,292,219 |

| Species | San Diego district | | | | | |
|------------------------------------|--------------------|----------------|-------------------|------------------|-------------------|------------------|
| | Off California | | Off Latin America | | Total | |
| | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | |
| Barracuda..... | 478,189 | \$15,506 | 34,547 | \$2,806 | 512,736 | \$18,012 |
| Cabrilla..... | | | 76,337 | 2,573 | 76,337 | 2,573 |
| Corbina..... | | | 290 | 23 | 290 | 23 |
| Flounders: | | | | | | |
| "California halibut"..... | 78,431 | 5,435 | 84,305 | 5,584 | 162,736 | 11,019 |
| "Ole"..... | 3,916 | 331 | | | 3,916 | 331 |
| Other..... | 22 | 1 | | | 22 | 1 |
| Grayfish..... | 56,289 | 320 | | | 56,289 | 320 |
| Herring..... | 4,665 | 62 | | | 4,665 | 62 |
| Horse mackerel..... | 600 | 6 | | | 600 | 6 |
| Kingfish..... | 1,643 | 34 | | | 1,643 | 34 |
| Mackerel..... | 9,338,361 | 55,504 | 969 | 21 | 9,339,330 | 55,525 |
| Marlin..... | 606 | 24 | | | 606 | 24 |
| Mullet..... | 19,827 | 800 | 2,390 | 116 | 22,217 | 916 |
| Perch..... | 349 | 14 | | | 349 | 14 |
| Pilchard or sardine..... | 532,591 | 3,350 | | | 532,591 | 3,350 |
| Pompano..... | 32 | 10 | 710 | 56 | 742 | 66 |
| Rock bass..... | 129,871 | 4,883 | 4,646 | 239 | 134,517 | 5,122 |
| Rockfishes..... | 556,964 | 18,174 | 36,813 | 1,417 | 593,777 | 19,591 |
| Rudderfish..... | 339 | 10 | | | 339 | 10 |
| Sablefish..... | 1,604 | 28 | | | 1,604 | 28 |
| Sculpin..... | 8,038 | 352 | | | 8,038 | 352 |
| Sea bass: | | | | | | |
| Black..... | 73,505 | 2,811 | 157,596 | 5,516 | 231,101 | 8,127 |
| White..... | 188,783 | 8,518 | 295,560 | 19,546 | 484,343 | 28,064 |
| Sheepshead..... | 7,685 | 223 | | | 7,685 | 223 |
| Skates..... | 700 | 8 | | | 700 | 8 |
| Smelt..... | 22,728 | 736 | 260 | 13 | 22,986 | 749 |
| Spanish mackerel..... | | | 1,185 | 46 | 1,185 | 46 |
| Swordfish..... | 234,787 | 17,685 | 40,063 | 2,844 | 274,850 | 20,529 |
| Tuna and tunalike fishes: | | | | | | |
| Bluefin..... | 7,302 | 419 | | | 7,302 | 419 |
| Bonito..... | 1,047,235 | 20,854 | 98,136 | 1,956 | 1,145,371 | 22,810 |
| Skipjack or striped tuna..... | | | 11,312,004 | 419,842 | 11,312,004 | 419,842 |
| Yellowfin..... | 6,867 | 431 | 35,039,278 | 1,566,308 | 35,046,145 | 1,566,739 |
| Whitefish..... | 42,108 | 1,673 | 8,597 | 347 | 50,705 | 2,020 |
| Yellowtail..... | 987,396 | 22,517 | 2,234,727 | 42,335 | 3,222,123 | 64,852 |
| Other fish..... | 300 | 12 | 729 | 37 | 1,029 | 49 |
| Total..... | 13,831,731 | 180,531 | 49,429,142 | 2,071,325 | 63,260,873 | 2,251,856 |
| SHELLFISH | | | | | | |
| Sea crawfish or spiny lobster..... | 73,788 | 10,578 | 648,132 | 87,461 | 721,920 | 98,039 |
| Turtles..... | | | 2,901 | 125 | 2,901 | 125 |
| Total..... | 73,788 | 10,578 | 651,033 | 87,586 | 724,821 | 98,164 |
| Grand total..... | 13,905,519 | 191,109 | 60,080,175 | 2,158,911 | 63,985,694 | 2,350,020 |

Fisheries of California, 1933—Continued

CATCH: BY WATERS

| Species | Off California † | | Off Latin America | |
|------------------------------------|--------------------|------------------|-------------------|------------------|
| | Pounds | Value | Pounds | Value |
| FISH | | | | |
| Anchovies..... | 317,292 | \$3,855 | | |
| Barracuda..... | 2,912,152 | 109,170 | 160,810 | \$13,431 |
| Cabrilla..... | | | 84,612 | 2,806 |
| Carp..... | 57,856 | 610 | | |
| Catfish..... | 172,463 | 19,849 | | |
| Cod †..... | 5,534,600 | 49,670 | | |
| Corbina..... | | | 290 | 23 |
| Flounders: | | | | |
| "California halibut"..... | 904,829 | 57,736 | 84,396 | 5,592 |
| "Sole"..... | 8,306,970 | 332,726 | | |
| Other..... | 1,074,732 | 43,179 | | |
| Flyingfish..... | 16,396 | 501 | | |
| Grayfish..... | 470,969 | 8,683 | 61 | 2 |
| Hake..... | 37,539 | 361 | | |
| Halibut..... | 321,664 | 16,459 | | |
| Hardhead..... | 156,687 | 7,694 | | |
| Herring..... | 601,445 | 3,031 | | |
| Horse mackerel..... | 1,010,850 | 11,921 | | |
| Kingfish..... | 564,266 | 12,153 | | |
| "Lingcod"..... | 1,088,955 | 35,360 | | |
| Mackerel..... | 69,613,930 | 420,127 | 969 | 27 |
| Marlin..... | 6,859 | 281 | | |
| Mullet..... | 21,201 | 868 | 2,813 | 133 |
| Perch..... | 214,902 | 8,511 | 112 | 11 |
| Pilchard or sardine..... | 509,797,481 | 1,505,117 | | |
| Pompano..... | 3,879 | 1,270 | 710 | 56 |
| Rock bass..... | 330,754 | 14,364 | 17,638 | 1,068 |
| Rockfishes..... | 4,743,698 | 151,584 | 43,912 | 1,635 |
| Rudderfish..... | 13,152 | 547 | | |
| Sablefish..... | 1,332,573 | 31,779 | | |
| Salmon..... | 4,569,808 | 265,701 | | |
| Sculpin..... | 68,425 | 3,806 | | |
| Sea bass: | | | | |
| Black..... | 119,582 | 4,740 | 329,712 | 13,314 |
| White..... | 828,142 | 46,426 | 334,950 | 22,074 |
| Shad..... | 1,157,489 | 27,853 | | |
| Sheepshead..... | 58,609 | 1,618 | | |
| Skates..... | 193,711 | 2,196 | | |
| Smelt..... | 729,442 | 27,430 | 260 | 13 |
| Spanish mackerel..... | | | 4,197 | 191 |
| Spittail..... | 17,509 | 199 | | |
| Squawfish..... | 727 | 25 | | |
| Striped bass..... | 485,926 | 26,764 | | |
| Suckers..... | 14,187 | 147 | | |
| Swordfish..... | 810,636 | 68,234 | 40,063 | 2,844 |
| Tal..... | | | 138 | 4 |
| Tomcod..... | 729 | 15 | | |
| Tuna and tunalike fish: | | | | |
| Albacore..... | 487 | 53 | | |
| Bluefin..... | 324,216 | 18,848 | 236,276 | 10,504 |
| Bonito..... | 1,967,244 | 33,836 | 284,955 | 6,396 |
| Skipjack or striped tuna..... | 10 | 1 | 16,687,298 | 613,090 |
| Yellowfin..... | 6,867 | 431 | 51,068,763 | 2,274,400 |
| Whitebait..... | 95,751 | 3,197 | | |
| Whitefish..... | 81,228 | 3,820 | 13,825 | 610 |
| Yellowtail..... | 1,233,286 | 31,283 | 2,665,602 | 56,353 |
| Other fish..... | 143,664 | 2,073 | 729 | 37 |
| Total..... | 622,535,759 | 3,406,090 | 72,063,091 | 3,024,608 |
| SHELLFISH | | | | |
| Crabs..... | 3,223,312 | 252,934 | | |
| Sea crawfish or spiny lobster..... | 380,475 | 53,353 | 669,430 | 91,102 |
| Shrimp..... | 2,068,750 | 31,498 | | |
| Abalone..... | 551,268 | 80,433 | | |
| Clams: | | | | |
| Hard..... | 21,040 | 5,321 | | |
| Pismo..... | 26,472 | 5,768 | | |
| Soft..... | 62,834 | 13,390 | | |
| Mussels, sea..... | 47 | 23 | | |
| Octopus..... | 31,521 | 2,146 | | |
| Oysters, market: | | | | |
| Eastern..... | 88,419 | 21,907 | | |
| Japanese..... | 26,824 | 6,706 | | |
| Native..... | 1,218 | 558 | | |

† The catch of cod was taken off Alaska.

NOTE.—The catch of pilchard by floating reduction plants off the coast of California is not included.

Fisheries of California, 1933—Continued

CATCH: BY WATERS—Continued

| Species | Off California | | Off Latin America | |
|---------------------|----------------|-------------|-------------------|-------------|
| | Pounds | Value | Pounds | Value |
| SHELLFISH—continued | | | | |
| Squid..... | 824, 543 | \$13, 939 | | |
| Turtles..... | | | 2, 901 | \$125 |
| Total..... | 7, 296, 723 | 487, 975 | 672, 331 | 91, 317 |
| WHALE PRODUCTS | | | | |
| Whale meat..... | 2, 214, 000 | 43, 242 | | |
| Whale oil..... | 2, 116, 890 | 41, 025 | | |
| Total..... | 4, 330, 890 | 84, 267 | | |
| Grand total..... | 634, 163, 372 | 3, 978, 352 | 72, 735, 422 | 3, 115, 925 |

Fisheries of the northern district of California, 1933

OPERATING UNITS: BY GEAR

| Item | Haul seines | Gill nets | | Lines | | Dip nets | Crab traps | Shovels | Total, exclusive of duplication |
|-------------------------|-------------|---------------|---------|--------------|--------|----------|------------|---------|---------------------------------|
| | | Drift, salmon | Other | Set and hand | Troll | | | | |
| Fishermen: | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| On vessels..... | | | | 22 | 30 | | | | 34 |
| On boats and shore..... | 21 | 203 | 18 | 67 | 230 | 11 | 21 | 16 | 455 |
| Total..... | 21 | 203 | 18 | 89 | 260 | 11 | 21 | 16 | 489 |
| Vessels, motor..... | | | | 8 | 14 | | | | 15 |
| Net tonnage..... | | | | 78 | 116 | | | | 132 |
| Boats: | | | | | | | | | |
| Motor..... | | | 15 | 50 | 202 | | 17 | | 202 |
| Other..... | 5 | 128 | | | | | | | 128 |
| Apparatus: | | | | | | | | | |
| Number..... | 5 | 128 | 15 | 265 | 1, 056 | 11 | 324 | 16 | |
| Length, yards..... | 1, 040 | | | | | | | | |
| Square yards..... | | 122, 131 | 11, 210 | | | | | | |
| Hooks..... | | | | 45, 560 | 4, 601 | | | | |

NOTE—The catch by paranzella nets was made entirely by fishermen from the San Francisco district.

CATCH: BY GEAR

| Species | Haul seines | | Gill nets | | Lines | | | |
|------------------|-------------|-------|-----------|---------|--------------|---------|-------------|----------|
| | | | | | Set and hand | | Troll | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | |
| Anchovies..... | | | 70 | \$2 | | | | |
| Flounders: | | | | | | | | |
| "Sole"..... | | | | | 920 | \$13 | | |
| Other..... | | | 3, 702 | 105 | | | | |
| Hallbut..... | | | | | 179, 470 | 9, 177 | 7, 999 | \$420 |
| "Lingcod"..... | | | | | 120, 829 | 2, 265 | 36, 022 | 455 |
| Perch..... | | | 4, 261 | 106 | | | | |
| Rockfishes..... | | | | | 30, 606 | 609 | 511 | 6 |
| Sablefish..... | | | | | 372, 114 | 9, 321 | | |
| Salmon..... | 18, 492 | \$850 | 423, 920 | 21, 192 | | | 2, 913, 635 | 163, 727 |
| Smelt..... | | | 26, 065 | 812 | | | | |
| Other fish..... | | | | | 21, 272 | 503 | 383 | 5 |
| Total..... | 18, 492 | 850 | 458, 018 | 22, 217 | 725, 213 | 21, 888 | 2, 958, 550 | 164, 613 |
| SHELLFISH | | | | | | | | |
| Octopus..... | | | | | 72 | 4 | | |
| Grand total..... | 18, 492 | 850 | 458, 018 | 22, 217 | 725, 285 | 21, 892 | 2, 958, 550 | 164, 613 |

NOTE.—The catch of pilchard by floating reduction plants off the coast of California is not included.

Fisheries of the San Francisco district of California, 1933—Continued

OPERATING UNITS: BY GEAR—Continued

| Item | Fyke nets | Dip nets | Bag nets, shrimp | Paran-zella nets | Beam trawls | Traps, crab | Har-poons, whal-ing | Rakes and tongs | Shov-els | Total, exclu-sive of dupli-cation |
|-------------------------|-----------|----------|------------------|------------------|-------------|-------------|---------------------|-----------------|----------|-----------------------------------|
| | Num-ber | Num-ber | Num-ber | Num-ber | Num-ber | Num-ber | Num-ber | Num-ber | Num-ber | Num-ber |
| Fishermen: | | | | | | | | | | |
| On vessels..... | | | 22 | 76 | | 2 | 16 | | | 388 |
| On boats and shore..... | 69 | 8 | 24 | | 18 | 246 | | 17 | 91 | 1,004 |
| Total..... | 69 | 8 | 46 | 76 | 18 | 248 | 16 | 17 | 91 | 1,392 |
| Vessels: | | | | | | | | | | |
| Steam..... | | | | | | | 2 | | | 2 |
| Net tonnage..... | | | | | | | 41 | | | 41 |
| Motor..... | | | 5 | 16 | | 2 | | | | 41 |
| Net tonnage..... | | | 31 | 228 | | 21 | | | | 821 |
| Sail..... | | | | | | | | | | 2 |
| Net tonnage..... | | | | | | | | | | 824 |
| Total vessels..... | | | 5 | 16 | | 2 | 2 | | | 45 |
| Total net tonnage..... | | | 31 | 228 | | 21 | 41 | | | 1,686 |
| Boats: | | | | | | | | | | |
| Motor..... | 37 | | 6 | | 18 | 244 | | 6 | | 564 |
| Other..... | 19 | | | | | | | 10 | 5 | 92 |
| Apparatus: | | | | | | | | | | |
| Number..... | 2,591 | 8 | 11 | 8 | 18 | 4,808 | 2 | 17 | 91 | |
| Length, yards..... | | | 7,312 | | | | | | | |
| Yards at mouth..... | | | | 133 | 120 | | | | | |

CATCH: BY GEAR

| Species | Purse seines | | Lampara and ring nets | | Haul seines | |
|---------------------------|--------------|-----------|-----------------------|---------|-------------|-------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | |
| Anchovies..... | | | 175,780 | \$1,994 | | |
| Carp..... | | | | | 44,333 | \$438 |
| Flounders..... | | | | | | |
| "California halibut"..... | | | 30 | 2 | | |
| Other..... | | | 192 | 4 | | |
| Hardhead..... | | | | | 138,840 | 6,724 |
| Herring..... | | | 3,420 | 17 | | |
| Kingfish..... | | | 9,865 | 395 | | |
| "Linzcod"..... | | | 153 | 5 | | |
| Mackerel..... | | | 218 | 6 | | |
| Perch..... | | | 272 | 10 | | |
| Pilchard or sardine..... | 35,605,615 | \$115,200 | 26,604,950 | 80,135 | | |
| Pompano..... | | | 59 | 27 | | |
| Rock bass..... | | | 5 | 1 | | |
| Rockfishes..... | | | 43 | 2 | | |
| Sea bass, white..... | | | 46 | 4 | | |
| Shad..... | | | 175 | 5 | | |
| Smelt..... | | | 13,986 | 561 | | |
| Splittail..... | | | | | 13,645 | 137 |
| Suckers..... | | | | | 13,934 | 139 |
| Tomcod..... | | | 40 | 1 | | |
| Whitebait..... | | | 26,123 | 784 | | |
| Total..... | 35,605,615 | 115,200 | 26,835,357 | 84,253 | 210,752 | 7,438 |

Fisheries of the San Francisco district of California, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Gill nets | | Lines | | | |
|---------------------------|-----------|--------|--------------|--------|---------|--------|
| | | | Set and hand | | Troll | |
| FISH | Pounds | Value | Pounds | Value | Pounds | Value |
| Anchovies..... | 9,315 | \$140 | | | | |
| Carp..... | 8,529 | 116 | 118 | \$2 | | |
| Catfish..... | 141 | 17 | 2,341 | 235 | | |
| Cod..... | | | 5,534,007 | 49,670 | | |
| Flounders: | | | | | | |
| "California halibut"..... | | | | 574 | 351 | \$28 |
| "Sole"..... | | | 7,341 | 294 | | |
| Other..... | 475 | 10 | 1,280 | 27 | | |
| Grayfish..... | | | 17,155 | 86 | | |
| Herring..... | 541,575 | 2,708 | | | | |
| "Lingcod"..... | 83 | 3 | 295,763 | 10,352 | 1,009 | 20 |
| Perch..... | 108,599 | 4,294 | 1,284 | 51 | | |
| Pilchard or sardine..... | 3,915 | 39 | | | | |
| Rockfishes..... | | | 362,771 | 12,084 | 184 | 4 |
| Sablefish..... | | | 78,190 | 1,905 | | |
| Salmon..... | 450,314 | 22,585 | | | 193,540 | 11,438 |
| Sculpin..... | | | 1,961 | 40 | | |
| Sea bass, white..... | 1,260 | 142 | | | | |
| Shad..... | 1,157,309 | 27,847 | | | | |
| Smelt..... | 299,631 | 11,994 | | | | |
| Splittail..... | 325 | 4 | | | | |
| Squawfish..... | 545 | 19 | 88 | 3 | | |
| Striped bass..... | 426,006 | 20,079 | 59,920 | 6,685 | | |
| Whitebait..... | 16,991 | 510 | | | | |
| Other fish..... | 36 | 1 | 1,251 | 25 | 25 | 1 |
| Total..... | 3,025,049 | 90,506 | 6,362,637 | 81,505 | 195,100 | 11,491 |
| SHELLFISH | | | | | | |
| Octopus..... | | | 8,568 | 600 | | |
| Grand total..... | 3,025,049 | 90,508 | 6,371,205 | 82,105 | 195,100 | 11,491 |

| Species | Fyke nets | | Dip nets | | Bag nets | |
|------------------|-----------|--------|----------|-------|-----------|----------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | |
| Carp..... | 4,876 | \$54 | | | | |
| Catfish..... | 169,981 | 19,697 | | | | |
| Hardhead..... | 17,847 | 960 | | | | |
| Smelt..... | | | 2,472 | \$121 | | |
| Splittail..... | 3,539 | 58 | | | | |
| Squawfish..... | 94 | 3 | | | | |
| Suckers..... | 253 | 8 | | | | |
| Whitebait..... | | | 1,568 | 78 | | |
| Other fish..... | 803 | 21 | | | | |
| Total..... | 197,393 | 20,701 | 4,040 | 199 | | |
| SHELLFISH | | | | | | |
| Shrimp..... | | | | | 1,494,768 | \$22,422 |
| Grand total..... | 197,393 | 20,701 | 4,040 | 199 | 1,494,768 | 22,422 |

| Species | Paranzella nets | | Beam trawls | | Traps | |
|---------------------------|-----------------|---------|-------------|---------|-----------|-----------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | |
| Flounders: | | | | | | |
| "California halibut"..... | 22,857 | \$1,929 | | | | |
| "Sole"..... | 3,965,998 | 163,553 | | | | |
| Other..... | 530,300 | 21,906 | | | | |
| Grayfish..... | 69,301 | 346 | | | | |
| Hake..... | 25,991 | 260 | | | | |
| Kingfish..... | 3,577 | 143 | | | | |
| "Lingcod"..... | 99,340 | 3,478 | | | | |
| Perch..... | 520 | 21 | | | | |
| Rockfishes..... | 291,515 | 10,327 | | | | |
| Sablefish..... | 45,667 | 1,142 | | | | |
| Skates..... | 138,222 | 1,382 | | | | |
| Tomcod..... | 251 | 5 | | | | |
| Other fish..... | 32,667 | 326 | | | | |
| Total..... | 5,226,186 | 204,818 | | | | |
| SHELLFISH | | | | | | |
| Crabs..... | 24,366 | 1,991 | | | 2,869,180 | \$234,466 |
| Shrimp..... | | | 593,184 | \$8,898 | | |
| Octopus..... | 22 | 2 | | | | |
| Total..... | 24,388 | 1,993 | 593,184 | 8,898 | 2,869,180 | 234,466 |
| Grand total..... | 5,250,574 | 206,811 | 603,184 | 8,898 | 2,869,180 | 234,466 |

Fisheries of the San Francisco district of California, 1933—Continued

CATCH: BY GEAR—Continued

| Species | Harpoons | | Rakes and tongs | | Shovels | |
|-----------------------|-----------|----------|-----------------|----------|---------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| SHELLFISH | | | | | | |
| Clams: | | | | | | |
| Hard..... | | | | | 16,522 | \$4,245 |
| Soft..... | | | | | 56,277 | 12,744 |
| Oysters, market: | | | | | | |
| Eastern..... | | | 58,419 | \$21,907 | | |
| Native..... | | | 1,218 | 558 | | |
| Total..... | | | 59,637 | 22,465 | 72,799 | 16,989 |
| WHALE PRODUCTS | | | | | | |
| Whale meat..... | 2,214,000 | \$43,242 | | | | |
| Whale oil..... | 2,116,890 | 41,025 | | | | |
| Total..... | 4,330,890 | 84,267 | | | | |
| Grand total..... | 4,330,890 | 84,267 | 59,637 | 22,465 | 72,799 | 16,989 |

Fisheries of the Monterey district of California, 1933

OPERATING UNITS: BY GEAR

| Item | Purse seines, sardine | Lampara and ring nets | | Gill nets | | | Lines | |
|-------------------------|-----------------------|-----------------------|--------|-----------------|---------------------------|---------|--------------|--------|
| | | Sardine | Squid | Drift, sea bass | Set, "California halibut" | Other | Set and hand | Troll |
| Fishermen: | Number | Number | Number | Number | Number | Number | Number | Number |
| On vessels..... | 334 | 234 | 50 | | | 2 | | 5 |
| On boats and shore..... | | 136 | 139 | 8 | 20 | 79 | 229 | 193 |
| Total..... | 334 | 370 | 189 | 8 | 20 | 81 | 229 | 208 |
| Vessels, motor..... | 33 | 21 | 8 | | | 1 | | 4 |
| Net tonnage..... | 1,293 | 229 | 67 | | | 20 | | 46 |
| Boats: | | | | | | | | |
| Motor..... | | 12 | 26 | 4 | 11 | 39 | 153 | 182 |
| Other..... | | | | | | 17 | 52 | |
| Apparatus: | | | | | | | | |
| Number..... | 33 | 33 | 34 | 4 | 11 | 82 | 980 | 442 |
| Length, yards..... | 10,960 | 10,101 | 6,507 | 11,408 | 54,733 | 111,999 | | |
| Square yards..... | | | | | | | | |
| Hooks..... | | | | | | | 137,972 | 2,596 |

| Item | Paran-zella nets | Trape | | Rakes and shovels | Abalone outfits | Total, exclusive of duplication |
|-------------------------|------------------|--------|---------|-------------------|-----------------|---------------------------------|
| | | Crab | Octopus | | | |
| Fishermen: | Number | Number | Number | Number | Number | Number |
| On vessels..... | 4 | | | | 58 | 538 |
| On boats and shore..... | 2 | 12 | 5 | 38 | 20 | 539 |
| Total..... | 6 | 12 | 5 | 38 | 78 | 1,077 |
| Vessels, motor..... | 1 | | | | 12 | 60 |
| Net tonnage..... | 21 | | | | 90 | 1,560 |
| Boats: | | | | | | |
| Motor..... | 1 | 5 | 3 | 7 | 4 | 208 |
| Other..... | | 5 | | 14 | | 63 |
| Apparatus: | | | | | | |
| Number..... | 1 | 47 | 51 | 38 | 16 | |
| Yards at mouth..... | 17 | | | | | |

Fisheries of the Monterey district of California, 1933—Continued

CATCH: BY GEAR

| Species | Purse seines | | Lampara and ring nets | | Gill nets | | Lines—Set and hand | |
|----------------------------------|--------------------|----------------|-----------------------|----------------|----------------|---------------|--------------------|---------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | |
| Anchovies | | | 90,643 | \$1,144 | 115 | \$17 | | |
| Barracuda | | | 5 | 1 | 24 | 2 | | |
| Flounders: | | | | | | | | |
| "California halibut" | 62 | \$4 | 343 | 24 | 18,364 | 1,312 | 2,640 | \$188 |
| "Sole" | | | 135 | 5 | 6,507 | 244 | 33,420 | 1,253 |
| Other | | | 865 | 31 | 13,585 | 406 | 19,488 | 687 |
| Grayfish | | | | | | | 2,285 | 12 |
| Hake | | | | | | | 492 | 5 |
| Herring | | | 48,260 | 203 | 3,335 | 33 | | |
| Horse mackerel | | | 31,198 | 1,567 | 549 | 28 | 1,732 | 87 |
| Kingfish | 80 | 2 | 44,581 | 1,252 | 94,444 | 2,650 | 47,867 | 1,344 |
| "Lingcod" | | | 175 | 6 | 3,525 | 127 | 164,698 | 5,943 |
| Mackerel | 255,933 | 3,744 | 790,217 | 11,559 | 921 | 14 | 305,683 | 4,472 |
| Perch | | | 18,617 | 532 | 25,433 | 714 | 11,049 | 277 |
| Pilchard or sardine | 198,802,431 | 618,279 | 55,496,755 | 106,108 | 12,408 | 99 | | |
| Pompano | | | 68 | 31 | 27 | 12 | | |
| Rock bass | | | | | 10 | 1 | | |
| Rockfishes | | | 1,151 | 40 | 2,935 | 80 | 1,612,494 | 44,580 |
| Sablefish | | | | | | | 233,880 | 3,448 |
| Sculpin | | | | | | | 2,070 | 21 |
| Sea bass: | | | | | | | | |
| Black | | | 30 | 1 | | | | |
| White | | | 199 | 15 | 4,930 | 365 | 385 | 29 |
| Shad | | | | | 5 | 1 | | |
| Skates | | | 490 | 7 | 2,604 | 36 | 14,978 | 207 |
| Smelt | 820 | 34 | 26,492 | 1,110 | 74,984 | 3,144 | 14,758 | 619 |
| Tuna and tunalike fishes, bonito | | | | | 10 | 1 | | |
| Whitebait | | | 940 | 52 | 669 | 37 | | |
| Other fish | | | 335 | 12 | | | 1,082 | 40 |
| Total | 199,059,326 | 622,063 | 56,551,499 | 183,700 | 265,404 | 9,323 | 2,469,001 | 63,212 |
| SHELLFISH | | | | | | | | |
| Crabs | | | | | 160,394 | 6,881 | | |
| Shrimp | | | | | 2 | 1 | | |
| Octopus | | | | | | 94 | 1,139 | 76 |
| Squid | 390 | 7 | 763,739 | 12,906 | 5,566 | | | |
| Total | 390 | 7 | 763,739 | 12,906 | 165,962 | 6,976 | 1,139 | 76 |
| Grand total | 199,059,716 | 622,070 | 57,315,238 | 196,606 | 431,366 | 16,299 | 2,470,140 | 63,288 |

| Species | Lines, troll | | Paranzella nets | | Traps | | Rakes and shovels | | Abalone outfits | |
|------------------------------------|----------------|---------------|-----------------|---------------|---------------|--------------|-------------------|--------------|-----------------|---------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | | | |
| Flounders: | | | | | | | | | | |
| "California halibut" | 184 | \$13 | 472 | \$33 | | | | | | |
| "Sole" | | | 529,378 | 20,079 | | | | | | |
| Other | | | 48,069 | 1,565 | | | | | | |
| Grayfish | | | 4,185 | 22 | | | | | | |
| Hake | | | 2,150 | 22 | | | | | | |
| Kingfish | | | 2,313 | 60 | | | | | | |
| "Lingcod" | 1,692 | 61 | 18,737 | 661 | | | | | | |
| Perch | | | 300 | 16 | | | | | | |
| Rockfishes | 2,127 | 65 | 25,399 | 909 | | | | | | |
| Sablefish | | | 4,635 | 68 | | | | | | |
| Salmon | 569,859 | 35,902 | | | | | | | | |
| Skates | | | 7,570 | 87 | | | | | | |
| Tuna and tunalike fishes, albacore | 407 | 43 | | | | | | | | |
| Other fish | | | 9,600 | 192 | | | | | | |
| Total | 574,269 | 36,084 | 652,798 | 23,705 | | | | | | |
| SHELLFISH | | | | | | | | | | |
| Crabs | | | 96 | 4 | 27,554 | \$1,182 | | | | |
| Shrimp | | | | | 796 | 177 | | | | |
| Abalone | | | | | | | | | 444,300 | \$69,732 |
| Clams, Pismo | | | | | | | 6,524 | \$1,398 | | |
| Octopus | | | | | 21,396 | 1,436 | | | | |
| Oysters, market, Japanese | | | | | | | 26,824 | 6,706 | | |
| Total | | | 96 | 4 | 49,746 | 2,795 | 33,348 | 8,104 | 444,300 | 69,732 |
| Grand total | 574,269 | 36,084 | 652,894 | 23,709 | 49,746 | 2,795 | 33,348 | 8,104 | 444,300 | 69,732 |

Fisheries of the San Pedro district of California, 1933

OPERATING UNITS: BY GEAR

| Item | Purse seines | | | Lampara and ring nets | | | Haul seines | Gill nets | | |
|-------------------------|--------------|----------|--------|-----------------------|----------|--------|-------------|------------------|---------------|--------|
| | Bar-racuda | Sar-dine | Tuna | Mack-erel | Sar-dine | Other | | Drift, barracuda | Set, sea bass | Other |
| | Number | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | | | | |
| On vessels..... | 178 | 825 | 477 | 453 | 279 | 4 | | 25 | 19 | 7 |
| On boats and shore..... | | | | 77 | | | 2 | 44 | 59 | 38 |
| Total..... | 178 | 825 | 477 | 530 | 279 | 29 | 2 | 69 | 78 | 45 |
| Vessels, motor..... | 19 | 80 | 47 | 48 | 28 | 1 | | 9 | 7 | 3 |
| Net tonnage..... | 533 | 3,439 | 2,252 | 899 | 501 | 5 | | 56 | 69 | 30 |
| Boats: | | | | | | | | | | |
| Motor..... | | | | 10 | | 6 | | 17 | 22 | 16 |
| Other..... | | | | | | | 1 | | 2 | 6 |
| Apparatus: | | | | | | | | | | |
| Number..... | 19 | 80 | 47 | 58 | 28 | 7 | 1 | 26 | 31 | 30 |
| Length, yards..... | 7,942 | 30,240 | 27,129 | 23,086 | 12,437 | 2,220 | 214 | | | |
| Square yards..... | | | | | | | | 240,237 | 167,487 | 26,441 |

| Item | Trammel nets | Lines | | Paran-zella nets | Traps, lobster | Har-poons, sword-fish | Rakes and shovels | Abalone outfits | Total, exclusive of duplication |
|-------------------------|--------------|--------------|--------|------------------|----------------|-----------------------|-------------------|-----------------|---------------------------------|
| | | Set and hand | Troll | | | | | | |
| | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | | | |
| On vessels..... | 21 | 344 | | 10 | 28 | 12 | | | 1,610 |
| On boats and shore..... | 78 | 338 | 67 | 14 | 174 | 69 | 42 | | 666 |
| Total..... | 99 | 682 | 67 | 24 | 202 | 81 | 42 | 4 | 2,276 |
| Vessels, motor..... | 8 | 40 | | 3 | 13 | 4 | | 1 | 184 |
| Net tonnage..... | 51 | 2,825 | | 41 | 108 | 30 | | 9 | 6,862 |
| Boats: | | | | | | | | | |
| Motor..... | 30 | 215 | 64 | 5 | 99 | 30 | | | 315 |
| Other..... | 1 | 28 | | | 24 | | 1 | | 49 |
| Apparatus: | | | | | | | | | |
| Number..... | 39 | 1,520 | 274 | 4 | 4,791 | 34 | 42 | 1 | |
| Square yards..... | 259,508 | | | | | | | | |
| Yards at mouth..... | | | | 67 | | | | | |
| Hooks..... | | 281,547 | 274 | | | | | | |

CATCH OFF CALIFORNIA: BY GEAR

| Species | Purse seines | | Lampara and ring nets | | Haul seines | | Gill nets | |
|---------------------------|--------------|----------|-----------------------|---------|-------------|-------|-----------|----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | |
| Anchovies..... | | | 41,369 | \$568 | | | | |
| Barracuda..... | 545,131 | \$28,167 | 770,415 | \$3,802 | | | 789,924 | \$30,846 |
| Flounders: | | | | | | | | |
| "California halibut"..... | 134 | 9 | 774 | 60 | | | 998 | 59 |
| " Sole "..... | | | 39 | 2 | | | 33 | 1 |
| Other..... | | | 9 | 1 | | | | |
| Flyingfish..... | 150 | 5 | | | | | 16,246 | 496 |
| Grayfish..... | 1,890 | 59 | 3,207 | 87 | | | 69,875 | 1,573 |
| Herring..... | | | | | | | | 8 |
| Horse mackerel..... | 51,094 | 718 | 914,124 | 9,290 | | | 3,933 | 46 |
| Kingfish..... | 145 | 3 | 193,915 | 3,397 | | | 235 | 4 |
| " Lingcod "..... | | | | | | | 30 | 1 |
| Mackerel..... | 756,439 | 4,558 | 54,189,423 | 306,124 | | | 191,616 | 1,445 |
| Mullet..... | | | | | 38 | \$2 | | 66 |
| Perch..... | 896 | 35 | 32,296 | 1,941 | | | 6,926 | 307 |
| Pilchard or sardine..... | 139,550,505 | 380,056 | 53,187,182 | 141,526 | | | 1,129 | 25 |
| Pompano..... | | | 3,672 | 1,185 | | | 21 | 5 |
| Rock bass..... | 15,197 | 749 | 23,105 | 967 | | | 4,705 | 203 |

Fisheries of the San Pedro district of California, 1935—Continued

CATCH OFF CALIFORNIA: BY GEAR—Continued

| Species | Purse seines | | Lampara and ring nets | | Haul seines | | Gill nets | |
|---------------------------|----------------------|-----------------|-----------------------|-----------------|----------------|------------|--------------------|----------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH—continued | | | | | | | | |
| Rockfishes..... | 36 | \$2 | 134 | \$5 | | | 315 | \$13 |
| Rudderfish..... | 227 | 9 | 4, 170 | 112 | | | 7, 482 | 384 |
| Sculpin..... | | | 100 | 4 | | | | |
| Sea bass: | | | | | | | | |
| Black..... | 522 | 20 | 980 | 36 | | | 2, 905 | 138 |
| White..... | 104, 868 | 6, 888 | 315, 665 | 15, 944 | | | 198, 640 | 13, 421 |
| Sheepshead..... | | | 50 | 2 | | | 152 | * |
| Skates..... | | | 90 | 1 | | | | |
| Smelt..... | 1, 142 | 34 | 163, 804 | 5, 377 | 15, 036 | \$337 | 66, 400 | 2, 701 |
| Tuna and tunalike fishes: | | | | | | | | |
| Bluefin..... | 277, 967 | 16, 077 | 38, 879 | 2, 346 | | | 62 | 5 |
| Bonito..... | 141, 433 | 2, 073 | 743, 802 | 10, 365 | | | 26, 377 | 418 |
| Yellowtail..... | 141, 079 | 5, 068 | 30, 215 | 943 | | | 895 | 42 |
| Other fish..... | 200 | 5 | 613 | 16 | | | 175 | 7 |
| Total..... | 141, 589, 055 | 442, 535 | 110, 658, 134 | 524, 091 | 15, 074 | 339 | 1, 380, 490 | 51, 718 |
| SHELLFISH | | | | | | | | |
| Squid..... | 10, 265 | 159 | 43, 655 | 757 | | | 310 | 4 |
| Grand total..... | 141, 599, 320 | 442, 694 | 110, 701, 789 | 524, 848 | 15, 074 | 339 | 1, 380, 800 | 51, 722 |

| Species | Trammel nets | | Lines | | | | Parantella nets | |
|------------------------------------|-----------------|----------------|--------------------|-----------------|-----------------|---------------|-----------------|----------------|
| | | | Set and hand | | Troll | | | |
| | | | Pounds | Value | Pounds | Value | | |
| FISH | | | | | | | | |
| Barracuda..... | | | 177, 607 | \$7, 378 | 150, 857 | \$5, 468 | | |
| Flounders: | | | | | | | | |
| "California halibut"..... | 407, 953 | \$27, 701 | 43, 655 | 2, 891 | 175 | 11 | 326, 832 | \$17, 091 |
| "Sole"..... | 5, 114 | 218 | 7, 408 | 311 | | | 656, 620 | 17, 982 |
| Other..... | 274 | 17 | 12, 904 | 1, 770 | | | 625 | 31 |
| Grayfish..... | 51, 177 | 1, 268 | 190, 271 | 4, 854 | 80 | 2 | 1, 894 | 37 |
| Hake..... | 150 | 1 | 5, 316 | 39 | | | | |
| Horse mackerel..... | | | 7, 620 | 179 | | | | |
| Kingfish..... | 100 | 2 | 165, 350 | 2, 864 | | | | |
| "Lingcod"..... | 35 | 1 | 12, 202 | 270 | | | | |
| Mackerel..... | | | 3, 784, 946 | 32, 699 | 273 | 2 | | |
| Perch..... | 24 | 2 | 3, 893 | 180 | | | | |
| Rock bass..... | 93 | 4 | 116, 212 | 5, 551 | 337 | 13 | 169 | 10 |
| Rockfishes..... | 408 | 14 | 1, 399, 808 | 46, 638 | | | 555 | 20 |
| Rudderfish..... | | | 934 | 32 | | | | |
| Sablefish..... | | | 487, 508 | 13, 093 | | | | |
| Salmon..... | | | | | 48 | 7 | | |
| Sculpin..... | 269 | 12 | 54, 489 | 3, 273 | | | 60 | 5 |
| Sea bass: | | | | | | | | |
| Black..... | 713 | 26 | 40, 927 | 1, 908 | | | | |
| White..... | 648 | 61 | 12, 668 | 1, 037 | 50 | 2 | | |
| Sheepshead..... | 1, 019 | 29 | 16, 399 | 463 | | | | |
| Skates..... | 6, 352 | 116 | 9, 625 | 161 | | | 7, 457 | 135 |
| Smelt..... | | | 4, 631 | 155 | | | | |
| Swordfish..... | | | 1, 064 | 94 | | | | |
| Tuna and tunalike fishes: | | | | | | | | |
| Albacore..... | | | 40 | 4 | | | | |
| Bluefin..... | | | 16 | 1 | | | | |
| Bonito..... | 620 | 7 | 6, 343 | 96 | 1, 324 | 22 | | |
| Skipjack or striped tuna..... | | | 10 | 1 | | | | |
| Whitefish..... | | | 37, 741 | 2, 071 | | | | |
| Yellowtail..... | 30 | 1 | 68, 817 | 2, 536 | 4, 854 | 176 | | |
| Other fish..... | 169 | 12 | 4, 750 | 193 | 85 | 2 | | |
| Total..... | 475, 148 | 29, 492 | 6, 673, 114 | 130, 762 | 158, 083 | 5, 705 | 994, 212 | 36, 211 |
| SHELLFISH | | | | | | | | |
| Sea crawfish or spiny lobster..... | 3, 852 | 569 | | | | | | |
| Octopus..... | | | 37 | 4 | | | | |
| Squid..... | 618 | 12 | | | | | | |
| Total..... | 4, 470 | 581 | 37 | 4 | | | | |
| Grand total..... | 479, 618 | 30, 073 | 6, 673, 151 | 130, 766 | 158, 083 | 5, 705 | 994, 212 | 36, 217 |

Fisheries of the San Pedro district of California, 1933—Continued

CATCH OFF CALIFORNIA: BY GEAR—Continued

| Species | Traps | | Harpoons | | Rakes and shovels | | Abalone outfits | |
|--|-----------------|----------------|-----------------|----------------|-------------------|---------------|-----------------|----------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | |
| Kingfish..... | 151 | \$3 | | | | | | |
| Marlin..... | | | 6, 253 | \$257 | | | | |
| Perch..... | 181 | 11 | | | | | | |
| Rock bass..... | 41, 050 | 1, 982 | | | | | | |
| Rockfishes..... | 455 | 17 | | | | | | |
| Sculpin..... | 1, 438 | 99 | | | | | | |
| Sheepshead..... | 33, 304 | 895 | | | | | | |
| Swordfish..... | | | 574, 755 | 50, 455 | | | | |
| Tuna and tunalike fishes, albacore..... | | | 40 | 6 | | | | |
| Whitefish..... | 1, 379 | 76 | | | | | | |
| Total..... | 77, 958 | 3, 083 | 581, 048 | 50, 718 | | | | |
| SHELLFISH | | | | | | | | |
| Crabs..... | 14, 818 | 682 | | | | | | |
| Sea crawfish or spiny lobster..... | 302, 835 | 42, 206 | | | | | | |
| Abalone..... | | | | | | | 106, 968 | \$10, 701 |
| Clams: | | | | | | | | |
| Hard..... | | | | | 4, 405 | \$1, 062 | | |
| Pismo..... | | | | | 19, 948 | 4, 370 | | |
| Mussels, sea..... | | | | | 47 | 23 | | |
| Octopus..... | 22 | 2 | | | | | | |
| Total..... | 317, 675 | 42, 890 | | | 24, 400 | 5, 455 | 106, 968 | 10, 701 |
| Grand total..... | 395, 633 | 45, 973 | 581, 048 | 50, 718 | 24, 400 | 5, 455 | 106, 968 | 10, 701 |

CATCH OFF LATIN AMERICA: BY GEAR

| Species | Purse seines | | Lampara nets | | Lines, set and hand | | Traps | |
|--|--------------------|-----------------|--------------|-----------|---------------------|-----------------|----------------|---------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | |
| Barracuda..... | 117, 572 | \$10, 320 | | | 8, 691 | \$605 | | |
| Cabrilla..... | | | | | 8, 275 | 233 | | |
| Flounders, "California halibut" Grayfish..... | 91 | 8 | | | 61 | 2 | | |
| Mullet..... | | | 423 | \$17 | | | | |
| Perch..... | | | | | 112 | 11 | | |
| Rock bass..... | 170 | 10 | | | 12, 822 | 819 | | |
| Rockfishes..... | | | | | 7, 099 | 218 | | |
| Sea bass: | | | | | | | | |
| Black..... | 13, 290 | 713 | | | 158, 826 | 7, 085 | | |
| White..... | 325 | 46 | | | 39, 065 | 2, 482 | | |
| Spanish mackerel..... | 410 | 8 | | | 2, 602 | 137 | | |
| Tai..... | | | | | 138 | 4 | | |
| Tuna and tunalike fishes: | | | | | | | | |
| Bluefin..... | 236, 276 | 10, 504 | | | | | | |
| Bonito..... | 154, 431 | 3, 492 | | | 32, 388 | 948 | | |
| Skipjack or striped tuna..... | 3, 006, 823 | 92, 730 | | | 2, 368, 471 | 100, 518 | | |
| Yellowfin..... | 2, 802, 603 | 113, 912 | | | 13, 226, 882 | 594, 180 | | |
| Whitefish..... | 50 | 3 | | | 5, 178 | 260 | | |
| Yellowtail..... | 15, 032 | 378 | | | 415, 843 | 13, 640 | | |
| Total..... | 6, 347, 073 | 232, 124 | 423 | 17 | 16, 286, 453 | 721, 142 | | |
| SHELLFISH | | | | | | | | |
| Sea crawfish or spiny lobster..... | | | | | | | 21, 298 | \$3, 731 |
| Grand total..... | 6, 347, 073 | 232, 124 | 423 | 17 | 16, 286, 453 | 721, 142 | 21, 298 | 3, 731 |

Fisheries of the San Diego district of California, 1933

OPERATING UNITS: BY GEAR

| Item | Lampara and ring nets | | Gill nets | | | Trammel nets | Lines | | Traps, lobster | Harpoons, swordfish and turtle | Total, exclusive of duplication |
|-------------------------|-----------------------|---------|------------------|---------------|--------|--------------|--------------|--------|----------------|--------------------------------|---------------------------------|
| | Mackerel | Sardine | Drift, barracuda | Set, sea bass | Other | | Set and hand | Troll | | | |
| | Number | Number | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | | | | | |
| On vessels..... | 133 | 41 | 3 | 3 | 14 | 3 | 639 | 15 | 27 | 29 | 710 |
| On boats and shore..... | 26 | 18 | 36 | 44 | 14 | 30 | 116 | 41 | 46 | 29 | 214 |
| Total..... | 159 | 59 | 39 | 47 | 14 | 33 | 755 | 41 | 61 | 58 | 924 |
| Vessels, motor..... | 14 | 4 | 1 | 1 | | 1 | 73 | | 5 | 7 | 81 |
| Net tonnage..... | 145 | 44 | 5 | 5 | | 7 | 4,503 | | 47 | 70 | 4,562 |
| Boats: | | | | | | | | | | | |
| Motor..... | 3 | 2 | 16 | 19 | 10 | 12 | 58 | 35 | 35 | 11 | 97 |
| Other..... | | | | 1 | 2 | | | | | | 2 |
| Apparatus: | | | | | | | | | | | |
| Number..... | 17 | 6 | 17 | 21 | 13 | 13 | 1,006 | 181 | 1,308 | 18 | |
| Length, yards..... | 5,610 | 2,550 | | | | | | | | | |
| Square yards..... | | | 137,003 | 110,061 | 17,180 | 128,713 | | | | | |
| Hooks..... | | | | | | | 55,418 | 181 | | | |

CATCH OFF CALIFORNIA: BY GEAR

| Species | Purse seines | | Lampara and ring nets | | Gill nets | | Trammel nets | |
|------------------------------------|--------------|-------|-----------------------|--------|-----------|---------|--------------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | |
| Barracuda..... | 5,609 | \$139 | 30,738 | \$787 | 222,154 | \$0,797 | | |
| Flounders: | | | | | | | | |
| "California halibut"..... | | | | | 28 | 2 | 76,682 | \$5,310 |
| "Sole"..... | | | | | | | 64 | 5 |
| Other..... | | | | | | | 22 | 1 |
| Grayfish..... | | | | | 26,141 | 127 | 14,348 | 99 |
| Herring..... | | | | | 4,665 | 62 | | |
| Horse mackerel..... | | | 600 | 6 | | | | |
| Kingfish..... | | | 13 | 1 | 172 | 4 | 35 | 1 |
| Mackerel..... | | | 8,187,781 | 47,716 | 95,750 | 578 | | |
| Mullet..... | | | | | 19,827 | 800 | | |
| Perch..... | 298 | 12 | | | 30 | 1 | | |
| Pilchard or sardine..... | | | 532,269 | 3,347 | 322 | 3 | | |
| Pompano..... | | | 32 | 10 | | | | |
| Rock bass..... | | | 1,542 | 36 | 1,212 | 45 | 3,333 | 172 |
| Rockfishes..... | | | | | 102 | 4 | 154 | 6 |
| Rudderfish..... | | | | | 339 | 10 | | |
| Sea bass: | | | | | | | | |
| Black..... | | | 180 | 4 | 3,566 | 114 | | |
| White..... | 236 | 8 | 182 | 7 | 163,790 | 7,291 | 260 | 28 |
| Sheepshead..... | | | | | | | 39 | 1 |
| Skates..... | | | | | | | 700 | 8 |
| Smelt..... | | | 60 | 2 | 22,396 | 724 | | |
| Tuna and tunalike fishes: | | | | | | | | |
| Bluefin..... | | | 82 | 5 | | | | |
| Bonito..... | | | 281,589 | 5,627 | 81,786 | 1,604 | 168 | 2 |
| Yellowtail..... | | | 4,689 | 87 | 2,496 | 75 | 140 | 3 |
| Total..... | 6,143 | 159 | 9,039,757 | 57,635 | 644,776 | 18,241 | 95,945 | 5,636 |
| SHELLFISH | | | | | | | | |
| Sea crawfish or spiny lobster..... | | | | | | | 1,511 | 202 |
| Grand total..... | 6,143 | 159 | 9,039,757 | 57,635 | 644,776 | 18,241 | 97,466 | 5,838 |

Fisheries of the San Diego district of California, 1933—Continued

CATCH OFF CALIFORNIA: BY GEAR—Continued

| Species | Lines | | | | Traps | | Harpoons | |
|------------------------------------|------------------|---------------|----------------|--------------|----------------|---------------|----------------|---------------|
| | Set and hand | | Troll | | Pounds | Value | Pounds | Value |
| FISH | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> |
| Barracuda..... | 204,799 | \$7,186 | 14,889 | \$597 | | | | |
| Flounders: | | | | | | | | |
| "California halibut"..... | 1,721 | 123 | | | | | | |
| "Sole"..... | 3,852 | 326 | | | | | | |
| Grayfish..... | 15,310 | 92 | 490 | 2 | | | | |
| Kingfish..... | 1,341 | 26 | | | 82 | \$2 | | |
| Mackerel..... | 1,054,830 | 7,210 | | | | | | |
| Marlin..... | | | | | | | 606 | \$24 |
| Perch..... | 21 | 1 | | | | | | |
| Rock bass..... | 99,211 | 3,461 | | | 24,573 | 1,169 | | |
| Rockfishes..... | 554,932 | 18,097 | | | 1,776 | 67 | | |
| Sablefish..... | 1,604 | 28 | | | | | | |
| Sculpin..... | 7,914 | 347 | | | 124 | 5 | | |
| Sea bass: | | | | | | | | |
| Black..... | 69,759 | 2,493 | | | | | | |
| White..... | 24,299 | 1,183 | 16 | 1 | | | | |
| Sheepshead..... | 3,224 | 64 | | | 4,422 | 158 | | |
| Smelt..... | 270 | 10 | | | | | | |
| Swordfish..... | | | | | | | 234,787 | 17,686 |
| Tuna and tunalike fishes: | | | | | | | | |
| Bluefin..... | 7,220 | 414 | | | | | | |
| Bonito..... | 570,976 | 11,377 | 112,716 | 2,244 | | | | |
| Yellowfin..... | 8,867 | 431 | | | | | | |
| Whitefish..... | 42,108 | 1,673 | | | | | | |
| Yellowtail..... | 971,177 | 22,155 | 8,894 | 107 | | | | |
| Other fish..... | 300 | 12 | | | | | | |
| Total..... | 3,641,735 | 76,709 | 137,005 | 3,041 | 30,977 | 1,401 | 235,393 | 17,709 |
| SHELLFISH | | | | | | | | |
| Sea crawfish or spiny lobster..... | | | | | 72,277 | 10,376 | | |
| Grand total..... | 3,641,735 | 76,709 | 137,005 | 3,041 | 103,254 | 11,777 | 235,393 | 17,709 |

NOTE.—The catch by purse seines was made entirely by fishermen from the San Pedro district.

CATCH OFF LATIN AMERICA: BY GEAR

| Species | Purse seines | | Lampara and ring nets | | Gill nets | | Trammel nets | |
|--------------------------------------|----------------|--------------|-----------------------|--------------|----------------|---------------|---------------|--------------|
| | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> |
| FISH | | | | | | | | |
| Barracuda..... | 1,661 | \$168 | | | 1,359 | \$92 | | |
| Flounders, "California halibut"..... | | | | | | | 81,261 | \$5,581 |
| Mullet..... | | | 2,227 | \$111 | 165 | 5 | | |
| Pompano..... | | | 710 | 56 | | | | |
| Rock bass..... | | | | | 338 | 10 | | |
| Sea bass: | | | | | | | | |
| Black..... | | | | | 11,623 | 424 | | |
| White..... | | | | | 175,464 | 11,082 | | |
| Smelt..... | | | | | 260 | 13 | | |
| Tuna and tunalike fishes: | | | | | | | | |
| Skipjack or striped tuna..... | 4,302 | 151 | | | | | | |
| Yellowfin..... | 183,206 | 7,328 | | | | | | |
| Yellowtail..... | 120 | 6 | | | 2,318 | 45 | | |
| Total..... | 189,289 | 7,653 | 2,937 | 167 | 191,525 | 11,671 | 81,261 | 5,581 |

Fisheries of the San Diego district of California, 1933—Continued

CATCH OFF LATIN AMERICA: BY GEAR—Continued

| Species | Lines | | | | Traps | | Harpoons | |
|--------------------------------------|-------------------|------------------|---------------|--------------|----------------|---------------|---------------|--------------|
| | Set and hand | | Troll | | Pounds | Value | Pounds | Value |
| FISH | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> | | | | |
| Barracuda..... | 24,654 | \$1,744 | 6,573 | \$502 | | | | |
| Cabrilla..... | 76,337 | 2,573 | | | | | | |
| Corbina..... | 290 | 23 | | | | | | |
| Flounders, "California halibut"..... | 41 | 3 | | | | | | |
| Mackerel..... | 969 | 21 | | | | | | |
| Rock bass..... | 4,308 | 229 | | | | | | |
| Rockfishes..... | 36,813 | 1,417 | | | | | | |
| Sea bass: | | | | | | | | |
| Black..... | 145,973 | 5,092 | | | | | | |
| White..... | 120,096 | 8,464 | | | | | | |
| Spanish mackerel..... | 1,185 | 46 | | | | | | |
| Swordfish..... | | | | | | | 40, (65) | \$2,844 |
| Tuna and tunalike fishes: | | | | | | | | |
| Bonito..... | 98,136 | 1,956 | | | | | | |
| Skipjack or striped tuna..... | 11,307,702 | 419,691 | | | | | | |
| Yellowfin..... | 34,856,072 | 1,558,860 | | | | | | |
| Whitefish..... | 8,597 | 347 | | | | | | |
| Yellowtail..... | 2,231,814 | 42,274 | 475 | 10 | | | | |
| Other fish..... | 729 | 37 | | | | | | |
| Total..... | 18,913,719 | 2,042,897 | 7,348 | 512 | | | 40,063 | 2,844 |
| SHELLFISH | | | | | | | | |
| Sea crawfish or spiny lobster..... | | | | | 648,132 | \$87,461 | | |
| Turtles..... | | | | | | | 2,901 | 125 |
| Total..... | | | | | 648,132 | 87,461 | 2,901 | 125 |
| Grand total..... | 18,913,719 | 2,042,897 | 7,348 | 512 | 648,132 | 87,461 | 42,964 | 2,969 |

NOTE.—The catch by purse seines was made entirely by fishermen from the San Pedro district.

HALIBUT FISHERY OF THE PACIFIC COAST ¹¹

The halibut fishery of the Pacific coast, which is prosecuted by United States (including Alaska) and Canadian vessels, ranks as one of the foremost fisheries of that section. During 1933, the total catch of halibut by vessels of both nationalities amounted to 45,951,000 pounds, valued at \$2,582,000. This is an increase of 6 percent in quantity and 48 percent in value as compared with the catch and its value in 1932. Of the total catch in 1933, 82 percent was taken by United States craft and 18 percent by Canadian craft. Considered according to ports of landing 48 percent was landed at ports in the State of Washington; 37 percent at Canadian ports; and 15 percent at ports in Alaska.

¹¹ These statistics were compiled from data collected by the International Fisheries Commission for Washington and British Columbia, and by Bureau agents for Alaska. The data for the Washington and Alaska landings, as well as those made by United States craft in British Columbia, are based on actual weight of the fares.

Halibut fishery of the Pacific coast, 1933

UNITED STATES OPERATING UNITS: BY FLEET CLASSIFICATION

| Item | Washing- ton fleet | Alaska fleet | Total |
|---|-----------------------|-----------------|-------|
| Regular halibut vessels: | | | |
| Number..... | 120 | 86 | 215 |
| Net tonnage..... | 3,712 | 1,578 | 5,290 |
| Crew..... | 885 | 451 | 1,336 |
| Dories..... | 142 | 96 | 229 |
| Skates of lines..... | 3,675 | 2,073 | 5,678 |
| Vessels in other fisheries but landing 1 or more fares of halibut: | | | |
| Number..... | 20 | 9 | 29 |
| Net tonnage..... | 432 | 79 | 511 |
| Crew..... | 110 | 33 | 143 |
| Dories..... | 20 | 9 | 29 |
| Skates of lines..... | 434 | 161 | 595 |
| Regular halibut boats: | | | |
| Number..... | | 27 | 27 |
| Crew..... | | 67 | 67 |
| Skates of lines..... | | 338 | 338 |
| Boats in other fisheries but landing 1 or more fares of halibut: | | | |
| Number..... | 1 | 85 | 86 |
| Crew..... | 3 | 135 | 138 |
| Skates of lines..... | 10 | 498 | 478 |

CATCH OF ALL SPECIES: BY UNITED STATES VESSELS AND BOATS

| Fleet classification | Landed in— | | | | | | Total | |
|---------------------------------|------------|-------------|---------------------|-----------|-----------|----------|------------|-------------|
| | Washington | | British Columbia | | Alaska | | | |
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| WASHINGTON FLEET | | | | | | | | |
| Regular vessels: | | | | | | | | |
| Halibut..... | 19,579,151 | \$1,277,314 | 2,035,711 | \$127,082 | 1,214,077 | \$63,622 | 22,828,939 | \$1,468,018 |
| Sablefish..... | 1,271,209 | 39,540 | 8,565 | 171 | 22,719 | 370 | 1,392,493 | 40,081 |
| “Lingcod”..... | 417,108 | 12,365 | | | | | 417,108 | 12,365 |
| Rockfishes..... | 180,333 | 6,286 | | | | | 180,333 | 6,286 |
| Total..... | 21,447,801 | 1,335,506 | 2,044,276 | 127,253 | 1,236,796 | 63,992 | 24,728,873 | 1,526,780 |
| Other vessels and boats: | | | | | | | | |
| Halibut..... | 763,174 | 49,450 | 4,130 | 115 | | | 767,304 | 49,565 |
| Sablefish..... | 7,867 | 202 | | | | | 7,867 | 202 |
| “Lingcod”..... | 51,021 | 1,261 | | | | | 51,021 | 1,261 |
| Rockfishes..... | 26,649 | 753 | | | | | 26,649 | 753 |
| Total..... | 848,711 | 51,666 | 4,130 | 115 | | | 852,841 | 51,781 |
| ALASKA FLEET | | | | | | | | |
| Regular vessels: | | | | | | | | |
| Halibut..... | 1,772,835 | 103,741 | 6,355,421 | 352,269 | 4,186,346 | 198,628 | 12,314,602 | 652,638 |
| Sablefish..... | 1,166 | 23 | 4,738 | 95 | 64,892 | 1,664 | 70,796 | 1,782 |
| “Lingcod”..... | 2,757 | 109 | | | | | 2,757 | 109 |
| Rockfishes..... | | | | | 1,078 | 22 | 1,078 | 22 |
| Total..... | 1,776,758 | 103,873 | 6,360,159 | 352,364 | 4,252,316 | 198,314 | 12,389,233 | 654,551 |
| Other vessels and boats: | | | | | | | | |
| Halibut..... | 12,061 | 926 | 362,903 | 16,738 | 1,379,345 | 56,060 | 1,764,309 | 73,724 |
| Sablefish..... | | | | | 7,155 | 126 | 7,155 | 126 |
| Total..... | 12,061 | 926 | 362,903 | 16,738 | 1,386,500 | 56,186 | 1,761,464 | 73,850 |
| COMBINED FLEETS | | | | | | | | |
| Regular vessels: | | | | | | | | |
| Halibut..... | 21,351,986 | 1,381,055 | 8,391,132 | 479,351 | 5,400,423 | 260,260 | 35,143,641 | 2,120,656 |
| Sablefish..... | 1,272,375 | 39,563 | 13,303 | 266 | 87,611 | 2,034 | 1,373,289 | 41,563 |
| “Lingcod”..... | 419,865 | 12,474 | | | | | 419,865 | 12,474 |
| Rockfishes..... | 180,333 | 6,286 | | | 1,078 | 22 | 181,411 | 6,308 |
| Total..... | 23,224,559 | 1,439,378 | 8,404,435 | 479,617 | 5,489,112 | 262,306 | 37,118,106 | 2,181,301 |
| Other vessels and boats: | | | | | | | | |
| Halibut..... | 775,235 | 50,376 | 367,033 | 16,853 | 1,379,345 | 56,060 | 2,521,618 | 123,269 |
| Sablefish..... | 7,867 | 202 | | | 7,155 | 126 | 15,022 | 228 |

Halibut fishery of the Pacific coast, 1933—Continued

CATCH OF ALL SPECIES: BY UNITED STATES VESSELS AND BOATS—Continued

| Fleet classification | Landed in— | | | | | | Total | |
|-----------------------------|---------------|--------------|------------------|--------------|---------------|--------------|---------------|--------------|
| | Washington | | British Columbia | | Alaska | | | |
| COMBINED FLEETS—con. | | | | | | | | |
| Other vessels and boats— | | | | | | | | |
| Continued. | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> |
| “Lingcod”..... | 51,021 | \$1,261 | | | | | 51,021 | \$1,261 |
| Rockfishes..... | 26,649 | 753 | | | | | 26,649 | 753 |
| Total..... | 860,772 | 52,592 | 367,033 | \$16,853 | 1,386,500 | \$56,186 | 2,614,305 | 125,631 |
| All vessels and boats: | | | | | | | | |
| Halibut..... | 22,127,221 | 1,431,431 | 8,758,165 | 496,204 | 6,779,768 | 316,310 | 37,665,154 | 2,243,945 |
| Sablefish..... | 1,280,242 | 39,765 | 13,303 | 266 | 94,766 | 2,160 | 1,388,311 | 42,191 |
| “Lingcod”..... | 470,886 | 13,735 | | | | 22 | 470,886 | 13,735 |
| Rockfishes..... | 206,982 | 7,039 | | | 1,078 | | 208,060 | 7,061 |
| Grand total..... | 24,085,331 | 1,491,970 | 8,771,468 | 496,470 | 6,875,612 | 318,492 | 39,732,411 | 2,306,932 |

CATCH OF HALIBUT: BY UNITED STATES AND CANADIAN VESSELS AND BOATS

[Expressed in thousands of pounds and thousands of dollars; that is, 000 omitted]

| Fleet classification | Landed in— | | | | | | Total | |
|------------------------------|-----------------|--------------|------------------|--------------|-----------------|--------------|-----------------|--------------|
| | Washington | | British Columbia | | Alaska | | | |
| WASHINGTON FLEET | | | | | | | | |
| Regular halibut vessels..... | <i>Quantity</i> | <i>Value</i> | <i>Quantity</i> | <i>Value</i> | <i>Quantity</i> | <i>Value</i> | <i>Quantity</i> | <i>Value</i> |
| Other vessels and boats..... | 19,579 | 1,277 | 2,036 | 127 | 1,214 | 64 | 22,829 | 1,468 |
| | 763 | 49 | 4 | (1) | | | 767 | 49 |
| Total..... | 20,342 | 1,326 | 2,040 | 127 | 1,214 | 64 | 23,596 | 1,517 |
| ALASKA FLEET | | | | | | | | |
| Regular halibut vessels..... | 1,773 | 104 | 6,355 | 353 | 4,187 | 196 | 12,315 | 653 |
| Other vessels and boats..... | 12 | 1 | 363 | 17 | 1,379 | 56 | 1,754 | 74 |
| Total..... | 1,785 | 105 | 6,718 | 370 | 5,566 | 252 | 14,069 | 727 |
| COMBINED FLEETS | | | | | | | | |
| Regular halibut vessels..... | 21,352 | 1,381 | 8,391 | 480 | 5,401 | 260 | 35,144 | 2,121 |
| Other vessels and boats..... | 775 | 60 | 367 | 17 | 1,379 | 56 | 2,521 | 123 |
| Total..... | 22,127 | 1,431 | 8,758 | 497 | 6,780 | 316 | 37,665 | 2,244 |
| British Columbia fleet..... | 20 | 1 | 8,266 | 337 | | | 8,286 | 338 |
| Grand total..... | 22,147 | 1,432 | 17,024 | 834 | 6,780 | 316 | 45,951 | 2,582 |

¹ Less than \$500.

VESSEL FISHERIES AT SEATTLE, WASH.

A total of 43,353,031 pounds of fishery products, valued at \$2,443,-235, were handled by Seattle wholesale dealers during 1933, exclusive of quantities received by transporting vessels or by rail from Alaska or Canada. This represents an increase of 3 percent in quantity and 36 percent in value as compared with the quantity and value of products handled during the preceding year. Of the total quantity handled 24,085,331 pounds, valued at \$1,491,970, were landed by fishing vessels, a decrease of less than one-half of 1 percent in quantity but an increase of 41 percent in value as compared with the previous year. Receipts by wholesale dealers from sources other than Alaska or Canada or from vessels in the halibut fleet amounted to 19,267,700 pounds, valued at \$951,265, which is an increase of 6 percent in quantity and 28 percent in value.

Fishery products landed by United States vessels at Seattle, Wash., 1933¹

BY FISHING GROUNDS

| Fishing grounds | Trips | Halibut | | | | Sablefish | | "Lingcod" | | Rockfishes | | Total | |
|----------------------------|---------------|---------------------|--------------------|---------------------|--------------------|------------------|------------------|-----------------|----------------|------------------|----------------|----------------------|--------------------|
| | | No. 1 | | No. 2 | | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| West of Cape Spencer..... | Number 481 | Pounds 8,991,764 | Value \$654,231 | Pounds 5,873,579 | Value \$329,759 | Pounds 48,412 | Value \$1,002 | Pounds 7,497 | Value \$201 | Pounds 10,023 | Value \$266 | Pounds 14,931,275 | Value \$985,459 |
| South of Cape Spencer..... | 728 | 2,643,128 | 208,712 | 4,618,750 | 238,729 | 1,231,830 | 38,763 | 463,389 | 13,534 | 196,959 | 6,773 | 9,154,056 | 506,511 |
| Total..... | 1,209 | 11,634,892 | 862,943 | 10,492,329 | 568,488 | 1,280,242 | 39,765 | 470,886 | 13,735 | 206,982 | 7,039 | 24,085,331 | 1,491,970 |

¹ Halibut fleet.

BY MONTHS

| | | | | | | | | | | | | | |
|----------------|-------|------------|---------|------------|---------|-----------|--------|---------|--------|---------|-------|------------|-----------|
| January..... | 11 | | | | | 500 | 31 | 42,758 | 2,335 | 32,305 | 1,762 | 75,563 | 4,128 |
| February..... | 127 | 1,684,671 | 81,731 | 785,300 | 23,875 | 6,061 | 113 | 29,718 | 569 | 6,540 | 129 | 2,512,290 | 106,417 |
| March..... | 134 | 1,503,943 | 100,068 | 873,718 | 44,399 | 20,516 | 570 | 19,024 | 577 | 10,337 | 346 | 2,427,538 | 145,960 |
| April..... | 138 | 849,249 | 68,268 | 1,093,953 | 56,106 | 13,977 | 342 | 75,588 | 1,542 | 17,966 | 392 | 2,050,733 | 126,650 |
| May..... | 156 | 1,443,628 | 92,786 | 1,378,880 | 58,272 | 4,381 | 62 | 73,767 | 1,262 | 18,195 | 304 | 2,918,851 | 152,686 |
| June..... | 150 | 1,323,451 | 93,308 | 1,438,703 | 59,144 | 72,427 | 2,069 | 56,030 | 1,024 | 25,998 | 484 | 2,916,609 | 156,049 |
| July..... | 110 | 1,031,763 | 82,008 | 1,229,595 | 68,171 | 101,355 | 2,804 | 19,290 | 383 | 9,569 | 191 | 2,391,572 | 153,557 |
| August..... | 135 | 1,216,477 | 101,300 | 1,543,180 | 100,646 | 108,636 | 2,889 | 24,750 | 586 | 19,463 | 396 | 2,917,506 | 205,817 |
| September..... | 81 | 1,032,970 | 99,328 | 874,149 | 64,415 | 276,630 | 8,331 | 3,244 | 98 | 3,358 | 79 | 2,190,351 | 172,251 |
| October..... | 90 | 1,042,703 | 99,743 | 914,469 | 71,517 | 369,963 | 12,227 | 18,469 | 635 | 4,517 | 183 | 2,350,121 | 184,305 |
| November..... | 59 | 506,037 | 44,403 | 355,382 | 21,943 | 292,189 | 9,755 | 42,598 | 1,498 | 14,158 | 623 | 1,210,364 | 78,222 |
| December..... | 18 | | | | | 13,607 | 552 | 65,650 | 3,226 | 44,576 | 2,150 | 123,833 | 5,928 |
| Total..... | 1,209 | 11,634,892 | 862,943 | 10,492,329 | 568,488 | 1,280,242 | 39,765 | 470,886 | 13,735 | 206,982 | 7,039 | 24,085,331 | 1,491,970 |

¹ This tabulation does not include fish received from Alaska or Canada or vessels in the halibut fleet.

Fishery products received by Seattle wholesale dealers, 1933: By months ¹

| Species | January | | February | | March | | April | | May | | June | |
|--------------------------------|---------|---------|----------|---------|---------|---------|---------|---------|---------|---------|-----------|---------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Flounders: | | | | | | | | | | | | |
| " Sole | 66,006 | \$1,673 | 42,891 | \$1,080 | 60,227 | \$1,723 | 58,487 | \$1,673 | 52,514 | \$1,214 | 38,177 | \$1,043 |
| Other | 4,119 | 42 | 1,100 | 14 | 2,030 | 31 | 2,438 | 37 | 2,963 | 30 | 499 | 2 |
| Halibut | | | | | 4,991 | 249 | 4,316 | 215 | 8,910 | 624 | 60,846 | 3,353 |
| Herring | 3,395 | 53 | 13,455 | 73 | 149,800 | 1,321 | 23,300 | 116 | | | | |
| " Lingcod" | 14,575 | 576 | 14,584 | 438 | 18,257 | 584 | 12,806 | 194 | 15,832 | 364 | 38,822 | 458 |
| Perch | 3,607 | 73 | 4,337 | 160 | 13,562 | 579 | 7,527 | 133 | 3,446 | 110 | 2,196 | 60 |
| Rockfishes | 18,917 | 479 | 15,767 | 571 | 9,836 | 287 | 10,136 | 227 | 8,677 | 259 | 9,956 | 241 |
| Sablefish | | | | | | | | | | | 5,510 | 193 |
| Salmon: | | | | | | | | | | | | |
| Blueback, red or sockeye | | | | | | | | | 490 | 34 | 23,179 | 1,623 |
| Chinook or king | 1,543 | 123 | 11,843 | 1,421 | 108,101 | 12,856 | 312,533 | 26,685 | 237,928 | 18,567 | 857,590 | 58,478 |
| Chum or keta | | | | | | | | | 155 | 6 | 2,738 | 66 |
| Humpback or pink | | | 34 | 2 | | | | | 4,985 | 272 | 16,156 | 601 |
| Silver or coho | 752 | 53 | | | | | | | 3,685 | 254 | 23,510 | 1,122 |
| Smelt | 14,781 | 512 | 10,328 | 227 | 33,050 | 641 | | | 10,005 | 600 | 15,970 | 928 |
| Steelhead trout | 6,175 | 556 | 7,886 | 774 | 4,934 | 588 | | | | | 450 | 32 |
| Sturgeon | | | | | | | | | | | | |
| Crabs | 47,683 | 3,280 | 55,350 | 3,474 | 60,799 | 3,220 | 43,869 | 2,858 | 75,544 | 3,955 | 44,385 | 2,452 |
| Octopus | 1,813 | 91 | 1,796 | 90 | 2,588 | 129 | 2,193 | 110 | 1,302 | 65 | 2,150 | 107 |
| Total | 183,366 | 7,511 | 179,371 | 8,324 | 458,175 | 22,208 | 477,605 | 32,248 | 426,396 | 26,354 | 1,142,106 | 70,749 |

| Species | July | | August | | September | | October | | November | | December | | Total | |
|--------------------------------|-----------|---------|-----------|---------|-----------|---------|-----------|---------|-----------|---------|----------|---------|------------|----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Flounders: | | | | | | | | | | | | | | |
| " Sole | 51,328 | \$1,505 | 25,816 | \$643 | 37,057 | \$1,202 | 47,525 | \$1,402 | 34,839 | \$1,143 | 33,670 | \$1,049 | 648,537 | \$15,350 |
| Other | 42 | 1 | 543 | 13 | 92 | 1 | 1,126 | 26 | 3,993 | 61 | 1,536 | 32 | 20,481 | 290 |
| Halibut | 23,380 | 1,403 | 30,563 | 1,377 | 413 | 31 | | | | | | | 133,419 | 7,252 |
| Herring | | | | | | | 1,365 | 14 | | | 653 | 19 | 191,868 | 1,596 |
| " Lingcod" | 12,822 | 252 | 29,844 | 471 | 21,439 | 571 | 12,253 | 457 | 32,852 | 1,179 | 9,565 | 409 | 233,651 | 5,953 |
| Perch | 2,129 | 60 | 4,984 | 128 | 3,899 | 97 | 4,470 | 120 | 3,766 | 75 | 4,454 | 178 | 58,379 | 1,773 |
| Rockfishes | 3,042 | 98 | 5,855 | 80 | 5,338 | 92 | 8,656 | 283 | 13,054 | 477 | 13,838 | 444 | 123,072 | 3,538 |
| Sablefish | | | 1,030 | 31 | 17,783 | 536 | 22,785 | 911 | 8,964 | 275 | | | 56,072 | 1,946 |
| Salmon: | | | | | | | | | | | | | | |
| Blueback, red or sockeye | 289,843 | 23,999 | 64,652 | 4,235 | 12,622 | 917 | | | | | | | 390,716 | 30,808 |
| Chinook or king | 1,246,072 | 72,430 | 1,729,151 | 139,695 | 436,642 | 27,252 | 133,084 | 7,161 | 52,915 | 3,134 | | | 5,127,402 | 367,802 |
| Chum or keta | 6,548 | 194 | 35,417 | 696 | 190,763 | 5,095 | 2,958,156 | 82,334 | 788,669 | 23,276 | 2,278 | 32 | 3,984,726 | 111,691 |
| Humpback or pink | 27,740 | 560 | 901,840 | 20,793 | 1,590,280 | 42,307 | 3,805 | 91 | | | | | 2,523,665 | 63,751 |
| Silver or coho | 253,259 | 12,856 | 922,645 | 51,495 | 1,497,164 | 83,552 | 1,583,619 | 95,122 | 351,806 | 22,873 | 128,059 | 5,511 | 4,758,479 | 272,337 |
| Smelt | 30,798 | 1,234 | 37,647 | 1,632 | 30,445 | 1,879 | 26,603 | 1,506 | 28,678 | 1,573 | 20,859 | 1,452 | 258,414 | 12,032 |
| Steelhead trout | 81,972 | 6,437 | 51,479 | 3,909 | 25,190 | 1,484 | 5,256 | 297 | 6,770 | 540 | 20,965 | 1,715 | 236,602 | 17,828 |
| Sturgeon | 868 | 39 | 962 | 77 | | | 132 | 16 | | | | | 2,412 | 164 |
| Crabs | 26,400 | 1,619 | 28,531 | 1,611 | 18,700 | 1,144 | 58,458 | 3,202 | 91,581 | 6,145 | 53,550 | 3,040 | 594,900 | 36,000 |
| Octopus | 357 | 18 | 1,220 | 61 | 2,337 | 117 | 3,580 | 143 | 3,821 | 1,53 | 1,748 | 70 | 24,905 | 1,165 |
| Total | 2,056,600 | 122,707 | 3,872,229 | 226,947 | 3,890,164 | 166,277 | 4,870,875 | 193,085 | 1,419,708 | 60,904 | 291,205 | 13,951 | 19,267,700 | 951,254 |

¹ This tabulation does not include fish received from Alaska or Canada or vessels in the halibut fleet.² 28,414 dozen.

LAKE FISHERIES¹²

The most recent catch statistics of the fisheries of the Great Lakes, including the international lakes of northern Minnesota, are those collected for the year 1932 and the most recent complete statistics on fisheries wholesale and manufacturing industries in the same region are for 1931. The yield of these fisheries in 1932 in the United States amounted to 83,744,389 pounds valued at \$4,389,061 to the fishermen, representing a decrease of 9 percent in quantity and 27 percent in value as compared with the catch and its value in 1931. Detailed statistics of the operating units and catch of the Lake fisheries for 1932 appear in "Fishery Industries of the United States, 1933", Appendix I to the Report of the Commissioner of Fisheries for the fiscal year 1934 while data on wholesale and manufacturing industries for 1931 are published in "Fisheries Industries of the United States, 1932", Appendix III to the Report of the Commissioner of Fisheries for the fiscal year 1933. A summary of these fisheries appears in the following tables.

Lake fisheries, 1932¹

OPERATING UNITS: BY STATES

| Item | New York | Pennsylvania | Ohio | Michigan | Indiana | Illinois | Wisconsin | Minnesota | Total |
|------------------------|----------|--------------|--------|----------|---------|----------|-----------|-----------|--------|
| | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| Fishermen: | | | | | | | | | |
| On vessels..... | 85 | 112 | 96 | 824 | 16 | 48 | 525 | ----- | 1,705 |
| On boats and shore: | | | | | | | | | |
| Regular..... | 98 | 24 | 601 | 1,704 | 10 | 2 | 368 | 252 | 3,059 |
| Casual..... | 130 | ----- | 249 | 870 | 49 | 10 | 638 | 222 | 2,168 |
| Total..... | 313 | 136 | 946 | 3,398 | 74 | 60 | 1,531 | 474 | 6,932 |
| Vessels: | | | | | | | | | |
| Steam..... | 5 | 12 | 7 | 51 | 1 | 1 | 29 | ----- | 106 |
| Net tonnage..... | 122 | 298 | 227 | 926 | 22 | 13 | 756 | ----- | 2,364 |
| Motor..... | 12 | 9 | 12 | 189 | 3 | 13 | 154 | ----- | 392 |
| Net tonnage..... | 87 | 95 | 138 | 1,818 | 41 | 163 | 1,693 | ----- | 4,055 |
| Total vessels..... | 17 | 21 | 19 | 240 | 4 | 14 | 183 | ----- | 498 |
| Total net tonnage..... | 209 | 393 | 365 | 2,744 | 63 | 196 | 2,449 | ----- | 6,419 |
| Boats: | | | | | | | | | |
| Motor..... | 71 | 7 | 251 | 879 | 51 | 7 | 244 | 114 | 1,624 |
| Other..... | 115 | 10 | 300 | 555 | 33 | ----- | 218 | 804 | 1,536 |

¹ The statistics of the catch from 1929 to 1932, inclusive, include data on crawfish, mussels, pearls and slugs which were omitted in most of the surveys prior to 1929. Data on the catch in Wisconsin for 1929 and in subsequent years also are not strictly comparable with data for previous years due to a more complete method of collection adopted in the more recent statistics. Data for the years 1929 to 1932 are shown on the basis of round weight whereas they formerly included dressed or partially dressed fish in some cases.

¹² The statistics of the catch presented herewith were obtained principally from the records of the various State fishery agencies and from the Dominion Bureau of Statistics, Ottawa, Canada. The data for the operating units (fishermen, vessels, boats, and gear) of the United States were obtained largely by Bureau agents in a special canvass; although State records in several instances were very helpful in this work. In all cases the statistics collected are for the calendar year, except for Lake of the Woods, Rainy Lake, and Lake Namakan in Minnesota, which are for two seasons. For Lake of the Woods the seasons are from June 2 to Nov. 1 and Dec. 1 to Apr. 1 and for Rainy and Namakan Lakes from May 15 to Nov. 1 and Dec. 1 to Apr. 1. The catches for these two seasons, in the order named, have been combined to constitute a year. The quantity of fish taken in these lakes between Jan. 1 and Apr. 1 is estimated at less than 3 percent of the total catch. For a clearer understanding of the statistics published in this section the reader is referred to the section in the latter part of the document entitled "Statistical survey procedure."

Lake fisheries, 1932—Continued

OPERATING UNITS: BY STATES—Continued

| Item | New York | Pennsylvania | Ohio | Michigan | Indiana | Illinois | Wisconsin | Minnesota | Total |
|----------------------------------|----------|--------------|----------|--------------|----------|----------|-------------|-------------|--------------|
| | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| Apparatus: | | | | | | | | | |
| Haul seines..... | 18 | | 123 | 141 | | | 50 | | 332 |
| Length, yards..... | 2, 190 | | 70, 485 | 49, 592 | | | 16, 829 | | 139, 098 |
| Gill nets: | | | | | | | | | |
| "Bull", 3 to 3½ inches..... | 584 | | | | | | | | 584 |
| Square yards..... | 233, 600 | | | | | | | | 233, 600 |
| "Shoal", 2¼ to 3¼ inches..... | 2, 009 | 4, 494 | 6, 001 | 9, 436 | 361 | 1, 390 | 10, 814 | 3, 614 | 38, 109 |
| Square yards..... | 314, 913 | 741, 840 | 919, 050 | 2, 160, 607 | 105, 750 | 288, 640 | 3, 070, 299 | 1, 196, 403 | 8, 797, 502 |
| "Shoal", 4 to 5¾ inches..... | 2, 751 | 3, 684 | 563 | 39, 453 | 440 | 1, 122 | 14, 366 | 1, 818 | 64, 197 |
| Square yards..... | 455, 088 | 933, 104 | 84, 460 | 11, 363, 279 | 129, 167 | 338, 668 | 4, 548, 867 | 685, 425 | 18, 538, 648 |
| "Shoal", 6 to 9¾ inches..... | 142 | | | 344 | 30 | | 40 | | 556 |
| Square yards..... | 26, 464 | | | 41, 649 | 6, 300 | | 6, 600 | | 80, 913 |
| "Shoal", 10 to 14 inches..... | 72 | | | | | | | | 72 |
| Square yards..... | 16, 800 | | | | | | | | 16, 800 |
| Trammel nets..... | | | 222 | | | | 4 | | 226 |
| Square yards..... | | | 9, 324 | | | | 413 | | 9, 737 |
| Lines: | | | | | | | | | |
| Hand..... | | | | 6 | | | | | 6 |
| Hooks..... | | | | 7 | | | | | 7 |
| Troll..... | | | | 41 | | | | | 41 |
| Hooks..... | | | | 46 | | | | | 46 |
| Trot..... | | | | 1, 783 | | 5 | 816 | 403 | 3, 049 |
| Hooks..... | 14, 050 | | | 670, 630 | | 500 | 236, 350 | 38, 930 | 960, 460 |
| Pound nets..... | | 45 | 18 | 1, 395 | 7 | | 294 | 73 | 1, 833 |
| Trap nets..... | 173 | 8 | 3, 875 | 3, 365 | | | 5 | | 7, 426 |
| Fyke nets..... | 131 | | 633 | 935 | | | 759 | 116 | 2, 574 |
| Crawfish pots..... | | | | | | | 2, 910 | | 2, 910 |
| Crowfoot bars..... | | | | 315 | 40 | | | | 360 |
| Picks..... | | | | 124 | | | 2 | | 126 |

CATCH: BY STATES

| Species | New York | | Pennsylvania | | Ohio | |
|-------------------------------|-------------|-----------|--------------|-----------|--------------|-------------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Blue pike..... | 558, 052 | \$27, 136 | 1, 606, 925 | \$72, 311 | 7, 784, 487 | \$311, 660 |
| Burbot..... | 22, 758 | 782 | 1, 839 | 19 | 246, 096 | 2, 465 |
| Carp..... | 78, 098 | 3, 747 | 7, 203 | 309 | 1, 789, 732 | 35, 792 |
| Catfish and bullheads..... | 67, 656 | 3, 181 | 2, 823 | 191 | 444, 802 | 18, 434 |
| Cisco..... | 27, 464 | 2, 746 | 82, 836 | 8, 283 | 49, 915 | 5, 090 |
| Eels..... | 43, 536 | 1, 306 | | | | |
| Goldfish..... | | | | | 44, 013 | 441 |
| Lake herring..... | 72, 793 | 5, 446 | | | | |
| Lake trout..... | 26, 975 | 3, 453 | 1, 051 | 126 | 15, 184 | 150 |
| Mooneye..... | | | 500 | 5 | 1, 355 | 106 |
| Pike or pickerel (jacks)..... | 14, 853 | 1, 202 | | | | |
| Rock bass..... | 1, 208 | 39 | | | | |
| Sauger..... | | | | | 3, 088, 883 | 122, 351 |
| Sheepshead..... | | | | | 2, 069, 439 | 41, 395 |
| Sturgeon..... | 27, 921 | 5, 825 | 13, 859 | 328 | | |
| Sucker "mullet"..... | 108, 467 | 4, 329 | 607 | 101 | | |
| Sunfish..... | 8, 084 | 256 | 13, 824 | 275 | 1, 094, 792 | 21, 893 |
| White bass..... | 2, 840 | 113 | | | | |
| Whitefish, common..... | 250, 120 | 43, 581 | 9, 252 | 369 | 240, 693 | 9, 328 |
| Yellow perch..... | 102, 296 | 6, 556 | 447, 398 | 67, 505 | 507, 084 | 78, 060 |
| Yellow pike..... | 23, 354 | 2, 508 | 329, 676 | 15, 694 | 9, 239, 058 | 329, 969 |
| | | | 17, 477 | 1, 746 | 1, 900, 396 | 184, 520 |
| Total..... | 1, 434, 475 | 112, 206 | 2, 535, 260 | 167, 262 | 28, 515, 829 | 1, 100, 584 |

Lake fisheries, 1932--Continued

CATCH: BY STATES--Continued

| Species | Michigan | | Indiana | | Illinois | |
|-------------------------------|------------|-----------|---------|--------|----------|--------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Rowfin | 2,948 | \$59 | | | | |
| Buffalofish | | | 1,910 | \$100 | | |
| Burbot | 19,575 | 241 | 3,962 | 220 | | |
| Carp | 2,141,439 | 72,206 | 3,320 | 119 | 200 | \$4 |
| Catfish and bullheads | 258,415 | 20,018 | | | | |
| Chubs | 948,429 | 68,531 | 184,698 | 16,473 | 467,445 | 28,704 |
| Goldfish | 4,520 | 90 | | | | |
| Lake herring | 3,442,460 | 86,099 | 123,050 | 5,076 | 85,580 | 3,444 |
| Lake trout | 6,813,110 | 630,361 | 98,391 | 8,307 | 280,606 | 21,877 |
| Pike or pickerel (jacks) | 60,584 | 4,813 | | | | |
| Rock bass | 16,292 | 605 | | | | |
| Sauger | 142,798 | 6,705 | | | | |
| Sheepshead | 75,039 | 2,257 | | | | |
| Smelt | 22,004 | 660 | | | | |
| Steelhead trout | | | 4,100 | 650 | | |
| Sucker "mullet" | 3,810,061 | 85,206 | 2,030 | 85 | | |
| Whitefish: | | | | | | |
| Common | 7,787,861 | 865,750 | 9,210 | 858 | 3,240 | 360 |
| Menominee | 141,598 | 13,895 | | | | |
| Yellow perch | 1,006,324 | 65,955 | 23,238 | 1,831 | 47,715 | 3,573 |
| Yellow pike | 1,759,108 | 213,474 | 6,430 | 793 | | |
| Mussel shells ¹ | 1,678,984 | 23,181 | 170,000 | 2,210 | | |
| Pearls and slugs ¹ | | 1,454 | | 170 | | |
| Total | 30,129,549 | 2,161,560 | 630,339 | 36,892 | 884,785 | 57,962 |

| Species | Wisconsin | | Minnesota | | Total | |
|-------------------------------|------------|---------|-----------|---------|------------|-----------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Blue pike | | | | | 9,947,464 | \$411,107 |
| Rowfin | | | | | 2,948 | 59 |
| Buffalofish | | | | | 1,910 | 100 |
| Burbot | 36,841 | \$553 | 45 | \$1 | 331,116 | 4,281 |
| Carp | 256,707 | 5,789 | 6,870 | 132 | 4,283,569 | 118,098 |
| Catfish and bullheads | 50,256 | 3,528 | 9,412 | 455 | 833,364 | 45,807 |
| Chubs | 2,430,070 | 134,014 | 27,870 | 560 | 4,056,512 | 248,272 |
| Cisco | | | | | 160,215 | 17,019 |
| Crappie | | | 1,018 | 193 | 1,018 | 193 |
| Eels | | | | | 43,536 | 1,306 |
| Goldfish | | | | | 48,833 | 531 |
| Lake herring | 2,839,066 | 26,941 | 5,123,435 | 54,251 | 11,686,374 | 181,257 |
| Lake trout | 2,909,113 | 206,185 | 532,467 | 47,282 | 10,661,712 | 919,591 |
| Mooneye | | | | | 15,684 | 155 |
| Pike or pickerel (jacks) | 76,178 | 4,790 | 219,586 | 5,413 | 372,556 | 16,324 |
| Rock bass | | | | | 17,500 | 644 |
| Sauger | | | 215,898 | 6,163 | 3,447,579 | 135,219 |
| Sheepshead | 167 | 6 | | | 2,158,504 | 43,986 |
| Smelt | 75,803 | 2,653 | | | 97,807 | 3,313 |
| Steelhead trout | 950 | 124 | | | 5,050 | 774 |
| Sturgeon | | | 1,384 | 355 | 29,912 | 6,281 |
| Sucker "mullet" | 1,042,315 | 23,998 | 120,871 | 1,470 | 6,192,360 | 137,256 |
| Sunfish | | | | | 8,084 | 256 |
| Tullibee | | | 1,296,668 | 16,104 | 1,296,668 | 16,104 |
| White bass | | | | | 252,895 | 9,810 |
| Whitefish: | | | | | | |
| Common | 548,739 | 36,314 | 176,862 | 11,214 | 9,730,504 | 1,101,642 |
| Menominee | 89,108 | 3,219 | 1,968 | 89 | 232,674 | 17,203 |
| Yellow perch | 682,663 | 41,633 | 41,630 | 1,661 | 11,472,500 | 466,902 |
| Yellow pike | 3,365 | 472 | 731,330 | 53,820 | 4,441,450 | 457,333 |
| Crawfish | 19,677 | 984 | | | 19,677 | 984 |
| Mussel shells ¹ | 45,930 | 239 | | | 1,894,914 | 25,630 |
| Pearls and slugs ¹ | | | | | | 1,624 |
| Total | 11,106,838 | 493,442 | 8,507,314 | 199,153 | 83,744,389 | 4,389,061 |

¹ From streams tributary to Lakes Michigan, Huron, and Erie.

Industries related to the fisheries of the Lake States

OPERATING UNITS SALARIES, AND WAGES, 1931

| Item | New York | Pennsylvania | Ohio | Michigan | Indiana | Illinois | Wisconsin | Minnesota | Total |
|---------------------------------|----------|--------------|-----------|-----------|----------|-------------|-----------|-----------|-------------|
| | Number | Number | Number | Number | Number | Number | Number | Number | Number |
| Transporting: | | | | | | | | | |
| Persons engaged..... | | | 16 | 4 | | | | | 19 |
| Vessels, motor..... | | | 8 | 1 | | | | | 9 |
| Net tonnage..... | | | 88 | 32 | | | | | 120 |
| Wholesale and manufacturing: | | | | | | | | | |
| Establishments..... | 16 | 8 | 41 | 57 | 3 | 55 | 35 | 15 | 230 |
| Persons engaged: | | | | | | | | | |
| Proprietors..... | 15 | 14 | 58 | 67 | 3 | 26 | 33 | 9 | 225 |
| Salaried employees..... | 22 | 7 | 63 | 54 | 1 | 249 | 48 | 27 | 471 |
| Wage earners: | | | | | | | | | |
| Average for season..... | 69 | 58 | 249 | 438 | 4 | 352 | 275 | 61 | 1,506 |
| Average for year..... | 54 | 53 | 215 | 193 | 4 | 318 | 150 | 47 | 1,034 |
| Paid to salaried employees..... | \$36,516 | \$24,225 | \$201,954 | \$157,875 | \$5,500 | \$765,402 | \$109,407 | \$49,010 | \$1,349,859 |
| Paid to wage earners..... | \$58,062 | \$59,860 | \$261,908 | \$183,872 | \$5,500 | \$479,437 | \$167,271 | \$44,640 | \$1,260,550 |
| Total salaries and wages..... | \$94,578 | \$84,085 | \$463,862 | \$341,747 | \$11,000 | \$1,244,839 | \$276,678 | \$93,650 | \$2,610,439 |
| Fishermen manufacturing..... | 2 | 2 | | 37 | 3 | 12 | 50 | | 106 |

PRODUCTS MANUFACTURED

| Item | New York | | Pennsylvania | | Ohio | | Michigan | |
|--|-----------------------|----------------------|-------------------|-------------------|----------------------|---------------------|----------------------|---------------------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| By manufacturing establishments: | | | | | | | 269,800 | \$67,591 |
| Chubs, smoked..... pounds..... | (¹) | (¹) | | | | | | |
| Eels, smoked..... do..... | (¹) | (¹) | | | | | | |
| Lake herring: | | | | | | | 2,052,300 | 62,433 |
| Salted..... do..... | | | | | | | (¹) | (¹) |
| Smoked..... do..... | (¹) | (¹) | | | (¹) | (¹) | | |
| Lake trout, smoked..... do..... | | | | | | | 21,245 | 4,570 |
| Pike: | | | | | | | | |
| Fresh fillets ² do..... | 284,678 | \$66,298 | 179,610 | \$36,558 | 1,244,370 | \$232,285 | | |
| Frozen fillets ³ do..... | 68,673 | 15,211 | 61,136 | 11,827 | 111,079 | 19,969 | | |
| Salmon, smoked..... do..... | (¹) | (¹) | | | 115,000 | 24,200 | (¹) | (¹) |
| Sturgeon, smoked..... do..... | (¹) | (¹) | | | 12,150 | 8,135 | | |
| Tullibee, smoked..... do..... | (¹) | (¹) | | | 142,500 | 28,230 | | |
| Whitefish, smoked..... do..... | (¹) | (¹) | | | 44,500 | 9,780 | 18,570 | 4,255 |
| Yellow perch: | | | | | | | | |
| Fresh fillets ⁴ do..... | 12,161 | 2,895 | 61,045 | 13,756 | 520,166 | 100,989 | | |
| Frozen fillets ⁵ do..... | (¹) | (¹) | 4,000 | 770 | 81,731 | 15,744 | | |
| Unclassified products: | | | | | | | | |
| Fillets, fresh and frozen ⁶ pounds..... | (¹) | (¹) | (¹) | (¹) | ⁸ 121,880 | ⁸ 19,376 | (¹) | (¹) |
| Smoked..... do..... | | | | | ⁹ 90,000 | ⁹ 24,440 | | |
| Miscellaneous ⁷ | ¹⁰ 298,000 | ¹⁰ 70,440 | ¹¹ 770 | ¹¹ 118 | | | ¹² 25,085 | ¹² 4,692 |
| Total..... | 663,512 | 154,844 | 306,561 | 63,029 | 2,483,376 | 483,148 | 2,387,009 | 133,541 |

¹ The production of this item is included under "Unclassified products."

² Data are for 1933.

³ The production of this item is included under "Miscellaneous."

⁴ Includes fresh fillets of lake trout and whitefish; and fresh and frozen fillets of sheepshead and white bass.

⁵ Includes smoked carp, chubs, lake herring, lake trout, paddlefish, or spoonbill cat and sablefish.

⁶ Both 1931 and 1933 data are included in these items.

⁷ Includes smoked chubs, ciscoes, eels, lake herring, salmon, sturgeon, tullibees, and whitefish; and frozen fillets of yellow perch.

⁸ Includes fresh fillets of sheepshead and white bass.

⁹ Includes smoked buffalofish, butterfish, carp, lake herring, sablefish, salmon, and whitefish; and pickled sea herring.

Industries related to the fisheries of the Lake States—Continued

PRODUCTS MANUFACTURED—Continued

| Item | New York | | Pennsylvania | | Ohio | | Michigan | |
|----------------------------------|----------|---------|--------------|--------|-----------|-----------|-----------|---------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| By fishermen: | | | | | | | | |
| Chubs, smoked.....pounds.. | | | | | | | 17,700 | \$5,040 |
| Lake herring: | | | | | | | | |
| Salted.....do..... | | | | | | | 249,600 | 6,911 |
| Smoked.....do..... | | | | | | | 6,250 | 1,222 |
| Lake trout, smoked.....do..... | | | | | | | 550 | 163 |
| Sturgeon roe, salted.....do..... | 45 | \$52 | | \$50 | | | | |
| Whitefish, smoked.....do..... | | | | | | | 5,500 | 1,290 |
| Total..... | 45 | 52 | 50 | 50 | | | 279,600 | 14,616 |
| Grand total..... | 663,557 | 164,896 | 306,611 | 63,079 | 2,483,376 | \$483,148 | 2,666,609 | 148,167 |

| Item | Indiana | | Illinois | | Wisconsin | | Minnesota | |
|---|------------------------------|---------------------------|------------------------------|---------------------------|------------------------------|---------------------------|------------------------------|---------------------------|
| | Quantity (¹) | Value (²) | Quantity (¹) | Value (²) | Quantity (¹) | Value (²) | Quantity (¹) | Value (¹) |
| By manufacturing establishments: | | | | | | | | |
| Chubs, smoked.....pounds.. | | | 905,476 | \$218,931 | 665,662 | \$119,346 | | |
| Eels, smoked.....do..... | | | 17,804 | 4,483 | | | | |
| Lake herring: | | | | | | | | |
| Salted.....do..... | | | | | 923,350 | 17,906 | | |
| Smoked.....do..... | | | (¹) | (¹) | 456,832 | 22,529 | (¹) | (¹) |
| Lake trout, smoked.....do..... | (²) | (²) | 13,803 | 3,534 | 160,290 | 28,847 | (¹) | (¹) |
| Pike: | | | | | | | | |
| Fresh fillets.....do..... | | | 984,927 | 202,965 | (¹) | (¹) | | |
| Frozen fillets.....do..... | | | (¹) | (¹) | | | | |
| Salmon, smoked.....do..... | | | 636,965 | 161,866 | 37,360 | 6,364 | (¹) | (¹) |
| Sturgeon, smoked.....do..... | | | (¹) | (¹) | | | | |
| Tullibee, smoked.....do..... | | | 325,536 | 65,165 | (¹) | (¹) | (¹) | (¹) |
| Whitefish, smoked.....do..... | | | (¹) | (¹) | 100,267 | 19,840 | (¹) | (¹) |
| Yellow perch: | | | | | | | | |
| Fresh fillets.....do..... | | | 103,241 | 24,077 | 63,500 | 13,715 | | |
| Unclassified products: | | | | | | | | |
| Fillets, fresh and frozen.....pounds..... | | | 109,218 | 20,171 | 55,000 | 10,100 | | |
| Smoked.....do..... | | | (¹) | (¹) | (¹) | (¹) | (¹) | (¹) |
| Miscellaneous..... | | | 2,713,986 | 49,240 | | 34,253 | 170,250 | \$33,664 |
| Total..... | | | 5,810,956 | 750,432 | | 272,900 | 170,250 | 33,664 |
| By fishermen: | | | | | | | | |
| Chubs, smoked.....pounds.. | 42,200 | \$10,550 | 53,000 | 10,800 | | | | |
| Lake herring, salted.....do..... | | | | | 1,027,000 | 25,675 | | |
| Lake trout, smoked.....do..... | 8,450 | 2,113 | | | | | | |
| Whitefish, smoked.....do..... | 790 | 188 | | | | | | |
| Total..... | 51,400 | 12,851 | 53,000 | 10,800 | 1,027,000 | 25,675 | | |
| Grand total..... | 51,400 | 12,851 | 5,863,956 | 761,032 | | 298,575 | 170,250 | 33,664 |

¹ The production of this item is included under "Unclassified."

² A small amount of smoked schub and lake trout produced in Indiana is included with Illinois.

³ Data are for 1933.

⁴ The production of this item is included under "Miscellaneous."

⁵ Includes fresh fillets of lake trout, sheepshead, white bass, and whitefish; and frozen fillets of pike.

⁶ Includes fresh fillets of pike, lake herring, and lake trout.

⁷ Both 1931 and 1933 data are included in these items.

⁸ Includes smoked buffalo fish, butterfish, carp, flounders, lake herring, mackerel, sablefish, shad, sturgeon, and whitefish; and pickled sea herring.

⁹ Includes pickled sea herring, smoked tullibees, spiced alewives, canned whitefish caviar, and mussel-shell products.

¹⁰ Includes smoked chubs, lake herring, lake trout, salmon, suckers, tullibees, and whitefish; and salted lake trout.

NOTE.—Unless otherwise indicated the data are for 1931. The total value of manufactured products for the Lake States was as follows: By manufacturing establishments, \$1,591,568; and by fishermen, \$63,644. Some of the above products may have been manufactured from products imported from another State or country; therefore they cannot be correlated directly with the catch within the State.

FISHERIES OF THE MISSISSIPPI RIVER AND TRIBUTARIES ¹³

The most recent complete catch statistics of the fisheries for the States of the Mississippi River and tributaries are those collected for the year 1931. The yield of fishery products in that year amounted to 82,382,523 pounds, valued at \$2,897,357, which was a decrease of 22 percent in quantity and 36 percent in value as compared with the quantity and value of the catch in 1922 when the most recent preceding survey was made. Detailed statistics of the fisheries of the Mississippi River and tributaries for 1931 appear in "Fishery Industries of the United States, 1932" by R. H. Fiedler, Appendix III to the Report of the Commissioner of Fisheries for the fiscal year 1933. A summary of these fisheries in 1931 as well as certain data for 1933 appear in the following tables.

Fisheries of the Mississippi River and tributaries, 1931

OPERATING UNITS: BY STATES

| Items | Ala- bama | Ar- kansas | Illinois | Indi- ana | Iowa | Kansas | Ken- tucky | Louis- iana | Minn- esota |
|-----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|----------------|----------------|
| Fishermen: | | | | | | | | | |
| On boats and shore: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| Regular..... | 104 | 1,463 | 708 | 20 | 245 | 35 | 89 | 1,402 | 160 |
| Casual..... | 131 | 1,524 | 1,318 | 1,735 | 648 | 88 | 440 | 3,108 | 578 |
| Total..... | 235 | 2,987 | 2,026 | 1,755 | 893 | 123 | 529 | 4,510 | 738 |
| Boats: | | | | | | | | | |
| Motor..... | 32 | 907 | 676 | 544 | 309 | 18 | 92 | 1,225 | 65 |
| Other..... | 190 | 2,359 | 957 | 1,189 | 457 | 123 | 420 | 2,369 | 446 |
| Apparatus: | | | | | | | | | |
| Haul seines..... | | 16 | 127 | 50 | 133 | | 24 | 377 | 113 |
| Length, yards..... | | 7,308 | 33,975 | 5,170 | 36,339 | | 2,057 | 85,166 | 49,968 |
| Anchor gill nets..... | | 4 | | | | | | 74 | 9 |
| Square yards..... | | 2,800 | | | | | | 17,400 | 9,999 |
| Trammel nets..... | | 31 | 28 | 3 | | 90 | | 78 | |
| Square yards..... | | 3,899 | 4,890 | 360 | | 9,026 | | 19,696 | |
| Lines: | | | | | | | | | |
| Trot..... | 449 | 5,327 | 1,312 | 320 | 1,158 | 17 | 627 | 5,757 | 186 |
| Hooks..... | 35,980 | 455,000 | 124,715 | 16,767 | 186,250 | 360 | 37,395 | 1,392,200 | 41,800 |
| Pound nets..... | | | | | 2 | | | | 27 |
| Fyke nets..... | 610 | 5,346 | 9,852 | 335 | 1,981 | 189 | 1,231 | 5,908 | 74 |
| Dip nets..... | | | 22 | | 10 | | | 159 | |
| Traps: | | | | | | | | | |
| Crawfish..... | | | | | | | | 18 | |
| Shrimp..... | | | | | | | | 88 | |
| Baskets: | | | 3,769 | | | | | | |
| Mussel dredges..... | | 426 | 14 | | | | | | |
| Yards at mouth..... | | 286 | 10 | | | | | | |
| Crowfoot bars..... | 168 | 1,038 | 840 | 1,092 | 464 | | 256 | 10 | 192 |
| Tongs..... | | 159 | | 72 | | | | 5 | |
| Rakes..... | | 70 | | | | | | | |
| Forks..... | | 102 | 33 | 1,278 | | 29 | | 5 | |
| Grabs..... | | | | | | | | 2,232 | |

¹³ For a clearer understanding of the statistics published in this section, the reader is referred to the section in the latter part of the document entitled "Statistical survey procedure."

Fisheries of the Mississippi River and tributaries, 1931—Continued

OPERATING UNITS: BY STATES—Continued

| Items | Mississippi | Missouri | Nebraska | Ohio | Oklahoma | South Dakota | Tennessee | Texas | Wisconsin | Total |
|----------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| Fishermen: | | | | | | | | | | |
| On boats and shore: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| Regular | 211 | 177 | | | 5 | | 327 | 5 | 262 | 5,153 |
| Casual | 198 | 170 | 259 | 49 | 19 | 67 | 206 | 41 | 112 | 10,731 |
| Total | 409 | 347 | 259 | 49 | 24 | 67 | 533 | 46 | 314 | 15,884 |
| Boats: | | | | | | | | | | |
| Motor | 138 | 84 | 30 | | | 2 | 138 | 6 | 160 | 4,426 |
| Other | 329 | 304 | 187 | 49 | 18 | 34 | 467 | 42 | 180 | 10,120 |
| Apparatus: | | | | | | | | | | |
| Haul seines | 16 | 47 | 12 | 2 | 2 | 11 | | | 83 | 1,013 |
| Length, yards | 6,885 | 5,668 | 906 | 180 | 60 | 1,048 | | | 20,149 | 255,779 |
| Anchor gill nets | 4 | 1 | | | 3 | | | | 6 | 101 |
| Square yards | 800 | 160 | | | 900 | | | | 13,488 | 45,637 |
| Trammel nets | 17 | 104 | 115 | | | | 52 | | | 518 |
| Square yards | 2,867 | 14,668 | 4,833 | | | | 3,560 | | | 63,799 |
| Lines: | | | | | | | | | | |
| Hand | | | | | | | 67 | | | 67 |
| Hooks | | | | | | | 464 | | | 67 |
| Trot | 847 | 516 | | 19 | 29 | 18 | 67 | 80 | 3 | 17,129 |
| Hooks | 72,155 | 34,600 | | 900 | 1,075 | 3,600 | 41,690 | 14,500 | 125 | 2,459,112 |
| Pound nets | | | | | | | | | 345 | 374 |
| Fyke nets | 2,591 | 1,872 | 296 | 76 | 85 | 68 | 1,735 | 70 | 222 | 32,541 |
| Dip nets | | | | | | | | | | 191 |
| Traps: | | | | | | | | | | |
| Crawfish | | | | | | | | | | 18 |
| Shrimp | 350 | | | | | | | | | 438 |
| Baskets | | | | | | | | | | 3,769 |
| Spears | | | | | | | 12 | | | 12 |
| Mussel dredges | | | | | | | | | | 440 |
| Yards at mouth | | | | | | | | | | 296 |
| Crowfoot bars | | | | | | | 230 | | 190 | 4,450 |
| Tongs | | 9 | | | | | | | | 245 |
| Rakes | | | | | | | | | | 70 |
| Forks | | | | | | | | | | 1,447 |
| Grabs | | | | | | | | | | 2,232 |

CATCH: BY STATES

| Species | Alabama | | Arkansas | | Illinois | | Indiana | |
|------------------------------|------------------|---------------|-------------------|----------------|-------------------|----------------|------------------|----------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | |
| Bowfin | | | 700 | \$28 | 8,308 | \$241 | | |
| Buffalofish | 21,330 | \$2,342 | 2,182,446 | 131,474 | 911,609 | 51,893 | 85,045 | \$8,158 |
| Carp | 11,000 | 1,118 | 808,206 | 27,268 | 4,878,744 | 128,221 | 157,641 | 10,162 |
| Catfish and bullheads | 81,200 | 8,850 | 1,077,343 | 93,150 | 647,696 | 68,890 | 35,370 | 5,302 |
| Crapple | 9,772 | 1,004 | 11,325 | 227 | | | | |
| Eels | | | | | 4,985 | 322 | | |
| Mooneye | | | | | 1,000 | 20 | | |
| Paddlefish or spoonbill cat | 3,958 | 338 | 93,200 | 2,159 | 104,846 | 5,480 | 16,492 | 1,724 |
| Quillback or "American carp" | 7,657 | 875 | 6,830 | 676 | 17,532 | 608 | 30,312 | 1,436 |
| Sheepshead | 45,909 | 4,972 | 676,358 | 29,877 | 177,709 | 11,321 | 38,740 | 3,711 |
| Sturgeon, shovelnose | 575 | 70 | | | 39,756 | 3,448 | 3,013 | 292 |
| Sucker "mullet" | 5,752 | 609 | 3,309 | 235 | 25,130 | 1,087 | 16,797 | 1,156 |
| White bass | | | | | 1,200 | 92 | | |
| Yellow pike | | | | | | | 4,550 | 693 |
| Total | 187,153 | 20,178 | 4,859,717 | 285,094 | 6,818,525 | 271,623 | 387,960 | 32,632 |
| SHELLFISH, ETC. | | | | | | | | |
| Mussel shells | 1,635,000 | 10,132 | 10,872,790 | 108,819 | 7,429,528 | 82,894 | 7,328,736 | 105,632 |
| Pearls | | | | 3,137 | | 190 | | 125 |
| Slags | | 2,444 | | 14,461 | | 11,835 | | 18,788 |
| Turtles: | | | | | | | | |
| Snapper | | | | | 14,577 | 691 | 500 | 25 |
| Soft shell | | | | | | | 400 | 20 |
| Total | 1,635,000 | 12,576 | 10,872,790 | 126,357 | 7,444,105 | 95,615 | 7,329,636 | 124,590 |
| Grand total | 1,822,153 | 32,754 | 15,732,507 | 411,451 | 14,262,630 | 367,238 | 7,717,596 | 157,222 |

Fisheries of the Mississippi River and tributaries, 1931—Continued

CATCH: BY STATES—Continued

| Species | Iowa | | Kansas | | Kentucky | | Louisiana | |
|------------------------------|------------------|----------------|----------------|---------------|------------------|---------------|-------------------|----------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | |
| Bowfin | 91,825 | \$3,759 | | | | | 5,715 | \$114 |
| Buffalofish | 746,615 | 59,705 | 24,325 | \$2,222 | 184,558 | \$14,429 | 8,784,314 | 263,261 |
| Carp | 1,594,244 | 80,134 | 117,489 | 10,956 | 113,461 | 8,124 | 204,743 | 4,127 |
| Catfish and bullheads | 467,340 | 48,593 | 770 | 111 | 131,777 | 17,043 | 6,602,987 | 528,579 |
| Eels | 325 | 15 | | | | | 200 | 6 |
| Garfish | | | | | | | 72,450 | 791 |
| Mooneye | 1,100 | 28 | | | 990 | 105 | | |
| Paddlefish or spoonbill cat | 9,400 | 638 | | | 18,322 | 1,617 | 495,544 | 21,508 |
| Pike or pickerel | 4,700 | 470 | | | | | | |
| Quillback or "American carp" | 60,450 | 1,339 | 100 | 11 | 11,355 | 984 | 20,700 | 431 |
| Sauger | | | | | 2,365 | 451 | | |
| Sheepshead | 343,449 | 17,619 | | | 52,560 | 6,762 | 1,976,600 | 39,577 |
| Sturgeon, shovelnose | 17,650 | 1,663 | 175 | 24 | 2,987 | 380 | | |
| Sucker "mullet" | 36,550 | 822 | | | 10,294 | 1,331 | | |
| Yellow pike | | | | | 70 | 18 | | |
| Total | 3,373,648 | 214,785 | 142,859 | 13,324 | 508,719 | 51,244 | 18,163,253 | 858,394 |
| SHELLFISH, ETC. | | | | | | | | |
| Crawfish | | | | | | | 29,248 | 292 |
| Shrimp | | | | | | | 38,503 | 2,423 |
| Mussel shells | 4,366,219 | 65,685 | 312,562 | 2,713 | 1,113,032 | 8,780 | 50,000 | 375 |
| Pearls | | 7,244 | | | | | | |
| Slugs | | 13,924 | | 636 | | 852 | | |
| Frogs | | | | | | | 872,651 | 130,612 |
| Terrapin | 19,100 | 377 | | | | | | |
| Turtles: | | | | | | | | |
| Snapper | 2,000 | 40 | | | | | 58,013 | 2,244 |
| Soft-shell | 17,000 | 340 | | | | | 1,700 | 34 |
| Total | 4,404,319 | 87,610 | 312,562 | 3,340 | 1,113,032 | 9,638 | 1,050,115 | 135,980 |
| Grand total | 7,777,967 | 302,395 | 455,421 | 16,673 | 1,621,751 | 60,882 | 19,213,368 | 994,374 |

| Species | Minnesota | | Mississippi | | Missouri | | Nebraska | |
|------------------------------|------------------|----------------|------------------|----------------|----------------|---------------|----------------|---------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | |
| Bowfin | 16,598 | \$282 | | | 17,000 | \$520 | | |
| Buffalofish | 257,431 | 15,092 | 1,511,126 | \$63,824 | 178,091 | 16,414 | 18,104 | \$1,813 |
| Carp | 2,151,119 | 97,756 | 225,276 | 6,730 | 433,117 | 33,355 | 63,032 | 6,305 |
| Catfish and bullheads | 53,804 | 4,841 | 635,049 | 42,384 | 91,430 | 15,467 | 34,174 | 5,135 |
| Eels | | | 250 | 20 | 1,055 | 53 | | |
| Minnows | | | | | 325 | 209 | | |
| Paddlefish or spoonbill cat | | | 158,821 | 5,879 | 40,103 | 2,917 | | |
| Quillback or "American carp" | 17,246 | 519 | 2,157 | 42 | 13,672 | 946 | | |
| Sheepshead | 152,545 | 7,938 | 106,844 | 2,576 | 38,186 | 3,773 | | |
| Sturgeon, shovelnose | 1,634 | 115 | 100 | 3 | 17,282 | 1,703 | | |
| Sucker "mullet" | 65,273 | 1,955 | | | 2,275 | 292 | | |
| Total | 2,715,650 | 128,498 | 2,639,623 | 121,458 | 833,636 | 75,670 | 145,310 | 16,253 |
| SHELLFISH, ETC. | | | | | | | | |
| Shrimp | | | 10,000 | 1,500 | | | | |
| Mussel shells | 782,630 | 7,827 | | | 94,000 | 1,193 | | |
| Pearls | | 157 | | | | | | |
| Slugs | | 1,174 | | | | 118 | | |
| Turtles, snapper | | | 100 | 3 | | | | |
| Total | 782,630 | 9,158 | 10,100 | 1,503 | 94,000 | 1,311 | | |
| Grand total | 3,498,280 | 137,656 | 2,649,723 | 122,961 | 927,636 | 76,981 | 145,310 | 16,253 |

Fisheries of the Mississippi River and tributaries, 1931—Continued

CATCH: BY STATES—Continued

| Species | Ohio | | Oklahoma | | South Dakota | | Tennessee | |
|-----------------------------------|----------------|--------------|---------------|--------------|----------------|---------------|------------------|----------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | | | |
| Black bass..... | | | | | | | 14,000 | \$1,680 |
| Buffalofish..... | 6,433 | \$662 | 21,605 | \$2,142 | 38,926 | \$3,894 | 478,592 | 34,247 |
| Carp..... | 14,370 | 1,643 | 4,268 | 425 | 52,836 | 2,642 | 247,841 | 9,594 |
| Catfish and bullheads..... | 4,380 | 811 | 4,935 | 695 | 13,500 | 3,528 | 271,753 | 24,750 |
| Crappie..... | | | | | 1,392 | 70 | 18,652 | 1,658 |
| Eels..... | | | | | | | 163 | 25 |
| Paddlefish or spoonbill cat..... | | | 5,332 | 533 | 400 | 40 | 5,034 | 301 |
| Quillback or "American carp"..... | 1,195 | 119 | 1,950 | 195 | 4,364 | 220 | 6,065 | 843 |
| Sheepshead..... | 1,318 | 224 | 1,550 | 155 | 697 | 70 | 197,670 | 10,465 |
| Sturgeon, shovelnose..... | 558 | 72 | | | | | 3,706 | 393 |
| Sucker "mullet"..... | 2,902 | 268 | | | 2,240 | 112 | 8,323 | 1,119 |
| Sunfish..... | | | | | | | 21,850 | 1,094 |
| White bass..... | | | | | | | 2,100 | 108 |
| Yellow pike..... | 325 | 60 | | | | | | |
| Total..... | 31,481 | 3,759 | 39,640 | 4,145 | 114,361 | 10,576 | 1,275,749 | 86,275 |
| SHELLFISH, ETC. | | | | | | | | |
| Mussel shells..... | 154,000 | 3,005 | | | | | 2,157,000 | 15,604 |
| Pearls..... | | | | | | | | 28 |
| Slugs..... | | 308 | | | | | | 1,724 |
| Frogs..... | | | | | | | 2,250 | 270 |
| Terrapin..... | | | | | | | 70 | 14 |
| Total..... | 154,000 | 3,313 | | | | | 2,159,320 | 17,640 |
| Grand total..... | 185,481 | 7,072 | 39,640 | 4,145 | 114,361 | 10,576 | 3,435,069 | 103,915 |

| Species | Texas | | Wisconsin | | Total | |
|-----------------------------------|----------------|--------------|------------------|---------------|-------------------|------------------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| FISH | | | | | | |
| Black bass..... | | | | | 14,000 | \$1,680 |
| Bowfin..... | | | 288,170 | \$4,355 | 428,316 | 9,299 |
| Buffalofish..... | 75,000 | \$2,190 | 268,001 | 13,528 | 15,772,451 | 687,288 |
| Carp..... | 6,900 | 138 | 777,474 | 23,800 | 11,891,781 | 455,399 |
| Catfish and bullheads..... | 47,800 | 3,824 | 65,539 | 5,825 | 10,266,847 | 877,798 |
| Crappie..... | | | | | 41,141 | 2,959 |
| Eels..... | | | | | 6,978 | 441 |
| Garfish..... | | | | | 72,450 | 791 |
| Minnows..... | | | | | 525 | 209 |
| Mooneye..... | | | | | 3,090 | 153 |
| Paddlefish or spoonbill cat..... | | | | | 961,462 | 43,134 |
| Pike or pickerel..... | | | | | 4,700 | 470 |
| Quillback or "American carp"..... | 500 | 10 | 66,353 | 2,032 | 268,438 | 11,286 |
| Sauger..... | | | | | 2,365 | 451 |
| Sheepshead..... | 10,300 | 206 | 84,409 | 3,692 | 3,904,844 | 142,938 |
| Sturgeon, shovelnose..... | | | | | 87,426 | 8,163 |
| Sucker "mullet"..... | | | 135,984 | 3,696 | 314,835 | 12,682 |
| Sunfish..... | | | | | 21,850 | 1,094 |
| White bass..... | | | | | 3,300 | 198 |
| Yellow pike..... | | | | | 4,945 | 771 |
| Total..... | 138,500 | 6,368 | 1,685,930 | 56,928 | 44,061,714 | 2,257,204 |
| SHELLFISH, ETC. | | | | | | |
| Crawfish..... | | | | | 29,248 | 292 |
| Shrimp..... | | | | | 48,503 | 3,923 |
| Mussel shells..... | | | 959,200 | 8,946 | 37,254,697 | 421,611 |
| Pearls..... | | | | 555 | | 11,436 |
| Slugs..... | | | | 2,012 | | 68,216 |
| Frogs..... | | | | | 874,901 | 130,882 |
| Terrapin..... | | | | | 19,170 | 391 |
| Turtles: | | | | | | |
| Snapper..... | | | | | 75,190 | 3,008 |
| Soft-shell..... | | | | | 19,100 | 394 |
| Total..... | | | 959,200 | 11,513 | 38,320,809 | 640,163 |
| Grand total..... | 138,500 | 6,368 | 2,645,130 | 68,441 | 82,382,523 | 2,897,357 |

Industries related to the fisheries of the Mississippi River and tributaries

OPERATING UNITS, SALARIES, AND WAGES, 1931

| Item | Arkansas | Illinois | Indiana | Iowa | Kentucky | Louisiana | Minnesota and North Dakota |
|--------------------------------------|-----------------|------------------|-----------------|--------------------|------------------|-----------------|----------------------------|
| Transporting: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| Persons engaged..... | 13 | | | | 11 | 5 | |
| Vessels, motor..... | 4 | | | | 2 | 2 | |
| Net tonnage..... | 69 | | | | 15 | 20 | |
| Wholesale and manufacturing: | | | | | | | |
| Establishments..... | 6 | 38 | 4 | 61 | 11 | 22 | 13 |
| Persons engaged: | | | | | | | |
| Proprietors..... | 3 | 42 | 1 | 52 | 8 | 24 | 11 |
| Salaried employees..... | 9 | 3 | 5 | 79 | 20 | 14 | 27 |
| Wage earners: | | | | | | | |
| Average for season..... | 152 | 331 | 140 | 2,500 | 261 | 70 | 112 |
| Average for year..... | 72 | 235 | 93 | 2,179 | 159 | 68 | 112 |
| Paid to salaried employees..... | \$11,417 | \$9,520 | \$6,820 | \$141,346 | \$33,159 | \$12,400 | \$55,200 |
| Paid to wage earners..... | \$53,503 | \$145,683 | \$51,444 | \$1,417,678 | \$81,643 | \$37,700 | \$81,500 |
| Total salaries and wages..... | \$64,920 | \$155,203 | \$58,264 | \$1,559,024 | \$114,802 | \$50,100 | \$136,700 |
| Fishermen manufacturing..... | | 4 | 2 | | | 200 | |

| Item | Mississippi | Missouri and Oklahoma | Nebraska and Kansas | Ohio and Pennsylvania | Tennessee | Wisconsin | Total |
|--------------------------------------|-----------------|-----------------------|---------------------|-----------------------|-----------------|-----------------|--------------------|
| Transporting: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| Persons engaged..... | | | | | | | 29 |
| Vessels, motor..... | | | | | | | 8 |
| Net tonnage..... | | | | | | | 104 |
| Wholesale and manufacturing: | | | | | | | |
| Establishments..... | 6 | 21 | 3 | 13 | 11 | 8 | 217 |
| Persons engaged: | | | | | | | |
| Proprietors..... | 7 | 24 | 3 | 17 | 9 | 3 | 204 |
| Salaried employees..... | 3 | 125 | 8 | 37 | 15 | 10 | 355 |
| Wage earners: | | | | | | | |
| Average for season..... | 26 | 326 | 52 | 175 | 90 | 38 | 4,275 |
| Average for year..... | 26 | 261 | 52 | 145 | 52 | 29 | 3,453 |
| Paid to salaried employees..... | \$16,000 | \$291,874 | \$17,400 | \$95,878 | \$34,884 | \$12,998 | \$738,896 |
| Paid to wage earners..... | \$22,382 | \$202,944 | \$45,580 | \$138,817 | \$38,177 | \$24,483 | \$2,341,534 |
| Total salaries and wages..... | \$38,382 | \$494,818 | \$62,980 | \$234,695 | \$73,061 | \$37,481 | \$3,080,430 |
| Fishermen manufacturing..... | 7 | 3 | | | | | 216 |

PRODUCTS MANUFACTURED

| Item | Indiana | | Iowa | | Kentucky, Illinois, and Missouri | |
|---|----------|-------|------------|------------------|----------------------------------|----------------|
| | Quantity | Value | Quantity | Value | Quantity | Value |
| By manufacturing establishments: | | | | | | |
| Salmon, smoked..... pounds | | | 240,000 | \$48,800 | | |
| Sturgeon, smoked..... do | | | 162,000 | 39,140 | | |
| Mussel-shell products: | | | | | | |
| Buttons ¹ gross | | | 13,698,445 | 3,139,905 | | (1) |
| Poultry feed ² tons | | | 7,575 | 59,639 | | (1) |
| Lime ³ do | | | 2,524 | 8,274 | | (1) |
| Unclassified ⁴ | | | | \$ 156,210 | | \$112,808 |
| Unclassified, smoked..... pounds | | | \$ 83,000 | \$ 14,440 | | |
| Total..... | | | | 3,466,408 | | 112,808 |

¹ The production of this item is included under unclassified products.

² Data are for 1933.

³ Includes the production of mussel-shell stucco, novelties, and colored shells and chips.

⁴ Includes the production of mussel-shell buttons, poultry feed, and lime.

⁵ Includes the production of smoked buffalo fish and tullibees.

Industries related to the fisheries of the Mississippi River and tributaries—Continued

PRODUCTS MANUFACTURED—Continued

| Item | Indiana | | Iowa | | Kentucky, Illinois, and Missouri | |
|------------------------------------|------------|------------|----------|--------------------|----------------------------------|----------------|
| | Quantity | Value | Quantity | Value | Quantity | Value |
| By fishermen: | | | | | | |
| Carp, smoked.....pounds..... | | | | | 667 | \$67 |
| Paddlefish roe, salted.....do..... | 450 | \$180 | | | 900 | 540 |
| Sheepshead, smoked.....do..... | | | | | 617 | 77 |
| Sturgeon: | | | | | | |
| Smoked.....do..... | | | | | 1,333 | 400 |
| Roe, salted.....do..... | | | | | 35 | 23 |
| Total..... | 450 | 180 | | | 3,552 | 1,116 |
| Grand total..... | 450 | 180 | | \$3,466,408 | | 113,926 |

| Item | Louisiana | | Minnesota and Nebraska | | Mississippi | | Tennessee, Ohio, and Pennsylvania | |
|---|---------------|--------------|------------------------|---------------|-------------|-----------|-----------------------------------|---------------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| By manufacturing establishments: | | | | | | | | |
| Salmon, smoked.....pounds..... | | | (1) | (1) | | | | |
| Sturgeon, smoked.....do..... | | | (1) | (1) | | | | |
| Whitefish smoked.....do..... | | | 255,000 | \$47,200 | | | | |
| Unclassified, smoked.....lbs..... | | | \$ 66,600 | \$ 19,793 | | | 291,500 | \$77,206 |
| Total..... | | | 321,600 | 66,993 | | | 291,500 | 77,206 |
| By fishermen: | | | | | | | | |
| Alligator hides.....pounds..... | 88,356 | \$7,363 | | | | | | |
| Paddlefish roe, salted.....do..... | | | | | 245 | \$92 | | |
| Total..... | 88,356 | 7,363 | | | 245 | 92 | | |
| Grand total..... | 88,356 | 7,363 | 321,600 | 66,993 | 245 | 92 | 291,500 | 77,206 |

¹ The production of this item is included under unclassified products.

² Includes the production of smoked eels, salmon, and sturgeon.

³ Includes the production of smoked buffalohead, butterfish, carp, chubs, lake trout, paddlefish, sablefish, salmon, and tullibees.

NOTE.—Unless otherwise indicated the data are for 1931. The total value of the manufactured products for the States of the Mississippi River and tributaries was as follows: By manufacturing establishments, \$3,723,414; and by fishermen, \$9,751. Some of the above products may have been manufactured from fishery products imported from another State or a foreign country; therefore, they cannot be correlated directly with the catch within the State.

LAKE PEPIN

Fisheries of Lake Pepin, 1933

OPERATING UNITS: BY GEAR

| Item | Haul seines | Anchor gill nets | Trot lines | Pound nets | Fyke nets | Total, exclusive of duplication |
|--------------------|-------------|------------------|------------|------------|-----------|---------------------------------|
| Fishermen: | | | | | | |
| Regular..... | Number 8 | Number 14 | Number 2 | Number 1 | Number 5 | Number 9 |
| Casual..... | 40 | 14 | 2 | 1 | 20 | 44 |
| Total..... | 48 | 14 | 2 | 2 | 26 | 53 |
| Boats: | | | | | | |
| Motor..... | 16 | 7 | | 2 | 17 | 23 |
| Other..... | 16 | 7 | 2 | 2 | 12 | 23 |
| Apparatus: | | | | | | |
| Number..... | 80 | 17 | 2 | 4 | 150 | |
| Length, yards..... | 4,875 | | | | | |
| Square yards..... | | 27,000 | | | | |
| Hooks..... | | | 600 | | | |

Fisheries of Lake Pepin, 1933—Continued

CATCH: BY GEAR

| Species | Haul seines | | Anchor gill nets | | Trot lines | |
|----------------------------|-------------|--------|------------------|-------|------------|-------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Bowfin..... | 10,300 | \$206 | | | | |
| Buffalofish..... | 9,700 | 388 | 8,000 | \$320 | | |
| Carp..... | 427,000 | 12,410 | 80,000 | 2,400 | 1,000 | \$30 |
| Catfish and bullheads..... | 8,450 | 845 | | | 3,000 | 300 |
| Sheepshead..... | 2,500 | 100 | | | 1,000 | 40 |
| Sucker "mullet"..... | 31,000 | 620 | | | | |
| Turtles: | | | | | | |
| Snapper..... | 1,000 | 20 | | | 200 | 4 |
| Soft-shell..... | 500 | 10 | | | | |
| Total..... | 490,450 | 14,599 | 88,000 | 2,720 | 5,200 | 374 |

| Species | Pound nets | | Fyke nets | | Total | |
|----------------------------|------------|-------|-----------|-------|---------|--------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Bowfin..... | | | | | 10,300 | \$206 |
| Buffalofish..... | 5,000 | \$200 | 19,000 | \$760 | 41,700 | 1,668 |
| Carp..... | 25,000 | 750 | 102,000 | 3,080 | 635,000 | 18,650 |
| Catfish and bullheads..... | 5,000 | 500 | 21,000 | 2,100 | 87,450 | 8,745 |
| Sheepshead..... | | | 5,500 | 220 | 9,000 | 360 |
| Sucker "mullet"..... | 5,000 | 100 | | | 36,000 | 720 |
| Turtles: | | | | | | |
| Snapper..... | | | | | 1,200 | 24 |
| Soft-shell..... | | | | | 500 | 10 |
| Total..... | 40,000 | 1,550 | 147,500 | 6,140 | 771,150 | 25,383 |

OPERATING UNITS: BY STATES

| Item | Minnesota | Wisconsin | Total for lake |
|------------------------|-----------|-----------|----------------|
| | Number | Number | Number |
| Fishermen: | | | |
| Regular..... | | 9 | 9 |
| Casual..... | 8 | 36 | 44 |
| Total..... | 8 | 45 | 53 |
| Boats: | | | |
| Motor..... | 2 | 20 | 22 |
| Other..... | 4 | 19 | 23 |
| Apparatus: | | | |
| Haul seines..... | 2 | 28 | 30 |
| Length, yards..... | 400 | 4,475 | 4,875 |
| Gill nets, anchor..... | | 17 | 17 |
| Square yards..... | | 27,000 | 27,000 |
| Lines, trot..... | 2 | | 2 |
| Hooks..... | 600 | | 600 |
| Pound nets..... | | 4 | 4 |
| Fyke nets..... | | 150 | 150 |

CATCH: BY STATES

| Species | Minnesota | | Wisconsin | | Total for lake | |
|----------------------------|-----------|-------|-----------|--------|----------------|--------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Bowfin..... | 500 | \$10 | 9,800 | \$196 | 10,300 | \$206 |
| Buffalofish..... | 200 | 8 | 41,500 | 1,660 | 41,700 | 1,668 |
| Carp..... | 65,000 | 1,950 | 570,000 | 16,700 | 635,000 | 18,650 |
| Catfish and bullheads..... | 3,650 | 365 | 33,800 | 3,380 | 37,450 | 3,745 |
| Sheepshead..... | 2,500 | 100 | 6,500 | 260 | 9,000 | 360 |
| Sucker "mullet"..... | 1,500 | 30 | 34,500 | 690 | 36,000 | 720 |
| Turtles: | | | | | | |
| Snapper..... | 200 | 4 | 1,000 | 20 | 1,200 | 24 |
| Soft-shell..... | | | 500 | 10 | 500 | 10 |
| Total..... | 73,550 | 2,467 | 697,600 | 22,916 | 771,150 | 25,383 |

LAKE KEOKUK

Fisheries of Lake Keokuk, 1933

OPERATING UNITS: BY GEAR

| Item | Haul seines | Trot lines | Fyke nets | Total, exclusive of duplication |
|--------------------|---------------|---------------|---------------|---------------------------------|
| Fishermen: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| Regular..... | 20 | 23 | 28 | 28 |
| Casual..... | 45 | 20 | 62 | 68 |
| Total..... | 65 | 20 | 85 | 96 |
| Boats: | | | | |
| Motor..... | 21 | 5 | 32 | 33 |
| Other..... | 32 | 18 | 50 | 54 |
| Apparatus: | | | | |
| Number..... | 30 | 60 | 555 | ----- |
| Length, yards..... | 3,400 | ----- | ----- | ----- |
| Hooks..... | ----- | 18,000 | ----- | ----- |

CATCH: BY GEAR

| Species | Haulsines | | Trotlines | | Fyke nets | | Total | |
|----------------------------|----------------|--------------|---------------|--------------|----------------|---------------|----------------|---------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Bowfin..... | 2,500 | \$50 | ----- | ----- | ----- | ----- | 2,500 | \$50 |
| Buffalofish..... | 14,000 | 555 | ----- | ----- | 61,800 | \$2,422 | 75,800 | 2,977 |
| Carp..... | 120,000 | 3,350 | 2,000 | \$50 | 91,000 | 2,680 | 213,000 | 6,080 |
| Catfish and bullheads..... | 5,000 | 500 | 9,000 | 900 | 74,500 | 7,450 | 88,500 | 8,850 |
| Sheepshead..... | 24,500 | 980 | 1,500 | 60 | 61,500 | 2,460 | 87,500 | 3,500 |
| Sturgeon, shovelnose..... | 875 | 72 | ----- | ----- | ----- | ----- | 875 | 72 |
| Sucker "mullet"..... | 300 | 6 | ----- | ----- | 1,600 | 82 | 1,900 | 38 |
| Turtles: | | | | | | | | |
| Snapper..... | 5,000 | 100 | ----- | ----- | 150 | 3 | 5,150 | 103 |
| Soft-shell..... | 2,000 | 40 | ----- | ----- | ----- | ----- | 2,000 | 40 |
| Total..... | 174,175 | 5,653 | 12,500 | 1,010 | 290,550 | 15,047 | 477,225 | 21,710 |

OPERATING UNITS: BY STATES

| Item | Illinois | Iowa | Total for lake |
|--------------------|---------------|---------------|----------------|
| Fishermen: | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| Regular..... | 7 | 21 | 28 |
| Casual..... | 28 | 40 | 68 |
| Total..... | 35 | 61 | 96 |
| Boats: | | | |
| Motor..... | 12 | 21 | 33 |
| Other..... | 22 | 32 | 54 |
| Apparatus: | | | |
| Haul seines..... | 7 | 23 | 30 |
| Length, yards..... | 700 | 2,700 | 3,400 |
| Lines, trot..... | 24 | 36 | 60 |
| Hooks..... | 7,200 | 10,800 | 18,000 |
| Fyke nets..... | 240 | 315 | 555 |

CATCH: BY STATES

| Species | Illinois | | Iowa | | Total for lake | |
|----------------------------|----------------|--------------|----------------|---------------|----------------|---------------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Bowfin..... | ----- | ----- | 2,500 | \$50 | 2,500 | \$50 |
| Buffalofish..... | 9,300 | \$372 | 66,500 | 2,605 | 75,800 | 2,977 |
| Carp..... | 76,000 | 2,280 | 137,000 | 3,800 | 213,000 | 6,080 |
| Catfish and bullheads..... | 42,700 | 4,270 | 45,800 | 4,580 | 88,500 | 8,850 |
| Sheepshead..... | 18,500 | 740 | 69,000 | 2,760 | 87,500 | 3,500 |
| Sturgeon, shovelnose..... | ----- | ----- | 875 | 72 | 875 | 72 |
| Sucker "mullet"..... | 1,000 | 20 | 900 | 18 | 1,900 | 38 |
| Turtles: | | | | | | |
| Snapper..... | 1,150 | 23 | 4,000 | 80 | 5,150 | 103 |
| Soft-shell..... | ----- | ----- | 2,000 | 40 | 2,000 | 40 |
| Total..... | 148,650 | 7,705 | 328,575 | 14,005 | 477,225 | 21,710 |

MISSISSIPPI RIVER BETWEEN LAKE PEPIN AND LAKE KEOKUK

Fisheries of the Mississippi River between Lake Pepin and Lake Keokuk, 1935

OPERATING UNITS: BY GEAR

| Item | Haul seines | Anchor gill nets | Trammel nets | Trot lines | Fyke nets | Dip nets | Total, exclusive of duplication |
|--------------------|---------------|------------------|---------------|---------------|---------------|---------------|---------------------------------|
| Fishermen: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| Regular..... | 50 | | | | 100 | 22 | 128 |
| Casual..... | 324 | 7 | 32 | 144 | 451 | 50 | 598 |
| Total..... | 374 | 7 | 32 | 144 | 551 | 72 | 721 |
| Boats: | | | | | | | |
| Motor..... | 135 | 7 | 32 | 20 | 330 | | 351 |
| Other..... | 125 | | | 126 | 181 | | 309 |
| Apparatus: | | | | | | | |
| Number..... | 160 | 7 | 32 | 144 | 10,960 | 72 | |
| Length, yards..... | 17,600 | | | | | | |
| Square yards..... | | 1,400 | 5,200 | | | | |
| Hooks..... | | | | 24,236 | | | |

CATCH: BY GEAR

| Species | Haul seines | | Anchor gill nets | | Trammel nets | | Trot lines | |
|----------------------------------|------------------|---------------|------------------|------------|---------------|--------------|----------------|--------------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| Bowfin..... | 56,000 | \$1,120 | | | | | | |
| Buffalofish..... | 257,000 | 10,287 | 4,000 | \$160 | 18,000 | \$720 | | |
| Carp..... | 1,346,300 | 39,864 | 6,000 | 180 | 19,000 | 570 | 40,700 | \$1,329 |
| Catfish and bullheads..... | 43,200 | 4,290 | 1,000 | 100 | 3,500 | 350 | 49,550 | 4,955 |
| Eels..... | 365 | 28 | | | | | 380 | 19 |
| Gizzard shad..... | 1,000 | 20 | | | | | | |
| Mooneye..... | 500 | 10 | | | | | | |
| Paddlefish or spoonbill cat..... | 2,950 | 260 | | | | | | |
| Pike or pickerel..... | 5,750 | 248 | | | | | | |
| Sheepshead..... | 284,300 | 11,393 | 3,500 | 140 | 8,500 | 340 | 30,250 | 1,247 |
| Sturgeon, shovelnose..... | 28,700 | 2,296 | 2,200 | 176 | 19,600 | 1,568 | | |
| Sucker "mullet"..... | 129,200 | 2,668 | | | 500 | 10 | | |
| Turtles: | | | | | | | | |
| Snapper..... | 16,650 | 333 | | | | | | |
| Soft-shell..... | 8,500 | 170 | | | | | | |
| Total..... | 2,180,415 | 72,987 | 16,700 | 756 | 69,100 | 3,558 | 120,850 | 7,550 |

| Species | Fyke nets | | Dip nets | | Total | |
|----------------------------------|------------------|---------------|----------------|--------------|------------------|----------------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Bowfin..... | 1,500 | \$30 | 12,000 | \$240 | 69,500 | \$1,390 |
| Buffalofish..... | 573,600 | 22,944 | 50,000 | 2,000 | 902,600 | 36,111 |
| Carp..... | 683,000 | 20,495 | 54,000 | 1,620 | 2,149,000 | 64,068 |
| Catfish and bullheads..... | 356,700 | 35,670 | 12,500 | 1,250 | 465,450 | 46,618 |
| Eels..... | | | | | 715 | 47 |
| Gizzard shad..... | | | | | 1,000 | 20 |
| Mooneye..... | | | | | 500 | 10 |
| Paddlefish or spoonbill cat..... | | | 3,000 | 240 | 5,950 | 500 |
| Pike or pickerel..... | | | | | 5,750 | 248 |
| Sheepshead..... | 291,500 | 11,720 | 82,000 | 3,280 | 700,050 | 28,120 |
| Sturgeon, shovelnose..... | | | 12,700 | 1,016 | 63,200 | 5,056 |
| Sucker "mullet"..... | 11,400 | 220 | | | 141,100 | 2,898 |
| Turtles: | | | | | | |
| Snapper..... | 300 | 6 | | | 16,950 | 339 |
| Soft-shell..... | | | | | 8,500 | 170 |
| Total..... | 1,918,000 | 91,085 | 226,200 | 9,646 | 4,531,265 | 185,583 |

Fisheries of the Mississippi River between Lake Pepin and Lake Keokuk, 1933—Con.

OPERATING UNITS: BY STATES

| Item | Illinois | Iowa | Minnesota | Wisconsin | Total |
|--------------------|------------|------------|-----------|------------|------------|
| | Number | Number | Number | Number | Number |
| Fishermen: | | | | | |
| Regular..... | 23 | 85 | ----- | 18 | 126 |
| Casual..... | 130 | 260 | 73 | 182 | 595 |
| Total..... | 153 | 345 | 73 | 150 | 721 |
| Boats: | | | | | |
| Motor..... | 73 | 163 | 28 | 97 | 361 |
| Other..... | 81 | 128 | 45 | 55 | 309 |
| Apparatus: | | | | | |
| Haul seines..... | 29 | 72 | 17 | 42 | 160 |
| Length, yards..... | 3,150 | 8,000 | 2,260 | 4,200 | 17,600 |
| Gill nets..... | ----- | ----- | ----- | 7 | 7 |
| Square yards..... | ----- | ----- | ----- | 1,400 | 1,400 |
| Trammel nets..... | ----- | 24 | ----- | 8 | 32 |
| Square yards..... | ----- | 4,133 | ----- | 1,067 | 5,200 |
| Lines, trot..... | 38 | 49 | 45 | 12 | 144 |
| Hooks..... | 5,800 | 8,700 | 8,136 | 1,600 | 24,236 |
| Fyke nets..... | 2,145 | 5,775 | 360 | 2,680 | 10,960 |
| Dip nets..... | ----- | 72 | ----- | ----- | 72 |

CATCH: BY STATES

| Species | Illinois | | Iowa | | Minnesota | |
|----------------------------------|----------------|---------------|------------------|---------------|----------------|---------------|
| | Pounds | Value | Pounds | Value | Pounds | Value |
| Bowfin..... | 2,500 | \$50 | 27,800 | \$556 | 4,500 | \$90 |
| Buffalofish..... | 106,800 | 6,672 | 445,200 | 17,808 | 42,800 | 1,719 |
| Carp..... | 351,600 | 10,563 | 919,700 | 27,640 | 230,400 | 6,466 |
| Catfish and bullheads..... | 107,950 | 10,795 | 231,650 | 23,165 | 38,100 | 3,780 |
| Eels..... | ----- | ----- | 715 | 47 | ----- | ----- |
| Gizzard shad..... | ----- | ----- | 1,000 | 20 | ----- | ----- |
| Mooneye..... | ----- | ----- | ----- | ----- | 500 | 10 |
| Paddlefish or spoonbill cat..... | 500 | 40 | 5,250 | 440 | ----- | ----- |
| Pike or pickeral..... | ----- | ----- | 5,550 | 240 | ----- | ----- |
| Sheepshead..... | 131,500 | 5,350 | 351,750 | 14,070 | 47,800 | 1,940 |
| Sturgeon, shovelnose..... | 10,200 | 816 | 41,500 | 3,320 | 3,000 | 240 |
| Sucker "mullet"..... | 7,000 | 134 | 47,600 | 1,084 | 31,700 | 634 |
| Turtles: | | | | | | |
| Snapper..... | 1,600 | 32 | 4,000 | 80 | 2,400 | 48 |
| Soft-shell..... | 500 | 10 | 2,750 | 55 | ----- | ----- |
| Total..... | 780,150 | 34,462 | 2,084,465 | 83,475 | 401,200 | 14,937 |

Fisheries of the Mississippi River between Lake Pepin and Lake Keokuk, 1933—Con.

CATCH: BY STATES—Continued

| Species | Wisconsin | | Total | |
|----------------------------------|-----------|--------|-----------|---------|
| | Pounds | Value | Pounds | Value |
| Bowfin..... | 34,700 | \$694 | 69,500 | \$1,390 |
| Buffalofish..... | 247,800 | 9,912 | 902,600 | 36,111 |
| Carp..... | 647,300 | 19,369 | 2,149,000 | 64,058 |
| Catfish and bullheads..... | 88,750 | 8,875 | 466,450 | 46,615 |
| Eels..... | | | 715 | 47 |
| Gizzard shad..... | | | 1,000 | 20 |
| Mooneye..... | | | 500 | 10 |
| Paddlefish or spoonbill cat..... | 200 | 20 | 5,950 | 500 |
| Pike or pickerel..... | 200 | 8 | 5,750 | 248 |
| Sheepshead..... | 169,000 | 6,760 | 700,050 | 28,120 |
| Sturgeon, shovelnose..... | 8,500 | 680 | 63,200 | 5,056 |
| Stucker "mullet"..... | 54,800 | 1,096 | 141,100 | 2,898 |
| Turtles: | | | | |
| Snapper..... | 8,950 | 179 | 10,950 | 339 |
| Soft-shell..... | 5,250 | 105 | 8,500 | 170 |
| Total..... | 1,265,450 | 47,698 | 4,531,265 | 185,582 |

FISHERIES OF ALASKA ¹⁴

The commercial catch of fishery products in Alaska during 1933, exclusive of whales, amounted to 627,395,274 pounds, valued at \$9,088,984, which is an increase of 5 percent in volume and 30 percent in value as compared with the catch in 1932. Of the total catch in 1933, 467,348,858 pounds, valued at \$7,498,037, consisted of salmon; 157,337,810 pounds, valued at \$1,443,414, other fish; and 2,708,606 pounds, valued at \$147,533, shellfish. In addition, 182 whales were taken. These fisheries gave employment to 8,656 fishermen, 1,283 persons on transporting craft, and 11,756 persons in fisheries wholesale and manufacturing industries—a total of 21,695 persons, which is an increase of 8 percent as compared with the number employed in 1932.

¹⁴ Statistics for the fisheries of Alaska are collected and compiled by the Alaska Division of this Bureau. A summary of these statistics appear in this section. For detailed figures the reader is referred to "Alaska Fisheries and Fur-Seal Industries in 1933" by Ward T. Bower, Appendix II to the Report of Commissioner of Fisheries for the fiscal year 1934.

Fisheries of Alaska, 1933

SUMMARY: BY DISTRICTS

| Item | Southeast Alaska | | Central Alaska | | Western Alaska | | Total | |
|--|------------------|---------------|----------------|---------------|----------------|---------------|---------------|---------------|
| | Number | Value | Number | Value | Number | Value | Number | Value |
| PERSONS ENGAGED | | | | | | | | |
| In fishing..... | 3, 879 | | 1, 888 | | 2, 889 | | 8, 656 | |
| In transporting..... | 537 | | 395 | | 351 | | 1, 283 | |
| In wholesale and manufacturing industries..... | 4, 742 | | 3, 131 | | 3, 883 | | 11, 756 | |
| Total..... | 9, 158 | | 5, 414 | | 7, 123 | | 21, 695 | |
| CRAFT EMPLOYED | | | | | | | | |
| Vessels fishing..... | 460 | | 41 | | 6 | | 507 | |
| Boats fishing..... | 1, 896 | | 1, 014 | | 1, 306 | | 4, 218 | |
| Vessels transporting..... | 121 | | 93 | | 78 | | 292 | |
| Scows, houseboats, pile drivers, etc..... | 234 | | 197 | | 187 | | 618 | |
| Total..... | 2, 711 | | 1, 345 | | 1, 579 | | 5, 635 | |
| CATCH | | | | | | | | |
| Fish: | <i>Pounds</i> | | <i>Pounds</i> | | <i>Pounds</i> | | <i>Pounds</i> | |
| Salmon..... | 169, 461, 046 | \$2, 671, 835 | 128, 249, 502 | \$2, 013, 780 | 169, 638, 310 | \$2, 812, 422 | 467, 348, 858 | \$7, 498, 037 |
| Other..... | 117, 921, 514 | 1, 242, 270 | 35, 569, 663 | 181, 531 | 3, 846, 633 | 19, 613 | 157, 337, 810 | 1, 443, 414 |
| Shellfish..... | 974, 187 | 48, 305 | 1, 734, 419 | 99, 228 | | | 2, 708, 606 | 147, 533 |
| Total..... | 288, 356, 747 | 3, 962, 410 | 165, 553, 584 | 2, 294, 539 | 173, 484, 943 | 2, 832, 035 | 627, 395, 274 | 9, 088, 984 |
| Whales..... | <i>Number</i> | | <i>Number</i> | | <i>Number</i> | | <i>Number</i> | |
| | | | 182 | | | | 182 | |
| WHOLESALE AND MANUFACTURING | | | | | | | | |
| Establishments..... | 79 | | 97 | | 48 | | 224 | |
| PRODUCTS AS PREPARED FOR MARKET | | | | | | | | |
| | <i>Pounds</i> | | <i>Pounds</i> | | <i>Pounds</i> | | <i>Pounds</i> | |
| Salmon..... | 109, 650, 847 | 10, 477, 788 | 72, 156, 995 | 7, 897, 181 | 81, 432, 367 | 11, 029, 103 | 263, 240, 209 | 29, 404, 072 |
| Herring..... | 44, 173, 201 | 806, 062 | 16, 705, 660 | 490, 837 | 2, 417, 315 | 105, 295 | 63, 296, 176 | 1, 402, 194 |
| Halibut..... | 14, 046, 850 | 724, 572 | 22, 061 | 1, 790 | | | 14, 068, 911 | 726, 362 |
| Cod..... | | | 251, 665 | 9, 757 | 86, 810 | 3, 150 | 338, 476 | 12, 907 |
| Trout..... | 37, 355 | 1, 815 | 3, 000 | 150 | | | 40, 355 | 1, 965 |
| Sablefish..... | 103, 095 | 4, 505 | | | | | 103, 095 | 4, 505 |
| Smelt..... | | | 500 | 50 | | | 500 | 50 |
| Flounder..... | 75, 000 | 1, 125 | | | | | 75, 000 | 1, 125 |
| Rockfish..... | 3, 533 | 119 | | | | | 3, 533 | 119 |
| Clam..... | 1, 992 | 361 | 521, 028 | 245, 977 | | | 523, 020 | 246, 338 |
| Shrimp..... | 309, 092 | 100, 340 | 9, 960 | 2, 042 | | | 319, 062 | 102, 382 |
| Crab..... | 219, 154 | 56, 331 | 351, 542 | 99, 249 | | | 570, 696 | 155, 580 |
| Whale..... | | | 3, 378, 125 | 68, 989 | | | 3, 378, 125 | 68, 989 |
| Total..... | 168, 620, 119 | 12, 173, 018 | 93, 400, 536 | 8, 816, 022 | 83, 936, 492 | 11, 137, 548 | 345, 957, 147 | 32, 126, 888 |

Fisheries of Alaska, 1933—Continued

OPERATING UNITS: BY DISTRICTS

| Item | Southeast Alaska | Central Alaska | Western Alaska | Total | Item | Southeast Alaska | Central Alaska | Western Alaska | Total |
|-------------------|------------------|----------------|----------------|---------------|--|------------------|----------------|----------------|---------------|
| | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | Apparatus—Continued: | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| Fishermen..... | 3, 879 | 1, 888 | 2, 889 | 8, 656 | Gill nets..... | 265 | 957 | 2, 124 | 3, 346 |
| Vessels fishing: | | | | | Yards..... | 49, 250 | 88, 900 | 313, 662 | 451, 812 |
| Steam..... | | 3 | | 3 | Beam trawls..... | 10 | 1 | | 11 |
| Net tonnage..... | | 217 | | 217 | Wheels..... | | | 280 | 280 |
| Motor..... | 460 | 38 | 6 | 504 | Lines: | | | | |
| Net tonnage..... | 6, 347 | 828 | 195 | 7, 370 | Hand lines (cod fishery)..... | | 83 | 27 | 60 |
| Boats fishing: | | | | | Trawl lines (cod fishery)..... | | 6 | 7 | 13 |
| Motor..... | 731 | 250 | 51 | 1, 032 | Troll lines (salmon fishery)..... | 3, 047 | | | 3, 047 |
| Other..... | 1, 165 | 764 | 1, 257 | 3, 186 | Skates of lines (halibut fishery)..... | 2, 341 | | | 2, 341 |
| Apparatus: | | | | | Crab pots..... | 732 | 280 | | 992 |
| Traps..... | 261 | 138 | 1 | 400 | Herring pounds..... | 4 | 12 | | 16 |
| Purse seines..... | 357 | 82 | 6 | 445 | Herring pound seines..... | 8 | 8 | | 16 |
| Yards..... | 115, 794 | 18, 216 | 2, 850 | 136, 860 | | | | | |
| Haul seines..... | 3 | 96 | 3 | 102 | | | | | |
| Yards..... | 600 | 19, 374 | 230 | 20, 204 | | | | | |

CATCH: BY DISTRICTS

[Estimated round weight and value to fishermen]

| Item | Southeast Alaska | | Central Alaska | | Western Alaska | | Total | |
|-------------------------------|------------------|--------------|----------------|--------------|----------------|---------------|---------------|---------------|
| | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> | <i>Pounds</i> | <i>Value</i> |
| FISH | | | | | | | | |
| Salmon: | | | | | | | | |
| Blueback, red or sockeye..... | 7, 073, 220 | \$146, 631 | 44, 521, 682 | \$857, 244 | 157, 730, 300 | \$2, 745, 537 | 209, 325, 102 | \$3, 749, 412 |
| Chinook or king..... | 8, 543, 780 | 194, 853 | 1, 914, 780 | 47, 075 | 1, 716, 800 | 21, 744 | 12, 174, 860 | 263, 672 |
| Chum or keta..... | 40, 926, 618 | 442, 409 | 20, 046, 636 | 221, 520 | 10, 056, 726 | 43, 208 | 71, 029, 980 | 707, 137 |
| Humpback or pink..... | 103, 132, 780 | 1, 718, 788 | 55, 956, 760 | 803, 225 | 112 | 3 | 159, 089, 652 | 2, 522, 014 |
| Silver or coho..... | 9, 784, 648 | 169, 156 | 5, 809, 744 | 84, 716 | 134, 872 | 1, 930 | 15, 729, 284 | 255, 802 |
| Herring..... | 102, 026, 831 | 510, 134 | 34, 945, 643 | 174, 728 | 3, 607, 933 | 18, 040 | 140, 550, 407 | 702, 902 |
| Halibut..... | 15, 607, 611 | 724, 572 | 24, 512 | 1, 790 | | | 16, 632, 123 | 726, 362 |
| Cod..... | | | 589, 368 | 4, 613 | 238, 700 | 1, 573 | 828, 053 | 6, 336 |
| Trout: | | | | | | | | |
| Dolly Varden..... | 35, 109 | 1, 391 | 9, 375 | 150 | | | 44, 484 | 1, 541 |
| Steelhead..... | 11, 585 | 424 | | | | | 11, 585 | 424 |
| Sablefish..... | 151, 610 | 4, 505 | | | | | 151, 610 | 4, 505 |
| Smelt..... | | | 750 | 50 | | | 750 | 50 |

| | | | | | | | | |
|-------------------------|--------------------|------------------|--------------------|------------------|--------------------|------------------|--------------------|------------------|
| Flounders..... | 83,333 | 1,125 | | | | | 83,333 | 1,125 |
| Rockfishes..... | 5,435 | 119 | | | | | 5,435 | 119 |
| Total..... | 287,382,560 | 3,914,105 | 163,819,165 | 2,195,311 | 173,484,943 | 2,832,035 | 624,686,668 | 8,941,461 |
| SHELLFISH | | | | | | | | |
| Crabs..... | 408,218 | 28,165 | 674,374 | 49,625 | | | 1,062,502 | 77,790 |
| Shrimp..... | 561,985 | 20,068 | 18,109 | 408 | | | 590,004 | 20,475 |
| Clams: | | | | | | | | |
| Butter..... | 3,984 | 72 | | | | | 3,984 | 72 |
| Razor..... | | | 1,041,936 | 49,195 | | | 1,041,936 | 49,195 |
| Total..... | 974,187 | 48,305 | 1,734,419 | 99,228 | | | 2,708,606 | 147,533 |
| Grand total..... | 288,356,747 | 3,962,410 | 165,553,584 | 2,294,539 | 173,484,943 | 2,832,035 | 627,395,274 | 9,068,984 |

NOTE.—In addition to the above, 182 whales were taken in Alaskan waters. The round weight and value to the fishermen cannot be determined, but the products amounted to 3,373,125 pounds, valued at \$68,969.

Industries related to the fisheries of Alaska, 1933

TRANSPORTING

| Item | Southeast Alaska | Central Alaska | Western Alaska | Total | Item | Southeast Alaska | Central Alaska | Western Alaska | Total |
|-----------------------|------------------|----------------|----------------|---------------|---|------------------|----------------|----------------|---------------|
| | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> | | <i>Number</i> | <i>Number</i> | <i>Number</i> | <i>Number</i> |
| Persons engaged..... | 537 | 395 | 351 | 1,283 | Vessels transporting—Continued: | | | | |
| Vessels transporting: | | | | | Motor..... | 121 | 92 | 69 | 282 |
| Steam..... | | 1 | 9 | 10 | Net tonnage..... | 3,556 | 2,919 | 4,507 | 10,962 |
| Net tonnage..... | | 1,724 | 16,643 | 18,367 | Scows, houseboats, pile drivers, etc..... | 234 | 197 | 187 | 618 |

Industries related to the fisheries of Alaska, 1933—Continued

WHOLESALE AND MANUFACTURING

| Item | Southeast Alaska | Central Alaska | Western Alaska | Total |
|---------------------------------------|------------------|-----------------|-----------------|------------------|
| Persons engaged..... | Number 4,742 | Number 3,131 | Number 3,883 | Number 11,756 |
| Establishments: | | | | |
| Handling fresh and frozen fish..... | 39 | 3 | | 42 |
| Curing fish..... | 31 | 57 | 36 | 124 |
| Canning fish..... | 42 | 43 | 23 | 108 |
| Manufacturing by-products..... | 7 | 5 | | 12 |
| Total (exclusive of duplication)..... | 79 | 97 | 48 | 224 |

PRODUCTS AS PREPARED FOR MARKET

| Item | Southeast Alaska | | Central Alaska | | Western Alaska | | Total | |
|----------------------------|------------------|----------|----------------|--------|----------------|-------|------------|----------|
| | Pounds | Value | Pounds | Value | Pounds | Value | Pounds | Value |
| FRESH | | | | | | | | |
| Salmon (for food)..... | 559,287 | \$30,601 | | | | | 559,287 | \$30,601 |
| Salmon (for bait)..... | 48,700 | 277 | 15,300 | \$100 | | | 64,000 | 377 |
| Herring (for bait)..... | 2,413,220 | 21,232 | 562,300 | 5,014 | | | 2,975,520 | 26,246 |
| Halibut..... | 8,280,476 | 415,833 | 22,061 | 1,790 | | | 8,282,537 | 417,623 |
| Trout..... | 27,822 | 1,377 | 1,500 | 120 | | | 29,322 | 1,497 |
| Sablefish..... | 8,990 | 271 | | | | | 8,990 | 271 |
| Smelt..... | | | 500 | 50 | | | 500 | 50 |
| Flounders..... | 75,000 | 1,125 | | | | | 75,000 | 1,125 |
| Rockfishes..... | 428 | 10 | | | | | 428 | 10 |
| Crabs: | | | | | | | | |
| Meat..... | 62,968 | 15,923 | 27,392 | 3,711 | | | 90,360 | 19,634 |
| Whole in shell..... | 30,090 | 665 | 28,710 | 951 | | | 58,800 | 1,616 |
| Clams, whole in shell..... | | | 120 | 25 | | | 120 | 25 |
| Shrimp: | | | | | | | | |
| Meat..... | 307,552 | 100,109 | 9,460 | 1,992 | | | 317,012 | 102,101 |
| Whole in shell..... | 1,540 | 231 | 500 | 50 | | | 2,040 | 281 |
| Total..... | 11,796,073 | 587,654 | 667,943 | 13,803 | | | 12,463,916 | 601,457 |
| FROZEN | | | | | | | | |
| Salmon (for food)..... | 4,236,252 | 221,382 | | | | | 4,236,252 | 221,382 |
| Herring (for bait)..... | 1,496,370 | 12,263 | | | | | 1,496,370 | 12,263 |
| Halibut..... | 5,786,374 | 308,739 | | | | | 5,786,374 | 308,739 |
| Trout..... | 9,533 | 438 | | | | | 9,533 | 438 |

| | | | | | | | | |
|-------------------------------|--------------------|-------------------|-------------------|------------------|-------------------|-------------------|--------------------|-------------------|
| Sablefish..... | 92,705 | 4,134 | | | | | 92,705 | 4,134 |
| Rockfishes..... | 3,105 | 109 | | | | | 3,105 | 109 |
| Total..... | 11,624,339 | 547,065 | | | | | 11,624,339 | 547,065 |
| Salmon: | CURED | | | | | | | |
| Mild-cured..... | 3,817,600 | 612,393 | | | 105,600 | \$10,435 | 3,923,200 | 622,828 |
| Pickled..... | 14,000 | 926 | 342,460 | 27,108 | 678,500 | 45,886 | 1,034,950 | 73,920 |
| Dried and dry-salted..... | | | 31,425 | 641 | 1,368,107 | 54,640 | 1,399,532 | 55,281 |
| Herring: | | | | | | | | |
| Pickled (for food): | | | | | | | | |
| Scotch-cure..... | 3,874,703 | 174,284 | 7,177,000 | 338,065 | 1,599,625 | 73,982 | 12,651,328 | 586,331 |
| Norwegian-cure..... | 31,250 | 1,750 | 6,000 | 450 | 253,700 | 11,819 | 290,950 | 14,019 |
| Roused..... | | | | | 509,790 | 17,474 | 509,790 | 17,474 |
| Spiced..... | 1,000 | 125 | | | | | 1,000 | 125 |
| Dry-salted..... | | | | | 54,200 | 2,020 | 54,200 | 2,020 |
| Cod: | | | | | | | | |
| Dry-salted..... | | | 36,620 | 1,067 | 45,810 | 1,350 | 82,430 | 2,417 |
| Pickled..... | | | 186,425 | 5,865 | 38,000 | 1,500 | 224,425 | 7,365 |
| Stockfish..... | | | 28,220 | 2,785 | 3,000 | 300 | 31,220 | 3,065 |
| Tongues..... | | | 400 | 40 | | | 400 | 40 |
| Trout, dried..... | | | 1,500 | 30 | | | 1,500 | 30 |
| Sablefish, pickled..... | 1,400 | 100 | | | | | 1,400 | 100 |
| Total..... | 7,739,953 | 789,578 | 7,810,040 | 376,051 | 4,656,332 | 219,406 | 20,206,325 | 1,385,035 |
| Salmon: | CANNED | | | | | | | |
| Blueback, red or sockeye..... | 3,894,048 | 573,847 | 23,255,232 | 3,336,299 | 77,504,304 | 10,723,142 | 104,653,584 | 14,633,288 |
| Chinook or king..... | 391,008 | 58,395 | 1,141,728 | 180,633 | 455,088 | 72,154 | 1,987,824 | 311,182 |
| Chum or keta..... | 20,393,328 | 1,729,485 | 9,978,192 | 868,067 | 1,250,352 | 115,254 | 31,621,872 | 2,712,806 |
| Humpback or pink..... | 70,944,624 | 6,729,582 | 33,817,824 | 3,144,137 | | | 104,762,448 | 9,873,719 |
| Silver or coho..... | 4,598,640 | 507,480 | 3,134,736 | 330,022 | | 69,888 | 7,803,264 | 845,019 |
| Miscellaneous fish..... | 3,360 | 520 | 9,000 | 2,647 | | 528 | 12,888 | 3,242 |
| Clams..... | 1,992 | 361 | 520,908 | 245,952 | | | 522,900 | 246,313 |
| Crabs..... | 126,096 | 39,743 | 295,440 | 94,587 | | | 421,536 | 134,330 |
| Total..... | 100,353,096 | 9,639,413 | 72,153,060 | 8,292,344 | 79,280,160 | 10,918,142 | 251,786,316 | 28,759,899 |
| Fertilizer: | BYPRODUCTS | | | | | | | |
| Salmon..... | 600,000 | 9,900 | 313,358 | 4,779 | | | 913,358 | 14,679 |
| Whale..... | | | 1,034,000 | 13,773 | | | 1,034,000 | 13,773 |
| Meal, herring..... | 17,634,860 | 277,611 | 4,496,000 | 71,911 | | | 22,030,860 | 349,522 |
| Oil: | | | | | | | | |
| Salmon..... | 150,000 | 3,000 | 117,750 | 2,748 | | | 267,750 | 5,748 |
| Herring..... | 18,821,798 | 318,797 | 4,464,360 | 75,397 | | | 23,286,158 | 394,194 |
| Whale..... | | | 2,260,125 | 53,066 | | | 2,260,125 | 53,066 |
| Sperm..... | | | 84,000 | 2,150 | | | 84,000 | 2,150 |
| Total..... | 37,106,658 | 609,308 | 12,769,593 | 223,824 | | | 49,876,251 | 833,132 |
| Grand total..... | 168,620,119 | 12,173,018 | 93,400,536 | 8,816,022 | 83,936,492 | 11,137,548 | 345,957,147 | 32,126,588 |

Norw.—Halibut products include all taken by the Alaska fleet, some of which were landed at other than Alaska ports. The total landings in Alaska in 1933 amounted to 6,779,798 pounds, valued at \$316,310 as compared with 4,562,983 pounds, valued at \$134,652, in 1932.

Supplementary table showing the pack of canned products in "Standard cases" ¹

| Item | Southeast Alaska | | Central Alaska | | Western Alaska | | Total | |
|-------------------------------|------------------|-------------|----------------|---------------|----------------|----------------|-------------|----------------|
| | Cases | Value | Cases | Value | Cases | Value | Cases | Value |
| Salmon: | | | | | | | | |
| Blueback, red or sockeye..... | 81, 128 | \$573, 847 | 484, 484 | \$3, 336, 299 | 1, 614, 673 | \$10, 723, 142 | 2, 180, 283 | \$14, 633, 288 |
| Chinook or king..... | 8, 146 | 58, 395 | 23, 786 | 180, 633 | 9, 481 | 72, 154 | 41, 413 | 311, 182 |
| Chum or keta..... | 424, 861 | 1, 729, 485 | 207, 879 | 868, 067 | 26, 049 | 115, 254 | 658, 789 | 2, 712, 806 |
| Humpback or pink..... | 1, 478, 013 | 6, 729, 682 | 704, 638 | 3, 144, 137 | ----- | ----- | 2, 182, 561 | 9, 873, 719 |
| Silver or coho..... | 95, 805 | 507, 480 | 65, 307 | 330, 022 | 1, 456 | 7, 517 | 162, 568 | 845, 019 |
| Miscellaneous fish..... | 70 | 520 | 187 | 2, 647 | 11 | 76 | 288 | 3, 242 |
| Clams..... | 133 | 361 | 34, 727 | 245, 952 | ----- | ----- | 34, 860 | 246, 313 |
| Crabs..... | 2, 627 | 39, 743 | 6, 155 | 94, 587 | ----- | ----- | 8, 782 | 134, 330 |
| Total..... | 2, 090, 781 | 9, 639, 413 | 1, 827, 063 | 8, 202, 344 | 1, 651, 670 | 10, 918, 142 | 5, 269, 514 | 28, 759, 899 |

¹ The pack of salmon, miscellaneous fish, and crabs has been converted to "standard cases" of forty-eight 1-pound cans, and clams to "standard cases" of 48 no. 1, 5-ounce cans.

Supplementary table showing the output of byproducts in tons and gallons

| Item | Southeast Alaska | | Central Alaska | | Western Alaska | | Total | |
|----------------------------|------------------|----------|----------------|----------|----------------|-------|-------------|-----------|
| | Quantity | Value | Quantity | Value | Quantity | Value | Quantity | Value |
| Fertiliser: | | | | | | | | |
| Salmon..... tons..... | 300 | \$9, 900 | 167 | \$4, 779 | ----- | ----- | 457 | \$14, 679 |
| Whale..... do..... | ----- | ----- | 517 | 13, 773 | ----- | ----- | 517 | 13, 773 |
| Meal, herring..... do..... | 8, 767 | 277, 611 | 2, 248 | 71, 911 | ----- | ----- | 11, 015 | 349, 522 |
| Oil: | | | | | | | | |
| Salmon..... gallons..... | 20, 000 | 3, 000 | 15, 700 | 2, 748 | ----- | ----- | 35, 700 | 5, 748 |
| Herring..... do..... | 2, 509, 573 | 318, 797 | 595, 248 | 75, 397 | ----- | ----- | 3, 104, 821 | 394, 194 |
| Whale..... do..... | ----- | ----- | 301, 350 | 53, 066 | ----- | ----- | 301, 350 | 53, 066 |
| Sperm..... do..... | ----- | ----- | 11, 200 | 2, 150 | ----- | ----- | 11, 200 | 2, 150 |
| Total..... | ----- | 609, 308 | ----- | 223, 824 | ----- | ----- | ----- | 833, 132 |

STATISTICAL SURVEY PROCEDURE

In order that those who use the statistical data contained in this report and previous reports of the Division of Fishery Industries may be informed as to the source of the figures and methods for their collection, it has been deemed advisable to outline in considerable detail the statistical survey procedure followed by the Division. This procedure has been developed over a period of years, and changes in method have been made at times where such changes have appeared to work toward general improvement. While the surveys in the several sections are not made in the same manner owing to varying facilities and records in different States, an attempt has been made to make the data collected by various methods in the producing areas comparable with respect to the same year as well as over a period of years. Throughout the entire plan it has been the intention to coordinate State and Federal fishery statistical work so that there will be as little duplication of effort as possible. The procedure will be discussed under two main heads—"Sectional surveys" and "Local and special surveys".

SECTIONAL SURVEYS

Statistical surveys of the fisheries and fishery industries of the various sections of the United States occupy by far the greatest part of the time of the statistical personnel of the Division. It is in the course of these surveys that the statistical and marketing agents visit the individual fishing localities of the various States to collect statistics of the volume of the catch of fish and its value, employment in fishing, quantity of fishing gear, number and classification of fishing and transporting craft, employment in wholesale and manufacturing establishments, and the volume and value of manufactured fishery products and byproducts. The various phases of these surveys are discussed in detail in the sections following.

History.—The first comprehensive statistical survey of the fisheries and fishery industries of the United States was made for the year 1880 by George Brown Goode, Assistant Director of the U. S. National Museum and associates, with the cooperation of the Commissioner of Fisheries and the Superintendent of the Tenth Census. Data for specific fisheries or restricted sections for years prior to 1880 were also collected in this early survey and recorded in Mr. Goode's reports. The survey for 1880, however, did not include the Mississippi River and tributaries. Periodic general surveys of a limited number of States or limited areas of the United States were made for various of the intervening years between 1880 and 1908 and from 1909 to 1928. In 1908 a survey of the entire United States was made. The next general survey of the entire United States was not made until 1931 although complete data for all sections excluding the Mississippi River and tributaries were collected for 1929 and 1930 and complete data on the catch and operating units of the fisheries were collected for 1932. In the latter survey, however, lack of sufficient funds prohibited collection of data on wholesale and manufacturing firms except those data collected as a part of the canned fishery products and byproducts surveys. In 1933 a complete general canvass was made of the New England, Middle Atlantic, Chesapeake, and Pacific States.

Following is a summary indicating the years for which statistics were collected on the fisheries and fishery industries in the various sections. Figures for the more recent years are available for free distribution from the Bureau in bulletin form, but figures for the earlier years are available only in the various printed reports of the Bureau. These reports are available for reference in the Bureau's library and at many public libraries.

In the New England States statistics on the catch of the marine fisheries and those conducted in the coastal rivers and bays of these States were collected for the years 1880, 1887, 1888, 1889, 1898, 1902, 1905, 1908, 1919, 1924, 1928, 1929, 1930, 1931, 1932, and 1933. For most of these years data on operating units and wholesale and manufacturing trade also were collected. In addition to the above, a partial statistical survey was made for the entire section in 1892; a partial survey of the fisheries in Maine, New Hampshire, and Massachusetts for the fiscal year 1897; the lobster fishery for 1900 and 1913; the oyster fishery for 1910; the shad and alewife fisheries for 1896; the menhaden industry for 1912; the fisheries of Massachusetts for 1879; and the fisheries of Connecticut for 1925 and 1926.

Statistics on the catch of the marine fisheries and those conducted in the coastal rivers and bays of the Middle Atlantic States were collected for the years 1880, 1887, 1888, 1889, 1890, 1891, 1897, 1898, 1901, 1904, 1908, 1921, 1926, 1929, 1930, 1931, 1932, and 1933. Data on operating units and wholesale and manufacturing trade also were collected for most of these years. In addition to these a statistical survey was made of the coastal fisheries of these States in 1915; catch in all States except New York, in 1892; the shad and alewife fisheries in 1896; the shad fisheries of the Delaware River in 1910; the shad fisheries of the Chesapeake Bay and tributaries in 1909; the menhaden industry in 1912; the lobster fisheries in 1900 and 1913; and the oyster fishery in 1911. The years for which statistics are available on the shad fishery of the Hudson River are given in the section entitled "Shad and alewife fisheries".

In the Chesapeake Bay States statistics on the catch of the marine fisheries and those conducted in coastal rivers and bays of these States were collected for the years 1880, 1887, 1888, 1890, 1891, 1897, 1901, 1904, 1908, 1920, 1925, 1929, 1930, 1931, 1932, and 1933. Data on operating units and wholesale and manufacturing trade also were collected for most of these years. In addition to the above, a statistical survey was made of the crab fishery for 1915; the oyster fishery and menhaden industry for 1912; and the shad and alewife fisheries for 1896, 1909, and 1915. The years for which statistics of the shad and alewife fisheries of the Potomac River are available are given in the section entitled "Shad and alewife fisheries".

In the South Atlantic and Gulf States statistics on the catch of the marine fisheries and those conducted in the coastal rivers and bays of these States were collected for the years 1880, 1888, 1889, 1890, 1897, 1902, 1908, 1918, 1923, 1927, 1928, 1929, 1930, 1931, and 1932. Data on operating units and wholesale and manufacturing trade also were collected for most of these years. In addition to the above, a statistical survey was made of the fisheries of these States, excluding Florida and Alabama, for 1887; the shad fishery of the South Atlantic States for 1910; the shad and alewife fisheries of the South Atlantic States for 1896; the sturgeon fishery of Florida for 1900; the menhaden

industry of the South Atlantic States for 1912; the shrimp fishery for 1916; the oyster fishery of the South Atlantic States for 1910; and the oyster fishery of the Gulf States for 1911.

In the Pacific Coast States statistics on the catch of the marine fisheries and those conducted in the coastal rivers and bays of these States were collected for the years 1880, 1888, 1892, 1895, 1899, 1904, 1908, 1915, and for all the years from 1922 to 1933, inclusive. These surveys have usually included data on operating units and wholesale and manufacturing trade. In addition to the above, statistics were obtained on the fisheries of California from 1918 to 1921, inclusive, and for the oyster fishery in 1912.

Statistics on the catch of the fisheries of the Great Lakes were collected for the years 1880, 1885, 1890, 1893, 1899, 1903, 1908, and for all the years from 1913 to 1932, inclusive. Statistics of the operating units and of the wholesale and manufacturing trade were collected for most of the years when canvasses were made from 1880 to 1908 and in 1917 and 1922 as well as in most of the years from 1926 to 1932, inclusive. In addition to the above a survey was made of the fisheries of Lake Ontario and of certain fisheries in other lakes for the year 1897.

Statistics of the catch of the fisheries of the Mississippi River and its tributaries were collected for the years 1894, 1899, 1903, 1908, 1922, and 1931. In addition figures have been obtained of the fisheries of Lakes Pepin and Keokuk for the years 1914 and 1917 and the years from 1927 to 1933, inclusive, and of the fisheries of the Mississippi River between Lakes Pepin and Keokuk for the years 1929 to 1933, inclusive.

Statistics also were collected on the fisheries of certain interior waters, other than the fisheries of the Great Lakes and the Mississippi River and its tributaries, for the years 1894, 1895, 1900, and 1902.

Statistical agents.—The statistics contained in this volume have been collected by a corps of trained statistical and marketing agents which comprises a part of the permanent staff of the Division of Fishery Industries of the Bureau. Most of these men have been with the Bureau for a period of 5 years or more. In the main they are college graduates and were recruited through civil-service examination. While in college, most of the men pursued biological or technical courses largely in fishery work which has especially suited them for coping with the many biological and technical aspects encountered in canvassing the fisheries. This training has been especially helpful in identification of the species which, because of the many local names applied to a particular species, causes considerable confusion.

Period covered.—In conducting the fishery statistical surveys, agents are dispatched to the districts to be surveyed as early in the calendar year as they can be spared from the tabulation and preparation for publication of their previous season's work. They collect statistics of fishery operations for the year preceding that in which they are working; and, since their field work occupies the greater part of the year, it is usually at least a year from the end of the calendar year for which they are collecting data until the figures are published. Most of the figures are collected for the calendar year. Where there are variations from this general practice, explanatory notes appear in the tables. Prior to 1930, statistics on the catch of oysters in the Atlantic and Gulf States were collected for the oyster season; that is, from September to

April, inclusive. Beginning with 1930 and down to the present, they have been collected on the basis of the calendar year.

Scope.—The scope of the coastal statistical surveys includes canvasses of the commercial fisheries of the oceans and bays and of the coastal rivers as far inland as commercial fishing is important. This usually coincides with the range of commercial fishing for anadromous species. Statistics of the fisheries of the Mississippi River cover canvasses of the fisheries of the Mississippi River proper as well as all of its tributaries wherein commercial fishing for either fish, crustaceans, or mollusks is prosecuted. Statistics of the fisheries of the Great Lakes cover canvasses of the fisheries prosecuted in the Lakes proper, adjacent bays, and the international lakes of northern Minnesota, as well as rivers which sustain a commercial fishery having outlets into these waters. Surveys for statistics of the wholesale and manufacturing fishery industries cover such plants located in the coastal, river, and lake areas adjacent to the waters mentioned above.

Methods of collection.—Several methods for the collection of fishery statistics are employed, each of which has been carefully studied to obtain the best results with the available personnel and funds. In most instances the agents obtain lists of the names of fishing vessels, names or numbers of motor boats, and names of owners of these craft from local customs officials. Also it often is possible to obtain the names of licensed commercial fishermen and occasionally some statistics on the catch from several of the State fishery agencies, from other State, county, or city agencies, or from private organizations.

With such preliminary records as are available for their guidance the agents then visit each fishing community in their field unless their preliminary records are so complete that personal visits in some areas may be eliminated. While it is impossible for the few agents available for this work to interview each fisherman in a given locality, the more important ones are visited and a sufficient number of those of lesser importance are interviewed to obtain reliable information on their production. In practice virtually all wholesale firms are visited as well as captains of fishing vessels (those of 5 net tons or over) and also most of the more important inshore fishermen.

In the Great Lakes and Pacific Coast States such exceptional cooperation has been obtained in recent years from the State fishery agencies in the collection of statistics that only fragmentary surveys need be made by the Bureau's agents to obtain the necessary data. Also the State fishery agencies in Maryland and Virginia recently have developed very complete statistical systems which greatly facilitate the Bureau's canvasses in these States.

As regards the fisheries of the Great Lakes and international lakes of northern Minnesota the Bureau obtains most of the catch statistics and usually the value of the catch direct from the records of the State fishery agency. To obtain data on the number of fishermen, boats, vessels, and gear the Bureau conducts such personal surveys among the fishermen as may be necessary to supplement the State records.

Bureau agents are stationed at Seattle, Wash., and Terminal Island, Calif., who survey the fisheries of the Pacific Coast States. As a rule they obtain figures on the volume of the catch from the records of the several State fishery agencies. In most cases the value of the catch is derived from dealers' records and sometimes from estimates

of prices. In Washington and Oregon the offshore fisheries are surveyed separately by the Bureau's agent to obtain the number of operating units, catch, and value of the catch. Statistics of the wholesale fishery industry for this section are obtained largely by personal interviews of the agents.

In the administration of the Alaska fisheries the Bureau obtains sworn statements concerning their activities from those prosecuting the fisheries in this area. These statements are compiled by the Alaska Division of this Bureau.

Statistics of the volume of the catch of fish of the Pacific Coast States are usually shown in weights as landed which may be in the round or dressed condition. Statistics on the volume of the catch of fish taken in the remainder of the United States are shown in round weight.

The figures in the tables for shellfish represent the weight of the meats in the case of univalve and bivalve mollusks and gastropods, and the round weight of crustaceans and such mollusks as squid and octopus.

Shore and vessel fisheries.—In general statistics of the shore fisheries as collected by the agents include data on the number of casual and regular fishermen; number and tonnage of motor and other fishing boats and accessory boats; kind and quantity of gear used, and the volume, value, and method of capture of each species caught by boats (for our purpose craft of less than 5 net tons capacity are called "boats") for each locality or group of localities. This method is not followed in some sections where the availability of data collected by the State fishery agencies obviates the necessity of detailed locality surveys.

Statistics of the vessel fisheries include data on the number of the crew, rig of vessel, net tonnage, kind and quantity of gear used, accessory boats carried and volume, value, and method of capture of each species caught by each vessel (for our purpose craft of 5 net tons capacity or more are called "vessels"). As in the shore fisheries the availability of figures collected by State fishery agencies may eliminate the necessity of our agents collecting these data for each vessel.

All persons engaged in commercial fishing operations are included as fishermen. For our purpose these have been divided into "regular" and "casual" fishermen. Regular fishermen are those who receive more than one-half of their annual income from fishing; and casual fishermen are those whose principal business is something other than fishing, and who receive less than one-half of their annual compensation from fishing.

The catch of fish is credited to the principal port of arrival and departure of the craft rather than its point of ownership, registration, documentation, or its port of landing. This accounts for catches of fish being shown in areas where they are not common since fishing vessels frequently fish in areas far from their principal fishing port.

Wholesale and manufacturing trade.—All persons or firms engaged in the wholesale buying and selling of fishery products or who produce manufactured fishery products are surveyed under this title. Where the business of fishing and wholesaling or manufacturing is combined, that part of the business devoted to either of the latter two phases is included in the wholesale and manufacturing survey and the part

devoted to fishing is included in the shore or vessel fisheries. If a wholesale business is conducted with no manufacturing and the business is so small that the full time of one man over the whole year or season is not required, it is then disregarded as a wholesale business. If commodities other than fishery products are handled the persons engaged, and salaries and wages paid, are prorated; and only that part concerned with fishery products is included. If such a firm requires less than the full time of one man over the whole year or season and if it does not manufacture, it is not included in the canvass. Retail firms that manufacture or whose wholesale business exceeds the retail part are included. Persons or firms engaged in the motor trucking of fishery products are included as wholesalers if they are engaged in wholesale buying and selling.

Buyers for a central firm are not canvassed as wholesale dealers unless they ship direct to the firms' customers from the buying point.

Fishermen or fishing concerns, except manufacturers, who do not buy fishery products are not included under this heading except that oyster-shucking firms are included provided shuckers are employed and irrespective of whether all or part of the oysters used are taken from the firms' privately owned beds.

Manufacturing concerns include those which prepare packaged fishery products; salted, spiced, smoked, dried, or otherwise cured fishery products; canned fishery products; or fishery byproducts.

Fishermen who manufacture are surveyed to obtain the number of persons so employed and the volume and value of the products prepared.

In collecting statistics of manufacturing firms the agents obtain data on the production for each plant in producing areas of products as marketed by the plant. Such products are usually "final" and in form for consumption; however, the products may be "intermediate" and require further processing before reaching the consumer markets. An outstanding example of an intermediate product is green-salted ground fish which almost invariably is further processed before final marketing. In reviewing the statistics of manufactured products it should be observed that intermediate products are not shown where they are prepared to the final stage in the original plant. An exception to this rule, however, is in the case of the production of mild-cured salmon which on account of its importance is shown in its entirety whether further processed in the producing plant or not. In this connection it should also be stated that several of the byproducts for which statistics are shown may be intermediate and the plants producing the final product are not surveyed by this Bureau. Outstanding among such products are marine-animal oils and fish scrap and meal.

Statistics of persons engaged in wholesale and manufacturing establishments are reported in three groups: Proprietors, salaried employees, and wage earners.

Proprietors represent those persons who devote their time to the conduct of the enterprise and receive their compensation in the form of profits. Managers of branch houses are not classified as proprietors.

Salaried employees usually include those persons paid by the week or month while wage earners usually consist of those paid on a per diem or piece-work basis. This, however, is not true in all cases,

since the distinction between these two classes depends primarily on the character of the work done rather than the unit of time employed for calculating rates of pay. In general, office employees are classified as salaried employees. Other employees, including plant workmen, are classed as wage earners. Plant foremen or superintendents are classified as salaried employees unless they are principally engaged in manual labor, in which case they are classified as wage earners. Active officers of corporations are classified as salaried employees. Statistics of wage earners are shown in two forms: The average number employed during the operating season; and the average number employed during the year (the monthly average for the year).

Transporting trade.—Statistics are obtained on the number of the crew and number of boats and vessels engaged in transporting fishery products from the fishing grounds to port or from port to port. However, if a craft is engaged in catching fish at any time of the year it is included as a fishing craft rather than as a transporter.

Publication of data.—Statistics of employment in the fisheries, craft and gear engaged, catch and value of catch, and certain data on industries related to the fisheries are summarized and published in bulletin form as soon as possible after completion of each survey. Later the figures in more detail are included in the annual reports of the Division.

LOCAL AND SPECIAL SURVEYS

Landings at certain important United States ports.—Statistics of the landings of aquatic products at the principal New England ports (Boston and Gloucester, Mass., and Portland, Maine) are obtained in a similar manner. An agent is permanently stationed at each of these ports. His duties include the obtaining of figures daily on the quantity of fish landed by each fishing vessel, the value of such fish landed, information concerning the date of departure and arrival of the vessel, and he also indicates the grounds from which the fish were taken and gear used in their capture. These data are forwarded the Bureau, where compilations are made. Products of American fisheries received duty free at Boston and Gloucester, Mass., and Portland, Maine, from the treaty coasts of Newfoundland, Magdalen Islands, and Labrador are included in the landings at these ports; however, they are not included in the catch in sectional fishery surveys of the New England States unless they represent a catch by United States vessels. Statistics of these landings are released monthly and annually in bulletin form and detailed data are published in the annual reports of this Division. Data on the landings at Boston and Gloucester, Mass., have been collected annually since 1893, and those for Portland, Maine, since June 1915. Some data are available for Boston and Gloucester prior to 1893.

Statistics of the landings of fish at Seattle, Wash., are collected by the Bureau's agent in that city. Landings are classified as those made by American fishing vessels and those received by Seattle wholesale dealers. The landings credited to United States fishing vessels are made by vessels operating distinctly as primary fishing units, usually in the offshore fisheries, while those credited as received by wholesale dealers are usually products of the shore fisheries collected mainly from points in Puget Sound and do not include fish received

from Alaska or Canada, or landings made by the halibut fleet. Statistics of these landings at Seattle are released monthly and annually in bulletin form and detailed data are published in the annual reports of this Division. Statistics of the landings by fishing vessels at Seattle have been collected since June 1915 and certain data on products received by Seattle wholesale dealers since December 1915.

Statistics on the landings of fish at New York City are obtained from J. H. Matthews, executive secretary of the Middle Atlantic Fisheries Association, New York, N. Y. During the years when one of the large trawling companies operated from Groton, Conn., the landings at that port were collected by the Bureau's agents and included with the landings at New York City in published data. Monthly bulletins including these landings are not issued. However, a summary of the year's activities is published in the annual reports of the Division. Statistics of the landings at either one or both of the above ports are available since 1922.

Statistics of the fishery products handled at the municipal wharf, Washington, D. C., are reported to the Bureau by agents of the Health Department in Washington. They are not published in bulletin form, but a summary of the year's activities is published in the annual reports of the Division. Data on products handled at the municipal wharf are available since 1921.

Atlantic mackerel fishery.—Statistics on the catch by the Atlantic mackerel fleet are obtained by combining the figures of mackerel landed at Boston and Gloucester, Mass., and Portland, Maine, with those obtained by Bureau agents, who in recent years have been stationed at other Atlantic ports where mackerel are landed. These agents obtain data on the volume of mackerel landed in a manner similar to that used to obtain figures on the landings by fishing vessels at the three New England ports. The figures include only the catches made by purse seine and drift gill net craft and are not complete for craft of under 5 net tons capacity using this type of gear. Statistics of this fishery appear only in the annual reports of this Division, although the landings at the principal New England ports appear in the monthly and annual bulletins published for those ports. Statistics of this fishery are available from 1905 to 1933, inclusive.

Shad and alewife fisheries.—Owing to the importance of the Hudson and Potomac Rivers in the production of shad, surveys for statistics of the catch, value of the catch, and operating units are made annually. On the Potomac River similar statistics also are obtained for the alewife fishery.

The survey in the Potomac River is conducted entirely by Bureau agents in a manner similar to that employed in the collection of the usual fishery statistics. The survey of the Hudson River shad fishery also is conducted by Bureau agents although considerable data is obtained from the New York State fishery agency for the catch in waters of this State.

Statistics of the shad and alewife fisheries are not published separately in bulletin form, but a summary of the year's activities is published in the annual reports of the division.

Statistics of the shad fishery of the Hudson River are available for 1896, 1897, 1898, 1901, 1904, 1910, and from 1915 to 1933, inclusive, while data for the shad fishery of the Potomac River are available

for 1896, 1901, 1904, 1909, 1915, and from 1919 to 1933, inclusive. Statistics of the alewife fishery of the Potomac River are available for 1896, 1909, 1915, and from 1919 to 1933, inclusive.

Pacific halibut fishery.—Statistics of the Pacific halibut fishery are obtained by the Bureau's agent in Seattle, aided by Bureau representatives in Alaska and the International Fisheries Commission. The fleet classification has been arbitrarily applied by including in the "Washington fleet" all United States and Alaska vessels that land more than half of their catch in that State. All other United States and Alaska vessels of the halibut fleet are included in the "Alaska fleet." Monthly and annual statistical bulletins are available on this fishery, being published along with the statistics of the landings of fishery products at Seattle, Wash., and detailed statistics are published in the annual reports of the Division. Statistics of the landings of halibut at Pacific coast ports have been collected since 1925.

Canned fishery products and byproducts.—Beginning in 1921, the Bureau has made annual surveys for statistics of the canned fishery products and byproducts industries of every section. These are begun the first week in January of each year for statistics of the production in the preceding year. The surveys usually occupy 6 to 9 week's time. During this period the Bureau obtains by mail, so far as possible, the production of canned fishery products or byproducts from each plant in the United States engaged in this business. Where it is impossible to obtain reports by mail the report is obtained by personal visit by the Bureau's agents. They obtain statistics of the production and value of the production for each commodity. Statistics of the canned fishery products and byproducts produced in Alaska are received on the same statements obtained by the Bureau that include statistics of their general fishery operations.

An annual statistical bulletin is issued on this trade, and detailed statistics of the output are published in the annual reports of the Division. In addition to the data obtained on the output of these products annually since 1921, data also usually were obtained prior to 1921 for the years the various sections were surveyed.

The value shown for canned products constitutes the gross amount received by the packer at the production point, no deductions being made for commission or expenses.

Packaged-fish trade.—Complete statistics of the annual production and value of fish packaged in the United States are obtained as a part of the survey for the statistics of the canned fishery products and byproducts industries. These statistics are released in bulletin form annually and detailed statistics are published in the annual reports of the Division. Statistics of the production of packaged fish are available for 1926 and the years from 1928 to 1933, inclusive.

Cold-storage holdings of fish.—An arrangement has been made with the Bureau of Agricultural Economics, Department of Agriculture, whereby statistics of the cold-storage holdings of the various species of fish, by sections of the United States, are furnished to this Bureau monthly. Included with statistics of the holdings are statements of the quantity of the various species of fish frozen and also the holdings of certain cured fish. Bulletins showing these statistics are issued monthly as well as annually, and detailed statistics are published in the annual reports of this Division. Statistics of cold-storage holdings of fishery products have been published since 1917 and data on quan-

tities of fish frozen, for the years from 1920 to 1925, inclusive, and from 1928 to 1933, inclusive.

Sponge market, Tarpon Springs.—A large proportion of the total output of sponges in Florida is handled through the sponge exchange at Tarpon Springs. In view of this, the Bureau has obtained from a representative of the exchange annual statistics of the quantity and value of the sponges, by variety classification, handled through it annually. Statistics of the quantity of sponges handled through the exchange are not published in bulletin form, but a summary of the year's activities is published in the annual reports of this Division. Statistics of the transactions on the sponge exchange are available for 1913, 1914, and for the years from 1917 to 1933, inclusive.

Foreign fishery trade.—Statistics of the foreign fishery trade are obtained from compilations made by the Bureau of Foreign and Domestic Commerce, Department of Commerce. Statistics of all known fishery products imported or exported have been assembled in one table and published annually in the reports of the Division in recent years. For earlier years they are available in the reports of the Bureau of Foreign and Domestic Commerce, the Bureau of Statistics, the Department of Commerce and Labor, and the Treasury Department.

PRACTICES AND TERMS

Certain practices and terms of importance used in the compilation of fishery statistics are explained below.

Days absent.—In computing "days absent" for vessels landing fares at the various ports, the day of departure and the day of arrival are included; thus, a vessel leaving port on the 8th of the month and returning on the 15th of the month will be shown as being absent 8 days.

Operating units.—Operating units as referred to in this document include persons engaged in the fisheries and fishing craft and gear employed.

Vessel.—The term "vessel" refers to a craft having a capacity of 5 net tons or more.

Boat.—The term "boat" refers to a craft having a capacity of less than 5 net tons.

Incidental catch.—The term "incidental catch" refers to the catch of certain species by a type of gear which ordinarily does not capture such species.

Percentages.—Percentages are usually shown as whole numbers. Fractions of percents are dropped if less than five-tenths, and the percentage is raised to the next higher integer if the fraction is greater than five-tenths. If the fraction is exactly five-tenths, the integer is raised or lowered to make it an even number.

Converting.—Many of the figures shown in the statistical tables published herewith have been reduced to thousands of pounds or dollars. In making these conversions the largest number from which a group of items is computed is raised or lowered to the nearest thousands place. If the number ends in an even 500, the thousands integer is raised or lowered to make it an even number. The individual items are changed to conform to the total thus obtained.

Confidential data.—The statistical data collected by the Division are confidential and are not released except by approval of the Washington

office. Statistics of production of wholesale and manufacturing firms are published only for commodities or geographical areas where the production of three or more concerns may be grouped. Every effort is made to publish only those figures which will not reveal individual enterprise.

CONVERSION FACTORS

It is the policy of the Bureau to show the detailed catch figures of all products in pounds for the sake of uniformity and for purposes of comparison. Following such a policy presents certain problems. In the case of fish there is little difficulty since in very rare instances are such products reported in units of measure other than pounds. For shellfish, however, the units of measure may be bushels, sacks, barrels, or thousands of shellfish, gallons of meats, etc. These many units make standardization difficult, but when coupled with the wide variation in the requirements or definition of some of these units in the various States the problem becomes even more complex.

All bivalve mollusks are reported in pounds of meats in the detailed catch tables presented in this report. In addition, there are included supplementary tables for most of the sections, which give data on the production in bushels. These supplementary tables also give the production of certain other shellfish, such as crabs, in number.

Oysters.—Probably the greatest problem in presentation of fishery statistics in uniform units of measure is in the case of oysters. Usually the production of oysters on the Atlantic and Gulf coasts is reported to Bureau agents in bushels; and prior to the data obtained for the year 1930 conversion from bushels to pounds of meats was effected on the basis of a uniform yield of 7 pounds of meats to the bushel. However, recent investigations have shown considerable variation from this figure. There follows a table which gives the results of these studies of the measures used for oysters in the various States and of the average yields per bushel. This table presents the factors that have been used in the oyster statistics given in this report.

Measures and yields of oysters ¹

| State | Capacity of State bushel | Variation from United States standard bushel | | Market oysters | |
|---------------------|--------------------------|--|----------------|------------------------|---------------------------|
| | | | | Yield per State bushel | Yield per standard bushel |
| | | | | Pounds of meats | Pounds of meats |
| | <i>Cubic inches</i> | <i>Cubic inches</i> | <i>Percent</i> | | |
| Massachusetts..... | 2, 150. 4 | | | 6. 57 | 6. 57 |
| Rhode Island..... | 2, 150. 4 | | | 6. 96 | 6. 96 |
| Connecticut..... | 2, 160. 4 | | | 6. 81 | 6. 81 |
| New York..... | 2, 150. 4 | | | 7. 00 | 7. 00 |
| New Jersey..... | 2, 257. 3 | +106. 9 | +5. 0 | 8. 70 | 8. 29 |
| Delaware..... | 2, 257. 3 | +106. 9 | +5. 0 | 7. 58 | 7. 50 |
| Maryland..... | 2, 801. 5 | +650. 1 | +30. 2 | 6. 57 | 5. 05 |
| Virginia..... | 3, 003. 4 | +853. 0 | +39. 7 | 6. 90 | 4. 72 |
| North Carolina..... | 2, 801. 9 | +651. 5 | +30. 3 | 5. 71 | 4. 38 |
| South Carolina..... | 4, 071. 5 | +1, 921. 1 | +49. 3 | 4. 76 | 2. 51 |
| Georgia..... | 2, 753. 4 | +603. 0 | +28. 0 | 5. 69 | 4. 45 |
| Florida..... | 3, 214. 1 | +1, 063. 7 | +49. 4 | 3. 29 | 2. 20 |
| Alabama..... | 2, 826. 2 | +676. 8 | +31. 4 | 2. 40 | 1. 53 |
| Mississippi..... | 2, 826. 2 | +676. 8 | +31. 4 | 2. 19 | 1. 67 |
| Louisiana..... | 2, 148. 4 | -2. 0 | -. 1 | 4. 14 | 4. 14 |
| Texas..... | 2, 700. 0 | +549. 6 | +26. 6 | 5. 05 | 4. 02 |

¹ Data for the New England, Middle Atlantic, and Chesapeake States are for 1933. Other data are for 1932.

Other mollusks.—The following table shows the conversion factors for various mollusks, other than oysters, used in this report.

*Average yields of certain mollusks in pounds of meats per bushel*¹

| State | Clams, hard | | Clams, soft | | Clams, surf | Clams, razor | Mussels, sea | Periwinkles and cockles | Scallops, bay | Scallops, sea | Conchs |
|---------------------|-------------|---------|-------------|---------|-------------|--------------|--------------|-------------------------|---------------|---------------|--------|
| | Public | Private | Public | Private | | | | | | | |
| Maine..... | 11 | | 15 | | | | 12 | 18 | | 6 | |
| Massachusetts..... | 11 | 11 | 15 | | 17 | 32 | 10 | 18 | 6 | 6 | |
| Rhode Island..... | 11 | 11 | 16 | | | | 13 | 18 | | | |
| Connecticut..... | 10 | | 14 | | | | | | 5.75 | 6 | |
| New York..... | 8 | 8 | 16 | | 12 | | 10 | | 5 | 6 | 18 |
| New Jersey..... | 8.96 | 8.96 | 20 | | 12.5 | | 13 | 11.85 | 6 | 6 | 18 |
| Delaware..... | 10 | 10 | | | | | | | | | |
| Maryland..... | 8 | 10 | | | | | | | | | |
| Virginia..... | 8 | | | | | | | | 5.5 | | |
| North Carolina..... | 8 | | | | | | | | | | |
| South Carolina..... | 8 | | | | | | | | | 6 | |
| Georgia..... | 8 | | | | | | | | | | |
| Florida..... | 8 | | | | | | | | 5.3 | | |

¹ Data for the New England, Middle Atlantic, and Chesapeake States are for 1933. Other data are for 1932.

Other conversion factors.—The principal other conversion factors that have been used in this report are as follows:

| | |
|---|---|
| Alewives..... | To convert number of fish to weight in pounds, multiply by 0.4. |
| Cod, large, salted..... | To convert to fresh-gutted weight, multiply by 1.90. |
| Cod, market, salted..... | To convert to fresh-gutted weight, multiply by 1.94. |
| Cod, scrod, salted..... | To convert to fresh-gutted weight, multiply by 1.98. |
| Crustaceans: | |
| Crabs, soft (New York, Maryland, and Virginia). | To convert number of crabs to weight in pounds, divide by 4. |
| Crabs, soft (North Carolina).... | To convert number of crabs to weight in pounds, divide by 3.63. |
| Crabs, soft (other States)..... | To convert number of crabs to weight in pounds, divide by 3. |
| Crabs, hard (North Carolina).... | To convert number of crabs to weight in pounds, divide by 4. |
| Crabs, hard (South Carolina and Georgia). | To convert number of crabs to weight in pounds, divide by 2. |
| Crabs, hard (Florida)..... | To convert number of crabs to weight in pounds, divide by 1.64. |
| Crabs, hard (Alabama and Texas). | To convert number of crabs to weight in pounds, divide by 1.72. |
| Crabs, hard (Mississippi)..... | To convert number of crabs to weight in pounds, divide by 1.92. |
| Crabs, hard (Louisiana)..... | To convert number of crabs to weight in pounds, divide by 1.86. |
| Crabs, hard (other States)..... | To convert number of crabs to weight in pounds, divide by 3. |
| Crabs, king..... | To convert number of crabs to weight in pounds, multiply by 4. |
| Crabs, rock..... | To convert number of crabs to weight in pounds, divide by 3. |
| Crabs, stone..... | To convert number of crabs to weight in pounds, multiply by 1.33. |
| Cusk, salted..... | To convert to fresh-gutted weight, multiply by 1.90. |
| Haddock, large, salted..... | To convert to fresh-gutted weight, multiply by 2.06. |

| | |
|-----------------------------|---|
| Haddock, scrod, salted..... | To convert to fresh-gutted weight, multiply by 2.10. |
| Hake, large, salted..... | To convert to fresh-gutted weight, multiply by 1.90. |
| Hake, small, salted..... | To convert to fresh-gutted weight, multiply by 1.98. |
| Halibut, salted..... | To convert to fresh-gutted weight, multiply by 2. |
| Herring, salted..... | To convert to round weight, multiply by 1.50. |
| Mackerel, salted..... | To convert to round weight, multiply by 1.35. |
| Menhaden..... | To convert number of fish to weight in pounds, multiply by 0.6. |
| Oil (east coast)..... | To convert gallons to pounds, multiply by 7.74. |
| Oil (west coast)..... | To convert gallons to pounds, multiply by 7.5. |
| Pollock, salted..... | To convert to fresh-gutted weight, multiply by 1.90. |
| Sponges, dried (Florida): | |
| Large wool..... | To convert number of bunches to weight in pounds, multiply by 3.5. |
| Medium wool..... | To convert number of bunches to weight in pounds, multiply by 1.75. |
| Small wool..... | To convert number of bunches to weight in pounds, multiply by 1. |
| Wool rags..... | To convert number of bunches to weight in pounds, multiply by 2.25. |
| Grass..... | To convert number of bunches to weight in pounds, multiply by 1. |
| Wire..... | To convert number of bunches to weight in pounds, multiply by 1.5. |
| Yellow..... | To convert number of bunches to weight in pounds, multiply by 1.25. |

COMMON AND SCIENTIFIC NAMES OF FISHERY PRODUCTS

In order to prevent misunderstanding in the use of common names employed in the tables and discussions, the following list of common and scientific names is given:

Common and scientific names of the commercial fishery products caught in the United States and Alaska

| Common name as shown in Bureau reports | Other common names | Scientific names |
|--|---|--|
| Albacore..... | Longfin tuna..... | <i>Germo alalunga</i> (Pacific coast). <i>Pomolobus pseudoharengus</i> . |
| Alewives..... | { Branch herring, wall-eyed or big-eyed herring. Blueback, glut herring..... | <i>Pomolobus aestivalis</i> . <i>Seriola</i> species. |
| Amberjack..... | | { <i>Engraulis mordax</i> . <i>Anchoviella delicatissima</i> . <i>Anchoviella compressa</i> . |
| Anchovies..... | | { <i>Pomacanthus arcuatus</i> . <i>Angelichtys isabelita</i> . <i>Sphyræna argentea</i> (Pacific coast). |
| Angelfish..... | | { <i>Sphyræna barracuda</i> (Atlantic coast). |
| Barracuda..... | | |

Common and scientific names of the commercial fishery products caught in the United States and Alaska—Continued

| Common name as shown in Bureau reports | Other common names | Scientific names |
|--|--|---|
| Black bass----- | { Smallmouth bass----- Largemouth bass----- | <i>Micropterus dolomieu.</i> <i>Micropterus salmoides.</i> |
| Bluefish----- | Tailor----- | <i>Pomatomus saltatrix.</i> |
| Blue pike----- | Pike perch, blue pickerel (Canada). | <i>Stizostedion glaucum.</i> |
| Blue runner or hard- tail. | Runner----- | <i>Caranx crysos.</i> |
| Bonito----- | | { <i>Sarda sarda.</i> <i>Sarda chiliensis.</i> |
| Bowfin----- | | <i>Amia calva.</i> |
| Buffalofish----- | | Ictiobus species. |
| Bullhead----- | | Ameiurus species. |
| Butterfish----- | Dollarfish----- | <i>Poronotus triacanthus.</i> |
| Burbot----- | Lawyer, ling----- | <i>Lota maculosa.</i> |
| Cabio----- | Coalfish, crab eater, co- bia. | <i>Rachycentron canadus.</i> |
| Cabrilla----- | Rock bass----- | <i>Epinephelus analogus</i> (Pa- cific coast). |
| Carp----- | German carp----- | <i>Cyprinus carpio.</i> |
| Catfish----- | | Siluridae species. |
| Cero----- | | <i>Scomberomorus regalis.</i> |
| Chubs----- | Tullibee in Canada; long- jaw, bluefin, blackfin in United States. | All <i>Leucichthys</i> except <i>artedi</i> (in Great Lakes). |
| Cigarfish----- | Scad----- | Decapterus species. |
| Cisco----- | Herring in Canada----- | <i>Leucichthys artedi</i> (Lake Erie only). |
| Cod----- | Codfish----- | { <i>Gadus macrocephalus</i> (Pa- cific coast). <i>Gadus callarias</i> (Atlantic coast). |
| Corbina----- | Orange mouth corbina--- | <i>Cynoscion xanthurum.</i> |
| Crappie----- | { White crappie----- Black crappie, strawberry bass, calico bass. | <i>Pomoxis annularis.</i> <i>Pomoxis sparoides.</i> |
| Crevalle----- | | <i>Caranx hippos.</i> |
| Croaker----- | Crocus, hardhead----- | <i>Micropogon undulatus.</i> |
| Cunner----- | Chogset, blue perch, berg- all. | <i>Tautoglabrus adspersus.</i> |
| Cusk----- | | <i>Brosimius brosme.</i> |
| Dolly Varden trout--- | Salmon trout, bull trout--- | <i>Salvelinus parkei.</i> |
| Dolphin----- | | <i>Coryphaena hippurus.</i> |
| Drum: | | |
| Black----- | | <i>Pogonias cromis.</i> |
| Red----- | Channel bass, redfish, spotted bass. | <i>Sciaenops ocellatus.</i> |
| Eels: | | |
| Common----- | | <i>Anguilla rostrata.</i> |
| Conger----- | | { <i>Leptocephalus conger.</i> <i>Gymnothorax</i> species. <i>Pleuronectidae</i> species. |
| Flounders----- | { Dabs, blackbacks, lemon sole, winter flounder, summer flounder. Halibut, "California"--- Sole----- | <i>Paralichthys californicus.</i> <i>Psetticthys melanostictus</i> (Pacific coast). |
| Flyingfish----- | | <i>Cysilurus californicus.</i> |
| Frigate mackerel--- | "Boo Hoo." | <i>Auzis thazard.</i> |
| Garfish----- | | (See sea gar.) |
| Gizzard shad----- | Nanny shad, mud shad--- | <i>Dorosoma cepedianum.</i> |

Common and scientific names of the commercial fishery products caught in the United States and Alaska—Continued

| Common name as shown in Bureau reports | Other common names | Scientific names |
|--|---|---|
| Goldeye..... | | Hiodon species. |
| Goldfish..... | Sand perch..... | <i>Carassius auratus</i> . |
| Goosefish..... | | <i>Lophius piscatorius</i> . |
| | { Dogfish..... | <i>Squalus sucklii</i> (Pacific coast.) |
| Grayfish..... | { Spiny dog..... | <i>Squalus acanthias</i> . |
| | { Smooth dog..... | <i>Mustelus mustelus</i> . |
| Groupers..... | "Sea bass"..... | { <i>Epinephelus</i> species. |
| Grunts..... | Margatefish, sailors choice (Key West). | { <i>Mycteroperca</i> species. |
| Haddock..... | | <i>Haemulon</i> species. |
| Hagfish..... | Slimefish..... | <i>Melanogrammus aeglefinus</i> . |
| | { Squirrel hake, Boston hake, ling, black hake, mud hake. | <i>Myxine glutinosa</i> . |
| Hake..... | { Merluccio..... | <i>Urophycis</i> species (Atlantic coast). |
| Halibut..... | | <i>Merluccius productus</i> (Pacific coast). |
| Hardhead..... | | <i>Hippoglossus hippoglossus</i> . |
| Harvestfish..... | Starfish, pappypish; butterfish (N. C.). | <i>Orthodon microlepidotus</i> (Pacific coast). |
| Herring: | | <i>Peprilus alepidotus</i> . |
| Lake..... | Herring..... | <i>Leucichthys artedi</i> (Great Lakes, except Erie). |
| Sea..... | | { <i>Clupea harengus</i> (Atlantic coast). |
| | | { <i>Clupea pallasii</i> (Pacific coast). |
| Herring smelt..... | Sea smelt..... | <i>Argentina silus</i> . |
| Hickory shad..... | Tailor shad..... | <i>Pomolobus mediocris</i> . |
| Hogfish..... | Capitaine, perro perro..... | <i>Lachnolaimus maximus</i> (Florida). |
| Horse mackerel..... | | <i>Trachurus symmetricus</i> (Pacific coast). |
| Jewfish..... | | <i>Promicrops itaiara</i> . |
| Kingfish..... | { Little roncador, croaker..... | <i>Scomberomorus cavalla</i> (Atlantic coast). |
| King whiting..... | Northern whiting, kingfish, seaming. | <i>Genyonemus lineatus</i> (California). |
| Ladyfish..... | Bonefish..... | <i>Menticirrhus</i> species. |
| Lake trout..... | | <i>Albula vulpes</i> . |
| Launce..... | Sand eel, lant, sand launce. | <i>Cristivomer namaycush</i> . |
| "Lingcod"..... | Cultus cod, blue cod, buffalo cod, ling. | <i>Ammodytes americanus</i> . |
| Mackerel..... | | { <i>Scomber scombrus</i> (Atlantic coast). |
| Marlin..... | Spearfish..... | <i>Scomber diego</i> (Pacific coast). |
| Menhaden..... | Mossbunker, pogy..... | <i>Tetrapturus mitsukurii</i> (Pacific coast). |
| Minnows..... | | <i>Brevoortia tyrannus</i> . |
| Mojarro..... | | Cyprinidae species. |
| Mooneye..... | Toothed herring..... | <i>Eucinostomus</i> species. |
| Moonfish..... | | Hiodon species. |
| | | { <i>Vomer setipinnis</i> . |
| | | { <i>Selene vomer</i> . |

Common and scientific names of the commercial fishery products caught in the United States and Alaska—Continued

| Common name as shown in Bureau reports | Other common names | Scientific names |
|--|-------------------------------------|--|
| Mullet..... | Jumping mullet..... | Mugil species. |
| Mummichog..... | Mayfish, killifish..... | Fundulus species. |
| Muttonfish..... | | <i>Lutianus analis</i> . |
| Paddlefish..... | Spoonbill cat..... | <i>Polyodon spathula</i> . |
| Parrotfish..... | | Scaridae species. |
| Perch (California)..... | | (See surf fishes) |
| Permit..... | Great pompano..... | <i>Trachinotus goodet</i> . |
| Pigfish..... | Hogfish (N. C.)..... | <i>Orthopristis chrysopterus</i> . |
| Pike or pickerel..... | Great Lakes pike..... | { <i>Esox reticulatus</i> . |
| Pilchard..... | Sardine..... | { <i>Esox lucius</i> . |
| Pilotfish..... | | { <i>Sardinia caerulea</i> . |
| Pinfish..... | | { <i>Naucrates ductor</i> . |
| Pollock..... | Bream, salt-water bream..... | { <i>Seriola zonata</i> . |
| | | { <i>Lagodon rhomboides</i> . |
| | | { <i>Pollachius virens</i> . |
| Pompano..... | | { Trachinotus species (Atlantic coast). |
| | | { <i>Palometa simillima</i> (Pacific coast). |
| Porgies..... | Porgee..... | Calamus species. |
| Porkfish..... | Sisi..... | <i>Anisotremus virginicus</i> |
| Quillback..... | Spearfish or skimfish..... | Carpiodes species. |
| Roach..... | Shiner..... | <i>Notemigonus crysoleucas</i> . |
| | { Redeye, goggle-eye..... | <i>Ambloplites rupestris</i> (Mississippi River to Atlantic seaboard). |
| Rock bass..... | { Groupers..... | <i>Paralabrax nebulifer</i> (Pacific coast). |
| Rockfishes..... | Rock cod..... | Sebestodes species (Pacific coast). |
| Rosefish..... | | <i>Sebastes marinus</i> . |
| Rudderfish..... | { Blue bass, greenfish..... | <i>Girella nigricans</i> (Pacific coast). |
| | { Halfmoon..... | <i>Medialuna californiensis</i> (Pacific coast). |
| Sablefish..... | Black cod..... | <i>Anaplopoma fimbria</i> |
| Salmon: | | |
| Atlantic..... | | <i>Salmo salar</i> (Atlantic coast). |
| Pacific: | | |
| Blueback, red or sock-eye. | | <i>Oncorhynchus nerka</i> . |
| Chinook or king. | Tyee, Columbia, Sacramento, spring. | <i>Oncorhynchus tshawytscha</i> . |
| Chum or keta. | Dog salmon..... | <i>Oncorhynchus keta</i> . |
| Humpback or pink. | | <i>Oncorhynchus gorbuscha</i> . |
| Silver or coho | | <i>Oncorhynchus kisutch</i> . |
| Steelhead..... | See steelhead trout. | |
| Sauger..... | Sand pike..... | <i>Stizostedion canadense</i> . |
| Sawfish..... | | <i>Pristis pectinatus</i> . |
| Sculpin..... | | Cottidae species. |
| Scup..... | Paugy or porgy, fair maid. | <i>Stenotomus</i> species. |
| | { Black jewfish or black sea bass. | <i>Stereolepis gigas</i> (Pacific coast). |
| Sea bass..... | { Black sea bass..... | <i>Centropristes striatus</i> (Atlantic coast). |
| | { White sea bass..... | <i>Cynoscion nobilis</i> (Pacific coast). |

Common and scientific names of the commercial fishery products caught in the United States and Alaska—Continued

| Common name as shown in Bureau reports | Other common names | Scientific names |
|--|---|---|
| Sea gar | Needlefish, billfish, houndfish. | Tylosurus species. |
| Sea robin | | Prionotus species. |
| Shad | American shad | <i>Alosa sapidissima</i> . |
| Sharks | | Carcharodon species; Mustelus species; Carcharhinus species; Sphyrna species. |
| | | <i>Archosargus probatocephalus</i> (Atlantic coast). |
| Sheepshead | Drum, fresh-water | <i>Aplodinotus grunniens</i> (Fresh-water). |
| | Redfish, flat head | <i>Pimelometopon pulcher</i> (Pacific coast). |
| Silver perch | Sand perch | <i>Bairdiella chrysura</i> . |
| Silversides | Spearing | Menidia species. |
| Skates | | Raja species. |
| Skipper | Billfish | <i>Scomberesox saurus</i> . |
| | | <i>Osmerus mordax</i> (Atlantic coast). |
| Smelts | | Argentinidae species (Pacific coast). |
| | Eulachon | <i>Thaleichthys pacificus</i> . |
| Snapper: | | |
| Mangrove | Gray snapper | <i>Lutianus griseus</i> . |
| Red | | <i>Lutianus blackfordii</i> . |
| Snook | Robalo, sergeantfish | <i>Centropomus undecimalis</i> . |
| Spadefish | Porgy (N. C.) | <i>Chaetodipterus faber</i> . |
| Spanish mackerel | | <i>Scomberomorus maculatus</i> . |
| Splittail | | <i>Pogonichthys macrolepidotus</i> . |
| Spot | Lafayette, goody | <i>Leiostomus xanthurus</i> . |
| Squawfish | Sacramento pike | <i>Ptychocheilus grandis</i> . |
| Squeteague: | | |
| Gray | Gray trout, weakfish, trout. | <i>Cynoscion regalis</i> . |
| Spotted | Spotted weakfish, spotted trout. | <i>Cynoscion nebulosus</i> . |
| Squirrelfish | | <i>Diplectrum formosum</i> . |
| Steelhead trout | Salmon trout | <i>Salmo gairdneri</i> . |
| Striped bass | Rockfish, rock | <i>Roccus lineatus</i> . |
| Sturgeon | | Acipenser species. |
| Sturgeon, shovelnose | | <i>Scaphirhynchus platyrhynchus</i> . |
| Sucker | Fresh-water mullet | Chatostomidae species. |
| | | Lepomis species. |
| Sunfish | | Centrarchidae species. |
| Surf fishes | | Embiotocidae species. |
| Swellfish | Puffer, swell toad, balloonfish, globefish. | <i>Spheroides maculatus</i> . |
| Swordfish | | <i>Xiphias gladius</i> . |
| Tai | | <i>Calumus brachysomus</i> . |
| Tautog | Blackfish, oysterfish | <i>Tautoga onitis</i> . |
| Tenpounder | Elops | <i>Elops saurus</i> . |
| Thimble-eyed mackerel | Bullseye | <i>Scomber colias</i> . |
| Tilefish | | <i>Lopholatilus chamaeleonticeps</i> . |
| | | <i>Microgadus tomcod</i> (Atlantic coast). |
| Tomcod | | <i>Microgadus proximus</i> (Pacific coast). |
| Tripletail | | <i>Lobotes surinamensis</i> . |

Common and scientific names of the commercial fishery products caught in the United States and Alaska—Continued

| Common name as shown in Bureau reports | Other common names | Scientific names |
|--|---|--|
| Tullibee | | (See chubs.) |
| Tuna and tunalike fishes: | | |
| Albacore | Longfin tuna | <i>Germo alalunga</i> . |
| Bluefin | { Tuna, leaping tuna (Pacific coast). "Horse mackerel" (Atlantic coast). | { <i>Thunnus thynnus</i> . |
| Bonito | | { <i>Sarda sarda</i> (Atlantic coast). <i>Sarda chiliensis</i> (Pacific coast). |
| Skipjack | Striped tuna | <i>Euthynnus pelayms</i> . |
| Yellowfin | | <i>Neothunnus macropterus</i> . |
| Turbot | { Greenland halibut American turbot | { <i>Reinhardtius hippoglossoides</i> (Off New England) <i>Balistes carolinensis</i> (Off Florida) |
| White bass | White lake bass | <i>Roccus chrysops</i> . |
| Whitebait | Small fry of any fish. | |
| Whitefish: | | |
| Common | | { <i>Coregonus clupeiformis</i> (Great Lakes). <i>Caulolatilus princeps</i> (Pacific coast). |
| Menominee | | <i>Prosopium quadrilaterale</i> |
| White perch | | <i>Morone americana</i> (Atlantic coast). |
| Whiting | Silver hake | <i>Merluccius bilinearis</i> . |
| Wolfish | | <i>Anarhichas lupus</i> . |
| Yellow perch | | <i>Perca flavescens</i> . |
| Yellow pike | Wall-eyed pike, pike perch, dore. | <i>Stizostedion vitreum</i> . |
| Yellowtail | | { <i>Ocyurus chrysurus</i> (Atlantic coast). <i>Seriola dorsalis</i> (Pacific coast). |
| Abalone | | <i>Halotis</i> species. |
| Clams: | | |
| Cockle | | <i>Cardium corbis</i> (Pacific coast). |
| Hard | { Butter Round clam, cherrystone, quahog, little neck | { <i>Saxidomus nuttall</i> . <i>Tivela stultorum</i> (Pacific coast). <i>Venus mercenaria</i> (Atlantic coast). <i>Venus mortoni</i> (Florida coast). |
| Pismo | | <i>Tivela stultorum</i> (Pacific coast). |
| Razor | | { <i>Enis</i> species (Atlantic coast). <i>Siliqua patula</i> (Pacific coast). |
| Surf | Skimmer | <i>Mactra solidissima</i> . |
| Cockles | Moonshell | <i>Natica heros</i> (Atlantic coast). |
| Conchs | | { <i>Strombus</i> species. <i>Busycon</i> species. |
| Coquina | Pompano shells | <i>Donax variabilis</i> . |

Common and scientific names of the commercial fishery products caught in the United States and Alaska—Continued

| Common name as shown in Bureau reports | Other common names | Scientific names |
|--|--|---|
| Crabs: | | |
| Hard..... | Hard-shell crab, blue crab. Dungeness crab..... | <i>Callinectes sapidus</i> . <i>Cancer magister</i> (Pacific coast). |
| Soft..... | Rock crab, hard crab..... | <i>Cancer irroratus</i> (Atlantic coast). |
| King..... | Soft-shelled crab, blue crab. | <i>Callinectes sapidus</i> . |
| Stone..... | Horseshoe crab..... | <i>Limulus</i> (Atlantic coast). |
| Crawfish: | | <i>Menippe mercenaria</i> . |
| Fresh-water..... | Crayfish..... | <i>Cambarus</i> species (Atlantic coast). <i>Astacus</i> species (Pacific coast). |
| Sea..... | Rock lobster, crayfish..... | <i>Panulirus argus</i> (Atlantic coast). <i>Panulirus interruptus</i> (Pacific coast). |
| Lobsters: | | |
| Common..... | | <i>Homarus americanus</i> (Atlantic coast). |
| Spiny..... | (See sea crawfish.) | |
| Mussels: | | |
| Sea..... | | <i>Mytilus californianus</i> (Pacific coast). <i>Mytilus edulis</i> . <i>Quadrula</i> species. <i>Lampsilis</i> species. |
| Fresh-water..... | | <i>Unio</i> species. <i>Symphynota</i> species. |
| Octopus..... | | <i>Octopus punctatus</i> (Pacific coast). |
| Oysters: | | |
| Eastern..... | | <i>Ostrea elongata</i> . |
| Western..... | Olympia..... | <i>Ostrea lurida</i> (Pacific coast). |
| Japanese (introduced). | Pacific..... | <i>Ostrea gigas</i> . |
| Periwinkles..... | | <i>Littorina</i> species. |
| Scallops: | | |
| Bay..... | | <i>Pecten irradians</i> (Atlantic coast). <i>Pecten aquisulcatus</i> (Pacific coast). |
| Sea..... | | <i>Pecten magellanicus</i> . <i>Peneus setiferus</i> . <i>Peneus brasiliensis</i> (Atlantic and Gulf coasts). <i>Pandalus</i> species (Pacific coast). |
| Shrimp..... | | <i>Pandalopsis</i> species (Pacific coast). <i>Crangon</i> species (Pacific coast). |
| Squid..... | | <i>Loligo opalescens</i> (Pacific coast). <i>Loligo pealei</i> (Atlantic coast). |
| Terrapin..... | Diamond-back terrapin..... | <i>Malaclemmys</i> species. |

Common and scientific names of the commercial fishery products caught in the United States and Alaska—Continued

| Common name as shown in Bureau reports | Other common names | Scientific names |
|--|------------------------------|---|
| Turtles: | | |
| Green..... | ----- | <i>Chelonia mydas.</i> |
| Loggerhead..... | ----- | <i>Thalassochelys carctta.</i> |
| Hawksbill..... | ----- | <i>Chelonia inbricata.</i> |
| Snapping..... | Hardshell, alligator turtle. | { <i>Chelydra serpentina.</i> <i>Macrochalys lacertina.</i> |
| Soft-shell..... | ----- | Trionyx species. |
| Frogs..... | ----- | <i>Rana</i> species. |
| Irish moss..... | ----- | <i>Chondrus crispus.</i> |
| Kelp..... | ----- | <i>Macrocystis</i> species; <i>Nereocystis</i> species; <i>Pelagophycus</i> species; <i>Alaria</i> species. |
| Sponges: | | |
| Glove..... | ----- | <i>Spongia graminea</i> (Hyatt) <i>Euspongia officianalis</i> (L.) |
| Grass..... | ----- | <i>Hippospongia equina cerebriformis.</i> |
| Sheepswool..... | ----- | <i>Hippospongia canaliculata gossypina.</i> |
| Yellow..... | ----- | <i>Hippospongia equina elastica.</i> |
| Trepang..... | Sea cucumber..... | <i>Cucumaris frondosa</i> ; <i>Thyone briareus.</i> |

PROGRESS IN BIOLOGICAL INQUIRIES, 1934¹

By ELMER HIGGINS, *Chief, Division of Scientific Inquiry*

[With the collaboration of investigators]

CONTENTS

| | Page |
|--|------|
| Introduction..... | 332 |
| Cooperative investigations..... | 334 |
| Publications..... | 335 |
| North and Middle Atlantic fishery investigations..... | 337 |
| Haddock..... | 338 |
| Mackerel..... | 334 |
| Cod..... | 346 |
| Flounders..... | 347 |
| Species in need of attention..... | 348 |
| Shore fishes of the Middle Atlantic States..... | 349 |
| Fishery investigations of the South Atlantic and Gulf coasts..... | 354 |
| Investigation of the spawning habits, larval development, and rate of growth of fishes..... | 354 |
| Survey of the fresh waters of Mississippi..... | 354 |
| Marine fishes of the Gulf coast..... | 354 |
| An investigation of the fishes and fish cultural possibilities of the fresh waters of Puerto Rico..... | 354 |
| Shrimp investigations..... | 355 |
| Pacific coast and Alaska fishery investigations..... | 359 |
| Columbia River salmon investigation..... | 359 |
| Puget Sound sockeye salmon investigation..... | 360 |
| Washington coho salmon investigation..... | 361 |
| Statistical study of salmon fisheries in Puget Sound region..... | 362 |
| Karluk River red salmon investigation..... | 363 |
| Chignik River red salmon investigation..... | 364 |
| Bristol Bay red salmon investigation..... | 364 |
| Pink salmon investigation..... | 364 |
| Herring investigation..... | 366 |
| Investigation concerning the protection of migratory fish at the Bonneville Dam..... | 368 |
| Great Lakes fishery investigations..... | 369 |
| Fishery statistics..... | 369 |
| Pike-perches..... | 369 |
| Yellow perch..... | 371 |
| Other outstanding problems..... | 371 |
| Uniform regulation and depletion..... | 371 |
| Oyster investigations..... | 372 |
| Experiments on oyster culture..... | 372 |
| Oil pollution investigations in Louisiana..... | 374 |
| Biology and cultivation of hard-shell clam (<i>Venus mercenaria</i>)..... | 375 |
| Rehabilitation of public oyster beds in North Carolina..... | 376 |
| Investigation of oysters at New River, N. C..... | 377 |
| Oyster investigations in Florida..... | 377 |
| Protozoan parasite of Virginia and Louisiana oysters..... | 378 |
| Oyster investigations in Washington..... | 379 |

¹Administrative Report No. 21, Appendix III to the Report of the U. S. Commissioner of Fisheries for 1935. Approved for publication, Nov. 7, 1935.

| | Page |
|---|------|
| Investigations on aquiculture..... | 380 |
| Pondfish culture..... | 381 |
| Trout culture..... | 382 |
| Fish diseases..... | 389 |
| Stream survey and stream improvement..... | 391 |
| Stream improvement in connection with stream surveys..... | 392 |
| Investigations in the Rocky Mountain regions in the interest of fish stocking..... | 392 |
| Cooperation with the Forest Service in stream and lake improvement..... | 394 |
| Cooperation with the University of Utah..... | 395 |
| Investigations in interior waters..... | 395 |
| Pollution studies..... | 395 |
| Power dam lakes and other impounded waters..... | 396 |
| Mussel propagation..... | 397 |
| Independent activities of the Fisheries biological laboratories..... | 397 |
| Woods Hole, Mass..... | 397 |
| Beaufort laboratory..... | 397 |

INTRODUCTION

The United States Bureau of Fisheries is the only Federal agency conducting research looking toward the development and conservation of the fishery resources in the United States. The major portion of research on conservation problems is conducted by the Division of Scientific Inquiry, for technological research is mainly promotive and contributes only indirectly to a solution of the problems of conservation by promoting a fuller use of fishery products. As a general rule, the State governments do not have facilities nor resources for the conduct of fishery research on an extensive scale. Many States, however, contribute to or cooperate effectively in work conducted by the United States Bureau of Fisheries.

The technical staff of the Division of Scientific Inquiry numbers 45 trained experts, with perhaps an equal number of temporary assistants at some time of the year, but the problems of the fisheries are as numerous as the species that are exploited. Some 30 important food and game fishes are the subjects of continuing research. To give adequate attention to the fisheries in all sections of the United States, both coastal and interior, and in Alaska, at least double this number of investigators would be required.

The aim of fishery investigations is to determine which species are showing depletion as a result of man's exploitation, what methods may be applied toward their conservation through the management of supplies on the basis of sustained yield, and the development of improved methods of cultivating aquatic animals. From a scientific standpoint research is conducted in the various fields of applied sciences, including fishery biology and aquiculture, and in the fundamental sciences of ecology, oceanography, limnology, physiology, systematic ichthyology, and biostatistics. Research projects during the past year were organized under three major branches: (1) Commercial fishery investigations in marine and fresh waters; (2) aquicultural investigations, including improvements in the propagation of fresh-water fish and the study of aquatic environments and pollution control; and (3) shellfishery investigations for the development of oyster farming and the control of shellfish pests. These projects are organized under seven distinct sections, each directed

by a responsible and experienced fishery biologist, and are so distributed as to cover each of the major geographical sections of the United States.

The problems of the fresh-water fisheries throughout the country at the present time are concerned chiefly with increasing hatchery output and stocking interior waters. The chief hatchery problems concerning output are proper diet and the control of disease. Through 10 years of research the Division has found cures for several bacterial diseases of hatchery fish, and has improved diets until 1 pound of trout can be produced from 3 pounds of food, with an increase in efficiency in feeding of about one-third.

The most outstanding problems in the stocking of interior waters are concerned with the development of a stocking policy. More attention must be given to placing the fish in suitable waters which contain an adequate natural food supply, and to producing larger fish better able to survive than heretofore in order to provide better angling. In this field the Bureau is cooperating with the United States Forest Service in devising better management of fishery resources comparable with the improved game-management program.

One of the most serious limitations on the maintenance of abundant stocks of food and game fishes in interior waters and the chief hindrance in stocking these waters with additional supplies is pollution from domestic and industrial sources. The destruction of fish life by polluting substances, already acute in many localities, is growing rapidly throughout the more densely settled and industrialized sections of the country. A temporary staff, working on emergency funds, has made great progress during the past year in determining the effect of stream pollution on aquatic life. The classification of polluting substances according to their biological effects and the determination of their potencies will permit the formulation of standards of water purity favorable to the production of an abundant fish supply and form the basis of an effective policy of stream purification and protection.

The second major activity of the Division is concerned with investigations of the commercial fisheries. During the past 10 years progress has been made in determining the causes of great changes in abundance of commercial fishes. During the period continued observation of the trend and condition of the fisheries has revealed dangers of depletion and abuses in commercial practice that menace the supply. It remains, however, to educate the public regarding conservation measures to be followed voluntarily by the trade or to be enforced by legislation. It also remains to keep continual check by scientific means from year to year on the changing supply and the effects of regulation.

One of the most significant and promising new undertakings of the Division during the past year is the inauguration of a comprehensive study of the fisheries of the Columbia River Basin. Attacking the problem of the fisheries of the basin as a whole, attention is being given to such matters as regulation of the commercial fishing in the lower estuary of the Columbia, which is believed to be depleting the stock; the protection of upstream migrants from loss at power dams; the reclamation or improvement of spawning areas to increase natural reproduction; improvement in the technique of artificial propa-

gation, where such is necessary to supplement natural propagation; the protection of downstream migrants from loss in irrigation works and at power dams; and the prevention of stream pollution. A competent staff has made a satisfactory beginning in this great undertaking, and if the management of the resource is based upon continued scientific studies we may be assured of its permanent productiveness.

The third branch of activity of the Division is concerned with the shellfish industry. Oysters are the most valuable single marine product and are the most susceptible to cultivation and management. The Bureau has assisted in continuing improvements in oyster farming, but there remains a vast field for service in directing oyster planting as a means of public relief and as conservation of the resources. Technical control and experiment are still required.

COOPERATIVE INVESTIGATIONS

For many years the Bureau has had valuable informal cooperation in fishery investigations from many institutions, both public and private, but the acceptance of formal cooperation was not legally authorized until the passage of an act of Congress approved May 21, 1930. The Bureau's program has always been closely correlated with work of various bureaus of the Department of Commerce from which it secures assistance of various kinds, chiefly relating to the promotive aspects of commercial fishery investigations. During the past year closer cooperation has been established with the United States Forest Service and the Bureau of Biological Survey of the Department of Agriculture, in relation to the development of fisheries of interior waters.

Reference has been made to fishery investigations conducted by the States. California has continued with the Bureau its cooperative investigations concerned with the trout supply, looking toward securing a more adequate supply of fish in streams and the more rational regulation of fishing within its borders. New York State has continued its cooperation in the conduct of studies of the nutritional requirements of trout to improve hatchery practices in feeding and rearing. Oregon has cooperated with a Bureau investigator in a study of fish diseases and hatchery practices. North Carolina, Connecticut, and Washington have assisted in investigations for the restoration of oyster beds in their coastal waters, and Georgia, Louisiana, and Texas have taken an active part in the study of the important shrimp fisheries investigations of the South Atlantic and Gulf coast. Despite the fact that State budgets have been reduced proportional to Federal budgets for these projects, this cooperation has been very effective and is greatly appreciated.

Another source of most valuable cooperation is provided by educational institutions, chiefly universities. Laboratory quarters for the Bureau's investigative staffs are provided by Harvard University as headquarters for North and Middle Atlantic fishery investigations, the University of Michigan as headquarters for the Great Lakes fishery investigations, the University of Utah for fishery investigations in the intermountain section, University of Missouri for investigation of interior waters, and Stanford University for California

trout investigations. The Wisconsin Natural History and Geologic Survey has cooperated with the Bureau in many ways. Yale University, Cornell University, the University of Washington, and the Oregon State Agricultural College have likewise provided quarters or other facilities for investigative work.

The value of this type of cooperation cannot be overestimated. The Bureau's investigators receive, in addition to actual laboratory and office quarters, the use of university libraries, advice, and assistance from the university faculties, and many other courtesies which stimulate a community of interest in technical problems of the fisheries. The universities thus contribute to research of practical value and application to their own communities; and their graduate students receive stimulation and advice in research problems similar to those of the Bureau and frequently part-time or temporary employment in Bureau projects, all of which contributes to the progress of aquatic biology in the United States.

Grateful acknowledgment of these various forms of cooperation is made as a rule in the section dealing with fishery investigations in the various localities.

PUBLICATIONS

Owing to the curtailed funds for printing the number of publications resulting from investigations of the staff or conducted under the supervision of the Division has been reduced. The list of papers published by the Bureau during 1934 follows:

DAVIDSON, FREDERICK A.

The homing instinct and age at maturity of pink salmon (*Oncorhynchus gorbuscha*). Bulletin No. 15, 13 pp., 10 figs.

DAVIS, H. S.

Care and diseases of trout. Investigational Report No. 22, 69 pp., 15 figs.

HIGGINS, ELMER.

Progress in biological inquiries, 1933. Appendix III, Report, Commissioner of Fisheries, 1934, pp. 313-383.

HILDEBRAND, SAMUEL F., and LOUELLA CABLE.

Reproduction and development of whittings or kingfishes, drums, spot, croaker, and weakfishes or sea trouts, family Sciaenidae, of the Atlantic coast of the United States. Bulletin No. 16, 57 pp., 44 figs.

SETTE, O. E.

Outlook for the mackerel fishery, 1934. Fishery Circular No. 17, 6 pp., 1 fig.

SETTE, O. E., and A. W. H. NIEDLER.

Statistics of the mackerel fishery off the east coast of North America, 1804 to 1930. Investigational Report No. 19, 48 pp., 6 figs.

The following papers were published by members of the staff of the Division of Scientific Inquiry or cooperating investigators during the year 1934 outside of the Bureau of Fisheries' series:

DAVIS, H. S.

The purpose and value of stream improvement. Transactions, American Fisheries Society, vol. 64, pp. 63-67.

Growth and heredity in trout. Transactions, American Fisheries Society, vol. 64, pp. 197-201.

DEASON, HILARY J.

Preliminary report on the growth rate, dominance, and maturity of the pike-perches (*Stizostedion*) of Lake Erie. Transactions, American Fisheries Society, vol. 63, pp. 348-360.

The development of fishes. The Fisherman, vol. 3, no. 11, pp. 1, 3, November.

- FIRTH, FRANK E.**
Anthias nicholst, a new fish taken off Virginia in the deep water trawl fishery. *Copeia*, no. 4, 1933 (1934).
Scyllorhinus retifer embryo in capsule. *Copeia*, no. 1, 1934.
- FISH, FREDERIC F.**
 Ulcer disease of trout. *Transactions, American Fisheries Society*, vol. 64, pp. 252-258.
- GALTSOFF, PAUL S.**
 The biochemistry of the invertebrates of the sea. *Ecological Monographs*, vol. 4, 1934, pp. 481-490.
 Factors governing the propagation of oysters and other marine invertebrates. *Proceedings of Fifth Science Congress*, pp. 4119-4120.
 The mystery of the ocean. *Scientific Monthly*, August 1934, pp. 172-175.
 The use of slag in oyster culture. *Fishing Magazine*, March 1934, pp. 11-12.
- HAZZARD, A. S.**
 Low water temperature, a limiting factor in the successful production of trout in natural waters. *Transactions, American Fisheries Society*, vol. 63.
 Limnological studies of the Strawberry Reservoir, Utah. [Abstract.] *Proceedings Utah Academy of Sciences, Arts and Letters*, vol. XI.
 The Rocky Mountain whitefish. *The Rocky Mountain Sportsman*, January.
 The rainbow and the steelhead. *The Rocky Mountain Sportsman*, March.
 Improving our fishing waters. *The Rocky Mountain Sportsman*, April.
 Progress of stream improvement in Utah. *The Rocky Mountain Sportsman*, November.
- HAZZARD, A. S., and M. J. MADSEN.**
 Studies of the food of the cutthroat trout. *Transactions, American Fisheries Society*, vol. 63.
- HIGGINS, ELMER.**
 A story of the shrimp industry. *Scientific Monthly*, vol. 38, pp. 429-443, May 1934.
 Fishery biology, its scope, development, and applications. *The Quarterly Review of Biology*, vol. 9, no. 3, pp. 275-291, September 1934.
- HILDEBRAND, SAMUEL F.**
 The capture of a young tarpon, *Tarpon atlanticus*, at Beaufort, North Carolina. *Copeia*, no. 1, 1934, p. 45.
- HILE, RALPH**
 Causes of variation in the growth rates of fishes. *The Fisherman*, vol. 3, no. 2, pp. 3-4, 10-11, March-April.
- HILE, RALPH, and HILARY J. DEASON.**
 Growth of the whitefish, *Coregonus clupeaformis* (Mitchill), in Trout Lake, Northeastern Highlands, Wisconsin. *Transactions, American Fisheries Society*, vol. 64.
- HILE, RALPH, and WM. R. DUDEN.**
 Methods for the investigation of the statistics of the commercial fisheries of the Great Lakes. *Transactions, American Fisheries Society*, vol. 63, pp. 292-305.
- HOPKINS, A. E.**
 Accessory hearts in the oyster. *Science*, vol. 80, pp. 411-412.
 Accessory hearts in the oyster, *Ostrea gigas*. *Biological Bulletin*, vol. LXVII, pp. 346-355.
- LOCKE, S. B., and A. S. HAZZARD.**
 The fish of Utah. Utah--resources and activities. State Department of Education.
- LORD, RUSSELL F.**
 Hatchery trout as foragers and game fish. *Transactions, American Fisheries Society*, vol. 64, pp. 339-345.
- NEEDHAM, P. R.**
 Quantitative studies of stream bottom foods. *Transactions, American Fisheries Society*, vol. 64, pp. 238-247.
 Notes on the food of trout. *California Fish and Game*, vol. 20, no. 2, pp. 119-127.
- NEEDHAM, P. R., and A. C. TAFT.**
 Observations on the spawning of steelhead trout. *Transactions, American Fisheries Society*, vol. 64, pp. 332-338.

NEARIT, R. A.

A convenient method for preparing celluloid impression of fish scales. *Journal du Conseil*, vol. 9, no. 3, pp. 373-376.

The fishery conservation problems of Long Island. *Fishing Gazette*, vol. 51, no. 8, pp. 27 and 29, and *Fishing Gazette*, vol. 51, no. 9, p. 41.

PEARSON, JOHN C.

Marking fish. *Louisiana Conservation Review*, vol. 4, no. 2, pp. 30-33, April.

PATTHEBCH, HERBERT F.

The role of copper in the setting, metamorphosis, and distribution of the American oyster, *Ostrea virginica*. *Ecological Monographs*, vol. 4, no. 1, pp. 47-107.

Scientific methods of oyster farming. *Scientific Monthly*, vol. 38, no. 2, pp. 118-128; *Atlantic Fisherman*, vol. 15, no. 4, pp. 17-18; *Fish and Oyster Reporter*, vol. 15, no. 8, pp. 2-3, 10-11.

Rehabilitation of public oyster beds in North Carolina. *Atlantic Fisherman*, vol. 15, no. 2, p. 14.

Oyster parasite threatens production in Florida. *Fishing Gazette*, vol. 51, no. 13, pp. 11 and 27.

TAFT, A. C.

California steelhead experiments. *Transactions, American Fisheries Society*, vol. 64, pp. 248-251.

VAN OOSTEN, JOHN.

On the deep trapnet in the State of Michigan. [A letter, Dec. 13, 1934.] *Michigan Tradesman*, no. 2674, p. 25, December.

The value of questionnaires in commercial fisheries regulations and surveys. *Transactions, American Fisheries Society*, vol. 64.

VAN OOSTEN, JOHN, H. J. DEASON, and FRANK W. JOHNS.

A microprojection machine designed for the study of fish scales. *Journal du Conseil*, vol. 9, no. 2, pp. 241-248.

WRIGHT, STILLMAN, and WILBUR M. TIDD.

Summary of limnological investigations in western Lake Erie in 1929 and 1930. *Transactions, American Fisheries Society*, vol. 63, pp. 271-285.

The following progress reports covering the more important investigations of the Division during the calendar year 1934 were prepared in the main by the investigators in charge of the various projects.

NORTH AND MIDDLE ATLANTIC FISHERY INVESTIGATIONS

The experience of the last several years of economic depression has brought into focus an aspect of the economy of our marine fisheries hitherto overlooked, but of great significance to the fundamentals of conservation of our marine resources. It is generally agreed among fishery biologists that few, if any, of our marine species can be reduced to extinction by overfishing. Rather, the effect of progressive depletion is to lower the level of abundance, to reduce the return per unit of fishing effort, to raise the cost of production, and to increase prices. Theoretically, this process might continue indefinitely so that the products become luxuries utilized only by the more affluent consumer. But this means practical extinction of the fishery as a commercial enterprise. Obviously, the industry must depend on reasonable volume, which, in the fisheries, can be had only while its products remain low enough in price to remain in the staple class. The optimum condition, of course, is found where the level of abundance is such as to provide a maximum continuing volume of yield. However, the experience of New England's fisheries in the past several years has brought to light another limitation; that is, the level of abundance must be such that fish can be pro-

duced at a cost that can meet price competition with meats. This was not demonstrated by depletion of fish to the point of raising costs of production but by the lowering of meat prices to such a point that fish could not compete successfully at the ordinary level of abundance and ordinary cost of production.

It is imperative, therefore, that there be developed means of observing the changing levels of abundance and bringing forces to bear to counteract trends in the dangerous direction. This is being done in the case of several of the most important species of this region, including haddock, mackerel, squeteague, and scup, as will appear in the following pages. But the staff and its facilities have been inadequate to cover other almost equally important species, among them the lobster, shad, flounders, sea bass, bluefish, and cod, not to mention some twenty-odd other species of lesser importance but constituting in the aggregate an important segment of the coastal and offshore marine fisheries.

Even the species receiving consideration cannot be accorded the amount of study that the problems connected therewith require for solution. The most serious handicap under which the staff is operating is the lack of a sea-going research ship equipped for trawling. It is badly needed to survey the stocks of young haddock, upon which the future of the trawling fleet depends; to operate experimental gear so as to develop means of saving young fish from needless destruction now attending certain methods of fishing; to catch live fish for tagging in order that their migrations may be followed; and to take observations on conditions of the sea water of our fishing grounds that determine the success or failure of future broods. Additional personnel is needed to perform field work and to assist in the analysis of the field data. The fisheries of the region provide products valued at \$25,000,000 annually as landed at the dock. The annual expenditure for the research on marine fishery problems of the region is only one one-hundredth of 1 percent of the value. This rate is too low to provide adequate insurance against unwitting over-exploitation on the one hand or unsound restrictions on the other hand.

As in former years, the staff, under the direction of O. E. Sette, has been provided with laboratory and library facilities by the Harvard Biological Laboratories and the Museum of Comparative Zoology at Harvard University, Cambridge, Mass., where its members have also benefited from consultation with members of the university, especially Henry B. Bigelow, professor of oceanography and director of the Woods Hole Oceanographic Institution, whose wealth of knowledge and experience relating to marine fisheries research has been ever at the disposal of the Bureau employees. It is a pleasure also to acknowledge the continued cooperation of fishermen and fishing companies in providing data essential to the progress of the work.

HADDOCK

In 1934 the fishing activities of the New England haddock fleet were extended over a much wider area than in previous years, yet the total haddock catch was little greater than in 1933. The landings at the principal ports (these include all landings except a rela-

tively small quantity at minor ports) reached about 135,000,000 pounds compared to 130,000,000 in 1933. However, for the first time in the history of the fishery the greater part of the catch came from the banks off the Nova Scotian coast (Sable Island, Banquereau), Browns, and La Have. The catch on Georges Bank (including South Channel and Nantucket Shoals), which in past years has averaged nearly 80 percent of the total United States landings, in 1934 declined to 40,000,000 pounds, the lowest in the history of the modern fishery, and but 30 percent of the total. The catch on the Nova Scotian banks by United States vessels reached 88,000,000 pounds.

The sharp decline in the Georges Bank fishery was the result of the continued poor fishing in this area, which caused practically all of the large trawlers and many of the large draggers to spend most of their time on the Nova Scotian banks. The great increase in running time to and from these more distant banks (Banquereau is about 625 miles from Boston, while Georges averages 180 miles) was more than compensated for by the greater catch per day's fishing.

The actual changes in the haddock population have been studied during 1934 along the lines described in previous annual reports. Investigation has continued under the direction of W. C. Herrington, assisted by J. R. Webster. Owing to a reduced staff and funds, field work has been confined almost entirely to the Boston Fish Pier where vessel interviews have provided information as to fishing time, fishing position, and other relevant data for practically all haddock trips landed. Extensive length-frequency data and scale samples were also obtained throughout the year from commercial catches made in each of the principal areas fished. F. E. Firth handled most of the work at the fish pier during the winter, and J. R. Webster or W. C. Neville during the summer, while Mr. Firth was assisting in the mackerel investigations.

Georges Bank and adjacent areas.—The haddock population in this area has been at a low level of abundance since 1929 and has shown little indication of recovery. In 1932 there was some improvement in the marketable stock caused by the growth to commercial size of the relatively numerous 1929 class. The gain was short-lived, however, for 1933 saw the level of abundance, as indicated by catch per day, again decline. This decline was caused by reduction in the stock through catch mortality (mortality due to the fishermen's catch) and natural mortality. The 1930 year class, which reached commercial size during 1933, was much too scanty to furnish sufficient contributions to the marketable stock to compensate for this mortality.

It has been mentioned that the 1934 haddock landings from Georges Bank amounted to only about one-half the quantity landed in 1933. However, this was due to the shift of a large part of the fishing effort to the Nova Scotian banks rather than to a decline in abundance on Georges. In fact, the catch per day for the first 8 months of 1934, as shown by our catch analysis, was some 15 to 20 percent higher than in 1933. (Figures for the balance of the year are not yet available.) There is some probability that this improvement in catch per day does not represent a proportional increase in

the marketable population. The number of boats on which the 1934 catch analysis is based was reduced to about one-half of the large otter trawlers, a reduction made necessary by a much curtailed staff. This reduction was made reluctantly after a careful analysis had demonstrated that during 1932 and 1933 the fluctuations in the average catch per day of the smaller group were representative of the entire fleet. But during 1934 the principal activities of the large trawlers were transferred to the Nova Scotian banks and the boats in question remained on Georges Bank only when fishing was relatively good. If catches were poor, the boats moved to the more distant banks. Consequently, the higher 1934 catch per day may be due partly to this selective fishery and not represent an equally great increase in abundance. Limited facilities have not yet permitted us to examine this possibility through the analysis of records of the smaller boats which regularly fish on Georges Bank.

There seems little doubt but that the size of the marketable stock of haddock on Georges Bank was somewhat greater in 1934 than in 1933, even though the increase was not as great as indicated by the catch-per-day figures. This improvement was due partly to the fairly good 1931 class which reached commercial size during the winter of 1933-34 and spring of 1934, and partly to the great decrease in the strain imposed on the stock by the reduced commercial catch. With only about half as much haddock caught out during 1934 as in the several preceding years, the population level was maintained and even somewhat increased with only moderate additions of young fish.

The success of the fishery on Georges Bank during any year—that is, the return for a given unit of fishing effort—is primarily dependent on two factors: (1) The rate at which the commercial stock declines as the result of catch and natural mortality (this is largely determined by the intensity of the fishery); and (2) the contribution to the commercial stock from the young haddock spawned 3 years earlier. Thus, the 1929 class boosted the catch in 1932; and the 1931 class, in 1934.

To make accurate forecasts for the fishery 1 or 2 years in advance it is necessary that the above-mentioned factors be known. With present facilities for observation it is possible to determine the rate of decline of a stock of fish once it has reached commercial size, providing it is possible to estimate the amount of fishing effort that will be expended in that area. However, we now have no means of determining how much an incoming year class will affect the commercial catch, for we have no observations by which we can evaluate the abundance of such year classes until they achieve commercial size and have been represented in the commercial landings for a period of nearly a year. Inasmuch as there are great fluctuations in the abundance of the different year classes, the annual contribution from this source is highly variable and is the most important factor in determining the fluctuating level of the fishery. The relative level of abundance in two general areas such as Georges Bank and the Nova Scotian Banks largely determines where the fishing fleet will expend their major efforts; consequently, it determines the intensity of the fishery in either area. Therefore, it follows that the level of abundance of the marketable stock during the course of any year is affected both directly and indirectly by the contribution of young

haddock to this stock. To estimate the size of this contribution a year or more in advance, it is necessary that it be possible to evaluate the abundance of haddock before they reach commercial size. For this work it is essential that a vessel be available for the collection of extensive samples of young haddock with small-meshed otter trawls.

An example of the limited reliability of forecasts based on present observations can be found in the 1933 annual report on haddock. That report discusses the probable course of the fishery on Georges Bank during 1934 and 1935. At that time it appeared that the catch per trawler day in 1934 would be somewhat less than in 1933. Actually the catch was somewhat better, due to the great reduction in fishing effort and to unexpectedly large contributions from the 1931 class, which it had been impossible to evaluate accurately until 1934, when it appeared in the commercial catch. Similarly, in the 1933 report the probability was expressed that the 1932 class was a failure and that consequently the fishery would continue to decline in 1935. Results of the fishery in the last part of 1934, however, indicate that the 1932 class is much larger than was supposed and may furnish a considerable contribution to the commercial stock. The abundance of this class cannot be determined from commercial-catch data until late in 1935. Consequently, at the present time it is impossible to make any predictions as to the 1935 fishery on Georges Bank except the general one that with fishing effort distributed as in 1934, the catch per day in 1935 probably will be considerably better than in the previous year.

Nova Scotian banks.—In 1934 nearly 90,000,000 pounds of haddock were landed from this area. This was about two-thirds of the entire United States catch and approximately twice as much as was ever before landed from these banks by the United States fleet. The actual level of abundance of commercial haddock, as shown by catch per day during the first 8 months of 1934, was about 10 percent higher than during the corresponding part of 1933. The increase was due to the 1929 class which came into the commercial fishery in the summer of 1933. (This same class on Georges Bank came into that fishery in the fall of 1931 owing to the faster growth rate in that area.)

The development of the fishery during the first 8 months of 1934 was much as anticipated in the 1933 annual report. At that time it was predicted that there would be an improvement in 1934 over 1933 in the catch per day during the spring and summer and a decline in the fall and winter. The spring and summer fishery has followed this trend although for various reasons the increase was not as great as was anticipated.

The fishery on the Nova Scotian banks should continue good in 1935 but somewhat below the 1934 level. However, this estimate can be no more than an approximation because of certain unevaluated factors. The 1931 year class will reach commercial size during the summer and fall of 1935 but it does not appear to be sufficiently abundant in this area to compensate fully for the decline in the present marketable stock owing to catch and natural mortality. Unfortunately, no reliable measure of the abundance of the 1931 year class is available due to the scarcity of information concerning the submarketable sizes. Our only basis for an estimate is a series

of length-frequency samples of unculled haddock obtained through the assistance of Messrs. Einar Sorenson and R. H. Blake, radio operators on two of the large otter trawlers.

Early life history.—One plankton and hydrographic cruise was made in April–May 1934, covering the area from Cape Cod and Nantucket Shoals to Browns and La Have Banks off the Nova Scotian coast. As in 1933 the trip was made possible through the kindness of the Woods Hole Oceanographic Institution in detailing the research ship *Atlantis* for this work. At each of the 64 stations oblique tows were made with a 2-meter stramin net and a ½-meter silk net. Vertical tows were made at some of the stations with a 1½-meter Helgoland larvae net or 1-meter silk net. Temperature and salinity data were obtained at each station and 629 drift bottles and 6 large drift floats released at stations and in strings. The 2-meter and ½-meter nets, used with current meters and the electrical inclinometer developed for this work, provided some rather definite data on the effects of clogging and change in towing speed. A cursory examination of the collections indicates that medium-sized haddock larvae were taken in the region southwest of Georges Shoals in greater numbers than at any stations on previous cruises. Drift bottle returns at the end of December 1934, totaled 64. About one-half of these were from the Bay of Fundy and but one-fifth from the United States coast west of Georges Bank. Four reports were received of positions of the drift floats, one float being reported twice. The surface drift as shown by these data appears similar to 1932.

During the past year L. A. Walford of the Harvard Graduate School completed the analysis of the egg and larvae collections of 1931 and 1932. The results show spawning concentrations on eastern and northern Georges and sometimes in the South Channel and clearly demonstrate the effect of water movements on the distribution of eggs and larvae. Evidence also is found that the southerly drift off Georges Bank such as observed in 1932 may be a factor in the failure of certain year classes.

Savings gear.—During the 5 years since 1929 the haddock population on Georges Bank has been held at a low level of abundance in spite of the contributions to the commercial stock from two fair year classes. The continued low level of abundance is the result of the intensive and destructive fishery that has developed in recent years. Although only those haddock weighing 1½ pounds or more are marketable, most of the commercial trawlers use nets which capture fish of all sizes from about one-fourth to one-half pound up. Nearly 2 years are required for the young haddock to grow from one-fourth pound to the marketable size of 1½ pounds. During this period they are constantly subject to decimation by the commercial fleet, and the millions that are thus destroyed constitute a complete economic and biologic loss and reduce the stock of fish on the banks to the same degree as the capture of equal numbers of large, commercially valuable fish.

An investigational report covering the results of the Bureau's savings gear experiments was submitted for publication during 1934. This work shows that the most practical method for reducing the destruction of undersized haddock is to increase the size of mesh used

in the otter trawls. A minimum cod-end mesh size of $4\frac{3}{4}$ inches (about $4\frac{1}{8}$ inches after use) will capture only about one-fifth as many undersized haddock as the 3-inch mesh that generally has been used in the commercial trawls in the past. At the same time this mesh will lose less than one-tenth of the haddock between 1.5 and 2.0 pounds, and none above that size. The report recommends that the industry adopt a minimum mesh size of not less than $4\frac{3}{4}$ inches to be used in any part of the otter trawls except certain portions of the cod-end; e. g., belly and after end.

It is important that the recommendations of this report be followed by definite action. The trawler operators in general have recorded themselves in favor of the adoption of large-meshed nets, but for various reasons there has not been much consistent progress toward the use of such gear. A move for definite Federal or international regulation appears necessary for real progress. It is not a matter that can be postponed indefinitely, for already there has been a trend toward marketing the smaller sizes of haddock and once this practice becomes established it will be very difficult to obtain the adoption of large-meshed gear that will reduce the catch of such fish. The capture of small haddock, even though marketable, is a very short-sighted and irrational practice, for on Georges Bank 1 year's growth will cause a 1-pound haddock to nearly double and a half-pound haddock to nearly treble in weight; thus not only reaching a size that is of more value per pound but also increasing the actual poundage of fish in the ocean available to the fishermen.

In the light of present knowledge of the species the most practicable action that can be taken for the conservation of the haddock fishery is to leave the fish in the ocean during their early years of rapid growth and concentrate the fishery on the larger sizes where natural mortality approximately neutralizes the increase in the weight of the stock through growth. Such action would increase materially the catch that might be taken annually from the fishing banks. The use of mesh with a minimum size of $4\frac{3}{4}$ inches, as recommended, will protect most of the young haddock on Georges Bank until the end of their third year. This by no means is the ultimate to be desired, for a fourth year on the bank would cause a weight increase in these fish of about 70 percent, an amount that is much greater than the loss in poundage due to natural mortality. However, the recommended mesh size of $4\frac{3}{4}$ inches appears to be immediately practicable, and if acted upon would constitute a very considerable contribution to the future of the fishery.

The results of the United States and Canadian haddock investigations were presented at the 1934 meeting of the North American Council on Fishery Investigations. After a discussion of these results the council approved the report of the Committee on Haddock Investigations. Following are some of the more important recommendations:

The committee endorses the opinion expressed at the general meeting of the council that the haddock problem has become of major importance to Canada and the United States (on the Atlantic coast); that immediate steps should be taken to adopt a common plan of investigation, which would adequately keep the situation under review and lead to the adoption of remedial measures.

Further, the committee urges the adoption of a joint program of intensified research into the biology of the haddock, * * *

The committee finds that existing facilities are quite inadequate to enable this broad program to be entered into. The prime essential is a suitably equipped research vessel. * * *

Additional personnel will be essential in both countries, as far as the United States part of the program is concerned at least two fish measurers or quay men would be required to collect statistics of commercial catches, and a technical assistant would be necessary to assist in laboratory work and sorting of material.

It is recognized that, even by making such provision, the authorities will not provide for a full completion of the program outlined. Some aspects, especially those of the early haddock stages, will receive quite inadequate attention in the absence of the provision of a suitable research vessel.

It has not yet been possible to make a start on any of the projects of the joint program because of the unavailability of additional personnel or facilities.

MACKEREL

The paramount influence on the stock of mackerel in the sea (and consequently the size of the catch) is the variability in the degree to which annual recruits of young mackerel serve to offset the decline due to mortality of the older stocks. Since 1925 continuous observation of the catch and its age composition has proved that certain year classes, such as those of 1923, 1928, 1930, and 1931, were so plentiful as to more than offset mortality of the older stocks. Certain others, those of 1924, 1927, and 1929, were so poor that they did not fully counterbalance mortality of the adult stock; and still others, those of 1925 and 1926, were practically nonexistent.

These variations in numerical strength of the different year classes have been the dominating cause of all the fluctuations that have occurred in the last 9 years and have formed the basis of annual predictions on yield since 1928.

Although the ultimate causes of fluctuations—the conditions responsible for inequality of the year classes—remain unknown, it is fairly certain that they cannot be controlled by man, for they come into play while the mackerel are very young and before commercial fishing has any effect on them. There remains, however, the problem of making the best possible use of such year classes as the conditions in the sea permit to survive. This is being done in part by the predictions already mentioned for they permit the members of the industry to foresee, in general, the prospects for the coming season. Thus, the disorganizing effects of fluctuating supply can be minimized to the extent that the affairs of their business can be altered to meet the situation.

These predictions take into account only the general level of abundance for the season as a whole. In many ways it would be more useful to have foresight of what is to happen during the parts of the season. For instance, if scarcity were destined to prevail during the latter part of the season, freezing and salting could be accelerated during the early parts and vice versa. This would benefit the fisherman by expanding the market during times of glut and benefit the buyer by avoiding to a greater degree the understocking or overstocking of frozen and salted mackerel.

The third and perhaps most important question to be solved before the most effective use can be made of this resource is the relative value, biologically speaking, of the yearling mackerel. Following the advent of successful year classes, these are caught in large quanti-

ties. They are fairly small ($\frac{1}{2}$ to 1 pound each), have poorer keeping properties than the larger mackerel, and command a lower price per pound. At the same time they are growing fast and are destined to double in weight during the next year. It is appropriate, therefore, to inquire whether a partial or total elimination of the yearling mackerel from the catch might not be more than repaid by the larger size and the greater value per pound of these fish in future years. Essentially it is a question of whether apparent mortality (rate of disappearance as measured by the catch per unit time of fishing) is greater than the increase in value due both to increase in poundage and price per pound.

The answer to this question is complicated by the fact that the rate of disappearance of yearling mackerel differs greatly from one year to another and that the differences are connected not with the numbers caught out during the yearling year but with the type of year class to which they belong. There appear to be at least two types of mackerel in the western Atlantic; one, which we have designated as "persistent"; the other as "transitory." The former is characterized by a low rate of annual disappearance, the latter by a high rate. They are distinguished by other peculiarities such as the time and locality of greatest concentration and differences in rate of growth.

The existence of the two types with their different rates of disappearance not only renders difficult a decision as to the merits of exploiting yearling mackerel but also causes predictions to be less assured and less definite. It is essential, therefore, to determine the basis for the differences in behavior of these two subdivisions of the population and especially to find means of distinguishing between them at an early stage, in order to foresee their subsequent decline.

Naturally, the work during 1934 consisted of collecting and analyzing data necessary for determining the rate of catching and the year class composition of the commercial run. This has been and must continue to be the only means of appraising the current condition of the fishery. In addition, a new technique was employed to analyze the size frequency distribution of the 9-year period to throw more light on the nature of the two types of year classes, and the catch statistics were reanalyzed to discover whether the periods of presence of the two types were sufficiently distinctive and regular to serve as a basis of within-the-season predictions.

The work continued, as in the past, under the direction of O. E. Sette. Field data were collected by F. E. Firth. They consisted of 1,413 interviews to obtain information on time and locality of catches, measurement of 39,605 mackerel, and collection of 1,467 scale samples.

The prediction for 1934 was that the abundance would be the same as in the previous year and that the catch by seiners would amount to approximately 54,000,000 pounds if fishing were unrestricted, and proportionately less if control of production were to take place under provisions of the mackerel code. If these were to be of equal extent to those employed by voluntary agreement, a catch of 28,000,000 pounds was to be expected. Actually, production control was exercised during a smaller portion of the season and the catch was ac-

cordingly larger, being about 35,000,000 pounds. The level of abundance has not yet been estimated, there being extended computations required to adjust for the effects of production control on the measure of abundance. However, it appears that the abundance was practically equal to or slightly above the predicted level.

During 1934 O. E. Sette served as administration member of the executive committee of the Atlantic Mackerel Fishing Industry Code. The provisions of the code were effective in preventing the dumping and wastage of mackerel such as had occurred during the previous season but were less successful in substantially increasing the income of mackerel fishermen.

Lack of sufficient personnel has prevented progress during the year on the vital matter of finding out the nature of the two types of mackerel and the means of distinguishing them at a sufficiently early stage to be of use in forecasts, as well as to throw light on the merits of exploiting yearling mackerel. This work would require large-scale tagging experiments at sea accompanied by examination of large numbers of mackerel from various areas and various portions of the season to discover whether there are reliable racial characters that distinguish the two types. Although it was impossible to include these particular items in the program, a small beginning was made on examination of mackerel for racial characters, but the numbers of individuals and the distribution of samples were far too limited to provide any conclusions.

If proper utilization of this important resource is to be attempted, it will be necessary to provide additional personnel both for field observations and laboratory analysis.

COD

Although the total catch of cod has not suffered a decline of the sort to cause concern over the future of this very important resource, there is good reason to suppose that the time may come in the not-too-distant future when heavier fishing intensity may have its effect. Even now the catch in southern New England, New York, and New Jersey waters is undergoing a severe decline, and we are ignorant of its cause or the remedies. Previous work has shown that this population is practically independent of the larger bodies of cod to be found from Georges Bank eastward to the Grand Banks, and it is possible that this southern segment of the species is undergoing a decline that should receive remedial attention.

For some years in the past the Bureau conducted large-scale tagging experiments on the cod which provided enlightenment on the relationships of the stock occupying the several grounds, showing that the southern segment was independent of all others, that the Maine coast was also primarily self-contained though acting as a feeder to the coastal waters of Nova Scotia and possibly to offshore grounds such as Georges Bank.

With the resignation of William C. Schroeder from the Bureau's service, this work ceased except for the analysis of returns currently received from tagging done during 1931 and 1932 which Mr. Schroeder has kindly continued to study and upon which he reports as follows:

Of 1,199 cod marked during 1931 with celluloid disks attached to the caudal peduncle and released off Mount Desert, Maine, 23.4 percent were recaptured

up to the end of 1932, 1.1 percent in 1933, and 0.3 percent in 1934. Of the total recaptures (24.8 percent), 90 percent were taken locally and 10 percent from distant localities. Of 1,481 cod marked during 1932 with celluloid strips inserted into the body cavity and released in the above-mentioned locality, 18.6 percent were returned up to the end of 1933 and 2.8 percent in 1934. Of the total returns of 21.4 percent, 91 percent were recaptured locally and 9 percent at distant points.

These experiments were undertaken in order to determine whether the young cod that predominate in the population along the shore migrate away from the shore as they grow older, thus serving to replenish the important fishing banks offshore. While these experiments have produced a higher proportion of distant recaptures than resulted from earlier experiments with less permanent tags, there does not appear to be any decided trend toward offshore recaptures as the fish grow older. To be sure, the fish involved in the experiment have not yet reached the age that is most common on the offshore grounds, and the experiment needs to run until they do reach such age before negative results become significant. However, the sharp reduction in returns threatens to reduce the numbers to the point of unreliability before this stage in the experiment is reached. It is believed that the sharp decline in returns from the 1931 releases is due to loss of the celluloid disks, and from the 1932 releases, to the belly tags becoming covered with mesenteric tissue so that they escape detection, especially on the offshore grounds where the fish are cleaned very hurriedly. Thus, it may be necessary to have further improvement in marking methods or to approach this question by another method, perhaps scale studies, in order to solve the problem.

At present the trawling fleet fishes mainly for haddock. Studies of the haddock resource (see above) make it appear certain that increased fishing will cause no considerable or sustained increase in the yield. This being the case, we can anticipate a more intense fishery for cod. In fact, there appears to be a trend in this direction already, and it is urgent, therefore, that studies of the condition of the cod resource be resumed. The most effective corrective measures are those that are applied before irreparable damage is done. Since the major portion of the fishery is carried on by a fleet of vessels which land their catches mainly at Boston, only a moderate outlay would be necessary to make the pertinent observations on the condition of this fishery along with those now made on haddock.

FLOUNDERS

The flounder fishery has assumed major importance during the last two decades. The various species of flounders are caught mainly by the smaller vessels of the otter trawl fleet commonly called "draggers." This type of boat has been increasing in numbers and concern is felt as to the capability of the resource to withstand the present intensity of fishing—so much so that several of the flounder-producing States have passed laws restricting the flounder fishery in various ways. Further restrictions are contemplated in at least one of these States. Although such restrictions are based on good intentions, they have not been preceded by any examination into the question of whether they will produce the desired results. No one knows whether the present sacrifices they impose will be compensated by future gains. In fact, little is known of the life history of the six species that make up the catch. Nothing is known of their rate of growth, the amount of migration from one region to another, or most important, to what degree the annual replacement compensates for removals by the commercial fishery and by natural mortality.

A single small experiment was undertaken in 1931 when some 4,000 winter flounders from Waquoit Bay were tagged and released. The returns in successive years from 1931 to 1934, inclusive, were 141, 64, 33, and 11. Unfortunately, we do not know what proportion of the decline in returns was due to loss of tags and what portions to mortality, but the results do not preclude the possibility that the rate of mortality of flounders is relatively high. The returns indicate that most of the Waquoit Bay stock returns there for spawning each winter, but during the summer they wander into the adjacent sounds and into the open sea beyond. During last season there was one return from Massachusetts Bay—the first indication that the flounders from south of Cape Cod may wander to its northern side. Previous distant returns were from Georges Bank to the east and from western Long Island to the west.

To formulate an effective conservation policy, it will be necessary to institute a series of observations similar to that employed in the case of the haddock and mackerel, whereby statistics are collected in such detail that an index of abundance could be computed in terms of catch per day's trawling. Field data for determining rate of growth and for arriving at the age composition of the catch from each important fishery area would need to be collected and studied, and additional tagging would have to be done to determine to what extent one locality is dependent on another for its stock. The distribution of the flounders is extensive so that observers would be needed at at least four ports. Thus, a staff of four members or more would be required for the field work alone and additional persons would be needed in the laboratory.

The need for undertaking this work is imperative. The total yield of flounders has already begun to decline (20 percent between 1929 and 1932) and several seasons must elapse before results from an investigation can be expected. If this fishery is allowed to retrograde too seriously before corrective measures are applied, either these will be more drastic than if applied earlier or the fishery will be stabilized at a lower level of yield than would otherwise need be the case.

SPECIES IN NEED OF ATTENTION

Lobster.—The lobster fishery has declined steadily in productivity since the time that earliest statistical evidence is available. The present annual catch is but one-third as large as in 1889, and, in spite of the ever-increasing number of lobster pots in operation, the present rate of decline is 10 percent per annum. Consequently, the annual decrease in abundance must be substantially more than 10 percent.

The decline has persisted for years in spite of State laws intended to preserve the resource. These laws consist mainly of a limit on the size of lobster which may be sold legally. It is imperative to determine, by a survey of the lobster population, whether such size limits protect a sufficient percentage of the stock. Furthermore, since the regulations on sale of undersized lobsters have been found extremely difficult to enforce, it is further necessary to find some more effective means of securing the necessary protection. It is desirable, therefore, to test lobster pots designed to catch only certain sizes and introduce such modifications as are indicated by the tests. If such pots prove

satisfactory, both from the standpoint of catching efficiency and from the standpoint of excluding the protected sizes from the catch, a definite improvement in regulation could be attained.

After an initial survey has determined the percentage of the population to be protected by size limits or other means, a continuous observation of the effects of new regulations would be necessary to determine their adequacy.

SHORE FISHES OF THE MIDDLE ATLANTIC STATES

The shore fishes of the Middle Atlantic States (New York, New Jersey, Pennsylvania, and Delaware) are of especial interest because they not only support an important commercial fishery but also provide a basis for a sport fishery the magnitude of which is not generally appreciated. Quite aside from its recreational value, this sport fishery has commercial aspects of great importance. The business of catering to the needs of anglers (boats and guide service, transportation, tackle, and bait) has reached astonishing proportions and has become an important item in the economics of seashore communities in these States.

Investigations of these shore fishes were continued under the direction of R. A. Nesbit, assisted by W. C. Neville. As in 1933, reduction of appropriations necessitated severe curtailment of activity. Field work was limited to a single visit to producing areas in October and to a 3-month observation of the catches made by the winter-trawl fishery. Prof. A. E. Parr, curator of the Bingham Oceanographic Foundation, continued his studies of the early life histories of these fishes in southern New Jersey.

Squeteague.—By the end of 1933 satisfactory methods of investigation had been developed and enough of the data collected between 1927 and 1933 had been analyzed to permit a summary of the principal scientific findings with regard to the life history of the squeteague. In 1934, additional scale and length frequency data, secured in October, were analyzed, and additional tag returns were received. Also, a large part of the earlier scale collections were examined according to the circulus spacing method for determining origins and subsequent migrations. The results confirm and strengthen the views expressed in the 1933 report.

Since the conservation recommendations which follow are based on these scientific findings, it is desirable to summarize them briefly:

(1) Not more than one-half of the squeteague taken north of Delaware Bay are the product of local spawning. This view is based on the observation that not more than one-half of the northern squeteague have scales of the type characteristic of northern juveniles. It is possible that even these northern juveniles are the product of southern spawning, migrating to northern waters early in the first summer of their lives. This suggestion is based on the observations of Professor Parr. Although he has found squeteague eggs in several New Jersey and New York localities each year from 1929 to 1934, the larvae are uniformly absent from the collections. This is in marked contrast to the presence of larvae in Virginia and North Carolina waters as reported by Hildebrand and Cable (1934) and by Pearson (unpublished manuscript).

(2) Nearly all (90 percent or more) of the squeteague taken in the Middle Atlantic States are 2 years of age or older, and nearly all have passed their yearling summer south of Delaware Bay. This view is based on age analysis of large samples collected at various localities in New York and New Jersey between 1928 and 1934 and on the observation that the second (yearling) growth zones of the scales of virtually all northern market squeteague are of the type characteristic of the corresponding growth zones of southern yearlings. The conclusions drawn from scale studies are confirmed by tagging experiments.

(3) Yearling squeteague predominate in the southern catches. This observation is based on age analysis of samples taken by Higgins and Pearson in 1925 in North Carolina and of samples taken during the present investigation in Virginia and North Carolina in 1928, 1929, 1931, 1933, and 1934.

(4) Squeteague grow much more rapidly in the North than in the South. For example, squeteague at the end of their third summer average only 11 inches in length and 0.43 pound in weight south of Delaware Bay. In southern New Jersey they average 12 inches in length and 0.56 pound in weight and in northern New Jersey 13 inches in length and 0.75 pound in weight.

For a number of years restriction of both the commercial and angling catch of squeteague has been urged in New York and New Jersey. In New York such proposals have invariably been rejected by the legislature. In New Jersey a number have been adopted, among them prohibition of purse seining within 2 miles of the shore, restriction of the length of gill nets, specification of minimum mesh size for gill nets, and imposition of closed seasons for gill-net fishing in certain areas. Restrictions are usually urged on the ground that abundance may be maintained or increased in the future by protecting spawning adults.

In the light of our present knowledge of squeteague life history, it is apparent that restriction of the New York and New Jersey catch of adult squeteague cannot be expected to influence future abundance sufficiently to compensate for the immediate sacrifices required. As has been pointed out, the northern spawning stock is not wholly self-perpetuating but depends on southern spawning for at least half of its annual increment even if it be assumed that northern juveniles result from northern spawning. This means that of the fry resulting from the spawning of 2 northern adults, on the average but 1 weak fish returns to the North 2 years later. Consequently, in order to add 1 more weak fish to the catch in 1937, it would be necessary to refrain from catching 2 in 1935, obviously an unprofitable bargain, even if the cost of enforcing restrictive legislation be left out of account.

It is necessary to consider other measures if the present unsatisfactory conditions in the Middle Atlantic States are to be improved. The yield of squeteague in these States has declined from an average of 17,000,000 pounds per annum for the years 1901, 1904, and 1908 (years for which statistics are available), to an average of 12,000,000 pounds for the years 1926, 1929, 1930, 1931, and 1932. It is highly probable that this decline is in large part due to an increase in the catch of the southern fishery (Maryland, Virginia, and North Caro-

lina) from an average of 11,500,000 pounds per annum in 1901 and 1908 to an average of 18,000,000 pounds per annum for the years 1925, 1929, 1930, 1931, and 1932. It has been seen that virtually the only squeteague available to the northern fishery are those which escape the intensive southern fishery to which they are subject during their yearling summer. If the increase in yield in the South were due to increased abundance rather than to an increase in the percentage toll taken by more intensive fishing, it is to be expected that a similar increase would have occurred in the northern fishery.

It does not necessarily follow, however, that the most effective use of the squeteague resource requires restriction of the southern fishery. It is unwise to postpone capture of fishes beyond the point where natural mortality removes more pounds of fish than are replaced by growth. Since the investigation has not progressed sufficiently to make possible reliable estimates of natural mortality, it cannot be said fairly that the yearling squeteague taken in the south would be more valuable if spared for an additional year or more.

Although there is insufficient evidence to justify restriction of that portion of the southern catch which is actually utilized, the present destruction of juvenile and yearling squeteague too small to be marketed cannot be too strongly condemned. As pointed out by Higgins and Pearson (1925), this destruction may amount to as much as 30 percent of all squeteague taken. There can be no reasonable objection on the part of southern fishermen to elimination of this waste, for enough of the benefit would accrue locally to compensate for any inconvenience brought about by changed methods of fishing. It is urgent, therefore, that experiments be undertaken at once to devise modifications of the gear to permit the escape of undersized fish.

Scup.—Continued investigation of this species by W. C. Neville during 1934 has been concerned with determining the effect, if any, on abundance of the increased exploitation to which this species has been subjected since about 1929. As stated in previous reports, the rapid development since that year of a winter trawl fishery off the Virginia capes for scup, sea bass, and fluke has resulted in an additional strain of approximately 25 percent on the general stock of scup. The analysis, therefore, has been mainly a study of the fluctuations in the catch of both the summer and winter fisheries to determine whether the changes are related to the increased fishing activity.

It has been demonstrated clearly from tagging experiments and from the ages and sizes of the fish in the catch that the summer and winter fisheries are drawing on the same stock of scup. Tagging experiments have demonstrated that scup migrate in the fall from the summer fishing grounds along the shores of southern New England, New York, and New Jersey to the winter fishing grounds off the Virginia capes, and in the spring make a return migration from the winter grounds to the summer area. Analysis of the catches has disclosed that broods of scup that have been conspicuous in the summer fishery have constituted a large part of the catch of the southern trawl fishery in the following winters.

Observation on the summer fishery during 1934 was not possible, through lack of funds, so that nothing definite is known as to what age groups are supporting that fishery at the present time and as

to the degree of success of reproduction in 1933 and 1934. Analysis of catch-record data supplied voluntarily by pound-net operators, however, indicates a high yield of scup in 1934 continuing the series of record total catches this particular fishery has experienced each summer since 1929 (for the State of New Jersey the pound-net catch rising from 2,400,000 pounds in 1929 to 3,500,000 pounds in 1933), following a period of scarcity from 1926 to 1928, inclusive (500,000 pounds in 1926 to 300,000 pounds in 1928). The record yields of 1929-33 were the result of successful spawning each season from 1927 to 1932, with the exception of 1929.

Observations on the southern trawl fishery during the past winter (1933-34) showed that the catch of approximately 2,500,000 pounds of scup landed principally at Virginia ports was the highest in the history of the fishery. As in most previous winters, the size and age composition of the landed catches was similar to that usually observed in the summer pound-net fishery, consisting, in the main, of small and medium-sized fish (7 to 9 inches in length, averaging one-fourth to one-half pound in weight).

It was disclosed further, in accordance with previous winters' observations, that the catch of this southern winter trawl fishery is not only dependent on abundance but also on the degree of availability, which in turn is affected by changes in the hydrographic conditions of the general fishing region. It is now known that changes in water temperatures affect the movement of the schools, causing changes in the amount and in the size of fish caught.

Thus, as indicated above, the total yield of scup by the summer and winter fisheries continues at record levels due to a series of successful spawnings. This condition cannot be expected to continue indefinitely. Sooner or later a series of spawning failures similar to those which caused the natural decline between 1926-28 will occur.

It is but prudent, therefore, to prepare for this condition by correcting existing abuses which may accelerate the inevitable decline or endanger the chances of natural recovery. The most conspicuous abuse is unnecessary destruction of young fish both in the winter trawl fishery and in the summer pound-net fishery. During the past winter the southern trawl fishery discarded at sea large quantities of small scup consisting in part of fish below legal limit and in part of small but legally marketable fish for which, under the economic conditions of last winter, no market could be found.

It is estimated that for a total catch of approximately 2,400,000 pounds of scup landed by vessels at Virginia ports in January, February, and March, more than 500,000 pounds or 20 percent were discarded at sea. Expressed in numbers of fish it means that of a total catch of approximately 4,700,000 scup, about 1,900,000 fish or 40 percent were destroyed and discarded. It is probable that an even larger proportion of the catch of vessels landing at ports in New York and New Jersey was discarded since the smaller sizes of fish are less readily marketable there.

Less definite figures are available for the pound-net fishery, particularly in recent years, but the waste is known to be great in the summers following the advent of large broods.

The advantages to be derived from reduction of this waste are as follows: (1) Since the fish now destroyed have no value, a clear

gain will be realized from those allowed to escape and grow to marketable size, even though reproduction continues to be successful for several years; and (2) when reproduction fails during a series of years (as experience indicates is inevitable), young fish spared will be doubly valuable, for they will increase the spawning reserve necessary to insure eventual natural recovery and they will augment the stock available to the fishery during the period of scarcity.

It is desirable, therefore, that experiments be conducted to determine the practicability of extending to the Middle Atlantic pound-net fishery the practice of sifting out small scup now generally used by operators of floating traps in Rhode Island. It is also desirable to reduce the waste of young fish in the winter trawl fishery by introducing changes in the mesh of the cod-ends of the trawls. Considerable experiment will be necessary to determine the correct sizes, for the problem is complicated by the fact that more than one species must be considered (in addition to scup, sea bass and fluke demand attention). Lack of funds has prevented experiments thus far, but it is urgent that they be undertaken at the earliest possible date.

Other species.—A number of important species have perforce been neglected during the course of Middle Atlantic investigations. Among them are sea bass, bluefish, and shad. Since bluefish and sea bass are taken to some extent by the summer pound-net fishery and sea bass constitutes an important part of the catches of the winter trawl fishery, advantage was taken of the opportunities to collect many data bearing on them at times when field observations were being made of the catches of squeteague, scup, and butterfish by these fisheries. Because of insufficient personnel it has been impossible to analyze the data collected and impossible to arrange for observations of the hand-line fisheries for bluefish and sea bass, and the lobster-pot fishery for sea bass. It is particularly urgent that investigations of sea bass be undertaken, for this species has been subjected to a greatly increased fishery strain through the rise of the lobster-pot fishery and the winter trawl fishery in recent years.

The shad fishery in the Delaware River has practically disappeared. In 1897 the yield of this fishery was 16,098,552 pounds, valued at \$436,546. In 1932 the yield had shrunk to 109,979 pounds, valued at \$15,469. Even if we assume that the yield in 1897 was excessive and that under proper management the fishery is incapable of a constant yield of more than half of the quantities caught in 1897, it is apparent that the destruction of this fishery is causing an annual loss of more than \$200,000 to the communities concerned and is depriving the consuming public of approximately 8,000,000 pounds of very desirable food. It is probable that pollution and obstruction of streams is in large part responsible, for in spite of very intensive fishing the yield in the less highly industrialized Chesapeake States has not diminished so severely. It is urgent, therefore, that rehabilitation be undertaken in accordance with the authorization provided by Public Law No. 121, Seventy-third Congress, March 10, 1934.

It is also desirable to seek means of restoration of the shad fishery in other localities where the destruction is not yet complete. The present yield in the Chesapeake States and in the Hudson River is

greatly reduced as compared with earlier years. Biological studies of shad life history, statistical study of fishing intensity, and pollution studies will be necessary to indicate the best procedure for rehabilitation and management of these fisheries.

FISHERY INVESTIGATIONS OF THE SOUTH ATLANTIC AND GULF COASTS

INVESTIGATION OF THE SPAWNING HABITS, LARVAL DEVELOPMENT, AND RATE OF GROWTH OF FISHES

The study of collections of young fish and field data collected principally on the coast of North Carolina was continued intermittently, as other duties permitted, by Dr. Samuel F. Hildebrand and Louella E. Cable. Additional specimens were obtained from Beaufort, N. C., and vicinity from towings made by Dr. James S. Gutsell in his study of the life history of shrimps. Several nearly complete and some partial series showing stages in the development were identified. Drawings, and in some instances descriptions, have been prepared.

SURVEY OF THE FRESH WATERS OF MISSISSIPPI

A general survey of the fresh waters of Mississippi was begun by Dr. Hildebrand in 1933. Although it was not found practicable to continue the field studies in 1934, considerable time was devoted to a study of the collections obtained the previous year. A special study of the minnows of the family Cyprinidae was made. Thirty species were recognized in the collection, and descriptions with notes on their life histories and habits have been prepared. Nearly all of these minnows are of importance because they constitute the chief food for many food and game fishes.

MARINE FISHES OF THE GULF COAST

Systematic studies of the fishes of the Gulf coast were continued by Isaac Ginsburg. Special attention was given to the families *Pleuronectidae*, *Gobiidae*, *Cyprinodontidae*, and *Syngnathidae*. In connection with this work the revision of a number of genera of American gobie is being prepared, which should be of help in placing the systematics of the *Gobiidae*, one of the most difficult families of American marine fishes, on a firm scientific foundation.

AN INVESTIGATION OF THE FISHES AND FISH CULTURAL POSSIBILITIES OF THE FRESH WATERS OF PUERTO RICO

An investigation of the status of the fishes of the fresh waters of Puerto Rico and the fish cultural possibilities existing there was undertaken by Dr. Samuel F. Hildebrand in cooperation with the Insular Department of Agriculture and Commerce. The principal fresh waters of Puerto Rico consist of 5 main streams with tributaries and 6 permanent reservoirs. Fishes in general are scarce in these waters. The indigenous fresh-water fishes consist almost wholly of representatives of salt-water families that in the course of time have

acquired a fresh-water habitat. These fishes disappear almost completely from the streams above an elevation of from 1,500 to 2,000 feet.

Food for fishes, consisting principally of crustaceans and insects, is abundant almost everywhere. The temperature of the waters at an elevation of 2,000 feet and higher is cool enough for rainbow trout, as it rarely exceeds 70° F. About 10,000 rainbow trout fry were hatched in the Loquillo National Forest from eggs received from the Division of Fish Culture through the United States Forest Service. Some 9,000 of these fry were liberated in streams within the forest while the remaining 1,000 were retained and fed principally on shrimps from the local streams. Whether the ones that were liberated have survived is not definitely known, but the ones retained and fed in captivity made fairly rapid growth.

Some bluegill sunfish (*Lepomis incisor*) and catfish (*Ameiurus melas*), presumably acquired through the New York Academy of Sciences, were introduced in two reservoirs in about 1913. These fishes have propagated and still are present in the two reservoirs in which they were introduced. A third species, one of the crappies (*Pomoxis*) also was introduced, and is said to have maintained itself, though it was not seen during the investigation. The introduction of temperate-zone fishes seems to have been so successful that more trout eggs and small bluegill sunfish (*Lepomis incisor*) and catfish (*Ameiurus nebulosus*) were sent during October and December 1934. The prospects that these American fishes will establish themselves are favorable, and it is hoped thereby to furnish sport and food for the people of Puerto Rico.

SHRIMP INVESTIGATIONS

The shrimp investigations have continued under the direction of Milton J. Lindner. As in the past, headquarters have been maintained in New Orleans, La., in offices furnished by the Louisiana Department of Conservation. Field stations have been located at the United States Fisheries Biological Station, Beaufort, N. C.; the Georgia Tidewater Commission, Brunswick, Ga.; and the San Patricio Canning Co., Aransas Pass, Tex. The Louisiana Department of Conservation; the Texas Game, Fish, and Oyster Commission; the Georgia Department of Game and Fish; and the San Patricio Canning Co. have continued their generous cooperation as in the past. Due to lack of funds, operation of the *Black Mallard* was interrupted on August 1, 1934. Operations of this vessel will be resumed for a brief period beginning January 1935.

In conjunction with the work of the Bureau the Louisiana Department of Conservation has been conducting a hydrographical survey of Louisiana waters under the direction of James Nelson Gowanloch.

The shrimp is the most valuable fishery resource of the South Atlantic and Gulf coasts. In 1932, the latest year for which statistics are available, the shrimp fishery, with a catch of over 88 million pounds and a value to the fishermen of over 2 million dollars, ranked sixth among those of the United States and Alaska. There are three species of shrimp that constitute the commercial catch, the common shrimp, *Penaeus setiferus*; the grooved shrimp, *P. brasiliensis*; and

the sea-bob, *Xiphopenaeus kroyeri*. The common shrimp is by far the most important because it comprises at least 95 percent of the commercial catch.

From a commercial standpoint, the shrimp fishery is in better condition than it has been in the last few years. Indications are that the pack of canned shrimp during the 1934-35 season will be larger than in the past several years with the market price holding up well. A closer organization of the canners and more cooperation between them was noted during 1934. The inauguration of the inspection system under the Pure Food and Drug Administration and the work of the shrimp section of the National Canners Association have resulted in a much improved pack of shrimp. The increase in the pack probably will cause a similar augmentation in the total landings of shrimp.

Because of the importance of the common shrimp, the major efforts of the investigation have been directed toward solving the problems presented by this species. The length measurements and gonad examinations show that the common shrimp reach maturity and spawn within 1 year. Not many spent shrimp are found and no shrimp which can be interpreted as being 2 or more years old has been taken. It is assumed from this that the shrimp die after spawning and that very few, if any, survive to spawn the following year. The spawning of the common shrimp occurs throughout the spring and summer in the open sea or Gulf, and the shrimp after hatching and passing through the larval stages move into the inside waters. These warm shallow sounds, bays, bayous, and rivers serve as nursery grounds. The young shrimp grow very rapidly and by June or July, depending upon the locality, attain sufficient size to appear in the commercial catch. By September almost the entire catch is composed of young shrimp from the spawning of the preceding spring and early summer. With the cooling of the waters in the early fall and winter there is a definite disappearance of the larger of the young shrimp from the inside waters and in some sections of the coast from the customary outside fishing grounds which, with few exceptions, are usually near the mouths of the passes.

At Beaufort, N. C., Dr. J. S. Gutsell has continued his studies of the structure and development of the ovarian eggs of the commercial species of shrimp. This study is of importance in determining just when spawning occurs and if a shrimp spawns more than once. Sections of many ovaries, including a number from the common shrimp with spermatophores attached, have been microscopically examined.

Because so few of the common shrimp are taken with spermatophores attached and because of the method of attachment, it seems certain that the spermatophores are not present on the female until they are ready or very nearly ready to spawn. Therefore, the large eggs found in the enlarged ovaries of all shrimp of this species with spermatophores attached are believed to be not only fully developed but also in a mature condition ready for emission and fertilization. Examination of such mature eggs reveals certain definite and distinctive structures in the yolk. These are not artifacts for under favorable conditions they may be seen in fresh eggs immediately after removal from newly killed shrimp. By proper fixation and staining they may be preserved and studied in sections which afford a superior picture of conditions within the ovary. One type of cytoplasmic body

does not appear until the eggs are fully grown. By means of it a substantially matured egg may be positively identified. This is of importance, for if examination of an ovary shows that this structure is present in the crop of large eggs it may be stated without any doubt that the shrimp is ready to spawn; thus, with proper sampling, the spawning season can be determined with precision.

In addition, Dr. Gutsell has examined a number of shrimp which appear to have recently spawned. From these studies he has secured ovarian characteristics which apparently are diagnostic of spent shrimp. This is a particularly important and valuable contribution. Dr. Gutsell is preparing a detailed report of these investigations which he expects to have completed before the end of the 1935 spawning season. The completion of the report is expected to terminate the study of this particular problem.

Mr. Anderson, during 1934, continued the sampling of the shrimp population along the South Atlantic coast aboard *Launch 58*. The territory sampled represents about 95 percent of the shrimp fishery on the Atlantic coast. For the purposes of this investigation, 9 outside and 2 inside stations were visited once each month and from 2 to 4 trawl hauls were taken at each station. The outside stations are from 35 to 50 miles apart and extend from Cape Romain, S. C., to Cape Canaveral, Fla. The stations were so chosen as to give a representative sample of the shrimp population at the more important fishing centers along this coast. At each station otter trawl hauls and plankton tows were made, surface and bottom temperatures recorded, and salinities determined. A representative sample of the shrimp obtained was measured, the development of gonads noted, and the numbers of each species of fish secured were recorded.

Confirmation of last year's data was obtained which indicated that the large shrimp disappear from the usual fishing grounds during the fall and winter along the Georgia and northern Florida coast. Similarly, confirmation was also secured for last year's data which indicated a progressive increase in size of the common shrimp from north to south along the Florida coast during the fall and winter. Four possible interpretations may be placed upon these phenomena:

1. There is a southward movement of the larger shrimp along the Florida coast during the fall and winter.
2. The larger Georgia and northern Florida shrimp move offshore during this period.
3. Due to heavy fishing in the summer and early fall, the larger shrimp are depleted along the Georgia and Florida coasts.
4. Combination of any two or all three of the above factors may be occurring.

As a possible means of interpreting these changes in the population of the shrimp on the South Atlantic coast, Mr. Lindner has been conducting racial studies at frequent intervals in four widely separated localities. Although these studies have not been completed, it has been found that there are differences in body proportions in the shrimp in different localities. A complicating factor arises in that these differences appear to be associated not only with locality but also with season of the year and age of the shrimp. An important discovery resulting from this study has furnished a means of distinguishing between 0 group and I group shrimp during the late summer and early fall.

Kenneth H. Mosher continued sampling the commercial catch of shrimp at Aransas Pass through the calendar year 1934. In 1933 the sampling was extended to include the commercial catch at four other Texas ports—Galveston, Palacios, Port Lavaca, and Port Isabel. The shrimp fishing at Palacios, Port Lavaca, and Port Isabel was so irregular that these three ports were disregarded in 1934, leaving only Aransas Pass and Galveston as regular sampling localities. During 1934, 20,516 shrimp were sexed, measured, and sexual maturity noted. Only 145 of the total measured were *P. brasiliensis* and only 71 were *Xiphopenaeus kroyeri*, or the two species taken together were only slightly over 1 percent of the total yearly sample. The remaining 20,300 shrimp were *P. setiferus*. In addition to the routine sampling, a number of the shrimp were weighed as well as measured in order to obtain the weight-length relationship.

After many unsuccessful attempts to keep shrimp alive for any length of time in various types of live boxes and aquaria, two pens were devised and placed in a small pond separated from Redfish Bay by a dike in which screened sluiceways were built to provide a current of water to the pond.

Early in December Mr. Lindner and Mr. Mosher initiated tagging experiments upon some shrimp placed in the pens. The celluloid disk or button tag was used. Although inclement weather caused the death of the shrimp a few weeks after tagging, the results were encouraging as it was evident that the shrimp were able to withstand the rigors of handling while being tagged. Mr. Mosher will continue these experiments as soon as the danger of freezing has passed.

Studies on the spawning and early growth of the commercial penaeid shrimps have been continued by John C. Pearson. A unique discovery during the year was the recognition of the eggs of a penaeid shrimp, probably *Penaeus setiferus*. These eggs, taken off the coast of Georgia on June 17, 1932, in surface plankton, appear to be the first penaeid shrimp eggs to be secured in a pelagic state. The eggs were spherical in shape and possessed an extremely thin delicate membrane. The diameter of the eggs ranged from 0.38 to 0.42 millimeter. Within each otherwise transparent egg rested a well-developed nauplius, the length of which ranged from 0.21 to 0.26 millimeter. The capture of these eggs at the surface in water of 30-foot depth would indicate that the eggs were spawned relatively close inshore and that they are strictly pelagic, floating at the surface of the sea.

Nauplii, zoeae, and mysis stages of the commercial penaeid shrimps have been abundantly secured in many plankton collections off the coasts of Louisiana, South Carolina, Georgia, and Florida. Zoeae and mysis stages of both the commercial shrimps, *Penaeus setiferus* and *P. brasiliensis*, are at hand and are being studied in detail with respect to morphological distinctions and oceanic distribution as to time and place. It is hoped that a report on the morphology and distribution of the young of these commercial shrimps can be completed during the coming year. Considerable material on the young stages of various noncommercial shrimps has also been collected with the studies of the commercial shrimps.

PACIFIC COAST AND ALASKA FISHERY INVESTIGATIONS

The major salmon and herring investigations have been continued through 1934 with some restrictions and limitations as to territory covered. During the year three new investigations were established; namely, the Columbia River salmon investigation, the Washington coho salmon investigation, and a detailed statistical study of the catches of all species of salmon from 1897 to 1933 in Puget Sound, the Fraser River, and Swiftsure Bank. The most severe limitation of activity has occurred in Alaska where the investigations have been confined mainly to the species of most economic importance; namely, the red salmon and the pink salmon.

COLUMBIA RIVER SALMON INVESTIGATION

In July 1934, the Division of Scientific Inquiry was allotted the sum of \$25,000 annually during a 10-year period for scientific studies and experiments relative to the salmon fisheries of the Columbia River. J. A. Craig, formerly engaged in a study of these fisheries, was placed in charge of the investigation. Mr. Craig is being assisted by A. J. Suomela, who has also had experience on the Columbia River.

The object of this investigation is to provide information and recommendations which will point the way to the conservation of these important fisheries and provide for their productivity in the future. With this purpose in mind, a comprehensive program for the investigation has been prepared.

The development of the lands in the Columbia River Basin has been attended with many activities, such as the construction of dams and irrigation diversions and the introduction of sources of pollution, which, coupled with the intensive commercial fishery, have placed a great strain on the salmon populations of this river. In order to obtain some measure of the effect of this strain on the abundance of these populations it is necessary to conduct a detailed analysis, on the basis of catch return per unit of fishing time and effort, of the catch statistics of the salmon fisheries. It is known that the "run" of blueback and chinook salmon into the Columbia River is composed of a great number of races, each spawning in different tributaries and having fairly definite times for entering the commercial catch. The statistical study should yield information regarding the status of these separate races, as well as that of the composite runs of these two species. The statistical analysis, therefore, becomes one of the most fundamental and important parts of the present program of investigation.

In order to secure exact information on (1) the location and extent of the spawning areas now available to the salmon, (2) the location and extent of the spawning areas which have been destroyed and the causes of their destruction, and (3) the general status of the salmon populations spawning in the Columbia River at the present time, a program of stream surveys has been undertaken. Field parties have been sent out to cover the entire Columbia River system as rapidly as possible in order to determine the above-mentioned facts. It is hoped to discover the present condition of these spawning areas and to bring out facts which will lead to their improvement and rehabilita-

tion. The results of these surveys will also be a means of indicating which of the tributaries are suitable for transplanting or restocking projects.

The program for the investigation also calls for studies of the life history, habits, and homing instincts of the several species of salmon. All of these facts must be known in order to conserve the fishery properly.

During the period from July 1934 to January 1935, inclusive, a large number of catch records have been collected and their analysis carried forward with the aim of preparing a report on the indices of abundance of the chinook and sockeye salmon from 1905 to the present time.

The stream-survey parties, although handicapped by adverse weather conditions during a great part of the time while in the field, have covered parts of the Wenatchee, Entiat, Methow, and Okanogan River systems as well as Wenatchee, Silver, and Osoyoos Lakes.

PUGET SOUND SOCKEYE SALMON INVESTIGATION

The activities of this investigation in 1934 were continued on the basis of the previous year's program. Thousands of sockeye salmon fingerlings from the Birdsvew hatchery were marked by the removal of two fins and liberated in each of three creeks tributary to the Skagit River. The object of this experiment was to determine something concerning the mechanism of the homing instinct in the sockeye salmon. If these fish are attracted to a stream by some property of the water in which they were reared they should enter that stream no matter whether they have traversed it before or not. On the other hand, if they return by some sense of geographical location they should return to the stream in which they were liberated even though they left it immediately.

With this theory in mind, 10,000 marked sockeyes were liberated in Bacon Creek, 10,000 in Day Creek, and 10,000 more in Diobsud Creek. All of these streams are tributary to the Skagit. Two enter this stream above Grandy Creek, where the fish were reared, and one below Grandy Creek. Different marks were used for each group of fish, and these fish were liberated at their natural time of migration. If it becomes evident that sockeyes will not return to a stream in which they were liberated at their time of migration and which they immediately left, additional experiments will be performed to find for what length of time it is necessary to hold fish in a stream in order that they become imbued with the urge to return to that stream.

Approximately 14,000 sockeye fingerlings were marked and liberated at the Quilcene hatchery. These fish were transferred from the Birdsvew hatchery in the eyed egg stage. The marking was for the purpose of determining whether or not sockeyes will return to Quilcene, which is not a sockeye stream, after being reared there. The returning adults will be detected in the commercial catch in Puget Sound and so provide a check on their survival if none appear at Quilcene.

Scale samples from the commercial catch at Bellingham, Wash., were collected, and marked fish were secured in cooperation with the British Columbia investigators during the past fishing season.

Statistics of the sockeye salmon fishery were collected and tabulated during the first half of the year. In July J. A. Craig, who was in charge of this problem, was transferred to the Columbia River investigation and Dr. G. A. Rounsefell was placed in charge. Dr. Rounsefell, in collaboration with G. B. Kelez, was engaged in preparing a preliminary report on the statistics of the salmon fisheries of Puget Sound during the last half of the year.

WASHINGTON COHO SALMON INVESTIGATION

In January 1934 a new investigation on the coho salmon of Puget Sound and the coastal district of Washington was started under the direction of George B. Kelez. The purpose of this investigation was the determination of the life history and state of abundance of this species, which not only contributes materially to the commercial fisheries of this region, but also forms a large part of the catches of the recreational salt-water fishing.

A survey was made of the location and extent of the fisheries for this species and the location and size of the streams in which they spawn. During the early spring, sampling was carried on for yearling fingerlings resident in the streams, and the presence of fry hatched during the year was noted. Collections of scales from adult fish were made at Bellingham, Wash., at regular intervals during the months of June to September (through the cooperation of the Puget Sound sockeye investigation). Collections of scales from both immature and mature fish in sport-fishing catches and from adult fish in the commercial catches landed in Seattle from Cape Flattery and Puget Sound were also made. Through the cooperation of several fishing resort owners, records were obtained as to the abundance of various sizes of cohos in inside waters during the past season.

A series of marking experiments was undertaken to secure information on life history, homing instinct, and migrations of the species; the fingerlings being supplied through the courtesy of the Washington State Department of Fisheries. Twenty-six thousand fry hatched at the Samish station in February were marked during May by the removal of the adipose and dorsal fins and liberated in Friday Creek, a tributary to the Samish River. At the same time 10,000 fry from the Skykomish station (which is located on an independent stream system) were moved to the Samish station and marked immediately by the removal of the adipose and left ventral fins. These fish were also liberated in Friday Creek in an attempt to determine the effect of transplanting the fry on their choice of the stream in which they will spawn. A portion of the Samish fry, from which those marked in May were taken, were held in the Samish ponds until November, and an additional lot of 26,000 were marked at that time by the removal of the dorsal and left ventral fins. The comparative returns from the two Samish experiments are expected to demonstrate whether or not the additional period of

pond rearing will contribute materially to the resistance of the fingerlings to disease and natural enemies. Because of the vigorous growth and low mortality exhibited by these fish during the period of pond rearing, the State department of fisheries has determined to continue this procedure next summer wherever practical at the rest of their stations.

STATISTICAL STUDY OF SALMON FISHERIES IN PUGET SOUND REGION

During the latter half of 1934, Dr. Rounsefell and Mr. Kelez, in charge of the red salmon and coho investigations on Puget Sound, collected a great mass of statistical data of the catches of salmon of all species by all types of gear in the Puget Sound region as well as from Swiftsure Bank and the Fraser River. This included daily lifts of salmon traps from 1897 to date, of daily seine boat deliveries both from Puget Sound and from Swiftsure Bank from 1912 to date, and individual gill-net catches of over half of all of the gill nets operating on the Fraser River from 1929 to date. Several important features of the fishery were so well covered by these data that a preliminary report was made entitled, "Abundance, seasonal occurrence, and development of the salmon fisheries of Swiftsure Bank, Puget Sound, and the Fraser River."

This report gives a brief summary of the history and development of each type of fishing gear, where it has been fished, and the numbers used. The earliest fishery was chiefly with gill nets in the Fraser. Later came traps and small inefficient purse seines in Puget Sound. The purse seines increased steadily in efficiency after the introduction of gasoline motors in 1903 until the present modern Diesel-powered boat was reached. The amount of fishing gear reached its peak during the World War, descended to a very low level from 1922 to 1924, and then increased slowly but steadily to date. The ocean fishing on Swiftsure Bank with purse seines began in 1911 and quickly reached large proportions.

Studies of the seasonal occurrence of each species of salmon showed that in the traps in Puget Sound the king salmon run very early, 50 percent being taken before the others have really commenced to appear. The red salmon run next, followed a month later by the pinks. The cohos and chums run later than the rest. In the purse seines in Puget Sound all species are taken somewhat later than in the traps. This is especially true of the chums. On Swiftsure Bank most of the species are taken at the same time as in Puget Sound, with the exception of the cohos. Large schools of feeding cohos in their third year that are only half as large as they would be if caught later in the season are purse seined early in the season. The number of cohos on the bank fall off as the catches of larger cohos increase inside of Puget Sound. In the Fraser River the season is somewhat more extended than in Puget Sound and the fish appear to run later.

A detailed study of abundance shows that the red salmon, although not as abundant in the "big" year cycle as formerly, appear on the whole to be increasing gradually since the low point reached during the period 1920 to 1925 resulting from the combination of overfishing during the World War, and the blockade of most of the spawning grounds in 1913 and 1914 by the slide in Hell's Gate Canyon.

The coho salmon are less abundant at present than at any time since the beginning of the fishery. It appears that they are in need of more stringent protective regulations.

The pink salmon, although maintaining a fair level of abundance since 1915 are in no wise as abundant as in 1911 and 1913, before the slide of 1913 in Hell's Gate Canyon completely wiped out the population spawning in the Thompson and Nicola Rivers and the tributaries of Seton and Anderson Lakes.

KARLUK RIVER RED SALMON INVESTIGATION

The Bureau of Fisheries early realized the necessity of obtaining information on the number of red salmon that should be permitted to spawn in the streams in order to produce returns that would provide the maximum surplus for the fishery. Since this involved a study of the complete returns from known spawning escapements, a stream had to be selected where both the portion of the run taken for commercial purposes and the portion escaping to the spawning grounds could be determined. The Karluk River on Kodiak Island was the first stream selected for this study since it supports a large red salmon population which is subject to commercial capture only in the vicinity of its mouth. Furthermore, the Karluk River is centrally located in a large red-salmon-producing area, and it was hoped that the fundamental facts secured from the study of its fishery could be applied generally to the fisheries of the other red-salmon-producing areas in Alaska. This investigation has been carried on continuously each year since its inception in 1921, and at the present time is being conducted by J. T. Barnaby, assisted by L. D. Townsend.

The information available to date on the returns from known spawning escapements in the Karluk River indicates that wide fluctuations may be expected in the rate of reproduction of its spawning populations. During the past year a complete analysis was made of the returns from the known escapements of 1921 to 1928. The escapements for these years are as follows: 1921, 1,500,000; 1922, 400,000; 1923, 694,579; 1924, 1,000,000; 1925, 1,620,927; 1926, 2,533,402; 1927, 872,538; and 1928, 1,093,817. The ratio of return to escapement for these years is as follows: 3.0 to 1, 5.6 to 1, 2.9 to 1, 0.8 to 1, 1.0 to 1, 0.6 to 1, 1.8 to 1, and 2.1 to 1, respectively. The smallest escapement, 400,000 fish, produced the greatest return per spawning fish, 5.6 to 1, whereas the largest escapement, 2,533,402 fish, produced the smallest return per fish, 0.6 to 1. In other words, there seems to be little opportunity of selecting a spawning escapement that will consistently produce a large population.

In 1926 a series of marking experiments were established for the purpose of determining the mortality of the red salmon during their life in the sea. These experiments have consisted of the marking of 50,000 seaward migrants each spring. The proportion of the adult salmon bearing the marks that have returned from each experiment indicates that although the mortality in the sea is considerable it seems to remain fairly constant from year to year. However, the rate of growth of the salmon in the ocean varies considerably and is responsible for the length of time the fish from different

brood years remain in the ocean before returning to spawn. Obviously, salmon staying an extra year in the ocean before returning to spawn are subject to a greater mortality since they are exposed to the factors which affect their mortality for an additional year's time. Hence the wide fluctuations in the rate of reproduction of the spawning populations may be due in part to fluctuations in their ocean mortality.

Evidence was secured during the past season which indicates that changes in the natural conditions in the streams and lakes during the spawning period and early development of the salmon can bring about wide fluctuations in their mortality. In fact, the relatively poor returns from the escapements of 1924 to 1928, inclusive, are probably due to the abnormal conditions which were observed on the spawning grounds in 1924 and 1926. Hence it is believed that the varying rate of reproduction of the spawning populations may be traced largely to the success or failure of the spawning salmon and the death rate of the ensuing young during their existence in fresh water. Therefore, the program for this investigation in the future will include an intensive study and evaluation of the factors which influence the success of each year's spawning population and the mortality of the brood from the egg stage to the time it migrates to the sea.

CHIGNIK RIVER RED SALMON INVESTIGATION

The activities of this investigation during the past year were greatly reduced owing to the transfer of its personnel, Messrs. H. B. Holmes and G. B. Kelez, to other Bureau investigations.

A temporary assistant was stationed at Chignik from May until October for the purpose of collecting fish-scale samples and the observation of the commercial catches for marked fish. The data collected have been filed for future use.

BRISTOL BAY RED SALMON INVESTIGATION

Although funds were not available for a biologist to carry on field work in Bristol Bay during the past year, scales of the 1934 red salmon populations in this area were collected through the cooperation of the Alaska Division of the Bureau. Scale samples and body measurements of the red salmon which spawn in the Bristol Bay region have been accumulating for a number of years. Funds have been requested for the development of this important field of work, and it is hoped that they will become available in the near future.

PINK SALMON INVESTIGATION

The pink salmon are found in practically all the waters of Alaska but are most abundant in the southeastern section. Millions of pink salmon spawn in this section each season in its hundreds of streams which vary greatly both in size and character of habitat. The Bureau of Fisheries through its conservation policy, which was made possible by the White law, has been able to build up the pink-salmon populations of southeastern Alaska to a state where they are fluctuating near their maximum level of abundance. In fact, the commer-

cial catch of pink salmon in this section during the past season was the greatest in the history of the fishery. This, however, does not imply that these populations of salmon are no longer in need of constant observation for past experience has shown that they are subject to serious injury due to changes in both economic and natural conditions regardless of their state of abundance. Hence to secure information in regard to the changing status of these salmon populations which will lead to their conservation so as to provide for a permanent and productive fishery is the primary aim of the pink-salmon investigation.

The activities of this investigation in 1934, as in past years, were confined to a study of the pink-salmon populations in southeastern Alaska. Owing to the varied environmental conditions under which the pink salmon reproduce in this section of the Territory, it is believed that the fundamental facts secured from the study of its fishery may be applied generally to pink-salmon fisheries of Alaska. Dr. Frederick A. Davidson is in charge of the investigation, assisted by S. J. Hutchinson.

A paper on the homing instinct and age of maturity of the pink salmon was published during the past year. The evidence to date indicates that the pink salmon have a great desire to return to spawn in the streams from whence they came, but that some straying of the adults may occur in localities where a number of streams are in close proximity to each other. The evidence thus far collected also indicates that the pink salmon have a complete 2-year life cycle, that is, they invariably return to spawn at the close of their second year of life. The results from the racial analysis of the pink salmon which have been completed to date, substantiate the evidence in regard to the age of maturity of the pink salmon. These results indicate that 2 distinct salmon populations are found in the streams, 1 that spawns in the odd years and 1 that spawns in the even years.

A continuous study is being made of the yearly fluctuations in the abundance of the pink salmon in southeastern Alaska beginning with the inception of the industry in 1895. The results from this study show that wide fluctuations may be expected in the yearly abundance of these salmon. Since the pink salmon have a complete 2-year life cycle, the number that return to spawn any year is the result of a single spawning brood and not of several broods as in the case of the red salmon. Hence a marked change in the mortality of any brood is directly reflected in the number of salmon that return 2 years later. Knowing the disastrous effects that lack of rainfall during the spawning period can have upon the success of a spawning population, it is not surprising to find these wide fluctuations in the yearly abundance of the pink salmon. The amount and time of occurrence of rainfall, however, are not the only factors which influence the mortality of the spawning salmon and subsequent brood. Hence a study is being made for the purpose of determining and evaluating all of the factors which influence the mortality of the salmon during their entire cycle of reproduction.

This study was started in the stream at Olive Cove, Alaska, but owing to handicaps encountered there it was transferred to the stream at Little Port Walter in the summer of 1934. The stream at Little Port Walter provides a very satisfactory location for this study.

It is only three-fourths mile in length, and the varied character of its stream bed provides for the study of practically all the types of spawning grounds found in the streams in southeastern Alaska. Furthermore, all of the commercial fishing which affects its salmon population is carried on just outside of the port and can readily be accounted for. Plans are being made for the construction of a permanent counting weir in the stream wherein both the adults that migrate into it to spawn and the resulting fry which migrate to the sea may be counted accurately. In this way both the mortality of the brood while in the streams and in the sea may be determined. It is hoped that funds will be available for a permanent observer at the stream for the purpose of making continuous observations of the environmental changes in the stream so as to determine the natural factors which influence the mortality of the brood. Once equipped with knowledge of the factors which influence the mortality of the broods, the Bureau will be able to develop further its conservation policy so as to provide for the most efficient utilization of the pink-salmon populations.

HERRING INVESTIGATION

With the exception of a few minor localities, the herring fisheries of Alaska may be grouped into four districts; namely, southeastern Alaska, Prince William Sound, Kodiak-Cook Inlet, and the Aleutian Islands. The fishery of southeastern Alaska is by far the most productive, and at the present time is supporting the majority of the fishing effort. Hence the activities of the herring investigation during the past year were confined mainly to a study of the fishery in this region. Dr. G. A. Rounsefell, who has been conducting the investigation since 1925, was transferred to the Puget Sound sockeye-salmon investigation in July owing to the need of a trained investigator for this work. E. H. Dahlgren, who has assisted Dr. Rounsefell for a number of years, was placed in charge of the investigation.

The stock of Alaska herring, being separated into a series of local races, presents a more complex problem than would be the case if it were a homogeneous population. The delineation of these races and a measure of the intermingling which takes place between stocks is of utmost importance. Estimates of changes in abundance and causes for such changes are dependent on this separation. Observations on the size and age compositions of the catch have clearly demonstrated the existence of dominant year classes, but the determination of the exact causes for such varying degrees of success of different spawning is yet to be solved.

A racial report now in press, by Dr. Rounsefell and Mr. Dahlgren, entitled "Races of Herring, *Clupea pallasii*, in southeastern Alaska", presents the progress already attained in the problem of the segregation of populations. The results of an analysis of vertebral counts and of growth rates, the comparison of proportional representations of year classes, and the recovery of tagged herring are presented in this paper. The separation of the herring populations in the vicinity of Petersburg and Noyes Island, and the localities east of Clarence Strait and south of Sumner Strait, including Wrangell, was made evident by comparisons of the means of the vertebral counts. Size

frequencies of age classes show that the herring of Noyes Island, Douglas Island-Icy Strait area, Affleck Canal, and Peril Strait are much slower growing than those of other localities. Age distributions support the analysis of vertebrae and growth rates, in separating the populations of Noyes Island, Peril Strait, and the Douglas Island-Icy Strait areas from neighboring localities. The recovery at Cape Ommaney of tags affixed to spawners at Sitka demonstrated the relation between that spawning ground and the summer feeding area. The failure to recover any of the tags affixed at Cape Bendel in Frederick Sound gives negative evidence to show lack of intermingling of these fish at the feeding grounds in lower Cape Ommaney with the other populations.

Further experiments in the tracing of migrations by tagging were carried out during the spring of 1934. Twenty-one thousand five hundred and sixty-one metal belly tags were affixed to spawners at two of the major spawning areas, 8,394 at Craig on the outside of Prince of Wales Island, and 13,167 at Sitka, on Baranof Island. On 18,368 of these, a tag modified from those of previous years was used, the change consisting of having the tag stamped from steel, then plated with nickel, and in increasing the size to about double that of the nickel tags used previously. The larger size makes the tagging easier and faster. Corrosion, which might be expected to occur when there are imperfections in the plating, did not take place in the body cavity of the fish, and tags recovered 6 months after tagging showed their original luster. The 3,193 others were tagged with pure nickel tags left over from the 1933 experiment.

By means of the electromagnets installed at the reduction plants, 153, or 2.1 percent, of the steel tags affixed at Craig were recovered as against six-tenths of 1 percent of the nickel tags from the same experiment. Recoveries were made largely from the Warren and Noyes Island areas, although a small percentage were reported taken from Cape Ommaney. The Noyes Island and Warren Island fish are thus shown to be of the same population. Four hundred and eighty-eight, or 4.4 percent, of the Sitka steel tags and 40, or 1.8 percent, of the nickel tags at Sitka were recovered. They were taken, as were those of the 1933 experiment, from the vicinity of Cape Ommaney. A small fraction of these recoveries were made at the Warren-Noyes area. The degree of intermingling of the Cape Ommaney and Warren-Noyes Island populations, first indicated in this experiment, cannot be measured accurately until a device is perfected to extract the tagged individual before being run through the reduction plant. Such a unit has been devised and will be given a trial during the coming year.

Biweekly samples of the catch taken during the 1934 season in southeastern Alaska indicate a great predominance of 4-year-old individuals. As a result of the population being so young a relatively small pack of cured fish was prepared, a condition which may be expected to continue during the 1935 season. If this year class is not too seriously depleted, it should yield a good pack in 1936 due to the increased size of the herring.

Samples of the catch in Prince William Sound were collected, adding an additional year's data from this region. Sampling at Dutch

Harbor, carried on in previous years through the cooperation of the Alaska Division, was discontinued for lack of funds to analyze the data.

INVESTIGATION CONCERNING THE PROTECTION OF MIGRATORY FISH AT THE
BONNEVILLE DAM

The future of the salmon, sea-run trout, and other anadromous fish of the Columbia River is endangered by the Bonneville Dam, which is being constructed by the War Department as a part of the Public Works program. This dam will obstruct the main Columbia below the tributaries that furnish the bulk of the spawning areas. The run of fish involved supply a great part of the sport fishing of the inland region of Oregon, Washington, and Idaho, as well as support a commercial fishery whose annual yield is valued at several million dollars. The task of designing fishways for the Bonneville Dam, therefore, becomes the most important undertaking of its kind ever attempted.

When the United States Army engineers started preparing detailed plans for the structures, they called upon the Bureau of Fisheries for recommendations relating to fishways. Such recommendations could not be furnished without a detailed study of the situation. In fact, the magnitude and importance of the task demanded consideration of new features in fishways. A portion of the funds allotted to the Bonneville project, therefore, was assigned to the Bureau of Fisheries for its investigations.

Harlan B. Holmes was placed in charge of the investigation. Henry F. Blood, a prominent engineer of Portland, Oreg., was employed to assist with the engineering phases. Other temporary employees have assisted with various biological and engineering phases of the work. The investigation has been conducted in close contact with the designing staff of the United States Army engineers, a committee representing the fishing interests, and the State fish and game departments.

The investigation has involved (1) an analysis of commercial fishery catch records and information from other sources for the purpose of estimating the time and magnitude of the runs of fish; (2) a familiarity with the physical features of the dam and power-house structures; (3) a historical study of the seasonal variations in the flow of the river, and the effects of these fluctuations upon the operation of fishways; (4) an extensive study of fishway principles including a survey of fishways now in use or previously tried, principles that have been proposed but never put into practice, and the development of new features; and (5) finally upon the basis of these investigations the presentations of detailed recommendations for fishway structures.

The investigation was started in November 1933, and some phases are not as yet completed. Tentative conclusions have been made available to the designing staff of the United States Army Engineers as rapidly as they become available. Formal recommendations for fishways for upstream migrants were submitted to the War Department in August 1934. Details and other features pertaining mainly to the passage of downstream migrants and the passage of fish dur-

ing the period of construction are being considered as rapidly as possible.

The fishways for upstream migrants recommended by the Bureau of Fisheries include 3 fish ladders, 5 fish locks, and 3 units of a new feature known as a "collecting system." The fish ladders are of special design to adapt them for wide ranges of variation in tailrace and forebay fluctuation. The fish lock, which is new in application to salmon and trout, embodies the essential principles of a navigation lock with minor differences to adapt it to the special use. The collecting system consists of a series of entrance weirs distributed at various points along the obstruction, all entrances communicating with a common passage which leads to the base of the fishway proper. The quantity of water flowing through the common passage is augmented by an auxiliary water supply, thus presenting as an attraction to the fish a much larger quantity of water than can be supplied by the fishway proper. The collection systems are recommended for use both with the fish locks and the fish ladders.

GREAT LAKES FISHERY INVESTIGATIONS

Owing to the continued curtailment of the Budget, no field work was conducted on the Great Lakes during the calendar year 1934. The Great Lakes research conducted under Dr. John Van Oosten was confined entirely to work in the laboratories furnished by the University of Michigan at Ann Arbor.

As in past years, the Bureau continued its cordial relations with the various Great Lakes States and provided them, whenever possible, with information and advice concerning their respective fisheries. Dr. Van Oosten represented the Bureau at various fisheries conferences on the Great Lakes as well as at the public hearing called by the National Recovery Administration at Muskegon, Mich., on April 25 to discuss the proposed Great Lakes fisheries code.

Much progress has also been made in what is often considered the more scientific aspect of fisheries research. The analysis of fisheries statistics begun on Lake Huron has been extended to Lake Michigan and Lake Erie. Life history studies of the whitefish, cisco, perch, yellow and blue pike-perch, and sauger of Lake Erie are nearing completion. It is planned to extend these important investigations to the other lakes of the Great Lakes chain.

FISHERY STATISTICS

The intensive statistical study of the commercial fisheries of the Great Lakes waters of the State of Michigan begun in 1933 was continued through 1934. Complete and detailed data are now available for the Lake Huron fisheries over the 5-year period, 1929-33. Similar statistical analyses have been started for the commercial fisheries of the Michigan waters of Lake Erie and Lake Michigan. In a brief preliminary publication, Dr. Ralph Hile and William R. Duden described the methods employed in these statistical investigations.

The statistical data on the commercial fisheries of Lake Huron have proved particularly valuable in their application to the problems relative to the use of the deep trap net for the capture of white-

fish. This new type of gear, which was introduced into Lake Huron in 1928, proved so very effective and its use expanded so rapidly during the succeeding years as to create grave fears concerning the future of the fishery. The Bureau's statistical data based on a study of the Lake Huron fisheries by geographical districts and over a period of 5 years, showed these fears to be well grounded. According to the Bureau's data, the history of the deep trap net fishery in a single fishing district may be described approximately as follows: The introduction of the deep trap net into a fishing area is followed by an immediate rise in whitefish production to a point far above the normal. This increased production continues over a period of 2 years, although a drop in the catch per unit effort may be detected in the second year of the use of the trap net. The third year sees a marked drop both in total production and in catch per unit effort.

The statistical evidence supplied by the Bureau that the use of the deep trap net occasions a rapid and severe depletion of the whitefish stock was directly instrumental in the securing of legislation regulating and restricting the use of this destructive gear. The publication of a complete report on the statistics of the commercial fisheries of Lake Huron is being delayed pending the analysis of the 1934 material.

PIKE-PERCHES

The racial and life history study of the three species of pike-perches (*Stizostedion*) of Lake Erie with which H. J. Deason has been occupied has progressed toward completion. Scale samples from 8,390 fish (2,812 yellow pike-perch, 3,441 blue pike-perch, and 2,137 saugers) have been mounted, aged, and measured during the course of the study. Lengths to the end of each year of life and annual increments of growth have been calculated and summarized in tabular form. Some conclusions of this work were given previously (*Progress in Biological Inquiries*, 1933), in which only a portion of the above material was employed.

Because of the necessity of correcting certain fundamental errors inherent in the application of the scale method of growth determinations to pike-perches, it was found desirable to devote some time to the basic assumptions of the scale theory insofar as they applied to these species of fish. This study dealt largely with the variations involved in the proportionate growth of the body of the fish and its scales. As a result of this study it was possible to secure more accurate data on the rate of growth of the various species.

The racial study of the pike-perches is concerned with the systematic relationship between the blue and yellow pike-perches, principally those of Lake Erie. A total of 1,038 preserved specimens have been examined and measured, which number is made up as follows: 449 blue pike-perch from Lake Erie, 52 blue pike-perch from Lake Ontario, 230 yellow pike-perch from Lake Erie, and 307 yellow pike-perch from Saginaw Bay in Lake Huron. The Lake Ontario and Lake Huron materials were employed in order to study the variation in a single species from different bodies of water. At the beginning of this study some 30 counts and measurements were made on each fish, but as the work progressed and it became increasingly evident

that some of the counts and measurements were valueless those proving not trenchant were dropped. Only 7 measurements and 2 counts (lateral line scales and number of vertebrae) were retained as indicative of possible racial differences.

Small average differences between the blue and yellow pike-perches of Lake Erie have been noted, but the percentage of overlap is great. Differences between the blue pike-perch of Lake Erie and those of Lake Ontario are as great as are the differences between the yellow pike-perch of Lake Erie and those of Lake Huron. In distinguishing the blue from the yellow pike-perch in Lake Erie the color and growth rate afford the best criteria. In Lake Erie these two species have a different distribution, constitute distinct fisheries units, and because of differences in growth rate and average size have different legal size limits. From the standpoint of law enforcement it would be most advantageous to have absolute morphological criteria which would differentiate the two species, but these apparently do not exist. Further work on the racial question must include the age determination of all specimens and a study of morphological variations with growth rate and with year classes.

YELLOW PERCH

F. W. Jobs continued the study of the life history of the yellow perch. Up to the present time scales of nearly 5,000 perch have been analyzed for age and growth rate. The perch material was exceptionally well adapted to an investigation of the growth relationship between the body of the fish and its scales. As a result of this work a better conception will be obtained concerning this most difficult problem that confronts nearly every investigator of the life history of fishes.

OTHER OUTSTANDING PROBLEMS

Although some of the most controversial of the practical problems on the Great Lakes, such as the proper mesh in trap nets and pound nets, destructiveness of chub nets to trout, regulation of deep trap nets, method of measuring mesh in gill nets, size limits of fish, and pollution, have been solved, much remains to be done. The controversial question of the destructiveness of hooks and bait nets to immature fish needs investigation. Further experimental work should be done on the proper mesh of gill nets employed in Lake Erie. A satisfactory method for measuring the mesh of heavy twine must be developed.

UNIFORM REGULATION AND DEPLETION

One of the most urgent needs on the Great Lakes today is concerted action by the various Commonwealths, both in this country and in Canada, to regulate and protect the commercial fisheries of the Great Lakes. Since 1927 the Bureau has cooperated with the various Commonwealths in some 10 interstate and international fisheries conferences. During this period the Bureau, in cooperation with various State conservation departments, has concentrated its scientific research on the Great Lakes on those controversial practical

problems that previously had prevented enactment of uniform regulation. In spite of the vast amount of scientifically collected data available to support the recommendations adopted at the conferences, little progress has been made in the enactment of the recommendations into laws.

The necessity of uniform regulation is obvious from the fact that, in many cases, fishermen from 2 States or 2 countries depend on the same population of fish. Lack of uniformity not only causes much unfair competition between fishermen of adjoining States but also fails completely to give the fish the necessary protection. Twelve species of fish have been either exterminated or seriously threatened. The blackfin bloater and salmon of Lake Ontario have been completely exterminated. The sturgeon, blackfin, chub, and Lake Erie cisco no longer support a commercial fishery. The other six species of chubs are severely depleted, and the whitefish is seriously threatened with commercial extinction. The final result of continued lack of uniform regulation appears to be obviously indicated by what has happened in the past.

OYSTER INVESTIGATIONS

The investigation of various problems relating to oyster culture was carried out in all of the principal centers of the oyster industry of the Atlantic and Pacific coastal States and the Gulf of Mexico. As compared with the previous year, the work was expanded by an experimental study of the oil-pollution problem in Louisiana waters in cooperation with the State Department of Conservation. Preliminary surveys made in 1933 clearly indicated the necessity of conducting a comprehensive investigation of this important problem.

The United States fisheries laboratories at Woods Hole, Mass., Beaufort, N. C., and Washington, D. C., served for laboratory investigations. The Bureau's field stations at Milford, Conn., and Olympia, Wash., were used as headquarters for field investigations and experiments on the cultivation of oysters. In both places the work was carried out in cooperation with the State authorities.

This work is being conducted under the supervision of Dr. Paul S. Galtsoff.

EXPERIMENTS ON OYSTER CULTURE

Experiments on oyster culture were continued by Dr. Galtsoff and V. L. Loosanoff at Milford, Conn., in cooperation with the Connecticut Shellfish Commission and the Connecticut Oyster Farms Co. During the summer the State's boat, the *Shellfish*, was used for field work in Long Island Sound, while during the cold season the research work was dependent upon the facilities offered by the Connecticut Oyster Farms Co. All of the laboratory work was carried out at the Osborn Zoological Laboratory, Yale University, and at the United States fisheries laboratory, Woods Hole, Mass. As in previous years, much needed information was supplied to the oystermen regarding the expected time of spawning and setting. On several occasions, at the request of the United States War Department, Mr. Loosanoff assisted in examining the oysters and oyster grounds in New Haven Harbor to determine the effect of the dredging opera-

tions conducted by the United States Corps of Engineers. On many occasions the oyster growers of the district called upon the laboratory at Milford for information and advice concerning various problems of oyster culture.

The major part of the work was a continuation of the investigation on the growth and fattening of oysters started by the Bureau in the spring of 1932. It consisted mainly of a study of environmental conditions and seasonal changes occurring in oysters. By comparing the temperatures of the water for the last 3 years, two points of great biological interest and significance become apparent: First, the duration of the hibernation period and the activity of the oysters vary, but within very narrow limits. During the years 1932, 1933, and 1934 the oysters hibernated 155, 160, and 149 days, respectively; second, the spawning period, that is, the number of days per year when the temperature of the water remained above 70° F., was 42 in 1932, 69 in 1933, and 59 in 1934. It is interesting to note that the shortest period of hibernation occurred in 1934 regardless of the temperature during this year. The spawning period of 1932 constituted only about 60 percent of that in 1933.

From the observations on phosphates it becomes evident that the utilization and regeneration of these nutrient salts in Long Island Sound is a definite cyclic process with a minimum amount present in the water during the warmest part of the year and a maximum amount occurring in winter. The temperature of the water and solar radiation are the two principal factors controlling the phosphate cycle, which in turn determines the fluctuations in the abundance of plankton. Analysis of river water showed that it is very poor in phosphates and contributes very little to the phosphate reserve of the Sound.

Observations on the growth of oysters show that they continue to increase in length and weight throughout the entire year, including the winter months. The rate of growth gradually decreases with age. In 1932 the average increase in length of 4-year-old oysters planted on experimental beds at Charles Island was about 2 cm, in 1933 it amounted to 0.9 cm, and during the last year it did not exceed 0.5 cm. Changes in the increase in total weight are even more striking. In 1932 the increase in total weight of 4-year-old oysters was 72 percent, in 1933 it was only 40 percent, and during the last year less than 10 percent. Thus, the self-inhibiting phase of growth—that is, the phase during which the rate of growth decreases with the increase in size or weight of the organism—is very pronounced in the 6-year-old oysters. The weight of the shells of oysters examined during the year constituted from 72.9 to 82.5 percent of total weight, the average, 77 percent, being almost identical with that obtained in 1932 and 1933.

During the year the weight of the meat constituted from 10 to 13 percent of total weight, the minimum occurring after spawning; that is, in August-September.

Observations on the glycogen and mineral content of the oysters were continued. Of the 199 samples of oyster meats collected during the course of the investigations 45 were analyzed for iron, copper, manganese, and zinc.

Observations on the spawning of oysters made by P. S. Galtsoff and J. F. Reppun at Woods Hole disclosed some interesting details of this process. It has been found that the eggs of an ovulating female pass through the gills and are discharged from the inhalent chamber instead of being expelled directly with the outgoing current of water passing through the cloaca. Because of this peculiar manner of shedding, the eggs are more uniformly distributed in the water and have a better chance to meet with the sperm. Experiments with artificial stimulation of spawning produced interesting results, showing that male oysters can be induced to shed sperm by a great variety of organic compounds, as, for instance, various hormones (thyrosin, theelin, two extracts of pituitary gland), sugars, peptone, egg albumen, sperm of various mollusks, and even yeast. The reaction of the female is, however, very specific; it can be induced only by the sperm of the oyster.

OIL-POLLUTION INVESTIGATIONS IN LOUISIANA

In an attempt to carry out a more comprehensive study of the oil-pollution problem in Louisiana, where a serious mortality of oysters occurred in 1932-33 coincident with the development of oil wells, the Bureau obtained from the Civil Works Administration an approval of a project to carry out both field and laboratory experiments in the affected territory. Unfortunately, out of the \$42,000 allotted for this purpose only \$3,000 were made available to the Bureau, and after the completion of a preliminary hydrographic survey of Timbalier and Terrebone Bays and adjacent bodies of water by R. O. Smith, the work in Louisiana was discontinued. Laboratory experiments were carried out, however, by H. F. Prytherch and R. O. Smith at Beaufort, N. C., and P. S. Galtsoff and V. Koehring at Woods Hole and Washington. Although the exact cause of the mortality of oysters had not been determined, the results of the laboratory experiments throw light on the possible effect of oil pollution on oysters. The report of this work has been submitted for publication.

The laboratory experiments were designed to test the effect of crude oil and oil-well bleed water on the vital activities of the oyster, especially on the rate of feeding, and on the growth of diatoms which constitute the principal food of the oyster. Using a method developed by Galtsoff in his previous studies on the physiology of feeding, hundreds of records were obtained showing the effect of the presence of oil on the rate of pumping of water through the gills. The results of these observations show that crude oil contains certain soluble in the sea water substances which inhibit the activity of the ciliated epithelium of the gills and reduce the amount of water which the oyster passes through the gills. This toxic effect is proportional to the concentration of the water soluble fraction of crude oil. It is of interest that repeated washing of one sample of oil seems to have no effect on toxicity of the extract, for samples of it obtained from oil that had been washed 28 times were as toxic as those prepared from a new sample. Oil extracts have no effect on the adductor muscle and oysters kept under oil or in running sea water which was allowed to pass through a layer of oil remain open for the same periods of time

as the controls. But the ciliary epithelium is very sensitive to the oil extract which inhibits its action. Since the oyster feeds on minute plants suspended in water, the rate of feeding is directly proportional to the volume of water pumped through the gills.

There is no doubt that the presence of oil decreases the rate of feeding of oysters, and oil absorbed by colloidal clay and deposited on mud bottom may exert its adverse effect on oysters long after all traces of it have disappeared from the surface of the water.

Oil-well bleed water has an effect on oysters similar to that of the oil extract.

Experiments with the diatom *Nitzschia dosteria*, grown under controlled laboratory conditions, show that the rate of propagation is decreased by oil floating on the surface of the experimental flask and by the addition of oil extract. From the results of the laboratory experiments an inference can be drawn that both factors, the decrease in the rate of feeding of oysters and the diminished production of diatoms, create conditions adverse to the cultivation of oysters in the oil-polluted areas.

BIOLOGY AND CULTIVATION OF HARD-SHELL CLAM (*VENUS MERCENARIA*)

While certain phases of the life history of *V. mercenaria* are quite well known, chiefly through the work of Kellogg and Belding, much is still left to be learned about this animal, and there is a growing demand for the information regarding the methods of its cultivation. The aim of the study being conducted by V. L. Loosanoff at Milford, Conn., is to obtain a broad knowledge of its life history, habits, and physiology, and to develop practical methods of propagation. It is expected that the results of this investigation will be instrumental in devising methods for increasing the natural supply and for the preservation of the clam fisheries.

At present, the study is confined to the following phases: Development and seasonal changes of gonads, spawning, development of the egg, growth, and feeding.

The work along the first line consists in histological and cytological studies of the material collected at regular intervals from the experimental clam beds near Milford, Conn. The study is not quite completed. It appears from the material examined so far that the primary gonads of young clams are of protandric nature, that is, the individuals first pass through the male phase. Whether the *Venus mercenaria* is strictly protandric or some of the individuals mature as females without the completion of a preliminary male phase will be established upon the examination of more material. Young clams 6 to 7 mm long already contain apparently functional spermatozoa.

Studies of the seasonal changes in the gonads of adult clams reveal processes which differ markedly from those observed in the oyster. In the latter, after spawning is completed the gonad tissue is absorbed, so that in the winter the gonads are in a well defined resting stage. In *Venus*, a certain amount of absorption is also evident, the principal degeneration of unspawned eggs taking place in December, but the gonad follicles do not shrink noticeably, and well developed oocytes are always present. In the male the situa-

tion is still more amazing. Active spermatozoa are present in every month of the year. Upon being placed in the water they swim actively and behave in normal fashion. One interesting observation has been made on the movements of spermatozoa. During the study of fertilization it has been observed that sperms of different clams do not behave similarly. While they all exhibit spiral movements, the spermatozoa of about 75 percent of males move in circles in an anti-clockwise direction, while the other 25 percent of the males produce sperms moving in the opposite direction. The sperms of one male usually all move in the same direction. On several occasions young oocytes were found present in adult males 6 to 8 years old. This apparently shows the potentiality for change of sex even in adult clams. However, gonads of truly bisexual character in an adult clam have been observed in one case only. Functional hermaphroditism has been observed only once.

A study of the spawning of clams and its physiology was conducted last summer. It has been found that it is very difficult to induce the spawning of clams, especially females, under the laboratory conditions. The critical spawning temperature is higher than that required for the oyster. Under the laboratory condition 25° C. was the minimum spawning temperature.

To study the growth of young clams three experimental beds representing different environmental conditions were established in different parts of Milford Harbor. Young clams from all of the beds were measured at regular intervals and the progress in growth recorded.

The feeding of clams depends upon the combined action of syphon, gill epithelium, and adductor muscles. At present only shell movements are being studied. The clams are kept under natural conditions in the large concrete tanks, 20 by 18 by 6 feet, in which the water is renewed by tidal action twice each day. Since the beginning of the experiment on October 17, 1934, over 160 records of shell movement have been obtained covering the range of temperature from 0° C. to 13° C.

A study of the glycogen content of hard-shell clams was begun last March. Samples of clams were collected every 2 weeks and the meats analyzed. The glycogen content varies greatly with physiological changes occurring in the clams at different seasons of the year. The maximum quantities of glycogen found in clams during the prespawning period constituted 10 to 11 percent of the meat weight (fresh basis). After spawning a very sharp drop takes place until at times it equals only 3.3 percent of the meat weight. Gradual recovery takes place after that and during the winter the glycogen content of the clams constitutes 6 to 9 percent of total weight. Grown under similar conditions clams always have a higher glycogen content than the oysters.

REHABILITATION OF PUBLIC OYSTER BEDS IN NORTH CAROLINA

In order to restock the depleted natural oyster beds of North Carolina, extensive operations for the transplantation of seed oysters have been carried out under the direction of Dr. H. F. Prytherch,

in cooperation with the North Carolina Department of Conservation and the Federal Emergency Relief Administration.

Over 825,000 bushels of seed oysters at an average cost of approximately 8 cents per bushel have been planted, and 78,567 bushels of old oyster shells have been scattered over the bottoms to provide a place of attachment for subsequent generations of this shellfish. Inspection of the replanted beds in Pamlico Sound during September 1934 showed that a high percentage of the seed oysters have survived transplanting and culling operations and, because of their rapid growth under more favorable conditions, would provide good market stock in approximately 2 years. At the recommendation of the Bureau all replanted areas have been closed by a recent ruling of the State Fisheries Board for a period of 2 years in order to enable the transplanted oysters to reach market size.

INVESTIGATION OF OYSTERS AT NEW RIVER, N. C.

A general survey was made during March by Dr. H. F. Prytherch of the oyster beds in the New River region to determine the cause of the poor condition of oysters during the previous winter. The results of the Bureau's investigation, briefly summarized, are as follows: (1) The poor condition of New River oysters is attributed to exceptionally dry weather and decreased river discharge during the period from October 1933 to February 1934 when precipitation was 75 percent below normal; (2) closure of the inlet has not interfered with oyster growth and reproduction, as sufficient salt water now passes into this region through two larger inlets via the Inside Route Canal; and (3) deepening of New River Inlet is not advisable as it might easily increase the salinity of the water to such an extent that oyster beds now producing a high-grade product would be overcrowded with seed oysters and also subject to severe attacks by the boring sponge.

OYSTER INVESTIGATIONS IN FLORIDA

A complete survey of the principal oyster-producing areas in northwest Florida was conducted by Dr. H. F. Prytherch in cooperation with the State Department of Conservation and Federal Emergency Relief Administration for the purpose of restoring and developing this natural resource as a work-relief project. The most important beds were found to be in a depleted condition and in need of restocking with seed oysters and shells. Experimental planting operations of the type successfully carried out by the Bureau in this region last year clearly demonstrate the advisability of employing such methods on a large scale. Biological studies were made of the spawning and setting of oysters in these waters, which show that the planting of old shells should be carried out during the period from April 1 to September 15.

In Choctawhatchee Bay over 12,000 barrels of shell have been distributed over barren bottoms to create new beds and to increase the acreage of areas which are producing high-grade oysters. The general procedure is to plant 100 to 300 barrels of shell per acre, over

which seed and adult oysters from the inshore reefs will be distributed in the spring to serve as spawners.

Destruction of oysters by parasitic flatworm.—A parasitic natural enemy of the oyster known as the "wafer" or so-called "leech" has become a serious menace to the continued natural production and private cultivation of this shellfish in Apalachicola Bay, Fla. Investigations conducted by Dr. Herbert F. Prytherch, in cooperation with the Florida Department of Conservation, show that this pest in the last 3 years has completely destroyed the seed and adult oysters on several of the best natural beds and is spreading rapidly to other valuable areas. The parasite is a turbellarian flatworm, *Stylochus inimicus*, measuring from one-half to three-fourths inch in diameter, which enters the shell of the oyster and feeds gradually upon the meat until it has killed its host. Since the fall of 1932 the parasite has destroyed the oysters on five natural beds comprising an area of 800 acres on which a crop estimated at 350,000 bushels, having an approximate value of \$175,000, has been lost. Recently this natural enemy has spread to other beds at a distance of 12 miles from the original outbreak and to a nearby oyster reef, which is the largest in this region and contains nearly half of the entire crop. The Bureau's investigations show that the rapid growth, reproduction, and spread of the leech in Apalachicola Bay is associated with an unusually dry period, a considerable decrease in river discharge, and an increase in the salinity of the coastal water of this region. It is desirable that extensive dredging operations be undertaken to destroy the leech on its principal breeding areas and that scientific studies be made of this parasite, which has been recently found on oyster beds from North Carolina to Texas.

PROTOZOAN PARASITE OF VIRGINIA AND LOUISIANA OYSTERS

Oyster planters in Mobjack Bay, Va., and in Terrebonne Parish, La., suffered a loss of their stock amounting to over a million dollars during the winters of 1929-30 and 1932-33, respectively. Cytological studies of oysters from both regions conducted by Dr. H. F. Prytherch show that they were heavily infested with a parasitic protozoan, which tentatively has been identified as one of the Haplosporidia. This parasite was particularly abundant in the tissue of the adductor muscle, mantle, and gills, and was frequently found in the muscle of the heart. Representative samples of oysters from the mortality areas showed in every case a heavy concentration of this protozoan, the number ranging from approximately 50 thousand to 1 and 2 million per oyster.

In the immediate vicinity of the fisheries laboratory at Beaufort, N. C., it has been observed for several years that a large number of oysters die at an age of 1½ to 2 years, following a period of abnormal shell growth, similar to that observed in the oyster mortalities in Virginia and Louisiana. Recent cytological studies have shown that these oysters are also heavily infected with the same protozoan parasite. In all three localities the mortality was highest in oysters grown under crowded conditions in which large numbers of this minute parasite were found. From a practical standpoint it is apparent that oysters should be planted sparsely to prevent reinfection and spread

of the parasite and also allow more favorable growing conditions for those which are heavily infected. Investigations are being continued at the Beaufort laboratory for the purpose of determining the life cycle, method of distribution, and proper identification of this micro-organism which appears to be a new natural enemy of the oyster of considerable economic importance.

OYSTER INVESTIGATIONS IN WASHINGTON

For a number of years the Olympia oyster industry has had difficulties due to unsatisfactory catches of seed oysters. Although the quality of the product has been normal the growers were unable, by use of the usual methods, to collect spat enough to fill the beds. For this reason an investigation of the spawning and setting habits of this species of oyster was undertaken by Dr. A. E. Hopkins. During four seasons accurate data have been collected on various phases of the problem, and the cumulative results are becoming constantly more significant. In addition, the rapid expansion of the Japanese, or "Pacific", oyster industry has involved numerous problems, such as propagation of this species in American waters, effect of environmental factors on fattening, and so forth.

Experimental studies have been carried on in typical oyster-growing areas of the most important bays near Olympia. Three times weekly during the breeding season 100 adult oysters (*Ostrea lurida*) were opened on each of the test beds, and the larvae from gravid specimens preserved for laboratory study. In this manner it was learned that larvae develop for about 10 days in the maternal brood chamber before being cast out into the open water. The data also show accurately the time and intensity of spawning in each locality throughout the season, as well as the total amount of spawning taking place. For example, in Oyster Bay in 1932 all of the oysters produced larvae once and 75 percent of them produced second broods. In 1933 only about 75 percent spawned as females, and in 1934 between 90 percent and 100 percent did so. These results indicate that there is wide variation between different seasons in the number of times spawning occurs, and consequently in the potential seed catch.

The time of beginning of spawning is clearly determined by temperature. During each season the first gravid specimens were found when the average daily water temperature reached 13° to 14° C. Records for the past 3 years show that the interval between spawning and setting varied between 39 and 51 days, of which the first 10 days represent development within the maternal brood chamber.

By careful analysis of the time and intensity of setting throughout the season it has been found that in certain bays periods of setting occur at intervals and are associated with the tidal cycles. This was found during previous years, and the 1934 results furnish further confirmation. Attempts have been made to determine in what manner the tidal cycle is effective, but the problem requires further experimentation. Although setting is heaviest during a run of extreme low tides, actual attachment of the larvae does not occur when the tide is low but appears to take place primarily at relatively high tide. Whether copper in the water is the effective agent for this species has not yet been determined.

Although at present it is impossible to predict with certainty when setting will begin, it is known that after the first larvae begin to attach there is still sufficient time for the growers to plant cultch before the most profuse setting takes place. By making constant observations and determining the exact date when setting begins one may be reasonably certain when the peak of the set will arrive, since this is so closely associated with the tidal cycle. By this method in 1933 and 1934 definite advice was given to oyster growers as to the time to plant cultch for best results. The seed catch of those who followed the advice has amply demonstrated the adequacy of the method.

Several years ago, on the basis of observations that larvae attach most abundantly to lower surfaces, a type of cement-coated cardboard spat collector—a modification of the egg-crate filler—was designed to furnish a large amount of horizontal surface. These collectors are now being used commercially in Puget Sound with very satisfactory results.

That the circulatory system of the oyster is more complicated than previously thought was shown by the discovery that an oyster possesses a pair of accessory hearts, or large, rhythmically pulsating blood vessels, within the mantle walls of the cloacal chamber. They appear to pump blood from the kidneys to the mantle and gills, where it is aerated. It is suggested that the organs are homologous to the branchial hearts of cephalopod mollusks. The accessory hearts are found in all three commercial species of oysters in the United States.

Further experiments were made on the adaptation of the Pacific oyster to changes in salinity. Adaptation of the feeding mechanism is relatively slow following a lowering of salinity, but rapid after a rise. Feeding ceases completely when salinity is reduced to 12-14 parts per thousand, and apparently adaptation does not occur, although growth of shell continues. It is probable that adaptation to salinities of 15-20 parts per thousand is never such as to permit the oyster to feed as rapidly as it does at 25-35 parts per thousand.

INVESTIGATIONS ON AQUICULTURE

Investigations in aquiculture were conducted in 1934 under the direction of Dr. H. S. Davis. These investigations, which were originally undertaken in connection with fish-cultural operations in the hatcheries, have been expanded to include field studies dealing with the various factors which affect fish in their natural environment. Opportunities for such field investigations have been greatly increased during the past year as a result of an allotment from the Public Works Administration for stream surveys and stream improvement in the national forests and parks.

The experimental hatcheries at Leetown, W. Va., and Pittsford, Vt., are strategically located for conducting investigations in the waters of the national forests. The Leetown station is within a short driving distance of both the George Washington and Monongahela National Forests, while the Pittsford station is situated within the boundaries of the recently enlarged Green Mountain National Forest.

Thus, both stations afford exceptionally favorable opportunities for conducting field studies and experiments in cooperation with the United States Forest Service.

POND-FISH CULTURE

Investigations in pond-fish culture during 1934 were limited to those conducted by O. Lloyd Meehan at the Natchitoches (La.) station. These investigations are designed primarily to throw light on problems which have arisen in connection with pond culture in the Southern States where conditions in many respects are quite different from those found at hatcheries in the North.

The investigations at the Natchitoches station were concerned chiefly with the use of fertilizers in pond culture. Since experiments in previous years have shown conclusively that proper fertilization of rearing ponds results in a marked increase in the production of fish, the experiments during 1934 were designed primarily to furnish a comparison of the value of different fertilizers for this purpose. The results, however, are inconclusive, since no correlations were found between any of the ecological groups of food organisms and the amount of the different elements composing the fertilizers. It is concluded that this was due to the fact that more fertilizer was used than was necessary. Consequently, there was at all times a surplus of food material available. Under such circumstances it is logical to assume that the production would be limited by other factors than food.

The results agree with those obtained in 1933 in failing to show that there is necessarily any correlation between the production of plankton in a pond and the production of fish as claimed by some authors. There is, however, a distinct correlation between the number of Chironomid larvæ per unit area of bottom sample and the production of fish. This indicates that fish production is dependent on bottom organisms rather than on plankton.

The experiments indicate that the number of fish produced per acre is not a good index of production in a pond after bass have reached a length of about $2\frac{1}{2}$ inches. The greater forage area needed by the larger fish and the change in food habits necessarily reduces the number of individuals. Consequently, with larger fish the weight of the fish rather than numbers should be taken as a criterion of production. This is especially important in the South owing to the early spawning season, the young bass usually reaching a length of $2\frac{1}{2}$ inches by the 1st of June. Bass can be reared successfully to this size without the use of forage minnows, but if they are to be kept in rearing ponds through the summer the need of forage fish is clearly indicated.

As a result of these experiments, Mr. Meehan believes that in the Gulf States at least it may be necessary to choose between heavily fertilized ponds which will carry a large number of bass to 2 or $2\frac{1}{2}$ inches and ponds producing an adequate supply of forage food for a longer growing season. In the heavily fertilized ponds the number of fish may run very high if the ponds are drained and the fish removed by the 1st of June. These ponds could be utilized during the remainder of the season for sunfish and catfish or other species that require a longer growing season and spawn later. The other alterna-

tive will be to supply a large amount of forage food to carry the bass through the summer until October or November, in which case there will be a smaller number of large bass.

An allotment from the Public Works Administration has made possible the construction of a number of bass ponds at the Leetown (W. Va.) experimental station, but these ponds were not completed in time to be used during the season of 1934.

TROUT CULTURE

Feeding experiments.—Owing to lack of sufficient funds the feeding experiments which have been carried on at the Pittsford and Leetown stations for several years were discontinued. However, at the Leetown station some preliminary studies were made on the growth of yearling and 2-year-old trout. These fish were fed a diet composed of 60 percent sheep liver, 20 percent salmon egg meal, and 20 percent meat meal. Each experimental lot consisting of 325 fish was held in a circular pool 22 feet in diameter. It was found that the rate of growth of both yearling brook and rainbow trout as well as that of 2-year-old rainbow and brown trout was considerably smaller than in the case of fingerlings. In feeding fingerlings the amount of food consumed increases steadily throughout the summer; but with the older fish on the diet used, the amount consumed remained unchanged in some instances for 2 months or more, the fish continuing to grow nevertheless.

It was found that both yearling brook and rainbow trout consumed a greater amount of food per day in proportion to their body weight than the 2-year-old fish. The experiment also showed that rainbow and brown trout could eat more of the diet fed without ill effects than the brook trout. In other words, the brook trout, when given an opportunity to eat all they could, suffered injurious effects as indicated by high mortality and loss of appetite. This indicates that brook trout are more easily overfed than brown or rainbow trout.

The nutrition studies have been continued at the Cortland, N. Y., hatchery by Dr. C. M. McCay and A. V. Tunison. Two lines of research were conducted during the past year: (1) Studies of the interrelationship between various foodstuffs, rate of growth, and mortality were continued; and (2) chemical-balance studies with trout have been initiated. The chemical-balance studies depend upon accurate chemical analyses of both the food ingested and the excreta.

The experiment to determine the relative ability of lake trout (*Cristivomer namaycush*), brown trout (*Salmo fario*), rainbow trout (*Salmo irideus*), and brook trout (*Salvelinus fontinalis*) to convert foodstuffs into body tissue were carried through the ninety-second week. All four species were maintained under similar conditions and fed the same diet, which consisted of 2 parts fresh beef liver and 1 part dry skim milk. The number of individuals was reduced from time to time so as to prevent overcrowding. The uniformity of the growth rates of the four species is surprising. Over long periods of time the growth tends to be strictly logarithmic. However, the rate changed to a lower one after the thirty-second

week. Under the conditions of the experiment the lake trout were the most efficient in converting food into body tissue, while the brown trout were the least efficient. It is evident that older fish are more efficient in the conversion of their food. The rate of growth and the maintenance requirements partly tend to counterbalance each other in the efficiency of food conversion.

Experimental feeding of trout fry has given some interesting results, from which the following conclusions can be drawn: (1) A mixture of 2 parts fresh beef liver and 1 part dry skim milk is a satisfactory diet for fry; (2) a higher percentage of milk tends to increase the mortality; (3) the mechanical method of feeding makes no difference in results when the fry are fed the same amount in percent of body weight; (4) different groups of fry yield uniform data for food conversion when the same diets are employed; (5) on a dry-food basis, the conversion of liver and milk is more efficient and economical than liver alone; (6) the most efficient conversion of food occurred in one of the groups of brook trout kept in warmer water; and (7) increasing the daily feedings from 4 to 6 gave a slightly better growth and conversion.

The large-scale experiments in the practical feeding of dry-feed mixtures of cottonseed meal, fish meal, dry skim milk, and salmon-egg meal supplemented with raw liver were continued. A different strain of fish was used in 1934 as well as several improvements in methods, but the results were practically the same as those obtained in 1933.

An attempt was made to produce the same brilliant coloration found in wild trout. Dried salmon eggs were extracted with 95 percent alcohol and the colored extract fed to brook trout. The fish fed the extract developed marked color at about the eighth week, while those fed the residue developed little, if any, color. Salmon oil produced a similar coloration when mixed with the dry food at a 10 percent level. However, in no case did the colors produced approach those of wild fish. The fat soluble histological pigments Sudan (III and IV) failed to produce color in the trout except that when fed at the 0.5 percent level the intestines and mesentery fat showed some pigment.

Some further work has been done on the feeding of preserved fresh meat. Liver preserved in both 1 and 2 percent formalin and in chlorinated lime was sealed in glass jars and kept at 10° C. for 2 years. This preserved liver was then used at a 25 percent level, to supplement a mixture of skim milk, fish meal, and cottonseed meal. After 14 weeks the trout had doubled their weight with no signs of being poisoned by the preserved meat.

The chemical balance studies have concerned themselves with the determination of the digestibility of fats. Two objectives were in mind: (1) to determine whether oils such as cottonseed and salmon were digested better by trout than a hard fat such as hydrogenated cottonseed oil (Crisco), and (2) to measure the relative digestibility of these fats by trout of two different ages and sizes. It was found that the oils were digested better than the solid fat, and that there was no real difference between the digestion and utilization of cottonseed oil and salmon oil. The melting point of the oil or fat is the essential characteristic rather than its origin. The trout seemed to utilize

about the same percent of the fats when the level in the diet was 7 and 25 percent. Fats were digested to about the same extent by trout weighing 2 grams and those weighing 100 grams.

Selective breeding.—Experiments in selective breeding of brook trout were conducted at the Pittsford (Vt.) station under the direction of R. F. Lord. At the beginning of the year, 43 lots of fingerling trout were being held in small compartments in hatchery troughs. Each lot was composed of the progeny of a single pair of selected fish. The parent fish were in their third year and averaged 12.9 inches in length with an average weight of 16.6 ounces in the case of the females, while the average weight of the males was 18.2 ounces and the length 13.6 inches. Such differences in the sizes of male and female brook trout of the same age are quite typical.

The young fish were kept under strict observation, and late in the spring those lots which did not reach the desired standard were discarded. The remaining lots were reduced to 400 fish each, which were kept segregated during the summer. The surplus fish from these lots were retained for general brood stock.

At the close of the season the various lots were carefully compared and only the very best lots with regard to growth, resistance to disease, and hereditary background were retained.

As indicated in previous reports, the results of these experiments in selective breeding have been very gratifying. There has been a remarkable increase in the egg production and rate of growth of selected fish and also an improvement in resistance to disease. Similar experiments with rainbow and brown trout are being carried on at the Leetown station, but there has not yet been time for these experiments to show material results.

Experiments in rearing exotic species.—The Pittsford brood stock of Montana grayling was stripped for the third time in the spring of 1934. A good hatch was obtained from these eggs, and about 6,000 fingerlings were retained for wintering. It is planned to use these fish to stock a pond which has recently been built in the Green Mountain National Forest. A stock of grayling derived from eggs shipped from Pittsford is also being reared at the Leetown station. These fish have done very well, and it is expected they will produce eggs in the spring of 1936.

Excellent results were obtained from a small lot of golden trout (*Salvelinus aureolus*) eggs from Lake Sunapee, N. H., which were shipped to Pittsford in the fall of 1933. The fingerlings were carried through the summer with very little loss and by fall many of them had reached a length of 6 inches or more. A tendency of the fish to crowd together at one end of the trough was overcome by excluding the light.

A stock of California golden trout (*Salmo aqua-bonita*) is also being reared at the Pittsford station. These fish are quite susceptible to furunculosis, and considerable losses have occurred from this disease; but in other respects they appear to be no more difficult to rear than other species of trout.

Field studies.—An investigation of the results of planting a certain number of rainbow trout fingerlings in a 1-mile section of a spring-fed stream near Leesburg, Va., was continued by E. W. Surber, in charge of the Leetown station. In addition to collecting data on the

ages, condition factors, and food of legal size rainbow trout removed from the stream, quantitative studies were made of the natural food present. For this purpose bottom samples were taken in the gravel riffles at 10 to 16 stations along the stream at monthly intervals.

During the season of 1934, a total of 94 legal-size fish were taken from the stream with hook and line. This is a production of 27.66 pounds of trout per acre. These results agree very closely with those obtained in 1933, when a production of approximately 30 pounds per acre was recorded. Measurements of the fish showed that they had made a very good growth and were in excellent condition.

Food studies again demonstrated the great dependence of rainbow trout on terrestrial insects during the summer periods, but these studies also showed greater dependence on aquatic forms than in the previous season. A total of 196 bottom samples on gravel riffles showed an average wet weight of 5.047 grams per square foot and 0.982 gram dry weight per square foot for the year. This amounted to an average standing crop of 485.8 pounds (wet weight) per acre in the gravel riffles of the stream. A study of the variability of sampling showed that the samples varied less than one-third from place to place in the numbers of *gammarus* and *fasciatus* (the predominant organisms) in individual samples during May, June, and July.

An attempt to obtain statistics on the number of trout caught during the season of 1934 in several streams in Vermont through cooperation with the anglers was far from successful. Mail boxes painted a brilliant red to attract attention were placed along the streams at locations where anglers usually left their cars. A conspicuous sign attached to each mail box called attention to the fact that the Bureau of Fisheries was attempting to secure figures on the number of fish taken annually from the stream in question as a guide to future stocking. The anglers were urged to cooperate by giving the desired information on cards provided for the purpose, which were to be deposited in the mail boxes. The boxes were all in position when the fishing season opened on May 1, 1934.

In spite of efforts to secure the cooperation of anglers with the least possible trouble on their part, the results were very disappointing. Only a small percentage of those fishing in the streams filled out reports during the first day or two of the season, and as the season progressed the number of cards deposited in the boxes steadily decreased until toward the end of the season almost no reports were received.

The best returns were received from the anglers on Furnace Brook, although it was evident that only a small percentage of those fishing the stream filed the reports. The data revealed that during the months of May and June fishing conditions were quite uniform, with an average catch of 7.4 trout per fishing attempt. In July fewer anglers reported that the average catch was 18 trout. This is not taken to mean that fishing conditions had improved but that only the more successful anglers reported.

Similar returns were obtained on the South Branch of Cold River, a small mountain stream with an average volume of about 12 to 15 cubic feet per second. This stream was stocked with 700 marked

yearling rainbow in September 1933. Only 24 reports were received from anglers for the month of May, and these reported 82 rainbows of the 1933 planting. Three reports were received later, which brought the total number of rainbows caught during the season to 87 fish. This is a return of approximately 12 percent. Since it was evident that only a small percentage of the anglers reported their catch, it is believed that the returns from this planting were quite satisfactory. One rainbow was reported which had been planted in 1932. These experiments are being repeated during the present season.

In cooperation with the Middlebury College an attempt was made to obtain similar data on waters stocked and controlled by the college. Special permits required of those fishing in streams on college property were issued with the understanding that each angler would report his daily catch in a space reserved for this purpose on the permit. Even under these circumstances it was found very difficult to get returns, and the reports received were too meager to throw much light on trout-stream production. They do show, however, that the average daily catch of legal-size fish was 9.4 per trip in 1933 and 7.4 per trip in 1934. The reports also show that the number of short trout taken and returned to the stream greatly exceed the number of those which had reached legal size.

California trout investigations.—Work has progressed rapidly on most of the projects initiated in 1932 when these investigations were started. Subsequent experience necessitated several modifications of the original program and the work now consists of two major projects, the Hot Creek brood stock experiment and the coastal stream steelhead studies. The former project is under the direction of Dr. P. R. Needham, who is also in charge of the field program as a whole. The latter project is under the supervision of A. C. Taft. Work on several minor projects is being carried on as time permits. Three assistants are supplied to the work by the California Division of Fish and Game.

The Hot Creek experimental project was greatly expanded this year with the construction of a series of 12 concrete raceways, 20 feet long by 4 feet wide. Two new ponds to hold brood stock were also constructed as well as a combination garage, meat house, and laboratory. Late in the fall a small experimental hatchery was completed with C. C. C. labor. Over \$3,000 was expended on these improvements. Over 67,000 small trout, consisting of 6 separate strains of rainbows and 1 lot of eastern brook trout, were placed in the ponds after construction work was completed. These fish are to form nuclei for rearing select domestic brood stocks for experimental purposes. An additional source of rainbow eggs is needed in California, and breeding experiments for high egg yields, rapid growth, coloration, and other genetically desirable characteristics, therefore, will be carried on largely with this species. One distinct advantage offered at the Hot Creek area is the tremendous natural production of shrimp, making it unnecessary to feed artificial food to many of the younger fish, thus reducing costs materially. Over a quarter of a pound of pure shrimp were taken from an area of 1 square foot in the watercress in one of the ponds. A fine series of springs supply about 30 second-feet of water at temperatures varying from 57° F. to 68° F. which permit rapid growth over most of the year.

Data from the coastal stream steelhead studies started in 1931 at Scott and Waddell Creeks and on the Klamath River are rapidly being augmented. The counting weir constructed on Waddell Creek in 1933 has been operated continuously since this time.

During the winter season of 1933-34 at Waddell Creek, the spawning migration consisted of 478 steelhead and 538 silver salmon of which 263 and 213 were females, respectively. The silver salmon migration was largely confined to the last 2 weeks of December, while the steelhead came in over a longer period which extended from January to the end of March. There was a considerable loss among the steelhead due to disease, which was probably furunculosis and which was associated with very low and warm water during March, April, and May. Most of these fish died after spawning. In addition to the adults moving upstream, 128 young fish varying in length from 90 to 295 centimeters were taken in the upstream trap during the months of December and January. These were undoubtedly fish which had spent the summer in the closed lagoon, and some of the larger were sexually mature males. They were given a distinctive mark and passed on upstream. Later, 70 of these fish were taken on their downstream migration with the other young steelhead.

There were small numbers of downstream migrating steelhead during December, January, and February, but the movement was greatest during March and April, and from length frequencies it appears that fish that had just completed their second year predominated. A total of 3,117 young steelhead were trapped and 2,452 of them marked by the removal of the adipose and right pectoral fins. During June, July, and August, a few fish of the year were taken in the trap each week.

The downstream migration of young silver salmon (*O. kisutch*) was much more concentrated than that of the young steelhead; that is, there was no scattering of fish previous and subsequent to the main movement. The migration was also later, coming in April and May. A total of 3,430 young salmon were taken, and 3,211 of them were marked by the removal of the adipose and right pectoral fins.

An interesting discovery was the fact that the two species of cottoids inhabiting the stream (*Cottus asper* and *C. gulosus*), apparently migrate downstream to spawn. During the period December to May, 3,357 of these fish were taken in the downstream trap and both males and females were approaching spawning condition as evidenced by the gonads. Although efforts were made to determine just where they spawned, neither eggs nor fry were found. These cottoids are very destructive to young trout and salmon and this annual period of migration may offer an opportunity to reduce their numbers in such streams as it is felt desirable to do so.

Work has been continued at Scott Creek on certain phases of the artificial propagation of steelhead and, in conjunction with Waddell Creek, on the homing instinct in these fish. The mouths of these two streams are separated by only 5 miles of coastline and the annual release of marked fish in each stream should make it possible to determine the degree of straying. It is becoming increasingly difficult to secure a supply of steelhead eggs for the hatcheries without encroaching on streams which are also heavily fished. An attempt is being made at Scott Creek to build up this stream for egg taking and

although water conditions have been so unfavorable during the past few years that parts of the stream have dried up completely, the run is being slowly increased through holding fish in ponds and planting them during the winter months. During the past year 10,054 of the fish planted were marked by the removal of both ventral fins. During the winter season 455 adult fish were taken in the trap and before release were tagged with celluloid disks and nickel wire. The tags were placed in the anterior base of the dorsal fin. Of the fish tagged the previous year approximately 7 percent returned this year as compared with a return of 13 percent from the previous year's tagging. The returns are quite obviously dependent on the age of the fish tagged as very few fish return to spawn the third time.

On the Klamath River 45,700 yearling trout were marked and planted during May. Part of these fish were from eggs taken in the Klamath River while 9,770 were from a possibly nonmigratory type of rainbow taken in Kosk Creek, a tributary of the Pit River. At Beaver Creek on the Klamath, arrangements have been made to install a trap at the egg-taking station to take downstream migrants. This will make it possible to correlate information obtained at Waddell Creek with that of the streams tributary to the Klamath.

Work was also continued on two minor projects, the Angora Lake and Truckee River marking experiments.

The Angora Lake project was started in 1933 with a plant of over 5,000 marked eastern brook trout. The objects of this experiment are to determine: (1) Total annual production in pounds of fish per acre of water as shown by anglers' catches; (2) survival rates from plants of various lots of marked trout of given sizes; and (3) the correlation between the actual production in fish and food conditions in the lake. The lake offers exceptional opportunities for obtaining complete returns from anglers, and the resort owner on the lake is receiving a small annual stipend to record and weigh all fish caught.

Returns for the fishing season of 1934 show that anglers took only 81 trout, having a total weight of 71.2 pounds. Since Angora Lake is, roughly, of about 5 acres area, these figures show that production in terms of fish landed by anglers was approximately 14 pounds per acre of water surface. The average catch per angler was 2.3 fish. Of species, 25 loch leven, 14 rainbow and 42 eastern brook trout were taken. The figures are based upon partially complete returns. None of the marked eastern brook trout planted in the previous summer entered the catches this year, due doubtless to the fact that their average size at planting was slightly under 2 inches.

Excellent returns have likewise been obtained from the Truckee River marking experiment which was started in 1932 with the planting of 40,000 trout. In 1933 an additional 40,000 were planted, and they entered into the catch in considerable numbers during the past summer. In one lot of 62 fish taken by anglers, 49 were fish of this marking.

Field studies of stream food problems were continued in Waddell Creek in the spring, though lack of time and the pressure of other matters prevented detailed work of the type carried on the year previously. Bottom food samples were taken to determine the bulk or weight of food which each group contributes to the total potential supply in this stream. It was found that caddisfly larvae and pupae,

which formed only 22.2 percent of the total number of organisms taken in one series of 13 samples, formed 43.9 percent of the total wet weight of all organisms, offering the most food by bulk of any group of aquatics. Mayfly nymphs, on the other hand, were most abundant, forming over 55 percent by number but only 28 percent by weight. Stonefly nymphs were third in weight of food at 12.2 percent and fourth in numbers at 7.8 percent. Aquatic truefly larvae and pupae (Diptera) formed 10.3 percent by number and only 7.9 percent by weight of total available foods.

Some work was started on the distribution and abundance of stream foods in relation to water temperatures and other ecological factors, but much further field and laboratory work will be necessary before results can be submitted on these phases of the problem.

FISH DISEASES

Investigations were conducted by Dr. Frederic F. Fish on the "ulcer disease" of trout which he has shown to differ in many respects from furunculosis with which it was formerly confused. The disease is characterized by definite thickening of the epithelium over a small area, producing an inconspicuous white patch which in general appearance closely resembles a small growth of fungus (*saprolignia*). This condition is best described as an "epithelial tuft." Eventually the skin becomes perforated and the small ulcer is formed. When a lesion of the "ulcer disease" develops on a fin, the soft tissue between the fin rays is destroyed leaving the rays projecting beyond the necrotic tissue. As the disease progresses the necrotic area advances in a more or less horizontal line toward the base of the fin. In advanced stages of the disease the fin may be entirely destroyed while the infection penetrates deep into the underlying tissue. There is no evidence, however, that the infection ever reaches the internal organs.

The causative agent of the disease is evidently a bacterium. Several species of bacteria have been isolated from diseased fish, but it has not yet been possible to demonstrate conclusively that any one of them is the specific etiological agent.

During the early part of the year Dr. Fish was stationed at the Leetown (W. Va.) station to investigate an epidemic of blue sac which was causing heavy losses among the brook and brown trout fry. In contrast with other investigators, Dr. Fish was unable to find any definite evidence that the disease was due to a specific infection. Cultures of the serum fluid gave negative results except in a few instances. Sections of the diseased tissue also failed to disclose any evidence of bacterial activity and there was nothing to indicate that the disease spread from one trough to another as is usually the case with bacterial infections.

During the summer Dr. Fish was detailed to cooperate with the Oregon State Game Commission in a study of fish-cultural problems which had arisen in connection with the operation of the State trout hatcheries. Field headquarters were located at the Oregon State College, Corvallis, and inspection trips were made to all but two minor State trout hatcheries and to many points where the Oregon Game Commission had been carrying on field work. The purpose

of these trips was to review the entire system of trout propagation and management in the State. Special attention was paid to the prevalence and control of disease among hatchery fish. At the end of the season a report was prepared on the operation of the individual trout hatcheries based on actual observations and on a questionnaire sent to all hatchery superintendents.

In order to facilitate the study of diseases which are becoming a serious problem in many trout and salmon hatcheries in the West it was decided to install a pathological laboratory at the Fisheries biological station in Seattle with Dr. Fish in charge. The need for such a laboratory is very evident since there are reasons for believing that in addition to most of the diseases which affect trout at Eastern hatcheries there are other diseases which are peculiar to this region or at least have not yet been found at Eastern hatcheries.

Experiments have been conducted looking toward the improvement of methods of treating fish in pools by allowing a very dilute chemical solution to flow into the pool at a uniform rate. Solutions of potassium permanganate, copper sulphate, and chlorine have been recommended for this purpose, but in no instance have the methods yet devised proved entirely satisfactory. It has been found extremely difficult in practice to treat fish in pools with such solutions without serious danger of killing large numbers of fish. In some instances very satisfactory results have been obtained, but in other cases heavy losses have occurred when using the same treatment. It is believed, however, that these difficulties can be eliminated largely by further experiment in the technique of treatment.

An experiment to test the value of iodine as a cure for furunculosis was carried out at the Pittsford station. For this purpose a lot of yearling California golden trout, which are quite susceptible to the disease, was divided into two equal lots and placed in small circular pools. Before beginning the experiment these fish had been showing a small daily mortality from furunculosis for some time. One lot was used as a control, the other lot was placed on a diet containing salmon oil and a solution of iodine according to the recommendations of E. R. Hewitt. The experiment was started August 10 and for the first 13 days the losses in both lots were approximately the same. The total losses on August 23 were 46 in the lot given iodine and salmon oil, and 47 in the controls. The supply of salmon oil was exhausted on that date and the iodine solution soon after, but the experimental lot was kept on a dry diet supplied by Mr. Hewitt for a month longer. During this period the loss among this lot was appreciably less than among the controls. The experiment was discontinued on September 22 at which time the losses among the lot on the Hewitt diet totaled 74 and among the controls 144.

While these results show approximately twice the mortality among the controls as among the fish kept on the Hewitt diet, it is by no means demonstrated that this difference was due to the inclusion of iodine or salmon oil in the food. As a matter of fact, the supply of both of these constituents was exhausted at about the time the fish began to show improvement, and the decreased mortality among the experimental lot may easily have been due to other causes. In previous experiments marked differences in mortality from furuncu-

losis have been observed frequently among fish on different diets. Whatever the explanation, these experiments evidently lend little support to the view that iodine is a specific cure for furunculosis.

STREAM SURVEY AND STREAM IMPROVEMENT

During the summer of 1934 the Bureau inaugurated an extensive series of stream survey and improvement work in the national forests and parks in cooperation with the United States Forest Service and the United States Park Service. This work was made possible by an allotment of \$127,500 from the Public Works Administration.

Stream surveys.—The stream surveys were conducted by 16 parties operating in forests and parks in various parts of the country. In most cases each party was composed of a biologist in charge, three assistant biologists, and a camp attendant. The equipment of each party consisted of a complete camping outfit, a 2½-ton pick-up truck, and the necessary apparatus for collecting physical, chemical, and geological data on lakes and streams. Because of climatic conditions, the time spent in the field varied greatly in different sections. One of the survey parties in the South, for instance, was in the field for over 8 months while those working at high altitudes in the West were in the field for only 3 months.

The primary purpose of the surveys is to provide an inventory of conditions in each lake and stream which affect the fish population. With this information at hand it will be possible to determine what species of fish is best adapted to each body of water and the number it can support most advantageously. This knowledge is essential for the development of a rational and systematic stocking policy which will make possible the most efficient utilization of forest waters. In the absence of such basic information, fish are frequently planted in waters to which they are not adapted, and in numbers which have no relation to the productive capacity of the stream.

In the East four parties surveyed completely or in part the Great Smoky Mountains National Park in North Carolina and Tennessee, and the following national forests: White Mountain and Green Mountain in New England, George Washington and Monongahela in Virginia and West Virginia, and Pisgah and Nantahala in North Carolina, South Carolina, and Georgia. The 12 parties operating in the Rocky Mountain region covered completely or in part the Glacier National Park and the following national forests: Wyoming, Challis, Sawtooth, Cache, Wasatch, Rapaho, Humboldt, Santa Fe, Carson, Coconino, Apache, Tonto, Crook, Mono, Inyo, Sequoia, Shasta, and Klamath.

A report including the important data collected by the survey parties and recommendations for stocking has been prepared for each forest. These reports give brief accounts of the physiography of the region followed by a description of the principal streams and lakes of each watershed, including the physical and chemical characteristics and accessibility. This is followed by a discussion of the biological characteristics of each body of water, including the fishes and other vertebrates, the kinds and relative abundance of food organisms, and the presence of aquatic plants. A summary

of the essential data required for developing a stocking program for each body of water is given in tabulated form, accompanied by stocking recommendations.

STREAM IMPROVEMENT IN CONNECTION WITH THE STREAM SURVEYS

Extensive work in stream improvement was carried on in the national forests in cooperation with the Forest Service. Except in a few instances, the work was done by C. C. C. labor under the supervision of men employed by the Bureau. Since there are few reliable data on which to base an estimate of the true value of stream improvement the work was primarily experimental.

It is evident that the nature of improvements which will prove of most value will vary widely with conditions and for that reason stream conditioning was carried on in as many different types of streams as possible. Very little work has been done heretofore on swift, mountain streams such as are found in the great majority of the national forests. Such streams are obviously more difficult to work with than sluggish, meadow streams which have been the type on which most improvement work has been done in the past. In some instances streams were selected for improvement that offered most serious obstacles to permanent installations for the purpose of subjecting the devices to the most rigorous tests. Preliminary reports indicate that most of these installations have withstood the effects of exceptionally heavy floods with little damage.

At the conclusion of the summer's work a memorandum was prepared on "Methods for the Improvements of Streams." This memorandum contains a discussion of the principles of stream improvement and detailed descriptions of the devices which have been found to be of greatest value.

INVESTIGATIONS IN THE ROCKY MOUNTAIN REGIONS IN THE INTEREST OF FISH STOCKING

The Bureau's investigations in this region by Dr. A. S. Hazzard are almost solely directed toward the betterment of sport fishing. Commercial fishing at present is restricted to the taking of whitefish in certain States and to limited sucker and carp fisheries in others. Angling (principally for trout) is becoming increasingly popular with the resident and tourist, while in the majority of areas the fishing is rapidly deteriorating. The rapid development of new roads and trails and the accompanying increase in angling have necessitated extensive stocking and environmental improvement.

Investigations during the past 4 years have demonstrated the need for intensive studies of the life histories of the principal native game and forage fish in order to determine their requirements. Such studies should indicate the types of waters to which they are best suited and how their environment can be improved. Similar studies of so-called "obnoxious" fishes such as the chub, bony-tail squawfish, and sculpin are also necessary in order to learn the extent of their competition with game fishes or their unsuspected value to these fishes. This information is necessary before intelligent efforts toward control can be made, if such control proves desirable.

Prior to 1934, the following investigations were undertaken: Survey of Grand Teton National Park, Wasatch Forest studies, initiation of Glacier National Park survey of food production in mountain streams and lakes, life history studies of the cutthroat trout and Rocky Mountain whitefish, also miscellaneous investigations in cooperation with various State fish and game departments.

During 1934, investigations in this region were expanded as a result of funds allotted from the Public Works Administration. Nine survey parties were assigned to the Intermountain States to study waters of certain national parks and forests. The following results were obtained:

The survey of Glacier Park was completed, thus furnishing the basis for a management plan which will maintain fishing in this popular recreation area. Valuable contributions to our knowledge of mountain lakes and streams will also be made as a result of this study. Papers by specialists concerning the physical and chemical conditions, aquatic plants, plankton, and fishes are in preparation for publication.

The survey party assigned to Idaho studied certain waters in the Salmon, Sawtooth, and Challis Forests. As a result of these studies detailed plans for stocking and otherwise improving the lakes and streams of this popular area have been furnished. The apparent failure of previous plantings of smelt and eastern landlocked salmon was demonstrated by this investigation. Also a planting in 1931 of pond weed seed (*P. pectinatus*) in Pettit and Stanley Lakes was shown to have failed. Numerous gill-net sets in Redfish, Pettit, Alturas, and Stanley Lakes indicated a great abundance of suckers and squawfish but no trout, other than a few native Dolly Varden. It is apparent, therefore, that the heavy plantings of various species made have failed to benefit the lakes. Studies of the life history of the squawfish and suckers should be made in order to effect their control.

In Utah a survey of the Cache Forest and certain waters of the Wasatch and Ashley Forests revealed the need for stream improvement and indicated that more intelligent planting would produce better fishing. Detailed plans for this work have been prepared based on data collected.

The survey party also cooperated with the Bureau of Animal Industry and the State fish and game department in determining the effects of the proposed copper-sulphate treatment in trout streams. A concentration of 1:500,000 was used for 24 hours in East Canyon Creek. The results indicated that wherever the concentration was strong enough to kill the objectionable snails it was deadly to trout, though not harmful to the other fish food.

In cooperation with the United States Forest Service, experiments in the fertilization of small, high mountain lakes were initiated. Since soil and water analyses of these lakes and a highly productive trout lake (Fish Lake) demonstrated a deficiency in carbonates, phosphates, and nitrates in the former, known quantities of these substances were introduced to determine the practicability of fertilization. Aquatic plants and fish food organisms from Fish Lake were also introduced. Detailed quantitative studies of chemistry and food

supply made prior to these alterations will be repeated later to determine the results.

The survey party in Nevada studied the streams and lakes in a portion of Humboldt Forest. Although difficult of access, these waters are becoming more heavily fished each year and stocking plans for them are in great demand. The superiority of eastern brook and cutthroat trout over the rainbow for these high waters was clearly proved by results of the survey. The need for pool improvement was also demonstrated.

In Arizona the survey party examined the waters of the Coconino, Tonto, Crook, Coronado, Apache, and Tusan Forests, and a portion of the San Carlos Indian Reservation. The enormous area covered was due to the scarcity of water in this region and the numerous roads. However, this very scarcity enhances the value of the public fishing water and is resulting in strenuous efforts of the fish and game commission and Forest Service to maintain the angling. The need for more systematic stocking, better regulation, stream-improvement work, and erosion control was demonstrated by the survey.

Portions of the Santa Fe and Carson Forests were studied by the New Mexico party. These areas are used extensively for recreation and the fishing has been declining in spite of the excellent work of the fish and game commission. The survey's results indicated a need for stricter regulation of fishing or more frequent planting of legal-sized fish. Recommendations for additional stream and lake improvement were also made.

The Colorado survey party worked in close cooperation with the Forest Service in carrying out the program for improvement of fishing in the Arapaho, Roosevelt, Pike, and San Isabel Forests. Recommendations for stream improvement and rearing pond construction were acted upon immediately during the past season. Detailed plans are being prepared which should assist in the stocking of these waters.

The Wyoming survey party studied a portion of the Wyoming Forest. The lower waters in this area, which are accessible by road, are heavily fished. It is the desire of the fish and game commission and the Forest Service to stock the higher waters according to their requirements in order to induce the anglers to fish there, thereby relieving the strain on the lower waters. Because of transportation difficulties only a limited number of waters could be studied this season. However, sufficient information was secured to enable some generalizations covering the species best adapted, and the intensity of planting which should be carried on in this region.

COOPERATION WITH THE FOREST SERVICE IN STREAM AND LAKE IMPROVEMENT

The West Fork of Rock Creek in the Deer Lodge Forest in Montana was selected as an experimental stream-improvement project. Careful counts of the fish and fish-food population were made in a section of this stream. Sixty-eight structures of various types were then installed. An accurate, large-scale map showing bottom types, location of counts, plant beds, and structure location was made. Careful studies in the future will be carried out to determine the benefit of the improvement work in this stream.

In each of the other seven States of this region the Bureau detailed an investigator to assist the Forest Service in stream-improvement work. It was his duty to prepare plans for the streams designated by the Forest Service and to furnish technical supervision. These men worked in 30 national forests in the region and planned 5,604 improvements, of which approximately half were completed this season. This work was recommended for 153.8 miles of stream, or approximately 36 structures per mile. The major part of the construction was accomplished by the C. C. C., although some was financed out of other Forest Service funds.

In order to determine the effectiveness of various devices in different types of streams, each structure was tagged and notes were taken concerning conditions before and after installation. A check of the work following the spring run-off should indicate the most successful structures for the streams of this region.

In addition to stream improvement, a limited amount of lake alteration was planned by these workers. Raising the outlet to increase the area and installing brush covers to furnish shelter for fish were carried out as experiments in a few selected lakes.

COOPERATION WITH THE UNIVERSITY OF UTAH

A number of advanced students registered for special work under the Bureau's direction have been pursuing certain basic fishery investigations. Life-history studies of several species of fish are in progress. A comparative study of some Utah trout streams and studies of the rate of food recovery in streams affected by floods are also being made. The collection of fishes which is being built up at the University Museum will be of great value both for the instruction of students and for future studies in this region by the Bureau or other investigators.

The impetus given to the study of fishes and their environment should aid greatly in the solution of the many problems confronting the agencies concerned with maintaining the sport fishing in this region.

INVESTIGATIONS IN INTERIOR WATERS

The extensive activities of the unit under the direction of Dr. M. M. Ellis have been made possible not only by the cooperative work of the various members of the staff, but also because of the cooperation of the Corps of Engineers of the United States Army, the Tennessee Valley Authority, and Division of Fish Culture of the United States Bureau of Fisheries. Active work was maintained throughout the year at the Columbia (Mo.) laboratories and at the substation, Fort Worth, Tex., and for considerable portion of the year on the floating laboratory, U. S. Quarterboat 348 and at the Neosho (Mo.) hatchery.

During the past 12 months the activities have followed three major lines.

POLLUTION STUDIES (F. P. 41 PROJECT)

During the course of the pollution studies streams and other bodies of water have been examined in 21 States, and a detailed survey of the

Mississippi River with reference to the effects of pollutant effluents on that stream has been made from Cairo, Ill., to Grand Rapids, Minn. The Mississippi River was chosen as presenting typical problems of a major stream receiving a variety of pollution effluents. Studies on other streams in different States have been made with reference to specific pollution problems as well. The effluents from some 15 types of industries representing over 80 industrial plants have been studied both in the field and in the laboratory. In addition, mine wastes, particularly from coal, lead, copper, iron, zinc, and arsenic mines, have been analyzed and assayed as pollutants. Oil pollution of fresh water has been given detailed attention in the States of Kansas, Oklahoma, Texas, and Louisiana, particularly during the months of October to January, inclusive; and these findings correlated with oil pollution already noted in our field work elsewhere.

Municipal effluents and garbage pollutants were given specific attention in the St. Louis, Davenport, and Minneapolis-St. Paul areas. During November a field party made an extensive study of the best sugar industry effluents in Nebraska, Colorado, and Kansas, and other field parties have studied the sulphur pollution in Louisiana, limestone wastes in Indiana, natural alkali pollution in North Dakota and Montana. All of this pollution work has had as a major objective the determination of: (1) The effects of these substances on fish life, particularly the major types of fresh-water fishes; (2) the effects of these substances on fish food, specifically those insects, crustaceans, and mollusks, which constitute a large part of the food of fresh-water fishes; and on the aquatic bacterial flora, particularly those forms concerned with the utilization of organic wastes and those forms which are part of the food chain series in fresh waters; and (3) the chemical and physical balances of the various substances occurring naturally in many of the fresh-water streams of the United States.

From these studies standards of water purity have been determined for fish life and measurements of pollution hazards made which can be applied quite generally to the fresh-water streams of the United States. New and practical methods for measurement of pollution hazards, relative toxicity, and importance of pollution effluents have been perfected in the course of this work. The work completed shows definitely the size of the task involved and the needs for continuation of studies along these lines.

POWER DAM LAKES AND OTHER IMPOUNDED WATERS

Studies of power dam lakes and other impounded waters have been carried forward, particularly in connection with the Tennessee Valley Authority projects in the Norris Dam area and in connection with the Fort Peck Dam which is being constructed in Montana. As a result of these studies, recommendations have been made to the Tennessee Valley Authority, and certain special lateral lakes are now being established in the Norris Dam area to insure the maintenance of fish food and young fish supplies in these areas.

The Fort Peck area was given a general survey and preparations made for the continuation of this work this spring, when the construction will have proceeded further.

In conjunction with the Corps of Engineers, United States Army, survey work, analyses, and experimental tests were made back of several of the new dams which have been constructed as part of the Mississippi River channel project and recommendations made concerning the effect of these constructions on water conditions, fish food, and other fisheries interests in the impounded waters held by these dams.

MUSSEL PROPAGATION

The mussel propagation experiments which were started at Fort Worth 2 years ago have been continued throughout the past year, and detailed data collected concerning the growth, mortality, and resistance of fresh-water mussels which are being grown under controlled conditions in close quarters in the experimental raceways at the Fort Worth project. The results of these experiments continue to be favorable to this method of artificial propagation, especially for certain species. As a corollary of the work at Fort Worth, a new shipping container has been devised, making it possible to ship live mussels long distances with very little loss.

INDEPENDENT ACTIVITIES OF THE FISHERIES BIOLOGICAL LABORATORIES

WOODS HOLE, MASS.

Lack of sufficient funds for the operation of the Woods Hole laboratory prevented the Bureau of Fisheries from providing the customary facilities for independent investigations at this laboratory.

Robert A. Nesbit, assisted by William C. Neville, ran parallel series of tests of methods of marking scup by insertion of tags into the coelomic cavity, in which it was found that the most successful method consisted of inserting flat celluloid strips $1\frac{1}{8}$ by $\frac{1}{8}$ inches. With this method 8 percent of the tags were lost during the first 3 weeks, and none was lost during the following 12 weeks of observation. By developing a marking technique in which losses are small and definitely ascertained, it becomes possible not only to trace the migrations of this species but also to determine what portion of the mortality is due to commercial fishing and what portion is due to natural causes, two items that are fundamental in formulating a conservative policy.

BEAUFORT LABORATORY

The Beaufort (N. C.) laboratory was severely damaged to the extent of over \$10,000 by the tropical hurricane of September 16, 1933. It was possible, however, with funds provided by the Public Works Administration, to restore the buildings and equipment by the following spring and continue operations for the propagation of the diamondback terrapin, and investigations concerning oil pollution and serious natural enemies of oysters in the South Atlantic and Gulf region. With additional funds provided by the Public Works Administration, a bridge has been constructed to connect the laboratory with the mainland, which will increase the protection of the Bureau's property in case of fire and tropical storms and will enable the gen-

eral public to inspect the station's exhibits of marine animals, fishing equipment, and methods developed for increasing the propagation of oysters and the diamondback terrapin.

Research.—Operation of the Beaufort laboratory was continued throughout the year under the direction of Dr. H. F. Prytherch and furnished facilities for the study of fishery problems of the South Atlantic and Gulf region. The chief investigations conducted here by the Bureau's staff, as reported in detail elsewhere, consist of (1) physiological effects of Louisiana crude petroleum and oil-well brines on the oyster; (2) studies of a protozoan parasite of the oyster associated with mortalities in Louisiana and Virginia; and (3) cytological studies of the ova and germ-cell cycle in two species of shrimp. The laboratory serves as headquarters for oyster investigations in the South Atlantic and Gulf States and, in cooperation with the conservation departments of North Carolina, South Carolina, and Florida, directs extensive operations for the rehabilitation of public oyster beds with funds provided by the Federal Emergency Relief Administration. Field investigations have also been made of the serious destruction of oysters in Apalachicola Bay, Fla., by a flat-worm parasite, *Stylochus inimicus*, commonly known as the "wafer" or "leech."

Laboratory facilities for marine research have been furnished to 27 investigators from other institutions who were engaged in the following studies: Dr. H. V. Wilson, University of North Carolina, the biology of sponge cells, with particular reference to the fine cytoplasmic reticula formed in metamorphosing larvae and regenerative masses; Irene Bolick, University of North Carolina, dissociation and reunion of sponge cells of several local species; Dr. A. S. Pearse, Duke University, assisted by H. W. Hatsel and E. D. Huntley, ecological studies of estuarine animals; Dr. Bert Cunningham, Duke University, effect of temperature on embryonic development of the diamondback terrapin; Dr. Paul L. Risley, State University of Iowa, transplantation of reproductive organs and sex differentiation in the terrapin; Dr. George G. Scott, City College of New York, comparative histology of marine fishes; Dr. and Mrs. B. G. Chitwood, United States Department of Agriculture, identification and distribution of free living marine nematodes; Dr. W. C. George, University of North Carolina, studies of chordate blood and regeneration in the ascidian *Styela*; Dr. Hoyt S. Hopkins, New York University, respiration in marine mollusks in relation to oxygen tension; Dr. W. E. Bullington, Randolph Macon College, studies of marine ciliates; Dr. Lowell E. Noland, University of Wisconsin, morphology and taxonomy of ciliate protozoa, particularly those of the order Peritricha; L. Lyndon Williams, Rensselaer Polytechnic Institute, tissue regeneration in hydroids and related forms; Reinard Harkema, Duke University, parasites of rodents occurring on the coast of North Carolina; Dr. Hugh H. Darby, Bartol Research Foundation, regeneration of chelipeds in *Alpheus* and *Uca*; G. Robert Lunz, Jr., Charleston Museum, Stomatopoda of the Beaufort region; Dr. J. Paul Visscher, Western Reserve University, rare species of barnacles.

In cooperation with the Federal Emergency Relief Administration, laboratory facilities were provided for the employment of eight college-trained women on a research project covering studies of the com-

plete life cycle of the black skimmer, *Rhynchops nigra*. The project was carried out under the supervision of Dr. Alice L. Brown, with the assistance of M. Waynick, A. L. Bason, A. S. Sherrill, B. Arnold, and L. Jarrett, of the North Carolina College for Women; I. Bolick, University of North Carolina; and R. L. Collie, of the North Carolina State Museum. Particular attention was given to the development of the embryo, growth, and morphology of the mandibles and the differentiation of sex. Material for these studies was obtained from the rookeries of the black skimmer on the Georgia coast and from several located in Pamlico Sound, N. C.

The facilities of the station were also utilized by the United States Chemical Warfare Service for tests of wood preservatives, and by the Bureau's Division of Fishery Industries for experiments on the durability of net twines treated with different preservatives. Cooperative tests were continued with the Woolsey Paint Co. in respect to the antifouling value of copper paints, and with the Tropical Paint & Oil Co. on the suitability of bakelite varnishes for marine use.

Terrapin culture.—Operations for the propagation of the diamondback terrapin, conducted in cooperation with the Division of Fish Culture, were unusually successful during the summer of 1934, when a record hatch of 12,446 young was obtained. In addition to this number approximately 1,000 eggs, embryos, and young terrapin were supplied to investigators from Duke University and the State University of Iowa for studies of development and possible control of sex in this species. During the spring of this year 10,445 young terrapin from the 1933 brood were distributed in southern coastal waters in cooperation with the North Carolina Department of Conservation, the South Carolina State Board of Fisheries, and the Georgia Department of Fish and Game.

The system of propagation has recently been modified and improved by simulating natural conditions, so as to permit hibernation of the young terrapin during the winter months in protected outdoor pens. This arrangement has not only reduced the cost and labor of their care in a heated rearing house, as required previously, but has considerably increased the number surviving the critical 9-month period from hatching to liberation. It was first tried with the 1933 brood, of which over 98 percent were grown to a suitable size for distribution. Though record low temperatures occurred during the hibernation period, the total losses of young terrapin in this brood amounted to only 179, as compared with losses ranging from 700 to over 3,000 during previous years.

The annual production of young diamondback terrapin at the Beaufort station hatchery since 1930 has been as follows: 1930, 5,778; 1931, 5,500; 1932, 11,086; 1933, 10,060; 1934, 10,445. These operations have been successful from a practical standpoint since 1912 and have provided over 70,000 terrapin for breeding purposes and restocking of coastal areas. Since the consumption and market demand for terrapin has increased considerably in the last 2 years, it is advisable that artificial propagation of this species be continued and expanded, and an effort made to develop State hatcheries for this purpose.

PROPAGATION AND DISTRIBUTION OF FOOD FISHES, FISCAL YEAR 1935¹

By GLEN C. LEACH, *Chief*, and M. C. JAMES, *Assistant Chief*, Division of
Fish Culture

CONTENTS

| | Page |
|---|------|
| Introduction..... | 401 |
| Species propagated..... | 402 |
| Production..... | 404 |
| Construction activities..... | 404 |
| Cooperation with other conservation agencies..... | 405 |
| Salvage operations..... | 406 |
| Assignments of fish eggs to States, Territories, and foreign countries..... | 407 |
| Transfer of eggs between stations..... | 408 |
| Station output..... | 409 |
| Egg collections..... | 415 |
| Notes on operations..... | 416 |
| Commercial species..... | 416 |
| Great Lakes species..... | 416 |
| Marine species..... | 417 |
| Anadromous species of the Atlantic coast..... | 417 |
| Game-fish propagation..... | 418 |
| Rocky Mountain territory..... | 418 |
| New England stations..... | 420 |
| Combination trout and pond-fish stations..... | 421 |
| Pond-fish stations..... | 423 |
| Mississippi River territory..... | 424 |
| Aquarium..... | 425 |
| Distribution operations..... | 425 |

INTRODUCTION

The output of fish and eggs from the Federal hatcheries again showed an upward trend in 1935, in contrast with the curtailment imposed during the fiscal year 1934. The resumption of work at one of the marine hatcheries, made possible by a slight increase in appropriations, together with enlargements and improvements effected by relief funds and labor at other hatcheries, was responsible for a distinct increase in production.

While there still appears to be uncertainty and confusion as to the best means of conserving and perpetuating other forms of wildlife, the operation of hatcheries remains at hand as a proven means of maintaining fish life particularly in interior waters. It is fully recognized that some species under certain conditions have failed to respond to fish-cultural methods, and it is also admitted that stocking may lack effectiveness when it is conducted on haphazard hit-or-miss lines. The fact remains, however, that each game fish planted

¹ Administrative Report No. 22, Appendix IV to the Report of the U. S. Commissioner of Fisheries for 1935. Approved for publication, Jan. 14, 1936.

in lake or stream is an increment to the normal fish resources of that body of water, and eggs of a commercial species fertilized, hatched, and planted as a byproduct of a commercial fishery are an asset of that commercial fishery.

Public acceptance of this fact has resulted in a flood of recommendations and requests urging the general establishment of new hatcheries in all parts of the country. Aside from the fact that funds are not available, either for construction or maintenance of such projects, it is essential to understand that natural limitations very frequently make it desirable to concentrate the propagation work in a few of the larger more efficient hatcheries. The conception of numerous local rearing ponds for the purpose of "finishing off" the hatchery products is sound, but these are much more modest projects than are visualized by the average sportsmen's group interested in securing a Federal or State hatchery for that section. However, the system of Federal hatcheries is as yet incomplete as far as certain sections of the country are concerned, and these sections will be inadequately stocked until some new hatcheries are built.

The following tables and statistics are presented as a listing, in part, of the raw material available for anglers of 1936, 1937, and subsequent years.

SPECIES PROPAGATED

Forty-three different species were propagated in sufficient numbers to warrant a separate listing, and several miscellaneous forms were handled in small numbers largely in an experimental way. Included in the output were pollock which were not propagated during the previous year. No pink or humpback salmon were produced, but a moderate number of cisco or lake herring eggs were secured and hatched. Many of the species shown on the following list are planted directly from the hatcheries, private applicants not participating in this distribution.

CATFISHES (SILURIDÆ):

- Catfish (*Leptops olivaris*).
- Spotted catfish (*Ictalurus punctatus*).
- Horned pout (*Ameiurus nebulosus*).

CARP (CYPRINIDÆ): Common carp (*Cyprinus carpio*).

BUFFALOFISH (CATOSTOMIDÆ): Common buffalo (*Ictiobus sp.*).

SHAD AND HERRING (CLUPEIDÆ):

- Shad (*Alosa sapidissima*).

SALMONS, TROUTS, AND WHITEFISHES (SALMONIDÆ):

- Common whitefish (*Coregonus clupeaformis*).
- Chinook, king, or quinnat salmon (*Oncorhynchus tshawytscha*).
- Chum salmon (*Oncorhynchus keta*).
- Coho salmon, silver salmon (*Oncorhynchus kisutch*).
- Red salmon, sockeye, or blueback salmon (*Oncorhynchus nerka*).
- Steelhead salmon (*Salmo gairdneri*).
- Atlantic salmon (*Salmo salar*).
- Landlocked salmon (*Salmo sebago*).
- Rainbow trout (*Salmo shasta*).
- Black-spotted trout, redthroat trout (*Salmo lewisi*).
- Loch Leven trout (*Salmo levenensis*).
- Lake trout, Mackinaw trout (*Cristivomer namaycush*).
- Brook trout (*Salvelinus fontinalis*).

GRAYLINGS (THYMALLIDÆ): Montana grayling (*Thymallus montanus*).

PIKES (ESOCIDÆ): Pike and pickerel (*Esox sp.*).

SUNFISHES (CENTRARCHIDAE):

- Crappie (*Pomoxis annularis* and *P. sparoides*).
- Largemouth black bass (*Micropterus salmoides*).
- Smallmouth black bass (*Micropterus dolomieu*).
- Rock bass (*Ambloplites rupestris*).
- Warmouth bass, goggle-eye (*Chaenobryttus gulosus*).
- Bluegill sunfish (*Lepomis incisor*).
- Green sunfish (*Lepomis cyanellus*).
- Redbreasted bream (*Lepomis auritus*).
- Red-eared sunfish (*Lepomis heros*).
- Common sunfish (*Lepomis gibbosus*).
- Rio Grande perch (*Herichthys cyanoguttatus*).

PERCHES (PERCIDAE):

- Pike perch (*Stizostedion vitreum*).
- Yellow perch, ringed perch (*Perca flavescens*).

WHITE BASSES (SERRANIDAE):

- White bass (*Rocuss chrysops*).
- White perch (*Morone americana*).

DRUMS (SCIAENIDAE): Fresh-water drum, lake sheepshead (*Aplodinotus grunniens*).

CODES (GADIDAE):

- Cod (*Gadus callarias*).
- Haddock (*Melanogrammus aeglefinus*).
- Pollock (*Pollachius virens*).

FLOUNDERS (PLEURONECTIDAE): Winter flounder, American flatfish.

MACKEREL (SCOMBRIDAE): Common mackerel (*Scomber scombrus*).

Summary, by species, of the output of fish and fish eggs during the fiscal year ending June 30, 1935

| Species | Eggs | Fry | Fingerlings | Total |
|----------------------|---------------|---------------|-------------|---------------|
| Catfish | 160,000 | | 16,486,300 | 16,646,300 |
| Buffalofish | 191,277,000 | 46,500 | 3,380,900 | 194,804,400 |
| Carp | 87,400,000 | 142,000 | 8,284,950 | 95,826,950 |
| Shad | | 11,420,000 | 3,350 | 11,433,350 |
| Whitefish | 50,000 | 39,239,000 | | 39,289,000 |
| Lake herring (cisco) | 320,000 | 2,320,000 | | 2,640,000 |
| Chinook salmon | | | 22,521,000 | 22,521,000 |
| Chum salmon | | 15,454,300 | | 15,454,300 |
| Silver salmon | 182,700 | | 2,097,000 | 2,279,700 |
| Sockeye salmon | | | 1,860,100 | 1,860,100 |
| Steelhead salmon | 50,000 | 47,300 | 1,153,400 | 1,250,700 |
| Atlantic salmon | 775,000 | | 85,000 | 860,000 |
| Landlocked salmon | | | 138,950 | 138,950 |
| Rainbow trout | 3,356,630 | 15,000 | 9,953,200 | 13,353,730 |
| Blackspotted trout | 3,851,160 | | 22,323,800 | 26,174,960 |
| Lock Leven trout | 9,052,160 | 7,138,745 | 5,721,265 | 21,912,200 |
| Lake trout | 2,306,000 | 1,662,500 | 106,450 | 4,074,950 |
| Brook trout | 10,159,900 | 2,080,000 | 16,299,000 | 28,538,900 |
| Grayling | 1,184,000 | | | 1,184,000 |
| Pike and pickerel | | | 36,900 | 36,900 |
| Crappie | | | 6,967,550 | 6,967,550 |
| Black bass: | | | | |
| Largemouth | | 560,750 | 2,545,250 | 3,106,000 |
| Smallmouth | | 1,147,500 | 239,000 | 1,387,400 |
| Rock bass | | | 236,400 | 236,400 |
| Warmouth bass | | | 36,600 | 36,600 |
| Sunfish | | | 10,603,300 | 10,603,300 |
| Pike perch | 510,075,000 | 5,620,000 | | 515,695,000 |
| Yellow perch | | 451,000,000 | 491,600 | 451,491,600 |
| White perch | | 1,920,000 | | 1,920,000 |
| Freshwater drum | | | 4,400 | 4,400 |
| White bass | | | 34,600 | 34,600 |
| Miscellaneous fishes | | | 2,043,700 | 2,043,700 |
| Mackerel | | 2,123,000 | | 2,123,000 |
| Cod | 1,679,043,000 | | | 1,679,043,000 |
| Haddock | 236,836,000 | | | 236,836,060 |
| Flatfish (flounder) | 356,311,000 | 918,518,000 | | 1,254,829,600 |
| Pollock | 309,304,000 | 95,782,000 | | 405,086,000 |
| Total | 3,381,794,450 | 1,550,246,595 | 133,683,955 | 5,071,725,000 |

PRODUCTION

An increase of approximately 1,800,000,000 in the output of fish and eggs over the previous year's figures was a sharp reversal in the downward trend of production. It is true that more extensive operations with the marine species were largely contributory to the increase, but the output of a number of other species was likewise augmented. Among these were the whitefish, chum salmon, silver salmon, Atlantic salmon, smallmouth bass, sunfish, and yellow perch. During the year only four hatcheries remained entirely out of productive operations, although several others were operated on a restricted basis in comparison with previous years.

There is always interest in the proportion of game fishes included in the total output. For 1935 the strictly game species were propagated to the number of 114,571,550, representing about 2.2 percent of the total. This compares with a larger output of 135,211,900 for 1934. However, the latter year shows the production of fingerlings and larger fish at a larger figure, 133,683,955, an increase of over 7,000,000. This, in part, accounts for the drop in the total output of game fish, since numerical production must inevitably decline when the fish are reared to a greater size before distribution. There is every reason to believe that the increased stocking value of the bigger fish compensates for the smaller numbers planted.

Indulgence in speculative computations will reveal that the output of game fish from Federal hatcheries would yield a prorated catch of 11 fish to each of the estimated 10,000,000 anglers in the country if all fish survived and were taken. Considering that in many sections practically all game fish taken are planted stock, it will be evident that there is little imminence of a "saturation point" in hatchery operations.

CONSTRUCTION ACTIVITIES

At the start of the year there were small unexpended balances in approximately 12 allotments for the repair and reconditioning of fish hatcheries. These allotments had been made by the Public Works Administration during the previous year, and the work was carried to a conclusion during the fiscal year 1935. A considerable proportion of a similar allotment for the continued development of the Leetown, W. Va., experimental hatchery remained available and was expended in part for the construction of bass and trout ponds, remodeling of one of the buildings, and construction of a new dwelling. Early in the year a new Public Works allotment of \$75,000 was received for the construction of a pond-fish hatchery at Harrison Lake, Va., 26 miles southeast of Richmond. Work was started during the winter and at the close of the year approximately 15 acres of ponds were virtually completed and other developments were well along to the point where the establishment could go into active operation during the fall of 1935. Arrangements were being made for the propagation of shad as well as pond fish at this point. Two dwellings as well as various service buildings comprised the structural improvements. The water supply, which is obtained from a lake, is fed through a canal. In general, the construction and improvements effected during the fiscal years 1934 and 1935 can be credited in part for the increased hatchery output attained during the latter year. Throughout 1935 labor was assigned to several of the Bureau's

stations by local F. E. R. A. offices. Lack of funds for materials made it impossible to perform major construction jobs by this means, but upkeep and maintenance work of real value was accomplished.

COOPERATION WITH OTHER CONSERVATION AGENCIES

The perfection of cooperative relationships with other Federal conservation bureaus, and with many of the State fish and game departments has become such a routine administrative matter that it hardly warrants special mention in this report. In view of the stress laid upon this feature, however, it is desirable to cite a few instances as examples of the possibilities of coordination in fish-cultural work.

The nature of the work performed by the National Park Service, Forest Service, the Office of Indian Affairs, the Bureau of Reclamation, and the Bureau of Biological Survey, gives them of necessity an interest in the Bureau's fish-cultural work. The stocking of streams and lakes under the control of the foregoing agencies is an important feature of their administration. Consequently, it has been necessary to maintain closer contact with these establishments, and it is felt that the Bureau has been of real service in numerous instances.

While these cooperative relationships are of more or less standing nature, there have been contacts with the newer emergency organizations which offer prospects of mutual value. The Agricultural Adjustment Administration, insofar as its work covers the utilization of marginal lands, has called upon the Bureau in several instances to plan a program for fish propagation and the provision of angling.

At the close of the year plans were being developed for one project in North Carolina whereby a hatchery would be constructed and turned over to the Bureau for subsequent operation with the object of providing fish for the surrounding territory.

The Bureau has been called upon to make a survey in the Tennessee Valley to aid the Tennessee Valley Authority in developing a program of conserving fishery resources and establishing a hatchery system.

The United States Army Engineers have given sympathetic consideration to the Bureau's recommendations with regard to developments in the Upper Mississippi River area. This has been based upon a hope that the dams comprising part of the 9-foot channel development might be modified so as to provide extensive propagating ponds for the production of fish native to that area.

With reference to the States, arrangements have been made for several additional States to review Federal applications for fish in order to assure that no undesirable species will be planted, and that the stocking will be in conformity with the natural demands and possibilities of the lakes and streams.

The State of Vermont has afforded generous help in a financial way in meeting the costs of operation at the extensive brook trout egg-producing plant maintained by the Bureau in the White Mountain National Forest, N. H.

Tennessee and West Virginia have furnished food for fish held at the Bureau's hatcheries and have handled the distribution of these fish when they were ready for release.

A number of the Western States have continued to supply helpful cooperation in the collecting of trout eggs and are compensated therefor by receiving a proportion of the eggs taken. There has

been a joint operation of trout and shad hatcheries in the State of South Carolina.

The State of Washington donated a site for a trout rearing station which was developed by the Bureau to furnish a supply of fish for the eastern part of the State where we lacked such facilities.

The mutual benefits of these relationships may be exemplified by citing the situation in the State of Indiana. This State has a limited mileage of trout waters but rather than incur the expense of a trout hatchery, it is allotted several hundred thousand trout fry from Federal hatcheries in other States. In return the State utilizes its facilities in distributing practically all of the fish produced at the Federal hatchery at Rochester, Ind.

The machinery through which these plans became effective is the National Planning Council of Federal and State Fisheries Authorities. The several regional meetings of the organization held during the year have brought about results fully justifying its formation.

The Bureau has been less active in sponsoring the establishment of fish rearing ponds or nurseries to be maintained and operated by private sportsmen's organizations. This movement is well developed now and while fish from Federal hatcheries are always available for suitable club nurseries, it is felt that the initiative should be taken by the groups which are to benefit most. We have continued to furnish assignments to nurseries which have been in operation for several years. Special effort has been made to render informational service by supplying technical data upon the problems of fish rearing, and acting in an advisory capacity. In this connection it is possible to be of as great service by pointing out the futility of attempting to rear fish under adverse conditions, as it is to lay out a rearing plant and formulate a program for its operation. The artificial propagation of fish is enough of a technical enterprise to require the supervision of technical men to give best results.

The extent of the cooperative relationships with private clubs operating rearing ponds and nurseries was practically unchanged so far as the number of organizations was concerned. Sixty-seven units received fish in comparison with 62 during the previous year. However, the total number of fish assigned was considerably reduced, allotments amounting to 1,750,000, a reduction of more than a million from the assignments made the previous year. This is attributed to increased rearing facilities at the Bureau's own stations, and the fact that better results are obtained by avoidance of overstocking of private rearing ponds.

SALVAGE OPERATIONS

By the expenditure of a few cents per thousand fish, it is possible to transfer tens of millions of fish which face sure destruction in the overflowed sloughs of the Upper Mississippi River, to living waters where they can survive. It has been the practice each year to salvage these bass, sunfish, crappie, perch and other game and commercial fish, as one of the chief activities in the administration of the Upper Mississippi Wild Life Refuge. The magnitude of the work varies according to river conditions, and in 1935 over 47,000,000 fish were rescued in comparison with 23,873,000 handled the previous year. Only 105,000 were used in restocking other waters, the balance being restored to the river. Fish raised in artificial controlled ponds are now being more widely used for general stocking purposes. Plans

have been prepared for an extensive series of hatchery ponds which will be required when the development of the 9-foot channel will, through the maintenance of constant water levels, have eliminated the necessity of salvaging fish.

Number and disposition of fish rescued, fiscal year 1935

| Locality and species | Delivered to applicants | Restored to original waters | Total number of fish rescued |
|----------------------------|-------------------------|-----------------------------|------------------------------|
| All stations: | | | |
| Black bass | 62,615 | 208,555 | 271,170 |
| Buffalofish | | 4,030,420 | 4,030,420 |
| Carp | | 8,423,000 | 8,423,000 |
| Catfish | 20,350 | 16,613,960 | 16,634,300 |
| Crappie | 348 | 6,963,652 | 6,966,000 |
| Drum | | 4,400 | 4,400 |
| Pike and pickerel | | 36,890 | 36,890 |
| Sunfish | 10,395 | 8,272,465 | 8,282,860 |
| White bass | | 34,600 | 34,600 |
| Yellow perch | | 423,145 | 423,105 |
| Miscellaneous fishes | | 2,043,700 | 2,043,700 |
| Total | 105,668 | 47,056,837 | 47,162,505 |
| Summary by stations: | | | |
| Fairport | | 658,388 | 658,388 |
| La Crosse | 24,764 | 3,372,336 | 3,397,500 |
| Lynxville | 1,990 | 4,445,280 | 4,447,270 |
| Bellevue | | 6,021,650 | 6,021,650 |
| Homer | 43,033 | 8,566,621 | 8,609,704 |
| Marquette | 36,831 | 23,886,494 | 23,922,325 |
| Total | 105,668 | 47,056,837 | 47,162,505 |

NOTE.—Carp listed herein are returned only to the Mississippi River where a commercial fishery for the species exists. None are planted in game fish waters.

ASSIGNMENTS OF FISH AND FISH EGGS TO STATES, TERRITORIES, AND FOREIGN COUNTRIES

The activity of the Bureau in assigning numerous fish and fish eggs to various State fish and game departments may be likened to a wholesale enterprise. These eggs are delivered both from hatchery brood stock and from collections obtained by trapping wild fish. They are the raw material used in operations at the State hatcheries. Many of them are furnished on an exchange basis or under cooperative arrangements whereby the recipients meet part of the costs of collections, while in other instances they are outright donations. The assignments of eggs and fish to the States in 1935, numbering 562,323,860, were more than in 1934 by 188,440,860.

It may be pointed out further that the Bureau also has an "export" business in eggs and fish. Each year there are several requests for advice and assistance in programs of establishing American fishes in foreign countries. Many of these plans do not materialize due to unsuitable conditions abroad. However, some shipments are occasionally furnished by the Bureau, and the table below shows the extent of this activity during 1935. The assignment of trout eggs to Canada was in compensation for Atlantic salmon eggs furnished by the Dominion. The shipment to Venezuela was especially noteworthy in that there was a high rate of survival (80 percent in case of one species) of varieties which are difficult to transport for long distances.

Assignment of fish and fish eggs to State fish commissions, fiscal year 1935

| States and species | Number | States and species | Number |
|--|-----------|---|-------------|
| Arizona: | | New Mexico: | |
| Brook trout..... | 10,000 | Blackspotted trout..... | 250,000 |
| Rainbow trout..... | 35,000 | Chinook salmon..... | 197,000 |
| California: Grayling..... | 5,000 | New York: Rainbow trout..... | 260,000 |
| Colorado: Rainbow trout..... | 102,000 | North Carolina: Rainbow trout..... | 582,000 |
| Connecticut: | | Ohio: | |
| Lake trout..... | 25,000 | Pike perch..... | 510,000,000 |
| Rainbow trout..... | 100,000 | Whitefish fry..... | 37,000,000 |
| Georgia: Rainbow trout..... | 200,000 | Oregon: | |
| Idaho: | | Blackspotted trout..... | 250,000 |
| Blackspotted trout..... | 25,000 | Silver salmon..... | 131,500 |
| Grayling..... | 370,000 | Steelhead salmon..... | 50,000 |
| Rainbow trout..... | 18,600 | Pennsylvania: Rainbow trout..... | 250,000 |
| Maine: | | South Carolina: Rainbow trout..... | 150,000 |
| Atlantic salmon..... | 775,000 | Tennessee: Rainbow trout..... | 550,000 |
| Brook trout..... | 1,078,000 | Utah: | |
| Massachusetts: Rainbow trout..... | 100,000 | Brook trout..... | 101,920 |
| Michigan: | | Grayling..... | 150,000 |
| Brook trout..... | 50,000 | Vermont: | |
| Grayling..... | 150,000 | Brook trout..... | 2,199,550 |
| Montana: | | Rainbow trout..... | 35,000 |
| Brook trout..... | 769,000 | Washington: | |
| Loch Leven trout..... | 1,896,000 | Blackspotted trout..... | 68,000 |
| Black bass..... | 42,870 | Brook trout..... | 167,000 |
| Catfish..... | 65,745 | Loch Leven trout..... | 50,000 |
| Crappie..... | 8,375 | Rainbow trout..... | 358,000 |
| Sunfish..... | 33,320 | West Virginia: Brook trout..... | 550,000 |
| Yellow perch..... | 3,980 | Wyoming: | |
| Nebraska: | | Blackspotted trout..... | 1,250,000 |
| Brook trout..... | 25,000 | Lake trout..... | 50,000 |
| Rainbow trout..... | 293,000 | Total | 582,323,880 |
| Nevada: | | | |
| Brook trout..... | 150,000 | | |
| Rainbow trout..... | 1,008,000 | | |
| New Hampshire: | | | |
| Lake herring..... | 320,000 | | |
| Rainbow trout..... | 25,000 | | |

Shipments of fish and fish eggs to foreign countries, fiscal year, 1935

| Country and species | Eggs | Yearlings, adults | Country and species | Eggs | Yearlings, adults |
|--|-----------|-------------------|----------------------------|-----------|-------------------|
| Canada: Blackspotted trout..... | 1,000,000 | ----- | Puerto Rico: | | |
| Cuba: | | | Rainbow trout..... | 50,000 | ----- |
| Black bass, largemouth..... | ----- | 60 | Catfish..... | ----- | 300 |
| Crappie..... | ----- | 114 | Sunfish..... | ----- | 1,800 |
| Warmouth bass..... | ----- | 128 | Venezuela: | | |
| Panama: | | | Rainbow trout..... | 50,000 | ----- |
| Black bass, largemouth..... | ----- | 325 | Bluegill sunfish..... | ----- | 105 |
| Carp..... | ----- | 11 | Carp..... | ----- | 130 |
| Catfish..... | ----- | 100 | Crappie..... | ----- | 30 |
| Crappie..... | ----- | 25 | Largemouth black bass..... | ----- | 235 |
| Sunfish..... | ----- | 250 | Yellow perch..... | ----- | 75 |
| Yellow perch..... | ----- | 75 | Total | 1,100,000 | 3,761 |

TRANSFER OF EGGS BETWEEN STATIONS

While the interhatchery transfer of fish eggs is more or less a routine administrative matter, the following tabulation is cited to demonstrate the extent of this practice. Many of the eggs so transferred are collected from wild fish at suitable locations, and the maintenance of expensive brood stock is avoided. With a sufficient return of fish to parental waters the wild producing stock maintains itself with adequate abundance.

Transfer of eggs between stations, fiscal year 1935

| Species | Number of eggs | From— | To— |
|-------------------------|------------------------|--------------------------------------|-----------------------------------|
| Blackspotted trout..... | 106,400 | Saratoga, Wyo..... | Bozeman, Mont. |
| | 53,200 | do..... | Hagerman, Idaho. |
| | 104,580 | do..... | Jackson Hole, Wyo. |
| | 53,200 | do..... | Springville, Utah. |
| | 2,000,000 | Yellowstone Park, Wyo..... | Jackson Hole, Wyo. |
| | 250,000 | do..... | Quilcene, Wash. |
| | 100,000 | do..... | Springville, Utah. |
| | 280,000 | Berkshire, Mass..... | LaCrosse, Wis. |
| | 1,100,000 | Berlin, N. H. (national forest)..... | Cape Vincent, N. Y., and suburbs. |
| | 253,825 | do..... | Erwin, Tenn. |
| Brook trout..... | 2,139,700 | do..... | St. Johnsbury, Vt. |
| | 700,000 | do..... | White Sulphur Springs, W. Va. |
| | 500,000 | do..... | Wytheville, Va. |
| | 400,000 | Craig Brook, Maine..... | Cape Vincent, N. Y. |
| | 850,000 | do..... | Erwin, Tenn. |
| | 600,000 | do..... | Leetown, W. Va. |
| | 50,000 | do..... | Manchester, Iowa. |
| | 360,700 | do..... | Nashua, N. H. |
| | 550,000 | do..... | White Sulphur Springs, W. Va. |
| | 500,000 | do..... | Wytheville, Va. |
| | 451,250 | Creede, Colo..... | Crawford, Nebr. |
| | 301,840 | do..... | Duluth, Minn. |
| | 150,300 | do..... | Hagerman, Idaho. |
| | 200,800 | do..... | Manchester, Iowa. |
| | 422,550 | do..... | Saratoga, Wyo. |
| | 100,450 | do..... | Spearfish, S. Dak. |
| | 1,010,400 | do..... | Springville, Utah. |
| | 859,000 | Leadville, Colo..... | La Crosse, Wis. |
| | 62,000 | do..... | Lake Mills, Wis. |
| | 50,000 | Pittsford, Vt..... | Leetown, W. Va. |
| 250,000 | do..... | Nashua, N. H. | |
| Chinook salmon..... | 202,800 | Clackamas, Oreg..... | Leadville, Colo. |
| | 100,000 | Butte Falls, Oreg..... | Clackamas, Oreg. |
| Landlocked salmon..... | 20,000 | Craig Brook, Maine..... | Nashua, N. H. |
| Loch Leven trout..... | 14,299,400 | Ennis, Mont..... | Bozeman, Mont. |
| | 95,000 | Leadville, Colo..... | Eagle Nest, N. Mex. |
| Rainbow trout..... | 183,000 | Ennis, Mont..... | Meadow Creek, Mont. |
| | 100,000 | Bourbon, Mo..... | Aquarium, Washington, D. C. |
| | 200,000 | do..... | Crawford, Nebr. |
| | 200,000 | do..... | Flintville, Tenn. |
| | 400,000 | do..... | Saratoga, Wyo. |
| | 567,000 | Manchester, Iowa..... | Bozeman, Mont. |
| | 209,000 | do..... | Creede, Colo. |
| | 92,000 | do..... | Duluth, Minn. |
| | 104,000 | do..... | Hagerman, Idaho. |
| | 107,000 | do..... | Leadville, Colo. |
| | 319,000 | do..... | La Crosse, Wis. |
| | 350,000 | do..... | Northville, Mich. |
| | 61,000 | do..... | Lake Mills, Wis. |
| | 209,000 | do..... | Spearfish, S. Dak. |
| | 201,000 | Neosho, Mo..... | Bozeman, Mont. |
| | 163,000 | do..... | Crawford, Nebr. |
| | 90,000 | do..... | Hagerman, Idaho. |
| | 100,000 | do..... | Leadville, Colo. |
| | 104,000 | do..... | Spearfish, S. Dak. |
| | 50,000 | Pittsford, Vt..... | Leetown, W. Va. |
| 58,000 | Springville, Utah..... | Hagerman, Idaho. | |
| 100,000 | do..... | Bear Lake, Utah. | |
| Steelhead salmon..... | 50,000 | Butte Falls, Oreg..... | Charlevoix, Mich. |

STATION OUTPUT

The following table, showing the producing hatcheries and the output by species of each, carries a total of 77 units of which 43 are main stations and 34 substations. The summary last year showed 83 stations and substations, but for administrative reasons a number of closely related substations such as field stations in the Upper Mississippi River and bass ponds around the Fort Belvoir, Va., station were consolidated as far as reports of output are concerned. Actual changes therefore involved the closure of one main station in Alaska, opening of two in the States, and the actual number of fish producing units was one more than last year, namely, 84. They

are located in 38 States, but State lines are entirely disregarded in distributing the output.

The carp listed in the following table comprise fish handled in rescue work, or eggs fertilized and hatched at stations located in that territory. The planting of carp is confined to waters which already support an important commercial fishery for the species and they are not distributed in other waters.

Stations and substations operated and the output of each, fiscal year 1935

| Stations and substations and species | Eggs | Fry | Fingerlings, yearlings, and adults | Total |
|---|-------------|-------------|------------------------------------|-------------|
| Baird, Calif.: Chinook salmon..... | | | 1,495,100 | 1,495,100 |
| Battle Creek, Calif.: Chinook salmon..... | | | 3,492,000 | 3,492,000 |
| Mill Creek, Calif.: Chinook salmon..... | | | 2,369,100 | 2,369,100 |
| Baker Lake, Wash.: | | | | |
| Blackspotted trout..... | | | 94,000 | 94,000 |
| Brook trout..... | | | 257,500 | 257,500 |
| Loch Leven trout..... | | | 80,500 | 80,500 |
| Rainbow trout..... | | | 48,400 | 48,400 |
| Sockeye salmon..... | | | 25,900 | 25,900 |
| Birdsview, Wash.: | | | | |
| Blackspotted trout..... | | | 294,400 | 294,400 |
| Brook trout..... | | | 180,000 | 180,000 |
| Loch Leven trout..... | | | 85,000 | 85,000 |
| Rainbow trout..... | | | 137,500 | 137,500 |
| Silver salmon..... | | | 413,000 | 413,000 |
| Sockeye salmon..... | | | 404,000 | 404,000 |
| Steelhead salmon..... | | | 603,000 | 603,000 |
| Mount Rainier, Wash.: | | | | |
| Blackspotted trout..... | | | 250,000 | 250,000 |
| Brook trout..... | | | 260,000 | 260,000 |
| Rainbow trout..... | | | 93,850 | 93,850 |
| Berkshire Trout Hatchery, Mass.: | | | | |
| Brook trout..... | | | 173,700 | 173,700 |
| Catfish..... | | | 4,730 | 4,730 |
| Smallmouth black bass..... | | 786,500 | | 786,500 |
| Boothbay Harbor, Maine: | | | | |
| Cod..... | 729,168,000 | | | 729,168,000 |
| Flatfish..... | | 774,000,000 | | 774,000,000 |
| Haddock..... | 120,401,000 | | | 120,401,000 |
| Rozeman, Mont.: | | | | |
| Blackspotted trout..... | 1,001,160 | | 3,408,010 | 4,409,170 |
| Brook trout..... | | | 470,410 | 470,410 |
| Loch Leven trout..... | 6,795,040 | | | 6,795,040 |
| Rainbow trout..... | 102,900 | | 423,700 | 526,600 |
| Ennis, Mont.: | | | | |
| Blackspotted trout..... | | | 308,555 | 308,555 |
| Loch Leven trout..... | 2,257,120 | 7,138,745 | 1,138,500 | 10,534,365 |
| Rainbow trout..... | | | 168,490 | 168,490 |
| Glacier Park, Mont.: | | | | |
| Blackspotted trout..... | | | 1,372,180 | 1,372,180 |
| Grayling..... | 509,000 | | | 509,000 |
| Rainbow trout..... | | | 85,060 | 85,060 |
| Miles City, Mont.: | | | | |
| Black bass, largemouth..... | | | 121,945 | 121,945 |
| Catfish..... | | | 87,530 | 87,530 |
| Crappie..... | | | 53,595 | 53,595 |
| Sunfish..... | | | 91,235 | 91,235 |
| Yellow perch..... | | | 5,805 | 5,805 |
| Cape Vincent, N. Y.: | | | | |
| Brook trout..... | | | 1,700 | 1,700 |
| Lake herring (cisco)..... | 320,000 | 2,320,000 | | 2,640,000 |
| Lake trout..... | 25,000 | 240,000 | 1,400 | 266,400 |
| Black bass, smallmouth..... | | | 73,240 | 73,240 |
| Whitefish..... | | 679,000 | | 679,000 |
| Barneveld, N. Y.: | | | | |
| Brook trout..... | | | 107,100 | 107,100 |
| Lake trout..... | | | 16,000 | 16,000 |
| Landlocked salmon..... | | | 500 | 500 |
| Loch Leven trout..... | | | 2,400 | 2,400 |
| Rainbow trout..... | | | 19,630 | 19,630 |
| Cortland, N. Y.: | | | | |
| Brook trout..... | | | 169,575 | 169,575 |
| Loch Leven trout..... | | | 36,980 | 36,980 |
| Rainbow trout..... | | | 1,610 | 1,610 |
| Watertown, N. Y.: | | | | |
| Brook trout..... | | | 46,630 | 46,630 |
| Lake trout..... | | | 66,050 | 66,050 |
| Loch Leven trout..... | | | 76,395 | 76,395 |
| Rainbow trout..... | | | 1,900 | 1,900 |

PROPAGATION AND DISTRIBUTION OF FOOD FISHES, 1935 411

Stations and substations operated and the output of each, fiscal year 1935—Continued

| Stations and substations and species | Eggs | Fry | Fingerlings, yearlings, and adults | Total |
|---------------------------------------|-------------|-------------|------------------------------------|-------------|
| Cape Vincent, N. Y.—Continued. | | | | |
| Rochester, N. Y.: | | | | |
| Black bass: | | | | |
| Largemouth..... | | | 35,236 | 35,235 |
| Smallmouth..... | | | 22,200 | 22,200 |
| Brook trout..... | | | 400 | 400 |
| Loch Leven trout..... | | | 4,740 | 4,740 |
| Rainbow trout..... | | | 30,000 | 30,000 |
| Clackamas, Oreg.: | | | | |
| Blackspotted trout..... | | | 75,000 | 75,000 |
| Brook trout..... | | | 385,000 | 385,000 |
| Chinook salmon..... | | | 2,543,900 | 2,543,900 |
| Loch Leven trout..... | | | 305,000 | 305,000 |
| Rainbow trout..... | | | 8,000 | 8,000 |
| Silver salmon..... | 1,200 | | 42,000 | 43,200 |
| Big White Salmon, Wash.: | | | | |
| Chinook salmon..... | | | 4,872,000 | 4,872,000 |
| Butte Falls, Oreg.: | | | | |
| Blackspotted trout..... | | | 447,240 | 447,240 |
| Chinook salmon..... | | | 3,825,000 | 3,825,000 |
| Silver salmon..... | 131,500 | | 137,500 | 269,000 |
| Steelhead salmon..... | 50,000 | | 352,000 | 402,000 |
| Little White Salmon, Wash.: | | | | |
| Chinook salmon..... | | | 3,872,180 | 3,872,180 |
| Chum salmon..... | | 199,000 | | 199,000 |
| Craig Brook, Maine: | | | | |
| Atlantic salmon..... | 775,000 | | 85,000 | 860,000 |
| Brook trout..... | 1,069,000 | 30,000 | 700,300 | 1,799,300 |
| Landlocked salmon..... | | | 103,800 | 103,800 |
| Smelt..... | 8,000,000 | | | 8,000,000 |
| Crawford, Nebr.: | | | | |
| Blackspotted trout..... | | | 483,000 | 483,000 |
| Brook trout..... | | | 328,930 | 328,930 |
| Loch Leven trout..... | | | 168,125 | 168,125 |
| Rainbow trout..... | | 15,000 | 603,415 | 618,415 |
| Black bass, largemouth..... | | | 26,400 | 26,400 |
| Catfish..... | | | 5,950 | 5,950 |
| Crappie..... | | | 2,340 | 2,340 |
| Sunfish..... | | | 11,100 | 11,100 |
| Yellow perch..... | | | 56,075 | 56,075 |
| Dexter, N. Mex.: | | | | |
| Black bass, largemouth..... | | | 146,440 | 146,440 |
| Catfish..... | | | 14,000 | 14,000 |
| Crappie..... | | | 800 | 800 |
| Sunfish..... | | | 58,375 | 58,375 |
| Duluth, Minn.: | | | | |
| Brook trout..... | | | 191,000 | 191,000 |
| Lake trout..... | | 1,422,500 | 23,000 | 1,445,500 |
| Loch Leven trout..... | | | 39,000 | 39,000 |
| Pike perch..... | | 5,620,000 | | 5,620,000 |
| Rainbow trout..... | | | 23,500 | 23,500 |
| Whitefish..... | | 1,560,000 | | 1,560,000 |
| Edenton, N. C.: | | | | |
| Black bass, largemouth..... | | | 41,100 | 41,100 |
| Crappie..... | | | 1,200 | 1,200 |
| Sunfish..... | | | 13,200 | 13,200 |
| White perch..... | | 1,920,000 | | 1,920,000 |
| Yellow perch..... | | 6,600,100 | | 6,600,100 |
| Erwin, Tenn.: | | | | |
| Black bass, largemouth..... | | | 1,437 | 1,437 |
| Brook trout..... | | | 428,870 | 428,870 |
| Rainbow trout..... | | | 56,080 | 56,080 |
| Rock bass..... | | | 1,600 | 1,600 |
| Sunfish..... | | | 230,000 | 230,000 |
| Fairport, Iowa: | | | | |
| Black bass, largemouth..... | | 52,500 | 71,200 | 123,700 |
| Buffalofish..... | 110,752,600 | 46,500 | 124,550 | 110,923,650 |
| Carp..... | 7,400,000 | 142,000 | 284,550 | 7,806,550 |
| Catfish..... | | | 20,125 | 20,125 |
| Crappie..... | | | 4,200 | 4,200 |
| Rock bass..... | | | 7,200 | 7,200 |
| Smallmouth black bass..... | | | 15,250 | 15,250 |
| Sunfish..... | | | 115,960 | 115,960 |
| Miscellaneous fishes..... | | | 8,700 | 8,700 |
| Flintville, Tenn.: | | | | |
| Loch Leven trout..... | | | 20,000 | 20,000 |
| Rainbow trout..... | | | 209,650 | 209,650 |
| Rock bass..... | | | 10,500 | 10,500 |
| Sunfish..... | | | 8,000 | 8,000 |
| Fort Belvoir, Va.: | | | | |
| Shad..... | | 11,430,000 | | 11,430,000 |
| Yellow perch..... | | 461,000,000 | | 461,000,000 |

Stations and substations operated and the output of each, fiscal year 1935—Continued

| Stations and substations and species | Eggs | Fry | Fingerlings, yearlings, and adults | Total |
|--------------------------------------|-------------|------------|------------------------------------|-------------|
| Gloucester, Mass.: | | | | |
| Cod | 949,875,000 | | | 949,875,000 |
| Flatfish | 300,452,600 | | | 300,452,600 |
| Haddock | 116,425,400 | | | 116,425,400 |
| Pollock | 309,304,000 | 95,782,000 | | 405,086,000 |
| Hagerman, Idaho: | | | | |
| Blackspotted trout | | | 215,900 | 215,900 |
| Brook trout | | | 384,500 | 384,500 |
| Loch Leven trout | | | 18,300 | 18,300 |
| Rainbow trout | 191,000 | | 789,900 | 980,900 |
| Steelhead salmon | | | 12,900 | 12,900 |
| Salmon, Idaho: | | | | |
| Blackspotted trout | | | 90,000 | 90,000 |
| Brook trout | | | 15,000 | 15,000 |
| Rainbow trout | | | 652,100 | 652,100 |
| La Crosse, Wis.: | | | | |
| Black bass: | | | | |
| Largemouth | | | 152,590 | 152,590 |
| Smallmouth | | | 560 | 560 |
| Brook trout | 75,000 | | 604,250 | 679,250 |
| Carp | | | 720,000 | 720,000 |
| Catfish | | | 297,900 | 297,900 |
| Crappie | | | 487,060 | 487,060 |
| Buffalofish | | | 270,000 | 270,000 |
| Drum | | | 200 | 200 |
| Lake trout | 2,281,000 | | | 2,281,000 |
| Loch Leven trout | | | 180,600 | 180,600 |
| Pike and pickerel | | | 8,100 | 8,100 |
| Rainbow trout | | | 254,700 | 254,700 |
| Sunfish | | | 537,125 | 537,125 |
| White bass | | | 5,100 | 5,100 |
| Yellow perch | | | 14,480 | 14,480 |
| Miscellaneous fishes | | | 630,000 | 630,000 |
| Bellevue, Iowa: | | | | |
| Black bass, largemouth | | | 40,300 | 40,300 |
| Carp | 44,000,000 | | 2,299,000 | 46,299,000 |
| Catfish | | | 647,000 | 647,000 |
| Crappie | | | 501,000 | 501,000 |
| Buffalofish | 77,625,000 | | 1,417,000 | 79,042,000 |
| Freshwater drum | | | 540 | 540 |
| Pike and pickerel | | | 1,485 | 1,485 |
| Sunfish | | | 572,000 | 572,000 |
| White bass | | | 24,150 | 24,150 |
| Yellow perch | | | 3,705 | 3,705 |
| Miscellaneous fishes | | | 515,000 | 515,000 |
| Homer, Minn.: | | | | |
| Black bass, largemouth | | | 129,600 | 129,600 |
| Buffalofish | | | 10,770 | 10,770 |
| Carp | | | 1,601,400 | 1,601,400 |
| Catfish | | | 279,000 | 279,000 |
| Crappie | | | 3,033,400 | 3,033,400 |
| Drum | | | 3,060 | 3,060 |
| Pike and pickerel | | | 7,365 | 7,365 |
| Sunfish | | | 3,599,165 | 3,599,165 |
| White bass | | | 1,810 | 1,810 |
| Yellow perch | | | 383,250 | 383,250 |
| Lake Mills, Wis.: | | | | |
| Black bass: | | | | |
| Largemouth | | | 76,550 | 76,550 |
| Smallmouth | | | 10,650 | 10,650 |
| Brook trout | | | 55,000 | 55,000 |
| Loch Leven trout | | | 57,500 | 57,500 |
| Rainbow trout | | | 46,000 | 46,000 |
| Lynxville, Wis.: | | | | |
| Black bass, largemouth | | | 5,170 | 5,170 |
| Buffalofish | | | 372,500 | 372,500 |
| Carp | | | 687,000 | 687,000 |
| Catfish | | | 1,428,000 | 1,428,000 |
| Drum | | | 600 | 600 |
| Pike and pickerel | | | 2,650 | 2,650 |
| Sunfish | | | 767,500 | 767,500 |
| White bass | | | 600 | 600 |
| Yellow perch | | | 4,900 | 4,900 |
| Miscellaneous fishes | | | 841,000 | 841,000 |
| Marquette, Iowa: | | | | |
| Buffalofish | 3,000,000 | | 1,185,900 | 4,185,900 |
| Black bass, largemouth | | | 54,000 | 54,000 |
| Carp | 36,000,000 | | 2,714,000 | 38,714,000 |
| Catfish | 160,000 | | 13,675,000 | 13,835,000 |
| Crappie | | | 2,733,000 | 2,733,000 |
| Pike and pickerel | | | 17,290 | 17,290 |
| Sunfish | | | 2,935,500 | 2,935,500 |

PROPAGATION AND DISTRIBUTION OF FOOD FISHES, 1935 413

Stations and substations operated and the output of each, fiscal year 1935—Continued

| Stations and substations and species | Eggs | Fry | Fingerlings, yearlings, and adults | Total |
|--------------------------------------|-------------|----------|------------------------------------|-------------|
| La Crosse, Wis.—Continued. | | | | |
| Marquette, Iowa—Continued. | | | | |
| White bass..... | | | 3, 000 | 3, 000 |
| Yellow perch..... | | | 15, 500 | 15, 500 |
| Miscellaneous fishes..... | | | 50, 000 | 50, 000 |
| Rochester, Ind.: | | | | |
| Black bass: | | | | |
| Largemouth..... | | | 88, 815 | 88, 815 |
| Smallmouth..... | | | 4, 790 | 4, 790 |
| Crappie..... | | | 1, 000 | 1, 000 |
| Rock bass..... | | | 3, 000 | 3, 000 |
| Sunfish..... | | | 20 | 20 |
| Yellow perch..... | | | 1, 000 | 1, 000 |
| Lamar, Pa.: | | | | |
| Brook trout..... | | | 23, 875 | 23, 875 |
| Rainbow trout..... | | | 14, 000 | 14, 000 |
| Leadville, Colo.: | | | | |
| Blackspotted trout..... | | | 364, 800 | 364, 800 |
| Brook trout..... | | | 2, 661, 150 | 2, 661, 150 |
| Loch Leven trout..... | | | 135, 000 | 135, 000 |
| Rainbow trout..... | | | 325, 000 | 325, 000 |
| Creede, Colo. | | | | |
| Blackspotted trout..... | | | 498, 000 | 498, 000 |
| Brook trout..... | 3, 809, 000 | | 1, 837, 750 | 5, 646, 750 |
| Loch Leven trout..... | | | 175, 000 | 179, 000 |
| Rainbow trout..... | 75, 000 | | 1, 008, 000 | 1, 083, 000 |
| Eagle Nest, N. Mex.: | | | | |
| Brook trout..... | | | 42, 000 | 42, 000 |
| Rainbow trout..... | | | 210, 500 | 210, 500 |
| Leetown, W. Va.: | | | | |
| Brook trout..... | | | 275, 530 | 275, 530 |
| Loch Leven trout..... | | | 26, 000 | 26, 000 |
| Rainbow trout..... | | | 210, 270 | 210, 270 |
| Smallmouth black bass..... | | 10, 000 | | 10, 000 |
| Louisville, Ky.: | | | | |
| Black bass: | | | | |
| Largemouth..... | | 210, 000 | 21, 610 | 231, 610 |
| Smallmouth..... | | 133, 000 | 5, 040 | 138, 040 |
| Crappie..... | | | 215 | 215 |
| Rock bass..... | | | 4, 250 | 4, 250 |
| Sunfish..... | | | 20, 400 | 20, 400 |
| Mammoth Spring, Ark.: | | | | |
| Black bass: | | | | |
| Largemouth..... | | 17, 000 | 326, 300 | 343, 300 |
| Smallmouth..... | | 210, 000 | 38, 250 | 248, 250 |
| Rock bass..... | | | 123, 600 | 123, 600 |
| Sunfish..... | | | 27, 300 | 27, 300 |
| Manchester, Iowa: | | | | |
| Brook trout..... | | | 186, 650 | 186, 650 |
| Loch Leven trout..... | | | 49, 500 | 49, 500 |
| Rainbow trout..... | | | 134, 870 | 134, 870 |
| Rock bass..... | | | 7, 500 | 7, 500 |
| Smallmouth black bass..... | | | 10, 590 | 10, 590 |
| Nashua, N. H.: | | | | |
| Brook trout..... | | | 188, 100 | 188, 100 |
| Landlocked salmon..... | | | 11, 000 | 11, 000 |
| Loch Leven trout..... | | | 8, 000 | 8, 000 |
| Rainbow trout..... | | | 20, 575 | 20, 575 |
| Catfish..... | | | 10, 000 | 10, 000 |
| Smallmouth black bass..... | | | 700 | 700 |
| Neosho, Mo.: | | | | |
| Black bass, largemouth..... | | | 39, 440 | 39, 440 |
| Catfish..... | | | 1, 700 | 1, 700 |
| Crappie..... | | | 24, 900 | 24, 900 |
| Rainbow trout..... | 853, 000 | | 5, 410 | 858, 410 |
| Rock bass..... | | | 53, 300 | 53, 300 |
| Sunfish..... | | | 77, 330 | 77, 330 |
| Yellow perch..... | | | 400 | 400 |
| Bourbon, Mo.: Rainbow trout..... | 1, 036, 000 | | | 1, 036, 000 |
| Natchitoches, La.: | | | | |
| Black bass, largemouth..... | | | 55, 735 | 55, 735 |
| Sunfish..... | | | 71, 755 | 71, 755 |
| Warmouth bass..... | | | 3, 600 | 3, 600 |
| Tishomingo, Okla.: | | | | |
| Black bass, largemouth..... | | | 77, 880 | 77, 880 |
| Catfish..... | | | 16, 200 | 16, 200 |
| Crappie..... | | | 106, 600 | 106, 600 |
| Rock bass..... | | | 29 | 29 |
| Sunfish..... | | | 116, 655 | 116, 655 |
| Warmouth bass..... | | | 24, 900 | 24, 900 |
| Yellow perch..... | | | 5, 600 | 5, 600 |

Stations and substations operated and the output of each, fiscal year 1935—Continued

| Stations and substations and species | Eggs | Fry | Fingerlings, yearlings, and adults | Total |
|--------------------------------------|---------------|--------------|------------------------------------|---------------|
| Northville, Mich.: | | | | |
| Black bass: | | | | |
| Largemouth..... | | | 2, 275 | 2, 275 |
| Smallmouth..... | | | 51, 000 | 51, 000 |
| Brook trout..... | | | 336, 300 | 336, 300 |
| Loch Leven trout..... | | | 270, 665 | 270, 665 |
| Rainbow trout..... | | | 365, 200 | 365, 200 |
| Steelhead salmon..... | | 47, 300 | | 47, 300 |
| Orangeburg, S. C.: | | | | |
| Black bass, largemouth..... | | | 258, 400 | 258, 400 |
| Catfish..... | | | 310 | 310 |
| Crappie..... | | | 1, 040 | 1, 040 |
| Shad..... | | | 3, 350 | 3, 350 |
| Sunfish..... | | | 160, 340 | 160, 340 |
| Warmouth bass..... | | | 7, 970 | 7, 970 |
| Pittsford, Vt.: | | | | |
| Blackspotted trout..... | | | 15, 000 | 15, 000 |
| Brook trout..... | | | 44, 700 | 44, 700 |
| Rainbow trout..... | | | 15, 900 | 15, 900 |
| Put in Bay, Ohio: | | | | |
| Pike perch..... | 510, 075, 000 | | | 510, 075, 000 |
| Whitefish..... | 50, 000 | 37, 000, 000 | | 37, 050, 000 |
| Quinalt, Wash.: | | | | |
| Blackspotted trout..... | | | 234, 520 | 234, 520 |
| Brook trout..... | | | 121, 010 | 121, 010 |
| Chinook salmon..... | | | 57, 800 | 57, 800 |
| Silver salmon..... | 50, 000 | | 1, 513, 500 | 1, 563, 500 |
| Sockeye salmon..... | | | 1, 183, 800 | 1, 183, 800 |
| Duckabush, Wash.: Chum salmon..... | | 13, 993, 300 | | 13, 993, 300 |
| Quilcene, Wash.: | | | | |
| Blackspotted trout..... | | | 250, 000 | 250, 000 |
| Brook trout..... | | | 59, 000 | 59, 000 |
| Chum salmon..... | | 1, 262, 000 | | 1, 262, 000 |
| Sockeye salmon..... | | | 250, 000 | 250, 000 |
| Steelhead salmon..... | | | 185, 500 | 185, 000 |
| National Forest of N. H.: | | | | |
| Brook trout..... | 2, 000, 000 | 2, 050, 000 | 489, 600 | 4, 539, 600 |
| Landlocked salmon..... | | | 635 | 635 |
| St. Johnsbury, Vt.: | | | | |
| Black bass, smallmouth..... | | | 9, 765 | 9, 765 |
| Landlocked salmon..... | | | 23, 100 | 23, 100 |
| San Marcos, Tex.: | | | | |
| Black bass, largemouth..... | | 176, 000 | 151, 140 | 327, 140 |
| Crappie..... | | | 5, 275 | 5, 275 |
| Sunfish..... | | | 46, 600 | 46, 600 |
| Fort Worth, Tex.: | | | | |
| Black bass, largemouth..... | | | 58, 000 | 58, 000 |
| Catfish..... | | | 3, 700 | 3, 700 |
| Crappie..... | | | 12, 800 | 12, 800 |
| Sunfish..... | | | 76, 900 | 76, 900 |
| Saratoga, Wyo.: | | | | |
| Blackspotted trout..... | | | 349, 470 | 349, 470 |
| Brook trout..... | | | 1, 157, 165 | 1, 157, 165 |
| Loch Leven trout..... | | | 229, 060 | 229, 060 |
| Rainbow trout..... | | | 121, 400 | 121, 400 |
| Spearfish, S. Dak.: | | | | |
| Black bass, largemouth..... | | | 14, 575 | 14, 575 |
| Blackspotted trout..... | | | 130, 200 | 130, 200 |
| Brook trout..... | | | 563, 350 | 563, 350 |
| Loch Leven trout..... | | | 715, 000 | 715, 000 |
| Rainbow trout..... | 50, 000 | | 945, 550 | 995, 550 |
| Springville, Utah: | | | | |
| Black bass, largemouth..... | | | 2, 110 | 2, 110 |
| Blackspotted trout..... | | | 640, 000 | 640, 000 |
| Brook trout..... | 101, 900 | | 249, 200 | 351, 100 |
| Loch Leven trout..... | | | 1, 514, 440 | 1, 514, 440 |
| Rainbow trout..... | 1, 068, 630 | | 427, 220 | 1, 495, 850 |
| Bear Lake, Utah: | | | | |
| Brook trout..... | | | 498, 100 | 498, 100 |
| Rainbow trout..... | | | 71, 420 | 71, 420 |
| Sockeye salmon..... | | | 263, 000 | 263, 000 |
| Marion, Ala.: | | | | |
| Black bass, largemouth..... | | | 172, 850 | 172, 850 |
| Sunfish..... | | | 201, 400 | 201, 400 |
| Tupelo, Miss.: | | | | |
| Black bass, largemouth..... | | 50, 500 | 181, 000 | 231, 500 |
| Sunfish..... | | | 225, 900 | 225, 900 |

PROPAGATION AND DISTRIBUTION OF FOOD FISHES, 1935 415

Stations and substations operated and the output of each, fiscal year 1935—Continued

| Stations and substations and species | Eggs | Fry | Fingerlings, yearlings, and adults | Total |
|--------------------------------------|------------|-------------|------------------------------------|-------------|
| Valdosta, Ga.: ¹ | | | | |
| Warm Springs, Ga.: | | | | |
| Black bass, largemouth..... | | 3,750 | 174,350 | 178,100 |
| Catfish..... | | | 2,970 | 2,970 |
| Crappie..... | | | 115 | 115 |
| Sunfish..... | | | 417,650 | 417,650 |
| Warmouth bass..... | | | 125 | 125 |
| White Sulphur Springs, W. Va.: | | | | |
| Black bass, largemouth..... | | 51,000 | 550 | 51,550 |
| Brook trout..... | | | 1,926,675 | 1,926,675 |
| Loch Leven trout..... | | | 381,630 | 381,630 |
| Rainbow trout..... | 221,000 | | 525,300 | 746,300 |
| Rock bass..... | | | 7,025 | 7,025 |
| Sunfish..... | | | 39,025 | 39,025 |
| Woods Hole, Mass.: | | | | |
| Flatfish..... | 35,859,000 | 144,518,000 | | 180,377,000 |
| Mackerel..... | | 2,123,000 | | 2,123,000 |
| Wytheville, Va.: | | | | |
| Black bass: | | | | |
| Largemouth..... | | | 7,900 | 7,900 |
| Smallmouth..... | | | 7,670 | 7,670 |
| Brook trout..... | | | 786,360 | 786,360 |
| Catfish..... | | | 1,725 | 1,725 |
| Rainbow trout..... | 795,000 | | 893,900 | 1,688,900 |
| Rock bass..... | | | 18,400 | 18,400 |
| Sunfish..... | | | 74,900 | 74,900 |
| Yellowstone Park, Wyo.: | | | | |
| Blackspotted trout..... | 2,850,000 | | 11,767,000 | 14,617,000 |
| Grayling..... | 675,000 | | | 675,000 |
| Moose, Wyo.: | | | | |
| Blackspotted trout..... | | | 1,386,000 | 1,386,000 |
| Brook trout..... | | | 92,200 | 92,200 |

¹ No output due to construction work.

NOTE.—Figures for "eggs" denote eggs planted or transferred to agencies outside of Bureau. Eggs transferred to other Bureau stations are shown separately by table on page 409.

EGG COLLECTIONS

As would be expected, the increased output of fish was based upon a larger collection of the raw material, namely, fish eggs. The attached tables give comparison of the 1935 and 1934 collections indicating in which species variations occurred.

Comparison of egg collections, fiscal years 1934 and 1935

| Species | 1934 | 1935 | Species | 1934 | 1935 |
|-------------------------|------------|------------|----------------------|---------------|---------------|
| Shad..... | 17,104,000 | 5,280,000 | Grayling..... | 2,118,400 | |
| Whitefish..... | 12,000,000 | 54,895,000 | Pike perch..... | 840,920,300 | 519,075,000 |
| Chinook salmon..... | 28,323,000 | 29,919,000 | Yellow perch..... | 6,450,000 | 14,000,000 |
| Chum salmon..... | 12,370,000 | 21,138,000 | White perch..... | 1,400,000 | 2,400,000 |
| Humpback salmon..... | 158,000 | | Cod..... | 1,037,262,000 | 1,753,129,000 |
| Silver salmon..... | 2,148,000 | 3,398,000 | Haddock..... | 291,754,000 | 284,483,000 |
| Sockeye salmon..... | 9,157,000 | 3,326,000 | Pollock..... | | 523,622,000 |
| Steelhead salmon..... | 905,000 | 1,352,000 | Winter flounder..... | 1,036,626,000 | 1,375,129,000 |
| Landlocked salmon..... | 35,000 | | Mackerel..... | 8,592,000 | 2,800,000 |
| Rainbow trout..... | 25,770,253 | 21,365,920 | Lake herring..... | | 5,620,000 |
| Blackspotted trout..... | 28,946,550 | 21,206,000 | Carp..... | | 83,250,000 |
| Loch Leven trout..... | 29,329,624 | 30,372,000 | Buffalofish..... | | 214,046,000 |
| Lake trout..... | 1,850,120 | 2,684,000 | | | |
| Brook trout..... | 23,331,758 | 29,265,500 | Total..... | 3,315,039,305 | 4,981,655,420 |

NOTES ON OPERATIONS

COMMERCIAL SPECIES

Pacific salmons.—While there was a moderate increase in the production of the less important species of Pacific salmon, the silver and chum, the output of chinook and sockeyes dropped. The distribution of the latter in particular was but a small fraction of the normal production of preceding years. This was due to cessation of operations in Alaska where sockeye eggs have been obtainable in large numbers. In addition, the run of sockeyes at the Quinault, Wash., station was negligible. This station is the main field of activity with this species outside of Alaska.

In the Columbia River territory operations proceeded in approximately normal scope, but a light run of fish in the Big and Little White Salmon Rivers curtailed the output of chinook salmon. Effort was made to augment the water supply at the main station (Clackamas, Oreg.) by digging a well without success. Some new rearing tanks were constructed. At the Butte Falls, Oreg., substation serving the Rogue River territory, chinook egg collections were good but the operations with steelhead trout were unsatisfactory due to the immaturity of the eggs when the fish entered the traps.

The Salmon, Idaho, substation was operated on a part-time basis, chiefly for the handling of rainbow trout eggs.

Total egg collections at the Birdsvew, Wash., station were below last year's take. A number of improvements, including completion of a dwelling, work upon Grandy Creek and the water-supply system, traps, etc., as well as minor jobs, were effected by relief labor. Sockeye salmon marking experiments were continued.

The Baker Lake substation was operated during the summer and fall months only and propagated only trout. A rock-slide arising from flood conditions blocked the water supply and Civilian Conservation Corps labor was obtained to remove it. The Mount Rainier, Wash., substation furnished about 600,000 trout for park waters. Effort is being made to develop egg-collecting fields in several lakes in the national park.

The most noteworthy item at the Quinault, Wash., station was the greatly reduced run of sockeye salmon. However, sufficient eggs of various species were obtained to utilize to capacity the rearing facilities of the station.

The Duckabush and Quilcene, Wash., substations as usual concentrated on the handling of chum salmon. Repeated damage to racks and traps by high water affected operations, but in spite of this a large collection of late-run chum salmon eggs was obtained.

In the California field the collection of chinook salmon eggs was concentrated at the Mill Creek substation. Of the 4,000,000 taken, 1,500,000 were transferred to the Baird station for incubation.

GREAT LAKES SPECIES

Comparable to last year, propagation of commercial species of the Great Lakes was confined to the Duluth, Minn., Put in Bay, Ohio, and Cape Vincent, N. Y., stations. The former succeeded in collecting a moderate number of lake trout and whitefish eggs during the regular open season. The quality of them was rather poor, due to lateness of the collecting season. A considerable number of game-

trout eggs were hatched, part of the fish being used in stocking rearing ponds constructed by the United States Forest Service. The State of Minnesota furnished a limited number of pike-perch eggs during the spring.

At the Cape Vincent, N. Y., station eggs of whitefish, lake trout, and lake herring, yielding slightly over 3,000,000 fry, were obtained from local fishermen. The main activities here are now concerned with the propagation of game fish, particularly smallmouth bass, and the administration of the three substations where trout are reared. This work has continued to be of increasing importance. The pike-perch hatchery on Lake Champlain at Swanton, Vt., was operated jointly by the States of Vermont and Connecticut.

The work at the Put in Bay, Ohio, station was closely tied in with that of the State hatchery. The Bureau collected 51,000,000 whitefish eggs, which were hatched and distributed from the State hatchery. During the spring over 500,000,000 pike-perch eggs were collected by the Bureau's force, with State facilities being used for hatching. The dock at the Bureau's station has disintegrated to the point where it is no longer safe for use. The Bureau's vessel *Shearwater* is now operated by the State, serving both hatcheries.

MARINE SPECIES

At the Boothbay Harbor, Maine, station one of the steam pumps and boilers was replaced by an electric pump, adding greatly to the efficiency and economy of operations. A new hot-water system was also required and installed as a result of this change. The flatfish propagation was conducted in the usual manner. However, a larger number of brood fish yielded a smaller number of eggs than during the previous year. Operations with the cod and haddock consisted of fertilizing and planting on the spawning grounds eggs taken in connection with commercial fisheries. The 729,000,000 cod eggs and the 120,000,000 haddock eggs thus handled were really a byproduct recovery.

At Woods Hole, Mass., a new water-supply suction line was installed. Due to shortage of funds, no cod were propagated, but 144,000,000 flounder fry were produced, together with a limited number of mackerel.

The Gloucester, Mass., station was reopened early in July after a 9-month closure. Considerable work was required in overhauling boilers, pumps, fish-cultural equipment, and in renovating the buildings, all of which were deteriorated. Everything was placed in readiness to permit propagation of pollock during the winter months. Part of the one-half billion eggs secured were planted on the spawning grounds directly from the fishing boats, the balance being incubated at the hatchery. In the spring the work of planting fertilized cod, haddock, and flounder eggs was taken up. Spawntakers placed aboard commercial fishing vessels encountered unusually favorable conditions, and large numbers of eggs were obtained. This station is of chief value in maintaining a local inshore fishery.

ANADROMOUS SPECIES OF THE ATLANTIC COAST

The propagation of commercial species native to east coast rivers was formerly much more extensive, but changing conditions accompanying development and industrialization have rendered such work

futile in many sections. Accordingly, the propagation of Atlantic salmon was confined to the Craig Brook, Maine, station, while work with the shad was centered on the Potomac River at the Fort Belvoir, Va. (formerly Fort Humphreys) station. It was again possible to secure 1,000,000 Atlantic salmon eggs from Canada, and these were distributed to various State hatcheries in Maine, with the exception of a limited number hatched at the Federal hatchery at Craig Brook. This procedure was followed for greater economy and efficiency in distribution.

The Fort Belvoir, Va., hatchery bettered last season's production of shad and resumed propagation of yellow perch with a very satisfactory production of over 400 million fry. This work does not conflict with shad operations but is rather supplementary and involves slight extra cost. Some minor improvements were made to the station buildings and grounds.

At the Edenton, N. C., station, yellow perch and white perch, as well as shad, were handled. Eight and a quarter million eggs of the former species were obtained. The commercial shad fishery from which the shad eggs are obtained was not particularly successful, and the output of fry amounted to only three and a half million. The station also propagated pond fish.

The Orangeburg, S. C., station again operated a small shad hatchery on the Edisto River as a cooperative project with the State of South Carolina.

GAME-FISH PROPAGATION

Artificial propagation of commercial species is largely a replenishment process, supplementing natural production. This is true with the game varieties also but in addition virtual extinction of certain species in certain sections is forestalled by the hatchery activities.

The Bureau's game fish hatcheries were operated with the conception that they were producing the basis of a great national recreation, and the areas devoted to recreation—the national parks and forests—were the subject of special attention. The numerical output of these varieties has been discussed elsewhere. The rising cost of the meats commonly used as fish food has constituted the most serious problem confronting this phase of our work. The requirements for larger fish necessitate feeding for long periods and rigid economy is essential if the game fish output is to be kept up in size and numbers. Distribution of the fish affords one of the best fields for cost reduction; and by using State facilities, as well as requiring private applicants to handle their allotments, production has been maintained at a high level. The findings of the Bureau's biologists engaged in aquicultural investigations have been applied at the hatcheries wherever practicable. The following summaries covering the work of the individual stations will, however, illustrate more fully the nature of the year's work.

ROCKY MOUNTAIN TERRITORY

At the close of the year all arrangements were being put into effect for the consolidation of the Rocky Mountain and Pacific coast territories to be administered by a regional director located at Seattle, Wash.

The activities at the Yellowstone Park substation were administered from the Salt Lake City headquarters. Inasmuch as the work covering the midsummer months extends into parts of two fiscal years, the data furnished here cover the summer of 1934. The total collection of black-spotted trout eggs exceeded 38,000,000, surpassing any previous season. Over 4,000,000 grayling eggs were also collected. The destruction of penned fish and pens by bears was halted by simply keeping dogs in the vicinity of the racks. The larger percentage of the output was retained for restocking in Yellowstone Park or in other national parks. A new water supply line was installed.

At the Springville, Utah, station the brood stock of rainbow trout yielded over 2,000,000 eggs, an increase of almost half a million over last year. A 1½-acre bass pond was completed with relief labor.

The Bear Lake, Utah, substation handled about 1,300,000 trout eggs. Some losses were experienced in rearing the fry and fingerlings. This was traced to excess nitrogen in the water and was corrected by aeration. The station hatched about 250,000 landlocked sockeye salmon known as "silver trout", for planting in local waters.

The Spearfish, S. Dak., station received over 10,000 hours of relief labor which was utilized in overhauling the pond system, renewing the water supply system, and numerous improvements to buildings and grounds which have greatly improved the efficiency and appearance of the plant. Approximately 7,300 cans of trout fingerlings were distributed providing for ample stocking of the territory served by this hatchery.

The Saratoga, Wyo., station was able to accomplish little in the way of major improvements, although a small Civilian Conservation Corps detail was used in building a wall along the creek through the station grounds. Fish-cultural work was conducted at a normal level with a good take of black-spotted trout eggs from station brood stock.

In the Colorado field, the Leadville station has partially completed a new rearing pond. This station together with the substation at Creede cooperated with private parties in the collection of trout eggs, and the two establishments handled over 12½ million eggs. The substation at Crystal Lake was used chiefly for rearing purposes. A 1½-acre earth rearing pond was constructed at Creede with Civilian Conservation Corps labor. Flood waters destroyed a dam in a storage reservoir at the Leadville station, affecting the power plant water supply.

At the Hagerman, Idaho, station a cold-storage room was installed, power being generated on the premises. Some work was done on overhauling the living quarters. While there was considerable loss on eggs shipped in from other points, operations as a whole were satisfactory and the rapid growth of the fish at this station makes it a valuable unit.

The Salmon, Idaho, substation was operated during the summer months for the rearing of fish derived from egg collections at Williams Lake. The take of over 3,500,000 eggs at this point exceeded all previous records.

Civilian Conservation Corps labor was available at the Crawford, Nebr., station for repairs and improvements to bass and trout ponds on the Fort Robinson Reservation.

The extensive operations in the Montana territory were highly successful.

Through assignments of relief workers and an assignment from the Civilian Conservation Corps, a limited amount of construction and repair work was carried on at the Bozeman, Mont., station. Such work consisted of repairing and graveling the roadways about the station and public highway, renewing the concrete floor in the hatchery building and equipping same with new troughs, constructing a fish-food-preparation house, and completing one apartment of a duplex cottage.

Distribution of trout eggs and fingerlings was made in the usual territory consisting of 15,022,430 eggs and 4,378,900 fingerlings, 1,946,674 more fingerlings being distributed during the year than the previous fiscal year.

At the Ennis substation repair work and construction was carried on by the regular personnel and with temporary help. A new concrete floor was put in the Meadow Creek hatchery; also 20 cedar troughs with wooden standards were set up in the building. At Blaine Springs 400 feet of supply pipe was put in, and a rearing pond 100 by 12 feet was constructed.

The Loch Leven egg collections totaled 25,511,655, which was approximately 3,000,000 less than the previous year. However, the take was ample to meet all requirements. The rainbow take during the spring was a few thousand larger than the previous year.

The Blaine Springs hatchery has been in operation for over a year and has proven satisfactory in every detail. With the water of a 53° temperature, a rapid growth among the fingerling trout is encouraged.

At the Miles City, Mont., substation relief workers were used in repairing a leak in the dam of Lake Keogh, which necessitated an excavation of gravel 60 feet long, 8 feet deep, and 5 feet wide and replacing the gravel with dirt that would pack and form a compact mass.

The total number of fingerlings harvested during the fall ranked among previous years, but the most outstanding feature of the harvest was that 121,525 fingerling bass were distributed.

The Dexter, N. Mex., station is the only Federal hatchery in this territory which is solely concerned with the production of pond fish. No major construction was performed, but considerable work was applied to improving ponds, particularly the drainage. Unfavorable weather during the spawning season resulted in a subnormal hatch of fry. Maintenance of adequate water levels in the ponds is always a problem at this station.

Among the miscellaneous activities was the establishment of rearing pools on a site near Spokane, Wash., donated by the State. Under a cooperative agreement with the adjacent State hatchery, a Bureau employee has been detailed to care for the trout during the rearing period.

The Jackson Hole, Wyo., station received fingerlings by transfer and also hatched eggs. During the spring months mortality was very heavy and other difficulties were experienced. However, a goodly number of fingerlings was available for stocking waters in the Teton National Park.

NEW ENGLAND STATIONS

The Nashua, N. H., station functioned in a normal manner and achieved an average production of three species of trout and land-locked salmon.

The Hartsville, Mass., station collected nearly 500,000 brook-trout eggs, sufficient for its own requirements and also providing for shipments of several hundred thousand to other stations. The cooperative collection of smallmouth bass fry from waters supplying reservoir lakes was most successful. Seven hundred and sixty-eight thousand fry were collected and distributed, 50 percent being at the disposal of the State of Connecticut in conformity with the existing agreement.

The East Orland, Maine, station took 6,860,000 brook-trout eggs from station stock. This was below the figure for the previous year due to the fact the brood stock is being reduced. Three hundred thousand landlocked salmon eggs were furnished by the State of Maine, and by exchange Canadian Atlantic salmon eggs to the number of 1,000,000 were obtained. An agreement was consummated with the State of Maine for the joint distribution of trout and salmon, and the output was moved at negligible cost to the Bureau. A few smelt eggs were handled. Some improvement and repair work was performed upon the ponds and water-supply system, including a start on the construction of two circular rearing ponds.

By continuation of the use of Civilian Conservation Corps labor and relief labor, the long-range development program for the York Pond, N. H., brook-trout station was brought nearer to completion. The nature of the work accomplished is too extensive for description here but one of the most important features was a new bulkhead at the outlet of York Pond and a new canal system for the power house. Realignment of rearing ponds, landscaping, and numerous minor jobs were prosecuted. Due to the disturbances arising from construction, the collections of brook-trout eggs were slightly less than last year. A cooperative arrangement in effect with the State of Vermont has resulted in financial aid to the program of rearing and distribution carried on from this station. The St. Johnsbury, Vt., station has been used as an auxiliary principally in handling the eggs destined for shipment from York Pond. The St. Johnsbury station enjoyed unusual success in the rearing of landlocked salmon during the year. The hatchery building was repainted.

The Pittsford, Vt., station, in addition to its strictly experimental work, produced some brook-trout eggs for shipment and distributed a considerable number of fingerling trout. It is rearing several varieties not native to New England, including grayling and golden trout, and has a stock of the rare Sunapee, or white trout.

COMBINATION TROUT AND POND-FISH STATIONS

At the Northville, Mich., station eight daphnia ponds were constructed for the production of bass food. Relief labor was used to complete the bass ponds and grade the station grounds. A satisfactory production of trout and bass was obtained. The Conservation Departments of Indiana, Michigan, and Ohio cooperated in the distribution of the output in their respective States. Five rearing projects handling about 450,000 trout were operated under the direction of this station.

At the Neosho, Mo., station the fall and summer collections of pond fish were approximately as during the previous year. The spring collections of largemouth bass were only 10 percent of those of

last year, due to almost constant rains, which broke up the bass schools, and to cool weather.

Ichthyophthirius developed at Neosho and became epidemic for the first time in 10 years. Selective breeding of rainbow trout has been continued along the lines previously reported. The distribution of trout was adequate to meet all requirements for that section.

At the Bourbon, Mo., substation the production of eyed rainbow eggs was approximately the same as that of last season and the hatching and rearing of the rainbow trout has been most successful. This substation is operated upon a cooperative basis by this Bureau in connection with the owners of the property. By agreement the Bureau receives 90 percent of all rainbow spawn.

Owing to a reduction in the brood stock the output of the Manchester, Iowa, station was considerably reduced from that of last year. Three thousand brood rainbow trout yielded 2,496,000 eggs. Consignments totaling 2,000,000 eyed eggs were shipped to applicants, State or Government establishments. The smallmouth black bass and rock bass produced a fair yield of eggs, and 10,860 large fingerling smallmouth bass, and 7,500 rock bass were distributed.

The Leetown, W. Va., station was engaged in numerous experimental activities which are discussed in the Report of the Division of Scientific Inquiry. However, over 600,000 fry, fingerlings, and adults were distributed. The hatch of trout eggs was poor due to carbon dioxide in the water supply. Other diseases were noted during the year and appropriate remedial measures applied. The station produced its first hatch of bass and has experimented with the rearing of the fry in trout ponds. Construction performed included the claying of the bottoms of seven bass ponds and the construction of one new pond. Concrete bottoms were placed in two long trout raceways. A dwelling for the director was completed during the year, and work was started on a fish culturist's cottage.

The new Lamar, Pa., station was in active operation although no fish were hatched. Trout fry transferred from other stations were reared, producing over 50,000 good-sized fish for distribution. Eight circular rearing ponds and four 50-foot raceways were constructed, and property lines were fenced. Several hundred feet of road were constructed.

The Flintville, Tenn., station has reduced its rainbow brood stock with consequent reduction in egg take. Surplus fish were turned over to the State. Production of pond fish was limited due to the fact that wild brood stock from Reelfoot Lake had not become fully acclimated.

Operations at the Erwin, Tenn., station were of a routine nature. Cooperative arrangements for the rearing of trout were in effect with sportsmen's organizations and the State.

At the White Sulphur Springs, W. Va., station painting of the station buildings was completed. A smaller number of eggs was secured from rainbow brood stock than was the case the previous year, but the percentage of the hatch was higher. Almost 100,000 pond fish were produced, an increase of 36,000. The station hatched trout eggs for the State of West Virginia, as has been done previously.

The total distribution of all species from the Wytheville, Va., station was slightly below the previous year, but as a whole the year's work was satisfactory. Large numbers of eggs and fish were assigned to the States of Virginia and North Carolina.

POND-FISH STATIONS

The hatcheries engaged in the culture of the warm-water or pond fishes are essentially a series of artificial ponds stocked with brood fish. The real fish-cultural work, therefore, lies in the management applied to these ponds. The output is somewhat at the mercy of weather and other uncontrollable factors; and, aside from fluctuations in production, a recital of station activities is largely a statement of routine.

At the Warm Springs, Ga., station reconditioning of buildings and grounds was continued with relief labor or funds. An old building was razed and the material used for other repairs. Two retaining walls were erected, and work was performed upon the ponds. The output of fish was slightly less than last year.

At Valdosta, Ga., construction work prohibited any production of fish. The construction activities included a reinforced concrete dam and necessary structures at the outlet giving complete control of the drainage wells. Eight thousand feet of ditches were excavated to facilitate drainage of the lake. Four holding pools were created by erecting dykes and concrete outlets in the bed of the lake. Other work comprised improvement of the holding shed and its water supply, fencing, roads, etc.

At the San Angelo, Tex., station, operated under the jurisdiction of the San Marcos, Tex., station, a dwelling, shop, and garage were completed and two ponds developed to the point where fry could be transferred to them for rearing.

At the Orangeburg, S. C., station two small mud ponds were completed. A number of the buildings were painted. The production of fish approximating a total of 500,000 was more than 100,000 in excess of last year's output. The cooperative shad work on the Edisto River gave very limited results, however.

At the Marion, Ala., station results in fish-cultural work this year were gratifying both as to the size and numbers of fish produced. A total of 434,280 fingerlings were delivered to applicants and 116,500 bass were planted in station rearing ponds. No fish were handled in the advanced fry stage this year. During the year 8 additional ponds were constructed covering a total of 28.73 acres, increasing the pond system to 51 ponds with a total water area of 100 acres. Another flowing well was brought in during the fall, and there is now sufficient water to supply ponds additional to those already constructed. Construction of buildings was continued during the first half of the year and there were completed during this period, one brick veneer dwelling, one large frame barn, two small barns, and one shipping shed. A large log cabin at Perry Lake, on the east side of the property and 1 mile from the office, was nearly completed. A telephone line was constructed to Marion, a distance of 7 miles.

Adverse weather conditions at the Louisville, Ky., station, with other unusual factors, caused a decrease in the output of smallmouth bass fry. During the year 133,000 smallmouth bass fry and 5,070 fingerlings were distributed, as well as 210,000 largemouth bass fry and 21,611 fingerlings. The latter were produced in increased numbers.

Considerable excavation of earth from upper portions of ponds, the uncovering of water supply and sewer lines for leaks; also other con-

struction work such as pouring a 4-inch concrete wall through embankments of ponds to prevent crayfish from tunneling from one pond to another, the widening of pond A, and other necessary work have been accomplished with labor furnished by the local welfare agency.

The Fairport, Iowa, station has become of increasing importance in the production of pond fish required to meet applications formerly filled with rescued fish. Relief labor was employed for a complete renovation of the pond system and other activities incidental to maintenance.

At the Tishomingo, Okla., substation the pond area has been increased during the past year by workers from the Federal Emergency Relief Administration so that at present there are 33 acres in ponds. Four miles of standard grade 40-foot highway was constructed to the station by the State. Interest is centered on the production of channel catfish at this hatchery and to date the work has been fairly successful.

At the Natchitoches, La., station the production has been approximately the same as during the previous year. Experiments are being conducted with fertilizers, their use and quantity per acre in order to procure data to determine the best methods of quantity production of pond fishes. The pond area of this station is being increased through the work of the Federal Emergency Relief Administration and this new area will be in production during the spring of 1936.

MISSISSIPPI RIVER TERRITORY

Fisheries matters in the Upper Mississippi Wild Life and Fish Refuge, and hatchery activities in the North Central States were supervised from the La Crosse, Wis., headquarters. This station hatched successfully over a million game trout and operated several bass ponds at the hatchery and in the adjacent territory. In addition, it supervised the rescue or salvage work in the refuge (discussed elsewhere) and cooperated in the collection of lake trout eggs in the Illinois waters of Lake Michigan. Three large trout rearing units in the national forests were stocked or supervised from this point, and the work also included supervision and stocking of private co-operating rearing ponds at 39 different locations in nearby States.

The construction of dams as a part of the 9-foot channel project for the Mississippi River is greatly changing conditions in this vicinity. Several semicontrolled ponds used for rearing bass have already been rendered useless and certain parts of the territory formerly covered by rescue crews are being flooded. As construction progresses these effects will be extended. The Bureau has developed plans for construction of propagating ponds adjacent to the new dams but no funds have been obtainable for the carrying out of these plans.

The Homer, Minn., substation affords shop and repair facilities for overhauling equipment and was used for this purpose as well as a center for rescue work.

The new Rochester, Ind., hatchery, in its second year of production, received the benefit of relief labor throughout the greater part of the year. This resulted in creation of additional ponds to the extent of more than 6 acres and the partial completion of another pond. Minor jobs required for the completion of the dwellings and other buildings were attended to. Due to drought conditions, the water supply was

inadequate to maintain the ponds throughout the season and the fish were distributed in July and August, thereby reducing the output of larger fish.

At the Lake Mills, Wis., substation there was a production of trout and pond fish commensurate with the facilities available.

AQUARIUM

The aquarium located under the main lobby in the Department of Commerce Building is maintaining its reputation as one of Washington's popular points of interest for visitors. The numerous organizations which gather for conventions in Washington, as a rule, include the aquarium in their itinerary of sightseeing.

The general exhibits were maintained in much the same manner as during the previous year and the display of trout has continued to be a point of particular interest. However, from time to time during the year it has been the practice to introduce more novel exhibits, one of which was a 500-pound green turtle. Another interesting novelty was a pair of piranha—the small man-eating fish of the Amazon Basin.

The number of specimens and species on display have varied from time to time, but as a rule from 1,500 to 2,000 individual specimens representing more than 50 different varieties are shown. Replacement specimens were obtained from several of the Bureau hatcheries, the majority being obtained from the La Crosse, Wis., territory.

The aquarium also was of great service in preparing and forwarding foreign shipments of live fish and eggs which were made to Puerto Rico, Panama, and Venezuela.

As usual the display of model hatching apparatus was kept in operation for the purpose of illustrating the methods followed in incubating various types of eggs.

DISTRIBUTION OPERATIONS

The large number of fish distributed in the waters of the national forests and national parks was to a large extent handled by those agencies themselves. In a number of instances sportsmen's clubs cooperated in distributing fish throughout a large territory. This was particularly evident in connection with the operations at the Spearfish, S. Dak., station. The trend is constantly toward the use of trucks for distribution purposes, and the fish cars are used only for the longer hauls.

The distribution cars engaged in the delivery of fish to applicants this year made 65 trips and carried an average of 250 pails per trip. The cars traveled 45,435 paid miles and 8,958 free miles. Detached messengers made 49,665 paid miles and 4,725 free miles in delivering fish to applicants. Truck deliveries during the year aggregated 40,351 miles. As formerly, the Bureau received free transportation and reduced rates from a number of railroads.

It is the Bureau's policy to notify applicants when their fish are ready for delivery and ask them to call at the hatchery to receive them. After the applicants have been notified and sufficient time has elapsed, if they do not call for their fish the same are planted in suitable waters. This course has been made necessary owing to shortage in distribution funds.

Summary, by States, of the distribution of fish, fiscal year 1935

| State and species | Number | State and species | Number |
|---|--------------------|----------------------------|---------------|
| Alabama: | | Maryland—Continued. | |
| Largemouth black bass..... | 206, 675 | Sunfish..... | 41, 000 |
| Loch Leven trout..... | 10, 000 | Rainbow trout..... | 17, 300 |
| Rainbow trout..... | 3, 250 | Massachusetts: | |
| Sunfish..... | 263, 925 | Brook trout..... | 275, 900 |
| Arkansas: | | Catfish..... | 1, 800 |
| Largemouth black bass..... | 197, 350 | Flatfish..... | 144, 518, 000 |
| Smallmouth black bass..... | 32, 550 | Mackerel..... | 2, 123, 000 |
| Rock bass..... | 52, 750 | Pollock..... | 95, 782, 000 |
| Sunfish..... | 21, 100 | Loch Leven trout..... | 3, 000 |
| California: Chinook salmon..... | 5, 681, 000 | Lake trout..... | 1, 750 |
| Colorado: | | Landlocked salmon..... | 2, 000 |
| Blackspotted trout..... | 816, 800 | Rainbow trout..... | 8, 375 |
| Brook trout..... | 5, 088, 400 | Smallmouth black bass..... | 341, 800 |
| Loch Leven trout..... | 302, 000 | Michigan: | |
| Rainbow trout..... | 1, 573, 000 | Brook trout..... | 389, 700 |
| Connecticut: | | Lake trout..... | 174, 000 |
| Brook trout..... | 8, 100 | Loch Leven trout..... | 21, 000 |
| Rainbow trout..... | 16, 700 | Largemouth black bass..... | 13, 300 |
| Smallmouth black bass..... | 7, 500 | Smallmouth black bass..... | 29, 850 |
| Delaware: Largemouth black bass..... | 3, 375 | Pike perch..... | 600, 000 |
| Florida: Sunfish..... | 500 | Rainbow trout..... | 293, 600 |
| Georgia: | | Sunfish..... | 9, 740 |
| Brook trout..... | 91, 000 | Whitefish..... | 1, 500, 000 |
| Catfish..... | 3, 910 | Yellow perch..... | 1, 500 |
| Largemouth black bass..... | 281, 300 | Minnesota: | |
| Rainbow trout..... | 40, 240 | Brook trout..... | 243, 900 |
| Sunfish..... | 269, 300 | Catfish..... | 279, 000 |
| Idaho: | | Carp..... | 1, 601, 000 |
| Blackspotted trout..... | 216, 000 | Buffalofish..... | 10, 770 |
| Brook trout..... | 378, 000 | Crappie..... | 3, 033, 000 |
| Largemouth black bass..... | 6, 100 | Lake trout..... | 119, 000 |
| Loch Leven trout..... | 18, 000 | Largemouth black bass..... | 18, 580 |
| Rainbow trout..... | 762, 000 | Smallmouth black bass..... | 3, 100 |
| Steelhead salmon..... | 12, 000 | Rainbow trout..... | 48, 800 |
| Illinois: | | Loch Leven trout..... | 61, 800 |
| Brook trout..... | 125 | Pike perch..... | 5, 020, 000 |
| Catfish..... | 1, 000 | Sunfish..... | 3, 265 |
| Largemouth black bass..... | 17, 100 | White bass..... | 1, 800 |
| Smallmouth black bass..... | 300 | Yellow perch..... | 383, 000 |
| Rainbow trout..... | 375 | Mississippi: | |
| Sunfish..... | 2, 600 | Largemouth black bass..... | 180, 175 |
| Indiana: | | Sunfish..... | 201, 000 |
| Brook trout..... | 28, 750 | Missouri: | |
| Catfish..... | 750 | Catfish..... | 400 |
| Largemouth black bass..... | 58, 750 | Crappie..... | 17, 450 |
| Smallmouth black bass..... | 8, 960 | Largemouth black bass..... | 94, 020 |
| Rainbow trout..... | 168, 500 | Rainbow trout..... | 3, 240 |
| Rock bass..... | 500 | Rock bass..... | 20, 900 |
| Sunfish..... | 4, 200 | Sunfish..... | 6, 240 |
| Yellow perch..... | 600 | Montana: | |
| Iowa: | | Blackspotted trout..... | 414, 420 |
| Catfish..... | 2, 000 | Brook trout..... | 139, 015 |
| Crappie..... | 20, 700 | Catfish..... | 87, 530 |
| Largemouth black bass..... | 39, 650 | Crappie..... | 53, 600 |
| Rainbow trout..... | 15, 500 | Largemouth black bass..... | 121, 950 |
| Loch Leven trout..... | 4, 000 | Loch Leven trout..... | 2, 252, 000 |
| Sunfish..... | 6, 700 | Sunfish..... | 91, 200 |
| Kansas: | | Yellow perch..... | 5, 800 |
| Crappie..... | 6, 800 | Nebraska: | |
| Largemouth black bass..... | 1, 925 | Brook trout..... | 274, 300 |
| Rock bass..... | 2, 100 | Catfish..... | 2, 800 |
| Sunfish..... | 2, 200 | Crappie..... | 1, 500 |
| Kentucky: | | Loch Leven trout..... | 138, 800 |
| Largemouth black bass..... | 223, 640 | Largemouth black bass..... | 11, 950 |
| Smallmouth black bass..... | 10, 500 | Rainbow trout..... | 609, 300 |
| Rock bass..... | 3, 350 | Sunfish..... | 1, 675 |
| Sunfish..... | 16, 900 | Yellow perch..... | 3, 000 |
| Louisiana: | | Nevada: | |
| Largemouth black bass..... | 24, 760 | Catfish..... | 200 |
| Sunfish..... | 13, 730 | Largemouth black bass..... | 800 |
| Maine: | | New Hampshire: | |
| Atlantic salmon..... | 85, 000 | Brook trout..... | 79, 400 |
| Brook trout..... | 1, 805, 100 | Catfish..... | 9, 400 |
| Flounder..... | 774, 000, 000 | Landlocked salmon..... | 7, 000 |
| Landlocked salmon..... | 238, 300 | Rainbow trout..... | 9, 200 |
| Smallmouth black bass..... | 33, 750 | Smallmouth black bass..... | 500 |
| Maryland: | | New Jersey: | |
| Brook trout..... | 58, 110 | Largemouth black bass..... | 32, 400 |
| Largemouth black bass..... | 1, 350 | Rainbow trout..... | 5, 000 |
| Smallmouth black bass..... | 550 | Sunfish..... | 75 |
| Loch Leven trout..... | 900 | Yellow perch..... | 500 |

PROPAGATION AND DISTRIBUTION OF FOOD FISHES, 1935 427

Summary, by States, of the distribution of fish, fiscal year 1935—Continued

| State and species | Number | State and species | Number |
|---|--------------|-----------------------------|-------------|
| New Mexico: | | Tennessee—Continued. | |
| Blackspotted trout..... | 36,000 | Crappie..... | 315 |
| Crappie..... | 800 | Largemouth black bass..... | 59,000 |
| Largemouth black bass..... | 146,450 | Smallmouth black bass..... | 385 |
| Rainbow trout..... | 12,000 | Loch Leven trout..... | 5,000 |
| Sunfish..... | 158,400 | Rainbow trout..... | 150,000 |
| New York: | | Rock bass..... | 9,150 |
| Brook trout..... | 443,500 | Sunfish..... | 37,700 |
| Lake trout..... | 322,000 | Texas: | |
| Landlocked salmon..... | 500 | Crappie..... | 5,335 |
| Loch Leven trout..... | 94,000 | Largemouth black bass..... | 241,800 |
| Lake herring..... | 2,640,000 | Sunfish..... | 45,600 |
| Rainbow trout..... | 19,200 | Rock bass..... | 750 |
| Smallmouth black bass..... | 73,200 | Utah: | |
| Whitefish..... | 679,000 | Black-spotted trout..... | 640,000 |
| North Carolina: | | Brook trout..... | 682,700 |
| Crappie..... | 2,500 | Largemouth black bass..... | 2,100 |
| Largemouth black bass..... | 58,000 | Loch Leven trout..... | 1,614,000 |
| Shad..... | 2,775,000 | Landlocked salmon..... | 18,200 |
| Yellow perch..... | 6,600,000 | Rainbow trout..... | 478,700 |
| White perch..... | 920,000 | Vermont: | |
| Sunfish..... | 30,325 | Atlantic salmon..... | 5,000 |
| North Dakota: Rainbow trout..... | 1,000 | Brook trout..... | 1,984,300 |
| Ohio: | | Landlocked salmon..... | 33,000 |
| Brook trout..... | 6,300 | Loch Leven trout..... | 2,000 |
| Catfish..... | 8,250 | Rainbow trout..... | 2,050 |
| Loch Leven trout..... | 9,500 | Smallmouth black bass..... | 49,765 |
| Largemouth black bass..... | 15,300 | Virginia: | |
| Smallmouth black bass..... | 32,400 | Catfish..... | 1,200 |
| Rainbow trout..... | 10,800 | Brook trout..... | 315,975 |
| Sunfish..... | 11,450 | Largemouth black bass..... | 15,150 |
| Whitefish..... | 37,000,000 | Smallmouth black bass..... | 8,830 |
| Oklahoma: | | Loch Leven trout..... | 2,800 |
| Catfish..... | 11,600 | Rainbow trout..... | 248,800 |
| Crappie..... | 90,200 | Rock bass..... | 19,300 |
| Largemouth black bass..... | 28,800 | Shad..... | 11,430,000 |
| Sunfish..... | 62,950 | Sunfish..... | 52,700 |
| Warmouth bass..... | 21,800 | Yellow perch..... | 451,000,000 |
| Yellow perch..... | 2,630 | Washington: | |
| Oregon: | | Black-spotted trout..... | 638,000 |
| Black-spotted trout..... | 85,000 | Brook trout..... | 697,500 |
| Brook trout..... | 385,000 | Chinook salmon..... | 8,554,000 |
| Chinook salmon..... | 5,364,000 | Landlocked salmon..... | 25,900 |
| Loch Leven trout..... | 305,000 | Loch Leven trout..... | 165,600 |
| Rainbow trout..... | 6,000 | Rainbow trout..... | 279,250 |
| Silver salmon..... | 362,000 | Silver salmon..... | 413,000 |
| Steelhead salmon..... | 489,000 | Sockeye salmon..... | 404,000 |
| Pennsylvania: | | Steelhead salmon..... | 523,000 |
| Brook trout..... | 345,000 | West Virginia: | |
| Catfish..... | 2,975 | Brook trout..... | 1,490,000 |
| Largemouth black bass..... | 2,100 | Largemouth black bass..... | 51,000 |
| Loch Leven trout..... | 221,200 | Smallmouth black bass..... | 10,000 |
| Rainbow trout..... | 215,800 | Loch Leven trout..... | 119,000 |
| Sunfish..... | 4,000 | Rainbow trout..... | 367,000 |
| Yellow perch..... | 9,300 | Rock bass..... | 200 |
| South Carolina: | | Sunfish..... | 22,000 |
| Catfish..... | 170 | Wisconsin: | |
| Crappie..... | 500 | Brook trout..... | 623,400 |
| Largemouth black bass..... | 223,000 | Catfish..... | 17,600 |
| Shad..... | 371,000 | Largemouth black bass..... | 188,300 |
| Sunfish..... | 137,000 | Smallmouth black bass..... | 5,310 |
| Warmouth bass..... | 8,000 | Loch Leven trout..... | 251,400 |
| South Dakota: | | Rainbow trout..... | 258,300 |
| Black-spotted trout..... | 122,200 | Sunfish..... | 28,300 |
| Brook trout..... | 324,000 | Wyoming: | |
| Catfish..... | 600 | Black-spotted trout..... | 871,000 |
| Largemouth black bass..... | 14,000 | Brook trout..... | 1,228,000 |
| Loch Leven trout..... | 600,500 | Catfish..... | 2,850 |
| Rainbow trout..... | 649,300 | Largemouth black bass..... | 7,350 |
| Tennessee: | | Loch Leven trout..... | 360,000 |
| Brook trout..... | 98,600 | Rainbow trout..... | 451,000 |
| Catfish..... | 5,225 | Yellow perch..... | 32,200 |

NOTE.— All buffalofish and carp shown in the above table are planted in commercial areas of the Mississippi River.