



Northeast Fisheries Science Center Reference Document 21-08

The International Sampling Program:  
Continent of Origin and  
Biological Characteristics of Atlantic Salmon  
Collected at West Greenland in 2019

October 2021



Northeast Fisheries Science Center Reference Document 21-08

# The International Sampling Program: Continent of Origin and Biological Characteristics of Atlantic Salmon Collected at West Greenland in 2019

by Timothy F Sheehan<sup>1</sup>, John Coyne<sup>2</sup>, Gareth D Davis<sup>3</sup>, Denise Deschamps<sup>4</sup>,  
Ruth Haas-Castro<sup>1</sup>, Patrick Quinn<sup>5</sup>, Louise Vaughn<sup>6</sup>, Rasmus Nygaard<sup>7</sup>, Ian R  
Bradbury<sup>8</sup>, Martha J Robertson<sup>8</sup>, Niall Ó Maoiléidigh<sup>6</sup>, Jon Carr<sup>9</sup>

<sup>1</sup> NOAA Fisheries Service, Northeast Fisheries Science Center, 166 Water Street, Woods Hole, MA 02543, USA

<sup>2</sup> Inland Fisheries Ireland, 3044 Lake Drive, Citywest Business Campus, Dublin 24, Ireland

<sup>3</sup> National Fisheries Service, Environment Agency, Bromholme Lane, Brampton, Huntingdon, England, United Kingdom, PE28 4NE

<sup>4</sup> Ministère des Forêts, de la Faune et des Parcs, Direction de l'expertise sur la faune aquatique, 880, chemin Sainte-Foy, Québec, Québec G1S 4X4, Canada

<sup>5</sup> Agri-Food and Biosciences Institute Newforge, Newforge Lane, Belfast, Northern Ireland, United Kingdom, BT9 5PX

<sup>6</sup> Marine Institute, Fisheries Ecosystems Advisory Services, The Farran Laboratory, Furnace, Newport, Co. Mayo, F28 PF65 Ireland

<sup>7</sup> Greenland Institute of Natural Resources, Kivioq 3, 3905 Nuussuaq, DK-3900 Nuuk, Greenland

<sup>8</sup> Fisheries and Oceans Canada, Science Branch, P.O. Box 5667, St. John's, Newfoundland and Labrador, A1C 5X1, Canada

<sup>9</sup> Atlantic Salmon Federation, 15 Rankine Mill Rd, Chamcook, NB, E5B 3A9, Canada

## **US DEPARTMENT OF COMMERCE**

National Oceanic and Atmospheric Administration

National Marine Fisheries Service

Northeast Fisheries Science Center

Woods Hole, Massachusetts

October 2021

## Northeast Fisheries Science Center (NEFSC) Reference Documents

**This series is a secondary scientific series** designed to assure the long-term documentation of and to enable the timely transmission of research results by Center and/or non-Center researchers, where such results bear upon the research mission of the Center (see the outside back cover for the mission statement). These documents receive internal scientific review, and most receive copy editing. The National Marine Fisheries Service does not endorse any proprietary material, process, or product mentioned in these documents.

If you do not have internet access, you may obtain a paper copy of a document by contacting the senior Center author of the desired document. Refer to the title page of the document for the senior Center author's name and mailing address. If there is no Center author, or if there is corporate authorship, contact the Center's Woods Hole Laboratory Library (166 Water St., Woods Hole, MA 02543-1026).

**Information Quality Act Compliance:** In accordance with section 515 of Public Law 106-554, the NEFSC completed both technical and policy reviews for this report. These pre-dissemination reviews are on file at the NEFSC Editorial Office.

This document may be cited as:

Sheehan TF, Coyne J, Davis DG, Deschamps D, Haas-Castro R, Quinn P, Vaughn L, Nygaard R, Bradbury IR, Robertson MJ, Maoiléidigh NO, Carr J. 2021. The International Sampling Program: Continent of Origin and Biological Characteristics of Atlantic Salmon Collected at West Greenland in 2019. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 21-08; 63 p.

# TABLE OF CONTENTS

Abstract .....	1
International Sampling Program .....	2
Continent and Region of Origin.....	5
Biological Characteristics of the Catches .....	7
Other Sampling .....	8
Acknowledgements.....	9
Table 1. Samplers participating in the 2019 sampling program by country, home institution, sampling period, and community/Northwest Atlantic Fisheries Organization (NAFO) division sampled. ....	10
Table 2. Evaluation of underreporting in 2019 of sampled communities at the Greenland Atlantic salmon ( <i>Salmo salar</i> ) fishery by community/Northwest Atlantic Fisheries Organization (NAFO) division.....	11
Table 3. Reported landings (kg) for the Greenland Atlantic salmon ( <i>Salmo salar</i> ) fishery (2002-2019) by Northwest Atlantic Fisheries Organization (NAFO) division as reported by the home rule government and the division-specific adjusted landings where the sampling teams observed more fish landed than were reported.....	12
Table 4. Reported tag recaptures (n = 1) at the Greenland Atlantic salmon ( <i>Salmo salar</i> ) fishery in 2019.....	14
Table 5. Reported tag recaptures (n = 142) from 2003-2018 at the Greenland Atlantic salmon ( <i>Salmo salar</i> ) fishery.....	15
Table 6. Reporting groups identified within the North Atlantic-wide Atlantic salmon ( <i>Salmo salar</i> ) single nucleotide polymorphism genetic baseline. ....	28
Table 7. The continental proportions of North American (NA) and European (E) Atlantic salmon ( <i>Salmo salar</i> ) caught at West Greenland in 2019 by Northwest Atlantic Fisheries Organization (NAFO) division. ....	29
Table 8. The estimated number, weighted by catch, of North American (NA) and European (E) Atlantic salmon ( <i>Salmo salar</i> ) caught at West Greenland by year from 1982-2019 and the proportion of the catch by weight. ....	30
Table 9. Bayesian proportional mean mixture composition estimates for the West Greenland Atlantic salmon ( <i>Salmo salar</i> ) fishery by Northwest Atlantic Fisheries Organization (NAFO) division sampled in 2019 using the single nucleotide polymorphism range-wide baseline. ....	32

Table 10. Annual mean fork lengths and whole weights by continent of origin (NA = North American and E = European) and sea age (1SW = 1 sea-winter, 2SW = 2 sea-winter, and PS = previous spawner) of Atlantic salmon ( <i>Salmo salar</i> ) caught at West Greenland from 1969-2019. ....	35
Table 11. Mean fork lengths (cm) and whole weight (kg) by sea age (1SW = 1 sea-winter and 2SW = 2 sea-winter), continent of origin and Northwest Atlantic Fisheries Organization (NAFO) division for Atlantic salmon ( <i>Salmo salar</i> ) caught at West Greenland in 2019 with corresponding standard deviation (S.D.). ....	36
Table 12. The river age (smolt age) composition (%) of Atlantic salmon ( <i>Salmo salar</i> ) by continent of origin (NA = North American and E = European) and Northwest Atlantic Fisheries Organization (NAFO) division caught at West Greenland in 2019.....	37
Table 13. River age distribution (%) for North American origin Atlantic salmon ( <i>Salmo salar</i> ) caught at West Greenland from 1968-2019. ....	38
Table 14. River age distribution (%) for European origin Atlantic salmon ( <i>Salmo salar</i> ) caught at West Greenland, 1968-2019. ....	40
Table 15. The sea age (1SW = 1 sea-winter, 2SW = 2 sea-winter, and Previous Spawners) composition of Atlantic salmon ( <i>Salmo salar</i> ) by continent of origin (NA = North American and E = European) and Northwest Atlantic Fisheries Organization (NAFO) division caught at West Greenland in 2019.....	42
Table 16. Sea age (1SW = 1 sea-winter, 2SW = 2 sea-winter, and PS = Previous Spawners) distribution (%) for North American and European origin Atlantic salmon ( <i>Salmo salar</i> ) caught at West Greenland from 1985-2019.....	43
Figure 1. Nominal catches and commercial quotas (metric tons, round fresh weight) of Atlantic salmon ( <i>Salmo salar</i> ) at West Greenland for 1960-2019 (top panel) and 2010-2019 (bottom panel). ....	44
Figure 2. Map of southwest Greenland showing communities at which Atlantic salmon ( <i>Salmo salar</i> ) have historically been landed. ....	45
Figure 3. Map showing total samples (gray circles) and analyzed subsamples (blue circles) during 2019 at the West Greenland Atlantic salmon ( <i>Salmo salar</i> ) fishery for the single nucleotide polymorphism analysis. ....	46
Figure 4. Map of sample locations for the single nucleotide polymorphism range-wide genetic baseline for North American (top) and European (bottom) regional groupings.....	48
Figure 5. The weighted proportions of North American and European Atlantic salmon ( <i>Salmo salar</i> ) caught at West Greenland from 1982-2019. ....	48
Figure 6. Proportions of unsampled adjusted landings of North American origin and European origin Atlantic salmon ( <i>Salmo salar</i> ) (left panels) and of sampled adjusted landings or North	

American origin and European origin Atlantic salmon (right panels) by North Atlantic Fisheries Organization (NAFO) division (top row represents division 1A and bottom row represents division 1F) sampled at West Greenland from 2005-2019. .... 49

Figure 7. The weighted numbers of North American and European Atlantic salmon (*Salmo salar*) caught at West Greenland from 1982-2019 (top) and 2010-2019 (bottom). .... 50

Figure 8. Bayesian estimates of mixture composition of samples from the West Greenland Atlantic salmon (*Salmo salar*) fishery for 2019 using the single nucleotide polymorphism (SNP) baseline overall and by Northwest Atlantic Fisheries Organization (NAFO) division. .... 51

Figure 9. Mean whole weight (kg) of European and North American 1 sea-winter (fish that have spent 1 winter at sea) Atlantic salmon (*Salmo salar*) sampled at West Greenland from 1969-2019 (top panel) and 2010-2019 (bottom panel). .... 52

## ABSTRACT

An Atlantic salmon (*Salmo salar*) mixed-stock fishery operating from August through October exists off the western coast of Greenland and primarily harvests 1 sea-winter (1SW) North American and European origin salmon destined to return to natal waters as 2 sea-winter (2SW) spawning adults. To collect data on the biological characteristics and origin of the harvest necessary for international stock assessment efforts, parties to the North Atlantic Salmon Conservation Organization's (NASCO) West Greenland Commission (WGC) agreed to participate in an international sampling program for the 2019 fishery. The sampling program was coordinated by the USA (NOAA Fisheries Service) and involved 7 samplers from 6 countries deployed among 4 communities (Sisimiut, Maniitsoq, Nuuk, and Qaqortoq) located on the west coast of Greenland. Reported landings in 2019 were 29.8 metric tons (t). Data on length, weight, and freshwater and marine age were collected from scale samples, and data on continent and region of origin were collected from genetic analysis of tissue samples. Since 2002 (with the exception of 2006, 2011, 2015, and 2018), unreported landings were identified by comparing the reported landings to the weight of the sampled harvest for each community. Unreported landings were not detected in 2019. In total, 1,340 salmon were observed by the sampling teams (approximately 13% by weight of the reported landings), and 1,119 of these were sampled for biological characteristics. As seen since the mid-1990s, a high proportion of the harvested stock was of North American origin (71.5%) with the balance of European origin (28.5%). Primary contributors to the sampled North American origin salmon were the Gaspé Peninsula, the Gulf of St. Lawrence, and the Labrador South reporting groups (65%). Of European origin salmon, 99% were from the United Kingdom/Ireland reporting group. No individuals were identified as having originated from the Kapisillit reporting group, which represents Greenland's only self-sustaining Atlantic salmon population. North American origin fish were primarily freshwater age 2 or 3 years (27% and 33% respectively). European origin fish were primarily freshwater age 2 (61%). Overall, 97% of the sampled fish were 1SW salmon. The mean length of North American 1SW salmon was 63.9 cm, and the mean whole weight was 2.93 kg; the mean length of European 1SW salmon was 63.4 cm, and the mean whole weight was 2.89 kg. Approximately 6,800 North American salmon (20.3 t) and 2,600 European salmon (8.1 t) were harvested, not taking into account any unreported catch. The sampling program was successful in adequately sampling the Greenland catch, both temporally and spatially, and provided essential input data to international stock assessment efforts that provide stock status and catch options for subsequent fishery management.

## INTRODUCTION

An important mixed-stock Atlantic salmon (*Salmo salar*) fishery exists off the western coast of Greenland. This fishery takes primarily 1 sea-winter (1SW, fish that have spent 1 winter at sea) North American and European origin salmon that would potentially return to natal waters as mature 2 sea-winter (2SW) spawning adults or older. Effective management of the resource on both continents requires annual collection of accurate landings data, continent and region of origin assignments, and biological characteristics data to assess the impact of the fishery on the contributing stock complexes. Data collected from the fishery are also required for use in assessment models which predict pre-fishery abundance of North American and European stocks to provide fishery managers with catch options required for setting harvest regulations.

Atlantic salmon were first documented off the coast of Greenland in 1780 and were targeted by a small local inshore gillnet fishery (Jensen 1990). During the early 1960s, the fishery developed an international presence; in 1965, vessels from Norway, Denmark, Sweden, and the Faroe Islands arrived and introduced an offshore drift gillnet fishery (ibid.). Reported catches increased to a high of 2,689 t in 1971 (Figure 1). Mark-recapture studies conducted during this period indicated that the Atlantic salmon caught in this fishery were of North American and European origin and were not uniformly distributed along the coast (Reddin et al. 2012). Because of concerns that this fishery would have deleterious impacts on the contributing stock complexes, a quota system was agreed upon and implemented in 1976 (Colligan et al. 2008), and since 1984, the North Atlantic Salmon Conservation Organization (NASCO) has established catch regulations.

Since 1969, a coordinated international sampling program has been conducted to obtain biological samples from the Greenland salmon fishery. From 1969-1981, research vessels were used to obtain samples. Since 1982, international teams of samplers have been deployed throughout West Greenland to obtain samples from fish processing plants (when a commercial fishery is allowed), local markets, and other vendors from individual communities landing salmon. The focus of this sampling program is to collect biological data and samples. Historically, lengths, weights, and scale samples were collected, and individual salmon were scanned for fin clips or external/internal tags. Beginning in 2002, tissue samples have been collected from fish for genetic stock identification.

The purpose of this paper is to:

- describe the international sampling program;
- present the results from the continent and region of origin analysis; and
- summarize the biological characteristics of the catch from West Greenland during the internal-use-only fishery of 2019.

## INTERNATIONAL SAMPLING PROGRAM

The West Greenland Commission (WGC) of NASCO has agreed to regulatory measures for the West Greenland fishery for all years from 1984 onward (except 1985, 1991, 1992, and 1996). Since 2006, these regulations have been applied as multiyear measures. The latest measure was established for the period of 2018-2020 (NASCO 2018; see WGC(18)11), and these regulations are set to continue in 2020 if the Framework of Indicators (FWI) developed and updated by the International Council for the Exploration of the Sea (ICES; ICES 2007, 2018) indicates no significant change, implying a reassessment of the catch advice would not be required.



From 2002-2011, the national quota for commercial landings of Atlantic salmon for export was set to 0 t by the Government of Greenland, but the internal-use-only fishery for personal and local consumption was unaffected. Selling of salmon to hotels, institutions, and local markets by licensed fishermen and an unlicensed fishery for private consumption was allowed. The internal-use-only fishery was without a quota limit, but landings of Atlantic salmon had been previously estimated at 20 t annually. The fishery generally operates during the months of August, September, and October, and from 2005-2014, the fishery opened on 1 August and closed on 31 October. The fishery is regulated according to the Government of Greenland Executive Order No. 5 of 21, 21 September 2018, an update from previous orders (Executive Order No. 12 of 1 August 2012 and Executive Order No. 21 of 10 August 2002). Starting in 2015, the Government of Greenland has delayed the opening of the fishery until 15 August with a closing date of 31 October.

From 2012-2014, the quota for commercial landings of Atlantic salmon for export remained at 0 t and hence no export of Atlantic salmon was allowed. However, in 2012, the Government of Greenland set a 35 t national quota for landing at fishing processing factories to provide a year-round supply of locally harvested Atlantic salmon within Greenland. The internal-use-only fishery for personal and local consumption remained unaffected and unrestricted by the quota for factory landings. A factory landings-only quota was again set to 35 t in 2013 but was then reduced to 30 t in 2014. From 2015-2017, the Government of Greenland unilaterally set a quota of 45 t for all components of its fishery, as a quota could not be agreed to by all parties of the WGC of NASCO (NASCO 2015). The regulatory measure stated that any harvest exceeding the quota within a year would be subtracted from the quota in the following year. Given overage of the 2015 harvest, the 2016 quota was set to 32 t by the Government of Greenland. The 2017 quota was set to 45 t as no overharvest was recorded in 2016.

A new multiannual regulatory measure for the Atlantic salmon fishery at West Greenland was adopted by members of NASCO's WGC in 2018 (NASCO 2018; see WGC(18)11). The measures applied to the 2018 fishery and as noted above will also apply to the 2020 fishery if the FWI indicate no significant change in the previously provided catch advice. Within the regulatory measure, the Government of Greenland agreed not to export wild Atlantic salmon or its products from Greenland and to prohibit landings and sales to fish processing factories. They also agreed to restrict the fishery from 15 August to no later than 31 October each year. Total allowable catch for all components of the fishery was set at 30 t, and any overharvest in a particular year would result in an equal reduction in the total allowable catch in the following year. The regulatory measure also set out a number of provisions aimed at improving the monitoring, management control, and surveillance of the fishery including a new requirement for all fishers to obtain a license to fish for Atlantic salmon, an agreement to collect catch and fishing activity data from all licensed fishers, and mandatory reporting requirements for all fishers. The measure also stated that as a condition of the license, all fishers will be required to allow samplers from the NASCO sampling program to take samples of their catches upon request.

Under NASCO's Statement of Cooperation on the West Greenland Fishery Sampling Program for 2019 (NASCO 2019; see WGC(19)06), parties to NASCO's WGC agreed to provide staff to sample Atlantic salmon catches from the West Greenland internal-use-only fishery during the 2019 season.

The objectives of the sampling program were to:

- continue the time series of data (1969-2018) on continent of origin and biological characteristics of the Atlantic salmon at the West Greenland fishery;

- provide data on mean weight, length, age, and continent of origin for use in the North American and European Atlantic salmon run-reconstruction models; and
- collect information on the recovery of internal and external tags.

As outlined in the sampling agreement, the European Union agreed to provide staff to sample the fishery for a minimum of 8 person-weeks (which would amount to 8 weeks of sampling), and the United States and Canada both agreed to a minimum of 2 person-weeks. Samplers from various countries involved in the program are outlined in Table 1.

The coordination of this effort was handled by the USA (NOAA Fisheries Service) with assistance from the Greenland Institute of Natural Resources (GINR). Individual samplers were deployed during the course of the fishing season to provide the best possible spatial and temporal coverage of the fishery. A total of 7 samplers were stationed in 4 communities located within 4 Northwest Atlantic Fisheries Organization (NAFO) divisions (Figure 2): Sisimiut (1B), Maniitsoq (1C), Nuuk (1D) and Qaqortoq (1F). Factory landings have not been authorized by the Government of Greenland since 2015, and therefore no factory landing samples were collected.

Reported landings in 2019 were 29.8 t (28.3 t for West Greenland and 1.4 t for East Greenland ICES Statistical Area XIV). In the past, underreporting of harvest was identified by comparing the reported landings to the sample data. From 2002-2018 (with the exception of 2006, 2011, 2015, and 2018), the sampling team documented more fish than reported in at least 1 division (ICES 2019). A documented salmon could be one that was either sampled, checked for an adipose clip only, or not sampled but seen. When this type of discrepancy occurs, the reported landings are adjusted to include the total weight of the fish documented as being landed during the sampling period, and the adjusted landings are included in all subsequent assessments. Considering that samplers are not stationed within a community throughout the entire fishing season and that there are numerous communities without samplers present, these adjusted landings should be considered minimum estimates.

In 2019, no such discrepancy was detected (Table 2). The time series of reported landings and adjusted landings for 2002-2019 are presented in Table 3. To provide the most reliable estimate of catch, which is necessary for estimating the potential fishery impacts on contributing stocks, it is important to continually improve the catch reporting procedures and the quality of the catch statistics. Factory landings and samples, when allowed, are not considered within this process since these landings are strictly regulated by the Government of Greenland (i.e., only licensed commercial fishers can land at designated factories) and are accounted for and reported by the factory managers to the Greenland Fisheries License Control Authority on a daily to weekly basis.

Landed fish were sampled at random, and when possible, the total catch was sampled. Individual fish were measured (fork length, mm) and weighed (gutted weight or whole weight, 0.01 kg). Scales were taken for age determination, and adipose fins were taken for DNA analysis for stock identification. Fish were also examined for fin clips, external marks, external tags, and internal tags. Adipose-clipped fish were sampled for microtags (coded wire tags).

Sampling teams observed 1,340 salmon. Of this total, 1,119 were sampled for biological characteristics representing 13% of the reported landings. Factory landings were not allowed by the Government of Greenland, and therefore no factory samples were obtained. A total of 85 fish were only checked for an adipose clip, and an additional 136 were documented as being landed but were not sampled or examined further. Biological characteristics data and samples were collected as follows:

- 1,117 fork lengths;
- 927 gutted weights;
- 125 whole weights;
- 1,049 scale samples; and
- 1,119 genetic samples.

In total, 20 adipose-clipped fish were documented. Of all the fish examined by the samplers, no internal or external tags were detected. A single spaghetti tag was provided directly by a fisher to the GINR (Table 2.4).

Tag recoveries at Greenland have been recorded from 1963 to the present time. In total, 5,508 tag recoveries were recorded and archived (Ó Maoiléidigh et al. 2018) from 1963-2009. A complete archive of tag recoveries from the contemporary sampling undertaken by this sampling program is also maintained and is provided in Table 2.5. A total of 142 tags have been recovered by the sampling program since 2003, not including the single recovery in 2019 listed in Table 2.4.

Non-factory sampling often occurs at a local market which is a centralized location where harvested salmon are present and available. Prior to any sampling, the sampler always obtains permission from the market manager or fisher. This arrangement has generally been successful for all samplers, although there have been issues in some years in Nuuk (Sheehan et al. 2013). Similar issues were also noted in 2014 when samplers were denied access to fish in Maniitsoq and Qaqortoq. No issues have occurred since 2015.

Since 2015, it has been a condition of the commercial fishing license to allow samplers access to landed salmon. However, given the commitment made by the Government of Greenland—in cooperation with the GINR—to sample Atlantic salmon from the city of Nuuk on a weekly basis during the 2019 fishing season (NASCO 2019; see WGC(19)06), a sampler was not deployed to Nuuk. Staff from the GINR were able to collect samples from Nuuk on two dates. This is the first time samples have been collected from Nuuk since 2011.

The limitation of the fishery to internal-use-only caused some practical problems for the sampling teams. However, the sampling program provided adequate representation of the Greenland catch, both temporally and spatially.

## CONTINENT AND REGION OF ORIGIN

Fin tissue samples were collected and preserved in *RNAlater*<sup>TM</sup>, an aqueous, nontoxic tissue and cell storage reagent that stabilizes and protects cellular RNA. A total of 1,119 usable samples were collected in 2019 from 4 communities in 4 NAFO divisions: Sisimiut in 1B (n = 371), Maniitsoq in 1C (n = 427), Nuuk in 1D (n = 68), and Qaqortoq in 1F (n = 253). Because of funding limitations, a subset of the tissue samples collected in 2019 was genetically analyzed (Figure 3). In total, 1,071 samples were processed from the 4 communities: Sisimiut (n = 365), Maniitsoq (n = 423), Nuuk (n = 36), and Qaqortoq in 1F (n = 247).

From 1969-2001, scale pattern analysis was used to make continent of origin determinations and estimate the proportion of the harvest originating from North American and European rivers (Reddin and Friedland 1999). From 2002-2016, DNA isolation and the subsequent microsatellite analyses were performed according to standardized protocols (King et al., 2001; Sheehan et al., 2010). A database of approximately 5,000 Atlantic salmon genotypes of known origin were used as a baseline to assign the samples to continent of origin.

Starting in 2017, a single nucleotide polymorphism range-wide baseline (SNP; Jeffery et al., 2018) providing 20 North American and 8 European reporting groups was used to determine continent and region of origin. The baseline has been revised, resulting in 21 North American and 10 European reporting groups (Table 6 and Figure 4; ICES 2019).

DNA extraction and genotyping of all fishery samples were carried out at the Aquatic Biotechnology Laboratory (Fisheries and Oceans Canada, Maritimes Region), and DNA was extracted with the Qiagen DNeasy Blood & Tissue 96-well extraction kit (Qiagen; [www.qiagen.com](http://www.qiagen.com)) following the guidelines of the manufacturer. DNA was quantified by using Quant-iT™ PicoGreen™ (Life Technologies; [www.thermofisher.com/us/en/home/brands/life-technologies.html](http://www.thermofisher.com/us/en/home/brands/life-technologies.html)) and diluted to a final concentration of 10 ng/μL in 10mM Tris (Qiagen Buffer EB). SNP genotyping of the 96 SNP loci was performed by using SNPtype assays (Fluidigm; [www.fluidigm.com](http://www.fluidigm.com)) per the manufacturer's protocols and as described in Jeffery et al. (2018). A Bayesian approach is used to estimate mixture composition or assign individuals to continent and region of origin. The approach uses the R package rubias (Anderson et al. 2008).

In total, 71.5% of the salmon sampled in 2019 were of North American origin, and 28.5% were of European origin (Figure 5). The NAFO division-specific continent of origin assignments are presented in Table 7. These findings show that high proportions of fish from the North American stock complex continue to contribute to the fishery. The variability in the recent stock complex contributions between divisions and the deviation from past trends (Figure 6) underscore the need to annually sample multiple NAFO divisions to achieve accurate estimates of continental contributions to the harvest.

Variations in the estimated weighted proportions and number of North American and European salmon harvested in the fishery from 1982-2019 are shown in Table 8 and in Figures 5 and 7. The 2019 North American weighted contribution (72%) is above the long-term mean (1982-2019, 69.3%) and lower than the previous 10-year mean (2009-2018, 79.2%). The European weighted contribution (28%) to the 2019 fishery was below the long-term mean (1982-2019, 30.7%) but above the previous 10-year mean (2009-2018, 20.8%). In terms of numbers of fish, the 2019 fishery caught approximately 6,800 North American salmon (20.3 t) and 2,600 European salmon (8.1 t). The 2019 total number of fish harvested (9,400) is below the previous 10-year mean (11,740) but is only 2.8% of the maximum estimate of 336,000 fish harvested in 1982.

The North American contributions to the West Greenland fishery, as in previous years (Bradbury et al. 2016; ICES 2017), are dominated by the Gaspé Peninsula, the Gulf of St. Lawrence, and the Labrador South reporting groups (Table 9 and Figure 8). These three groups accounted for 65% of the North American contributions in 2019. The Northeast Atlantic contributions were dominated by the United Kingdom/Ireland reporting group with 99% of the European contributions in 2019.

From North America, there are smaller (0.1–5.4%), contributions to the harvest for a number of other reporting groups (Labrador Central, Quebec City Region, Ungava, St. Lawrence N. Shore Lower, Western Newfoundland, Maine United States, Lake Melville, Anticosti, Newfoundland 2, Newfoundland 1, Eastern Nova Scotia, and Northern Newfoundland, Table 9 and Figure 8). Within the European contributions, all other reporting groups were estimated to contribute 0-0.2% to the overall harvest. The update results support the previous conclusion by ICES (2017) that stocks from the Northern North-East Atlantic Commission (NEAC) do not contribute a significant amount to the harvest at West Greenland. Further, the variation in NAFO division-specific region of origin assignments highlight the variation of region-specific contributions across years and NAFO divisions.

## BIOLOGICAL CHARACTERISTICS OF THE CATCHES

Biological characteristics (length, weight, and age) were recorded for all sampled fish. An overall decrease in mean whole weight of both European and North American 1SW salmon occurred between 1969 and 1995 (Table 10 and Figure 9). This trend was reversed in 1996 when mean weights began to increase, although evidence suggests these trends may be partially explained by annual variation in the timing of the sampling program (ICES 2011, 2015). In 2019, the mean length of North American 1SW salmon was 63.9 cm, and the mean whole weight was 2.93 kg; the mean length of European 1SW salmon was 63.4 cm, and the mean whole weight was 2.89 kg. The North American 1SW fork length estimate was approximately equal to the 2018 value (63.8 cm) and below the previous 10-year average (2009-2018, 65.6 cm). The European 1SW mean fork length was also approximately equal to the 2018 value (63.9 cm) and was slightly below the previous 10-year average (2009-2018, 64.4 cm). The North American 1SW whole weight was approximately equal to the 2018 value (2.91 kg) and was below the previous 10-year average (2009-2018, 3.28 kg). The European 1SW whole weight was approximately equal to the 2018 value (2.93 kg) and below the previous 10-year average (2009-2018, 3.15 kg). A summary of the mean fork lengths and whole weights in the 2019 fishery by sea age, continent of origin, and NAFO division is presented in Table 11. Note that the weight data have not been adjusted for date of capture, and hence may not represent an actual change in mean weight over the time series because fish sampled later in the fishing season have had additional time to grow compared to fish sampled early in the season (ICES 2011, 2015).

The smolt age distribution of the sampled catch by continent of origin and NAFO division is presented in Table 12. The smolt age distributions by origin for all North American and European origin salmon caught (1968-2019) are provided in Tables 13 and 14.

The mean smolt age of the 2019 North American origin samples was 3.3 years. Although age-1 smolts historically represent a small proportion of the catch (previous 10-year mean of 0.7%, 2009-2018), the 2019 value (0.6%) remains among the lowest in the time series. There has been a consistent trend over the past 2 decades of decreased contributions of age-1 smolts as the overall (1968-2019) mean contribution of age-1 smolts equals 2.3%. This is indicative of the relatively minor contributions of the more southerly North American populations as age-1 smolt natural and hatchery production is restricted to the southern end of the range (ICES 2004). The percentage of age-2 smolts of North American origin in the 2019 fishery (26.9%) decreased from the 2018 value (29.8%) and is slightly below the previous 10-year mean (2009-2018, 29.0%). Age-3 and older smolts accounted for 72.5% of the 2019 harvest of North American salmon, which is slightly above the previous 10-year mean (70.3%, 2009-2018) and the overall mean for the 49-year time series (1968-2019 excluding data gaps in 1977 and 1993-1994, 66.6%).

The mean smolt age of the European salmon in 2019 was 2.3 years. The percentage of age-1 smolts (7.5%) decreased from the 2018 value (13.7%) and is slightly below the previous 10-year mean of 9.8% (2009-2018). The percentage of age-2 smolts (60.5%) in the 2019 fishery is slightly lower than in 2018 (62.1%) and the previous 10-year mean (2009-2018, 61.4%). The contribution of age-3 and older European origin smolts (32.0%) is slightly greater than the previous 10-year mean (2009-2018, 28.8%).

The sea age distribution of the sampled catch by continent of origin and NAFO division is presented in Table 15. As expected, the 1SW age group was dominant (96.5%) in the 2019 fishery. This value is within the range of historical values (Table 16). Concerns have been raised over recent difficulty with discerning winter annuli from apparent “checks” in the marine zone of

Atlantic salmon multi-sea winter scales. Care should be taken to properly discern true marine annuli from growth checks, and we note that further study of this phenomenon is warranted.

## **OTHER SAMPLING**

The International Sampling Program at West Greenland provides a unique opportunity for researchers to obtain samples, above what is normally collected by the program, in support of Atlantic salmon research efforts at minimal additional costs. In recent years, the Sampling Program Coordinator has received inquiries from researchers requesting the collection of a variety of sample types from the Atlantic salmon harvested at Greenland. The Program Coordinator reviews all requests received. If the request is reasonable and will not detract from the primary tasks of the samplers, the Program Coordinator will work with the individual researchers and the samplers to facilitate the collection of the requested samples. The objective of this section is to provide an overview of the purpose of these additional samples collected by the sampling program. A generic title and the sample requester have been identified for each.

### **Disease Sampling**

*Jon Carr (Atlantic Salmon Federation)*

No disease samples were collected in 2019 from Atlantic salmon harvested at Greenland. One sampler was prepared to collect samples but was unable to because of a combination of bad weather, low landings, and market demand for landed salmon within the community. Landed salmon were quickly sold, thereby not leaving enough time for the sampler to conduct the detailed disease sampling. Disease samples were last collected in 2017.

Samples consisting of gill, spleen, liver, heart, kidney, and pyloric caeca were collected from 30 individual fish from a single community (Maniitsoq) in 2017 and kidney samples were collected from a single community (Paamiut) in 2016. The goal of the study was to assess the presence and abundance of a broad range of infectious agents, including viruses, bacteria, and microparasites known or suspected to cause disease in salmon worldwide. Samples were evaluated using the Fluidigm Biomark HT-qPCR platform and assay panel to quantify the presence and relative loads of 47 infectious agents in preserved tissues (Miller et al. 2014; 2016).

Infection profiles did not differ significantly between years within continental stock groupings, so data from 2016 and 2017 were pooled. Nine infective agent species were detected (1 bacterium, 3 viruses, 5 parasites), with greater richness among the North American origin salmon versus the European origin salmon sampled. All agents detected in the European origin group (*Parvicapsula pseudobranchicola*, *Tetracapsuloides bryosalmonae*, *Paranucleospora theridion*, *Candidatus Piscichlamydia salmonis*, *Piscine orthoreovirus* (PRV)) were also detected in the North American origin group, which hosted four additional agents (*Ichthyophonus hoferi*, *Sphaerothecum destruens*, ISAV, viral encephalopathy and retinopathy). A manuscript describing these results and the results from similar sampling on wild adult returns and escaped farmed salmon from a number of eastern Canadian rivers has been accepted for publication (Teffer et al. 2020).

### **Sea Lice Sampling**

*Mark Fast (Atlantic Veterinary College, University of Prince Edward Island, Canada)*

*Helene Fjørtoft (Norwegian University of Science and Technology, Norway)*  
*Kim Præbel (UiT The Arctic University of Norway)*

Live sea lice were collected and preserved in RNAlater™ from Atlantic salmon harvests at Greenland. A total of 135 samples were collected from 3 communities in 2019. Samples were split evenly and provided to 3 different researchers in support of 3 different research projects. The projects are investigating the genomics of the Atlantic sea lice as it may relate to the ecology and drug resistance of the species, as well as the role farm/wild interactions may play into sea lice epidemiology. Sample processing and analysis continues for all studies.

## **ACKNOWLEDGEMENTS**

We would like to acknowledge the Greenland Institute of Natural Resources and the fishers and residents of Greenland who provided access to their fish. We would also like to thank the various laboratories and agencies for supporting the program, providing the samplers, and for the funding necessary to support the sampling at Greenland. Funding support for the samplers was provided by NOAA Fisheries Service (for Ruth Haas-Castro), Fisheries and Oceans Canada, Moncton (for Denise Deschamps), Inland Fisheries Ireland (for John Coyne), Marine Institute (for Louise Vaughn), Centre for Environment, Fisheries and Aquaculture Science (for Gareth Davies) and Agri-Food and Biosciences Institute (for Patrick Quinn). Fisheries and Oceans Canada provided funding to support the genetic processing and continent/region of origin analysis. Fisheries and Oceans Canada (Newfoundland and Labrador Region) conducted the aging of all scale samples collected and maintains the master sampling database. Reference to trade names does not imply endorsement by any collaborating agency or government.

**Table 1. Samplers participating in the 2019 sampling program by country, home institution, sampling period, and community/Northwest Atlantic Fisheries Organization (NAFO) division sampled.**

<b>Sampler</b>	<b>Country</b>	<b>Home Institution</b>	<b>Sampling Period</b>	<b>Community (NAFO Division)</b>
John Coyne	Ireland	Inland Fisheries	16 Aug - 29 Aug	Maniitsoq (1C)
Gareth Davies	UK (England and Wales)	Environment Agency	16 Aug – 29 Aug	Qaqortoq (1F)
Louise Vaughn	Ireland	Marine Institute	25 Aug – 07 Sep	Sisimiut (1B)
Denise Deschamps	Canada	Ministère des Forêts, de la Faune et des Parcs du Québec Canada	31 Aug – 18 Sep	Maniitsoq (1C)
Patrick Quinn	Ireland	Agri-Food and Biosciences Institute	03 Sep – 14 Oct	Qaqortoq (1F)
Rasmus Nygaard	Greenland	Greenland Institute of Natural Resources	10 and 20 Sep	Nuuk (1D)
Ruth Haas-Castro	USA	NOAA Fisheries Service	14 Sep – 26 Sep	Sisimiut (1B)



**Table 2. Evaluation of underreporting in 2019 of sampled communities at the Greenland Atlantic salmon (*Salmo salar*) fishery by community/Northwest Atlantic Fisheries Organization (NAFO) division. The total number of salmon documented by the sampling teams (salmon that have been sampled, seen but not sampled, and seen and checked for an adipose fin clip only) is converted to a total whole weight (WW) based on a conversion factor of 1.11 and compared to the reported landings for each community. Gutted weight is denoted as GW.**

<b>Community (NAFO Division)</b>	<b># sampled</b>	<b>Additional # seen</b>	<b>Ave. sampled GW (kg)</b>	<b>Ave. converted WW (kg)</b>	
Sisimiut (1B)	371	31	2.58	2.87	
Maniitsoq (1C)	427	26	2.59	2.87	
Nuuk (1D)	68	0	2.65	2.95	
Qaqortoq (1F)	253	164	2.85	3.16	
<b>Total</b>	<b>1,119</b>	<b>221</b>	<b>2.65</b>	<b>2.95</b>	

<b>Community</b>	<b>Est. WW sampled/seen (kg)</b>	<b>Reported Commercial landings (kg)</b>	<b>Adjusted Commercial landings (kg)</b>	<b>Difference (kg)</b>	<b>Difference as % of reported landings</b>
Sisimiut (1B)	1,153	2,306	2,306	0	0%
Maniitsoq (1C)	1,300	3,487	3,487	0	0%
Nuuk (1D)	200	6,751	6,751	0	0%
Qaqortoq (1F)	1,320	1,768	1,768	0	0%
<b>Total</b>	<b>3,973</b>	<b>14,312</b>	<b>14,312</b>	<b>0</b>	<b>0%</b>

**Table 3. Reported landings (kg) for the Greenland Atlantic salmon (*Salmo salar*) fishery (2002–2019) by Northwest Atlantic Fisheries Organization (NAFO) division as reported by the home rule government and the division-specific adjusted landings where the sampling teams observed more fish landed than were reported. Landings from International Council for the Exploration of the Seas Statistical Area XIV (East Greenland) are not included in the assessment but amounted to 1.4 t in 2019. Shaded cells indicate that sampling took place in that year and division.**

Year		NAFO Division						Total
		1A	1B	1C	1D	1E	1F	
2002	Reported	14	78	2,100	3,752	1,417	1,661	9,022
	Adjusted						2,408	9,769
2003	Reported	619	17	1,621	648	1,274	4,516	8,694
	Adjusted			1,782	2,709		5,912	12,312
2004	Reported	3,476	611	3,516	2,433	2,609	2,068	14,712
	Adjusted				4,929			17,209
2005	Reported	1,294	3,120	2,240	756	2,937	4,956	15,303
	Adjusted				2,730			17,276
2006	Reported	5,427	2,611	3,424	4,731	2,636	4,192	23,021
	Adjusted							
2007	Reported	2,019	5,089	6,148	4,470	4,828	2,093	24,647
	Adjusted						2,252	24,806
2008	Reported	4,882	2,210	10,024	1,595	2,457	4,979	26,147
	Adjusted				3,577		5,478	28,627
2009	Reported	195	6,151	7,090	2,988	4,296	4,777	25,496
	Adjusted				5,466			27,975
2010	Reported	17,263	4,558	2,363	2,747	6,766	4,252	37,949
	Adjusted		4,824		6,566		5,274	43,056
2011	Reported	1,858	3,662	5,274	7,977	4,021	4,613	27,407
	Adjusted							
2012	Reported	5,353	784	14,991	4,564	3,993	2,951	32,636
	Adjusted		2,001				3,694	34,596

**Table 3. continued. Reported landings (kg) for the Greenland Atlantic salmon (*Salmo salar*) fishery (2002–2019) by Northwest Atlantic Fisheries Organization (NAFO) division as reported by the home rule government and the division-specific adjusted landings where the sampling teams observed more fish landed than were reported. Landings from International Council for the Exploration of the Seas Statistical Area XIV (East Greenland) are not included in the assessment but amounted to 1.4 t in 2019. Shaded cells indicate that sampling took place in that year and division.**

Year		NAFO Division						Total
		1A	1B	1C	1D	1E	1F	
2013	Reported	3,052	2,359	17,950	13,356	6,442	3,774	46,933
	Adjusted		2,461				4,408	47,669
2014	Reported	3,626	2,756	13,762	19,123	14,979	3,416	57,662
	Adjusted						4,036	58,282
2015	Reported	751	8,801	10,055	17,966	4,170	14,134	55,877
	Adjusted							
2016	Reported	763	1,234	7,271	4,630	4,492	7,265	25,655
	Adjusted		1,499					25,920
2017	Reported	1,114	1,665	9,335	6,858	3,219	5,563	27,754
	Adjusted		1,942					28,031
2018	Reported	2,434	5,684	13,726	8,202	4,214	4,788	39,048
	Adjusted							
2019	Reported	776	3,036	4,351	8,027	4,822	7,321	28,333
	Adjusted							

**Table 4. Reported tag recaptures (n = 1) at the Greenland Atlantic salmon (*Salmo salar*) fishery in 2019. Northwest Atlantic Fisheries Organization (NAFO) division refers to NAFO or International Council for the Exploration of the Sea statistical areas. No tags were recovered from sampled fish by the sampling team, but 1 tag was provided directly by a fisher or consumer to the Greenland Institute of Natural Resources. In previous years, tags provided directly by a fisher or consumer were sometimes from historical recoveries. Empty cells identify incomplete recapture or release information.**

<b>Tag type</b>	<b>Tag code (Seq. code)</b>	<b>Release country</b>	<b>River released</b>	<b>Place released</b>	<b>Release year</b>	<b>Recapture Community (NAFO Division)</b>	<b>Recapture year</b>
Spaghetti	blue (S1011)	Canada	Hunt River		2017	Maniitsoq (1C)	2019

**Table 5. Reported tag recaptures (n = 142) from 2003-2018 at the Greenland Atlantic salmon (*Salmo salar*) fishery. NAFO Division/ICES Area refers to Northwest Atlantic Fisheries Organization or International Council for the Exploration of the Sea statistical areas. Empty cells identify incomplete recapture or release information.**

	TAG INFORMATION		RELEASE INFORMATION						RECAPTURE INFORMATION						
<i>YEAR ENTERED</i>	TAG TYPE	TAG CODE (SEQ. CODE)	COUNTRY	RIVER	DATE	LIFE STAGE	COMMUNITY (AREA)	NAFO DIVISION/ICES AREA	YEAR	ENV. NO.	DATE	LENGTH (CM)	WEIGHT (KG)	GW OR WW	EXACT OR EST.
2003	carlin	green (C58283)	UK(Scot)	North Esk	April - May 2002	smolt	East Greenland	XIV	2003		27-Oct-03	80.0			est.
2003	carlin	green (C51949)	UK(Scot)	North Esk	Apr-June 2001	smolt	Qaqortoq	1F	2003	4579	11-Sep-03	84.0	6.36	GW	exact
2003	cwt	04 47 58	Ireland	Ballynahinch	18-Feb-02	smolt	Qaqortoq	1F	2003	4896	26-Aug-03	67.5	3.24	GW	exact
2003	cwt	22 42 36	UK(E&W)	Severn (Teme)	14-Mar-02	smolt	Qaqortoq	1F	2003	4478	4-Sep-03	65.4	2.54	GW	exact
2003	cwt	04 47 34	Ireland	Parteen	11-Apr-02	smolt	Qaqortoq	1F	2003	4287	21-Aug-03	56.2	1.78	GW	exact
2003	cwt	01 47 74	Ireland	Screebe	11-Apr-02	smolt	Maniitsoq	1C	2003	6017	1-Sep-03	66.1	3.62	WW	exact
2003	cwt	04 47 39	Ireland	Delphi	23-Apr-02	smolt	Nuuk	1D	2003	69	13-Aug-03	69.0	3.42	GW	exact
2003	cwt	01 47 80	Ireland	Burrishoole	30-Apr-02	smolt	Qaqortoq	1F	2003	4874	26-Aug-03	66.7	3.46	GW	exact
2003	cwt	01 47 76	Ireland	Burrishoole	30-Apr-02	smolt	Qaqortoq	1F	2003	4366	29-Aug-03	66.4	3.38	GW	exact
2003	cwt	01 47 82	Ireland	Burrishoole	30-Apr-02	smolt	Qaqortoq	1F	2003	4451	3-Sep-03	57.8	1.96	GW	exact
2003	cwt	01 42 22 (102/117)	UK(E&W)	Dee	May-02	smolt	Qaqortoq	1F	2003	4141	14-Aug-03	62.3	2.34	GW	exact

	TAG INFORMATION		RELEASE INFORMATION						RECAPTURE INFORMATION						
<i>YEAR ENTERED</i>	TAG TYPE	TAG CODE (SEQ. CODE)	COUNTRY	RIVER	DATE	LIFE STAGE	COMMUNITY (AREA)	NAFO DIVISION/ICES AREA	YEAR	ENV. NO.	DATE	LENGTH (CM)	WEIGHT (KG)	GW OR WW	EXACT OR EST.
2003	streamer	green (NW20837)	Canada	NW Miramichi	2-Jun-02	smolt	Qaqortoq	1F	2003	4744	22-Aug-03	65.8	2.56	GW	exact
2003	streamer	clear (A02249)	Canada	SW Miramichi	4-Jun-02	smolt	Qaqortoq	1F	2003	4156-4190	15-Aug-03				est.
2003	streamer	green (NW32274)	Canada	SW Miramichi	May-June 2001	smolt	Maniitsoq	1C	2003	4474	Sep-03	65.8	2.56	GW	exact
2003	VIE	right eye orange	USA	Penobscot or Dennys	April - May 2002	smolt	Nuuk	1D	2003	104	14-Aug-03	61.0	2.40	GW	exact
2003	VIE	left eye orange	USA	Penobscot or Dennys	April - May 2002	smolt	Qaqortoq	1F	2003	4209	15-Aug-03	66.5	3.40	GW	exact
2003	VIE	left eye orange	USA	Penobscot or Dennys	April - May 2002	smolt	Qaqortoq	1F	2003	4236	18-Aug-03	64.8	2.50	GW	exact
2004	anchor	blue, YY 979	Canada	Miramichi	Jul-Oct 03	adult	Nuuk	1D	2004		17-Oct-04	84.0			est.
2004	anchor	A14601	Canada	Restigouche	May-June	smolt	Nuuk	1D	2004	572	3-Sep-04	65.3	3.40	WW	exact
2004	anchor	blue	Canada				Nuuk	1D	2004	316	17-Aug-04	60.0	2.22	GW	exact
2004	cwt	47 01 65	Ireland	Shannon	27-Mar-03	parr	Nuuk	1D	2004	291	17-Aug-04	61.0	2.50	GW	exact
2004	PIT	00302243	Canada	Miramichi	17-May-03	smolt	Qaqortoq	1F	2004		6-Oct-04				est.
2004	VIE	right eye pink	USA	Penobscot	22-Apr-03	smolt	Maniitsoq	1C	2004	6087	14-Sep-04	65.2	3.28	GW	exact

	TAG INFORMATION		RELEASE INFORMATION						RECAPTURE INFORMATION						
<i>YEAR ENTERED</i>	TAG TYPE	TAG CODE (SEQ. CODE)	COUNTRY	RIVER	DATE	LIFE STAGE	COMMUNITY (AREA)	NAFO DIVISION/ICES AREA	YEAR	ENV. NO.	DATE	LENGTH (CM)	WEIGHT (KG)	GW OR WW	EXACT OR EST.
2004	VIE	right eye pink	USA	Penobscot	22-Apr-03	smolt	Maniitsoq	1C	2004	6315	21-Sep-04	65.3	2.84	WW	exact
2004	VIE	left eye red	USA	Penobscot	1-May-03	smolt	Maniitsoq	1C	2004		25-Sep-04				est.
2004	VIE	left eye yellow	USA	Dennys	9-May-03	smolt	Nuuk	1D	2004	137	14-Aug-04	62.5	2.82	GW	exact
2004	VIE	left eye yellow	USA	Dennys	9-May-03	smolt	Nuuk	1D	2004	362	17-Aug-04	64.4	3.52	WW	exact
2005	streamer	A43223	Canada	SW Miramichi	May	smolt	Sisimiut	1B	2005		20-Oct-05	74.0			est.
2005	streamer	A34346	Canada	SW Miramichi	May/June 04	smolt	Qaqortoq	1F	2005			70.0			est.
2005	VIE	right eye green	USA	Penobscot	12-Apr-04	smolt	Nuuk	1D	2005	140	20-Aug-05	69.0	3.48	GW	exact
2005	VIE	right eye orange	USA	Penobscot	29-Apr-04	smolt	Maniitsoq	1C	2005	6023	13-Sep-05	68.0	3.86	GW	exact
2005	VIE	right eye orange	USA	Penobscot	29-Apr-04	smolt	Maniitsoq	1C	2005	6024	13-Sep-05	71.0	4.36	GW	exact
2005	VIE	center jaw red	unknown				Nuuk	1D	2005	186	23-Aug-05	6.40	2.24	GW	exact
2006	carlin	green, 908.009	USA	Penobscot	1-May-96	smolt	Uummanaq	1A	2006		Sep-06	70-80			est.
2006	carlin	blue, YY12,172	Canada	SW Miramichi	3-Aug-05	adult	Qaqortoq	1F	2006		26-Sep-06	92.0			est.

	TAG INFORMATION		RELEASE INFORMATION						RECAPTURE INFORMATION						
<i>YEAR ENTERED</i>	<b>TAG TYPE</b>	<b>TAG CODE (SEQ. CODE)</b>	<b>COUNTRY</b>	<b>RIVER</b>	<b>DATE</b>	<b>LIFE STAGE</b>	<b>COMMUNITY (AREA)</b>	<b>NAFO DIVISION/ICES AREA</b>	<b>YEAR</b>	<b>ENV. NO.</b>	<b>DATE</b>	<b>LENGTH (CM)</b>	<b>WEIGHT (KG)</b>	<b>GW OR WW</b>	<b>EXACT OR EST.</b>
2006	carlin	blue, YY09,968	Canada	SW Miramichi	22-Aug-05	adult	Qasigiannguit	1A	2006		27-Oct-06				est.
2006	carlin	blue, YY10,805	Canada	NW Miramichi	1-Sep-05	adult	Sisimiut (1B)	1B	2006		18-Oct-06				est.
2006	cwt	23 40 61 (01123)	Spain	Tea (Galicia)	14-Nov-03	parr	Nuuk	1D	2006	385	28-Aug-06	68.0	2.68	GW	exact
2006	streamer	clear, A78113	Canada	SW Miramichi	10-May-05	smolt	Maniitsoq	1C	2006		Sep-06				est.
2006	streamer	clear, A48507	Canada	Miramichi	30-May-05	smolt	Nuuk	1D	2006	376	28-Aug-06	65.7	2.60	GW	exact
2006	streamer	clear, A63913	Canada	Restigouche	1-Jun-05	smolt	Nuuk	1D	2006	81	12-Aug-06	58.0	1.76	GW	exact
2006	streamer	clear, A73298	Canada	Margaree	7-Jun-05	smolt	Paamuit (1E)	1E	2006			52.6			est.
2006	VIE	right eye yellow	USA	Dennys	6-Apr-05	smolt	Nuuk	1D	2006	337	28-Aug-06	65.5	3.30	GW	exact
2007	carlin	blue, YY16,697	Canada	SW Maramichi	Sept/Oct 06	adult	Nuuk	1D	2007		23-Sep-07	75.0			est.
2008	cwt	23 31 34 (17383)	Spain	R. Asón (Cantabria)	3-Nov-05	parr	Nuuk	1D	2007						est.
2007	cwt	23 41 08 (13574)	Spain	Ulla	March-06	smolt	Nuuk	1D	2007	295	19-Aug-07	64.5	2.76	GW	exact
2007	streamer	clear, VI 0822	Canada	Cains	May/June 06	smolt	Maniitsoq	1C	2007		5-Oct-07				est.



	TAG INFORMATION		RELEASE INFORMATION						RECAPTURE INFORMATION						
<i>YEAR ENTERED</i>	TAG TYPE	TAG CODE (SEQ. CODE)	COUNTRY	RIVER	DATE	LIFE STAGE	COMMUNITY (AREA)	NAFO DIVISION/ICES AREA	YEAR	ENV. NO.	DATE	LENGTH (CM)	WEIGHT (KG)	GW OR WW	EXACT OR EST.
2007	VIE	right eye green	USA	Penobscot	May-06	smolt	Paamiut	1E	2007	10163	29-Aug-07	63.5	1.98	GW	exact
2007	VIE	right eye red	USA	Penobscot	May-06	smolt	Nuuk	1D	2007	510	5-Sep-07	62.0	3.24	WW	exact
2008	carlin	464,784	USA	Penobscot	7-May-87	smolt	Narsaq	1F	2008			69.0			est.
2008	cwt	62 01 05 (03239)	UK(Scot)	North Esk	05-Apr-07	smolt	Sisimiut	1B	2008	2499	30-Sep-08	62.9	3.10	GW	exact
2008	cwt	unk	unknown				Qaqortoq	1F	2008	4090	28-Aug-08	67.9	2.94	GW	exact
2008	PIT	unk	unknown				Maniitsoq	1C	2008		1-Oct-08	70.0			est.
2008	streamer	clear, B05324	Canada	Restigouche	May/June	smolt	Sisimiut	1B	2008	2119	6-Sep-08	62.8	2.68	GW	exact
2009	carlin	green, 829.816	USA	Penobscot	29-Apr-91	smolt	Narsaq	1F	2009		23-Sep-09				est.
2009	carlin	blue, YY16,182	Canada	SW Miramichi	21-Sep-06	adult	Narsaq	1F	2009		20-Oct-09				est.
2009	carlin	green, NJ-063966	Norway	Alta	4-Jun-07	smolt	Qaqortoq	1F	2009		12-Aug-09				est.
2009	carlin	light green, NK-073312	Norway	Figgjo	15-Apr-08	smolt	65 37 N, 37 27 W	XIV	2009		12-Aug-09	40.0			est.
2009	carlin	light green, NY 069745	Norway	Eira	5-May-08	smolt	Tasiilaq	XIV	2009		3-Oct-09	61.0			est.
2009	carlin	light blue, YY17,656	Canada	SW Miramichi	16-Jul-08	adult	Sisimiut	1B	2009		15-Oct-09	75.0			est.

	TAG INFORMATION		RELEASE INFORMATION						RECAPTURE INFORMATION						
<i>YEAR ENTERED</i>	TAG TYPE	TAG CODE (SEQ. CODE)	COUNTRY	RIVER	DATE	LIFE STAGE	COMMUNITY (AREA)	NAFO DIVISION/ ICES AREA	YEAR	ENV. NO.	DATE	LENGTH (CM)	WEIGHT (KG)	GW OR WW	EXACT OR EST.
2009	carlin	light blue, YY24,460	Canada	SW Miramichi	2-Sep-08	adult	Sisimiut	1B	2009		2-Oct-09	88.0			est.
2009	cwt	42 1 32 18 1 (3585)	UK(E&W)	River Frome	24-Apr-08	smolt	Sismiut	1B	2009	2603	6-Oct-09	67.9	4.18	WW	exact
2009	cwt	47/05/37	Ireland	Bundorragha River	28-Apr-08	smolt	Sismiut	1B	2009	2553	2-Oct-09	67.3	4.40	WW	exact
2009	streamer	clear, B06584	Canada	Restigouche	17-May-08	smolt	Ivittuut	1E	2009		7-Sep-09				est.
2009	streamer	clear, B17418	Canada	Restigouche	28-May-08	smolt	Qaqortoq	1F	2009		14-Oct-09	70.0			est.
2010	cwt	47 05 61	Ireland	Bundorragha	28-Apr-09	smolt	Nuuk	1D	2010	11	10-Sep-10	665	3.62	WW	exact
2010	cwt	47 05 62	Ireland	Bundorragha	28-Apr-09	smolt	Nuuk	1D	2010	129	16-Sep-10	669	4.08	WW	exact
2010	cwt	Agency tag #13	Canada	St-Jean (Quebec, Gaspé)		smolt		1B	2010	2069	6-Sep-10	671	3.2	GW	exact
2010	cwt	59 01 84 (06829)	Norway	Dale	5/30/2009	smolt	Qaqortoq	1F	2010	4044	16-Aug-10	640	2.70	GW	exact
2010	cwt	47 05 62	Ireland	Bundorragha	28-Apr-09	smolt	Qaqortoq	1F	2010	4061	17-Aug-10	640	2.78	GW	exact
2010	cwt	47 05 60	Ireland	Bundorragha	28-Apr-09	smolt	Qaqortoq	1F	2010	4220	23-Aug-10	650	2.50	GW	exact
2010	VIE	REG	USA	Penobscot	4/13 - 4/21/09	smolt	Nuuk	1D	2010	95	8-Sep-10	682	4.74	WW	exact

	TAG INFORMATION		RELEASE INFORMATION						RECAPTURE INFORMATION						
<i>YEAR ENTERED</i>	TAG TYPE	TAG CODE (SEQ. CODE)	COUNTRY	RIVER	DATE	LIFE STAGE	COMMUNITY (AREA)	NAFO DIVISION/ICES AREA	YEAR	ENV. NO.	DATE	LENGTH (CM)	WEIGHT (KG)	GW OR WW	EXACT OR EST.
2010	streamer	clear, B19964	Canada	Restigouche	5/21/2009	smolt	Ukivit	1E	2010		27-Aug-10	650	2.43	GW	est.
2010	streamer	clear, B47437	Canada	SW Miramichi	5/20/2009	smolt	Kangilinnguit	1E	2010		19-Sep-10	640	4.00	WW	est.
2011	carlin	YY25,646 (blue)	Canada	Miramichi	Jun-Sep 2010	adult	Nuuk	1D	2011	301	12-Aug-11	817	4.66	GW	exact
2011	carlin	YY30,149 (blue)	Canada	Miramichi	Jul-Oct 2010	adult	Maniitsoq	1C	2011		26-Oct-11	950	9.20	GW	est.
2011	streamer	B-47437 (clear)	Canada	SW Miramichi	May-Jun 2009	smolt	Itissaaq	1E	2010		19-Sep-10	640	4.00	WW	est.
2011	streamer	B-19964 (clear)	Canada	Restigouche	May-Jun 2009	smolt	Paamiut	1E	2010		Sep-10	650	2.43	GW	exact
2011	acoustic	Vemco 57948	Canada	Riviere St Jean	Jun-10	kelt	Nuuk	1D	2011	514	22-Sep-11	850	6.16	GW	exact
2011	PIT	na	unknown				Nuuk	1D	2011	158	26-Sep-11	693	4.50	WW	exact
2012	carlin	YY34,105 (light blue)	Canada	NW Miramichi River	10/9/2011	adult	Nanortalik	1F	2012			87	5.50	WW	est.
2012	spaghetti	A-01698 (red)	Canada	Campbellton River	5/11/2012	adult		1D	2012		11-Aug-12	57			est.
2012	carlin	YY 32,569 (light blue)	Canada	SW Miramichi River	8/26/2011	adult	Nuuk	1D	2012		8-Oct-12	94	9.14	WW	est.
2012	carlin	YY35,191 (light blue)	Canada	SW Miramichi River	10/8/2011	adult	Nuuk	1D	2012		24-Oct-12	85	3.50	WW	est.

	TAG INFORMATION		RELEASE INFORMATION						RECAPTURE INFORMATION						
<i>YEAR ENTERED</i>	<b>TAG TYPE</b>	<b>TAG CODE (SEQ. CODE)</b>	<b>COUNTRY</b>	<b>RIVER</b>	<b>DATE</b>	<b>LIFE STAGE</b>	<b>COMMUNITY (AREA)</b>	<b>NAFO DIVISION/ ICES AREA</b>	<b>YEAR</b>	<b>ENV. NO.</b>	<b>DATE</b>	<b>LENGTH (CM)</b>	<b>WEIGHT (KG)</b>	<b>GW OR WW</b>	<b>EXACT OR EST.</b>
2012	carlin	R 695532 S (light green)	Sweden	Lagan	4/24/2011	smolt	Qaqortoq	1F	2012		27-Oct-12	75	5.00	WW	est.
2012	carlin	YY35,639 (light blue)	Canada	SW Miramichi River	9/24/2011	adult	Aasiaat	1B	2012		12-Oct-12	75	12.00	WW	est.
2013	carlin	NL 083810 (green)	Norway	Imsa	15-Mar-12	smolt	Sulussugutip allanngua (btwn Maniitsoq and Napasq)	1C	2013				3.20	GW	est.
2013	carlin	H7 (front) Return to MAFF (back) (green)	UK(E&W)	Ouse	1975	smolt	Aasiaat	1B							
2013	carlin	YY37,601	Canada	Miramichi	9/24/2012	kelt	Aasiaat	1B	2013		20-Oct-13		10.50	WW	est.
2014	carlin	light blue (YY31.575)	Canada	Northwest Miramichi	8/12/2013	adult	Sisimuit	1B	2014			850	13.90	WW	est.
2014	carlin	dark blue (RDH W40190)	Canada	East River	5/10/1979	smolt			1970's						
2014	carlin	dark blue (RDH X41376)	Canada	S. John River	4/23/1981	smolt	Kaangaamiut area	1C	1987-1988						
2014	carlin	dark blue (RDH X74055)	Canada	LeHave	5/12/1981	smolt	Kaangaamiut area	1C	1987-1988						

	TAG INFORMATION		RELEASE INFORMATION						RECAPTURE INFORMATION						
<i>YEAR ENTERED</i>	<b>TAG TYPE</b>	<b>TAG CODE (SEQ. CODE)</b>	<b>COUNTRY</b>	<b>RIVER</b>	<b>DATE</b>	<b>LIFE STAGE</b>	<b>COMMUNITY (AREA)</b>	<b>NAFO DIVISION/ ICES AREA</b>	<b>YEAR</b>	<b>ENV. NO.</b>	<b>DATE</b>	<b>LENGTH (CM)</b>	<b>WEIGHT (KG)</b>	<b>GW OR WW</b>	<b>EXACT OR EST.</b>
2014	carlin	dark blue (RDH Y5714)	Canada	NW Miramichi	10/15/1992	adult	Kaangaamiut area	1C	1987-1988						
2014	carlin	dark blue (RDH Y7326)	Canada	SW Miramichi	8/23/1992	adult	Kaangaamiut area	1C	1987-1988						
2014	carlin	dark blue (RDH Z42712)	Canada	New ALbany	5/2/1983	smolt	Kaangaamiut area	1C	1987-1988						
2014	carlin	light blue (YY34,811)	Canada	NW Miramichi	6/29/2013	adult	Qarajat Iluami	1D	2014		10/16/2014	730	1/4/1900		est.
2014	carlin	light blue (YY37,601)	Canada	SW Miramichi	9/24/2012	adult	Aasiaat area	1B	2013		10/20/2013				
2014	floy	yellow (A-00814)	Canada		6/1/2014	adult	Narsaq	1F	2014		10/27/2014				
2014	carlin	black (RFP2792)	France				Arsuk Area	1E							
2014	carlin	light blue/light green (58232)	Norway	Figgjo	5/18/1977	smolt	Arsuk Area	1E	2000-2001						
2014	carlin	light green (98925)	Norway	Drammenselva	5/6/1986	smolt	Qeqertarsuatsiaat	1D	1988-1989						
2014	carlin	green (24404)	UK(Scot)	North Esk	5/8/1981	smolt	Kaangaamiut area	1C	1987-1988						

	TAG INFORMATION		RELEASE INFORMATION						RECAPTURE INFORMATION						
<i>YEAR ENTERED</i>	<b>TAG TYPE</b>	<b>TAG CODE (SEQ. CODE)</b>	<b>COUNTRY</b>	<b>RIVER</b>	<b>DATE</b>	<b>LIFE STAGE</b>	<b>COMMUNITY (AREA)</b>	<b>NAFO DIVISION/ ICES AREA</b>	<b>YEAR</b>	<b>ENV. NO.</b>	<b>DATE</b>	<b>LENGTH (CM)</b>	<b>WEIGHT (KG)</b>	<b>GW OR WW</b>	<b>EXACT OR EST.</b>
2014	carlin	green (USA 145,063)	USA	Union	5/3/1979	smolt	Qeqertarsuatsiaat	1D	1988-1989						
2014	carlin	green (USA 217175)	USA	Penobscot	5/7/1980	smolt	Qeqertarsuatsiaat	1D	1988-1989						
2014	carlin	green (USA 24630)	USA	Penobscot	5/9/1984	smolt	Kaangaamiut area	1C	1987-1988						
2014	carlin	green (USA 289697)	USA	Penobscot	5/4/1981	smolt	Kaangaamiut area	1C	1987-1988						
2014	carlin	green (USA 291510)	USA	Penobscot	5/4/1981	smolt	Kaangaamiut area	1C	1987-1988						
2014	carlin	green (USA 398,712)	USA	Penobscot	5/9/1986	smolt	Qeqertarsuatsiaat	1D	1988-1989						
2014	carlin	green (USA 398,917)	USA	Penobscot	5/9/1986	smolt	Qeqertarsuatsiaat	1D	1988-1989						
2014	carlin	green (USA-CTR 167,495)	USA	Conneticut	4/25/1977	smolt	Sisimiut	1B	1978-1982						
2015	carlin	322,343 (green)	USA	Penobscot	8-May-86	smolt	Paamuit	1E							
2015	carlin	846,920 (green)	USA	Penobscot	29-Apr-91	smolt	Paamuit	1E							
2015	carlin	42501 (green)	Canada				Paamuit	1E							

	TAG INFORMATION		RELEASE INFORMATION						RECAPTURE INFORMATION						
<i>YEAR ENTERED</i>	<b>TAG TYPE</b>	<b>TAG CODE (SEQ. CODE)</b>	<b>COUNTRY</b>	<b>RIVER</b>	<b>DATE</b>	<b>LIFE STAGE</b>	<b>COMMUNITY (AREA)</b>	<b>NAFO DIVISION/ICES AREA</b>	<b>YEAR</b>	<b>ENV. NO.</b>	<b>DATE</b>	<b>LENGTH (CM)</b>	<b>WEIGHT (KG)</b>	<b>GW OR WW</b>	<b>EXACT OR EST.</b>
2015	carlin	AA 26325 (light green)	Canada	Musquodoboit	1985	smolt	Nanortalik	1F							est.
2015	carlin	R 799099 S (light green)	Sweden	Nissan	4/14/2014	smolt	Qaqortoq	1F	2015		20-Sep-15	65	2.55	GW	est.
2015	carlin	MSA 01,153 (blue)	Canada	Miramichi	7/11/2014	adult	Paamiut	1E	2015		23-Oct-15	74	4.18	GW	est.
2016	cwt	01 42 87	UK(E&W)	Dee	May-15	smolt	Paamuit	1E	2016	10079	09/21/16	625	2.36	GW	exact
2016	cwt	07 47 14	Ireland	Corrib	9-Apr-15	smolt	Qaqortoq	1F	2016	4086	08/23/16	577	2.10	GW	exact
2016	carlin	blue (A59055)	Canada	LaHave	5/21/1974	smolt	Arsuk Area	1E							1975-1980
2016	carlin	blue (G48113)	Canada	St. John River	4/30/1973	smolt	Arsuk Area	1E							1975-1980
2016	carlin	blue (RHD M97851)	Canada	St. John River	5/3/1979	smolt	Arsuk Area	1E							1975-1980
2016	carlin	Green (DD20701)	Canada	Saint Mary's	5/25/1989	smolt	Arsuk Area	1E							1975-1980
2016	carlin	Green (BB62280)	Canada	Middle River	5/26/1987	smolt	Arsuk Area	1E							1975-1980
2016	carlin	Brown (B334255)	Norway	Imsa	5/16/1977	smolt	Arsuk Area	1E							1975-1980
2016	carlin	Dark Green (W1346)	UK(Scot)	North Esk	5/26/1977	smolt	Arsuk Area	1E							1975-1980

	TAG INFORMATION		RELEASE INFORMATION						RECAPTURE INFORMATION						
<i>YEAR ENTERED</i>	<b>TAG TYPE</b>	<b>TAG CODE (SEQ. CODE)</b>	<b>COUNTRY</b>	<b>RIVER</b>	<b>DATE</b>	<b>LIFE STAGE</b>	<b>COMMUNITY (AREA)</b>	<b>NAFO DIVISION/ ICES AREA</b>	<b>YEAR</b>	<b>ENV. NO.</b>	<b>DATE</b>	<b>LENGTH (CM)</b>	<b>WEIGHT (KG)</b>	<b>GW OR WW</b>	<b>EXACT OR EST.</b>
2016	carlin	Light Green (40825)	UK(Scot)	North Esk	5/22/1982	smolt	Arsuk Area	1E	1975-1980						
2016	carlin	Green (USA 15,812)	USA	Penobscot	5/7/1974	smolt	Arsuk Area	1E	1975-1980						
2016	carlin	Green (USA 61 466)	USA	Penobscot	5/5/1983	smolt	Arsuk Area	1E	1975-1980						
2016	carlin	Light blue (YY00,898)	Canada	Southwest Miramichi	9/18/2003	adult	Narsaq area	1F	2004						
2016	radio	white (360 027)	USA	Androscoggin	5/14/2015	smolt	Kangaamiut	1C	2016						
2017	spaghetti	green (AR3284)	Canada				Qaqortoq	1F	2017	4004	08/23/17	795	4.72	GW	exact
2017	VIE	right eye green	USA	Penobscot	5/2/2016	smolt	Qaqortoq	1F	2017	4021	08/24/17	650	2.90	GW	exact
2017	VIE	left eye red	USA	Penobscot	4/28/2016	smolt	Qaqortoq	1F	2017	4031	08/24/17	671	3.08	GW	exact
2017	carlin	blue (YY41, 797)	Canada	Southwest Miramichi	7/14/2016	adult	Sisimiut	1B	2017	2162	09/23/17	856	6.78	GW	exact
2017	cwt	470763	Ireland	Burrishoole	3/5/2016	smolt	Sisimiut	1B	2017	2082	09/17/17	646	2.91	GW	exact
2017	cwt	470766	Ireland	Bundorragha	4/29/2016	smolt	Maniitsoq	1C	2017	6385	09/29/17	634	2.97	GW	exact
2017	carlin	YY42964	Canada	Northwest Miramichi	10/2/2016	adult	Qaqortoq	1F					8.50	WW	est.
2017	carlin	blue (RDH W95477)	Canada	Tobique River	5/2/1980	smolt	Arsuk	1E	circa 2010			570	3.1	WW	est.



	TAG INFORMATION		RELEASE INFORMATION						RECAPTURE INFORMATION						
<i>YEAR ENTERED</i>	TAG TYPE	TAG CODE (SEQ. CODE)	COUNTRY	RIVER	DATE	LIFE STAGE	COMMUNITY (AREA)	NAFO DIVISION/ ICES AREA	YEAR	ENV. NO.	DATE	LENGTH (CM)	WEIGHT (KG)	GW OR WW	EXACT OR EST.
2017	carlin	light blue (YY42,764)	Canada	Northwest Miramichi	7/19/2016	adult	Sisimiut	1B	10/19/2017			820	8	WW	est.
2018	carlin	blue (X87060 RDH)	Canada	Middle River	5/27/1981	smolt	Arsuk	1E	1975						
2018	spaghetti	blue (AR4535)	Canada				Nanortalik	1F	2018		9/19/18		5.8	WW	est.

**Table 6. Reporting groups identified within the North Atlantic-wide Atlantic salmon (*Salmo salar*) single nucleotide polymorphism genetic baseline. See Figure 4 for reporting group locations.**

North America		Europe	
Reporting Group	Code	Reporting Group	Code
Anticosti	ANT	Baltic Sea	BAL
Avalon Peninsula	AVA	Barents-White Seas	BAR
Burin Peninsula	BPN	United Kingdom/Ireland	BRI
Eastern Nova Scotia	ENS	European Broodstock	EUB
Fortune Bay, Newfoundland	FTB	France	FRN
Gaspé Peninsula	GAS	Greenland	GL
Gulf of St. Lawrence	GUL	Iceland	ICE
Inner Bay of Fundy	IBF	Northern Norway	NNO
Labrador Central	LAC	Southern Norway	SNO
Labrador South	LAS	Spain	SPN
Lake Melville	MEL		
Newfoundland 1	NF1		
Newfoundland 2	NF2		
Northern Newfoundland	NNF		
St. Lawrence North Shore – Lower	QLS		
Quebec City Region	QUE		
St. John River & Aquaculture	SJR		
Ungava Bay	UNG		
Maine, United States	USA		
Western Newfoundland	WNF		
Western Nova Scotia	WNS		

**Table 7. The continental proportions of North American (NA) and European (E) Atlantic salmon (*Salmo salar*) caught at West Greenland in 2019 by Northwest Atlantic Fisheries Organization (NAFO) division.**

NAFO Division	Fishing dates	Number		Totals	Percentages	
		NA	E		NA	E
1B	Aug 26 – Sep 24	314	51	365	86.0	14.0
1C	Sep 09 – Sep 17	249	174	423	58.9	41.1
1D	Sep 10 and Sep 20	17	19	36	47.2	52.8
1F	Aug 16 – Sep 13	186	61	247	75.3	24.7
TOTAL		766	305	1071	71.5	28.5

**Table 8. The estimated number, weighted by catch, of North American (NA) and European (E) Atlantic salmon (*Salmo salar*) caught at West Greenland by year from 1982-2019 and the proportion of the catch by weight. Numbers are rounded to the nearest hundred fish. Continent of origin assignments were based on scale characteristics until 1995, scale characteristics and DNA until 2001, and DNA only from 2002 onwards. No samples were collected in 1993 or 1994.**

	Proportion weighted by catch		Numbers of salmon caught	
	NA	E	NA	E
1982	57	43	192,200	143,800
1983	40	60	39,500	60,500
1984	54	46	48,800	41,200
1985	47	53	143,500	161,500
1986	59	41	188,300	131,900
1987	59	41	171,900	126,400
1988	43	57	125,500	168,800
1989	55	45	65,000	52,700
1990	74	26	62,400	21,700
1991	63	37	111,700	65,400
1992	45	55	46,900	38,500
1993	-	-	-	-
1994	-	-	-	-
1995	67	33	21,400	10,700
1996	70	30	22,400	9,700
1997	85	15	18,000	3,300
1998	79	21	3,100	900
1999	91	9	5,700	600
2000	65	35	5,100	2,700
2001	67	33	9,400	4,700
2002	69	31	2,300	1,000

**Table 8, continued. The estimated number, weighted by catch, of North American (NA) and European (E) Atlantic salmon (*Salmo salar*) caught at West Greenland by year from 1982-2019 and the proportion of the catch by weight. Numbers are rounded to the nearest hundred fish. Continent of origin assignments were based on scale characteristics until 1995, scale characteristics and DNA until 2001, and DNA only from 2002 onwards. No samples were collected in 1993 or 1994.**

	Proportion weighted by catch		Numbers of salmon caught	
	NA	E	NA	E
2003	64	36	2,600	1,400
2004	72	28	3,900	1,500
2005	74	26	3,500	1,200
2006	69	31	4,000	1,800
2007	76	24	6,100	1,900
2008	86	14	8,000	1,300
2009	89	11	7,000	800
2010	80	20	10,000	2,600
2011	93	7	7,500	600
2012	79	21	7,800	2,100
2013	82	18	11,500	2,700
2014	72	28	12,800	5,400
2015	79	21	13,500	3,900
2016	64	36	5,100	3,300
2017	74	26	6,100	2,200
2018	80	20	10,600	2,600
2019	72	28	6,800	2,600

**Table 9. Bayesian proportional mean mixture composition estimates for the West Greenland Atlantic salmon (*Salmo salar*) fishery by Northwest Atlantic Fisheries Organization (NAFO) division sampled in 2019 using the single nucleotide polymorphism range-wide baseline. Regions correspond to reporting groups identified in Table 6 and Figure 4. Mean estimates provided with 95% credible interval in parentheses. Estimates of mixture contributions not supported by significant individual assignments ( $P > 0.8$ ) are represented as zero. Credible intervals with a lower bound of zero (gray font), or close to zero, indicate little support for the mean assignment value.**

<b>Reporting Group</b>	<b>ROO</b>	<b>NAFO 1B</b>	<b>NAFO 1C</b>	<b>NAFO 1D</b>	<b>NAFO 1F</b>	<b>Overall</b>
Baltic Sea	EUR	0.0	0.0	0.0	0.0	0.0
Barents-White Seas	EUR	0.0	0.0	0.0	0.0	0.0
European Broodstock	EUR	0.0	0.0	0.0	0.0	0.0
France	EUR	0.0	0.2 (0.0, 0.9)	0.0	0.0	0.1 (0.0, 0.3)
Greenland	EUR	0.0	0.0	0.0	0.0	0.0
Iceland	EUR	0.0	0.0	0.0	0.0	0.0
Northern Norway	EUR	0.0	0.0	0.0	0.0	0.0
Southern Norway	EUR	0.0	0.0	0.0	0.0	0.0
Spain	EUR	0.5 (0.0, 1.6)	0.0	0.0	0.4 (0.0, 1.5)	0.2 (0.0, 0.6)
<b>United Kingdom/Ireland</b>	<b>EUR</b>	<b>13.5 (10.2, 17.2)</b>	<b>40.7 (36.1, 45.4)</b>	<b>51.7 (35.8, 67.3)</b>	<b>24.0 (18.9, 29.5)</b>	<b>28.2 (25.6, 31.0)</b>
<b>Anticosti</b>	<b>NA</b>	<b>0.0</b>	<b>1.5 (0.5, 2.9)</b>	<b>0.0 (0.0, 0.0)</b>	<b>1.6 (0.4, 3.7)</b>	<b>0.9 (0.4, 1.7)</b>
Avalon Peninsula	NA	0.0	0.0	0.0	0.0	0.0
Burin Peninsula	NA	0.0	0.0	0.0	0.0	0.0
Eastern Nova Scotia	NA	0.0	0.0	0.0	0.9 (0.1, 2.5)	0.4 (0.1, 0.9)
Fortune Bay	NA	0.0	0.0	0.0	0.0	0.0
<b>Gaspé Peninsula</b>	<b>NA</b>	<b>20.1 (15.7, 24.7)</b>	<b>15.3 (11.8, 19.2)</b>	<b>24.8 (12.2, 40.1)</b>	<b>20.8 (15.4, 26.7)</b>	<b>18.6 (16.1, 21.2)</b>
<b>Gulf of St. Lawrence</b>	<b>NA</b>	<b>19.2 (14.9, 23.8)</b>	<b>12.1 (8.9, 15.6)</b>	<b>2.8 (0.0, 10.8)</b>	<b>14.3 (9.8, 19.3)</b>	<b>14.2 (12.0, 16.6)</b>
Inner Bay of Fundy	NA	0.0	0.0	0.0	0.0	0.0
<b>Labrador Central</b>	<b>NA</b>	<b>7.0 (3.8, 10.9)</b>	<b>5.0 (2.8, 7.6)</b>	<b>7.3 (0.3, 18.1)</b>	<b>3.3 (1.3, 6.2)</b>	<b>5.4 (3.9, 7.2)</b>
<b>Labrador South</b>	<b>NA</b>	<b>19.1 (14.6, 23.9)</b>	<b>11.8 (8.6, 15.3)</b>	<b>0.0</b>	<b>12.6 (8.7, 17.2)</b>	<b>13.5 (11.4, 15.8)</b>
<b>Lake Melville</b>	<b>NA</b>	<b>1.6 (0.3, 3.7)</b>	<b>1.5 (0.5, 3.1)</b>	<b>0.0</b>	<b>0.0</b>	<b>1.5 (0.8, 2.6)</b>
<b>Maine, United States</b>	<b>NA</b>	<b>1.7 (0.6, 3.4)</b>	<b>1.4 (0.5, 2.8)</b>	<b>0.0</b>	<b>3.2 (1.4, 5.8)</b>	<b>1.9 (1.2, 2.9)</b>
<b>Newfoundland 1</b>	<b>NA</b>	<b>0.6 (0.1, 1.6)</b>	<b>0.0</b>	<b>0.0</b>	<b>2.1 (0.5, 4.3)</b>	<b>0.7 (0.2, 1.4)</b>
<b>Newfoundland 2</b>	<b>NA</b>	<b>0.8 (0.1, 2.1)</b>	<b>0.0</b>	<b>0.0</b>	<b>0.9 (0.1, 2.5)</b>	<b>0.9 (0.4, 1.6)</b>
Northern Newfoundland	NA	0.0	0.0	0.0	0.4 (0.0, 1.5)	0.1 (0.0, 0.4)
<b>Quebec City Region</b>	<b>NA</b>	<b>2.6 (0.7, 5.0)</b>	<b>1.9 (0.7, 3.7)</b>	<b>0.0</b>	<b>3.5 (1.1, 6.8)</b>	<b>2.3 (1.3, 3.7)</b>
St. John River & AQ	NA	0.0	0.0	0.0	0.0	0.0

<b>Reporting Group</b>	<b>ROO</b>	<b>NAFO 1B</b>	<b>NAFO 1C</b>	<b>NAFO 1D</b>	<b>NAFO 1F</b>	<b>Overall</b>
St. Lawrence N. Shore Lower	NA	4.4 (2.4, 7.0)	2.3 (1.0, 4.1)	7.8 (1.2, 18.8)	2.9 (1.1, 5.5)	3.7 (2.6, 5.0)
Ungava	NA	6.6 (4.3, 9.4)	2.1 (1.0, 3.7)	0.0	6.1 (3.4, 9.4)	4.6 (3.4, 5.9)
Western Newfoundland	NA	2.2 (0.9, 4.1)	3.0 (1.5, 5.1)	0.0	2.3 (0.7, 4.6)	2.3 (1.4, 3.4)
Western Nova Scotia	NA	0.0	0.0	0.0	0.0	0.0



**Table 10. Annual mean fork lengths and whole weights by continent of origin (NA = North American and E = European) and sea age (1sw = 1 sea-winter, 2sw = 2 sea-winter, and ps = previous spawner) of Atlantic salmon (*Salmo salar*) caught at West Greenland from 1969-2019. No samples were collected in 1977, 1993, or 1994. The 2017 European previous spawner value is based on two fish, and the 2019 European previous spawner value is based on one fish. Note that the mean fork lengths and weights have not been corrected to adjust for the annual variation in the timing of the sampling program.**

	Whole weight (kg)									Fork length (cm)					
	Sea age and origin						All sea ages			Sea age and origin					
	1SW		2SW		PS		NA	E	Total	1SW		2SW		PS	
NA	E	NA	E	NA	E	NA				E	NA	E	NA	E	NA
1969	3.12	3.76	5.48	5.80	-	5.13	3.25	3.86	3.58	65.0	68.7	77.0	80.3	-	75.3
1970	2.85	3.46	5.65	5.50	4.85	3.80	3.06	3.53	3.28	64.7	68.6	81.5	82.0	78.0	75.0
1971	2.65	3.38	4.30	-	-	-	2.68	3.38	3.14	62.8	67.7	72.0	-	-	-
1972	2.96	3.46	5.85	6.13	2.65	4.00	3.25	3.55	3.44	64.2	67.9	80.7	82.4	61.5	69.0
1973	3.28	4.54	9.47	10.00	-	-	3.83	4.66	4.18	64.5	70.4	88.0	96.0	61.5	-
1974	3.12	3.81	7.06	8.06	3.42	-	3.22	3.86	3.58	64.1	68.1	82.8	87.4	66.0	-
1975	2.58	3.42	6.12	6.23	2.60	4.80	2.65	3.48	3.12	61.7	67.5	80.6	82.2	66.0	75.0
1976	2.55	3.21	6.16	7.20	3.55	3.57	2.75	3.24	3.04	61.3	65.9	80.7	87.5	72.0	70.7
1977	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1978	2.96	3.50	7.00	7.90	2.45	6.60	3.04	3.53	3.35	63.7	67.3	83.6	-	60.8	85.0
1979	2.98	3.50	7.06	7.60	3.92	6.33	3.12	3.56	3.34	63.4	66.7	81.6	85.3	61.9	82.0
1980	2.98	3.33	6.82	6.73	3.55	3.90	3.07	3.38	3.22	64.0	66.3	82.9	83.0	67.0	70.9
1981	2.77	3.48	6.93	7.42	4.12	3.65	2.89	3.58	3.17	62.3	66.7	82.8	84.5	72.5	-
1982	2.79	3.21	5.59	5.59	3.96	5.66	2.92	3.43	3.11	62.7	66.2	78.4	77.8	71.4	80.9
1983	2.54	3.01	5.79	5.86	3.37	3.55	3.02	3.14	3.10	61.5	65.4	81.1	81.5	68.2	70.5
1984	2.64	2.84	5.84	5.77	3.62	5.78	3.20	3.03	3.11	62.3	63.9	80.7	80.0	69.8	79.5
1985	2.50	2.89	5.42	5.45	5.20	4.97	2.72	3.01	2.87	61.2	64.3	78.9	78.6	79.1	77.0
1986	2.75	3.13	6.44	6.08	3.32	4.37	2.89	3.19	3.03	62.8	65.1	80.7	79.8	66.5	73.4
1987	3.00	3.20	6.36	5.96	4.69	4.70	3.10	3.26	3.16	64.2	65.6	81.2	79.6	74.8	74.8
1988	2.83	3.36	6.77	6.78	4.75	4.64	2.93	3.41	3.18	63.0	66.6	82.1	82.4	74.7	73.8
1989	2.56	2.86	5.87	5.77	4.23	5.83	2.77	2.99	2.87	62.3	64.5	80.8	81.0	73.8	82.2
1990	2.53	2.61	6.47	5.78	3.90	5.09	2.67	2.72	2.69	62.3	62.7	83.4	81.1	72.6	78.6
1991	2.42	2.54	5.82	6.23	5.15	5.09	2.57	2.79	2.65	61.6	62.7	80.6	82.2	81.7	80.0
1992	2.54	2.66	6.49	6.01	4.09	5.28	2.86	2.74	2.81	62.3	63.2	83.4	81.1	77.4	82.7
1995	2.37	2.67	6.09	5.88	3.71	4.98	2.45	2.75	2.56	61.0	63.2	81.3	81.0	70.9	81.3
1996	2.63	2.86	6.50	6.30	4.98	5.44	2.83	2.90	2.88	62.8	64.0	81.4	81.1	77.1	79.4
1997	2.57	2.82	7.95	6.11	4.82	6.9	2.63	2.84	2.71	62.3	63.6	85.7	84.0	79.4	87.0
1998	2.72	2.83	6.44	-	3.28	4.77	2.76	2.84	2.78	62.0	62.7	84.0	-	66.3	76.0
1999	3.02	3.03	7.59	-	4.20	-	3.09	3.03	3.08	63.8	63.5	86.6	-	70.9	-
2000	2.47	2.81	-	-	2.58	-	2.47	2.81	2.57	60.7	63.2	-	-	64.7	-
2001	2.89	3.03	6.76	5.96	4.41	4.06	2.95	3.09	3.00	63.1	63.7	81.7	79.1	75.3	72.1
2002	2.84	2.92	7.12	-	5.00	-	2.89	2.92	2.90	62.6	62.1	83.0	-	75.8	-
2003	2.94	3.08	8.82	5.58	4.04	-	3.02	3.10	3.04	63	64.4	86.1	78.3	71.4	-
2004	3.11	2.95	7.33	5.22	4.71	6.48	3.17	3.22	3.18	64.7	65.0	86.2	76.4	77.6	88.0
2005	3.19	3.33	7.05	4.19	4.31	2.89	3.31	3.33	3.31	65.9	66.4	83.3	75.5	73.7	62.3
2006	3.10	3.25	9.72	-	5.05	3.67	3.25	3.26	3.24	65.3	65.3	90.0	-	76.8	69.5
2007	2.89	2.87	6.19	6.47	4.94	3.57	2.98	2.99	2.98	63.5	63.3	80.9	80.6	76.7	71.3
2008	3.04	3.03	6.35	7.47	3.82	3.39	3.08	3.07	3.08	64.6	63.9	80.1	85.5	71.1	73.0
2009	3.28	3.40	7.59	6.54	5.25	4.28	3.48	3.67	3.50	64.9	65.5	84.6	81.7	75.9	73.5
2010	3.44	3.24	6.40	5.45	4.17	3.92	3.47	3.28	3.42	66.7	65.2	80.0	75.0	72.4	70.0
2011	3.30	3.18	5.69	4.94	4.46	5.11	3.39	3.49	3.40	65.8	64.7	78.6	75.0	73.7	76.3
2012	3.34	3.38	6.00	4.51	4.65	3.65	3.44	3.40	3.44	65.4	64.9	75.9	70.4	72.8	68.9
2013	3.33	3.16	6.43	4.51	3.64	5.38	3.39	3.20	3.35	66.2	64.6	81.0	72.8	69.9	73.6
2014	3.25	3.02	7.60	6.00	4.47	5.42	3.39	3.13	3.32	65.6	63.6	86.0	78.7	73.6	83.5
2015	3.36	3.13	7.52	7.10	4.53	3.81	3.42	3.18	3.37	65.6	64.4	84.1	82.5	74.2	67.2
2016	3.18	2.79	7.77	5.18	4.03	4.12	3.32	2.89	3.18	65.2	62.6	85.1	76.0	72.2	70.9
2017	3.42	3.31	6.50	3.69	4.94	8.00	3.50	3.36	3.46	66.6	64.8	85.1	72.4	76.7	81.8
2018	2.91	2.93	9.27	5.59	4.53	-	2.97	3.00	2.97	63.8	63.9	87.5	76.3	77.1	-
2019	2.93	2.89	6.62	6.27	4.01	2.76	3.01	2.83	2.96	63.9	63.4	78.4	76.8	72.1	6.21

**Table 11. Mean fork lengths (cm) and whole weight (kg) by sea age (1sw = 1 sea-winter and 2sw = 2 sea-winter), continent of origin and Northwest Atlantic Fisheries Organization (NAFO) division for Atlantic salmon (*Salmo salar*) caught at West Greenland in 2019 with corresponding standard deviation (S.D.). Table does not include salmon of unknown age, origin, fork length, or weight.**

NAFO Div.	1 SW		2 SW		Previous spawners		Fork length (cm) (S.D.)	All sea ages		No.	
	Fork length (cm) (S.D.)	Whole weight (kg) (S.D.)	Fork length (cm) (S.D.)	Whole weight (kg) (S.D.)	Fork length (cm) (S.D.)	Whole weight (kg) (S.D.)		Fork length (cm) (S.D.)	Whole weight (kg) (S.D.)		
North American and European											
1B	63.4 (3.4)	2.82 (0.64)	75.8 (16.2)	5.97 (4.36)	75.9 (8.3)	4.56 (1.42)	63.8 (4.2)	354	2.87 (0.78)	353	63.4 (3.4)
1C	63.1 (3.2)	2.90 (0.52)	68.1 -	3.81 -	64.5 (4.0)	2.74 (0.28)	63.1 (3.2)	423	2.91 (0.52)	423	63.1 (3.2)
1F	63.9 (3.8)	2.98 (0.62)	77.9 (11.2)	6.58 (2.95)	70.5 (8.9)	3.83 (1.96)	65.0 (5.8)	244	3.19 (1.22)	242	63.9 (3.8)
All Areas	63.4 (3.4)	2.89 (0.59)	76.8 (11.1)	6.27 (2.95)	71.7 (8.8)	3.95 (1.68)	63.8 (4.4)	1021	2.96 (0.83)	1018	63.4 (3.4)
North American											
1B	63.6 (3.4)	2.82 (0.66)	64.3 -	2.89 -	75.9 (8.3)	4.56 (1.42)	63.9 (4.1)	303	2.87 (0.74)	302	63.6 (3.4)
1C	63.9 (3.2)	3.00 (0.56)	68.1 -	3.81 -	65.8 (4.7)	2.73 (0.40)	64.0 (3.2)	249	3.00 (0.56)	249	63.9 (3.2)
1F	64.6 (3.6)	3.03 (0.63)	82.5 (8.9)	7.72 (2.73)	70.5 (8.9)	3.83 (1.96)	65.8 (5.9)	183	3.27 (1.27)	183	64.6 (3.6)
All Areas	63.9 (3.4)	2.93 (0.62)	78.4 (10.7)	6.62 (3.08)	72.1 (8.7)	4.01 (1.70)	64.4 (4.4)	735	3.01 (0.87)	734	63.9 (3.4)
European											
1B	62.4 (3.4)	2.78 (0.53)	87.2 -	9.05 -	- -	- -	62.9 (4.9)	51	2.90 (1.02)	51	62.4 (3.4)
1C	62.0 (2.8)	2.77 (0.42)	- -	- -	62.1 -	2.76 -	62.0 (2.8)	174	2.77 (0.42)	174	62.0 (2.8)
1F	62.0 (3.7)	2.81 (0.59)	70.9 (11.6)	4.88 (2.68)	- -	- -	62.6 (4.9)	61	2.95 (0.99)	59	62.0 (3.7)
All Areas	63.4 (3.4)	2.89 (0.59)	76.8 (11.1)	6.27 (2.95)	- -	- -	62.3 (3.7)	286	2.83 (0.70)	284	63.4 (3.4)

**Table 12. The river age (smolt age) composition (%) of Atlantic salmon (*Salmo salar*) by continent of origin (NA = North American and E = European) and Northwest Atlantic Fisheries Organization (NAFO) division caught at West Greenland in 2019.**

NAFO Division	Origin	River age (%)							Total No.
		1	2	3	4	5	6	7	
1B	NA	0.7	26.0	30.6	26.7	14.2	1.4	0.3	288
	E	8.3	56.3	29.2	6.3	0.0	0.0	0.0	48
		1.8	30.4	30.4	23.8	12.2	1.2	0.3	336
1C	NA	0.4	25.3	36.1	24.5	12.9	0.8	0.0	249
	E	5.8	70.5	19.1	4.6	0.0	0.0	0.0	173
		2.6	43.8	29.1	16.4	7.6	0.5	0.0	422
1F	NA	0.6	30.5	30.5	24.3	14.1	0.0	0.0	177
	E	11.7	35.0	35.0	16.7	1.7	0.0	0.0	60
		3.4	31.6	31.6	22.4	11.0	0.0	0.0	237
All Areas	NA	0.6	26.9	32.5	25.4	13.7	0.8	0.0	713
	E	7.5	60.5	24.2	7.5	0.4	0.0	0.0	281
		2.5	36.4	30.2	20.3	10.0	0.6	0.0	994

**Table 13. River age distribution (%) for North American origin Atlantic salmon (*Salmo salar*) caught at West Greenland from 1968-2019. Table does not include salmon of unknown age or origin. Because of rounding, not all rows add to 1.0. No samples were collected in 1977, 1993, or 1994.**

YEAR	1	2	3	4	5	6	7	8
1968	0.3	19.6	40.4	21.3	16.2	2.2	0	0
1969	0	27.1	45.8	19.6	6.5	0.9	0	0
1970	0	58.1	25.6	11.6	2.3	2.3	0	0
1971	1.2	32.9	36.5	16.5	9.4	3.5	0	0
1972	0.8	31.9	51.4	10.6	3.9	1.2	0.4	0
1973	2.0	40.8	34.7	18.4	2.0	2.0	0	0
1974	0.9	36	36.6	12.0	11.7	2.6	0.3	0
1975	0.4	17.3	47.6	24.4	6.2	4.0	0	0
1976	0.7	42.6	30.6	14.6	10.9	0.4	0.4	0
1977	-	-	-	-	-	-	-	-
1978	2.7	31.9	43.0	13.6	6.0	2.0	0.9	0
1979	4.2	39.9	40.6	11.3	2.8	1.1	0.1	0
1980	5.9	36.3	32.9	16.3	7.9	0.7	0.1	0
1981	3.5	31.6	37.5	19.0	6.6	1.6	0.2	0
1982	1.4	37.7	38.3	15.9	5.8	0.7	0	0.2
1983	3.1	47.0	32.6	12.7	3.7	0.8	0.1	0
1984	4.8	51.7	28.9	9.0	4.6	0.9	0.2	0
1985	5.1	41.0	35.7	12.1	4.9	1.1	0.1	0
1986	2.0	39.9	33.4	20.0	4.0	0.7	0	0
1987	3.9	41.4	31.8	16.7	5.8	0.4	0	0
1988	5.2	31.3	30.8	20.9	10.7	1.0	0.1	0
1989	7.9	39.0	30.1	15.9	5.9	1.3	0	0
1990	8.8	45.3	30.7	12.1	2.4	0.5	0.1	0
1991	5.2	33.6	43.5	12.8	3.9	0.8	0.3	0
1992	6.7	36.7	34.1	19.1	3.2	0.3	0	0
1993	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-
1995	2.4	19.0	45.4	22.6	8.8	1.8	0.1	0
1996	1.7	18.7	46.0	23.8	8.8	0.8	0.1	0
1997	1.3	16.4	48.4	17.6	15.1	1.3	0	0
1998	4.0	35.1	37.0	16.5	6.1	1.1	0.1	0
1999	2.7	23.5	50.6	20.3	2.9	0.0	0	0
2000	3.2	26.6	38.6	23.4	7.6	0.6	0	0
2001	1.9	15.2	39.4	32.0	10.8	0.7	0	0
2002	1.5	27.4	46.5	14.2	9.5	0.9	0	0
2003	2.6	28.8	38.9	21.0	7.6	1.1	0	0
2004	1.9	19.1	51.9	22.9	3.7	0.5	0	0
2005	2.7	21.4	36.3	30.5	8.5	0.5	0	0

**Table 13, continued. River age distribution (%) for North American origin Atlantic salmon (*Salmo salar*) caught at West Greenland from 1968-2018. Table does not include salmon of unknown age or origin. Because of rounding, not all rows add to 1.0. No samples were collected in 1977, 1993, or 1994.**

<b>YEAR</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>
2006	0.6	13.9	44.6	27.6	12.3	1.0	0	0
2007	1.6	27.7	34.5	26.2	9.2	1.0	0	0
2008	0.9	25.1	51.9	16.8	4.7	0.6	0	0
2009	2.6	30.7	47.3	15.4	3.7	0.4	0	0
2010	1.6	21.7	47.9	21.7	6.3	0.8	0	0
2011	1.0	35.9	45.9	14.4	2.8	0	0	0
2012	0.3	29.8	39.4	23.3	6.5	0.7	0	0
2013	0.1	32.6	37.3	20.8	8.6	0.6	0	0
2014	0.4	26.0	44.5	21.9	6.9	0.4	0	0
2015	0.1	31.6	40.6	21.6	6.0	0.2	0	0
2016	0.1	21.3	43.3	26.8	7.3	1.1	0	0
2017	0.3	31.0	41.6	19.6	7.2	0.3	0	0
2018	0.5	29.8	38.4	24.1	6.5	0.7	0	0
2019	0.6	26.9	32.5	25.4	13.7	0.8	0	0
10 year mean	0.5	28.7	41.1	22.0	7.2	0.6	0.0	0.0
Overall mean	2.3	31.1	39.6	18.9	6.9	1.0	0.1	0.0

**Table 14. River age distribution (%) for European origin Atlantic salmon (*Salmo salar*) caught at West Greenland, 1968-2019. Table does not include salmon of unknown age or origin. Because of rounding, not all rows add to 1.0. No samples were collected in 1993 and 1994.**

YEAR	1	2	3	4	5	6	7	8
1968	21.6	60.3	15.2	2.7	0.3	0	0	0
1969	0	83.8	16.2	0	0	0	0	0
1970	0	90.4	9.6	0	0	0	0	0
1971	9.3	66.5	19.9	3.1	1.2	0	0	0
1972	11.0	71.2	16.7	1.0	0.1	0	0	0
1973	26.0	58.0	14.0	2.0	0	0	0	0
1974	22.9	68.2	8.5	0.4	0	0	0	0
1975	26.0	53.4	18.2	2.5	0	0	0	0
1976	23.5	67.2	8.4	0.6	0.3	0	0	0
1977	-	-	-	-	-	-	-	-
1978	26.2	65.4	8.2	0.2	0	0	0	0
1979	23.6	64.8	11.0	0.6	0	0	0	0
1980	25.8	56.9	14.7	2.5	0.2	0	0	0
1981	15.4	67.3	15.7	1.6	0	0	0	0
1982	15.6	56.1	23.5	4.2	0.7	0	0	0
1983	34.7	50.2	12.3	2.4	0.3	0.1	0.1	0
1984	22.7	56.9	15.2	4.2	0.9	0.2	0	0
1985	20.2	61.6	14.9	2.7	0.6	0	0	0
1986	19.5	62.5	15.1	2.7	0.2	0	0	0
1987	19.2	62.5	14.8	3.3	0.3	0	0	0
1988	18.4	61.6	17.3	2.3	0.5	0	0	0
1989	18.0	61.7	17.4	2.7	0.3	0	0	0
1990	15.9	56.3	23.0	4.4	0.2	0.2	0	0
1991	20.9	47.4	26.3	4.2	1.2	0	0	0
1992	11.8	38.2	42.8	6.5	0.6	0	0	0
1993	-	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-
1995	14.8	67.3	17.2	0.6	0	0	0	0
1996	15.8	71.1	12.2	0.9	0	0	0	0
1997	4.1	58.1	37.8	0.0	0	0	0	0
1998	28.6	60.0	7.6	2.9	0.0	1.0	0	0
1999	27.7	65.1	7.2	0	0	0	0	0
2000	36.5	46.7	13.1	2.9	0.7	0	0	0
2001	16.0	51.2	27.3	4.9	0.7	0	0	0
2002	9.4	62.9	20.1	7.6	0	0	0	0
2003	16.2	58.0	22.1	3.0	0.8	0	0	0
2004	18.3	57.7	20.5	3.2	0.2	0	0	0
2005	19.2	60.5	15.0	5.4	0	0	0	0

**Table 14, continued. River age distribution (%) for European origin Atlantic salmon (*Salmo salar*) caught at West Greenland from 1968-2019. Table does not include salmon of unknown age or origin. Because of rounding, not all rows add to 1.0. No samples were collected in 1993 or 1994.**

<b>YEAR</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>
2006	17.7	54.0	23.6	3.7	0.9	0	0	0
2007	7.0	48.5	33.0	10.5	1.0	0	0	0
2008	7.0	72.8	19.3	0.8	0.0	0	0	0
2009	14.3	59.5	23.8	2.4	0.0	0	0	0
2010	11.3	57.1	27.3	3.4	0.8	0	0	0
2011	19.0	51.7	27.6	1.7	0	0	0	0
2012	9.3	63.0	24.0	3.7	0	0	0	0
2013	4.5	68.2	24.4	2.5	0	0	0	0
2014	4.5	60.7	30.8	4.0	0	0	0	0
2015	9.2	54.9	28.8	5.8	1.2	0	0	0
2016	2.5	63.3	29.6	4.3	0.3	0	0	0
2017	10.0	73.0	15.4	1.7	0	0	0	0
2018	13.7	62.1	19.0	5.2	0	0	0	0
2019	7.5	60.5	24.2	7.5	0.4	0	0	0
10 year mean	9.1	61.5	25.1	4.0	0.3	0.0	0.0	0.0
Overall mean	16.2	61.1	19.4	3.0	0.3	0.0	0.0	0.0



**Table 15. The sea age (1SW = 1 sea-winter, 2SW = 2 sea-winter, and Previous Spawners) composition of Atlantic salmon (*Salmo salar*) by continent of origin (NA = North American and E = European) and Northwest Atlantic Fisheries Organization (NAFO) division caught at West Greenland in 2019. Table does not include salmon with unknown age or origin. Because of rounding, not all rows add to 100.**

NAFO	Origin	Sea age composition (%)			Total No.
		1SW	2SW	Previous Spawners	
1B	NA	97.0	0.3	2.6	303
	E	98.0	2.0	0.0	51
		97.2	0.6	2.3	354
1C	NA	98.8	0.4	0.8	249
	E	99.4	0.0	0.6	174
		99.1	0.2	0.7	423
1F	NA	90.2	4.4	5.5	183
	E	93.4	6.6	0.0	61
		91.0	4.9	4.1	244
All areas	NA	95.9	1.4	2.7	735
	E	97.9	1.7	0.3	286
		96.5	1.5	2.1	1021



**Table 16. Sea age (1SW = 1 sea-winter, 2SW = 2 sea-winter, and PS = Previous Spawners) distribution (%) for North American and European origin Atlantic salmon (*Salmo salar*) caught at West Greenland from 1985-2019. Table does not include salmon of unknown age or origin. Not all rows add to 100 because of rounding errors. No samples were collected in 1993 or 1994.**

	North American			European		
	1SW	2SW	PS	1SW	2SW	PS
1985	92.5	7.2	0.3	95.0	4.7	0.4
1986	95.1	3.9	1.0	97.5	1.9	0.6
1987	96.3	2.3	1.4	98.0	1.7	0.3
1988	96.7	2.0	1.2	98.1	1.3	0.5
1989	92.3	5.2	2.4	95.5	3.8	0.6
1990	95.7	3.4	0.9	96.3	3.0	0.7
1991	95.6	4.1	0.4	93.4	6.5	0.2
1992	91.9	8.0	0.1	97.5	2.1	0.4
1993	-	-	-	-	-	-
1994	-	-	-	-	-	-
1995	96.8	1.5	1.7	97.3	2.2	0.5
1996	94.1	3.8	2.1	96.1	2.7	1.2
1997	98.2	0.6	1.2	99.3	0.4	0.4
1998	96.8	0.5	2.7	99.4	0.0	0.6
1999	96.8	1.2	2.0	100.0	0.0	0.0
2000	97.4	0.0	2.6	100.0	0.0	0.0
2001	98.2	2.6	0.5	97.8	2.0	0.3
2002	97.3	0.9	1.8	100.0	0.0	0.0
2003	96.7	1.0	2.3	98.9	1.1	0.0
2004	97.0	0.5	2.5	97.0	2.8	0.2
2005	92.4	1.2	6.4	96.7	1.1	2.2
2006	93.0	0.8	5.6	98.8	0.0	1.2
2007	96.5	1.0	2.5	95.6	2.5	1.5
2008	97.4	0.5	2.2	98.8	0.8	0.4
2009	93.4	2.8	3.8	89.4	7.6	3.0
2010	98.2	0.4	1.4	97.5	1.7	0.8
2011	93.8	1.5	4.7	82.8	12.1	5.2
2012	93.2	0.7	6.0	98.0	1.6	0.4
2013	94.9	1.4	3.7	96.6	2.4	1.0
2014	91.3	1.1	7.6	96.1	2.4	1.5
2015	97.0	0.7	2.3	98.2	1.2	0.6
2016	93.5	2.5	4.0	95.5	3.5	1.0
2017	92.5	1.5	6.0	93.1	5.7	1.2
2018	97.4	0.4	2.2	97.4	2.6	0.0
2019	95.9	1.4	2.7	97.9	1.7	0.3



**Figure 1. Nominal catches and commercial quotas (metric tons, round fresh weight) of Atlantic salmon (*Salmo salar*) at West Greenland for 1960–2019 (top panel) and 2010–2019 (bottom panel). Total reported landings from 2010–2019 are displayed by landings type. No quotas were set from 2010–2011, but from 2012–2014, an annual quota was set and applied to factory landings only. Starting in 2015, a single quota was set for all components of the fishery.**








Figure 2. Map of southwest Greenland showing communities at which Atlantic salmon (*Salmo salar*) have historically been landed. Northwest Atlantic Fisheries Organization (NAFO) divisions (1A-1F) are also shown. In 2019, samples were obtained from Sisimiut (1B), Maniitsoq (1C), Nuuk (1D), and Qaqortoq (1F).




**Figure 3. Map showing total samples (gray circles) and analyzed subsamples (blue circles) during 2019 at the West Greenland Atlantic salmon (*Salmo salar*) fishery for the single nucleotide polymorphism analysis. Sample locations from north to south are Sisimiut, Maniitsoq, Nuuk and Qaqortoq located in Northwest Atlantic Fisheries Organization divisions 1B, 1C, 1D, and 1F, respectively.**



**Figure 4. Map of sample locations for the single nucleotide polymorphism range-wide genetic baseline for North American (top) and European (bottom) regional groupings. See Table 6 for location abbreviations.**



**Figure 5. The weighted proportions of North American and European Atlantic salmon (*Salmo salar*) caught at West Greenland from 1982-2019. Proportions were weighted by the estimated numbers of salmon by origin for each division according to the adjusted landings.**



**Figure 6. Proportions of unsampled adjusted landings of North American origin and European origin Atlantic salmon (*Salmo salar*) (left panels) and of sampled adjusted landings of North American origin and European origin Atlantic salmon (right panels) by North Atlantic Fisheries Organization (NAFO) division (top row represents division 1A and bottom row represents division 1F) sampled at West Greenland from 2005–2019. Year-division combinations with data identify when and where sampling occurred. Division 1A 2005 value is from 1 sample.**







Figure 7. The weighted numbers of North American and European Atlantic salmon (*Salmo salar*) caught at West Greenland from 1982–2019 (top) and 2010–2019 (bottom). Numbers are rounded to the nearest hundred fish. In 2019, it is estimated that approximately 6,800 North American origin and 2,600 European origin fish were harvested.





**Figure 8. Bayesian estimates of mixture composition of samples from the West Greenland Atlantic salmon (*Salmo salar*) fishery for 2019 using the single nucleotide polymorphism (SNP) baseline overall and by Northwest Atlantic Fisheries Organization (NAFO) division. Reporting Groups are identified in Table 6 and Figure 4.**



**Figure 9. Mean whole weight (kg) of European and North American 1 sea-winter (fish that have spent 1 winter at sea) Atlantic salmon (*Salmo salar*) sampled at West Greenland from 1969-2019 (top panel) and 2010-2019 (bottom panel). These data have not been adjusted for the period of sampling, and it is known that salmon grow quickly during the period of feeding and while in the fishery at West Greenland. Caution is urged when interpreting these uncorrected data.**

## REFERENCES CITED

- Anderson EC, Waples RS, Kalinowski ST. 2008. An improved method for predicting the accuracy of genetic stock identification. *Can J Fish Aquat Sci.* 65(7):1475-1486.
- Bradbury IR, Hamilton LC, Sheehan TF, Chaput G, Robertson MJ, Dempson JB, Reddin DG, Morris V, King TL, Bernatchez L. 2016. Genetic mixed-stock analysis disentangles spatial and temporal variation in composition of the West Greenland Atlantic Salmon fishery. *ICES J Mar Sci.* 73(9):2311-2321.
- Colligan M, Sheehan T, Pruden J, Kocik J. 2008. The challenges posed by international management of Atlantic salmon: balancing commercial, recreational and societal interests - The North Atlantic Salmon Conservation Organization (NASCO). In: Schechter MG, Leonard NJ, Taylor WW, editors. *International governance of fisheries ecosystems: learning from the past, finding solutions for the future.* American Fisheries Society; p. 458.
- [ICES] International Council for the Exploration of the Sea. 2004. Report of the Working Group on North Atlantic Salmon (WGNAS). 29 March-8 April 2004. Halifax, Canada: ICES. 293 p. Report No.: CM 2004/ACFM:20.
- [ICES] International Council for the Exploration of the Sea. 2007. Study Group on Establishing a Framework of Indicators of Salmon Stock Abundance (SGEFISSA). 27-30 November 2006. Halifax, Canada: ICES. 71 p. Report No.: CM 2007/DFC:01.
- [ICES] International Council for the Exploration of the Sea. 2011. Report of the Working Group on North Atlantic Salmon (WGNAS). 22-31 March 2011. Copenhagen, Denmark: ICES. 286 p. Report No.: CM 2011/ACOM:09.
- [ICES] International Council for the Exploration of the Sea. 2015. Report of the Working Group on North Atlantic Salmon (WGNAS). 17-26 March 2015. Moncton, Canada: ICES. 332 p. Report No.: CM 2015/ACOM:09.
- [ICES] International Council for the Exploration of the Sea. 2017. Report of the Working Group on North Atlantic Salmon (WGNAS). 29 March-7 April 2017. Copenhagen, Denmark: ICES. 296 p. Report No.: CM 2017/ACOM:20.
- [ICES] International Council for the Exploration of the Sea. 2018. Report of the Working Group on North Atlantic Salmon (WGNAS). 4-13 April 2018. Woods Hole, MA (US): ICES. 383 p. Report No.: CM 2018/ACOM:21.
- [ICES] International Council for the Exploration of the Sea. 2019. Working Group on North Atlantic Salmon (WGNAS). *ICES Scientific Reports* 1:16. ICES. 368 pp. Available from: <http://doi.org/10.17895/ices.pub.4978>
- Jeffery NW, Wringe BF, McBride M, Hamilton LC, Stanley RRE, Bernatchez L, Bentzen P, Kent M, Clément M, Gilbey J, Sheehan TF, Bradbury IR. 2018. Range-wide regional assignment

- of Atlantic salmon (*Salmo salar*) using genome wide single-nucleotide polymorphisms. *Fish Res.* 206:163-175.
- Jensen JM. 1990. Atlantic salmon at Greenland. *Fish Res.* 10:29-52.
- King TL, Kalinowski ST, Schill WB, Spidle AP, Lubinski BA. 2001. Population structure of Atlantic salmon (*Salmo salar* L.): a range-wide perspective from microsatellite DNA variation. *Mol Ecol.* 10:807-821.
- Miller KM, Gardner IA, Vanderstichel R, Burnley T, Schulze AD, Li S, Tabata A, Kaukinen KH, Ming TJ, Ginther NG. 2016. Report on the performance evaluation of the Fluidigm BioMark platform for high-throughput microbe monitoring in salmon. Ottawa, Canada: DFO Canadian Science Advisory Secretariat. Research Document No.: 2016/038.
- Miller KM, Teffer AK, Tucker S, Li, S., Schulze AD, Trudel M, Juanes F, Tabata A, Kaukinen KH, Ginther NG, et al. 2014. Infectious disease, shifting climates, and opportunistic predators: Cumulative factors potentially impacting wild salmon declines. *Evol Appl.* 7:812-855.
- [NASCO] North Atlantic Salmon Conservation Organization. 2015. Report of the Thirty-Second Annual Meetings of the Commissions. 2-5 June 2015. Happy Valley-Goose Bay, Canada.
- [NASCO] North Atlantic Salmon Conservation Organization. 2018. Report of the Thirty-Fifth Annual Meeting of the West Greenland. 12-15 June 2018. Portland, Maine (US).
- [NASCO] North Atlantic Salmon Conservation Organization. 2019. Report of the Thirty-Sixth Annual Meeting of the West Greenland. 5-7 June 2019. Tromsø, Norway.
- Ó Maoiléidigh N, White J, Hansen LP, Jacobsen JA, Potter T, Russell I, Reddin D, Sheehan TS. 2018. Fifty years of marine tag recoveries from Atlantic salmon. ICES. 121 p. Cooperative Research Report No. 343.
- Reddin DG, Friedland KD. 1999. A history of identification to continent of origin of Atlantic salmon (*Salmo salar* L.) at west Greenland, 1969-1997. *Fish Res.* 43:221-235.
- Reddin DG, Hansen LP, Bakkestuen V, Russell I, White J, Potter ECE, Sheehan TF, Ó Maoiléidigh N, Dempson JB, Smith GW, Isaksson A, Fowler M, Jacobsen JA, Mork KA, Amiro P. 2012. Distribution and biological characteristics of Atlantic salmon (*Salmo salar*) at Greenland based on the analysis of historical tag recoveries. *ICES J Mar Sci.* 69(9):1589–1597.
- Sheehan TF, Legault CM, King TL, Spidle AP. 2010. Probabilistic-based genetic assignment model: assignments to subcontinent of origin of the West Greenland Atlantic salmon harvest. *ICES J Mar Sci.* 67:537–550.

Sheehan TF, Assunção MGL, Deschamps D, Laughton B, Ó Cuaig M, Nygaard R, King TL, Robertson MJ, Ó Maoiléidigh N. 2013. The International Sampling Program: Continent of origin and biological characteristics of Atlantic salmon collected at West Greenland in 2012. July 2013. 25 p. US Department of Commerce Northeast Fisheries Science Center. Reference Document No. 13-20.

Teffer A, Carr J, Tabata A, Schulze A, Bradbury I, Deschamps D, Gillis CA, Brunson E, Mordecai G, Miller K. 2020. A molecular assessment of infectious agents carried by Atlantic salmon at sea and in three eastern Canadian rivers, including aquaculture escapees and North American and European origin wild stocks. *Facets*. 5:234-263.

## **Procedures for Issuing Manuscripts in the Northeast Fisheries Science Center Reference Document (CRD) and the Technical Memorandum (TM) Series**

The mission of NOAA's National Marine Fisheries Service (NMFS) is "stewardship of the nation's ocean resources and their habitat." As the research arm of the NMFS's Greater Atlantic Region, the Northeast Fisheries Science Center (NEFSC) supports the NMFS's mission by "conducting ecosystem-based research and assessments of living marine resources, with a focus on the Northeast Shelf, to promote the recovery and long-term sustainability of these resources and to generate social and economic opportunities and benefits from their use." Results of NEFSC research are largely reported in primary scientific media (e.g., anonymously peer-reviewed scientific journals). However, to assist itself in providing data, information, and advice to its constituents, the NEFSC occasionally releases its results in its own series.

*NOAA Technical Memorandum NMFS-NE* – This series is issued irregularly. The series typically includes: data reports of long-term field or lab studies of important species or habitats; synthesis reports for important species or habitats; annual reports of overall assessment or monitoring programs; manuals describing program-wide surveying or experimental techniques; literature surveys of important species or habitat topics; proceedings and collected papers of scientific meetings; and indexed and/or annotated bibliographies. All issues receive internal scientific review, and most issues receive technical and copy editing.

*Northeast Fisheries Science Center Reference Document* – This series is issued irregularly. The series typically includes: data reports on field and lab studies; progress reports on experiments, monitoring, and assessments; background papers for, collected abstracts of, and/or summary reports of scientific meetings; and simple bibliographies. Issues receive internal scientific review, and most issues receive copy editing.

### **CLEARANCE**

All manuscripts submitted for issuance as CRDs must have cleared the NEFSC's manuscript/abstract/webpage review process. If your manuscript includes material from another work which has been copyrighted, you will need to work with the NEFSC's Editorial Office to arrange for permission to use that material by securing release signatures on the "NEFSC Use-of-Copyrighted-Work Permission Form."

For more information, NEFSC authors should see the NEFSC's online publication policy manual, "Manuscript/Abstract/Webpage Preparation, Review, & Dissemination: NEFSC Author's Guide to Policy, Process, and Procedure."

### **STYLE**

The CRD series is obligated to conform with the style contained in the current edition of the United States Government Printing Office Style Manual; however, that style manual is silent on many

aspects of scientific manuscripts. The CRD series relies more on the CSE Style Manual. Manuscripts should be prepared to conform with both of these style manuals.

The CRD series uses the Integrated Taxonomic Information System, the American Fisheries Society's guides, and the Society for Marine Mammalogy's guide for verifying scientific species names.

For in-text citations, use the name-date system. A special effort should be made to ensure all necessary bibliographic information is included in the list of references cited. Personal communications must include the date, full name, and full mailing address of the contact.

## **PREPARATION**

Once your document has cleared the review process, the Editorial Office will contact you with publication needs—for example, revised text (if necessary) and separate digital figures and tables if they are embedded in the document. Materials may be submitted to the Editorial Office as email attachments or intranet downloads. Text files should be in Microsoft Word, tables may be in Word or Excel, and graphics files may be in a variety of formats (JPG, GIF, Excel, PowerPoint, etc.).

## **PRODUCTION AND DISTRIBUTION**

The Editorial Office will perform a copy edit of the document and may request further revisions. The Editorial Office will develop the inside and outside front covers, the inside and outside back covers, and the title and bibliographic control pages of the document.

Once the CRD is ready, the Editorial Office will contact you to review it and submit corrections or changes before the document is posted online. A number of organizations and individuals in the Northeast Region will be notified by e-mail of the availability of the document online.