

Northeast Fisheries Science Center Technical Memorandum 452

Research Track Assessment Report for Atlantic Sea Scallops 2025

October 2026

Northeast Fisheries Science Center Technical Memorandum 452

Research Track Assessment Report for Atlantic Sea Scallops 2025

Report prepared by NOAA Fisheries, Northeast Fisheries Science Center

NOAA Fisheries, Northeast Fisheries Science Center

US DEPARTMENT OF COMMERCE

National Oceanic and Atmospheric Administration

National Marine Fisheries Service

Northeast Fisheries Science Center

Woods Hole, Massachusetts

October 2026

Editorial Notes

Information Quality Act Compliance: In accordance with section 515 of Public Law 106-554, the Northeast Fisheries Science Center (NEFSC) completed both technical and policy reviews for this report. These pre-dissemination reviews are on file at the NEFSC Editorial Office.

Species Names: The NEFSC Editorial Office's policy on the use of species names in all technical communications is generally to follow the American Fisheries Society's lists of scientific and common names for fishes, mollusks, and decapod crustaceans and to follow the Society for Marine Mammalogy's guidance on scientific and common names for marine mammals. Exceptions to this policy occur when there are subsequent compelling revisions in the classifications of species, resulting in changes in the names of species.

Statistical Terms: The NEFSC Editorial Office's policy on the use of statistical terms in all technical communications is generally to follow the International Standards Organization's handbook of statistical methods.

This document may be cited as:

> Northeast Fisheries Science Center. 2026. Atlantic Sea Scallop Research Track Assessment Report 2025. Northeast Fish Sci Tech Memo 452. 103p.

TABLE OF CONTENTS

SUMMARY REPORT OF THE ATLANTIC SCALLOP TRACK STOCK ASSESSMENT PEER REVIEW 1
Introduction 1
TERM OF REFERENCE 1 3
TERM OF REFERENCE 2 6
TERM OF REFERENCE 3 8
TERM OF REFERENCE 4 13
TERM OF REFERENCE 5 19
TERM OF REFERENCE 6 21
TERM OF REFERENCE 7 23
TERM OF REFERENCE 8 26
Literature Cited 27
Appendix 1 – Terms of Reference for Atlantic Scallop Research Track Stock Assessment 28
Appendix 2 – Initial agenda for Atlantic Scallop Research Track Assessment Peer Review meeting, April 21 – 24, 2025. 30
Appendix 3 – Materials provided or referenced during the Atlantic Sea Scallop Research Track Stock Assessment Peer Review meeting 34
Appendix 4 – Meeting attendees at the Atlantic Sea Scallop Research Track Stock Assessment Peer Review meeting 36
2025 ATLANTIC SEA SCALLOP RESEARCH TRACK ASSESSMENT 38
ATLANTIC SEA SCALLOP RESEARCH TRACK WORKING GROUP (WG) 38
ACKNOWLEDGEMENTS 38
TERMS OF REFERENCE 39
EXECUTIVE SUMMARY 41
INTRODUCTION 47
Distribution 47
Life History 47
Growth 48
Shell height to weight relationships 48
Natural mortality 49
Ecosystem and Climate Influences 50
Landings, fishing effort, discards 53
SURVEY DATA 61

Fishing mortality, recruitment, and stock biomass 68
Reference points 75
PROJECTIONS 81
GeoSAMS 83
RESEARCH RECOMMENDATIONS 83
BACKUP ASSESSMENT APPROACH 90
ADDITIONAL INVESTIGATIONS 93
REFERENCES 96
List of Working Papers 103

SUMMARY REPORT OF THE ATLANTIC SCALLOP RESEARCH TRACK STOCK ASSESSMENT PEER REVIEW

Panel summary report prepared by Panel Members (April 21–24, 2025):
Thomas Miller (Chair), MAFMC SSC
Yong Chen, independent contractor with CIE
Noel Cadigan, independent contractor with CIE
Martin Cryer, independent contractor with CIE

INTRODUCTION

The Northeast Region Coordinating Council (NRCC)[1] has developed an enhanced stock assessment process to improve the quality of assessments. The process involves two tracks of assessment work: 1) a management track that includes routine updates of previously approved assessment methods to support regular management actions (e.g., annual catch limits), and 2) a research track that allows comprehensive research and development of improved assessments on a stock-by-stock or topical basis. The research track assessment process allows for a more thorough review of information available and for the evaluation of different assessment approaches than would be possible in a standard stock assessment process where the results are immediately used for management advice. This Panel reviewed the Research Track Assessment for the Atlantic Sea Scallop stock

The most recent stock assessment for Atlantic Sea Scallop was a management track assessment in 2021 that was based on the benchmark assessment from SARC 65 in 2017 (NEFSC 2018). Assessments for Atlantic Sea Scallop have used a catch at size analysis (CASA) framework since 2007. Materials for this assessment were prepared by a Workgroup (WG), chaired by Dr. Patrick Sullivan. The WG included staff from NOAA Fisheries, the regional management councils, academic institutions and, because the species distribution of Atlantic scallop extends into Canada, a scientist from Fisheries and Oceans Canada (DFO).

The Atlantic Sea Scallop Research Track Assessment Peer Review took place in Woods Hole, MA during April 21–24, 2025. All members of the Review Panel and all but one of the presenters were present in person at NEFSC for the peer review. The Review Panel greatly appreciates the availability of these scientists, as the in-person format greatly facilitated discussion and interactions. A hybrid option was available for agency and academic scientists, fishers and members of the public to participate, which helped ensure the process was open and transparent.

[1] Atlantic States Marine Fisheries Commission (ASFMC), Greater Atlantic Regional Fisheries Office (GARFO), Mid-Atlantic Fishery Management Council (MAFMC), New England Fishery Management Council (NEFMC), and Northeast Fisheries Science Center (NEFSC).

Periods for public comment were set aside on each day of the meeting. This report was prepared by the members of the Review Panel, which included four independent, international scientists selected by the Center for Independent Experts (CIE). Yong Chen and Thomas Miller (chair) are both members of the Mid-Atlantic Fishery Management Council's Scientific and Statistical Committee (SSC). Noel Cadigan (Canada) and Martin Cryer (New Zealand) provided additional independent perspectives of the stock assessment. All four Panel members have extensive experience with size-structured stock assessment modeling approaches to invertebrate fisheries.

The Research Track Assessment Report and 14 supporting working papers were provided to the Panel on the NEFSC data portal (https://apps-nefsc.fisheries.noaa.gov/saw/sasi.php) on April 7, 2025. Appendix 3 lists the material provided to the Review Panel. Prior to the meeting, members of the Panel met with Brian Hooper (NEFSC's Stock Assessment Process Lead) and Kristan Blackhart (Chief, NEFSC Population Dynamics Branch) on April 16, 2025 to review and discuss the meeting agenda, reporting requirements, meeting logistics and the overall process. The Panel Chair is grateful for discussions with Cate O'Keefe, executive Director of the New England Fishery Management Council. The Review Panel is grateful to Jason Boucher, Toni Chute, Kiersten Curti, Alex Hansell, Liz Brooks, Chris Legault and Emily Liljestrand who served as rapporteurs. Appendix 4 lists the attendees for the meeting.

Panel members and the Chair drafted this Summary Report in a Google Doc. A preliminary summary of the Panel's conclusions and recommendations was provided at the end of the meeting to the Atlantic Scallop Research Track Workgroup (WG) and members of the NEFSC Population Dynamics Branch. Subsequent to the meeting, the Panel Chair compiled and edited this Summary Report with assistance from the CIE Panelists before submission of a draft report to the WG. The scope of the WG review of the draft was limited to suggesting corrections for errors of fact or requesting that Panel recommendations be clarified. Additionally, each of the CIE Panelists will submit separate reviewer reports to the CIE.

Terms of Reference (ToRs) for the WG, and by extension the Review Panel, are provided in Appendix 1. We note that the specific ToRs deviate from the standard ToRs provided to a research track WG in that the assessment was charged with provision of biological reference points and a stock status specification statement.

The Review Panel noted that in several ways, the Atlantic Scallop Research Track Review was non-standard. For example, it was unusual that it was required to provide stock status advice. This may have led the WG to adopt a more conservative approach to changes to the data analysis, model exploration and projection methodologies than might have occurred otherwise. Perhaps as a result the Atlantic Scallop Research Track Assessment is risk averse—in some ways more like a management track assessment than is typical. As a result, the Review Panel viewed the Atlantic Scallop Research Track Assessment to be an opportunity that was missed to more fully explore novel ways to analyze the data, and to more fully explore model configurations or even alternative

model structures. In the Review Panel's experience, it is typical for Research Track Assessments to proceed from the previously approved assessment in a series of steps that may first update data, then assess changes in model fit based on alternative assumptions or model structures before arriving at the Research Track WG's final recommended model. Somewhat unusually, the WG for this Atlantic Scallop Research Track Assessment presented the recommended model first, and provided only limited evaluation of alternatives during the review. Based on discussions during the review meeting, the Review Panel believes that it is highly likely that modeling decisions were made carefully and in a structured order during data analyses and model development; however, the Review Panel was frequently unable to point to specific evidence that supported this belief. Finally, the information provided to the Review Panel was not comprehensive, which might limit its ability to better complete its task. Descriptions of evidence to support decisions made by the WG were often lacking from the written materials, and key citations were missing. The Review Panel notes that the WG may have tried to balance criticism that they provided too much information for SARC 65. Yet, the Review Panel notes that they are asked to provide evidence-based reviews of the work that was completed. This task is made more difficult when the analyses are not presented to the Review Panel in a logical, structured and documented sequence or when documentation is incomplete and the pathway to the final assessment is opaque.

The Panel's detailed evaluation of the WG's response to the nine ToRs is provided in subsequent sections below. The Panel agreed that ToRs 1, 2, 7, 8 and 9 were fully met and ToRs 3–6 were partially met.

TERM OF REFERENCE 1

Identify relevant ecosystem and climate influences on the stock. Characterize the uncertainty in the relevant sources of data and their link to stock dynamics. Consider findings, as appropriate, in addressing other TORs. Report how the findings were considered under impacted TORs.

The Review Panel considers that work presented by the WG fully met this ToR. The Review Panel believes that the WG have clearly identified and documented ecosystem impacts on the abundance, distribution and composition of the Atlantic Sea Scallop population in the Mid Atlantic, Georges Bank and Gulf of Maine.

Distributional analyses from dredge surveys conducted between 1999–2023 demonstrate that Atlantic scallop have been excluded from shallower, inshore waters and from waters in the southern regions of the Mid-Atlantic by warming temperatures. Size distributions show survival of scallops 2 years and older have declined substantially, such that scallops larger than 70 mm shell height are largely absent from waters off of Virginia (Fig. 1). This recent truncation of the size distribution occurred during a time when there was essentially no fishing, implying a substantial increase in natural mortality rates (M) in this region. Water temperatures > 18°C represent a significant stress on Atlantic scallop, particularly for individuals older than 2 years. Data on oceanic temperatures in the Southern New England / Mid-Atlantic region developed using the GLORYS Ocean Reanalysis system (https://www.mercator-ocean.eu/en/ocean-science/glorys/) indicated increases in the proportion of suitable habitats that experienced temperatures > 18°C, particularly in summer and autumn months (Fig. 2). The spatial extent and duration of heat stress events increased from 1993 to 2023, particularly in September and October. This warming can be expected to be a consistent and, perhaps, an increasing feature of the areas of potential scallop habitat in the short-term future.




Figure 1. Dredge survey shell heights off of Virginia showing reduced numbers of larger scallops over time. The black solid line is for 1998–2000, the blue dashed line is from 2001–2008 and the red dashed line for 2009–2018. Credit: Figure 1.1 in SCA_RT_WP_TOR1a Ecosystem Influences.pdf




Figure 2. Mean seasonal heat stress area as proportion of SAMS areas. The blue lines show a linear regression with grey shading showing the 95% confidence interval. (Winter = Jan-Mar, Spring = Apr-Jun, Summer = Aug-Sept, Fall = Oct-Dec. Credit: Figure 1.12 2025_SCA_RT_WP_TOR1b Climate Influences.pdf

It is uncertain if the high fall bottom temperatures thought to be harmful to scallop survival will persist in the medium- to long-term. Reliable forecasts of the relevant bottom temperatures are not available. However, the Review Panel agreed that it was appropriate that the assessment team included consideration of ecosystem and climate influences when setting reference points and forecast procedures. Time-varying M has been included in CASA scallop assessment models to account for changes in survival rates due to changes in the ecosystem.

Predation of Atlantic Sea Scallops by the sea star *Astropecten americanus* appears to be reducing or excluding sea scallops from the deeper waters of the Mid-Atlantic. Elevated levels of the sea star *Asterias vulgaris* and concomitantly increased numbers of sea scallop "clappers", an indicator of recent natural mortality, were observed on surveys of the northern edge of Georges Bank in 2024, in both U.S. and Canadian waters. Off-colored "gray meats", potentially caused by protists in the genus *Apicomplexa*, have been most commonly observed on Georges Bank, and in

particular in Closed Area I. This may have contributed to the increased natural mortality observed in the Georges Bank closed areas during 2011–13. Other parasitic infections and impacts on scallop mortality rates were described by the assessment team.

The WG presented information on growth of scallops under this ToR, but the Review Panel felt it was more appropriate to discuss this work under ToR 3.

The Review Panel finds that the WG completed work required to understand ecosystem effects on Atlantic Sea Scallop. The Review Panel encourages the WG to continue this work in preparation for future assessments. In particular, the Review Panel recognizes opportunities to "close the loop" by considering whether information on time-varying natural mortality derived from the assessment model can be analyzed to further elucidate the importance and extent of ecosystem changes on Atlantic Sea Scallop population dynamics. Such analyses could examine disease and parasitism incidence or direct environmental factors such as temperature.

The Review Panel fully supports the regional approach taken by the WG in developing the assessment models. There is compelling evidence that the dynamics of Atlantic Sea Scallops in these two different regions are experiencing different patterns of natural and fishing mortality and other life history processes (e.g., growth). The Review Panel encourages the WG to initiate research to explore whether biological factors exist that underpin the regional differences observed in the assessment model results. This work could involve modern molecular approaches to stock definitions that have shown the ability to detect substructure in populations at this scale (Clucas et al., 2019; Reid et al., 2023). Other lines of evidence from coupled physical-biological models may also be helpful indicators of restrictions on connectivity between regions (Tian et al., 2009). Such work is important to assess whether multiple stocks exist within this region.

TERM OF REFERENCE 2

Estimate catch from all sources including landings and discards. Describe the spatial and temporal distribution of landings, discards, and fishing effort. Characterize the uncertainty in these sources of data.

The Review Panel considers that the work presented by the WG fully met this ToR. The Review Panel believes that the landings, discards and incidental mortality have been appropriately characterized, and where possible, sources of uncertainty have been identified and quantified.

The WG fully documented data sources related to catches from different sectors of the fishery over time. Data from 1975 onwards are used in the assessment, but the WG assembled data on long term landings for Atlantic Sea Scallop since 1887. The WG group documented well the history of management actions that have changed fishing patterns and also reviewed how catches have been recorded and reported historically. Atlantic Sea Scallop catches have been reported as a part of the NEFSC Catch Accounting and Monitoring System (CAMS) since 2020.

Fishery catches are recorded from the Mid-Atlantic, Southern New England, Georges Bank and the Gulf of Maine. Southern New England and Gulf of Maine catches remain a minor component of the fishery. However, there have been increased landings from the Gulf of Maine in recent years. Landings are reported by five market categories, defined by the number of "meats" (the adductor muscle) per pound of scallop. During 1975–2000 landings were mostly of the 20–30 or greater market categories. However, since 2000, approximately 75% of landings are U10 and 10–20 market categories.

The Review Panel discussed with the WG the potential importance of unreported landings. There is a small allowance for scallop landings in some groundfish fisheries, but the allowance is small (20 Lbs./trip) and WG believed that overall, this source of landings is likely small currently. However, information on incidental and unreported catches early in the history of the fishery are not available.

Catches are considered a census and thus estimates of uncertainty are not relevant.

There are no recognized recreational fisheries for scallops, which is a source of removals often associated with large degrees of uncertainty.

Discard levels of Atlantic Sea Scallop, and its associated uncertainty, have been estimated from the Northeast Fishery Observer Program since 1989. The approaches to estimating discard mortality have changed as the catch recording methodology has changed (e.g., SBRM to CAMS). Discard mortality averaged 5.2% of landings for 1989–2023 but has increased recently (5.9% for 2010–2023). Strong regional differences exist in the percentage of scallops discarded. Discarding and discard mortality is not included in the CASA, or SYM models. If discards remain at low levels, this may be acceptable. However, the evidence that catches are increasingly dominated by large scallops (U10 and 10–20 count) suggest that increased targeting is occurring, which may be related to an increase in discards. Similarly, the importance of high levels of uncertainty in some years and regions may warrant increased attention to discarding and discard mortality in the future.

The WG considered incidental mortality in the Atlantic Sea Scallop fishery. Estimates of incidental mortality in published research have ranged from 15–20% on hard bottom in Canadian waters (Caddy, 1973) to more recent estimates of 2.5% for the Mid-Atlantic and 8% for Georges Bank (Ferraro et al., 2017). Incidental mortality estimates are derived from video surveys conducted after an area had been dredged. Ferraro et al. (2017) noted that the regional differences in incidental mortality likely reflect a difference in the character of the substrate with higher mortality on Georges Bank reflective of the presence of harder sediments generally. However, these estimates incorporate only the direct, physical damage to scallops of dredging and based on experience from New Zealand, may under-estimate longer term mortality rates. The CASA assessment models use incidental mortality rates of 6 and 11% for the Mid-Atlantic and Georges Bank regions respectively. Similar values are used for the SYM and SAMS models.

TERM OF REFERENCE 3

Present the survey data used in the assessment (e.g., indices of relative or absolute abundance, recruitment, state surveys, age-length data, application of catchability and calibration studies, etc.) and provide a rationale for which data are used. Describe the spatial and temporal distribution of the data. Characterize the uncertainty in these sources of data.

The Review Panel concluded that the WG partially met this ToR. The Review Panel believes that the work presented during the review meeting represents our best understanding of trends in population abundance, distribution and composition. The Review Panel indicated a lack of documentation provided limited its ability to fully understand the challenges in interpreting survey time series.

The dredge survey, which is the longest fishery-independent survey used in the assessment, was well described. Data are available from 1979 to present for the NEFSC dredge survey. Sampling was conducted from the R/V Albatross IV (1979–1989, 1993–2007) and the R/V Hugh R. Sharp (2008–2022). The Virginia Institute of Marine Science has been conducting a dredge survey in the Mid-Atlantic since 2005 using a commercial vessel. This has become the principal survey in the Mid-Atlantic since 2014, and it was expanded in 2023 to survey Georges Bank. Vessel calibration studies involving the three principal vessels involved have indicated that tow length is a more important factor than vessel identity. Accordingly, dredge survey catches are standardized for tow length and depth, but not vessel.

Dredge catches are reported in numbers per tow and biomass per tow with associated CVs. Plots of the dredge survey time series are shown in Fig. 3.




Figure 3. Dredge time series for numbers (left) and biomass (right) (dots), including lowess smoothers (lines) for a) Georges Bank and b) Mid-Atlantic. Credit. Figure 3.2. 2025_SCA_RT_TOR3 Survey Data.pdf

Atlantic Sea Scallops are also surveyed using two optical methods. A drop camera survey has been conducted since 2003 by researchers at the University of Massachusetts Dartmouth. A weighted frame that provides a structure for lights and cameras is lowered to the bottom, and images are captured. The frame is then raised off the bottom while the vessel drifts a short distance before the frame is lowered back to sediment. Four such samples are collected at each location. Individual locations together form a 1.5 nautical mile (nm.) or 3 nm. sampling grid. Images are subsequently annotated to estimate the number and size of scallops observed at each sampling location. Digital still cameras were added in 2009 and the main camera was changed to digital format in 2017. These changes altered the field of view and the size-selectivity of the survey. Annotations have been calibrated in tank experiments. Drop camera surveys have been conducted in the Mid-Atlantic and Georges Bank. Annual surveys have a common footprint in each region, and data falling outside of this footprint were not used in the analysis. GAMs were used to create abundance and density maps throughout each common footprint area each year. Estimates of uncertainty were also calculated.

The second optical survey is a towed, habitat mapping camera system, termed Habcam. Habcam collects approximately 6 images/second while it is towed at 5–7 knots approximately 2 m above the bottom. Four different versions of the towed body have been used during the survey time period. There is a structural autocorrelation in sequential Habcam images, and thus only 1 image in 50 or 1 in 100 are annotated. This is equivalent to one image every 25–50 m. Data from each image are interpolated to maps of larger areas using zero-inflated general additive models to remove spatial trends, followed by ordinary kriging.

Estimates from Habcam have been used to estimate the survey capture efficiency of the dredge survey. The best estimate of the dredge survey q was 0.4 on sand and 0.27 in areas of gravel or cobble. The assessment uses these, and other calibration results, to expand the dredge estimates of relative abundance to estimates of absolute abundance. Beginning in 2015, a dredge efficiency estimate of 0.13 was applied to high-density scallop beds to account for reduced dredge efficiency. The reduced capture efficiency is observed when scallop density is ≥ 2 scallops m^{-2}.

The Review Panel noted a general lack of documentation in material provided to address this ToR. The Review Panel understands that three principal surveys have been reviewed previously by an independent CIE review (Center for Independent Experts, 2015). The Review Panel further recognizes that the previous assessment received criticism during the SARC 62 review process over the amount of material provided, and accordingly material provided for this review was abbreviated. However, the Review Panel noted that the lack of documentation, particularly for the complex, non-standard camera-based surveys, limited its ability to fully review and understand the survey time series presented. For example, no diagnostics were provided from the geostatistical models used to expand both optical surveys and it was unclear whether improvements to the analytical approaches recommended by the 2015 CIE (op. cit.) review had been adopted wholly or in part. There were large differences in survey estimates during 2015–2018 (Fig. 4) and the reasons for these differences should have been described and documented.




Figure 4. Comparison of survey estimates, 2003–2023, for Georges Bank (left), and Mid-Atlantic (right). Numbers are on the top row and biomass on the bottom row. Credit: 2025_SCA_RT_TOR3_Survey Data.pdf

The Review Panel noted that CVs were not shown in any of the fishery-independent survey plots and requested that they be presented. The Review Panel believes that the uncertainty associated with survey estimates should be routinely shown on plots, unless there are strong compelling reasons not to do so. The lack of uncertainty estimates in survey plots (Figures 3 and 4) makes interpreting the reliability of trends more difficult.

The Review Panel was concerned that the reported CVs of the Habcam survey may be too small. This concern was also raised in the 2015 CIE review of the Atlantic Sea Scallop survey programs (Center for Independent Experts, 2015). It was not clear to the Review Panel whether or how the concerns of this earlier CIE review for Habcam survey methodologies, and indeed for the other surveys, have been evaluated and, where appropriate, incorporated.

The Review Panel suggests that size distributions associated with each survey should have been included in the assessment report. These data, when plotted as a ridge plot, allow reviewers to assess the ability of the survey to follow the progression of size classes through the sub populations over years. Information on how effective sample sizes were calculated was also missing.

As will be discussed under ToR 4, the model struggles to estimate survey catchabilities. Accordingly, a fuller discussion of the foundation of survey catchability estimates including data collection and analytical approaches used would have been helpful.

Growth and Mortality

Growth of Atlantic Sea Scallops can be inferred from the presence of annuli laid down in the shell. However, transition of annuli counts to age can be problematic because of uncertainty in the timing of the formation of the first annulus. Each annulus is assumed to be a check mark produced by spawning. Age at first spawning is inconsistent among scallops, with most spawning for the first time at age-2, but a not inconsequential number are believed to spawn for the first time at age-3. For this reason, the assessment used annuli ring widths as an indication of recent growth.

Hart and Chute (2009b) document a mixed effects modeling approach that allows for estimation of the von Bertalanffy parameters, k and L_∞. These methods were briefly summarized in the Assessment Report. Regional models for the Mid-Atlantic and Georges Bank were fit separately with and without covariates. The subsequent stock assessment used a number of growth transition matrix "blocks." The number of blocks differed between the Mid-Atlantic and Georges Bank.

As with other sections for this ToR, the Review Panel found that the documentation of the methods employed was incomplete. The Review Panel finds that details of the sample sizes, model diagnostics and fit that describe how the increment width data were transformed to stochastic growth matrices were not provided. There was no discussion of the adequacy of the sampling to support the estimation of the growth transition matrices.

The number of size transition matrix blocks for the Mid-Atlantic seemed high to the Review Panel and lacked justification regarding how growth periods were separated.

Assuming that the first growth increment was formed at age-2, the WG developed regional empirical estimates of natural mortality using life history invariants based on the resultant estimate of t_{max}. The best estimate of natural mortality in the Mid-Atlantic was M=0.4 based on an oldest aged shell of 14 years. This was considerably higher than the estimate of M=0.25 used in recent past assessments. Similarly, the best estimate of natural mortality for Georges Bank was M=0.27, based on an oldest aged shell of 19 years. This is higher than the estimate of M = 0.2 in previous recent assessments. The Review Panel supported this approach and these estimates.

TERM OF REFERENCE 4

Use the appropriate assessment approach to estimate annual fishing mortality, recruitment and stock biomass (both total and spawning stock) for the time series, and estimate their uncertainty. Compare the time series of these estimates with those from the previously accepted assessment(s). Evaluate a suite of model fit diagnostics (e.g., residual patterns, sensitivity analyses, retrospective patterns), and (a) comment on likely causes of problematic issues, and (b), if possible and appropriate, account for those issues when providing scientific advice and evaluate the consequences of any correction(s) applied.

The Review Panel concluded that the WG partially met this ToR. The Review Panel believes that the regional models presented during the review meeting provide a reliable foundation for generating management advice. However, as with ToR 3, the Review Panel noted a lack of documentation provided limits the Panel's full understanding of the performance of the assessment models.

The 2025 Research Track Assessment for Atlantic Sea Scallop continued to use the Catch at Size Analysis (CASA, Sullivan et al., 1990; NEFSC, 2018) that has been used in previous assessments. At its core, this model projects a size-structured population forward in time based on a growth transition matrix providing the probability of moving from one size class. The growth transition matrices were determined outside the CASA model. Inputs to the CASA model include the initial size, landings by size category, survey indices and size compositions (including diagnostics about effective sample sizes).

The 2025 Research Track Assessment developed three regional CASA models to assess the population dynamics of Atlantic Sea Scallop: the Mid-Atlantic; Georges Bank Open; and Georges Bank Closed. The regions follow earlier regional designations and reflect substantial differences in fishery, life history and natural mortality dynamics in each region (NEFSC, 2018). Each model considered scallop populations initiated at 5 mm shell height and used shell height to meat weight conversions from 2001–2023. Each model integrated data from 1975–2023. Commercial catch data was available throughout the modeled period. Commercial size composition data came either from port samples or from observer data. Fishery selectivity patterns differed by region and over time. Most were logistic in form, but dome-shaped selectivity functions were used for the Mid-Atlantic region for 1975–1979 and for Georges Bank Closed region for 2019 and 2022–2023. The Review Panel thought that more detailed information should have been provided to justify the selectivity forms chosen and for fixing those forms in the model. Information from up to six different fishery-independent surveys were incorporated depending on region and time period: NEFSC unlined dredge; NEFSC lined dredge; NEFSC winter bottom trawl; University of Massachusetts Dartmouth, School for Marine Science and Technology (SMAST) large drop camera survey; SMAST digital drop camera survey; and the Habcam survey. All survey data, except the unlined dredge and winter trawl surveys, were analyzed and corrected

for catchability external to the model such that they were essentially survey estimates of absolute biomass. Effective sample sizes for each survey were tuned so that the medians of assumed values were similar to expected values based on goodness of fit. Logistic survey selectivity curves were used in model fitting. Only size classes > 40 mm shell height were used to fit the CASA models.

As noted under ToR 3, multiple size transition matrices were used in implementing CASA for this assessment. In particular, the Mid-Atlantic regional model used four different size transition matrices, with each block being applied for short time periods. For example, the size transition model varied from Model 2 (1978) to Model 3 (1979) to Model 4 (1980–1981) to Model 3 (1982–83) and back to Model 2 (1984–1989). The statistical methodology that produced these abrupt changes is documented by the WG, but questions over how biologically plausible such changes in growth remain unaddressed.

Natural mortality was modeled by a very flexible function that admitted a logistic transition from a juvenile to an adult natural mortality rate with individual, independent yearly deviations. The average (for all years) juvenile and adult mortality rates were fixed for the Mid-Atlantic and Georges Bank Open regions; the mean mortality rate was estimated for the Georges Bank Closed region. The logistic transition point (i.e., L_{50}, the length at which M was 50% of the juvenile M plus 50% of the adult M) was fixed at 75 mm shell height.

Beta distributed priors were used to estimate catchabilities for the lined dredge, SMAST large camera, SMAST digital camera and Habcam surveys in each regional model. The survey abundance data in the model were multiplied by 0.5 and the assumed prior mean catchability was 0.5 for each survey; hence, effectively, the priors for surveys had a mean of 1.0. The CVs for the catchability priors were survey specific. The mean value was a consequence of the choice of beta distribution to fit catchabilities. Estimated catchabilities > 0.5 indicate the survey indices are greater than the model-estimated abundance.

The WG presented model fits for “base case” models in each region. The Review Panel interpreted these results as the WGs preferred model. Subsequent to the diagnostic consideration of model fits to the base model configuration, the WG presented a limited number of additional model runs that explored different assumptions or starting conditions.

A number of model outputs were included in the 2025 Research Track Assessment report. In most cases the regional models fit commercial catch data well—conditioned on the assumption that catch is well described (CV = 0.05). Survey size distributions were also generally well described by the model. However, there was a consistent pattern that the estimated survey catchabilities were often > 0.5, particularly for both SMAST surveys and the Habcam survey. This pattern suggests that unscaled values of these surveys are higher than expected from the abundances predicted by the model. In other words, either the model tends to underestimate biomass relative to the absolute abundance surveys or the surveys all overestimate biomass (to differing degrees).

The Review Panel felt that model results for the three regions, combined with the results presented on bottom water temperatures under ToR 1, fully support the WG's decision to use a higher rate of natural mortality in the Mid-Atlantic than had been used previously, and that has been used in Georges Bank.

Overall, the Review Panel was felt that documentation of the modeling choices made by the WG was incomplete such that it struggled to understand how the WG moved from the SARC 65 assessment (NEFSC 2018) to the current assessment and how the "base model" was identified. Moreover, the Review Panel felt that only limited model diagnostics were provided. The Panel concluded that additional information on modelling choices and assumptions about growth and fishery selectivity should have been provided.

The Review Panel expressed concern over the shifts in estimated survey catchability values from their prior means. The Review Panel could not determine whether the shift in these values represent features of the survey data streams, or constraints in other parts of the model structure. Additional likelihood profiling might have been useful to identify the data sets or penalties which were most influential in causing the model to estimate biomass lower than the surveys suggested. In addition, the only real source of process error is the annual deviations permitted in M (Fig. 5).

The extent to which the clear patterns in the temporal distributions in M in the three regions evident in Fig. 5 may reflect underlying variation in M or may reflect the model altering its understanding of M to fit other input time series. If the pattern is not caused by variation in the true value of M, the Review Panel asks whether there are attributes of camera-based surveys that lead to biases in estimated abundances, or is high precision in other data inputs constraining the way the model can fit the data so as to increase estimates of survey catchabilities? Does the lined dredge provide a reliable index of absolute abundance based on the efficiency values assumed in the inputs for the CASA model?

The Review Panel felt that the presentation of model fits to size composition made it difficult to assess whether model misspecification had occurred. The Review Panel requests that future assessments present size-composition fits as ridge plots, which facilitates comparing fits across years. There was concern expressed by the Review Panel that some abundance pulses were not well tracked as they moved through the size compositions of the population.




Figure 5. Size-specific annual estimates of natural mortality (M) for Atlantic Sea Scallop in three separate modeled regions: A) Mid-Atlantic, B) Georges Bank Closed, and C) Georges Bank Open. Credit: 2025 Atlantic Sea Scallop Research Track Assessment WG.




Figure 6. Trends in CASA estimated (solid black line) and observed survey-based estimates of annual abundance for the Georges Bank Open Base Case. Credit Fig 4.73 from 2025_SCA_RT_WP_TOR4a Fishing Mortality, Recruitment and Stock Biomass.pdf

The Review Panel considered closely the patterns in the lined dredge survey in the Georges Bank Open model fits (Fig. 6). The model clearly does not "believe" the short-term peaks in abundance evident in the lined dredge survey time series. This is a situation where knowledge of the survey CVs would have been helpful, but the Review Panel also expressed interest in understanding whether patchiness in survey catches within strata are driving the oscillating patterns in the survey time series. Analyses conducted by members of the WG during the Review Panel meeting indicated that extreme patchiness is indeed a feature of dredge survey catches. The Review Panel recommends the WG consider using geometric means as a foundation for calculating survey indices rather than arithmetic means as currently used.

The Review Panel felt that lack of fit to the model should continue to be discussed regionally. For example, the issue raised above related to survey catchability is particularly evident in the Georges Bank Open model runs. The Review Panel asks whether there is something specific about the spatial distribution of scallop in this region that is responsible for the model fit observed? However, the Review Panel also appreciated the WGs attempt to compare model-derived estimates across regions to identify potential broader patterns of change (Fig. 7).




Figure 7. Trends in biomass, fishing mortality rate, size-specific natural mortality rates and recruitment for the three regions modeled in the 2025 Atlantic Sea Scallop Research Track Assessment. Credit 2025 Atlantic Sea Scallop Research Track Assessment WG.

TERM OF REFERENCE 5

Update or redefine Status Determination Criteria (SDC; point estimates or proxies for BMSY, BTHRESHOLD, FMSY and MSY reference points) and provide estimates of those criteria and their uncertainty, along with a description of the sources of uncertainty. If analytic model-based estimates are unavailable, consider recommending alternative measurable proxies for reference points. Compare estimates of current stock size and fishing mortality to existing, and any redefined, SDCs. Provide stock status based on updated reference points.

The Review Panel felt that the information provided by the WG partially met this ToR. The Review Panel believes that the approach to establishing reference points is reasonable, and appropriately includes uncertainty in input parameters, thereby producing reference point distributions. The Review Panel believes that the reference points produced are appropriate for providing management advice. However, as with ToR 3, the lack of documentation provided limited the Panel's full understanding of the details of how reference points were determined.

Biological Reference Points were estimated in the SYM program which uses stochastic equilibrium analyses, detailed in Hart (2013), and are not derived directly from CASA. The peer review paper on which the method is based was provided at the request of the Review Panel. Separate SYM models were presented for the Mid-Atlantic and Georges Bank regions. The Review Panel was provided initially with distributions of M and stock-recruitment relationships that are used as input into the analysis for each region. The Review Panel requested and subsequently received information on estimates of recruits per spawner, an input to the projection analysis.

The WG noted further that initial simulations for the Mid-Atlantic, which used the full stock recruitment relationship, resulted in non-credible reference points. Investigations of the reasons for this outcome identified a recent (2016–2021) pattern of consistently negative residuals in the stock recruitment pattern, that was not evident in similar data for Georges Bank (Fig. 8, provided by the WG to the Review Panel during the meeting). Recruitment in the Mid-Atlantic during 2016–2021 had been 59% lower than estimated in the stock-recruitment model. Accordingly, recruitment in the Mid-Atlantic SYM model was reduced by a factor of 0.41. Additionally, natural mortality rates have been higher for adult Mid-Atlantic scallops in recent years (Fig. 5), and so an elevated M was also used.




Figure 8. Trends in recruitment estimated in the SAMS model developed for the Mid-Atlantic and Georges Bank for the 2025 Atlantic Sea Scallop Research Track Assessment. The Review Panel added the solid line at zero for clarity. Credit 2025 Atlantic Sea Scallop Research Track Assessment WG.

The calculated Mid-Atlantic yield curve was very flat, suggesting F_{MSY} is not well defined for this region. Indeed, this is reflected in the high value for F_{MSY} for the Mid-Atlantic. In contrast the yield curve for Georges Bank was well defined. The estimated reference points for the two regions were combined appropriately to produce an estimate of the biological reference point for the two regions combined. The resulting reference points for the two regions are presented below.

Table 1. Select biological reference points for Atlantic Sea Scallop derived from the SAMS model developed for the 2025 Atlantic Sea Scallop Research Track Assessment

Region	MSY (mt of meats)	F_{MSY}	B_{MSY} (mt of meats)	$B_{threshold}$ (mt of meats)	B_{2023} (mt of meats)	F_{2023}
Mid-Atlantic	7,941	1.56	15,909		20,556	0.06
Georges Bank	22,706	0.36	83,414		49,400	0.47
Combined	28,402	0.49	93,282	46,641	69,956	0.33

The Review Panel is concerned about the reliability of the combined region reference point. There is strong evidence in the material presented to the Panel that there needed to be different spatial regions in the CASA model to fully represent the range of dynamics present throughout the stock range. The different modelling regions appear to be experiencing substantially different environmental conditions, different rates and temporal patterns of natural mortality, growth, and recent patterns of exploitation. Thus, it seems to the Review Panel that reference points are region-specific and that combining them, although done so in a mathematically correct way, introduces additional and unquantified risk. For example, a single reference point based on the entire region risks not identifying overfishing that could be occurring in Georges Bank. The Review Panel questions whether a single reference point for the entire region is appropriate.

TERM OF REFERENCE 6

Define and document methods for producing projections; provide justification for assumptions of fishery selectivity, fecundity, mortality and recruitment; comment on the reliability of resulting projections considering the effects of uncertainty and sensitivity to projection assumptions. Compare the results of SAMS and GeoSAMS and comment on their appropriateness for use in management.

The Review Panel felt that the information provided by the WG partially met this ToR. The Review Panel believes that the approach to providing projections for setting fishery quotas is reasonable and appropriately includes uncertainty in input parameters. However, as with ToR 3, the Review Panel noted a lack of documentation provided to it limits the Panel's full understanding of the details of how projections were made.

Projections are obtained through a computer simulation application called SAMS, which was initially developed in 1999. This program simulates the dynamics of Atlantic Sea Scallop in 24 separate geographic subareas; 7 in the Mid-Atlantic, 12 on Georges Bank, and five in the Gulf of Maine. Recruitment in each sub region is modeled stochastically with the log-transformed mean and covariance for recruitment in each area matching that observed in NEFSC dredge survey time series. Mean recruitment is then scaled to a region-wide Beverton-Holt stock recruitment relationship, making the SAMS model more comparable to the SYM reference points model. Within each region, separate dynamic models track annual patterns in growth, mortality, and exploitation. A fleet dynamics model distributes fishing activity among the different spatial regions. The model has been used in largely the same format since 1999. The simulation program predates application of CASA to Atlantic Sea Scallops. Although the scallop dynamics within SAMS are broadly similar to those in CASA, the two programs do not have any formal connection. The Review Panel requested evidence of the extent to which the dynamics of the two models are similar. During the meeting, the WG presented information from earlier comparisons of CASA and SAMS that show a high degree of coherence between the two models (Fig. 9).




Figure 9. Performance of model forecasted biomasses from CASA (solid black line) and SAMS (colored, dashed lines). The coherence of the forecasts between the models is taken as evidence that they are reliable for management use. Credit 2025 Atlantic Sea Scallop Research Track Assessment WG.

The WG presented evidence that there is a systemic bias in the performance of the SAMS model based on comparisons of 1-yr forward projections and observed values. The WG expressed belief that the bias—particularly in the Mid-Atlantic—results from the low natural mortality rates that were used in projections. Public comment at the meeting highlighted strong concerns over whether this makes the model useful in projections.

The Review Panel noted that the documentation for SAMS was particularly deficient and incomplete. The Panel strongly recommends additional details are provided in the future.

The Review Panel accepts the WG's explanation that the bias in model performance is likely to be related to the existence of high rates of natural mortality being present in both the Mid-Atlantic region and Georges Bank. This explanation is considerably strengthened by the results of the CASA model, which indeed suggest higher natural mortality rates in both regions than previously estimated. Accordingly, the Review Panel was surprised that SAMS was not updated for the 2025 Research Track Assessment to verify if the purported role for increased natural mortality does indeed reconcile the discrepancy between 1-yr forward projections and observations. The Review Panel also noted that the increases in base rates and deviations in natural mortality would be expected to increase variability among SAMS simulations.

The Review Panel was asked to comment on the development of geoSAMS, which was not ready to be reviewed at the 2025 Research Track. The Review Panel felt that insufficient information was provided to permit a fuller evaluation of the geoSAMS platform. The Review Panel appreciated that the finer spatial resolution of geoSAMS may lead to better performance of the fleet demand model. However, the Review Panel notes that the higher resolution may challenge the parameterization of recruitment, growth and natural mortality in the model; indeed, there may be a spatial mismatch between the scales at which the fleet demand model operates and those of the other population processes.

TERM OF REFERENCE 7

Review, evaluate, and report on the status of research recommendations from the last assessment peer review, including recommendations provided by the prior assessment working group, peer review panel, and SSC. Identify new recommendations for future research, data collection, and assessment methodology. If any ecosystem influences from TOR 1 could not be considered quantitatively under that or other TORs, describe next steps for development, testing, and review of quantitative relationships and how they could best inform assessments. Prioritize research recommendations.

The Review Panel felt that the WG fully met this ToR. The WG provided detailed responses to research recommendations from SARC 65, the 2018 Management Track Assessment and from annual SSC reports. Impressive progress has been made in some areas, such as the development and application of Generalized Random Tessellation Stratified (GRTS) methods for survey design for Georges Bank and the Mid-Atlantic. Incorporation of environmental data into

this Research Track Assessment is another example of where substantial progress has been made. The WG acknowledged transparently that there were other areas in which progress has been more limited. The Review Panel recognizes the challenge of addressing multiple research recommendations obtained from multiple venues and from different perspectives.

The Review Panel was uncertain over the extent to which the WG had responded to research recommendations received from the 2015 CIE review of the Atlantic Sea Scallop surveys. For example, this Review Panel report has already raised concerns over the small CVs associated with the Habcam survey under ToR 3, a point that was also raised in the 2015 CIE review. It was not clear to the Review Panel whether the 2015 CIE review was not included in the WGs comprehensive compilation of review recommendations because it predated SARC 65, but the Review Panel recommends the findings of this review be re-visited.

The Review Panel found that the specific research recommendations developed by the 2025 Research Track Assessment for Atlantic Sea Scallop to be limited in scope.

The Review Panel supports the effort to expand aging of scallop both to fill in historical gaps in the record, but also to assess whether age-based assessment models are feasible. The Review Panel recognizes the challenge of the indeterminacy of the age of first ring formation may represent, however, the panel felt that the power that age-based assessment models may provide make this assessment approach worthwhile.

The Review Panel strongly supports the recommendation to re-evaluate the current size-based approach to the assessment of sea scallop. New size-based approaches have been developed and applied to other species (e.g., Cao et al., 2017), some of which have a state-space structure, and more appropriately incorporate time-varying M and other model process errors as stochastic processes, and environmental covariates, which the 2025 Research Track Assessment have shown are important. Another factor motivating the change is that CASA is written in ADMB, a nonlinear estimation program that was state of the art when CASA was first coded. Today, ADMB is no longer being developed or supported, and thus its stability will come into question over the next 5–10 years. Some members of the Review Panel are already struggling in their own work to use ADMB code that once compiled successfully, but now no longer does. Additionally, the emerging generation of assessment analysts are more likely to code in TMB or RTMB and so the institutional knowledge of ADMB is waning. All these factors suggest that the WG should be actively exploring new approaches to assessing Atlantic Sea Scallop, so that they will be prepared by the time this species is next subject to a Research Track Assessment.

The Review Panel noted the concerns over the estimation of survey q's in the current model under ToR 4. The Panel believes that this likely results from constraints in other areas of the model leaving changes in q as one of the few pathways the model optimization can explore, although the panel has no specific evidence to support this belief. We view the substantial deviations in q from their priors to be a significant source of uncertainty in our understanding of the 2025 Research

Track assessment model performance, and a source of risk in interpreting model results. The Review Panel recommends strongly that additional work be conducted to explore alternative hypotheses regarding model structure and data inputs that may be driving the shifts in survey q's evident in the results.

The Review Panel discussed concerns over the reliability and consistency of annotation of optical images that impact the survey abundance and biomass time series derived from these images. The Review Panel discussed the extent to which best practices are followed in the analysis of images, which would see multiple annotators scoring the same image for some or all samples (Holmes et al., 2025). Practices such as the interspersion of test images into the analytical workflow to assess reliability of estimates are standard in routine aging laboratories and could perhaps be evaluated for optical surveys. Discussion with WG members indicated that some efforts to assess the reliability of image annotation occurs, but the lack of documentation of the QA/QC of annotation in the 2025 Research Track Assessment Report challenges the Review Panels full understanding of the processes followed. WG members expressed hope in further advancements in machine learning and AI algorithms may help solve these challenges. Even were this to be achieved, questions over reliability of image annotation would remain. The Review Panel recommends that those conducting and using the optical surveys evaluate how the reliability of extracting data from survey images can be improved. The Review Panel notes the importance of this action, given the discrepancies between the assumed and estimated catchabilities for the optical surveys.

The Review Panel has previously noted concerns over spatial autocorrelation that characterizes many optical surveys as well as the highly contagious distribution of Atlantic Sea Scallop themselves. The Review Panel recommends that the performance of the CASA or future assessment models be assessed when fitted using survey indices that downweight the highly contagious distribution of scallops in some stations by considering use of geometric means rather than arithmetic means.

The Review Panel supports the development of regional assessment models to characterize the dynamics of Atlantic Sea Scallops. The Review Panel noted in ToR 5 the incongruency of having spatial assessment models that are required to reflect the differences in the dynamics of scallop "populations" in the different regions and then combining results to produce a single management reference point. Explorations of the appropriate scale of stock definitions in this species seems appropriate. The Review Panel recommends research to identify and, where possible, characterize differences in productivity among the different spatial regions. Genetic analysis of stock structure using modern molecular approaches (e.g., radSeq or other approaches) appear warranted.

The 2025 Research Track Assessment clearly demonstrated the importance of environmental and ecosystem factors in understanding stock dynamics. Given that adult sea

scallops have greatly limited motility, the Review Panel suggests that development of habitat suitability models and joint species distribution models may be helpful to understanding the future trajectory of scallops in different sub regions. The Review Panel suggests that this might be particularly relevant in understanding the potential for recovery of scallop populations in closed areas in MAB.

TERM OF REFERENCE 8

Develop a backup assessment approach to providing scientific advice to managers if the proposed assessment approach does not pass peer review or the approved approach is rejected in a future management track assessment.

The Review Panel view this ToR as having been fully met.

The Research Track WG provided a hierarchical approach to addressing the challenge in which either the length-based model (CASA), or the SYM model or biological reference points become unavailable. The WG provides a logical sequence of decision-making of how management advice could continue to be provided. The Review Panel questioned how biological reference points could be estimated from catch and area swept data. The Review Panel suggested that data poor approaches in which catch and area swept information provide a foundation of biological reference point proxies. The Review Panel noted that Plan B does not include consideration of a case in which the SAMS model that is used to set catch advice fails. SAMS relies on estimates of M from CASA. While in the short term, SAMS could continue to be used in the absence of recent estimates from CASA, a point would be reached in which the input parameters to SAMS from prior CASA runs become unreliable. In such circumstances, analysts would be able to provide biological reference points, and stock status advice, but would not be able to provide catch advice.

Additional Material Considered by the Review Panel

The work presented related to fisheries and surveys in the Gulf of Maine contributed to the Review Panel's understanding of the extent, pattern and scale of fisheries for Atlantic Sea Scallop. The Review Panel supports further work to extend our knowledge of scallop fisheries in the Gulf of Maine such that future assessments may be able to present a unified analysis covering the entire range of scallops in US waters.

LITERATURE CITED

Caddy, J. F. 1973. Underwater Observations on Tracks of Dredges and Trawls and Some Effects of Dredging on a Scallop Ground. Journal of the Fisheries Research Board of Canada, 30: 173-+.

Cao, J., Chen, Y., and Richards, A. E. 2017. Improving assessment of *Pandalus* stocks using a seasonal size-structured assessment model with environmental variables. Part I: Model description and application. Canadian Journal of Fisheries and Aquatic Science, 74: 349-362.

Center for Independent Experts. 2015. Summary Report of the Review of Sea Scallop Survey Methodology and Their Integration for Stock Assessment and Fishery Management.

Clucas, G. V., Kerr, L. A., Cadrin, S. X., Zemeckis, D. R., Sherwood, G. D., Goethel, D., Whitener, Z., et al. 2019. Adaptive genetic variation underlies biocomplexity of Atlantic Cod in the Gulf of Maine and on Georges Bank. PLOS ONE, 14.

Ferraro, D. M., Trembanis, A. C., Mille, D. C., and Rudders, D. B. 2017. Estimates of Sea Scallop (Placopecten magellanicus) Incidental Mortality from Photographic Multiple before-after-Control-Impact Surveys. Journal of Shellfish Research, 36: 615-626.

Hart, D. R. 2013. Quantifying the tradeoff between precaution and yield in fishery reference points. ICES Journal of Marine Science, 70: 591-603.

Hart, D. R., and Chute, A. S. 2009. Estimating von Bertalanffy growth parameters from growth increment data using a linear mixed-effects model, with an application to the sea scallop. ICES Journal of Marine Science, 66: 2165-2175.

Holmes, S. J., McGregor, V. L., Underwood, M. J., and Wieczorek, A. M. 2025. Developments in scampi surveys: 2023-24. p. 19. Ed. by Fisheries New Zealand. Fisheries New Zealand, Wellington, NZ.

NEFSC 2018. 65th Northeast Regional Stock Assessment Workshop (65th SAW) Assessment Report. Ed. by N.M.F.S. United States. Northeast Fisheries Science Center, Woods Hole, MA.

Reid, K., Hoey, J. A., Gahagan, B. I., Schondelmeier, B. P., Hasselman, D. J., Bowden, A. A., Armstrong, M. P., et al. 2023. Spatial and temporal genetic stock composition of river herring bycatch in southern New England Atlantic herring and mackerel fisheries. Canadian Journal of Fisheries and Aquatic Sciences.

Sullivan, P. J., Lai, H. L., and Gallucci, V. F. 1990. A Catch-at-Length Analysis That Incorporates a Stochastic-Model of Growth. Canadian Journal of Fisheries and Aquatic Sciences, 47: 184-198.

Tian, R. C., Chen, C. S., Stokesbury, K. D. E., Rothschild, B. J., Cowles, G. W., Xu, Q. C., Hu, S., et al. 2009. Modeling the connectivity between sea scallop populations in the Middle Atlantic Bight and over Georges Bank. Marine Ecology Progress Series, 380: 147-160.

Appendix 1 – Terms of Reference for Atlantic Scallop Research Track Stock Assessment

1. Identify relevant ecosystem and climate influences on the stock. Characterize the uncertainty in the relevant sources of data and their link to stock dynamics. Consider findings, as appropriate, in addressing other TORs. Report how the findings were considered under impacted TORs.

2. Estimate catch from all sources including landings and discards. Describe the spatial and temporal distribution of landings, discards, and fishing effort. Characterize the uncertainty in these sources of data.

3. Present the survey data used in the assessment (e.g., indices of relative or absolute abundance, recruitment, state surveys, age-length data, application of catchability and calibration studies, etc.) and provide a rationale for which data are used. Describe the spatial and temporal distribution of the data. Characterize the uncertainty in these sources of data.

4. Use the appropriate assessment approach to estimate annual fishing mortality, recruitment and stock biomass (both total and spawning stock) for the time series, and estimate their uncertainty. Compare the time series of these estimates with those from the previously accepted assessment(s). Evaluate a suite of model fit diagnostics (e.g., residual patterns, sensitivity analyses, retrospective patterns), and (a) comment on likely causes of problematic issues, and (b), if possible and appropriate, account for those issues when providing scientific advice and evaluate the consequences of any correction(s) applied.

5. Update or redefine Status Determination Criteria (SDC; point estimates or proxies for BMSY, BTHRESHOLD, FMSY and MSY reference points) and provide estimates of those criteria and their uncertainty, along with a description of the sources of uncertainty. If analytic model-based estimates are unavailable, consider recommending alternative measurable proxies for reference points. Compare estimates of current stock size and fishing mortality to existing, and any redefined, SDCs. Provide stock status based on updated reference points.

6. Define and document methods for producing projections; provide justification for assumptions of fishery selectivity, fecundity, mortality and recruitment; comment on the reliability of resulting projections considering the effects of uncertainty and sensitivity to projection assumptions. Compare the results of SAMS and GeoSAMS and comment on their appropriateness for use in management.

7. Review, evaluate, and report on the status of research recommendations from the last assessment peer review, including recommendations provided by the prior assessment

working group, peer review panel, and SSC. Identify new recommendations for future research, data collection, and assessment methodology. If any ecosystem influences from TOR 1 could not be considered quantitatively under that or other TORs, describe next steps for development, testing, and review of quantitative relationships and how they could best inform assessments. Prioritize research recommendations.

8. Develop a backup assessment approach to providing scientific advice to managers if the proposed assessment approach does not pass peer review or the approved approach is rejected in a future management track assessment.

Appendix 2 – Initial agenda for Atlantic Scallop Research Track Assessment Peer Review meeting, April 21–24, 2025.

DRAFT AGENDA*
*All times are approximate at the discretion of the Review Panel Chair. This meeting is open to the public

Day 1: Monday April, 21 2025

Time	Topic	Presenter(s)	Notes
9:00 - 9:05 a.m.	Welcome & Logistics	Brian Hooper	
9:05 - 9:15 a.m.	Introductions & Agenda	Tom Miller	
9:15 - 9:40 a.m.	Overview of Research Track Process and Scallop Research Track	Kristan Blackhart	
9:40 - 10:10 a.m.	Management History and Fishery Practices	Jonathon Peros	
10:10 - 10:30 a.m.	Summary of Community Engagement Session	Jonathon Peros	
10:30 - 10:45 a.m.	BREAK		
10:45 - 11:30 a.m.	Scallop Life History	Dvora Hart	
11:30 - 12:00 p.m.	Term of Reference (TOR) 1 Overview	Dvora Hart	Ecosystem
12:00 - 1:00 p.m.	LUNCH		
1:00 - 1:30 p.m.	Temperature Influences	Joseph Caracappa	Ecosystem
1:30 - 2:00 p.m.	Shell Disease and Nematodes	Dave Rudders	Ecosystem
2:00 - 2:45 p.m.	TOR 1 Question and Answer (Q&A)	Review Panel	Ecosystem
2:45 - 3:00 p.m.	BREAK		
3:00 - 3:30 p.m.	TOR 2	Jessica Blaylock	Catch
3:30 - 4:00 p.m.	TOR 2 Q&A	Review Panel	Catch
4:00 - 4:15 p.m.	Public Comment	Public	

Time	Topic	Presenter(s)	Notes
4:15 - 5:00 p.m.	Discussion & Summary	Review Panel	Conclusions, Recommendations, & Final Wrap-up for TORs 1 & 2
5:00 p.m.	ADJOURN		

Day 2: Tuesday April 22, 2025

Time	Topic	Presenter(s)	Notes
9:00 – 9:05 a.m.	Welcome & Logistics Agenda	Brian Hooper Tom Miller	
9:05 - 9:25 a.m.	Gulf of Maine Surveys	Jonathon Peros	
9:25 - 10:10 a.m.	TOR 3	Dvora Hart	Surveys
10:10 - 10:45 a.m.	TOR 3 Q&A	Review Panel	Surveys
10:45 - 11:00 a.m.	BREAK		
11:00 - 12:00 p.m.	Mid-Atlantic Catch At Size Analysis (CASA) Model	Jui-Han Chang	Models
12:00 - 1:00 p.m.	LUNCH		
1:00 - 2:00 p.m.	Georges Bank Closed Areas CASA Model	Jui-Han Chan	Models
2:00 - 3:00 p.m.	Georges Bank Open Areas CASA Model	Jui-Han Chan	Models
3:00 - 3:15 p.m.	BREAK		
3:15 - 3:45 p.m.	TOR 4 Q&A	Review Panel	Models
3:45 - 4:00 p.m.	Public Comment	Public	
4:00 - 5:00 p.m.	Discussion & Summary	Review Panel	Conclusions, Recommendations, & Final Wrap-up for TORs 3 & 4
5:00 p.m.	ADJOURN		

Day 3: Wednesday April 23, 2025

Time	Topic	Presenter(s)	Notes
9:00 - 9:05 a.m.	Welcome & Logistics Agenda	Brian Hooper Tom Miller	
9:05 - 10:00 a.m.	Review Homework	Review Panel	If needed
10:00 - 10:45 a.m.	TOR 5	Dvora Hart	Biological Reference Points (BRPs)
10:45 - 11:00 a.m.	BREAK		
11:00 - 12:00 p.m.	TOR 5 Q&A	Review Panel	BRPs
12:00 - 1:00 p.m.	LUNCH		
1:00 - 1:45 p.m.	TOR 6	Dvora Hart	Projections
1:45 - 2:30 p.m.	TOR 6 Q&A	Review Panel	Projections
2:30 - 3:00 p.m.	TOR 8	Patrick Sullivan	Backup Approach
3:00 - 3:15 p.m.	BREAK		
3:15 - 3:45 p.m.	TOR 8 Q&A	Review Panel	Backup Approach
3:45 - 4:00 p.m.	Public Comment	Public	
4:00 - 5:00 p.m.	Discussion & Summary	Review Panel	Conclusions, Recommendations, & Final Wrap-up for TORs 5, 6, & 8
5:00 p.m.	ADJOURN		

Day 4: Thursday April 24, 2025

Time	Topic	Presenter(s)	Notes
9:00 - 9:05 a.m.	Welcome & Logistics Agenda	Brian Hooper Tom Miller	
9:05 - 10:30 a.m.	Review Homework	Review Panel	If needed
10:30 - 10:45 a.m.	BREAK		
11:00 - 11:30 a.m.	TOR 7	Jonathon Peros	Research Recommendations
11:30 - 11:45 a.m.	TOR 7 Q&A	Review Panel	Research Recommendations
11:45 - 12:00 p.m.	Public Comment	Public	
12:00 - 1:00 p.m.	Discussion & Summary	Review Panel	Conclusions, Recommendations, & Final Wrap-up TOR 7; any remaining issues
1:00 p.m.	ADJOURN		

Appendix 3 – Materials provided or referenced during the Atlantic Sea Scallop Research Track Stock Assessment Peer Review meeting

Working papers and presentations were available on a NEFSC website (https://apps-nefsc.fisheries.noaa.gov/saw/sasi.php) by selecting the species and year of assessment.

Working Papers and Background Documentation:

Report (Available on April 7)
2025_SCA_RT_Working Group Summary Report_rev.pdf

Background (generally available by April 7)
2025_SCA_RT_WP_Sea Scallop Life History Parameters.pdf
2025_SCA_RT_WP_Sea Scallop_TOR1a Ecosystem Influences.pdf
2025_SCA_RT_WP_Sea Scallop_TOR1b Climate Influences.pdf
2025_SCA_RT_WP_Sea Scallop_TOR2 Landings, Fishing Effort, Discards.pdf
2025_SCA_RT_WP_Sea Scallop_TOR3 Survey Data.pdf
2025_SCA_RT_WP_Sea Scallop_TOR4a Fishing Mortality, Recruitment and Stock Biomass.pdf
2025_SCA_RT_WP_Sea Scallop_TOR4b CASA Technical Description.pdf
2025_SCA_RT_WP_Sea Scallop_TOR5 reference points.pdf
2025_SCA_RT_WP_Sea Scallop_TOR6 projections.pdf
2025_SCA_RT_WP_Sea Scallop_TOR7 Research Recommendations.pdf
2025_SCA_RT_WP_Sea Scallop_TOR8 Plan B Backup Approach.pdf
2025_SCA_RT_WP_Sea Scallop_TOR9a Gulf of Maine Scallop Resources.pdf
2025_SCA_RT_WP_Sea Scallop_TOR9b Scallop Community Engagement Meeting.pdf
2025_SCA_RT_WP_Sea Scallop_TOR9c Management History.pdf

Presentations (generally available the day before the presentation)
SCA_Presentation_Atlantic Sea Scallop RT Review Intro – 2025.pdf
SCA_Presenation_Gulf of Maine.pdf
SCA_Presentation_Management History and Fishery Practices.pdf
SCA_Presentation_Summary of Community Engagement Meeting.pdf
SCA_Presentation_TOR 7_Research recommendations.odf
SCA_Presentation_TOR1_Nematodes and Shell Blisters.pdf
SCA_Presentation_TOR1_Overview.pdf
SCA_Presentation_TOR1_Temperature Influences.pdf
SCA_Presentation_TOR2_rev.pdf
SCA_Presentation_TOR3.pdf
SCA_Presentation_TOR4_CASA_rev.pdf
SCA_Presentation_TOR5.pdf
SCA_Presentation_TOR6.pdf
SCA_Presentation_TOR8.pdf
SCA_Presentation_life history.pdf

Figures (Generally available during meeting in response to Review Panel questions)
2025_SCA_RT_Homework_Additional figures.pdf
2025_SCA_RT_Howework_PlotAllStat.pdf
2025_SCA_RT_Homework_mortality_bubble_plots.pdf

Tables (Generally available during meeting in response to Review Panel questions)
2025_SCA_RT_Homework_TOR2.pdf

Appendix 4 – Meeting attendees at the Atlantic Sea Scallop Research Track Stock Assessment Peer Review meeting

GARFO - Greater Atlantic Regional Fisheries Office
MAFMC - Mid Atlantic Fisheries Management Council
NEFMC - Mid Atlantic Fisheries Management Council
NEFSC - Northeast Fisheries Science Center
SSC - Science and Statistical Committee

~~~~~~

Peer Review Panel
Thomas Miller, Chair - MAFMC SSC
Noel Cadigan, CIE
Yong Chen, CIE
Martin Cryer, CIE

Research Track Working Group
Patrick Sullivan, Chair - Cornell University
Jessica Blaylock, NEFSC
Jui-Han Chang, NEFSC
Adam Delargy, SMAST
Dvora Hart, NEFSC
David Keith, Canada
Amber Lisi, SMAST
Jonathon Peros, NEFMC
David Rudders, Virginia Institute of Marine Science

Attendees
Kiara Acevedo Martinez, NEFSC
Evan Balzano ,Maine Coast Fishermen's Association
Andrew Beet, NEFSC
Kristan Blackhart, NEFSC
Jason Boucher, NEFSC
Liz Brooks, NEFSC
Russell Brown, NEFSC
Connor Buckley, NEFMC
Steve Cadrin, SMAST
Joseph Caracappa, NEFSC
Toni Chute, NEFSC
Jamie Cournane, NEFMC
Kiersten Curti, NEFSC
Rachel Feeney, NEFMC
Corrin Flora, North Carolina Division of Marine Fisheries
Benjamin Galuardi, GARFO
Maxwell Grezlik, NEFSC
Melanie Griffin, Massachusetts Division of Marine Fisheries
Alex Hansell, NEFSC
~~~~~~

Eric Hansen, Hansen Scalloping Inc
Amanda Hart, NEFSC
Cameron Hodgdon, NEFSC
Brian Hooper, NEFSC
Carl Huntsberger, University of Maine
Kim Hyde, NEFSC
Emily Keiley, GARFO
Scott Large, NEFSC
Chris Legault, NEFSC
Emily Liljestrand, NEFSC
Nancy McHugh, NEFSC
Michael McManus, NEFSC
Drew Minkiewicz, Fisheries Survival Fund
Adelle Molina, NEFSC
Dana Morton, NEFSC
Robert Murphy, NEFSC
Chandler Nelson, NEFMC
Cate O'Keefe, NEFMC
Emily O'Toole, Coonamessett Farm Foundation
Stephanie Owen, NEFSC
John Pappalardo, Cape Cod Commercial Fishermen's Alliance
Jonathon Peros, NEFMC
Ted Platz, Ocean Harvest, Inc
John Quinn, Quinn Fisheries
Paul Rago, MAFMC SSC
Sally Roman, Virginia Institute of Marine Science
Sefatia Romeo Theken, MA Department of Fish and Game (FWE)
Melissa Smith, Maine Department of Maine Resources
Laura Solinger, NEFSC
Bridget St Amand, NEFSC
Kevin Stoksbury, SMAST
Michele Traver, NEFSC
Samuel Truesdell, NEFSC
Abigail Tyrell, NEFSC
Kelly Whitmore, Massachusetts Division of Marine Fisheries
Renee Zobel, New Hampshire Fish and Game

2025 ATLANTIC SEA SCALLOP RESEARCH TRACK ASSESSMENT

The Sea Scallop Research Track Working Group met 16 times (January 20, 2023; March 6, 2023; March 23, 2023; April 12, 2023; May 25, 2023; July 24, 2023; November 3, 2023; January 22, 2024; March 18, 2024; April 12, 2024; June 21, 2024; August 21, 2024; February 25, 2025; March 10, 2025; March 25, 2025; and April 4, 2025) in the development of this assessment. In addition, the Working Group had an in-person/virtual meeting with constituents at a community engagement meeting on December 19, 2024. This report was prepared by the members of the Atlantic Sea Scallop Research Track Working Group

ATLANTIC SEA SCALLOP RESEARCH TRACK WORKING GROUP (WG)

Patrick Sullivan (Cornell University, WG Chair)
Deborah Hart (NEFSC Population Dynamics, Assessment Lead)
Jui-Han Chang (NEFSC Population Dynamics, Co-Assessment Lead)
Jessica Blaylock (NEFSC)
Adam Delargy (University of Massachusetts, Dartmouth)
David Keith (Department of Fisheries and Oceans, Canada)
Amber Lisi (University of Massachusetts, Dartmouth)
Jonathon Peros (NEFMC)
David Rudders (VIMS, College of William and Mary)

Other participants involved in one or more of the meetings:
Larry Alade (NEFSC), Kristan Blackhart (NEFSC), Russ Brown (NEFSC), Connor Buckley (NEFMC), Joseph Caracappa (NEFSC), Alex Dunn (NEFSC), Glen Gawarkiewicz (WHOI), Melanie Griffin (MA DMF and NEFMC), Brian Hooper (NEFSC), Carl Huntsburger (ME DMR), Kim Hyde (NEFSC), Shannah Jaburek (GARFO), Chris Legault (NEFSC), Chris Kellogg (NEFMC), Scott Large (NEFSC), Drew Minkiewicz (Fisheries Survival Fund), Jose Montanez (MAFMC), Chandler Nelson (NEFMC), Cate O'Keefe (NEFMC), Tori Pilger (NEFSC), Paul Rago (retired NEFSC), Michele Traver (NEFSC), Abby Tyrell (NEFSC).

ACKNOWLEDGEMENTS

The WG would like to acknowledge the many people responsible for data collection, including survey groups, at-sea observers and port agents. The WG also appreciated input from Atlantic sea scallop constituents throughout the process.

TERMS OF REFERENCE

The purpose of the Atlantic Sea Scallop Research Track Assessment is to evaluate new and updated datasets and models with the ultimate goal of improving Atlantic sea scallop stock analysis. To accomplish this there are nine Terms of Reference (TORs) to be examined:

1. *Identify relevant ecosystem and climate influences on the stock. Characterize the uncertainty in the recent sources of data and their link to stock dynamics. Consider findings, as appropriate, in addressing other Terms of Reference. Report how the findings were considered under impacted Terms of Referenc*e.

2. *Estimate catch from all sources including landings and discards. Describe the spatial and temporal distribution of landings, discards, and fishing effort. Characterize the uncertainty in these sources of data.*

3. *Present the survey data used in the assessment (e.g., indices of relative or absolute abundance, recruitment, state surveys, age-length data, application of catchability and calibration studies, etc.) and provide a rationale for which data are used. Describe the spatial and temporal distribution of the data. Characterize the uncertainty in these sources of data.*

4. *Use the appropriate assessment approach to estimate annual fishing mortality, recruitment and stock biomass (both total and spawning stock) for the time series) and estimate their uncertainty. Compare the time series of these estimates with those from the previously accepted assessment(s). Evaluate a suite of model fit diagnostics (e.g., residual patterns, sensitivity analyses, retrospective patterns), and (a) comment on likely causes of problematic issues, and (b), if possible and appropriate, account for those issues when providing scientific advice and evaluate the consequences of any correction(s) applied.*

5. *Update or redefine Status Determination Criteria (SDC; point estimates or proxies for BMSY, BTHRESHOLD, FMSY and MSY reference points) and provide estimates of those criteria and uncertainty, along with a description of the sources of uncertainty. If analytic model-based estimates are unavailable, consider recommending alternative measurable proxies for reference points. Compare estimates of current stock size and fishing mortality to existing, and any redefined, SDCs. Provide stock status based on updated reference points.*

6. *Define and document methods for producing projections; provide justification for assumptions of fishery selectivity, fecundity, mortality and recruitment; comment on the reliability of resulting projections considering the effects of uncertainty and sensitivity to projection assumptions. Compare the results of SAMS and GeoSAMS and comment on their appropriateness for use in management.*

7. *Review, evaluate, and report on the status of research recommendations from the last assessment peer review, including recommendations provided by the prior assessment working group, peer review panel, and SSC. Identify new recommendations for future research, data collection, and assessment methodology. If any ecosystem influences from Term of Reference 1 could not be considered quantitatively under that or other Terms of Reference, describe next steps for development, testing and review of quantitative relationships and how they could best inform assessments. Prioritize research recommendations.*

8. *Develop a backup assessment approach to providing scientific advice to managers if the proposed assessment approach does not pass peer review or the approved approach is rejected in a future management track assessment.*

EXECUTIVE SUMMARY

*Term of Reference (TOR) #1: Identify relevant ecosystem and climate influences on the stock. Characterize the uncertainty in the recent sources of data and their link to stock dynamics. Consider findings, as appropriate, in addressing other Terms of Reference. Report how the findings were considered under impacted Terms of Referenc*e.

Elevated adult natural mortality and reduced adult abundance was first observed in the far south of the scallops' range, off of Virginia, in the early 2000s, and these phenomena have been gradually expanding northward and to inshore areas throughout the Mid-Atlantic region, likely due to increasing bottom temperatures. This is addressed in the Mid-Atlantic CASA model by estimating natural mortality by year and size.

Food supply (phytoplankton), temperature, pH, predators, invasive species, and parasites and diseases can all affect scallop populations. Growth and weight at shell height tend to be greater at shallower depths, probably due to increased food supply.

The sea star *Astropecten americanus* appears to be reducing or excluding sea scallops from the deeper water of the Mid-Atlantic and is spreading northward and into shallower waters. *Cancer* crabs are a major predator of juvenile sea scallops, and may be acting as agents of density dependence, explaining elevated natural mortality rates associated with large year classes.

The invasive colonial tunicate *Didemnum vexillum* has been observed in portions of Georges Bank and Gulf of Maine and can exclude scallops from settling in the areas that they have colonized. Since 2015, a substantial portion of the scallops in the southern Mid-Atlantic have been infected with larvae of the nematode *Sulcascaris sulcata*. The effects of these parasites on the scallops is unclear, but they do cause difficulty in marketing the scallop meats.

The prevalence of shell blister disease, likely caused by *Polydora* sp. worms, has been increasing in recent years, perhaps associated with warming temperatures. These blisters are associated with higher scallop mortality rates. Gray meat disease has been observed mostly in Georges Bank and can be caused by infection by Apicomplexa protists. High mortality in the early 2010s in portions of some of the closed areas of Georges Bank has been attributed to gray meat disease.

TOR #2: Estimate catch from all sources including landings and discards. Describe the spatial and temporal distribution of landings, discards, and fishing effort. Characterize the uncertainty in these sources of data.

Sea scallop landings in the US increased substantially after the mid-1940s, with peaks occurring around 1960, 1978, 1990, 2004, and 2020. Maximum US landings were 29,109 mt meats

in 2004. US landings during 2001–2012 were all over 20,000 mt and landings in 2019 were 27,649 mt. Landings in 2023 were 12,699 mt meats.

Efforts in the US sea scallop fishery generally increased from the mid-1970s to about 1991, and then decreased during the 1990s, first because of low catch rates, and later as a result of effort reduction measures. Total effort since 2007 has been variable and decreasing slightly, with some shifts between regions.

On average, discards were 5.2% of total landings for 1989–2023 and 5.9% for 2010–2023. Scallop dredges likely kill and injure some scallops that are contacted by the dredge but not caught. We estimated incidental mortality to be 10% of fully recruited fishing mortality on Georges Bank and 5% in the Mid-Atlantic.

TOR #3: Present the survey data used in the assessment (e.g., indices of relative or absolute abundance, recruitment, state surveys, age-length data, application of catchability and calibration studies, etc.) and provide a rationale for which data are used. Describe the spatial and temporal distribution of the data. Characterize the uncertainty in these sources of data.

Long-term scallop surveys throughout Georges Bank and the Mid Atlantic Bight have been conducted using three methods and by multiple survey groups. The Northeast Fisheries Science Center (NEFSC) conducted regular scallop dredge surveys since 1975, and with a consistent lined dredge since 1979. The Virginia Institute of Marine Science (VIMS) has conducted a similar dredge survey focused on specific areas since 1999 and region-wide surveys annually from 2014. The NEFSC also has conducted a towed camera survey (Habcam) since 2011. The Habcam Group and Coonamessett Farm Foundation (CFF) have also conducted Habcam surveys annually since 2005. Lastly, the University of Massachusetts Dartmouth, School for Marine Science and Technology (SMAST) have been conducting a drop camera survey that focused on specific regions from 1999 and resource-wide surveys from 2003. This Term of Reference describes the surveys and collates annual estimates of scallop abundance and biomass from each of these surveys. These estimates were used to fit the models implemented in this Research Track.

TOR #4: Use the appropriate assessment approach to estimate annual fishing mortality, recruitment and stock biomass (both total and spawning stock) for the time series) and estimate their uncertainty. Compare the time series of these estimates with those from the previously accepted assessment(s). Evaluate a suite of model fit diagnostics (e.g., residual patterns, sensitivity analyses, retrospective patterns), and (a) comment on likely causes of problematic issues, and (b), if possible and appropriate, account for those issues when providing scientific advice and evaluate the consequences of any correction(s) applied.

Size-structured catch-at-size-analysis models were used to assess sea scallops in three regions: Georges Bank Open, Georges Bank Closed, and Mid-Atlantic, from 1975 to 2023. The CASA models were tuned to catch, commercial shell heights from port and at-sea observer sampling, and survey data for both abundance and shell height. The population in CASA was modeled with 5 mm shell height bins, starting at 0–5 mm. Recruitment was empirically estimated at age 1 (0–35 mm; with no assumed stock-recruitment relationship). Scallops larger than 40 mm were used in tuning. Growth in CASA models was advanced using growth transition matrices estimated from growth increment data outside of CASA. Natural mortality in the CASA model can be estimated by year and size for juveniles and adults, as well as annually without varying by size (Hart and Chang 2022). Natural mortality was estimated for juveniles and adults for the Mid-Atlantic, annually for the Georges Bank Closed model, and for juveniles only for the Georges Bank Open model. AIC and our knowledge of these stocks were used as the primary tools for model selection.

The base case models for all three regions converged and fit catch and survey indices and size composition data without apparent residual patterns. The CASA estimated abundance and biomass generally aligned well with survey observations, although CASA biomass tended to be below the dredge survey estimates for early years in the Georges Bank open model. Compared to previous assessments, estimated abundance and biomass from the current CASA model are more in line with the dredge survey, especially for the Mid-Atlantic model. No serious retrospective patterns were observed in any of the three stocks.

The average estimated natural mortality from all sizes and years for the Mid-Atlantic, Georges Bank closed, and Georges Bank open stocks were 0.42, 0.27, and 0.3, respectively. These estimates were higher than those from the previous assessment, which ranged between 0.22 to 0.27 (NEFSC 2020). The models estimated elevated adult natural mortality in the Mid-Atlantic in recent years and elevated juvenile natural mortality in the Georges Bank open areas during large recruitment events.

Whole-stock fishing mortality generally increased from 1975 to 1992, then declined strongly from 1992 to 1995, and has remained fairly low but has increased somewhat in recent years. Abundance and biomass were low in all three areas until the mid-1990s, then rapidly increased. Abundance and biomass have rapidly declined in recent years in the Mid-Atlantic. Whole stock biomass, abundance, and fishing mortality in the terminal year 2023 were 69,956 mt meats, 5,112 million scallops, and 0.33, respectively.

TOR #5: Update or redefine Status Determination Criteria (SDC; point estimates or proxies for BMSY, BTHRESHOLD, FMSY and MSY reference points) and provide estimates of those criteria and uncertainty, along with a description of the sources of uncertainty. If analytic model-based estimates are unavailable, consider recommending alternative measurable proxies for reference points. Compare estimates of current stock size and fishing mortality to existing, and any redefined, SDCs. Provide stock status based on updated reference points.

Reference points were calculated using the Stochastic Yield Model (SYM) separately in the Mid-Atlantic and Georges Bank and then combined. F_{MSY} was estimated as 0.36 on Georges Bank, 1.56 in the Mid-Atlantic, and 0.49 for the combined resource. B_{MSY} was estimated as 83414 mt meats for Georges Bank, 15909 mt meats for the Mid-Atlantic, and 93282 mt meats for the combined resource.

Combined biomass in 2023 from the three CASA models was 69956 mt meats. This is below the target biomass (B_{MSY}) but above $B_{threshold}$ = 46641 mt meats, so the stock is not overfished. Fishing mortality for the combined stock in 2023 was 0.33, which is below F_{MSY}; hence overfishing is not occurring. However, if sea scallops were managed as two separate stocks (Mid-Atlantic and Georges Bank), overfishing would have been occurring on Georges Bank in 2023.

TOR #6: Define and document methods for producing projections; provide justification for assumptions of fishery selectivity, fecundity, mortality and recruitment; comment on the reliability of resulting projections considering the effects of uncertainty and sensitivity to projection assumptions. Compare the results of SAMS and GeoSAMS and comment on their appropriateness for use in management.

A spatial forecasting model (the Scallop Area Management Simulator, SAMS) is used in sea scallop management. In the model the resource is divided into a number of subareas; in the current configuration, there are 7 subareas in the Mid-Atlantic, 12 on Georges Bank, and 5 in the Gulf of Maine.

Population dynamics is modeled similarly in SAMS to CASA and SYM, that is, populations are size-structured and use stochastic growth transition matrices. However, population dynamics are modeled for each subarea in SAMS, and so are on a finer scale than the other models. SAMS can simulate the rotational management used in the fishery, including closing chosen areas, and then reopening them to area-specific quotas.

Scallopers are allocated a number of "open" area days, which they can fish in any area that is not closed or under special access. SAMS allocates fishing effort to the open subareas by assuming effort is proportional to LPUE. LPUE is estimated using a linear regression between mean exploitable biomass and observed LPUE in the open areas.

TOR # 7: Review, evaluate, and report on the status of research recommendations from the last assessment peer review, including recommendations provided by the prior assessment working group, peer review panel, and SSC. Identify new recommendations for future research, data collection, and assessment methodology. If any ecosystem influences from Term of Reference 1 could not be considered quantitatively under that or other Terms of Reference, describe next steps for development, testing and review of quantitative relationships and how they could best inform assessments. Prioritize research recommendations.

Previous Research Recommendations

The working group reviewed research recommendations from the 2018 benchmark stock assessment and peer review in (SAW/SARC 65); the 2020 management track assessment and peer-review, and the New England Fishery Management Council's Science and Statistical Committee (SSC) meetings in 2020, 2021, 2022, 2023, and 2024. The following (here and in more detail below and in the Working Paper) is a catalogue of research recommendations from the most recent stock assessments and SSC input. The working group notes that not all SSC comments and recommendations are within the purview of the working group.

Previous Research Recommendations

- Coordinate survey programs
- Dredge efficiency
- Time-varying mortality
- Gulf of Maine fishery
- Unreported landings
- Spatially explicit methods (GEOSAMS)
- Disease
- VMS data
- LPUE forecasting
- Thermal impacts
- Incidental mortality
- Area specific growth rates
- Gonad-based estimates of SSB
- Bridging old to new models
- Dynamic selectivity in SYM
- Recruitment assumptions in SAMS
- Survey re-stratification
- Factors affecting scallop stock dynamics
- Evaluate projection model
- Increased natural mortality in some regions

New Research Recommendations

- Age scallops
- Spatially explicit methods (GEOSAMS)
- Environmental impacts
- State-space methods
- Review of growth assumptions
- Predator-prey relationships
- Stock structure
- AI for scallop images

TOR #8: Develop a backup assessment approach to providing scientific advice to managers if the proposed assessment approach does not pass peer review or the approved approach is rejected in a future management track assessment.

The WG identified the parts of the process where a potential failure to achieve an acceptable result might occur. Failure could occur in the application of the CASA assessment model used to estimate population abundance or in the SYM determination of reference points, or both. Swept area survey estimates of density expanded to suitable habitat areas is an alternative

method to CASA for deriving abundance. Deterministic calculation of reference points is an alternative method to SYM. Failure of both suggests reverting to a swept area estimate with average catch relative to this biomass to determine appropriate exploitation rates.

Additional discussions are provided on the scallop resource in the Gulf of Maine, and results from the scallop community engagement meeting.

INTRODUCTION

Distribution

The Atlantic sea scallop, *Placopecten magellanicus*, is a bivalve mollusk that occurs on the eastern North American continental shelf from Cape Hatteras to the Gulf of St. Lawrence and Newfoundland, typically on firm sand and gravel bottoms. Major aggregations in US waters occur in the Mid-Atlantic from Virginia to Long Island at depths of 35 to 80 m, on Georges Bank, including the Great South Channel and Nantucket Shoals, at depths from 40 to 160 m, and, to a lesser extent, in the Gulf of Maine, mainly in relatively shallow waters (Hart and Chute 2004). This assessment focuses on the two main portions of the sea scallop stock and fishery, Georges Bank in the north and the Mid-Atlantic in the south. Results for Georges Bank and the Mid-Atlantic are combined to evaluate the stock as a whole.

Life History

Sea scallops feed by filtering phytoplankton, microzooplankton, and detritus particles. Sexes are separate and fertilization is external. Larvae spend 5−7 weeks in the water column before settling. Major predators include sea stars (e.g., *Asterias* spp. and *Astropecten americanus*) and *Cancer* spp. crabs (Hart and Chute 2004, Hart 2006, Shank et al. 2012). Scallops fully recruit to the NEFSC dredge survey at 40 mm SH, and to the current commercial fishery at around 90–110 mm SH, although scallops as small as 70 mm were landed during the 1980s and 1990s.

Sexual maturity commences at age 2, corresponding to shell heights (SH) of about 35–75 mm. Sea scallops > 40 mm that are reliably detected in the surveys used in this assessment are thus all considered mature individuals. However, individuals younger than 4 years may contribute little to total egg production, but fecundity increases rapidly with age (MacDonald and Thompson 1985; Hart and Chute 2004, Hennen and Hart 2012).

Spawning generally occurs in late summer or early autumn throughout the sea scallops' range. Spring spawns or minor dribble spawns at other times can also occur. The spring spawn is often strong in the Mid-Atlantic Bight (DuPaul et al. 1989) and spring spawns on Georges Bank have also been observed and may be becoming stronger with increases in winter temperatures (Almeida et al. 1994, Dibacco et al. 1995, Thompson et al. 2014). The timing of spawning has no direct effect on a size-based stock assessment.

Growth

Sea scallop growth can be inferred using visible "rings" laid down on the shell. These rings have been confirmed as annual marks, although the year one ring is typically missing (Stevenson and Dickie 1954, Merrill et al. 1966, Hart and Chute 2009a, Chute et al. 2012). Obtaining absolute age from shell rings can be problematic for some scallops since early rings may be obscure, especially on older scallops (Claereboudt and Himmelman, 1996). For this reason, Hart and Chute (2009b) treated the distance between rings as annual growth increments, with age unknown. They introduced a method to estimate von Bertalanffy growth parameters from such data which includes (individual) random effects on the parameters L_∞ and K. This allows not only estimation of mean von Bertalanffy coefficients, but also their variability among individuals in the population. This method was updated for this assessment using shells collected between 1982 and 2023. Growth has varied with time. For this reason, the time series was split into four growth time blocks in the Mid-Atlantic, and two each in the Georges Bank Open and Closed areas. Estimates are given in the table below. More details can be found in the Life History Working Paper.

Growth Table. Estimated von Bertalanffy growth parameters by region (MA=Mid-Atlantic, GBCl=Georges Bank Closed, GBOp=Georges Bank Open) and year. The SD columns are estimates of the variability of the parameter among individuals, whereas the SE columns give the standard error of the population mean parameter.

Region	Period	L_∞	K	SDL	SDK	SEL	SEK
MA	1990–99	125.3	0.570	7.1	0.13	0.35	0.003
MA	1975–1978; 1984–89; 2000–03	130.3	0.570	7.5	0.13	0.36	0.003
MA	1979; 1982–3; 2004–8;2019–20	134.3	0.570	7.9	0.13	0.37	0.003
MA	2009–12; 2021–22	136.9	0.570	8.2	0.13	0.37	0.003
GBCl	1986–97; 2015–22	142.7	0.444	9.6	0.12	0.34	0.002
GBCl	1975–85; 1998–2014	147.5	0.444	10.3	0.12	0.35	0.002
GBOp	1987–90; 2011–22	140.7	0.455	8.1	0.11	0.16	0.002
GBOp	1975–86; 1991–2010	143.4	0.455	8.5	0.11	0.17	0.002

Shell height to weight relationships

Shell height-meat weight relationships allow conversion from numbers of scallops at a given size to meat weights. They are expressed in the form $W = \exp(\alpha + \beta \ln(H))$, where W is meat weight in grams and H is shell height in mm.

Shell height/meat weight data have been collected during annual NEFSC sea scallop surveys since 2001, and in recent years, also by VIMS dredge surveys. These data have been used in scallop assessments since 2007 and were updated for this assessment. Parameters were estimated using mixed-effect generalized linear models, similar to Hennen and Hart (2012). The baseline shell height to meat weight relationships used in the CASA model are given in the table below. Relationships for Georges Bank Open and Georges Bank Closed did not significantly differ, so a single relationship for both was used.

Meat weights depend on factors which affect feeding and metabolic rates, including depth and location. Meat weights decrease with depth, probably because of reduced food (phytoplankton) supply. Depth and subarea have a significant effect on the shell height/meat weight relationship (Hennen and Hart 2012). In this assessment, covariate-adjusted shell height/meat weight relationships were used to calculate survey biomass, and the simple weight shell height relationship was used in the models (CASA, SYM and SAMS), where depth is not explicit.

Meat weights for scallops in the commercial fishery may differ from those predicted from research survey data for a number of reasons. First, the shell height-meat weight relationship varies seasonally, in part due to the reproductive cycle, so that meat weights collected during the dredge surveys during the late spring and early summer may differ from those in the rest of year. Additionally, commercial fishers concentrate on speed and often leave some meat on the shell during shucking (Naidu 1987, DuPaul et al. 1990). On the other hand, meats in fishery catches may gain weight due to water uptake during storage on ice. Finally, fishers may target areas with relatively large meat weight at shell height, which may increase commercial meat weights compared to those collected on research vessels.

Observer data was used to adjust meat weights for seasonal variation and for commercial practices. Annual commercial meat weight anomalies were computed based on the seasonal patterns of landings together with the mean monthly commercial meat weight at shell height. More details on shell height to meat weight conversions can be found in the Life History Parameters working paper.

Shell height to meat weight table $W = \exp(\alpha + \beta \ln(SH))$

Region	α	β
MA	-9.8076	2.784
GB	-10.262	2.851

Natural mortality

The 2018 and 2020 assessments assumed adult natural mortalities of $M = 0.2$ on Georges Bank and 0.25 in the Mid-Atlantic, although in some cases, M was estimated by year in the CASA models (Hart and Chang 2022). We used estimators based on longevity (maximum observed age) to obtain baseline estimates of M of 0.4 for the Mid-Atlantic and 0.27 for Georges Bank (see the Life History Parameters working paper for details). All three CASA models estimate M by year, although only juvenile M in the Georges Bank Open model and not by size in the Georges Bank Closed model. See TOR-4 for more details.

Ecosystem and Climate Influences

*TOR #1: Identify relevant ecosystem and climate influences on the stock. Characterize the uncertainty in the recent sources of data and their link to stock dynamics. Consider findings, as appropriate, in addressing other Terms of Reference. Report how the findings were considered under impacted Terms of Referenc*e.

Ecosystem influences

The most important environmental driver of the U.S. sea scallop fishery in recent years has been increasing bottom temperatures in the Mid-Atlantic Bight area that has reduced the productivity of scallops in this region. As will be discussed later in this assessment, recruitment in this region has been very low during the last 6 years of the assessment time period (2018–2023), and adult natural mortality has been increasing. This is most evident spatially, with few adult scallops observed in 2023 in the southern and inshore portions of the Mid-Atlantic. Increased adult mortality was first observed in the southernmost surveyed area, off of Virginia, in the early 2000s, and has been slowly spreading northward since then. Over half of all U.S. sea scallop landings were from the Mid-Atlantic region between 2000–2019, but only about 5% of landings were from this region in 2023. There is evidence that bottom temperatures are increasing rapidly, and in some areas may be exceeding the scallops' upper temperature tolerance during the early autumn. Further details regarding this phenomenon and an exploratory analysis can be found in the Caracappa et al. (2025) working paper.

There is strong evidence that food supply (mostly phytoplankton, but also detritus and microzooplankton) affects growth and reproductive output of sea scallops (MacDonald and Thompson 1985,1986ab, MacDonald et al. 1987, Zang et al. 2022). Phytoplankton supply declines with increasing depth (Shumway et al. 1987), explaining observed declines in growth, and weight at size (Langton et al. 1987, Barber et al. 1988, Hart and Chute 2009b, Hennen and Hart 2012). For survey data, these effects are taken into account by using shell height to meat or gonad weight relationships that depend on depth. The SAMS model uses area-specific growth and allometric relationships, based on data from surveys and shell growth analysis. For example, scallops in the relatively deep water Hudson Canyon area are relatively slow growing with smaller meats at size, and scallops in the shallow inshore Mid-Atlantic area are faster growing with larger meats at size. The CASA and SYM models, which operate on a region-wide level, cannot explicitly take geographic variations in growth and weights into account. However, the different growth periods built into the CASA model can reflect changes in growth and weights at size over time, possibly due to environmental conditions, and due to shifts of the distributions of scallops to, e.g., deeper or shallower areas.

Ocean acidification (OA) may affect scallop populations. In particular, there have been several studies demonstrating that OA can reduce the growth and survival of larval scallops

(Talmage and Gobler 2009, White et al. 2013, Andersen et al. 2013), and swimming ability (Schalkhausser et al. 2013). Pousse et al. (2023) showed that OA reduces feeding rates and growth of juvenile sea scallops. Similarly, Lagos et al. (2023) showed that OA reduced growth and shell thickness in the Peruvian scallop *Argopecten purpuratus*. Cooley et al. (2015) and Rheuban et al. (2018) used a forward projection model similar to a non-spatial version of SAMS together with forecasts of future ocean warming and acidification. They predicted that OA will reduce sea scallop yields in the second half of this century, due to reduced growth and larval survival.

Predators can affect scallop distribution and mortality. In particular, the sea star *Astropecten americanus* appears to be reducing or excluding sea scallops from the deeper water of the Mid-Atlantic (Hart 2006, Shank et al. 2012). This sea star consumes a wide variety of small invertebrates, including early post-settlement sea scallops (Franz and Worley 1982, Adebola et al. 2022), and is commonly observed at very high densities in the Mid-Atlantic outer shelf. *A. americanus* appear to be limited by cold water temperatures, and for that reason, are only common in areas where winter minimum bottom temperatures remain above 5° C (Franz et al. 1981). *A. americanus* densities have been monitored in dredge surveys since 2000, and there is evidence that their distribution is expanded northward and inshore, likely due to warming temperatures (NEFSC 2018). Elevated abundance of the sea star *Asterias vulgaris* and concomitantly increased numbers of sea scallop "clappers", an indicator of natural mortality, were observed on surveys of the northern edge of Georges Bank in 2024, in both U.S. and Canadian waters.

Cancer spp. crabs (*Cancer irroratus* and *C. borealis*) are important predators on juvenile sea scallops (shell heights < 90 mm, Elner and Jamison 1979, Nadeau et al. 2009, Hart and Shank 2011) and may be agents of density dependence. Wong et al. (2005) seeded juvenile scallops in experimental plots at densities of 1, 6, or 69 m^2. Scallop density in the high-density sites declined markedly due to both predation, primarily by *Cancer* spp. crabs, and dispersal, resulting in final densities of about 1 m^2 regardless of treatment. Predation rates of *Cancer* crabs on juvenile sea scallops were greater when scallops are more common than alternative prey species and increase with increasing scallop density (Barbeau et al. 1998, Wong and Barbeau 2005). This is accounted for in this assessment by directly estimating juvenile natural mortality by year.

The invasive colonial ascidian *Didemnum vexillum* has been observed in portions of Georges Bank and Gulf of Maine. This tunicate can rapidly spread over gravel/cobble substrate, often completely covering it. Scallop larvae cannot settle on *D. vexillum* (Morris et al. 2009), so this tunicate turns preferred settlement substrate into unavailable habitat for juvenile scallops. Consistent with this, Habcam survey data indicate that sea scallops are less common in areas dominated by *D. vexillum* (Working Paper TOR1a Figure 1.2, Kaplan et al. 2017). Additionally, this tunicate is more common in fished areas than in an adjacent closed area. This may be due to fishing activity dispersing the tunicates by both direct transport and fragmentation (Morris and Carman 2012). *D. vexillum* appears to be restructuring the benthic invertebrate community in areas that it dominates (Kaplan et al. 2018). Fortunately, *D. vexillum* has been observed to date only in

relatively small portions of Georges Bank, mainly on the northern edge near the Hague line border with Canada.

Off-colored "gray meats" have been most commonly observed on Georges Bank, and in particular in Closed Area I (Levesque et al. 2016). They have been associated with infections with Apicomplexa protists (Inglis et al. 2016). However, Siemann et al. (2019) suggested that gray meats are more of a symptom of poor condition and can occur due to multiple causes. Gray meats due to Apicomplexa infections may have contributed to the increased natural mortality observed in the Georges Bank closed areas during 2011–13. The variable natural mortality included in the Georges Bank Closed Areas model is one method to account for this.

Since 2015, a substantial portion of the scallops in the Mid-Atlantic have been infected with larvae of the nematode *Sulcascaris sulcata*, particularly in the south (Rudders et al. 2023; Working Paper TOR1a Figures 1.3, 1.4). They create brown or orange lesions in the scallop meat, caused by the immune response in the scallops. The adult nematode lives in the gastrointestinal track of sea turtles, primarily loggerheads, who acquire the nematodes by consuming infected scallops or other mollusks. These nematodes are associated with scallop and sea turtle populations around the world (Lichtenfels et al. 1978, 1980; Lester et al. 1980). It is likely that this outbreak of nematodes was due, at least in part, to the large 2013 scallop year class in the Mid-Atlantic, which may have increased the amount of scallops consumed by the turtles. Increases in the loggerhead turtle population off the northeast US may also be a contributing factor. It is unclear whether infections by the nematodes increases the natural mortality of the scallops. Lightly infected scallops do not appear to be seriously negatively affected by the nematodes, but heavy infections may increase the scallops' mortality. However, the nematodes have had a clear effect on the fishery, as heavily infected scallops are difficult to market (even though there is no public health risk).

Sea scallop blister disease occurs when the shell underneath the gonad gets colonized by a fouling organism, most likely *Polydora* spp. polychaete worms. The scallop then puts shell material over this area, creating a blister (Working Paper TOR1a Figure 1.5). Shell blisters are associated with reduced growth, yield (Working Paper TOR1a Figure 1.6), fecundity, deteriorated physical condition, mortality, inhibited water filtration due to reduced shell-cavity volume, and weakened and/or deformed shell (Kent, 1979, Lauckner 1983, Diez et al., 2011). Blisters contain anaerobic metabolites like hydrogen sulfide, which can reduce quality and marketability (Handley and Bergquist 1997). Two decades ago, blister disease was limited to scallops in the deeper water in the southern half of the Mid-Atlantic, suggesting that it is limited by cold temperatures. However, blister disease is becoming more common throughout the Mid-Atlantic and has also been observed on Georges Bank and the Bay of Fundy, likely due to increasing bottom temperatures.

Landings, fishing effort, discards

TOR #2: Estimate catch from all sources including landings and discards. Describe the spatial and temporal distribution of landings, discards, and fishing effort. Characterize the uncertainty in these sources of data.

Management history

The sea scallop fishery in the US Exclusive Economic Zone (EEZ) is managed under the Atlantic Sea Scallop Fishery Management Plan (FMP), first implemented on May 15, 1982. A detailed account of scallop fishery management actions and applications of science into management advice is contained in the Working Paper for TOR2.

From 1982 to 1993, the primary management control was a minimum average meat weight requirement for landings, commonly known as the "meat count" requirement. In 1984, Georges Bank was divided into US and Canadian EEZs; prior to this time, US and Canadian vessels fished on both sides of the current boundary.

Amendment 4 of the Sea Scallop Plan (NEFMC 1993), implemented in 1994, changed the management strategy from meat count regulation to limited access, effort control and gear regulations for the entire US EEZ. Limited access permits were issued to vessels with a history in the fishery; no new permits have been issued since. Restrictions were gradually made on days-at-sea (DAS), minimum ring size, and crew limits. DAS have been reduced from over 200 in fishing year 1994 to 120 during 1999–2003 to 24 in open areas in 2018–2022, and finally to 20 in 2023. Note that the scallop "fishing year" used by management was historically March-February, but this was changed to April-March starting in April 2018; however, unless otherwise stated, this assessment uses calendar years.

The minimum size of the rings in the dredge bag was gradually increased from 76 mm (3") in 1994 to 83 mm (3.25") in 1995, 89 mm (3.5") during 1996–2004 and 102 mm (4") since December 2004. The minimum size of the twine top mesh has also been increased from 15 to 25 cm (6" to 10") since December 2004; while this measure is mainly to reduce bycatch of finfish, it also likely allows some small swimming scallops to escape. Finally, the crew size on a vessel has been restricted, usually to seven persons, to incentivize fishing for larger scallops (since most scallops must be shucked at sea by regulation, a limited crew can process a greater weight of large scallops than smaller ones per unit time). In addition to these measures, three large areas on Georges Bank and Nantucket Shoals were closed to groundfish and scallop fishing in December 1994 (Murawski et al. 2000, Almeida et al. 2005). Scallop biomass increased about 20-fold in these areas between 1994–2004 (Hart and Rago 2006). Two areas in the Mid-Atlantic were closed to scallop fishing in April 1998 for three years to similarly increase scallop biomass and mean weight.

Sea scallops were formally declared overfished in 1997, and Amendment 7 was implemented during 1998 with more stringent DAS limitations and a mortality schedule intended to rebuild the stocks within ten years. Subsequent analyses conducted in 1999, using an early version of the Scallop Area Management Simulator (SAMS) model that took into account for the first time the rebuilding effects of closed areas, indicated that the stocks would rebuild with less severe effort reductions than called for in Amendment 7, so the DAS schedule was thus modified. A combination of closures, effort reduction, gear and crew restrictions led to a rapid increase in biomass (Murawski et al. 2000, Hart and Rago 2006, Hart et al. 2013), and sea scallops were declared to be rebuilt in 2001.

Prior to 2004, there were a number of *ad hoc* area management measures, including the Georges Bank and Mid-Atlantic closures in 1994 and 1998, limited reopenings of portions of the Georges Bank areas between June 1999 and January 2001, and reopening of the first Mid-Atlantic rotational areas in 2001. A new set of regulations was implemented as Amendment 10 (NEFMC 2003) during 2004. This amendment formalized an area-based management system (Hart 2003, Hart and Rago 2006, O'Keefe and NEFMC PDT 2022), with provisions and criteria for new rotational closures, and separate allocations (in DAS or total allowable catch [TAC]) for reopened closed areas and general open areas. The three Georges Bank closed areas were divided into access areas, where fishing is periodically permitted, and long-term closures, where no scallop fishing is permitted. Some of the long-term closures were reopened to scallop fishing in 2018.

Unlike the Georges Bank closures, which applied to all scallop and groundfish fishing, the Mid-Atlantic rotational areas are specific to the scallop fishery. Two areas (Hudson Canyon South and Virginia Beach) were closed in 1998 and then reopened in 2001. Although the small Virginia Beach closure in the far south of the scallops range was unsuccessful, scallop biomass built up in Hudson Canyon access area while it was closed, and it supported substantial landings during 2001–2007. This area was again closed in 2008, reopened in 2011 and closed for a third time in 2014, and reopened in 2015. A third rotational closure, the Elephant Trunk access area east of Delaware Bay, was closed in 2004 after extremely high densities of small scallops were observed in surveys during 2002 and 2003. About 30,000 mt of scallops worth about $500 million were landed from that area after it was reopened in 2007. It was closed again in December 2012 after high numbers of small scallops were again observed in surveys. A portion of this area was reopened in 2015, and the remainder was reopened in 2017. A fourth access area, Delmarva, directly south of the Elephant Trunk area, was initially closed in 2007, reopened in 2009, closed in 2012 and reopened in 2014. In 2016, the Delmarva, Elephant Trunk, and Hudson Canyon areas were combined into the Mid-Atlantic access area (MAAA), which remained open to scallop fishing in subsequent years. However, a decrease in scallop productivity in the region led managers to revert the Delmarva portion of the MAAA to open bottom in 2018 and revert the entire remaining MAAA back to open bottom in 2022. In the same year, the new New York Bight rotational area was implemented and closed to protect growth of scallop in the area.

Most landings have come from about 350 vessels with limited access permits. Two types of allocation are given to each of these vessels. The first are trips (with a trip limit, typically 12,000–18,000 lbs or 5443–8165 kg meats) to specified rotational access areas that had been closed to scallop fishing in the past. The second are DAS, which can be used in areas outside the closed and access areas. Vessels fishing under DAS are restricted to a 7-man crew and must shuck their scallops at sea to limit their processing power.

The remainder of landings come from vessels operating under "General Category" permits that are restricted to a certain amount per trip, with a maximum of one trip per day. This limit was originally 181 kg (400 lbs) meats, which was increased to 272 kg (600 lbs) meats in 2011. The trip limit in accessed areas was increased to 363 kg (800 lbs) meats in fishing year 2022. Landings from these vessels were less than 1% of total landings in the late 1990s but increased to about 10% of landings during 2007–2009 and currently constitute about 6% of total landings. This type of permit had been open access but was converted to an individual transferable quota (ITQ) fishery in March 2010.

The primary port in the sea scallop fishery is New Bedford, MA, which accounted for over 75% of landings in 2022 and 2023. With scallop distribution and fishing effort shifting north over time, other principal ports have changed in recent years, with Point Judith, RI and Gloucester, MA, becoming more important at the expense of Cape May, NJ, and Seaford, VA. Lesser amounts of scallops are landed in many ports from North Carolina to Maine. Toothless offshore (New Bedford style) scallop dredges are the main gear type in all regions, although otter trawls are used to some extent in the Mid-Atlantic, and a small fraction of the catch in the Gulf of Maine comes from divers. A typical limited access vessel tows two 4–4.6 m dredges, but some limited access vessels are restricted to a single 3.2 m dredge (termed a "small dredge permit"). Most general category vessels also use a single 3.2 m or smaller dredge, but some use otter trawls in the Mid-Atlantic. Most bycatch of sea scallops from other fisheries occurs in otter trawls, where the target species are squid, flounders, and other groundfish, although this is relatively small compared to the directed fishery. Recreational catch is negligible.

Landings

Prior to 1994, landings and effort data were collected from dealer data and port interviews by port agents. Since 1994, commercial data have been available as dealer reports (DR) and in vessel trip report (VTR) logbooks. DR give landings, but not area fished, and have reported landings by market category since 1998. VTR data contain information about area fished, fishing effort, and retained catches of sea scallops. A standardized area allocation (AA) method (Wigley et al. 2008) for matching DR to VTRs and assigning areas to landings was used to allocate landings to region from 1994–2019. In 2020, the Catch Accounting and Monitoring System (CAMS; O'Keefe et al. 2023) became the single comprehensive data source for all northeast commercial

catch (landings and discards), consolidating information from several databases (including DR, VTR) and the AA process.

US landings data were stratified by region into four groups: Mid-Atlantic (statistical areas ≥ 600 and 539), Southern New England (statistical areas 533–534, and 536–538), Georges Bank (statistical areas 520–526, 541–543, 561–562, and 551–552), and the Gulf of Maine (statistical areas 463–464, 467, 500, and 511–515). Landings from unknown areas were prorated to each region in proportion to their known landings. Note that landings from Block Island (statistical area 539) are now included with Mid-Atlantic to be consistent with how surveys are considered in the assessment; this differs from previous assessments where landings from Block Island were included with Southern New England. Data were also stratified by gear into three groups: trawls (otter trawls and beam trawls), dredge (scallop dredge, rakes, clam/quahog dredge, other dredge), and other (all other gears). All 1964–2023 landings data were updated to incorporate any data corrections (data retrieval date 07/17/2024); 1887–1963 landings are from the 2018 stock assessment (NEFSC 2018).

Most landings of sea scallops only retain the adductor muscle, or "meat", although there is a small market for roe-on and whole scallops. If not otherwise specified, landings in this assessment will be in terms of meat weight.

Sea scallop landings in the US increased substantially after the mid-1940s, with peaks occurring around 1960, 1978, 1990, 2004, and 2020. Maximum US landings were 29,109 mt meats in 2004. US landings during 2001–2012 were all over 20,000 mt and landings in 2019 were 27,649 mt; the maximum in the 20th century was 17,246 mt in 1990. Landings in 2023 were 12,699 mt meats.

Landings from the Georges Bank and Mid-Atlantic regions have dominated the fishery since 1964 (Figure 2.1). US Georges Bank landings had peaks during the early 1960s, around 1980 and 1990, but declined precipitously in 1993 and remained low through 1998. Landings from Georges Bank during 1999–2004 were fairly steady, averaging almost 5,000 mt annually, and then increased in 2005–2006, primarily due to reopening of portions of the groundfish closed areas to scallop fishing. Georges Bank landings again increased in 2012–2013, this time mainly due to shift of open effort from the Mid-Atlantic to Georges Bank, to take advantage of large year classes in the latter region but declined in the next few years. Georges Bank landings peaked again in 2018 and 2019 at over 15,000 mt, the largest values in the time series from this region, driven by the exceptionally large 2012-year class. In most recent years, landings from this region have remained high (above 10,000 mt).

Prior to the mid-1980s, Mid-Atlantic landings were generally lower than those on Georges Bank. Mid-Atlantic landings during 1962–1982 averaged less than 1,800 mt per year. An upward trend in both recruitment and landings became evident in the Mid-Atlantic in the mid-1980s and continued into the early 2000s. Landings in this region peaked in 2004 at 23,565 mt before

gradually declining since, reflecting declining recruitment and increasing natural mortality, and concomitant effort shift onto Georges Bank. Landings from the Mid-Atlantic in 2023 were under 700 mt, reminiscent of levels in the early 1970s.

Landings from other areas (Gulf of Maine and Southern New England) were minor in comparison. Gulf of Maine landings represented over 7% of total US landings in 2022 and 2023, increasing from less than 3% in the prior two decades. Maximum landings in the Gulf of Maine were 1,616 mt during 1980 but trended downward afterwards through 2009, when landings were just 71 mt. Gulf of Maine landings have been increasing in the most recent years; landings in 2022 were the highest since 1981 at 1,096 mt.

Dealer data (landings) have been reported by market categories (under 10 meats per pound, 10–20 meats per pound, 20–30 meats per pound etc.) since 1998. These data indicate a trend towards larger sea scallops in landings. While nearly half the landings in 1998 were in the smaller market categories (more than 30 meats per pound), 75% or more of recent landings were below 20 count and about 99% were below 30 count. In the late 1990s, Mid-Atlantic and Georges Bank scallop landings consisted of a rather uniform distribution of different market sizes. Starting in the early 2000s, the 20–30 count (15–23 g) market category dominated landings. Around 2005, the 10–20 count (23–45 g) market category became dominant indicating the presence of older larger scallops in the catch, a result of rotational management implementation. The largest scallops (U10, >45 g) also became more prevalent during that time, starting in the Mid-Atlantic and shifting to Georges Bank as the spatial distribution of the stock and effort changed. Market categories in the Gulf of Maine do not show this pattern as strongly.

Fishing effort and LPUE

Fishing effort (measured as days absent) increased in the Mid-Atlantic starting in the mid-1970s, peaking in the early 2000s and decreased on average since then. In contrast, the importance of Georges Bank has increased in recent years as effort shifted to that region. The Gulf of Maine appears to have had significant effort prior to 2000 but this is primarily due to the presence of vessels that were targeting sea urchins but also landed sea scallops. Overall, effort from the Gulf of Maine and Southern New England regions is minimal relative to Georges Bank and the Mid-Atlantic.

Landings per unit effort (LPUE) were computed as landings per day fished, where days fished represent the time in days that the gear is fishing. This was obtained from the port interview records from larger vessels prior to 1994 and from at-sea observers on large dredge limited access vessels since 1994. LPUE showed a general downward trend from the beginning of the time series to around 1998, with occasional spikes upward due to strong recruitment events. LPUE increased considerably since then as the stock recovered. Note the close correspondence in most years between the LPUE in the Mid-Atlantic and Georges Bank, probably reflecting the mobility of the fleet; if one area has higher catch rates, it is fished harder until the rates are equalized. Although

comparisons of LPUE before and after the change in data collection procedures during 1994 need to be made cautiously, there is no clear break in the LPUE trend in 1994.

Fishing effort (days fished) was computed as the quotient of Landings/LPUE. Effort is thus in units of days fished on large dredge limited access vessels; general category vessels, which usually only fish with one small dredge, would need to fish for several days to account for a single "day fished" under this metric. Efforts in the US sea scallop fishery generally increased from the mid-1970s to about 1991, and then decreased during the 1990s, first because of low catch rates, and later as a result of effort reduction measures. Effort increased in the Mid-Atlantic during 2000–2007, initially due to reactivation of latent effort among limited access vessels, and then due to increases in general category effort. Total effort since 2007 has been variable and decreasing slightly, with some shifts between regions.

Data from vessel monitoring systems (VMS) that are required on all scallop vessels fishing on Georges Bank and the Mid-Atlantic can give detailed spatial information on fishing activity. The VMS gives the positions of the vessels every half hour, from which average speed can be calculated. These data were then filtered to eliminate times where the vessels were not fishing (either steaming or simply shucking scallops without gear in the water) (Wigley and Palmer 2008), and the resulting spatial distributions were plotted.

Discards and discard mortality

Sea scallops are sometimes discarded on directed scallop trips because they are too small to be economically profitable to shuck, or because of high grading, particularly during access area trips. Ratios of discard to total catch (by weight) have been recorded by sea samplers aboard commercial vessels since 1992, although sampling intensity on non-access area trips was low until 2003.

Discarded sea scallops may suffer mortality on deck due to shell damage, high temperatures, or desiccation. There may also be mortality after they are thrown back into the water from physiological stress, or from increased predation due to shock and inability to swim (Veale et al. 2000, Jenkins and Brand 2001). Murawski and Serchuk (1989) estimated that about 90% of tagged scallops were still living several days after being tagged and placed back in the water. Total discard mortality (including mortality on deck) is uncertain but has been estimated as 20% in previous assessments (e.g., NEFSC 2010, NEFSC 2018). However, discard mortality may be higher in the Mid-Atlantic during the summer due to high water and deck temperatures, and likely strongly depends in both regions on crew practices; scallops returned to the water promptly have much higher chances of survival than ones left on deck for longer periods.

Discard estimates for 1989–2023 are presented in Table 2.3 in the TOR-2 working paper. Discards for 1989–2017 were calculated using the Standardized Bycatch Reporting Methodology (SBRM; Wigley et al. 2008). This approach uses a combined d/k_{all} ratio estimator (Cochran 1977),

where d is discarded pounds of sea scallops and k_{all} is kept pounds of all species, calculated from the Northeast Fisheries Observer Program (NEFOP) data. Discard weight was derived by multiplying the d/k_{all} ratio of each fleet by the corresponding dealer or VTR landings. Additional details regarding the discard estimates are available in previous assessment reports for 1989–2009 (Appendix B2 in NEFSC 2014) and 2010–2017 (Appendix A4 in NEFSC 2018). Discards for 2018–2023 are from CAMS, which provide improved estimates. The original (SBRM) approach separated observer information from catch and then matched discard rates to unobserved trips based on stratification, such that different gear table definitions sometimes led to situations where observed strata did not exist in the catch or vice versa. In contrast, CAMS joins the observer and catch information prior to estimation, which leads to improved discard estimates.

On average, discards were 5.2% of total landings for 1989–2023 and 5.9 % for 2010–2023. By region, most discarding occurred in Southern New England (12.9% average for 2010–2023), especially during 2014–2020 when discards were close to 20% of landings, ranging up to 27.2 % in 2020. Discarding in the Mid-Atlantic varied over time with low periods representing less than about 5% of landings, and high periods (1994, 2001–2004, 2016, 2023) representing about 10–15 % of landings. Discarding in Georges Bank was low (<5%) or moderate (5–10%) in most years, with a few years (e.g., 2010, 2020) when discarding was >10%, with a peak of 20% in 2000. The smallest proportion of discarding occurred in the Gulf of Maine region where discards were typically less than 4% until 2021, with a little higher in 2022 and 2023 (4.9% and 5.5%, respectively).

Incidental mortality

Scallop dredges likely kill and injure some scallops that are contacted but not caught, primarily due to damage (e.g., crushing) caused to the shells by the dredge. Caddy (1973) estimated that 15–20% of the scallops remaining in the track of a dredge were killed. Murawski and Serchuk (1989) estimated that less than 5% of the scallops remaining in the track of a dredge suffered non-landed mortality. Caddy's study was done in a relatively hard bottom area in Canada, while the Murawski and Serchuk study was in sandy bottom off the coast of New Jersey. It is possible that the difference in indirect mortality estimated in these two studies was due to different bottom types (Murawski and Serchuk 1989). A 2017 study estimated somewhat lower incidental mortality rates of 2.5% in the Mid-Atlantic and 8% on Georges Bank (Ferraro et al. 2017). Two other unpublished studies presented during working group meetings suggest similar rates (Bochenek et al., Smolowitz et al.).

To use the above estimates to relate landed and non-landed fishing mortality in stock assessment calculations, it is necessary to know the efficiency *e* of the dredge (the probability that a fully recruited scallop in the path of a dredge is captured). Similarly, the fraction of scallops that suffer mortality among sea scallops in the path of the dredge but not caught is denoted by *c*. The ratio *R* of scallops in the path of the dredge that were caught to those killed but not caught is:

$$R = \frac{e}{c(1-e)} \quad (1)$$

If scallops suffer direct (i.e., landed) fishing mortality at rate F_L, then the rate of indirect (non-landed) fishing mortality will be (Hart 2003):

$$F_I = F_L/R = \frac{1}{e}F_L c(1-e) \quad (2)$$

Assuming c = 0.025 and e = 0.6 for the Mid-Atlantic and c = 0.08 and e = 0.5 for Georges Bank gives estimates of F_I of about 0.02 for the Mid-Atlantic and 0.08 for Georges Bank. Using an estimate of c = 0.04 from Murawski and Serchuk (1989) for the Mid-Atlantic and c = 0.12 from Caddy (1973) for Georges Bank gives estimates of incidental mortality of 0.03 and 0.12, respectively. The CASA model does not explicitly model discarding or discard mortality of small scallops, although it is included in the SYM and SAMS model. Because of this, the working group agreed to set c = 0.05 in the Mid-Atlantic and c = 0.1 on Georges Bank in the SYM and SAMS models, and c = 0.06 and 0.11, respectively, in the CASA model.

The above calculations assume that the scallop is fully selected to the gear. If that is not the case, equation 2 becomes:

$$F_I(h) = \frac{1}{e}F_L c(1 - e * q(h)) \quad (3)$$

where $q(h)$ is the catchability of commercial gear on a scallop of shell height h. We took $q(h)$ to be of the form:

$$q(h) = q_0 s(h)$$

where q_0 is taken as 0.5 on Georges Bank and 0.6 in the Mid-Atlantic, and $s(h)$ is commercial selectivity as estimated by the CASA model.

Commercial shell height data

Since most sea scallops are shucked at sea, it has often been difficult to obtain reliable commercial size compositions. Port samples of shells brought in by scallopers have been collected, but there are questions about whether the samples were representative of the landings and catch. Port samples taken during the meat count era (mid-1980s through 1993) often appear to be selected for their size rather than being randomly sampled, and the size composition of port samples from 1992–1994 differed considerably from those collected by at-sea observers during this same period. For this reason, commercial size compositions from port samples after 1984 when meat count regulations were in force are not used in this assessment.

Sea samplers (observers) have collected shell heights of kept scallops from commercial vessels since 1992, and discarded scallops since 1994. Although these data are likely more reliable than that from port sampling, they still must be interpreted cautiously for years prior to 2003

(except for the access area fisheries) due to limited observer coverage. Except for 2006, observer coverage rates have been over 5% since 2003, and have been over 10% in some years.

Shell heights from port and sea sampling data indicate that sea scallops between 70–90 mm often made up a considerable portion of the landings during 1975–1998, but sizes selected by the fishery have increased since then, so that scallops less than 90 mm were rarely taken since 2002.




Figure 2.1 U.S. sea scallop landings (mt meats) for the Mid-Atlantic (MA), Georges Bank (GB), Southern New England (SNE), and Gulf of Maine (GOM) regions for 1965–2023. Landings where the area was unknown were prorated.

SURVEY DATA

TOR #3: Present the survey data used in the assessment (e.g., indices of relative or absolute abundance, recruitment, state surveys, age-length data, application of catchability and calibration studies, etc.) and provide a rationale for which data are used. Describe the spatial and temporal distribution of the data. Characterize the uncertainty in these sources of data.

Dredge surveys

Sea scallop dredge surveys were conducted by NEFSC in 1975 and annually after 1977 to measure the abundance and size composition of sea scallops in the Georges Bank and Mid-Atlantic regions, with a consistent lined gear since 1979. The Virginia Institute of Marine Science (VIMS) conducted intensive dredge surveys of selected regions on commercial vessels since 1999, and region-wide dredge surveys of the Mid-Atlantic Bight since 2014. Except for the first few years, VIMS surveys have towed one 8’ survey dredge identical to the NEFSC dredge, and one commercial dredge. All VIMS data collected from the survey dredge for fully covered strata (original or post-stratified) were treated in the same way as NEFSC tows. The partially randomized

grid design was treated as random when calculating variances. This likely slightly overstates the true sample variance.

Based on dredge survey estimates, biomass and abundance on Georges Bank were generally low until around 1995 (Figures 3.1 and 3.2). Very large increases were observed during 1995–2000 after implementation of closures and effort reduction measures and have remained high since.

In the Mid-Atlantic Bight, dredge abundance and biomass indices were at low levels during 1979–1997, and then increased rapidly during 1998–2003 due to area closures, reduced fishing mortality, changes in fishery selectivity, and strong recruitment. Biomass was relatively stable during 2003–2008, but then declined, in part due to poor recruitment and fishing down of rotational areas. Survey shell height frequencies show a trend to larger shell heights in both regions since 1995.

Drop camera survey

Region-wide drop camera surveys have been conducted by the School for Marine and Technology since 2003 (Stokesbury et al. 2004, Bethony and Stokesbury 2018). The survey coverage of the drop camera has varied throughout the Mid-Atlantic and Georges Bank regions over the years. Therefore, the coverage was standardized to a common footprint, where scallop density was predicted within unsurveyed areas within the common footprint using a generalized additive modelling approach with latitude, longitude, year, and depth as covariates. The drop camera survey used a systematic grid sampling design, and sizes of grids were used in different regions and years to reflect funding and management interests at the time. The standard error of mean density estimates from the survey were calculated assuming a stratified random design, whereby each grid location (typically with four drops) was treated as a stratum.

The primary cameras on this survey from 2003–2015 were standard definition (SD) “large” video cameras; surveys since then have used much higher resolution digital cameras. For the purposes of CASA modeling, the SD and digital camera surveys are considered to be separate time series.

The drop camera survey estimated Georges Bank abundance to be between 2 and 5 billion scallops from 2003 to 2014 (Figure 3.2), before higher abundances were estimated from 2015 to 2018 with a peak of 8.21 billion scallops in 2017. Drop camera abundance estimates on Georges Bank have been between 4 and 6 billion scallops since 2019. Biomass estimates on Georges Bank fluctuated between 50,000 and 110,000 mt from 2003 to 2015 and peaked at 167,297 metric tons in 2018. Biomass estimates then declined to between 50,000 and 100,000 metric tons from 2019.

The drop camera survey estimated Mid-Atlantic abundance at 10.39 billion scallops in 2003, and this was followed by substantial decline to 2.74 billion scallops by 2011. Abundance

increased again and reached 9.90 billion estimated individuals by 2015. The drop camera only surveyed the full Mid Atlantic again in 2017 and 2019, but these estimates followed the declining trend of abundance observed by the other surveys at these times. The drop camera Mid Atlantic biomass estimates followed a similar pattern to the abundance estimates, with 105,169 mt estimated in 2003, around 43,295 mt in 2012, 149,269 mt in 2017 and declining biomass estimates after this point.

Habcam towed camera survey

The Habitat Mapping Camera (Habcam) is an underwater towed digital camera system. The camera(s) take rapid-fire photos of the sea floor (typically 5–6/sec) as it is towed at speeds between 5–7 knots at roughly 2 m above the bottom. Four Habcam vehicles have been used, with most of the data presented here from the V3 and V4 systems, both of which have two digital still cameras. Because of the large number of images collected, only subsets were examined for sea scallop measurements and counts, most commonly, about 2% of the images taken (roughly one every 30 m), corresponding to about 100,000 annotated images per year. Data from the tracks were interpolated to the survey domain using a regression kriging approach (Chang et al. 2017).

The Habcam surveys estimated Georges Bank abundance to be around 4 billion individuals from 2011 to 2013 (Figure 3.2). Abundance then increased to 8 and 10 billion over the following three years due to a record strong year class. After these peak years of abundance, abundance declined back to slightly less than 4 billion scallops from 2019 onwards. Habcam estimated Georges Bank scallop biomass to be 102,676 metric tons in 2011 and then estimated a decline to 47,351 metric tons by 2013. This was then followed by years of higher biomass, with a peak of 148,920 metric tons in 2016. Georges Bank biomass was estimated to be around 50,000 metric tons from 2019 onwards.

The Habcam surveys estimated the Mid Atlantic scallop abundance to be around 5 billion from 2012 to 2014. A sharp increase to 15.8 billion scallops was observed in 2015, due to a strong year class. This peak value was followed by a rapid decline in abundance to around 2 billion individuals from 2020 onwards. The Habcam biomass estimates from the Mid Atlantic followed a similar pattern to the abundance estimates, with 49,186 metric tons estimated in 2012, a peak of 121,781 metric tons in 2016 and 17,934 metric tons estimated in 2023.

Scallops in the Deep Southeast Portion of Nantucket Lightship Closed Area

The very large 2012-year class on Georges Bank included very dense settlement in relatively deep water in the southeast portion of Nantucket Lightship Closed Area, where previously there had been very few scallops. Scallops in this area grew very slowly, and were nicknamed “Peter Pan” scallops, because they did not want to grow up.


In 2014, many of these scallops were below the 40 mm threshold for inclusion in the survey gear abundance estimates and were only partially selected to the drop camera survey due to the lower-resolution standard definition video camera used in this survey prior to 2016. For these reasons, the Habcam survey was used to estimate abundance of these scallops. The peak abundance in 2015 was about 11 billion scallops, according to the Habcam survey, an extraordinary number that exceeds the number of scallops in the whole resource for some years. Maximum biomass was in 2017 at about 63000 mt meats. Estimates for scallops in this area have declined across all

surveys from these peaks, with all surveys estimating less than 50 million scallops above 40 mm shell height in this area in 2023.

Gulf of Maine surveys

Scallop surveys have been conducted in the Gulf of Maine federal waters using the SMAST drop camera and the Maine Department of Marine Resources (DMR) unlined survey dredge since 2009. The data from these surveys are not currently used in the CASA or SYM models, but they are detailed in the Gulf of Maine working paper.

(a) Georges Bank



(b) Mid-Atlantic




Figure 3.1. Dredge time series for numbers (left) and biomass (right) (dots), including lowess smoothers (lines) for (a) Georges Bank and (b) Mid-Atlantic.




Figure 3.2. Comparison of survey estimates, 2003–2023, for Georges Bank (left), and Mid-Atlantic (right). Numbers are on the top row, and biomass on the bottom row.

FISHING MORTALITY, RECRUITMENT, AND STOCK BIOMASS

TOR #4: Use the appropriate assessment approach to estimate annual fishing mortality, recruitment and stock biomass (both total and spawning stock) for the time series) and estimate their uncertainty. Compare the time series of these estimates with those from the previously accepted assessment(s). Evaluate a suite of model fit diagnostics (e.g., residual patterns, sensitivity analyses, retrospective patterns), and (a) comment on likely causes of problematic issues, and (b), if possible and appropriate, account for those issues when providing scientific advice and evaluate the consequences of any correction(s) applied.

Model Configuration

Catch-at-size-analysis (CASA, Sullivan et al. 1990, Hart et al. 2013, Hart and Chang 2022) is the primary assessment estimation model; it has been used since 2007 (NEFSC 2007, 2010, 2014, 2018, and 2020). Separate models were configured for Georges Bank's open and closed areas, whereas the Mid-Atlantic was assessed using a single CASA model (Figure 4.1). The Deep Southeast Nantucket Lightship scallops (DSENLS) were not included in the CASA modeling due to their unique location and growth patterns.

All three CASA models, Mid-Atlantic, Georges Bank open, and Georges Bank closed, were run from 1975 to 2023. The CASA models were tuned to catch, unlined dredge, lined dredge, winter bottom trawl (Mid-Atlantic only), drop camera (split into SD and digital camera time series), and Habcam surveys. Details of catch and survey information can be found in TOR 2 and 3 working papers. The lined dredge, Habcam, and drop camera digital camera surveys were assumed to have flat selectivity for scallops 40+ mm. Selectivities of the drop camera "large" SD camera and unlined dredge were fixed at experimentally determined values. Selectivity for the winter bottom trawl survey was modeled using an ascending logistic curve with parameters fixed at values estimated by the SARC-59 assessment (NEFSC 2014). Beta prior probabilities are used to incorporate knowledge regarding absolute scale from the surveys. Priors on survey catchability were applied to four broad-scale scallop-targeted surveys, including lined dredge, large SD drop camera, digital drop camera, and Habcam surveys.

The population in CASA was modeled with 5 mm shell height bins, starting at 0–5 mm. Recruitment was estimated at age 1 (0–35 mm), with no assumed stock-recruitment relationship. As in previous assessments, only scallops larger than 40 mm were used in tuning because smaller scallops were not fully selected by any of the surveys. Growth in CASA models was advanced using growth transition matrices estimated from growth increment data outside of CASA. Natural mortality in the CASA model can be estimated by year and size, as well as annually but constant with size (Hart and Chang 2022). The size-varying natural mortality by year was estimated as the sum of juvenile and adult mortality

components (mean natural mortality plus annual deviance for each component) with a logistic curve to partition the natural mortalities between juveniles and adults. The natural mortality estimation options were explored and evaluated for each stock. AIC and our knowledge of these stocks were used as the primary tools for model selection.

Base Case Model Results

The base case model for all three stocks converged and fit catch, survey indices, and size composition data without apparent residual patterns. The CASA estimated abundance and biomass generally aligned well with survey observations (Figures 4.2–4.4), although the estimates tended to be below the dredge survey estimates for early years in the Georges Bank open model.

For the Mid-Atlantic model, size-specific natural mortality was estimated for both juveniles and adults, with the mean natural mortality fixed at 0.4 based on likelihood profiling and life history parameters (see Life History WP). As a result, the mean natural mortality for all years and sizes increased from 0.27 from the previous assessment (NEFSC 2020) to 0.42 for the Mid-Atlantic stock. For the Georges Bank closed model, natural mortality was estimated by year without varying by size, and the average natural mortality by year was 0.27, which also increased compared to the previous assessment (0.23; NEFSC 2020). For the Georges Bank open model, the mean natural mortality for juveniles and adults was not estimated and was fixed at 0.27 based on the estimate from the Georges Bank closed stock, and only juvenile natural mortalities were estimated. The mean estimated natural mortalities were 0.3 for all years and sizes, whereas the mean was 0.22 for the updated SARC-65 assessment (NEFSC 2020). The models estimated elevated adult natural mortality in Mid-Atlantic in recent years and elevated juvenile natural mortality in Georges Bank during large recruitment events.

Fishery selectivities were logistic for most years and models, except for the domed shaped selectivities for the earliest period in the Mid-Atlantic model and some recent years in Georges Bank closed. The doming in the Mid-Atlantic was due to fishing being concentrated in a relatively small area during these early years; in the Georges Bank closed area, it was due to fishing that was concentrated in the Nantucket Lightship area which contained mostly intermediate sized scallops, while most of the large scallops occurred in other areas that were lightly fished or were closed. Fishery selectivity in general has shifted strongly towards large scallops since 1998 stock wide.

Whole-stock fishing mortality generally increased from 1975 to 1992, then declined strongly from 1992 to 1995, and has remained low but increased somewhat in recent years (Fig 4.3). The trend in whole-stock abundance and biomass is more or less the reverse of the trend in fishing mortality but with higher variations (Fig 4.2). Whole stock biomass, abundance, and fishing mortality in the terminal year 2023 were 69,956 mt meats, 5112

million scallops, and 0.33, respectively, not including the DSENLS scallops in Georges Bank. Retrospective scores for the entire sea scallop stock were mild (Mohn's =0.17 for biomass and -0.09 for fishing mortality), and individual stocks showed minor to moderate retrospective patterns; none of these were serious enough to require a retrospective adjustment.

Historical retrospective comparisons

The current CASA model estimated abundance, biomass, and fishing mortality were compared to those from the last five assessments (SARC-45/NEFSC 2007, SARC-50/NEFSC 2010, SARC-59/NEFSC 2014, SARC-65/NEFSC 2018, and updated SARC-65/NEFSC 2020) and lined dredge survey (Figure 4.5). While the model performance in Georges Bank was relatively stable and slightly improved by this assessment, with less bias in the earlier years' abundance and biomass estimates, noticeable improvements were made for the Mid-Atlantic model, which no longer produces estimates that are trended low compared to the dredge survey.




Figure 4.1. Charts of U.S. sea scallop grounds, showing the Mid-Atlantic and Georges Bank stock areas. Georges Bank was split into Open, Closed, and Deep Southeast Nantucket Lightship (DSENLS) portions.




Figure 4.2. Comparison of CASA model estimated biomass with expanded estimates from the lined dredge (red triangle), SMAST large camera (blue cross), HabCam (green dots), and SMAST digital camera (light green x) for Mid-Atlantic areas.




Figure 4.3. Comparison of CASA model estimated biomass with expanded estimates from the lined dredge (red triangle), SMAST large camera (blue cross), HabCam (green dots), and SMAST digital camera (light green x) for Georges Bank closed areas.




Figure 4.4. Comparison of CASA model estimated biomass with expanded estimates from the lined dredge (red triangle), SMAST large camera (blue cross), HabCam (green dots), and SMAST digital camera (light green x) for Georges Bank open areas.




Figure 4.5. Comparison of base case CASA model estimates of abundance (top), biomass (middle), and fishing mortality (bottom) to previous CASA model estimates for Georges Bank and Mid-Atlantic sea scallops.

REFERENCE POINTS

TOR #5: Update or redefine Status Determination Criteria (SDC; point estimates or proxies for BMSY, BTHRESHOLD, FMY and MSY reference points) and provide estimates of those criteria and uncertainty, along with a description of the sources of uncertainty. If analytic model-based estimates are unavailable, consider recommending alternative measurable proxies for reference points. Compare estimates of current stock size and fishing mortality to existing, and any redefined, SDCs. Provide stock status based on updated reference points.

Introduction

Per recruit reference points were used as proxies for F_{MSY} and B_{MSY} in scallop assessments prior to 2010. The per recruit reference point F_{MAX}, the fully recruited fishing mortality rate that generates maximum yield-per-recruit, was used as a proxy for F_{MSY}. The biomass reference point was defined as the product of B_{MAX} (biomass per recruit at $F = F_{MAX}$) and median numbers of recruits. As selectivity has shifted to larger scallops, yield per recruit curves have become increasingly flat, particularly in the Mid-Atlantic, making per-recruit reference points unstable. Additionally, recruitment has been stronger during periods when biomass has been high, suggesting that spawner-recruit relationships should be included. Finally, risk-based reference points are needed to calculate Acceptable Catch Levels/Allowable Biological Catch (ACLs/ABCs) and target fishing mortalities.

The stochastic yield model (SYM) (Hart 2013) was developed to address these issues; it has been used to calculate reference points in scallop assessments since 2010 (NEFSC 2010). It uses Monte-Carlo simulations to propagate the uncertainty in per recruit and stock-recruit calculations while calculating yield curves, F_{MSY} and B_{MSY}. Separate SYM models were run; one each for the Mid-Atlantic and Georges Bank, and their results combined to obtain whole stock reference points. Because of its limited time series and minor contribution to landings and biomass, Gulf of Maine scallops were not included in these calculations. Georges Bank reference points are used as proxy reference points for Gulf of Maine scallops.

Methods

The SYM model combines per-recruit calculations with stock-recruit relationships to estimate yield curves, as discussed in Beverton and Holt (1957) and Shepherd (1982). The SYM approach treats both the per-recruit and the stock-recruit relationships as being uncertain and takes this uncertainty into account.

Although the SYM model is separate from CASA, efforts were made to make the two models as compatible as possible. In particular, growth was modeled using stochastic growth matrices based on the most recent growth period. However, because SYM (unlike

CASA) uses a stock-recruit relationship, the Georges Bank open and closed areas were combined, so that two SYM models were used, one for Mid-Atlantic and one for Georges Bank.

Uncertainties in the SYM model can be divided into uncertainty in the per recruit models, and that from the stock recruit relationship. Per recruit calculations depend on a number of parameters that each carry a level of uncertainty, including shell height/meat weight parameters, fishery selectivity, cull size and the fraction of discards that survive, the incidental mortality rate, and the natural mortality rate *M*. Each of these was modeled by specifying a distribution, together with parameters for that distribution, typically a mean and variance. A more detailed discussion of all these parameters can be found in Hart (2013). The form of the distributions used are the same as in Hart (2013), but point estimates of some of the parameters, such as natural mortality, selectivity and shell height/meat weight, have been updated as discussed previously.

Of all the input parameters to the per recruit modeling, by far the most uncertain is natural mortality. Natural mortality was modeled as an inverse gamma distribution (i.e., $1/\gamma$, where γ is a gamma distribution). This makes sense for several reasons. First, in many methods to estimate natural mortality, such as maximum longevity or "clapper" ratios for scallops, the uncertain quantity is in the denominator. Secondly, inverse gamma distributions are skewed to the right, which makes sense for natural mortality. For example, it is possible that the true natural mortality is twice or more than its point estimate, but it cannot be zero or less. For this assessment, the mean *M* was taken as 0.27 on Georges Bank, consistent with the mean natural mortality in this area used in the CASA models. The Mid-Atlantic model presents more complex issues because of its recent declines in productivity. To take this into account, the base Mid-Atlantic SYM model used the mean natural mortality at size from the CASA model in the last five years (2019−2023). This resulted in *M* values ranging from 0.36 for a 50 mm scallop to around 0.56 for large adults. Uncertainties in natural mortalities were set at $\sigma = 0.08$ for both regions.

Beverton-Holt stock-recruit curves were fitted to spawning stock biomass *B* (using meat weight) and recruitment estimates from base CASA model runs

$$R = \frac{sB}{h + B} \qquad (1)$$

assuming square-root-normal errors, where *s* is the expected asymptotic recruitment, and *h* is the spawning stock biomass where the expected recruitment is half its asymptotic value. Standard errors of the stock-recruit parameters and their correlation were estimated using the delta method.

For this and other recent assessments, CASA estimates recruits as one-year olds. These were advanced to age two using the CASA estimated Ms by size and year. The SYM model thus starts its per recruit calculations at 50 mm, approximately two years old, and the stock-recruit curve (Equation 1) was estimated in terms of two-year-old recruits. Meat weight was used as the primary surrogate for spawning stock biomass (SSB), although there was some exploration of using gonad weight instead of meat weight.

Recruitment in the Mid-Atlantic has been poor in recent years, likely due to shifts in environmental conditions (TOR 1). Actual recruitment during the last five years in this area has been on average 41% of that predicted by the stock-recruitment relationship. For that reason, recruitment in the base Mid-Atlantic SYM model was reduced by a factor of 0.41 from the prediction of the stock-recruit relationship; sensitivities were performed without this adjustment.

At each iteration of the simulation model, parameter values were drawn from their corresponding distributions, and per recruit and yield curves were calculated. This was repeated 100,000 times and the results of each iteration were stored. The stock-recruit parameters were simulated as correlated square-root normals.

For each run, equilibrium recruitment at fishing mortality F is given by:

$$R = s - h/b(F) \quad (2)$$

where $b(F)$ is SSB per recruit at fishing mortality F, and s and h are as in equation (1). Total yield $Y(F)$ is therefore:

$$Y(F) = y(F)R = y(F)[s - h/b(F)] \quad (3)$$

where $y(F)$ is yield per recruit at fishing mortality F.

Mean yield curves calculated by this method can be disproportionately influenced by outliers, both in cases where the population collapses at zero fishing and where predicted yields are unrealistically high (Hart 2013). For this reason, we used the 10% trimmed mean of the 100,000 runs to obtain point estimates of yield at each fishing mortality, where the highest and lowest 10% of the runs were removed. The probabilistic FMSY was taken as the fishing mortality that maximizes the trimmed mean yield curve; MSY and BMSY are the trimmed mean yield and SSB curves at FMSY. In some previous assessments, the median was used instead because of instability when using the trimmed mean. This was not observed in this assessment, and it was judged that the trimmed mean is a better measure of expected yield than the median.

Results

Distributions of MSY and BMSY are somewhat less variable but still have considerable uncertainty. The trimmed mean yield curve shows that Georges Bank is currently estimated to be much more productive and had a much lower FMSY than the Mid-Atlantic. The reference point estimates for the combined stock are FMSY = 0.49 and BMSY = 93282 mt meats (Table 5.1).

If no adjustments for recent conditions were made in the Mid-Atlantic (i.e. M = 0.4 and no adjustment for poor recent recruitment), MSY for the Mid-Atlantic would be around 25000 mt meats. Setting M to its estimated value in the last five years in the Mid-Atlantic but not adjusting recruitment reduces the Mid-Atlantic MSY to about 20000 mt meats. The working group judged that adjusting both M and recruitment to reflect recent conditions is more credible.

Comparing these reference points to CASA estimates gives stock status. Combined biomass in 2023 from the three CASA models is 69956 mt meats. This is below the target biomass of B_{MSY} = 93282 but above $B_{threshold}$ = 46641 mt meats, so the stock is not overfished. Combined fully recruited fishing mortality in 2023 was 0.33. This is below F_{MSY} = 0.49 so overfishing is not occurring. It should be noted, however, that fully recruited fishing mortality in both the Georges Bank Closed and Georges Bank Open area in 2023 was about 0.47, which is above the Georges Bank F_{MSY} = 0.36. Thus, if sea scallops were managed as two separate stocks (Mid-Atlantic and Georges Bank), overfishing would be occurring on Georges Bank.

Table 5.1. Summary of reference points and 2023 biomass and fishing mortality for Georges Bank, Mid-Atlantic and combined.

Region	MSY	*F*MSY	*B*MSY	*B*threshold	*B*2023	*F*2023
GB	22706	0.36	83414		49400	0.47
MA	7941	1.56	15909		20556	0.06
Combined	28402	0.49	93282	46641	69956	0.33




Figure 5.1. Trimmed mean yield curves for the Mid-Atlantic, Georges Bank and combined. The green vertical line indicates the combined FMSY = 0.49.




Figure 5.2. Stacked plot showing CASA estimated biomass over time for the Mid-Atlantic, Georges Bank Closed and Open, and Habcam estimates for the "Peter Pan" southeastern Nantucket Lightship area. The black dashed line indicates the target biomass B_{MSY} and the red dashed line the threshold biomass $B_{MSY}/2$. Note that these reference points are only valid for the last 5 years.




Figure 5.3. Estimated fishing mortality in Georges Bank Open, Closed and in the Mid-Atlantic, compared with *F* reference points for the Mid-Atlantic (red horizontal dashed line), Georges Bank (blue horizontal dashed line), and overall (black horizontal dashed line). Note that these reference points are only applicable for the last five years.

PROJECTIONS

TOR #6: Define and document methods for producing projections; provide justification for assumptions of fishery selectivity, fecundity, mortality and recruitment; comment on the reliability of resulting projections considering the effects of uncertainty and sensitivity to projection assumptions. Compare the results of SAMS and GeoSAMS and comment on their appropriateness for use in management.

Because of the sedentary nature of sea scallops, fishing mortality can vary considerably in space even in the absence of area specific management (Hart 2001). Rotational management and long-term closures exacerbate this heterogeneity. Projections that ignore spatial variation can be unrealistic and misleading. For example, suppose 80% of the stock biomass is in areas closed to fishing (as has occurred in some years in Georges Bank). A stock projection that ignored the closure and assumed an overall *F* of 0.2 would forecast landings nearly equal to the entire stock biomass in the areas open to fishing. Thus, using a non-spatial forecasting model could lead to unsustainable harvest levels under area

management. For these reasons, a spatial forecasting model (the Scallop Area Management Simulator, SAMS) was developed for use in sea scallop management. Various versions of SAMS have been used since 1999. In the current version of SAMS used here, the resource is divided into 24 separate subareas, 7 in the Mid-Atlantic, 12 on Georges Bank, and 5 in the Gulf of Maine.

Population dynamics is modeled similarly in SAMS to CASA and SYM, that is, populations are size-structured and use stochastic growth transition matrices. However, processes can be modeled at a finer spatial scale in SAMS compared to the other models. Specifically, each area has its own stochastic growth transition matrix derived from the shell increments collected in that area from the most recent growth period. Similarly, each SAMS area has its own shell height to meat weight relationship, based on data from that area. Natural and fishing mortality and recruitment are also area specific. Fishing mortality in a subarea can either be explicitly specified in each area, as would be the case for closed areas or "access areas", which are managed by an area-specific quota, or calculated using a simple fleet dynamics model which assumes fishing effort is proportional to estimated LPUE. Adult M is based on that used for CASA in the most recent period, except in the two southernmost areas in the Mid-Atlantic, Virginia and Delmarva, which are set at 4 and 0.6, respectively.

Projected recruitment is modeled stochastically with the log-transformed mean and covariance for recruitment in each area matching that observed in NEFSC dredge survey time series. Mean recruitment is then scaled to a region-wide Beverton-Holt stock recruitment relationship, making the SAMS model more comparable to the SYM reference points model. Thus, the recruitment is multiplied by $s_r(B_r)/s_r(B_{\text{mean}})$, where s_r is the (mean) regional stock recruit relationship (for two years old) used in the SYM model, B_r is the current regional biomass, and B_{mean} is the mean overall biomass.

Initial conditions are based on surveys, which are bootstrapped parametrically. The model is run 1000 times, with different initial conditions, and recruitment. Natural mortality is also varied among the runs, following its distribution in the SYM models.

Because the "open" areas are managed through days at sea, LPUE needs to be estimated in these areas. This is done using a linear regression between mean exploitable biomass in the open areas and observed LPUE in these areas. The SAMS model gives projections of mean open area exploitable biomass, which then are translated into projected LPUE using the regression.

Mid-Atlantic projected biomass has been consistently overestimated. These projections assumed $M = 0.25$ for the Mid-Atlantic (except the two most southern areas). This is further evidence that M has been underestimated in this area. Additionally, recruitment in the Mid-Atlantic has been well below average. In future years, Mid-Atlantic

recruitment will be reduced, consistent with the SYM model. Georges Bank projections showed a much smaller overestimation. The increase in *M* from 0.2 to 0.27 will reduce this bias. Additionally, recruitment on Georges Bank during these years has been somewhat below average, although there is no evidence of an environmental shift as was observed in the Mid-Atlantic.

GeoSAMS

While SAMS was an advancement over non-spatial models, it is still spatially crude compared to the scale that the fishery operates. For example, SAMS assumes that effort and fishing mortality are spatially uniform inside each SAMS area, whereas in reality, fishing mortality can be highly heterogeneous because scallopere are attracted to areas with highest catch rates (Hart 2001). In addition, it is difficult to reconfigure SAMS to a new spatial configuration; a more flexible approach was desired.

To resolve these issues, work began on a GeoSAMS, a more spatially explicit version of the SAMS model. In GeoSAMS, population dynamics occurs at each survey location in the initial year, using methods very similar to the SAMS model. After each model year, the estimates at the survey points are interpolated to a fine scale grid, using a regression kriging methodology similar to that used for the Habcam survey. This allows for estimates of biomass and landings from any desired area. Unfortunately, funding for the GeoSAMS project was halted in early 2025, when it was nearing but not quite operational status.

RESEARCH RECOMMENDATIONS

TOR #7: Review, evaluate, and report on the status of research recommendations from the last assessment peer review, including recommendations provided by the prior assessment working group, peer review panel, and SSC. Identify new recommendations for future research, data collection, and assessment methodology. If any ecosystem influences from Term of Reference 1 could not be considered quantitatively under that or other Terms of Reference, describe next steps for development, testing and review of quantitative relationships and how they could best inform assessments. Prioritize research recommendations.

The following summarizes the working group's response to previous research recommendations. Bulleted items are the research recommendations, and the sub-bullets are the working group's responses. Additional discussion on methods for including ecosystem influences and new research recommendations are provided in the TOR7 Working Paper.

Surveys

- Further investigate methods for better survey coordination between the various survey programs, including survey design, timing, and standardized data formatting for easier sharing. (2018 Benchmark Assessment)
    - The New England Fisheries Management Council (NEFMC) and Northeast Fisheries Science Center (NEFSC) formed a Scallop Survey Working Group (SSWG) to address these topics. The SSWG developed recommendations to address four Terms of Reference, including survey spatial coverage, sampling intensity and frequency, data standardization, storage, and access, potential impacts from the development of offshore wind, and data needs to support future stock assessments.
- Investigate changes in dredge efficiency and saturation due to high scallop densities or high bycatch rates. (2018 Benchmark Assessment)
    - Ongoing research at the Virginia Institute of Marine Science. Not completed.
- Improve training of annotators used in optical surveys and develop standardized QA/QC procedures for data collected from imagery. (2018 Benchmark Assessment)
    - Standardized QA/QC procedures have been developed for NEFSC Habcam, along with a detailed training and annotation manual that were implemented in 2022. SMAST Drop Camera program also uses standardized training sets, and a QA/QC process. Substantial progress has been made.
- Investigate the use of software for automated annotation of imagery from optical surveys. (2018 Benchmark Assessment)
    - Work is underway at NEFSC. AI algorithms continue to improve rapidly. NEFSC staff have submitted a publication in this area (focused on sand dollars, Vanaki et al. 2025), although staff capacity has been a challenge to achieve full implementation.
    - Other survey groups that work with optical tools (SMAST Drop Cam and Coonamessett Farm Foundation Habcam) have ongoing projects in this area.
    - A similar recommendation is made by the working group for future research recommendations.
- Consideration of surveys in the GOM region be included in the ongoing Scallop Survey Working Group (SSWG) and NEFSC-supported scallop survey redesign efforts. (2021 SSC Report)
    - There are two recommendations from the SSWG report that are particularly relevant to surveys of the Gulf of Maine region. First, the Northern Gulf of

Maine management area and Gulf of Maine resource area should be included in regular survey coverage. Second, the effort should be made to match appropriate sampling tools, designs, and methods with specific conditions of survey areas. The Gulf of Maine is heavily fished using fixed gear (mainly lobster pots with vertical lines), which presents challenges for sampling with mobile gear (either dredge, or a towed camera system). Of the two mobile sampling techniques, dredge sampling is generally towed over shorter distances than Habcam, which runs transects over larger areas on Georges Bank and the Mid-Atlantic. There is no dedicated federal funding for surveys of the Gulf of Maine. However, surveys have been regularly funded though the Scallop Research Set-Aside Program. The results of these surveys are presented in the Gulf of Maine appendix. The SSWG final report is available at this page: https://www.nefmc.org/committees/scallop-survey-working-group

  - A sub-group of the SSC reviewed and recommended the use of Generalized Random Tessellation Stratified (GRTS) method for survey design for Georges Bank and the Mid-Atlantic. The sub-group report is available at this page: https://www.nefmc.org/library/september-2024-ssc-report. Extension of this effort to the Gulf of Maine is planned.

- Continued survey data collection and analyses of scallop populations in the Gulf of Maine to support the future development of region-specific reference points and growth estimates (2018 Benchmark).
  - The Scallop RSA continues to support annual surveys in the Gulf of Maine. A working paper was prepared as part of this assessment to summarize the state of survey information. Gulf of Maine specific reference points will require more years of data.

Gonad-based estimates of SSB and reference points:

- Further work to develop gonad-based estimates of SSB and reference points. (2020 management track assessment)
  - Gonad-based estimates were explored in this research track. Ultimately, the working group did not recommend transitioning to this approach because the gonad data were very noisy, particularly in the Mid-Atlantic with its strong spring spawn. While in principle gonad-based estimates are better reflective of SSB, more research is required because of the data issues.

Scallop Forecasting Model (Scallop Area Management Simulator or SAMS), LPUE, and VMS data:

- Further refine and test methods for forecasting LPUE. (2018 Benchmark Assessment)
    - Limited progress, and further review of LPUE models is needed. Some development of spatial choice models for scallop fishermen, which have a role in LPUE forecasts. There have also been some changes to the LPUE model in annual management actions.

- Develop a spatially explicit methodology for forecasting the abundance and distribution of sea scallops by incorporating spatial data from surveys, landings, and fleet effort (aka GEOSAMS). (2018 Benchmark Assessment)
    - Progress has been made. The working group received a presentation on the development of GEOSAMS in January. Funding for this work was halted due to a budget cut that supported a part-time programmer.

- Revive and streamline previously developed methods for interpreting VMS data. (2018 Benchmark Assessment)
    - Limited progress. The Scallop Plan Development has developed a standard way to categorize fishing activity using speed filters to mask steaming and other non-fishing activities. There was discussion regarding including depth in the filtering process, and the working group noted that there may be more sophisticated methods to explore, such as Hidden Markov models.

- The SAMS model seems to be having some difficulty capturing some of the recent stock changes. Recommendation to investigate new treatments for recruitment assumptions in the SAMS model. A revamp of the SAMS model to allow for more spatial estimation would be another fruitful area to explore. The SSC recommends a review of the SAMS model in the next management track assessment and supports NEFSC's development of a geostatistical SAMS model for the 2024 research track assessment. (2020 SSC Report)
    - The primary performance problems in SAMS has been overestimation of abundance and biomass, particularly in the Mid-Atlantic, due to the decline in productivity of this region. Estimates of *M* have increased and recruitment decreased in the Mid-Atlantic in this assessment which should eliminate or reduce the overestimation issue with the SAMS model.

- The working group discussed the practical importance of tracking and reviewing the performance of forecasts for individual areas, since these are used in making fishery allocations.
- As discussed above, work had begun developing the GEOSAMS model but are on pause due to budget cuts.

- The SSC recommends that ongoing research on potential drivers of changes in sea scallop stock dynamics (e.g., changing ocean conditions, including ocean acidification and warming) be included in the upcoming review of the SAMS model and in the 2024 research track assessment for scallops. (2021 SSC Report)
  - New sources of information include the working paper on thermal conditions (TOR 1), and revised CASA models (TOR 4).
  - There is ongoing work at the NEFSC's Milford Lab on ocean acidification.
  - Future work is needed to incorporate this work into SAMS and reviewing the results of the forecasts.

- The SSC recommends continued evaluation of the performance of the projection model and the need for a more holistic evaluation of changes in stock dynamics, a synoptic evaluation of potential drivers (e.g., changing ocean conditions, including ocean warming), and revision of model assumptions to account for these changes. (2022 SSC Report)
  - See responses regarding the SAMS model above.
  - Some annual ad-hoc increases in *M* were made for management purposes in estimation areas at the southern end of the range (Virginia Beach, Delmarva). A retrospective-type analysis on historical area-specific SAMS projections is recommended; a quantitative measure of past bias could be used as a tool to inform future projections. (2024 SSC Report)
  - Summary of SAMS performance over time is shown in Figure 4.2 in TOR 6. Future work can focus on plots by sub-areas, possibly later this year.

Environmental Drivers and Stock Dynamics:

- Continued investigation of discard mortality, particularly during warm water periods, by incorporating environmental data. (2018 Benchmark Assessment)
  - Thermal mortality has not yet been adequately addressed; the models still assume 20% discard mortality but this is probably too low in Mid-Atlantic because of high surface water temperatures during the summer. Discard mortality remains an uncertainty, and further work is needed.

- The SSC recommends investigation of the environmental drivers affecting the productivity of this stock, such as those influencing recruitment and natural mortality. (2022 SSC Report)
- Research on the implications of expected future increases in bottom temperature on natural mortality and how these impacts on scallop survival will likely continue to extend northward. The SSC suggests that a thorough examination of the impacts of ecosystem and climate change on population dynamics is critical during the ongoing Sea Scallop Research Track Stock Assessment. (2023 SSC Report)
- The SSC supports continued monitoring of changes in the dynamics of this stock (i.e., recruitment, growth, and natural mortality) and research to understand the role of environmental drivers as this represents the most pressing concern facing the future of this fishery. (2024 SSC Report)
    - These three recommendations are partially addressed in TOR 1; however, additional work is needed. The working group noted the importance of integrating changing environmental and oceanographic information into scallop science and management, and the need for cross-branch collaboration with the Ecosystem Dynamics and Assessment Branch (EDAB). Additional collaborative and trans-disciplinary research programs are needed to advance this work.

Stock Assessment Model, Aging, Growth, Selectivity:

- Analyze past juvenile scallop mortality events and develop better methods to model time-varying mortality in the assessment models. (2018 Benchmark Assessment)
    - There has been continuing work on this topic (e.g., Hart and Chang 2022) that was presented in this research track assessment (TOR 4). The CASA models include age/time varying estimates of natural mortality.
- Continue development of scallop ageing methods and examination of scallop growth processes including density dependent effects. (2018 Benchmark Assessment)
    - Work examining density-dependent reproductive output was recently published in a peer-reviewed journal (Kowaleski et al. 2024).
    - More work should be conducted to examine the estimation methods of growth transition matrices, enabling these matrices to better capture spatial and temporal changes in growth so that these changes can be incorporated into CASA models.

- Investigate methods to better estimating biomass and abundance variances from Habcam optical surveys including development of Bayesian geostatistical methods. (2018 Benchmark Assessment)
    - A peer-reviewed paper was published on this topic (Duskey et al. 2023), but it is not ready for full implementation at this time.

- Investigate and estimate current and historical unreported landings and effects of spatially heterogeneous fishing mortality on mortality estimates. (2018 Benchmark Assessment)
    - No recent progress on this topic. Unreported fishing mortality early in the assessment time series is suspected, although there are no obvious solutions to address the challenge. Hart (2001, 2003) presents earlier peer-reviewed work on the effects of spatially heterogeneous fishing mortality.

- Investigate and parameterize sub-lethal effects of disease, parasites, or discarding on mortality, growth, and landings. (2018 Benchmark Assessment)
    - There is ongoing work at the Virginia Institute of Marine Science looking at shell disease and nematodes; see Rudders et. al. (2019). An Investigation into the Scallop Parasite Outbreak on the Mid-Atlantic Shelf: Transmission Pathways, Spatio-Temporal Variation of Infection and Consequences to Marketability. RSA Award Number: NA16NMF4540043. VIMS Marine Resource Report No. 2019–02.
    - Multiple survey groups as well as at-sea observers are tracking scallop condition.

- Continue improvements of observer recordings for vessel fishing behavior, including deck loading and shucking dynamics in responses to disease or poor scallop health. (2018 Benchmark Assessment)
    - Observer protocols were modified a few years ago to better track scallop health and meat condition. If gray meats or parasites are present, observers must resample meat weight at least twice per watch and weigh affected meats separately from clean meats.
    - Observers not currently checking for shell blister disease.

- Continue investigating the extent of incidental fishing mortality, particularly on hard bottom habitats. (2018 Benchmark Assessment)
    - A recently published paper examined these issues (Ferraro et al. 2017). However, this work did not focus on hard bottom habitats, so some uncertainty remains.

- Continued development of the SYM model that models selectivity dynamically as a function of full recruitment fishing mortality. (2020 Management Track)
    - Not complete; work is still underway at the NEFSC.

- Transitioning the CASA model into the "next generation" of assessments would also be a natural progression; a state-space model with parameters that can be connected to environmental variables could result in more accurate estimates of biomass and fishing mortality rate and reference points with an improved understanding of uncertainty and the relationship between scallops and the ecosystem. (2024 SSC Report)
    - This recommendation was made while the research track process was ongoing. The working group recommended updating the CASA models for this research track at the start of this process in 2023. This SSC recommendation is carried forward in future research recommendations below, and in the working group's response to the TOR 7 on quantitative assessment of ecosystem influences. The working group notes that CASA is attempting to capture process error in time-varying natural mortality.
    - DFO Canada is using a spatial delay-difference state-space model to assess the Atlantic sea scallop resource in some areas (McDonald et al. 2021).

BACKUP ASSESSMENT APPROACH

TOR #8: Develop a backup assessment approach to providing scientific advice to managers if the proposed assessment approach does not pass peer review or the approved approach is rejected in a future management track assessment.

Introduction

The WG identified the parts of the process where a potential failure to achieve an acceptable result might occur (Working Paper TOR8, Figure 1).

1. If the Catch at Size Analysis model (CASA) fails to perform for one or more of the three management areas. The CASA model estimates the number of fully recruited scallops in the population for each management region (Mid-Atlantic, Georges Bank closed, and Georges Bank open).
2. If the Stochastic Yield Model (SYM) fails to perform for either or both two modeled regions (Mid-Atlantic or the combined Georges Bank area). The SYM combines the Mid-Atlantic and Georges Bank results to derive a stockwide result. Using a Monte Carlo simulation, candidate reference points are provided, including recommendations for biological reference points of F_{MSY}, B_{MSY}, and MSY. SYM accounts for uncertainty in the CASA model estimates and addresses the additional

issues of unaccounted mortality during the fishing process (e.g. bycatch mortality of discards and incidental mortality caused by gear on bottom). The FMSY and BMSY from the SYM model are used in status determination.

We note that a failure in any one of the three components of Part 1, or any of the two components of Part 2 will affect the calculation of biological reference points that are used in conjunction with the New England Fishery Management Council's harvest control rule to develop Over Fishing Limit (OFL) and Acceptable Biological Catch (ABC) estimates. Broadly speaking, any failure would require a backup plan, but the approach to remedy the situation would vary for each part of the process.

Methods

CASA Backup

The scallop dredge surveys are designed in such a way that the number of scallops observed by area of bottom covered by a tow and corrected for catchability can provide a measure of average scallop density by area (Miller et al. 2019). This density by area can be expanded as a stratified mean to get the number of scallops for each region of interest. These population estimates can be fed into the SYM to calculate reference points. Similar calculations can be made using optical survey methods (See TOR 3 working paper).

SYM Backup

SYM was outlined in (Hart 2013) and has not, in substance, changed since its use in 2014 or during the 2018 Stock Assessment Workshop (65th SAW), when it was approved for use except for changes in parameters. The research track working group is recommending changes to M and recruitment in SYM in this assessment.

Reference points could be calculated deterministically using means or medians or trimmed means of the various CASA output parameters rather than through stochastic simulation. However, stochastic estimates of F_{MSY} are lower than deterministic estimates because only the stochastic reference points take into account the risk that F_{MSY} has been overestimated due to parameter misestimation. So conservative use of such reference points is recommended.

Model Free Backup

Should the failure of approval for any or all the model steps in Parts 1 and 2 occur in such a way that status determination remains unknown there are other options that can be considered. For example, the surveys could still be used to determine standing stock biomass. In this scenario, a conservative determination of fishing mortality would be determined by using the realized catch over the last five years along with an effort

determination from the LPUE to come up with an appropriate Acceptable Biological Catch (ABC) for the fishery. However, under such a scenario the productivity of the stock may remain unknown and consequently certain status determination metrics such as the Overfishing Limit (OFL) would remain unavailable.

Discussion

These backup plans are only for consideration when the main assessment approach, CASA, is unable to converge in any formulation, is not completed, or is rejected during peer review of the research track or management track process. The WG has provided these backup plans taking into consideration possible reasons why this might happen. There are also some reasons this backup approach might not work.

As with any other assessment, a more practical concern is that the approaches for all three assessment regions rely on the scallop dredge surveys and possibly the optical surveys. If these surveys are unable to be completed successfully in each region, the approaches could fail. Missing surveys are a different problem from zero captures during a survey. Obviously, as the amount of zero captures increases, so does the risk to the stock. Methods exist to fill in the gaps under certain sets of assumptions regarding total coverage by the existing survey methods.

Conclusions and Recommendation

The WG recommends that if a backup plan is needed, the adjusted area survey measures (e.g. dredge, drop camera, Habcam) of fully recruited scallop density expanded by survey strata and then summing over strata can be used to arrive at an estimate of scallop abundance. These abundance estimates can be combined with similar estimates for the other management region or with the CASA estimates for that region to be processed by the SYM to arrive at the appropriate reference point recommendations. If the SYM should fail, then deterministic calculations using means, medians or trimmed means should replace the stochastic calculations. Should model steps prove to be inadequate, then a model free approach should be used to determine an interim ABC.

ADDITIONAL INVESTIGATIONS

The WG provides additional discussions on the scallop resource in the Gulf of Maine and results from the scallop community engagement meeting The WG also prepared a working paper on historical and current management actions. These additional investigations were incorrectly labeled as TOR 9 in working papers and are not an official TOR of the research track assessment.

Gulf of Maine Scallop Resource

While the majority of the Atlantic sea scallop resource in the U.S. is found on Georges Bank and in the Mid-Atlantic, sea scallops also occur in the Gulf of Maine (GOM) in both state and federal waters. The federal scallop resource in the GOM is managed by the New England Fishery Management Council and NOAA Fisheries and had supported a growing directed fishery in recent years.

There have been several changes to the management of the scallop resource in the GOM since the last sea scallop benchmark assessment (2018) and management track assessment (2020). This working paper builds on earlier efforts to catalogue fishery dependent and independent data in the Gulf of Maine region that were presented in the previous two benchmark assessments (NEFSC 2014, 2018). Over the long-term, the working group recommends that the scallop population in the Gulf of Maine be included in stock assessment models and included in calculations for status determination.

Scallops in this region are currently not considered in the CASA (TOR 4) and SYM (TOR 5) models, and reference points for the area have not been developed. To develop catch advice for the directed fishery, the SAMS projection model is used with recent year's survey data. When setting specifications for the overall fishery, scallops that are surveyed in the Gulf of Maine (inclusive of the NGOM management unit) are counted toward the legal limits (OFL and ABC).

In the absence of reference points and a stock assessment model for the Gulf of Maine, the OFL and ABC estimates for the Gulf of Maine have been derived using the Georges Bank reference points from the 2020 management track assessment ($F_{OFL}=F_{MSY}=0.46$, $F_{ABC}=0.32$); this assessment updates these estimates to $F_{OFL} = F_{MSY} = 0.36$ and $F_{ABC} = 0.29$. This approach was recommended by the NEFMC's SSC in October 2021. Catch limits for the NGOM management area are set using target fishing mortality rates below the ABC proxy, and by regulation must be between F=0.15 and F=0.25.

Scallop Community Engagement Meeting

The Atlantic Sea Scallop Research Track Working Group (Working Group) held a hybrid community engagement session on Wednesday, December 18, 2024 from 10:00 a.m. to 12:00 p.m. in New Bedford, MA and via webinar. The New England Fishery Management Council (Council) and NOAA Fisheries' Northeast Fisheries Science Center (NEFSC) jointly hosted the session. This meeting provided an opportunity for community members, including members of the scallop fishery, researchers, and the public, to engage in discussions about the current state of the scallop fishery and ongoing research efforts. Approximately 85 individuals attended, comprising Scallop Research Track Working Group members, NEFSC and NEFMC staff and council members, fishermen, industry representatives, and other community members.

Dr. Dvora Hart, lead stock assessment scientist for scallops, and Dr. Pat Sullivan, Chair of the Research Track Working Group, presented an overview of the research track assessment. They outlined the methodology for incorporating environmental and biological data into stock assessment models, including advancements in scientific techniques aimed at better understanding recruitment patterns, environmental influences, and predator

impacts on scallops. The presentations were followed by a structured discussion that encouraged community members to share their observations and feedback.

Dr. Hart, Dr. Sullivan and Mr. Jonathan Peros (NEFMC) fielded questions about the presentation. Members of the community raised several questions, particularly concerning the availability and use of environmental data. One participant inquired about the availability of data that may show warming temperatures, to which Dr. Hart responded that temperature and other environmental factors are measured regularly and made publicly accessible. She noted that there was unusually strong stratification in the Mid-Atlantic in 2024, with cooler bottom temperatures, despite warming long-term trends.

Questions about phytoplankton sampling were also addressed, with Dr. Hart explaining that these kinds of data are collected using various methods, including satellite imagery, physical sampling and from in-situ sensors, including a sensor on the Habcam V4 vehicle. Unrelated to the focus of this outreach meeting, an attendee questioned the rationale behind the Council's suggestion to add four additional days-at-sea (DAS), asking if this implied an increase in scallop abundance. It was clarified that the decision, made by the NEFMC, is multifaceted and not necessarily linked to a rise in scallop numbers.

A participant raised concerns about the models used for assessments, asking how much they rely on scientific versus experiential data. Dr. Hart answered that the models are primarily science-based, though there is some room for judgment in interpreting data. She acknowledged the importance of incorporating fishermen’s observations and supported the idea of smaller, informal meetings to facilitate communication between scientists and fishermen. This sentiment was echoed by another attendee, who expressed frustration about the perceived lack of responsiveness to fishermen’s input.

Dr. Sullivan and Dr. Hart reiterated that the New England Fishery Management Council (NEFMC) makes management decisions, and the research track focuses solely on the scientific basis for these policies. They encouraged participants to continue providing input to both the research track team and to managers such as the NEFMC.

REFERENCES

Almeida F, Valentine P, Reid R, Arlen L, Auster P, Cross J, Guida V, Lindholm J, Link J, Packer D (2005) Symposium Abstract: The Effectiveness of Marine Protected Areas on Fish and Benthic Fauna: The Georges Bank Closed Area II Example. In AFS Symp 41, p. 589. American Fisheries Society.

Almeida F, T Sheehan, Smolowitz R (1994) Atlantic sea scallop, *Placopecten magellanicus*, maturation on Georges Bank during 1993. NEFSC Ref Doc 94-13

Adebola T, Hart D, Chigbu P (2022). Bathymetric trends in the body size, and diet of *Astropecten americanus* in the northwest Atlantic Ocean. Estuarine, Coast Shelf Sci, 269:107814

Andersen S, Grefsrud ES, Harboe T (2013) Effect of increased pCO2 on early shell development in great scallop (*Pecten maximus* Lamarck) larvae. Biogeosciences Discuss, 10(2).

Barbeau MA, Scheibling RE, Hatcher BG (1998) Behavioural responses of predatory crabs and sea stars to varying density of juvenile sea scallops. Aquaculture, 169:87-98.

Barber BJ, Getchell R, Shumway S, Schick D (1988) Reduced fecundity in a deep-water population of the giant scallop *Placopecten magellanicus* in the Gulf of Maine, USA. Mar Ecol Prog Ser, 42:207-212.

Bethoney, ND, Stokesbury KD (2018) Methods for image-based surveys of benthic macroinvertebrates and their habitat exemplified by the drop camera survey for the Atlantic Sea Scallop. J Visualized Exp:JoVE, 137:57493.

Beverton RJH, Holt SJ (1957) On the Dynamics of Exploited Fish Populations. Chapman and Hall, London.

Caddy JF (1973) Underwater observations on tracks of dredges and trawls and some effects of dredging on a scallop ground. J Fish Res Bd Can, 30:173-180.

Chang, JH, Shank, BV, Hart, DR (2017). A comparison of methods to estimate abundance and biomass from belt transect surveys. Limn Oceanog:Methods, 15(5):480-494.

Chute, AS, Wainright, SC, Hart, DR (2012). Timing of shell ring formation and patterns of shell growth in the sea scallop *Placopecten magellanicus* based on stable oxygen isotopes. Journal of Shellfish Research, 31:649-662.

Claereboudt MR, Himmelman JH (1996) Recruitment, growth and production of giant scallops (*Placopecten magellanicus*) along an environmental gradient in Baie des Chaleurs, eastern Canada. Mar Biol, 124(4):661-670

Cochran WG (1977) Sampling Techniques, 3rd ed. Wiley & Sons, New York. 428p

Cooley SR, Rheuban JE, Hart DR, Luu V, Glover DM, Hare JA, Doney SC (2015) An integrated assessment model for helping the United States sea scallop (*Placopecten magellanicus*) fishery plan ahead for ocean acidification and warming. PLoS One, 10(5):e0124145

Dibacco C, Robert G, Grant J (1995) Reproductive cycle of the sea scallop, *Placopecten magellanicus* (Gmelin, 1791), on northeastern Georges Bank. J Shellfish Res, 14: 59-69

DuPaul WD, Kirkley JE, Schmitzer AC (1989) Evidence of a semiannual reproductive cycle for the sea scallop, *Placopecten magellanicus* (Gmelin, 1791), in the Mid-Atlantic region. J Shellfish Res 8:173-178

DuPaul WD, Fisher RA, Kirkley JE (1990) An evaluation of at-sea handling practices: Effects on sea scallop meat quality, volume and integrity. Contract report to Gulf and South Atlantic Fisheries

Duskey E, Hart D, Chang J-H, Sullivan, PJ (2023). Partitioning spatial dynamics in abundance of marine fisheries stocks between fine- and broad-scale variation: A Bayesian approach. Fish Res, 267:106816.

Diez ME Radashevsky VI, Orensanzm JM, Cremonte F (2011) Spionid polychaetes (Annelida: Spionidae) boring into shells of molluscs of commercial interest in northern Patagonia, Argentina. Ital J Zool, 78:497-504.

Elner RW, Jamieson GS (1979) Predation of sea scallops, *Placopecten magellanicus*, by the rock crab, *Cancer irroratus*, and the American lobster, *Homarus americanus*. J Fish Res Board Can, 36:537-543.

Ferraro DM, Trembanis AC, Miller DC, Rudders DB (2017) Estimates of sea scallop (*Placopecten magellanicus*) incidental mortality from photographic multiple before-After-Control-Impact Surveys. J Shellfish Res, 36(3):615-626

Franz DR, Worley EK (1982) Seasonal variability of prey in the stomachs of *Astropecten americanus* (Echinodermata Asteroidea) from off Southern New England, USA. Estuar Coast Shelf Sci, 14(4):355-368

Franz DR, Worley EK, Merrill AS (1981) Distribution patterns of common seastars of the middle Atlantic continental shelf of the northwest Atlantic (Gulf of Maine to Cape Hatteras). Biol Bull, 160(3):394-418.

Guderley H, Labbé-Giguere S, Janssoone X, Bourgeois M, Pérez HM, Tremblay I (2009) Thermal sensitivity of escape response performance by the scallop *Placopecten magellanicus*: impact of environmental history. J Exp Mar Bio Ecol, 377(2):113-119.

Handley SJ, Bergquist PR (1997) Spionid polychaete infestations of intertidal Pacific oysters Crassostrea gigas (Thunberg), Mahurangi Harbour, Northern New Zealand. Aquaculture, 153:191-205.

Hart DR (2001). Individual-based yield-per-recruit analysis, with an application to the Atlantic sea scallop, *Placopecten magellanicus*. Can J Fish Aquat Sci, 58(12):2351-2358.

Hart DR. (2003). Yield-and biomass-per-recruit analysis for rotational fisheries, with an application to the Atlantic sea scallop (*Placopecten magellanicus*). Fish Bull, 101(1):44-58.

Hart DR (2006). Effects of sea stars and crabs on sea scallop *Placopecten magellanicus* recruitment in the Mid-Atlantic Bight (USA). Mar Ecol Prog Ser, 306:209-221.

Hart DR (2013) Quantifying the tradeoff between precaution and yield in fishery reference points. ICES J Mar Sci, 70(3):591-603

Hart DR, Chute AS (2004) Essential fish habitat source document: Sea scallop, *Placopecten magellanicus*, life history and habitat characteristics, 2nd ed. NOAA Technical Memorandum NMFS NE-189.

Hart DR, Chute AS (2009a) Verification of Atlantic sea scallop, *Placopecten magellanicus*, shell growth rings by tracking cohorts in fishery closed areas. Can J Fish Aquat Sci, 66:751-758

Hart DR, Chute AS (2009b) Estimating von Bertalanffy growth parameters from growth increment data using a linear mixed-effects model, with an application to the sea scallop *Placopecten magellanicus*. ICES J Mar Sci, 66(10):2165-2175.

Hart DR, Chang JH (2022) Estimating natural mortality for Atlantic Sea scallops (*Placopecten magellenicus*) using a size-based stock assessment model. Fish Res, 254:106423.

Hart DR, Jacobson LD, Tang J (2013). To split or not to split: assessment of Georges Bank sea scallops in the presence of marine protected areas. Fish Res, 144:74-83.

Hart DR, Rago PJ (2006). Long-term dynamics of US Atlantic sea scallop *Placopecten magellanicus* populations. N Am J Fish Manage, 26(2):490-501.

Hart DR, Shank, BV (2011). Mortality of sea scallops *Placopecten magellanicus* in the Mid-Atlantic Bight: Comment on Stokesbury et al. (2011). Mar Ecol Prog Ser, 443: 293-297.

Hennen DR, Hart DR (2012) Shell height-to-weight relationships for Atlantic sea scallops (*Placopecten magellanicus*) in offshore US waters. J Shellfish Res, 31(4):1133-1144.

Inglis SD, Kristmundsson Á., Freeman MA, Levesque M, Stokesbury K. (2016). Gray meat in the Atlantic sea scallop, *Placopecten magellanicus*, and the identification of a known pathogenic scallop apicomplexan. J Invertebrate Path, 141: 66-75.

Jacobson, LD, Stokesbury KD, Allard MA, Chute A, Harris BP, Hart D, Jaffarian T, Marino, II MC, Nogueira JI and Rago P. (2010) Measurement errors in body size of sea scallops (*Placopecten magellanicus*) and their effect on stock assessment models. Fishery Bulletin, 108:233-248.

Jenkins SR, Brand AR (2001) The effect of dredge capture on the escape response of the great scallop, *Pecten maximus* (L.): implications for the survival of undersized discards. J Exp Mar Bio Ecol, 266(1):33-50.

Kaplan KA, Hart DR, Hopkins K, Gallagher S, York A, Taylor R, Sullivan PJ (2017) Evaluating the interaction of the invasive tunicate *Didemnum vexillum* with the Atlantic sea scallop *Placopecten magellanicus* on open and closed fishing grounds of Georges Bank. ICES J Mar Sci, 74(9):2470-2479

Kaplan KA, Hart DR, Hopkins K, Gallager S, York A, Taylor R, Sullivan PJ (2018) Invasive tunicate restructures invertebrate community on fishing grounds and a large protected area on Georges Bank. Biol Invasions, 20:87-103

Kent RML (1979) The influence of heavy infestations of Polydora ciliate on the flesh content of *Mytilus edulis*. J Mar Biol Assoc UK, 59:289-297.

Kowaleski KR, Roman SA, Mann R, Rudders DB. (2024). Extreme population densities reduce reproductive effort of Atlantic sea scallops in high-density recruitment events. Mar Ecol Prog Ser, 746:67-85.

Lagos NA, Benıtez S, Duarte C, Lardies MA and others (2023) Effects of temperature and ocean acidification on shell characteristics of *Argopecten purpuratus*: implications for scallop aquaculture in an upwelling-influenced area. Aquacult Environ Interact, 8:357-370.

Langton RW, Robinson WE, Schick D (1987). Fecundity and reproductive effort of sea scallops *Placopecten magellanicus* from the Gulf of Maine. Mar Ecol Prog Ser, 37:19-25.

Lauckner G (1983) Diseases of Mollusca: Bivalvia. In: Kinne O (Ed), Diseases of marine animals, Vol II, Introduction, Bivalvia to Scaphopoda. Biologische Anstalt Helgoland, Hamburg.

Levesque MM, Inglis SD, Shumway SE, Stokesbury KD (2016) Mortality assessment of Atlantic sea scallops (*Placopecten magellanicus*) from gray-meat disease. J Shellfish Res, 35(2):295-305.

Lester RJG, Blair D, Heald D (1980) Nematodes from scallops and turtles from Shark Bay, Western Australia. Mar Freshwater Res, 31: 713-717

Lichtenfels JR, Bier JW, Madden PA (1978) Larval anisakid (*Sulcascaris*) nematodes from Atlantic molluscs with marine turtles as definitive hosts. Trans Am Microscopical Soc, 97:199-207.

Lichtenfels JR, Sawyer TK, Miller GC (1980) New hosts for larval *Sulcascaris* sp.(Nematoda, Anisakidae) and prevalence in the calico scallop (*Argopecten gibbus*). Trans Am Microscopical Soc, 99:448-451.

O'Keefe C, and the NEFMC Scallop PDT (2022) Evaluation of the Atlantic Sea Scallop Rotational Management Program. Report to the New England Fishery Management Council.

O'Keefe C, Fuglebakk E, Holmes S, Tingley G (2023) Summary Report of the Catch Accounting and Management System (CAMS). Report to the Northeast Fisheries Science Center.

MacDonald BA, Thompson RJ (1985) Influence of temperature and food availability on the ecological energetics of the giant scallop *Placopecten magellanicus*. II. Reproductive output and total production. Mar Ecol Prog Ser, 25: 295-303

MacDonald BA, Thompson RJ (1986a) Influence of temperature and food availability on the ecological energetics of the giant scallop *Placopecten magellanicus*. Mar Biol, 93(1): 37-48

MacDonald BA, Thompson RJ (1986b) Production, dynamics and energy partitioning in two populations of the giant scallop *Placopecten magellanicus* (Gmelin). J Exp Mar Biol Ecol, 101: 285-299

MacDonald BA, Thompson RJ, Bayne BL (1987) Influence of temperature and food availability on the ecological energetics of the giant scallop *Placopecten magellanicus*. Oecologia, 72(4):550-556.

McDonald RR, Keith DM, Sameoto JA, Hutchings JA, Flemming JM. 2021. Explicit incorporation of spatial variability in a biomass dynamics assessment model, ICES J Mar Sci, 78:3265–3280.

Merrill AS, Posgay JA, Nichy F (1966) Annual marks on shell and ligament of sea scallop (*Placopecten magellanicus*). Fish Bull, 65:299-311

Miller TJ, Hart DR, Hopkins K, Vine NH, Taylor R, York AD, Gallager SM (2019) Estimation of the capture efficiency and abundance of Atlantic sea scallops (*Placopecten magellanicus*) from paired photographic–dredge tows using hierarchical models. Can J Fish Aquat Sci, 76(6):847-855

Morris Jr JA, Carman MR, Hoagland KE, Green-Beach ER, Karney RC (2009) Impact of the invasive colonial tunicate *Didemnum vexillum* on the recruitment of the bay scallop (*Argopecten irradians*) and implications for recruitment of the sea scallop (*Placopecten magellanicus*) on Georges Bank. Aquat Invasions, 4(1):207-211

Morris JA, Carman MR (2012) Fragment reattachment, reproductive status, and health indicators of the invasive colonial tunicate *Didemnum vexillum* with implications for dispersal. Biol Invasions, 14(10):2133-2140

Murawski SA, Brown R, Lai HL, Rago PJ, Hendrickson L. (2000) Large-scale closed areas as a fishery-management tool in temperate marine systems: the Georges Bank experience. Bull Mar Sci, 66(3):775-798

Murawski SA, Serchuk FM (1989) Environmental effects of offshore dredge fisheries for bivalves. ICES ICES CM 1989/K27. 12 pp.

Naidu KS (1987) Efficiency of meat recovery from Iceland scallops (*Chlamys islandica*) and sea scallops (*Placopecten magellanicus*) in the Canadian offshore fishery. J Northw Atl Fish Sci 7:131-136

Nadeau, M, Barbeau MA, Brêthes JC (2009). Behavioural mechanisms of sea stars (*Asterias vulgaris* Verrill and *Leptasterias polaris* Müller) and crabs (*Cancer irroratus* Say and *Hyas araneus* Linnaeus) preying on juvenile sea scallops (*Placopecten magellanicus* (Gmelin)), and procedural effects of scallop tethering. J Exp Mar Bio Ecol, 374(2):134-143

NEFMC (1993) Amendment #4 and supplemental environmental impact statement to the sea scallop fishery management plan. New England Fisheries Management Council, Saugus, MA

NEFMC (2003) Final Amendment 10 to the Atlantic sea scallop fishery management plan with a supplemental environmental impact statement, regulatory impact review, and regulatory flexibility analysis. New England Fisheries Management Council, Newburyport, MA

NEFSC (2007) 45th Northeast Regional Stock Assessment Workshop (45th SAW) Assessment Report. US Dept Commer, Northeast Fish Sci Cent Ref Doc 07-16, Woods Hole, MA

NEFSC (2010) 50th Northeast Regional Stock Assessment Workshop (50th SAW) Assessment Report. US Dept Commer, Northeast Fish Sci Cent Ref Doc 10-17, 844p, Woods Hole, MA

NEFSC (2014) 59th Northeast Regional Stock Assessment Workshop (59th SAW) Assessment Report. US Dept Commer, Northeast Fish Sci Cent Ref Doc 14-09, 782p, Woods Hole, MA

NEFSC (2018) 65th Northeast Regional Stock Assessment Workshop (65th SAW) Assessment Report. US Dept Commer, Northeast Fish Sci Cent Ref Doc 18-11, Woods Hole, MA, https://doi.org/10.25923/zapm-ga75

NEFSC (2020) Sea Scallop Management Track Assessment Fall 2020. https://doi.org/10.25923/6g5w-c636

Pousse E, Poach ME, Redman DH, Sennefelder G, Hubbard W, Osborne K, Munroe D, Hart D, Hennen D, Dixon MS, Li Y (2023) Juvenile Atlantic sea scallop, *Placopecten magellanicus*, energetic response to increased carbon dioxide and temperature changes. PLOS Climate, 2(2):e0000142

Rheuban JE, Doney SC, Cooley SR, Hart DR (2018). Projected impacts of future climate change, ocean acidification, and management on the US Atlantic sea scallop (*Placopecten magellanicus*) fishery. PloS one, 13(9):e0203536.

Rudders DB, Roman SA, Fisher R, McDowell J (2023). Observations on a reemerging epizootic of the sea scallop, *Placopecten magellanicus*, Resource. J Shellfish Res, 42(1):51-60.

Schalkhausser B, Bock C, Stemmer K, Brey T, Pörtner HO, Lannig G (2013) Impact of ocean acidification on escape performance of the king scallop, *Pecten maximus*, from Norway. Mar Bio, 160(8):1995-2006.

Shank BV, Hart DR, Friedland KD (2012). Post-settlement predation by sea stars and crabs on the sea scallop in the Mid-Atlantic Bight. Mar Ecol Prog Ser, 468:161-177.

Shumway SE, Selvin R, Schick DF (1987) Food resources related to habitat in the scallop *Placopecten magellanicus* (Gmelin, 1791): a qualitative study. J Shellfish Res, 6(2): 89-95

Siemann, LA, Garcia, LM, Huntsberger, CJ, Smolowitz, RJ. (2019). Investigating the Impact of Multiple Factors on Gray Meats in Atlantic Sea Scallops (*Placopecten magellanicus*). J Shellfish Res, 38(2):233-243.

Stevenson JA, Dickie LM (1954) Annual growth rings and rate of growth of the giant scallop *Placopecten magellanicus* (Gmelin) in the Digby Area of the Bay of Fundy. J Fish Res Board Can, 11:660-671

Shepherd JG (1982) A versatile new stock-recruitment relationship for fisheries, and the construction of sustainable yield curves. ICES J Mar Sci, 40(1): 67-75.

Stokesbury KDE, Harris BP, Marino MC, Nogueira JI (2004) Estimation of sea scallop abundance using a video survey in off-shore USA waters. J Shellfish Res, 23:33-44.

Sullivan PJ, Lai HL, Gallucci VF (1990) A catch-at-length analysis that incorporates a stochastic model of growth. Can J Fish Aquat Sci, 47:184-198.

Talmage SC, Gobler CJ (2009) The effects of elevated carbon dioxide concentrations on the metamorphosis, size, and survival of larval hard clams (*Mercenaria mercenaria*), bay scallops (*Argopecten irradians*), and Eastern oysters (*Crassostrea virginica*). Limnol Oceanogr, 54(6):2072-2080

Thompson KJ, Inglis SD, Stokesbury KDE (2014) Identifying spawning events of the sea scallop *Placopecten magellanicus* on Georges Bank. J Shellfish Res, 33(1):77-87

Vanaki S, Hart DR, Chang JH (2025) How many dollars are in the sea? Estimating sand dollar (*Echinarachnius parma*) abundance using an iteratively trained convolutional neural network and generalized additive models. Ecol Informatics, 90:103311.

White MM, McCorkle DC, Mullineaux LS, Cohen AL (2013) Early exposure of bay scallops (*Argopecten irradians*) to high CO2 causes a decrease in larval shell growth. PloS One, 8(4):e61065

Wigley SE, Hersey P, Palmer JE (2008) A description of the allocation procedure applied to the 1994 to 2007 commercial landings data. NEFSC Ref Doc 08-18

Wong MC, Barbeau MA (2005) Prey selection and the functional response of sea stars (*Asterias vulgaris* Verrill) and rock crabs (*Cancer irroratus* Say) preying on juvenile sea scallops (*Placopecten magellanicus* (Gmelin)), and blue mussels (*Mytilus edulis* Linnaeus). J Exp Mar Biol Ecol, 327:1-21

Wong MC, Barbeau MA, Hennigar AW, Robinson SMC (2005) Protective refuges for seeded juvenile scallops (*Placopecten magellanicus*) from sea star (*Asterias* spp.) and crab (*Cancer irroratus* and *Carcinus maenas*) predation. Can J Fish Aquat Sci, 62:1766-1781

Zang Z, Ji R, Hart DR, Chen C, Zhao L, Davis CS (2022). Modeling Atlantic sea scallop (*Placopecten magellanicus*) scope for growth on the Northeast US Shelf. Fish Oceanography, 31(3):271-290.

List of Working Papers

- Sea Scallop Life History Parameters
- TOR1a Ecosystem Influences
- TOR1b Climate Influences
- TOR2 Landings, Fishing Effort, Discards
- TOR3 Survey Data
- TOR4a Fishing Mortality, Recruitment, and Stock Biomass
- TOR4b CASA Technical Description
- TOR5 Reference Points
- TOR6 Projections
- TOR7 Research Recommendations
- TOR8 Plan B Backup Approach
- TOR 9a Gulf of Maine Scallop Resource
- TOR 9b Scallop Community Engagement Meeting
- TOR 9c Management History

Procedures for Issuing Manuscripts in the Northeast Fisheries Science Center Reference Document (CRD) and the Technical Memorandum (TM) Series

The mission of NOAA's National Marine Fisheries Service (NMFS) is "stewardship of the nation's ocean resources and their habitat." As the research arm of the NMFS's Greater Atlantic Region, the Northeast Fisheries Science Center (NEFSC) supports the NMFS's mission by "conducting ecosystem-based research and assessments of living marine resources, with a focus on the Northeast Shelf, to promote the recovery and long-term sustainability of these resources and to generate social and economic opportunities and benefits from their use." Results of NEFSC research are largely reported in primary scientific media (e.g., anonymously peer-reviewed scientific journals). However, to assist itself in providing data, information, and advice to its constituents, the NEFSC occasionally releases its results in its own series.

NOAA Technical Memorandum NMFS-NE – This series is issued irregularly. The series typically includes: data reports of long-term field or lab studies of important species or habitats; synthesis reports for important species or habitats; annual reports of overall assessment or monitoring programs; manuals describing program-wide surveying or experimental techniques; literature surveys of important species or habitat topics; proceedings and collected papers of scientific meetings; and indexed and/or annotated bibliographies. All issues receive internal scientific review, and most issues receive technical and copy editing.

Northeast Fisheries Science Center Reference Document – This series is issued irregularly. The series typically includes: data reports on field and lab studies; progress reports on experiments, monitoring, and assessments; background papers for, collected abstracts of, and/or summary reports of scientific meetings; and simple bibliographies. Issues receive internal scientific review, and most issues receive copy editing.

CLEARANCE

All manuscripts submitted for issuance as CRDs must have cleared the NEFSC's manuscript/abstract/webpage review process. If your manuscript includes material from another work which has been copyrighted, you will need to work with the NEFSC's Editorial Office to arrange for permission to use that material by securing release signatures on the "NEFSC Use-of-Copyrighted-Work Permission Form."

STYLE

The CRD series is obligated to conform to the style contained in the current edition of the United States Government Printing Office Style Manual; however, that style manual is silent on many aspects of scientific manuscripts. The CRD series relies more on the CSE Style Manual. Manuscripts should be prepared to conform to both style manuals.

The CRD series uses the Integrated Taxonomic Information System, the American Fisheries Society's guides, and the Society for Marine Mammalogy's guide for verifying scientific species names.

For in-text citations, use the name-date system. A special effort should be made to ensure all

necessary bibliographic information is included in the list of references cited. Personal communications must include the date, full name, and full mailing address of the contact.

PREPARATION

Once your document has cleared the review process, the Editorial Office will contact you with publication needs—for example, revised text (if necessary) and separate digital figures and tables if they are embedded in the document. Materials may be submitted to the Editorial Office as email attachments or intranet downloads. Text files should be in Microsoft Word, tables may be in Word or Excel, and graphics files may be in a variety of formats (JPG, GIF, Excel, PowerPoint, etc.).

PRODUCTION AND DISTRIBUTION

The Editorial Office will perform a copy edit of the document and may request further revisions. The Editorial Office will develop the inside and outside front covers, the inside and outside back covers, and the title and bibliographic control pages of the document.

Once the CRD is ready, the Editorial Office will contact you to review it and submit corrections or changes before the document is posted online. Several organizations and individuals in the Northeast Region will be notified by e-mail of the availability of the document online.