



NOAA Technical Report NMFS SSRF-783

Biomass and Density of Macrobenthic Invertebrates on the U. S. Continental Shelf off Martha's Vineyard, Mass., in Relation to Environmental Factors

Don Maurer and Roland L. Wigley

July 1984

Old Woman Creek Estuary
ODNR - Div. of Natural Areas
2514 Cleveland Rd., E.
Huron, Ohio 44839
(419) 433-4601

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service

NOAA TECHNICAL REPORTS

National Marine Fisheries Service, Special Scientific Report—Fisheries


The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for optimum use of the resources. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, development and enforcement of domestic fisheries regulations, surveillance of foreign fishing off United States coastal waters, and the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing service and economic analysis programs, and mortgage insurance and vessel construction subsidies. It collects, analyzes, and publishes statistics on various phases of the industry.

The Special Scientific Report—Fisheries series was established in 1949. The series carries reports on scientific investigations that document long-term continuing programs of NMFS, or intensive scientific reports on studies of restricted scope. The reports may deal with applied fishery problems. The series is also used as a medium for the publication of bibliographies of a specialized scientific nature.

NOAA Technical Reports NMFS SSRF are available free in limited numbers to governmental agencies, both Federal and State. They are also available in exchange for other scientific and technical publications in the marine sciences. Individual copies may be obtained from D822, User Services Branch, Environmental Science Information Center, NOAA, Rockville, MD 20852. Recent SSRF's are:

722. Gulf menhaden, *Brevoortia patronus*, purse seine fishery: Catch, fishing activity, and age and size composition, 1964-73. By William R. Nicholson. March 1978, iii + 8 p., 1 fig., 12 tables.
723. Ichthyoplankton composition and plankton volumes from inland coastal waters of southeastern Alaska, April-November 1972. By Chester R. Mattson and Bruce L. Wing. April 1978, iii + 11 p., 1 fig., 4 tables.
724. Estimated average daily instantaneous numbers of recreational and commercial fishermen and boaters in the St. Andrew Bay system, Florida, and adjacent coastal waters, 1973. By Doyle F. Sutherland. May 1978, iv + 23 p., 31 figs. 11 tables.
725. Seasonal bottom-water temperature trends in the Gulf of Maine and on Georges Bank, 1963-75. By Clarence W. Davis. May 1978, iv + 17 p., 22 figs., 5 tables.
726. The Gulf of Maine temperature structure between Bar Harbor, Maine, and Yarmouth, Nova Scotia, June 1975-November 1976. By Robert J. Pawlowski. December 1978, iii + 10 p., 14 figs., 1 table.
727. Expendable bathythermograph observations from the NMFS/MARAD Ship of Opportunity Program for 1975. By Steven K. Cook, Barclay P. Collins, and Christine S. Carty. January 1979, iv + 93 p., 2 figs., 13 tables, 54 app. figs.
728. Vertical sections of semimonthly mean temperature on the San Francisco-Honolulu route: From expendable bathythermograph observations, June 1966-December 1974. By J. F. T. Saur, L. E. Eber, D. R. McLain, and C. E. Dorman. January 1979, iii + 35 p., 4 figs., 1 table.
729. References for the identification of marine invertebrates on the southern Atlantic coast of the United States. By Richard E. Dowds. April 1979, iv + 37 p.
730. Surface circulation in the northwest Gulf of Mexico as deduced from drift bottles. By Robert F. Temple and John A. Martin. May 1979, iii + 13 p., 8 figs., 4 tables.
731. Annotated bibliography and subject index on the shortnose sturgeon, *Acipenser brevirostrum*. By James G. Hoff. April 1979, iii + 16 p.
732. Assessment of the Northwest Atlantic mackerel, *Scomber scombrus*, stock. By Emory D. Anderson. April 1979, iv + 13 p., 9 figs., 15 tables.
733. Possible management procedures for increasing production of sockeye salmon smolts in the Naknek River system, Bristol Bay, Alaska. By Robert J. Ellis and William J. McNeil. April 1979, iii + 9 p., 4 figs., 11 tables.
734. Escape of king crab, *Paralithodes camtschatica*, from derelict pots. By William L. High and Donald D. Worlund. May 1979, iii + 11 p., 5 figs., 6 tables.
735. History of the fishery and summary statistics of the sockeye salmon, *Oncorhynchus nerka*, runs to the Chignik Lakes, Alaska, 1888-1956. By Michael L. Dahlberg. August 1979, iv + 16 p., 15 figs., 11 tables.
736. A historical and descriptive account of Pacific coast anadromous salmonid rearing facilities and a summary of their releases by region, 1960-76. By Roy J. Wahle and Robert Z. Smith. September 1979, iv + 40 p., 15 figs., 25 tables.
737. Movements of pelagic dolphins (*Stenella* spp.) in the eastern tropical Pacific as indicated by results of tagging, with summary of tagging operations, 1969-76. By W. F. Perrin, W. E. Evans, and D. B. Holts. September 1979, iii + 14 p., 9 figs., 8 tables.
738. Environmental baselines in Long Island Sound, 1972-73. By R. N. Reid, A. B. Frame, and A. F. Draxler. December 1979, iv + 31 p., 40 figs., 6 tables.
739. Bottom-water temperature trends in the Middle Atlantic Bight during spring and autumn, 1964-76. By Clarence W. Davis. December 1972, iii + 13 p., 10 figs., 9 tables.
740. Food of fifteen northwest Atlantic gadiform fishes. By Richard W. Langton and Ray E. Bowman. February 1980, iv + 23 p., 3 figs., 11 tables.
741. Distribution of gammaridean Amphipoda (Crustacea) in the Middle Atlantic Bight region. By John J. Dickinson, Roland L. Wigley, Richard D. Brodeur, and Susan Brown-Leger. October 1980, iv + 46 p., 26 figs., 52 tables.
742. Water structure at Ocean Weather Station V, northwestern Pacific Ocean, 1966-71. By D. M. Husby and G. R. Seckel. October 1980, 18 figs., 4 tables.
743. Average density index for walleye pollock, *Theragra chalcogramma*, in the Bering Sea. By Loh-Lee Low and Ikuo Ikeda. November 1980, iii + 11 p., 3 figs., 9 tables.

NOAA Technical Report NMFS SSRF-783



**Biomass and density
of Macrobenthic Invertebrates
on the U. S. Continental Shelf
off Martha's Vineyard, Mass.,
in Relation to Environmental
Factors**

Don Maurer and Roland L. Wigley

July 1984

U.S. DEPARTMENT OF COMMERCE

Malcolm Baldrige, Secretary

National Oceanic and Atmospheric Administration

John V. Byrne, Administrator

National Marine Fisheries Service

William G. Gordon, Assistant Administrator for Fisheries

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

CONTENTS

Introduction	1
Materials and methods	2
Field and laboratory	2
Analysis	2
Environmental setting	2
Physiography	2
Sediment composition	2
Physical oceanography	4
Circulation	4
Bottom temperature	4
Results	5
Faunal composition	5
Total faunal biomass and density	5
Polychaetous annelids	6
Mollusca	6
Scaphopoda	6
Gastropoda	6
Pelecypoda	8
Crustacea	8
Decapoda	8
Cumacea	8
Isopoda	8
Amphipoda	8
Echinodermata	8
Ophiuroidea	8
Holothuroidea	8
Echinoidea	9
Asteroidea	9
Mud patch fauna	9
Discussion	9
Ecological relationships	9
Depth	9
Temperature	9
Sediment and related environmental variables	10
Faunal assemblage	11
Acknowledgments	12
Literature cited	12
Appendix	14

Figures

1. Location of bottom-sampling stations for fauna and sediment off Martha's Vineyard, Mass., June 1962.	3
2. Bottom-sediment types off Martha's Vineyard, Mass., from samples collected at stations, June 1962.	3
3. Bottom temperature off Martha's Vineyard, Mass., from measurements taken at stations, June 1962.	4
4. Quantitative distribution of biomass for all macrobenthic invertebrates combined off Martha's Vineyard, Mass., from samples collected at stations, June 1962.	5
5. Quantitative distribution of individuals for all macrobenthic invertebrates combined off Martha's Vineyard, Mass., from samples collected at stations, June 1962.	6
6. Summary of average biomass and number of individuals of major taxa in relation to depth, temperature, median sediment size, and silt-clay, off Martha's Vineyard, Mass., from samples collected June 1962.	7

Table

1. Comparison of wet weight biomass of macrobenthos in terms of bathymetric stratum of the U.S. northeastern Atlantic	11
---	----

Appendix Tables

1. Station location and environmental variables off Martha's Vineyard, Mass., June 1962. 14

2. List of macrobenthic invertebrates collected off Martha's Vineyard, Mass., June 1962. 16

3. Wet weight biomass of major taxa of macrobenthic invertebrates per station off Martha's Vineyard, Mass., June 1962. 17

4. Number of individuals per m² of major taxa of macrobenthic invertebrates per station off Martha's Vineyard, Mass., June 1962. 19

Biomass and Density of Macrobenthic Invertebrates on the U.S. Continental Shelf off Martha's Vineyard, Massachusetts, in Relation to Environmental Factors¹

DON MAURER² and ROLAND L. WIGLEY³

ABSTRACT

The mean density and mean biomass of macrobenthic invertebrates on the U.S. continental shelf off Martha's Vineyard, Mass., were 3,008/m² and 245.7 g/m², respectively. The latter estimate was considerably higher than values from the North Sea, Scotian Shelf, and Middle-Atlantic Bight. Molluscs (pelecypods and gastropods) and echinoderms (echinoids and ophiuroids) greatly influenced patterns of total biomass distribution. The ocean quahog, *Arctica islandica*, was the dominant species in terms of biomass. Total density was dominated by crustaceans (amphipods), polychaetous annelids, molluscs (small pelecypods), and echinoderms (ophiuroids).

Mean density of mollusca was positively associated with sediment size. Mean biomass and density of crustacea were negatively associated with depth, grain size, and bottom temperature, whereas the same parameters for the Echinodermata were positively associated with those environmental factors.

Three faunal assemblages emerged which were analogous to those described from earlier studies on Georges Bank (sand fauna, silt-sand fauna, muddy-basin fauna). The fauna from the Mud Patch most closely resembled the silty-sand fauna.

INTRODUCTION

Research on the northeastern U.S. continental shelf has accelerated because of a wide variety of human activities ranging from fishing and recreation to transportation, mineral recovery, waste disposal, and oil and gas exploration (Grosslein et al. 1979). These activities are not always compatible, and managers face important decisions attempting to reconcile diverse uses of this valuable resource—the continental shelf. Before decisions can be made, the resource must be assessed in a variety of ways.

Any assessment of the shelf should involve benthic invertebrates, an important component of the shelf ecosystem (Mills and Fournier 1979). The benthos are important in their own right and as a measure of the health of the ecosystem, and play a critical role in trophic relationships providing a major source of energy to economically and ecologically important groundfish (Cohen et al.⁴). In addition, the benthos play a supporting role in nutrient exchange, providing a mechanism for flux of nutrients initially trapped in sediments to the water column (Zeitzschel 1980). The purpose of this account is to report on the distribution of biomass and density of macrobenthic invertebrates off Martha's Vineyard.

Georges Bank off southern New England has been the scene of intensive fishing activity for over 300 yr (Wigley 1961). The Georges Bank area is bounded on the southwest by Martha's

Vineyard and Nantucket Island. Directly south of Martha's Vineyard is an area about 80-100 m deep termed the Mud Patch, consisting of fine-grain sediment. This sediment type is relatively rare on the northeastern shelf which normally consists of sand. The origin of the Mud Patch is uncertain, but it may be an active site of deposition (Milliman et al.⁵). If so, then processes controlling deposition of fine-grain sediment must be considered in assessing their effect on benthic biota. Presumably, benthos living in the Mud Patch would be exposed to relatively higher levels of particulate contaminants (trace metals, hydrocarbons) than benthos living on the surrounding, relatively dynamic, sand bottoms. Assuming this relationship, bioaccumulation of contaminants by the benthos might provide opportunity for biomagnification through food webs. The trophic relationship between demersal fish and benthos is well documented for Georges Bank (Wigley 1965).

In the early 1960's benthic research was conducted on the shelf south of Martha's Vineyard and Nantucket I. Some of these data were reported (Wigley 1963; Wigley and McIntyre 1964; Wigley and Stinton 1973), but the largest portion was placed in a data file report pending further analysis and synthesis (Maurer and Wigley⁶). The report contains maps of biomass and density distribution. With the advent of gas and oil exploration, together with other diverse activities on Georges Bank, documentation of the benthos in and around the Mud Patch would seem to be of special interest to man-

¹Contribution No. 13 from the Southern California Ocean Studies Consortium.

²Southern California Ocean Studies Consortium, California State University, Long Beach, 1250 Bellflower Blvd., Long Beach, CA 90840.

³35 Wilson Road, Woods Hole, MA 02543.

⁴Cohen, E. B., M. D. Grosslein, M. P. Sissenwine, and F. Steimle. 1979. An energy budget of Georges Bank. Workshop on Multispecies Approaches to Fisheries Management, St. John's, Newfoundland, 26-30 November 1979, p. 1-37. National Marine Fisheries Service Woods Hole Lab., Woods Hole, Mass.

⁵Milliman, J. D., M. H. Bothner, and C. M. Parmenter. 1980. Sesson in New England Shelf and Slope Waters, 1976-1977. In J. M. Aaron (editor), Environmental geological studies in the Georges Bank area, United States northeastern Atlantic Outer Continental Shelf, 1975-1977, p. 1-1 to 1-73. Final Report submitted to the Bureau of Land Management, U.S.G.S., Woods Hole, Mass.

⁶Maurer, D., and R. L. Wigley. 1981. Distribution of biomass and density of macrobenthic invertebrates on the U.S. Continental Shelf off Martha's Vineyard. Lab. Ref. Doc. 81-15 (unpubl.), 97 p. Northeast Fisheries Center, NMFS, NOAA, Woods Hole, Mass.

agers because of its proximity to the coast and its potential as a depositional sink of contaminants. Although there are a number of review documents dealing with benthos off southern New England (Pratt 1973; TRIGOM-PARC⁷; Wigley and Theroux^{8,9}; Maurer¹⁰), no study has featured the fauna of the Mud Patch.

MATERIALS AND METHODS

Field and Laboratory

Samples were collected 11-20 June 1962 by the National Marine Fisheries Service vessel RV *Delaware* at 64 stations south of Martha's Vineyard, Mass. (Fig. 1, Appendix Table 1). Stations were spaced at intervals of 16 km on a grid pattern with eight north-south transects at right angles to the depth contours. At each station (except station 7) two quantitative bottom samples were collected with a Smith-McIntyre grab. This instrument effectively sampled a 0.1 m² area of bottom to a depth of 10-17 cm. At sea, grab samples for macrobenthic studies were washed through a 1.0 mm mesh screen. Macrobenthos remaining on the screen after washing were removed and preserved in a solution of neutral Formalin.

In the laboratory macrobenthos were sorted, identified, counted, and weighed. External moisture was removed from specimens by blotting. Shells, internal skeletons, and exoskeletons were included in the values expressed as wet weight biomass (g/m²).

Sediment samples were collected with a Dietz-LaFond grab at each station and at two localities equally spaced between stations along the cruise. The locations of sediment samples used in conjunction with biological analyses are depicted in Figure 1. Terminology follows the Wentworth Particle Size Classification (Twenhofel and Tyler 1941), and nomenclature follows the classification of Shepard (1954) and Emery (1960). Determinations were made of median sediment size (ϕ), percent sand, percent silt, percent clay, percent Kjeldahl nitrogen, percent organic carbon measured, and carbon/nitrogen ratio of the sediments (Hathaway¹¹). Standard sieving procedures were used to measure sediment size. Based on sediment analyses, a composite sediment-type map was made (Fig. 2). Appendix Table 1 lists environmental data per station.

Analysis

Wet weight and number per taxon by station and gear were punched on cards and a computer listing was prepared. Based on the listing, the average biomass and density per station of major taxa (Amphipoda, Pelecypoda, Asteroidea, etc.) were determined and distribution maps plotted by computer (Maurer and Wigley footnote 6). In some cases, maps were also made of particularly important genera and species. Correlation coefficients (R) were computed for average weight and number transformed ($\log_e(N + 1)$) of major taxa in relation to environmental variables.

ENVIRONMENTAL SETTING

Physiography

The Georges Bank area off New England is a submerged northeast extension of the Atlantic Coastal Plain (Aaron¹²). The Bank, which encompasses about 42,000 km², is covered by up to 200 m of water. The study area for this account (Fig. 1) lies immediately southwest of the Bank.

The study area encompasses about 130 km² and extends across the continental shelf to the upper portion of the continental slope. Bottom topography is moderately smooth. Water depths increase gradually and rather uniformly from shore outward to the shelf break, about 120 m. The average gradient of the continental slope off the Middle Atlantic Bight varies from 2° to 7° (Milliman 1973); beyond the shelf break in the study area, the depth gradient is relatively steep, averaging 4° (Wigley and Stinton 1973). The most distinguishing feature on the shelf break is the number of gullies and canyons that transect the slope (Fig. 1).

Sediment Composition

Six major sediment types occurred in the study area (Fig. 2). Sand, silty sand, and sandy silt occurred over a large area, whereas gravel-sand, sand-silt-clay, and silt were much less widespread. Sand with some gravel (stations 1, 45, 47) covered more than half the area, mainly in shallow water (0 to 60-80 m) except in the eastern sector and in a narrow (6 km) band parallel to and just below the outer periphery of the continental shelf. In shallow water the sands were silt free and occasionally mixed with large quantities of shell (mollusks and echinoderm plates). Admixtures of silt occurred with the sand over most of the remaining area.

A large area (80 × 100 km) of fine-grain sediment (Mud Patch) occurred in the southwestern sector (Fig. 2). A relatively circular area of sand-silt-clay near its center was surrounded by an inner band of sandy silt which grades to an outer band of silty sand. Illite is normally the most important clay mineral, and organic carbon is higher here than in the surrounding sand (Appendix Table 1). This is the largest known natural area of fine-grain sediment on the Middle Atlantic Shelf. Sediments on the continental slope were dominated by silt and clay.

⁷TRIGOM-PARC. 1974. A socio-economic and environmental inventory of the North Atlantic Region, Vol. 1, Book 3, 198 p. Report to the Bureau of Land Management, South Portland, Me.

⁸Wigley, R. L., and R. B. Theroux. 1976. Macrobenthic invertebrate fauna of the Middle Atlantic Bight region. Part II: Faunal composition and quantitative distribution. Northeast Fisheries Center, NMFS, Woods Hole, Mass., 395 p.

⁹Wigley, R. L., and R. B. Theroux. Reconnaissance survey of the quantitative distribution of macrobenthic invertebrates in the offshore New England region. Manuscr. in prep. Northeast Fisheries Center, NMFS, Woods Hole, Mass.

¹⁰Maurer, D. 1982. Review of benthic invertebrates of Georges Bank in relation to gas and oil exploration with emphasis on management implications. Report to Northeast Fisheries Center, Woods Hole, Mass., and Sandy Hook Laboratory, Highlands, N.J., 329 p.

¹¹Hathaway, J. C. (editor). 1971. Data file, continental margin program, Atlantic Coast of the United States. Vol. 2. Sample collection and analytical data. Ref. No. 71-15, 496 p. Woods Hole Oceanographic Institution, Woods Hole, Mass.

¹²Aaron, J. M. 1980. A summary of environmental geologic studies in the Georges Bank area, United States northeastern Atlantic outer continental shelf, 1975-1977. Executive Summary of the Final Report submitted to the Bureau of Land Management. U.S.G.S., Woods Hole, Mass., 22 p.




Figure 1.—Location of stations of bottom samples for fauna (o) and sediment (x) off Martha's Vineyard, Mass., June 1962. Stations 1, 2, 6, and 7 are from the 1977 BLM study.




Figure 2.—Bottom-sediment types off Martha's Vineyard, Mass., from samples collected at stations (o), June 1962.

In the fine-grain sediments southwest of Nantucket I., values of 6.0-10 mg/g organic carbon and 0.8-1.2 mg/g nitrogen were recorded (New England OCS¹³). Highest concentrations (15,000 µg/l) of seston were measured on the shelf south of Nantucket Shoals during the winter (Milliman et al. footnote 5; Aaron et al. 1980).

Physical Oceanography

Circulation.—The major pattern of circulation on Georges Bank was described earlier by Bigelow (1927) and Bumpus (1973, 1976) and has been updated by Butman et al. (1980). The data of Butman et al. showed a clockwise circulation pattern around the Bank. Flow on the southern flank was consistently toward the southwest. West of the Great South Channel and south of Nantucket I., mean flow was measured to the west-northwest along contours of local bathymetry (Dorkins 1980).

On the eastern side of Great South Channel, mean flow was consistently toward the north at 5-10 cm/s at all depths (Butman et al. 1980). At a station just south of Great South Channel, there was a westerly flow similar to that along the southern

flank. On the western side of Great South Channel, little net flow was measured.

Great South Channel has been historically considered a major exit for Gulf of Maine waters (Schlitz in press). However, recent water current measurements indicate that mean flow within the channel is directed mainly toward the Gulf of Maine (Butman et al. 1980). Schlitz et al. (1977) showed that a permanent front exists across the channel from Nantucket Shoals toward Georges Bank, separating the Gulf of Maine from the shelf to the south. According to Schlitz (in press), a small amount of water flows southward through Nantucket Shoals, but this probably does not contribute a significant volume.

Bottom temperature.—When the study was conducted in June 1962, a cell of cold bottom water (6.1°-6.9°C) extended in an east-west band from the New York Bight eastward to long. 69°30'W (Fig. 3). This cell occurred at depths of 40-80 m, roughly the mid-shelf region. The cold cell was bounded on the north by higher coastal water temperatures (<12°C) and on the south by values of 10-12°C near the shelf break. Long-term (1940-66) annual maximum and minimum bottom-water temperatures near the shelf break were 16°-17°C and 1°-2°C, respectively (Colton and Stoddard 1973). However, the annual range here is normally 2°C. Offshore shelf waters, particularly in shallow portions, may range from 3°C in February-March to 14°C in September-November (Wigley and Stinton 1973).

¹³New England OCS Environmental Benchmark. 1978. Draft final report, Vol. V, Sects. G-J. Energy Resources Co., Cambridge, Mass., 224 p.




Figure 3.—Bottom temperature (°C) off Martha's Vineyard, Mass., from measurements taken at stations (o), June 1962.

RESULTS

Faunal Composition

A total 214 taxa were identified (Appendix Table 2). Since some taxa were identified only to phylum (Porifera, Nemertea) and since some taxa identified to genus may include several species, the total number of taxa is conservative. Of the number of taxa, ~24.3% were molluscs, 27.1% arthropods, 24.3% polychaetous annelids, 10.7% echinoderms, and 13.6% miscellaneous taxa, including sipunculids, coelenterates, nemerteans, ectoprocts, ascidiaceans, pogonophorans, and hemichordates. Biomass (wet weight g/m^2) and density (no./ m^2) are presented in Appendix Tables 3 and 4. About 190,000 individuals and 15,500 g of specimens were collected.

Total Faunal Biomass and Density

Biomass and density were distributed among the major taxa as follows:

	Biomass (g/m^2)	Density (no./ m^2)
Annelida	10.9%	20.3%
Mollusca	56.7%	5.9%
Crustacea	4.1%	62.5%
Echinodermata	21.3%	7.0%
Misc. taxa	7.0%	4.3%

Average biomass and density per station for total fauna were $245.7 g/m^2$ and $3,008$ individuals/ m^2 , respectively. Biomass in the shallow central stations generally ranged from 100 to $999 g/m^2$ (Fig. 4). Biomass decreased to a range of 1-100 g/m^2 in slightly deeper water and then increased again to 100-999 g/m^2 between 60 and 100 m. Below 100 m on the western side of the study area, biomass again decreased to $<100 g/m^2$, but biomass was still in the 100-999 g/m^2 range on the eastern side to almost 200 m. Molluscs, polychaetes, and shallow-water echinoderms contributed heavily to the highest values of biomass.

The density pattern for total fauna was more irregular than for biomass (Fig. 5). A range of 1,000 to $>3,000 m^2$ occurred in shallow water. A major portion of the area contained densities of 1,000-2,999/ m^2 . Lowest densities (1-999/ m^2) were recorded below ~ 100 m. Crustacea, primarily amphipods, and polychaetes contributed greatly to the highest density values.

Stations were grouped into ranges of depth, temperature, median sediment size, and silt-clay (Maurer and Wigley footnote 6). Mean biomass and density were calculated for those ranges and summarized. A depth range of 40-80 m, temperature 6.0° - $7.9^\circ C$, ϕ 2.0-3.99, and 0.6% silt-clay comprised the main distribution of biomass (Fig. 6). The principal distribution of density was accounted for by a depth range of 0-100 m, temperature 6.0° - $9.9^\circ C$, ϕ 1.0-3.99, and 0-40% silt-clay. Mean biomass and density declined with depth but the correlations were low ($R = -0.20$).




Figure 4.—Quantitative distribution of biomass (g/m^2) for all macrobenthic invertebrates combined off Martha's Vineyard, Mass., from samples collected at stations (o), June 1962.




Figure 5.—Quantitative distribution of individuals (no./m²) for all macrobenthic invertebrates combined off Martha's Vineyard, Mass., from samples collected at stations (o), June 1962.

Polychaetous Annelids

Many species of polychaetes were identified (Appendix Table 2). The number of species is a conservative estimate because recent work with a 0.5 mm mesh sieve has yielded higher numbers of species (Maurer and Leathem 1980). Mean biomass of polychaetes ranged from 0.1-39 g/m² throughout most of the area to >100 g/m² at a few stations (Maurer and Wigley footnote 6). A biomass range of 40-99 g/m² was measured directly south of Nantucket I. and Nantucket Shoals.

Polychaetes were more evenly distributed throughout environmental ranges than any of the major taxa (Fig. 6). Mean biomass and density were negatively associated with temperature ($R = -0.39$, $R = -0.39$). A number of polychaete species showed marked differential distribution. Maldanids and *Scalibregma inflatum* occurred in shallow water and sand. The latter species showed the same relationship over Georges Bank (Maurer and Leathem 1980). In contrast, terebellids and *Sternapis scutata* occurred in deeper water with fine-grain sediment.

Mollusca

In terms of mean biomass and density, Mollusca contained Scaphopoda (0.1 and 0.6%, respectively), Gastropoda (4.8 and 15.2%), and Pelecypoda (95.1 and 84.2%). Biomass ranged from 0.1 to >100 g/m² with the highest values directly south of Nantucket Shoals and southwest of Nantucket I. (Maurer and Wigley footnote 6). A depth range of 20-80 m, 6.0°-9.9°C, ϕ 2.0-3.9, and 0-20% silt-clay comprised the main distribution

of biomass (Fig. 6). Density distribution of Mollusca was very different from biomass distribution because the most numerous molluscan species were smaller than the main contributors to biomass. Densities ranging from 100 to >400/m² occurred in the central part of the study area between 60 and 100 m. This central area was surrounded by densities ranging from 1 to 99/m². The main distribution of density for molluscs encompassed 80-100 m, 7.0°-10.9°C, ϕ 3.0-5.9, and 20-100% silt-clay (Fig. 6). Mean density of mollusca was positively associated with grain size ($R = 0.44$).

Scaphopoda.—The scaphopods *Cadulus pandionis*, *C. verrilli*, and *Dentalium occidentale* were sampled only at deep stations (Maurer and Wigley footnote 6). Scaphopods occurred in relatively low biomass (0.05-0.08 g/m²) and density (3.3-7.5/m²) in water deeper than 200 m. Their distribution mainly encompassed 10.0°-12°C, ϕ 4.0-5.9, and 40-80% silt-clay (Fig. 6).

Gastropoda.—Although biomass and density of gastropods ranged from 0.1 to >100 g/m² and from 1 to >400/m², respectively, there were only a few stations that contained >100 g/m² or >400/m² (Maurer and Wigley footnote 6). Biomass and density patterns of gastropods were very different (Fig. 6). Biomass patterns were mainly influenced by large-to-intermediate size, shallow water, sand-dwelling taxa (*Colus*, *Nassarius*, *Buccinum*, *Neptunea*, *Lunatia*), and density was influenced by smaller, deep-water, silt-sand and sandy-silt species (*Alvania carinata*). There was a significant negative association between density of gastropods and temperature ($R = -0.37$).

	Depth (m)							Temperature (°C)						Median Sediment Size ϕ					% Silt - Clay						
	20	40	60	80	100	200	>200	6	7	8	9	10	11	12	13	1	2	3	4	5	0	20	40	60	80
Porifera	1							2						<1					<1						
Coelenterata	26							53						43					41						
Nemertea	3							4						1					2						
Annelida	29 43 46 23							30 40 25 28						25 29 31					32 25						
Pogonophora	<1							<1						<1					<1						
Sipunculida	3 4							2 4						2					3						
Mollusca	204 223 247							187 252 122 156						183 241					260 50						
Gastropoda	30							41						18					13						
Pelecypoda	118 192 367 137							134 186 236 214						225 175 200					216 163 176 271						
Scaphapoda	<1							<1						<1					<1						
Crustacea	12 19 14							13 15 13 12						13 12 13					15 11						
Cumacea	2084							2127						2745					3203 1088						
Isopoda	45 34							18 21 25 38						40 25					33						
Amphipoda	9 18 12							12 14 10 10						12 10 12					13 10						
Decapoda	1941							2063						2680					3101 1059						
Echinodermata	35 157 72							63						35					101 93 77 77						
Holothuroidea	45 51 64 439							228 435 332 410 223						164 196 323 503					353 196 338 514						
Echinoidea	28 28 121							56 41						56 56					56 71 61						
Ophiuroidea	8							9						25					9 23						
Asteroidea	24 40							73 104						32					32 157						
Ectoprocta	24 32							28 35 17						17 17					19 15 22						
Enteropneusta	563 375							129 314 314 394 212						159 287 328					319 310 356						
Ascidiacea	12							5						5 5 7					8 12						
Miscellaneous	6 4							5 9						5 8					6 8						
Total	8							244						6					<1						
	220							2						3					3						
	<1 <1							<1						<1					1						
	4 4 5							4						3 4					6						
	10 5							8						11					4 9						
	106 52 32 36							51						112 55					50 50 48						
	25 31							20 57 55						18 50 14					44 14 13						
	354 103 107 108 157							325 104 516						142 115 93 116					103 99 153						
	276 323 354 214							300 372 258						252 378 151					336 197 160						
	3223 3130 3442							3145 4464 3483 2114						3606 3440 3440					4247 2286 1763						

Figure 6.—Summary of average biomass (g/m²) and number of individuals (no./m²) of major taxa in relation to depth, temperature, median sediment size, and silt-clay, off Martha's Vineyard, Mass., from samples collected June 1962. Biomass above line; number below line.

Pelecypoda.—Mean biomass of pelecypods ranged from 0.1 to >100 g/m². Highest values were recorded directly south of Nantucket Shoals and Nantucket I. (Maurer and Wigley footnote 6). A range of 20-80 m, 6.0°-9.9°C, ϕ 2.0-3.9, and 0-20% silt-clay comprised the main biomass distribution (Fig. 6).

Density patterns differed from biomass patterns because of size differences. A range of 40-200 m, 6.0°-10.9°C, ϕ 3.0-5.9, and 20-100% silt-clay comprised the main density distribution. There was a significant negative association between pelecypod biomass and temperature and depth, and a positive association between density and sediment size. Species of bivalves showed marked differential distribution. *Arctica islandica* and *Cerastoderma pinnulatum* mainly occurred at depths shallower than 80 m and in sand with <20% silt-clay. In contrast, *Cuspidaria striata*, *C. perrostrata*, and *Batharca pectunculoides* occurred almost exclusively at depths >150 m in sediment with >50% silt-clay. Molluscan biomass distribution was chiefly influenced by the distribution of the pelecypod *A. islandica*. Small species of bivalves contributed significantly to the density pattern of Mollusca.

Crustacea

In terms of biomass and density, the Crustacea contained Decapoda (5.7 and 0.2%, respectively), Cumacea (1.3 and 2.1%), Isopoda (5.0 and 1.0%), and Amphipoda (88 and 96.7%). Mean biomass and density of crustaceans ranged from 0.1 to >50 g/m² and from 1 to >1,000/m². A depth range of 20-80 m, 6.0°-9.9°C, ϕ 1.0-3.9, and 0-20% silt-clay comprised the main distribution of crustacean biomass and density (Fig. 6). Mean biomass and density were negatively associated with depth ($R = -0.53$, $R = -0.56$), sediment size ($R = -0.31$, $R = -0.30$), and bottom temperature ($R = -0.48$, $R = -0.50$).

Decapoda.—Mean biomass and density of decapods ranged from 0.1 to 49 g/m² and from 10 to 25/m². Their distribution was sporadic (Maurer and Wigley footnote 6), but grab samples are not the most effective way to collect large crustaceans. *Crangon septemspinosa*, *Pagurus pubescens*, *Cancer* spp., and *Pandalus* spp. were collected frequently. *Crangon septemspinosa* occurred at depths <20 m to >200 m. In contrast, *Hyas* sp., *Euprognatha* sp., *Munida* sp., and *Geryon* sp. were generally collected deeper than 100 m.

Cumacea.—Mean biomass and density of cumaceans ranged from 0.1 to 1.0 g/m² and from 1.0 to >400/m². Cumaceans occurred widely and were relatively evenly spread, occurring mainly at lower densities of 1-100/m² (Maurer and Wigley footnote 6). Characteristic species included *Diastylis polita*, *D. quadrispinosa*, *Eudorella emarginata*, *Leptostyllis* sp., *Eudorellopsis* sp., and *Leptocuma* sp. A depth range of 20 m, 7.0°-8.9°C, ϕ 3.0-3.9, and 0-10% silt-clay comprised the main biomass distribution, whereas the main density distribution was most accurately contained in a depth range of 20-40 m, 6.0°-8.9°C, ϕ 2.0-3.9, and 0-20% silt-clay (Fig. 6). Mean biomass and density of cumaceans were negatively associated with depth ($R = -0.36$, $R = -0.52$), and density with temperature ($R = -0.29$).

Isopoda.—Mean biomass and density ranged from 0.1 to 10 g/m² and from 1 to >100/m². The greatest numbers of isopods

occurred between 40 and 60 m throughout most of the study area, but between 60 and 80 m directly south of Nantucket Shoals (Maurer and Wigley footnote 6). Biomass and density patterns were essentially the same. Characteristic species were *Cirolana polita*, *Chiridotea tuftsi*, *Ptilanthura tenuis*, *Edotea triloba*, and *Calathura* sp. Mean density of isopods was negatively associated with depth ($R = -0.38$), sediment size ($R = -0.38$), and temperature ($R = -0.26$).

Amphipoda.—Mean biomass and density of amphipods ranged from 0.1 to >50 g/m² and from 1 to >1,000/m². The shallow water portion mostly contained densities >1,000/m². There was a sharp decline in biomass and density below 100 m (Maurer and Wigley footnote 6). Characteristic amphipod taxa were *Leptocheirus pinguis*, *Unciola irrorata*, *Caprella* spp., *Coropium* spp., ampelescids (including *Ampelisca compressa*, *A. macrocephala*, and *Byblis serrata*), phoxocephalids, photids, and haustoriids. A depth range of 20-80 m, 6.0°-9.9°C, ϕ 1.0-3.9, and 0-40% silt-clay comprised the main distribution of biomass. These ranges were the same for density except for depth with a range of 20-100 m (Fig. 6). The distribution of amphipod biomass and density greatly influenced the same patterns for combined Crustacea. Mean biomass and density were negatively associated with depth ($R = -0.53$, $R = -0.54$), sediment size ($R = -0.31$, $R = -0.31$), and temperature ($R = -0.49$, $R = -0.51$).

Echinodermata

In terms of biomass and density, echinoderms contained Ophiuroidea (21.7 and 78.6%, respectively), Holothuroidea (52.9 and 5.9%), Echinoidea (19.7 and 13.8%), and Asteroidea (5.7 and 1.7%). Mean biomass and density ranged from 0.1 to >100 g/m² and from 1 to >1,000/m². Biomass and density patterns of echinoderms were generally dissimilar (Fig. 6). Mean biomass and density were positively associated with depth ($R = 0.27$, $R = 0.25$), sediment size ($R = 0.44$, $R = 0.36$), and temperature ($R = 0.33$, $R = 0.42$).

Ophiuroidea.—Mean biomass and density of ophiuroids ranged from 0.1 to >100 g/m² and from 1 to >500/m². However, in general, biomass values of 0.1-49 g/m² were most common, as were densities of 1 to >500/m². Characteristic species were *Amphilimna olivacea*, *Amphioplus abditus*, *A. fragilis*, *Axiognathus squamatus*, and *Ophiura sarsi*. In terms of biomass, ophiuroids were mainly restricted to depths >80 m, 10.0°-12.9°C, ϕ 4.0-5.9, and 20-100% silt-clay, and for density 7.0°-12.9°C and ϕ 3.0-5.9 (Fig. 6). Mean biomass and density were positively associated with depth ($R = 0.44$, $R = 0.40$), sediment size ($R = 0.36$, $R = 0.40$), and temperature ($R = 0.54$, $R = 0.48$).

Holothuroidea.—Mean biomass and density of holothurians ranged from 0.1 to >100 g/m² and from 1 to 399/m². Holothurians were almost exclusively collected in the deeper portion of the study area below 80 m (Maurer and Wigley footnote 6). Characteristic species were *Synapta* sp., *Astichopus* sp., *Molpadia* sp., and *Havelockia scabra*. A depth range of 40-100 m, 6.0°-7.9°C, ϕ 3.0-4.9, and 20-80% silt-clay comprised the main biomass distribution, whereas a depth range of 80-100 m and 6.0°-10.9°C comprised the main density distribution (Fig. 6).

Echinoidea.—Mean biomass and density of echinoids ranged from 0.1 to >100 g/m² and 0.1 to 399/m². The distribution of echinoids was primarily influenced by the distribution of two species (*Echinarachnius parma* and *Schizaster fragilis*) (Maurer and Wigley footnote 6). *Echinarachnius parma* influenced echinoid biomass and density in shallower water, sand, and low percent silt-clay, and *S. fragilis* was influenced in deeper water, sandy-silt, and high silt-clay.

Asteroidea.—Mean biomass and density of asteroids ranged from 0.1 to >50 g/m² and from 1-49/m². Their distribution was sporadic (Maurer and Wigley footnote 6), but grab samples are not the most effective way to collect large asteroids. Characteristic species were *Asterias vulgaris*, *Leptasterias* sp., *Porania* sp., *Henricia* sp., *Astropecten americanus*, and *Astropecten* sp. *Asterias vulgaris* represented a shallow-water species, and *Astropecten americanus* was a deeper water species. Mean biomass and density were positively associated with sediment size ($R = 0.32$, $R = 0.26$).

Mud Patch Fauna

Earlier in this report, the southwest quadrant of the Martha's Vineyard-Nantucket Shoals area was referred to as the Mud Patch (Fig. 2). The following stations generally comprised the Mud Patch: Stations 23-25, 33-36, 38-42, 49-53, and 57-59. Average biomass for these stations was 123.7 g/m², and density was 1,536.2/m², which is markedly lower than respective values of 245.7 g/m² and 3,008/m² for the entire Martha's Vineyard-Nantucket Shoals area.

Based on two BLM stations (1 and 2) in the shallow water and clean sand of Martha's Vineyard-Nantucket Shoals area (Fig. 1), the dominant polychaete species (in abundance and frequency of occurrence) were *Exogone hebes*, *Spiophanes bombyx*, *Parapionosyllis longicirrata*, *Sphaerosyllis erinaceus*, *Owenia fusiformis*, *Scalibregma inflatum*, *Aricidea catherinae*, and *Spio pettiboneae* (Maurer and Leathem¹⁴).

The Mud Patch was characterized by anthozoans, small-to-medium size bivalves (*Bathyarca pectunculoides*, *Nuculana acuta*, *Yolida sapatilla*, *Nucula* spp.), small-to-medium size decapods (*Catapagurus sharreri*, *Hyas coarctatus*, *Euprognatha rastellifera*, *Munida iris*), and a variety of generally small echinoderms (*Amphilimna olivacea*, *Amphioplus macilentus*, *Amphiura otteri*, *Havelockia scabra*, *Schizaster fragilis*). Based on two BLM stations (6 and 7) in sediment containing 32-37% silt-clay near the Mud Patch (Maurer and Leathem footnote 14), dominant species were *Paraonis gracilis*, *Cossura longocirrata*, *Ninoe nigripes*, *Aricidea suecica*, *Nephtys incisa*, *Tharyx annulosus*, *Terebellides stroemii*, *Maldanidae*, and *Cirratulidae* in the deeper water and silty-sand and sandy-silt. Robert Reid¹⁵ reported high numbers of the amphipod *Ampelisca agassizi* from the Mud Patch in some 1980 collections together with other species cited above. This fauna contains a high proportion of selective and non-selective deposit feeders, both surface and buried, representing a typical soft-bottom community. Echinoderms were particularly important as con-

tributors to biomass in the Mud Patch. Because of their general mode of feeding, the biota of the Mud Patch would have considerable potential for ingesting pollutants associated with deposition of fine-grain sediment.

DISCUSSION

Ecological Relationships

A summary of mean biomass and number of individuals of major taxa in relation to depth, temperature, median sediment size, and silt-clay is presented in Figure 6. Also contained in Figure 6 are the range (biomass and number of individuals) of each taxon per environmental variable, together with values of average biomass and density plotted for a specific mid-range.

Depth.—Biomass and density of polychaetes off Martha's Vineyard were not significantly associated with depth. However, for the Georges Bank area, the density of polychaetes increased significantly with depth down to ~80 m with some indication of reduction in biomass with depth (Maurer and Leathem footnote 14). Off Martha's Vineyard, biomass and density of molluscs combined were not significantly associated with depth. However, the density of scaphopods increased significantly with increasing depth and the biomass of pelecypods decreased with depth. Biomass pattern of pelecypods with depth was dominated by the distribution of *A. islandica* ranging between 40 and 60 m. For Georges Bank, density of combined molluscs increased with water depth (Maurer¹⁶).

For combined crustaceans off Martha's Vineyard, mean biomass decreased with depth. Mean density of cumaceans and isopods was negatively associated with depth. Mean biomass of amphipods decreased with increasing depth. Density pattern of combined crustaceans was dominated by the distribution of amphipods, whereas biomass pattern of combined crustaceans was dominated by decapods. For Georges Bank, mean biomass of amphipods decreased in deeper water (Maurer footnote 16).

For combined echinoderms off Martha's Vineyard, mean biomass and density increased with depth. In another study, the bathymetric distribution of echinoderms off the northern Oregon coast was not as uniform, as low wet weights were obtained from depths of 86-139 m and 1,189-1,234 m (Alton 1972). Mean biomass and density of ophiuroids increased with depth off Martha's Vineyard. For Georges Bank, mean density of combined echinoderms showed no significant relationship with depth, but biomass decreased significantly in deeper water (Maurer footnote 16). These relationships reflected the fact that although density was relatively regular throughout the depth range, larger species (sea stars and echinoids) occurred in relatively shallow water, with smaller species (brittle stars) in deeper water.

Mean biomass and number of individuals of combined taxa were high between 0 and 100 m depths. Biomass and density decreased with increasing depth, although these relationships were not statistically significant. Off southern New England, there was a marked reduction in density and biomass with

¹⁴Maurer, D., and W. Leathem. 1980. Ecological distribution of polychaetous annelids of Georges Bank. CMS-1-80. College of Marine Studies, Univ. of Delaware, Lewes, Del., 181 p.

¹⁵Robert Reid, Northeast Fisheries Center Sandy Hook Laboratory, National Marine Fisheries Service, NOAA, Highlands, NJ 07732, pers. commun. 1981.

¹⁶Maurer, D. 1982. Review of benthic invertebrates of Georges Bank in relation to gas and oil exploration with emphasis on management implications. Report to Northeast Fisheries Center, NMFS, Woods Hole, Mass., and Sandy Hook Laboratory, Highlands, N.J., 329 p.

depth (Wigley and Theroux 1981). Total biomass was dominated by molluscs (pelecypods and gastropods) and echinoderms (echinoids and ophiuroids), and density pattern was dominated by amphipods, polychaetes, small bivalves, and ophiuroids. According to Parsons et al. (1977), a number of studies have demonstrated decreased macrofaunal numbers and biomass with increased water depth. Abrupt faunal discontinuities tended to occur at 100-300 m. These changes generally corresponded to the vertical distribution of other environmental factors, such as organic carbon, nitrogen, and total particulates which decrease rapidly with increasing depth. Depth-related decreases in macrobenthos are more closely linked to suspended living biomass than to total particulate matter (Parsons et al. 1977).

Parsons et al. (1977) cited examples of higher mean biomass in regions of higher phytoplankton production. This relationship suggests that benthic biomass in inshore coastal areas may be strongly influenced by sedimentation of organic matter produced during a bloom. This process undoubtedly contributes to macrobenthos at shallow depths of continental shelves. The shallow depths of Georges Bank and Nantucket Shoals fit these conditions.

Data from other studies on benthic biomass with depth are presented in Table 1. Mean biomass was generally highest in the 50-99 m stratum. Moreover, mean biomass was higher off New England than the New York Bight. Off the Scotian Shelf, wet weight biomass was 24 g/m² at 0-90 m depth and 22.1 g/m² at 90-180 m (Mills 1980). The inverse relationship between benthic biomass and depth is probably more related to the geographic position of zones of primary productivity and sedimentation on shelves rather than to absolute values of depth per se. This relationship has been recognized for some time (Rowe 1971; Sokolova 1972).

Another aspect related to depth involves substratum stability. Sediments in shallow coastal waters and the inner shelf are subject to vigorous current action. Wave base to 80-85 m depth is not uncommon during the winter on Georges Bank (Aaron et al. 1980). Accordingly during times of high energy flow, sediment stability decreases, impeding colonization by many infaunal organisms. Specialized species such as haustoriid amphipods and rapidly burrowing bivalves commonly dominate these sites. Nantucket Shoals represents an area of high seasonal sediment instability.

Temperature.—Although there were some quantitative relationships between biomass and density of various taxa and bottom temperature, the presence of a cold-water cell bounded by warmer water on the north and south made it difficult to in-

terpret distribution according to temperature (Fig. 3). In this case, there was a shallow-water (0-50 m) and deep-water (80-100 m) zone both containing water ranging from 7° to 12°C. Although the maximum biomass of certain taxa (Ophiuroidea) was associated with shallow water and deep water (Maurer and Wigley footnote 6), their maximum biomass was similar according to temperature (11.0°-12.9°C). The Coelenterata, another deep-water taxon, had its highest biomass in 11.0°-11.9°C (Maurer and Wigley footnote 6), temperatures normally associated with depths of about 40 m for this time of year. Thus it is important to bear in mind the relative position of the cold-water cell in relation to depth when comparing distribution according to temperature.

Off Atlantic City, N.J., Boesch et al.¹⁷ concluded that temperature was the principal hydrographic factor affecting macrobenthic distribution. Temperatures were more variable on the inner and central shelf and more constant on the outer shelf. Differences in temperature regime were probably the prime cause of the sharper faunal change at the outer shelf/shelf-break transition.

Sediment and related environmental variables.—Off Martha's Vineyard, the pattern of biomass and number of individuals were relatively even throughout a range of median sediment size and percent silt-clay (Fig. 6). For Georges Bank, density of infauna increased significantly with percent gravel (Maurer and Leathem footnote 14). In addition, there were some significant relationships between the density of several dominant polychaete species and sediment parameters (percent silt-clay, percent silt, percent carbon, percent nitrogen, microbial biomass, and bacterial biomass) (Maurer and Leathem footnote 14).

Off Martha's Vineyard, polychaetes were ubiquitous in regard to sediment type and were major contributors to both average density and biomass of all benthic organisms in each sediment type (Maurer and Wigley footnote 6). This apparent lack of correlation with a sediment type may partly be due to differences in sieve size, wherein some of the smaller taxa known to occur abundantly on the shelf were probably missed with a 1.0 mm mesh net. In a related study off southern New England, greatest amounts of polychaetes were found in shell and sand-gravel (750 and 555/m², respectively), somewhat lesser amounts in sand, silty sand, gravel, and silt (433, 331, 289, and 118/m²), and lowest (23 and 9/m²) in sand-shell and clay (Wig-

¹⁷Boesch, D., J. N. Kraeuter, and D. K. Serafy. 1977. Benthic ecological studies: Megabenthos and macrobenthos. In Middle Atlantic Outer Continental Shelf Environmental Studies, Chap. 6, 111 p. Draft Rep. to Bureau of Land Management.

Table 1.—Comparison of wet-weight biomass (g/m²) of macrobenthos (Annelida, Mollusca, Crustacea, Echinodermata) in relation to bathymetric stratum at locations off the U.S. northeastern Atlantic coast.

Depth (m)	Georges Bank ¹ (0.5 mm sieve)	Southern New England ² (1.0 mm sieve)	Martha's Vineyard/ Nantucket Shoals (1.0 mm sieve)	New York Bight ³	
				(1.0 mm sieve)	(0.5 mm sieve)
25-49	140.6	308.6	230.4	121.0	78.0
50-99	460.0	230.1	314.3	156.4	96.6
100-199	24.2	60.4	75.6	27.2	33.2

¹Maurer, see text footnote 16.

²Wigley and Theroux, see text footnote 8.

³Boesch et al., see text footnote 17.

ley and Theroux 1981). Wet-weight biomass values between 20 and 30 g/m² occurred in four sediment types: Shell, silty sand, gravel, and sand in order of decreasing amounts. Silt and sand-gravel contained 7 and 116 polychaetes/m², respectively, and sand-shell and clay had the smallest biomass with 1.7 and 0.45 g/m². In general, there were no correlations between density of annelids and sediment organic carbon for the southern New England area of the New York Bight. The highest biomass values (45.4 and 37.4 g/m²) occurred in sediment with 1.5-1.9% and 2.0-2.9% organic content, respectively.

Off Martha's Vineyard, mean density of combined Mollusca was positively associated with sediment size (ϕ) as was that of pelecypods. For Georges Bank, mean density of combined molluscs increased with percent carbon in the sediment (Maurer footnote 16). Franz (1976) reported three molluscan-sediment groups in northeastern Long Island Sound. One group consisted of very fine sand and contained species similar to that of the Mud Patch. A second and third group consisted of medium sand and coarse sand and contained molluscan species very similar to the sand bottom off Martha's Vineyard. Based on mean grain diameter, sorting, silt-clay content, and fauna, Driscoll and Brandon (1973) identified four facies in Buzzards Bay. In addition, the density of particular molluscan-sediment relationships emerged with certain feeding types. Similar relationships have been reported elsewhere (Maurer 1967a, b). Because of the variety of feeding types in many major taxa, it is difficult to correlate major taxa with sedimentary variables. This exercise is most accurately accomplished at the species level.

Mean biomass and density of combined crustaceans were negatively associated with sediment size off Martha's Vineyard. This pattern was primarily influenced by amphipods and, to a lesser extent, isopods. For Georges Bank, mean density and biomass of amphipods were positively associated with percent sand, and mean biomass declined with percent gravel (Maurer footnote 16). The number of amphipod species increased significantly with percent sand and decreased with percent gravel, percent silt, percent silt-clay, percent carbon, and percent nitrogen. In Long Island Sound, there was a strong correlation in summer between Shannon-Weaver diversity of benthic amphipods and sediment texture, with diversity increasing due to decreasing species dominance, and, most importantly, to increasing species richness as sediments became coarser (Biernbaum 1979). Increased sediment instability caused by winter storms resulted in marked diversity decrease. The response to seasonal sediment stability by the benthic biota, including amphipods, must be a critical feature influencing recruitment, maintenance, and production on shallow portions of Georges Bank, including Nantucket Shoals in this study area.

Mean biomass and density of combined echinoderms were positively associated with sediment type off Martha's Vineyard. This relationship was recorded for ophiuroids and holothurians. According to Tyler and Banner (1977) there was a significant relationship off the Bristol Channel between the density of adult dominant ophiuroids and percent fine material with a weaker relationship between density and organic matter in the sediment. They concluded that distribution of both larvae and adults correlated with the energy distribution of the hydrodynamic regime. In view of the hydrodynamic regime off Martha's Vineyard influencing, on the one hand, deposition in the Mud Patch, and, on the other hand, extensive scouring on

Nantucket Shoals, their findings might be applied to echinoderms and the entire benthic community in the study area.

The association between benthic animals and sediment is not a simple causal relationship (Rhoads 1974). Sediment composition and associated physical properties (grain size, sorting, porosity, mechanical strength) are primarily controlled by geologic processes. In turn, geologic and physical oceanographic processes exert considerable control over chemical properties of sediment (nutrients, oxygen tension, geochemistry). Finally, chemical properties catalyze and interact with biological properties of sediment (algal sheaths, feces, organic film, bacterial and fungal slime). Because of these properties, sediment provides a substratum for colonization, a medium in which reside temporarily or permanently, material for tube and burrow construction, and a source of nutrition. Thus quantitative relationships between benthos and sediment parameters deserve attention; however, because of the varied histories and origins of the sediment parameters, the relationships are not always immediately obvious in terms of their ecological significance. Without quantitative chemical measures of sediment properties (Johnson 1974), sediment might be considered an integrative environmental factor to which the benthos are responding.

Faunal assemblage

Since identification to species level was not always possible, it was not feasible to quantitatively define communities in the study area. Accordingly, the less formal term "assemblage" was used here to designate a recurring group of organisms living within broadly defined and repetitive environmental conditions. Based on studies in the Gulf of Maine and Georges Bank, four major benthic assemblages were tentatively outlined (Wigley¹⁸): Sand fauna, silty-sand fauna, gravel fauna, and muddy basin. Pratt (1973) elaborated on Wigley's scheme and suggested that these assemblages extend along the Middle Atlantic Bight. Characteristic species were recommended for the sand assemblage off the Delmarva Peninsula (Maurer et al. 1976) which confirmed Wigley's (1968) and Pratt's (1973) projections. Examination of sediment data (Appendix Table 1), species list (Appendix Table 2), and the distribution maps presented here indicates that almost half the study area contained the sand fauna (*Echinarachnius parma*, *Crangon septemspinosa*, *Chiridotea tuftsi*, *Pagurus acadianus*, *Leptocuma minor*, *Haustoriidae*, *Phoxocephalus holbolli*, *Paraphoxus* sp., *Lunatia heros*, *Nassarius trivittatus*, *Spisula solidissima*, *Molgula* spp.).

The fauna of the southwestern quadrat and south central portion of this study—the Mud Patch—compares well with the silty-sand fauna recognized earlier by Wigley (1968) from other areas on Georges Bank. The silty-sand bottom and Mud Patch both contained *Havelockia scabra*, ampeliscids, *Dichelopandalus leptocerus*, *Diastylis* spp., *Edotea triloba*, *Scalibregma inflatum*, *Nephtys incisa*, *Cerianthus*, *Nucula* spp., *Nuculana* sp., *Amphioplus* spp., and *Amphilimna olivacea*. A muddy-basin fauna between fishing banks was also identified earlier (Wigley 1968). The deeper stations in this study (7-10, 21, 22, 37, 38, 52, 53, 56, 57), contained species that are in common with the muddy-basin fauna. For example,

¹⁸Wigley, R. L. 1958. Bottom ecology. In Annual Report, U.S.D.I., Bur. Comm. Fish., Woods Hole Laboratory, Woods Hole, Mass., p. 55-58.

Schizaster fragilis, *Ophiura sarsi*, *Ophiura robusta*, *Amphiura otteri*, *Cadulus* spp., *Dentalium* sp., *Sternaspis scutata*, *Amphitrite* sp., *Onuphis* spp., and *Leanira* sp. were characteristic of muddy-basin fauna and the deep stations off Martha's Vineyard.

In the most comprehensive benthic survey of the U.S. Middle Atlantic Shelf, five faunal zones were recognized (Boesch et al. footnote 17). Faunal changes were mainly gradual rather than abrupt. The faunal zones included: Inner shelf (to 30 m), central shelf (38-50 m), outer shelf (50-100 m), shelf break (100-200 m), and continental slope (>200 m). According to Boesch et al. (footnote 17), the inner and central shelf assemblages were relatively similar, and outer shelf assemblages contained both inshore and offshore species overlapping in distribution. In contrast, shelf break and continental slope assemblages were more discrete.

Comparison of the faunal assemblages of the Martha's Vineyard study with Boesch et al. (footnote 17) is difficult for several reasons. The study by Boesch et al. used a 0.5 mm sieve, and their quantitative (cluster analysis) determination of species and site groups emphasized polychaetes and peracarid crustaceans which were missed or deemphasized in earlier studies with coarser sieves.

Another difficulty lies in the presence of the Mud Patch off Martha's Vineyard. Transects from the east side of the study area (Fig. 2) would be more comparable to transects studied by Boesch et al. (footnote 17). The Mud Patch affords the opportunity for colonization in shallower depths by species normally encountered in mud bottoms at deeper depths further out on the central and outer shelf. Coincidental with this expansion into shallower water is the response of deeper dwelling mud-bottom species to a different temperature regime. Species considered characteristic of a zone normally <11°C (Fig. 3) and with a smaller seasonal range would be colonizing a site with a wider temperature fluctuation. It might be expected that the Mud Patch would consist of a fauna containing both inner and outer shelf components.

Based on bathymetry and sediment type, the fauna off Martha's Vineyard could be conveniently arranged into the faunal zones proposed by Boesch et al. (footnote 17). The added complication of the Mud Patch must also be considered. These qualitative comparisons are primarily offered as suggestions for testing rather than as formal community designations.

ACKNOWLEDGMENTS

This account was supported in part by a contract with the National Marine Fisheries Service Ocean Pulse Program, and was monitored by John Pearce who provided considerable support and encouragement.

LITERATURE CITED

- AARON, J. M., B. BUTMAN, M. H. BOTHNER, and R. E. SYLVESTER.
1980. Environmental conditions relating to potential geologic hazards. U.S. Northeastern Atlantic Continental Margin. U.S.G.S. Misc. Field Studies Map MF-1193, 3 sheets.
- ALTON, M. S.
1972. Bathymetric distribution of the echinoderms off the northern Oregon coast. In A. T. Pruter and D. L. Alverson (editors), *The Columbia River Estuary and adjacent ocean waters: Bioenvironmental studies*, p. 475-537. Univ. Wash. Press, Seattle.
- BIERNBAUM, C. K.
1979. Influence of sedimentary factors on the distribution of benthic amphipods of Fishers Island Sound, Connecticut. *J. Exp. Mar. Biol. Ecol.* 38:201-223.
- BIGELOW, H. B.
1927. Physical oceanography of the Gulf of Maine. U.S. Bur. Fish., Bull. 40:511-1027.
- BUMPUS, D. F.
1973. A description of the circulation on the continental shelf of the east coast of the United States. *Prog. Oceanogr.* 6:111-157.
1976. Review of the physical oceanography of Georges Bank. *Int. Comm. Northwest Atl. Fish. Res. Bull.* 12:119-134.
- BUTMAN, B., M. A. NOBLE, R. C. BEARDSLEY, J. A. VERMERSCH, R. A. LIMEBURNER, B. MAGNELL, and R. J. SCHLITZ.
1980. The mean circulation on Georges Bank as measured by moored current meters. *Int. Counc. Explor. Sea ICES C.M./C:33, Hydrogr. Comm.*, 1-9.
- COLTON, J. B., Jr., and R. R. STODDARD.
1973. Bottom-water temperatures on the continental shelf, Nova Scotia to New Jersey. U.S. Dep. Commer., NOAA Tech. Rep. NMFS CIRC-376, 55 p.
- DORKINS, C. A.
1980. Flow along the continental shelf south of Nantucket Island. *Int. Counc. Explor. Sea ICES C.M./C:33, Hydrogr. Comm.*, 13 p.
- DRISCOLL, E. G., and D. E. BRANDON.
1973. Mollusc-sediment relationships in northwestern Buzzards Bay, Massachusetts, U.S.A. *Malacologia* 12:13-46.
- EMERY, K. O.
1960. The sea off Southern California. A modern habitat of petroleum. John Wiley & Sons, N.Y., 366 p.
- FRANZ, D.
1976. Benthic molluscan assemblages in relation to sediment gradients in northeastern Long Island Sound, Connecticut. *Malacologia* 15:377-399.
- GROSSLEIN, M. D., B. E. BROWN, and R. C. HENNEMUTH.
1979. Research, assessment, and management of a marine ecosystem in the northwest Atlantic, a case study. In J. Cairns, Jr., G. P. Patil, and W. E. Waters (editors), *Environmental biomonitoring, assessment, prediction, and management: certain case studies and related quantitative issues*, p. 289-357. International Co-Operative Publ. House, Fairland, Md.
- JOHNSON, R. G.
1974. Particulate matter at the sediment-water interface in coastal environments. *J. Mar. Res.* 32:313-330.
- MAURER, D.
1967a. Filtering experiments on marine pelecypods from Tomales Bay, California. *Veliger* 9:305-309.
1967b. Mode of feeding and diet, and synthesis of studies on marine pelecypods from Tomales Bay, California. *Veliger* 10:72-76.
- MAURER, D., P. KINNER, W. LEATHEM, and L. WATLING.
1976. Benthic faunal assemblage off the Delmarva Peninsula. *Estuarine Coastal Mar. Sci.* 4:163-177.
- MAURER, D., and W. LEATHEM.
1980. Dominant species of polychaetous annelids of Georges Bank. *Mar. Ecol., Prog. Ser.* 3:135-144.
- MILLIMAN, J. D.
1973. Marine geology 10. In S. B. Saila (editor), *Coastal and offshore environmental inventory, Cape Hatteras to Nantucket Shoals*. Univ. Rhode Island Mar. Publ. Ser. 2, 91 p.
- MILLS, E. L.
1980. The structure and dynamics of shelf and slope ecosystems off the northeast Coast of North America. In K. R. Tenore and B. C. Coull (editors), *Marine benthic dynamics*, p. 25-47. Belle W. Baruch Symp. Mar. Sci., No. 11. Univ. S.C. Press, Columbia.
- MILLS, E. L., and R. O. FOURNIER.
1979. Fish production and the marine ecosystems of the Scotian Shelf, Eastern Canada. *Mar. Biol. (Berl.)* 54:101-108.
- PARSONS, T. R., M. TAKAHASHI, and B. HARGRAVE (editors).
1977. Benthic communities. In *Biological oceanographic processes*, 2d ed., p. 176-264. Pergamon Press, N.Y.
- PRATT, S. D.
1973. Benthic fauna 5. In S. B. Saila (editor), *Coastal and offshore environmental inventory, Cape Hatteras to Nantucket Shoals*. Univ. Rhode Island Mar. Publ. Ser. 2, 70 p.
- RHOADS, D. C.
1974. Organism-sediment relations on the muddy sea floor. *Oceanogr. Mar. Biol. Annu. Rev.* 12:263-300.

- ROWE, G. T.
1971. Benthic biomass in the Pisco, Peru upwelling. *Invest. Pesq.* 35:129-135.
- SCHLITZ, R. J.
In press. Processes of upwelling along the northern side and contributions to the nutrient supply of Georges Bank. *Can. J. Fish. Aquat. Sci.*
- SCHLITZ, R. J., W. R. WRIGHT, and T. LAUGHTON.
1977. The Cape Cod Front and Great South Channel. [Abstr.] *EOS, Trans. Am. Geophys. Union* 58(9).
- SHEPARD, F. P.
1954. Nomenclature based on sand-silt-clay ratios. *J. Sediment. Petrol.* 24:151-158.
- SOKOLOVA, M. N.
1972. Trophic structure of deep-sea macrobenthos. *Mar. Biol. (Berl.)* 16:1-12.
- TWENHOFEL, W. H., and S. A. TYLER.
1941. *Methods of study of sediments.* McGraw-Hill, N.Y., 183 p.
- TYLER, P. A., and F. T. BANNER.
1977. The effect of coastal hydrodynamics on the echinoderm distribution in the sublittoral of Oxwich Bay, Bristol Channel. *Estuarine Coastal Mar. Sci.* 5:293-308.
- WIGLEY, R. L.
1961. Benthic fauna of Georges Bank. *Trans. 26th North Am. Wildl. Nat. Resour. Conf.*, p. 310-317.
1963. Pogonophora on the New England continental slope. *Science (Wash., D.C.)* 141:358-359.
1965. Density-dependent food relationships with reference to New England groundfish. *Int. Comm. Northw. Atl. Fish., Spec. Publ.* 6:501-513.
1968. Benthic invertebrates of the New England fishing banks. *Underwater Nat.* 5(1):8-13.
- WIGLEY, R. L., and A. D. MCINTYRE.
1964. Some quantitative comparisons of offshore meiobenthos and macrobenthos south of Martha's Vineyard. *Limnol. Oceanogr.* 9:485-493.
- WIGLEY, R. L., and F. C. STINTON.
1973. Distribution of macroscopic remains of recent animals from marine sediments off Massachusetts. *Fish. Bull., U.S.* 71:1-40.
- WIGLEY, R. L., and R. B. THEROUX.
1981. Macrobenthic invertebrate fauna of the Middle Atlantic Bight Region: Faunal composition and quantitative distribution. *U.S.G.S. Prof. Pap.* 529, Chap. N1-198.
- ZEITZSCHEL, B.
1980. Sediment-water interactions in nutrient dynamics. *In* K. R. Tenore and B. C. Coull (editors), *Marine benthic dynamics*, p. 195-218. *Belle W. Baruch Symp. Mar. Sci.*, No. 11. Univ. S.C. Press, Columbia.

Appendix Table 1. Station Location and Environmental Variables off Martha's Vineyard, Massachusetts.

Station	Station location		Water depth m	Water temperature °C	Type	Bottom Sediment Parameters						
						Median	Composition			Nitrogen	Organic	Carbon
	Lat. (N)	Long. (W)				diameter φ	Sand %	Silt %	Clay %	(Kjeldahl) %	carbon %	Nitrogen
1	40°58'	69°30'	46	8.6	Sand-gravel	--	--	--	--	--	--	--
2	40°51'	69°31'	46	8.6	Sand	1.00	100.0	0.0	0.0	0.003	0.050	17.0
3	40°40'	69°31'	51	7.7	Sand	2.35	100.0	0.0	0.0	0.010	0.11	11.1
4	40°30'	69°29'	62	7.1	Sand	2.54	100.0	0.0	0.0	0.019	0.13	6.9
5	40°21'	69°30'	76	9.4	Sand	3.47	81.0	12.7	6.3	0.070	0.46	6.6
6	40°10'	69°31'	91	8.6	Sand	2.23	90.0	10.0	0.0	0.034	0.26	7.7
7	40°00'	69°30'	128	11.9	Sand	2.79	84.5	10.2	5.3	0.031	0.24	7.8
8	39°57'	69°30'	183	11.9	Sand	2.89	86.0	9.8	4.2	0.022	0.12	5.5
9	39°56'	69°45'	201	8.9	Sand	2.70	85.0	10.7	4.2	0.035	0.22	6.3
10	40°00'	69°45'	139	11.7	Silty-Sand	3.85	57.0	33.4	9.5	0.037	0.61	7.0
11	40°10'	69°45'	95	7.3	Silty-Sand	3.70	69.0	23.6	7.5	0.064	0.41	6.5
12	40°20'	69°46'	79	6.7	Sand	3.48	85.5	9.2	5.4	0.059	0.41	7.0
13	40°30'	68°45'	73	7.2	Sand	2.63	95.0	1.9	3.2	0.026	0.17	6.6
14	40°40'	69°45'	59	7.5	Sand	2.24	100.0	0.0	0.0	0.004	0.03	7.9
15	40°50'	69°45'	37	9.2	Sand	2.05	100.0	0.0	0.0	0.003	0.02	8.0
16	40°46'	70°00'	38	8.1	Sand	2.67	100.0	0.0	0.0	0.018	0.03	4.6
17	40°39'	69°59'	49	7.5	Sand	3.27	81.0	10.3	8.7	0.072	0.39	5.5
18	40°30'	70°00'	73	6.8	Sand	3.65	67.0	25.3	7.7	0.082	0.55	6.8
19	40°20'	49°59'	91	6.7	Sand	3.79	62.0	31.4	6.7	0.069	0.48	6.9
20	40°10'	70°00'	117	11.4	Sand	4.12	45.0	44.8	10.2	0.080	0.58	7.2
21	40°00'	70°00'	165	(11.4)	Sand	3.35	66.0	23.9	10.1	0.058	0.48	8.4
22	40°03'	70°15'	183	10.8	Silty-sand	2.18	66.8	23.6	9.6	0.060	0.59	9.9
23	40°10'	70°15'	113	11.6	Silty-sand	4.31	36.2	53.4	10.4	0.092	0.70	7.6
24	40°20'	70°15'	90	7.8	Silty-sand	4.12	47.0	38.9	14.1	0.133	0.92	6.9
25	40°30'	70°15'	70	6.6	Sand	3.83	58.8	30.7	10.5	0.076	0.52	6.8
26	40°40'	70°15'	51	7.2	Sand	3.37	85.9	10.2	3.8	0.048	0.19	4.0
27	40°50'	70°15'	44	8.1	Sand	3.43	86.1	9.3	4.5	0.035	0.20	5.7
28	41°00'	70°15'	33	9.4	Sand	3.35	90.7	4.2	4.7	0.033	0.13	4.0
29	41°11'	70°16'	27	12.2	Sand	1.35	100.0	0.0	0.0	0.011	0.07	6.9
30	41°10'	70°30'	38	8.3	Sand	2.03	100.0	0.0	0.0	0.018	0.11	6.2
31	41°00'	70°30'	48	7.5	Sand	2.81	79.5	14.1	6.4	0.044	0.28	6.3
32	40°50'	70°30'	59	6.9	Sand	3.30	67.5	24.1	3.5	0.073	0.51	7.0
33	40°40'	70°30'	62	6.7	Silty-Sand	3.48	60.0	25.3	14.7	0.137	0.84	6.2
34	40°30'	70°30'	73	7.5	Sandy-silt	4.54	33.2	49.0	17.8	0.148	1.03	6.9
35	40°20'	70°30'	97	10.0	Sandy-silt	4.67	12.3	72.3	15.4	0.146	1.12	7.6

Appendix Table 1 (cont.)

Station	Station location		Water depth m	Water temperature °C	Type	Bottom Sediment Parameters						
						Median diameter φ	Composition			Nitrogen (Kjeldahl)	Organic carbon	Carbon Nitrogen
	%	%				%	%	%	%			
36	40°10'	70°30'	128	12.1	Silty-sand	2.26	82.8	10.2	7.0	0.046	0.41	8.9
37	40°04'	70°29'	220	9.4	Sand	1.79	88.8	6.2	5.0	0.038	0.28	7.5
38	40°02'	70°44'	194	10.8	Silty-sand	3.61	53.6	31.8	14.6	0.113	0.63	5.6
39	40°10'	70°45'	132	11.7	Silty-sand	5.02	14.5	66.4	19.1	0.183	1.05	5.7
40	40°20'	70°46'	106	9.2	Sand-silt-clay	5.01	14.0	61.6	24.5	0.212	1.46	6.9
41	40°30'	70°45'	79	6.7	Sandy-silt	4.86	19.3	62.6	18.1	0.194	1.03	5.3
42	40°40'	70°45'	66	6.4	Silty-sand	3.42	55.8	29.7	14.5	0.112	0.69	6.2
43	40°50'	70°45'	55	6.8	Sand	2.37	85.5	7.8	6.7	0.054	0.22	4.1
44	41°00'	70°45'	51	6.7	Sand	1.48	100.0	0.0	0.0	0.008	0.07	8.7
45	41°10'	70°45'	38	8.6	Sand-gravel	1.00	100.0	0.0	0.0	0.006	0.06	10.6
46	41°10'	71°00'	40	9.7	Sand	1.59	100.0	0.0	0.0	0.007	0.08	11.2
47	41°00'	71°00'	51	6.4	Sand-gravel	2.53	85.1	6.8	8.1	-	-	-
48	40°50'	71°00'	59	6.1	Sand	2.30	83.4	8.5	7.7	0.046	0.23	5.0
49	40°40'	71°00'	70	6.3	Sandy-silt	5.09	16.4	61.0	22.6	0.220	1.09	5.0
50	40°30'	71°00'	84	6.3	Clayey-silt	5.84	3.1	69.3	27.6	0.245	1.24	5.0
51	40°21'	71°00'	99	9.4	Sandy-silt	4.77	31.2	49.0	19.8	0.150	0.94	6.3
52	40°10'	71°00'	146	10.8	Silty-sand	2.35	70.0	17.5	11.5	0.067	0.45	6.7
53	40°06'	71°00'	179	10.8	Silty-sand	4.77	24.1	54.1	21.8	0.124	1.04	8.4
54	39°59'	71°00'	366	(6.1)	Silt	5.16	25.2	66.8	8.0	0.112	0.88	7.9
55	39°56'	71°00'	567	(6.1)	Silt	4.39	42.7	36.3	21.0	0.160	1.17	7.3
56	40°03'	71°16'	183	10.3	Sand	2.18	86.0	8.4	5.6	0.050	0.21	4.1
57	40°10'	71°15'	110	10.8	Silty-sand	2.68	66.0	23.6	10.4	0.099	0.74	7.5
58	40°20'	71°15'	91	9.0	Silty-sand	3.29	52.0	35.3	12.7	0.136	0.81	5.9
59	40°30'	71°15'	77	6.7	Silty-sand	2.95	69.5	19.5	11.0	0.086	0.32	3.7
60	40°40'	71°15'	62	6.2	Sand	2.38	76.6	16.1	7.3	0.060	0.29	4.9
61	40°50'	71°15'	62	6.4	Sand	3.00	87.9	6.5	5.6	0.044	0.20	4.5
62	41°01'	71°16'	48	6.7	Sand	1.63	100.0	0.0	0.0	0.009	0.06	7.1
63	41°10'	71°15'	38	8.1	Sand	2.14	100.0	0.0	0.0	0.003	0.06	7.9
64	41°00'	71°30'	55	6.9	Sand	2.38	89.4	6.1	4.5	0.042	0.24	5.6

Porifera	Amphipoda (Cont.)	Pelecypoda (Cont.)
	<i>Casco bigelowi</i>	<i>Phacoides filiosus</i>
	<i>Corophium</i> sp.	<i>Placopecten magellanicus</i>
Cnidaria	<i>Dulichia</i> sp.	<i>Siliqua costata</i>
Hydrozoa	<i>Eriopisa elongata</i>	<i>Spisula solidissima</i>
<i>Hydractinia echinata</i>	<i>Harpinia propinqua</i>	<i>Tellina agilis</i>
Anthozoa	Haustoriidae	<i>Thracia</i> sp.
<i>Cerianthus</i> sp.	<i>Hippomedon serratus</i>	<i>Thyasira ferruginosa</i>
<i>Edwardsia</i> sp.	<i>Lembos</i> sp.	<i>Thyasira gouldi</i>
<i>Epizoanthus americanus</i>	<i>Leptocheimius pinguis</i>	<i>Thyasira ovata</i>
<i>Pennatulaculeata</i>	<i>Orchomenella groenlandica</i>	<i>Thyasira trisinuata</i>
<i>Stylatula elegans</i>	<i>Paraphoxus</i> sp.	<i>Venericardia borealis</i>
	<i>Photis macrocoxa</i>	<i>Yoldia sapotilla</i>
Nemertea	<i>Phoxocephalus holbolli</i>	Scaphopoda
	<i>Ptotomedia fasciata</i>	<i>Cadulus pandionis</i>
Annelida	<i>Siphonocetes smithianus</i>	<i>Cadulus vermilli</i>
<i>Aglaphamus circinata</i>	<i>Stenopleustes gracilis</i>	<i>Dentalium occidentale</i>
<i>Ammotrypane aulogaster</i>	<i>Unciola irrorata</i>	Gastropoda
<i>Amphitrite</i> sp.	<i>Unciola leucopsis</i>	<i>Alvania carinata</i>
<i>Ancistrostylis</i> sp.	Decapoda	<i>Buccinum undatum</i>
<i>Aphrodita hastata</i>	<i>Axius serratus</i>	<i>Colus stimpsoni</i>
<i>Arabella iricolor</i>	<i>Cancer borealis</i>	<i>Crepidula plana</i>
<i>Aricidea jeffreysii</i>	<i>Cancer irroratus</i>	<i>Crucibulum striatum</i>
<i>Asychis biceps</i>	<i>Catapagurus sharperi</i>	<i>Cylichna alba</i>
<i>Brada</i> sp.	<i>Crangon septemspinosa</i>	<i>Epitonium dallianum</i>
<i>Capitella</i> sp.	<i>Bythocaris nana</i>	<i>Lunatia heros</i>
<i>Ceratocephale loveni</i>	<i>Dichelopandalus leptocerus</i>	<i>Lunatia triseriata</i>
<i>Chaetozone</i> sp.	<i>Europrognatha rastellifera</i>	<i>Nassarius trivittatus</i>
<i>Chone infundibuliformis</i>	<i>Hyas coarctatus</i>	<i>Neptunea</i> sp.
<i>Coosura longocirrata</i>	<i>Munida iris</i>	<i>Polinices</i> sp.
<i>Drilonereis longa</i>	<i>Pagurus acadianus</i>	<i>Retusa gouldi</i>
<i>Eunice pennata</i>	<i>Pagurus arcuatus</i>	<i>Scaphander</i> sp.
<i>Flabelligera</i> sp.	<i>Pagurus politus</i>	
<i>Glycera robusta</i>	<i>Pontophilus brevirostris</i>	Echinodermata
<i>Glycera tessellata</i>	Isopoda	Asteroidea
<i>Goniada brumnea</i>	<i>Calathura</i> sp.	<i>Asterias vulgaris</i>
<i>Goniada maculata</i>	<i>Chiridotea tuftsi</i>	<i>Astropecten americanus</i>
<i>Harmothoe extenuata</i>	<i>Cirolana polita</i>	<i>Astropecten</i> sp.
<i>Hyalinoecia tubicola</i>	<i>Edotea triloba</i>	<i>Henricia sanguinolenta</i>
<i>Laonice cirrata</i>	<i>Ptilanthura tenuis</i>	<i>Leptasterias tenera</i>
<i>Leanira</i> sp.	Cumacea	<i>Porania</i> sp.
<i>Lumbrineris fragilis</i>	<i>Diastylis polita</i>	Echinoidea
<i>Lumbrineris tenuis</i>	<i>Diastylis quadrispinosa</i>	<i>Schizaster fragilis</i>
<i>Melinna cristata</i>	<i>Eudorella emarginata</i>	<i>Echinarachnius parma</i>
<i>Nephtys buccera</i>	<i>Eudorellopsis</i> sp.	Ophiuroidea
<i>Nephtys incisa</i>	<i>Leptocoma minor</i>	<i>Amphilima olivacea</i>
<i>Nereis pelagica</i>	<i>Leptostylis</i> sp.	<i>Amphioplus abditus</i>
<i>Ninoe nigripes</i>	<i>Petalosarsia declivis</i>	<i>Amphioplus macilentus</i>
<i>Notocirrus</i> sp.	Mysidacea	<i>Amphiura fragilis</i>
<i>Omphis conchylega</i>	<i>Bathymysis renoculata</i>	<i>Amphiura otteri</i>
<i>Omphis opalina</i>	<i>Erythrope erythroptalma</i>	<i>Axiognathus squamatus</i>
<i>Omphis quadricuspis</i>	<i>Hypererythrope caribbaea</i>	<i>Ophiura robusta</i>
<i>Orbinia ornata</i>	<i>Mysis mixta</i>	<i>Ophiura sarsi</i>
<i>Owenia</i> sp.	<i>Neomysis americana</i>	Holothuroidea
<i>Paradiopatra</i> sp.	Cirripedia	<i>Caudina arenata</i>
<i>Paramphinoe pulchella</i>	<i>Balanus</i> sp.	<i>Chiridota</i> sp.
<i>Paraonis neopolitana</i>	Pycnogonida	<i>Cucumaria frondosa</i>
<i>Phyllococe mucosa</i>	<i>Achelia spinosa</i>	<i>Havelockia scabra</i>
<i>Prionospio</i> sp.	<i>Paranymphe spinosum</i>	<i>Molpadia oolitica</i>
<i>Scalibregma inflatum</i>	Mollusca	<i>Psolus fabricii</i>
<i>Sphaerodorium gracilis</i>	Amphineura	<i>Stereoderma unisemita</i>
<i>Spio</i> sp.	<i>Chaetoderma nitidulum</i>	Bryozoa
<i>Spirochaetopterus</i> sp.	Pelecypoda	<i>Dendrobeania murrayana</i>
<i>Spiophanes bombyx</i>	<i>Anomia</i> sp.	<i>Electra hastingsae</i>
<i>Sternaspis scutata</i>	<i>Arctica islandica</i>	<i>Electra pilosa</i>
<i>Sthenelais limicola</i>	<i>Astarte undata</i>	<i>Haplota clavata</i>
<i>Streblosoma spiralis</i>	<i>Bathyarca pectunculoides</i>	<i>Hippothoa hyalina</i>
<i>Tharyx</i> sp.	<i>Cerastoderma pinnulatum</i>	<i>Semparia chelata</i>
Sipunculida	<i>Crenella glandula</i>	Ascidacea
<i>Golfingia catherinae</i>	<i>Cuspidaria perrostrata</i>	<i>Bostrichobranchus</i>
<i>Golfingia elongata</i>	<i>Cuspidaria striata</i>	<i>pilularis</i>
<i>Golfingia margaritacea</i>	<i>Ensis directus</i>	<i>Ciona intestinalis</i>
<i>Golfingia mirata</i>	<i>Hiatella</i> sp.	<i>Cnemidocarpa mollis</i>
<i>Golfingia (Phascoioides)</i> sp.	<i>Lyonsia arenosa</i>	<i>Heterostigma singulare</i>
<i>Onchseoma steenstrupi</i>	<i>Lyonsia hyalina</i>	<i>Molgula citrina</i>
<i>Phascolion strombi</i>	<i>Macao calcarea</i>	<i>Molgula complanata</i>
Arthropoda	<i>Mesodesma arctatum</i>	<i>Molgula siphonalis</i>
Amphipoda	Mytilidae	Pogonophora
<i>Aeginina longicornis</i>	<i>Nucula proxima</i>	<i>Siboglinum atlanticum</i>
<i>Ampelisca compressa</i>	<i>Nucula tenuis</i>	<i>Siboglinum ekmani</i>
<i>Ampelisca macrocephala</i>	<i>Nuculana acuta</i>	
<i>Anonyx</i> sp.	<i>Pandora gouldiana</i>	Hemichordata
<i>Byblis serrata</i>	<i>Pandora inflata</i>	Enteropneusta
<i>Caprella</i> sp.	<i>Periploma papyratium</i>	<i>Balanoglossus</i> sp.
	<i>Phacoides blakeanus</i>	

Appendix Table 3. Wet Weight Biomass (g/m²) of Major Taxa of Macrobenthic Invertebrates per Station off Martha's Vineyard, Massachusetts.

Station Number	PORIFERA	COELENTERATA	NEMERTEA	ANNELIDA	POGONOPHORA	SIPUNCULA	MOLLUSCA	Gastropoda	Pelecypoda	Scaphapoda
1	12.0	1.1	8.2	22.6	-	-	45.2	0.5	44.7	-
2	2.9	-	-	3.1	-	-	-	-	-	-
3	-	-	0.2	0.5	-	-	928.4	-	928.4	-
4	-	18.7	3.1	68.5	-	11.9	934.8	392.7	542.1	-
5	-	0.9	1.7	45.4	-	1.2	272.5	0.1	272.4	-
6	-	1.9	0.2	32.5	-	-	47.5	-	47.5	-
8	-	6.2	-	11.7	-	<0.1	-	-	-	-
9	0.9	10.9	0.1	14.8	-	0.1	6.8	-	5.7	1.1
10	-	10.5	0.3	4.7	-	0.1	3.0	-	3.0	-
11	-	11.6	8.9	10.6	-	2.1	6.0	0.1	5.9	-
12	-	5.0	1.5	26.7	-	0.1	1413.0	0.1	1412.9	-
13	-	0.3	3.8	118.8	-	0.2	65.6	1.0	64.6	-
14	-	0.1	0.2	13.3	-	5.4	0.2	0.1	0.1	-
15	-	-	0.1	6.8	-	-	0.1	-	0.1	-
16	-	-	-	2.6	-	-	1.4	-	1.4	-
17	-	0.6	0.4	44.9	-	3.6	545.2	9.1	536.1	-
18	-	0.1	0.6	29.4	-	0.1	496.3	0.2	469.1	-
19	-	-	1.4	8.8	-	0.1	1.2	-	1.2	-
20	-	342.3	-	1.3	-	0.1	0.8	-	0.8	-
21	-	-	-	4.9	-	0.1	6.4	-	6.4	-
22	-	-	-	5.7	-	4.1	0.2	-	0.2	-
23	-	1.9	2.0	15.6	-	0.2	2.2	-	2.2	-
24	-	3.8	20.0	15.3	-	-	0.3	-	0.3	-
25	-	-	1.3	20.8	-	-	9.4	0.1	9.3	-
26	-	0.1	2.5	73.4	-	0.1	7.2	0.1	7.1	-
27	-	0.1	3.1	72.7	-	-	77.8	-	77.8	-
28	-	0.1	1.8	83.9	-	11.0	933.1	1.0	932.1	-
29	-	5.2	8.5	60.2	-	-	44.4	0.1	44.3	-
30	-	-	4.1	3.4	-	11.9	808.0	-	808.0	-
31	-	2.1	0.3	35.7	-	10.1	23.2	10.6	12.6	-
32	-	-	0.4	15.9	-	-	3.4	0.1	3.3	-
33	-	-	0.3	31.3	-	-	7.0	0.1	6.9	-
34	-	2.0	0.1	15.9	-	1.4	5.3	0.2	5.1	-
35	-	7.4	2.0	21.9	-	3.7	4.2	0.3	3.9	-
36	-	18.7	0.7	15.3	-	-	0.9	-	0.9	-
37	-	-	2.7	2.6	-	-	1.6	0.1	1.5	-
38	-	1.4	-	24.1	-	0.1	0.3	0.1	0.1	0.1
39	-	-	1.1	10.0	-	-	3.6	0.1	3.5	-
40	-	5.3	0.1	2.4	-	-	12.5	0.1	12.4	-
41	-	5.6	-	10.3	-	-	0.1	-	0.1	-
42	-	0.9	0.1	32.6	-	-	0.1	0.1	-	-
43	-	-	0.1	22.4	-	-	23.0	0.1	22.9	-
44	-	-	0.2	44.0	-	-	0.8	0.7	0.1	-
45	-	-	-	37.6	-	-	-	-	-	-
46	-	-	0.1	11.8	-	-	1.6	0.3	1.3	-
47	-	-	1.0	10.3	-	-	-	-	-	-
48	-	-	0.1	49.2	-	-	1326.5	0.5	1326.0	-
49	-	0.5	0.1	38.5	-	-	11.1	0.2	10.9	-
50	-	1.5	-	48.1	-	-	3.1	0.1	3.0	-
51	-	17.8	0.1	53.8	-	0.1	9.5	0.1	9.4	-
52	-	9.9	0.1	25.3	-	-	13.9	0.1	13.8	-
53	0.2	2.9	0.1	8.4	-	0.1	6.6	-	6.6	-
54	-	0.3	-	13.2	0.1	13.4	1.2	0.2	0.9	0.1
55	-	-	-	8.3	0.2	-	2.2	0.1	2.1	-
56	-	1.3	1.1	8.7	-	-	4.3	0.1	4.2	-
57	-	1.8	-	-	-	0.1	-	-	-	-
58	-	7.9	0.6	19.2	-	-	15.3	1.0	14.3	-
59	-	8.8	0.1	23.4	-	-	5.1	-	5.1	-
60	-	-	0.2	114.7	-	6.8	24.3	0.2	24.1	-
61	-	-	0.2	40.5	-	0.1	0.9	0.2	0.7	-
62	-	0.1	-	14.8	-	7.6	7.9	7.8	0.1	-
63	-	-	-	28.2	-	0.1	-	-	-	-
64	-	-	-	21.0	-	13.1	622.5	0.1	622.4	-

Appendix Table 3 (cont.)

Station Number	CRUSTACEA	Cumacea	Isopoda	Amphipoda	Decapoda	ECHINODERMATA	Holothuroidea	Echinoidea	Ophiuroidea	Asteroidea	ECTOPROCTA	ENTEROPNEUSTA	Ascidiacea	TOTAL
1	18.9	-	-	3.6	15.3	-	-	-	-	-	51.6	-	-	159.6
2	2.2	-	0.1	2.0	0.1	-	-	-	-	-	1.1	-	-	9.3
3	1.2	0.1	0.1	1.0	-	2.7	-	2.7	-	-	-	-	0.1	933.1
4	9.2	0.2	0.1	8.8	0.1	11.8	-	11.8	-	-	4.7	-	1.9	1064.6
5	52.5	0.2	4.3	40.9	7.1	1.8	-	0.9	-	0.9	-	-	-	376.0
6	18.9	0.1	0.1	12.5	6.2	7.4	3.5	-	3.9	-	-	-	4.0	112.4
8	0.2	0.1	-	0.1	-	105.3	-	-	105.3	-	-	-	-	123.4
9	0.7	-	0.1	0.6	-	1.1	-	-	1.1	-	-	-	1.2	36.6
10	0.2	0.1	-	0.1	-	86.4	-	38.4	34.1	13.9	0.1	-	-	105.3
11	1.9	0.1	0.7	1.1	-	71.4	25.0	-	46.4	-	-	-	7.1	119.6
12	16.9	0.1	0.1	16.7	-	12.3	12.2	-	0.1	-	-	10.2	-	1485.7
13	32.3	0.7	3.2	27.9	0.5	0.5	0.5	-	-	-	-	-	1.5	223.0
14	2.4	0.1	0.2	2.1	-	20.9	-	20.7	0.1	0.1	0.1	-	0.1	42.7
15	4.0	-	-	3.9	0.1	-	-	-	-	-	-	-	-	11.0
16	4.5	0.1	0.1	4.3	-	0.6	-	0.6	-	-	0.1	-	7.7	16.9
17	14.8	0.1	0.2	14.5	-	7.8	-	7.8	-	-	-	-	3.1	620.4
18	13.5	0.1	1.1	12.3	-	-	-	-	-	-	-	-	0.2	513.2
19	8.4	-	-	8.4	-	615.9	562.4	-	21.6	31.9	-	-	0.5	636.3
20	0.2	0.1	-	0.1	-	15.3	-	-	12.7	2.6	-	-	-	360.0
21	0.2	0.1	-	0.1	-	284.0	-	251.6	32.3	0.1	-	-	-	295.6
22	0.1	-	-	0.1	-	86.6	10.2	-	76.4	-	-	-	-	96.7
23	0.3	0.1	0.1	0.1	-	38.5	12.8	-	15.6	10.1	-	-	11.0	71.7
24	2.1	-	-	2.1	-	188.9	175.5	-	13.4	-	-	-	-	230.4
25	5.6	0.2	-	1.9	3.5	-	-	-	-	-	-	-	-	37.1
26	12.2	0.2	0.1	11.9	-	-	-	-	-	-	0.1	-	-	95.6
27	20.0	0.2	1.1	18.7	-	2.4	-	2.4	-	-	-	-	-	176.1
28	7.8	0.1	2.6	4.9	0.2	4.7	-	4.7	-	-	0.1	-	-	1042.5
29	9.8	0.2	0.2	8.3	1.1	-	-	-	-	-	16.5	-	22.4	167.0
30	15.5	1.1	1.3	13.1	-	9.1	-	9.1	-	-	-	-	2.8	854.8
31	63.4	0.2	0.7	62.5	-	7.1	-	6.3	-	0.8	4.0	-	0.2	146.1
32	0.4	0.1	-	0.3	-	0.5	-	-	-	0.5	-	-	-	20.6
33	1.1	0.2	0.1	1.8	-	455.0	454.9	0.1	-	-	-	-	-	494.7
34	6.1	0.1	0.1	5.8	0.1	216.0	211.5	-	4.5	-	-	-	-	246.8
35	1.6	0.1	-	1.5	-	108.7	66.8	-	41.9	-	-	-	-	149.5
36	0.3	0.1	-	0.2	-	33.9	-	-	33.9	-	-	-	37.6	107.4
37	0.1	-	-	0.1	-	244.6	8.4	177.1	59.1	-	-	-	0.2	251.8
38	2.9	-	2.4	0.5	-	8.4	8.2	-	0.2	-	-	-	-	37.2
39	0.1	-	-	0.1	-	74.5	-	32.9	41.6	-	-	-	7.2	96.5
40	5.0	0.1	0.1	4.8	-	58.7	2.4	23.0	33.3	-	-	-	7.6	91.6
41	5.8	0.1	-	5.7	-	37.5	-	-	1.0	36.5	-	-	5.6	64.9
42	3.2	0.1	-	3.1	-	-	-	-	-	-	-	-	-	36.9
43	14.9	0.1	0.1	14.7	-	0.1	-	-	0.1	-	-	-	-	60.5
44	23.2	0.1	2.6	20.5	-	-	-	-	-	-	-	-	-	68.2
45	16.4	0.1	0.1	16.2	-	-	-	-	-	-	-	-	49.7	103.7
46	23.3	0.3	0.4	22.6	-	0.1	-	0.1	-	-	-	-	5.4	42.3
47	13.0	0.3	3.9	8.8	-	-	-	-	-	-	-	-	1.0	25.3
48	41.0	0.4	0.8	37.7	2.1	0.1	-	-	-	0.1	-	-	-	1416.9
49	3.7	-	-	3.7	-	0.1	-	-	-	0.1	-	-	-	54.0
50	0.2	0.1	-	0.1	-	183.2	138.7	-	11.9	32.6	-	-	32.8	268.9
51	3.1	0.1	0.1	2.9	-	33.5	15.8	-	17.7	-	-	5.3	10.1	133.3
52	0.1	-	-	0.1	-	10.2	3.9	-	6.3	-	-	2.0	-	61.5
53	0.1	-	-	0.1	-	61.0	23.7	-	37.3	-	-	-	-	79.4
54	0.3	-	0.1	0.2	-	3.4	2.6	-	0.8	-	-	-	-	31.9
55	0.1	-	-	0.1	-	10.3	-	-	10.3	-	-	0.1	-	21.2
56	0.2	-	0.1	0.1	-	17.0	-	-	8.7	8.3	-	-	-	32.6
57	0.2	0.1	-	0.1	-	31.7	10.2	-	21.5	-	-	-	8.8	42.6
58	0.3	-	-	0.3	-	72.8	-	-	22.2	50.6	-	-	3.3	119.4
59	0.6	-	-	0.6	-	2.8	-	-	2.8	-	-	-	9.7	50.5
60	26.6	0.1	0.1	26.4	-	-	-	-	-	-	-	-	15.1	187.7
61	69.7	0.4	0.1	69.2	-	0.1	-	-	-	0.1	-	-	-	111.5
62	17.6	0.1	3.7	13.8	-	0.1	-	0.1	-	-	-	-	1.7	49.8
63	5.6	0.1	0.3	5.2	-	56.3	-	56.2	0.1	-	-	-	-	90.2
64	12.8	0.4	0.1	12.1	0.2	0.5	-	0.5	-	-	-	-	0.4	670.3

Appendix Table 4. Number of Individuals per m² of Major Taxa Per Station.

Station Number	FORIFERA	COELENTERATA	NEMERTEA	ANNELIDA	POGONOPHORA	SIPUNCULIDA	MOLLUSCA	Gastropoda	Pelecypoda	Scaphopoda	CRUSTACEA	Cumacea	Isopoda	Amphipoda	Decapoda	ECHINODERMATA	Holothuroidea	Echinoidea	Ophiuroidea	Asteroidea	ECTOPROCTA	ENTEROPNEUSTA	ASCIDACEA	TOTAL
1	11	11	11	1,145	-	-	22	11	11	-	1,192	-	-	1,171	21	-	-	-	-	-	1,932	-	-	4,324
2	11	-	-	53	-	-	-	-	-	-	73	-	31	31	11	-	-	-	-	-	11	-	-	148
3	-	-	32	42	-	-	11	-	11	-	170	11	21	138	-	21	-	21	-	-	-	-	111	287
4	-	144	21	1,360	-	41	37	11	26	-	2,379	82	36	2,251	10	135	-	135	-	-	11	-	111	4,139
5	-	16	16	1,129	-	11	179	16	163	-	10,957	47	21	10,868	21	32	-	11	-	21	-	-	-	12,340
6	-	10	62	955	-	-	279	-	279	-	8,970	16	27	8,916	11	225	36	-	189	-	-	-	62	10,563
8	-	21	-	572	-	62	114	-	114	-	31	10	-	21	-	489	-	-	489	-	-	-	-	1,289
9	11	140	11	483	-	11	166	-	123	43	222	-	11	211	-	32	-	-	32	-	-	-	11	1,087
10	-	51	10	170	-	10	61	-	61	-	36	10	-	26	-	379	-	10	339	30	10	-	-	727
11	-	23	28	500	-	28	491	12	479	-	564	35	11	518	-	1,096	81	-	1,015	-	-	-	23	2,753
12	-	10	15	678	-	10	159	10	149	-	9,353	10	10	9,333	-	41	31	-	10	-	-	51	-	10,317
13	-	10	10	1,346	-	20	255	10	245	-	19,083	71	10	18,992	10	30	10	-	20	-	-	-	41	20,795
14	-	10	26	820	-	31	20	10	10	-	371	37	37	297	-	412	-	392	10	10	10	-	10	1,710
15	-	-	10	26	-	-	21	-	21	-	524	-	-	514	10	-	-	-	-	-	-	-	-	581
16	-	-	-	456	-	-	21	-	21	-	631	16	32	583	-	11	-	11	-	-	11	-	196	1,326
17	-	10	21	1,082	-	26	115	47	68	-	3,031	62	62	2,907	-	172	-	172	-	-	-	-	31	4,488
18	-	10	15	651	-	10	308	40	268	-	3,030	20	10	3,000	-	-	-	-	-	-	-	-	10	4,034
19	-	-	20	394	-	51	230	-	230	-	2,708	-	-	2,708	-	861	200	-	651	10	-	-	10	4,274
20	-	20	-	41	-	18	27	-	27	-	84	12	-	72	-	344	-	10	324	10	-	-	-	534
21	-	-	-	172	-	20	153	-	153	-	25	10	-	15	-	335	-	15	310	10	-	-	-	705
22	-	-	-	199	-	61	71	-	71	-	41	-	-	41	-	416	10	-	406	-	-	-	-	788
23	-	35	20	379	-	101	81	-	81	-	50	20	20	10	-	576	10	-	556	10	-	-	51	1,293
24	-	20	30	318	-	-	25	-	25	-	111	-	-	111	-	283	91	-	192	-	-	-	-	787
25	-	-	25	838	-	-	576	20	556	-	343	20	-	313	10	-	-	-	-	-	-	-	-	1,782
26	-	10	15	1,184	-	10	722	20	702	-	1,312	66	10	1,236	-	-	-	-	-	-	10	-	-	3,263
27	-	31	10	750	-	-	357	-	357	-	1,999	56	20	1,923	-	15	-	15	-	-	-	-	-	3,162
28	-	10	10	1,225	-	21	82	10	72	-	1,737	10	222	1,484	21	16	-	16	-	-	10	-	-	3,111
29	-	20	21	867	-	-	82	20	62	-	1,326	36	42	1,238	10	22	-	22	-	-	31	-	541	2,910
30	-	-	16	993	-	48	37	-	37	-	1,409	595	56	758	-	22	-	22	-	-	-	-	49	2,574
31	-	10	10	833	-	10	239	56	183	-	3,883	137	15	3,731	-	20	-	10	-	10	10	-	10	5,025
32	-	-	25	247	-	-	101	10	91	-	76	10	-	66	-	10	-	-	-	10	-	-	-	459
33	-	-	15	490	-	-	457	354	121	-	152	51	20	81	-	20	10	10	-	-	-	-	-	1,152
34	-	30	10	227	-	20	162	51	111	-	853	35	10	788	20	112	56	-	56	-	-	-	-	1,414
35	-	30	20	379	-	40	1,030	404	626	-	131	20	-	111	-	576	71	-	505	-	-	-	-	2,206
36	-	76	10	394	-	-	30	-	30	-	50	10	-	40	-	424	-	-	424	-	-	-	333	1,317
37	-	-	10	61	-	-	131	10	121	-	30	-	-	30	-	368	40	10	318	-	-	-	20	620

Appendix Table 4 (cont.)

Station Number	FORIFERA	COELENTERATA	NEMERTEA	ANNELIDA	POGONOPHORA	SIPUNCULIDA	MOLLUSCA	Gastropoda	Pelecypoda	Schaphopoda	CRUSTACEA	Cumacea	Isopoda	Amphipoda	Decapoda	ECHINODERMATA	Holothuroidea	Echinoidea	Ophiuroidea	Asteroidea	ECTOPROCTA	ENTEROPNEUSTA	ASCIDACEA	TOTAL
38	-	42	-	965	-	62	82	10	62	10	139	-	10	129	-	20	10	-	10	-	-	-	-	1,310
39	-	-	10	177	-	-	91	20	71	-	20	-	-	20	-	717	-	10	707	-	-	-	106	1,121
40	-	51	10	61	-	-	570	30	540	-	166	10	10	146	-	1,606	20	778	808	-	-	-	51	2,512
41	-	30	-	182	-	-	51	-	51	-	1,393	40	-	1,353	-	30	-	-	20	10	-	-	81	1,767
42	-	10	10	1,020	-	-	133	102	31	-	1,571	10	-	1,561	-	-	-	-	-	-	-	-	-	2,744
43	-	-	10	763	-	-	46	31	15	-	1,960	10	10	1,940	-	10	-	-	10	-	-	-	-	2,789
44	-	-	10	959	-	-	46	20	26	-	2,800	31	102	2,667	-	-	-	-	-	-	-	-	-	3,815
45	-	-	-	841	-	-	-	-	-	-	5,623	10	10	5,603	-	-	-	-	-	-	-	-	98	6,562
46	-	-	31	619	-	-	58	21	37	-	4,381	68	21	4,292	-	16	-	16	-	-	-	-	73	5,178
47	-	-	20	444	-	-	-	-	-	-	3,075	30	81	2,964	-	-	-	-	-	-	-	-	20	3,559
48	-	-	16	1,804	-	-	198	73	125	-	3,651	276	16	3,338	21	16	-	-	-	16	-	-	-	5,685
49	-	15	10	1,116	-	-	177	81	96	-	813	-	-	813	-	30	-	-	-	30	-	-	-	2,161
50	-	10	-	333	-	-	252	10	242	-	40	20	-	20	-	125	15	-	101	10	-	-	51	812
51	-	51	20	273	-	10	394	30	364	-	111	10	30	71	-	747	45	-	702	-	-	30	71	1,707
52	-	77	10	979	-	-	280	10	270	-	10	-	-	10	-	461	10	-	461	-	-	10	-	1,837
53	10	10	10	328	-	20	273	-	273	-	25	-	-	25	-	222	10	-	212	-	-	-	-	898
54	-	20	-	803	51	197	91	25	51	15	25	-	10	15	-	35	10	-	25	-	-	-	-	1,222
55	-	-	-	432	36	-	30	10	20	-	10	-	-	10	-	20	-	-	20	-	-	10	-	538
56	-	51	10	503	-	-	133	10	123	-	116	-	10	106	-	72	-	-	62	10	-	-	-	885
57	-	10	-	259	-	36	71	-	71	-	66	20	-	46	-	550	10	-	540	-	-	-	15	1,007
58	-	35	15	374	-	-	576	10	566	-	51	-	-	51	-	697	-	-	687	10	-	-	61	1,809
59	-	35	10	490	-	-	56	-	56	-	15	-	-	15	-	126	-	-	126	-	-	-	354	1,086
60	-	-	15	938	-	15	525	20	505	-	1,555	50	10	1,495	-	-	-	-	-	-	-	-	126	3,174
61	-	-	15	756	-	10	122	20	102	-	3,442	162	30	3,250	-	10	-	-	-	10	-	-	-	4,355
62	-	12	-	871	-	53	62	52	10	-	4,980	12	72	4,896	-	10	-	10	-	-	-	-	54	6,042
63	-	-	-	380	-	21	-	-	-	-	1,929	73	26	1,830	-	135	-	125	10	-	-	-	-	2,465
64	-	-	-	439	-	63	83	10	73	-	3,681	177	10	3,483	11	10	-	10	-	-	-	-	10	4,286